Manuscript Title

Bullying Victimization and Brain Development: A Longitudinal Structural Magnetic Resonance Imaging

Study from Adolescence to Early Adulthood

Running Title

Bullying and Brain Development: An sMRI Study

Authors

Michael Connaughton, PhD.[1]; Orla Mitchell, MSc.[1]; Emer Cullen, MSc.[1]; Michael O'Connor, MSc.[1]; Tobias

Banaschewski, M.D., Ph.D.[2]; Gareth J. Barker, Ph.D.[3]; Arun L.W. Bokde, Ph.D.[4]; Rüdiger Brühl, Ph.D.[5];

Sylvane Desrivières, Ph.D.[6]; Herta Flor, Ph.D.[7,8]; Hugh Garavan, Ph.D.[9]; Penny Gowland, Ph.D.[10]; Antoine

Grigis, Ph.D.[11]; Andreas Heinz, M.D., Ph.D.[12, 13]; Herve Lemaitre, Ph.D[11,14]; Jean-Luc Martinot, M.D.,

Ph.D.[15]; Marie-Laure Paillère Martinot, M.D., Ph.D.[15,16]; Eric Artiges, M.D., Ph.D.[15,17]; Frauke Nees,

Ph.D.[2,18]; Dimitri Papadopoulos Orfanos, Ph.D.[8]; Luise Poustka, M.D.[19]; Michael N. Smolka, M.D.[20]; Sarah

Hohmann, PhD.[2]; Nathalie Holz, PhD.[2]; Nilakshi Vaidya, M.Sc.[21]; Henrik Walter, M.D., Ph.D.[12]; Gunter

Schumann, M.D. [20, 22]; Robert Whelan, Ph.D.[23]; Darren Roddy, MD., PhD. [1]*

Corresponding author

Dr Darren Roddy, Senior Clinical Lecturer in Psychiatry, Royal College of Surgeons, Dublin 2

Darrenroddy@rcsi.com

Affiliations

1.   Department of Psychiatry, Royal College of Surgeons in Ireland, Dublin 2, Ireland.

2.   Department of Child and Adolescent Psychiatry and Psychotherapy, Central Institute of Mental

Health, Medical Faculty Mannheim, Heidelberg University, Square J5, 68159 Mannheim, Germany;

German Center for Mental Health (DZPG), partner site Mannheim-Heidelberg-Ulm.

3.   Department of Neuroimaging, Institute of Psychiatry, Psychology & Neuroscience, King's College

London, United Kingdom.

4.   Discipline of Psychiatry, School of Medicine and Trinity College Institute of Neuroscience, Trinity

College Dublin, Dublin, Ireland.

5.   Physikalisch-Technische Bundesanstalt (PTB), Braunschweig and Berlin, Germany.

6.   Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology &

Neuroscience, King's College London, United Kingdom.

7.   Institute of Cognitive and Clinical Neuroscience, Central Institute of Mental Health, Medical

Faculty Mannheim, Heidelberg University, Square J5, Mannheim, Germany.

8.   Department of Psychology, School of Social Sciences, University of Mannheim, 68131 Mannheim,

Germany.

9.   Departments of Psychiatry and Psychology, University of Vermont, 05405 Burlington, Vermont, USA.

10.   Sir Peter Mansfield Imaging Centre School of Physics and Astronomy, University of Nottingham, University Park, Nottingham, United Kingdom.

11.   NeuroSpin, CEA, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France.

12.   Department of Psychiatry and Psychotherapy CCM, Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Berlin, Germany.

13.   German Center for Mental Health (DZPG), Berlin-Potsdam, Germany.

14.   Institut des Maladies Neurodégénératives, UMR 5293, CNRS, CEA, Université de Bordeaux, 33076 Bordeaux, France.

15.   Institut National de la Santé et de la Recherce Médicale, INSERM U A10 "Trajectoires développementales & psychiatrie", University Paris-Saclay, Ecole Normale Supérieure Paris-Saclay, CNRS; Centre Borelli, Gif-sur-Yvette, France.

16.   AP-HP. Sorbonne Université, Department of Child and Adolescent Psychiatry, Pitié-Salpêtrière Hospital, Paris, France.

17.   Psychiatry Department, EPS Barthélémy Durand, Etampes, France.

18.   Institute of Medical Psychology and Medical Sociology, University Medical Center Schleswig Holstein, Kiel University, Kiel, Germany.

19.   Department of Child and Adolescent Psychiatry, Center for Psychosocial Medicine, University Hospital Heidelberg, Heidelberg, Germany.

20.   Department of Psychiatry and Psychotherapy, Technische Universität Dresden, Dresden, Germany.

21.   Centre for Population Neuroscience and Stratified Medicine (PONS), Department of Psychiatry and Psychotherapy, Charité Universitätsmedizin Berlin, Germany.

22.   Centre for Population Neuroscience and Precision Medicine (PONS), Institute for Science and Technology of Brain-inspired Intelligence (ISTBI), Fudan University, Shanghai, China.

23.   School of Psychology and Global Brain Health Institute, Trinity College Dublin, Ireland.

Data Availability

Code Availability

https://github.com/mconnaug/Bullying_Brain_Development

Abstract

This study investigated associations between bullying victimization and brain development using longitudinal structural MRI data from the IMAGEN cohort (n = 2,094; 1,009 females) across three time points (∼14, ∼19, and ∼22 years). A data-driven analysis revealed that higher bullying victimization was significantly associated with accelerated volumetric growth in subcortical and limbic regions, including the putamen (β = 0.12, 95% CI: 0.10–0.15), amygdala (β = 0.07, 95% CI: 0.05–0.09), hippocampus (β = 0.06, 95% CI: 0.04–0.08), and anterior cingulate cortex (caudal: β = 0.05, 95% CI: 0.03–0.07; rostral: β = 0.06, 95% CI: 0.04–0.08). In contrast, bullying victimization was also significantly associated with reduced volumetric growth in the cerebellum (β = –0.09, 95% CI: –0.11 to –0.07), entorhinal cortex (β = –0.10, 95% CI: –0.13 to –0.07), and insula (β = –0.08, 95% CI: –0.11 to –0.06). Exploratory analyses indicated that females exhibited more pronounced changes in emotional processing regions, while males showed greater changes in motor and sensory areas. Overall, the findings indicate that bullying victimization is associated with widespread structural differences in brain development from adolescence to early adulthood, with sex-specific trajectories.

Introduction

Bullying victimization, characterized by targeted and often chronic peer aggression—whether physical, verbal, or relational—is a common experience during childhood and adolescence, predominantly perpetrated by school peers (1). Persistent exposure to bullying is strongly associated with increased risk for depression, anxiety, suicidality, and impaired cognitive and social functioning (2-4). These adverse outcomes often extend into adulthood, raising concerns that bullying may be associated with disruptions in core neurodevelopmental processes during a critical period of brain maturation (5).

Adolescence is a particularly sensitive period for brain development, marked by extensive biological and psychological transformations (6). This phase involves substantial reorganization of neural circuits supporting executive function, emotion regulation, social cognition, and stress responsivity (7, 8). These changes are especially pronounced in frontolimbic regions (e.g., prefrontal cortex, anterior cingulate cortex), parietotemporal areas (e.g., superior temporal sulcus, temporoparietal junction), and subcortical structures (e.g., amygdala, hippocampus, striatum) (7, 8). While this plasticity supports adaptive development, it may also render the adolescent brain particularly vulnerable to adverse environmental exposures such as bullying victimization (5). Yet, how bullying victimization is associated with changes in brain development over time remains poorly understood.

Neurobiologically, experiences of bullying victimization may influence brain development through multiple interacting pathways (5). It has been linked to stress-induced neuroendocrine reactivity, neuromodulation, and limbic system dysregulation, which may collectively drive widespread structural brain alterations (5). These effects are mediated by multiple interconnected systems, including dysregulation of the hypothalamic-pituitary-adrenal (HPA) axis, heightened inflammatory responses, altered dopaminergic and serotonergic signaling, oxytocin pathway disruption, and imbalances in autonomic nervous system activity (9-13). Together, these systems can lead to sustained cortisol release and other neurochemical changes that result in downstream alterations in neurodevelopment, particularly in circuits involved in emotional, social, and cognitive functioning (5).

While cross-sectional structural MRI studies have offered evidence that bullying-related stress may alter brain structure (14), our understanding of its association with brain development over time remains limited. To date, only two longitudinal structural MRI studies have begun to address this question. Menken et al. (2023), using data from the ABCD study, found that children aged 9–11 exposed to bullying exhibited developmental differences,

including steeper increases in hippocampal and entorhinal cortex volumes, along with accelerated cortical thinning in several frontal and temporal regions (15). Quinlan et al. (2020), using the IMAGEN cohort, reported that adolescents exposed to peer victimization from ages 14 to 19 exhibited steeper declines in left putamen volume, which predicted higher anxiety symptoms in early adulthood (16). However, their analysis focused on nine bilateral regions of interest, potentially overlooking broader patterns of neurodevelopmental change. While both studies were pioneering in establishing early longitudinal links between bullying victimization and brain development, each was limited to two imaging time points, restricting the ability to capture non-linear, region-specific developmental trajectories.

Indeed, brain development across adolescence is rarely linear (7, 8, 17). Cortical and subcortical structures often follow quadratic growth patterns, including periods of acceleration, deceleration, and stabilization (7, 8, 17). Two-time-point designs constrain our ability to capture such inflection points and may obscure meaningful developmental variability. In contrast, three-time-point longitudinal designs allow for more accurate mapping of non-linear, region-specific developmental trajectories, which is particularly important as the field moves toward the development of normative brain growth charts (18). Such charts may help identify neurodevelopmental shifts that contribute to diverging mental health outcomes, particularly in adolescents exposed to environmental stressors like bullying victimization.

To address this gap, this study aims to measure the association between bullying victimization on cortical and subcortical brain development using structural MRI data collected at three time points: ages 14, 19, and 22. Unlike prior work limited to two time points and a small set of predefined regions, our study leverages three MRI time points and a whole-brain approach to capture more nuanced, region-specific trajectories. We hypothesize that increased bullying victimization is associated with widespread variation in brain development, particularly in regions involved in emotion regulation and social processing. Specifically, we expect to observe distinct patterns of volumetric change, such as decreased cortical and increased subcortical volume change across adolescence and early adulthood. In addition, we will conduct exploratory analyses to examine potential sex differences in bullying-related brain development. Emerging evidence suggests that males and females differ in both neurobiological stress responses and developmental trajectories (19, 20). By examining sex differences, we aim to clarify whether the neurodevelopmental associations of bullying victimization vary by sex, contributing to a more nuanced understanding of brain maturation.

Materials and Methods

Participants

This study used data from the IMAGEN project, a European multicenter research initiative investigating how various factors influence brain development and mental health in adolescents. For a comprehensive description of the project's methodology, refer to Schumann et al. (21). Participants were assessed at eight sites across England, Ireland, France, and Germany. Initial data were collected at age 14, with follow-up assessments conducted at ages 16, 19, and 22. This study specifically analyzes data from ages 14, 19, and 22; data from age 16 were excluded, as MRI scans were not conducted at that time point. A detailed overview of recruitment procedures and inclusion/exclusion criteria is presented in eTable 1, with further details available in Schumann et al. (21).

Written informed consent was obtained from participants and their parent/guardian prior to enrollment. The IMAGEN study was approved by ethics committees at each site, including King's College London, University of Nottingham, Trinity College Dublin, University of Heidelberg, Technische Universität Dresden, Commissariat à l'Energie Atomique et aux Energies Alternatives, and University Medical Center.

Bullying victimization

Bullying victimization was assessed using items adapted from the revised Olweus Bully/Victim Questionnaire (22) at all three study timepoints (~14, ~19, and ~22 years). Participants responded to four items evaluating the frequency of victimization over the past six months, each rated on a 5-point Likert scale:

1. "I was bullied at school (a student/peer said or did nasty or unpleasant things to me)."
2. "I was called mean names, made fun of, or teased in a hurtful way by a student/peer."
3. "A student/peer left me out of things on purpose, excluded me from their group of friends, or completely ignored me."
4. "I was hit, kicked, pushed or shoved around, or locked indoors by a student/peer."

Response options ranged from 1 ("never") to 5 ("three or more times a week"). Scores from the four items were summed and standardized (z-scores) to generate bullying victimization scores at ages 14, 19, and 22, with higher scores indicating increased frequency of self-reported bullying victimization. The composite Olweus Bully/Victim Questionnaire score demonstrated excellent internal consistency in the current sample (Cronbach's $\alpha = 0.89$).

MRI Acquisition and Protocol

Structural magnetic resonance imaging (MRI) data were acquired across eight IMAGEN sites in Europe, all using 3T MRI systems (Siemens: 5 sites; Philips: 2 sites; General Electric: 1 site). High-resolution anatomical scans were obtained using a 3D T1-weighted magnetization-prepared rapid gradient echo (MPRAGE) sequence, aligned with the Alzheimer's Disease Neuroimaging Initiative (ADNI) protocol. Full details of the IMAGEN MRI acquisition and quality control procedures, including scanner standardization protocols, are available in Schumann et al. (21) and subsequent publications (23, 24), and are accessible via the project's Standard Operating Procedures (https://imagen-europe.org/). In brief, T1-weighted anatomical images were acquired using a 3D MPRAGE sequence (voxel size = 1.1 × 1.1 × 1.1 mm³; TR = 2300 ms; TE = 2.9 ms). Full acquisition parameters are provided in eTable 2. All structural MRI scans underwent IMAGEN's centralized quality control procedure, which included manual inspection for artifacts, data quality, and head motion (see Supplemental Material). Following these procedures, 50 scans were excluded (24 at timepoint 1, 17 at timepoint 2, 9 at timepoint 3), resulting in a final dataset of 4,555 structural MRI scans.

All MRI images were processed using FreeSurfer's *recon-all* pipeline (version 5.3.0) for full cortical reconstruction and subcortical segmentation (25, 26). This automated process included skull stripping, intensity normalization, Talairach transformation, surface reconstruction, topology correction, and anatomical labeling. Brain parcellation was performed using the Desikan-Killiany-Tourville (DKT) atlas, producing 88 cortical and subcortical regions of interest, from which volume metrics were extracted, along with total gray matter volume (27).

Prior to statistical analyses, outliers were identified as values exceeding ±3 standard deviations from the mean (28). These were visually inspected across timepoints to assess consistency. In line with best practices, only values deemed implausible or inconsistent with developmental trajectories were removed to reduce the influence of artefactual data (29).

Statistical Analysis

Mixed-effects modeling was used to investigate associations between bullying victimization and longitudinal brain development from adolescence to early adulthood. Analyses were conducted in R (version 4.1.1) using the *lme4* package (version 1.1-27.1) (30). Model fitting followed a structured, top-down selection procedure comprising three sequential

stages: (1) determining the optimal developmental trajectory for each region of interest (ROI), (2) specifying the random effects structure, and (3) evaluating the fixed effects of bullying victimization. A complete summary of all models tested is provided in eTable 3. Model selection was guided by a combination of the Akaike Information Criterion (AIC) and likelihood ratio tests (LRT). A model was considered to show improved fit if it demonstrated a reduction in AIC greater than 10 and an LRT p-value < 0.05, in line with established criteria for mixed model comparison (31-33). All mixed-effects models were run with mean-centered continuous variables.

Step 1: Determining the Optimal Brain Region Developmental Trajectory

To characterize normative patterns of brain development, we first evaluated whether a linear or quadratic representation of time (indexed as months since baseline scan) provided the best fit for each brain region of interest (ROI). This analysis was performed separately for all 88 ROI volumes, including cortical and subcortical structures. Consistent with top-down model selection procedures, both models were initially specified with covariates only— sex, pubertal development status (PDS), socioeconomic status (SES), stressful life events (SLE), and intracranial volume (ICV)—excluding any bullying victimization terms at this stage. See supplemental material for a detailed description of covariates. All models included random intercepts and slopes for participants and random intercepts for scan site. In line with best practices for multisite neuroimaging studies, scan site was modeled as a random effect to account for variance across imaging locations (34). A quadratic time term ($time^2$) was retained only when its inclusion significantly improved model fit, allowing for modeling of nonlinear neurodevelopmental trajectories commonly observed during adolescence (18).

Step 2: Specifying the Random Effects Structure

Following identification of the optimal developmental trajectory, we next determined the most appropriate random effects structure. For both the linear and quadratic models, we compared random intercepts-only models (R1) with models that also included random slopes for time (R2). Scan site was consistently modeled as a random intercept to adjust for multisite acquisition variability. These comparisons were evaluated using the same AIC and LRT thresholds as in Step 1. The selected structure allowed for adequate modeling of within-subject variation in brain development.

Step 3: Testing Fixed Effects of Bullying Victimization

After finalizing the model structure, we evaluated the fixed effects of bullying
victimization using a series of nested models. The null model (F0) included all covariates and
time terms but excluded any bullying-related predictors. The simple effects model (F1) added
the main effect of bullying victimization frequency, while the interaction model (F2) further
included the bullying victimization score × time term to examine whether the association
between bullying and brain development varied over time. All models were estimated using
maximum likelihood (ML) for model fit comparison, and final parameter estimates were
derived using restricted maximum likelihood (REML), which provides unbiased estimates of
variance components by accounting for the loss of degrees of freedom when estimating fixed
effects (30).

An unstructured variance-covariance matrix was used for random effects, permitting
unrestricted estimation of correlations between intercepts and slopes and thus accommodating
individual heterogeneity in developmental trajectories (35). To control for multiple
comparisons, we applied a false discovery rate (FDR) correction at q < 0.05 using the *stats*
package in R (version 4.1.1) (36, 37). This correction was applied to all p-values associated
with the bullying victimization score main effects and bullying victimization score × time
interactions in the optimally fitted models. Only results that survived FDR correction are
reported.

Sex-Specific Associations Between Bullying Victimization and Brain Development

Exploratory sex-specific analyses were conducted using the optimal models identified
during the model selection procedure described above. To test for sex-dependent effects,
additional interaction terms for bullying victimization score × sex and bullying victimization
score × time × sex were added to these models (F3). As in previous steps, model fit was
evaluated using AIC and LRT, and FDR correction was applied to p-values associated with
both interaction terms. Only associations that remained statistically significant after correction
are reported. These analyses aimed to explore potential sex differences in neurodevelopmental
responses to bullying victimization during adolescence.

Results

Descriptive Statistics

299   Full demographic and clinical characteristics are presented in Table 1. Of the 4,555
300   available MRI scans, 43 were excluded due to missing bullying questionnaire data (Time 1 =

11, Time 2 = 30, Time 3 = 2). The final analytic sample comprised 4,512 scans from 2,094 participants (females: 2,171; males: 2,341), spanning an age range of 13.23 to 25.11 years across three time points (see eFigure 1). Bullying victimization showed modest but statistically significant rank-order stability across waves (T1–T2: ρ = 0.24, p < .000001; T2–T3: ρ = 0.20, p < .000001; T1–T3: ρ = 0.19, p < .000001). Attrition analyses were conducted using baseline data and are reported in the Supplementary Material.

Table 1: Demographic and Site Characteristics Across Three Time Points.

Variable Name	Time 1	Time 2	Time 3
N	2052	1328	1132
Sex (female %)	1009 (49.17%)	632 (47.59%)	530 (46.81%)
Age	14.44 (0.39)	19.01 (0.78)	22.56 (0.65)
OB/VQ score (range)	4-20	0-16	0-16
SES (mean)	0.71 (1.09)	-	-
Pubertal Status (mean)	3.60 (0.71)	-	-
Ethnicity (White %)	89.4%	-	-
Number of Scans at IMAGEN Site			
London	251	168	153
Northampton	343	211	169
Dublin	192	122	116
Berlin	252	118	155
Hamburg	261	181	153
Mannheim	245	155	140
Paris	253	204	124
Dresden	255	169	122

Legend: N = total number of participants assessed at each respective time point. The Olweus Bully/Victim Questionnaire (OB/VQ) is used to measure bullying victimization and bullying experiences. Pubertal status was assessed via self-report using the Pubertal Development Scale (PDS). Socioeconomic status (SES) was indexed using the family stresses subsection of the Development and Well-Being Assessment (DAWBA). Data collection for Time 1 occurred between 2008–2010, for Time 2 between 2013–2015, and for Time 3 between 2016–2019.

Bullying Victimization and Brain Development

Model selection fit statistics are presented in the supplemental material (eTables 1–5), with the final models derived from the top-down selection procedure shown in eTable 6. Full model results are provided in eTables 7–10 and eFigure 2.

Significant associations between bullying victimization scores and brain volume development were identified in 30 regions, indicating consistent structural differences across adolescence and early adulthood (Figures 1 and 2). These included widespread volume reductions in cortical areas such as the orbitofrontal cortex, superior and rostral middle frontal gyri, precuneus, precentral and postcentral gyri, insula, entorhinal cortex, temporal pole, and regions within the parietal, occipital, and cerebellar cortices. In contrast, increased volumes were observed in several limbic and subcortical structures, including the amygdala, hippocampus, parahippocampal gyrus, putamen, caudate, nucleus accumbens, and frontal pole. Additional volume reductions were also found in the thalamus, pallidum, and ventral diencephalon.

Significant bullying victimization-by-time interactions were identified in approximately 16 regions, indicating that bullying victimization frequency was associated with altered neurodevelopmental trajectories (Figures 1 and 2). Higher levels of victimization frequency were linked to accelerated volumetric growth in regions such as the amygdala, hippocampus, putamen, anterior cingulate cortex, and banks of the superior temporal sulcus. Conversely, reduced volumetric growth was observed in the insula, entorhinal cortex, cerebellum, and visual and parietal cortices (e.g., cuneus, lingual gyrus, superior parietal lobule). These findings suggest that bullying may influence not only regional brain structure but also the pace of brain maturation from adolescence into early adulthood.

Sex Differences in the Associations of Bullying Victimization on Brain Development

Exploratory analyses revealed significant three-way interactions between bullying victimization, sex, and time on brain volume development (eTables 11–12). These findings indicate that the association between bullying victimization and age-related brain volume trajectories varies by sex (Figure 3).

Higher bullying victimization frequency over time was linked to greater volume increases in females compared to males in several subcortical and limbic regions, including the bilateral parahippocampal gyrus, right caudate, left putamen, and brainstem. Conversely, males

342 showed relatively greater volume increases in regions such as the left supramarginal gyrus,
343 right pars orbitalis, right temporal pole, left ventral diencephalon, and left cerebellar cortex.

344   These results point to sex-specific patterns in the association between bullying
345 victimization and brain development, with distinct regional trajectories emerging across
346 adolescence and early adulthood.

Discussion

This three-time-point longitudinal neuroimaging study of bully victimization examined
the association between bullying victimization and brain development across adolescence and
early adulthood. The principal findings are: (1) bullying victimization frequency is
significantly associated with altered development in a wide range of cortical and subcortical
brain structures, and (2) these alterations show notable sex-specific patterns.

The novelty of this study lies in capturing widespread non-linear developmental brain
changes not observed in previous studies, highlighting that the neurodevelopmental
associations of bullying victimization may be more widespread than previously known (15,
16). These findings are robust across variables such as sex, age, MRI scanner site,
socioeconomic status, pubertal status, and other negative life events, suggesting that the
observed brain differences may be uniquely attributed to bullying victimization.

Association Between Bullying Victimization and Brain Development
Habit formation, emotional salience and stress regulation: striatal and subcortical systems

Subcortical structures appear to be particularly sensitive to bullying victimization, with
significant volumetric increases observed in the structures of the basal ganglia, including the
caudate, putamen, and nucleus accumbens. These regions are integral to motor control,
emotional regulation, and reward processing (38). The observed differences in the volumes of
the caudate and putamen corroborate previous findings (16). Meanwhile, the discovery of
increased volumes in the nucleus accumbens and pallidum represents novel contributions to
the field.

Prolonged bullying may trigger neuroplastic adaptations as the brain attempts to cope.
Enlarged dorsal striatal structures (caudate and putamen), involved in automatic responses and
attention, may underlie increased striatum-dependent ("habit") learning in bullied individuals
(5). These individuals often rely on coping behaviors shaped by past threats, which may be

maladaptive in safe contexts, contributing to social distress and difficulty adapting to new
environments. Victimized adolescents also show a shift toward striatal-dependent memory
processing, associated with cognitive inflexibility and anxiety (39). This type of memory
primarily involves the ventral striatum (nucleus accumbens), which plays a central role in
negative emotional processing and is linked to internalizing and externalizing symptoms in
bullied youth (40). Thus, its enlargement may reflect a heightened bias toward emotionally
salient memory encoding. Though less studied, the pallidum also shows volumetric disruptions
in this context and has been associated with depression, anxiety, and OCD (41-43). As part of
the cortico-striato-thalamo-cortical (CSTC) circuitry, the pallidum is critical for emotional
regulation, and its enlargement may reflect CSTC disruption, contributing to stress sensitivity
and emotional dysregulation (41, 44).

A novel finding, that contradicts our hypothesis was the association between increased
bullying exposure and reduced ventral diencephalon volume. This region, which includes the
hypothalamus, is essential for regulating the neuroendocrine stress response (45). Reduced
volume here may impair hormonal regulation, compromising the body's ability to manage
stress effectively (46, 47), and increasing susceptibility to anxiety, mood disorders, and other
stress-related conditions (48).

Emotional reactivity and memory biases: limbic regions

Significant differences were also observed in the limbic system, particularly the
amygdala and hippocampus. Enlarged hippocampal volume aligns with prior findings (15),
while increased amygdala volume was not previously reported, possibly due to methodological
differences such as the binary classification of victimization status in earlier studies. It is
plausible that amygdala enlargement reflects heightened emotional reactivity to chronic stress
(49). Victimized individuals often show increased neural responses to emotional stimuli,
suggesting enhanced stress sensitivity (50, 51), which may contribute to elevated risk for
anxiety, depression, and related disorders (52, 53).

Hippocampal enlargement may reflect neuroplastic adaptations to prolonged stress,
such as increased neurogenesis or dendritic branching in response to emotionally salient
memories (45). These neuroplastic changes may underpin the negative emotional memory
biases and increased false memory recall observed in individuals who have experienced
bullying victimization (46). Specifically, violent and aggressive false memories have been
shown to be positively associated with bullying victimization (54). This heightened memory
processing demand could help explain the observed structural increases. Despite these findings,

mixed results regarding hippocampal volume call for further research to clarify its role and
long-term implications in bullying-related neurodevelopment.
410

Executive control, emotional regulation, and social cognition: Frontal, temporal, parietal,
and occipital lobes
Alterations were also noted in various cortical areas across the frontal, temporal,
parietal, and occipital lobes. In the frontal lobe, changes were observed in regions such as the
medial orbitofrontal cortex, superior frontal gyrus, and frontal pole. Previous research has
linked functional and structural alterations in these areas to bullying victimization, suggesting
that victims may have difficulties in regulating emotions and making decisions under stress
(14). Our findings support these previous studies, reinforcing the idea that bullying impacts
critical areas involved in executive functioning (5).
The temporal regions, including the superior temporal gyrus and parahippocampal
gyrus, showed changes that could affect memory and emotional association. Studies have
indicated that victims of bullying often exhibit altered temporal lobe structures, contributing to
difficulties in processing and recalling emotionally charged memories (5). Our results are
consistent with these findings, suggesting a common pathway through which bullying affects
memory and emotional processing (5).
Significant volume reductions were found in the insula, which are novel findings not
previously reported in the literature. Prior research has highlighted significant differences in
insula activity in the context of bullying victimization, but structural changes had not been
documented until now (55). The insula is a deep cortical structure critical for emotional
processing and interoceptive awareness (5). Alterations in this structure have been linked to
heightened sensitivity to emotional stimuli and social rejection, a common feature of bullying
victimization (5).
433

Environmental interpretation via perceptual, predictive, and integrative systems: parietal,
occipital and cerebellum regions
Alterations in the parietal and occipital lobes, such as the precuneus and cuneus, may
influence how victims process and respond to visual and spatial cues (56, 57). This can impact
their social interactions and stress responses (5). Previous studies have found that these regions,
when affected by bullying, can alter how individuals perceive and react to their environment,
potentially leading to social withdrawal and heightened anxiety (5). Our findings align with

these studies, suggesting that visual and spatial processing deficits may contribute to the social

challenges faced by bullied individuals (5).

Our study revealed a novel link between bullying victimization frequency and reduced

cerebellar volume, suggesting that prolonged social stress may impair cerebellar development.

While traditionally associated with motor coordination, the cerebellum also plays a key role in

social and cognitive processes essential for social interaction (58). Poor motor skills, governed

by cerebellar function, are strong predictors of being bullied (59), as individuals with less

refined motor abilities may struggle with social coordination and integration. Beyond motor

control, the cerebellum contributes to higher-order functions such as prediction, error-based

learning, and emotional recognition (58). Disruptions in these mechanisms may impair

interpretation of social cues, leading to negative emotional biases common in bullied

individuals, who often misread neutral or ambiguous interactions as hostile, thus increasing

their vulnerability to interpersonal difficulties and psychopathology (58, 60).

In summary, our findings indicate that bullying victimization is linked to widespread

cortical alterations affecting cognitive and emotional functioning. These structural changes

may contribute to the psychological and behavioral difficulties seen in victims and support a

neurobiological cycle in which bullying exacerbates vulnerability to further victimization and

long-term mental health risks.

<u>Sex-Specific Differences in Brain Volume Development</u>

This study identified distinct sex-specific patterns of brain volume changes associated

with bullying victimization frequency, suggesting that males and females may exhibit different

neurodevelopmental adaptations to similar social stressors. Females demonstrated relatively

greater volume increases in subcortical and limbic regions—specifically the bilateral

parahippocampal cortex, right caudate, left putamen, and brainstem—while males showed

increased volume predominantly in a combination of cortical and subcortical regions, including

the left supramarginal gyrus, right pars orbitalis, right temporal pole, left ventral diencephalon,

and left cerebellar cortex.

These divergent patterns may reflect differences in the nature of bullying typically

experienced by each sex. Females are more commonly exposed to relational bullying—such as

social exclusion, manipulation, and rumour-spreading (61)—which preferentially engages

limbic and paralimbic networks involved in emotional memory and social cognition (62-65).

In contrast, males more frequently encounter physical and overt forms of bullying, including

verbal aggression and threats, which may more strongly recruit sensorimotor and salience-detection circuits, consistent with volumetric changes observed in regions such as the cerebellum and ventral diencephalon (66-68) .

Together, they underscore the importance of considering sex as a moderating factor in the neural associations of bullying victimization. Although these insights are promising, further research is needed to elucidate the underlying mechanisms and their long-term neurodevelopmental relevance.

Neurobiological Mechanisms Linked to Brain Volume Alterations in Bullying Victimization

The region-specific brain volume changes associated with bullying victimization likely reflect broader neurobiological mechanisms triggered by chronic social stress. One key system implicated is the hypothalamic–pituitary–adrenal (HPA) axis, which becomes activated in response to psychological stress and elevates glucocorticoid levels—a pattern observed in bullied children (69-72). These glucocorticoids disproportionately affect multiple brain regions, including the prefrontal cortex (PFC) and hippocampus (73). Sustained cortisol exposure can impair synaptic plasticity, promote dendritic atrophy, and even lead to neurodegeneration, which may underlie the reduced PFC volume observed in bullied youth (74). Interestingly, although chronic stress is typically associated with hippocampal atrophy (75), this study and previous research found that bullying victimization is associated with increased hippocampal volume development during early life (15). This may reflect an early adaptive response, where initial stress exposure temporarily enhances dendritic complexity or glial activity before volume declines with prolonged stress (76, 77). Similarly, increased volume in the ventral diencephalon, including the hypothalamus, may reflect stress-induced plasticity within neuroendocrine circuits (78). Chronic stress enhances corticotropin-releasing hormone (CRH) production, increases excitatory input, and reduces inhibitory control in the paraventricular nucleus, sustaining HPA axis activation (78). These adaptations may support short-term homeostasis under prolonged social threat and contribute to the observed volumetric increase.

Beyond the HPA axis, bullying may also activate the locus coeruleus–norepinephrine (LC-NE) system, a brainstem arousal network responsive to threat (79). Sustained norepinephrine release is associated with increased amygdala excitability and synaptic activity (80), which over time may drive structural plasticity—such as dendritic hypertrophy (81)—

potentially contributing to the amygdala volume increases observed in bullied adolescents. A

concurrent reduction in medial PFC volume—essential for top-down regulation of the LC-NE

system—could reflect a shift toward reflexive, emotion-driven processing (82), potentially

explaining heightened threat sensitivity in bullying victims (83).

Additionally, volumetric increases in the caudate, putamen, and nucleus accumbens

may result from stress-induced changes in dopaminergic signalling within mesostriatal and

mesolimbic pathways (84). This dopaminergic hyperactivity has been linked to structural

plasticity, including dendritic growth and increased spine density in medium spiny neurons—

cellular changes that may underlie the volumetric expansion of striatal regions (85). These

changes may reflect an adaptive response to repeated exposure to socially salient stressors,

enhancing the salience of threat cues and reinforcing habitual coping behaviors (86). Over time,

this plasticity could contribute to inflexible behavioral patterns and heightened sensitivity to

social stress observed among individuals exposed to bully victimization (39)

While these findings point to region-specific developmental adaptations, the

neurobiological mechanisms underlying altered brain development in bullied youth are

inherently complex and likely involve the dynamic interplay of hormonal, neuroimmune, and

neurotransmitter systems. Continued investigation into these pathways is essential, as

developing a neurobiological framework is critical for understanding the long-term impact of

bullying victimization.

<u>Limitations</u>

This study has several limitations that warrant consideration. Bullying victimization in

this study was measured from age 14 onward, potentially missing developmental changes from

earlier experiences. As a result, early influences on brain development may be

underrepresented, including critical periods affected by prior victimization. Future research

should incorporate earlier and broader longitudinal data to better capture these effects. Another

limitation of the current study is that we were unable to distinguish between school-based and

out-of-school bullying experiences, as participant educational status was not consistently

recorded. Future research should explore how the context of bullying may differentially impact

developmental trajectories. While the questionnaire included items that could reflect

cyberbullying, it was not specifically measured, which may have led to an underestimation of

victimization. Future studies should use dedicated cyberbullying measures, such as those by

Aricak et al. (87), to better assess its impact on brain development. Ethnicity was not included

as a covariate due to imbalanced representation across sites. While scan site served as a proxy

- with 89.4% of participants aligning with the dominant local ethnic group - this does not fully

capture ethnic variation. More diverse and balanced samples are needed to examine ethnicity's

role in bully victimization and neurodevelopment. As IMAGEN is a community-based cohort,

clinical diagnoses of pre-existing mental health conditions were not collected. This makes it

difficult to determine whether observed brain changes are driven by bullying victimization

alone or influenced by earlier psychiatric vulnerabilities. With only three timepoints per

participant, higher-order nonlinear models (e.g., cubic, biquadratic) were not feasible due to

the risk of overfitting and model instability (88, 89). Linear and quadratic trajectories, however,

are well-established in adolescent brain development (17), making them a statistically

appropriate and theoretically grounded choice. The sparse longitudinal sampling also limited

our ability to reliably test whether within-person changes in bullying victimization were

paralleled by changes in cortical developmental trajectories, which will be an important focus

for future studies using denser longitudinal designs better suited to capturing dynamic within-

person victimization trajectories.

<u>Conclusion</u>

This longitudinal MRI study demonstrates that bullying victimization is associated with

widespread structural differences in brain development from adolescence to early adulthood,

with distinct sex-specific patterns. These findings extend prior work by highlighting

neurodevelopmental alterations across circuits involved in stress, emotional learning, and

social cognition. While causality cannot be inferred, the results underscore bullying

victimization as a salient social experience linked to long-term variation in brain maturation

and provide a foundation for future neuroimaging research to explore the underlying

mechanisms driving mental health vulnerability.

royalities from Hogrefe, Kohlhammer, CIP Medien, Oxford University Press; the present work

is unrelated to these relationships. Dr Barker has received honoraria from General Electric

Healthcare for teaching on scanner programming courses. Dr Poustka served in an advisory or

consultancy role for Roche and Viforpharm and received speaker's fee by Shire. She received

royalties from Hogrefe, Kohlhammer and Schattauer. The present work is unrelated to the

above grants and relationships. The other authors report no biomedical financial interests or

potential conflicts of interest.

Acknowledgments

This work received support from the following sources: the European Union-funded

FP6 Integrated Project IMAGEN (Reinforcement-related behaviour in normal brain function

and psychopathology) (LSHM-CT- 2007-037286), the Horizon 2020 funded ERC Advanced

Grant 'STRATIFY' (Brain network based stratification of reinforcement-related disorders)

(695313), Horizon Europe 'environMENTAL', grant no: 101057429, UK Research and

Innovation (UKRI) Horizon Europe funding guarantee (10041392 and 10038599), Human

Brain Project (HBP SGA 2, 785907, and HBP SGA 3, 945539), the Chinese government via

the Ministry of Science and Technology (MOST). The German Center for Mental Health

(DZPG), the Bundesministerium für Bildung und Forschung (BMBF grants 01GS08152;

01EV0711; Forschungsnetz AERIAL 01EE1406A, 01EE1406B; Forschungsnetz IMAC-Mind

01GL1745B), the Deutsche Forschungsgemeinschaft (DFG project numbers 458317126

[COPE], 186318919 [FOR 1617], 178833530 [SFB 940], 386691645 [NE 1383/14-1],

402170461 [TRR 265], 454245598 [IRTG 2773]), the Medical Research Foundation and

Medical Research Council (grants MR/R00465X/1 and MR/S020306/1), the National

Institutes of Health (NIH) funded ENIGMA-grants 5U54EB020403-05, 1R56AG058854-01

and U54 EB020403 as well as NIH R01DA049238, the National Institutes of Health, Science

Foundation Ireland (16/ERCD/3797). NSFC grant 82150710554. Further support was provided

by grants from: - the ANR (ANR-12-SAMA-0004, AAPG2019 - GeBra), the Eranet Neuron

(AF12-NEUR0008-01 - WM2NA; and ANR-18-NEUR00002-01 - ADORe), the Fondation de

France (00081242), the Fondation pour la Recherche Médicale (DPA20140629802), the

Mission Interministérielle de Lutte-contre-les-Drogues-et-les-Conduites-Addictives

(MILDECA), the Assistance-Publique-Hôpitaux-de-Paris and INSERM (interface grant), Paris

Sud University IDEX 2012, the Fondation de l'Avenir (grant AP-RM-17-013), the Fédération

pour la Recherche sur le Cerveau.

 Figure Titles and Legends

Figure 1: Magnitude of Effect Sizes for Bullying victimization and Bullying victimization-by-

Age on Cortical Brain Regions.

Figure 1 legend: This figure presents standardized effect sizes (β) for the associations between

bullying victimization score and the bullying victimization score-by-time interaction on

cortical brain volumes. The top panel displays the effect sizes for the main effect of bullying

victimization scores, while the bottom panel illustrates the effect sizes for the interaction

between bullying victimization and time. The color scale represents standardized β values, with

warmer shades indicating stronger positive or negative effects. These visualizations highlight

cortical regions where bullying victimization and its interaction with time are significantly

linked to changes in brain volume development.

Figure 2: Magnitude of Effect Sizes for Bullying victimization and Bullying victimization-by-

Age on Subcortical Brain Regions.

Figure 2 legend: This figure displays standardized effect sizes (β) for the associations between
bullying victimization score and bullying victimization score-by-time on subcortical brain
volumes. The left panel illustrates the effect sizes for the main effect of bullying victimization
scores, while the right panel shows the effect sizes for the interaction between bullying
victimization and time. The color scale reflects standardized β values, with warmer colors
indicating larger effect sizes in either the positive or negative direction. These visualizations
highlight subcortical regions where bullying victimization and its interaction with time are
significantly associated with volume development.
633
Figure 3: Sex Differences in the Impact of Peer Victimization on Brain Development.

Figure 3 legend: The analysis had peer victimization as a sum score, but for visualization
purposes, these scores were grouped into four categories. "Male Low" represents males in the
lowest quartile of bullying victim scores, shown in red. "Male High" represents males in the
highest quartile of bullying victim scores, shown in blue. "Female Low" represents females in
the lowest quartile of bullying victim scores, shown in dark green. "Female High" represents
females in the highest quartile of bullying victim scores, shown in purple. The shaded region
around each line represents the 95% confidence interval for the predicted values.

References

1.   Olweus D, Pellegrini AD. Bullying at school. Aggressive behavior. 1996;22(2):150-4.
2.   Arseneault L. Annual Research Review: The persistent and pervasive impact of being
bullied in childhood and adolescence: implications for policy and practice. Journal of Child
Psychology and Psychiatry. 2018;59(4):405-21.
3.   Copeland WE, Wolke D, Angold A, Costello EJ. Adult psychiatric outcomes of
bullying and being bullied by peers in childhood and adolescence. JAMA psychiatry.
2013;70(4):419-26.
4.   Lereya ST, Copeland WE, Costello EJ, Wolke D. Adult mental health consequences of
peer bullying and maltreatment in childhood: two cohorts in two countries. Lancet Psychiatry.
2015;2(6):524-31.
5.   Palamarchuk IS, Vaillancourt T. Integrative brain dynamics in childhood bullying
victimization: cognitive and emotional convergence associated with stress psychopathology.
Frontiers in integrative neuroscience. 2022;16:782154.
6.   Spear LP. Neurodevelopment during adolescence. Neurodevelopmental mechanisms in
psychopathology. 2003:62-83.
7.   Tamnes CK, Herting MM, Goddings AL, Meuwese R, Blakemore SJ, Dahl RE, et al.
Development of the Cerebral Cortex across Adolescence: A Multisample Study of Inter-
Related Longitudinal Changes in Cortical Volume, Surface Area, and Thickness. J Neurosci.
2017;37(12):3402-12.
8.   Mills KL, Goddings AL, Herting MM, Meuwese R, Blakemore SJ, Crone EA, et al.
Structural brain development between childhood and adulthood: Convergence across four
longitudinal samples. Neuroimage. 2016;141:273-81.
9.   Danese A, J Lewis S. Psychoneuroimmunology of Early-Life Stress: The Hidden
Wounds of Childhood Trauma? Neuropsychopharmacology. 2017;42(1):99-114.
10.   Vaillancourt T, Hymel S, McDougall P. The Biological Underpinnings of Peer
Victimization: Understanding Why and How the Effects of Bullying Can Last a Lifetime.
Theory Into Practice. 2013;52(4):241-8.
11.   Hostinar CE, Gunnar MR. Social support can buffer against stress and shape brain
activity. AJOB neuroscience. 2015;6(3):34-42.
12.   McLaughlin KA, Sheridan MA, Lambert HK. Childhood adversity and neural
development: deprivation and threat as distinct dimensions of early experience. Neurosci
Biobehav Rev. 2014;47:578-91.
13.   Tottenham N, Galván A. Stress and the adolescent brain: Amygdala-prefrontal cortex
circuitry and ventral striatum as developmental targets. Neurosci Biobehav Rev. 2016;70:217-
27.
14.   Ke T, De Simoni S, Barker E, Smith P. The association between peer-victimisation and
structural and functional brain outcomes: A systematic review. JCPP Advances.
2022;2(2):e12081.
15.   Menken MS, Rodriguez Rivera PJ, Isaiah A, Ernst T, Cloak CC, Chang L. Longitudinal
alterations in brain morphometry mediated the effects of bullying victimization on cognitive
development in preadolescents. Dev Cogn Neurosci. 2023;61:101247.
16.   Quinlan EB, Barker ED, Luo Q, Banaschewski T, Bokde AL, Bromberg U, et al. Peer
victimization and its impact on adolescent brain development and psychopathology. Molecular
Psychiatry. 2020;25(11):3066-76.
17.   Vijayakumar N, Allen NB, Youssef G, Dennison M, Yücel M, Simmons JG, et al. Brain
development during adolescence: A mixed-longitudinal investigation of cortical thickness,
surface area, and volume. Hum Brain Mapp. 2016;37(6):2027-38.

18.   Bethlehem RAI, Seidlitz J, White SR, Vogel JW, Anderson KM, Adamson C, et al. Brain charts for the human lifespan. Nature. 2022;604(7906):525-33.

19.   Cservenka A, Stroup ML, Etkin A, Nagel BJ. The effects of age, sex, and hormones on emotional conflict-related brain response during adolescence. Brain and cognition. 2015;99:135-50.

20.   Whittle S, Yücel M, Yap MB, Allen NB. Sex differences in the neural correlates of emotion: evidence from neuroimaging. Biological psychology. 2011;87(3):319-33.

21.   Schumann G, Loth E, Banaschewski T, Barbot A, Barker G, Büchel C, et al. The IMAGEN study: reinforcement-related behaviour in normal brain function and psychopathology. Molecular psychiatry. 2010;15(12):1128-39.

22.   Olweus D. Revised Olweus bully/victim questionnaire. Journal of Psychopathology and Behavioral Assessment. 1996.

23.   Burt KB, Whelan R, Conrod PJ, Banaschewski T, Barker GJ, Bokde ALW, et al. Structural brain correlates of adolescent resilience. Journal of Child Psychology and Psychiatry. 2016;57(11):1287-96.

24.   Sibilia F, Jost-Mousseau C, Banaschewski T, Barker GJ, Büchel C, Desrivières S, et al. The relationship between negative life events and cortical structural connectivity in adolescents. IBRO Neuroscience Reports. 2024;16:201-10.

25.   Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C, et al. Whole brain segmentation: automated labeling of neuroanatomical structures in the human brain. Neuron. 2002;33(3):341-55.

26.   Fischl B, van der Kouwe A, Destrieux C, Halgren E, Ségonne F, Salat DH, et al. Automatically parcellating the human cerebral cortex. Cereb Cortex. 2004;14(1):11-22.

27.   Desikan RS, Ségonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, et al. An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. Neuroimage. 2006;31(3):968-80.

28.   Iglewicz B, Hoaglin DC. Volume 16: how to detect and handle outliers: Quality Press; 1993.

29.   Massara P, Asrar A, Bourdon C, Ngari M, Keown-Stoneman CDG, Maguire JL, et al. New approaches and technical considerations in detecting outlier measurements and trajectories in longitudinal children growth data. BMC Med Res Methodol. 2023;23(1):232.

30.   Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using lme4. Journal of Statistical Software. 2015;67(1):1 - 48.

31.   Bohon C, Welch H. Quadratic relations of BMI with depression and brain volume in children: Analysis of data from the ABCD study. Journal of psychiatric research. 2021;136:421-7.

32.   Burnham KP, Anderson DR. Multimodel inference: understanding AIC and BIC in model selection. Sociological methods & research. 2004;33(2):261-304.

33.   Lewis F, Butler A, Gilbert L. A unified approach to model selection using the likelihood ratio test. Methods in ecology and evolution. 2011;2(2):155-62.

34.   Bayer JMM, Dinga R, Kia SM, Kottaram AR, Wolfers T, Lv J, et al. Accommodating site variation in neuroimaging data using normative and hierarchical Bayesian models. NeuroImage. 2022;264:119699.

35.   Verbeke G. Linear mixed models for longitudinal data. Linear mixed models in practice: Springer; 1997. p. 63-153.

36.   Team RC. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing; 2024.

37.   Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. Journal of the Royal statistical society: series B (Methodological). 1995;57(1):289-300.

38.   Young CB, Reddy V, Sonne J. Neuroanatomy, basal ganglia. StatPearls [Internet]:
StatPearls Publishing; 2023.
39.   Lago T, Davis A, Grillon C, Ernst M. Striatum on the anxiety map: Small detours into
adolescence. Brain research. 2017;1654:177-84.
40.   Telzer EH, Fowler CH, Davis MM, Rudolph KD. Hungry for inclusion: Exposure to
peer victimization and heightened social monitoring in adolescent girls. Development and
psychopathology. 2020;32(4):1495-508.
41.   Zhang S, Zhang Y, Ma W, Qi Z, Wang Y, Tao Q. Neural correlates of negative emotion
processing in subthreshold depression. Social Cognitive and Affective Neuroscience.
2022;17(7):655-61.
42.   Chang CH, Grace AA. Amygdala-ventral pallidum pathway decreases dopamine
activity after chronic mild stress in rats. Biol Psychiatry. 2014;76(3):223-30.
43.   Boedhoe PS, Schmaal L, Abe Y, Ameis SH, Arnold PD, Batistuzzo MC, et al. Distinct
subcortical volume alterations in pediatric and adult OCD: a worldwide meta-and mega-
analysis. American Journal of Psychiatry. 2017;174(1):60-9.
44.   Zhang W, Provensi G, Maffei A. New challenges and perspectives in emotion
regulation and processing. Frontiers in Behavioral Neuroscience. 2022;16:1006627.
45.   De Kloet ER, Joëls M, Holsboer F. Stress and the brain: from adaptation to disease.
Nature reviews neuroscience. 2005;6(6):463-75.
46.   Esmaeel TEZ, Abdelghany HS, Mounir SM, Rasekh AI, Mahmoud HAA, Allam
MFAB. The utility of volumetric MRI in assessment of volume changes in ventral
diencephalon in autistic children. Egyptian Journal of Radiology and Nuclear Medicine.
2023;54(1):173.
47.   Ozgen Saydam B, Yildiz BO. Polycystic ovary syndrome and brain: an update on
structural and functional studies. The Journal of Clinical Endocrinology & Metabolism.
2021;106(2):e430-e41.
48.   Jadambaa A, Thomas HJ, Scott JG, Graves N, Brain D, Pacella R. The contribution of
bullying victimisation to the burden of anxiety and depressive disorders in Australia.
Epidemiology and Psychiatric Sciences. 2020;29:e54.
49.   Forster GL, Simons RM, Baugh LA. Revisiting the role of the amygdala in
posttraumatic stress disorder. The amygdala-where emotions shape perception, learning and
memories: InTechOpen, London; 2017. p. 67585.
50.   Swartz JR, Carranza AF, Knodt AR. Amygdala activity to angry and fearful faces
relates to bullying and victimization in adolescents. Social Cognitive and Affective
Neuroscience. 2019;14(10):1027-35.
51.   Yamamoto T, Toki S, Siegle GJ, Takamura M, Takaishi Y, Yoshimura S, et al.
Increased amygdala reactivity following early life stress: a potential resilience enhancer role.
BMC Psychiatry. 2017;17(1):27.
52.   Ye Z, Wu D, He X, Ma Q, Peng J, Mao G, et al. Meta-analysis of the relationship
between bullying and depressive symptoms in children and adolescents. BMC Psychiatry.
2023;23(1):215.
53.   Dantchev S, Hickman M, Heron J, Zammit S, Wolke D. The independent and
cumulative effects of sibling and peer bullying in childhood on depression, anxiety, suicidal
ideation, and self-harm in adulthood. Frontiers in psychiatry. 2019;10:651.
54.   Vannucci M, Nocentini A, Mazzoni G, Menesini E. Recalling unpresented hostile
words: False memories predictors of traditional and cyberbullying. European Journal of
Developmental Psychology. 2012;9(2):182-94.
55.   Cubillo A. Neurobiological correlates of the social and emotional impact of peer
victimization: A review. Frontiers in psychiatry. 2022;13:866926.

56.   Cavanna AE, Trimble MR. The precuneus: a review of its functional anatomy and
behavioural correlates. Brain. 2006;129(3):564-83.
57.   Grill-Spector K, Malach R. The human visual cortex. Annu Rev Neurosci.
2004;27(1):649-77.
58.   Pierce JE, Péron J. The basal ganglia and the cerebellum in human emotion. Social
cognitive and affective neuroscience. 2020;15(5):599-613.
59.   Bejerot S, Plenty S, Humble A, Humble MB. Poor motor skills: A risk marker for bully
victimization. Aggressive Behavior. 2013;39(6):453-61.
60.   Franzen M, de Jong PJ, Veling W, aan het Rot M. Victims of bullying: Emotion
recognition and understanding. Frontiers in psychology. 2021;12:729835.
61.   Useche SA, Valle-Escolano R, Valle E, Colomer-Pérez N. Gender differences in
teenager bullying dynamics and predictors of peer-to-peer intimidation. Heliyon.
2023;9(9):e20243.
62.   Aminoff EM, Kveraga K, Bar M. The role of the parahippocampal cortex in cognition.
Trends Cogn Sci. 2013;17(8):379-90.
63.   Bhanji JP, Delgado MR. The social brain and reward: social information processing in
the human striatum. Wiley Interdiscip Rev Cogn Sci. 2014;5(1):61-73.
64.   Delgado MR, Nearing KI, Ledoux JE, Phelps EA. Neural circuitry underlying the
regulation of conditioned fear and its relation to extinction. Neuron. 2008;59(5):829-38.
65.   Ulrich-Lai YM, Herman JP. Neural regulation of endocrine and autonomic stress
responses. Nature Reviews Neuroscience. 2009;10(6):397-409.
66.   Ben-Shabat E, Matyas TA, Pell GS, Brodtmann A, Carey LM. The Right
Supramarginal Gyrus Is Important for Proprioception in Healthy and Stroke-Affected
Participants: A Functional MRI Study. Front Neurol. 2015;6:248.
67.   Manto M, Bower JM, Conforto AB, Delgado-García JM, da Guarda SN, Gerwig M, et
al. Consensus paper: roles of the cerebellum in motor control--the diversity of ideas on
cerebellar involvement in movement. Cerebellum. 2012;11(2):457-87.
68.   Martinez-Ferre A, Martinez S. Molecular Regionalization of the Diencephalon.
Frontiers in Neuroscience. 2012;Volume 6 - 2012.
69.   Ouellet-Morin I, Cantave C, Paquin S, Geoffroy MC, Brendgen M, Vitaro F, et al.
Associations between developmental trajectories of peer victimization, hair cortisol, and
depressive symptoms: a longitudinal study. Journal of Child Psychology and Psychiatry.
2021;62(1):19-27.
70.   Vaillancourt T, Duku E, Becker S, Schmidt LA, Nicol J, Muir C, et al. Peer
victimization, depressive symptoms, and high salivary cortisol predict poorer memory in
children. Brain and Cognition. 2011;77(2):191-9.
71.   Kliewer W, Sosnowski DW, Noh H, McGuire K, Wright AW. Peer victimization and
cortisol production in children and adolescents: A systematic review. Journal of Applied
Biobehavioral Research. 2019;24(4):e12172.
72.   Murphy F, Nasa A, Cullinane D, Raajakesary K, Gazzaz A, Sooknarine V, et al.
Childhood Trauma, the HPA Axis and Psychiatric Illnesses: A Targeted Literature Synthesis.
Frontiers in Psychiatry. 2022;Volume 13 - 2022.
73.   Quax RA, Manenschijn L, Koper JW, Hazes JM, Lamberts SWJ, van Rossum EFC, et
al. Glucocorticoid sensitivity in health and disease. Nature Reviews Endocrinology.
2013;9(11):670-86.
74.   Krasner H, Ong CV, Hewitt P, Vida TA. From Stress to Synapse: The Neuronal
Atrophy Pathway to Mood Dysregulation. Int J Mol Sci. 2025;26(7).
75.   Vellapandian C. Hypothalamic-Pituitary-Adrenal (HPA) Axis: Unveiling the Potential
Mechanisms Involved in Stress-Induced Alzheimer's Disease and Depression. Cureus.
2024;16(8):e67595.

76.　　Lucassen PJ, Pruessner J, Sousa N, Almeida OF, Van Dam AM, Rajkowska G, et al.
Neuropathology of stress. Acta Neuropathol. 2014;127(1):109-35.

77.　　LaDage LD, McCormick GL, Robbins TR, Longwell AS, Langkilde T. The effects of
early-life and intergenerational stress on the brain. Proc Biol Sci. 2023;290(2011):20231356.

78.　　Herman JP, Tasker JG. Paraventricular Hypothalamic Mechanisms of Chronic Stress
Adaptation. Front Endocrinol (Lausanne). 2016;7:137.

79.　　Suárez-Pereira I, Llorca-Torralba M, Bravo L, Camarena-Delgado C, Soriano-Mas C,
Berrocoso E. The Role of the Locus Coeruleus in Pain and Associated Stress-Related
Disorders. Biological Psychiatry. 2022;91(9):786-97.

80.　　Giustino TF, Ramanathan KR, Totty MS, Miles OW, Maren S. Locus Coeruleus
Norepinephrine Drives Stress-Induced Increases in Basolateral Amygdala Firing and Impairs
Extinction Learning. J Neurosci. 2020;40(4):907-16.

81.　　Padival M, Quinette D, Rosenkranz JA. Effects of repeated stress on excitatory drive
of basal amygdala neurons in vivo. Neuropsychopharmacology. 2013;38(9):1748-62.

82.　　Arnsten AFT, Raskind MA, Taylor FB, Connor DF. The effects of stress exposure on
prefrontal cortex: Translating basic research into successful treatments for post-traumatic stress
disorder. Neurobiology of Stress. 2015;1:89-99.

83.　　Kellij S, Lodder GMA, van den Bedem N, Güroğlu B, Veenstra R. The Social
Cognitions of Victims of Bullying: A Systematic Review. Adolescent Research Review.
2022;7(3):287-334.

84.　　Baik J-H. Stress and the dopaminergic reward system. Experimental & Molecular
Medicine. 2020;52(12):1879-90.

85.　　Fasano C, Bourque MJ, Lapointe G, Leo D, Thibault D, Haber M, et al. Dopamine
facilitates dendritic spine formation by cultured striatal medium spiny neurons through both
D1 and D2 dopamine receptors. Neuropharmacology. 2013;67:432-43.

86.　　McEwen BS, Gianaros PJ. Central role of the brain in stress and adaptation: links to
socioeconomic status, health, and disease. Ann N Y Acad Sci. 2010;1186:190-222.

87.　　Aricak T, Siyahhan S, Uzunhasanoglu A, Saribeyoglu S, Ciplak S, Yilmaz N, et al.
Cyberbullying among Turkish adolescents. Cyberpsychology & behavior. 2008;11(3):253-61.

88.　　Singer JD, Willett JB. Applied longitudinal data analysis: Modeling change and event
occurrence. New York, NY, US: Oxford University Press; 2003. xx, 644-xx, p.

89.　　Vijayakumar N, Mills KL, Alexander-Bloch A, Tamnes CK, Whittle S. Structural brain
development: A review of methodological approaches and best practices. Developmental
Cognitive Neuroscience. 2018;33:129-48.