Title

Comparative risk-assessment of highly pathogenic avian influenza H5 viruses spread

in French broiler and layer sectors.

Authors

Claire Hautefeuille[1*], Facundo Muñoz[1], Gwenaëlle Dauphin[2], Mathilde Paul[3], Marisa

Peyre[1] and Flavie Goutard[1]

[1]CIRAD, UMR ASTRE, F-34398 Montpellier, France.

ASTRE, Univ Montpellier, CIRAD, INRAE, Montpellier, France

[2] CEVA Animal Health, Libourne, France

[3] IHAP, Université de Toulouse, INRAE, ENVT, Toulouse, France

* Corresponding author

E-mail: claire.hautefeuille@cirad.fr

Abstract

Since 2015, French poultry production is threatened almost every year by a

reintroduction of highly pathogenic avian influenza H5 viruses. The duck sector was

the most concerned by this crisis but other sectors such as broiler, layer and turkey

were also affected by outbreaks. The objective of this work was to assess the risk of

highly pathogenic avian influenza H5 virus transmission from one farm to another

within the French broiler and layer production network.

This study used the WOAH risk assessment framework. After drawing up a scenario

tree of virus transmission from one farm to another, data were collected through a

literature review or through experts' elicitation. Three questionnaires were developed

according to the experts' field of expertise: avian influenza, broiler and layer sectors.

The experts' estimates were combined using a beta distribution weighted by their

confidence level. A Monte Carlo iteration process was used to combine the different

probabilities of the scenario tree and to assess the transmission risk.

In the broiler sector, the highest transmission probabilities were observed if the

exposed farm was an indoor broiler farm and the source a broiler farm (indoor or

free-range). The high transmission probability between broiler farms integrated within

the same association suggests that integration is an important risk factor. Person

movement, transport of feed and manure management were the pathways with the

highest transmission probabilities between two integrated indoor broiler farms with

good biosecurity levels. In the layer sector, the highest transmission probabilities

were observed if the source farm was a free-range farm and the exposed farm a

production farm (indoor or free-range). The pathways with the highest transmission

probabilities were egg transport and person movement. The sensitivity analysis

showed that the exposed farm's biosecurity had a significant impact on the

transmission probability.

Our results provide an insight on the role of each type of farms in the virus spread
within the French broiler and layer production sectors and will be useful for the
implementation of control measures such as movement restriction or vaccination.

48 Introduction

Since 2015, France has faced several epizootic waves of highly pathogenic avian
influenza (HPAI) viruses [1–3]. The main poultry production sector affected by these
waves has been the fattening duck production sector – for the production of duck
liver-based delicacies called "foie-gras" [4,5]. During these waves, while the initial
introduction of HPAI viruses has likely been due to wild birds [1,6], the large spread
of the viruses between poultry farms was linked with the poultry production systems.
The structure of the fattening duck sector was one of the main factors facilitating the
spread of HPAI [6]. Indeed, in this sector, live birds' movements occur between the
different production stages: rearing, breeding and force-feeding. It has been
demonstrated that these movements have played a major role in the spread of HPAI
viruses [7,8].

Other poultry production sectors exist in France: the meat sector (broiler, meat duck,
turkey, guinea fowl) and layer sector [9]. In the broiler production sector, birds enter
the farm as day-old chicks and leave the farm to go to the slaughterhouse. In the
layer sector, production is organized in two stages: pullet (from day-old chick to
ready-to-lay layers) and layer (from ready-to-lay layers to spent hen). In between
these two stages, live bird movements take place to transport ready-to-lay layers
from the pullet farm to the layer farm. The same movement pattern is observed for
breeders of all species. A live bird movement occurs between the growth stage farm

and the fertile adult stage farm. Even if there are no live bird movements between

broiler or layer farms, other avian influenza (AI) virus pathways such, as fomite or

human movements, may exist [10]. Identified fomite movements may occur through

birds picked-up for slaughter, feed delivery, egg collection, manure and litter

management, dead bird management and shared equipment [11–15]. Movements

involving direct contacts between humans and birds are identified as presenting the

higher risk of AI virus spread (veterinarian, staff working on multiple premises,

company staff, technical advisors etc.) [16,17].

Few risk assessment studies looked at the impact of the integration level (integrated

or independent) or farming conditions (indoor vs. free-range) on the risk of AI viruses

introduction in a farm [13,15,18]. Moreover, a few modelling studies collected data on

the frequency of fomite and human movements between farms [11,14]. Nevertheless,

these studies never took a close look at the origin of these movements, i.e. type of

farm from which the movement originated.

So far, no study has been conducted on the risk of AI introduction into a chicken farm

from another farm within the chicken production sector in France, even though these

sectors have also been strongly affected by outbreaks (329 outbreaks in chicken

farms during the 2020-22 period) [2]. The aim of this study was to assess the risk of

spread of AI virus within two other French poultry production sectors: broiler and

layer. This risk assessment study has been conducted considering different types of

farms as described in a previous study [9]. These types of farms were defined

according to the sector (broiler and layer)), production type (i.e. future breeders,

breeders, production), the farming conditions (i.e. indoor and free-range) and the

integration level of the farm. For this purpose, this study assesses the risk of

transmission from one farm of a defined type to another farm of the same or different

type within the same production sector. It did not consider the transmission risk

between farms from different poultry sectors and not the risk of introduction through

environmental pathways. The current study focuses on the relative transmission risk,

i.e. to identify the type of farms with the highest risk of virus transmission through

indirect contacts (human and fomites movements). This study will allow us to identify

pathways and farm types most at risk of AI virus transmission.

Materials and methods

Risk assessment model

This study focused on the assessment of the risk of introduction described by the

WOAH framework [19,20]. The aim of the study was to evaluate the relative risk of

viral spread between different types of farms. We assumed that once the virus is

introduced into the flock, the probability of a bird being infected, as well as the risk of

exposure, is close to 100%, and we did not assess the consequences of AI spread.

We considered it highly likely that bird movements from an infected farm would

introduce the virus and cause an infection on a free farm. Consequently, we focused

solely on the risk of introduction through indirect pathways within a poultry production

network. Four pathways were identified in a previous literature scoping review [10]:

human with a direct contact with birds; trucks transporting live birds; trucks

transporting feeds, manure and dead birds; shared equipment in direct contact with

birds (Figure 1). Two other pathways that are known to represent a very low risk

were not considered: human and shared equipment with no direct contact with birds

or manure [11,15,21]. Environmental pathways such as air-born or rodent as

mechanical vector transmission pathways were assumed homogeneous across farm

types and excluded from the study. Geographical proximity between farms was not

considered either.

Figure 1: The transmission pathways considered in this risk-assessment study.

The assessment was conducted for the broiler and layer sectors on several types of

farms defined in Table 1 and adapted from the description provided by Hautefeuille et

al. [9]. Considering the low number of breeder farms and future layers farms, we did

not include the integration level for these types of farms.

Table 1: Listing of the different types of farms studied according to the production

type, the farming conditions and the integration level of farms in the French broiler

and layer production sector

Farming type	Production type	Integration level
Indoor	Broiler	integrated in the same farmers' association
		integrated in another farmers' association
		independent
Free-range	Broiler	integrated in the same farmers' association
		integrated in another farmers' association

		independent
Selection	Breeder	-
	Future breeder	-

Data source

The probability of virus introduction from each pathway was estimated using scenario

trees as recommended by EFSA [22]. Two scenario trees were built (Figure 2): one

to estimate the probability of the non-detection of an infected farm (Figure 2 and

Table 2) and one to estimate the probability of viral transmission from the source

farm to the exposed farm through a given pathway (Figure 3 and Table 3). The first

step of this study was to identify, in the scenario trees, the branches for which

estimates of risk probabilities could be obtained from the French legislation. The

second step was to generate data for each pathway on the transmission risk, the

frequency of occurrence for one type of farm and the probability that this pathway

exists between two different types of farms. An expert elicitation survey was

conducted to obtain the data not available from the scientific literature.

Figure 2: Scenario tree to assess the risk of not detecting an infected farm

Table 2: Probability that an infected farm (source) is not detected

Node	Parameters estimates	Branch of node	Data sources or justification

Farm routinely tested	Probability that a farm is routinely tested against HPAI	Yes No	French legislation [23]
Time of infection	Estimation of the proportion of farms infected for more or less than 3 days	Less than 3 days More than 3 days	Incubation period [24]
Clinical signs	Probability that a trained farmer detects clinical signs of HPAI infection before 3 days of infection	Yes No	Expert elicitations
	Probability that a trained farmer detects clinical signs of HPAI infection after 3 days of infection	Yes No	Expert elicitations
Farm tested if clinical signs detected	Probability that a farm is tested if clinical signs are detected	Yes No	Expert elicitations
Farm reported if tested positive	Probability that a farm is reported if tested positive	Yes No	French legislation [23]

148 Figure 3: Scenario tree to assess the risk of virus transmission from an infected farm

149 (source) to a free farm (exposed) through a given pathway.

150 Table 3: Probability that a silently-infected farm (source farm) transmits the virus to a

151 free farm (exposed farm) for a given pathway

Node	Parameters estimates	Branch of node	Data sources or justification

Biosecurity level of source farm (infected farm)	Estimation of the proportion of farms of the same type of farms than source farm in the different biosecurity levels	Very low Low Good High Very High	Expert elicitations
Virus release from source farm (infected farm)	Probability that the virus is released from source farm according to the studied pathway	Yes No	Expert elicitations
Link between the source farm and the exposed farm	Probability that the studied pathway occurred in the exposed farm	Yes No	Expert elicitations or production cycle values
	Probability that the studied pathway went during the previous week in the source farm	Yes No	Expert elicitations
Biosecurity level of exposed farm (free of infection)	Estimation of the proportion of farms of the same type of farms than the exposed farm in the different biosecurity levels	Very low Low Good High Very High	Expert elicitations
Virus introduction in the exposed farm	Probability that the virus is introduced in the exposed farm according to the studied pathway	Yes No	Expert elicitations

Fomite (i.e. person, truck or material) disinfection outside farms was considered to be

similar for all types of farms and not sufficient to stop the virus spread.

Once the virus is introduced in the exposed farm (free of infection), it was considered

that the probability that the virus enters in contact with birds varies according to the

pathway. A farm consists of several areas to meet biosecurity criteria as explained in

the introduction of the biosecurity grid: public area for visitors, professional area

where only the persons and vehicles authorized to work on the farm can go, and the

production unit (i.e. the building where the birds are raised with a possible outdoor

area). For all studied pathways, the area into which the virus is introduced (Table 4

and Figure 4) and the probability that a virus is transmitted from one area to another

(Table 5) were defined.

Table 4: List of the different studied pathways and farm area of virus introduction

according to this pathway

Pathways	Farms concerned	Area of the farm where the virus is introduced
Human with a direct contact with birds	All	Direct contact with birds
Chick transport	Broiler Future breeder	
Birds' pick-up transport (pullet and to slaughter)	All	Professional area
Feed delivery	All	
Manure management	All	

Dead bird management	All	Public area
Shared equipment	All	Production unit
Egg	Layer, Breeder	Professional area

168 Figure 4: Schematic representation of the different areas composing a broiler farm
169 (based on[25])

171 Table 5: Probability that a virus coming into contact with birds after introduction into a

172 free farm of a given biosecurity level

Node	Parameters estimates	Branch of node	Data sources or justification
Virus presents in public are	Probability that the virus in introduced in the professional area	Yes No	Expert elicitations
Virus presents in professional area	Probability that the virus in introduced in the production unit	Yes No	Expert elicitations
Virus presents in production unit	Probability that the virus enters in contact with birds	Yes No	Expert elicitations

Experts' elicitation survey

The aim of this survey was to collect data on the transmission risk for each type of farms (8 farm types for broiler and 7 for layer production) and for each pathway (9 different pathways). We created questionnaires for three different profiles: one for experts on AI epidemiology and poultry farm biosecurity, one for experts on French broiler sector and one for experts on French layer sector.

Experts were recruited according to their experience on the different questionnaire topics. The survey on AI was initially sent to scientists from national French research centres who are currently working and publishing on AI in France. Surveys about the broiler and the layer sectors were initially sent to people who represent these industries at a national level. Following this initial recruitment, a "snowball" recruitment was conducted, in which experts who participated in the survey were asked to recommend other experts to fill out the questionnaire. Experts filled the survey independently without being in contact with each other. The survey was not anonymous so that experts could be contacted for clarification, but all their answers were analysed and presented anonymously.

The questionnaire was designed and deployed online using the platform Survey Monkey (www.surveymonkey.com). For each questionnaire a pilot survey was sent to two experts in the corresponding field in order to test and optimize the survey. To make the questionnaires shorter, some simplifications were decided and validated by these two experts (Table 6).

Table 6: Simplification made according to the probability and the pathway concerned and justification validated by two experts of the poultry production sector.

Probability concerned	Pathway concerned	Simplification made	Justification
Probability of virus introduction and release	Pullet transport Birds to slaughter transport	Merge of the two pathways as pick-up transport	Pick-up of birds is conducted in a similar way for pullet and birds to slaughter.
Probability that the studied pathway went during the previous week in farm A (i.e. Probability of pathway origin)	Manure Dead birds	Merge of the two pathways	Management similar for both pathways
	Shared equipment	No consideration of the integration level of farms	Management similar regardless the integration level

The experts were emailed a description of the study objectives and of what was expected for their participation with the link to the online survey. The beginning of the online survey included instructions and contact information of the administering researchers. The questionnaires are available in French (Supplementary file 1).

Each questionnaire was designed to take about one hour to be filled by experts. Questionnaires started with a description of the experts and questions on their expertise level. The first questionnaire on AI epidemiology was divided in four sections: risk of AI virus introduction in a farm, risk of AI virus release from an infected farm, risk associated with wild birds, and risk of an infected farm going undetected. To simplify the questionnaire, risk was assessed per farm biosecurity

level rather than farm type. Grids usually used for biosecurity evaluation process are

developed to assess the biosecurity at one farm level in a very detailed way and in

the aim to improve the biosecurity of this one farm. Thus, it was necessary to develop

a specific less detailed biosecurity grid to order farms within one farming type

between five biosecurity levels defined from very low to very high. This biosecurity

grid was built based on previous works on the biosecurity evaluation process [7,26]

and guidelines developed by the French poultry technical institute (ITAVI) [25]

(Supplementary file 2). This grid was validated by two experts outside of the list of

experts identified for the risk assessment and currently working on French poultry

farms biosecurity. For the five defined biosecurity levels, experts were asked about

the probability of introduction or release of the virus for each risk pathway (poultry

production related or wild birds related). At the end, the experts were asked to define

the probability that an infected farm would go undetected. The other two

questionnaires were designed to collect data on broiler and layer sectors. To assess

the biosecurity level of a studied farm type, experts were asked to distribute 100

farms of this type between the five different levels in order to have the percentage of

farms per biosecurity level (Supplementary file 1). The second section aimed to

identify for each introduction pathway and for each type of farm, the frequency that

this pathway occurred and the probability that this pathway links the studied farm with

another type of farms. The questionnaire concluded on the capacity of a farmer to

detect an AI infection based on the type of farm. Regarding the probabilities of virus

introduction or release according to the several pathways, experts were asked for the

most probable, along with the minimum and maximum values. For question

concerning probabilities of pathway occurrence, experts were asked for a range of

probabilities (Null: 0%, Negligeable: >0 – 20%, Low: 20-40%, Moderate: 40-60%,

High: 60-80% and Very high: >80%). For question concerning frequencies, experts

were asked to select on possibility from a range between "more than seven times a

week" and "less than once every two months". For each question, experts were

asked to provide their confidence level with a score from 1 to 5 (Table 7) and they

could add comments. Finally, for the three questionnaires, the experts were asked

some open-ended questions regarding factors that might have influenced their

answers. All the answers provided by the experts are available in Supplementary file

1.

Table 7: Scoring of confidence levels

Level of confidence	Score
Certain	5
Very confident	4
Moderately confident	3
Not confident	2
Uncertain	1

244 Data analysis

Data collected from the SurveyMonkey platform were saved in a database template

developed in Microsoft Excel® version 2007. The data from these two questionnaires

were used to elicit Beta distributions characterising the ensemble of experts'

assessments on each probability of interest. For each probability of interest, the

parameters of the Beta distribution were determined in terms of expected value and

variance of the distribution, which were in turn calculated from the expert's responses

as follows. For the question asking for the most probable, minimum and maximum

values of the target probabilities, the expected value was set at the average of the

most probable values weighted by the expert confidence level and the variance was

calculated using a formula (1) balancing a maximal variance V_m and an average

variance V_i between experts.

$$var = V_i + \frac{(V_m - V_i)^3}{V_m{}^2}\,(1)$$

The maximal variance V_m corresponds to the variance of a Uniform distribution with

support over the maximal interval (i.e. the smallest interval containing all the intervals

reported by the experts). The average variance V_i, by contrast, is the average of the

variances of the Uniform distributions over each of the reported intervals. This

variance calculation allows the representations of divergent opinions without reducing

compatible assessments. For questions requiring only one interval of values (i.e.

21% to 40%, 41% to 60%, etc.), such as questions on between farm contact

probabilities, the centre of the interval was considered as the most probable, and the

boundaries as the minimum and the maximum value for each expert. The analysis

was conducted with R (version 4.0.2).

The frequency of occurrence and the origin of some pathways did not need to be

assessed by experts as these values depend on the production cycle. The egg

transport was considered to enter into a given breeder farm every day and the

possible origin of this truck was always breeder farms. The frequency of chick and

pullet delivery and slaughter take off was defined by the production cycle of the

different types of production. As it appears that experts could not provide an

occurrence frequency for shared equipment, it was arbitrarily considered to occur

once a month and validated with the experts who validated the questionnaire.

A Monte Carlo iterative process was carried out on the scenario trees described

above, in order to combine the different probabilities and to calculate the

transmission probability from one type of farm to another. For each iteration, the

probability of realisation of the event for all branches of the scenario tree were

randomly sampled from the Beta distribution previously elicited and all probabilities

were multiplied to obtain the transmission probability. The simulation process used

50 000 iterations.

In addition, we estimated the relative transmission probabilities between farm types

with respect to the pair that exhibited the highest mean transmission probability. The

target pair of source and exposed farm types was deemed to be at very high

transmission risk when the sampled transmission probability was greater than that of

the reference pair in 40% or more of the Monte Carlo iterations. For percentages

between 25 and 40%, the pair was considered to be at high transmission risk. For

percentages between 15 and 25%, 5 and 15% and below 5%, the pair was

considered to be at medium, low and very low transmission risk respectively.

Sensitivity analysis

A sensitivity analysis was conducted using the one-at-time method applied on the

biosecurity level of the source and the exposed farms and on the pathways [27]. This

analysis was applied only for two types of farms with the most extreme daily

transmission probabilities: one representing the highest and one the lowest. This

sensitivity analysis is based on the distribution of the transmission probabilities on 10

000 iterations.

Results

Experts' description

At the end of the process, six experts responded to the questionnaire on AI epidemiology: three had more than ten years of experience, one had over five years and two had more than three years. Seven experts, all with over ten years of experience, responded to the questionnaire on the broiler sector. Five experts, each with over ten years of experience, answered the questionnaire on the layer sector.

Probability of avian influenza virus transmission

The survey results for the broiler production network indicated that transmission probabilities were highest either from indoor integrated production farms (i.e. not selection farms) to indoor broiler farms (regardless of integration level), or from indoor or free-range production farms to independent indoor farms. These farm pairs were considered to be at very high transmission risk (Table 8). One of the highest median transmission probabilities was observed between independent indoor broiler farms, which served as the reference for relative comparison. The lowest probabilities were found either from selection farms to free-range farms (regardless of the level of integration) or when the exposed farm was a selection farm. High transmission probabilities were also found from free-range or independent indoor farms to integrated indoor farms, and between free-range independent farms. Finally, while transmission probability was of medium level from production to free-range farms, it was low from selection to indoor farms.

For the layer production network, the transmission probabilities were the highest

between production farms (pullet, indoor or free-range layer) (Table 9). One of the

highest median transmission probabilities was between independent free-range layer

farms, which were used as the reference farms for the relative risk comparison.

Transmission probabilities were the lowest between selection farms. Finally,

transmission probabilities were medium from production to pullet farms, from pullet to

production farms and from indoor to production farms (except for independent indoor

farms).

Table 8: Daily transmission probabilities of HPAI H5 virus between an infected source farm and an exposed farm (free of infection)

according to the type of farms in the French broiler production network (median, Q25 and Q75 values of 50 000 iterations).

Percentage of probabilities below the one estimated between independent indoor broiler farms: Dark brown: 50-60% (very high risk

of transmission), brown: 60-75% (high risk), medium brown: 75-85% (medium risk), light brown: 85-95% (low risk), no colour: over

95% (very low risk).

Values in %		Exposed farm (free of infection)					
		Indoor broiler		Free-range broiler		Selection broiler	
Source farm (infected)		Integrated	Independent	Integrated	Independent	Breeder	Future breeder
Indoor broiler	Integrated in the same farmers' association	1.2 [0.5 - 3]	1.0 [0.4 - 2.5]	0.2 [0.1 - 0.4]	0.3 [0.1 - 0.6]	0.004 [0.0005 - 0.02]	0.001 [0.00008 - 0.01]
	Integrated in another farmers' association	0.8 [0.3 - 2]	1.0 [0.4 - 2.5]	0.2 [0.1 - 0.5]	0.3 [0.1 - 0.6]	0.004 [0.0005 - 0.02]	0.001 [0.00008 - 0.01]
	Independent	0.6 [0.2 - 1.4]	1.2 [0.5 - 2.8]	0.2 [0.1 - 0.5]	0.3 [0.2 - 0.7]	0.005 [0.0006 - 0.03]	0.001 [0.00009 - 0.01]
Free-range broiler	Integrated in the same farmers' association	0.5 [0.2 - 1.5]	1.0 [0.4 - 2.4]	0.3 [0.2 - 0.8]	0.3 [0.1 - 0.7]	0.004 [0.0005 - 0.03]	0.001 [0.00008 - 0.01]
	Integrated in another farmers' association	0.5 [0.2 - 1.5]	1.0 [0.4 - 2.4]	0.3 [0.1 - 0.6]	0.3 [0.2 - 0.7]	0.004 [0.0006 - 0.03]	0.001 [0.00009 - 0.01]
	Independent	0.6 [0.3 - 1.7]	1.2 [0.5 - 2.8]	0.3 [0.1 - 0.5]	0.4 [0.2 - 1]	0.005 [0.0007 - 0.03]	0.001 [0.0001 - 0.01]
Selection broiler	Breeder	0.04 [0.009 - 0.2]	0.06 [0.01 - 0.3]	0.01 [0.003 - 0.04]	0.02 [0.005 - 0.05]	0.005 [0.0005 - 0.03]	0.001 [0.00008 - 0.01]
	Future Breeder	0.04 [0.008 - 0.2]	0.06 [0.01 - 0.3]	0.01 [0.002 - 0.04]	0.01 [0.004 - 0.04]	0.002 [0.0002 - 0.02]	0.002 [0.0001 - 0.02]

Table 9: Daily transmission probabilities of HPAI H5 virus between an infected source farm and an exposed farm (free of infection) according to the type of farms in the French layer production network (median, Q25 and Q75 values of 50 000 iterations). Percentage of probabilities below the one estimated between independent free-range layer farms: Dark brown: 50-60% (very high risk of transmission), brown: 60-75% (high risk), medium brown: 75-85% (medium risk), light brown: 85-95% (low risk), no colour: over 95% (very low risk)

Values in % Source farm (infected)		Exposed farm (free of infection)						
		Pullet	Indoor layer		Free-range layer		Selection layer	
			Integrated	Independent	Integrated	Independent	Breeder	Future breeder
Pullet		0.2 [0.08 - 0.5]	0.1 [0.04 - 0.3]	0.1 [0.05 - 0.3]	0.1 [0.05 - 0.3]	0.1 [0.04 - 0.3]	0.006 [0.001 - 0.03]	0.003 [0.0005 - 0.02]
Indoor layer	Integrated	0.1 [0.06 - 0.4]	0.2 [0.09 - 0.6]	0.2 [0.08 - 0.5]	0.2 [0.08 - 0.6]	0.2 [0.09 - 0.6]	0.006 [0.001 - 0.03]	0.004 [0.0006 - 0.02]
	Independent	0.1 [0.06 - 0.4]	0.2 [0.07 - 0.5]	0.3 [0.1 - 0.7]	0.2 [0.09 - 0.6]	0.2 [0.09 - 0.6]	0.006 [0.001 - 0.03]	0.004 [0.0006 - 0.02]
Free-range layer	Integrated	0.2 [0.09 - 0.5]	0.3 [0.1 - 0.8]	0.3 [0.1 - 0.9]	0.4 [0.2 – 0.9]	0.3 [0.1 - 0.8]	0.008 [0.002 - 0.03]	0.005 [0.0008 - 0.02]
	Independent	0.2 [0.07 - 0.5]	0.3 [0.1 - 0.7]	0.3 [0.1 - 0.8]	0.3 [0.1 - 0.7]	0.4 [0.1 – 0.9]	0.008 [0.001 - 0.03]	0.005 [0.0008 - 0.02]
Selection layer	Breeder	0.008 [0.002 - 0.03]	0.006 [0.001 - 0.02]	0.007 [0.002 - 0.03]	0.007 [0.002 - 0.02]	0.008 [0.002 - 0.03]	0.006 [0.0007 - 0.04]	0.003 [0.0004 - 0.02]
	Future breeder	0.007 [0.001 - 0.03]	0.006 [0.001 - 0.02]	0.006 [0.001 - 0.02]	0.006 [0.002 - 0.02]	0.006 [0.002 - 0.02]	0.003 [0.0004 - 0.02]	0.004 [0.0005 - 0.02]

Sensitivity analysis

The sensitivity analysis was conducted on two farm types: one with the highest transmission probability (integrated indoor broiler farm, Figure 5 and Figure 6 Figure 5) and one with the lowest (future breeder farm for broiler production), both as source (infected) and exposed (free) farms (Supplementary file 3). In both cases, the biosecurity level has a strong influence on transmission probability: it was high when both source and exposed farms had low biosecurity and low when both had high biosecurity (Figure 5 Figure 5).

The biosecurity level of the exposed farm had a greater influence than that of the source farm. For all pathways, variations in the source farm's biosecurity (infected) had low impact on transmission probabilities (Figure 6 A). In contrast, variations in the biosecurity level of the exposed farm significantly influence transmission probabilities (Figure 6 B). Indeed, probabilities were high when the exposed farms had a very low to good level (from 1 to 3 out of 5), and very low when the level was high (above 4), for all pathways except person movement.

Figure 5: Distribution of the transmission probabilities across all **pathways** according to the biosecurity level of the source farm (x axis) and to the biosecurity level of the exposed farm (colours) with integrated indoor broiler farm as exposed and source farms. Threshold at 1/100 (black), 1/1000 (red) and 1/10 000 (orange).

Figure 6: Distribution of the transmission probabilities (A) across all **biosecurity levels of the exposed farm (free)** according to the pathways (x axis) and to the biosecurity level of the source farm (infected) (colours) and (B) across all **biosecurity level of the source farm (infected)** according to the pathways (x axis) and to the biosecurity level of the exposed farm (free) (colours) with integrated indoor broiler

367 farm as exposed and source farms. Threshold at 1/100 (black), 1/1000 (red) and 1/10
368 000 (orange).

370 Farm biosecurity and pathways in the broiler sector

Most integrated broiler farms (indoor and free-range) were scored with a good

biosecurity level (median value of 45% of indoor and 37% of free range)

(Supplementary file 4). Independent indoor farms were scored with a higher

biosecurity level (42% at good level) than independent free-range broiler farms (30%

at good level). In both systems, independent broiler farms were estimated to have

lower biosecurity than integrated farms. Among integrated indoor broiler farms with

good biosecurity, the highest transmission probabilities were linked to person

movement and transport of feed (median value of 0.23% and 0.03% respectively).

These pathways were the most frequent with median occurrence frequency of 21%

for person movement and 30% for transport of feed. When considering an integrated

indoor broiler farm as an exposed farm, the probability of contact through person and

feed movement was high if the source farm belongs to the same farmers'

association, medium if from another farmers' association and low otherwise (

Figure 7). Integrated free-range broiler farms had the lowest transmission

probabilities as exposed farm despite similar biosecurity scores than integrated

indoor broiler farms. The pathways with the highest transmission probability between

two integrated free-range broiler farms at good biosecurity level were material, feed

and manure movement (respectively median value of 0.02% and 0.01%). Person

movement had less impact on the probability of transmission in free-range farms than

in indoor farms, associated with a lower occurrence frequency. For both integrated

indoor and free-range farms, transport of birds from hatchery (chicks) or to slaughter

and dead bird management had the least impact on transmission probabilities.

Movement of birds from hatchery and to slaughter had the lowest frequency of

occurrence (associated with the production cycle). Dead bird management was

considered to have a high frequency of occurrence, but this pathway is the only one

that stops in the public area of the farm.

Figure 7:   Probability (median, 50% and 80% quantile central intervals) of indirect

contact in exposed integrated indoor broiler farms (free of infection) by pathway and

type of source farm (infected) (A): Indoor broiler farm integrated in the same farmer

association, B) Indoor broiler farm integrated in another farmer association, C) Indoor

broiler farm independent, D) Free-range broiler farm, E) Breeder farm)

404   Farm biosecurity and pathways in the layer sector

Biosecurity levels were estimated to range from good to very good in indoor layer

farms and from low to good in free-range farms, regardless of integration level. Pullet

farms were considered with a good biosecurity level (Supplementary file 4). Due to

their production, layer birds transport occurs at a lower frequency than broiler

production. The pathways with the highest transmission probabilities were for both

egg transport, person movement and shared equipment. Egg transport had the

highest occurrence frequency in both indoor and free-range layer farms (median

values of 45% and 30% respectively). Person movements and shared equipment

were just as frequent as the other pathways but they are both associated with direct

contact with birds.

Farm biosecurity and pathways in selection farms

For both broiler and layer sectors, experts considered the biosecurity levels of selection farms to have high to very high biosecurity levels (Supplementary file 4), implying very low transmission probabilities, except for person movement. While breeder farm results were similar across sectors, egg transport and manure management appeared to have higher transmission probabilities in broiler than in layer sector (Supplementary file 3 Figure 3). If the transmission probabilities are very low in selection farms at high and very high biosecurity levels, they are closed to those observed in production farms for pathways such as person movement, transport of feed and eggs, and manure management if the biosecurity level of the farms reached a good level.

Discussion

This study analyzed the risk of virus transmission within French production networks, i.e. the risk that a virus released by an infected farm is introduced into a free farm across all possible indirect pathways and farm types. Most AI virus risk assessments have focused on risk factors related to the production network and to wild birds [28,29]. Some studies addressed the risk of introduction or release of AI viruses in farms with various production types [14,15,18,21], and a few explored the risk of virus transmission between farms types [18,30]. However, none considered the integration level of the farm or the variation of transmission risk across farm types.

This study used scenario trees model and a combination of literature data and experts' elicitation partly based on the DELPHI method [31]. Unlike classic DELPHI

method (reaching consensus) we used the difference of opinion among experts to

estimate uncertainty about transmission probabilities. Hence, our study is subject to

potential biases. To reduce the bias, we implemented six different tools: (1) different

questionnaires to segment topics, (2) expert confidence level included into the

statistical analysis, (3) option for experts to skip questions, (4) a comment section for

each group of questions, (5) a guide to help the experts understand the questions

and (6) a biosecurity grid to harmonize expert understanding. The expert elicitation

was conducted in 2020. Even if the number of worldwide outbreaks increased

drastically since 2021 [32] the experts were already aware of the threat because of

2015-16 and 2017-18 epizootics.

A key innovation of this study is the inclusion of biosecurity levels in the risk

assessment. While biosecurity classifications developed for poultry farms generally

provide a farm-level biosecurity score [26,33,34], few studies have assessed level of

biosecurity by farm types (breeder, broiler or layer) [35] or integration level in the

network [36,37] and, none have used this classification to assess the risk of disease

transmission within a given farm type. Using an approach based on farm biosecurity

enabled us to study a larger number of farm types and to generalize our results to

any French poultry farm. The farm biosecurity level of an exposed farm (i.e. a farm

that is free of infection) has a major impact on the risk of transmission of the AI virus

between farms regardless of the pathway and of the biosecurity level of the source

farm (i.e. a farm that is already infected), as previously highlighted in several studies

[38–41]. In our study, farms dedicated to producing birds destined for reproduction

(future breeders and breeder farms) mainly received a score of 5, indicating very high

biosecurity, while production farms were scored 3, indicating good biosecurity. Fewer

than 5% of the production farms were considered to have a very high level of

biosecurity, in line with the results of a study conducted in New Zealand [33]. Another

study in South Korea that measured the risk of HPAI for farms linked by livestock

transport vehicles reported a low risk for breeder chicken farms [18]. In the present

study the proportion of broiler farms with a low biosecurity level (level 2) was higher

among independent farms than in integrated farms. Nonetheless, the differences

observed need to be interpreted with caution as expert confidence was higher for

integrated farms than for independent farms. Similar results were observed in layer

sector except for independent indoor farms. However, only three experts out of five

provided data for this type of farm (mean confidence of 3.7 out of 5) illustrating the

lack of knowledge concerning this minority farm type. A French study provided

information on biosecurity as a function of farm type (including broiler and layer) and

location, but did not investigate independent farms [42].

This study highlighted that farm network organisation increases the occurrence

probability and frequency of transmission pathways between farms within this

network and thus the probability of viral transmission. First, this probability was

highest between broiler farms that belong to the same farmers' association. The

probability of contact was highest between these farms, particularly via contact

pathways such as people in direct contact with birds and feed transport trucks. This

reflects the French poultry network in France where poultry farmers' associations

share feed manufacturer, technicians, and often veterinarians. The role of the poultry

network in the spread of AI between farms was already highlighted in previous

studies [11,13,16]. Second, egg transport plays a major role in the probability of

transmission between layer farms [18,30]. Third, the existence of many of pathways

between breeder farms increases the risk of transmission of the virus. Indeed, our

results show that if the biosecurity level of a breeder farm that is an exposed farm is

only 'good', the probability of transmission is similar to that of production farms. This

result was confirmed by previous studies [35,38]. A study of HPAI H5 outbreaks in

France between 2020 and 2022 showed that at least 9% of outbreaks occurred in a

breeder farm [2].

The study focused only on indirect transmission pathways (i.e., humans or fomites)

rather than direct pathways (i.e., live domestic birds). Live domestic bird movement is

known to present high risk of virus transmission, especially in the duck sector [8].

However, this particular pathway is absent in broiler farms and only used at very

infrequent intervals in layer and breeder farms (about once every 12-13 months).

Improved truck cleaning and disinfection process would considerably reduce the risk

of viral transmission within the network of farms. French studies detected the

presence of the viral genome in half (in 2017) and in one third (in 2021) of trucks

tested after disinfection [43,44]. In the present study, fomites management (i.e. a

person, truck or equipment) outside farms was considered similar across farm types

and insufficient to affect risk transmission. Disinfection of fomites inside the farm is

included in the biosecurity classification. Based on poultry production experts, we

considered that the frequency of occurrence of shared equipment was the same for

all farms (once a month), regardless of the type or the network of farms, as this

occurrence depends on management and location of the farm. While this impacts the

relative role of shared equipment to viral transmission in farms with a low biosecurity

level especially, it does not affect inter-farm comparisons.

Our study focused on the transmission pathways that have been identified as being

the most at risk of transmission in the literature [10], but other transmission

pathways, such as wind or vermin (i.e. mechanical vectors such as rodents or

insects) are also mentioned. In France air-borne transmission is considered limited

[45], even though some viral genome have been detected in air and dust in infected

farms [46,47]. Other studies conducted in the USA suggest that airborne

transmission might play a role in the spread of AI viruses [48,49]. Virus transmission

through the movement of vermin between farms is possible but difficult to distinguish

from other environmental factors [28,30,50]. These last pathways are often linked to

farm proximity. High poultry density is known to have an impact in the spread of AI

viruses [51]. Our study assumed that environmental transmission is constant across

farm types regardless of their level of biosecurity. While this impacts geographic

differentiation, future work could integrate spatial data.

Finally, this work provides a complete picture of the risk of AI viruses spreading

within French broiler and layer production networks including the risk of transmission

of the virus between any type of farms. To mitigate this risk, management measures

should focus on biosecurity and controlling very frequent movements, especially of

people, of feed, manure and dead birds. Network managers should be aware of intra-

association movements and take the necessary actions, such as cleaning and

disinfection, between visits to farms, to reduce this risk. The risk-based farm typology

developed for this study can inform control measures such as restriction of

movement or vaccination.

Acknowledgements

The authors gratefully acknowledge all the experts who answered the questionnaire

and the reviewers for their constructive comments.

Funding

This work was co-funded by Ceva Santé Animale (https://www.ceva.com) and Cirad

(https://www.cirad.fr/) within the framework of a public private partnership PhD

funding (Cifre funding n° 2017-1177 from Association Nationale de la Recherche et

de la Technologie (ANRT: https://www.anrt.asso.fr/fr/accueil-anrt)) and within the

BioFluARN project framework, co-funded by the French government via the Banque

Publique d'Investisement France (BPIFrance: https://www.bpifrance.fr/) as part of the

"Plan de Relance" and the "Programme d'investissements d'avenir". The funders had

no role in study design, data collection and analysis, decision to publish, or

preparation of the manuscript.

547 Conflict of interest disclosure

548 The authors declare that they have no financial conflict of interest with the content of

549 this article.

551 References

1. Briand F-X, Schmitz A, Ogor K, Prioux AL, Guillou-Cloarec C, Guillemoto C, et
al. Emerging highly pathogenic H5 avian influenza viruses in France during
winter 2015/16: phylogenetic analyses and markers for zoonotic potential.
Eurosurveillance. 2017;22: 30473. doi:10.2807/1560-7917.ES.2017.22.9.30473

2. Lambert S, Durand B, Andraud M, Delacourt R, Scoizec A, Le Bouquin S, et al.
Two major epidemics of highly pathogenic avian influenza virus H5N8 and H5N1
in domestic poultry in France, 2020–2022. Transboundary and Emerging
Diseases. 2022. doi:10.1111/tbed.14722

3. Napp S, Majó N, Sánchez-Gónzalez R, Vergara-Alert J. Emergence and spread
of highly pathogenic avian influenza A(H5N8) in Europe in 2016-2017.
Transboundary and Emerging Diseases. 2018;65: 1217–1226.
doi:10.1111/tbed.12861

4. Bronner A, Niqueux E, Schmitz A, Bouquin SL, Huneau-Salaün A, Guinat C, et
al. Description de l'épisode d'influenza aviaire hautement pathogène en France
en 2016-2017. Bulletin épidémiologique, santé animale et alimentation. 2017;79:
13–17.

5. Le Bouquin S, Scoizec A, Niqueux E, Schmitz A, Briand F-X. L'épisode
d'influenza aviaire en France en 2015-2016 – Situation épidémiologique au 30
juin 2016. Bulletin épidémiologique, santé animale et alimentation. 2016;75: 7.

6. Guinat C, Artois J, Bronner A, Guérin JL, Gilbert M, Paul MC. Duck production
systems and highly pathogenic avian influenza H5N8 in France, 2016–2017.
Scientific Reports. 2019;9. doi:10.1038/s41598-019-42607-x

7. Delpont M, Blondel V, Robertet L, Duret H, Guerin J-L, Vaillancourt J-P, et al.
Biosecurity practices on foie gras duck farms, Southwest France. Preventive
Veterinary Medicine. 2018;158: 78–88. doi:10.1016/j.prevetmed.2018.07.012

8. Guinat C, Durand B, Vergne T, Corre T, Rautureau S, Scoizec A, et al. Role of
Live-Duck Movement Networks in Transmission of Avian Influenza, France,
2016-2017. Emerging infectious diseases. 2020;26: 472–480.
doi:10.3201/eid2603.190412

9. Hautefeuille C, Azzouguen B, Mouchel S, Dauphin G, Peyre M. Evaluation of
vaccination strategies to control an avian influenza outbreak in French poultry
production networks using EVACS tool. Preventive Veterinary Medicine. 2020.

10. Hautefeuille C, Dauphin G, Peyre M. Knowledge and remaining gaps on the role
of animal and human movements in the poultry production and trade networks in
the global spread of avian influenza viruses - A scoping review. PLoS ONE.
2020;15. doi:10.1371/journal.pone.0230567

11. Dorea FC, Vieira AR, Hofacre C, Waldrip D, Cole DJ. Stochastic Model of the
Potential Spread of Highly Pathogenic Avian Influenza from an Infected
Commercial Broiler Operation in Georgia. Avian Diseases. 2010;54: 713–719.
doi:10.1637/8706-031609-ResNote.1

12. Hagenaars TJ, Boender GJ, Bergevoet RHM, van Roermund HJW. Risk of
poultry compartments for transmission of highly pathogenic avian influenza.
PLoS ONE. 2018;13. doi:10.1371/journal.pone.0207076

13. Leibler JH, Carone M, Silbergeld EK. Contribution of Company Affiliation and
Social Contacts to Risk Estimates of Between-Farm Transmission of Avian
Influenza. PLOS ONE. 2010;5: e9888. doi:10.1371/journal.pone.0009888

14. Longworth N, Mourits MCM, Saatkamp HW. Economic Analysis of HPAI Control
in the Netherlands I: Epidemiological Modelling to Support Economic Analysis.
Transboundary and Emerging Diseases. 2014;61: 199–216.
doi:10.1111/tbed.12021

15. Singh M, Toribio JA, Scott AB, Groves P, Barnes B, Glass K, et al. Assessing
the probability of introduction and spread of avian influenza (AI) virus in

commercial Australian poultry operations using an expert opinion elicitation.
PLoS ONE. 2018;13.

16. Dent JE, Kiss IZ, Kao RR, Arnold M. The potential spread of highly pathogenic
avian influenza virus via dynamic contacts between poultry premises in Great
Britain. BMC Veterinary Research. 2011;7: 59. doi:10.1186/1746-6148-7-59

17. Sharkey KJ, Bowers RG, Morgan KL, Robinson SE, Christley RM.
Epidemiological consequences of an incursion of highly pathogenic H5N1 avian
influenza into the British poultry flock. Proceedings of the Royal Society of
London B: Biological Sciences. 2008;275: 19–28. doi:10.1098/rspb.2007.1100

18. Yoon H, Lee I, Kang H, Kim K-S, Lee E. Big data-based risk assessment of
poultry farms during the 2020/2021 highly pathogenic avian influenza epidemic
in Korea. PLoS One. 2022;17: e0269311. doi:10.1371/journal.pone.0269311

19. WOAH. Handbook on Import Risk Analysis for Animals and Animal Products.
2nd ed. 2010.

20. WOAH. Handbook on Import Risk Analysis for Animals and Animal Products.
2nd ed. 2010.

21. Ssematimba A, Hagenaars TJ, de Wit JJ, Ruiterkamp F, Fabri TH, Stegeman
JA, et al. Avian influenza transmission risks: Analysis of biosecurity measures
and contact structure in Dutch poultry farming. Preventive Veterinary Medicine.
2013;109: 106–115. doi:10.1016/j.prevetmed.2012.09.001

22. EFSA Panel on Animal Health and Welfare (AHAW). Guidance on Risk
Assessment for Animal Welfare: Guidance on risk assessment for animal
welfare. EFSA Journal. 2012;10: 2513. doi:10.2903/j.efsa.2012.2513

23. Légifrance. Arrêté du 25 septembre 2023 relatif aux mesures de surveillance, de
prévention, de lutte et de vaccination contre l'influenza aviaire hautement
pathogène (IAHP). Sep 25, 2023. Available:
https://www.legifrance.gouv.fr/loda/id/JORFTEXT000048110961/2025-01-27/

24. Swayne D. Epidemiology of Avian Influenza in Agricultural and Other Man-Made
Systems. Avian Influenza. John Wiley & Sons; 2008. pp. 59–86.

25. ITAVI. Arbre de decision: Toutes les fiches relatives aux mesures de biosécurité
IA. Mar 2019 [cited 27 May 2020]. Available: http://influenza.itavi.asso.fr/

26. Gelaude P, Schlepers M, Verlinden M, Laanen M, Dewulf J. Biocheck.UGent: A
quantitative tool to measure biosecurity at broiler farms and the relationship with
technical performances and antimicrobial use. Poultry Science. 2014;93: 2740–
2751. doi:10.3382/ps.2014-04002

27. Saltelli A, Annoni P. How to avoid a perfunctory sensitivity analysis.
Environmental Modelling & Software. 2010;25: 1508–1517.
doi:10.1016/j.envsoft.2010.04.012

28. Gustafson L, Jones R, Dufour-Zavala L, Jensen E, Malinak C, McCarter S, et al.
Expert elicitation provides a rapid alternative to formal case-control study of an
H7N9 avian influenza outbreak in the United States. Avian Diseases. 2018;62:
201–209. doi:10.1637/11801-011818-Reg.1

29. Grabkowsky BJ, Windhorst H-W. Investigation and quantification of prevalent
risk factors for the introduction of avian influenza in European poultry farms.
World's   Poultry   Science   Journal.   2010;66:   381–398.
doi:10.1017/S0043933910000486

30. Scott AB, Toribio J-ALML, Singh M, Groves P, Barnes B, Glass K, et al. Low-
and High-Pathogenic Avian Influenza H5 and H7 Spread Risk Assessment
Within and Between Australian Commercial Chicken Farms. Front Vet Sci.
2018;5. doi:10.3389/fvets.2018.00063

31. Hasson F, Keeney S, McKenna H. Research guidelines for the Delphi survey
technique. Journal of Advanced Nursing. 2000; 8.

32. Shi J, Zeng X, Cui P, Yan C, Chen H. Alarming situation of emerging H5 and H7
avian influenza and effective control strategies. Emerging Microbes & Infections.
2023;12: 2155072. doi:10.1080/22221751.2022.2155072

33. Greening SS, Mulqueen K, Rawdon TG, French NP, Gates MC. Estimating the
level of disease risk and biosecurity on commercial poultry farms in New
Zealand.   New   Zealand   Veterinary   Journal.   2020;68:   261–271.
doi:10.1080/00480169.2020.1746208

34. Van Limbergen T, Dewulf J, Klinkenberg M, Ducatelle R, Gelaude P, Méndez J,
et al. Scoring biosecurity in European conventional broiler production. Poultry
Science. 2018;97: 74–83. doi:10.3382/ps/pex296

35. Van Steenwinkel S, Ribbens S, Ducheyne E, Goossens E, Dewulf J. Assessing
biosecurity practices, movements and densities of poultry sites across Belgium,
resulting in different farm risk-groups for infectious disease introduction and
spread.   Preventive   Veterinary   Medicine.   2011;98:   259–270.
doi:10.1016/j.prevetmed.2010.12.004

36. Indrawan D, Cahyadi ER, Daryanto A, Hogeveen H. The role of farm business
type on biosecurity practices in West Java broiler farms. Preventive Veterinary
Medicine. 2020;176. doi:10.1016/j.prevetmed.2020.104910

37. Wei H, Aengwanich W. Biosecurity Evaluation of Poultry Production Cluster
(PPCs) in Thailand. International Journal of Poultry Science. 2012;11: 582–588.
doi:10.3923/ijps.2012.582.588

38. Duvauchelle A, Huneau-Salaün A, Balaine L, Rose N, Michel V. Risk factors for
the introduction of avian influenza virus in breeder duck flocks during the first 24
weeks   of   laying.   Avian   Pathology.   2013;42:   447–456.
doi:10.1080/03079457.2013.823145

39. Guinat C, Comin A, Kratzer G, Durand B, Delesalle L, Delpont M, et al.
Biosecurity risk factors for highly pathogenic avian influenza (H5N8) virus

683  infection in duck farms, France. Transboundary and Emerging Diseases. 2020.
684  doi:10.1111/tbed.13672

685 40. McQuiston JH, Garber LP, Porter-Spalding BA, Hahn JW, Pierson FW,
686  Wainwright SH, et al. Evaluation of risk factors for the spread of low
687  pathogenicity H7N2 avian influenza virus among commercial poultry farms.
688  Journal of the American Veterinary Medical Association. 2005;226: 767–772.
689  doi:10.2460/javma.2005.226.767

690 41. Nishiguchi A, Kobayashi S, Yamamoto T, Ouchi Y, Sugizaki T, Tsutsui T. Risk
691  Factors for the Introduction of Avian Influenza Virus into Commercial Layer
692  Chicken Farms During the Outbreaks Caused by a Low-Pathogenic H5N2 Virus
693  in Japan in 2005. Zoonoses and Public Health. 2007;54: 337–343.
694  doi:https://doi.org/10.1111/j.1863-2378.2007.01074.x

695 42. Delpont M, Guinat C, Guérin J-L, Le Ieu E, Vaillancourt J-P, Paul MC.
696  Biosecurity measures in French poultry farms are associated with farm type and
697  location. Preventive Veterinary Medicine. 2021;195: 105466.
698  doi:10.1016/j.prevetmed.2021.105466

699 43. Huneau-Salaün A, Scoizec A, Thomas R, Martenot C, Schmitz A, Pierre I, et al.
700  Avian influenza outbreaks: evaluating the efficacy of cleaning and disinfection of
701  vehicles and transport crates. Poultry Science. 2022;101.
702  doi:10.1016/j.psj.2021.101569

703 44. Huneau-Salaün A, Scoizec A, Thomas R, Le Bouquin S. Cleaning and
704  disinfection of crates and trucks used for duck transport: field observations
705  during the H5N8 avian influenza outbreaks in France in 2017. Poultry Science.
706  2020. doi:10.1016/j.psj.2019.10.015

707 45. Guinat C, Rouchy N, Camy F, Guérin JL, Paul MC. Exploring the wind-borne
708  spread of highly pathogenic avian influenza H5N8 during the 2016-2017
709  epizootic in France. Avian Diseases. 2019;63: 235–237. doi:10.1637/11881-
710  042718-ResNote.1

711 46. Scoizec A, Niqueux E, Thomas R, Daniel P, Schmitz A, Le Bouquin S. Airborne
712  detection of H5N8 highly pathogenic avian influenza virus genome in poultry
713  farms, France. Frontiers in Veterinary Science. 2018;5.
714  doi:10.3389/fvets.2018.00015

715 47. Filaire F, Lebre L, Foret-Lucas C, Vergne T, Daniel P, Lelievre A, et al. Highly
716  Pathogenic Avian Influenza A(H5N8) Clade 2.3.4.4b Virus in Dust Samples from
717  Poultry Farms, France, 2021. Emerging Infectious Diseases. 2022;28: 1446–
718  1450. doi:10.3201/eid2807.212247

719 48. Nguyen XD, Zhao Y, Lin J, Purswell JL, Tabler T, Voy B, et al. Modeling long-
720  distance airborne transmission of highly pathogenic avian influenza carried by
721  dust particles. Sci Rep. 2023;13: 16255. doi:10.1038/s41598-023-42897-2

722 49. Zhao Y, Richardson B, Takle E, Chai L, Schmitt D, Xin H. Airborne transmission
723  may have played a role in the spread of 2015 highly pathogenic avian influenza

outbreaks in the United States. Scientific Reports. 2019;9. doi:10.1038/s41598-
019-47788-z

50. Velkers FC, Blokhuis SJ, Veldhuis Kroeze EJB, Burt SA. The role of rodents in
avian influenza outbreaks in poultry farms: A review. Veterinary Quarterly.
2017;37: 182–194. doi:10.1080/01652176.2017.1325537

51. Bauzile B, Durand B, Lambert S, Rautureau S, Fourtune L, Guinat C, et al.
Impact of palmiped farm density on the resilience of the poultry sector to highly
pathogenic avian influenza H5N8 in France. Vet Res. 2023;54: 56.
doi:10.1186/s13567-023-01183-9

(A) Variation on biosecurity levels of the exposed farm (free)

(B) Variation on biosecurity levels of the source farm (infected)