Title:

Assessing the risk of highly pathogenic avian influenza H5 viruses spread within

French broiler and layer production networks.

Authors:

Claire Hautefeuille[1*], Facundo Muñoz[1], Gwenaëlle Dauphin[2], Mathilde Paul[3], Marisa

Peyre[1] and Flavie Goutard[1]

[1] CIRAD, UMR ASTRE, F-34398 Montpellier, France.

ASTRE, Univ Montpellier, CIRAD, INRAE, Montpellier, France

[2] CEVA Animal Health, Libourne, France

[3] IHAP, Université de Toulouse, INRAE, ENVT, Toulouse, France

*   Corresponding author

E-mail: claire.hautefeuille@cirad.fr (CH)

## Abstract

Since 2015, French poultry production is threatened almost every year by a reintroduction of highly pathogenic avian influenza H5 viruses. The duck sector was the most concerned by this crisis but other sectors such as broiler, layer and turkey were also affected by outbreaks. The objective of this work was to assess the risk of highly pathogenic avian influenza H5 virus transmission from one farm to another within the French broiler and layer production network.

This study used the WOAH risk assessment framework. After drawing up a scenario tree of virus transmission from one farm to another, data were collected through a literature review or through experts' elicitation. Three questionnaires were developed according to the experts' field of expertise: avian influenza, broiler and layer sectors. The experts' estimates were combined using a beta distribution weighted by their confidence level. A Monte Carlo iteration process was used to combine the different probabilities of the scenario tree and to assess the transmission risk.

In the broiler sector, the highest transmission probabilities were observed if the exposed farm was an indoor broiler farm and the source a broiler farm (indoor or free-range). The high transmission probability between broiler farm integrated within the same company highlighted the role of integration in this probability. In the layer sector, the highest transmission probabilities were observed if the source farm was a free-range farm and the exposed farm a production farm (indoor or free-range). The sensitivity analysis showed that the exposed farm's biosecurity had a significant impact on the transmission probability.

43    Our results provide an insight on the role of each type of farms in the virus spread

44    within the French broiler and layer production network and will be useful for the

45    implementation of control measures such as movement restriction or vaccination.

# Introduction

Since 2015, France has faced several epizootic waves of highly pathogenic avian

influenza (HPAI) viruses [1–3]. The main poultry production sector affected by these

waves has been the fattening duck production sector – for the production of duck liver-

based delicacies called "foie-gras" [4,5]. During these waves, while the initial

introduction of HPAI viruses has likely been due to wild birds [1,6], the large spread of

the viruses between poultry farms was linked with the poultry production systems. The

structure of the fattening duck sector was one of the main factors facilitating the spread

of HPAI [6]. Indeed, in this sector, live birds' movements occur between the different

production stages: rearing, breeding and force-feeding. It has been demonstrated that

these movements have played a major role in the spread of HPAI viruses [7,8].

Other poultry production sectors exist in France: the meat sector (broiler, meat duck,

turkey, guinea fowl) and layer sector [9]. In the broiler production sector, birds enter

the farm as day-old chicks and leave the farm to go to the slaughterhouse. In the layer

sector, production is organized in two stages: pullet (from day-old chick to ready-to-lay

layers) and layer (from ready-to-lay layers to spent hen). In between these two stages,

live bird movements take place to transport ready-to-lay layers from the pullet farm to

the layer farm. The same movement pattern is observed for breeders of all species. A

live bird movement occurs between the growth stage farm and the fertile adult stage

farm. Even if there are no live bird movements between broiler or layer farms, other

avian influenza (AI) virus pathways such, as fomite or human movements, may exist

[10]. Identified fomite movements may occur through birds pick-up to slaughter, feed

delivery, eggs collection, manure and litter management, dead bird management and

shared equipment [11–15]. Movements involving direct contacts between humans and

birds are identified as presenting the higher risk of AI virus spread (veterinarian, staff

working on multiple premises, company staff, technical advisors etc.) [16,17].

Few risk assessment studies looked at the impact of the integration level (integrated

or independent) or farming conditions (indoor vs. free-range) on the risk of AI viruses

introduction in a farm [13,15,18]. Moreover, a few modelling studies collected data on

the frequency of fomite and human movements between farms [11,14]. Nevertheless,

these studies never took a close look at the origin of these movements, i.e. type of

farm from which the movement originated.

So far, no study has been conducted on the risk of AI introduction into a chicken farm

from another farm within the chicken production sector in France, even though these

sectors have also been strongly affected by outbreaks (329 outbreaks in chicken farms

during the 2020-22 period) [2]. The aim of this study was to assess the risk of spread

of AI virus within two other French poultry production sectors: broiler and layer. This

risk assessment study has been conducted considering different types of farms as

described in a previous study [9]. These types of farms were defined according to the

sector (broiler and layer)), production type (i.e. future breeders, breeders, production),

the farming conditions (i.e. indoor and free-range) and the integration level of the farm.

For this purpose, this study assesses the risk of transmission from one farm of a

defined type to another farm of the same or different type within the same production

sector. It did not consider the transmission risk between farms from different poultry

sectors. The current study focuses on the relative transmission risk, i.e. to identify the

type of farms with the highest risk of virus transmission through indirect contacts

(human and fomites movements). This study will allow us to identify pathways and farm

types most at risk of AI virus transmission.

# Materials and methods

## Risk assessment model

This study used the risk analysis methodology described by the WOAH [19] to conduct

a partial risk assessment of the likelihood of an AI virus entering a flock. As the aim of

the study was to measure the relative risk of viral spread between different types of

farms, we focused on the risk of introduction. We assumed that once the virus is

introduced into the flock, the probability of a bird being infected, as well as the risk of

exposure, is close to 100%, and we did not assess the consequences of AI spread.

The virus introduction through the live bird movement pathway from an infected farm

to a free farm was considered very high. Since we have assumed that the introduction

of at least one infected bird into a free farm makes the farm infected, our risk

assessment study has focused solely on the risk of introduction through indirect

pathways within a poultry production network. These pathways were identified in a

previous literature scoping review [10]: human with a direct contact with birds; trucks

transporting live birds; trucks transporting feeds, manure and dead birds; shared

materials in direct contact with birds (Figure 1). Two other pathways known to

represent a very low risk and thus were not considered: human and shared materials

with no direct contact with birds or manure [11,15,20]. Environmental pathways such

as air-born or rodent as mechanical vector transmission pathways were not assessed

as the study was conducted under the hypothesis that these environmental pathways

would be similar between all types of farms. Geographical proximity between farms

was not considered either.

Figure 1: The transmission pathways considered in this risk-assessment study.

The assessment was conducted for the broiler and layer sectors on several types of

farms defined in Table 1 and adapted from the description provided by Hautefeuille et

al. [9]. Considering the low number of breeder farms and future layers farms, we did

not include the integration level for these types of farms.

Table 1: Listing of the different types of farms studied according to the production type,

the farming conditions and the integration level of farms in the French broiler and layer

production sector

| Production type | Farming conditions | Integration level |
|---|---|---|
| Broiler | Indoor | integrated with a hatchery |
| | | integrated without any hatchery |
| | | independent |
| | Free-range | integrated with a hatchery |
| | | integrated without any hatchery |
| | | independent |
| Breeder broiler | Indoor | - |
| Future breeder broiler | | - |
| Future layer | Indoor | - |

| | | integrated |
|---|---|---|
| Layer | Indoor | independent |
| | Free-range | integrated |
| | | independent |
| Breeder layer | Indoor | - |
| Future breeder layer | | - |

## Data source

The probability of virus introduction from each pathway was calculated using scenario

trees (Supplementary file 1), as recommended by EFSA [21]. The first step of this study

was to identify, in the scenario trees, the branches for which estimates of risk

probabilities could be obtained from the French legislation (Supplementary file 1). The

second step was to generate data for each pathway on the transmission risk, the

frequency of occurrence for one type of farm and the probability that this pathway exists

between two different types of farms. An expert elicitation survey was conducted to

obtain the data not available from the scientific literature.

## Experts' elicitation survey

The aim of this survey was to collect data on the transmission risk for each type of

farms (8 farm types for broiler and 7 for layer production) and for each pathway (9

different pathways). We decided to create a survey with three questionnaires, each of

which was a reasonable length for respondents while still collecting all of the necessary

data. Each questionnaire was intended for one type of expert profile: one for experts

on AI epidemiology and poultry farm biosecurity, one for experts on French broiler

sector and one for experts on French layer sector. In our study, experts were defined

as persons with relevant experience on one of the different topics studied.

Experts were recruited according to their experience on the different questionnaire

topics. Experts requested at first for the questionnaire on AI were scientists from

diverse national French research centres currently working and publishing on AI in

France. Experts requested at first for the questionnaire on broiler or layer sector are

national representatives of the French broiler or layer sector. Following this initial

recruitment, a "snowball" recruitment was conducted, in which experts who participated

in the survey were asked to recommend other experts to fill out the questionnaire.

Experts filled the survey independently without being in contact with each other. The

questionnaire was completed between June 1 and July 31, 2020.

The questionnaire was designed online using the tool Survey Monkey

(www.surveymonkey.com). For each questionnaire a pilot survey was sent to two

experts working on the questionnaire's field in order to test the survey and optimize it.

In the aim to reduce the length of each questionnaire, some simplifications were

decided and validated with these two experts (Table 2).

Table 2: Simplification made according to the probability and the pathway concerned

and justification validated by two experts of the poultry production sector.

| Probability concerned | Pathway concerned | Simplification made | Justification |
| --- | --- | --- | --- |

| Probability of virus introduction and release | Pullet transport Birds to slaughter transport | Merging of the two pathways as pick-up transport | Pick-up of birds is conducted in a similar way for pullet and birds to slaughter. |
|---|---|---|---|
| Probability that the studied pathway occurred during the previous week in farm A (i.e. Probability of pathway origin) | Manure Dead birds | Merge of the two pathways | Management similar for both pathways |
| | Shared materials | No consideration of the integration level of farms | Management similar regardless the integration level |

The experts were emailed a description of the study objectives and of what was

expected for their participation with the link to the online survey. The beginning of the

online survey included instructions and contact information of the administering

researchers. As this study was based solely on expert elicitation and did not involve

animal experimentation, it was not necessary to consult an animal research ethics

committee. Completion of the questionnaire constituted authorization to use the data

provided and informed consent. The survey was not anonymous in order to contact the

experts for clarification, but all their answers were analysed and presented

anonymously. Personal information was retained to identify the experts, but was not

used for the analysis presented in this work. The questionnaires are available in French

upon request.

Each questionnaire was designed to take about one hour to be filled by experts.

Questionnaires started with a description of the experts and questions on their

expertise level. The first questionnaire on AI epidemiology was divided in four sections:

risk of AI virus introduction in a farm, risk of AI virus release from an infected farm, risk

associated with wild birds, and risk of an infected farm going undetected. To simplify

the questionnaire, risk was assessed per farm biosecurity level rather than farm type.

To do so a biosecurity grid was built based on previous works on the biosecurity

evaluation process [7,22] and guidelines developed by the French poultry technical

institute (ITAVI) [23]. This grid was validated by two experts outside of the list of experts

identified for the risk assessment and currently working on French poultry farms

biosecurity. Five biosecurity levels were defined from very low to very high, and are

described in the biosecurity grid (Supplementary file 2). For the five defined biosecurity

levels, experts were asked about the probability of introduction or release of the virus

for each risk pathway (poultry production related or wild birds related). At the end, the

experts were asked to define the probability that an infected farm would go undetected.

The second questionnaire aimed to collect data on broiler sector and had four sections:

biosecurity level of poultry farms, frequency of between farm movements and material,

and detection level of AI virus infection. To assess the biosecurity level of a studied

farm type, experts were asked to distribute 100 farms of this type between the five

different levels in order to have the percentage of farms per biosecurity level

(Supplementary file 2). The second section aimed to identify for each introduction

pathway and for each type of farm, the frequency that this pathway occurred and the

probability that this pathway links the studied farm with another type of farms. The

questionnaire concluded on the capacity of a farmer to detect an AI infection based on

the type of farm. For question concerning probabilities, experts were asked for the most

probable, the minimum and the maximum values. For each question, experts were

asked to provide their confidence level with a score from 1 to 5 (Table 3) and they

could add comments. Finally, for the three questionnaires, the experts were asked

some open-ended questions regarding factors that might have influenced their

answers.

Table 3: Matching between confidence level and score

| Level of confidence | Score |
|---|---|
| Certain | 5 |
| Very confident | 4 |
| Moderately confident | 3 |
| Not confident | 2 |
| Uncertain | 1 |

## 208 **Data analysis**

Data collected from the SurveyMonkey platform were saved in a database template

developed in Microsoft Excel® version 2007. The data from these two questionnaires

were used to elicit Beta distributions characterising the ensemble of experts'

assessments on each probability of interest. For each probability of interest, the

parameters of the Beta distribution were determined in terms of expected value and

variance of the distribution, which were in turn calculated from the expert's responses

as follows. For the question asking for the most probable, minimum and maximum

values of the target probabilities, the expected value was set at the average of the most

probable values weighted by the expert confidence level and the variance was

calculated using a formula (1) balancing a maximal variance $V_m$ and an average

variance $V_i$ between experts.

$$var = V_i + \frac{(V_m - V_i)^3}{V_m^{\,2}} \qquad (1)$$

The maximal variance $V_m$ corresponds to the variance of a Uniform distribution with

support over the maximal interval (i.e. the smallest interval containing all the intervals

reported by the experts). The average variance $V_i$, by contrast, is the average of the

variances of the Uniform distributions over each of the reported intervals. This variance

calculation allows the representations of divergent opinions without reducing

compatible assessments. For questions requiring only one interval of values (i.e. 21%

to 40%, 41% to 60%, etc.), such as questions on between farm contact probabilities,

the centre of the interval was considered as the most probable, and the boundaries as

the minimum and the maximum value for each expert. The analysis was conducted

with R (version 4.0.2).

The frequency of occurrence and the origin of some pathways did not need to be

assessed by experts as these values depend on the production cycle. The egg

transport was considered to enter into a given breeder farm every day and the possible

origin of this truck was always breeder farms. The frequency of chick and pullet delivery

and slaughter take off was defined by the production cycle of the different types of

production. As it appears that experts could not provide an occurrence frequency for

shared material, it was arbitrarily considered to occur once a month.

A Monte Carlo iterative process was carried out on the scenario trees described above,

in order to combine the different probabilities and to calculate the transmission

probability from one type of farm to another. The probability of realisation of the event

for all branches of the scenario tree were randomly sampled from the Beta distribution

previously elicited. The simulation process used 10 000 iterations.

## 244    Sensitivity analysis

A sensitivity analysis was conducted using the one-at-time method applied on the

biosecurity level of the source and the exposed farms and on the pathways [24]. This

analysis was applied only for two types of farms with the most extreme daily

transmission probabilities: one representing the highest and one the lowest. This

sensitivity analysis is based on the distribution of the transmission probabilities on 10

000 iterations.

# 252   Results

## 253    Experts' description

At the end of the process, six experts responded to the questionnaire on AI

epidemiology. Among these experts, three had more than ten years of experience on

AI, one more than five years and the last two more than three years. Seven experts

responded to the questionnaire on the broiler sector and all of them had more than 10

years of experience on the broiler sector. Five experts responded to the questionnaire

on the layer sector and all of them had over 10 years of experience on the layer sector.

## Probability of avian influenza virus transmission

Based on the survey results, we found that, for the broiler production network, the highest transmission probabilities are observed when the source farm is a production farm (i.e. not selection) (indoor or free-range) and the exposed farm is an indoor broiler farm, regardless of the integration level (Table 4). Transmission probabilities are lowest when the exposed farm is a selection farm or when the source farm is a selection farm and the exposed farm a free-range farm (without consideration of the integration level).

For the layer production network, the highest transmission probabilities are observed when the source farm and the exposed farm are production farms (pullet, indoor or free-range layer) (Table 5). Transmission probabilities are lowest when the source farm and/or the exposed farm are selection farms.

Table 4: Daily transmission probabilities of HPAI H5 virus between an infected source farm and an exposed farm (free of infection) according to the type of farms in the French broiler production network (median, Q25 and Q75 values of 10 000 iterations). Percentage of transmission probabilities below the transmission probability between independent indoor broiler farms: Dark brown: 50-60%, brown: 60-75%, medium brown: 75-85%, light brown: 85-95%, no colour: over 95%.

| Values in % | | Exposed farm (free of infection) | | | | | | | |
| --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| | | Indoor broiler | | | Free-range broiler | | | Selection | |
| | | Integrated with hatchery | Integrated without hatchery | Independent | Integrated with hatchery | Integrated without hatchery | Independent | Breeder | Future breeder |
| Source farm (infected) | | | | | | | | | |
| Indoor broiler | Integrated with hatchery | 1.2 [0.5 - 3] | 0.8[0.3 - 2] | 1.0 [0.4 - 2.5] | 0.2 [0.1 - 0.4] | 0.2 [0.1 - 0.4] | 0.3 [0.1 - 0.6] | 0.007 [0.0009 - 0.04] | 0.005 [0.0005 - 0.04] |
| | Integrated without hatchery | 0.8 [0.3 - 2] | 1.2 [0.5 - 3.1] | 1.0 [0.4 - 2.5] | 0.2 [0.1 - 0.5] | 0.2 [0.1 - 0.5] | 0.3 [0.1 - 0.6] | 0.007 [0.0009 - 0.04] | 0.005 [0.0005 - 0.04] |
| | Independent | 0.6 [0.2 - 1.4] | 0.6 [0.2 - 1.4] | 1.2 [0.5 - 2.8] | 0.2 [0.1 - 0.5] | 0.3 [0.1 - 0.5] | 0.3 [0. 2 - 0.7] | 0.008 [0.001 - 0.05] | 0.006 [0.0006 - 0.04] |
| | Integrated with hatchery | 0.6 [0.2 - 1.5] | 0.5 [0.2 - 1.5] | 1.0 [0.4 - 2.4] | 0.4 [0.2 - 0.8] | 0.3 [0.1 - 0.6] | 0.4 [0.2 - 0.7] | 0.008 [0.001 - 0.05] | 0.006 [0.0006 - 0.04] |

| | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| Free-range broiler | Integrated without hatchery | 0.5 [0.2 - 1.5] | 0.5 [0.2 - 1.5] | 1.0 [0.4 - 2.4] | 0.3 [0.1 - 0.6] | 0.4 [0.2 - 0.8] | 0.4 [0.2 - 0.7] | 0.007 [0.001 - 0.04] | 0.006 [0.0006 - 0.04] |
| | Independent | 0.6 [0.3 - 1.7] | 0.7 [0.3 - 1.7] | 1.2 [0.5 - 2.8] | 0.3 [0.1 - 0.6] | 0.3 [0.1 - 0.6] | 0.5 [0.2 - 1] | 0.009 [0.001 - 0.05] | 0.007 [0.0007 - 0.05] |
| Selection | Breeder | 0.05 [0.009 - 0.2] | 0.05 [0.009 - 0.2] | 0.07 [0.02 - 0.3] | 0.01 [0.003 - 0.04] | 0.01 [0.003 - 0.04] | 0.02 [0.005 - 0.05] | 0.006 [0.0007 - 0.04] | 0.004 [0.0004 - 0.03] |
| | Future breeder | 0.04 [0.008 - 0.2] | 0.04 [0.008 - 0.2] | 0.06 [0.01 - 0.3] | 0.01 [0.002 - 0.04] | 0.01 [0.003 - 0.04] | 0.01 [0.004 - 0.04] | 0.003 [0.0003 - 0.02] | 0.006 [0.0006 - 0.04] |

Table 5: Daily transmission probabilities of HPAI H5 virus between an infected source farm and an exposed farm (free of infection) according to the type of farms in the French layer production network (median, Q25 and Q75 values of 10 000 iterations). Percentage of transmission probabilities below the transmission probability between independent free-range layer farms: Dark brown: 50-60%, brown: 60-75%, medium brown: 75-85%, light brown: 85-95%, no colour: over 95%

| Values in % | | Exposed farm (free of infection) | | | | | | |
| --- | --- | --- | --- | --- | --- | --- | --- | --- |
| | | Pullet | Indoor layer | | Free-range layer | | Selection | |
| Source farm (infected) | | | Integrated | Independent | Integrated | Independent | Breeder | Future breeder |
| Pullet | | 0.2 [0.06 - 0.5] | 0.1 [0.05 - 0.3] | 0.1 [0.05 - 0.4] | 0.1 [0.05 - 0.3] | 0.1 [0.04 - 0.3] | 0.007 [0.001 - 0.03] | 0.005 [0.0007 - 0.02] |
| Indoor layer | Integrated | 0.1 [0.04 - 0.4] | 0.2 [0.09 - 0.6] | 0.2 [0.08 - 0.5] | 0.2 [0.09 - 0.6] | 0.2 [0.09 - 0.6] | 0.007 [0.001 - 0.03] | 0.005 [0.0008 - 0.02] |
| | Independent | 0.1 [0.04 - 0.4] | 0.2 [0.07 - 0.5] | 0.3 [0.1 - 0.7] | 0.2 [0.09 - 0.6] | 0.2 [0.09 - 0.6] | 0.007 [0.001 - 0.03] | 0.005 [0.0008 - 0.02] |
| Free-range layer | Integrated | 0.2 [0.06 - 0.6] | 0.3 [0.1 - 0.8] | 0.4 [0.1 - 0.9] | 0.4 [0.2 - 1] | 0.4 [0.2 - 0.9] | 0.009 [0.002 - 0.04] | 0.006 [0.001 - 0.03] |
| | Independent | 0.2 [0.06 - 0.6] | 0.3 [0.1 - 0.8] | 0.4 [0.1 - 0.9] | 0.3 [0.1 - 0.8] | 0.4 [0.2 - 1] | 0.009 [0.00187 - 0.04] | 0.006 [0.0011 - 0.03] |
| Selection | Breeder | 0.005 [0.0008 - 0.02] | 0.009 [0.002 - 0.03] | 0.009 [0.002 - 0.03] | 0.008 [0.002 - 0.03] | 0.008 [0.002 - 0.03] | 0.006 [0.0008 - 0.04] | 0.004 [0.0004 - 0.02] |
| | Future breeder | 0.004 [0.0007 - 0.02] | 0.007 [0.002 - 0.03] | 0.007 [0.002 - 0.03] | 0.006 [0.002 - 0.02] | 0.006 [0.002 - 0.02] | 0.004 [0.0004 - 0.03] | 0.004 [0.0005 - 0.03] |

## Sensitivity analysis

The sensitivity analysis was conducted on two types of farms: one with the highest transmission probability (integrated indoor broiler farm) (

Figure 2 and Figure 3) and one with the lowest (future breeder farm for broiler production) as source (infected) and exposed (free) farms (Supplementary file 3).

The distribution of transmission probabilities for all pathways in both cases highlighted the importance of the farm biosecurity level on transmission probability. Indeed, it showed that the transmission probabilities are high for all pathways when biosecurity levels of both source and exposed farms are low, but have a lower impact when the biosecurity levels are high (Figure 2).

The biosecurity level of the source farm has less influence on the probability of transmission than the biosecurity level of the exposed farm on the transmission probability, as showed by the distribution of transmission probabilities. For all pathways, the variation in the biosecurity level of the source farm (infected) had a low impact on the variation of the transmission probabilities (Figure 3 A). On the other hand, the variation in the biosecurity level of the exposed farm had a significant impact on the variation in transmission probabilities (Figure 3 B). Indeed, transmission probabilities were high in exposed farms with biosecurity levels below (3 on 5), while they were very low in farms with high biosecurity level (above 4), for all pathways except person movement for both farm types and material movement for future breeder farms. The distributions of transmission probabilities were very similar between the two types of farms for all pathways except for the shared material. As their respective total transmission probabilities were different, this result reinforces the impact of biosecurity level on the transmission.

Figure 2: Distribution of the transmission probabilities across all **pathways** according

to the biosecurity level of the source farm (x axis) and to the biosecurity level of the

exposed farm (colours) with integrated indoor broiler farm as exposed and source

farms. Threshold at 1/100 (black), 1/1000 (red) and 1/10 000 (orange).

Figure 3: Distribution of the transmission probabilities (A) across all **biosecurity levels**

**of the exposed farm (free)** according to the pathways (x axis) and to the biosecurity

level of the source farm (infected) (colours) and (B) across all **biosecurity level of the**

**source farm (infected)** according to the pathways (x axis) and to the biosecurity level

of the exposed farm (free) (colours) with integrated indoor broiler farm as exposed and

source farms. Threshold at 1/100 (black), 1/1000 (red) and 1/10 000 (orange).

## Farm biosecurity and pathways in the broiler sector

The majority of integrated broiler farms (indoor and free-range) were scored with a

good biosecurity level (3 in 5) (53% of indoor and 45% of free range) (Supplementary

file 4**)**. Independent indoor broiler farms were scored with a higher biosecurity level

(43% at good level) than independent free-range broiler farms (42% at low level). For

both indoor and free-range farms, independent broiler farms were estimated to have a

lower biosecurity level than integrated farms. The pathways which generated the

highest transmission probabilities between two integrated indoor broiler farms with a

good biosecurity level are person movement, feed transport and manure management

(respective median value of 0.23%, 0.03% and 0.03%). The most frequent

transmission pathways for integrated broiler farms were person movement (median

occurrence frequency of 21%) and feed transport (median occurrence frequency of

30%). When considering an integrated indoor broiler farm as an exposed farm, the

probability of contact through person and feed movement was high if the source farm

was an indoor broiler farm integrated in the same farmers' association, medium if the

source farm was an indoor broiler farm integrated in another farmers' association and

low if the source farm was an independent indoor broiler farm, a free range broiler farm

or a selection farm (Figure 4).

The production farms with the lowest transmission probability were integrated free-

range broiler farms as exposed farm even if their biosecurity levels were equivalent to

biosecurity levels of integrated indoor broiler farms. The pathways with the highest

transmission probability between integrated free-range broiler farm at good biosecurity

level were material, feed and manure movement (respectively median value of 0.02%

and 0.01%). Person movement had a lower impact on the probability of transmission

in free-range farms than in indoor farms. The occurrence frequency of person

movement between integrated free-range broiler farms (median value of 62%) was

lower than between integrated indoor broiler farms (median value of 78%). For both

types of farms (integrated indoor and free-range), transport of birds from hatchery

(chicks) or to slaughter and dead bird management were the pathways with the lowest

impact on the transmission probabilities. Movement of birds from hatchery and to

slaughter had the lowest frequency of occurrence (associated with the production

cycle). Dead bird management was considered to have a high frequency of

occurrence, but this pathway is the only one that stops in the public area of the farm

(Supplementary file 1).

Figure 4: Frequency of pathway occurrence in integrated indoor broiler farm according

to the different types of pathways and the different type of farms as pathway source

(Broiler I int SA: Indoor farm integrated in the same farmer association, Broiler I int OA:

Indoor farm integrated in another farmer association, Broiler I ind: Indoor farm

independent, Broiler FR: Free-range broiler farm)

## Farm biosecurity and pathways in the layer sector

Indoor layer farms were estimated to have between very good biosecurity level and

good biosecurity level. Free-range farms were estimated to have between good

biosecurity level. Pullet farms were considered with a good biosecurity level (3 in 5)

(50%) (Supplementary file 4).

Due to their production, layer birds transport occurs at a lower frequency than broiler

production. Moreover, the pathway with the highest impact on transmission

probabilities was egg transport. When comparing the layer production farm with the

highest transmission probability (independent free-range layer farms with a low

biosecurity level) and the farm with the lowest transmission probability (integrated

indoor layer farms with a good biosecurity level), the pathways with the highest

transmission probabilities were for both egg transport, person movement and shared

material. Nonetheless, independent free-range farms had higher transmission

probabilities than integrated indoor farms (median transmission probability for egg

transport of 0.09% and 0.59% respectively). According to the five experts of the layer

sector, the transport of eggs was the pathway with the highest frequency of occurrence

for both indoor and free-range layer farms (median values of 45% and 30%

respectively). Person movements and shared material were just as frequent as the

other pathways but they are both associated with direct contact with birds.

## Farm biosecurity and pathways in selection farms

For both broiler and layer sectors, experts considered the biosecurity levels of selection

farms to be high to very high (4 to 5 out of 5), implying very low transmission

probabilities, except for person movement. As with future breeder farms (Figure 4),

shared material has no impact on transmission probability. If the results for breeder

farms in both productions were similar, eggs transport and manure management

appeared to have higher transmission probabilities in broiler production than in layer

production. If the transmission probabilities are very low at high and very high

biosecurity levels, they are closed to those observed in production farm for person

movement, feed and eggs transport, and manure management if the biosecurity level

of the breeder farms reached a good level (3 in 5).

# Discussion

This study looked at the risk of virus transmission within French production networks,

i.e. the risk that a virus released by an infected farm be introduced into a free farm by

all possible indirect pathways and for all types of farms described in the network

analysis. Most of risk assessment studies conducted on the risk of AI virus introduction

into a poultry farm are based on risk factors related to the production network and wild

birds [25,26]. Several risk assessment studies have been conducted on the risk of

introduction or release of AI viruses in different production types [14,15,18,20]. Few

studies looked at the risk of virus transmission between farms depending on farm types

[18,27]. None of them considered the farm integration levels and the risk of virus

transmission from one farm type of another (e.g. from an indoor broiler farm to a

breeder farm).

Another originality of this study is for the inclusion of biosecurity levels in the risk

assessment process. Biosecurity classifications developed for poultry farms generally

provide a farm-level biosecurity score [22,28,29]. Few studies have assessed

biosecurity level according to the farm type (breeder, broiler or layer) [30] or to the level

of farm integration [31,32] yet none have used this classification to assess the risk of

disease transmission per farm type. Our description of biosecurity levels was largely

based on previous work done in France [7,23] and it was adapted and validated with

poultry farm biosecurity specialists in France. The biosecurity criteria were illustrated

with a scheme (Supplementary file 1) and the experts expressed no difficulties to

understand the grid and the biosecurity classification. This approach based on farm

biosecurity enabled to study a large number of farm types. The risk assessment

conducted on biosecurity level can be used to assess the risk of virus transmission

between farms in any other French poultry production network. Likewise, the

developed biosecurity grid can be used to assess the biosecurity level of any type of

French poultry farm.

The farm biosecurity level of the exposed farm (free of infection) has a major impact

on the AI virus between-farm transmission risk regardless the pathway and the

biosecurity level of the source farm (infected). The importance of farm biosecurity on

AI virus introduction risk has been highlighted in several studies [33–36]. In this study,

farms dedicated to reproductive birds (future breeder and breeder farms) were mainly

scored 5, indicating very high biosecurity, while production farms scored 3, indicating

good biosecurity. Less than 5% of the production farms were considered to be at a

very high biosecurity level. A study conducted in New Zealand found a similar result

[28]. Another study in South Korea that measured the risk of HPAI for farms linked by

livestock vehicle movement showed a low risk for 73% of breeder chicken farms [18].

Both free-range and indoor integrated broiler farms were considered with an overall

good biosecurity level. In contrast, indoor layer farms were considered having a higher

biosecurity (level 3 (good) to level 4 (high)) than free-range layer farms (level 2 (low)

to level 3 (good)). As biosecurity measures to limit contact between domestic birds and

wild birds were included in the biosecurity grid to assess the farm biosecurity level, the

impact of free-range was considered in this assessment. This result is interesting as

risk assessment studies in other countries have shown that free-range farms are more

at risk of virus introduction by wild birds [26,37]. A serological study conducted in

France showed that free-range farms were less infected by HPAI than indoor farms,

and that biosecurity measures applied in these farms contributed to reduce the risk of

virus introduction and/or exposure [38]. Wild birds are known to have a role in HPAI

virus introduction and long-distance spread but their role in short distance

dissemination is limited [39,40]. The proportion of broiler farms with a low biosecurity

level (level 2) was higher in independent farms (40% of indoor and 42% of free-range)

that in integrated farms (15% of indoor and 22% of free-range). This result is difficult

to compare with other studies because these studies were conducted in countries with

very different production context (Indonesia and Thailand) [31,32]. Moreover, the

difference observed must be considered carefully as expert confidence was higher for

integrated farms (mean of 4.2 and 3.9 out of 5 for indoor and free-range farms

respectively) than for independent farms (mean of 3 and 3.4 out of 5 for indoor and

free-range farms respectively). Similar results were observed for independent free-

range layer farms but not for independent indoor layer farms for which the proportion

of farms with high biosecurity level (mean value of 43%) was higher than the proportion

of with good biosecurity level (mean value of 37%). Nonetheless, only 3 experts out of

5 provided data for this type of farms (mean confidence of 3.7 out of 5) which illustrated

the lack of knowledge for this minority farm type. This work is based on expert opinion

and highlight the need for more detailed data on farm biosecurity. A study carried out

in France provided information on biosecurity as a function of farm type (including

broiler and layer) and location, but did not investigate independent farms [41].

Moreover, this biosecurity level assessment has been conducted at the national level

and do not consider farm spatial heterogeneity in France.

This risk assessment study aimed to investigate all the links between all farm types in

the French broiler and layer production networks and thereby identify the type of farms

and links associated with the highest AI transmission risk. The links studied only

concerned indirect transmission pathways (i.e., humans or fomite) and not direct

transmission pathways (i.e., live domestic birds). Live domestic bird movement is

known to present high risk of virus transmission, especially in the duck sector where

live duck movement occurs between different growing stages [8]. Nonetheless, this

pathway is inexistant in broiler farms or at very low frequency in layer and breeder

farms (about once every 12-13 months). This is the reason why this study focused on

indirect contact which are much more likely to occur during an AI epidemic. In the

current study, we considered that fomite (i.e. person, truck or material) management

outside of farms was similar for all types of farms and was not sufficient to affect the

viral transmission. The disinfection of fomites inside farms is included in the biosecurity

classification. If the impact of fomite disinfection outside of farms is the same for all

types of farms, this impact will not change the relative risk. Nonetheless, a good

cleaning and disinfection process at the truck level would considerably reduce the

transmission risk between farms. Studies conducted in France detected the presence

of the viral genome in half (in 2017) and in a third (in 2021) of trucks tested after

disinfection [42,43].

Person movement was the pathway with the highest transmission probabilities. This

pathway is associated with direct contact with birds and high occurrence frequency.

The other pathways including direct contact with potentially infectious substances such

as manure or dead birds or shared material in direct contact with birds were associated

with high transmission probabilities. Previous works obtained similar results [11,15,20].

Feed transport was also associated with high transmission probabilities. Our study

focused on the transmission pathways described as the most at risk of transmission in

the literature [10]. Nonetheless, other pathways such as transmission by wind or

vermin (i.e. mechanic vectors such as rodents or insects) are also studied in the

literature. Air-borne transmission was found to have a limited role in AI viruses

transmission in France [44], even though this transmission is possible at short distance

as some virus genome was found in air samples of infected farms [45]. Dust seems to

have also a role in viral dispersal [46]. Virus transmission through vermin movements

between farms is possible but difficult to distinguish from other environmental factors

[25,27,47]. These last pathways can be included in an environmental transmission

pathway, which is mainly due to high proximity between farms. High poultry density is

known to have an impact in the spread of AI viruses [48]. Our study compared the

relative transmission risk from one farm to another and was conducted under the

hypothesis that environmental transmission pathway will occur in the same way for all

types of farms regardless their biosecurity level. Environmental transmission could

have an impact on short distance spread and should be considered for studies

integrating geographical information, which is not the case for the current work.

In the broiler sector, the transmission risk was higher when exposed farms (free) were

indoor farms compared to other type of farms. This is mainly due to the higher

frequency that a pathway occurs in this type of farm compared to free-range farms, as

previously identified [30]. This transmission risk was the highest between broiler farms

integrated within the same farmers' association. The probability of contact through a

given pathway was the highest between these farms, especially for person in direct

contact with birds and feed truck pathways. This result can be easily explained by the

integration structure in France. Indeed, poultry farmers' associations are based on a

feed manufacturer and technicians and veterinarians are often employed by this farmer

association. The role of the integration of in-between farm spread of AI was highlighted

in previous studies [11,13,16]. Contact frequency was higher between farms of the

same type (i.e. indoor, free-range and breeder) than between farms of different types.

According to experts, fewer contacts occurred between integrated and independent

farms, especially for indoor broiler farms. In the layer sector, the transmission

probability was higher for free-range layer farms as source (infected), related to the

farm biosecurity level. This result confirmed that egg transport have a major role in the

transmission probability [18,27]. Even if breeder farms have a very high biosecurity

level, the high frequency of pathways increase the risk of virus transmission between

breeder farms. Indeed, our result showed that if the biosecurity level of a breeder farm

as exposed farm is reduced to a good biosecurity level, the transmission probability is

similar to the one for production farms. This transmission risk in breeder farms is

confirmed by previous studies [30,33]. The study of HPAI H5 outbreaks between 2020

and 2022 in France showed that at least 9% of outbreaks occurred in a breeder farm

[2].

This work provides a full picture of the risk of AI virus spread within French broiler and layer production networks considering the virus transmission risk between any farm types. To mitigate this risk, management measures should focus on biosecurity, especially for the movements that enter the farm. A careful attention should be paid to movement with a high occurrence frequency such as person movement, feed, manure and dead bird transport. Integrator should be aware of the increased risk between farm within the same integration and take actions, such as cleaning and disinfection process between farms, to reduce this risk. This work provides a typology of farms based on the transmission risk which will be useful for the implementation of control measures such as movement restriction or vaccination.

## Acknowledgements

The authors gratefully acknowledge all the experts who answered the questionnaire.

## Funding

This work was co-funded by Ceva Santé Animale (https://www.ceva.com ) and Cirad (https://www.cirad.fr/ ) within the framework of a public private partnership PhD funding (Cifre funding n° 2017-1177 from Association Nationale de la Recherche et de la Technologie (ANRT: https://www.anrt.asso.fr/fr/accueil-anrt )) and within the BioFluARN project framework, co-funded by Banque Publique d'Investisement France (BPIFrance: https://www.bpifrance.fr/ ). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

## Conflict of interest disclosure

The authors declare that they have no financial or non-financial conflicts of interest in relation to the content of the article.

# References

1. Briand F-X, Schmitz A, Ogor K, Prioux AL, Guillou-Cloarec C, Guillemoto C, et al. Emerging highly pathogenic H5 avian influenza viruses in France during winter 2015/16: phylogenetic analyses and markers for zoonotic potential. Eurosurveillance. 2017;22: 30473. doi:10.2807/1560-7917.ES.2017.22.9.30473

2. Lambert S, Durand B, Andraud M, Delacourt R, Scoizec A, Le Bouquin S, et al. Two major epidemics of highly pathogenic avian influenza virus H5N8 and H5N1 in domestic poultry in France, 2020–2022. Transboundary and Emerging Diseases. 2022. doi:10.1111/tbed.14722

3. Napp S, Majó N, Sánchez-Gónzalez R, Vergara-Alert J. Emergence and spread of highly pathogenic avian influenza A(H5N8) in Europe in 2016-2017. Transboundary and Emerging Diseases. 2018;65: 1217–1226. doi:10.1111/tbed.12861

4. Bronner A, Niqueux E, Schmitz A, Bouquin SL, Huneau-Salaün A, Guinat C, et al. Description de l'épisode d'influenza aviaire hautement pathogène en France en 2016-2017. Bulletin épidémiologique, santé animale et alimentation. 2017;79: 13–17.

5. Le Bouquin S, Scoizec A, Niqueux E, Schmitz A, Briand F-X. L'épisode d'influenza aviaire en France en 2015-2016 – Situation épidémiologique au 30 juin 2016. Bulletin épidémiologique, santé animale et alimentation. 2016;75: 7.

6. Guinat C, Artois J, Bronner A, Guérin JL, Gilbert M, Paul MC. Duck production systems and highly pathogenic avian influenza H5N8 in France, 2016–2017. Scientific Reports. 2019;9. doi:10.1038/s41598-019-42607-x

7. Delpont M, Blondel V, Robertet L, Duret H, Guerin J-L, Vaillancourt J-P, et al. Biosecurity practices on foie gras duck farms, Southwest France. Preventive Veterinary Medicine. 2018;158: 78–88. doi:10.1016/j.prevetmed.2018.07.012

8. Guinat C, Durand B, Vergne T, Corre T, Rautureau S, Scoizec A, et al. Role of Live-Duck Movement Networks in Transmission of Avian Influenza, France, 2016-2017. Emerging infectious diseases. 2020;26: 472–480. doi:10.3201/eid2603.190412

9.　Hautefeuille C, Azzouguen B, Mouchel S, Dauphin G, Peyre M. Evaluation of
vaccination strategies to control an avian influenza outbreak in French poultry
production networks using EVACS tool. Preventive Veterinary Medicine. 2020.

10.　Hautefeuille C, Dauphin G, Peyre M. Knowledge and remaining gaps on the role
of animal and human movements in the poultry production and trade networks in
the global spread of avian influenza viruses - A scoping review. PLoS ONE.
2020;15. doi:10.1371/journal.pone.0230567

11.　Dorea FC, Vieira AR, Hofacre C, Waldrip D, Cole DJ. Stochastic Model of the
Potential Spread of Highly Pathogenic Avian Influenza from an Infected
Commercial Broiler Operation in Georgia. Avian Diseases. 2010;54: 713–719.
doi:10.1637/8706-031609-ResNote.1

12.　Hagenaars TJ, Boender GJ, Bergevoet RHM, van Roermund HJW. Risk of poultry
compartments for transmission of highly pathogenic avian influenza. PLoS ONE.
2018;13. doi:10.1371/journal.pone.0207076

13.　Leibler JH, Carone M, Silbergeld EK. Contribution of Company Affiliation and
Social Contacts to Risk Estimates of Between-Farm Transmission of Avian
Influenza. PLOS ONE. 2010;5: e9888. doi:10.1371/journal.pone.0009888

14.　Longworth N, Mourits MCM, Saatkamp HW. Economic Analysis of HPAI Control
in the Netherlands I: Epidemiological Modelling to Support Economic Analysis.
Transboundary and Emerging Diseases. 2014;61: 199–216.
doi:10.1111/tbed.12021

15.　Singh M, Toribio JA, Scott AB, Groves P, Barnes B, Glass K, et al. Assessing the
probability of introduction and spread of avian influenza (AI) virus in commercial
Australian poultry operations using an expert opinion elicitation. PLoS ONE.
2018;13.

16.　Dent JE, Kiss IZ, Kao RR, Arnold M. The potential spread of highly pathogenic
avian influenza virus via dynamic contacts between poultry premises in Great
Britain. BMC Veterinary Research. 2011;7: 59. doi:10.1186/1746-6148-7-59

17.　Sharkey KJ, Bowers RG, Morgan KL, Robinson SE, Christley RM.
Epidemiological consequences of an incursion of highly pathogenic H5N1 avian
influenza into the British poultry flock. Proceedings of the Royal Society of London
B: Biological Sciences. 2008;275: 19–28. doi:10.1098/rspb.2007.1100

18.　Yoon H, Lee I, Kang H, Kim K-S, Lee E. Big data-based risk assessment of poultry
farms during the 2020/2021 highly pathogenic avian influenza epidemic in Korea.
PLoS One. 2022;17: e0269311. doi:10.1371/journal.pone.0269311

19.　OIE. Chapter 2.1. Import risk Analysis. Terrestrial Animal Health Code. 2019.
Available:
https://www.oie.int/fileadmin/Home/eng/Health_standards/tahc/current/chapitre_i
mport_risk_analysis.pdf

20.　Ssematimba A, Hagenaars TJ, de Wit JJ, Ruiterkamp F, Fabri TH, Stegeman JA,
et al. Avian influenza transmission risks: Analysis of biosecurity measures and

contact structure in Dutch poultry farming. Preventive Veterinary Medicine.
2013;109: 106–115. doi:10.1016/j.prevetmed.2012.09.001

21. EFSA Panel on Animal Health and Welfare (AHAW). Guidance on Risk
Assessment for Animal Welfare: Guidance on risk assessment for animal welfare.
EFSA Journal. 2012;10: 2513. doi:10.2903/j.efsa.2012.2513

22. Gelaude P, Schlepers M, Verlinden M, Laanen M, Dewulf J. Biocheck.UGent: A
quantitative tool to measure biosecurity at broiler farms and the relationship with
technical performances and antimicrobial use. Poultry Science. 2014;93: 2740–
2751. doi:10.3382/ps.2014-04002

23. ITAVI. Arbre de decision: Toutes les fiches relatives aux mesures de biosécurité
IA. Mar 2019 [cited 27 May 2020]. Available: http://influenza.itavi.asso.fr/

24. Saltelli A, Annoni P. How to avoid a perfunctory sensitivity analysis. Environmental
Modelling & Software. 2010;25: 1508–1517. doi:10.1016/j.envsoft.2010.04.012

25. Gustafson L, Jones R, Dufour-Zavala L, Jensen E, Malinak C, McCarter S, et al.
Expert elicitation provides a rapid alternative to formal case-control study of an
H7N9 avian influenza outbreak in the United States. Avian Diseases. 2018;62:
201–209. doi:10.1637/11801-011818-Reg.1

26. Grabkowsky BJ, Windhorst H-W. Investigation and quantification of prevalent risk
factors for the introduction of avian influenza in European poultry farms. World's
Poultry Science Journal. 2010;66: 381–398. doi:10.1017/S0043933910000486

27. Scott AB, Toribio J-ALML, Singh M, Groves P, Barnes B, Glass K, et al. Low- and
High-Pathogenic Avian Influenza H5 and H7 Spread Risk Assessment Within and
Between Australian Commercial Chicken Farms. Front Vet Sci. 2018;5.
doi:10.3389/fvets.2018.00063

28. Greening SS, Mulqueen K, Rawdon TG, French NP, Gates MC. Estimating the
level of disease risk and biosecurity on commercial poultry farms in New Zealand.
New Zealand Veterinary Journal. 2020;68: 261–271.
doi:10.1080/00480169.2020.1746208

29. Van Limbergen T, Dewulf J, Klinkenberg M, Ducatelle R, Gelaude P, Méndez J,
et al. Scoring biosecurity in European conventional broiler production. Poultry
Science. 2018;97: 74–83. doi:10.3382/ps/pex296

30. Van Steenwinkel S, Ribbens S, Ducheyne E, Goossens E, Dewulf J. Assessing
biosecurity practices, movements and densities of poultry sites across Belgium,
resulting in different farm risk-groups for infectious disease introduction and
spread. Preventive Veterinary Medicine. 2011;98: 259–270.
doi:10.1016/j.prevetmed.2010.12.004

31. Indrawan D, Cahyadi ER, Daryanto A, Hogeveen H. The role of farm business
type on biosecurity practices in West Java broiler farms. Preventive Veterinary
Medicine. 2020;176. doi:10.1016/j.prevetmed.2020.104910

32. Wei H, Aengwanich W. Biosecurity Evaluation of Poultry Production Cluster
(PPCs) in Thailand. International Journal of Poultry Science. 2012;11: 582–588.
doi:10.3923/ijps.2012.582.588

33. Duvauchelle A, Huneau-Salaün A, Balaine L, Rose N, Michel V. Risk factors for
the introduction of avian influenza virus in breeder duck flocks during the first 24
weeks    of    laying.    Avian    Pathology.    2013;42:    447–456.
doi:10.1080/03079457.2013.823145

34. Guinat C, Comin A, Kratzer G, Durand B, Delesalle L, Delpont M, et al. Biosecurity
risk factors for highly pathogenic avian influenza (H5N8) virus infection in duck
farms,    France.    Transboundary    and    Emerging    Diseases.    2020.
doi:10.1111/tbed.13672

35. McQuiston JH, Garber LP, Porter-Spalding BA, Hahn JW, Pierson FW,
Wainwright SH, et al. Evaluation of risk factors for the spread of low pathogenicity
H7N2 avian influenza virus among commercial poultry farms. Journal of the
American    Veterinary    Medical    Association.    2005;226:    767–772.
doi:10.2460/javma.2005.226.767

36. Nishiguchi A, Kobayashi S, Yamamoto T, Ouchi Y, Sugizaki T, Tsutsui T. Risk
Factors for the Introduction of Avian Influenza Virus into Commercial Layer
Chicken Farms During the Outbreaks Caused by a Low-Pathogenic H5N2 Virus
in    Japan    in    2005.    Zoonoses    and    Public    Health.    2007;54:    337–343.
doi:https://doi.org/10.1111/j.1863-2378.2007.01074.x

37. Scott AB, Toribio J-A, Singh M, Groves P, Barnes B, Glass K, et al. Low
Pathogenic Avian Influenza Exposure Risk Assessment in Australian Commercial
Chicken Farms. Front Vet Sci. 2018;5: 68. doi:10.3389/fvets.2018.00068

38. Huneau-Salaün A, Niqueux E, Protino J, Guyot M. Situation sanitaire favorable
vis-a-vis de l'influenza aviaire des poulets label rouge et biologiques en France
en 2018. Bulletin épidémiologique, santé animale et alimentation. 2020 [cited 27
Jan 2021]. Available: https://be.anses.fr/sites/default/files/O-042_2020-11-30_IA-
LRbio_Huneau_MaqF.pdf

39. Briand F-X, Niqueux E, Schmitz A, Martenot C, Cherbonnel M, Massin P, et al.
Multiple independent introductions of highly pathogenic avian influenza H5 viruses
during the 2020–2021 epizootic in France. Transboundary and Emerging
Diseases. 2022;69: 4028–4033. doi:10.1111/tbed.14711

40. Guinat C, Agui CV, Vaughan TG, Scire J, Pohlmann A, Staubach C, et al.
Disentangling the role of poultry farms and wild birds in the spread of highly
pathogenic avian influenza virus H5N8 in Europe. bioRxiv; 2021. p.
2021.10.22.465255. doi:10.1101/2021.10.22.465255

41. Delpont M, Guinat C, Guérin J-L, Le leu E, Vaillancourt J-P, Paul MC. Biosecurity
measures in French poultry farms are associated with farm type and location.
Preventive    Veterinary    Medicine.    2021;195:    105466.
doi:10.1016/j.prevetmed.2021.105466

42. Huneau-Salaün A, Scoizec A, Thomas R, Martenot C, Schmitz A, Pierre I, et al. Avian influenza outbreaks: evaluating the efficacy of cleaning and disinfection of vehicles and transport crates. Poultry Science. 2022;101. doi:10.1016/j.psj.2021.101569

43. Huneau-Salaün A, Scoizec A, Thomas R, Le Bouquin S. Cleaning and disinfection of crates and trucks used for duck transport: field observations during the H5N8 avian influenza outbreaks in France in 2017. Poultry Science. 2020. doi:10.1016/j.psj.2019.10.015

44. Guinat C, Rouchy N, Camy F, Guérin JL, Paul MC. Exploring the wind-borne spread of highly pathogenic avian influenza H5N8 during the 2016-2017 epizootic in France. Avian Diseases. 2019;63: 235–237. doi:10.1637/11881-042718-ResNote.1

45. Scoizec A, Niqueux E, Thomas R, Daniel P, Schmitz A, Le Bouquin S. Airborne detection of H5N8 highly pathogenic avian influenza virus genome in poultry farms, France. Frontiers in Veterinary Science. 2018;5. doi:10.3389/fvets.2018.00015

46. Filaire F, Lebre L, Foret-Lucas C, Vergne T, Daniel P, Lelièvre A, et al. Highly Pathogenic Avian Influenza A(H5N8) Clade 2.3.4.4b Virus in Dust Samples from Poultry Farms, France, 2021. Emerging infectious diseases. 2022;28: 1446–1450. doi:10.3201/eid2807.212247

47. Velkers FC, Blokhuis SJ, Veldhuis Kroeze EJB, Burt SA. The role of rodents in avian influenza outbreaks in poultry farms: A review. Veterinary Quarterly. 2017;37: 182–194. doi:10.1080/01652176.2017.1325537

48. Bauzile B, Durand B, Lambert S, Rautureau S, Fourtune L, Guinat C, et al. Impact of palmiped farm density on the resilience of the poultry sector to highly pathogenic avian influenza H5N8 in France. Vet Res. 2023;54: 56. doi:10.1186/s13567-023-01183-9

(A) Variation on biosecurity levels of the exposed farm (free)

(B) Variation on biosecurity levels of the source farm (infected)