Bimodal nonlinear dendrites in PV+ basket cells drive distinct memory-related oscillations

Alexandra Tzilivaki[1,2,3 *], Matthew Evan Larkum[1,2,3,4], Dietmar Schmitz[1,2,3,5,6, * #]

[1]Charité-Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität Berlin, and Berlin Institute of Health, Neuroscience Research Center, 10117 Berlin, Germany

[2]Einstein Center for Neurosciences, Chariteplatz 1, 10117 Berlin, Germany

[3]NeuroCure Cluster of Excellence, Chariteplatz 1, 10117 Berlin, Germany

[4]Department of Biology, Humboldt Universität zu Berlin, Berlin 10117, Germany

[5]German Center for Neurodegenerative Diseases (DZNE), 10117 Berlin, Germany

[6]Bernstein Center for Computational Neuroscience, Humboldt-Universität zu Berlin, Philippstrasse. 13, 10115 Berlin, Germany

* Correspondence: AT alexandra.tzilivaki@charite.de , DS dietmar.schmitz@charite.de

Lead contact author: DS dietmar.schmitz@charite.de

Abstract

Neuronal oscillations are crucial for organizing distinct memory stages and behavioral states, yet the precise cellular mechanisms through which interneurons shape these dynamics remain incompletely understood. We propose and computationally explore a novel link between the nonlinear dendritic integration modes of parvalbumin-positive fast-spiking basket cells (PV+ FSBCs) and the modulation of hippocampal oscillations. Employing biophysical circuit level modeling, we test the hypothesis that PV+ FSBCs can flexibly adapt their firing patterns and influence rhythms, independent of gross changes in synaptic input quantity, by selectively engaging either their supralinear or sublinear dendritic branches. Specifically, supralinear dendrites promote high-frequency oscillations and reduce the excitation/inhibition (E/I) balance in the circuit, whereas sublinear dendrites enhance slow oscillatory power and elevate the E/I balance. This bimodal dendritic strategy provides PV+ FSBCs with an energy efficient mechanism to regulate circuit and oscillatory dynamics without necessitating large scale increases in synaptic drive. Our findings thereby uncover a previously unrecognized role for PV+ FSBC dendritic computations in regulating memory-related oscillations, offering new, experimentally testable hypotheses into the subcellular mechanisms that govern hippocampal rhythm generation.

Main

Activity in the mammalian hippocampus is characterized by distinct frequency bands that are essential for various memory processes[1]. Neuronal activity commonly correlates with local field potential (LFP) oscillations, including slow (theta 3–10 Hz), and fast (gamma range 30–200 Hz) bands [1–5]. Experimental and computational work has linked these slow and fast components to different stages of memory processing and distinct behavioral states. In particular, slow oscillations are associated with memory encoding, spatial navigation, and sensory processing[4], while fast oscillations are implicated in memory recall and consolidation [4,6]. These components frequently co-occur through oscillatory coupling, a phenomenon considered as a signature of working memory (**Figure 1a**) [5,7–11]. To understand

the cellular mechanisms behind these oscillatory events, significant efforts have been made
to delineate the effects of various types of both excitatory and inhibitory hippocampal
populations. Among them, PV+ interneurons have emerged as particularly crucial for
orchestrating both slow and fast oscillations, as well as their coupling. [9,12–17]. Nevertheless,
prevailing research has largely focused on the activation or silencing of PV+ interneurons,
often through optogenetic or broad computational manipulations. Consequently, the specific
subcellular properties, such as dendritic processing, that can empower these interneurons to
exert precise control over distinct oscillatory states and memory processes, remain largely
unexplored.

A long-standing dogma postulated that GABAergic interneurons, including PV+ cells,
possess passive, purely linear dendrites incapable of nonlinear synaptic integration. This
view suggested that their influence was determined only by the total quantity of synaptic
inputs they receive, implying that circuit regulation during oscillations depended on
aggregate synaptic drive without considering complex, potentially energy-efficient, nonlinear
dendritic processing. Challenging this paradigm, recent experimental work has demonstrated
that hippocampal PV+ interneurons, exhibit sophisticated nonlinear dendritic computations
that significantly augment their computational power and can maximize their input-output
(I/O) transformations. [18–21]. Specifically, in an earlier *in vitro* work, Katona and colleagues
reported that PV+ interneuron dendrites could express nonlinear summation[22]. Subsequent
biophysical computational modeling of hippocampal and cortical PV+ FSBCs (one of the 3
major subtypes of PV+ interneurons), proposed that their dendritic trees distinctly feature
coexisting supralinear and sublinear domains[23] (**Figures 1a, S2a-d**). This predicted bimodal
nonlinear processing, where PV+ FSBCs trees can leverage both nonlinear integration
types, highlighted morphological and functional distinctions despite similar active properties.
Supralinear branches typically possess larger volumes and lower input resistance and are
capable of local generation of spikes. The smaller, high-resistance sublinear branches do not
generate local spike events[23,24,26]. Notably, the core principle of this theoretical prediction,
that individual PV+ neurons can possess dendrites with distinct integration rules, was later
experimentally supported. Using *in vitro* 2-photon glutamate uncaging, Cornford and
colleagues 21 showed the coexistence of both supralinear and (sub)linear integration within
the CA1 PV+ cells whose somata locate in striatum pyramidale layer and their dendritic
branches can reach striatum oriens and lacunosum moleculare layers. This experimental
work, confirmed that PV+ cell dendrites are not functionally uniform, and is in strong
agrrement with the modeling predictions [23]. Thus, it further emphasized the potentially
prominent role of dendritic nonlinearities in PV+ regulation in circuit dynamics across
hippocampus. Although further experimental work is needed to charachterize the dendritic
processing rules in other types of intereurons in the hippocampus and beyond, recent work
in the L2/3 PV+ and SOM+ populations also indicates diverse nonlinear dendritic
strategies[25].

Building on these insights, the present theoretical study was motivated by fundamental
questions regarding the local circuit level impact of hippocampal PV+ FSBCs dendritic
nonlinearities during memory-related oscillations: How might PV+FSBCs leverage such
properties to influence circuit dynamics? What advantages could bimodal, as opposed to
unimodal, dendritic integration confer? And what are the distinct consequences of
preferentially activating supralinear versus sublinear dendritic domains? Given the
established role of PV+ FSBCs in both slow and fast oscillations, [12,27,28]we employed
advanced biophysical modeling to test our central hypothesis: that the supralinear and
sublinear dendrites of PV+ FSBCs may differentially modulate their activity and,
consequently, orchestrate both spiking activity and LFPs during mnemonic oscillations.
Specifically, we posited that PV+ FSBCs could exploit their bimodal dendritic architecture to

selectively shape fast and slow oscillatory activity within a hippocampal microcircuit (**Figures**
1-4). Such a mechanism would enable them to fine tune the E/I balance and modulate both
slow and fast oscillatory signals without necessitating gross changes in synaptic input
quantity (**Figures 3-4)**, thereby offering a flexible and energy-efficient strategy for engaging
with distinct memory-related rhythms.

To test our hypothesis, we utilized an established biologically plausible hippocampal
microcircuit model [29], that incorporates compartmental biophysical models of PCs with
simplified morphologies [29] and PV+ FSBCs with anatomically reconstructed bimodal
nonlinear dendritic trees [23] , mimicking the experimental phenotype of [21] (**Tables S1–S5,**
Figures. 1a, S1, S2). Both neuronal types were extensively calibrated based on
electrophysiological data (**Tables S1-S5, Figure S1**). In light of the limited experimental
validation currently available for dendritic integration mechanisms in other hippocampal
interneuron types (e.g., SOM+ or VIP+), and to avert speculative assumptions, here we
focus in hippocampal PV+FSBCs with their characteristic bimodal nonlinear dendritic
architecture (**Figure 1a**) as the sole inhibitory component within the microcircuit under
investigation.

Upon activation with a theta-like input (as per [30,31]), our model recapitulated the
experimentally observed coupling of slow and fast oscillations (**Figure 1a**) [9,10,32], generating
a slow oscillatory component alongside a fast oscillatory component (**Figure. 1a, d**). To
systematically investigate how PV+ FSBCs dendritic integration modes influence network
dynamics, we initially designed the following distinct activation conditions while ensuring that
the total number of synaptic contacts remained constant. 1. *Supralinear Activation (shown in*
purple): To understand the effects of supralinear dendrites in these cells, we restricted all
synaptic targeting exclusively to supralinear branches (**Figures. 1b,1d,1f,2b**). 2. *Sublinear*
Activation (shown in cyan): Similarly, to examine the role of sublinear dendritic processing,
we constrained all synaptic inputs to sublinear branches (**Figures. 1c,1e,1g, 2c**), and 3.
Bimodal Activation (shown in grey): Synapses onto PV+ FSBCs trees were distributed
across both supralinear and sublinear dendritic branches (**Figure 2a**). This represents a
mixed scenario where both types of dendrites can be targeted. In all three conditions, the
total number of synaptic inputs remained the same and identical (**Figures 1-3**), and all other
parameters and simulation settings were kept unchanged to ensure a controlled comparison
of how supralinear vs sublinear dendrites shape PV+ FSBCs firing patterns and network
oscillatory behavior. Our model predicts that supralinear dendritic activation in PV+ FSBCs
preferentially amplifies the fast oscillatory component of the *in silico* LFP signals, whereas
sublinear dendritic activation enhances the slow LFP component (**Figures 1-2**). Specifically,
when supralinear PV+ dendrites are activated, the LFP exhibits much higher fast peak-
frequency values as well as higher fast peak-frequency power values compared to sublinear
activation conditions (**Figures, 1-2**). In sharp contrast, PV+ FSBCs sublinear dendritic
activation significantly reduces the fast peak-frequency power relative to supralinear
activation (**Figures. 1-2**), while simultaneously increasing slow LFP power values (**Figures.**
1g, 2f). These predictions suggest that PV+ FSBCs can leverage their complex dendritic
architecture to selectively regulate distinct oscillatory components, dynamically shaping
hippocampal network activity based on input spatial targeting rather than input amount.

To assess the robustness of the differential impact of supralinear versus sublinear dendritic
activation on LFP oscillations, we performed multiple sensitivity analyses on synaptic
parameters and input characteristics. In all cases, supralinear activation enhanced both peak
power and peak frequency power of the fast LFP component, whereas sublinear activation
consistently resulted in stronger slow oscillatory power compared to supralinear activation
(**Figure, S3**).

Figure 1. Supralinear and Sublinear branches of PV+ FSBCs trees, differentially modulate memory-related hippocampal oscillations under the same amount of synaptic activation. (a) *Schematic Illustration of the hippocampal microcircuit model. PCs and PV+ FSBCs populations are activated by a theta (4 Hz) input for 1000 msec. An LFP electrode is positioned nearby the PCs somata. Upon activation, the model exhibits slow (bandpassed at 3-10 Hz) and fast (bandpassed at 30-200 Hz) LFP bands. PV+ FSBCs models include anatomical reconstructions and are equipped with bimodal nonlinear trees (See Glossary) as per* [33,34]. **(b-c).** *The same Number of Synapses are activated either in the Supralinear (shown in purple, **b**) or in the Sublinear (shown in cyan, **c**) branches of the PV+ FSBCs dendritic trees. **(d-g).** Power Spectrum Density (PSD) plots under the distinct synaptic placement configurations. When only the supralinear branches receive synapses (supralinear activation), the fast oscillatory component is massively enhanced in its frequency and power compared to the sublinear activation scenario (**d,e**). In contrast, an increase in slow oscillatory power is predicted when synapses are restricted to sublinear branches (**f-g**). of the PV+ FSBCs trees. Data from 30 random simulation trials.*

Figure 2. LFP frequency and power are regulated by PV+ FSBCs nonlinear dendritic activation. (a-c) Schematic illustration of the activation protocols and representative traces of the evoked LFP bandpassed at slow (3-10 Hz) and fast (30-200 Hz) frequences. The same number of synapses targets either both types of nonlinear dendrites (bimodal activation) (**a**), or only supralinear (**b**) or only sublinear (**c**). (**d-f**) Comparison of the peak fast-frequency (**d**) and the peak powers of the fast (**e**) and slow (**f**) LFP components. Data from 30 random simulation trials. Synaptic activation of supralinear dendrites of the PV+ FSBCs bimodal trees results in a higher peak fast-frequency and increased fast-frequency peak power, while reducing the peak power of slow oscillations compared to sublinear dendrite activation. Statistical comparisons across multiple groups were performed using the Kruskal-Wallis test followed by a post-hoc correction for multiple comparisons, suitable for data with unequal variance.

To elucidate the mechanisms underlying the observed LFP modulations, particularly given
that the total number of synapses remained constant across all three activation protocols
(bimodal, supralinear, and sublinear, shown in our Figures 1-2), we analyzed the
corresponding spiking activity of both PCs and PV+FSBCs (**Figure 3a-c**). Our analysis
revealed that supralinear activation significantly increased PV+ FSBCs firing rates compared
to both bimodal and sublinear activation scenarios (**Figure 3a, c**). This heightened inhibitory
activity, in turn, suppressed PCs firing, as evidenced by the lower PCs firing frequencies
observed under supralinear activation (**Figure 3b, c**). Conversely, sublinear dendritic
activation diminished PV+ FSBC firing rates, leading to a disinhibition of PCs and
consequently higher PCs firing frequencies relative to the supralinear condition (**Figure 3**).
The bimodal activation protocol, with synapses distributed across both dendritic types,
yielded intermediate firing rates for both PV+ FSBCs and PCs, indicative of a intermediate
interplay between the effects of the two distinct nonlinear integration modes (**Figure 3c**).
These findings collectively suggest that the nonlinear dendritic properties of PV+ FSBCs
facilitate a differential modulation of network activity: supralinear dendritic inputs enhance
PV+ FSBCs firing to stronger suppress PCs activity, while inputs solely to sublinear
branches, result in reduced, compared to supralinear, PV+ FSBC firing, thereby disinhibiting
PCs. Such dynamic regulation of the network's E/I balance underscores the functional
significance of the supralinear and sublinear nature of PV+ FSBCs trees in shaping

memory-

hippocampal circuit excitability and related oscillations.

Figure 3 Supralinear and sublinear PV+ FSBCs dendrites differentially modulate the excitation/inhibition (E/I) balance in the microcircuit model under identical synaptic activation conditions. (a) Representative raster plots depicting the spiking activity of PCs (blue) and PV+ FSBCs (brown) across theta cycles when synaptic inputs are selectively targeted to either supralinear dendrites (upper panel) or sublinear dendrites (lower panel) of PV+ FSBCs. (b) Representative spike trains of PCs (left) and PV+ FSBCs (right). When PV+ FSBCs receive synaptic input in their supralinear branches, their firing rates increase more profoundly across theta cycles, leading to a stronger decrease in PCs spiking. Conversely, when synaptic inputs are restricted to sublinear dendrites, PV+ FSBCs firing rates are reduced, resulting in increased PCs activity. (c) Activation of a fixed number of synapses on supralinear dendrites (shown in purple) significantly increases the firing frequency of PV+ FSBCs, leading to a reduction in the activity of PCs when compared to bimodal (supralinear and sublinear, shown in grey) or purely sublinear activation. Conversely, activation of randomly selected sublinear dendrites (shown in cyan) decreases the firing frequency of PV+ FSBCs, resulting in higher disinhibition of the PCs. Data are based on 30 random simulation trials. Pairwise comparisons were performed using the Mann–Whitney U test.

These observations demonstrate that PV+ FSBCs can dynamically adapt their firing patterns depending on whether synaptic input is received by their supralinear or sublinear dendritic branches, despite the total synaptic input remaining constant. Supralinear branches, characterized by larger volumes and lower input resistance [33], facilitate enhanced forward propagation of synaptic signals (**Figure S2g**). In contrast, sublinear branches, which have smaller volumes and higher input resistance [33], limit forward propagation (**Figure. S2f**), leading to a decrease in PV+ FSBCs firing rates. Consequently, this differential activation modulates the E/I balance in the network, directly shaping the oscillatory activity observed in the LFP. These findings indicate that PV+ FSBCs can potentially use the bimodal nonlinear trees to differentially process incoming synaptic inputs, even when the total synaptic input remains unchanged. Such a mechanism may not only modulate PV+ FSBCs firing responses at the cellular level (**Figure 3**) but also regulate oscillations **(Figure 1-2)**, thereby influencing hippocampal mnemonic functions.

It is well established that increasing the number of synaptic inputs onto a neuron can elevate its firing rate, which in turn can influence the frequency and power of oscillations. From this perspective, even sublinear dendrites, despite lacking the ability to generate dendritic spikes, can still drive high-frequency output if provided with sufficient synaptic input. Indeed, recent studies have shown that interneurons with purely sublinear dendrites can contribute to gamma oscillations when strongly activated[35]. Our model aligns with this to some extent: activation of sublinear PV+ FSBCs dendrites supports fast oscillations too (see also **Figure S4**), but their frequency and power are substantially lower compared to activation of supralinear dendrites. This raises a key conceptual question: what would be the functional advantage of PV+ FSBCs possessing both sublinear and supralinear dendrites, if either could potentially support high-frequency activity given enough input?

We propose that nonlinear dendritic diversity within the same cell may offer a computationally and energetically efficient mechanism for dynamically modulating PV+ FSBCs output and hippocampal oscillations (**Figure 4**). Activating large numbers of synapses is energetically costly. In contrast, bimodal nonlinear dendritic integration may

enable PV+ FSBCs to achieve distinct functional outcomes, such as differential modulation
of firing rates, E/I balance, and oscillatory states, without requiring an increase in synaptic
load.

In our simulations (**Figure 4**), we systematically examine the impact of different synaptic
input onto either supralinear or sublinear dendrites across multiple levels. The results show a
clear dichotomy in the response depending on which dendritic type is targeted: Increasing
synaptic input onto supralinear dendrites progressively enhances PV+ FSBCs firing, strongly
suppresses excitatory PCs activity, and strengthens high-frequency oscillations (**Figure 4a,**
right side; b–f, purple curves). In sharp contrast, increasing synaptic input onto sublinear
dendrites leads to comparatively reduced PV+ FSBCs output, stronger disinhibition of PCs,
and an increase in slow (theta) oscillatory power (**Figure 4a, left side; b–f, cyan curves**).
Notably, even at the highest levels of synaptic input, sublinear activation fails to replicate the
fast oscillatory dynamics achieved through supralinear activation. Together, these findings
reveal that PV+ FSBCs can exploit their nonlinear bimodal dendritic architecture to
adaptively shape cellular and circuit activity depending on input spatial location, rather than
synaptic quantity alone. This supports a view in which dendritic nonlinearities may provide
an energy-efficient, and flexible mechanism for interneuron-mediated control of memory
related oscillatory states.

Figure 4. Dendritic integration mode governs PV+ FSBCs mediated modulation of
hippocampal oscillations irrespectively of synaptic input amount. *(**a**) Schematic*
summary illustrating how synaptic input targeting different dendritic integration modes in PV+
FSBCs. Sublinear (left cyan) vs. Supralinear (right purple), leads to distinct Cellular and LFP
effects: Activation of sublinear dendrites reduces PV+ FSBCs firing, increases PCs activity,
and enhances slow oscillations. In contrast, activation of supralinear dendrites increases
*PV+ FSBCs output, suppresses PCs, and strengthens fast oscillatory components. (**b–f**)*
Simulations upon increasing synaptic input number to either supralinear (purple) or sublinear
(cyan) dendrites reveal a consistent divergence in cellular (firing frequency) and LFP

responses: PV+ FSBCs firing rates increase more when their supralinear dendrites are
*targeted in comparison to sublinear targeting (**b**). In contrast, PCs firing is less reduced*
*when different number of synapses are placed in sublinear PV+ FSBCs dendrites (**c**). Peak*
slow frequency power is enhanced with sublinear activation and reduced with supralinear
*input (**d**). Peak fast-frequency power and frequency is enhanced with supralinear activation*
*and remain low under sublinear input, even when the number of synapses is high (**e–f**). Data*
from 30 simulation trials show mean and std values.

We further explored how the nonlinear integration strategies of PV+ FSBCs can impact their
cellular responses and circuit effects. (**Figure. S4a-b**). Specifically, we repeated our
simulations pipeline but with PV+ FSBCs models that are equipped with purely supralinear
or purely sublinear trees. Leveraging the distinct morphological features of supralinear and
sublinear branches in our bimodal FSBCs reconstructions, we adjusted the dendritic
morphologies as per [33]. When PV+ FSBCs were equipped with purely supralinear dendritic
trees, we observed enhanced fast frequency and power compared to both bimodal and
purely sublinear tree configurations. This configuration also maximized their firing (**Figure**
S4a, c-h). Conversely, PV+ FSBCs with purely sublinear dendritic trees present a reduced
spike rate which increased the PCs activity (**Figure. S4h**). Sublinear trees exhibited
increased slow power and decreased fast power and lower fast power compared to
supralinear trees (**Figure. S4b-g**). Crucially, these simulations of 'one-mode' integration are
informative beyond direct comparison with the bimodal case (**Figures 1-4**), as they delineate
the functional consequences should PV+ FSBCs, or other PV+ subtypes thereof,
predominantly operate via one of these fundamental linear/nonlinear strategies, aligning with
all (often diverse) experimental observations of dendritic computations in PV+ interneurons
across areas[22,25].

These simulations underscore that PV+ FSBCs may take advantage of their bimodal
nonlinear dendritic trees by dynamically utilizing their spiking-supralinear or their passive
linear/ sublinear dendrites to regulate the E/I balance within the network, thereby
orchestrating distinct oscillatory behaviors under equivalent synaptic input conditions.
Importantly, activating either only supralinear or only sublinear branches of a bimodal tree
(**Figs.1-2**) exhibits similar phenotypic responses to configurations of purely supralinear or
purely sublinear trees (**Fig.S3**). This further supports the notion that the bimodal nonlinear
nature of PV+ FSBCs dendrites allows for flexible modulation of network activity and
oscillatory patterns crucial for hippocampal mnemonic functions.

Apart from the type of dendritic integration, the spatial arrangement of synapses, along the
selected dendritic branch, influences the firing characteristics[36]. It's well established that
supralinear dendrites tend to enhance clustered synaptic allocation, whereas sublinear
dendrites prefer dispersed activation. For PV+ FSBCs, it has been observed that dispersed
synaptic activation leads to higher firing frequencies compared to clustered [20,33,37]. This
phenomenon is influenced by the co-existence of sublinear along with supralinear branches,
coupled with their small size and the presence of A-type potassium channels[33,37,38], which
discourages preference for clustered inputs. To investigate how synaptic clustering affects
the oscillatory behavior of our network, we simulated the activation of the same number of
synapses as in our disperse protocols (as in **Figure. S4**) but now clustered in a few
randomly chosen dendrites (**Figure. S5**). As anticipated, clustered activation of sublinear
PV+ FSBC trees significantly reduced their firing rates while maximizing the E/I balance in
the network. Conversely, clustering synapses in supralinear trees enhanced PV+ FSBC
firing rates compared to both bimodal and sublinear clustering configurations, while also
decreasing the firing rates of the PCs (**Figure. S5j**). At the oscillatory level, akin to our

simulations with dispersed synaptic activation, clustered synapses in supralinear PV+ FSBC trees enhanced fast LFP components, whereas synaptic clustering in sublinear trees boosted the slow LFP power (**Figure. S5a-i**). These findings underscore the dual impact of dendritic morphology and synaptic spatial arrangement on neuronal firing patterns and network oscillatory dynamics. They highlight the intricate mechanisms by which PV+ FSBCs can dynamically modulate their responses and contribute to the regulation of network excitability and oscillatory activity crucial for hippocampal function. An important consideration for future experimental and computational work is whether supralinear and sublinear dendrites receive synaptic inputs from distinct pathways which may also be temporally segregated. Such compartmentalization of inputs could profoundly influence inhibitory activity across different oscillatory and behavioral states (see Discussion).

Finally, we analyzed the Phase-Amplitude Coupling (PAC),[39] to check the slow-fast oscillation coupling in the LFP across all cases (**Figure. S6**). Indeed, the LFP response aligns with experimental findings, indicating that slow activity entrains hippocampal networks to slow-fast coupling (**Figure S6a**)[5,8,40–44]. Activation of either supralinear or sublinear branches did not alter the coupling between the two oscillatory components (**Figure. S6b-f**). However, coupling was enhanced when synapses were clustered on fully sublinear dendritic trees of PV+ FSBCs. Conversely, coupling was reduced when similar clustering occurred on supralinear PV+ FSBCs dendritic trees (**Figure. S6g-j**). These results indicate that as the power and frequency of the fast component decrease, it becomes more synchronized with the slower component.

Discussion

1. Bimodal Dendritic Integration in PV+ FSBCs: A Mechanism for Rhythm-Specific Modulation?

While the role of interneurons in memory-related hippocampal oscillations is well established [45–48], the subcellular mechanisms through which they influence these rhythms remain poorly understood. PV+ FSBCs, a major PV+ interneuron subtype, have recently been implicated in fast gamma and sharp wave-ripple (SWRs) oscillations[49], and dendritic supralinearity has been proposed as a mechanism for enhancing fast activity [50] However, no direct mechanistic framework has previously explained how different, and possibly diverse dendritic integration modes such as the bimodal profile explored here could contribute to selective control of slow vs. fast memory-related oscillations. Our computational predictions serve in addressing a key knowledge gap by proposing a novel framework where the dendritic integration modes, rather than merely the synaptic input quantity, may constitute a critical determinant of the oscillatory regime that PV+ FSBCs support. Specifically, activating supralinear dendrites significantly promotes PV+ FSBCs firing, decreases the E/I balance, and amplifies fast oscillatory power and frequency. Conversely, engaging sublinear dendrites leads to less prominent PV+ FSBCs output, consequent disinhibition of PCs, and an enhancement of slow oscillations (as detailed across **Figures 1–4** and **Suppl. Figures 4-5**). Notably, these distinct dynamic outcomes manifest even when the total synaptic input to PV+ FSBCs is held constant (**Figures 1-3**), strongly suggesting that these interneurons possess an intrinsic, spatially defined computational mechanism for precise oscillatory tuning.

2. Implications for Memory Encoding, Consolidation, and Recall

Memory formation involves distinct temporal phases such as encoding, consolidation, and recall, each linked to specific oscillatory states (e.g., theta for encoding, gamma and SWRs for consolidation, theta-gamma coupling for recall). Our results suggest that PV+ FSBCs dendritic integration modes are ideally suited to switch the hippocampal network between these functional states: In one hand, sublinear dendritic activation, which enhances slow power and disinhibits PCs, may support learning and encoding, when increased excitability is beneficial for storing novel information. On the other hand, supralinear activation, which may lead to more pronounced suppression of the PCs and enhances high-frequency synchronization, may facilitate consolidation and recall, by reinforcing stored traces and refining the temporal coordination of spike timing. This dual-mode mechanism could presumably enable PV+ FSBCs to fine-tune their engagement with both rhythms depending on which type of nonlinear dendritic domain is driven. Thus, the functional value of dendritic nonlinearities in hippocampal PV+ FSBCs may not be merely electrical, but cognitive[51,52]: it could offer a subcellular lever for dynamically tuning inhibition in alignment with memory demands.

3. A Framework for Energy-Efficient and Dendritic Specific Computation

Beyond functional flexibility, energy efficiency emerges as a critical computational advantage. While both supralinear and sublinear dendrites can support PV+ FSBCs firing, our simulations (**Figure 4**) highlight that supralinear branches achieve fast-frequency modulation more effectively and at a lower synaptic cost. This supports the view that dendritic spikes are not essential for fast rhythms per se, (in agreement with [35]) but may offer a more economical and spatially targeted solution. This prediction serves as a new idea with implications for how neural systems balance performance and metabolic constraints at the cellular and network level computations[51]. Importantly, these findings refine our understanding of PV+ FSBCs contributions: their nonlinear bimodal dendritic architecture can enable input-specific, rhythm-specific, and energy-aware modulation of activity at the circuit level. Such a perspective complements classical models where inhibition is adjusted by input quantity alone and opens the door to studying location-sensitive memory modulation at the dendritic level.

4. Future Directions: Generalizing Across Interneuron Types and Circuits

While our simulations establish a robust computational foundation for understanding PV+ FSBCs, they concurrently pave the way for broader experimental investigations. The hippocampus alone harbors over 20 distinct interneuron subtypes, each characterized by unique morphological, developmental, and connectivity profiles. Significant heterogeneity exists even within major inhibitory classes; for instance, the PV+ interneuron family, in addition to PV+ FSBCs, includes other subtypes such as the bistratified and axoaxonic cells. Each subtype exhibits substantial variations in its morphological, molecular, and intrinsic properties, as well as distinct connectivity patterns (e.g., perisomatic, dendritic, or axonal targeting) within local circuits. [45,46]. Encouragingly, recent experimental work has begun to reveal nonlinear dendritic properties in other inhibitory populations beyond the hippocampus [25]. Nevertheless, for most hippocampal inhibitory subtypes, the specific dendritic integration modes, and the functional relevance of their input targeting remain largely uncharacterized. Our current modeling framework is designed for modularity and can be readily adapted to incorporate such emerging experimental data. As new biophysical and morphological profiles for other interneuron classes and subtypes are available, our model can be expanded. This will enable simulations to predict how diverse dendritic architectures and input targeting strategies contribute to distinct oscillatory roles across various inhibitory cell

types. To facilitate and guide these future experimental endeavors, we subsequently outline
several methodological approaches for validating our model's predictions:

First, *ex vivo* whole-cell patch-clamp recordings combined with two-photon glutamate
uncaging in brain slices can be employed to functionally identify distinct dendritic integration
profiles in interneurons, like the approaches used in Cornford et al., 2019 for investigating
the diverse (supralinear, and linear/sublinear) integration properties in hippocampal PV+
dendrites[21]. Additionally, voltage-sensitive dye imaging and two-photon fluorescence imaging
could help visualize dendritic activity during synaptic stimulation, building on prior work by
Katona et al., 2011[22]. *In vivo*, photoactivatable fluorescent proteins (e.g., PA-GFP or PA-
mCherry), driven by interneuron-specific promoters, can be used for precise localization and
targeting of dendritic subtypes. To assess dendritic function during memory-related
oscillations, *in vivo* two-photon voltage imaging in awake behaving animals could reveal
whether supralinear and sublinear dendrites are differentially engaged across theta and
gamma rhythms. Simultaneous LFP recordings with dendritic imaging could further clarify
how these dendritic domains contribute to oscillatory signatures, similar to what shown in the
pioneering work of Liao et al.,[53]. Moreover, optogenetic activation of input pathways mapped
to specific dendritic regions would allow causal testing of their roles in modulating
interneurons firing and network dynamics. Finally, mapping input selectivity onto distinct
dendrites (i.e. supralinear and sublinear) could be achieved through *in vitro* mGRASP-based
methods, or *in vivo* monosynaptic rabies virus tracing. These approaches could reveal
whether specific afferent sources (e.g., from entorhinal cortex, medial septum, or locus
coeruleus) preferentially target distinct dendritic domains. Together with synergistic modeling
efforts, such integrative strategies would offer a powerful path toward uncovering how
dendritic processing in interneurons can dynamically and energy efficiently regulate the
temporal architecture of memory processing in the hippocampus and potentially across
broader cortical circuits.

Limitations of the Study

Our microcircuit deliberately focuses on PV+ FSBCs, as this is the interneuron subtype with
the most detailed experimental data on dendritic integration, but it does not include the full
diversity of hippocampal inhibitory cells. Furthermore, our model utilizes a generic rhythmic
input, as it remains an important open question whether distinct afferent pathways
preferentially target the supralinear versus sublinear dendritic domains of PV+ neurons.

Author contributions: AT and DS conceived the study. AT designed and ran the
simulations, analyzed the data and prepared the figures. MEL and DS provided senior
conceptual input on the study. DS funded and supervised the project. AT wrote the
manuscript. All authors edited and approved the final version of the manuscript.

Acknowledgements: A.T. was supported by the DFG with the SFB1315-2 TP A01 Brenda
Milner Award and the Einstein Center for Neurosciences Berlin Fellowship. M.E.L. was
supported by the European Research Council (101055340, ERC AdG, Cortical Coupling).
D.S. was supported by the Einstein Foundation Berlin, the European Research Council
(ERC) under the Europeans Union's Horizon 2020 research and innovation program
(BrainPlay grant, agreement no. 810580), the German Research Foundation (Deutsche

Forschungsgemeinschaft [DFG], SFB-958 – project 184695641, project 431572356, FOR
3004 – project 415914819, SFB 1315 – project 327654276 and under Germany's Excellence
Strategy – Exc-2049-390688087 NeuroCure), and the Federal Ministry of Education and
Research (BMBF, SmartAge – project 01GQ1420B). We thank very much Dr. Michalis
Pagkalos, Dr. Nikolaus Maier, Dr. John Tukker, Dr. Daniel Parthier, Dr. Spiros Chavlis and Dr.
Jeremie Sibille for critical discussions, proofreading and comments. We also thank Prof. Dr.
Nelson Rebola, Dr. Yann Zerlaut, and Dr. Yiota Poirazi, for comments on an earlier version of
the work.

Code and Data availability

540 All codes/scripts and datasets required to reproduce the results and figures, as well as all
541 statistical analyses, are accessible in the ModelDB database.
542 https://modeldb.science/2018008 *Access code:* 280621Interneurons! They are also
543 available on GitHub: https://github.com/AlexandraTzilivaki/Tzilivakietal2024 For questions
544 contact alexandra.tzilivaki@charite.de

Competing interests

547 The authors declare no competing interests.

<div align="center">

Methods

</div>

Model implementation and availability

Simulations were conducted using NEURON (v7.6)[54] on a High-Performance Computing
Cluster, utilizing 111 CPU cores on a 64-bit CentOS Linux operating system.

All codes/scripts and datasets required to reproduce the results and figures, as well
as all statistical analyses, are accessible in the ModelDB database.

Please visit:

https://modeldb.science/2018008 Access code: 280621Interneurons!

They are also freely available on GitHub:
https://github.com/AlexandraTzilivaki/Tzilivakietal2024

Neuronal populations

1. PV+ FSBCs

Two multi-compartmental biophysical models of CA3 PV+ FSBCs were employed,
adopted from Tzilivaki et al. (2019)[33]. These models include detailed anatomical
reconstructions of somata and dendritic trees taken from Neuromorpho database
(originally published in Tukker et al. 2007[55]), (**Figures 1, S1-2**). Both FSBCs feature
bimodal nonlinear dendritic branches, characterized by both supralinear and sublinear
branches. They are equipped with fast voltage-dependent sodium channels (gnafin),
delayed rectifier potassium channels (gkdrin), slow inactivation potassium channels

570	(gslowin), slow calcium-dependent potassium channels (gkcain), A-type potassium
571	channels in proximal and distal dendritic regions (gkadin, gkapin), h-currents (ghin), and
572	L-, N-, and T-type voltage-activated calcium channels (gcal, gcan, and gcat,
573	respectively). These models have been extensively validated against experimental data
574	and accurately capture the intrinsic features and electrophysiological responses of
575	hippocampal PV+ FSBCs (See original reference[33] and Supplementary Information). The
576	mean dendritic diameter and total dendritic length were measured for the supralinear and
577	sublinear dendrites of the two FSBCs morphological reconstructions. Dendritic volume
578	was calculated as per Tzilivaki et al. (2019)[33] using the following formula:

$$V = \pi * \left(\frac{diameter}{2}\right)^2 * length \ (\mu m^3)$$

579	To causally manipulate the dendritic morphologies of FSBCs to generate fully supralinear or
580	sublinear trees (**Figure S2**), we performed causal manipulations according to the approach
581	described in Tzilivaki et al. (2019)[33]. We fixed the diameter and length of all dendrites to
582	create trees with average supralinear or sublinear dendritic volume (**Figure S1B**), which
583	dictated the integration mode as shown in the original modeling publication. Dendritic Input
584	Resistance was calculated using the following formula:

$$Rin = \frac{DV}{I} \ (\frac{mV}{nA})$$

| 586 | where I = −100 pA injected in each dendritic branch and DV is the generated |
| 587 | IPSP. |

| 589 | For more detailed information about the models, please refer to the relevant |
| 590 | publication[33] and **Tables S1-2, 4-5**. |

| 592 | 2. Pyramidal Cells (PCs) |

593	Biophysically relevant hippocampal PC models with reduced morphology (n = 20) were
594	adopted from Hadler et al. (2024)[29]. These models consist of somata and proximal, distal,
595	and basal dendritic branches. PCs include a Ca2+ pump and buffering mechanism, Ca2+
596	activated slow AHP and medium AHP potassium (K+) currents, an HVA L-type calcium
597	(Ca2+) current, an HVA R-type Ca2+current, an LVA T-type Ca2+ current, an h-current, a
598	fast sodium (Na+) current, a delayed rectifier K+ current, a slowly inactivating K+ M-type
599	current, and a fast inactivating K+ A-type current. These current mechanisms were non-
600	uniformly distributed along the somatodendritic compartments. They were validated based
601	on *in vitro* data to replicate the electrophysiological profile and basic dendritic architecture of
602	hippocampal CA3 PCs[29,56]. The PC models do not include detailed nonlinear dendritic trees,
603	as this was beyond the scope of this study. Incorporating realistic anatomical reconstructions
604	for PCs would have significantly increased computational complexity and reduced simulation
605	speed. For more information, see Hadler et al. (2024)[29] and **Tables S3-5**.

| 606 | **Synaptic properties** |

607	The PC models were equipped with AMPA, NMDA, and γ-aminobutyric acid type A (GABAA)
608	synapses, while the FSBCs included Ca2+ permeable AMPA (CP-AMPA), NMDA, GABAA,
609	and autaptic GABAA synapses. The synaptic conductance values for every connection type
610	were calibrated based on experimental[57–60] and modelling studies[33] and are listed in **Table**
611	**S5**

To ensure the robustness of our finding, we repeated the simulations upon increased or
decreased synaptic conductance values (15% change) (see sensitivity analysis **Figure S2**)

Hippocampal Microcircuit

To study the effect of PV+ FSBCs dendritic processing in circuit level function, we
implemented an experimentally constrained microcircuit model of the CA3. Typically, for
such research questions, small microcircuit modelling is the best choice as it provides
enough detail to study subcellular mechanisms at the circuit level while keeping the
computational demands low. Our hippocampal microcircuit configuration was adopted from
Hadler et al. (2024)[29]. The model consists of 20 PCs and 2 FSBCs. In each random
simulation trial (n = 30), each PC contacted up to seven (7) randomly chosen PCs with one
AMPA and one NMDA synapse activation per contact. Each PV+ FSBC received synaptic
input from fifteen (15) randomly chosen PCs in each simulation trial. Additionally, each PC
received thirteen (13) feedback inhibitory GABAergic inputs from each PV+ FSBC per
simulation trial. Each FSBC formed five (5) GABAergic synapses per simulation trial and was
self-inhibited through autapses. For further details, see the Simulation Paradigms chapter
and Hadler et al. (2024)[29].

Simulation Paradigms

Input

The microcircuit is activated by a theta entrained presynaptic population as per Turi
et al., 2018[61]. The input was modeled as an artificial presynaptic population (N=22)
using NEURON's VecStim function. The spike times of the presynaptic population
were generated using a sinusoidal theta like filter that was applied so to account for
theta like modulated spike times.
Thata like probability formula (as per [61]):

$$p(t) = (\sin\left(2.0 * pi * \frac{f_{theta*spike}}{1000.0} + phi_{theta}\right) + 1.0)/2.0$$

Where:
• spike = the spike time in msec (float)
• f_theta = theta-cycle frequency in Hz. (float)
For the simulations shown in **Figures 1-4** and **Supplementary Figures 4-6**,
f_theta=4 Hz. For the sensitivity analysis simulations as shown in **Supplementary**
Figure 3, we changed the input the theta frequency to f_theta=5Hz.
• phi_theta= theta cycle phase in radians.
For the simulations shown in **Figures 1-4** and **Supplementary Figures 4-6**,
phi_theta=0 (equal to 0 radians). For the sensitivity analysis simulations as shown in
Supplementary Figure 3, we shifted the input phase so phi_theta=0.5 (equal to 180
radians).
647
648 If the probability p(t) is greater than 0.7 a spike is generated for the specific artificial
649 neuron (n=22). Every input neuron has its own theta modulated spike train.
650
Each PC received input from 5 artificial presynaptic neurons, while each PV+ FSBC
received input from 7 artificial presynaptic neurons. Although the input was sufficient
to activate the PC population due to its reduced morphology, 7 artificial presynaptic
neurons were insufficient to evoke spiking activity in the PV+ FSBC models, given
their realistic complex anatomical reconstructions. This subthreshold activation was

selected to ensure that PV+ FSBCs were primarily engaged in the network due to the
local inputs they received from PCs.
658
1. **Simulations**
The microcircuit model was simulated for approximately 12,000 milliseconds (ms)
with a time step of 0.1 ms. The first 200 ms were excluded from the analysis to allow
the model to reach an equilibrium state. For the dispersed protocols, synapses to
FSBCs (both input and local PC-to-FSBC synapses) were randomly assigned to
dendrites, meaning one randomly chosen dendrite received one pair of NMDA and
CP-AMPA synapses. For the clustered protocol (**Supplementary Figure S4**), the
same total amount of synapses was placed in 4 randomly chosen dendrites (cluster
size: 5-7 synapses (per dendrite) Data represent the results (mean and standard
deviation values (std)) of thirty (30) random simulation trials for each protocol. In
every trial, the total number of synaptic contacts and the connectivity ratios remained
identical, but different randomly chosen neurons (from both PC and PV+ FSBC
populations) were connected to different random neurons. Additionally, in each trial,
different dendrites from both PCs and PV+ FSBCs were randomly chosen and
synapses were activated at different parts of the chosen dendrites. This approach
ensured that the results reflected the diversity, especially of the PV+ FSBCs trees, in
bimodal activation protocols. Furthermore, two different anatomical reconstructions
were used for our models to account for PV+ FSBCs morphological variability (see
relevant chapter).
678
2. **LFP simulation and Spectral analysis**

To record the Local Field Potential (LFP), an *in silico* electrode was simulated based on
NEURON's extracellular function, following the modelling approach of Vladimirov et al.
(2013)[62]. The electrode was placed close to the PCs somata and remained in the exact
same position throughout the execution of all protocols and simulation trials. The
sampling frequency was set at 10 kHz. Despite the microcircuit model's small size, it
efficiently generated fast and slow oscillations comparable to experimental
observations[5,10,39,42] while maintaining low computational complexity and demands.

The *in silico* LFP datasets were band-passed at two respective bands: slow (3 – 10 Hz)
and fast (30-200 Hz). Slow and fast Peak-Frequency powers and peak Fast frequency
were determined using custom-made MATLAB scripts (MATLAB, The MathWorks Inc.,
Natick, MA; Torrence and Compo 1998) utilizing the p-welch function (0.4 Hz resolution)
and visualized as Power-Frequency plots. Phase-Amplitude Coupling (PAC) analysis
was conducted using Comodulogram generation and the calculation of the modulation
index (MI) metric that were adopted from Tort et al. (2008)[32]. Wavelet phase was
calculated at 15 levels from 1-15 Hz, and the amplitude at 50 levels from 30-180 Hz. The
MI was obtained by measuring the divergence of the observed amplitude distribution
from the uniform distribution.

Statistical Analysis

Statistical analyses for multigroup comparisons were performed using the Kruskal-Wallis
test, followed by a post-hoc correction for multiple comparisons (multcompare function
Matlab). For pairwise comparisons where data exhibited unequal variance, p-values
were calculated using the Mann-Whitney U test.

Supplementary Information

Figure S1. Electrophysiological calibration and responses of the models. a-c. *FS BCs*
responses. ***a.*** *Frequency- Injected current (FI) curve upon current clamp simulation in the*
cell bodies of the two multicompartmental FSBCs (1 sec duration). ***b.*** *Firing profiles of the*
two FSBCs after depolarizing current injections in the cell bodies (300 pA, 1 sec). FSBCs
exhibit the typical high-frequency firing pattern. ***c.*** *CP-AMPA (−70 mV) and NMDA (+60 mV)*
currents upon stimulation as per [60]. Traces represents the mean of the two
multicompartmental FSBCs. ***d.*** *A three-step voltage clamp of voltage changes from −70 mV*
to 10 mV (duration 1 msec) and back to −70 mV was used to produce inhibitory current.

During the validation of this current, the reversal potential of Cl– was adjusted from –80 to
–16 mV, in order to reproduce the experimental set up of [63], Mean trace of the two FSBCs
responses. ***e-h*** *Pyramidal Cell responses.* ***e.*** *FI curve of the Pyramidal Cell model same*
approach as in a. ***f.*** *Firing profile of the Pyramidal Cell under current injection (300 pA for 1*
sec) at the cell body. The model successfully represents the typical experimental phenotype
shown in [56]. ***g.*** *AMPA (-70 mV voltage clamp) and NMDA (60 mV voltage clamp) currents of*
the Pyramidal Cell model mimic the experimental data of [58]. ***h.*** *Inhibitory current calibration*
based on experimental data from [64].

Figure S2*. **Mechanisms of bimodal nonlinear dendritic integration in***
Multicompartmental Models of Hippocampal FSBCs. ***a.****Illustration of the morphological*
characteristics of supralinear and sublinear dendrites in bimodal FSBC models. Supralinear
dendrites (purple) are larger, whereas sublinear dendrites (blue) are longer and thinner. ***b-c****.*
Discriminative features between dendrite types: supralinear dendrites have larger volumes
(b) and lower input resistance (c) compared to sublinear dendrites. Statistical significance
was determined using the Mann–Whitney U test. ***d-e****. Supralinear dendrites (purple) can*
generate local sodium-dependent spikes, whereas sublinear dendrites (blue) cannot (d).
Blocking active sodium conductance in FSBC dendrites completely abolishes the supralinear
mode (e). ***f-g****. Forward propagation efficiency in bimodal nonlinear dendrites of FSBCs:*
Current injection (100 pA) at randomly selected dendrites and recording at the soma show
that sublinear branches (f) propagate signals less effectively compared to supralinear
branches (g). Panels a,d,e were adopted from [33]

*Figure S3. Robustness/Sensitivity Analysis on Synaptic and Input Parameters. **a-c**. 15% increase in the synaptic conductance values (applied to both input and network synapses for/from both PCs and FSBCs) do not alter the enhancement of fast peak frequency and power observed with supralinear activation. **d-f**. 15% reduction of the synaptic conductance values (for both input and network synapses involving PCs and FSBCs) also maintain the observed increase in fast peak frequency and power upon supralinear activation. **g-i**. Modifications to input parameters (input phase shifted 180°, peak frequency 5 Hz) similarly do not affect the enhancement of fast peak frequency and power driven by supralinear activation. Statistical analyses for multigroup comparisons were performed using the Kruskal-Wallis test, followed by a post-hoc correction for multiple comparisons.*

Figure S4. Differential Modulation of Slow and Fast LFP Components by Supralinear
and Sublinear FSBCs Dendritic Trees. a. *Activation of FSBCs equipped with purely*
supralinear dendritic trees, showcasing representative LFP traces bandpassed at slow (3-10
Hz) and high (30-200 Hz) frequencies. ***b***. *Similar to* ***a*** *but displaying activation of FSBCs with*
purely sublinear dendritic trees. ***c-d***. *Power Spectrum Density (PSD) plots of the LFP evoked*
when FSBCs are equipped with purely supralinear (c) or purely sublinear (d) dendritic trees,
highlighting differences in frequency response. ***e-g***. *Comparative analysis of the peak fast-*
frequency (e) and peak power of fast (f) and slow (g) oscillations for FSBCs with bimodal

(grey), supralinear (purple), or sublinear (blue) dendritic trees. Data are derived from 30
*random simulation trials. **h**. Firing activity of the PCs and FSBCs populations within the*
microcircuit network across 30 random simulation trials. Activation of supralinear FSBC
dendritic trees results in a decreased E/I balance compared to both bimodal (control) and
sublinear trees. Statistical analyses for multigroup comparisons were conducted using the
Kruskal-Wallis test followed by a post-hoc correction for multiple comparisons. Paired
comparisons and p-values were calculated using the Mann-Whitney U test for data with
unequal variance.

Figure S5. Impact of Clustered Synaptic Activation on LFP and E/I Balance in Bimodal,
Supralinear, and Sublinear FSBC Dendritic Configurations. a. Activation of FSBCs with
bimodal nonlinear dendrites (control configuration), showing representative LFP traces for
slow (3-10 Hz) and fast (30-200 Hz) frequencies. Synapses are clustered in a few randomly
chosen supralinear or sublinear branches. **b**. Activation of FSBCs with purely supralinear
dendritic trees, showing LFP traces under clustered synaptic activation. **c**. Activation of
FSBCs with purely sublinear dendritic trees, similar to conditions in a and b, illustrating the

effect of clustered synaptic activation on LFP traces. **d-f**. Power Spectrum Density (PSD)
plots for LFPs under clustered synaptic conditions in FSBCs with bimodal (d), supralinear
(e), and sublinear (f) dendritic configurations. **g-i**. Comparison of peak frequency (g) and
peak power for the fast LFP component (30-200 Hz) (h), and peak power for the slow LFP
component (i), across dendritic configurations. **j**. Firing activity of PC and FSBC populations
in the microcircuit network from 30 random simulation trials. Clustering in supralinear FSBC
dendritic trees decreases the E/I balance in the network compared to bimodal (control) or
sublinear trees. Statistical analyses for multigroup comparisons were conducted using the
Kruskal-Wallis test followed by a post-hoc correction for multiple comparisons. Paired
comparisons and p-values were calculated using the Mann-Whitney U test for data with
unequal variance.

Figure S6. Slow-Fast Oscillation Coupling in the Microcircuit Model Under Various
***Dendritic and Synaptic Configurations in FSBCs. a-e**. Representative comodulograms*
illustrating the slow-fast coupling for the protocols detailed in Figures 2 and S3. These
visualizations provide insights into the phase-amplitude coupling dynamics under different
synaptic and dendritic configurations. Data from 30 random simulation trials are represented.

f. Coupling analysis shows that synaptic distribution in a dispersed configuration across
bimodal, purely supralinear, or purely sublinear FSBC dendritic trees does not affect slow-
fast oscillation coupling. g-i. Similar to a-e but showcasing comodulograms for a clustered
synaptic arrangement (detailed in Figure S4), using the same number of synapses as in the
dispersed experiments. j. MI indicates that slow-fast oscillation coupling decreases when
synapses are clustered in purely sublinear dendritic trees compared to other configurations.
Multigroup comparisons were performed using the Kruskal-Wallis test followed by a post-hoc
correction for multiple comparisons for multi-group data with unequal variance. This
statistical approach was chosen to accommodate the diversity in the data from 30 random
simulation trials.

Table S1. Active properties of Fast Spiking Basket Cell (FSBC) Models

Ion channel (S/cm^2)	Soma	Axon	Proximal dendrites	Distal Dendrites	*values as per [33,65,67]
Na$_v$	0.396	1.296	0.018	0.014	
H$_v$	0.00001	X	x	X	
Kdr$_v$	0.0432	0.144	0.0009	0.009	

Table S2. Passive properties of FSBCs

	Soma	Axon	Proximal dendrites	Distal Dendrites
Leak conductance (g_pas)[68]	1.315e-4 S/cm^2	3.55e-6 S/cm^2	1.315e-4 S/cm^2	1.34e-5 S/cm^2
Resting Membrane Potential (e_pas)[68,69]	-68 mV	-68 mV	-68 mV	-68 mV
Membrane capacitance (cm)[68]	1.2 uf/cm^{-2}	1.2 uf/cm^{-2}	1.2 uf/cm^{-2}	1.2 uf/cm^{-2}
Axial Resistance (Ra) [68]	172 ohm/ cm	172 ohm/ cm	142 ohm/ cm	142 ohm/ cm
		X	0.0002	0.0002
Calcium buffering dynamics [33,63,68]	Yes	No	Yes	Yes

Table S3. Passive Parameters and Active Conductance Values of the Pyramidal Cell (PC) Model

Mechanism	Soma	Proximal Apical Dendrite(n=1)	Distal Apical Dendrites (n=2)	Basal Dendrites (n=2)
Leak conductance [S/cm2]	0,0002	0,0002	0,0002	0,0002
Na+ conductance [S/cm2]	0,0105	0,0084	0,0084	0,0084
Delayed rectifier K+ conductance [S/cm2]	0,00086	0,00086	0,00086	0,00086
Proximal A-type K+	0,0075	0,015	-	0,0075

conductance [S/cm2]				
Distal A-type K+ conductance [S/cm2]	-	-	0,04875	-
M-type K+ conductance [S/cm2]	0.06	0.06	-	0.06
Ih conductance [S/cm2]	0.00005	0.0001	-	0.00005
L-type Ca2+ conductance [S/cm2]	0.0007	0.00003	-	0.00003
R-type Ca2+ conductance [S/cm2]	0.0003	0.00003	-	-
T-type Ca2+ conductance [S/cm2]	0.00005	0.0001	-	0.0001
Ca2+ -dependent sAHP K+ conductance [S/cm2]	0.0015	0.001	-	0.0005
Ca2+-dependent mAHP K+ conductance [S/cm2]	0,9	0,03	-	0,08
Membrane capacitance Cm [μF/cm2]	1	1	1	1
Membrane resistance Rm [Ohm cm2]	6000	6000	6000	6000
Axial resistance Ra [Ohm cm]	150	150	150	150

893 *values as per [29]

Table S4. Electrophysiological properties of the PCs and FSBCs models

	Value
Rheobase (pA)	200 (PC) / 150 +/- 10 (FSBC)
Input Resistance (MΩ)	142 (PC) / 97+/- 30 (FSBC)
Spike Threshold (mV)	-43,11 (PC) / -37 +/-1 (FSBC)
Spike Overshoot (mV)	30 (PC) / 48 +/- 1 (FSBC)
Resting Membrane Potential (mV)	-70 (PC) / -68 (FSBC)

Table S5. Synaptic conductance weight values of the PC and FSBC models.

Synapse Type	Pyramidal model	FSBC model
INPUT AMPA	0,00272	-
INPUT NMDA	0,0053	0.00032
INPUT CA++ PERMEABLE (CP)-AMPA	-	0.00075
AMPA	0,0017	-
CP-AMPA	-	0,0034
NMDA	0,0051	0.00144
GABA	0,0068	0,007
Autapse GABA	-	0,007

*values based on[29,33,57,58,60,64,70]

Glossary

Fast-Spiking Basket Cells (FSBCs)

GABAergic inhibitory interneurons that primarily target the perisomatic region of pyramidal
cells and other GABAergic neurons. Most FSBCs express parvalbumin (PV+) and are
characterized by their ability to fire at high frequencies.

PV+ neuron	
Parvalbumin-positive interneuron.	
SST+ neuron	
Somatostatin-positive interneuron.	
VIP+ neuron	
Vasoactive Intestinal Polypeptide-positive interneuron.	
Bimodal Dendritic Trees	
Dendritic trees in which different branches exhibit distinct nonlinear modes of excitatory synaptic integration: specifically, supralinear and sublinear. These integration modes coexist within the same neuron and are governed by differences in the morphological and/or active properties of individual dendritic branches.	
EPSPs (Excitatory Postsynaptic Potential)	
Depolarizing electrical signals generated in the postsynaptic neuron upon the activation of excitatory synaptic input.	
Dendritic Spikes	
Local, nonlinear depolarizations in dendrites that reflect supralinear summation of excitatory postsynaptic potentials (EPSPs). Often modeled as a sigmoidal function of integrated EPSPs.	
Memory Encoding	
The cognitive process by which new information is initially acquired and stored—commonly referred to as learning.	
Memory Consolidation	
The process by which newly encoded information is stabilized and stored for long-term retrieval.	
Memory Recall	
The retrieval of previously encoded and consolidated information.	
E/I Balance (Excitation/Inhibition Balance)	
The ratio of excitatory to inhibitory activity within a local circuit or network. An increase in inhibition lowers the E/I balance, while a reduction in inhibition raises it.	
Neuronal Oscillations	
Rhythmic fluctuations in electrical activity within the brain, typically measured via local field potentials (LFPs), that are associated with various behavioral and cognitive states.	

References:

1. Buzsáki, G., and Draguhn, A. (2004). Neuronal Oscillations in Cortical Networks. Science (1979) *304*, 1926–1929. https://doi.org/10.1126/science.1099745.

2.   Griffiths, B.J., and Jensen, O. (2023). Gamma oscillations and episodic memory.
Trends Neurosci *46*, 832–846. https://doi.org/10.1016/j.tins.2023.07.003.

3.   Jackson, J., and Skinner, F.K. (2018). Hippocampus, Theta, Gamma, and Cross-
Frequency Coupling. In Encyclopedia of Computational Neuroscience (Springer New
York), pp. 1–11. https://doi.org/10.1007/978-1-4614-7320-6_30-2.

4.   Lisman, J.E., and Jensen, O. (2013). The Theta-Gamma Neural Code. Neuron *77*,
1002–1016. https://doi.org/10.1016/j.neuron.2013.03.007.

5.   Tort, A.B.L., Scheffer-Teixeira, R., Souza, B.C., Draguhn, A., and Brankačk, J. (2013).
Theta-associated high-frequency oscillations (110–160Hz) in the hippocampus and
neocortex. Prog Neurobiol *100*, 1–14.
https://doi.org/10.1016/j.pneurobio.2012.09.002.

6.   Buzsáki, G., and Wang, X.-J. (2012). Mechanisms of Gamma Oscillations. Annu Rev
Neurosci *35*, 203–225. https://doi.org/10.1146/annurev-neuro-062111-150444.

7.   Jackson, J., Goutagny, R., and Williams, S. (2011). Fast and Slow Gamma Rhythms
Are Intrinsically and Independently Generated in the Subiculum. The Journal of
Neuroscience *31*, 12104–12117. https://doi.org/10.1523/JNEUROSCI.1370-11.2011.

8.   Zhang, X., Zhong, W., Brankačk, J., Weyer, S.W., Müller, U.C., Tort, A.B.L., and
Draguhn, A. (2016). Impaired theta-gamma coupling in APP-deficient mice. Sci Rep *6*,
21948. https://doi.org/10.1038/srep21948.

9.   Wulff, P., Ponomarenko, A.A., Bartos, M., Korotkova, T.M., Fuchs, E.C., Bähner, F.,
Both, M., Tort, A.B.L., Kopell, N.J., Wisden, W., et al. (2009). Hippocampal theta
rhythm and its coupling with gamma oscillations require fast inhibition onto
parvalbumin-positive interneurons. Proceedings of the National Academy of Sciences
106, 3561–3566. https://doi.org/10.1073/pnas.0813176106.

10. Schomburg, E.W., Fernández-Ruiz, A., Mizuseki, K., Berényi, A., Anastassiou, C.A.,
Koch, C., and Buzsáki, G. (2014). Theta Phase Segregation of Input-Specific Gamma
Patterns in Entorhinal-Hippocampal Networks. Neuron *84*, 470–485.
https://doi.org/10.1016/j.neuron.2014.08.051.

11. Lisman, J.E., and Jensen, O. (2013). The Theta-Gamma Neural Code. Neuron *77*,
1002–1016. https://doi.org/10.1016/j.neuron.2013.03.007.

12. Tzilivaki, A., Tukker, J.J., Maier, N., Poirazi, P., Sammons, R.P., and Schmitz, D.
(2023). Hippocampal GABAergic interneurons and memory. Neuron *111*, 3154–3175.
https://doi.org/10.1016/j.neuron.2023.06.016.

13. Topolnik, L., and Tamboli, S. (2022). The role of inhibitory circuits in hippocampal
memory processing. Nat Rev Neurosci *23*, 476–492. https://doi.org/10.1038/s41583-
022-00599-0.

14. Cardin, J.A., Carlén, M., Meletis, K., Knoblich, U., Zhang, F., Deisseroth, K., Tsai, L.-
H., and Moore, C.I. (2009). Driving fast-spiking cells induces gamma rhythm and
controls sensory responses. Nature *459*, 663–667.
https://doi.org/10.1038/nature08002.

15. Pelkey, K.A., Chittajallu, R., Craig, M.T., Tricoire, L., Wester, J.C., and McBain, C.J.
(2017). Hippocampal gabaergic inhibitory interneurons. Physiol Rev *97*, 1619–1747.
https://doi.org/10.1152/physrev.00007.2017.

16.   Allen, K., and Monyer, H. (2015). Interneuron control of hippocampal oscillations. Curr
Opin Neurobiol *31*, 81–87. https://doi.org/10.1016/j.conb.2014.08.016.

17.   Lucas, E.K., and Clem, R.L. (2018). GABAergic interneurons: The orchestra or the
conductor in fear learning and memory? Brain Res Bull *141*, 13–19.
https://doi.org/10.1016/j.brainresbull.2017.11.016.

18.   Katona, G., Kaszás, A., Turi, G.F., Hájos, N., Tamás, G., Vizi, E.S., and Rózsa, B.
(2011). Roller Coaster Scanning reveals spontaneous triggering of dendritic spikes in
CA1 interneurons. Proc Natl Acad Sci U S A *108*, 2148–2153.
https://doi.org/10.1073/pnas.1009270108.

19.   Judák, L., Chiovini, B., Juhász, G., Pálfi, D., Mezriczky, Z., Szadai, Z., Katona, G.,
Szmola, B., Ócsai, K., Martinecz, B., et al. (2022). Sharp-wave ripple doublets induce
complex dendritic spikes in parvalbumin interneurons in vivo. Nat Commun *13*, 6715.
https://doi.org/10.1038/s41467-022-34520-1.

20.   Tzilivaki, A., Kastellakis, G., Schmitz, D., and Poirazi, P. (2022). GABAergic
Interneurons with Nonlinear Dendrites: From Neuronal Computations to Memory
Engrams. Neuroscience *489*, 34–43.
https://doi.org/10.1016/j.neuroscience.2021.11.033.

21.   Cornford, J.H., Mercier, M.S., Leite, M., Magloire, V., Häusser, M., and Kullmann, D.M.
(2019). Dendritic NMDA receptors in parvalbumin neurons enable strong and stable
neuronal assemblies. Elife *8*. https://doi.org/10.7554/eLife.49872.

22.   Katona, G., Kaszás, A., Turi, G.F., Hájos, N., Tamás, G., Vizi, E.S., and Rózsa, B.
(2011). Roller Coaster Scanning reveals spontaneous triggering of dendritic spikes in
CA1 interneurons. Proc Natl Acad Sci U S A *108*, 2148–2153.
https://doi.org/10.1073/pnas.1009270108.

23.   Tzilivaki, A., Kastellakis, G., and Poirazi, P. (2019). Challenging the point neuron
dogma: FS basket cells as 2-stage nonlinear integrators. Nat Commun *10*, 3664.
https://doi.org/10.1038/s41467-019-11537-7.

24.   Tran-Van-Minh, A., Cazé, R.D., Abrahamsson, T., Cathala, L., Gutkin, B.S., and
DiGregorio, D. a (2015). Contribution of sublinear and supralinear dendritic integration
to neuronal computations. Front Cell Neurosci *9*, 67.
https://doi.org/10.3389/fncel.2015.00067.

25.   Morabito, Z. et al. (2024). A dendritic substrate for temporal diversity of cortical
inhibition. bioRxiv.

26.   Hu, H., and Vervaeke, K. (2018). Synaptic Integration in Cortical Inhibitory Neuron
Dendrites. Neuroscience *368*, 115–131.
https://doi.org/10.1016/J.NEUROSCIENCE.2017.06.065.

27.   Chamberland, S., and Topolnik, L. (2012). Inhibitory control of hippocampal inhibitory
neurons. Front Neurosci *6*, 1–13. https://doi.org/10.3389/fnins.2012.00165.

28.   Klausberger, T., and Somogyi, P. (2008). Neuronal Diversity and Temporal Dynamics:
The Unity of Hippocampal Circuit Operations. Science (1979) *321*.

29.   Hadler, M.D., Tzilivaki, A., Schmitz, D., Alle, H., and Geiger, J.R.P. (2024). Gamma
oscillation plasticity is mediated via parvalbumin interneurons. Sci Adv *10*.
https://doi.org/10.1126/sciadv.adj7427.

30. Park, K., Lee, J., Jang, H.J., Richards, B.A., Kohl, M.M., and Kwag, J. (2020).
Optogenetic activation of parvalbumin and somatostatin interneurons selectively
restores theta-nested gamma oscillations and oscillation-induced spike timing-
dependent long-term potentiation impaired by amyloid β oligomers. BMC Biol *18*, 7.
https://doi.org/10.1186/s12915-019-0732-7.

31. Turi, G.F., Li, W.K., Chavlis, S., Pandi, I., O'Hare, J., Priestley, J.B., Grosmark, A.D.,
Liao, Z., Ladow, M., Zhang, J.F., et al. (2019). Vasoactive Intestinal Polypeptide-
Expressing Interneurons in the Hippocampus Support Goal-Oriented Spatial Learning.
Neuron *101*, 1150-1165.e8. https://doi.org/10.1016/j.neuron.2019.01.009.

32. Tort, A.B.L., Kramer, M.A., Thorn, C., Gibson, D.J., Kubota, Y., Graybiel, A.M., and
Kopell, N.J. (2008). Dynamic cross-frequency couplings of local field potential
oscillations in rat striatum and hippocampus during performance of a T-maze task.
Proceedings of the National Academy of Sciences *105*, 20517–20522.
https://doi.org/10.1073/pnas.0810524105.

33. Tzilivaki, A., Kastellakis, G., and Poirazi, P. (2019). Challenging the point neuron
dogma: FS basket cells as 2-stage nonlinear integrators. Nat Commun *10*, 3664.
https://doi.org/10.1038/s41467-019-11537-7.

34. Cornford, J.H., Mercier, M.S., Leite, M., Magloire, V., Häusser, M., and Kullmann, D.M.
(2019). Dendritic NMDA receptors in parvalbumin neurons enable strong and stable
neuronal assemblies. Elife *8*. https://doi.org/10.7554/eLife.49872.

35. Kriener, B., Hu, H., and Vervaeke, K. (2022). Parvalbumin interneuron dendrites
enhance gamma oscillations. Cell Rep *39*, 110948.
https://doi.org/10.1016/j.celrep.2022.110948.

36. Larkum, M.E., and Nevian, T. (2008). Synaptic clustering by dendritic signalling
mechanisms. Curr Opin Neurobiol *18*, 321–331.
https://doi.org/10.1016/j.conb.2008.08.013.

37. Hu, H., Gan, J., and Jonas, P. (2014). Interneurons. Fast-spiking, parvalbumin[+]
GABAergic interneurons: from cellular design to microcircuit function. Science *345*,
1255263. https://doi.org/10.1126/science.1255263.

38. Hu, H., and Vervaeke, K. (2018). Synaptic Integration in Cortical Inhibitory Neuron
Dendrites. Neuroscience *368*, 115–131.
https://doi.org/10.1016/J.NEUROSCIENCE.2017.06.065.

39. Tort, A.B.L., Komorowski, R., Eichenbaum, H., and Kopell, N. (2010). Measuring
Phase-Amplitude Coupling Between Neuronal Oscillations of Different Frequencies. J
Neurophysiol *104*, 1195–1210. https://doi.org/10.1152/jn.00106.2010.

40. Scheffzük, C., Kukushka, V.I., Vyssotski, A.L., Draguhn, A., Tort, A.B.L., and Brankačk,
J. (2011). Selective Coupling between Theta Phase and Neocortical Fast Gamma
Oscillations during REM-Sleep in Mice. PLoS One *6*, e28489.
https://doi.org/10.1371/journal.pone.0028489.

41. Scheffer-Teixeira, R., Belchior, H., Caixeta, F. V., Souza, B.C., Ribeiro, S., and Tort,
A.B.L. (2012). Theta Phase Modulates Multiple Layer-Specific Oscillations in the CA1
Region. Cerebral Cortex *22*, 2404–2414. https://doi.org/10.1093/cercor/bhr319.

42. Belluscio, M.A., Mizuseki, K., Schmidt, R., Kempter, R., and Buzsáki, G. (2012).
Cross-Frequency Phase–Phase Coupling between Theta and Gamma Oscillations in

the Hippocampus. The Journal of Neuroscience *32*, 423–435.
https://doi.org/10.1523/JNEUROSCI.4122-11.2012.

43.   Park, K., Lee, J., Jang, H.J., Richards, B.A., Kohl, M.M., and Kwag, J. (2020).
Optogenetic activation of parvalbumin and somatostatin interneurons selectively
restores theta-nested gamma oscillations and oscillation-induced spike timing-
dependent long-term potentiation impaired by amyloid β oligomers. BMC Biol *18*, 7.
https://doi.org/10.1186/s12915-019-0732-7.

44.   Canolty, R.T., Edwards, E., Dalal, S.S., Soltani, M., Nagarajan, S.S., Kirsch, H.E.,
Berger, M.S., Barbaro, N.M., and Knight, R.T. (2006). High Gamma Power Is Phase-
Locked to Theta Oscillations in Human Neocortex. Science (1979) *313*, 1626–1628.
https://doi.org/10.1126/science.1128115.

45.   Tzilivaki, A., Tukker, J.J., Maier, N., Poirazi, P., Sammons, R.P., and Schmitz, D.
(2023). Hippocampal GABAergic interneurons and memory. Neuron *111*, 3154–3175.
https://doi.org/10.1016/j.neuron.2023.06.016.

46.   Topolnik, L., and Tamboli, S. (2022). The role of inhibitory circuits in hippocampal
memory processing. Nat Rev Neurosci *23*, 476–492. https://doi.org/10.1038/s41583-
022-00599-0.

47.   Allen, K., and Monyer, H. (2015). Interneuron control of hippocampal oscillations. Curr
Opin Neurobiol *31*, 81–87. https://doi.org/10.1016/j.conb.2014.08.016.

48.   Klausberger, T., and Somogyi, P. (2008). Neuronal Diversity and Temporal Dynamics:
The Unity of Hippocampal Circuit Operations. Science (1979) *321*.

49.   Judák, L., Chiovini, B., Juhász, G., Pálfi, D., Mezriczky, Z., Szadai, Z., Katona, G.,
Szmola, B., Ócsai, K., Martinecz, B., et al. (2022). Sharp-wave ripple doublets induce
complex dendritic spikes in parvalbumin interneurons in vivo. Nat Commun *13*, 6715.
https://doi.org/10.1038/s41467-022-34520-1.

50.   Memmesheimer, R.-M. (2010). Quantitative prediction of intermittent high-frequency
oscillations in neural networks with supralinear dendritic interactions. Proceedings of
the National Academy of Sciences *107*, 11092–11097.
https://doi.org/10.1073/pnas.0909615107.

51.   Zerlaut, Y., and Tzilivaki, A. (2025). Interneuronal modulations as a functional switch
for cortical computations: mechanisms and implication for disease. Front Cell
Neurosci *18*. https://doi.org/10.3389/fncel.2024.1479579.

52.   Tzilivaki, A., Kastellakis, G., Schmitz, D., and Poirazi, P. (2022). GABAergic
Interneurons with Nonlinear Dendrites: From Neuronal Computations to Memory
Engrams. Neuroscience *489*, 34–43.
https://doi.org/10.1016/j.neuroscience.2021.11.033.

53.   Liao, Z., Gonzalez, K.C., Li, D.M., Yang, C.M., Holder, D., McClain, N.E., Zhang, G.,
Evans, S.W., Chavarha, M., Yi, J., et al. (2024). Functional architecture of intracellular
oscillations in hippocampal dendrites. bioRxiv, 2024.02.12.579750.

54.   Hines, M.L., and Carnevale, N.T. (1997). The NEURON Simulation Environment.
Neural Comput *9*, 1179–1209. https://doi.org/10.1162/neco.1997.9.6.1179.

55.   Tukker, J.J., Fuentealba, P., Hartwich, K., Somogyi, P., and Klausberger, T. (2007).
Cell type-specific tuning of hippocampal interneuron firing during gamma oscillations

in vivo. The Journal of Neuroscience *27*, 8184–8189.
https://doi.org/10.1523/{JNEUROSCI}.1685-07.2007.

56.   Sun, Q., Sotayo, A., Cazzulino, A.S., Snyder, A.M., Denny, C.A., and Siegelbaum,
S.A. (2017). Proximodistal Heterogeneity of Hippocampal CA3 Pyramidal Neuron
Intrinsic Properties, Connectivity, and Reactivation during Memory Recall. Neuron *95*,
656-672.e3. https://doi.org/10.1016/j.neuron.2017.07.012.

57.   Banke, T.G., and McBain, C.J. (2006). GABAergic Input onto CA3 Hippocampal
Interneurons Remains Shunting throughout Development. The Journal of
Neuroscience *26*, 11720–11725. https://doi.org/10.1523/JNEUROSCI.2887-06.2006.

58.   Fukushima, F., Nakao, K., Shinoe, T., Fukaya, M., Muramatsu, S., Sakimura, K.,
Kataoka, H., Mori, H., Watanabe, M., Manabe, T., et al. (2009). Ablation of NMDA
Receptors Enhances the Excitability of Hippocampal CA3 Neurons. PLoS One *4*,
e3993. https://doi.org/10.1371/journal.pone.0003993.

59.   Kohus, Z., Káli, S., Rovira-Esteban, L., Schlingloff, D., Papp, O., Freund, T.F., Hájos,
N., and Gulyás, A.I. (2016). Properties and dynamics of inhibitory synaptic
communication within the CA3 microcircuits of pyramidal cells and interneurons
expressing parvalbumin or cholecystokinin. J Physiol *594*, 3745–3774.
https://doi.org/10.1113/JP272231.

60.   Wang, H.-X., and Gao, W.-J. (2009). Cell type-specific development of NMDA
receptors in the interneurons of rat prefrontal cortex. Neuropsychopharmacology *34*,
2028–2040. https://doi.org/10.1038/npp.2009.20.

61.   Turi, G.F., Li, W.-K., Chavlis, S., Pandi, I., O'Hare, J., Priestley, J.B., Grosmark, A.D.,
Liao, Z., Ladow, M., Zhang, J.F., et al. (2019). Vasoactive Intestinal Polypeptide-
Expressing Interneurons in the Hippocampus Support Goal-Oriented Spatial Learning.
Neuron *101*, 1150-1165.e8. https://doi.org/10.1016/j.neuron.2019.01.009.

62.   Vladimirov, N., Tu, Y., and Traub, R.D. (2013). Synaptic gating at axonal branches,
and sharp-wave ripples with replay: a simulation study. European Journal of
Neuroscience *38*, 3435–3447. https://doi.org/10.1111/ejn.12342.

63.   Bacci, A., Rudolph, U., Huguenard, J.R., and Prince, D.A. Cellular/Molecular Major
Differences in Inhibitory Synaptic Transmission onto Two Neocortical Interneuron
Subclasses.

64.   Netsyk, O., Hammoud, H., Korol, S. V., Jin, Z., Tafreshiha, A.S., and Birnir, B. (2020).
Tonic GABA -activated synaptic and extrasynaptic currents in dentate
gyrus granule cells and CA3 pyramidal neurons along the mouse
hippocampal dorsoventral axis. Hippocampus *30*, 1146–1157.
https://doi.org/10.1002/hipo.23245.

65.   Goldberg, J.H., Yuste, R., and Tamas, G. (2003). Ca2+ imaging of mouse neocortical
interneurone dendrites: contribution of Ca2+-permeable AMPA and NMDA receptors
to subthreshold Ca2+dynamics. J Physiol *551*, 67–78.
https://doi.org/10.1113/jphysiol.2003.042598.

66.   Goldberg, J.H., and Yuste, R. (2005). Space matters: Local and global dendritic Ca2+
compartmentalization in cortical interneurons. Trends Neurosci *28*, 158–167.
https://doi.org/10.1016/j.tins.2005.01.005.

67.   Goldberg, J.H., and Yuste, R. (2005). Space matters: Local and global dendritic Ca2+
1153   compartmentalization in cortical interneurons. Trends Neurosci *28*, 158–167.
1154   https://doi.org/10.1016/j.tins.2005.01.005.

1155   68.   Nörenberg, A., Hu, H., Vida, I., Bartos, M., and Jonas, P. (2010). Distinct nonuniform
1156   cable properties optimize rapid and efficient activation of fast-spiking GABAergic
1157   interneurons. Proc Natl Acad Sci U S A *107*, 894–899.
1158   https://doi.org/10.1073/pnas.0910716107.

1159   69.   Bacci, A., Rudolph, U., Huguenard, J.R., and Prince, D.A. Cellular/Molecular Major
1160   Differences in Inhibitory Synaptic Transmission onto Two Neocortical Interneuron
1161   Subclasses.

1162   70.   Kohus, Z., Káli, S., Rovira-Esteban, L., Schlingloff, D., Papp, O., Freund, T.F., Hájos,
1163   N., and Gulyás, A.I. (2016). Properties and dynamics of inhibitory synaptic
1164   communication within the CA3 microcircuits of pyramidal cells and interneurons
1165   expressing parvalbumin or cholecystokinin. J Physiol *594*, 3745–3774.
1166   https://doi.org/10.1113/JP272231.

1167