Bimodal dendritic processing in basket cells drives distinct memory-related
oscillations

Alexandra Tzilivaki[1,2,3, #], Matthew Evan Larkum[1,2,3,4], Dietmar Schmitz[1,2,3,5,6,#]

[1]Charité-Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-
Universität Berlin, and Berlin Institute of Health, Neuroscience Research Center, 10117
Berlin, Germany

[2]Einstein Center for Neurosciences, Chariteplatz 1, 10117 Berlin, Germany

[3]NeuroCure Cluster of Excellence, Chariteplatz 1, 10117 Berlin, Germany

[4]Department of Biology, Humboldt Universität zu Berlin, Berlin 10117, Germany

[5]German Center for Neurodegenerative Diseases (DZNE), 10117 Berlin, Germany

[6]Bernstein Center for Computational Neuroscience, Humboldt-Universität zu Berlin,
Philippstrasse. 13, 10115 Berlin, Germany

Correspondence: AT alexandra.tzilivaki@charite.de , DS dietmar.schmitz@charite.de

Classification: Biological Sciences, Neuroscience

Keywords: Interneurons, Learning and memory, Hippocampal Oscillations, Dendrites,
Biophysical model

Abstract

Hippocampal oscillations regulate distinct memory stages and behavioral states, yet the
mechanisms by which interneurons contribute to these rhythms remain elusive. Here, we
establish a novel link between the bimodal nonlinear dendritic processing of PV+ fast-spiking
basket cells (FSBCs) and hippocampal oscillations. Our model predicts that FSBCs can
flexibly adjust their activity without changes in synaptic input quantity, leveraging their
supralinear and sublinear dendritic branches to selectively shape network dynamics.
Supralinear dendritic activation enhances high-frequency oscillations, while sublinear
dendritic activation amplifies slow oscillatory power, dynamically modulating the
excitation/inhibition (E/I) balance. These findings reveal a previously unrecognized role of
FSBC dendritic integration in memory-related oscillatory dynamics, offering new insights into
the cellular mechanisms governing hippocampal rhythm generation.

Significance Statement

Hippocampal oscillations play a crucial role in memory processes, yet how interneurons
dynamically regulate these rhythms remains unresolved. Our study reveals that fast-spiking
basket cells (FSBCs) exploit bimodal nonlinear dendritic processing to selectively shape
slow and fast oscillatory components, independently of synaptic input quantity. This energy-
efficient mechanism allows FSBCs to fine-tune the excitation-inhibition balance, modulating
network dynamics essential for learning and consolidation. By challenging the conventional
view of interneuron function, our findings redefine how inhibitory circuits contribute to
hippocampal rhythm generation, providing new insights into the cellular basis of memory-
related oscillations.

Main

Hippocampal oscillations exhibit a variety of distinct frequency bands (Buzsáki and Draguhn, 2004). For example, neuronal activity often associates with a slow local field potential (LFP) component, known as theta activity, which ranges from 3 to 10 Hz (Buzsáki and Draguhn, 2004; Jackson and Skinner, 2018; Lisman and Jensen, 2013). Additionally, hippocampal activity includes fast-frequency oscillations spanning from 30 to 200 Hz (Buzsáki and Draguhn, 2004; Griffiths and Jensen, 2023; Jackson and Skinner, 2018; Lisman and Jensen, 2013; Tort et al., 2013). Both experimental and computational studies have linked these slow and fast components to different stages of hippocampal memory formation and distinct behavioral states (Buzsáki, 1989; Buzsáki and Wang, 2012; Lisman and Jensen, 2013). Specifically, slow 3-10 Hz oscillations correlate with memory encoding, spatial navigation, and sensory processing (Lisman and Jensen, 2013). Conversely, fast oscillations are associated with memory encoding and consolidation (Buzsáki and Wang, 2012; Lisman and Jensen, 2013). Notably, these fast and slow components often occur together in a phenomenon known as oscillation coupling (Jackson et al., 2011; Schomburg et al., 2014; Tort et al., 2013; Wulff et al., 2009; Zhang et al., 2016; Zhong et al., 2017) which has been proposed as a signature of working memory processing and various behavioral states (**Figure.1a**) (Lisman and Jensen, 2013; Tort et al., 2013). To understand the cellular mechanisms behind these oscillatory events, significant efforts have been made to delineate the effects of various subclasses of both excitatory and inhibitory hippocampal populations. Among these, PV+ GABAergic interneurons play a crucial role in both slow and fast oscillations as well as in coupled oscillatory activity (Allen and Monyer, 2015; Cardin et al., 2009; Lucas and Clem, 2018; Pelkey et al., 2017a; Topolnik and Tamboli, 2022; Tzilivaki et al., 2023; Wulff et al., 2009). However, most research has focused on manipulating PV+ interneurons through optogenetic and computational approaches, primarily investigating their activation or silencing during oscillatory activity. While these studies have significantly advanced our understanding of the importance of PV+ interneurons in hippocampal oscillations, they have provided little insight into the specific subcellular properties that enable these interneurons to exert such precise control over distinct oscillatory states and memory processes.

For many years, a prevailing dogma dominated the field of interneuron research, suggesting that interneurons possessed passive, linear dendrites incapable of integrating synaptic inputs in a nonlinear manner. According to this view, interneurons were thought to exert inhibition based purely on the quantity of synaptic inputs they received. This simplistic framework assumed that the inhibitory regulation of a network's activity across distinct oscillatory activity depends solely on the synaptic input amount that interneurons receive, disregarding the potential influence of potential complex nonlinear dendritic integration rules. Contrary to this linear model, recent experimental findings have revealed that hippocampal PV+ interneurons exhibit complex nonlinear dendritic computations, significantly enhancing their computational capabilities and shaping their input-output (I/O) transformations (Chiovini et al., 2014; Cornford et al., 2019; Hu and Vervaeke, 2018a; Judák et al., 2022; Katona et al., 2011; Morabito, 2024; Tzilivaki et al., 2019, 2022). Nonlinear dendrites allow neurons to function as intrinsic computational units, enabling them to differentially respond to incoming synaptic inputs (Tran-Van-Minh et al., 2015). While nonlinear dendritic integration has long been recognized in pyramidal cells and is known to optimize their role in memory formation, its presence in interneurons has only recently gained attention. To date, most available data come from studies on hippocampal PV+ interneurons (Cornford et al., 2019; Katona et al., 2011), a highly diverse class that includes distinct subtypes, such as aspiny fast-spiking basket cells (FSBCs), which primarily target the somata of pyramidal neurons and other interneurons (Pelkey et al., 2017b; Topolnik and Tamboli, 2022; Tzilivaki et al., 2023).

In the case of hippocampal FSBCs, biophysical computational modeling has proposed that their dendritic trees exhibit two distinct modes of nonlinear integration, both coexisting within the same dendritic tree (Tzilivaki et al., 2019, 2022). Specifically, multicompartmental biophysical simulations indicate that FSBCs possess two types of nonlinear dendrites: supralinear dendrites, capable of generating local dendritic spikes, and sublinear dendrites, which integrate excitatory synaptic input passively (**Figures. 1b-c, S2a-d**). This model suggests that FSBCs engage in bimodal nonlinear dendritic processing, leveraging both supralinear and sublinear integration mechanisms. Although both types of nonlinear dendrites coexist within the same dendritic tree (**Figure. 1b-c, S2**) and share similar active properties, they differ in morphology and function (Tzilivaki et al., 2019). Supralinear branches exhibit larger volumes and lower input resistance, allowing them to generate local sodium spikes (**Figure. S2a-e**). In contrast, sublinear branches found to have smaller volumes and higher input resistance. Given their small volume and high input resistance (**Figure S2**), sublinear dendrites are structurally constrained from generating spikes. When coincident synaptic inputs arrive at these branches, they produce large, rapidly rising excitatory postsynaptic potentials (EPSPs). However, the presence of A-type potassium channels facilitates rapid membrane repolarization, effectively preventing local spike initiation.

Notably, the theoretical prediction of bimodal nonlinear dendritic integration in FSBCs, a major subclass of hippocampal PV+ interneurons, was later supported experimentally. Specifically, studies in CA1 PV+ interneurons provided direct evidence of bimodal nonlinear dendritic processing, demonstrating the presence of both supralinear and sublinear dendrites (Cornford et al., 2019). This discovery challenges traditional assumptions about interneuron function, emphasizing the critical role of dendritic nonlinearities in shaping hippocampal network dynamics. While further research is needed to fully unravel the dendritic processing patterns of other interneuron subtypes, existing evidence strongly supports the idea that the complex nonlinear dendritic behavior of PV+ FSBCs serves functions beyond the traditional view of simply regulating network excitation.

A fundamental question that arises—and one that inspired this study—is: How do FSBCs leverage their bimodal nonlinear dendrites at the network level during memory encoding and recall related oscillations? Why are these cells equipped with two distinct types of nonlinear dendrites rather than just one? What are the network-level implications of supralinear and sublinear activation during different memory-related oscillations? Given the involvement of FSBCs in both slow (e.g., theta-frequency) and fast oscillations (Chamberland and Topolnik, 2012; Klausberger and Somogyi, 2008; Tzilivaki et al., 2023), we employed state-of-the-art biophysical modeling to test our hypothesis: that the supralinear and sublinear branches of FSBCs differentially modulate their activity and, consequently, orchestrate local field potentials (LFPs) during mnemonic functions. Specifically, we hypothesized that FSBCs exploit their bimodal dendritic processing to selectively shape fast and slow oscillatory activity within a hippocampal circuit **(Figures 1-2)**. This mechanism could allow FSBCs to fine-tune the excitation-inhibition (E/I) balance and modulate both slow and fast oscillatory signals without requiring changes in synaptic input quantity **(Figure 3)**. Such a strategy would maximize the flexibility of this interneuron subtype, enabling it to engage with distinct memory-related rhythms in an energy-efficient manner.

To test our hypothesis, we utilized a published biologically plausible hippocampal microcircuit model (Hadler et al., 2024), which incorporates compartmental biophysical models of pyramidal cells (PCs) with simplified morphologies (Hadler et al., 2024) and FSBCs with anatomically reconstructed bimodal nonlinear dendritic trees (Tzilivaki et al., 2019) (**Tables S1–S5, Figures. 1a, S1, S2**). Both neuronal types were extensively calibrated based on

electrophysiological data (**Tables S1-S5, Figure S1**). Given that, to our knowledge,
experimental evidence on dendritic integration in other interneuron subtypes (e.g., SOM+ or
VIP+) remains insufficient, we sought to avoid assumptions. Therefore, we exclusively
modelled FSBCs with bimodal nonlinear dendritic trees as the only inhibitory component of
our microcircuit model.

Upon activation with a theta-like input (as per Park et al., 2020; Turi et al., 2019), our model
recapitulated the experimentally observed coupling of slow and fast oscillations (Schomburg
et al., 2014; Tort et al., 2008; Wulff et al., 2009), generating a slow oscillatory component
alongside a fast oscillatory component (**Figure. 1a, d**). To systematically investigate how
FSBC dendritic integration modes influence network dynamics, we designed three distinct
activation conditions while ensuring that the total number of synaptic inputs remained
constant across all simulations: 1. *Bimodal Activation* (Control Condition): Synapses onto
FSBCs were randomly distributed across both supralinear and sublinear dendritic branches
(**Figures. 1a, d, 2a**). This represents a balanced scenario where inputs arrive at both
dendritic types. 2. *Supralinear Activation*: To understand the effects of supralinear dendrites
in these cells, we restricted all synaptic inputs exclusively to supralinear branches (**Figures.**
1e, 2b). 3. *Sublinear Activation*: Similarly, to examine the role of sublinear dendritic
processing, we constrained all synaptic inputs to only sublinear branches (**Figures. 1f, 2c**).
In all three conditions, the total number of synaptic inputs remained the same and identical,
and all other network parameters and simulation settings were kept unchanged to ensure a
controlled comparison of how the two types of nonlinear dendrites shape FSBCs firing
patterns and network oscillatory behavior.

Our model predicts that supralinear dendritic activation in FSBCs preferentially amplifies the
fast oscillatory component of our LFP signal, whereas sublinear dendritic activation
enhances the slow LFP (**Figures 1-2**). Specifically, when supralinear dendrites are activated,
the network exhibits higher fast peak-frequency values as well as higher fast peak-frequency
power values compared to both bimodal and sublinear activation conditions (**Figures. 1e,**
2b, 2d, 2e). In sharp contrast, sublinear FSBCs dendritic activation significantly reduces the
fast peak-frequency power relative to both bimodal and supralinear activation (**Figure. 2d**),
while simultaneously increasing slow LFP power values (**Figures. 1f, 2c, 2e, 2f**). These
predictions suggest that FSBCs leverage their bimodal dendritic architecture to selectively
regulate distinct oscillatory components, dynamically shaping hippocampal network activity
based on input distribution.

To assess the robustness of the differential impact of supralinear versus sublinear dendritic
activation on LFP oscillations, we performed a sensitivity analysis on synaptic parameters
and input characteristics. In all cases, supralinear activation enhanced both peak power and
peak frequency power of the fast LFP component, whereas sublinear activation consistently
resulted in stronger slow oscillatory power compared to supralinear activation (**Figure, S3**).

Figure 1. Supralinear and Sublinear branches of FSBCs differentially modulate memory-related hippocampal oscillations. *Schematic Illustration of the hippocampal microcircuit model. Pyramidal Cells (PCs) and FSBCs populations are activated by a theta (4 Hz) input for 1000 msec. An LFP electrode is positioned nearby the PCs somata. Upon activation the model exhibits slow (bandpassed at 3-10 Hz) and fast (bandpassed at 30-200 Hz) LFP bands. The arrow indicated the in-silico electrode position. FSBCs are equipped with bimodal nonlinear trees as per* (Cornford et al., 2019; Tzilivaki et al., 2019). ***(b-c).*** *Representative input-output curves from supralinear (**b**) and sublinear (**c**) dendritic branches in FSBC models, in response to synaptic stimulation. Increasing numbers of synapses (from 1 to 20 with step☐=☐1) are uniformly distributed in every stimulated branch and are activated with a single pulse. The y-axis shows the amplitude of the dendritic EPSP caused by synaptic activation while the x-axis shows the expected EPSP amplitude that would result from the linear summation of synaptic EPSPs. The dashed line indicates linear summation. Insets show representative traces. Panels were adopted from Tzilivaki et al., 2019.* ***(d-f).*** *Power Spectrum Density (PSD) plots under 3 synaptic placement configurations: when both supralinear and sublinear dendrites of the FSBCs receive synapses (bimodal activation) (**d**), when only the supralinear branches receive synapses (supralinear activation) (**e**), and when only the sublinear branches receive synapses (sublinear activation) (**f**). The number of activated synapses remain identical for all 3 configurations. Activation of the supralinear dendrites (**e**) supports the fast LFP signature while sublinear activation (**f**) promotes slow LFP power. Data from 30 random simulation trials.*

Figure 2. LFP frequency and power are regulated by FSBC nonlinear dendritic activation. *(a-c) Schematic illustration of the activation protocols and representative traces of the evoked LFP bandpassed at slow (theta 3-10 Hz) and fast (30-200 Hz) frequences. The same number of synapses targets either both types of nonlinear dendrites (bimodal activation) (a) or only supralinear (b) or only sublinear (c). (d-f) Comparison of the peak fast-frequency (d) and the peak powers of the fast (e) and slow (f) LFP components. Data from 30 random simulation trials. Synaptic activation of supralinear dendrites of the FSBCs bimodal trees results in a higher peak fast-frequency and increased fast-frequency peak power, while reducing the peak power of slow oscillations compared to sublinear dendrite activation. Statistical comparisons across multiple groups were performed using the Kruskal-Wallis test followed by a post-hoc correction for multiple comparisons, suitable for data with unequal variance.*

Given that the total number of synapses was identical across all three activation conditions
(bimodal, supralinear, and sublinear), we sought to understand the factors driving the
observed LFP modulation. To achieve this, we analyzed the spiking activity of both pyramidal
cells (PCs) and FSBCs under these conditions (**Figure. 3a-c**). Our analysis showed that
under supralinear activation, FSBCs exhibited significantly higher firing rates compared to
the bimodal and sublinear conditions (**Figure. 3a, c**). This increase in FSBC activity led to a
suppression of PC firing, as observed in the lower firing frequencies of PCs under this
condition (**Figure. 3b, c**). Conversely, under sublinear activation, FSBC firing rates were
reduced, resulting in disinhibition of the PCs and subsequently higher PC firing frequencies
compared to the supralinear condition (**Figure. 3a, b, c**). The bimodal activation condition, in
which synapses were distributed across both dendritic types, displayed intermediate firing
rates for both FSBCs and PCs, reflecting a balance between the effects observed in the
supralinear and sublinear conditions (**Figure. 3c**). These results demonstrate that dendritic
nonlinearities in FSBCs enable differential modulation of network activity, with supralinear
dendrites suppressing PC activity via enhanced FSBCs firing and sublinear dendrites
facilitating PC activity through reduced FSBC firing. This dynamic regulation of the E/I
balance highlights the functional significance of FSBCs' bimodal nonlinear dendritic
architecture in shaping hippocampal network oscillatory dynamics.

Figure 3 **Supralinear and sublinear FSBC dendrites differentially modulate the E/I**
balance in the microcircuit under identical synaptic activation conditions. (a)
Representative raster plots depicting the spiking activity of pyramidal cells (PCs) (blue) and
FSBCs (brown) across theta cycles when synaptic inputs are selectively targeted to either
supralinear dendrites (upper panel) or sublinear dendrites (lower panel) of FSBCs. (b)
Representative spike trains of pyramidal cells (left) and FSBCs (right). When FSBCs receive
synaptic input in their supralinear branches, their firing rates increase across theta cycles,
leading to a decrease in PC spiking. Conversely, when synaptic inputs are restricted to

sublinear dendrites, FSBC firing rates decrease, resulting in increased pyramidal cell activity.
(c) *Activation of a fixed number of synapses on randomly selected supralinear dendrites*
significantly increases the firing frequency of FSBCs, leading to a reduction in the activity of
PCs when compared to bimodal (supralinear and sublinear) or purely sublinear activation.
Conversely, activation of randomly selected sublinear dendrites decreases the firing
frequency of FSBCs, resulting in disinhibition of the PCs. Data are based on 30 random
simulation trials. Pairwise comparisons were performed using the Mann–Whitney U test.

These observations demonstrate that FSBCs can dynamically adapt their firing patterns
depending on whether synaptic input is received by their supralinear or sublinear dendritic
branches, despite the total synaptic input remaining constant. Supralinear branches,
characterized by larger volumes and lower input resistance (Tzilivaki et al., 2019), facilitate
enhanced forward propagation of synaptic signals (**Figure S2g**). In contrast, sublinear
branches, which have smaller volumes and higher input resistance (Tzilivaki et al., 2019),
limit forward propagation (**Figure. S2f**), leading to a decrease in FSBCs firing rates.
Consequently, this differential activation modulates the E/I balance in the network, directly
shaping the oscillatory activity observed in the LFP. These findings indicate that PV+ FSBCs
leverage their bimodal nonlinear dendrites to differentially process incoming synaptic inputs,
even when the total synaptic input remains unchanged. This mechanism not only modulates
FSBC firing responses at the cellular level (**Figure 3**) but also plays a crucial role in
determining the network's oscillatory state, thereby influencing hippocampal rhythm
dynamics.

We further explored how altering the dendritic integration profile of FSBCs, specifically by
manipulating them to possess purely supralinear or sublinear trees, impacts network
behavior compared to the control (bimodal) trees (**Figure. S4a-b**). Leveraging the distinct
morphological features of supralinear and sublinear branches in our bimodal FSBCs
reconstructions, we adjusted the dendritic morphologies as per (Tzilivaki et al., 2019). When
FSBCs were equipped with purely supralinear dendritic trees, we observed enhanced fast
frequency and power compared to both bimodal and sublinear tree configurations. This
configuration also maximized the firing response of FSBCs (**Figure S4a, c-h**). Conversely,
FSBCs with purely sublinear dendritic trees present a reduced spike rate which increased
the PCs activity (**Figure. S4h**). Sublinear FSBC trees exhibited increased slow power and
decreased fast power and lower fast power compared to supralinear trees (**Figure. S4b-g**).

These simulations underscore that FSBCs take advantage of their bimodal nonlinear
dendritic trees by dynamically utilizing their supralinear and sublinear dendrites to regulate
the E/I balance within the network, thereby orchestrating distinct oscillatory behaviors under
equivalent synaptic input conditions. Importantly, activating either only supralinear or only
sublinear branches of a bimodal tree (**Figs.1-2**) exhibits similar phenotypic responses to
hypothetical configurations of purely supralinear or purely sublinear trees (**Fig.S3**). This
further supports the notion that the bimodal nonlinear nature of FSBCs dendrites allows for
flexible modulation of network activity and oscillatory patterns crucial for hippocampal
mnemonic functions.

Apart from the type of dendritic integration, the spatial arrangement of synapses also
influences the firing characteristics (Kastellakis et al., 2015). It's well-established that
supralinear dendrites tend to enhance clustered synaptic allocation, whereas sublinear
dendrites prefer dispersed activation. For FSBCs, it has been observed that dispersed
synaptic activation leads to higher firing frequencies compared to clustered (Hu et al., 2014;
Tzilivaki et al., 2019, 2022). This phenomenon is influenced by the co-existence of sublinear

along with supralinear branches, coupled with their small size and the presence of A-type potassium channels(Hu and Vervaeke, 2018b; Hu et al., 2014; Tzilivaki et al., 2019), which discourages preference for clustered inputs. To investigate how synaptic clustering affects the oscillatory behavior of our network, we simulated the activation of the same number of synapses as in our disperse protocols (as in **Figure. S4**) but now clustered in a few randomly chosen dendrites (**Figure. S5**). As anticipated, clustered activation of sublinear FSBC trees significantly reduced their firing rates while maximizing the E/I balance in the network. Conversely, clustering synapses in supralinear trees enhanced FSBC firing rates compared to both bimodal and sublinear clustering configurations, while also decreasing the firing rates of the PCs (**Figure. S5j**). At the oscillatory level, akin to our simulations with dispersed synaptic activation, clustered synapses in supralinear FSBC trees enhanced fast LFP components, whereas synaptic clustering in sublinear trees boosted the slow LFP power (**Figure. S5a-i**). These findings underscore the dual impact of dendritic morphology and synaptic spatial arrangement on neuronal firing patterns and network oscillatory dynamics. They highlight the intricate mechanisms by which FSBCs can dynamically modulate their responses and contribute to the regulation of network excitability and oscillatory activity crucial for hippocampal function. An important consideration for future experimental studies is whether supralinear and sublinear dendrites receive synaptic inputs from distinct hippocampal or cortical regions. Such compartmentalization or segregation of inputs could profoundly influence FSBC activity across different oscillatory and behavioral states (see Discussion).

Finally, we analyzed the Phase-Amplitude Coupling (PAC),(Tort et al., 2010) to check the slow-fast oscillation coupling in the LFP across all cases (**Figure. S6**). Indeed, the LFP response aligns with experimental findings, indicating that slow activity entrains hippocampal networks to slow-fast coupling (**Figure .S6a**)(Belluscio et al., 2012; Canolty et al., 2006; Park et al., 2020; Scheffer-Teixeira et al., 2012; Scheffzük et al., 2011; Tort et al., 2013; Zhang et al., 2016). Activation of either supralinear or sublinear branches did not alter the coupling between the two oscillatory components (**Figure. S6b-f**). However, coupling was enhanced when synapses were clustered on fully sublinear dendritic trees of FSBCs. Conversely, coupling was reduced when similar clustering occurred on supralinear FSBC dendritic trees (**Figure. S6g-j**). These results indicate that as the power and frequency of the fast component decrease, it becomes more synchronized with the slower component.

Discussion

While the role of interneurons in hippocampal memory-related oscillations is well established (Allen and Monyer, 2015; Klausberger and Somogyi, 2008; Pelkey et al., 2017b; Topolnik and Tamboli, 2022; Tzilivaki et al., 2023), a significant gap remains in understanding the cellular and subcellular mechanisms that enable these neurons to flexibly control memory-related rhythms. Recent studies have highlighted the involvement of PV+ interneuron dendrites in fast oscillations (Chiovini et al., 2014; Judák et al., 2022; Kriener et al., 2022), and suggest the potential advantage of supralinear dendritic integration (Memmesheimer, 2010), yet a direct link between interneuronal dendritic integration and its impact on slow and fast hippocampal memory-related rhythms remains unexplored. In particular, our biophysical modeling predictions reveal that hippocampal FSBCs, a prominent subtype of PV+ interneurons, exhibit distinct firing patterns that modulate the E/I balance independently of synaptic input quantity. Specifically, activation of supralinear dendritic branches in bimodal FSBCs (**Figures 1-2**) reduces the E/I balance (**Figure. 3**), thereby promoting fast oscillatory activity associated with memory consolidation (**Figures. 1-2**). In contrast, activation of

sublinear dendritic branches increases the E/I balance (**Figure. 3**), leading to an enhanced
slow LFP signature (**Figures. 1- 2**), which is linked to memory encoding. These new findings
redefine our understanding of FSBCs contributions to hippocampal memory oscillations, by
shedding light on how these interneurons exploit their intricate and highly adaptive nonlinear
dendritic trees. This work also sets the foundation for future experimental efforts aimed at
linking dendritic processing at the subcellular level to its broader effects on cellular and
network computations, particularly under the influence of GABAergic inhibition.

Despite significant progress in understanding interneuron function, we are still far from fully
linking subcellular properties of interneurons not only to memory states but also to
behavioral states, such as learning during spatial navigation and memory consolidation
during sleep. To comprehensively investigate how interneuron dendrites influence memory-
related oscillations, it is essential to explore the diversity of hippocampal interneurons. Within
the hippocampal formation alone, over 20 distinct interneuron subtypes have been identified,
each classified based on biophysical, morphological, developmental, and connectivity
profiles. Beyond FSBCs, other PV+ interneuron subclasses and different inhibitory neuron
classes (e.g., SOM+ and VIP+ interneurons) are known to participate in distinct memory-
related rhythms (Ascoli et al., 2008; Pelkey et al., 2017a; Sanchez-Aguilera et al., 2021).
However, a major gap remains in characterizing the dendritic integration mechanisms of
these diverse interneuron subtypes. To bridge this gap, experimental approaches such as
glutamate uncaging, dendritic patch clamp recordings and detailed biophysical modeling are
needed to determine the precise dendritic processing mechanisms and their corresponding
biophysical underpinnings. Expanding our knowledge in this area will enable the refinement
of computational models, such as the one presented in this study, to provide a detailed
framework for understanding how dendritic function across inhibitory and excitatory neuronal
subtypes shapes the oscillatory activity of the networks during different memory stages.

Additionally, future studies should investigate whether specific interneuron dendritic
compartments receive distinct synaptic inputs from different hippocampal and cortical areas,
as this could provide a critical link between dendritic function, interneuron activity, and
network oscillations associated with learning and memory. A critical unresolved question is
where and when nonlinear dendrites receive inputs during distinct behavioral and memory
stages. Specifically, in the case of bimodal nonlinear dendritic trees in PV+ FSBCs, it
remains unknown whether supralinear and sublinear branches are activated by different
axons originating from local hippocampal neurons or extrahippocampal sources, such as the
entorhinal cortex, septum, or locus coeruleus (Fernández-Ruiz et al., 2021; Pelkey et al.,
2017b; Topolnik and Tamboli, 2022; Tzilivaki et al., 2023). Given that functional connectivity
patterns within and beyond the hippocampus are thought to underlie distinct behavioral
states (e.g., active navigation vs. sleep-related consolidation), we believe it would be of great
interest to map the precise origins of these inputs. Our modeling results indicate that PV+
FSBCs respond differently depending on whether their supralinear or sublinear dendritic
branches are targeted—regardless of the total synaptic input. This raises the intriguing
possibility that PV+ FSBCs might exploit their bimodal nonlinear dendrites to interpret inputs
from different sources across distinct memory stages (encoding/learning and consolidation)
and behavioral states. Could supralinear branches preferentially process inputs from one
brain region while sublinear branches receive signals from another, dynamically tuning
FSBC responses during learning versus consolidation? Could the activation of inhibitory
sublinear branches facilitate a shift in the network into a "learning mode" by promoting
disinhibition? Similarly, could inhibitory supralinear activation function as a mechanism to
transition the network into a "consolidation mode" for processing and stabilizing learned
information? To address these questions, future experimental studies should employ
advanced, *in vivo* approaches to identify the origins of synaptic inputs targeting supralinear

and sublinear dendrites during distinct memory stages. Computational models, like the one
presented here, can then be refined to incorporate this knowledge, allowing for predictions of
how connectivity patterns and input sources influence dendritic computations in PV+ FSBCs.
Ultimately, unraveling these mechanisms will provide a deeper understanding of how
complex dendritic architectures contribute to hippocampal network function across different
memory and behavioral states.

Author contributions: AT and DS conceived the study. AT designed and ran the
simulations, analyzed the data, and prepared the figures. MEL and DS provided senior
conceptual input on the study and methodology. AT wrote the manuscript. DS funded and
supervised the project. All authors edited and approved the final version of the manuscript.

Acknowledgements: A.T. was supported by the DFG with the SFB1315-2 TP A01 Brenda
Milner Award and the Einstein Center for Neurosciences Berlin Fellowship. M.E.L. was
supported by the European Research Council (101055340, ERC AdG, Cortical Coupling).
D.S. was supported by the Einstein Foundation Berlin, the European Research Council
(ERC) under the Europeans Union's Horizon 2020 research and innovation program
(BrainPlay grant, agreement no. 810580), the German Research Foundation (Deutsche
Forschungsgemeinschaft [DFG], SFB-958 – project 184695641, project 431572356, FOR
3004 – project 415914819, SFB 1315 – project 327654276 and under Germany's Excellence
Strategy – Exc-2049-390688087 NeuroCure), and the Federal Ministry of Education and
Research (BMBF, SmartAge – project 01GQ1420B). We thank Michalis Pagkalos, Nikolaus
Maier, John Tukker, and Jeremie Sibille for helpful discussions, comments and proofreading,
Yann Zerlaut, Nelson Rebola, and Yiota Poirazi, for comments on an earlier version of the
work and Daniel Parthier for helpful discussions on the analysis.

Code and Data availability

All codes/scripts and datasets required to reproduce the results and figures, as well as all
statistical analyses, are accessible in the ModelDB database.
https://modeldb.science/2018008 *Access code:* 280621Interneurons! They are also
available on Github: https://github.com/AlexandraTzilivaki/Tzilivakietal2024

Competing interests

The authors declare no competing interests.

References:

Allen, K., and Monyer, H. (2015). Interneuron control of hippocampal oscillations. Curr. Opin.
Neurobiol. *31*, 81–87.

Ascoli, G.A., Alonso-Nanclares, L., Anderson, S.A., Barrionuevo, G., Benavides-Piccione,
R., Burkhalter, A., Buzsáki, G., Cauli, B., DeFelipe, J., Fairén, A., et al. (2008). Petilla
terminology: Nomenclature of features of GABAergic interneurons of the cerebral cortex.
Nat. Rev. Neurosci. *9*, 557–568.

Belluscio, M.A., Mizuseki, K., Schmidt, R., Kempter, R., and Buzsáki, G. (2012). Cross-
Frequency Phase–Phase Coupling between Theta and Gamma Oscillations in the

Hippocampus. J. Neurosci. *32*, 423–435.

Buzsáki, G. (1989). Two-stage model of memory trace formation: A role for "noisy" brain
states. Neuroscience *31*, 551–570.

Buzsáki, G., and Draguhn, A. (2004). Neuronal Oscillations in Cortical Networks. Science
(80-.). *304*, 1926–1929.

Buzsáki, G., and Wang, X.-J. (2012). Mechanisms of Gamma Oscillations. Annu. Rev.
Neurosci. *35*, 203–225.

Canolty, R.T., Edwards, E., Dalal, S.S., Soltani, M., Nagarajan, S.S., Kirsch, H.E., Berger,
M.S., Barbaro, N.M., and Knight, R.T. (2006). High Gamma Power Is Phase-Locked to Theta
Oscillations in Human Neocortex. Science (80-.). *313*, 1626–1628.

Cardin, J.A., Carlén, M., Meletis, K., Knoblich, U., Zhang, F., Deisseroth, K., Tsai, L.-H., and
Moore, C.I. (2009). Driving fast-spiking cells induces gamma rhythm and controls sensory
responses. Nature *459*, 663–667.

Chamberland, S., and Topolnik, L. (2012). Inhibitory control of hippocampal inhibitory
neurons. Front. Neurosci. *6*, 1–13.

Chiovini, B., Turi, G.F., Katona, G., Kaszás, A., Pálfi, D., Maák, P., Szalay, G., Szabó, M.F.,
Szabó, G., Szadai, Z., et al. (2014). Dendritic Spikes Induce Ripples in Parvalbumin
Interneurons during Hippocampal Sharp Waves. Neuron *82*, 908–924.

Cornford, J.H., Mercier, M.S., Leite, M., Magloire, V., Häusser, M., and Kullmann, D.M.
(2019). Dendritic NMDA receptors in parvalbumin neurons enable strong and stable
neuronal assemblies. Elife *8*.

Fernández-Ruiz, A., Oliva, A., Soula, M., Rocha-Almeida, F., Nagy, G.A., Martin-Vazquez,
G., and Buzsáki, G. (2021). Gamma rhythm communication between entorhinal cortex and
dentate gyrus neuronal assemblies. Science (80-.). *372*.

Griffiths, B.J., and Jensen, O. (2023). Gamma oscillations and episodic memory. Trends
Neurosci. *46*, 832–846.

Hadler, M.D., Tzilivaki, A., Schmitz, D., Alle, H., and Geiger, J.R.P. (2024). Gamma
oscillation plasticity is mediated via parvalbumin interneurons. Sci. Adv. *10*.

Hu, H., and Vervaeke, K. (2018a). Synaptic Integration in Cortical Inhibitory Neuron
Dendrites. Neuroscience *368*, 115–131.

Hu, H., and Vervaeke, K. (2018b). Synaptic Integration in Cortical Inhibitory Neuron
Dendrites. Neuroscience *368*, 115–131.

Hu, H., Gan, J., and Jonas, P. (2014). Interneurons. Fast-spiking, parvalbumin[+] GABAergic
interneurons: from cellular design to microcircuit function. Science *345*, 1255263.

Jackson, J., and Skinner, F.K. (2018). Hippocampus, Theta, Gamma, and Cross-Frequency
Coupling. In Encyclopedia of Computational Neuroscience, (New York, NY: Springer New
York), pp. 1–11.

Jackson, J., Goutagny, R., and Williams, S. (2011). Fast and Slow Gamma Rhythms Are
Intrinsically and Independently Generated in the Subiculum. J. Neurosci. *31*, 12104–12117.

Judák, L., Chiovini, B., Juhász, G., Pálfi, D., Mezriczky, Z., Szadai, Z., Katona, G., Szmola,
B., Ócsai, K., Martinecz, B., et al. (2022). Sharp-wave ripple doublets induce complex
dendritic spikes in parvalbumin interneurons in vivo. Nat. Commun. *13*, 6715.

Kastellakis, G., Cai, D.J., Mednick, S.C., Silva, A.J., and Poirazi, P. (2015). Synaptic

clustering within dendrites: An emerging theory of memory formation. Prog. Neurobiol. *126*,
19–35.

Katona, G., Kaszás, A., Turi, G.F., Hájos, N., Tamás, G., Vizi, E.S., and Rózsa, B. (2011).
Roller Coaster Scanning reveals spontaneous triggering of dendritic spikes in CA1
interneurons. Proc. Natl. Acad. Sci. U. S. A. *108*, 2148–2153.

Klausberger, T., and Somogyi, P. (2008). Neuronal Diversity and Temporal Dynamics: The
Unity of Hippocampal Circuit Operations. Science (80-.). *321*.

Kriener, B., Hu, H., and Vervaeke, K. (2022). Parvalbumin interneuron dendrites enhance
gamma oscillations. Cell Rep. *39*, 110948.

Lisman, J.E., and Jensen, O. (2013). The Theta-Gamma Neural Code. Neuron *77*, 1002–
1016.

Lucas, E.K., and Clem, R.L. (2018). GABAergic interneurons: The orchestra or the
conductor in fear learning and memory? Brain Res. Bull. *141*, 13–19.

Memmesheimer, R.-M. (2010). Quantitative prediction of intermittent high-frequency
oscillations in neural networks with supralinear dendritic interactions. Proc. Natl. Acad. Sci.
107, 11092–11097.

Morabito, Z. et al. (2024). A dendritic substrate for temporal diversity of cortical inhibition.
BioRxiv.

Park, K., Lee, J., Jang, H.J., Richards, B.A., Kohl, M.M., and Kwag, J. (2020). Optogenetic
activation of parvalbumin and somatostatin interneurons selectively restores theta-nested
gamma oscillations and oscillation-induced spike timing-dependent long-term potentiation
impaired by amyloid β oligomers. BMC Biol. *18*, 7.

Pelkey, K.A., Chittajallu, R., Craig, M.T., Tricoire, L., Wester, J.C., and McBain, C.J. (2017a).
Hippocampal gabaergic inhibitory interneurons. Physiol. Rev. *97*, 1619–1747.

Pelkey, K.A., Chittajallu, R., Craig, M.T., Tricoire, L., Wester, J.C., and McBain, C.J. (2017b).
Hippocampal gabaergic inhibitory interneurons. Physiol. Rev. *97*, 1619–1747.

Sanchez-Aguilera, A., Wheeler, D.W., Jurado-Parras, T., Valero, M., Nokia, M.S., Cid, E.,
Fernandez-Lamo, I., Sutton, N., García-Rincón, D., de la Prida, L.M., et al. (2021). An
update to Hippocampome.org by integrating single-cell phenotypes with circuit function in
vivo. {PLoS} Biol. *19*, e3001213.

Scheffer-Teixeira, R., Belchior, H., Caixeta, F. V., Souza, B.C., Ribeiro, S., and Tort, A.B.L.
(2012). Theta Phase Modulates Multiple Layer-Specific Oscillations in the CA1 Region.
Cereb. Cortex *22*, 2404–2414.

Scheffzük, C., Kukushka, V.I., Vyssotski, A.L., Draguhn, A., Tort, A.B.L., and Brankačk, J.
(2011). Selective Coupling between Theta Phase and Neocortical Fast Gamma Oscillations
during REM-Sleep in Mice. PLoS One *6*, e28489.

Schomburg, E.W., Fernández-Ruiz, A., Mizuseki, K., Berényi, A., Anastassiou, C.A., Koch,
C., and Buzsáki, G. (2014). Theta Phase Segregation of Input-Specific Gamma Patterns in
Entorhinal-Hippocampal Networks. Neuron *84*, 470–485.

Topolnik, L., and Tamboli, S. (2022). The role of inhibitory circuits in hippocampal memory
processing. Nat. Rev. Neurosci. *23*, 476–492.

Tort, A.B.L., Kramer, M.A., Thorn, C., Gibson, D.J., Kubota, Y., Graybiel, A.M., and Kopell,
N.J. (2008). Dynamic cross-frequency couplings of local field potential oscillations in rat
striatum and hippocampus during performance of a T-maze task. Proc. Natl. Acad. Sci. *105*,
20517–20522.

Tort, A.B.L., Komorowski, R., Eichenbaum, H., and Kopell, N. (2010). Measuring Phase-
Amplitude Coupling Between Neuronal Oscillations of Different Frequencies. J.
Neurophysiol. *104*, 1195–1210.

Tort, A.B.L., Scheffer-Teixeira, R., Souza, B.C., Draguhn, A., and Brankačk, J. (2013).
Theta-associated high-frequency oscillations (110–160Hz) in the hippocampus and
neocortex. Prog. Neurobiol. *100*, 1–14.

Tran-Van-Minh, A., Cazé, R.D., Abrahamsson, T., Cathala, L., Gutkin, B.S., and DiGregorio,
D. a (2015). Contribution of sublinear and supralinear dendritic integration to neuronal
computations. Front. Cell. Neurosci. *9*, 67.

Tzilivaki, A., Kastellakis, G., and Poirazi, P. (2019). Challenging the point neuron dogma: FS
basket cells as 2-stage nonlinear integrators. Nat. Commun. *10*, 3664.

Tzilivaki, A., Kastellakis, G., Schmitz, D., and Poirazi, P. (2022). GABAergic Interneurons
with Nonlinear Dendrites: From Neuronal Computations to Memory Engrams. Neuroscience
489, 34–43.

Tzilivaki, A., Tukker, J.J., Maier, N., Poirazi, P., Sammons, R.P., and Schmitz, D. (2023).
Hippocampal GABAergic interneurons and memory. Neuron *111*, 3154–3175.

Wulff, P., Ponomarenko, A.A., Bartos, M., Korotkova, T.M., Fuchs, E.C., Bähner, F., Both,
M., Tort, A.B.L., Kopell, N.J., Wisden, W., et al. (2009). Hippocampal theta rhythm and its
coupling with gamma oscillations require fast inhibition onto parvalbumin-positive
interneurons. Proc. Natl. Acad. Sci. *106*, 3561–3566.

Zhang, X., Zhong, W., Brankačk, J., Weyer, S.W., Müller, U.C., Tort, A.B.L., and Draguhn, A.
(2016). Impaired theta-gamma coupling in APP-deficient mice. Sci. Rep. *6*, 21948.

Zhong, W., Ciatipis, M., Wolfenstetter, T., Jessberger, J., Müller, C., Ponsel, S., Yanovsky,
Y., Brankačk, J., Tort, A.B.L., and Draguhn, A. (2017). Selective entrainment of gamma
subbands by different slow network oscillations. Proc. Natl. Acad. Sci. *114*, 4519–4524.

a *Microcircuit model of hippocampal oscillations*

PCs

FSBCs

LFP

b *FSBC supralinear dendrites*

Actual EPSP amplitude (mV)

Expected EPSP amplitude (mV)

40 mV / 20 ms

c *FSBC sublinear dendrites*

Actual EPSP amplitude (mV)

Expected EPSP amplitude (mV)

40 mV / 20 ms

d "Bimodal activation"
Synapses to supralinear and sublinear FSBC dendrites

Fast LFP band

Power (a.u.)

Slow LFP band

Frequency (Hz)

e "Supralinear Activation"
Synapses to supralinear FSBC dendrites

Fast LFP band

Power (a.u.)

Slow LFP band

Frequency (Hz)

f "Sublinear Activation"
Synapses to sublinear FSBC dendrites

Fast LFP band

Power (a.u.)

Slow LFP band

Frequency (Hz)

a "Bimodal activation"
Synapses to supralinear and sublinear FSBC dendrites

50 μV, 100 msec

100 μV, 40 msec

b "Supralinear Activation"
Synapses to supralinear FSBC dendrites

50 μV, 100 msec

100 μV, 40 msec

c "Sublinear Activation"
Synapses to sublinear FSBC dendrites

50 μV, 100 msec

100 μV, 40 msec

d

Peak Fast-Frequency (Hz)

0.9949
3.25083-08   5.7119e-08

Bimodal Activation   Supralinear Activation   Sublinear Activation

e

Peak Fast-Frequency Power (mV/Hz) ×10⁻⁴

0.0012
3.8935 e-05   7.0103e-15

Bimodal Activation   Supralinear Activation   Sublinear Activation

f

Peak Slow-Frequency Power (mV/Hz) ×10⁻⁴

0.4976
0.0027   2.7705e-05

Bimodal Activation   Supralinear Activation   Sublinear Activation