The oocyte zinc transporter *Slc39a10/Zip10* is a regulator of zinc sparks

during fertilization in mice.

Atsuko Kageyama[1], Narumi Ogonuki[2], Takuya Wakai[3], Takafumi Namiki[1,4], Yui

Kawata[1], Manabu Ozawa[5], Yasuhiro Yamada[6], Toshiyuki Fukada[7], Atsuo Ogura[2],

Rafael A. Fissore[8], Naomi Kashiwazaki[1,9], Junya Ito[1,9,10*]

[1] Laboratory of Animal Reproduction, School of Veterinary Medicine, Azabu University,

Sagamihara, 252-5201, Japan.

[2] Bioresource Engineering Division, Bioresource Research Center, RIKEN, Tsukuba, 305-0074,

Japan.

[3] Department of Animal Science, Graduate School of Environmental and Life Science,

Okayama University, Okayama, 700-8530, Japan.

[4] Present address: Department of Life Science Frontiers, Center for iPS Cell Research and

Application (CiRA), Kyoto University, Kyoto, 606-8507, Japan.

[5] Laboratory of Reproductive Systems Biology, Center for Experimental Medicine and Systems

Biology, The Institute of Medical Science, The University of Tokyo, Tokyo 108-8639, Japan.

[6] Department of Molecular Pathology, Graduate School of Medicine, The University of Tokyo,

Tokyo 113-0033, Japan.

[7] Laboratory of Molecular and Cellular Physiology, Faculty of Pharmaceutical Sciences,

Tokushima Bunri University, Tokushima 770-8514, Japan.

[8] Department of Veterinary and Animal Sciences, University of Massachusetts, Amherst

01003-9292, the United States.

[9] Graduate School of Veterinary Science, Azabu University, Sagamihara, 252-5201, Japan.

[10] Center for Human and Animal Symbiosis Science, Azabu University, Sagamihara, 252-5201,

Japan.

*Correspondence:

Junya Ito, Laboratory of Animal Reproduction, Graduate School of Veterinary Sciences, Azabu

University, Sagamihara, 252-5201, Japan, itoj@azabu-u.ac.jp

29
30
31

32 **Abstract**

33 In all vertebrates studied to date, a rise(s) in intracellular calcium is indispensable

34 for successful fertilization and further embryonic development. Recent studies

35 demonstrated that zinc is ejected to the extracellular milieu, the 'zinc spark', and

36 follows the first few calcium rises of fertilization. However, the role of the zinc

sparks in fertilization and development, and the supporting influx mechanism(s) are unknown. In this study in mouse oocytes, we investigated the role of zinc transporters *Zip6/Slc39a6* and *Zip10/Slc39a10*. Zip10 mRNA or ZIP10 was expressed throughout folliculogenesis in oocyte or the plasma membrane, respectively. ZIP6 was also expressed in nuclear localization in oocytes and granulosa cells throughout folliculogenesis. The number of ovulated oocytes was examined in oocyte-specific *Zip6* (*Zip6$^{d/d}$*: *Zip6$^{flox/flox}$ Gdf9$^{Cre/+}$*) and *Zip10* (*Zip10$^{d/d}$*: *Zip10$^{flox/flox}$ Gdf9$^{Cre/+}$*) knockout mice, and no change was observed for either strain. *Zip10$^{d/d}$* oocytes matured and formed normal metaphase II spindles, but had lower labile zinc levels as suggested by the zinc indicator, Fluozin-3 intensity. The levels of zinc fluorescence intensity in the *Zip6$^{d/d}$* group were not different from the *Zip6$^{f/f}$*. Fertilization-induced calcium oscillations were present in both *Zip6$^{d/d}$* and *Zip10$^{d/d}$* oocytes, but zinc sparks were observed in *Zip6$^{d/d}$* but not in *Zip10$^{d/d}$* oocytes. Despite other events of egg activation proceeding normally in *Zip10$^{d/d}$* oocytes, embryo development into blastocysts was compromised. We show here for the first time that the zinc transporter ZIP10 contributes to zinc homeostasis in oocytes and embryos, highlighting the pivotal role of this divalent cation in early development.

Classification

Major Classification: Biological Sciences

Minor Classification: Developmental Biology

Keyword: mammals, fertilization, zinc spark, embryo development, zinc signaling

Significance statement

Recent studies reported that the ejection of zinc into the extracellular milieu also called the 'zinc spark', occurs during mammalian fertilization including humans and all vertebrates. However, the role and molecules involved in producing these sparks remain unclear. We demonstrated for the first time that a zinc transporter, *Slc39a10/Zip10,* is required for the zinc sparks of fertilization in mice. In addition,

ZIP10-dependent zinc signaling is not involved in the zona reaction or the block

of polyspermy but it is in embryonic development. Taken together, we show here

a zinc transporter-dependent signaling event has pivotal roles in fertilization and

embryonic development in mice. Our findings will contribute to improving

embryonic development in all mammals.

Introduction

In mammalian fertilization, the sperm factor is released into the ooplasm

to activate the oocyte (1). It is now believed that phospholipase C zeta (PLCζ) (1,

2) is the sperm factor and its induces in the ooplasm repetitive increases in the

intracellular Ca^{2+}, termed 'calcium oscillations.' These oscillations play an

essential role in triggering the oocyte activation events, such as cortical granule

exocytosis, the block of polyspermy, meiotic resumption and exit from MII arrest

(3-7). A calcium increase is highly conserved in many species as the trigger of

egg activation and is vital for successful fertilization.

In recent years, zinc ions as well as calcium ions, are thought to play

important roles during fertilization (8-10). Zinc ion is an essential trace element

and the second most abundant transition ion in the human body after iron (11).

Zinc homeostasis is essential for optimal metabolic function in the reproductive

process in mammals (12), therefore zinc deficiency causes abnormalities such

as fetal teratogenicity, long gestation periods, problematic labor, low birth weight

and weak offspring (13, 14). Zinc also has important role in the male and female

germ cells (9, 10). Especially in the mouse female germ cells, zinc-deficient

condition caused profound defects in oocyte maturation, cumulus cell expansion

and ovulation cycle (15). Several studies have been conducted in mice to

understand how zinc regulates oocyte maturation. Zinc was reported to be a

critical regulator of meiosis throughout oocyte maturation, including maintaining

release from the first and second meiotic arrest (16). In addition, zinc levels of

oocytes were found to increase by 50% as the oocyte progressed through

meiosis (17, 18).

In 2011, Kim et al. reported the importance of zinc during mammalian

fertilization (19). Zinc is predominantly stored in vesicles that are symmetrically

arranged along the oocyte cortex at the GV stage, and zinc-stored vesicles
(cortical granules) are located away from the spindle and form a hemispherical
pattern at the MII stage (19-24). After the penetration of sperm, it has been
induced a 'zinc spark' that releases billions of zinc ions from the cortical granules,
and the total zinc content of the oocytes decreases by 10-20% in at the end of
fertilization in mice (19, 25). In addition, these events immediately follow a series
of calcium oscillations (19). Zinc spark has also been observed in the cattle, non-
human primate, human and Xenopus laevis oocytes (19, 22, 26, 27), suggesting
that this phenomenon is a highly conserved event in vertebrate. More surprisingly,
full-term development of mouse embryos has been reported by chelation of Zn^{2+}
ions without Ca^{2+} release, suggesting the depletion of zinc ions in oocytes may
be sufficient for oocyte activation. Zinc has also been reported to impact the
sperm during fertilization in mice. Zinc accumulation in the zona pellucida
increases fibril binding along the glycoprotein matrix and decreases the number
of sperm that can reach the fertilized oocytes (24, 28, 29). After the zinc sparks,
the released zinc affects the forward motility of sperm to prevent their passage
through the zona matrix (29). Zinc spark is associated with the release of
ovastacin (*Astl*, the official gene name), which is required for the zona reaction,
and functions as a polyspermy block mechanism that is initiated only a few
minutes after fertilization (29).

The transport of zinc ions into and out of cells is regulated by zinc
transporters. In mammals, 14 zinc ion importers, called as ZIP (ZIP1-ZIP14),
have been identified (30-32). Previous study showed that mammalian oocytes
regulate zinc uptake through two maternally derived and cortically distributed zinc
transporters, ZIP6 and ZIP10 (21). Further, they reported that targeted disruption
using *Zip6-* and *Zip10-* specific morpholino injection or antibody incubation
induced alteration of the intracellular labile zinc content, spontaneous resumption
of meiosis from the PI arrest and premature arrest at a telophase I-like state (21).
It is clear from these reports that ZIP6 and ZIP10 are involved in zinc transport in
oocytes, but the mechanisms are not clear. ZIP6 and ZIP10 form and function in
a complex (33), regulate glycogen synthase kinase (GSK-3) and other enzymes
in breast and kidney cancer cells, and regulate epithelial-mesenchymal transition

(EMT) (33, 34). ZIP10 is also involved in many life phenomena such as hair
follicles, skin epidermis and atopic dermatitis (35, 36). In addition, ZIP10 is also
an essential gene for embryonic hematopoiesis (37).

However, while functional analysis of the zinc transporters in various cells and
tissues has been conducted, the function of these zinc transporters in oocytes
has not been elucidated.

In this study, we generated oocyte specific *Zip6* and *Zip10* conditional
knockout mice and examined the function of ZIP10 in the oocytes and the role of
zinc signaling during fertilization and embryonic development. This study
provides clues that elucidate its role in fertilization, which is still largely unknown.
In addition, this is the first report confirming the function of the zinc transporter in
oocytes, which will contribute to future research.

Results

Zinc transporters, ZIP6, and ZIP10, are expressed in mouse oocytes
through follicular development

First, using *in situ* hybridization and immunofluorescent staining, we
examined the expression of ZIP10 during follicular development. As shown in Fig.
1, *Zip10* mRNA was expressed from the primordial oocyte. It continued to be
expressed in oocytes of primary, secondary, and antral follicles. ZIP10 protein
was also expressed in the plasma membrane of primordial oocytes (Fig.1; Arrow).
We also confirmed the expression of ZIP10 protein at the plasma membrane of
oocytes of primary, secondary, and antral follicles. Although ZIP6 was also
expressed in oocytes throughout folliculogenesis (Fig.1; Arrow), it displayed
nuclear localization in oocytes and granulosa cells of primary, secondary, and
antral follicles.

Phenotype of oocyte-specific *Zip6$^{d/d}$* and *Zip10$^{d/d}$* female mice.

To elucidate the roles of *Zip6* and *Zip10* in the mouse oocytes, oocyte-
specific *Zip6* (*Zip6$^{d/d}$*) and *Zip10* (*Zip10$^{d/d}$*) knockout mice were generated (Fig.
S1*A-C*). *Zip6$^{f/f}$* and *Zip10$^{f/f}$* mice were used as controls, respectively. We
examined the number of ovulated oocytes or defects in oocyte maturation. After

superovulation and ovulation, the number of ovulated oocytes collected from the

oviduct in $Zip10^{d/d}$ mice was 22.1 ± 2.5 oocytes/mouse (Fig. 2*A*), which was

equivalent to that in $Zip10^{f/f}$ mice (18.7 ± 2.9 oocytes/mouse, p > 0.05). The

numbers of ovulated oocytes in $Zip6^{d/d}$ and $Zip6^{f/f}$ mice were also equivalent (28.7

± 3.6 and 32.8 ± 3.3 oocytes/mice, respectively, p > 0.05) (Fig. S2*A*). As for

oocyte maturation, the rate of oocytes with a first polar body at 10h, 12h, and 14h

in the $Zip10^{d/d}$ group was not different than that of $Zip10^{f/f}$ group ($Zip10^{d/d}$: 58.7%,

61.9%, and 79.4%, $Zip10^{f/f}$: 50.8%, 54.1% and 70.5%, respectively, p > 0.05, Fig.

2*B*). In addition, $Zip10^{d/d}$ oocytes display intact metaphase spindles just as in the

$Zip10^{f/f}$ groups (Fig. 2*C*).

Next, we examined the levels of labile zinc in germinal vesicle (GV), MII,

and two pronuclei (2PN) zygotes by comparing the fluorescence intensity

following loading with the dye, FluoZin-3AM (Fig. 2*D*). The FluoZin-3AM

fluorescence intensity in GV and MII oocytes of the $Zip10^{d/d}$ group was lower than

the $Zip10^{f/f}$ group (p < 0.05). After fertilization, the fluorescence intensity in $Zip10^{f/f}$

zygotes decreased dramatically, and also decreased for the $Zip10^{d/d}$ group,

although it remained higher than for $Zip10^{f/f}$ oocytes. When compared within

group, the fluorescence intensity in $Zip10^{f/f}$ between GV and MII oocytes was

significantly different (GV vs. MII and MII vs 2PN; p < 0.05, respectively), but the

stages of $Zip10^{d/d}$ oocytes were not different despite clear trends (p > 0.05). The

levels of zinc fluorescence intensity in the $Zip6^{d/d}$ group were not different from

the controls $Zip6^{f/f}$, which decreased markedly at fertilization (Fig. S2*B*; p > 0.05).

The zinc sparks of $Zip10^{d/d}$ oocytes were suppressed after fertilization or

artificial activation.

To determine if the absence of ZIP6 and ZIP10 influenced the detection of

Zn^{2+} sparks associated with fertilization, we monitored extracellular zinc and

intracellular calcium during fertilization of $Zip6^{d/d}$ and $Zip10^{d/d}$ oocytes (Fig. S3

and 3*A*). $Zip10^{f/f}$ oocytes displayed the expected calcium oscillations following

fertilization, and a zinc spark followed the first calcium rise (Fig. 3*A* upper side,

movies 1 and 2). In contrast, $Zip10^{d/d}$ oocytes did not release zinc ions

immediately after the first calcium spike, despite mounting normal calcium

oscillations (Fig. 3*A* lower side, movies 3 and 4). In *Zip6$^{d/d}$* oocytes, a zinc spark

occurred immediately after the first intracellular calcium rise at fertilization, just as

in control *Zip6$^{f/f}$* oocytes (Fig. S3). The extracellular zinc sparks were examined

following artificial oocyte activation of mouse oocytes with ionomycin. The zinc

sparks occurred immediately after the intracellular calcium rise in *Zip10$^{f/f}$* oocytes

(Fig. 4*B* upper side, movies 5 and 6), however, *Zip10$^{d/d}$* oocytes did not release

zinc ions after the calcium spike (Fig. 3*B* lower side, movies 7 and 8).

Zip10$^{d/d}$ mouse oocytes can be fertilized but not developed to the blastocyst.

Zip6$^{d/d}$ and *Zip10$^{d/d}$* oocytes then were used for IVF, and we examined the

rates of fertilization, polyspermy, and embryo development. Successful

fertilization was confirmed at 6 h after IVF by the presence of pronuclei (Fig. 4*A*

and *B*). *Zip10$^{d/d}$* oocytes were fertilized at rates like those observed for control

oocytes (Fig. 4*A*). Consistent with this, the zona reaction triggered by fertilization

and examined using a ZP2 antibody was similar in *Zip10$^{d/d}$* and *Zip10$^{f/f}$* oocytes

(Fig. 4*D*). We also examined the localization of ovastacin, whose expression in

the cortex and loss post-fertilization was comparable between the two groups (Fig.

4*E*). As for the expression of JUNO, it had the same expression than between

null and control oocytes and the temporal dynamics of its disappearance from the

cortex after fertilization was similar for both *Zip10$^{f/f}$* and *Zip10$^{d/d}$* groups (Fig. 4*F*).

We next examined the development to the blastocyst stage, and whereas

Zip10$^{f/f}$ zygotes developed at the expected rates, approximately 75.0%, a smaller

fraction of *Zip10$^{d/d}$* zygotes did and only 32.7% reached this stage ($p < 0.05$) (Fig.

4*B*). As shown in Fig. 4*C*, the total cell numbers of blastocysts were also lower

for *Zip10$^{d/d}$* embryos (51.6 ± 2.1 cells) than for those of *Zip10$^{f/f}$* ones (72.7 ± 2.1

cells, $p < 0.05$). *Zip6$^{d/d}$* oocytes did not display any alterations in the rate of

fertilization or development (Fig. S2*C* and *D*; $p > 0.05$).

Discussion

The zinc sparks associated with fertilization were first discovered in the

mouse. Similar observations followed in other mammalian species and

amphibians. However, the underlying mechanism(s) and biological role of zinc

sparks are not elucidated. In this study, we have examined the role of the zinc

transporters ZIP6 and ZIP10 using conditionally gene deficient mice, and queries

their contribution to the zinc sparks of fertilization.

First, we confirmed the expression of zinc transporter in mouse oocytes.

Our results are consistent with previous studies (21, 38). ZIPs, including ZIP6

and ZIP10, are transmembrane proteins (39-42). In primordial follicles, the

ooplasmic staining of ZIP10 we anticipate correspond to ooplasmic vesicular

sites. ZIP10 expression shifted to the plasma membrane in primary and antral

follicle phase. A similar localization shit of ZIP10 to the oocyte surface was

reported during oocyte maturation, which is much later than reported here by.

Furthermore, in that study, ZIP10 was detected in the nuclear/nucleolar positions

of oocytes of all follicular stages (38), which we did not observe. On the other

hand, ZIP6 was expressed at the nuclear/nucleolar regions and in granulosa cells.

This localization of ZIP6 was consistent with that in a previous study (38).

However, our results failed to notice that ZIP6 shifts to the at plasma membrane

in antral follicle oocytes (21, 38). The results indicate that ZIP6 may be fulfilling a

distinct function within the oocytes compared to ZIP10.

To assess the role of these transporters in mice, we generated oocyte-

specific *Zip6 or Zip10* knockout mice (Fig. S1; *Zip6*$^{d/d}$ and *Zip10*$^{d/d}$, respectively).

Previous reports noted that the disruption of these transporters using specific

morpholinos or incubation in function-blocking antibodies induced the change of

intracellular labile zinc quota into mice oocytes (21). The amount of free zinc ions

in those mouse oocytes was measured using Fluozin-3AM, zinc indicator. The

amount of labile zinc ions in *Zip10*$^{d/d}$ oocytes was significantly lower than in the

Zip10$^{f/f}$ (Fig. *2D*). We failed to observe any effect on zinc levels of in *Zip6*$^{d/d}$

oocytes (Fig. *S2B*). The results indicate that ZIP10 is mostly responsible for the

uptake of zinc ions in mouse oocytes.

Zinc insufficient GV oocytes do not maintain the meiotic arrest at PI. This

is suggested to happen by premature activation of the MOS-MAPK pathway in

the presence of low zinc (16). In addition, low ooplasmic zinc accelerates meiotic

progression that contributes to the extrusion of large polar bodies (18). The early

mitotic inhibitor 2 (EMI2), a zinc-binding APC/C proteasome inhibitor, is also an

essential component of the cytoplasmic factor (CSF) that initiates entry into MII
phase. EMI2 is a zinc binding protein, and when zinc is reduced, the activity of
the APC/C proteasome is stimulated, increasing the degradation of CCNB1,
reduced MPF activity leading to early release of meiosis arrest (8, 18, 43-45).
Previous studies reported that targeted disruption of *Zip10* using morpholino
injections and function-inhibiting antibodies during meiotic maturation perturbed
meiosis progression and resulted in cell cycle arrest at the telophase I-like state
(21). Based on these findings we hypothesized that the reduced state of free zinc
ions in *Zip10$^{d/d}$* mouse oocytes may resemble the state of zinc-deficient oocytes.
We investigated the maturation progression and spindle organization in oocyte
matured in vitro from the GV to MII stage. We failed to observe any differences
(Fig. *2A, B* and *C*). We also did not observed abnormalities in oocytes from *Zip6$^{f/f}$*
mice (Fig. *S2A*; P > 0.05). We speculate that the disparate outcomes observed
in previous studies may be attributed to the presence of trace amounts of zinc
ions in media or in the oocytes.

Surprisingly, the amount of zinc ions in *Zip10$^{d/d}$* 2PN zygote was higher
than in *Zip10$^{f/f}$* and despite some differences in fluorescence intensity between
Zip10$^{d/d}$ GV, MII and 2PN zygote, the labile zinc concentrations were not
significantly different between these groups (Fig. *2D*). Therefore, we monitored
zinc sparks and calcium oscillations in the *Zip10*-KO oocytes. Several studies
reported the important of IP$_3$Rs in mammalian oocytes (46, 47), which is essential
for egg activation because its inhibition precludes Ca^{2+} oscillations (48-50). It has
been reported that a putative Zinc-finger motif in a helical linker (LNK) domain
near the C-terminus of *IP$_3$R1* plays a role in *IP$_3$R1* function (51, 52), Recently,
Akizawa et al., reported that both deficient and excessive zinc ions compromise
IP$_3$R1 sensitivity, diminishing and terminating calcium oscillations (53). Our
results showed that a single zinc spark occurs immediately after first calcium rise
of oscillations in *Zip10$^{f/f}$* oocytes, as reported by Kim et al (19) (Fig. *3A*), and
similar results are obtained in *Zip6$^{d/d}$* and *Zip6$^{f/f}$* mouse oocytes (Fig. *S3*).
However, in *Zip10$^{d/d}$* oocytes, despite the presence calcium elevations, zinc
sparks failed to occur. Similarly, *Zip10$^{d/d}$* oocytes activated with ionomycin
(artificial oocyte activation) did not show a zinc spark (Fig. *3B*). In this study,

calcium oscillations occurred in oocytes with low zinc state. In addition, there was
no difference in the amplitude frequency of calcium ions in oocytes of both groups
within the observation time. Akizawa et al., produced zinc deficiency state with
TPEN, a strong chelator of intracellular labile zinc ions (53). On the other hand,
we did not completely remove intracellular labile zinc ions from mouse oocytes.
We speculate that zinc ions existed in $Zip10^{d/d}$ mouse oocytes induce Ca^{2+}
release without compromising IP_3R1 sensitivity. However, this study
demonstrated that the accumulation of intracellular zinc ions mediated by ZIP10
is essential for inducing zinc sparks in mouse oocytes.

The total cellular zinc ions content required for meiosis in the oocyte
substantially increases (a 50% increase) from the prophase I arrest to the arrest
at metaphase of meiosis II (17, 18). The zinc is stored in undefined cortical
granules, and an average of 10^6 zinc atoms are released from these vesicles at
the time of fertilization (exocytosis) (19-23, 25). There are several possibilities
that might explain why $Zip10$ conditional KO mice have low zinc sparks. The most
logical is that these oocytes experience reduced zinc loading of cortical of zinc
vesicles, leading to the absence of sparks. Our results suggest that in oocytes,
zinc sparks and exocytosis require adequate concentrations of zinc ions in the
cortical secretory vesicles, which is not accomplished when ZIP10 is not present.

Zinc sparks have been reported to contribute to the rapid and permanent
block mechanisms to prevent polyspermy. The former is achieved by zinc in the
extracellular milieu decelerating the forward motility of sperm (29). Secondly, the
release of labile pools of zinc at the same time that the zinc metalloendopeptidase
ovastacin is released with the cortical granules (29, 54, 55), cause ZP2 protein
cleavage inducing a complete, although belated block to polyspermy (24, 28, 29).
However, the physiological significance of the zinc sparks' contributions to
polyspermy have not been thoroughly tested, and $Zip10^{d/d}$ mice, which are almost
devoid of them, offer a great model. Our results showing normal rates of
monospermic fertilization in $Zip10^{d/d}$ oocyte suggest that contributions of Zinc to
polyspermy control is negligible (Fig. *4A*). Our results also show that the
ovastacin undergoes normal release in $Zip10^{f/f}$ oocytes after fertilization (Fig. *4E*
and 4*D*), confirming previous studies of a minor role of the ZP2 to polyspermy.

We observed the disappearance of JUNO after fertilization (56) *Zip10^{d/d},*
suggesting that gamete fusion took place (Fig. *4F*). Together, these results
support the view that zinc spark is not directly involved in the polyspermy rejection
mechanism.

Interestingly, there were no differences in the rates of fertilization
(pronucleus formation), but development to the blastocyst stage was significantly
reduced in *Zip10*-null embryos (Fig. *4B*). These embryos seem to gradually arrest
after the 2-cell stage (data not shown) and lose viability. It is well known that zinc
levels play a role in embryonic development. Zinc insufficiency caused altered
chromatin structure in the nuclei of blastomeres that displayed decreased global
transcription, causing arrest in embryonic development (57). The abnormal zinc
homeostasis, particularly during the one-cell stage, inhibits the activation of the
embryonic genome that occurs around the two-cell stage in mice due to reduce
the translational capacity through the inhibition of ribosomal RNA synthesis by
RNA polymerase I (10, 58). Our results also show that *Zip10^{d/d}* blastocysts had
significantly lower cell numbers compared to the blastocysts from *Zip10^{f/f}* mice
(Fig. *4C*). Our results suggest that Zip10^{d/d} mice oocytes can be fertilizable, but
their developmental potential is decreased. The lack of ZIP10-mediated zinc
influx during the folliculogenesis of *Zip10^{d/d}* may compromise gene expression
during these early stages of oocyte development. *Zip10^{d/d}* mouse oocytes thus
may have compromised developmental potential from the outset. Future studies
should assess the transcriptomic profile of *Zip10^{d/d}* mouse oocytes.

In conclusion, we elucidated that ZIP10 is required for zinc ion uptake into
oocytes and the intracellular zinc ions regulated by ZIP10 was important for zinc
sparks and progression of embryonic development. Currently, the many infertility
patients exist in the world (the survey of WHO). It is reported that approximately
half of the adult females are deficient in zinc in their serum (59, 60). Zinc has
been reported to play essential roles in many physiological functions including
reproductive functions. The findings will help elucidate the role of zinc signaling
in the fields of fertilization/embryogenesis. Furthermore, the development of zinc-
focused assisted reproductive technologies and fertilization/embryogenesis
media could contribute to improve the developmental potential of oocytes and

embryos in other mammalian species.

Materials and Methods

360 All chemicals and reagents were purchased from Sigma-Aldrich (St. Louis, MO,
361 USA) unless otherwise stated.

Animals

364   Animals were housed in the barrier facility at Azabu University. The
deletion of the ZIP10 gene in the whole body is known to result in embryonic
lethality (37). The conditional knockout mice were generated by Cre-loxP system.
The following mouse strains were used: $Zip10^{flox/flox}$ ($Zip10^{f/f}$) mouse <B6;129-
Slc39a10 <tm1.1Tfk>> (RBRC06221) (35, 61, 62) was provided by the RIKEN
BRC through the National BioResource Project of the MEXT/AMED, Japan.
$Zip6^{flox/flox}$ ($Zip6^{f/f}$) mouse <B6;129-Slc39a6> was generated by AdAMS. $Gdf9^{Cre/+}$
mice <Stock Tg (GDF9-iCre) 5092Coo/J, Stock No: 011062> were delivered from
Jackson Laboratory (63). Crlj: C57BL/6J female mice (6-8 weeks old) and Crlj:
C57BL/6J male mice (8-12 weeks old) were purchased from Charles River
Laboratories Japan (Yokohama, Japan). The mice were housed under controlled
lighting conditions (daily light period, 0600-1800 h). The study was approved by
the Animal Experimentation Committees of Azabu University and were performed
in accordance with the committees' guidelines (200318-13 and 230309-1).
Oocyte-specific gene knockout ($Zip10^{f/f}$, $Gdf9^{Cre/+}$; $Zip10^{d/d}$) mice were generated
by crossing $Gdf9^{Cre/+}$ male mice with $Zip10^{f/f}$ female mice (Fig. S1A-E). Similarly,
$Zip6^{d/d}$ mice were generated (Fig. S1A-E).

In situ hybridization (ISH)

383   ISH was performed as previously described with some modifications (64).
Briefly, fixed ovaries were paraffin-embedded and paraffin sections (6 μm) were
mounted on MAS-coated slides (Matsunami Glass Industries, Osaka, Japan)
under RNase-free conditions. Sense or antisense digoxigenin (DIG)-labeled RNA
probes for $Slc39a10$ ($Zip10$) were purchased from Genostaff. The sections were
deparaffinized, rehydrated, and post-fixed in 10% neutral buffered formalin (NBF)

for 30 min at 37°C, followed by the treatment with 0.2% hydrogen chloride and 5

µg/ml proteinase K (FUJIFILM Wako Pure Chemical, Osaka, Japan) for 10 min

at 37°C, respectively. Hybridization was performed with DIG-labeled probes (250

ng/ml) in a humidified chamber at 60°C overnight. The slides were washed after

hybridization, then treated with blocking reagent (Genostaff) for 15 min and

alkaline phosphatase-conjugated anti-DIG antibody (1:2,000; Roche Diagnostics,

Basel, Switzerland) for 1 h at room temperature. The signals were detected by 4-

nitro-blue tetrazolium/5-bromo-4-chloro-3-indolyl phosphate (NBT/BCIP, Roche

Diagnostics) in a humidified container for 12 h at 4°C. The sections were

counterstained with Kernechtrot solution (Muto Pure Chemicals, Tokyo, Japan).

Signals detected by the sense probe were used as a control for background levels.

Immunofluorescent staining (IF) for ovary

IF was performed as previously described with some modifications (64).

Fixed ovaries were paraffin-embedded and the sections (6 µm) were

deparaffinized, hydrated and conducted for antigen retrieval by autoclaving in 10

mM sodium citrate buffer (pH = 6.0) for 5 min. The sections were further incubated

on ice for 30 min. After blocking with Bloking One Histo (06349-64, NACALAI

TESQUE Inc, Kyoto, Japan) for 1 h, the slides were incubated with primary

antibody for Rabbit anti-mouse-Zip10 (1:200; (62)), Rabbit anti-SLC39A6 (1:200,

HPA042377, Sigma), Rabbit anti-Foxl2 (1:300; (65, 66)), Rat anti-ZP2 (1:100, sc-

32752, Santa Crus Biotechnology, Dallas, Texas, USA) in Can Get Signal

immunostain (TOYOBO, Tokyo, Japan) overnight at 4°C. The slides were

incubated with Alexa Fluor 488 donkey anti-rabbit IgG (H+L), Alexa Fluor 594

donkey anti-rabbit IgG (H+L) or Alexa Fluor 488 donkey anti-rat IgG (H+L)

conjugated secondary antibodies (Jackson Immuno Research Laboratories,

West Grove, PA, USA) diluted 1:500 in Can Get Signal immunostain (TOYOBO)

for 1 h, and mounted with ProLong Glass Antifade Mountant with NucBlue Stain

(P36981, Thermo Fisher Scientific, Waltham, MA, USA). Micrographs were

captured by BZ-X700 microscopy (Keyence, Osaka, Japan).

Oocyte preparation

Germinal vesicle (GV) oocytes were collected in the manner described in
our previous study (67) with some modifications. Ovaries were collected from
female mice 48 h after intraperitoneally injection with 5 IU equine chorionic
gonadotropin (eCG) (PMS; Nippon Zenyaku Kogyo, Fukushima, Japan). They
were placed in a 35 mm culture dish containing MEMα (no nucleosides, powder;
Gibco/Thermo Fisher Scientific, Tokyo, Japan) medium with 26 mM $NaHCO_3$, 75
mg/l penicillin, 50 mg/l streptomycin sulfate, 5% (v/v) heat-treated fetal calf serum
(FCS), and 10 ng/ml epidermal growth factor (EGF). Cumulus-oocyte complexes
(COCs) were released from the antral follicles by gentle puncturing with a needle.
To obtain metaphase II (MII) oocytes, the mice were intraperitoneally
injected with 5 IU eCG followed by injection with 5 IU human chorionic
gonadotropin (hCG) (Gonatropin; ASKA Pharmaceutical, Tokyo, Japan) at 48 h
later. COCs were collected from the oviductal ampulla 14-16 h after hCG injection.
These oocytes were used in the following experiments.
435

Count of ovulated oocytes

437   COCs-MII were obtained by superovulation treatment. The cumulus cells
438 were removed from the COCs-MII with hyaluronidase (1 mg/ml) and gentle
439 pipetting. All collected oocytes were counted and calculated as the number of
440 ovulations.
441

In vitro maturation (IVM) of GV oocytes

IVM were conducted in the manner described in our previous study (68)
with some modifications. Ovaries were placed in a 35 mm culture dish containing
MEMα medium with 26 mM $NaHCO_3$, 75 mg/l penicillin, 50 mg/l streptomycin
sulfate, 5% (v/v) FCS, and 10 ng/ml EGF. COCs-GV were released from the
antral follicles by gentle puncturing with a needle. The COCs-GV were washed 3
times and cultured in 500 µl of same medium in 4-well dish at 37°C in an
atmosphere of 5% CO_2 in air for 10, 12 and 14 h. After the culture, cumulus cells
were removed from the COCs with hyaluronidase (1 mg/ml) and gentle pipetting.
The extrusion of 1st polar body was evaluated at each time.
452

IF for oocytes/preimplantation embryos

The α-tubulin and JUNO (PE anti-mouse FR4) were performed with some modification of the methods of Inoue et al (69). The oocytes or embryos were fixed in 4% PFA for 30 min at room temperature, washed in PBS containing 1% polyvinyl alcohol (PBS/PVA). They were permeabilized by treatment of 0.5% Triton X-100 for 15 min and washed 2 times in 1% BSA/PBS/PVA followed by blocked for 20 min in same medium. They were incubated for overnight at 4° with primary antibodies to rabbit anti-α-tubulin (1:200; 11H10, cell Signaling Technology) or rat anti-mouse FR4 (JUNO; 1:250; 12A5 BioLegend), wash 3 times in 1% BSA/PBS/PVA. Primary antibodies were detected using Alexa Fluor 488 donkey anti-rabbit IgG (H+L) (1:250) or Alexa Fluor 488 donkey anti-rat IgG (H+L) (1:250) for 1 h at room temperature. After staining, all samples mounted in VECTASHIELD Mounting Medium with DAPI (H-1200; Vector Laboratories, California, USA), imaged using BZ-X700 microscopy (Keyence, Osaka, Japan).

The ovastacin staining was performed with some modifications of the methods of Burkart et al (54). oocytes or embryos were fixed in 4% PFA for overnight at 4 °C, washed in PBS containing 0.3% polyvinylpyrrolidone (PVP), and then blocked in 0.3% BSA/0.1 M glycine (three times for 10 min) followed by permeabilization in 0.2% Triton X-100 for 15 min (70). Samples were then incubated for overnight at 4°C with Rabbit polyclonal anti-ovastacin (54), 1:200, gifted antibody), washed with 0.3% PVP/0.1% Tween (three times for 10 min), and incubated for 1 h at room temperature with Alexa Fluor 488 anti-rabbit secondary antibody (1:500) followed by staining and mounting with VECTASHIELD Mounting Medium with DAPI. Samples were imaged using TCS SP5 II confocal microscope (Leica Microsystems, Wetzlar, Germany).

In vitro fertilization (IVF)

IVF and sperm collection were conducted with some modification of the method described in our previous study (71). In brief, ovulated COCs-MII were preincubated for 1 h in 80 µl human tubal fluid (HTF) droplets supplemented with 1.25 mM reduced glutathione (GSH). Frozen-thawed sperm suspensions were

suspended in 200 µl preincubation medium (HTF containing 0.4 mM methyl-β-
cyclodextrin) and 0.1 mg/ml PVA, but without bovine serum albumin, and were
incubated at 37 °C under 5% CO_2 in humidified air for 1 h. At the time of
insemination, preincubated spermatozoa were transferred into the droplets with
oocytes at final concentrations of 2.0×10^6 sperm/ml. After 6 h, oocytes were
separated from spermatozoa and cumulus cells using a fine glass pipette and
transferred into 50 µl KSOMaa medium. They were cultured at 37 °C under 5%
CO_2 in humidified air for approximately 24-96 h. After IVF at 96 h, the blastocysts
were fixed in 4% PFA for 30 min at room temperature. Following three times
washed in PBS/PVA for 30 min each. Nuclear DNA was stained and mounted in
VECTASHIELD Mounting Medium with DAPI. Cell numbers were determined by
visually inspecting nuclei stained with DAPI using an inverted fluorescence
microscope.
497
Zinc measurements
FluoZin-3AM staining was conducted using the method described in our
previous study (72). The obtained GV oocytes, cumulus cells were removed from
the COCs with gentle pipetting. The obtained MII oocytes, cumulus cells were
removed from the COCs with hyaluronidase (1 mg/ml) and gentle pipetting.
Oocytes with a polar body were defined as MII. After IVF for 6h, two pronucleus
stage embryos (2PN) were collected. GV, MII and 2PN were loaded in 50 µl
medium that was suitable for each stage containing the amyl ester of the
membrane permeant zinc-specific fluorophore, FluoZin-3AM (2 µM; F24195,
Thermo Fisher Scientific, excitation 494 nm/emission 516 nm) for 1 h in
humidified CO_2 [5% (v/v) in air] at 37 °C followed by washing three times in
medium and then observation with TCS SP5 II confocal microscope. The pixel
intensity per unit area after background subtraction was determined in GV, MII
and 2PN within the circle (white circle) and ImageJ image processing software.
512
Measurement of Ca^{2+} and Zinc spark during fertilization
Zip10^{f/f} and *Zip10^{d/d}* female mice (7-12weeks) were treated for
superovulation with eCG and hCG as same procedure. COCs were collected at

14-16 h after hCG injection and removed cumulus cells with hyaluronidase. The

zona pellucida (ZP) was removed by brief treatment in acidic Tyrode's solution.

The ZP-free MII were cultured in HTF containing Calbryte^{TM} 590 AM (10 µM,

excitation 581 nm/emission 593 nm; AAT Bioquest, CA, USA) for 10-20 min in

humidified CO_2 [5% (v/v) in air] at 37 °C. In case of IVF, ZP-free MII were allowed

to settle in 5 µl calcium-free HTF medium drop on dish to attach oocytes. After

attached, HTF medium that contained BSA and membrane impermeant zinc-

specific fluorophore, FluoZin-3 (20 µM; F24194, Thermo Fisher Scientific,

excitation 494 nm/emission 516 nm) was slowly added to a final volume of 45 µl,

taking care not to disturb oocytes, to start imaging. In case ionomycin treatment,

after calcium labeling, ZP-free MII were allowed to settle in 45 µl calcium-free

HEPES-CZB medium drop on dish, to start imaging. Imaging was performed on

confocal microscope using 488 nm (Zn^{2+}) and 555 nm (Ca^{2+}) excitation (Nikon

solutions, Tokyo, Japan) for max 50 min every 4 seconds. Capacitated frozen-

thawed sperm or 5 µM ionomycin were added to ZP-free MII at 2 min after

imaging start. Imaging analysis was performed by defining regions of interest

(ROIs) and measuring fluorescence intensity over time using NIS-Elements

(Nikon). The intracellular ROIs were drawn as the entire interior area of the cell.

The extracellular ROIs were defined as a ring around the perimeter of cell. The

ring thickness was conserved for all data analyses. The same methos was used

for the *Zip6$^{f/f}$* and *Zip6$^{d/d}$ mice*.

Western blotting

Western blotting was carried out as described (73) with some

modifications. Thirty MII oocytes or 2PN embryos were lysed in Laemmli sample

buffer (Bio-Rad Laboratories, Tokyo, Japan) with 5% 2-mercaptoethanol.

Samples were separated on 8% Bis-Tris gels by SDS-PAGE and transferred to

PVDF membranes (Bio-Rad). The PVDF membranes were blocked in 10% skim

milk (FUJIFILM Wako) in Tris-buffered saline (TBS) with 0.1% Tween-20

(Yoneyama yakuhin kogyo, Osaka, Japan) and probed with primary antibody to

rat anti-ZP2 (1:1,000) or monoclonal mouse anti-β-actin (1:5,000; A5316, Sigma-

Aldrich) for overnight at 4°C. The membranes were incubated with secondary

antibody; HRP-conjugated anti-rat IgG (1:5000; Cell Signaling Technology) or

HRP-conjugated anti-mouse IgG (1:5,000; Cell Signaling Technology) for 1 h at

room temperature. After washing of the membranes, immunoreactive proteins

were visualized using ECL^{TM} Western Blotting Analysis System (Cytiva Global

Life Sciences Technologies, Tokyo, Japan), according to the manufacture's

recommendation. The membranes were exposed by ImageQuant LAS 4000.

Statistical analysis

Values from three or more times were used for evaluation of statistical

significance. Statistical analysis was performed using Statcel 3 software (OMS

Ltd., Saitama, Japan). The fertility, the total numbers of oocytes collected from

each mouse, the percentage of first polar extrusions, the fluorescence intensity

of Fluozin-3 AM and the cell count of blastocysts were evaluated statistically by

Student's t-test analysis. The rate of fertilization and embryo development was

analyzed using chi square tests. Values are shown as means ± S.E.M, and

significant differences were considered at p values < 0.05.

Acknowledgments and funding sources

We thank the members of the Laboratory of Animal Reproduction, School of

Veterinary Medicine, Azabu University for technical help. *Zip10$^{flox/flox}$* (*Zip10$^{f/f}$*)

mouse <B6;129-Slc39a10 <tm1.1Tfk>> (RBRC06221) was provided by the

RIKEN BRC through the National BioResource Project of the MEXT/AMED,

Japan. We also thank Dr. Austin J. Cooney (The University of Texas at Austin,

Austin, United States) for providing GDF9-iCre mice. This research was

supported by Grants-in-Aid for Scientific Research from the Japan Society for the

Promotion of Science (JSPS) (KAKENHI, 21H02384 and 20H05373 to J.I.,

22H04268 and 24H02601 to A.K.), JSPS KAKENHI Grant Number JP 15H04584

(AdAMS) to JI, the Sasakawa Scientific Research Grant from The Japan Science

Society (2019-4044) to A.K. and Mishima Kaiun Memorial Foundation (A-135) to

A.K. This work was supported by JSPS KAKENHI Grant Number JP 22H04922

(AdAMS). This study was also supported by Center for Diversity, Equity &

Inclusion, Azabu University (A.K). This research was partially supported by the
Center for Human and Animal Symbiosis Science, Azabu University and a
research project grant awarded by the Azabu University Research Services
Division to J.I.

References

1. S. Yoon, R. A. Fissore, Release of phospholipase C zeta and [Ca2+]i
oscillation-inducing activity during mammalian fertilization. *Reproduction.* **134**,
695-704 (2007).

2. C. M. Saunders *et al.,* PLC zeta: A sperm-specific trigger of ca(2+)
oscillations in eggs and embryo development. *Development.* **129**, 3533-3544
(2002).

3. Z. Xu, G. S. Kopf, R. M. Schultz, Involvement of inositol 1,4,5-trisphosphate-
mediated Ca2+ release in early and late events of mouse egg activation.
Development. **120**, 1851-1859 (1994).

4. K. T. Jones, Mammalian egg activation: From Ca2+ spiking to cell cycle
progression. *Reproduction.* **130**, 813-823 (2005).

5. T. Ducibella, R. M. Schultz, J. Ozil, Role of calcium signals in early
development. *Semin. Cell. Dev. Biol.* **17**, 324-332 (2006).

6. J. Ito, J. Parrington, R. A. Fissore, PLCzeta and its role as a trigger of
development in vertebrates. *Mol. Reprod. Dev.* **78**, 846-853 (2011).

7. H. Sugita *et al.,* Molecular mechanism of oocyte activation in mammals: Past,
present, and future directions. *Biomolecules.* **14**, 359 (2024).

8. T. Suzuki *et al.,* Full-term mouse development by abolishing Zn2+-dependent
metaphase II arrest without Ca2+ release. *Development.* **137**, 2659-2669
(2010).

9. D. Allouche-Fitoussi, H. Breitbart, The role of zinc in male fertility. *Int. J. Mol.*
Sci. **21**, 7796 (2020).

10. A. Kageyama, J. Terakawa, J. Ito, N. Kashiwazaki, Roles of zinc signaling in
mammalian reproduction. Metallomics Research. **2**, 2022.

11. S. Lubna, R. Ahmad, Clinical and biochemical understanding of zinc
interaction during liver diseases: A paradigm shift. *J. Trace. Elem. Med. Biol.*
77, 127130 (2023).

12. A. M. Pascua *et al.,* Reproductive hormones influence zinc homeostasis in
the bovine cumulus-oocyte complex: Impact on intracellular zinc concentration
and transporters gene expression. *Theriogenology.* **146**, 48-57 (2020).

13. A. E. Favier, The role of zinc in reproduction. hormonal mechanisms. *Biol.*
Trace. Elem. Res. **32**, 363-382 (1992).

14. R. S. Bedwal, A. Bahuguna, Zinc, copper and selenium in reproduction.
Experientia. **50**, 626-640 (1994).

15. X. Tian, F. J. Diaz, Zinc depletion causes multiple defects in ovarian
function during the periovulatory period in mice. *Endocrinology.* **153**, 873-886
(2012).

16. B. Y. Kong *et al.,* Zinc maintains prophase I arrest in mouse oocytes
through regulation of the MOS-MAPK pathway. *Biol. Reprod.* **87**, 1-12 (2012).

17. A. M. Kim, S. Vogt, T. V. O'Halloran, T. K. Woodruff, Zinc availability
regulates exit from meiosis in maturing mammalian oocytes. *Nat. Chem. Biol.* **6**,
674-681 (2010).

18. M. L. Bernhardt, A. M. Kim, T. V. O'Halloran, T. K. Woodruff, Zinc
requirement during meiosis I-meiosis II transition in mouse oocytes is
independent of the MOS-MAPK pathway. *Biol. Reprod.* **84**, 526-536 (2011).

19. A. M. Kim *et al.,* Zinc sparks are triggered by fertilization and facilitate cell
cycle resumption in mammalian eggs. *ACS. Chem. Biol.* **6**, 716-723 (2011).

20. E. L. Que *et al.,* Quantitative mapping of zinc fluxes in the mammalian egg
reveals the origin of fertilization-induced zinc sparks. *Nat. Chem.* **7**, 130-139
(2015).

21. B. Y. Kong *et al.,* Maternally-derived zinc transporters ZIP6 and ZIP10 drive
the mammalian oocyte-to-egg transition. *Mol. Hum. Reprod.* **20**, 1077-1089
(2014).

22. E. L. Que *et al.,* Bovine eggs release zinc in response to parthenogenetic
and sperm-induced egg activation. *Theriogenology.* **127**, 41-48 (2019).

23. Y. Jo *et al.,* Spire localization via zinc finger-containing domain is crucial for
the asymmetric division of mouse oocyte. *Faseb. J.* **33**, 4432-4447 (2019).

24. E. L. Que *et al.,* Zinc sparks induce physiochemical changes in the egg
zona pellucida that prevent polyspermy. *Integr. Biol. (Camb).* **9**, 135-144 (2017).

25. H. C. Lee *et al.,* Zinc exocytosis is sensitive to myosin light chain kinase
inhibition in mouse and human eggs. *Mol. Hum. Reprod.* **26**, 228-239 (2020).

26. F. E. Duncan *et al.,* The zinc spark is an inorganic signature of human egg
activation. *Sci. Rep.* **6**, 24737 (2016).

27. J. F. Seeler *et al.,* Metal ion fluxes controlling amphibian fertilization. *Nat.*
Chem. **13**, 683-691 (2021).

28. S. Aonuma, M. Okabe, M. Kawaguchi, Y. Kishi, Zinc effects on mouse
spermatozoa and in-vitro fertilization. *J. Reprod. Fertil.* **63**, 463-466 (1981).

29. K. Tokuhiro, J. Dean, Glycan-independent gamete recognition triggers egg
zinc sparks and ZP2 cleavage to prevent polyspermy. *Dev. Cell.* **46**, 627-640.e5
(2018).

30. T. Fukada, T. Kambe, Molecular and genetic features of zinc transporters in
physiology and pathogenesis. *Metallomics.* **3**, 662-674 (2011).

31. T. Takagishi, T. Hara, T. Fukada, Recent advances in the role of
SLC39A/ZIP zinc transporters in vivo. *Int. J. Mol. Sci.* **18**, 2708 (2017).

32. T. Kambe, T. Tsuji, A. Hashimoto, N. Itsumura, The physiological,
biochemical, and molecular roles of zinc transporters in zinc homeostasis and
metabolism. *Physiol. Rev.* **95**, 749-784 (2015).

33. K. M. Taylor *et al.,* Zinc transporter ZIP10 forms a heteromer with ZIP6
which regulates embryonic development and cell migration. *Biochem. J.* **473**,
2531-2544 (2016).

34. D. Pal, U. Sharma, S. K. Singh, R. Prasad, Association between ZIP10
gene expression and tumor aggressiveness in renal cell carcinoma. *Gene.* **552**,
195-198 (2014).

35. B. Bin *et al.,* Requirement of zinc transporter ZIP10 for epidermal
development: Implication of the ZIP10-p63 axis in epithelial homeostasis. *Proc.*
Natl. Acad. Sci. U. S. A. **114**, 12243-12248 (2017).

36. K. Nakajima *et al.,* Possible involvement of zinc transporter ZIP10 in atopic
dermatitis. *J. Dermatol.* **47**, e51-e53 (2020).

37. X. He *et al.,* The zinc transporter SLC39A10 plays an essential role in
embryonic hematopoiesis. *Adv. Sci. (Weinh).* **10**, e2205345 (2023).

38. Y. Chen *et al.,* Zinc dynamics regulate early ovarian follicle development. *J.*
Biol. Chem. **299**, 102731 (2023).

39. D. J. Eide, The SLC39 family of metal ion transporters. *Pflugers. Arch.* **447**,
796-800 (2004).

40. T. Kambe, Y. Yamaguchi-Iwai, R. Sasaki, M. Nagao, Overview of
mammalian zinc transporters. *Cell. Mol. Life. Sci.* **61**, 49-68 (2004).

41. M. Lu, D. Fu, Structure of the zinc transporter YiiP. *Science.* **317**, 1746-
1748 (2007).

42. G. Schmitt-Ulms *et al.,* Evolutionary descent of prion genes from the ZIP
family of metal ion transporters. *PLoS One.* **4**, e7208 (2009).

43. M. L. Bernhardt *et al.,* A zinc-dependent mechanism regulates meiotic
progression in mammalian oocytes. *Biol. Reprod.* **86**, 114 (2012).

44. S. Shoji *et al.,* The zinc-binding region (ZBR) fragment of Emi2 can inhibit
APC/C by targeting its association with the coactivator Cdc20 and UBE2C-
mediated ubiquitylation. *FEBS. Open. Bio.* **4**, 689-703 (2014).

45. M. Ohe *et al.,* Emi2 inhibition of the anaphase-promoting
complex/cyclosome absolutely requires Emi2 binding via the C-terminal RL tail.
Mol. Biol. Cell. **21**, 905-913 (2010).

46. R. A. Fissore *et al.,* Differential distribution of inositol trisphosphate receptor
isoforms in mouse oocytes. *Biol. Reprod.* **60**, 49-57 (1999).

47. J. Parrington *et al.,* Expression of inositol 1,4,5-trisphosphate receptors in
mouse oocytes and early embryos: The type I isoform is upregulated in oocytes
and downregulated after fertilization. *Dev. Biol.* **203**, 451-461 (1998).

48. S. Miyazaki, M. Ito, Calcium signals for egg activation in mammals. *J.*
Pharmacol. Sci. **100**, 545-552 (2006).

49. S. Miyazaki *et al.,* Block of Ca2+ wave and Ca2+ oscillation by antibody to
the inositol 1,4,5-trisphosphate receptor in fertilized hamster eggs. *Science.*
257, 251-255 (1992).

50. Z. Xu, C. J. Williams, G. S. Kopf, R. M. Schultz, Maturation-associated
increase in IP3 receptor type 1: Role in conferring increased IP3 sensitivity and
Ca2+ oscillatory behavior in mouse eggs. *Dev. Biol.* **254**, 163-171 (2003).

51. G. Fan *et al.,* Gating machinery of InsP3R channels revealed by electron
cryomicroscopy. *Nature.* **527**, 336-341 (2015).

52. N. Paknejad, R. K. Hite, Structural basis for the regulation of inositol
trisphosphate receptors by ca(2+) and IP(3). *Nat. Struct. Mol. Biol.* **25**, 660-668
(2018).

53. H. Akizawa, E. Lopes, R. A. Fissore, Zn(2+) is essential for ca(2+)
oscillations in mouse eggs. *bioRxiv.* (2023).

54. A. D. Burkart *et al.,* Ovastacin, a cortical granule protease, cleaves ZP2 in
the zona pellucida to prevent polyspermy. *J. Cell. Biol.* **197**, 37-44 (2012).

55. C. Schmitz *et al.,* The E-modulus of the oocyte is a non-destructive measure
of zona pellucida hardening. *Reproduction.* **162**, 259-266 (2021).

56. E. Bianchi, B. Doe, D. Goulding, G. J. Wright, Juno is the egg izumo
receptor and is essential for mammalian fertilization. *Nature.* **508**, 483-487
(2014).

57. B. Y. Kong *et al.,* The inorganic anatomy of the mammalian preimplantation
embryo and the requirement of zinc during the first mitotic divisions. *Dev. Dyn.*
244, 935-947 (2015).

58. G. F. Chanfreau, Zinc'ing down RNA polymerase I. *Transcription.* **4**, 217-
220 (2013).

59. A. S. Prasad, Zinc deficiency in women, infants and children. *J. Am. Coll.*
Nutr. **15**, 113-120 (1996).

60. H. Yokokawa *et al.,* Serum zinc concentrations and characteristics of zinc
deficiency/marginal deficiency among japanese subjects. *J. Gen. Fam. Med.* 21,
248-255 (2020).

61. S. Hojyo *et al.,* Zinc transporter SLC39A10/ZIP10 controls humoral
immunity by modulating B-cell receptor signal strength. *Proc. Natl. Acad. Sci. U.*
S. A. **111**, 11786-11791 (2014).

62. T. Miyai *et al.,* Zinc transporter SLC39A10/ZIP10 facilitates antiapoptotic
signaling during early B-cell development. *Proc. Natl. Acad. Sci. U. S. A.* **111**,
11780-11785 (2014).

63. Z. Lan, X. Xu, A. J. Cooney, Differential oocyte-specific expression of cre
recombinase activity in GDF-9-iCre, Zp3cre, and Msx2Cre transgenic mice.
Biol. Reprod. **71**, 1469-1474 (2004).

64. T. Namiki *et al.,* Uterine epithelial Gp130 orchestrates hormone response
and epithelial remodeling for successful embryo attachment in mice. *Sci. Rep.*
13, 854-y (2023).

65. J. Cocquet *et al.,* Evolution and expression of FOXL2. *J. Med. Genet.* **39**,
916-921 (2002).

66. J. C. Polanco *et al.,* Sox10 gain-of-function causes XX sex reversal in mice:
Implications for human 22q-linked disorders of sex development. *Hum. Mol.*
Genet. **19**, 506-516 (2010).

67. J. Ito *et al.,* Inositol 1,4,5-trisphosphate receptor 1, a widespread Ca2+
channel, is a novel substrate of polo-like kinase 1 in eggs. *Dev. Biol.* **320**, 402-
413 (2008).

68. M. Kamoshita, K. Fujiwara, J. Ito, N. Kashiwazaki, Highly successful
production of viable mice derived from vitrified germinal vesicle oocytes. *PLoS*
One. **16**, e0248050 (2021).

69. A. Inoue *et al.,* Maternal H3K27me3 controls DNA methylation-independent
imprinting. *Nature.* **547**, 419-424 (2017).

70. B. Baibakov *et al.,* Sperm binding to the zona pellucida is not sufficient to
induce acrosome exocytosis. *Development.* **134**, 933-943 (2007).

71. A. Kageyama *et al.,* A combined treatment with progesterone, anti-inhibin
serum, and equine chorionic gonadotropin improves number of ovulated
oocytes in young C57BL/6J mice. *J. Reprod. Dev.* **69**, 223-226 (2023).

72. A. Kageyama *et al.,* Dynamic changes of intracellular zinc ion level during
maturation, fertilization, activation, and development in mouse oocytes. *Anim.*
Sci. J. **93**, e13759 (2022).

73. J. Ito *et al.,* Phosphorylation of inositol 1,4,5-triphosphate receptor 1 during
in vitro maturation of porcine oocytes. *Anim. Sci. J.* **81**, 34-41 (2010).

Figure legends

Fig. 1. Expression of ZIP6 and ZIP10 in mouse ovary. (A) In situ hybridization

in the mouse ovary showed ZIP10 expression in oocyte and granulosa cell from

primordial, primary, secondary and antral follicle. (B) Immunofluorescent staining

for ZIP10 (green) in the mouse ovary showed ZIP10 expression in oocyte

membrane. Arrow indicates primordial follicular oocyte. (C) Immunofluorescent

staining for ZIP6 (green) in the mouse ovary showed ZIP6 expression in oocyte

nucleus and granulosa cells. Arrow indicates primordial follicular oocyte. (D)

Immunofluorescent staining showed ZP2 (green; zona pellucida) and FOXL2

(red; granulosa cells) in the mouse ovary. ZP2 was not stain in primordial follicle,

but primary, secondary and antral follicles stained. FOXL2 observed at granulosa

cells of all stage follicles. The scale bar represents 20 μm of primordial-secondary

follicle and 150 μm of antral follicle (A-D).

Fig. 2 Number of collected oocytes and dynamics of zinc ion in _Zip10$^{d/d}$_

mice. (A) The results of average number of oocytes in each group. Data

represents the average ± SEM. These experiments were repeated at least thrice.

Statistical differences were calculated according to student _t_-test (p > 0.05; no

significant difference). (B) The percentage of extrusion of first polar body at 10,

12 and 14 h after IVM. These experiments were repeated at least thrice.

Statistical differences were calculated according to student _t_-test (p > 0.05; no

significant difference). (C) The morphology of spindle and chromosome

organization in _Zip10$^{f/f}$_ and _Zip10$^{d/d}$_ MII oocytes at 14 h after IVM. Anti–α-tubulin

antibody (green) was used to stain the spindles. Chromosomes were stained with

DAPI (blue). The scale bar represents 10 μm. (D) Comparison with the

fluorescence intensity of intracellular zinc ion in GV, MII and 2PN. The upper

images showed the fluorescence of intracellular zinc ion of oocyte or embryo

treated with FluoZin-3AM. Representative images are shown. The white dotted

circles indicate the positions of oocytes and embryos. Scale bars denote 10 μm.

The lower part showed the fluorescence intensity of zinc ions in oocytes or

embryos. Data represent the average ± SE of the experiments. For each

experiment, 10–20 oocytes/embryos were stained and used for the measurement

in each stage of the experiment, and these experiments were repeated three

times. Statistical differences were calculated according to the welch's _t_-test.

Different letters represent significant differences (p < 0.05).

Fig. 3 Measurement of calcium spike and zinc spark in _Zip10$^{d/d}$_ mice. (A) The

representative images of before or after IVF in mouse oocytes. Left side images

showed before fertilization. Right side images showed after fertilization. The

oocytes increased calcium ion and released zinc ion shortly after fertilization. The white dotted circles indicate the positions of oocytes. Successful fertilization was confirmed by simultaneously monitoring intracellular calcium oscillations with Calbryte 590 AM and excellular zinc ions with FluoZin-3 every 4 s. Capacitated frozen-thawed sperm was added to ZP-free MII at 2 min after imaging start. Orange line showed calcium ion and dark blue line showed zinc ion. Intracellular calcium increases immediately before a zinc spark. (B) The representative images of an MII egg activated with 5 μM ionomycin followed by monitoring of intracellular calcium oscillations with Calbryte 590 AM and extracellular zinc using 20 μM FluoZin-3. The ionomycin was added to ZP-free MII at 2 min after imaging start. Orange line showed calcium ion and dark blue line showed zinc ion. Intracellular calcium increases immediately before a zinc spark.

Fig. 4 Presence of a mechanism to prevent multisperm fertilization in Zip10 cKO mice. (A) The percentages of oocytes with each number of PN at 6 h after insemination. Yellow region showed other including degeneration, degression and fragmentation. Gray region showed unfertilization, namely MII oocytes. Orange showed 2PN2PB, namely embryo possessed one female and male pronucleous (2PN) and second polar body (2PB). Blue region showed multisperm fertilization (3PN2PB). (B) The percentage of fertilized oocytes and developmental embryos. Data represent the average ± SE of the experiments. These experiments were repeated at least thrice. Statistical differences were calculated according to the chi-square test. Different letters represent significant differences ($p < 0.05$). (C) The cell number of blastocyst derived from IVF. Blastocysts were fixed, immunostained, and physically flattened between a slide and coverslip. Photographs represent a single plane of focus. Nuclei are indicated by DAPI staining. The count used inverted fluorescence microscope. These experiments were repeated three times, and each group counted total 46 embryos. Statistical differences were calculated according to the student t-test ($p < 0.05$; significant difference). (D) Western blot of oocytes from $Zip10^{f/f}$ and $Zip10^{d/d}$ mice at 0 or 6 h after insemination using rat anti-ZP2 antibody. Intact ZP2 and the cleaved C-terminal fragment of ZP2 measured 120-130 kD and

undetected, respectively. Expression level of β-actin serves as a protein loading

control. Molecular mass is indicated at the left. (E) MII oocytes and 2PN embryos

from *Zip10^{f/f}* and *Zip10^{d/d}* mice were imaged by confocal microscopy after staining

with rabbit anti-ovastacin (green). Chromosomes were stained with DAPI (blue).

The scale bar represents 10 μm. (F) MII oocytes and 2PN embryos from *Zip10^{f/f}*

and *Zip10^{d/d}* mice were imaged by BZ-X700 microscopy after staining with rat

anti-mouse FR4 (JUNO; green). Chromosomes were stained with DAPI (blue).

The scale bar represents 10 μm.

Supplement Figure legends

Fig. S1 Generation of oocyte specific *Zip6* and *Zip10* conditional knockout

mouse. (A) Schematic of mating pattern of *Zip6* and *Zip10* conditional knockout

mice. (B) Strategy used to develop floxed *Zip6* and *Zip10* allele. The mouse ZIP6

locus is shown at top. A loxP site (black triangle) sequence insert into a region

before and after exon 3 of *Zip6* to produce the *Zip6* floxed allele with loxP sites

flanking exons 3. In the presence of Cre recombinase there is further

recombination resulting in the inactivated *Zip6* allele lacking exons 3 (*Zip6* Δflox).

The mouse ZIP10 locus is shown at top. A loxP site (black triangle) sequence

insert into a region between exons 5 and 6, and a loxP site sequence between

exons 8 and 9 of *Zip10* to produce the *Zip10* floxed allele with loxP sites flanking

exons 6-8. In the presence of Cre recombinase there is further recombination

resulting in the inactivated *Zip10* allele lacking exons 6-8 (*Zip10* Δflox). (C)

Genotyping of flox, Δflox and iCre was performed by genomic PCR. To elucidate

the roles of ZIP10 in the mouse oocytes, we crossed *Zip10^{flox/flox}* (*Zip10^{f/f}*) mice

with Growth Differentiation factor 9 (*Gdf9*)-Cre transgenic mice to generate

Zip10^{flox/flox} Gdf9-Cre (*Zip10^{d/d}*) mice. *Zip10^{f/f}* mice were used as control. Similarly,

we crossed *Zip6^{flox/flox}* (*Zip6^{f/f}*) mice with *Gdf9*-Cre transgenic mice to generate

Zip6^{flox/flox} Gdf9-Cre (*Zip6^{d/d}*) mice. *Zip6^{f/f}* mice were used as control. The tail tips

of 2 weeks old mice were cut. The samples were lysed in 100 μl of DirectPCR

Lysis Reagent (Viagen Biotech) with proteinase K (0.1 mg/ml) at 50°C for

overnight. To inactivate proteinase K, the samples were incubated at over 80 °C

for 1 h. The genotypes confirmed by polymerase chain reaction (PCR), as

suggested by Miyai et al., (2014) (62) and RIKEN BRC using

CAAGGCCAGCCAAAATTCTA (A-F) and GCTTTCCTCCCATCCTGATT (A-R)

to detect Wild-type (292 bp) and *Zip10* flox (420 bp), A-F and

GTGGCATGCGTGGAAGTTAG (B-R) to detected *Zip10* flox null (550 bp) alleles.

the genotyping of *Zip6* flox was confirmed using

CCAGCATTGCCCTCTGTAAGAGTC (*Slc39a6*_F) and

GCCTAAAGAAGATACACTGACACGACG (*Slc39a6*_R) to detect Wild-type (353

bp) and *Zip6* flox (387 bp), GGACCGGTTGCATAGAGGAG (*Slc39a6*_Null_F)

and GAGGCAGGCGGATTTCTGAG (*Slc39a6*_Null_R) to detected *Zip6* flox null

(500 bp) alleles. Similarly, we detected *Gdf9* iCre/+ alleles (200 bp) using

CAGGTTTTGGTGCACAGTCA (21218) and GGCATGCTTGAGGTCTGATTAC

(25494) to suggest by Jackson Laboratory. MII oocytes were used in order to

detect oocyte expression.

Fig. S2 Number of collected oocytes, dynamics of zinc ion and percentage

of fertilization in *Zip6^{d/d}* mice. (A) The results of average number of oocytes in

each group. Data represents the average ± SEM. These experiments were

repeated at least thrice. Statistical differences were calculated according to

student *t*-test (p > 0.05; no significant difference). (B) Comparison with the

fluorescence intensity of intracellular zinc ion in MII and 2PN. Data represent the

average ± SE of the experiments. For each experiment, 10–20 oocytes/embryos

were stained and used for the measurement in each stage of the experiment, and

these experiments were repeated three times. Statistical differences were

calculated according to student *t*-test (p > 0.05; no significant difference). (C) The

percentages of oocytes with each number of PN at 6 h after insemination. Yellow

region showed other including degeneration, degression and fragmentation. Gray

region showed unfertilization, namely MII oocytes. Orange showed 2PN2PB,

namely embryo possessed one female and male pronucleous (2PN) and second

polar body (2PB). Blue region showed multisperm fertilization (3PN2PB). (D) The

percentage of fertilized oocytes and developmental embryos. Data represent the

average ± SE of the experiments. These experiments were repeated at least thrice. Statistical differences were calculated according to the chi-square test (p > 0.05).

Fig. S3 Measurement of calcium spike and zinc spark in *Zip6$^{d/d}$* mice. The representative images of before or after IVF in mouse oocytes. Left side images showed before fertilization. Right side images showed after fertilization. The oocytes increased calcium ion and released zinc ion shortly after fertilization. The white dotted circles indicate the positions of oocytes. Successful fertilization was confirmed by simultaneously monitoring intracellular calcium oscillations with Calbryte 590 AM and excellular zinc ions with FluoZin-3 every 4 s. Capacitated frozen-thawed sperm was added to ZP-free MII at 2 min after imaging start. Orange line showed calcium ion and dark blue line showed zinc ion. Intracellular calcium increases immediately before a zinc spark.

937 **Fig. 1**

939 | Primordial follicle | Primary follicle | Secondary follicle | Antral follicle |

Fig. 2

966 A

B

968 18.7 ± 2.9   22.1 ± 2.5

973 $p > 0.05$

976 D

984 * $P < 0.05$
a-c: $P < 0.05$

985 d: $P > 0.05$

Fig. 4

A

B

C

D

E

F

* *p* < 0.05

* *p* < 0.05