Symbiont loss and gain, rather than co-diversification shapes

honeybee gut microbiota diversity and function

Aiswarya Prasad[1], Asha D Pallujam[2], Rajath Siddaganga[3], Ashwin Suryanarayanan[3], Florent

Mazel[1], Axel Brockmann[3], Sze H Yek[2,4] and Philipp Engel[1], *

[1]Department of Fundamental Microbiology, University of Lausanne, Switzerland

[2]School of Science, Monash University Malaysia, Malaysia

[3]National Centre for Biological Sciences-Tata Institute of Fundamental Research, Bengaluru, Karnataka,

India

4 Institute for Tropical Biology and Conservation, Universiti Malaysia Sabah, Malaysia

*To whom correspondence should be addressed

e-mail: philipp.engel@unil.ch

Keywords

Abstract

Studying gut microbiota evolution across animals is crucial for understanding symbiotic

interactions but is hampered by the lack of high-resolution genomic data. Honeybees, with their

specialized gut microbiota and well-known ecology, offer an ideal system to study this evolution.

Using shotgun metagenomics on 200 honeybee workers from five species, we recovered

thousands of metagenome-assembled genomes, identifying several novel bacterial species. While

microbial communities were mostly host-specific, we found both specialist and generalist

bacteria, even among closely related species, with notable variation between host species. Some

generalists emerged host-specific only at the strain level, suggesting recent host switches.

Unexpectedly, we found no evidence of codiversification between hosts and symbionts. Instead,

symbiont gains, losses, and replacements led to functional differences, such as the ability to

degrade pollen-derived pectin. Our results provide new insights into gut microbiota evolution

and uncover the functional potential of the previously underexplored gut microbiota of these

important pollinators.

Introduction

Animals harbor specialized microbial communities in their gut, which can influence health and

disease susceptibility through diverse functional capabilities[1]. These communities are often host-

specific, where the gut microbiota of individuals of the same host species are more similar in

composition to each other than to those of different host species[2]. In some animal groups, it has

also been observed that more closely related host species harbor more similar gut microbiota in

composition[3]. This host specificity at the community level emerges from the host-restricted

distribution of individual community members. Dispersal limitation and host-filtering processes

have been suggested to contribute to the host-specific distribution of community members in the

animal gut[2,4].

Some gut bacteria (*e.g.*, in termites[5] and hominids[6–10]) appear to have codiversified with their

hosts, as their phylogenies are congruent with that of the host. However, only a small fraction of

the analyzed bacteria showed signs of codiversification in these studies. On the other hand,

examples of generalist gut symbionts exist as well[11,12]. This variation in the distribution patterns

of different gut bacteria suggests they do not follow the same evolutionary trajectory across

various animal species. Indeed, a recent study[13] showed that host specificity could vary across

bacterial lineages in the animal gut (*i.e.*, there are generalists and specialists), and this may not be

explained by microbial phenotype alone[14], with host ecology playing an important role[2]. This

raises the question of why generalist and specialist gut bacteria exist and whether specialist

bacteria primarily evolved *via* repeated switches between hosts or through codiversification with

hosts.

Our current understanding of the evolution and diversification of gut microbial communities

across hosts is limited by at least four technical and conceptual factors. (1) Previous studies have

focused on highly diverse and variable systems across host individuals, making it challenging to

understand whether microbiota change by acquisition of new members or by *in situ* evolution.

(2) Significant differences in the relatedness and ecology of the compared populations hinder the

identification of the drivers of community change over evolutionary time scales and the

prediction of their functional consequences. (3) Samples from different animals often come from

different geographic regions, making geography an important confounding factor that could

explain the host-restricted distribution observed for some gut symbionts. (4) Most comparative

studies are based on 16S rRNA analysis, which lacks the resolution to determine the distribution

of closely related bacterial species, strains and functional gene content across hosts[15].

Comparative studies at a high genetic resolution of the microbiota of closely related animals with

similar phenotypes found in various geographic locations are needed to overcome these

limitations. The gut microbiota of honeybee workers is a promising model offering the combined

advantage of a simple yet specialized microbial community shared among related host species

with similar ecology, social behavior, and overlapping geographic ranges[16–23]. Prominent

honeybee gut microbiota members include Lactobacillus, Bombilactobacillus, Bifidobacterium,

and several proteobacteria. These bacterial lineages from honeybees are usually monophyletic

but belong to host-specific species, of which several can coexist simultaneously in the same host

individual[24–26]. Thus, the predominant members of the honeybee gut microbiota have likely been

acquired in a common ancestor of social bees and subsequently diversified in honeybees by

separation into different host species and distinct ecological niches in the gut[16,27,28]. Co-

diversification has been suggested as the primary mode of evolution, but formal tests of co-

phylogeny have not yet been conducted. So far, most comparative studies are based on 16S

rRNA amplicon sequencing with low genetic resolution or relatively few isolate genomes and

biased towards isolates from the managed Western honeybee *Apis mellifera,* providing limited

quantitative insights into the distribution of divergent strains and species across hosts. Shotgun

metagenomics allows the sequencing of all genomes in a given sample, characterization of the

distribution of community members at high resolution (down to the strain level) across hosts, and

assessment of the functional consequences of differences among hosts. However, this approach

has only been used to study the Western honeybee (*Apis mellifera*)[20,29–31] and the Asian

honeybee (*Apis cerana*)[32,33].

In this study, we applied shotgun metagenomics to individual worker bees of five different

species of honeybees representing the most abundant and prevalent species of honeybees

described so far, sampled across peninsular Malaysia and South India. From a total of 200

shotgun metagenomes from individual worker bee guts, we generated a comprehensive dataset of

metagenome-assembled genomes (MAGs), expanding the catalog of the bacterial genomes

available from these important pollinators and model for microbiome research. We reveal

striking differences in host specificity across host and bacterial species and identify previously

unrecognized but prevalent members of the bee gut microbiota that expand the functional

repertoire of these communities in specific host species. Overall, our results show that ancient

gut microbiota-host interactions can be maintained in the absence of co-diversification.

Results

An expanded catalog of bacterial genomes from the gut of five honeybee species

We shotgun sequenced (35M ±19M Illumina paired-end reads per sample) the hindguts of 200 individual worker bees from 40 different colonies of five honeybee species covering all three major clades of honeybees Micrapis (dwarf honeybee): *A. florea* and *A. andreniformis*, Megapis (giant honeybee): *A. dorsata*, Apis: *A. mellifera* and *A. cerana* sampled in South India and various locations across peninsular Malaysia **(Fig. 1A, Supplementary Table S1, Supplementary Fig. S1**).

We recovered a total of 1,959 high- and medium-quality metagenome-assembled genomes (MAGs, *i.e.* completeness score > 50% and contamination <5%) (**Supplementary Fig. S2**), including 156 from *A. florea*, 163 from *A. andreniformis*, 869 from *A. dorsata*, 324 from *A. mellifera* and 413 from *A. cerana*) (**Fig. 1B, Supplementary Table S2**). The MAGs were discriminated into 150 species spanning 23 genera (prevalence > 10%) using 95% ANI (average nucleotide identity) clustering (**Supplementary Fig. S3**). A majority of these species belonged to genera previously reported to be predominant in the gut microbiota of honeybees, including *Lactobacillus* (formerly Firm5), *Bombilactobacillus* (formerly Firm4), *Bifidobacterium*, *Gilliamella*, *Snodgrassella*, *Frischella*, *Bartonella* and *Apibacter* (constituted by 91 / 150 species clusters), but also lesser known genera such as *Dysgonomonas* (family Dysgonomonadaceae), *Saezia* (family Burkholderiaceae), *Pectinatus* (family Selenomonadaceae), *Entomomonas* (family Pseudomonadacea), *Pluralibacter* (family Enterobacteriaceae), WRHT01 (Desulfovibrionaceae family), CALYQQ01 (family Enterobacteriaceae) and CALYQJ01 (family Lactobacillaceae) (constituted by 21 / 150 species). Around 47% of the identified bacterial

species (71 / 150 species) have not been cultured to date, with many belonging to the

predominant genera (41 / 71 species) and with genomic information currently lacking in public

databases (**Fig. 1B**). The established collection of MAGs massively expands the catalog of

bacterial genomes from the gut environment of honeybees.

Most honeybee species harbor host-specific microbial communities in their gut

We assessed the taxonomic diversity of the gut microbiota across the five honeybee species

using our collection of MAGs. Between 7 - 56M reads from each sample mapped to these MAGs

(**Supplementary Table S3**), providing sufficient depth of bacterial reads for MAG recovery

from individual guts. Rarefaction curves show that our sampling effort prioritizing broad

sampling from different colonies of each host species (**Fig. 1A**) was sufficient to cover most of

the species-level diversity in each host species (**Supplementary Fig. S4**).

Prevalence of the identified genera varied greatly across host species (**Fig. 1C**). *Lactobacillus*,

Bombilactobacillus, and *Bifidobacterium* were consistently present across all five honeybee

species (**Fig. 1B**). All other genera were only found in a subset of the five host species or

occurred sporadically throughout. Notably, samples of *A. dorsata* contained several additional

genera in high prevalence, such as *Dysgonomas* (90.9%), *Pectinatus* (70.5%), and *Saezia*

(90.9%), which were rarely found in the other host species.

Comparison of the species-level bacterial diversity revealed that *A. dorsata* contained the most

diverse microbiota, followed by *A. mellifera*, *A. cerana* and then the two dwarf honeybees (**Fig.**

1C). This pattern holds when considering only species of the shared genera (*i.e.*, genera with a

prevalence of at least 10% in each host), except that *A. dorsata* and *A. mellifera* were not

significantly different from each other (**Supplementary Fig S5**). Interestingly, the bacterial

biomass per individual gut was the highest in *A. mellifera*, followed by *A. dorsata*, *A. cerana* and

the dwarf bees, and hence, it cannot necessarily explain the differences in alpha-diversity (**Fig.**

1E). Most bacterial species displayed significant strain level diversity as measured by the

percentage of single nucleotide variants (SNVs) per bacterial species within each bee gut with

the possible co-occurrence of several closely related strains (**Supplementary Fig. S6**). The

western honeybee *A. mellifera* hosted significantly more strains than other hosts within most

bacterial species, while dwarf bees harbored the least strain-level diversity (**Fig. 1F**).

Taxonomic composition was host-specific at the species level as suggested by β-diversity

analyses (PERMANOVA Sorenen distance Ω^2=0.197, p=0.001, N=195), but with no distinction

between the two dwarf bee species (**Fig. 1E**) and no influence of geography (Ω^2=0.009, p=0.078,

N=131 excluding two host species sampled unequally across locations) (**Supplementary Fig.**

S7-9). We found no evidence for phylosymbiosis, as there was no correlation between bacterial

community distance and host divergence (Spearman's p-value = 0.49, mean q-value =

0.619±0.204 across 1000 mantel tests, **Fig. 1H**). For example, *A. andreniformis* and *A. mellifera*

are the most distant host species but have much smaller distances in bacterial community

composition than other pairs of host species.

Host specificity of individual community members varies across bacteria and host species

Specificity at the community level emerges because of the host-specific distribution of individual

community members[13]. We calculated the specificity per bacterial species based on Rohde's

Index[34], which considers the prevalence of each community member across samples from

different hosts (**Fig. 2A, Supplementary Table S4**).

Overall, we found large differences in the specificity of individual bacterial species both within

and between genera. For example, the genus Bombilactobacillus mainly consisted of generalist

species found across all five bee species, while the genera Lactobacillus or Bifidobacteria

contained specialists and generalists (**Fig. 2A**). Some species showed an intermediate level of

specificity as they were shared only among a subset of the five honeybee species and were more

prevalent in one host than another (*e.g.*, one of the *Snodgrassella* species). Few shared genera

consisted of only specialist species (*Apibacter* and Commensalibacter). Notably, there was a

clear difference in the specificity of bacterial species across hosts. The microbiota of *A. dorsata*

and *A. mellifera* were mainly composed of specialist species, while the microbiota of the smaller

bee species (*A. cerana*, *A. florea*, and *A. andreniformis*) was primarily composed of shared

species (**Fig. 2A**)

Bacterial species may be shared between host species but exhibit host specificity at the strain

level. To look for such strain-level host specificity, we measured the extent of shared SNVs

between samples for each bacterial species shared between two or more host species using

popANI[35]. We found both bacterial species that segregate and species that do not segregate by

host species in terms of strain-level composition (**Fig. 2B, Fig. 2C**). This pattern was observed

across all analyzed genera (**Fig. 2B, Fig. 2C**). Moreover, in some cases, we also found that

samples segregated by sampling location (**Fig. 2B, Fig. 2C**). Overall, 75% of the species were

host-specific at either strain or species level.

Host switches and species gains and losses interrupted the host-specific evolution of gut

microbiota members of honeybees

Codiversification has been described in other host-associated microbiomes as an important driver

of host-specificity patterns[6,9,10]. To test for evidence of codiversification in the honeybee gut

microbiota, we constructed core genome phylogenies for all prevalent genera using the collection

of MAGs assembled from the five honeybee species (**Fig. 3**). We used genus-level phylogenies

to test for codiversification because divergence time within genera was roughly equivalent to

divergence times in the *Apis* genus (see methods). While most honeybee-derived MAGs from a

given bacterial genus were monophyletic relatives to bacteria found in bumble bees and stingless

bees, they belonged to distinct clades corresponding to the species clusters determined by ANI

(see above). We identified several novel lineages of bacteria, especially in the giant honeybee

Apis dorsata, that have not been isolated or represented in public databases before. Among these

are several species of the genus *Dysgonomonas* forming several monophyletic clades, related to

species isolated previously from cockroaches[36] and human stool[37], among other sources. In

agreement with the observed differences in host specificity, a subset of the identified species-

level clades were exclusive to one host species (24% - 47% of all nodes), while others were

mixed. To test for codiversification, we used the Hommola cospeciation test, a generalization of

the Mantel test for arbitrary host-parasite interaction graphs[38]. We then carried out second-order

permutation tests (N=100) with shuffled tip labels to account for the rate of false discoveries as

described before[39]. Overall, we found that in all nine tested genera (trees containing multiple

nodes containing host-specific clades from three or more host species), we detected no more

significant codiversifying nodes than expected by chance (**Supplementary Table S5 and**

Supplementary Table S6). Overall, we show that honeybees and their gut bacteria did not

codiversify despite high host specificity. Instead, as seen in the phylogenies (**Fig. 3**), deep-

branching host-specific clades are interrupted by recent generalist clades, and deep-branching

generalist clades comprise adjacent host-specific clades (and show more host-specificity at the

strain level), indicative of a history of intense shuffling of the associations.

Functional differences in the gut microbiota across honeybees

To test whether the host specificity observed at the taxonomic level translates into functional

host specificity of the gut microbiota, we performed a functional analysis of the MAGs. Of

4,759,622 prokaryotic ORFs in our MAG database, more than half (2,685,895 ORFs) were

assigned a KO identifier. The rarefaction curve analysis of these KOs indicated that most of the

functional diversity across the analyzed samples was recovered with our MAG database

(**Supplementary Fig. S10**). Overall, the number of KOs per sample followed a similar trend as

the taxonomic diversity, with *A. dorsata* having the most and *A. florea* having the fewest KOs

(**Fig. 2A**).

Multivariate analysis of normalized KO counts (and the Robust Aitchison distance) showed that

the functional composition of the gut microbiota of five honeybee species was distinct, even

between *A. florea* and *A. andreniformis,* which overlapped in composition at the taxonomic level

(Ω^2=0.312, p=0.001). To understand which bacteria drive these differences, we performed a

similar analysis using counts of all KO families in each high- and medium-quality MAG across

all samples. As expected, MAGs of bacterial species from the same genus clustered across all

host species, suggesting that they occupy similar functional niches and that the overall functional

capacity of the bee microbiota is conserved across hosts. Yet, there were some important

differences. Some of the genera only found in *A. dorsata* formed distinct clusters (*e.g.*,

Dysgonomonas), suggesting that these bacteria occupy unique functional niches in the

microbiota of this host. Other genera (*e.g.*, Saezia and Entomomonas) clustered with community

members prevalent in other honeybee species (*e.g.*, Snodgrassella) but absent from *A. dorsata*,

indicating potential functional replacements. Finally, although MAGs from the same genus

mostly overlapped in terms of their functional profile, we found some cases where they clustered

by host of origin, suggesting host-specific differences in the functional repertoire (and

potentially, adaptation) within these genera (**Supplementary Fig. S11**).

Since pollen-derived glycan (*i.e.*, pectin and hemicellulose) breakdown has been described as

one of the significant catabolic functions of the gut microbiota *of A. mellifera*[30,40], we specifically

looked at the Carbohydrate-active enzyme (CAZyme) repertoire of the microbiota across the five

host species. We detected 19,562 CAZyme families among the prokaryotic ORFs in our MAG

database. Overall, the composition of these gene families was host-specific (**Supplementary**

Fig. S12). While all five honeybee species harbored a similar set of CAZymes for pectin,

hemicellulose, and cellulose degradation, *A. dorsata* contained a more extensive and diverse set

of such gene families, especially many of which are involved in pectin degradation (**Fig. 4D**). To

understand which bacteria drove these differences, we considered the number of genes belonging

to CAZyme families from each bacterial species. Previously recognized members contributed

most families to degrade pollen-derived glycans, specifically *Gilliamella* and *Bifidobacterium*

(**Fig. 4E**), except in the giant honeybee *A. dorsata, where* the genus *Dysgonomonas* also

contributed key enzyme families involved in pectin degradation such as PL1, PL11, GH28,

GH78, GH128, or CE12. Interestingly, the CAZyme genes of Dysgonomonas were often

organized in large genomic islands reminiscent of specialized PULs (polysaccharide utilization

loci) found in gut symbionts of mammals[41], suggesting that this community member expands the

functional capability to degrade plant-derived fibers in this particular honeybee species (**Fig. 4F**).

Discussion

Many animals harbor host-specific bacterial communities in their gut[2–4,13]. However, it is still unclear whether and why host specificity varies among closely related host species or bacterial lineages, which evolutionary processes underlie the observed patterns of specificity and their functional consequences. Our comparative shotgun metagenomic analysis of the gut microbiota of five honeybee species provides novel insights into these questions. It advances our understanding of the evolutionary processes that govern gut microbiota composition across closely related animal species.

We find that the community composition of the gut microbiota of three of the five analyzed honeybee species is host-specific, confirming previous results based on amplicon sequencing[16] or less comprehensive metagenomic datasets[32]. However, in contrast to *A. mellifera*, *A. cerana* and *A. dorsata*, the gut microbiota of the two phylogenetically ancestral dwarf honeybee species, *A. florea* and A. *andreniformis,* were indistinguishable from each other, suggesting that host specificity is not absolute but can vary between closely related host species. Our genome-level data, coupled with our broad sampling design, allowed us to probe further than community-level patterns to identify the host-specificity of individual clades of community members.

We found that 75% of the bacterial species detected were host-specific at the bacterial species or strain level. First of all, we found that host specificity emerges at different taxonomic levels, for example, at the genus (such as *Saezia* and *Dysgonomonas* in *A. dorsata*), species (such as many *Gilliamella*, *Snodgrassella* and *Lactobacillus* species), and strain level (such as *Bombilactobacillus mellis-36_2*). Considering that all of these bacteria belong to clades specific

to social bees, this suggests that the host-specific association between different gut symbionts

and their honeybee hosts has been established across different phases in their evolutionary

history. These findings also highlight that host-specific associations can be overlooked without

strain-level/genomic resolution.

Another key finding of our analysis is that the degree of host-specificity varies across bacteria

(i.e., there are generalist and specialist species). This has also been observed in other

systems[13,14,42] and linked to bacterial phenotypes, such as aerotolerance and spore-formation, that

affect dispersal ability[13,43–46]. However, in addition to previous work, we observed unexpected

high variability in host specificity between closely related bacteria, i.e., species belonging to the

same genera, suggesting that microbial traits, such as aerotolerance, usually conserved within

genera, cannot explain variation in specificity patterns. Instead, we find host identity to be an

important determinant of the specificity of individual community members. Most of the highly

host-specific bacterial species were found in *A. dorsata* and *A. mellifera,* while the communities

of *A. cerana* and the two dwarf honeybees were dominated by shared bacterial species. *A.*

mellifera and *A. dorsata* guts may harbor more specific ecological niches than the other bee

species, facilitating the colonization of additional 'specialist' bacteria. This is consistent with the

observation that *A. cerana* and the two dwarf bees harbor significantly fewer bacterial species

per individual bee than *A. dorsata* and *A. mellifera*. The smaller body and colony size of these

bees may limit carrying capacity and niche diversity in the gut, as proposed previously[16,32].

Another possibility is that the host control over their gut microbiota is stronger in some bee

species than in others, as observed in other animal species[47]. Still, the reason for this variation is

unclear.

In contrast to studies of other social bees[16,27,48,49], we did not find any evidence for co-

diversification in the honeybee gut microbiota. This might be because previous analyses did not

(1) apply extensive tests and corrections[39], (2) did not use genome-scale phylogenies, or (3) did

not sample the full breadth of microbiota diversity as they relied on isolate genomes. Our results

also contrast with codiversification findings in mammals and, in particular, hominids[5–10].

However, even in hominids, only 16%-36%[6,8,10] of all tested bacterial clades showed statistically

significant signals of codiversification, suggesting that the evolution of only a minority of the

microbiota is driven by codiversification and that codiversification might be the exception rather

than the rule in both mammals and social bees.

Overall, our results show that ancient gut microbiota-host interactions can be maintained in the

absence of codiversification. Instead, host switches and symbiont gains and losses between

related species have been frequent, suggesting that they govern compositional changes in gut

microbiota over evolutionary time. This is surprising given the social lifestyle of bees that

facilitate within-host species transmission mode. In contrast, ecological opportunities for

spillover and invasion of the gut microbiota are not uncommon, maybe due to the sympatric

distribution of *Apis* species. Furthermore, since all five honeybee species have similar dietary

preferences, gut structure, and physiology and lack an adaptive immune system, the barrier to

colonization between native and non-native bacteria may be low. This is supported by

experiments with microbiota-free bees, which showed that bacterial strains isolated from

honeybees can colonize a non-native honeybee species and sometimes even outcompete native

bacterial strains[16,28].

The host-specific nature of the microbiota of these honeybees translates into host-specific

microbial functional potential. In particular, both the specific repertoire of conserved members

and the gain of new members with unique potential significantly impacted microbiota functions.

For example, honeybee-specific lineages of *Dysgonomonas*, a genus of polysaccharolytic

Bacteroidetes also present in the gut of cockroaches, termites, and humans[36,37,50], greatly

expanded the repertoire of pectin and hemicellulose degradation genes in the gut microbiota of

A. dorsata without necessarily replacing another gut symbiont. A. dorsata may be distinct in its

broad exposure to more diverse microbes compared to other honeybees, given its highly

migratory nature[51]. Interestingly, differences in diet may not explain differences in microbiome

diversity, considering that *A. dorsata* tends to forage on fewer mass-flowering plants[52] , and

plant-pollinator network analysis suggests that *A.florea*, *A. cerana*, and *A. dorsata* show

substantial overlap in the flowers they visit[53]. In summary, the honeybee gut microbial

community shows a highly dynamic evolution despite the presence of a few conserved lineages.

Our results form the basis for more detailed functional characterization of diverse bacterial gut

symbionts previously not recognized as important members of the honeybee gut microbiome. It

vastly expands our knowledge of bee microbiota obtained chiefly from the managed western

honeybee. More generally, it advances our understanding of host specificity in gut microbiota-

host associations by showing that even ancient gut microbiota-host interactions can be

maintained in the absence of co-diversification via dynamic gain, losses and exchanges of

microbes. This history of frequent turnover can have a critical impact on microbiome functions

and suggest that apparently conserved ancient gut microbiota–host associations are actually

susceptible to change over evolutionary time.

Methods

Sampling and storage

All honeybee workers (10-50 individuals per colony) collected from colonies across locations described in the metadata sheet (**Supplementary Table S1**) were stored in small ventilated boxes and processed for storage within the day. We stored the bees in absolute ethanol at -20°C until dissection and DNA isolation, sufficiently preserving the sample for sequencing the microbiome[54]. Our sampling strategy prioritized covering many colonies across locations to capture the diversity of each host species rather than many individuals from the same colony, which has been done before[29,32].

Dissection and DNA extraction

As the hindgut contains most of the bacterial biomass, we aimed to decrease the ratio of DNA from the host cells to bacterial cells. Hence, we removed the gut tissue by carefully opening the cuticle, removing the stinger and venom glands, and, finally, cutting the tissue near the pylorus, removing the malpighian tubules and sections, including the midgut and above. Then, the tissue was transferred to a bead-beating tube containing 478 µL of sterile PBS and 2.0 mm sterile glass beads and subjected to bead-beating for 30s at 6 m/s in FastPrep-24 5G (MP Biomedicals). Next, 0.1 mm zirconia beads along with 250 µL of 2x CTAB, 2 µL of β-mercaptoethanol and 20 µL of proteinase K (20 mg/mL) was added and bead-beating was carried out for 30s at 6 m/s in FastPrep-24 5G (MP Biomedicals). The homogenate was incubated for 1h at 56°C. Next, 750 µL of Phenol-chloroform isoamyl alcohol (25:24:1, v/v) at pH 8.0 was added and mixed thoroughly, followed by centrifugation for 10 min at 16,000 xg. The upper aqueous layer was transferred to a new tube, and 500 µL of chloroform was added and mixed thoroughly. The mixture was

centrifuged for 10 min at 16,000 xg, and the 300 μL of the upper phase was mixed with 900 μL

of chilled absolute ethanol and incubated at -20°C overnight. Next, the precipitated DNA was

pelleted by centrifugation for 30 mins at 16,000 xg at 4°C. After two washes with 70% ethanol,

it was resuspended in 50 μL of nuclease-free water. Next, it was treated with RnaseA and

purified using magnetic beads (Clean NGS). DNA concentration was measured using Qubit™

(Thermo Fischer Scientific). Bacterial absolute abundance was measured using qPCR with

primers targeting the 16S rRNA region as described earlier[55].

Metagenomic sequencing

Library preparation was carried out using the Nextera XT library kit (Illumina), according to

manufacturer instructions, with the number of PCR cycles chosen to account for the amount of

DNA in samples. Illumina sequencing was carried out using Novaseq (2x 150 bp) with a target

of 10-50 million microbial reads. Since there was a significant difference between host species in

the total gut microbial content as determined by qPCR using universal primers, first, a run of

sequencing was carried out without normalization of the amount of samples added to the library

to estimate the percentage of host DNA and microbial DNA in the library made from each

sample. This was useful in ensuring that there was enough DNA from the bacterial community in

all the samples, including those where the library might have been dominated by host DNA due

to lower bacterial biomass. Further sequencing was then carried out after pool correction, where

the library was adjusted in such a way when possible that at least 15 million microbial reads

were obtained (based on expectation from the first shallow sequencing), even for the samples

with the lowest percentage of microbial DNA. All library preparation and sequencing was done

at the Lausanne Genomic Technologies facility (GTF), UNIL. Details about all further

bioinformatic analysis, including the code used, can be found on GitHub

(https://github.com/Aiswarya-prasad/honeybee-cross-species-metagenomics) and Zenodo

(https://zenodo.org/doi/10.5281/zenodo.13732977) in the repositories.

Mapping to databases

To assess the amount of bacterial, host, and other reads in each sample, reads were mapped to

host and MAG databases. The host database comprised draft or complete genomes of all the four

host species (GCA_000184785.2, GCA_000469605.1, GCA_001442555.1, GCF_003254395.2)

except *Apis andreniformis* since no genome was available in public databases for this host

species. A database was compiled using representative MAGs identified for each 95% ANI

cluster or bacterial species with the highest quality score (calculated as described by dRep[56]

using various MAG characteristics, including checkM contamination and completeness).

Trimmed and quality-filtered reads were mapped to the host database using BWA-mem[57]

followed by the MAG database (and *vice versa* to account for non-specific mapping). Three

samples were removed from the downstream analysis as their metagenomic reads did not map

the host that they were labeled as (**Supplementary Table S2**) and might have faced cross-

contamination during sample processing. Samples for which relatively few reads mapped to the

MAG database often were also low in bacterial biomass as measured by qPCR (**Supplementary**

Fig. S13). Indicating that the sequencing and workflow to detect microbial diversity in each

sample was otherwise robust. Initial taxonomic profiling of the reads using mOTUs[58] detected

the expected core marker genes from various honeybee gut microbiota members.

Assembly and binning of MAGs

Raw reads were trimmed and quality filtered before assembly using Trimmomatic[59] and bbtools.

For some samples that had a large number of reads (19/150 samples) due to which assemblies

were not successful, bbnorm was used to down-sample regions of high depth and for samples

with too many host reads due to which assemblies failed (3/150 samples), host-filtered reads

were used. The detailed workflow used for assembly and binning into MAGs is described in

GitHub (https://github.com/Aiswarya-prasad/honeybee-cross-species-metagenomics). Briefly,

contigs were assembled using metaSpades[60]. Filtered reads were mapped against assembled

scaffolds (all against all samples), and MetaBAT2 used the coverage information[61] to bin

scaffolds into MAGs. The quality and completeness of resulting MAGs were assessed using

CheckM[62] and low ($< 50\%$ completeness or $>5\%$ contamination), medium (50-90%

completeness and $<5\%$ contamination), high ($> 90\%$ completeness and $<5\%$ contamination).

MAGs were then clustered into groups averaging 95% average nucleotide identity (ANI) using

dRep[56].

Taxonomic annotation was carried out using GTDB-tk[63]. No manual curation of MAGs was

performed; however, Kaiju[64] (nr database) and Kraken2[65] (custom database including bacteria,

viruses and honeybee) classification of scaffolds screened by random sampling further confirmed

that scaffolds from the same MAG matched each other and the taxonomic assignment of the

MAG. Each cluster of MAGs was named to reflect the GTDB species name or, in cases where

none was assigned, a unique identifier, including the genus name. Combined with the inspection

of ANI heatmaps (**Supplementary Fig. S13**), a final set of species names was assigned to each

of the 95% ANI clusters.

Taxonomic profiling

InStrain[35] was used according to its recommended workflow for taxonomic profiling at the

bacterial species and strain level. Trimmed and quality-filtered reads were competitively mapped

to the database of representative MAGs using the tool bowtie2[66], which presents mapping scores

in a format compatible with InStrain. Among many approaches to detecting microbial species in

a dataset, this stands out in its appropriate filters applied to mapped reads, including read ANI,

which helps differentiate genomes of closely related species (<95% ANI) and breadth of the

genome mapped to avoid falsely identifying species as present due to high coverage of a region

shared with others or horizontally transferred. Moreover, genome coverage of a given species

can be misleading when comparing communities of varying diversity, as an equally abundant

species would have higher coverage in a community with low diversity. A species was

considered present if it had a median coverage of at least 1x across all scaffolds and its genome

(comprising all scaffolds) was at least 50% covered by reads. Relative coverage, when used, was

obtained by dividing the coverage of each species by that of the sum of all and qPCR copy

numbers of the bacterial 16S rRNA region were used to estimate absolute abundance for

comparison between samples where mentioned. For samples from India (n=50), qPCR values

were interpolated based on the median of the host species. The underlying data (presence-

absence, relative abundance etc.) and normalization approach are explicitly indicated for each

downstream analysis section. After filtering samples as mentioned earlier and removing samples

with fewer than 1 million filtered reads mapping to the representative database of MAGs, a total

of 195 samples were included in the analysis (*A. mellifera* 34, *A. cerana* 45, *A. dorsata* 44, *A.*

florea 42 and *A. andreniformis* 30).

Strain-level composition profiles for each bacterial species in the representative MAG database

were generated per sample using InStrain in database mode. The percentage of variant sites was

calculated as the percentage ratio of the number of SNVs to the total length of the MAG covered.

A site is counted as an SNV if the site is covered at least 5x and the allele is present at a

frequency of at least 0.05. Regions mapping with a read ANI of <0.92 (might be cross-mapping

of reads from a different species) or coverage of <1x were not considered. To compare the strain

profile of samples, the popANI measure, as comprehensively described by InStrain[35], was used.

Briefly, a value from 0 to 1 was calculated for each species for each pair of samples in which the

genome was covered at least by 5x and at least half of the genome was covered in both samples.

A position of the genome was considered if there was more than one allele with a frequency of at

least 0.05. If both samples share no alleles in that position, it is counted as a popANI

substitution. The value for that pair of samples was then calculated as the ratio of positions

containing popANI substitutions and the number of positions compared. This value was used to

create a matrix of Jaccard distances between all pairs of samples.

Phylogeny construction and codiversification test

To gain a rough estimate of the taxonomic level at which to expect codiversification, we

computed a core genome phylogeny using the nucleotide sequence of 120 bacterial marker genes

for all medium and high-quality MAGs identifying core genes using GTDB-tk, aligning in a

codon-aware manner using MACSE[67] and constructed using IQ-TREE[68]. From this tree, the

median substitutions per site for MAGs of the same species was 0.014, for different species of

the same genus was 0.39, and for different genera was 2.65 (**Supplementary Table S7**). Bacteria

evolve at a rate in the order of magnitude of 10^{-3} substitutions per site per million years[69]. Since

the *Apis* clade is about 30 million years old[70], gut microbes that diverged during this timeline

could be as divergent as different species within a genus or closer, confirming that genus-level

trees are appropriate for these tests.

OrthoFinder[71] was used to infer orthologous gene families of all medium and high-quality MAGs

for each genus along with previously published genomes of the same genus from other

environments as outgroup (details of genomes used in metadata files included in the Github

(https://github.com/Aiswarya-prasad/honeybee-cross-species-metagenomics) and Zenodo

(https://zenodo.org/doi/10.5281/zenodo.13732977) repository. Phylogenies were constructed

using IQ-TREE using the amino acid sequences of genus-level single-copy core genes aligned by

MAFFT[72]. Since medium-quality MAGs were included in phylogenies, core genes can be

randomly missing in some MAGs. Naively defining core genes as those present in all MAGs

would result in the exclusion of most orthologs. Hence, as implemented in OrthoFinder, when

<1000 single-copy orthologs are present in all MAGs, it iteratively identifies the orthologs that

are found once in N-1 MAGs until either considering one less MAG does not double the number

of orthologs or more than half the MAGs are dropped. Core single-copy orthologs identified

were in agreement with those orthologs identified as core by mOTUpan[73], which uses a Bayesian

approach informed by the completeness score of MAGs. The number of orthologous gene

families (OG) used for core genome phylogenies and the minimum percentage of MAGs of the

genus in which each OG was detected (*e.g.*, 100% means all OGs were detected in all MAGs and

50% means every OG was found in 50% or more of the MAGs): Apibacter - 1096 (94.5%),

Apilactobacillus - 704 (81.8%), Bartonella - 705 (81.5%), Bifidobacterium - 118 (84.0%),

Bombilactobacillus - 194 (89.4%), CALYQJ01 - 883 (89.5%), CALYQQ01 - 1221 (84.6%),

Commensalibacter - 1120 (96.6%), Dysgonomonas - 223 (86.5%), Entomomonas - 1144

(91.7%), Frischella - 984 (85.9%), Gilliamella - 189 (88.0%), Pectinatus - 1143 (97.3%),

Pluralibacter - 1419 (100.0%), Saezia - 1000 (92.3%), Snodgrassella - 799 (91.5%), WRHT01 -

1897 (100.0%)]. Further information about the parameters and best-fit model chosen to compute

the phylogenies can be found in the log files included in the corresponding directory containing

IQ-TREE results in the Zenodo (https://zenodo.org/doi/10.5281/zenodo.13732977) repository.

A host phylogeny inferred based on complete mtDNA sequences of honeybee species rooted by

a bumble species published before[74] was used for codiversification tests. Tests were carried out

using the Hommola cospeciation test[75] (implemented in Python) on all nodes containing at least

7 MAGs from at least three host species and with a maximum tip-tip distance not exceeding 1 for

100 permutations. This was carried out on trees with tip labels artificially permuted randomly

(N=100) to estimate the number of nodes displaying significant results by chance, as described in

an earlier study[39]. Thresholds were set at various levels for comparison of the number of

significant nodes in the original tree and the median number of significant nodes in tip label

permuted trees: relaxed ($p<0.05$), medium ($p<0.05$ and $r>0.75$) and strict ($p<0.01$ and $r>0.75$).

Gene functional annotation and profiling

Open reading frames (ORFs) were identified on all assembled scaffolds of each sample using

Prodigal[76]. ORFs marked partial or shorter than 300 bp were removed. Further, those ORFs

detected on scaffolds identified as being of eukaryotic origin by Whokaryote[77] or unclassified as

bacteria by Kaiju (nr database) were also excluded. The remaining ORFs were then annotated

using DRAM[78] to obtain KEGG ortholog (KO) identities. Finally, the gene catalog comprising

all these ORFs was assigned to CAZyme families as implemented in Cayman[79].

Trimmed and filtered reads from each sample were mapped back to their respective assemblies

using bowtie2. Low-quality (Q<20) alignments were removed, and coverage per gene/ORF was

calculated using BEDtools[80]. Genes were considered detected if >5 reads were mapped to it and

> 90% of the bases in the gene were covered by at least a read. The abundance of each gene was

then calculated in terms of reads per kilobase per million (RPKM) calculated as follows:

$RPKM = \frac{\# \, Mapped \, reads}{\left(\frac{Gene \, length}{1000}\right) * \#Total \, reads * 10^6}$. For KOs and CAZymes, summed RPKMs of constituent

genes were used. To estimate the average count of CAZymes of interest (all families other than

GT and AA families) in each genus per individual, the RPKM of all CAZymes in each genus

was summed per sample, and then the average of this sum across samples was converted to a

relative proportion to represent the relative contribution of each genus on average in each host

species and plotted. Only genera with a prevalence >0.05 in the respective host were considered.

Statistical tests and bioinformatic analyses

All details about the specific parameters used for each tool and their versions, along with conda

environment specifications, are included in the Github (https://github.com/Aiswarya-

prasad/honeybee-cross-species-metagenomics) and Zenodo

(https://zenodo.org/doi/10.5281/zenodo.13732977) repositories. In general, bioinformatic

workflows were implemented using Snake make (v7.28.1) to parallelize jobs on a computing

cluster at UNIL, preprocessing and filtering using Python scripts, visualization using ggplot with

several related packages in R and iTOL for phylogenetic trees and arrangement and decoration of

plots using Adobe Illustrator. Significance in boxplots was tested using the Wilcoxon Rank Sum

test (significant if p<0.05). Distance matrices were calculated using the method indicated in their

respective plots- The package Betapart[81] was used to estimate the turnover component of the

Jaccard and Sorensen distances. Multivariate analyses were visualized using PcoA or NMDS

and tested using the adonis2 function implemented in the R package Vegan. The effect size was

adjusted for sample size to get Ω^2 using the adonis_OmegaSq function from the package

Russel88/MicEco (https://zenodo.org/badge/latestdoi/83547545). To test for phylosymbiosis, we

minimized bias due to pseudo-replication by performing 1000 permutations. Each permutation

included five randomly selected samples, one of each host species, tested for correlation between

bacterial and host distances using a Mantel test (as implemented in the Vegan package). The p-

values were adjusted using the FDR method.

Acknowledgments

We thank the following people and organizations for supporting honeybee sampling and access. The Nature Inspired team (John Chan, Steven Wong, & JC Tan), Mr. HoneyBee Farm, The Orang Asli community (Temengor, Perak), Sherwani Honey Bee, Mr. Wee, Mr. Steve, and MY Bee Saviour. The Cayman team (Quinten Ducarmon, Nicolai Karcher, and Christian Schudoma) for their support with early access to their software and custom annotation of our dataset. The Lausanne Genomics Technology Facility (GTF) team at the University of Lausanne. This work was supported by a Spirit grant (grant number 189496), an SNSF Consolidator grant ('GLOBEE,' grant number 213860), and the NCCR Microbiomes, a National Centre of Competence in Research (grant number 180575), funded by the Swiss National Science Foundation.

Author contributions

Conceptualization – PE, SHY, AP, FM; Methodology – AP, FM, PE; Software – AP; Validation – AP; Formal analysis – AP; Investigation – ADP, AP, RS, AS; Resources – PE, SHY, AB; Data curation – AP; Writing - Original Draft – AP; Review and Editing – PE, FM, AP, AB, SHY; Visualization – AP; Supervision – PE, SHY, FM, AB; Project administration – PE, SHY, AB; Funding acquisition – PE, SHY

Competing interests: All authors declare they have no competing interests.

Data availability: Raw metagenomic data has been deposited to the NCBI Sequence Read Archive (SRA) under the Project ID PRJNA1157353. The collection of MAGs, some important

intermediate files, and tables can be found in the Zenodo

(https://zenodo.org/doi/10.5281/zenodo.13732977) repository.

Code availability: Code and details of parameters and software used are available at

https://github.com/Aiswarya-prasad/honeybee-cross-species-metagenomics archived at

https://zenodo.org/doi/10.5281/zenodo.13732977

Figures

Figure 1. Community-level patterns in the honeybee gut microbiome across taxonomic scales. (A) Apis bee phylogeny with tips indicating extant species and the colored tips species included in this study. Pictures of bees in [82] are to-scale. Numbers and text next to species names indicate the sampling design. (B) Barplot showing the number of MAGs recovered from this study for which no isolate genomes of the same species (>95% ANI) were found in GTDB, those that were represented by MAGs or SAGs in GTDB and those that are similar to published genomes from isolated bacteria represented in GTDB. (C) Heatmap with the prevalence of each

genus within each host species showing genera with prevalence > 10% excluding the following three: *Klebsiella* (*K. pneumonia*) – 10% in *A. mellifera*; Pantoea (P. dispersa) – 17% in *A. mellifera*; *Melissococcus* (*M. plutonius*) – 13% in *A. andreniformis*). (D) Boxplot of the number of bacterial species in each honeybee gut across host species. (E) Boxplot of the number of 16S rRNA gene copies per 0.1μL of individual bee gut homogenate across host species. (F) Boxplot of the percentage of variable sites per individual of each host species per genus (four most prevalent genera shown), with each box including points represented by all species of the respective genus found in individuals of that host species. (G) PCoA plot based on the turnover component of Sorensen distance across host species. (H) Scatter plot of microbiota dissimilarity (as the turnover component of Sorenson distance) as a function of host species divergence based on 16S+12S rRNA[83] with white points with thick black outlines marking the median of values for the comparisons of all pairs of samples from a given pair of host species.

Figure 2. Host-specificity of individual gut microbial lineages across honeybee hosts. (A)

Scatter-pie plot of Rohde's index of specificity with each circle representing a bacterial species

with the size representing the number of bees in which the species was found, and the pie chart

showing the proportion of samples belonging to each honeybee host. The adjacent boxplot shows

Rohde's specificity index for each species found in each honeybee host. (B) Distribution of species richness per genus across three categories of specificity levels: host-specific at the species level (grey), host-specific at the strain level, but the species is detected across hosts (dark blue) or not specific even at the strain level (light blue). (C) NMDS plots from a strain level generalist and specialist bacterial species are included as an example. The Jaccard dissimilarity was calculated in terms of strain level composition (popANI) tested using PERMANOVA (significant if the p-value is < 0.05). Only bacterial species in at least five samples and at least two host species and samples with enough coverage, the bacterial species to recover SNPs, were included.

Figure 3. Bacterial genus-level core genome phylogenies of MAGs. Core genome phylogenies of MAGs from various genera found in more than one host species with the color at the leaf tip indicating the host species from which the MAG was recovered and solid grey boxes the set of genomes included to represent outgroup species of the genus from environments other than honeybee guts. The scale bar represents 0.1 amino-acid substitutions per site. The phylogenies are maximum likelihood trees inferred by IQ-TREE (see Methods).

Figure 4. Functional potential of honeybee gut microbiota across host species. (A) Boxplot

of the number of KEGG ortholog (KO) families detected per individual of each host species

Wilcox test ($p < 0.05$) indicated on the plot. (B) PCoA of KO family composition across all samples was visualized using the Robust Aitchison distance on a matrix of reads per kilobase per million (RPKM) summed across genes belonging to the respective KO family. Each point on the plot represents an individual bee gut. PERMANOVA test indicated on the plot ($p < 0.05$). (C) PCoA of all KO families from genes annotated within MAGs across all samples from each host species plotted on the same scale across host species using the Robust-Aitchison distance on the RPKM matrix of KOs of MAGs such that each point on the plot represents a MAG recovered from an individual bee. (D) Dotplot of the mean number of genes from each CAZyme family with a potential role in pectin, hemicellulose, or cellulose degradation per individual of each host species. (E) Barplot of the proportion of genes from CAZyme families contributed by each bacterial genus. (F) Loci within Dysgonomonas MAGs encoding putative polysaccharide utilization loci (PULs).

References

1. Levin, D. *et al.* Diversity and functional landscapes in the microbiota of animals in the wild. *Science* **372**, eabb5352 (2021).

2. Mallott, E. K. & Amato, K. R. Host specificity of the gut microbiome. *Nat. Rev. Microbiol.* **19**, 639–653 (2021).

3. Lim, S. J. & Bordenstein, S. R. An introduction to phylosymbiosis. *Proc. R. Soc. B* (2020) doi:10.1098/rspb.2019.2900.

4. Mazel, F. *et al.* Is Host Filtering the Main Driver of Phylosymbiosis across the Tree of Life? *mSystems* **3**, e00097-18 (2018).

5. Arora, J. *et al.* Evidence of cospeciation between termites and their gut bacteria on a geological time scale. *Proc. R. Soc. B Biol. Sci.* **290**, 20230619 (2023).

6. Sanders, J. G. *et al.* Widespread extinctions of co-diversified primate gut bacterial symbionts from humans. *Nat. Microbiol.* **8**, 1039–1050 (2023).

7. Moeller, A. H. *et al.* Cospeciation of gut microbiota with hominids. *Science* **353**, 380–382 (2016).

8. Rühlemann, M. C. *et al.* Functional host-specific adaptation of the intestinal microbiome in hominids. *Nat. Commun.* **15**, 326 (2024).

9. Good, B. H. Limited codiversification of the gut microbiota with humans. 2022.10.27.514143 Preprint at https://doi.org/10.1101/2022.10.27.514143 (2023).

10. Suzuki, T. A. *et al.* Codiversification of gut microbiota with humans. *Science* **377**, 1328–1332 (2022).

11. Song, S. J. *et al.* Comparative Analyses of Vertebrate Gut Microbiomes Reveal Convergence between Birds and Bats. *mBio* **11**, 10.1128/mbio.02901-19 (2020).

12. Groussin, M. *et al.* Unraveling the processes shaping mammalian gut microbiomes over evolutionary time. *Nat. Commun.* **8**, 14319 (2017).

13. Mazel, F., Guisan, A. & Parfrey, L. W. Transmission mode and dispersal traits correlate with host specificity in mammalian gut microbes. *Mol. Ecol.* **33**, e16862 (2024).

14. Raulo, A. *et al.* Social and environmental transmission spread different sets of gut microbes in wild mice. *Nat. Ecol. Evol.* **8**, 972–985 (2024).

15. Nayfach, S. & Pollard, K. S. Toward Accurate and Quantitative Comparative Metagenomics. *Cell* **166**, 1103–1116 (2016).

16. Kwong, W. K. *et al.* Dynamic microbiome evolution in social bees. *Sci. Adv.* **3**, e1600513 (2017).

17. Niode, N. J., Adji, A., Rimbing, J., Tulung, M. & Tallei, T. E. Composition and diversity of bacteria from giant Asian honeybee Apis dorsata gut. *Biodiversitas J. Biol. Divers.* **22**, (2021).

18. Ganeshprasad, D. N., Jani, K., Shouche, Y. S. & Sneharani, A. H. *Gut Bacterial Inhabitants of Open Nested Honey Bee, Apis Florea*. https://www.researchsquare.com/article/rs-225332/v1 (2021) doi:10.21203/rs.3.rs-225332/v1.

19. Gruneck, L., Gentekaki, E., Khongphinitbunjong, K. & Popluechai, S. Distinct gut microbiota profiles of Asian honey bee (Apis cerana) foragers. *Arch. Microbiol.* **204**, 187 (2022).

20. Wu, J. *et al.* Honey bee genetics shape the strain-level structure of gut microbiota in social transmission. *Microbiome* **9**, 225 (2021).

21. Gruneck, L., Khongphinitbunjong, K. & Popluechai, S. Gut microbiota associated with two species of domesticated honey bees from Thailand. *Symbiosis* **83**, 335–345 (2021).

22. Khan, K. A. *et al.* Gut microbial diversity in Apis cerana indica and Apis florea colonies: a comparative study. *Front. Vet. Sci.* **10**, (2023).

23. Ge, Y., Jing, Z., Diao, Q., He, J.-Z. & Liu, Y.-J. Host Species and Geography Differentiate Honeybee Gut Bacterial Communities by Changing the Relative Contribution of Community Assembly Processes. *mBio* **12**, e00751-21 (2021).

24. Ellegaard, K. M. *et al.* Genomic changes underlying host specialization in the bee gut symbiont *Lactobacillus* Firm5. *Mol. Ecol.* **28**, 2224–2237 (2019).

25. Sarton-Lohéac, G. *et al.* Deep Divergence and Genomic Diversification of Gut Symbionts of Neotropical Stingless Bees. *mBio* **0**, e03538-22 (2023).

26. Ellegaard, K. M. *et al.* Extensive intra-phylotype diversity in lactobacilli and bifidobacteria from the honeybee gut. *BMC Genomics* **16**, 284 (2015).

27. Li, Y., Leonard, S. P., Powell, J. E. & Moran, N. A. Species divergence in gut-restricted bacteria of social bees. *Proc. Natl. Acad. Sci.* **119**, e2115013119 (2022).

28. Zhang, W. *et al.* Genomic features underlying the evolutionary transitions of *Apibacter* to honey bee gut symbionts. *Insect Sci.* **29**, 259–275 (2022).

29. Ellegaard, K. M. & Engel, P. Genomic diversity landscape of the honey bee gut microbiota. *Nat. Commun.* **10**, 446 (2019).

30. Engel, P., Martinson, V. G. & Moran, N. A. Functional diversity within the simple gut microbiota of the honey bee. *Proc. Natl. Acad. Sci.* **109**, 11002–11007 (2012).

31. Regan, T. *et al.* Characterisation of the British honey bee metagenome. *Nat. Commun.* **9**, 4995 (2018).

32. Ellegaard, K. M., Suenami, S., Miyazaki, R. & Engel, P. Vast Differences in Strain-Level Diversity in the Gut Microbiota of Two Closely Related Honey Bee Species. *Curr. Biol.* **30**, 2520-2531.e7 (2020).

33. Wu, Y. *et al.* Genetic divergence and functional convergence of gut bacteria between the Eastern honey bee Apis cerana and the Western honey bee Apis mellifera. *J. Adv. Res.* **37**, 19–31 (2022).

34. Rohde, K. & Rohde, P. P. How to measure ecological host specificity. *Vie Milieu* (2008).

35. Olm, M. R. *et al.* inStrain profiles population microdiversity from metagenomic data and sensitively detects shared microbial strains. *Nat. Biotechnol.* **39**, 727–736 (2021).

36. Vera-Ponce de León, A., Jahnes, B. C., Duan, J., Camuy-Vélez, L. A. & Sabree, Z. L. Cultivable, Host-Specific Bacteroidetes Symbionts Exhibit Diverse Polysaccharolytic Strategies. *Appl. Environ. Microbiol.* **86**, e00091-20 (2020).

37. Bilen, M. *et al.* Dysgonomonas massiliensis sp. nov., a new species isolated from the human gut and its taxonogenomic description. *Antonie Van Leeuwenhoek* **112**, 935–945 (2019).

38. Hommola, K., Smith, J. E., Qiu, Y. & Gilks, W. R. A Permutation Test of Host–Parasite Cospeciation. *Mol. Biol. Evol.* **26**, 1457–1468 (2009).

39. Moeller, A. H., Sanders, J. G., Sprockett, D. D. & Landers, A. Assessing co-diversification in host-associated microbiomes. *J. Evol. Biol.* **n/a**, (2023).

40. Zheng, H. *et al.* Division of labor in honey bee gut microbiota for plant polysaccharide digestion. *Proc. Natl. Acad. Sci.* **116**, 25909–25916 (2019).

41. Wardman, J. F., Bains, R. K., Rahfeld, P. & Withers, S. G. Carbohydrate-active enzymes (CAZymes) in the gut microbiome. *Nat. Rev. Microbiol.* **20**, 542–556 (2022).

42. Gile, G. H. Protist symbionts of termites: diversity, distribution, and coevolution. *Biol. Rev.* **99**, 622–652 (2024).

43. Tung, J. *et al.* Social networks predict gut microbiome composition in wild baboons. *eLife* **4**, e05224 (2015).

44. Kearney, S. M. *et al.* Endospores and other lysis-resistant bacteria comprise a widely shared core community within the human microbiota. *ISME J.* **12**, 2403–2416 (2018).

45. Valles-Colomer, M. *et al.* The person-to-person transmission landscape of the gut and oral microbiomes. *Nature* **614**, 125–135 (2023).

46. Moeller, A. H., Suzuki, T. A., Phifer-Rixey, M. & Nachman, M. W. Transmission modes of the mammalian gut microbiota. *Science* **362**, 453–457 (2018).

47. Wilde, J., Slack, E. & Foster, K. R. Host control of the microbiome: Mechanisms, evolution, and disease. *Science* **385**, eadi3338 (2024).

48. Koch, H., Abrol, D. P., Li, J. & Schmid-Hempel, P. Diversity and evolutionary patterns of bacterial gut associates of corbiculate bees. *Mol. Ecol.* **22**, 2028–2044 (2013).

49. Powell, E., Ratnayeke, N. & Moran, N. A. Strain diversity and host specificity in a specialized gut symbiont of honeybees and bumblebees. *Mol. Ecol.* **25**, 4461–4471 (2016).

50. Yang, Y. *et al.* Dysgonomonas macrotermitis sp. nov., isolated from the hindgut of a fungus-growing termite. *Int. J. Syst. Evol. Microbiol.* **64**, 2956–2961 (2014).

51. Dyer, F. C. & Seeley, Th. D. Colony migration in the tropical honey beeApis dorsata F. (Hymenoptera: Apidae). *Insectes Sociaux* **41**, 129–140 (1994).

52. Raffiudin, R. *et al.* The effect of land cover on the foraging behavior and pollen in the honey of the giant bee Apis dorsata in Sumatra. *Front. Bee Sci.* **2**, (2024).

53. Stewart, A. B., Sritongchuay, T., Teartisup, P., Kaewsomboon, S. & Bumrungsri, S. Habitat and landscape factors influence pollinators in a tropical megacity, Bangkok, Thailand. *PeerJ* **6**, e5335 (2018).

54. Hammer, T. J., Dickerson, J. C. & Fierer, N. Evidence-based recommendations on storing and handling specimens for analyses of insect microbiota. *PeerJ* **3**, e1190 (2015).

55. Kešnerová, L. *et al.* Disentangling metabolic functions of bacteria in the honey bee gut. *PLOS Biol.* **15**, e2003467 (2017).

56. Olm, M. R., Brown, C. T., Brooks, B. & Banfield, J. F. dRep: a tool for fast and accurate genomic comparisons that enables improved genome recovery from metagenomes through de-replication. *ISME J.* **11**, 2864–2868 (2017).

57. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* **25**, 1754–1760 (2009).

58. Ruscheweyh, H.-J. *et al.* Cultivation-independent genomes greatly expand taxonomic-profiling capabilities of mOTUs across various environments. *Microbiome* **10**, 212 (2022).

59. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

60. Nurk, S., Meleshko, D., Korobeynikov, A. & Pevzner, P. A. metaSPAdes: a new versatile metagenomic assembler. *Genome Res.* **27**, 824–834 (2017).

61. Kang, D. D. *et al.* MetaBAT 2: an adaptive binning algorithm for robust and efficient genome reconstruction from metagenome assemblies. *PeerJ* **7**, e7359 (2019).

62. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res.* **25**, 1043–1055 (2015).

63. Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database. *Bioinformatics* **36**, 1925–1927 (2020).

64. Menzel, P., Ng, K. L. & Krogh, A. Fast and sensitive taxonomic classification for metagenomics with Kaiju. *Nat. Commun.* **7**, 11257 (2016).

65. Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2. *Genome Biol.* **20**, 257 (2019).

66. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**, 357–359 (2012).

67. Ranwez, V., Harispe, S., Delsuc, F. & Douzery, E. J. P. MACSE: Multiple Alignment of Coding SEquences Accounting for Frameshifts and Stop Codons. *PLOS ONE* **6**, e22594 (2011).

68. Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol. Biol. Evol.* **32**, 268–274 (2015).

69. Ochman, H., Elwyn, S. & Moran, N. A. Calibrating bacterial evolution. *Proc. Natl. Acad. Sci.* **96**, 12638–12643 (1999).

70. Su, Y.-C., Chiu, Y.-F., Warrit, N., Otis, G. W. & Smith, D. R. Phylogeography and species delimitation of the Asian cavity-nesting honeybees. *Insect Syst. Divers.* **7**, 5 (2023).

71. Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome Biol.* **20**, 238 (2019).

72. Katoh, K., Misawa, K., Kuma, K. & Miyata, T. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res.* **30**, 3059–3066 (2002).

73. Buck, M., Mehrshad, M. & Bertilsson, S. mOTUpan: a robust Bayesian approach to leverage metagenome-assembled genomes for core-genome estimation. *NAR Genomics Bioinforma.* **4**, lqac060 (2022).

74. Carr, S. M. Multiple mitogenomes indicate Things Fall Apart with Out of Africa or Asia hypotheses for the phylogeographic evolution of Honey Bees (Apis mellifera). *Sci. Rep.* **13**, 9386 (2023).

75. The scikit-bio development team. scikit-bio: A Bioinformatics Library for Data Scientists, Students, and Developers. (2020).

76. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* **11**, 119 (2010).

77. Pronk, L. J. U. & Medema, M. H. Whokaryote: distinguishing eukaryotic and prokaryotic contigs in metagenomes based on gene structure. *Microb. Genomics* **8**, 000823 (2022).

78. Shaffer, M. *et al.* DRAM for distilling microbial metabolism to automate the curation of microbiome function. *Nucleic Acids Res.* **48**, 8883–8900 (2020).

79. Ducarmon, Q. R. *et al.* Large-scale computational analyses of gut microbial CAZyme repertoires enabled by Cayman. 2024.01.08.574624 Preprint at https://doi.org/10.1101/2024.01.08.574624 (2024).

80. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841–842 (2010).

81. Baselga, A. & Orme, C. D. L. betapart: an R package for the study of beta diversity. *Methods Ecol. Evol.* **3**, 808–812 (2012).

82. Kotthoff, U., Wappler, T. & Engel, M. Miocene honey bees from the Randeck Maar of southwestern Germany (Hymenoptera, Apidae). *ZooKeys* **96**, 11–37 (2011).

83. Ghonche-Golan, S., Nazemi-Rafie, J. & Rezapanah, M. The relationship study among Apis spp. using mitochondrial markers, Procrustes coordinates and residuals of geometric morphometric method. *Biologia (Bratisl.)* **77**, 2519–2530 (2022).