Vaginal microbiota as a predictor of preterm birth: an observational cohort study in a high-risk population

Authors

Laura Lesimple[1,2], Jessica Rousseau[3], Céline Plainvert[1,4], Luce Landraud[5,6], Nathalie Grall[6,7], François Goffinet[3,8], Pierre-Yves Ancel[3,9], Christophe Pannetier[2], Claire Poyart[1,4], Laurent Mandelbrot[6,10]*, Asmaa Tazi[1,4]*, The Inspire Consortium

Affiliations

[1] Université Paris Cité, Inserm U1016, CNRS UMR 8104, Institut Cochin, Team Bacterial Pathogenesis and Innate Immune Signaling, Fédération Hospitalo-Universitaire Prem'Impact, Paris, France.

[2] BforCure, Montreuil, France.

[3] Université Paris Cité, CRESS, Obstetrical Perinatal and Pediatric Epidemiology Research Team, EPOPE, French Institute for Medical Research and Health INSERM, INRAE, Paris, France.

[4] Department of Bacteriology, National Reference Center for Streptococci, University Hospitals Assistance Publique - Hôpitaux de Paris, Hôpital Cochin, Fédération Hospitalo-Universitaire Prem'Impact, Paris, France.

[5] Department of Microbiology, University Hospitals Assistance Publique - Hôpitaux de Paris, Hôpital Louis Mourier, Colombes, France.

[6] Université Paris Cité, IAME INSERM U1137, Paris, France.

[7] Department of Bacteriology, University Hospitals Assistance Publique - Hôpitaux de Paris, Hôpital Bichat, Paris, France.

[8] Department of Obstetrics and Gynecology, Port-Royal Maternity, University Hospitals Paris

Centre Cochin Port Royal, AP-HP, Paris, France.

[9] URC-CIC P1419, University Hospitals Paris Centre Cochin Port Royal, AP-HP, Paris,

France.

[10] Department of Obstetrics and Gynecology, University Hospitals Assistance Publique -

Hôpitaux de Paris, Hôpital Louis Mourier, Colombes, France.

* Laurent Mandelbrot and Asmaa Tazi contributed equally

Corresponding author

Pr Asmaa Tazi

HUPC Cochin, Service de Bactériologie, 27 rue du Faubourg St Jacques, 75014, Paris,

France.

E-mail: asmaa.tazi@aphp.fr

Phone: +33 1 58 41 34 14 - Fax: +33 1 58 41 15 48

Short title: Vaginal microbiota signatures of preterm birth

The authors report no conflict of interest.

At a glance

Why was this study conducted?

Despite the growing evidence of an association between vaginal microbiota composition and

preterm birth, data remain elusive, because most studies were relatively small to examine

cofounding factors. Moreover, longitudinal studies focusing on high-risk populations are

lacking to transpose microbiological data into a clinical perspective.

Key findings

In a prospective cohort of over 1,800 pregnant women, using microbiological cultures of

vaginal samples at admission and multivariable analyses adjusting for demographic and

obstetrical factors, we identified lactobacilli depletion and enterobacteria as strong vaginal

microbiota signatures of preterm labor and preterm rupture of membranes as compared to

controls. Importantly, lactobacilli depletion constituted an independent risk factor for

spontaneous preterm birth.

What does this add to what is known?

We confirm the implication of vaginal microbiota in preterm birth and highlight the

importance of lactobacilli in this outcome. Accurate characterization of vaginal microbiota at

admission for preterm labor as well as preterm premature rupture of membranes could

improve the management of these complications of pregnancy.

Abstract

Background: Increasing evidence link vaginal microbiota composition and preterm birth. However, the impact of the vaginal microbiota on preterm labor and premature rupture of membranes and on their outcome remain elusive, mainly because most studies are relatively small-sized.

Objectives: To identify microbiota signatures of preterm labor, premature rupture of membranes, and spontaneous preterm birth.

Study Design: We conducted an observational, prospective, longitudinal cohort study from August 2018 until June 2023 in 3 maternity wards from university hospitals in the Paris, France, area. Women with singleton pregnancies were categorized in 3 groups including a control group, and 2 groups of increasing risk of preterm birth: preterm labor and preterm premature rupture of membranes. Demographic, clinical data, past medical and obstetrical history and pregnancy outcome were collected. Vaginal swabs were collected at admission and were analyzed using microbiological cultures. The association between bacterial species, the cohort groups, and spontaneous preterm birth was studied in univariate analyses. Adjusted odds ratio (aOR) and 95% confidence intervals (95% CI) were calculated in multivariable analyses.

Results: A total of 1,848 women were included, of whom 1,048 (56.7%) in the control group, 417 (22.6%) with preterm labor, and 383 (20.7%) with premature rupture of membranes. Together with demographic features such as ethnicity and obstetrical history, several vaginal microbiota signatures were identified in high-risk groups. In multivariable analysis, preterm labor was correlated with lactobacilli depletion (aOR 1.49, 95% CI 1.07-2.07), enterobacteria (aOR 1.91, 95% CI 1.40-2.62) and *Gardnerella vaginalis* (aOR 4.50, 95% CI 1.80-13.02), and premature rupture of membranes with lactobacilli depletion (aOR 2.10, 95% CI 1.52-

2.89) and enterobacteria (aOR 2.39, 95% CI 1.75-3.26). Finally, lactobacilli depletion was a

risk factor for spontaneous preterm birth (aOR 1.68, 95% CI 1.07-2.68).

Conclusions: In this large cohort, *G. vaginalis* was associated with preterm labor, while

lactobacilli depletion and enterobacteria were risk factors for preterm labor and premature

rupture of membranes. In addition, lactobacilli depletion was correlated with spontaneous

preterm birth. Further studies are needed to address the benefits of adapted antibiotics and

probiotics to restore a healthy microbiota for preventing preterm birth.

Abstract word count: 342

Manuscript word count: 2970

Key words : Vaginal microbiota, dysbiosis, pregnancy outcomes, preterm birth, premature

preterm rupture of membrane.

Introduction

Preterm birth (PTB), defined as birth before 37 weeks of gestation, is a frequent and

severe adverse pregnancy outcome which concerns approx. 7% pregnancies in industrialized

countries and accounts for 14 millions annual births worldwide.[1] As the leading causes of

death in children under the age of 5 years, PTB and its sequelae represent one of the most

prevalent conditions in the Global Burden of Disease analysis and are therefore considered as

a major worldwide health issue.[2,3] Up to 40% of spontaneous PTB (sPTB) are preceded by

preterm premature rupture of membranes (PPROM), increasing the risk of intrauterine

infection and *in utero* demise,[4,5] and disruption of the vaginal microbiota is increasingly

considered to be a contributing factor.[6,7]

The composition of the vaginal microbiota is highly dynamic, changing with age,

hormonal fluctuations, gestational status, ethnicity, sexual behavior, and lifestyle.[8–10]

Microbiota studies based on sequencing of vaginal samples from reproductive-age women

have identified five types of vaginal community structures designated community state types

(CSTs).[10] Four of these are dominated by one *Lactobacillus* species while the last one,

designated CST-IV, is characterized by a more diverse microbiota dominated by anaerobic

bacteria such as *Gardnerella vaginalis* and *Prevotella* spp. Although the distribution of these

CSTs largely varies according to ethnicity,[11] it is generally believed that a healthy vaginal

microbiota is dominated by lactobacilli with a low bacterial diversity.[12,13] Conversely, a

decrease in lactobacilli abundance is correlated to an increase of microbiota diversity

frequently referred to as vaginal dysbiosis and typically exemplified by the CST-IV. In the

context of pregnancy, increasing evidence links preterm labor (PTL), PPROM, and PTB with

low-lactobacilli microbiota characterized by a high diversity and an enrichment in

G. vaginalis and various anaerobes, whereas microbiota dominated by lactobacilli and more

specifically by *Lactobacillus crispatus* (CST-I) have been found associated with term

birth.[6,14–16]

Despite growing evidence of an association between vaginal microbiota and

pregnancy outcomes, incorporating these observations into routine patient management

remains a challenge. Studies that have attempted to address using conventional

microbiological methods whether pregnancy outcomes could be correlated with specific

microbial signatures other than vaginal dysbiosis are lacking.[17] In addition, the impact of the

vaginal microbiota on PTL and PPROM and on their outcome remain elusive, mainly because

most studies are relatively small-sized.[6]

Here, our goal was to use conventional microbial techniques to identify microbiota

signatures of PTL and PPROM along with microbiological risk factors for sPTB in high-risk

women hospitalized for PTL and PPROM.

Materials and methods

Study design

We conducted an observational, prospective, longitudinal cohort study from August

2018 until June 2023 in 3 maternity wards, of which 2 of level 3 and 1 of level 2b, from

university hospitals in the Paris, France, area. The total annual number of births at these 3

centers was approximately 11,000. Women were categorized in 3 groups including a control

group and 2 groups at high risk of PTB, referred to as the PTL and PPROM groups. The

eligibility criteria for maternal participation in the study included all pregnant women aged

>18 years who responded to the following inclusion criteria: *i)* PTL: hospitalization for

threatened PTB with intact membranes; and *ii)* PPROM: rupture of membranes before 37

weeks of gestation and before the onset of labor contractions. The control group consisted of

pregnant women who delivered at term with no history of PTL and no premature rupture of

membranes. These women were enrolled at the time of *Streptococcus agalactiae* (Group B

Streptococcus, GBS) antenatal screening between 34 and 38 weeks of gestation, as

recommended by the French guidelines for the prevention of perinatal infectious disease.[18]

Women who did not speak or understand French and multiple pregnancies were excluded.

Demographic, clinical data, past medical and obstetrical history, and pregnancy

outcome were collected. Maternal place of birth was collected as a reflect of ethnicity, given

that reporting ethnicity itself is forbidden in France.

Sampling and laboratory procedures

Vaginal swabs were sampled at admission at the hospital or at GBS antenatal

screening and sent to the laboratory in Amies transport medium (ESwab™ system, Copan

Diagnostics, Italy). Specimens were immediately processed for analysis by conventional

culture methods using non-selective rich media incubated at 37°C in aerobic atmosphere with

5% CO_2. In addition, a Columbia agar with 5% horse blood, nalidixic acid and colistin

(bioMérieux, Marcy l'Etoile, France) for the selective isolation of anaerobes and Gram-

positive cocci and a Granada agar (Fisher Scientific, Illkirch, France) for the selective

isolation of GBS were incubated at 37°C under anaerobic atmosphere. All plates were

examined after 16h and 44h of incubation.

All different morphotypes of bacterial and yeast colonies were identified using matrix

assisted laser desorption ionization - time of flight (MALDI-TOF) mass spectrometry (Bruker

Daltonics, Billerica, Massachussetts, USA). The density of vaginal colonization was

estimated semi-quantitatively as follows: colonies on a single quadrant, low; colonies on more

than two quadrants, high.

In addition, the overall vaginal microbiota composition was assessed by direct

microscopic examination of the vaginal samples after Gram staining and determination of the

Nugent score in 2 out of 3 maternity wards according to routine practices.[19]

Primary outcomes and studied factors

The primary outcomes were the microbiological signatures of PTL and PPROM and

the microbiological risk factors for sPTB defined as spontaneous onset of labor before 37

weeks of gestation. The studied factors were the microbial species isolated in the vaginal

samples at admission or antenatal GBS screening. A specific attention was paid to bacteria of

the *Lactobacillus* genus, and to species potentially associated with adverse outcomes

including enterobacteria (*Escherichia coli*, *Klebsiella pneumoniae*, and others),

Staphylococcus aureus, GBS, *Streptococcus anginosus*, *G. vaginalis*, strict anaerobes

(*Prevotella* spp., *Bacteroides* spp., and others), and fungi. Unless otherwise specified, only

the presence or absence of the bacterial species, not their quantity, was considered in the

statistical analyses.

Microbiological data were adjusted to known sPTB risk factors and to confounding

variables such as demographics (country of birth), body mass index (BMI), gravidity, parity

and pregnancy-related risk factors (history of preterm delivery, gestational diabetes,

hypertensive disease, and tobacco use).

Statistical analyses

197   All data were collected in a computerized database and analyzed using the GraphPad

198 Software v8.3.0 (Dotmatics, Boston, Massachusetts, USA) and RStudio v1.1.447 (Posit,

199 Boston, Massachusetts, USA). Firstly, we assessed the prevalence of the studied factors

according to the cohort groups as a proportion of the total number of participants in each group. Mann-Whitney tests were used when appropriate and categorical data were compared using chi-squared or Fisher's exact tests. Odds ratios (OR) and 95% confidence intervals (95% CI) were calculated in univariate analysis. Secondly, to study the microbiological signatures of PTL and PPROM and the microbiological risk factors for sPTB, multivariable logistic regression models were used for prognostic covariates. Microbiological signatures of PTL and PPROM were determined using the control group as a reference. Microbiological risk factors for sPTB were determined using women in the PTL and PPROM groups who delivered at term as the reference. Adjusted OR (aOR) were calculated using known sPTB risk factors and confounding variables in multivariable analyses. Two models were used to examine enterobacteria, with either enterobacteria as a single variable or the two most common species e.g., *E. coli* and *K. pneumoniae*, as independent variables. A *P* value < 0.05 was considered significant. Missing data were excluded.

Ethics declaration

The reported study was approved by the French ethics committee on November, 2017 (identifier 2017 – A02755-48; NCT03371056). All individuals enrolled in the cohort provided signed informed consent that included authorization for the collection of medical information and biological samples. All collected data were anonymized and integrated in a secured electronic database.

Results

Study population

A total of 1,848 women were included, of whom 1,048 (56.7%) in the control group, 417 (22.6%) in the PTL group, and 383 (20.7%) in the PPROM group. The characteristics of

the study population are summarized in Table 1. Compared to controls, PTL and PPROM

were characterized by a higher proportion of women born in Sub Saharan Africa, history of

preterm deliveries, and obesity. Eventually, 283 (57.6%) and 58 (13.4%) women with PTL

and PPROM delivered at 37 weeks of gestation or more, respectively.

Vaginal microbiota signatures of PTL and PPROM

A Nugent score was determined for 1,435 (77.7%) vaginal samples and indicated that

bacterial vaginosis was uncommon, below 5% whatever the cohort group considered (Table

2). Nevertheless, the distribution of the Nugent scores varied significantly between PTL,

PPROM and controls, Nugent scores $\geq$ 7 being more frequent in PTL (4.8% vs. 1.9%,

P=0.008) and intermediate scores being overrepresented in PPROM (18.7% *vs*. 12.8% in

controls, P=0.01). Accordingly, microbial cultures of the vaginal samples revealed significant

differences between the cohort groups, most importantly in the presence and abundance of

lactobacilli, enterobacteria, and anaerobes, including *G. vaginalis* (Table 2). Conversely, the

presence and abundance of *S. aureus*, GBS and *Candida* spp. did not vary according to the

cohort groups.

Compared to controls, microbial signatures of PTL included a higher proportion of

samples negative for lactobacilli and positive for enterobacteria and anaerobes (Figure 1).

Enterobacteria were more frequent and also more abundant (Table 2), as assessed by the

number of samples with numerous bacteria compared to the total number of positive sample

(68.5% in PTL *vs*. 45.8% in controls, P=0.0001).

Microbial signatures of PPROM included a higher proportion of samples negative for

lactobacilli and positive for enterobacteria but not for anaerobes (Figure 1). Significant

differences were also observed considering their abundance (Table 2). Lactobacilli were more

frequently in low quantities (14.5% in PPROM *vs*. 3.8% in controls, *P*<0.0001) whereas

enterobacteria were more abundant (62.8% in PPROM *vs*. 45.8% in controls, *P*=0.004).

Microbial risk factors for PTL and PPROM

Demographical and clinical characteristics such as ethnicity and BMI were associated

with pregnancy outcomes. Since they are also presumed to impact microbiota composition,

we tested their association with vaginal microbiota composition in our study population and

confirmed their correlation with the Nugent score and with the presence and abundance of

lactobacilli and enterobacteria (Supplementary Tables 1 and 2). The place of birth but not the

BMI also correlated with anaerobes and *G. vaginalis* (Supplementary Table 1).

Next, to identify the microbiological factors associated with PTL and PPROM, we

performed univariate (Table 3) and then multivariable logistic regression analyses adjusting

for ethnicity and BMI into account, showing that PTL was strongly associated with maternal

origin from French overseas territories and Sub Saharan Africa, with a history of preterm

delivery, and also with the absence of lactobacilli (aOR: 1.49, 95% CI 1.07-2.07), the

presence of enterobacteria (aOR: 1.91, 95% CI 1.40-2.62) and of *G. vaginalis* (aOR: 4.50,

95% CI 1.80-13.02) (Figure 2). For PPROM, the strongest risk factors were again maternal

origin from French overseas territories and Sub Saharan Africa and also vaginal microbiota

depleted in lactobacilli (aOR 2.10, 95% CI 1.52-2.89) and including enterobacteria (aOR

2.39, 95% CI 1.77-3.26) (Figure 2).

Overall, the absence of lactobacilli and the presence of enterobacteria in vaginal swab

cultures were both correlated with PTL and PPROM, whereas *G. vaginalis* was only a

signature of PTL. Similar results were obtained with the second model of multivariable

logistic regression including *E. coli* and *K. pneumoniae* instead of enterobacteria

(Supplementary Figure 1), showing that vaginal colonization by *E. coli* was strongly

associated with PTL (aOR 2.08, 95% CI 1.46-2.96) and PPROM (aOR 2.35, 95% CI 1.63-3.33).

Vaginal microbiota risk factors for spontaneous preterm birth

Finally, we sought to determine whether the vaginal microbiota at admission for PTL or PPROM could be predictive of sPTB, which affected 41.0% (119/290) and 95.4% (209/219) pregnancies in each group, respectively. In univariate analysis, we found that in addition to expected clinical factors such as gestational diabetes, the absence of lactobacilli and the presence of enterobacteria were associated with a higher risk of spontaneous preterm delivery (Table 4). In multivariable analysis, the absence of lactobacilli was the only microbiological risk factor for sPTB with an aOR of 1.68, 95% CI 1.07-2.68 (Figure 3).

Comment

Principal Findings

In this prospective cohort study of over 1,800 pregnant women, we identified using simple microbiological cultures of vaginal samples distinct microbiological signatures of PTL and PPROM. PTL was strongly associated with the presence of *G. vaginalis* and less significantly with the absence of lactobacilli, whereas PPROM was correlated with a depletion in lactobacilli and with the presence of enterobacteria. Importantly, the composition of microbiota on admission, particularly a depletion in lactobacilli, was predictive of sPTB.

Results in the Context of What is Known

Our results support previous findings regarding the importance of a healthy lactobacilli-rich microbiota for pregnancy outcome. Reduced *Lactobacillus* spp. abundance and increased vaginal bacterial diversity were identified as risk factors for PPROM in a study

that analyzed the vaginal microbiome of 86 women.[21] A meta-analysis conducted by Gudnadottir *et al.* in which the vaginal microbiome was assessed before the onset of labor reported an increased risk of preterm delivery in women with low-lactobacilli microbiota.[15] Lastly, in a recent longitudinal cohort study of 175 pregnant women comprising 40 PTB, the vaginal microbiome of pregnancies resulting in PTB was found to be genetically more diverse, a diversity which was driven by *Gardnerella* spp.[20] In our study, enterobacteria but not *G. vaginalis* were identified as a risk factor for PPROM. This discrepancy is likely related to the methodology used, as conventional culture techniques are less sensitive than metagenomics for the detection of anaerobes.

Clinical Implications

The role of bacterial vaginosis and *G. vaginalis* in PTL and PTB has long been suspected.[22,23] Increasing evidence now supports a direct role for *G. vaginalis* in the pathophysiology of PTL is now accumulating. *G. vaginalis* and other anaerobes such as *Prevotella* spp. are capable of degrading cervical mucus, promoting ascending intrauterine infections, local inflammation, uterine motility, and PTL.[24,25] It is now generally accepted that an antibiotic treatment, mainly based on metronidazole or clindamycin, should be given to women presenting with PTL and diagnosed with symptomatic bacterial vaginosis. In many countries, national guidelines also recommend screening for bacterial vaginosis and treatment of women with a history of PTB, although this strategy has not been shown to be effective in reducing the risk of PTB.[26–29] Here we show that *G. vaginalis* is a risk factor for PTL but more importantly, it is the absence of lactobacilli that appears to be predictive of sPTB. These observations, in line with numerous metagenomic studies highlighting the protective role of lactobacilli and in particular *L. crispatus*, *L. gasseri*, and *L. jensenii*,[16,30,31] argue for a comprehensive management of bacterial vaginosis in pregnancy aimed at restoring a healthy

microbiota. Such management would likely combine antibiotics targeting anaerobes with

probiotics, as proposed for recurrent vaginosis.[28]

Our results also raise questions about the management of vaginal colonization by

enterobacteria during PTL and PPROM. We identified *E. coli*, and more generally

enterobacteria, as strongly associated with PTL and PPROM in univariate and multivariable

analyses and also, although only in univariate analysis, for sPTB. However, in contrast to

GBS, there are currently no consensus guidelines for the screening and the administration of

antibiotics against enterobacteria during pregnancy, even in the case of PPROM. In the

absence of evidence of infection, strategies vary widely between countries, ranging from no

treatment to broad-spectrum antibiotics such as 3[rd] generation cephalosporins.[32] Our

observations also support the administration of prophylaxis tailored to the pathogenic species

present in the vaginal microbiota at admission for PTL and PPROM, although further studies

are needed to determine the benefits of such a strategy.

Research Implications

Interventional studies to assess the efficacy and tolerability of antibiotics alone or in

combination with probiotics to reduce the risk of PTB and PPROM in high-risk populations

are still lacking. Randomized control trials should take into account that a particular probiotic

strain may not be protective for all women according to their CST. The benefit of

personalized targeted antibiotic prophylaxis directed against the specific pathogenic species

e.g., GBS, enterobacteria, *Candida* spp. and others, found in the vaginal flora in reducing the

risk of PTB, especially in the case of PPROM, should also be investigated. There is also an

important need to investigate the association between the vaginal microbiota and the risk of

early-onset neonatal sepsis.[33]

Strengths and Limitations

This is a large cohort of over 1,800 patients, including over 400 PTL and nearly 400

PPROM along with low-risk controls. The sample size allowed adjustment for important

confounders of the outcome measures, including demographic factors such as ethnicity, BMI,

and history of PTB. The conventional culture approach is widely available in microbiology

laboratories and thus applicable to routine clinical settings, in contrast to metagenomics.

Hence, we were able to demonstrate that the vaginal microbiota is an independent risk factor

for PTL, and PPROM, as well as for sPTB. However, there are limitations that need to be

considered when interpreting our results. First, the microbiota analyses were performed using

simple vaginal swab cultures. This leads to three main issues: (i) fastidious species such as

Atopobium spp., or Mollicutes could not be studied; (ii) since culture is less sensitive than

metagenomics, the reported species were most likely the dominant ones and minor species

were overlooked; (iii) quantitative analysis could not be performed and ultimately only the

presence or absence of major species was studied. The second major limitation is the lack of

identification of lactobacilli at the species level. Consequently, the vaginal microbiotas were

not categorized according to CSTs, and the protective role of the different *Lactobacillus*

species could not be addressed.

Conclusions

In this large multicentric cohort study focusing on pregnancies at high risk of PTB, we

used bacterial culture to highlight novel distinct vaginal microbiota signatures of adverse

pregnancy outcomes. We identified *G. vaginalis* as the strongest risk factor for PTL,

enterobacteria as a risk factor for PPROM, and lactobacilli depletion as a risk factor for PTL,

PPROM, and sPTB. Further studies are needed to address the potential of interventions aimed

at restoring a healthy microbiota in reducing these outcomes.

Transparency declaration

Ethics statement

The reported study was reviewed by the French ethics committee who granted approval on November, 2017 (identifier 2017 – A02755-48; NCT03371056). All individuals enrolled in the cohort provided signed informed consent that included authorization for the collection of medical information and biological samples. All collected data were anonymized and integrated in a secured electronic database.

Conflicts of interest

The authors have no conflicts of interest to disclose.

Funding

This work was supported by the Programme d'Investissements d'avenir and bpifrance (Structuring R&D Project for Competitiveness—PSPC): # DOS0053477 SUB et DOS0053473 AR). L.L. is a doctoral fellow supported by the Association Nationale de la Recherche et Technologie (Cifre # 2019 / 1510). The funders had no role in the design and conduct of the study.

Acknowledgements

We thank all the contributors to this study, collectively called the InSPIRE Consortium. The full list can be found in the accompanying Supplementary Information.

Part of the results were previously presented in April 2023 at the ECCMID scientific meeting (Copenhagen, Denmark), in September 2023 at the ESPBC congress (Haarlem, The

Netherlands), and in January 2025 at the SMFM 2025 pregnancy meeting (Denver, Colorado,

USA).

Access to data

All authors had full access to data. Data will be made available upon requests directed to the

corresponding author.

Authors' contributions

Conceptualization: C.Po., L.M., and A.T. Methodology: J.R., P.-Y.A., L.M., and A.T. Formal

analysis: L.Le., J.R., P.-Y.A., L.M., and A.T. Investigation: L.Le., J.R., and A.T. Resources:

C.Pl., L.La, N.G., F.G., and L.M. Data curation: L.Le, J.R., C.Pl, L.La, N.G., and A.T.

Writing – Original draft: L.Le and A.T. Writing – Review and Editing: all authors.

Supervision: C.Pa, C.Po., L.M. and A.T. Funding acquisition: C.Pa, C.Po and L.M.

References

1.   Ohuma EO, Moller AB, Bradley E, et al. National, regional, and global estimates of

preterm birth in 2020, with trends from 2010: a systematic analysis. *Lancet Lond Engl.*

2023;402(10409):1261-1271. doi:10.1016/S0140-6736(23)00878-4

2.   Harrison MS, Goldenberg RL. Global burden of prematurity. *Semin Fetal Neonatal*

Med. 2016;21(2):74-79. doi:10.1016/j.siny.2015.12.007

3.   GBD 2016 Causes of Death Collaborators. Global, regional, and national age-sex

specific mortality for 264 causes of death, 1980-2016: a systematic analysis for the Global

Burden   of   Disease   Study   2016.   *Lancet   Lond   Engl.*   2017;390(10100):1151-1210.

doi:10.1016/S0140-6736(17)32152-9

4.   Goldenberg RL, Culhane JF, Iams JD, Romero R. Epidemiology and causes of

preterm birth. *Lancet Lond Engl*. 2008;371(9606):75-84. doi:10.1016/S0140-6736(08)60074-4

5.   Lorthe E. Épidémiologie, facteurs de risque et pronostic de l'enfant. RPC : rupture prématurée des membranes avant terme CNGOF. *Gynécologie Obstétrique Fertil Sénologie*. 2018;46(12):1004-1021. doi:10.1016/j.gofs.2018.10.019

6.   Bayar E, Bennett PR, Chan D, Sykes L, MacIntyre DA. The pregnancy microbiome and preterm birth. *Semin Immunopathol*. 2020;42(4):487-499. doi:10.1007/s00281-020-00817-w

7.   Schuster HJ, Bos AM, Himschoot L, et al. Vaginal microbiota and spontaneous preterm birth in pregnant women at high risk of recurrence. *Heliyon*. 2024;10(10):e30685. doi:10.1016/j.heliyon.2024.e30685

8.   Greenbaum S, Greenbaum G, Moran-Gilad J, Weintraub AY. Ecological dynamics of the vaginal microbiome in relation to health and disease. *Am J Obstet Gynecol*. 2019;220(4):324-335. doi:10.1016/j.ajog.2018.11.1089

9.   Song SD, Acharya KD, Zhu JE, et al. Daily Vaginal Microbiota Fluctuations Associated with Natural Hormonal Cycle, Contraceptives, Diet, and Exercise. Rao K, ed. *mSphere*. 2020;5(4):e00593-20. doi:10.1128/mSphere.00593-20

10.   Ravel J, Gajer P, Abdo Z, et al. Vaginal microbiome of reproductive-age women. *Proc Natl Acad Sci*. 2011;108(supplement_1):4680-4687. doi:bayar

11.   France MT, Ma B, Gajer P, et al. VALENCIA: a nearest centroid classification method for vaginal microbial communities based on composition. *Microbiome*. 2020;8(1):166. doi:10.1186/s40168-020-00934-6

12.   Romero R, Hassan SS, Gajer P, et al. The composition and stability of the vaginal microbiota of normal pregnant women is different from that of non-pregnant women. *Microbiome*. 2014;2(1):4. doi:10.1186/2049-2618-2-4

13. MacIntyre DA, Chandiramani M, Lee YS, et al. The vaginal microbiome during pregnancy and the postpartum period in a European population. *Sci Rep*. 2015;5(1):8988. doi:10.1038/srep08988

14. Fettweis JM, Serrano MG, Brooks JP, et al. The vaginal microbiome and preterm birth. *Nat Med*. 2019;25(6):1012-1021. doi:10.1038/s41591-019-0450-2

15. Gudnadottir U, Debelius JW, Du J, et al. The vaginal microbiome and the risk of preterm birth: a systematic review and network meta-analysis. *Sci Rep*. 2022;12(1):7926. doi:10.1038/s41598-022-12007-9

16. Baud A, Hillion KH, Plainvert C, et al. Microbial diversity in the vaginal microbiota and its link to pregnancy outcomes. *Sci Rep*. 2023;13(1):9061. doi:10.1038/s41598-023-36126-z

17. Gorczyca K, Kozioł MM, Kimber-Trojnar Ż, et al. Premature rupture of membranes and changes in the vaginal microbiome – Probiotics. *Reprod Biol*. 2024;24(3):100899. doi:10.1016/j.repbio.2024.100899

18. HAS. Prévention anténatale du risque infectieux bactérien néonatal précoce. 2001. https://www.has-sante.fr/upload/docs/application/pdf/prevention_antenatale_du_risque_infectieux_bacterien_-_rec.pdf

19. Nugent RP, Krohn MA, Hillier SL. Reliability of diagnosing bacterial vaginosis is improved by a standardized method of gram stain interpretation. *J Clin Microbiol*. 1991;29(2):297-301. doi:10.1128/jcm.29.2.297-301.1991

20. Liao J, Shenhav L, Urban JA, et al. Microdiversity of the vaginal microbiome is associated with preterm birth. *Nat Commun*. 2023;14(1):4997. doi:10.1038/s41467-023-40719-7

21. Brown RG, Al-Memar M, Marchesi JR, et al. Establishment of vaginal microbiota

composition in early pregnancy and its association with subsequent preterm prelabor rupture of the fetal membranes. *Transl Res*. 2019;207:30-43. doi:10.1016/j.trsl.2018.12.005

22.   Hillier SL, Martius J, Krohn M, Kiviat N, Holmes KK, Eschenbach DA. A Case–Control Study of Chorioamnionic Infection and Histologic Chorioamnionitis in Prematurity. *N Engl J Med*. 1988;319(15):972-978. doi:10.1056/NEJM198810133191503

23.   Eschenbach DA, Gravett MG, Chen KC, Hoyme UB, Holmes KK. Bacterial vaginosis during pregnancy. An association with prematurity and postpartum complications. *Scand J Urol Nephrol Suppl*. 1984;86:213-222.

24.   Howe L, Wiggins R, Soothill PW, Millar MR, Horner PJ, Corfield AP. Mucinase and sialidase activity of the vaginal microflora: implications for the pathogenesis of preterm labour. *Int J STD AIDS*. 1999;10(7):442-447. doi:10.1258/0956462991914438

25.   Lewis WG, Robinson LS, Gilbert NM, Perry JC, Lewis AL. Degradation, Foraging, and Depletion of Mucus Sialoglycans by the Vagina-adapted Actinobacterium *Gardnerella vaginalis*. *J Biol Chem*. 2013;288(17):12067-12079. doi:10.1074/jbc.M113.453654

26.   Collège national des gynécologues et obstétriciens français. Prévention de la prématurité spontanée et de ses conséquences. Published online 2016. https://cngof.fr/app/pdf/RPC//RPC%20DU%20CNGOF/2016/RPC_2016_Prmaturit_spontane .pdf?x29325

27.   British Association for Sexual Health and HIV. Bacterial Vaginosis. Published online 2012. https://www.bashh.org/resources/21/bacterial_vaginosis_2012/

28.   Farr A, Swidsinski S, Surbek D, et al. Bacterial Vaginosis: Guideline of the DGGG, OEGGG and SGGG (S2k-Level, AWMF Registry No. 015/028, June 2023). *Geburtshilfe Frauenheilkd*. 2023;83(11):1331-1349. doi:10.1055/a-2169-8539

29.   Subtil D, Brabant G, Tilloy E, et al. Early clindamycin for bacterial vaginosis in pregnancy (PREMEVA): a multicentre, double-blind, randomised controlled trial. *The*

Lancet. 2018;392(10160):2171-2179. doi:10.1016/S0140-6736(18)31617-9

30.   Tabatabaei N, Eren A, Barreiro L, et al. Vaginal microbiome in early pregnancy and

subsequent risk of spontaneous preterm birth: a case–control study. *BJOG Int J Obstet*

Gynaecol. 2019;126(3):349-358. doi:10.1111/1471-0528.15299

31.   Callahan BJ, DiGiulio DB, Goltsman DSA, et al. Replication and refinement of a

vaginal microbial signature of preterm birth in two racially distinct cohorts of US women.

Proc Natl Acad Sci. 2017;114(37):9966-9971. doi:10.1073/pnas.1705899114

32.   Werter DE, Dehaene I, Gurney L, Vargas Buján M, Kazemier BM. Differences in

clinical practice regarding screening and treatment of infections associated with spontaneous

preterm birth: An international survey. *Eur J Obstet Gynecol Reprod Biol*. 2021;266:83-88.

doi:10.1016/j.ejogrb.2021.09.009

33.   Ben M'Barek I, Landraud L, Desfrere L, et al. Contribution of vaginal culture to

predict early onset neonatal infection in preterm prelabor rupture of membranes. *Eur J Obstet*

Gynecol Reprod Biol. 2021;261:78-84. doi:10.1016/j.ejogrb.2021.04.016

Figure legends.

Figure 1. Vaginal microbiota composition at admission for preterm labor (PTL) and

preterm rupture of membranes (PPROM).

Vaginal microbiota composition was characterized by vaginal swab cultures and compared

between controls (n=1,048) and women admitted for PTL (n=417) or PPROM (n=383). The

percentage of vaginal samples positive for selected bacterial group or species are represented.

Error bars indicate the upper limit of the 95% confidence interval. ** $p<0.01$; *** $p<0.001$,

ns: not significant.

Figure 2. Multivariable analysis of risk factors for preterm labor (PTL) and preterm

premature rupture of membranes (PPROM).

Error bars indicate the upper and lower limit of the 95% confidence interval (CI); aOR:

adjusted odds ratio.

Figure 3. Multivariable analysis of risk factors for spontaneous preterm birth.

Error bars indicate the upper and lower limit of the 95% confidence interval (CI); aOR:

adjusted odds ratio.

Table 1. Cohort demographic and clinical characteristics.

	Control N=1,048	PTL N=417	PPROM N=383	*P* value
Maternity ward, n (%)				NT
Center 1	677 (64.6)	114 (27.3)	192 (50.1)	
Center 2	371 (35.4)	203 (48.7)	137 (35.8)	
Center 3	0 (0)	100 (24.0)	54 (14.1)	
Age, median (IQR), years	33 (30-36)	32 (29-35)	33 (30-36)	<0.0001
Gestational age at inclusion, median (IQR), weeks	36 (35-36)	29 (25-32)	32 (27-35)	<0.0001
Maternal place of birth, n (%)				<0.0001
France	556 (53.2)	147 (35.9)	116 (30.7)	
France, overseas territories[a]	29 (2.8)	25 (6.1)	24 (6.3)	
Europe, other countries	79 (7.6)	22 (5.4)	20 (5.3)	
North Africa	199 (19.0)	84 (20.5)	106 (28.0)	
Sub-Saharan Africa	89 (8.5)	98 (24.0)	77 (20.4)	
Asia	35 (3.3)	10 (2.4)	19 (5.0)	
Others	58 (5.6)	23 (5.6)	16 (4.2)	
Missing	*3 (0.3)*	*8 (1.0)*	*5 (1.3)*	
Gravidity, n (%)				0.06
1	311 (29.7)	129 (30.9)	123 (32.1)	
2	336 (32.1)	122 (29.3)	92 (24.0)	
≥3	400 (38.2)	166 (39.8)	168 (43.9)	
Missing	*1 (0.1)*	*0 (0)*	*0 (0)*	
Parity, n (%)				0.001
0	469 (44.8)	220 (52.8)	190 (49.6)	
1	378 (36.1)	129 (30.9)	102 (26.6)	
≥2	200 (19.1)	68 (16.3)	91 (23.8)	
Missing	*1 (0.1)*	*0 (0)*	*0 (0)*	
History of preterm delivery, n (%)	109 (10.4)	95 (22.8)	67 (17.5)	<0.0001
Missing	*3 (0.3)*	*0 (0)*	*0 (0)*	
Body mass index, median (IQR)	22.4 (20.4 - 25.7)	23.0 (20.3 - 26.7)	24.5 (21.7 - 28.7)	<0.0001
Underweight (<18.5), n (%)	57 (5.4)	38 (9.2)	19 (5.1)	
Normal weight (18.5-24.9), n (%)	670 (64.0)	228 (55.2)	179 (47.6)	
Overweight (25-29.9), n (%)	211 (20.2)	90 (21.8)	102 (27.1)	
Obese (≥30), n (%)	109 (10.4)	57 (13.8)	76 (20.2)	
Missing	*1 (0.1)*	*4 (1.0)*	*7 (1.8)*	
Gestational diabetes, n (%)	194 (18.5)	69 (16.5)	82 (21.4)	0.21
Missing	*1 (0.1)*	*0 (0)*	*0 (0)*	
Hypertensive disease, n (%)	41 (3.9)	15 (3.6)	4 (1.0)	0.02
Missing	*1 (0.1)*	*0 (0)*	*0 (0)*	
Smoking during pregnancy, n (%)	56 (5.4)	36 (8.7)	25 (6.6)	0.06

	Control N=1,048	PTL N=417	PPROM N=383	*P* value
Missing	*3 (0.3)*	*3 (0.7)*	*3 (0.8)*	
Fetal growth restriction, n (%)	27 (2.6)	29 (7.0)	27 (7.0)	<0.0001
Missing	*1 (0.1)*	*0 (0)*	*0 (0)*	
Gestational age at delivery, median (IQR), weeks	40.0 (39.0-40.7)	38.1 (35.1-39.3)	35.1 (30.6-36.4)	<0.0001
Extremely preterm (<28 weeks), n (%)	0 (0)	33 (7.9)	57 (15.0)	
Very preterm (28-31 weeks), n (%)	0 (0)	24 (5.8)	66 (17.3)	
Moderately preterm (32-36 weeks), n (%)	0 (0)	96 (23.0)	204 (53.5)	
Term (≥ 37 weeks), n (%)	1048 (100)	264 (63.3)	54 (14.2)	
Missing	*0 (0)*	*4 (1.0)*	*2 (0.5)*	
Mode of delivery				<0.0001
Vaginal, spontaneous	672 (64.4)	290 (71.2)	219 (57.6)	
Vaginal, induced	317 (30.4)	98 (24.1)	105 (27.6)	
Planned Caesarean	55 (5.2)	19 (4.7)	56 (14.8)	
Missing	*4 (0.4)*	*10 (2.4)*	*3 (0.8)*	
Interval between ROM and delivery, median (IQR), days	0 (0-1)	0 (0-1)	5 (1-17)	<0.0001
Missing	*0 (0)*	*36 (8.6)*	*13 (3.4)*	
Antibiotics between inclusion and delivery[b], n (%)	138 (13.2)	126 (30.4)	313 (81.7)	<0.0001
Missing	*0 (0)*	*2 (0.5)*	*0 (0)*	
Live births, n (%)	1046 (100)	411 (100)	378 (99.2)	NT
Missing	*2 (0.2)*	*6 (1.4)*	*2 (0.5)*	

[a] Including territories in South America (French Guiana), the Atlantic Ocean (Guadeloupe,
Martinique, Saint-Martin, Saint-Barthélémy, Saint-Pierre-et-Miquelon), the Indian Ocean
(Reunion Island, Mayotte), the Pacific Ocean (New Caledonia, French Polynesia, the Wallis
and Futuna Islands), and the French Southern and Antarctic Territories.
[b] Including intrapartum antibiotic prophylaxis and excluding non-systemic antibiotic therapy.
IQR: interquartile range; NT: not tested; PPROM: preterm premature rupture of membranes;
PTL: threatened preterm birth or preterm labor; ROM: rupture of membranes.

Table 2. Vaginal microbiology at inclusion.

Variable, n (%)	Control N=1,048	PTL N=417	*P* value[a]	PPROM N=383	*P* value[b]
Nugent score			0.01		0.048
Normal (0-3)	713 (85.3)	270 (85.7)		225 (79.2)	
Intermediate (4-6)	107 (12.8)	30 (9.5)		53 (18.7)	
Bacterial vaginosis (≥7)	16 (1.9)	15 (4.8)		6 (2.1)	
Missing	*212 (20.2)*	*102 (24.5)*		*99 (25.8)*	
Microbiological cultures					
***Lactobacillus* spp.**			<0.0001		<0.0001
Numerous	845 (80.8)	281 (67.5)		201 (54.9)	
Rare	33 (3.2)	16 (3.8)		34 (9.3)	
Absent	168 (16.0)	119 (28.6)		131 (35.8)	
Missing	*2 (0.2)*	*1 (0.2)*		*17 (4.4)*	
Enterobacteria, all species			<0.0001		<0.0001
Numerous	77 (7.3)	85 (20.4)		81 (21.2)	
Rare	91 (8.7)	39 (9.4)		48 (12.6)	
Absent	880 (84.0)	293 (70.3)		253 (66.2)	
Missing	*0 (0)*	*0 (0)*		*1 (0.3)*	
Escherichia coli			<0.0001		<0.0001
Numerous	52 (5.0)	59 (14.1)		51 (13.5)	
Rare	67 (6.4)	33 (7.9)		41 (10.8)	
Absent	929 (88.6)	325 (77.9)		286 (75.7)	
Missing	*0 (0)*	*0 (0)*		*5 (1.3)*	
Klebsiella pneumoniae			<0.0001		<0.0001
Numerous	13 (1.2)	21 (5.0)		22 (5.8)	
Rare	12 (1.1)	6 (1.4)		8 (2.1)	
Absent	1023 (97.6)	390 (93.5)		348 (92.1)	
Missing	*0 (0)*	*0 (0)*		*5 (1.3)*	
Staphylococcus aureus			0.07		0.64
Numerous	24 (2.3)	19 (4.6)		9 (2.4)	
Rare	25 (2.4)	10 (2.4)		6 (1.6)	
Absent	985 (95.3)	387 (93.0)		363 (96.0)	
Missing	*14 (1.3)*	*1 (0.2)*		*5 (1.3)*	
Group B *Streptococcus*			0.91		0.86
Numerous	125 (11.9)	53 (12.7)		43 (11.3)	
Rare	32 (3.1)	12 (2.9)		10 (2.6)	
Absent	891 (85.0)	352 (84.4)		328 (86.1)	
Missing	*0 (0)*	*0 (0)*		*2 (0.5)*	
Streptococcus anginosus			0.48		0.09
Numerous	26 (2.5)	15 (3.6)		17 (4.5)	
Rare	12 (1.1)	4 (1.0)		2 (0.5)	
Absent	1010 (96.4)	398 (95.4)		363 (95.0)	
Missing	*0 (0)*	*0 (0)*		*1 (0.3)*	
Anaerobes[c]			0.003		0.11
Numerous	113 (10.8)	72 (17.3)		53 (13.9)	

Variable, n (%)	Control N=1,048	PTL N=417	*P* value[a]	PPROM N=383	*P* value[b]
Rare	15 (1.4)	5 (1.2)		2 (0.5)	
Absent	919 (87.8)	340 (81.5)		326 (85.6)	
Missing	*1 (0.1)*	*0 (0)*		*2 (0.5)*	
Gardnerella vaginalis			<0.0001		0.06
Numerous	72 (6.9)	66 (15.8)		40 (10.5)	
Rare	10 (1.0)	4 (1.0)		2 (0.5)	
Absent	965 (92.2)	347 (83.2)		339 (89.0)	
Missing	*1 (0.1)*	*0 (0)*		*2 (0.5)*	
Candida spp.			0.97		0.67
Numerous	149 (14.3)	61 (14.7)		53 (14.0)	
Rare	31 (3.0)	12 (2.9)		8 (2.1)	
Absent	865 (82.8)	341 (82.4)		318 (83.9)	
Missing	*3 (0.3)*	*3 (0.7)*		*4 (1.0)*	

[a] Statistical analysis comparing PTL to control.

[b] Statistical analysis comparing PPROM to control.

[c] including *Gardnerella vaginalis.*

PTL: preterm labor; PPROM: preterm premature rupture of membranes.

554 **Table 3.** Univariate analysis of factors associated with preterm labor and preterm premature rupture of membranes.

	PTL (N=417)			PPROM (N=383)		
	OR	95% CI	*P* value	OR	95% CI	*P* value
Place of birth						
France	1 (ref)	-	<0.0001	1 (ref)	-	<0.0001
France, overseas territories[a]	3.26	1.85-5.74	-	3.97	2.23-7.06	-
Europe, other countries	1.05	0.63-1.75	-	1.21	0.71-2.06	-
North Africa	1.60	1.17-2.18	-	2.55	1.87-3.48	-
Sub-Saharan Africa	4.16	2.97-5.85	-	4.15	2.88-5.97	-
Asia	1.08	0.52-2.23	-	2.60	1.44-4.71	-
Gravidity						
1	1 (ref)	-	0.57	1 (ref)	-	0.01
2	0.88	0.65-1.17	-	0.69	0.51-0.94	-
≥3	1.00	0.76-1.32	-	1.06	0.81-1.40	-
Parity						
0	1 (ref)	-	0.02	1 (ref)	-	0.002
1	0.73	0.56-0.94	-	0.67	0.51-0.88	-
≥2	0.72	0.53-1.00	-	1.12	0.83-1.52	-
History of preterm delivery	2.53	1.87-3.43	<0.0001	1.82	1.31-2.53	0.0003
Body mass index						
Normal weight (18.5-24.9)	1 (ref)	-	0.004	1 (ref)	-	<0.0001
Underweight (<18.5)	1.96	1.27-3.03	-	1.25	0.72-2.15	-
Overweight (25-29.9)	1.25	0.94-1.67	-	1.81	1.36-2.41	-
Obese (≥30)	1.54	1.08-2.1	-	2.61	1.86-3.65	-

	PTL (N=417)			PPROM (N=383)		
Gestational diabetes	0.87	0.64-1.1	0.37	1.20	0.90-1.60	0.22
Hypertension	0.92	0.50-1.67	0.78	0.26	0.09-0.73	0.006
Smoking during pregnancy	1.68	1.09-2.60	0.02	1.24	0.76-2.02	0.38
Vaginal swab cultures						
Absence of lactobacilli	2.09	1.60-2.74	<0.0001	2.91	2.22-3.82	<0.0001
Presence of						
- Enterobacteria	2.22	1.70-2.90	<0.0001	2.67	2.04-3.49	<0.0001
- *Escherichia coli*	2.21	1.64-2.98	<0.0001	2.51	2.86-3.40	<0.0001
- *Klebsiella pneumoniae*	2.83	1.62-4.94	0.0001	3.53	2.05-6.08	<0.0001
- *Staphylooccus. aureus*	1.51	0.94-2.42	0.09	0.83	0.46-1.50	0.54
- Group B *Streptococcus*	1.05	0.77-1.44	0.76	0.92	0.66-1.28	0.61
- *Streptococcus anginosus*	1.27	0.72-2.23	0.41	1.39	0.79-2.44	0.25
- Anaerobes[b]	1.63	1.19-2.21	0.002	1.21	0.86-1.70	0.27
- *Gardnerella vaginalis*	2.37	1.69-3.34	<0.0001	1.46	0.99-2.16	0.06
- *Candida* spp.	1.03	0.76-1.39	0.86	0.92	0.67-1.27	0.62

[a] Including territories in South America (French Guiana), the Atlantic Ocean (Guadeloupe, Martinique, Saint-Martin, Saint-Barthélémy, Saint-Pierre-et-
Miquelon), the Indian Ocean (Reunion Island, Mayotte), the Pacific Ocean (New Caledonia, French Polynesia, the Wallis and Futuna Islands), and the French
Southern and Antarctic Territories.

[b] including *Gardnerella vaginalis.*

CI: confidence interval; OR: Odds ratio; PPROM: preterm premature rupture of membranes; PTL: preterm labor.

Table 4. Risk factors for spontaneous preterm delivery.

Variable, n (%)	Term births N=181	Preterm births N=328	OR	95% CI	*P* value
Gestational age at admission, weeks					
32-36	64 (35.4)	135 (41.2)	1 (ref)	-	0.15
28-31	56 (30.9)	76 (23.2)	0.64	0.41-1.01	-
< 28	61 (33.7)	117 (35.7)	0.91	0.59-1.40	-
Place of birth					
France	76 (42.9)	108 (33.4)	1 (ref)	-	0.08
France, overseas territories[a]	12 (6.8)	17 (5.3)	1.00	0.45-2.21	-
Europe, other countries	12 (6.8)	19 (5.9)	1.11	0.51-2.43	-
North Africa	32 (18.1)	82 (25.4)	1.80	1.09-2.98	-
Sub Saharan Africa	30 (16.9)	65 (20.1)	1.52	0.90-2.57	-
Asia	5 (2.8)	18 (5.6)	2.53	0.90-7.12	-
Others	10 (5.6)	14 (4.3)	NT	NT	NT
Missing	*4 (2.2)*	*5 (1.5)*	-	-	-
Gravidity					
1	62 (34.3)	115 (35.1)	1 (ref)	-	0.31
2	60 (33.1)	89 (27.1)	0.80	0.51-1.25	-
≥3	59 (32.6)	124 (37.8)	1.13	0.73-1.76	-
Parity					
0	102 (56.4)	177 (54.0)	1 (ref)	-	0.85
1	53 (29.3)	99 (30.2)	1.08	0.71-1.63	-
≥2	26 (14.4)	52 (15.9)	1.15	0.68-1.96	-
History of preterm delivery	35 (19.3)	61 (18.6)	0.95	0.60-1.51	0.84
Body mass index					
Normal weight (18.5-24.9)	117 (65.0)	159 (49.2)	1 (ref)	-	0.005
Underweight (<18.5)	13 (7.2)	27 (8.4)	1.53	0.76-3.09	-
Overweight (25-29.9)	34 (18.9)	82 (25.4)	1.77	1.11-2.83	-
Obese (≥30)	16 (8.9)	55 (17.0)	2.53	1.38-4.64	-
Missing	*1 (0.6)*	*5 (1.5)*	-	-	-
Gestational diabetes	19 (10.5)	72 (22.0)	2.40	1.39-4.13	0.001
Hypertension	3 (1.7)	3 (0.9)	0.55	0.11-2.74	0.67
Smoking during pregnancy	12 (6.7)	23 (7.1)	1.07	0.52-2.20	0.86
Missing	*1 (0.6)*	*4 (1.2)*	-	-	-
Vaginal microbiological cultures					
Lactobacilli, absence	41 (22.7)	108 (33.6)	1.73	1.14-2.63	0.01
Missing	*0 (0)*	*7 (2.1)*	-	-	-
Presence of					

Variable, n (%)	Term births N=181	Preterm births N=328	OR	95% CI	*P* value
- Enterobacteria	45 (24.9)	111 (33.8)	1.55	1.03-2.32	0.04
- *Escherichia coli*	32 (17.7)	83 (25.3)	1.58	1.00-2.49	0.05
- *Klebsiella pneumoniae*	7 (3.9)	27 (8.2)	2.23	0.95-5.23	0.06
- *Staphylococcus aureus*	10 (5.5)	13 (4.0)	0.71	0.30-1.64	0.42
- Group B *Streptococcus*	28 (15.5)	41 (12.5)	0.78	0.46-1.31	0.35
- *Streptococcus anginosus*	9 (5.0)	20 (6.1)	1.24	0.55-2.79	0.60
- Anaerobes[b]	32 (17.7)	58 (17.7)	1.00	0.62-1.61	0.92
- *Gardnerella vaginalis*	30 (16.6)	49 (14.9)	0.88	0.54-1.45	0.62
- *Candida* spp.	28 (15.5)	56 (17.1)	1.13	0.69-1.85	0.64

[a] Including territories in South America (French Guiana), the Atlantic Ocean (Guadeloupe,
Martinique, Saint-Martin, Saint-Barthélémy, Saint-Pierre-et-Miquelon), the Indian Ocean (Reunion
Island, Mayotte), the Pacific Ocean (New Caledonia, French Polynesia, the Wallis and Futuna
Islands), and the French Southern and Antarctic Territories.

[b] Including *Gardnerella vaginalis*.

CI: confidence interval. NT: not tested; OR: Odds ratio.

Supplementary Information

Supplementary Table 1. Vaginal microbiota characteristics according to the place of

birth.

Supplementary Table 2. Vaginal microbiota characteristics according to body mass

index.

Supplementary Figure 1. Second model of multivariable analysis of risk factors for

preterm labor (PTL) and preterm premature rupture of membranes (PPROM).

In this model, *Escherichia coli* and *Klebsiella pneumoniae* were used instead of

enterobacteria as variables. Error bars indicate the upper and lower limit of the 95%

confidence interval (CI); aOR: adjusted odds ratio.

Supplementary File 1. Composition of the Inspire consortium.

Figure 1. Vaginal microbiota composition at admission for preterm labor (PTL) and

preterm rupture of membranes (PPROM).

Vaginal microbiota composition was characterized by vaginal swab cultures and compared

between controls (n=1,048) and women admitted for PTL (n=417) or PPROM (n=383). The

percentage of vaginal samples positive for selected bacterial group or species are represented.

Error bars indicate the upper limit of the 95% confidence interval. ** $p < 0.01$; *** $p < 0.001$,

ns: not significant.

Figure 2. Multivariable analysis of risk factors for preterm labor (PTL) and preterm premature rupture of membranes (PPROM).

Error bars indicate the upper and lower limit of the 95% confidence interval (CI); aOR: adjusted odds ratio.

Figure 3. Multivariable analysis of risk factors for spontaneous preterm birth.

Error bars indicate the upper and lower limit of the 95% confidence interval (CI); aOR: adjusted odds ratio.

607