Vaginal microbiota as a predictor of preterm birth : an observational cohort study

Authors

Laura Lesimple[1,2], Jessica Rousseau[3], Céline Plainvert[1,4], Luce Landraud[5,6], Nathalie Grall[7], François Goffinet[3,8], Pierre-Yves Ancel[3,9], Christophe Pannetier[2], Claire Poyart[1,4], Laurent Mandelbrot[6,10]*,, Asmaa Tazi[1,4]*

Affiliations

[1]Institut Cochin, Inserm U1016, CNRS UMR 8104, Université Paris Cité, Paris, France.

[2]BforCure, Montreuil, France.

[3]Obstetrical, Perinatal, and Pediatric Epidemiology Team and Biostatistics, Sorbonne Paris Cité Research Center, INSERM U1153, Université Paris Cité, Paris, France.

[4]Department of Bacteriology, National Reference Center for Streptococci, University Hospitals Assistance Publique - Hôpitaux de Paris, Hôpital Cochin, Paris, France.

[5]Department of Microbiology, University Hospitals Assistance Publique - Hôpitaux de Paris, Hôpital Louis Mourier, Colombes, France.

[6]Université Paris Cité, IAME INSERM U1137, Paris, France.

[7]Department of Bacteriology, University Hospitals Assistance Publique - Hôpitaux de Paris, Hôpital Bichat, Paris, France.

[8]Department of Obstetrics and Gynaecology, Port-Royal Maternity, University Hospitals Paris Centre Cochin Port Royal, AP-HP, Paris, France.

[9]URC-CIC P1419, University Hospitals Paris Centre Cochin Port Royal, AP-HP, Paris, France.

[10]Department of Obstetrics and Gynaecology, University Hospitals Assistance Publique - Hôpitaux de Paris, Hôpital Louis Mourier, Colombes, France.

26   ***** Laurent Mandelbrot and Asmaa Tazi contributed equally

27   2

Corresponding author

Dr Asmaa Tazi

HUPC Cochin, Service de Bactériologie, 27 rue du Faubourg St Jacques, 75014, Paris, France.

E-mail: asmaa.tazi@aphp.fr

Phone: +33 1 58 41 34 14 - Fax: +33 1 58 41 15 48

Short title: Vaginal microbiota signatures of preterm birth

At a glance

Why was this study conducted?

Despite the growing number of scientific studies highlighting the link between vaginal microbiota composition and adverse pregnancy outcomes, data remain controversial, mostly because studies are relatively small-sized and do not investigate cofounding factors such demographics and medical and obstetrical history. Moreover, microbiota data are difficult to transpose into a real-life perspective where clinical management relies primarily on clinical risk factors and vaginal swab cultures.

Key findings

We included over 2,400 pregnant women in a large longitudinal cohort study and combined culturomics at admission for preterm labor or preterm rupture of membranes and multivariable analyses in which demographic and obstetrical data were studied. We identified lactobacilli depletion and enterobacteria as strong vaginal microbiota signatures of preterm labor and preterm rupture of membranes. Importantly, both signatures constituted independent risk factors for preterm birth.

What does this add to what is known?

These findings highlight the importance of vaginal microbiota together with demographic and clinical features as risk factors for adverse pregnancy outcome and preterm birth. They also underline the need for an accurate characterization of vaginal microbiota at admission for preterm labor and preterm rupture of membranes for a better management of these complications of pregnancy.

Abstract

Background: Preterm birth (birth before 37 weeks of gestational age) is a frequent and severe adverse pregnancy outcome. Despite the growing number of scientific studies highlighting the link between vaginal microbiota composition and adverse pregnancy outcomes, data remain controversial.

Objectives: To identify microbiota signatures of preterm labor and premature rupture of outer membranes; to determine microbiological risk factors for preterm birth.

Study Design: We conducted an observational, prospective, longitudinal cohort study from August 2018 until June 2023 in 3 maternity wards from university hospitals in the Paris, France, area. Women were categorized in 4 groups including a control group, and 3 groups of increasing risk of intrauterine infection and preterm birth: prelabor rupture of membranes at term, preterm labor, and preterm premature rupture of membranes. Demographic, clinical data, past medical and obstetrical history and pregnancy outcome were collected. Vaginal swabs were collected at admission and were analyzed using culturomics. The association between bacterial species and the cohort groups and eventually preterm birth was studied in univariate analyses. Adjusted odds ratio (aOR) and 95% confidence intervals (95% CI) were calculated in multivariable analyses.

Results: A total of 2,476 women were included, of whom 1,068 (43.1%) in the control group, 477 (19.3%) with term premature rupture of outer membranes, 495 (20.0%) with preterm labor, and 436 (17.6%) with preterm premature rupture of outer membranes. Together with demographic features such as ethnicity and obstetrical history, several vaginal microbiota signatures were identified as correlated with pregnancy outcome. In multivariable analysis, prelabor rupture of membranes at term was associated with enterobacteria (aOR 1.97, 95% CI 1.46-2.65) and *Gardnerella vaginalis* (aOR 5.19, 95% CI 2.22-13.78); preterm labor with lactobacilli depletion (aOR 1.49, 95% CI 1.08-2.06), enterobacteria (aOR 1.86, 95% CI 1.36-

2.53) and *G. vaginalis* (aOR 4.62, 95% CI 1.86-13.34); preterm premature rupture of membranes with lactobacilli depletion (aOR 2.04, 95% CI 1.49-2.80) and enterobacteria (aOR 2.38, 95% CI 1.74-3.24). Finally, together with gestational diabetes, lactobacilli depletion and enterobacteria both represented risk factors for preterm birth, especially in singleton pregnancies (aOR 1.54, 95% CI 1.05-2.28 and aOR 1.62, 95% CI 1.11-2.36, respectively).

Conclusions: In this large cohort study, we identified on the one hand, *G. vaginalis* as associated with prelabor rupture of membranes at term and preterm labor, and on the other hand, lactobacilli depletion and enterobacteria as risk factors for preterm labor, preterm premature rupture of outer membranes, and preterm birth, emphasizing the importance of a healthy microbiota in pregnancy outcome. Further studies are needed to address the benefits of adapted antibiotic prophylaxes and probiotics aiming at restoring a healthy microbiota in reducing the risk of preterm birth.

Introduction

Preterm birth (PTB), defined as birth before 37 weeks of gestation, is a frequent and severe adverse pregnancy outcome which concerns more than 10% pregnancies and accounts for 14 millions annual births worldwide.[1] As the leading causes of death in children under the age of 5 years, PTB and its sequelae represent one of the most prevalent conditions in the Global Burden of Disease analysis and are therefore considered as a major worldwide health issue.[2,3] Up to 40% of PTB are preceded by preterm premature rupture of membranes (PPROM), increasing the risk of intrauterine infection, *in utero* demise and prematurity.[4,5] Nevertheless, causes of PTB and PPROM are still poorly understood. Beside multiples factors including ethnicity, obstetrical history, and lifestyle, perturbations of the vaginal microbiota are increasingly recognized as possible contributing factors.[6,7]

The composition of the vaginal microbiota is highly dynamic, changing with age, hormonal fluctuations, gestational status, ethnicity, sexual behavior, and lifestyle.[8–10] Microbiota studies based on sequencing of vaginal samples from reproductive-age women have identified five types of vaginal community structures designated community state types (CSTs).[10] Four of these are dominated by one *Lactobacillus* species while the last one, designated CST-IV, is characterized by a more diverse microbiota dominated by anaerobic bacteria such as *Gardnerella vaginalis, Prevotella* spp, and *Fannyhessea vaginae* – formerly known as *Atopobium vaginae*. Although the distribution of these CSTs largely varies according to ethnicity,[11] it is generally believed that a healthy vaginal microbiota is dominated by lactobacilli with a low bacterial diversity, particularly during pregnancy.[12,13] Conversely, a decrease in lactobacilli abundance is correlated to an increase of microbiota diversity typically exemplified by the CST-IV. Low-lactobacilli microbiota can furthermore be associated to clinical symptoms known as bacterial vaginosis and with higher susceptibility to sexually transmitted or urinary tract infections.[14] In the context of pregnancy, increasing evidence links preterm labor (PTL),

PPROM, and PTB with low-lactobacilli microbiota characterized by a high diversity and an enrichment in *G. vaginalis* and various anaerobes, whereas microbiota dominated by lactobacilli and more specifically by *Lactobacillus crispatus* (CST-I) have been found associated to term birth.[6,15–17]

However, despite the growing number of scientific studies that highlight the link between vaginal microbiota composition and pregnancy outcomes, data remain controversial. Most studies are relatively small-sized and do not investigate cofounding factors such as ethnicity. Moreover, microbiota data are difficult to transpose into a real-life perspective where clinical management relies primarily on clinical risk factors and vaginal swab cultures. Here, we conducted a large prospective observational cohort study involving more than 2,400 pregnant women and used simple culturomics and multivariable analyses that took demographic and clinical data into account to identify microbiota signatures of PTL and PPROM, and eventually to determine microbiological risk factors for PTB.

Materials and methods

Study design

We conducted an observational, prospective, longitudinal cohort study from August 2018 until June 2023 in 3 maternity wards, of which 2 of level 3 and 1 of level 2b, from university hospitals in the Paris, France, area. The total annual number of births at these 3 centers is approximately 11,000. Women were categorized in 4 groups including a control group and 3 groups of increasing risk of intrauterine infection and PTB. The eligibility criteria for maternal participation in the study included all pregnant women aged >18 years who responded to the following inclusion criteria : *i)* prelabor rupture of membranes at term (TPROM) e.g., rupture of membranes after 37 weeks of gestation and more than 12 hours before labor onset; *ii)* preterm labor (PTL) e.g., hospitalization for threatened PTB or PTL with intact membranes; and *iii)*

PPROM e.g., rupture of membranes before 37 weeks of gestation. The control group was

constituted by pregnant women who delivered at term with no complication. In this latter group,

women were proposed to be enrolled at the time of *Streptococcus agalactiae* (Group B

Streptococcus, GBS) antenatal screening between 34 and 38 weeks of gestational age (GA) as

recommended by the French guidelines for the prevention of perinatal infectious disease.[18]

Women who did not speak or understand French were excluded.

Demographic, clinical data, past medical and obstetrical history, and pregnancy outcome were

collected. Notably, maternal place of birth was collected as a reflect of ethnicity, given that

reporting ethnicity itself is forbidden in France.

Sampling and laboratory procedures

Vaginal swabs were sampled at admission or at GBS antenatal screening for the control group

and sent to the laboratory in Amies transport medium (ESwab™ system, Copan Diagnostics,

Italy). Specimens were immediately processed for analysis by conventional culture methods

using non-selective rich media incubated at 37°C in aerobic atmosphere with 5% CO_2. In

addition, a Columbia agar with 5% horse blood, nalidixic acid and colistin (bioMérieux, Marcy

l'Etoile, France) for the selective detection of anaerobes and Gram-positive cocci and a Granada

agar (Fisher Scientific, Illkirch, France) for the selective detection of GBS were incubated at

37°C under anaerobic atmosphere. Lastly, a Drigalski and a Candida agar plates (bioMérieux,

Marcy l'Etoile, France) for the selective detection of Gram-negative rods and yeasts,

respectively, were incubated at 37°C under aerobic atmosphere. All plates were examined after

16h and 44h of incubation. All different morphotypes of bacterial and yeast colonies were

identified using matrix assisted laser desorption ionization - time of flight (MALDI-TOF) mass

spectrometry (Bruker Daltonics, Billerica, Massachussetts, USA). The density of vaginal

colonization was estimated semi-quantitatively as follows: colonies on a single quadrant, low;

colonies on more than two quadrants, medium to high. In addition, the overall vaginal microbiota composition was assessed by direct microscopic examination of the vaginal samples after Gram staining and determination of the Nugent score in 2 out of 3 maternity wards according to routine practices.[19]

Primary outcomes and studied factors

The primary outcomes were PROM at admission and PTB in the PTL and PPROM groups. The studied factors were demographics (country of birth as a reflect of ethnicity, body mass index (BMI)), pregnancy-related risk factors (gestity, parity, history of preterm delivery, multiple pregnancy, gestational diabetes, hypertensive disease, tobacco use), and vaginal microbiota composition assessed by culturomics. A specific attention was paid to bacteria of the *Lactobacillus* genus, and to species potentially associated with adverse outcomes including enterobacteria (*Escherichia coli*, *Klebsiella pneumoniae*, and others), *Staphylococcus aureus*, GBS, *Streptococcus anginosus*, *G. vaginalis*, strict anaerobes (*Prevotella* spp., *Bacteroides* spp., and others), and fungi of the *Candida* genus.

Statistical analyses

All data were collected in a computerized database and analyzed using the GraphPad Software v8.3.0 (Dotmatics, Boston, Massachusetts, USA) and RStudio v1.1.447 (Posit, Boston, Massachusetts, USA). Firstly, we assessed the prevalence of the studied factors according to the cohort groups as a proportion of the total number of participants. Secondly, to study the risk factors associated to each cohort group and the risk factors for PTB, a multivariable logistic regression model was used for prognostic covariates. For univariate statistical analyses, Mann-Whitney tests were used when appropriate, and categorical data were compared using chi-squared or Fisher's exact tests. Odds ratios and 95% CI were calculated in univariate analyses.

Adjusted odds ratios were then calculated using any variables remaining associated with the

outcome with a p-value <0.2 from the univariate analysis, excepting microbiological variables

which were all included in the analyses. Two models were used to study enterobacteria. In the

first one, enterobacteria were analyzed as a single variable. In the second one, the two most

represented species e.g., *E. coli* and *K. pneumoniae*, were specifically analyzed. A p-value <

0.05 was considered significant. Missing data were excluded.

Ethics declaration

The reported study was reviewed by the French ethics committee who granted approval on

November, 2017 (identifier 2017 – A02755-48; NCT03371056). All individuals enrolled in the

cohort provided signed informed consent that included authorization for the collection of

medical information and biological samples. All collected data were anonymized and integrated

in a secured electronic database.

Results

Study population

A total of 2,476 women were included, of whom 1,068 (43.1%) in the control group, 477

(19.3%) in the TPROM group, 495 (20.0%) in the PTL group, and 436 (17.6%) in the PPROM

group. The characteristics of the study population are summarized in Table 1. A number of

demographical and clinical characteristics varied according to the group at admission. For

instance, women born in Sub Saharan Africa were overrepresented in the PTL and PPROM

groups. In addition, the PTL and PPROM groups were characterized by a higher proportion of

multiple pregnancies, history of preterm deliveries, and obesity. Eventually, 283 (57.6%) and

58 (13.4%) women of the PTL and PPROM groups delivered at term.

Vaginal microbiota signatures of preterm labor and premature rupture of membranes

A Nugent score was determined for 2,018 (81.5%) vaginal samples and indicated that bacterial vaginosis was uncommon, below 5% whatever the cohort group considered (Table S1). However, intermediate scores were overrepresented in the PPROM group (60/330, 18.2% *vs.* 110/851, 12.9% in controls, p=0.02). Culturomics of the vaginal samples revealed significant differences according to the cohort groups, most importantly in the abundance of lactobacilli, enterobacteria, and anaerobes, including *G. vaginalis* (Figure 1 and Table S1). Overall, lactobacilli were less frequently isolated in the PTL (357/493, 72.4%) and PPROM (274/418, 65.6%) groups compared with controls (894/1066, 83.9%). Conversely, vaginal samples positive for enterobacteria and anaerobes were overrepresented in the high-risk groups. For instance, enterobacteria were isolated in more than 25% women of the TPROM, PTL and PPROM groups *vs.* 16% in controls. *E. coli* was the most frequent species and showed the strongest significant differences. Similarly, *G. vaginalis* was cultured in 13.7% (65/474), 16.8% (83/495), and 10.6% (46/434) samples from the TPROM, PTL and PPROM groups, respectively, *vs.* 7.8% (83/1,067) controls. The abundance of *S. aureus*, GBS and *Candida* spp. did not vary significantly according to the cohort groups (Table S1).

Demographical and clinical characteristics such as ethnicity and BMI are linked with pregnancy outcomes and are also presumed to impact microbiota composition. Therefore, we tested their association with the species of the vaginal microbiota and found that the place of birth (Table S2) and the BMI (Table S3) were both correlated with the presence and abundance of lactobacilli.

Demographic, clinical and microbiological factors associated with PTL and PROM

Next, we addressed the association between demographics, clinical factors, and vaginal microbiota with PTL and PROM using univariate (Table S4) then multivariable logistic

regression analyses (Figures 2 and S1). Altogether, the factors the most associated with

TPROM were a North African or Asian origin, nulliparity, and a vaginal sample positive for

enterobacteria (aOR: 1.97, 95% CI 1.46-2.65) or *G. vaginalis* (aOR: 5.19, 95% CI 2.22-13.78).

PTL was strongly associated with the maternal place of birth and most-importantly with birth

in French overseas territories and Sub-Saharan Africa, with nulliparity, a multiple pregnancy,

a history of preterm delivery, and smoking during pregnancy. Considering vaginal microbiota,

the absence of lactobacilli (aOR: 1.49, 95% CI 1.08-2.06) and the presence of enterobacteria

(aOR: 1.86, 95% CI 1.36-2.53) or *G. vaginalis* (aOR: 4.62, 95% CI 1.86-13.34) were all

associated with PTL. Lastly, women from all areas outside Europe were more frequently

represented in the PPROM group in comparison with controls, together with nulliparous

women, women with a multiple pregnancy, a history of preterm delivery, and overweight.

Considering vaginal microbiota composition, the absence of lactobacilli (aOR 2.04, 95% CI

1.49-2.80) and the presence of enterobacteria (aOR 2.38, 95% CI 1.74-3.24) in vaginal swab

cultures were both strongly correlated with PPROM.

Altogether, the absence of lactobacilli in vaginal swab cultures was a risk factor for PTL and

PPROM but not for TPROM, enterobacteria were associated with TPROM, PTL, and PPROM,

and *G. vaginalis* was a signature of TPROM and PTL but not of PPROM. Similar results were

obtained with the second model of multivariable logistic regression where *E. coli* and *K.*

pneumoniae were included instead of enterobacteria (Figure S1), showing that vaginal

colonization by *E. coli* was strongly associated with TPROM (aOR 1.91, 95%CI 1.37-2.67),

PTL (aOR 2.02, 95% CI 1.43-2.86), and PPROM (aOR 2.39, 95% CI 1.69-3.38).

Vaginal microbiota with enterobacteria or lacking lactobacilli are risk factors for preterm

birth

We next focused on the PTL and PPROM groups, with the highest risk of adverse pregnancy

outcomes, and we sought to determine whether the vaginal microbiota at admission could be

predictive of PTB. Firstly, we analyzed in univariate and multivariable analyses the

demographical, clinical and microbiological risk factors for spontaneous PTB (510/826, 61.7%,

Table S5). This first analysis showed that the outcome was highly correlated with multiple

pregnancy (aOR 4.80, 95% CI 2.72-9.03, p<0.0001). Whereas vaginal microbiota lacking

lactobacilli and including enterobacteria were both associated with PTB in univariate analysis,

only enterobacteria remained correlated with PTB in multivariable analysis (aOR 1.64, 95%CI

1.15-2.37, Figure 3A). Similar results were obtained in the second model of multivariable

analysis where *E. coli* was associated with preterm birth (aOR 1.52, 95% CI 1.03-2.29, Figure

S2A). In a second analysis focusing on singleton pregnancies which represented 86.7%

(716/826) of women admitted for PTL or PPROM, the absence of lactobacilli and the presence

of enterobacteria both remained significantly correlated with PTB in multivariable analysis

(aOR 1.54, 95% CI 1.05-2.28 and 1.62, 95% CI 1.11-2.36, respectively, Figures 3B and S2B,

Table S6).

Comment

Principal Findings

In this cohort study of over 2,400 pregnant women from various ethnicities recruited in three

French maternity wards, we identified using simple culturomics of vaginal samples distinct

microbiological signatures of PTL, TPROM, and PPROM. Notably, we identified, using

multivariable logistic regression models, that vaginal microbiota, together with demographic

and clinical features, was correlated with these complications of pregnancy. For instance, PTL

was strongly associated with the presence of *G. vaginalis* and less significantly with the absence

of lactobacilli, TPROM with the presence of *G. vaginalis* and enterobacteria, whereas PPROM

was correlated with a depletion in lactobacilli and with the presence of enterobacteria. Importantly, microbiota composition at admission, and particularly a depletion in lactobacilli and the presence of enterobacteria were predictive of PTB.

Results in the Context of What is Known

These results are in perfect agreement with those already published in the literature. Considering PTB, a meta-analysis conducted by Gudnadottir *et al.* in which the vaginal microbiome was assessed before the onset of labor reported an increased risk of preterm delivery in women with low-lactobacilli microbiota.[16] In a recent longitudinal cohort study of 175 pregnant women comprising 40 PTB, the vaginal microbiome of pregnancies that ended preterm was found to be genetically more diverse, a diversity which was driven by *Gardnerella* spp.[20] Lastly, the analysis of the vaginal microbiota at delivery of a subset of 749 women from our cohort study using shotgun metagenomics found that PTB was correlated to the overall diversity of the microbiota.[17] In our study, the role of lactobacilli depletion and the correlated increased microbiota diversity as risk factors for PTB varied according to the studied population. For instance, we identified lactobacilli depletion as a risk factor for PTB only in women with a singleton pregnancy. These divergences are likely due to the specificities of our cohort which included high-risk pregnancies and multiple pregnancies, which appeared as the strongest risk factor for PTB.

Considering PPROM, Brown *et al.* analyzed the vaginal microbiome of 86 women and identified reduced *Lactobacillus* spp. abundance and increasing vaginal bacterial diversity as a risk factor.[21] Similarly, Yan et al. analyzed the vaginal microbiota of 112 women and found an increased abundance of *G. vaginalis* and other anaerobes in women presenting with PPROM. In line with the study of Brown *et al.*, we found that a depletion in lactobacilli was strongly associated with PPROM both in univariate and multivariable analysis. Conversely, *G. vaginalis*

was not identified as a risk factor for PPROM. This discrepancy is likely related to the

methodology that was used in our study where microbiota was assessed by culture, which is

less sensitive than metagenomics, especially for anaerobes.

Clinical Implications

The role of bacterial vaginosis and *G. vaginalis* in PTL and PTB has been suspected for a long

time.[22,23] Growing evidence regarding the direct role of *G. vaginalis* in the pathophysiology of

PTL is now accumulating. *G. vaginalis* and other anaerobes such as *Prevotella* spp. are able to

degrade the cervical mucus, favoring ascending intra-uterine infections, local inflammation,

uterine motility, and PTL.[24,25] It is now generally recognized that an antibiotic treatment, mainly

based on metronidazole or clindamycin, should be administered to women presenting with PTL

and diagnosed with symptomatic bacterial vaginosis. In many countries, national guidelines

also recommend screening for bacterial vaginosis and treatment of women with a history of

PTB, although this strategy has not proven to be efficient in reducing the risk of PTB.[26–28] Here

we show that *G. vaginalis* is a risk factor for PTL but more importantly that it is the absence of

lactobacilli and the presence of enterobacteria that seeme predictive of PTB. These

observations, in line with numerous metagenomic studies which underline the protective role

of lactobacilli and more especially of *L. crispatus*, *L. gasseri*, and *L. jensenii,*[17,29,30] argue for a

comprehensive management of bacterial vaginosis in pregnancy aiming at restoring a healthy

microbiota. Such management would presumably associate antibiotics targeting anaerobes to

probiotics, as proposed for recurrent vaginosis.[28]

Our results also raise questions about the management of vaginal colonization by enterobacteria

during TPROM, PTL and PPROM. We identified *E. coli*, and more generally enterobacteria as

strongly associated with TPROM, PTL and PPROM in univariate and multivariable analyses.

Both were also identified as risk factors for PTB. However, in contrast to GBS, there is currently

no consensus guidelines for the screening and the administration of antibiotics against

enterobacteria during pregnancy, not even in case of PROM. In the absence of signs of

infection, strategies largely vary between countries and range from no treatment to large

spectrum antibiotics such as 3rd generation cephalosporins.[31] Our observations also argue in

favor of administering a prophylaxis adapted to the pathogenic species present in the vaginal

microbiota at admission for TPROM, PTL, and PPROM, although further studies are needed

to ascertain the benefits of such a strategy.

Research Implications

Interventional studies aiming at assessing the role of antibiotics alone or in combination with

probiotics in reducing the risk of PTB and PPROM in high-risk populations and women who

experience bacterial vaginosis should be undertaken. These studies should rely on randomized

control trials that would take cofounding variables such as ethnicity, BMI, and gestational age

at intervention into account. Indeed, a given probiotic strain would not necessarily be protective

for all women, a property that might be linked to women's CSTs. In addition, the benefits of

personalized targeted antibiotic prophylaxes directed against the specific pathogenic species

e.g., GBS, enterobacteria, *Candida* spp. and others, found in the vaginal flora in reducing the

risk of PTB especially in case of PPROM should also be investigated.

Strengths and Limitations

This is a large cohort study where over 2,400 women were enrolled, including almost 500 PTL

and 500 PPROM. In addition to this major strength, our study also investigated important

cofounding variables of the outcome measures including demographical factors such as

ethnicity, BMI, history of PTB, and others. Hence, we could demonstrate that the vaginal

microbiota is an independent risk factor for TPROM, PTL, and PPROM, and also for PTB.

However, there are limitations that must be taken into account in the interpretation of our

results. First of all, microbiota analyses were carried out using simple culturomics and not

metagenomics. This leads to three main issues: (i) fastidious species such as *Atopobium* spp.,

or mollicutes could not be studied; (ii) culturomics being less sensitive than metagenomics,

cultured species were most likely the predominant ones and minor species have probably been

overlooked; (iii) although a semi-quantitative analysis of the cultured species was initially

performed, it rapidly came out because of differences in their growth characteristics, that this

analysis was biased and not representative of the microbiota equilibrium. Eventually, only the

presence or absence of major species was investigated. The second main limitation is the

absence of identification of lactobacilli at the species level. Consequently, the vaginal

microbiota were not categorized according to CSTs, and the protective role conferred by the

different species could not be addressed.

Conclusions

In this large multicentric cohort study, we underlined using culturomics distinct vaginal

microbiota signatures of adverse pregnancy outcomes. We identified *G. vaginalis* and

lactobacilli depletion as independent and strong risk factors for TPROM and PTL, and of

PPROM, respectively. Besides, enterobacteria were identified as risk factors for these three

situations and for PTB. Further studies are needed to address the potential of interventional

studies aiming at restoring a healthy microbiota in reducing these outcomes.

Transparency declaration

Ethics statement

The reported study was reviewed by the French ethics committee who granted approval on

November, 2017 (identifier 2017 – A02755-48; NCT03371056). All individuals enrolled in the

cohort provided signed informed consent that included authorization for the collection of

medical information and biological samples. All collected data were anonymized and integrated

in an secured electronic database.

Conflicts of interest

The authors have no conflicts of interest to disclose.

Funding

This work was supported by the Programme d'Investissements d'avenir and bpifrance

(Structuring R&D Project for Competitiveness—PSPC): # DOS0053477 SUB et DOS0053473

AR) .L.L. is a doctoral fellow supported by the Association Nationale de la Recherche et

Technologie (Cifre # 2019 / 1510). The funders had no role in the design and conduct of the

study.

Acknowledgements

We thank all the contributors to this study, collectively called the InSPIRE Consortium. The

full list can be found in the accompanying Supplementary Information file 1.

Part of the results were previously presented in April 2023 at the ECCMID scientific meeting

(Copenhagen, Denmark).

Access to data

All authors had full access to data. Data will be made available upon requests directed to the

corresponding author.

Authors' contributions

Conceptualization: C.Po., L.M., and A.T. Methodology: J.R., P.-Y.A., L.M., and A.T. Formal analysis: L.Le., J.R., P.-Y.A., L.M., and A.T. Investigation: L.Le., J.R., and A.T. Resources: C.Pl., L.La, N.G., F.G., and L.M. Data curation: L.Le, J.R., C.Pl, L.La, N.G., and A.T. Writing – Original draft: L.Le and A.T. Writing – Review and Editing: all authors. Supervision: C.Pa, C.Po., L.M. and A.T. Funding acquisition: C.Pa, C.Po and L.M.

References

1.   Ohuma EO, Moller AB, Bradley E, et al. National, regional, and global estimates of preterm birth in 2020, with trends from 2010: a systematic analysis. *Lancet Lond Engl*. 2023;402(10409):1261-1271. doi:10.1016/S0140-6736(23)00878-4

2.   Harrison MS, Goldenberg RL. Global burden of prematurity. *Semin Fetal Neonatal Med*. 2016;21(2):74-79. doi:10.1016/j.siny.2015.12.007

3.   GBD 2016 Causes of Death Collaborators. Global, regional, and national age-sex specific mortality for 264 causes of death, 1980-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Lond Engl*. 2017;390(10100):1151-1210. doi:10.1016/S0140-6736(17)32152-9

4.   Goldenberg RL, Culhane JF, Iams JD, Romero R. Epidemiology and causes of preterm birth. *Lancet Lond Engl*. 2008;371(9606):75-84. doi:10.1016/S0140-6736(08)60074-4

5.   Lorthe E. Épidémiologie, facteurs de risque et pronostic de l'enfant. RPC : rupture prématurée des membranes avant terme CNGOF. *Gynécologie Obstétrique Fertil Sénologie*. 2018;46(12):1004-1021. doi:10.1016/j.gofs.2018.10.019

6.   Bayar E, Bennett PR, Chan D, Sykes L, MacIntyre DA. The pregnancy microbiome and preterm birth. *Semin Immunopathol*. 2020;42(4):487-499. doi:10.1007/s00281-020-00817-w

7.   Schuster HJ, Bos AM, Himschoot L, et al. Vaginal microbiota and spontaneous preterm

birth in pregnant women at high risk of recurrence. *Heliyon*. 2024;10(10):e30685.

doi:10.1016/j.heliyon.2024.e30685

8.   Greenbaum S, Greenbaum G, Moran-Gilad J, Weintraub AY. Ecological dynamics of

the vaginal microbiome in relation to health and disease. *Am J Obstet Gynecol*.

2019;220(4):324-335. doi:10.1016/j.ajog.2018.11.1089

9.   Song SD, Acharya KD, Zhu JE, et al. Daily Vaginal Microbiota Fluctuations Associated

with Natural Hormonal Cycle, Contraceptives, Diet, and Exercise. Rao K, ed. *mSphere*.

2020;5(4):e00593-20. doi:10.1128/mSphere.00593-20

10.   Ravel J, Gajer P, Abdo Z, et al. Vaginal microbiome of reproductive-age women. *Proc*

Natl Acad Sci. 2011;108(supplement_1):4680-4687. doi:10.1073/pnas.1002611107

11.   France MT, Ma B, Gajer P, et al. VALENCIA: a nearest centroid classification method

for vaginal microbial communities based on composition. *Microbiome*. 2020;8(1):166.

doi:10.1186/s40168-020-00934-6

12.   Romero R, Hassan SS, Gajer P, et al. The composition and stability of the vaginal

microbiota of normal pregnant women is different from that of non-pregnant women.

Microbiome. 2014;2(1):4. doi:10.1186/2049-2618-2-4

13.   MacIntyre DA, Chandiramani M, Lee YS, et al. The vaginal microbiome during

pregnancy and the postpartum period in a European population. *Sci Rep*. 2015;5(1):8988.

doi:10.1038/srep08988

14.   Waltmann A, Thomas C, Duncan JA. The role of the genital microbiota in the

acquisition and pathogenesis of sexually transmitted infections. *Curr Opin Infect Dis*.

2023;36(1):35-48. doi:10.1097/QCO.0000000000000893

15.   Fettweis JM, Serrano MG, Brooks JP, et al. The vaginal microbiome and preterm birth.

Nat Med. 2019;25(6):1012-1021. doi:10.1038/s41591-019-0450-2

16.   Gudnadottir U, Debelius JW, Du J, et al. The vaginal microbiome and the risk of preterm

birth: a systematic review and network meta-analysis. *Sci Rep*. 2022;12(1):7926. doi:10.1038/s41598-022-12007-9

17. Baud A, Hillion KH, Plainvert C, et al. Microbial diversity in the vaginal microbiota and its link to pregnancy outcomes. *Sci Rep*. 2023;13(1):9061. doi:10.1038/s41598-023-36126-z

18. HAS. Prévention anténatale du risque infectieux bactérien néonatal précoce. Published 2001. https://www.has-sante.fr/upload/docs/application/pdf/prevention_antenatale_du_risque_infectieux_bacterien_-_rec.pdf

19. Nugent RP, Krohn MA, Hillier SL. Reliability of diagnosing bacterial vaginosis is improved by a standardized method of gram stain interpretation. *J Clin Microbiol*. 1991;29(2):297-301. doi:10.1128/jcm.29.2.297-301.1991

20. Liao J, Shenhav L, Urban JA, et al. Microdiversity of the vaginal microbiome is associated with preterm birth. *Nat Commun*. 2023;14(1):4997. doi:10.1038/s41467-023-40719-7

21. Brown RG, Al-Memar M, Marchesi JR, et al. Establishment of vaginal microbiota composition in early pregnancy and its association with subsequent preterm prelabor rupture of the fetal membranes. *Transl Res*. 2019;207:30-43. doi:10.1016/j.trsl.2018.12.005

22. Hillier SL, Martius J, Krohn M, Kiviat N, Holmes KK, Eschenbach DA. A Case–Control Study of Chorioamnionic Infection and Histologic Chorioamnionitis in Prematurity. *N Engl J Med*. 1988;319(15):972-978. doi:10.1056/NEJM198810133191503

23. Eschenbach DA, Gravett MG, Chen KC, Hoyme UB, Holmes KK. Bacterial vaginosis during pregnancy. An association with prematurity and postpartum complications. *Scand J Urol Nephrol Suppl*. 1984;86:213-222.

24. Howe L, Wiggins R, Soothill PW, Millar MR, Horner PJ, Corfield AP. Mucinase and

sialidase activity of the vaginal microflora: implications for the pathogenesis of preterm labour.

Int J STD AIDS. 1999;10(7):442-447. doi:10.1258/0956462991914438

25. Lewis WG, Robinson LS, Gilbert NM, Perry JC, Lewis AL. Degradation, Foraging, and

Depletion of Mucus Sialoglycans by the Vagina-adapted *Actinobacterium Gardnerella*

vaginalis. *J Biol Chem*. 2013;288(17):12067-12079. doi:10.1074/jbc.M113.453654

26. Collège national des gynécologues et obstétriciens français. Prévention de la prématurité

spontanée et de ses conséquences. Published online 2016.

https://cngof.fr/app/pdf/RPC//RPC%20DU%20CNGOF/2016/RPC_2016_Prmaturit_spontane

.pdf?x29325

27. British Association for Sexual Health and HIV. Bacterial Vaginosis. Published online

2012. https://www.bashh.org/resources/21/bacterial_vaginosis_2012/

28. Farr A, Swidsinski S, Surbek D, et al. Bacterial Vaginosis: Guideline of the DGGG,

OEGGG and SGGG (S2k-Level, AWMF Registry No. 015/028, June 2023). *Geburtshilfe*

Frauenheilkd. 2023;83(11):1331-1349. doi:10.1055/a-2169-8539

29. Tabatabaei N, Eren A, Barreiro L, et al. Vaginal microbiome in early pregnancy and

subsequent risk of spontaneous preterm birth: a case–control study. *BJOG Int J Obstet*

Gynaecol. 2019;126(3):349-358. doi:10.1111/1471-0528.15299

30. Callahan BJ, DiGiulio DB, Goltsman DSA, et al. Replication and refinement of a

vaginal microbial signature of preterm birth in two racially distinct cohorts of US women. *Proc*

Natl Acad Sci. 2017;114(37):9966-9971. doi:10.1073/pnas.1705899114

31. Werter DE, Dehaene I, Gurney L, Vargas Buján M, Kazemier BM. Differences in

clinical practice regarding screening and treatment of infections associated with spontaneous

preterm birth: An international survey. *Eur J Obstet Gynecol Reprod Biol*. 2021;266:83-88.

doi:10.1016/j.ejogrb.2021.09.009

Figure legends.

Figure 1. Vaginal microbiota composition at admission for prelabor rupture of membranes at term (TPROM), threatened preterm birth and preterm labor (PTL), and preterm rupture of membranes (PPROM).

Vaginal microbiota composition was characterized by culturomics and compared between controls (n=1,068) and women admitted for TPROM (n=477), PTL (n=495), or PPROM (n=436). The percentage of vaginal samples positive for each bacterial group or species are represented. Error bars indicate the upper limit of the 95% confidence interval. * p<0.05; *** p<0.001.

Figure 2. Multivariable analysis of risk factors for prelabor rupture of membranes at term (TPROM), threatened preterm birth and preterm labor (PTL), and preterm premature rupture of membranes (PPROM).

Adjusted odds ratios were calculated using any variables associated with the outcome with a p-value <0.2 from the univariate analysis, excepting microbiological variables which were all included. *S. aureus*, *S. anginosus* and *Candida* spp. did not show significant associations with the outcomes and are not represented. aOR: adjusted odds ratio; CI: confidence interval.

Figure 1.

Figure 1. Vaginal microbiota composition at admission for prelabor rupture of

membranes at term (TPROM), threatened preterm birth and preterm labor (PTL), and

preterm rupture of membranes (PPROM).

Vaginal microbiota composition was characterized by culturomics and compared between

controls (n=1,068) and women admitted for TPROM (n=477), PTL (n=495), or PPROM

(n=436). The percentage of vaginal samples positive for each bacterial group or species are

represented. Error bars indicate the upper limit of the 95% confidence interval. * $p < 0.05$; ***

$p < 0.001$.

Figure 2.

C - PPROM

		aOR (95% CI; p-value)
Place of birth	France	1
	France, overseas territories	3.16 (1.65-5.99; 0.0005)
	Europe	1.06 (0.59-1.83; 0.84)
	North Africa	1.96 (1.38-2.79; 0.0002)
	Sub Saharan Africa	3.50 (2.30-5.33; <0.0001)
	Asia	2.62 (1.38-4.88; 0.003)
Parity	0	1
	1	0.51 (0.34-0.78; 0.002)
	≥ 2	0.54 (0.31-0.91; 0.02)
Pregnancy with multiples		9.66 (5.22-18.61; <0.0001)
History of preterm delivery		1.76 (1.17-2.64; 0.007)
Body mass index	18.5 – 24.9	1
	< 18.5	1.35 (0.71-2.48; 0.34)
	25.0 – 29.9	1.58 (1.13-2.20; 0.007)
	≥ 30	2.14 (1.43-3.19; 0.0002)
Gestational diabetes		0.88 (0.62-1.23; 0.46)
Vaginal microbiological cultures	Absence of lactobacilli	2.04 (1.49-2.80; <0.0001)
	Enterobacteria	2.38 (1.74-3.24; <0.0001)
	Group B *Streptococcus*	0.89 (0.60-1.29; 0.54)
	Anaerobes	0.88 (0.41-1.78; 0.74)
	G. vaginalis	1.42 (0.64-3.30; 0.40)

Figure 2. Multivariable analysis of risk factors for prelabor rupture of membranes at term (TPROM), threatened preterm birth and preterm labor (PTL), and preterm premature rupture of membranes (PPROM).

Adjusted odds ratios were calculated using any variables associated with the outcome with a p-value <0.2 from the univariate analysis, excepting microbiological variables which were all included. *S. aureus*, *S. anginosus* and *Candida* spp. did not show significant associations with the outcomes and are not represented. aOR: adjusted odds ratio; CI: confidence interval.

A

		aOR (95% CI; p-value)
Gestational age at admission (weeks)	≥ 32	1
	28-31	0.54 (0.37-0.80; 0.002)
	<28	0.69 (0.47-1.02; 0.06)
Place of birth	France	1
	France, overseas territories	0.92 (0.46-1.84; 0.82)
	Europe	1.14 (0.58-2.31; 0.70)
	North Africa	1.34 (0.88-2.03; 0.17)
	Sub Saharan Africa	1.56 (0.98-2.49; 0.06)
	Asia	2.67 (1.11-7.14; 0.04)
Pregnancy with multiples		4.80 (2.72-9.03; <0.0001)
Body mass index	18.5 – 24.9	1
	< 18.5	1.22 (0.65-2.30; 0.54)
	25.0 – 29.9	1.25 (0.83-1.89; 0.28)
	≥ 30	1.31 (0.82-2.11; 0.26)
Gestational diabetes		1.92 (1.24-3.02; 0.004)
Vaginal microbiological cultures	Absence of lactobacilli	1.39 (0.96-2.03; 0.08)
	Enterobacteria	1.64 (1.15-2.37; 0.007)
	Group B *Streptococcus*	0.86 (0.55-1.35; 0.52)
	Anaerobes	2.63 (0.86-9.83; 0.11)
	G. vaginalis	0.40 (0.10-1.29; 0.15)

B – Singleton pregnancies

		aOR (95% CI; p-value)
Gestational age at admission (weeks)	≥ 32	1
	28-31	0.51 (0.34-0.77; 0.001)
	<28	0.71 (0.47-1.06; 0.09)
Place of birth	France	1
	France, overseas territories	0.82 (0.40-1.67; 0.58)
	Europe	1.27 (0.62-2.67; 0.52)
	North Africa	1.32 (0.85-2.04; 0.22)
	Sub Saharan Africa	1.43 (0.88-2.34; 0.16)
	Asia	2.64 (1.09-7.13; 0.04)
Body mass index	18.5 – 24.9	1
	< 18.5	1.09 (0.57-2.12; 0.79)
	25.0 – 29.9	1.24 (0.81-1.90; 0.32)
	≥ 30	1.27 (0.78-2.08; 0.33)
Gestational diabetes		1.78 (1.14-2.83; 0.01)
Vaginal microbiological cultures	Absence of lactobacilli	1.54 (1.05-2.28; 0.03)
	Enterobacteria	1.62 (1.11-2.36; 0.01)
	Group B *Streptococcus*	0.88 (0.55-1.41; 0.59)
	Anaerobes	2.58 (0.84-9.72; 0.12)
	G. vaginalis	0.37 (0.10-1.22; 0.12)

Figure 3. Multivariable analysis of risk factors for preterm birth in women admitted for

threatened preterm birth, preterm labor, and preterm premature rupture of membranes

for (A) all pregnancies and (B) singleton pregnancies.

Adjusted odds ratios were calculated using any variables associated with the outcome with a p-

value <0.2 from the univariate analysis, excepting microbiological variables which were all

included. *S. aureus*, *S. anginosus* and *Candida* spp. did not show significant associations with

the outcomes and are not represented. aOR: adjusted odds ratio; CI: confidence interval.

Table 1. Cohort demographic and clinical characteristics.

	Control n=1,068	TPROM n=477	PTL n=495	PPROM n=436	p-value
Maternity ward, n (%)					
Center 1	682 (63.9)	262 (54.9)	246 (49.7)	163 (37.4)	NT
Center 2	386 (36.1)	215 (45.1)	144 (29.1)	216 (49.5)	NT
Center 3	0	0	105 (21.2)	57 (13.1)	NT
Age, median (IQR), years	33 (30-36)	33 (30-36)	32 (29-35)	33 (30-36)	<0.0001
Gestational age at inclusion, median (IQR), weeks	36 (35-36)	39 (38-40)	29 (26-32)	32 (27-35)	<0.0001
Maternal place of birth, n (%)					
France	565 (53.1)	211 (46.2)	185 (38.4)	142 (33.2)	<0.0001
France, overseas territories[a]	29 (2.7)	13 (2.9)	30 (6.2)	25 (5.8)	0.001
Europe, other countries	81 (7.6)	44 (9.6)	26 (5.4)	24 (5.6)	0.04
North Africa	206 (19.3)	111 (24.3)	95 (19.7)	115 (26.9)	0.004
Sub-Saharan Africa	89 (8.4)	28 (6.1)	106 (22.0)	84 (19.6)	<0.0001
Asia	35 (3.3)	29 (6.3)	12 (2.5)	20 (4.7)	0.01
Others	60 (5.6)	21 (4.6)	28 (5.8)	18 (4.2)	0.58
Missing	*3 (0.3)*	*20 (4.2)*	*13 (2.6)*	*8 (1.8)*	NT
Gestity, n (%)					
1	317 (29.7)	215 (45.1)	156 (31.5)	148 (33.9)	<0.0001
2	341 (32.0)	127 (26.6)	146 (29.5)	102 (23.4)	0.006
≥3	409 (38.3)	135 (28.3)	193 (39.0)	186 (42.7)	<0.0001
Missing	*1 (0.1)*	*0*	*0*	*0*	NT
Parity, n (%)					
0	477 (44.7)	317 (66.5)	267 (53.9)	223 (51.1)	<0.0001
1	385 (36.1)	105 (22.0)	150 (30.3)	115 (26.4)	<0.0001
≥2	205 (19.2)	55 (11.5)	78 (15.8)	98 (22.5)	<0.0001
Missing	*1 (0.1)*	*0*	*0*	*0*	NT
Multiple pregnancy, n (%)	19 (1.8)	1 (0.2)	78 (15.8)	53 (12.2)	<0.0001
History of preterm delivery, n (%)	111 (10.4)	18 (3.8)	100 (20.2)	73 (16.7)	<0.0001
Missing	*3 (0.3)*	*0*	*0*	*0*	NT
Body mass index, median (IQR)	22.4 (20.4 - 25.7)	22.4 (20.2 - 25.2)	22.9 (20.3 - 26.6)	24.5 (21.5 - 28.8)	<0.0001
Underweight (<18.5), n (%)	57 (5.3)	36 (7.5)	42 (8.6)	23 (5.4)	0.05
Normal weight (18.5-24.9), n (%)	685 (64.2)	314 (66.0)	278 (56.7)	204 (47.8)	<0.0001
Overweight (25-29.9), n (%)	213 (20.0)	88 (18.5)	103 (21.0)	112 (26.2)	0.02
Obese (≥30), n (%)	112 (10.5)	38 (8.0)	67 (13.7)	88 (20.6)	<0.0001
Missing	*1 (0.1)*	*1 (0.2)*	*5 (1.0)*	*9 (2.1)*	NT
Gestational diabetes, n (%)	197 (18.5)	79 (16.6)	78 (15.8)	94 (21.6)	0.10
Missing	*1 (0.1)*	*0*	*0*	*0*	NT

	Control n=1,068	TPROM n=477	PTL n=495	PPROM n=436	p-value
Hypertensive disease, n (%)	41 (3.8)	6 (1.3)	22 (4.4)	5 (1.1)	0.001
Missing	*1 (0.1)*	*0*	*0*	*0*	NT
Smoking during pregnancy, n (%)	57 (5.4)	27 (5.7)	37 (7.5)	29 (6.7)	0.36
Missing	*3 (0.3)*	*0*	*4 (0.8)*	*3 (0.7)*	NT
Fetal growth restriction, n (%)	29 (2.7)	5 (1.0)	46 (9.3)	39 (8.9)	<0.0001
Missing	*1 (0.1)*	*0*	*0*	*0*	NT
Gestational age at delivery, median (IQR), weeks	40.0 (39.0-40.7)	39.7 (39.0-40.6)	37.7 (34.3-39.1)	35.0 (30.6-36.4)	<0.0001
Extremely preterm (<28 weeks), n (%)	0	0	37 (7.6)	65 (15.0)	<0.0001
Very preterm (28-31 weeks), n (%)	0	0	34 (6.9)	76 (17.5)	<0.0001
Moderately preterm (32-36 weeks), n (%)	0	0	137 (27.9)	235 (54.1)	<0.0001
Term (≥ 37 weeks), n (%)	1066 (100)	477 (100)	283 (57.6)	58 (13.4)	<0.0001
Missing	*2 (0.2)*	*0*	*4 (0.8)*	*2 (0.5)*	NT
Mode of delivery, n (%)					
Vaginal, spontaneous	764 (71.7)	324 (68.0)	305 (62.1)	246 (56.8)	<0.0001
Vaginal, induced	118 (11.1)	86 (18.0)	87 (17.7)	56 (12.9)	0.0002
Planned Caesarean	61 (5.7)	5 (1.0)	38 (7.8)	60 (13.9)	<0.0001
Caesarean in labor	123 (11.5)	62 (13.0)	61 (12.4)	71 (16.4)	0.09
Missing	*2 (0.2)*	*0*	*4 (0.8)*	*3 (0.7)*	NT
Interval between ROM and delivery, median (IQR), days	0 (0-1)	1 (1-2)	0 (0-1)	4 (1-17)	<0.0001
Missing	*21 (2.0)*	*3 (0.6)*	*42 (8.5)*	*13 (2.9)*	NT
Antibiotics between inclusion and delivery[b], n (%)	139 (13.0)	209 (43.8)	146 (29.6)	353 (81.0)	<0.0001
Missing	*0*	*0*	*2 (0.4)*	*0*	NT
Live births, n (%)	1083 (100)	478 (100)	561 (98.9)	481 (99.0)	NT
Missing	*3 (0.3)*	*0*	*6 (1.2)*	*2 (0.4)*	NT

[a] Including territories in South America (French Guiana), the Atlantic Ocean (Guadeloupe, Martinique, Saint-Martin, Saint-Barthélémy, Saint-Pierre-et-Miquelon), the Indian Ocean (Reunion Island, Mayotte), the Pacific Ocean (New Caledonia, French Polynesia, the Wallis and Futuna Islands), and the French Southern and Antarctic Territories.

[b] Including intrapartum antibiotic prophylaxis and excluding non-systemic antibiotic therapy.

IQR: interquartile range; NT: not tested; PPROM: preterm premature rupture of membranes; PTL: threatened preterm birth or preterm labor; ROM: rupture of membranes; TPROM: prelabor rupture of membranes at term.