Unveiling the hidden interactome of CRBN molecular glues with chemoproteomics

Kheewoong Baek[1,2,†], Rebecca J. Metivier[1,2,†], Shourya S. Roy Burman[1,2], Jonathan W.

Bushman[1,2], Ryan J. Lumpkin[1,2], Dinah M. Abeja[1], Megha Lakshminarayan[1], Hong Yue[1,2],

Samuel Ojeda[1,2], Alyssa L. Verano[1,2], Nathanael S. Gray[3], Katherine A. Donovan[1,2,*], Eric S.

Fischer[1,2,4,**]

1. Department of Cancer Biology, Dana–Farber Cancer Institute, Boston, Massachusetts 02215,

USA

2. Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical School,

Boston, Massachusetts 02115, USA

3. Department of Chemical and Systems Biology, ChEM-H and Stanford Cancer Institute,

Stanford Medical School, Stanford University, Stanford, CA, 94305, USA 4. Lead Contact

[†] These authors contributed equally

[*] Correspondence: kdonovan@crystal.harvard.edu (KAD); Eric_Fischer@dfci.harvard.edu (ESF)

SUMMARY

Targeted protein degradation and induced proximity refer to strategies that leverage the recruitment of proteins to facilitate their modification, regulation or degradation. As prospective design of glues remains challenging, unbiased discovery methods are needed to unveil hidden chemical targets. Here we establish a high throughput affinity purification mass spectrometry workflow in cell lysates for the unbiased identification of molecular glue targets. By mapping the targets of 20 CRBN-binding molecular glues, we identify 298 protein targets and demonstrate the utility of enrichment methods for identifying novel targets overlooked using established methods. We use a computational workflow to estimate target confidence and perform a biochemical screen to identify a lead compound for the new non-ZF target PPIL4. Our study provides a comprehensive inventory of targets chemically recruited to CRBN and delivers a robust and scalable workflow for identifying new drug-induced protein interactions in cell lysates.

Keywords: IMiD, molecular glue, E3 ligase, cereblon, targeted protein degradation, proteomics, degrader, ubiquitin, PROTAC, PPIL4

INTRODUCTION

Targeted Protein Degradation (TPD) represents a promising therapeutic approach to

remove disease-associated proteins from cells[1,2]. The process of TPD involves hijacking the

ubiquitylation machinery for the covalent attachment of ubiquitin molecules to a desired protein

of interest, which in turn leads to degradation by the proteasome[3,4]. Ubiquitin-mediated TPD

utilizes two types of small molecules, molecular glues[5] and heterobifunctional degraders (also

known as PROteolysis Targeting Chimeras, or PROTACs)[3], both of which chemically induce

ternary complex formation between a protein target and a ubiquitin E3 ligase, followed by

proximity-driven ubiquitylation and subsequent degradation.

Despite the rapid growth of TPD as a therapeutic strategy, the discovery and

development of effective degraders remains challenging. Heterobifunctional degraders rely on

a linker to connect two binding warheads: one for the ligase and one for the protein of

interest[3,6,7]. Although this modular design offers the flexibility to target any protein with a known

binder, the resulting molecules often possess poor drug-like properties due to their large size.

Molecular glues present a possible alternative due to their small size and improved drug-like

properties. However, as they lack binding to their target protein and instead enhance protein-

protein interactions (PPI) between a ligase and substrate, rational design of molecular glues is

far more challenging[8]. Over the last decade, the discovery of new molecular glue degraders has

largely relied on serendipity through phenotypic screening of large libraries of molecules and

retrospective identification of their degradation targets. Although FDA-approved molecular glue

degraders exist, including the immunomodulatory drugs (IMiDs) thalidomide, lenalidomide, and

pomalidomide, they have all been characterized as molecular glues in retrospect after their

serendipitous discovery. IMiD molecules bind to the CUL4-RBX1-DDB1-CRBN (CRL4[CRBN]) E3

ligase[5,9-13] creating a favorable surface for new proteins (neo-substrates) to bind for induced

degradation. Since this discovery, significant efforts into the design and screening of new IMiD

analogs have revealed up to 50 neo-substrates in the public domain, all carrying a glycine

containing β-hairpin structural degron[5,11,14-22]. Remarkably, computational modeling of the

AlphaFold2 (AF2) structures available in the Protein Data Bank (PDB) suggest that we are just

scraping the surface of what is targetable by these molecules[14].

Given the mechanism of action (MoA) of degraders, global chemoproteomics has proven

to be an effective tool for the identification of protein degradation targets[19,23-25]. Using this

approach, the target space of degraders for multiple therapeutic target classes have been

extensively mapped for tractable targets, including kinases[23,26,27], bromo-domains[28-30], HDACs[31]

and zinc finger (ZF) transcription factors[19]. Although this method has greatly expanded the

repertoire of known targets, limited sensitivity has restricted the ability to identify proteins with

low expression levels without screening libraries of cell lines or using target enrichment

methods. This approach also remains blind to a key aspect of these molecules: the identification

of proteins that are recruited to the ligase but do not ultimately get degraded. Such "non-

degrading glue" targets may be subject to poor lysine accessibility, lack of degradative ubiquitin

chain formation[32], high deubiquitinase activity, poor proteasome access, or other resistance

mechanisms. However, these targets still represent important therapeutic targets if these factors

can be overcome to convert silent molecular glues into molecular glue degraders or functional

modulators of the target.

Methods to identify chemically induced protein-protein interactions include

immunoprecipitation mass spectrometry (IP-MS)[33] and proximity labeling approaches coupled to

mass spectrometry[34,35]. IP-MS approaches have been employed for the identification of direct

protein interactions, whereas proximity labeling approaches are commonly employed for the

mapping of proximity interactomes in cells and in vivo[34,36-38], enabling the identification of protein

interactions within a 10-30 nm radius of the epitope tagged protein of interest[34,36,37,39]. Although

these in-cell methods have demonstrated successful identification of chemically induced

interactions, they often require extensive fine tuning of various factors including noise,

sensitivity, variability and scalability.

In this study, we establish a simple, robust and sensitive workflow to facilitate high

throughput discovery of degrader-induced protein-protein interactions and develop them into

selective tools and therapeutic candidates. We use this method to build a comprehensive

inventory of 298 distinct protein targets recruited to CRBN including many new zinc finger (ZF)

transcription factors and proteins from new target classes, including RNA-recognition motif

(RRM) domain proteins. We evaluate the binding potential of these targets through structural

alignment with IMiD-bound CRBN and performed biochemical and structural validation studies

on a series of non-ZF targets. We then screened a library of ~6000 IMiD analogs against a

novel non-ZF target, PPIL4, identifying a selective lead degrader molecule, thereby presenting a

blueprint for the effective discovery of novel molecular glue degraders.

RESULTS

Unbiased identification of degrader-induced interactors in-lysate

To establish a workflow for the identification of chemically induced protein-protein

interactions, we set out to simplify traditional IP-MS methods. We hypothesized that we could

create a controlled environment with reduced biological variability and enhanced scalability by

establishing a workflow in cell lysate using spiked in recombinant protein as the bait. Our

workflow harnesses small molecule degrader-induced ternary complex formation in cell lysate

using recombinant FLAG-tagged CRBN in complex with DDB1 excluding the BPB domain (ΔB),

which prevents CUL4 interaction to inhibit ubiquitylation of the recruited target, with a small

molecule degrader. After incubation, we enrich with a highly selective antibody for the FLAG

epitope tag followed by label free quantitative proteomic assessment to identify interactors

(**Figure 1A**). To benchmark and explore the viability of this approach for identification of protein-

protein interactions, we selected two representative degrader molecules that have been

thoroughly profiled in published reports – pomalidomide, a IMiD molecular glue[19,40] and SB1-G-

187, a kinase-targeted heterobifunctional degrader[26] (**Figure 1B**). We profiled these two

molecules across two different cell lines, MOLT4 and Kelly, selected for their orthogonal

expression profiles and broad coverage of known CRBN neo-substrates including IKZF1/3

(MOLT4) and SALL4 (Kelly)[19]. The pomalidomide screen revealed 11 different enriched proteins

across these two cell lines (9 in MOLT4 and 4 in Kelly cells), which revealed three novel targets,

ASS1, ZBED3 and ZNF219 (**Figure 1C, Figure S1A-B, Table S1**). We then validated

recruitment of these three novel targets to CRBN using dose response immunoblot or TR-FRET

analysis (**Figure S1C-D**).

To assess the overlap of these enriched targets with published degradation data for

pomalidomide, we performed a hit comparison with publicly available global proteomics data

(http://proteomics.fischerlab.org), which includes ten independent pomalidomide treatments

spanning HEK293T, Kelly and MOLT4 cell lines (**Figure 1D**). Like our enrichment data, the

global degradation data also maps 11 targets as degradable, however only 4 of these targets

(IKZF1, IKZF3, ZFP91 and SALL4) overlap with those that we see enriched in this dataset. The

SB1-G-187 kinase degrader screen identified 18 enriched targets across the two cell lines (16 in

MOLT4 and 7 in Kelly cells) including multiple non-protein kinase targets which raised the

question of how these proteins are being recruited to CRBN by a kinase degrader (**Figure 1E,**

Figure S1E-F, Table S1). Assessment of the non-kinase targets revealed that several are

known to form complexes with different kinases, such as TAB1 and TAB2 which form a

functional kinase complex with MAP3K7[41,42], and UNC119 which binds to myristoylated SRC to

regulate cellular localization[43]. This data suggests that these non-kinases are being recruited to

CRBN through piggybacking on their kinase binding partners. Of the other recruited non-kinase

targets, ZBED3 is also identified in the pomalidomide treatment suggesting recruitment through

the IMiD handle of the degrader and SDR39U1 was reported as a non-kinase target in a

chemoproteomics profiling study of kinase inhibitor probes[44]. Next, we assessed the differences

and overlap in hits between publicly available degradation data in MOLT4 and Kelly cells for

kinase-targeted heterobifunctional degrader, SB1-G-187, and enrichment data (**Figure 1E**). We

found 6 overlapping hits - all protein kinases - including CDK1, IRAK1, LCK, LYN, MAP3K7 and

SRC. Like the pomalidomide data, we observe similar numbers of proteins identified in either

degradation data (15) or enrichment data (12), demonstrating that these two methods

complement each other to expand the target scope of these molecules. Together, the data

collected for these two degrader molecules demonstrate the value of our workflow for identifying

chemically induced protein-protein interactions invisible to degradation assays, while also

highlighting opportunities for improving sensitivity.

Mapping the interactomes of IMiD molecular glue degraders

Next, using the functional enrichment method as a basis, we set out to optimize and

address the critical need for sensitivity and high throughput. IP-MS experiments typically require

labor-intensive sample preparation steps which create a significant source of variability and lead

to high background and false positive rates while also placing limits on the number of samples

that can be prepared in parallel. To address these limitations, we automated the enrichment and

sample preparation procedures to enable effective mapping of interaction targets across

libraries of molecules at scale. We incorporated a cost effective Opentrons OT-2 liquid handling

platform to automate the sample preparation process from addition of all immunoprecipitation

components to tryptic digestion for 96 samples in parallel (**Figure 2A**). To address throughput

and depth of the proteomics workflow, we took advantage of recent updates in instrumentation

(timsTOF Pro2, Bruker) and acquisition methods (diaPASEF)[45] that allow for significant

improvements in sensitivity (**Figure 2A**). In contrast to the data-dependent acquisition (DDA)

data collected in our proof-of-concept analysis (**Figure 1**), diaPASEF measures peptides by

systematically sampling all precursor ions within a specified m/z range regardless of their

abundance which enhances the reproducibility and depth of peptide coverage to allow for more

accurate and robust quantification of peptides in complex samples. Comparison of the average

numbers of proteins and peptides quantified between the DDA collection (**Figure 1A**) and

diaPASEF collection (**Figure 2A**) revealed a >5-fold increase in proteins and a ~9-fold increase

in peptides quantified (**Figure S2A-B**) confirming a significant improvement in depth and

sensitivity.

Work over the last several years has led to the identification of a growing list of ~50

neo-substrates that are recruited to CRBN by IMiD analogs for chemically induced

degradation[14]. Validated targets include a large number of C_2H_2 zinc finger (ZF) transcription

factors such as IKZF1/3[5], ZFP91[18], or SALL4[15,19], but only a few non-ZF proteins such as G1 to

S phase transition protein 1 (GSPT1)[11] and casein kinase 1 alpha (CK1α)[12,17]. These targets do

not possess any similarity, but instead all share a common structural CRBN binding motif

consisting of an 8-residue loop that connects the two strands of a β-hairpin and has a glycine at

the sixth position (G-loop)[11,12,18]. Remarkably, a recently reported analysis of available

AlphaFold2 (AF2) predicted structures for proteins in the human proteome uncovered over

2,500 proteins that harbor a G-loop potentially compatible with IMiD-recruitment to CRBN, with

C_2H_2 ZF proteins revealing themselves as the most prevalent domain class, aligning with the

dominance that this class has amongst the experimentally confirmed targets[14]. Due to the

extensive range of proteins that are predicted to be chemically recruitable to CRBN, we asked

how many of these proteins are already targeted by existing chemistry, but not yet identified due

to lack of sensitivity of existing methods. To explore the range of proteins chemically recruited to

CRBN, we screened a curated library of 20 different IMiD analogs through our automated

lysate-based IP workflow. We assembled this library to incorporate a broad range of IMiD-based

scaffolds including the parental FDA-approved IMiDs (thalidomide, lenalidomide,

pomalidomide)[46,47], where there is a high value to identifying new targets for drug repurposing

efforts. We included a series of IMiD analogs that are undergoing clinical trials (CC-220, CC-

92480, CC-90009)[48-50] and molecules that have demonstrated promiscuity (FPFT-2216, CC-

122)[51,52]. Finally, we included a series of in-house synthesized scaffolds developed in the

context of targeting Helios (IKZF2)[53] or part of an effort to diversify IMiDs with the addition of

fragments on an extended linker (**Figure 2B**). We screened this library at 1 µM concentration

across MOLT4 and Kelly cell lines (including a second 5 µM concentration for pomalidomide)

and identified proteins that were enriched in the degrader compared to DMSO control IP

treatment (**Figure 2C, Figures S3-4, Tables S2-3**). Using significance cutoffs of fold change

(FC) >1.5, P-value <0.001 and combining the data from both cell lines, we identified a total of

298 enriched proteins (**Table S4**). We rationalized that the likelihood of observing the same

proteins enriched as false positives across multiple treatments with similar IMiD analog

molecules is low and therefore used 'frequency of enrichment' as a measure of confidence. We

observed 102 proteins enriched in at least three independent IPs, and each of the top 5 proteins

(PATZ1, ZBED3, WIZ, IKZF2 and ASS1) enriched in more than 20 independent IPs across the

database (**Figure 2D, Table S4**). Surprisingly, although published reports have confirmed

degradation of PATZ1, WIZ and IKZF2, none of these top 5 enriched proteins regularly feature

amongst those proteins that we commonly see reported in existing unbiased screens of IMiD-

based molecules indicating the orthogonal data generated by this profiling method, identifying

targets that might otherwise be overlooked. ZBED3 and ASS1 showed frequent enrichment

across our database without any prior reporting of degradation, even at concentrations up to 5

µM of pomalidomide (**Figure S2C, Table S4**), suggesting the first reported examples of targets

that are chemically glued to CRBN but lack productive degradation, thus emphasizing the

benefit of alternative binding focused approaches for target identification. Also, important to note

is that the new targets identified in this study are not only targets of new IMiD analogs but are

also identified as targets of IMiDs in clinical trials and with FDA approval.

To assess the fraction of newly identified IMiD targets, we compared the 298 enriched

proteins to a list of literature reported targets and discovered an overlap of only 28 targets. We

identified 270 novel targets and found only 22 targets were reported in the literature but not

identified as hits in our study (**Figure 2E, Table S4**)[14]. Considering the prevalence of C_2H_2 ZF

transcription factors amongst reported IMiD targets, we asked whether this dominance holds

true across our extended list of targets. To assess this, we extracted superfamily, family and

domain information from curated databases including InterPro[54,55], Uniprot[56] and Superfamily[57]

to categorize the targets based on studied features (**Figure 2F-G, Table S4**). Of the 298 targets

identified, after C-terminal domain classification, the C_2H_2 ZF superfamily represents the largest

segment, accounting for >14% of the targets in the top 10 enriched superfamilies. This is

followed by RNA-recognition motif domain proteins (RRM, >13%) and nucleotide-binding alpha-

beta plait domain superfamilies (α-β plait domain, >12%). Notably, protein kinase-like domain

proteins also features on this top 10 list of superfamilies (kinase-like domain, >6.4%) which

aligns with our knowledge that kinases can be targets of IMiD molecular glues (eg, CSNK1A1 or

WEE1)[12,17,58] and suggests that molecular glues may be a viable alternative to PROTACs, which

are currently widely explored for kinase targeting. Exploration of the top 10 domain

classifications across the dataset shows a similar trend with C_2H_2 ZF, RRM, ZF, protein kinases

and BTB/POZ domains showing the highest representation across the targets identified (**Figure**

2G).

This dataset builds upon previous identifications of protein kinases as targets of IMiD

molecules[17,22,59], and further extends the kinase list adding CDK7, IRAK1 and TBK1 as novel

putative molecular glue targets. It also broadens the scope of tractable targets by introducing

multiple new families as targetable by CRBN-based molecular glues, illustrating the extensive

potential of these molecules. Through the application of our unbiased target enrichment

workflow, we have significantly increased the number of experimentally detected IMiD targets,

expanding beyond the C_2H_2 ZF protein family to a wide range of protein families including

protein kinases and proteins involved in RNA metabolism.

Zinc finger transcription factors enriched among targets

To validate the 270 previously unreported targets, we sought to establish a

computational screening pipeline to score the compatibility of targets for recruitment to CRBN.

Structural studies on IMiD-mediated CRBN neo-substrates, both natural and designed, have

established the common G-loop motif that is recognized by the CRBN-IMiD complex[12,60]. We

used MASTER[61] to mine the AF2 database[62] for proteins containing G-loops with similar

backbone architecture to the G-loop in known neo-substrate CSNK1A1 (PDB: 5FQD, aa35-42)

resulting in a set of 46,040 loops across 10,926 proteins (**Figure 3A**).

Due to structural constraints, not all these proteins are compatible with CRBN. To

identify CRBN-compatible proteins, we first extracted domains containing the G-loops based on

domain definitions from DPAM, a tool that parses domains from AF models based upon

predicted aligned errors (PAE) and evolutionary classification[63]. Next, we aligned the domains to

our reference CSNK1A1-IMiD-CRBN-DDB1ΔB structure based on the G-loop and calculated a

clash score. We used the van der Waals force term for interchain contacts in Rosetta's low-

resolution mode[64] to obtain a side-chain independent clash estimate. Out of 16 known neo-

substrates with validated G-loops (**Table S5, Figure S5**), 15 had clash scores below 2, while

ZNF654[40] had a score of 172, indicating a minor clash. The clash was caused by a low

confidence region in the AF2 structure and could be resolved by relaxing the complex with

Rosetta (**Figure S6A**)[65]. On the other hand, a protein with no evidence supporting it being a

neo-substrate, PAAF1, had a major clash with a score of 1,551 which could not be resolved by

relaxation (**Figure S6B**). Based on these examples, and analysis of the clash scores of all hit

proteins containing a clear structural G-loop in AF2 (**Figure S6C)**, we filtered out domains with

scores greater than 200 resulting in a list of 14,189 loops across 6,018 proteins with

nonexistent, or marginal clashes with CRBN (**Figure 3A, Table S5**).

Of the 298 total enriched candidates,199 were found to have a clear structural G-loop

with 162 having a clash score below 200. Given the high proportion of ZF proteins identified as

targets across this enrichment database (**Figure 2F**), we mapped the fold change in enrichment

for proteins with an annotated ZF domain across all 20 degraders for both MOLT4 (**Figure 3B**)

and Kelly cells (**Figure 3C**). Across these two cell lines, we identified 19 previously reported and

28 new neo-substrates as chemically recruited to CRBN. We then used our G-loop database to

inform on which of these targets have a tractable G-loop and found that only five of the 57

identified targets do not contain a structural G-loop (**Figure 3C, Table S5**). Given what we know

about the recruitment and binding of CRBN neo-substrates, targets usually bind through a

dominant structural hairpin. Since we do not have validated degron information for all these ZF

targets, we assumed that the G-loop with the lowest clash score has the highest likelihood to

bind and therefore proceeded with evaluation of a single G-loop for each target. To gauge how

the clash scores for these ZF targets compare to all hits in the G-loop database, we compared

the clash scores for our ZF targets to those of all hits (**Figure S6C**) demonstrating a pronounced

trend towards lower scores for ZF targets suggesting fewer unfavorable interactions (**Figure**

S5A-B). Notably, when we explored the ZF hits with higher clash scores (>10) and >3 hit

frequency, we realized that almost all of these have a reported association with at least one of

the validated hits – ZMYND8 (cs 455, binds to ZNF687), and RNF166 (cs 17, binds to

ZNF653/ZBTB39/ZNF827) – which also offers the possibility that these proteins could be

collateral targets, recruited via piggybacking on their binding partners, the direct binders (**Table**

S4). Finally, we compared ZF targets across the two cell lines as an additional means for

validation, and found 10 overlapping proteins, 6 of which are novel recruited targets (ZBED3,

MNAT1, MTA2, ZBTB44, TRIM28) (**Figure 3D**).

There are many factors to take into consideration when looking to predict target

degradability, such as ternary complex formation[26,31,66] and target ubiquitylation[12,32,67-69], and

multiple studies have placed an emphasis on exploring their role in driving productive

degradation[70,71]. For degrader-induced degradation to occur, a ternary complex consisting of

ligase-degrader-target needs to form for proximity-mediated ubiquitin transfer to the target

protein. Because ternary complex formation is necessary for successful protein degradation, we

set out to explore the relationship between ternary complex formation and degradation for ZF

targets identified in this study. We focused our evaluation on the parental IMiD molecules which

have been subjected to degradation target profiling using unbiased global proteomics analysis

across a panel of four cell lines (SK-N-DZ, Kelly, MM.1S, hES)[19]. Comparison of the enriched

ZF targets to the published degradation data shows a consistent trend across the three IMiDs

where only ~30% of the enriched targets that were quantified in global proteomics studies were

degraded, with ~ 60% of the targets quantified but not reported as degraded (**Figure 3E, Table**

S5). The data was then grouped to allow a global comparison of the enriched versus degraded

IMiD targets. The comparison revealed that of the 31 ZF targets enriched across these three

molecular glues, only 11 of the 29 proteins quantified in global proteomics experiments were

found to be degraded (**Figure 3F, Table S5**). 18 proteins were quantified in global proteomics

but were not identified as degraded. This prompted us to question whether these targets were

resistant to degradation by IMiDs and their analogs, or if they were not identified as degraded

due to experimental limitations such as inadequate sensitivity to detect minor changes in protein

abundance, rapid protein turnover or suboptimal experimental conditions. We found that

although several of the targets (WIZ, PATZ1, ZNF687, ZMYM2 and HIC2) were not reported as

degraded in Donovan et al.[19], they have since been reported as degraded in other published

studies[40,72,73] confirming that IPs provide a complementary approach able to overcome

limitations in sensitivity. The absence of degradation data for the remaining targets could imply

that these targets are resistant to degradation, or similar to the above proteins, the appropriate

degradation experiment has yet to be performed. These data demonstrate that our IP workflow

provides a significant advantage over global proteomics analysis by enabling selective isolation

and enrichment of targets that may be below the change in abundance threshold for consistent

identification with global proteomics approaches.

IMiD derived molecular glues recruit hundreds of non-zinc finger proteins

316   The largest target class of CRBN neo-substrates today are ZF containing proteins,

however, of the ~20,000 proteins in the human proteome, ZF containing proteins only make up

a relatively small proportion with about ~1700 ZF proteins reported[74]. So far, only a handful of

targets are reported to lack a ZF motif, which includes GSPT1[11], CK1α[12,17], PDE6D[75], and

RAB28[75]. With this in mind, we examined our list of targeted proteins with a focus on those that

do not contain a reported ZF domain and found 251 non-ZF proteins enriched across the IP

dataset (**Figure 4A, Table S4**). These non-ZF proteins include a wide range of families such as

protein kinases (IRAK1, TBK1, CDK7), RNA recognition motif proteins (ELAVL1, PPIL4, CSTF2,

RBM45), metabolic enzymes (ASS1, PAICS, ACLY, CS, ACADVL), translational proteins

(MARS1, ETF1, EEF1E1, EIF4B) and more spanning different biological pathways. To assist in

establishing confidence in some of these targets, we performed a comparison of the non-ZF

targets enriched in the two tested cell lines and found 39 targets were identified in both MOLT4

and Kelly cells, including the four above mentioned targets (**Figure 4A-B**). We then assessed

the AF2 structures of each of these 39 proteins and found that almost all of them (33/39) contain

a structural G-loop (**Figure 4B, Figure S5, Table S5**).

Given the large number of non-ZF targets identified in this study and the lack of

emphasis in the public domain with regards to non-ZF CRBN neo-substrates, we selected a

series of non-ZF proteins for further experimental validation. Firstly, to demonstrate that these

neo-substrates are directly recruited to CRBN, we examined ternary complex formation using

recombinant purified proteins. Using two of the more promiscuous molecular glues,

pomalidomide and FPFT-2216, we tested previous reported degradation targets PDE6D,

RAB28 and DTWD1, along with a newly discovered target PPIL4. Indeed, PDE6D, DTWD1,

and PPIL4 formed compound dependent ternary complex with CRBN at varying effective

concentrations (**Figure 4C**). However, RAB28, which was previously reported to be degraded

by IMiDs[19] and FPFT-2216[75], did not show any evidence for direct binding to CRBN using

purified proteins. Since RAB28 has previously been reported as a CRBN neo-substrate and

consistently scored across our enrichment study, we explored whether there was any evidence

suggesting that RAB28 could be a collateral target. Exploration of protein-protein interaction

databases including BioPlex[33] and STRING-DB[76] revealed that RAB28 is known to bind to two

validated IMiD-CRBN neo-substrates PDE6D and ZNF653 (**Table S4**), suggesting that RAB28

is likely an indirectly recruited target. These data demonstrate that in addition to identifying

direct binders, we can also identify indirect binding partners that may be simultaneously

recruited together with direct binding neo-substrates. Given that targeted protein degradation

requires not only recruitment to CRBN, but also CRBN mediated ubiquitin transfer for

degradation, we also monitored whether the recruited proteins can be ubiquitylated by

CRL4^{CRBN}. In vitro ubiquitylation assays showed robust ubiquitin modification on all 3 recruited

non-ZF proteins in the presence of pomalidomide or FPFT-2216 (**Figure 4D**). In addition, all

three of these targets were degraded in response to IMiD treatment as observed by global

proteomics analysis (**Figure S6D, Table S5**). Using structural G-loop alignments, we then

assessed the potential for each of these three proteins to bind to IMiD-CRBN and found that all

three proteins had a G-loop with a clash score of <200 (**Figure 4E**). However, the aligned clash

score for DTWD1 was relatively high and approaching the upper 200 threshold (cs 198). We

performed relaxation with Rosetta and found that this reduced the clash score to 1.58 by

allowing minor shifts in the overall conformation while retaining the structural G-loop (**Figure**

S6E). This process demonstrates that in some cases, clash scores can be relieved through

minor structural rearrangements using Rosetta relax.

To expand our understanding of the recruitment of non-ZF targets, we determined

cryo-EM structures of CRBN-DDB1ΔB-FPFT-2216 bound to PPIL4 and PDE6D, respectively

(**Figure 4F, Figure S7, Table 1**). The complex structures were both refined to a global

resolution of around 3.4 Å and the quality of the resulting maps were sufficient to dock the

complex components, but the flexibly tethered PPIL4 resulted in a lower local resolution. We

were able to observe PPIL4 engagement with FPFT-2216-CRBN via its Gly278 harboring G-

loop as expected from the G-loop alignment, as well as for PDE6D via its Gly28 G-loop.

Furthermore, overall density allowed fitting of FPFT-2216 in bulk although the reduced

resolution in that region did not permit exact positioning of the molecule. Nevertheless, we were

able to see that the glutarimide ring engages CRBN's binding pocket in a similar manner to

other IMiD molecular glues. The triazole interacts with the backbone of the G-loop, and the

methoxythiophene moiety potentially contacts both the PPIL4 backbone of the G-loop and

Arg273. This suggests that the triazole and the methoxythiophene moieties could provide

specificity elements to FPFT-2216 mediated neo-substrate recruitment. The methoxythiophene

moiety also engaged Arg23 of PDE6D, indicating that FPFT-2216 might derive specificity in

engaging an arginine residue from its neo-substrates. Analysis of the non-ZF targets of FPFT-

2216 revealed several other proteins harboring an arginine or a lysine residue at this sequence

location (PDE6D, SCYL1, RBM45, PPIL4). Finally, we compared the experimental structure to

the AF2 predicted G-loop aligned structure of PPIL4 (**Figure S6F**). The G-loop aligned structure

of PPIL4 presented a clash score of 3.38, which showed the C-terminal region of CRBN around

residue Arg373 to be clashing with PPIL4's loop harboring residue Val250. Although the low

resolution permitted only backbone level fitting of PPIL4, we observed that the cryo-EM

structure revealed a minor shift in the RRM domain of PPIL4 to accommodate this minor clash

suggested in the G-loop aligned structure while retaining overall conformational similarity of the

G-loop (**Figure S6F**). Meanwhile, PDE6D retained overall similar conformation with minor shifts

that did not alter the interaction with CRBN (**Figure S6G**).

These data demonstrate that RRM domain containing proteins represent a new class

of proteins targetable through CRBN dependent molecular glues. Using structural modeling we

increase confidence in these new targets while also providing a reminder that structural analysis

and AF2 predicted structures are static models and although they provide excellent structural

guidance, we need to keep in mind that proteins in solution are flexible and dynamic.

Discovery of new and selective molecular glue for PPIL4

While the proteomics-based screening workflow identifies novel putative CRBN targets

and provides initial chemical matter, it does not necessarily provide the best starting point for

developing a chemical probe or therapeutic due to the limited number of molecules screened.

We hypothesized that this limitation could be overcome by following up proteomics screening

with a target centric screen of a larger CRBN binder library to identify the optimal chemical

starting point. To test this, we set out to identify PPIL4 targeting molecular glues with improved

selectivity and lacking the triazole moiety. We employed an IMiD molecular glue library

consisting of ~6000 compounds of various IMiD analogs that were either synthesized in-house

or purchased externally. We screened this library against PPIL4 using TR-FRET to measure

compound-induced PPIL4 recruitment to CRBN (**Figure 5A**). TR-FRET ratios were obtained by

incubating the library with GFP fused CRBN-DDB1ΔB, biotinylated PPIL4, and Tb-labeled

streptavidin that binds to the biotinylated PPIL4. The library was compared relative to the

positive control, whereby the 520/490 ratio of FPFT-2216 at 10 µM was normalized as 1, and

compounds were tested at 1.66 µM or 3.33 µM to find hits with equal or improved efficacy in

directly recruiting PPIL4 to CRBN-DDB1ΔB. We were able to narrow down the library to two

molecules that performed similar or better than FPFT-2216 (**Figure 5B)**. These lead compounds

were subject to a full titration to assess recruitment efficacy by TR-FRET. Ultimately, after

recognizing one of the two hits was due to autofluorescence, we were able to identify a

molecule, Z6466608628, that produced a higher 520/490 ratio, and a better EC50 of 0.34 µM

compared to FPFT-2216, measured at 1.05 µM in this experiment (**Figure 5C-D**).

To test the efficacy and selectivity of our lead compound, we first performed IP-MS in

comparison with FPFT-2216 in Kelly cell lysate. While FPFT-2216 recruited many proteins,

Z6466608628 selectively recruited PPIL4, along with its binding partner DHX40 (**Figure 5E,**

Table S5). We then performed global proteomics in MOLT4 cells to confirm that Z6466608628

can induce selective downregulation of PPIL4 (**Figure 5F, Table S5**). These data collectively

demonstrate the complete workflow, starting from the identification of a novel non-ZF target

PPIL4 in a chemoproteomics screen, to the discovery of a new PPIL4 selective molecular glue

that would serve as an excellent lead molecule for structural optimization.

DISCUSSION

Targeted protein degradation and induced proximity are part of a rapidly expanding field focused on the development of small molecules that leverage induced neo-protein-protein interactions to drive pharmacology. In this study, we develop and showcase a new workflow for high sensitivity, unbiased target identification of degraders and non-degrading molecular glues, identifying more than 290 targets recruited to CRBN by IMiD-like molecules. We demonstrate that this new approach to target identification can reveal critical insights and new targets that are missed by traditional screening methodologies and provide a blueprint from discovery to optimization and structure guided design of new molecular glue degraders.

Thalidomide and its derivatives, lenalidomide and pomalidomide (IMiDs), have had a checkered past. These molecules have been in use for a variety of indications, on and off, since the 1950's and have experienced perhaps the greatest turnaround in drug history. From devastating birth defects to effective hematological cancer therapy, and more recently, significant investment in utilizing these molecules for TPD-based therapeutics. While a decade of research has slowly uncovered around 50 reported neo-substrates of IMiD's, thousands of proteins harbor G-loops that have the potential for recruitment to CRBN by IMiD molecules. Our simple, cost effective and highly scalable unbiased screening workflow combines whole cell lysate with recombinant Flag-CRBN and degraders to enrich target binders from the complex proteome. Through an IMiD-analog diversity screen across two cell lines, we mapped a significant expansion of the neo-substrate repertoire by identifying 298 proteins recruited to CRBN, with 270 of these being novel targets. Unlike many current high throughput screening workflows that focus on the end point – degradation, this workflow allows us to explore the fundamental first step of proximity induced degradation – recruitment, where we are now able to identify targets that are directly or indirectly recruited to CRBN. This sensitive workflow sheds light onto a previously unchartered element of the molecular glue mechanism of action and establishes insights into how and why certain molecular glues may exhibit higher efficacy than

others. Surprisingly, we discovered targets recruited to CRBN that are resistant to degradation,

demonstrating the first examples of targets being glued to CRBN without productive

degradation. Exploration of two of these targets, ASS1 and ZBED3, does not offer any clues as

to why they are not degraded since both have reported ubiquitylated sites[77]. Numerous

possibilities exist, from these targets being tightly preoccupied by other binding partners,

geometric constraints leading to inaccessibility of lysines, removal of ubiquitylation by

deubiquitinases, or preclusion of the catalytic sites due to size and shape preventing active

ubiquitylation. It is important to note that the non-degrading functions of these molecular glues

may have interesting degradation-independent pharmacology that have not yet been

investigated, thus providing an opportunity for future experimental research.

The comprehensive G-loop database provided us with prefiltered insights as to whether

these targets have the potential to be recruited to CRBN through the currently established

mechanism of G-loop binding. However, although most targets identified in this study do have a

structural G-loop, we do have numerous instances of proteins that do not harbor a G-loop.

Some of these targets do have a structurally similar hairpin motif but are lacking the 'essential'

glycine in position six. Whereas other targets did not have this structural motif at all. These

findings indicate the potential for alternative recruitment mechanisms such as proteins

piggybacking on a direct G-loop carrying target. This concept of collateral (or bystander)

targeting was also demonstrated in a study exploring HDAC degradability, where it was found

that both HDACs, and their known complex binding partners can be degraded[31]. Alternatively,

and perhaps more intriguingly, the potential for recruitment of proteins through a distinct

structural motif suggesting there may be new binding mechanisms that are pending discovery.

The potential capacity for IMiDs to yield interfaces favorable for recruitment of various structural

motifs would considerably expand the diversity CRBN neo-substrates and broaden therapeutic

applications.

Amongst the targets identified in this study, we not only discovered many new C_2H_2 ZF

transcription factor targets but also extended targets beyond C_2H_2 ZF proteins, into additional

classes of proteins such as those containing RNA recognition motif (RRM) domain and kinase

domains, confirming that CRBN is an incredibly versatile ligase and very well suited to hijacking

for TPD applications. We reveal 251 non-ZF targets, a dramatic increase in the breadth and

number of proteins targeted by CRBN from the currently reported targets of less than a dozen.

Direct binding data using TR-FRET on a selection of these targets validates their direct binding

mechanism, and structural characterization further corroborates this binding while validating the

generated G-loop alignment database as a tool to assist prioritization of targets using clash

score assessment. Using the accumulative data, we selected a novel non-ZF neo-substrate,

PPIL4, for additional screening to illustrate the utility of this workflow for prioritization efforts.

After a biochemical ternary complex recruitment screen of around 6,000 IMiD analogs, we

selected a single hit compound and used chemoproteomics to confirm selective recruitment of

PPIL4 to CRBN. Genomic studies have reported that PPIL4 is essential for brain specific

angiogenesis and has implications in intracranial aneurysms[78], and is known to regulate the

catalytic activation of the spliceosome[79]. Thus, this new molecular glue could be of great interest

to target the splicing pathway, in relation to intracranial aneurysms, or in other contexts.

We believe our strategic workflow and comprehensive data package, along with outlining

specific applications of these, provides a valuable resource for the chemical biology, drug

discovery and induced proximity communities. Importantly, the workflow is neither limited to

CRBN nor to TPD, but rather can be applied to any induced proximity application. We expect

the enrichment workflow will provide a blueprint for expansion into target identification for

induced proximity platforms as well as further expansion of targets for protein degraders beyond

molecular glues. Through initial scouting efforts on heterobifunctional degraders and additional

ligases we are confident there are many novel discoveries to be made with already existing

chemistry and we envision this as an evolving resource where we will continue to release data

as it becomes available.

SIGNIFICANCE

Degraders and molecular glues are small molecules that can target and promote the

degradation of specific proteins providing a novel approach for modulating protein function.

Currently available unbiased methods to identify targets of degraders, although successful in

identifying transient and/or degraded targets, are limited in sensitivity and ability to identify direct

binders of these molecules, prohibiting identification of targets that have weak expression

changes or are glued and not degraded. Here, we develop an automatable high throughput

method for the identification of chemically-induced binders. We demonstrate the ability to

comprehensively identify new targets by identifying 298 neo-substrates of CRBN, significantly

expanding the repertoire of actionable targets. We then used structural and biochemical

characterization alongside a computational structural alignment workflow to validate hit targets

and selected one novel target, PPIL4, to perform a focused biochemical screen for the

identification of a new lead molecule. CRBN is the most targeted ligase in the TPD field, with

molecules FDA approved and more in clinical trials it is important that we understand the

complete cellular and molecular impact of targeting this ligase. The findings presented in this

study, open a new and complementary avenue for target identification and create a valuable

data resource mapping a wide range of neo-substrates of the CRBN ligase. Through expansion

of the range of CRBN targets, we not only enhance our knowledge of newly druggable targets

and offer new avenues for therapeutic development, but we also enhance our understanding of

the molecular mechanisms and cellular pathways that are influenced by existing and future IMiD

molecules providing opportunities for improved drug design.

ACKNOWLEDGEMENTS

We thank S. Dixon-Clarke, M. Hunkeler, T. Levitz, Y. Xiong, J. Che, T. Zhang and members of the Fischer and Gray labs for helpful discussions, reagents, and support. We thank the Harvard Cryo-EM center for Structural Biology for support on data collection. Financial support for this work was provided by the National Institutes of Health (R01CA214608 and R01CA262188 (both to E.S.F.). K.B. is a Damon Runyon Fellow supported by the Damon Runyon Cancer Research Foundation (DRG-2514-24). Figures 1A, 2A and 5A were created in Biorender.

AUTHOR CONTRIBUTIONS

K.B. designed experiments, performed structural work and biochemical assays, analyzed the data, interpreted results, wrote the manuscript. R.J.M. initiated the study, designed proteomics experiments, performed proteomics experiments, analyzed the data. S.S.R.B. performed computational alignment analysis, interpreted results. J.W.B. initiated the study and performed biochemical experiments. R.J.L. wrote proteomics analysis code. D.M.A performed proteomics. M.L. performed immunoblots. H.Y. performed TR-FRET screen. S.O. performed TR-FRET screen. A.L.V. synthesized molecules. N.S.G. supervised experiments. K.A.D. conceived the study, designed experiments, analyzed the data, interpreted the results, wrote the manuscript and supervised the study. E.S.F. conceived the study, interpreted results, supervised and funded the study. All authors read, edited and approved the final manuscript.

DECLARATION OF INTERESTS

E.S.F. is a founder, scientific advisory board (SAB) member, and equity holder of Civetta Therapeutics, Proximity Therapeutics, Stelexis Biosciences, and Neomorph, Inc. (also board of directors). He is an equity holder and SAB member for Avilar Therapeutics, Photys Therapeutics, and Ajax Therapeutics and an equity holder in Lighthorse Therapeutics and Anvia Therapeutics. E.S.F. is a consultant to Novartis, EcoR1 capital, Odyssey and Deerfield. The

552 Fischer lab receives or has received research funding from Deerfield, Novartis, Ajax, Interline,

553 Bayer and Astellas. K.A.D receives or has received consulting fees from Kronos Bio and

554 Neomorph Inc. N.S.G. is a founder, science advisory board member (SAB) and equity holder in

555 Syros, C4, Allorion, Lighthorse, Inception, Matchpoint, Shenandoah (board member), Larkspur

556 (board member) and Soltego (board member). The Gray lab receives or has received research

557 funding from Novartis, Takeda, Astellas, Taiho, Jansen, Kinogen, Arbella, Deerfield,

558 Springworks, Interline and Sanofi.

 FIGURE TITLES AND LEGENDS

Figure 1 | Proof of concept for target enrichment in-lysate.

(**A**) Schematic representation of the first-generation enrichment-based quantitative proteomics workflow established for target enrichment and identification. (**B**) Chemical structures of degraders – Pomalidomide (molecular glue) and SB1-G-187 (heterobifunctional). (**C**) Scatterplots depicting relative protein abundance following Flag-CRBN-DDB1ΔB enrichment from in-lysate treatment with 1 µM Pomalidomide and recombinant Flag-CRBN-DDB1ΔB spike in. Left: MOLT4 cells and Right: Kelly Cells. Scatterplots display fold change in abundance to DMSO. Significant changes were assessed by moderated t-test as implemented in the limma package[80] with $\log_2$ FC shown on the y-axis and negative $\log_{10}$ P-value on the x-axis. (**D**) Venn diagram showing unique and overlapping hits for Pomalidomide found in our enrichment study and in publicly available whole cell degradation data. (**E**) As in **C**, but with 1 µM SB1-G-187 treatment. (**F**) As in **D**, but with SB1-G-187 treatment.

Figure 2 | Unveiling and mapping CRBN recruited neo-substrates.

(**A**) Schematic representation of the second-generation enrichment-based quantitative

proteomics workflow established for target enrichment and identification. (**B**) Chemical

structures of the 20 CRBN-based degraders profiled in this study. (**C**) Scatterplot depicting

relative protein abundance following Flag-CRBN-DDB1ΔB enrichment from in-lysate treatment

with degrader and recombinant Flag-CRBN-DDB1ΔB spike in. Scatterplot displays fold change

in abundance to DMSO. Significant changes were assessed by moderated t-test as

implemented in the limma package[80] with $\log_2$ FC shown on the y-axis and negative $\log_{10}$ P-

value on the x-axis. Scatterplots for all 21 treatments across MOLT4 and Kelly cells can be

found in separate PDF's "**Figures S3-4**", representative example for a single treatment

(Pomalidomide, 1 μM) is displayed here. (**D**) The number of independent IPs for which

enrichment was observed for each target. Inset, the top 20 frequently enriched target proteins.

(**E**) Venn diagram showing unique and overlapping hits found in our enrichment study and in

published literature. (**F**) Donut chart representing the proportions of enriched proteins contained

within the Top 10 different superfamily categories. (**G**) Donut chart representing the proportions

of enriched proteins contained within the Top 10 different domain categories.

Figure 3 | Structural alignment and assessment of ZF CRBN neo-substrates.
(**A**) Schematic representation of the computational workflow established for AF2 G-loop binding
compatibility with CRBN-IMiD. (**B**) Heatmap displaying the log2 fold change (log2 FC) of
significant (P-value <0.001) molecular glue dependent ZF targets in MOLT4 cells. White space
in the heatmap corresponds to log2FC = 0 or no quantification. Previously reported targets are
marked with a blue dot, newly reported targets are marked with a green dot and targets with a
structural G-loop are marked with a gray dot. Significant changes were assessed by moderated
t-test as implemented in the limma package[80]. (**C**) As in **B**, but with Kelly cells. (**D**) Venn
diagram showing unique and overlapping ZF hits comparing MOLT4 and Kelly cell targets. (**E**)

604   Stacked bar plot showing the proportion of targets complexed and degraded by each of the

605   indicated IMiD molecules. "not seen" indicates enriched targets were not quantified in global

606   proteomics studies. "seen, degraded" indicates enriched targets quantified and reported as

607   degraded in global proteomics. "seen, not degraded" indicates enriched targets were quantified

608   but not degraded in global proteomics [19]. (**F**) Pie chart displays the IMiD-grouped data from **E**.

Figure 4 | Assessment and validation of CRBN non-ZF neo-substrates.

(**A**) Venn diagram showing unique and overlapping non-ZF hits comparing MOLT4 and Kelly cell targets. (**B**) Heatmap displaying the log2 fold change (log2 FC) for the 39 overlapping hits from **A**. White space in the heatmap corresponds to log2FC = 0 or no quantification. Previously reported targets are marked with a blue dot, newly reported targets are marked with a green dot and targets with a structural G-loop are marked with a gray dot. Significant changes were assessed by moderated t-test as implemented in the limma package[80]. (**C**) TR-FRET with titration of FPFT-2216 or pomalidomide to N-terminally biotinylated FL PDE6D, DTWD1, PPIL4 or RAB28 at 20 nM, incubated with terbium-streptavidin at 2 nM to monitor binding to GFP-CRBN-DDB1ΔB at 200 nM. Values were determined by technical replicates of n=2. (**D**) Immunoblots of ubiquitylation assay establishing PDE6D, DTWD1 and PPIL4 as FPFT-2216 and pomalidomide-induced neo-substrates of CRBN. (**E**) Structural G-loop alignment of AF2 PPIL4, PDE6D, and DTWD1 with CRBN-DDB1ΔB (PDB ID 5QFD, 6UML). Corresponding clash score is displayed. (**F**) Cryo-EM 3D reconstruction of PPIL4-RRM bound in ternary complex with FPFT-2216-CRBN-DDB1 FL, and PDE6D bound with FPFT-2216-CRBN-DDB1ΔB. Maps are postprocessed with DeepEMhancer[81]. Inset of each shows the potential binding mode of action of FPFT-2216 engaging PPIL4 or PDE6D via neo-substrate G-loop and its interacting residues.

Figure 5 | High throughput IMiD analog library screen for improved hit molecules for PPIL4. (**A**) Schematic of the high throughput TR-FRET screening workflow used to screen >6,000 IMiD analogs. (**B**) TR-FRET: Normalized 520/490 ratio for each of the >6,000 compounds derived from IMiD molecules combined from the Gray/Fischer laboratories and those purchased from Enamine with GFP-CRBN-DDB1ΔB at 50 nM, biotinylated PPIL4 at 20 nM, and terbium-streptavidin at 2 nM. (**C**) TR-FRET: titration of FPFT-2216 and lead compounds to GFP-CRBN-DDB1ΔB at (200 nM), biotinylated PPIL4 at 20 nM, and terbium-streptavidin at 2 nM. Values were determined by technical replicates of n=2. (**D**) Chemical structures of FPFT-2216 alongside new lead compound from **B** and **C**. (**E**) Scatterplot depicting relative protein abundance following Flag-CRBN-DDB1ΔB enrichment from Kelly cell in-lysate

treatment with FPFT-2216 (left) and Z6466608626 (right) and recombinant Flag-CRBN-

DDB1ΔB spike in. Scatterplot displays fold change in abundance to DMSO. Significant changes

were assessed by moderated t-test as implemented in the limma package[80] with $\log_2$ FC shown

on the y-axis and negative $\log_{10}$ P-value on the x-axis. (**F**) Scatterplot depicting relative protein

abundance following Z6466608626 treatment in MOLT4 cells. Significant changes were

assessed by moderated t-test as implemented in the limma package[80] with $\log_2$ FC shown on

the y-axis and negative $\log_{10}$ P-value on the x-axis.

Table 1. Data collection and refinement statistics for cryo-EM datasets, related to Figures 4 and S7

	CRBN-DDB1 FPFT-2216 PPIL4 RRM	CRBN-DDB1ΔB FPFT-2216 PDE6D Consensus refine	CRBN-DDB1ΔB FPFT-2216 PDE6D Local refine
Microscope	Talos Arctica	Talos Arctica	
Voltage (kV)	200	200	
Camera	Gatan K3	Gatan K3	
Magnification (×)	36,000	36,000	
Pixel size (Å)	1.1	1.1	
Total electron exposure (e⁻/Å²)	54	51.2	
Number of frames	40	50	
Defocus range (μm)	−2.0 to −0.8	−2.0 to −0.8	
Data collection software	SerialEM4.1b		
Micrographs collected	4,170	4,474	
Total extracted particles	3,198,055	3,393,589	
EMDB accession code	EMD-XXXXX	EMD-XXXXX	EMD-XXXXX
PDB accession code	PDB-XXXX	PDB-XXXX	
Final particles used	219,802	515,636	515,636
Map resolution (Å, FSC 0.143)	3.5	3.3	3.4
FSC threshold	0.143	0.143	0.143
Model composition			
Protein residues	1211	1260	
Ligands	2	2	
Refinement package	phenix.real_space_refine		
Model-to-map CC	0.72	0.78	
Model-to-map FSC (Å, FSC 0.5)	3.8	3.5	
Mean *B* factors (Å²)			
Protein	63.69	98.29	
Ligand	69.37	118.85	
Water	–	–	
Bond root-mean-square deviation (RMSD)			
Lengths (Å)	0.005	0.005	

	CRBN-DDB1 FPFT-2216 PPIL4 RRM	CRBN-DDB1ΔB FPFT-2216 PDE6D Consensus refine	CRBN-DDB1ΔB FPFT-2216 PDE6D Local refine
Angles (°)	0.615	0.797	
Validation			
MolProbity score	2.05	1.53	
Clash score	11.78	6.37	
Rotamer outliers (%)	1.27	0.11	
CaBLAM outliers (%)	2.76	1.54	
Ramachandran plot			
Favored (%)	94.26	96.96	
Allowed (%)	5.74	3.04	
Disallowed (%)	0.00	0.00	

653
654
655

RESOURCES AVAILABILITY

Lead Contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the Lead Contact, Eric Fischer (Eric_Fischer@DFCI.HARVARD.EDU).

Materials Availability

Small molecules described in this study will be made available on request, upon completion of a

Materials Transfer Agreement.

Supplemental information.

Table S1. Table reporting Log2 Fold Change and P-value for all proteins quantified in CRBN-

DDB1ΔB IP-MS experiments in the presence of pomalidomide (1 μM) or SB1-G-187 (1 μM) in

MOLT4 or Kelly cell lysate. Statistics generated through moderated t-test in limma package[80].

Table related to Figures 1 and S1.

Table S2. Table reporting Log2 Fold Change and P-value for all proteins quantified in CRBN-

DDB1ΔB IP-MS experiments in the presence of IMiD analogs in MOLT4 cell lysate. Statistics

generated through moderated t-test in limma package[80]. Table related to Figure 2-4 and S3.

Table S3. Table reporting Log2 Fold Change and P-value for all proteins quantified in CRBN-

DDB1ΔB IP-MS experiments in the presence of IMiD analogs in Kelly cell lysate. Statistics

generated through moderated t-test in limma package[80]. Table related to Figure 2-4 and S4.

Table S4. Table reporting frequency of "hit" for all proteins identified as a hit across the IMiD

library IP-MS screens in MOLT4 and Kelly cell lysates. Table reports target

domain/family/superfamily classifications and interactome information. Table related to Figures

2, 4 and S2.

Table S5. Table reporting the G-loop details for all proteins in the human proteome that were

identified as having a domain carrying a G-loop. Table reports IP and global proteomics data for

PPIL4 studies. Table related to Figures 3-5, S5-6.

683
684

 SUPPLEMENTAL FIGURES AND LEGENDS

Figure S1. Validation of enriched targets from lysate IPs, related to Figure 1, Table S1

(**A**) Box plot depicting all quantified peptides for each of the enriched targets from MOLT4 cells comparing DMSO control and Pomalidomide treatments. (**B**) As in **A** but for Kelly cells. (**C**) Flag-CRBN IP experiments were performed in the presence of increasing concentration of pomalidomide in both MOLT4 and Kelly cells. Following elution, ASS1 protein levels were assessed by immunoblot. (**D**) TR-FRET: titration of IMiD analogs to GFP-CRBN-DDB1ΔB at 200 nM, ZBED3$_{39-108}$ or ZNF219$_{53-110}$ at 20 nM, and terbium-streptavidin at 2 nM. Values were determined by technical replicates of n=2. (**E**) Box plot depicting all quantified peptides for each

of the enriched targets from MOLT4 cells comparing DMSO control and SB1-G-187 treatments.

(**F**) As in **E** but for Kelly cells.

Figure S2. Quantitative proteomics for exploration of targets recruited by IMiD analogs,

related to Figure 2

(**A**) Histogram displaying number of unique proteins quantified in DDA and diaPASEF IP-MS

experiments for MOLT4 and Kelly cells. (**B**) Histogram displaying number of unique peptides

quantified in DDA and diaPASEF IP-MS experiments for MOLT4 and Kelly cells. (**C**) Scatterplot

depicting relative protein abundance following 5 µM Pomalidomide treatment in MOLT4 cells.

Significant changes were assessed by moderated t-test as implemented in the limma package[80]

with $\log_2$ FC shown on the y-axis and negative $\log_{10}$ P-value on the x-axis.

Figure S3. Scatterplots of IP-MS with molecular glues, related to Figure 2, 3, S4, Table S2,

S3, S4

Scatterplots depicting relative protein abundance following Flag-CRBN-DDB1 enrichment from

in-lysate treatment with degrader and recombinant Flag-CRBN-DDB1 spike in. Scatterplot

displays fold change in abundance for each of the 20 molecules relative to DMSO in MOLT4

715   cells. Significant changes were assessed by moderated t-test as implemented in the limma

716   package[80] with $\log_2$ FC shown on the y-axis and negative $\log_{10}$ P-value on the x-axis.

Figure S4. Scatterplots of IP-MS with molecular glues, related to Figure 2, 3, S4, Table S2, S3, S4

Scatterplots depicting relative protein abundance following Flag-CRBN-DDB1 enrichment from in-lysate treatment with degrader and recombinant Flag-CRBN-DDB1 spike in. Scatterplot displays fold change in abundance for each of the 20 molecules relative to DMSO in Kelly cells.

725 Significant changes were assessed by moderated t-test as implemented in the limma package[80]

726 with $\log_2$ FC shown on the y-axis and negative $\log_{10}$ P-value on the x-axis.

Figure S5. Structural alignments of G-loop neo-substrates, related to Figure 3, 4, S5,

Table S5

Structural G-loop alignment showing alignment of AF2 structures by the G-loop for a subset of

the neo-substrate targets identified as hits in this study with CRBN (PDB ID: 5FQD).

Corresponding clash scores are displayed.

Figure S6. Resolving clashes from G-loop alignments, related to Figure 3, 4, S5, Table S5

(**A**) G-loop aligned AF2 structures for ZNF654 demonstrating the clash score and structural shift

before and after relaxation with Rosetta. (**B**) As in **A**, but for PAAF1. (**C**) Violin plot depicting the

distribution of class scores for a single G-loop from each of the identified targets in this dataset

(left), and those for ZF proteins only (right). (**D**) Scatterplot depicting relative protein abundance

following FPFT-2216 treatment in MOLT4 cells. Significant changes were assessed by

moderated t-test as implemented in the limma package[80] with $\log_2$ FC shown on the y-axis and

negative $\log_{10}$ P-value on the x-axis. (**E**) As in **A**, but for DTWD1. (**F**) Overlay of G-loop aligned

AF2 structure of PPIL4 and structure obtained by cryo-EM. (**G**) Same as in (**F**) but with PDE6D.

Figure S7. Cryo-EM processing schematic of neosubstrate-molecular glue-CRBN-DDB1,

related to Figure 4, Table 1

(**A**) Cryo-EM processing schematic of CRBN-DDB1 (FL) with PPIL4 RRM domain and FPFT-

2216. **(B)** Cryo-EM processing schematic of CRBN-DDB1ΔB with PDE6D and FPFT-2216.

754
755 **REFERENCES**
756

1. Bondeson, D.P., and Crews, C.M. (2017). Targeted protein degradation by small
molecules. Annual review of pharmacology and toxicology *57*, 107-123.
2. Neklesa, T.K., Winkler, J.D., and Crews, C.M. (2017). Targeted protein degradation by
PROTACs. Pharmacology & therapeutics *174*, 138-144.
10.1016/j.pharmthera.2017.02.027.
3. Sakamoto, K.M., Kim, K.B., Kumagai, A., Mercurio, F., Crews, C.M., and Deshaies, R.J.
(2001). Protacs: chimeric molecules that target proteins to the Skp1–Cullin–F box
complex for ubiquitination and degradation. Proceedings of the National Academy of
Sciences *98*, 8554-8559.
4. Komander, D., and Rape, M. (2012). The ubiquitin code. Annual review of biochemistry
81, 203-229.
5. Kronke, J., Udeshi, N.D., Narla, A., Grauman, P., Hurst, S.N., McConkey, M., Svinkina, T.,
Heckl, D., Comer, E., Li, X., et al. (2014). Lenalidomide causes selective degradation of
IKZF1 and IKZF3 in multiple myeloma cells. Science *343*, 301-305.
10.1126/science.1244851.
6. Winter, G.E., Buckley, D.L., Paulk, J., Roberts, J.M., Souza, A., Dhe-Paganon, S., and
Bradner, J.E. (2015). Phthalimide conjugation as a strategy for in vivo target protein
degradation. Science *348*, 1376-1381.
7. Bondeson, D.P., Mares, A., Smith, I.E., Ko, E., Campos, S., Miah, A.H., Mulholland, K.E.,
Routly, N., Buckley, D.L., and Gustafson, J.L. (2015). Catalytic in vivo protein knockdown
by small-molecule PROTACs. Nature chemical biology *11*, 611.
8. Wang, B., Cao, S., and Zheng, N. (2024). Emerging strategies for prospective discovery of
molecular glue degraders. Current Opinion in Structural Biology *86*, 102811.
9. Ito, T., Ando, H., Suzuki, T., Ogura, T., Hotta, K., Imamura, Y., Yamaguchi, Y., and Handa,
H. (2010). Identification of a primary target of thalidomide teratogenicity. Science *327*,
1345-1350. 10.1126/science.1177319.
10. Fischer, E.S., Bohm, K., Lydeard, J.R., Yang, H., Stadler, M.B., Cavadini, S., Nagel, J.,
Serluca, F., Acker, V., Lingaraju, G.M., et al. (2014). Structure of the DDB1-CRBN E3
ubiquitin ligase in complex with thalidomide. Nature *512*, 49-53. 10.1038/nature13527.
11. Matyskiela, M.E., Lu, G., Ito, T., Pagarigan, B., Lu, C.C., Miller, K., Fang, W., Wang, N.Y.,
Nguyen, D., Houston, J., et al. (2016). A novel cereblon modulator recruits GSPT1 to the
CRL4(CRBN) ubiquitin ligase. Nature *535*, 252-257. 10.1038/nature18611.
12. Petzold, G., Fischer, E.S., and Thoma, N.H. (2016). Structural basis of lenalidomide-
induced CK1alpha degradation by the CRL4(CRBN) ubiquitin ligase. Nature *532*, 127-130.
10.1038/nature16979.
13. Lu, G., Middleton, R.E., Sun, H., Naniong, M., Ott, C.J., Mitsiades, C.S., Wong, K.K.,
Bradner, J.E., and Kaelin, W.G., Jr. (2014). The myeloma drug lenalidomide promotes the
cereblon-dependent destruction of Ikaros proteins. Science *343*, 305-309.
10.1126/science.1244917.

14. Oleinikovas, V., Gainza, P., Ryckmans, T., Fasching, B., and Thomä, N.H. (2024). From
thalidomide to rational molecular glue design for targeted protein degradation. Annual
Review of Pharmacology and Toxicology *64*, 291-312.
15. Matyskiela, M.E., Couto, S., Zheng, X., Lu, G., Hui, J., Stamp, K., Drew, C., Ren, Y., Wang,
M., Carpenter, A., et al. (2018). SALL4 mediates teratogenicity as a thalidomide-
dependent cereblon substrate. Nature chemical biology *14*, 981-987. 10.1038/s41589-
018-0129-x.
16. Sievers, Q.L., Gasser, J.A., Cowley, G.S., Fischer, E.S., and Ebert, B.L. (2018). Genome-
wide screen identifies cullin-RING ligase machinery required for lenalidomide-
dependent CRL4(CRBN) activity. Blood *132*, 1293-1303. 10.1182/blood-2018-01-821769.
17. Kronke, J., Fink, E.C., Hollenbach, P.W., MacBeth, K.J., Hurst, S.N., Udeshi, N.D.,
Chamberlain, P.P., Mani, D.R., Man, H.W., Gandhi, A.K., et al. (2015). Lenalidomide
induces ubiquitination and degradation of CK1alpha in del(5q) MDS. Nature *523*, 183-
188. 10.1038/nature14610.
18. An, J., Ponthier, C.M., Sack, R., Seebacher, J., Stadler, M.B., Donovan, K.A., and Fischer,
E.S. (2017). pSILAC mass spectrometry reveals ZFP91 as IMiD-dependent substrate of
the CRL4(CRBN) ubiquitin ligase. Nature communications *8*, 15398.
10.1038/ncomms15398.
19. Donovan, K.A., An, J., Nowak, R.P., Yuan, J.C., Fink, E.C., Berry, B.C., Ebert, B.L., and
Fischer, E.S. (2018). Thalidomide promotes degradation of SALL4, a transcription factor
implicated in Duane Radial Ray syndrome. Elife *7*. 10.7554/eLife.38430.
20. Yamamoto, J., Suwa, T., Murase, Y., Tateno, S., Mizutome, H., Asatsuma-Okumura, T.,
Shimizu, N., Kishi, T., Momose, S., and Kizaki, M. (2020). ARID2 is a pomalidomide-
dependent CRL4CRBN substrate in multiple myeloma cells. Nature chemical biology *16*,
1208-1217.
21. Renneville, A., Gasser, J.A., Grinshpun, D.E., Jean Beltran, P.M., Udeshi, N.D., Matyskiela,
M.E., Clayton, T., McConkey, M., Viswanathan, K., and Tepper, A. (2021). Avadomide
induces degradation of ZMYM2 fusion oncoproteins in hematologic malignancies. Blood
cancer discovery *2*, 250-265.
22. Li, L., Xue, W., Shen, Z., Liu, J., Hu, M., Cheng, Z., Wang, Y., Chen, Y., Chang, H., and Liu,
Y. (2020). A cereblon modulator CC-885 induces CRBN-and p97-dependent PLK1
degradation and synergizes with volasertib to suppress lung cancer. Molecular Therapy-
Oncolytics *18*, 215-225.
23. Huang, H.T., Dobrovolsky, D., Paulk, J., Yang, G., Weisberg, E.L., Doctor, Z.M., Buckley,
D.L., Cho, J.H., Ko, E., Jang, J., et al. (2018). A Chemoproteomic Approach to Query the
Degradable Kinome Using a Multi-kinase Degrader. Cell chemical biology *25*, 88-99.e86.
10.1016/j.chembiol.2017.10.005.
24. Sathe, G., and Sapkota, G.P. (2023). Proteomic approaches advancing targeted protein
degradation. Trends in pharmacological sciences *44*, 786-801.
25. King, E.A., Cho, Y., Hsu, N.S., Dovala, D., McKenna, J.M., Tallarico, J.A., Schirle, M., and
Nomura, D.K. (2023). Chemoproteomics-enabled discovery of a covalent molecular glue
degrader targeting NF-κB. Cell chemical biology *30*, 394-402. e399.

26.   Donovan, K.A., Ferguson, F.M., Bushman, J.W., Eleuteri, N.A., Bhunia, D., Ryu, S., Tan, L.,
839   Shi, K., Yue, H., and Liu, X. (2020). Mapping the Degradable Kinome Provides a Resource
840   for Expedited Degrader Development. Cell.

841   27.   Bondeson, D.P., Smith, B.E., Burslem, G.M., Buhimschi, A.D., Hines, J., Jaime-Figueroa, S.,
842   Wang, J., Hamman, B.D., Ishchenko, A., and Crews, C.M. (2018). Lessons in PROTAC
843   Design from Selective Degradation with a Promiscuous Warhead. Cell chemical biology
844   *25*, 78-87.e75. 10.1016/j.chembiol.2017.09.010.

845   28.   Nowak, R.P., DeAngelo, S.L., Buckley, D., He, Z., Donovan, K.A., An, J., Safaee, N.,
846   Jedrychowski, M.P., Ponthier, C.M., Ishoey, M., et al. (2018). Plasticity in binding confers
847   selectivity in ligand-induced protein degradation. Nature chemical biology *14*, 706-714.
848   10.1038/s41589-018-0055-y.

849   29.   Li, Y.-D., Ma, M.W., Hassan, M.M., Hunkeler, M., Teng, M., Puvar, K., Lumpkin, R.,
850   Sandoval, B., Jin, C.Y., and Ficarro, S.B. (2023). Template-assisted covalent modification
851   of DCAF16 underlies activity of BRD4 molecular glue degraders. bioRxiv, 2023.2002.
852   2014.528208.

853   30.   Hsia, O., Hinterndorfer, M., Cowan, A.D., Iso, K., Ishida, T., Sundaramoorthy, R.,
854   Nakasone, M.A., Imrichova, H., Schätz, C., and Rukavina, A. (2024). Targeted protein
855   degradation via intramolecular bivalent glues. Nature *627*, 204-211.

856   31.   Xiong, Y., Donovan, K.A., Eleuteri, N.A., Kirmani, N., Yue, H., Razov, A., Krupnick, N.M.,
857   Nowak, R.P., and Fischer, E.S. (2021). Chemo-proteomics exploration of HDAC
858   degradability by small molecule degraders. Cell chemical biology *28*, 1514-1527. e1514.

859   32.   Huang, H.-T., Lumpkin, R.J., Tsai, R.W., Su, S., Zhao, X., Xiong, Y., Chen, J., Mageed, N.,
860   Donovan, K.A., and Fischer, E.S. (2024). Ubiquitin-specific proximity labeling for the
861   identification of E3 ligase substrates. Nature Chemical Biology, 1-10.

862   33.   Huttlin, E.L., Ting, L., Bruckner, R.J., Gebreab, F., Gygi, M.P., Szpyt, J., Tam, S., Zarraga,
863   G., Colby, G., and Baltier, K. (2015). The BioPlex network: a systematic exploration of the
864   human interactome. Cell *162*, 425-440.

865   34.   Cho, K.F., Branon, T.C., Udeshi, N.D., Myers, S.A., Carr, S.A., and Ting, A.Y. (2020).
866   Proximity labeling in mammalian cells with TurboID and split-TurboID. Nature Protocols
867   *15*, 3971-3999.

868   35.   Roux, K.J., Kim, D.I., Burke, B., and May, D.G. (2018). BioID: a screen for protein-protein
869   interactions. Current protocols in protein science *91*, 19.23. 11-19.23. 15.

870   36.   Kim, D.I., KC, B., Zhu, W., Motamedchaboki, K., Doye, V., and Roux, K.J. (2014). Probing
871   nuclear pore complex architecture with proximity-dependent biotinylation. Proceedings
872   of the National Academy of Sciences *111*, E2453-E2461.

873   37.   Kim, D.I., Jensen, S.C., Noble, K.A., Kc, B., Roux, K.H., Motamedchaboki, K., and Roux, K.J.
874   (2016). An improved smaller biotin ligase for BioID proximity labeling. Molecular biology
875   of the cell *27*, 1188-1196.

876   38.   Kido, K., Yamanaka, S., Nakano, S., Motani, K., Shinohara, S., Nozawa, A., Kosako, H., Ito,
877   S., and Sawasaki, T. (2020). AirID, a novel proximity biotinylation enzyme, for analysis of
878   protein–protein interactions. Elife *9*, e54983.

879   39.   Martell, J.D., Deerinck, T.J., Sancak, Y., Poulos, T.L., Mootha, V.K., Sosinsky, G.E.,
880   Ellisman, M.H., and Ting, A.Y. (2012). Engineered ascorbate peroxidase as a genetically
881   encoded reporter for electron microscopy. Nature biotechnology *30*, 1143-1148.

40. Sievers, Q.L., Petzold, G., Bunker, R.D., Renneville, A., Slabicki, M., Liddicoat, B.J.,
Abdulrahman, W., Mikkelsen, T., Ebert, B.L., and Thoma, N.H. (2018). Defining the
human C2H2 zinc finger degrome targeted by thalidomide analogs through CRBN.
Science *362*. 10.1126/science.aat0572.
41. Roh, Y.S., Song, J., and Seki, E. (2014). TAK1 regulates hepatic cell survival and
carcinogenesis. Journal of gastroenterology *49*, 185-194.
42. Shibuya, H., Yamaguchi, K., Shirakabe, K., Tonegawa, A., Gotoh, Y., Ueno, N., Irie, K.,
Nishida, E., and Matsumoto, K. (1996). TAB1: an activator of the TAK1 MAPKKK in TGF-β
signal transduction. Science *272*, 1179-1182.
43. Garivet, G., Hofer, W., Konitsiotis, A., Klein, C., Kaiser, N., Mejuch, T., Fansa, E., Alsaabi,
R., Wittinghofer, A., and Bastiaens, P.I. (2019). Small-molecule inhibition of the UNC-Src
interaction impairs dynamic Src localization in cells. Cell Chemical Biology *26*, 842-851.
e847.
44. Golkowski, M., Brigham, J.L., Perera, B.G.K., Romano, G.S., Maly, D.J., and Ong, S.-E.
(2014). Rapid profiling of protein kinase inhibitors by quantitative proteomics.
Medchemcomm *5*, 363-369.
45. Meier, F., Brunner, A.-D., Frank, M., Ha, A., Bludau, I., Voytik, E., Kaspar-Schoenefeld, S.,
Lubeck, M., Raether, O., and Bache, N. (2020). diaPASEF: parallel accumulation–serial
fragmentation combined with data-independent acquisition. Nature methods *17*, 1229-
1236.
46. Knight, R. (2005). IMiDs: a novel class of immunomodulators. (Elsevier), pp. 24-30.
47. D'Amato, R.J., Loughnan, M.S., Flynn, E., and Folkman, J. (1994). Thalidomide is an
inhibitor of angiogenesis. Proc Natl Acad Sci U S A *91*, 4082-4085.
48. Lonial, S., Popat, R., Hulin, C., Jagannath, S., Oriol, A., Richardson, P.G., Facon, T., Weisel,
K., Larsen, J.T., and Minnema, M.C. (2022). Iberdomide plus dexamethasone in heavily
pretreated late-line relapsed or refractory multiple myeloma (CC-220-MM-001): a
multicentre, multicohort, open-label, phase 1/2 trial. The Lancet Haematology *9*, e822-
e832.
49. Richardson, P.G., Trudel, S., Popat, R., Mateos, M.-V., Vangsted, A.J., Ramasamy, K.,
Martinez-Lopez, J., Quach, H., Orlowski, R.Z., and Arnao, M. (2023). Mezigdomide plus
dexamethasone in relapsed and refractory multiple myeloma. New England Journal of
Medicine *389*, 1009-1022.
50. Surka, C., Jin, L., Mbong, N., Lu, C.-C., Jang, I.S., Rychak, E., Mendy, D., Clayton, T.,
Tindall, E., and Hsu, C. (2021). CC-90009, a novel cereblon E3 ligase modulator, targets
acute myeloid leukemia blasts and leukemia stem cells. Blood, The Journal of the
American Society of Hematology *137*, 661-677.
51. Gemechu, Y., Millrine, D., Hashimoto, S., Prakash, J., Sanchenkova, K., Metwally, H.,
Gyanu, P., Kang, S., and Kishimoto, T. (2018). Humanized cereblon mice revealed two
distinct therapeutic pathways of immunomodulatory drugs. Proc Natl Acad Sci U S A
115, 11802-11807. 10.1073/pnas.1814446115.
52. Hagner, P.R., Man, H.-W., Fontanillo, C., Wang, M., Couto, S., Breider, M., Bjorklund, C.,
Havens, C.G., Lu, G., and Rychak, E. (2015). CC-122, a pleiotropic pathway modifier,
mimics an interferon response and has antitumor activity in DLBCL. Blood, The Journal
of the American Society of Hematology *126*, 779-789.

53. Wang, E.S., Verano, A.L., Nowak, R.P., Yuan, J.C., Donovan, K.A., Eleuteri, N.A., Yue, H.,
Ngo, K.H., Lizotte, P.H., and Gokhale, P.C. (2021). Acute pharmacological degradation of
Helios destabilizes regulatory T cells. Nature chemical biology *17*, 711-717.

54. Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen,
J., Mitchell, A., and Nuka, G. (2014). InterProScan 5: genome-scale protein function
classification. Bioinformatics *30*, 1236-1240.

55. Paysan-Lafosse, T., Blum, M., Chuguransky, S., Grego, T., Pinto, B.L., Salazar, G.A.,
Bileschi, M.L., Bork, P., Bridge, A., and Colwell, L. (2023). InterPro in 2022. Nucleic acids
research *51*, D418-D427.

56. UniProt: the universal protein knowledgebase in 2023. (2023). Nucleic acids research
51, D523-D531.

57. Gough, J., Karplus, K., Hughey, R., and Chothia, C. (2001). Assignment of homology to
genome sequences using a library of hidden Markov models that represent all proteins
of known structure. Journal of molecular biology *313*, 903-919.

58. Razumkov, H., Jiang, Z., Baek, K., You, I., Geng, Q., Donovan, K., Tang, M.T., Metivier,
R.J., Mageed, N., and Seo, P. (2024). Discovery of CRBN-dependent WEE1 Molecular
Glue Degraders from a Multicomponent Combinatorial Library. bioRxiv, 2024.2005.
2004.592550.

59. Zhao, M., Hu, M., Chen, Y., Liu, H., Chen, Y., Liu, B., and Fang, B. (2021). Cereblon
modulator CC-885 induces CRBN-dependent ubiquitination and degradation of CDK4 in
multiple myeloma. Biochemical and biophysical research communications *549*, 150-156.

60. Furihata, H., Yamanaka, S., Honda, T., Miyauchi, Y., Asano, A., Shibata, N., Tanokura, M.,
Sawasaki, T., and Miyakawa, T. (2020). Structural bases of IMiD selectivity that emerges
by 5-hydroxythalidomide. Nature communications *11*, 4578.

61. Zhou, J., and Grigoryan, G. (2015). Rapid search for tertiary fragments reveals protein
sequence–structure relationships. Protein Science *24*, 508-524.

62. Varadi, M., Bertoni, D., Magana, P., Paramval, U., Pidruchna, I., Radhakrishnan, M.,
Tsenkov, M., Nair, S., Mirdita, M., and Yeo, J. (2024). AlphaFold Protein Structure
Database in 2024: providing structure coverage for over 214 million protein sequences.
Nucleic acids research *52*, D368-D375.

63. Zhang, J., Schaeffer, R.D., Durham, J., Cong, Q., and Grishin, N.V. (2023). DPAM: A
domain parser for AlphaFold models. Protein Science *32*, e4548.

64. Leman, J.K., Weitzner, B.D., Lewis, S.M., Adolf-Bryfogle, J., Alam, N., Alford, R.F.,
Aprahamian, M., Baker, D., Barlow, K.A., and Barth, P. (2020). Macromolecular modeling
and design in Rosetta: recent methods and frameworks. Nature methods *17*, 665-680.

65. Maguire, J.B., Haddox, H.K., Strickland, D., Halabiya, S.F., Coventry, B., Griffin, J.R.,
Pulavarti, S.V.K., Cummins, M., Thieker, D.F., and Klavins, E. (2021). Perturbing the
energy landscape for improved packing during computational protein design. Proteins:
Structure, Function, and Bioinformatics *89*, 436-449.

66. Rodriguez-Rivera, F.P., and Levi, S.M. (2021). Unifying catalysis framework to dissect
proteasomal degradation paradigms. ACS Central Science *7*, 1117-1125.

67. Barroso-Gomila, O., Merino-Cacho, L., Muratore, V., Perez, C., Taibi, V., Maspero, E.,
Azkargorta, M., Iloro, I., Trulsson, F., and Vertegaal, A.C. (2023). BioE3 identifies specific
substrates of ubiquitin E3 ligases. Nature Communications *14*, 7656.

68.   Mukhopadhyay, U., Levantovsky, S., Carusone, T.M., Gharbi, S., Stein, F., Behrends, C., and Bhogaraju, S. (2024). A ubiquitin-specific, proximity-based labeling approach for the identification of ubiquitin ligase substrates. Science Advances *10*, eadp3000.

69.   Liwocha, J., Li, J., Purser, N., Rattanasopa, C., Maiwald, S., Krist, D.T., Scott, D.C., Steigenberger, B., Prabu, J.R., and Schulman, B.A. (2024). Mechanism of millisecond Lys48-linked poly-ubiquitin chain formation by cullin-RING ligases. Nature Structural & Molecular Biology *31*, 378-389.

70.   Bai, N., Riching, K.M., Makaju, A., Wu, H., Acker, T.M., Ou, S.-C., Zhang, Y., Shen, X., Bulloch, D.N., and Rui, H. (2022). Modeling the CRL4A ligase complex to predict target protein ubiquitination induced by cereblon-recruiting PROTACs. Journal of Biological Chemistry *298*.

71.   Békés, M., Langley, D.R., and Crews, C.M. (2022). PROTAC targeted protein degraders: the past is prologue. Nature Reviews Drug Discovery *21*, 181-200.

72.   Yamanaka, S., Horiuchi, Y., Matsuoka, S., Kido, K., Nishino, K., Maeno, M., Shibata, N., Kosako, H., and Sawasaki, T. (2022). A proximity biotinylation-based approach to identify protein-E3 ligase interactions induced by PROTACs and molecular glues. Nature Communications *13*, 183.

73.   Yu, H.H., Reitsma, J.M., Sweredoski, M.J., Moradian, A., Hess, S., and Deshaies, R.J. (2019). Single subunit degradation of WIZ, a lenalidomide-and pomalidomide-dependent substrate of E3 ubiquitin ligase CRL4CRBN. BioRxiv, 595389.

74.   Cassandri, M., Smirnov, A., Novelli, F., Pitolli, C., Agostini, M., Malewicz, M., Melino, G., and Raschellà, G. (2017). Zinc-finger proteins in health and disease. Cell death discovery *3*, 1-12.

75.   Teng, M., Lu, W., Donovan, K.A., Sun, J., Krupnick, N.M., Nowak, R.P., Li, Y.-D., Sperling, A.S., Zhang, T., and Ebert, B.L. (2021). Development of PDE6D and CK1α degraders through chemical derivatization of FPFT-2216. Journal of medicinal chemistry *65*, 747-756.

76.   Szklarczyk, D., Kirsch, R., Koutrouli, M., Nastou, K., Mehryary, F., Hachilif, R., Gable, A.L., Fang, T., Doncheva, N.T., and Pyysalo, S. (2023). The STRING database in 2023: protein–protein association networks and functional enrichment analyses for any sequenced genome of interest. Nucleic acids research *51*, D638-D646.

77.   Hornbeck, P.V., Zhang, B., Murray, B., Kornhauser, J.M., Latham, V., and Skrzypek, E. (2015). PhosphoSitePlus, 2014: mutations, PTMs and recalibrations. Nucleic acids research *43*, D512-D520.

78.   Barak, T., Ristori, E., Ercan-Sencicek, A.G., Miyagishima, D.F., Nelson-Williams, C., Dong, W., Jin, S.C., Prendergast, A., Armero, W., and Henegariu, O. (2021). PPIL4 is essential for brain angiogenesis and implicated in intracranial aneurysms in humans. Nature medicine *27*, 2165-2175.

79.   Schmitzová, J., Cretu, C., Dienemann, C., Urlaub, H., and Pena, V. (2023). Structural basis of catalytic activation in human splicing. Nature *617*, 842-850.

80.   Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015). limma powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res *43*, e47. 10.1093/nar/gkv007.

81.   Sanchez-Garcia, R., Gomez-Blanco, J., Cuervo, A., Carazo, J.M., Sorzano, C.O.S., and
Vargas, J. (2021). DeepEMhancer: a deep learning solution for cryo-EM volume post-
processing. Communications biology *4*, 874.
1016