Alternative lipid synthesis in response to phosphate limitation promotes antibiotic

tolerance in Gram-negative ESKAPE pathogens

Roberto Jhonatan Olea-Ozuna[1], Melanie J. Campbell[2], Samantha Y. Quintanilla[1], Sinjini

Nandy[1], Jennifer S. Brodbelt[2], and Joseph M. Boll[1, #]

Affiliations:

[1]Department of Biological Sciences, University of Texas at Dallas, Richardson, TX, USA

[2]Department of Chemistry, University of Texas at Austin, Austin, TX, USA

[#]Corresponding Author:

Email: joseph.boll@utdallas.edu

Short Title: Membrane remodeling in response to phosphate limitation promotes antibiotic

tolerance

Keywords: Gram-negative; phospholipid; antibiotic tolerance; aminolipid; membrane;

bilayer

Abstract

The Gram-negative outer membrane protects bacterial cells from environmental toxins such as antibiotics. The outer membrane lipid bilayer is asymmetric; while glycerophospholipids compose the periplasmic facing leaflet, the surface layer is enriched with phosphate-containing lipopolysaccharides. The anionic phosphates that decorate the cell surface promote electrostatic interactions with cationic antimicrobial peptides such as colistin, allowing them to penetrate the bilayer, form pores, and lyse the cell. Colistin is prescribed as a last-line therapy to treat multidrug-resistant Gram-negative infections.

Acinetobacter baumannii is an ESKAPE pathogen that rapidly develops resistance to antibiotics and persists for extended periods in the host or on abiotic surfaces. Survival in environmental stress such as phosphate scarcity, represents a clinically significant challenge for nosocomial pathogens. In the face of phosphate starvation, certain bacteria encode adaptive strategies, including the substitution of glycerophospholipids with phosphorus-free lipids. In bacteria, phosphatidylethanolamine, phosphatidylglycerol, and cardiolipin are conserved glycerophospholipids that can form lipid bilayers, particularly in the presence of other lipids. Here, we demonstrate that in response to phosphate limitation, conserved regulatory mechanisms induce alternative lipid production in *A. baumannii*. Specifically, phosphate limitation induces formation of three lipids, including amine-containing ornithine and lysine aminolipids. Mutations that inactivate aminolipid biosynthesis exhibit fitness defects relative to wild type in colistin growth and killing assays. Furthermore, we show that other Gram-negative ESKAPE pathogens accumulate aminolipids under phosphate limiting growth conditions, suggesting aminolipid biosynthesis may represent a broad strategy to overcome cationic antimicrobial peptide-mediated killing.

Author Summary

Gram-negative ESKAPE pathogens, including A*cinetobacter baumannii*, are responsible for a dramatic increase in the morbidity and mortality of patients in healthcare settings over the past two decades. Infections are difficult to treat due to antibiotic resistance and tolerance; however, broadly conserved mechanisms that promote antibiotic treatment failure have not been extensively studied. Herein, we identify an alternative lipid biosynthesis pathway that is induced in phosphate starvation that enables Gram-negative ESKAPE pathogens, including *A. baumannii*, *Klebsiella pneumoniae*, and *Enterobacter cloacae* to build lipid bilayers in the absence of glycerophospholipids, which are the canonical bilayers lipid. Replacement of the anionic phosphate in the lipid headgroup with zwitterionic ornithine and lysine promote survival against colistin, a last resort antimicrobial used against Gram-negative infections. These studies suggest that ESKAPE pathogens can remodel their bilayers with phosphate free lipids to overcome colistin treatment and that aminolipid biosynthesis could be targeted to improve antimicrobial treatment.

Introduction

Introduction

The Gram-negative cell envelope consists of a symmetrical bilayer of

glycerophospholipids in the inner membrane, while the outer membrane exhibits an

asymmetrical composition, with glycerophospholipids in the periplasmic leaflet and

lipopolysaccharide enriched in the outer leaflet (1). The intricate organization underscores

the remarkable complexity of bacterial membrane architecture, crucial for microbial survival

in various environments. However, under specific stress conditions such as nutrient

limitation, temperature fluctuations or exposure to antimicrobial agents, certain bacteria

activate alternative lipid biosynthesis pathways or modify existing lipids to adapt and ensure

cellular viability (2). One example is aminolipids, which contain amino acid headgroups like

lysine, glycine, glutamine, and serine-glycine, with ornithine being the most common (3,4).

Ornithine lipids (OLs) are phosphorus-free and found exclusively in bacteria; they are absent

in archaea or eukaryotes (5). Their basic structure comprises a 3-hydroxylated fatty acid

linked by an amide bond to the α-amino group of ornithine and a second fatty acid attached

by an ester bond to the 3-hydroxyl group of the first fatty acid (6). Although OLs are found

in both the inner and outer lipid bilayers of Gram-negative bacteria, they are enriched in the

outer membrane (7–10). OL biosynthesis is catalyzed by two acyltransferases, OlsB and

OlsA, or by the bifunctional acyltransferase OlsF (11–13). In some Gram-negative pathogens

such as *Pseudomonas aeruginosa* or *Vibrio cholerae*, OLs are exclusively formed under

phosphate limiting conditions (14,15), indicating the presence of a specific regulatory

mechanism. The importance of aminolipids transcends basic physiology, especially in the

context of antibiotic resistance. ESKAPE pathogens, a group of pathogens that include

Acinetobacter baumannii, are notorious for their ability to overcome antibiotic treatment and

cause hospital-acquired infections (16). Aminolipid synthesis has been implicated in increased bacterial fitness under antimicrobial stress (9,17,18), potentially contributing to pathogen persistence in clinical settings. Additionally, there is a notable relationship between membrane lipid remodeling and resistance to colistin, a last-resort antibiotic that is used against multi-drug resistant Gram-negative infections (19–22). Chemical modifications to the lipid A domain of lipopolysaccharide or enrichment of amino acid-containing glycerophospholipids have been associated with colistin resistance (23–25), highlighting the importance of understanding lipid metabolism in combating antibiotic resistance.

In this study, we demonstrate that *A. baumannii* produced two aminolipids in limiting phosphate growth conditions, including lysine lipids (LLs) and OLs. OL and LL synthesis is dependent on the *olsB* and *olsA* genes, and *olsB* expression is regulated transcriptionally by the response regulator, PhoR. Additionally, mutants deficient in aminolipid synthesis exhibit increased colistin susceptibility relative to wild type. We also found that other Gram-negative ESKAPE pathogens, including *Klebsiella pneumoniae* and *Enterobacter cloacae*, accumulate aminolipids under phosphate limited growth conditions. These findings suggest a broad survival strategy among ESKAPE pathogens that could promote survival during antibiotic treatment.

Results

Phosphate limitation induces lipid membrane composition modifications in *A. baumannii*

The membrane lipid composition of diverse *A. baumannii* isolates was analyzed, including strains ATCC 17978, ATCC 19606, and AB5075, cultivated in complex lysogeny broth (LB) medium supplemented with ^{32}P-orthophosphoric acid. Labelled cells were collected at mid-logarithmic growth phase, lipids were extracted using the Bligh and Dyer method, and separated by hydrophobicity using two-dimensional thin-layer chromatography (TLC), as previously (20,26,27). We also prepared lipid extracts from *Escherichia coli* K-12 strain W3110 for comparison, which served as a well-characterized standard (28). TLC analysis showed conserved glycerophospholipid enrichment corresponding to known structures, including phosphatidylethanolamine (PE), phosphatidylglycerol (PG), and cardiolipin (CL) (**Figure S1A**). Additionally, *A. baumannii* strains also produced two distinct lipid species, including lyso-PE (20) and an unknown phospholipid, denoted as UPL1, that could be a CL derivative, mono-lyso CL (29). The chemical structures of known phospholipids are shown in **Figure S1B**.

A. baumannii strain ATCC 17978 was cultured in minimal medium with either 1 mM (excess) or 50 μM (limiting) phosphate, conditions commonly used to distinguish between phosphate-sufficient and phosphate-limited growth (6, 13–15, 29, 30). Growth in 1 mM phosphate induced a lipid composition almost equivalent to growth in complex LB medium (**Figure 1 and Figure S1A**). One notable change was that UPL1 was absent, and another unidentified lipid, denoted as UPL2, was formed. To explore lipid biosynthesis under phosphate-limiting conditions, membrane lipid profiles were analyzed after growth in

minimal media supplemented with 50 μM (limiting) phosphate concentrations (**Figure 1**).

Limited phosphate availability impacted relative lipid levels, suggesting decreased

phosphate-containing lipid synthesis. However, unlike PE and PG glycerophospholipids,

relative CL levels did not change significantly, suggesting its production may provide a

strategy to maintain membrane charge stability. Concomitant production of three unknown

lipids, referred to as unknown lipids 1 (U1), 2 (U2), and 3 (U3), were produced in phosphate

limiting growth. Ninhydrin staining revealed that U1 and U2 contained free amines, like PE.

Increased ratios of ninhydrin-stained unknown aminolipids relative to PE showed that under

phosphate-limiting conditions, potential phosphate-free aminolipids were produced. The

total lipid quantifications after lipids were extracted and separated on TLC is shown in

Figure 1: Lipid composition of *A. baumannii* strain ATCC 17978 in excess and limiting phosphate concentrations. Strains were grown in basal media 2 (BM2) with excess (1mM) or limiting (50 μM) phosphate. Cells were collected, lipids were extracted using the Bligh and Dyer method and separated using 2-dimensional thin-layer chromatography. Total lipids were stained with sulfuric acid (left). Aminolipids were stained using ninhydrin (right). Specific lipids are labelled: PE, phosphatidylethanolamine; PG, phosphatidylglycerol; CL, cardiolipin; LPE, lyso-PE; U1, unknown lipid 1; U2, unknown lipid 2; UPL1, unknown phospholipid 1. Red letters denote aminolipids that provide the focus of the study.

Figure S2A.

Growth in limiting phosphate concentrations slowed growth in ATCC 17978 relative

to excess phosphate (**Figure S2B**), and microscopic analysis revealed that cells elongated

and increased their surface area when phosphate is limiting (**Figure S2C and S2D**), a

response previously reported in other Gram-negative bacteria (32). Additionally, while the

composition of lipooligosaccharide (LOS) fractions remained consistent across limiting

phosphate conditions, the relative level of LOS was decreased under phosphate limitation

(**Figure S2E**).

Aminolipids synthesized during phosphate limitation are OLs and LLs

One dimensional TLC showed three lipids (PE, U1, and U2) stained with ninhydrin

(**Figure S3A**). After separation of lipids based on hydrophobicity, individual bands

corresponding with U1 and U2 were scraped from the TLC plates and extracted using the

Bligh and Dyer method. U1 and U2 bands were analyzed by liquid chromatography-mass

spectrometry (LC-MS) and structurally characterized using tandem mass spectrometry

(MS/MS) (**Figure 2**). The elution profiles obtained for the U1 and U2 bands are displayed in

Figures 2A and **2D,** respectively. To determine the composition of aminolipids, a data-

dependent acquisition method was used to isolate and activate the most abundant ions

detected in each chromatographic peak with higher energy collisional dissociation (HCD) in

negative-ionization mode. HCD of precursor *m/z* 621.52 found in the U1 extract resulted in

the loss of the headgroup, which was observed at *m/z* 131.08, and corresponded to

deprotonated ornithine (**Figure 2B**). The most abundant fragment ion (*m/z* 367.30) was used

to identify the acyl chain connected to the headgroup as 16:0 (number of carbon atoms:

double bonds). Further, the complementary ion of *m/z* 253.22 confirmed the identity of the

fatty acid connected at the 3-hydroxyl position of the first fatty acid as 16:1. Following this

analysis, lipids containing a double bond were subsequently targeted in a second LC run in

the positive-ionization mode using 193 nm ultraviolet photodissociation (UVPD) for

MS/MS. UVPD is an alternative fragmentation method that utilizes high-energy photons to

activate and dissociate the selected lipid precursor ions, allowing localization of the double

bonds within the fatty acyl chains. UVPD of precursor m/z 623.53 produced two fragment

ions separated by 24 Da that originate from cleavages adjacent to the double bond. This pair

of diagnostic ions localizes the double bond to the 9th position (**Figure 2C**). This LC-MS/MS

strategy identified 49 OLs, including double bond isomers, and 10 unknown lipids in the U1

extract (**Table S1**) and a total of 24 OLs and 16 unknown lipids in the U2 extract (**Table S2**).

Figure 2: Structural analysis of the lipids produced during phosphate limitation. A. LC-MS trace of U1 lipid extract in negative-ionization mode. **B.** HCD (NCE 22) mass spectrum of m/z 621.52 ([M-H]$^-$), an ornithine lipid found in the U1 extract. **C.** UVPD (4 pulses at 2 mJ/pulse) mass spectrum of m/z 623.53 ([M+H]$^+$). **D.** LC-MS trace of U2 lipid extract in negative-ionization mode. **E.** HCD (NCE 22) mass spectrum of m/z 635.54 ([M-H]$^-$), a lysine lipid found in U2 extract. **F.** UVPD (4 pulses at 2 mJ/pulse) mass spectrum of m/z 637.47 ([M+H]$^+$). The selected precursor ions are labeled with asterisks in B, C, E, and F.

HCD of the unknown lipids yielded similar fragmentation to that observed for OLs, but the

fragmentation patterns were distinguished by the release of a deprotonated headgroup that

corresponded to either a lysine or monomethylated ornithine headgroup (*m/z* 145.10) (**Figure**

2E). While the aminolipid head group could be either lysine or monomethylated ornithine,

the absence of an ortholog for OlsG, the enzyme responsible for OL methylation (33), and

the presence of *olsG* only in certain planctomycete genomes, strongly suggests that OL

methylation is unlikely to occur in *A. baumannii*. Therefore, the identified lipid is denoted

herein as a LL.

After confirming that *A. baumannii* strain ATCC 17978 produces OLs and LLs, we

explored if other *A. baumannii* isolates form these aminolipids during phosphate limitation.

One-dimensional TLC stained with ninhydrin revealed that diverse *A. baumannii* strains,

including ATCC 17978, ATCC 19606, AB5075, AYE, and the environmental isolate, *A.*

baylyi, were also capable of aminolipid biosynthesis in response to phosphate limitation

(**Figure S3B**). Together, these studies suggest that *Acinetobacter* can form lipid bilayers with

not only glycerophospholipids, but also membranes enriched with OLs and LLs when

phosphate availability is limited.

Differentially regulated genes during phosphate starvation

To determine genes that regulate aminolipid biosynthesis, total RNA was isolated,

rRNA was depleted, and transcripts were sequenced from *A. baumannii* strain ATCC 17978

after growth in minimal media supplemented with 1 mM (excess) and 50 µM (limiting)

phosphate. Using a 3-fold cutoff for the weighted proportions fold change and an FDR p-

value correction < 0.05, 67 upregulated genes and 109 downregulated genes were found

(**Figure 3A**). Many down-regulated genes were involved in iron uptake, such as those coding

Figure 3: The *olsB* gene is required for ornithine and lysine lipid formation in phosphate limited growth conditions. A. Volcano plot of differentially regulated genes in excess and limiting phosphate. Lines indicate 3.5-fold cutoffs. **B.** 2D thin-layer chromatography stained with sulfuric acid showing wild type and Δ*olsB* strains grown in phosphate limiting (50 µM) conditions. Specific lipids are labelled: PE, phosphatidylethanolamine; PG, phosphatidylglycerol; CL, cardiolipin; OL, ornithine lipid; LL, lysine lipid; U3, unknown lipid 3; U4, unknown lipid 4. OL and LL aminolipids are labelled in red. **C.** Proposed ornithine lipid (OL) and lysine lipid (LL) biosynthesis pathways in *A. baumannii*.

for ferric uptake regulators (Fur) and components of the ABC transporters for iron.

Additionally, genes involved in lysine degradation and aminoacid metabolism were down-

regulated, indicating potential changes in nitrogen metabolism under phosphate-limited

conditions. Upregulated genes were associated with TAT-dependent proteins export, which

could be involved in the transport of key lipids or membrane proteins, as well as

phospholipases, which may play a role in membrane remodeling. Furthermore, genes related

to regulation and phosphate transporters were induced, likely in response to changes in

phosphate availability. A complete list of significant ($P < 0.05$) up- and down-regulated genes

is included in **Table S3**. The RNA-sequencing results were validated using qRT-PCR

analysis to measure differential expression between three specific genes in excess and

limiting phosphate concentrations (**Figure S4A)** and via phenotypic characterization of

several mutants described below. Validation of the dataset suggested that phosphate

limitation induces a broad transcriptional reprogramming, affecting processes such as lipid

biosynthesis, membrane integrity, and nutrient uptake, which are crucial for bacterial survival

under stress conditions.

One notable up-regulated gene (*A1S_0889*) showed a $\log_2$ fold change of 8.57 and

31% identity and 51% similarity (91% coverage) to *P. aeruginosa olsB*. While not induced

in phosphate starvation, *A. baumannii* also encodes a putative *olsA* orthologue (*A1S_2990*)

with 43% identity and 61% similarity (78% coverage) to *P. aeruginosa olsA*. OlsBA has been

characterized in several bacteria and is involved in OLs biosynthesis (8,11,12,14,34).

Notably, *A. baumannii olsA* is located at a distinct site on the chromosome relative to *olsB*

and is not transcriptionally regulated by phosphate concentrations.

olsB* and *olsA* genes are required for aminolipids synthesis in *A. baumannii

Differential gene expression analysis indicated changes in gene dosage in response to

excess/limiting phosphate concentration. Specifically, *A1S_0889* expression increased in

response to phosphate limitation, whereas *A1S_2990* expression did not change. To confirm

that *A1S_0889* was an *olsB* orthologue, we generated Δ*olsB* in *A. baumannii* strain ATCC

17978, by deleting codons 34 and 214. TLC analysis revealed that under phosphate

limitation, aminolipids were absent in the *olsB*-deficient mutant and a mutant carrying only

the empty vector but wild-type accumulated OLs and LLs (**Figure 3B**). Furthermore, *olsB*

complementation from a non-native promoter restored OL and LL formation. Based on the

lipid migration patterns in wild type, the lipids are OLs and LLs.

Additionally, we generated Δ*olsB* (*HMPREF0010_01383*) using recombineering in

strain ATCC 19606 and also analyzed the *olsB::tn* and *olsA::tn* mutants from the AB5075

transposon mutant library (35), which contain insertions in the respective genes,

ABUW_3039 and *ABUW_0502*. Lipid analysis when the mutants were cultivated under

phosphate limitation showed that the wild-type strains ATCC 19606 and AB5075 produced

OLs and LLs, while the *olsB* and *olsA* mutants did not (**Figure S4B and S4C**). The *olsA*::*tn*

mutant lipid profile did not show the expected accumulation of lyso-aminolipids, consistent

with findings in other Gram-negative Δ*olsA* mutants (30). This suggests that lyso-aminolipids

are tightly regulated and rapidly degraded within the cell. These data suggested that ornithine

and lysine lipid biosynthesis in *A. baumannii* is dependent on OlsB and OlsA. We will also

note that we observed an unknown lipid, labelled as U4 in figure 3B and figure S4B that may

indicate an OlsA-dependent lipid.

Together, the data supports a model suggesting that aminolipid synthesis occurs in at

least two distinct steps within *A. baumannii* (**Figure 3C**), consistent with previous work in

other Gram-negative bacteria (11,12). Initially, ornithine and lysine undergo acylation in an

OlsB-dependent reaction, leading to the formation of lyso-OL and lyso-LL. Subsequently, in

a second step facilitated by OlsA, lyso-OL and lyso-LL are further acylated at the hydroxy

position, yielding OL and LL, respectively.

Microscopic analysis and comparison of LOS fractions and levels between

aminolipid-producing and aminolipid-deficient *A. baumannii* strains revealed no significant

morphological differences (**Figure S4D and S4E**) or changes in LOS fractions and relative

levels (**Figure S2E**). However, it is noteworthy that the core fraction of LOS from *A.*

baumannii strains ATCC 19606 differs from of the ATCC 17978 or AB5075 strains.

Additionally, a faint band in the core fraction was observed in the *olsA*::*tn* transposon mutant

of AB5075, which was not present in the wild type strain.

Mutants deficient in aminolipids synthesis show differential growth rate under

phosphate limitation

To assess growth rates under phosphate limitation, wild-type and *olsB* mutant strains

were grown in minimal medium with limiting phosphate concentrations (**Figure S4F**). The

optical density at 600 nm (OD_{600}) was monitored over time. While growth of strain ATCC

17978 $\Delta olsB$ was not impacted, the growth rate in strains ATCC 19606 and AB5075 was

reduced in the *olsB* or *olsA* T101 mutants, suggesting there are strain-dependent effects on

aminolipid biosynthesis that impact fitness.

Aminolipid biosynthesis mutants are defective in colistin tolerance

Changes in lipid composition could alter the physicochemical properties of the

bilayers, particularly the charge. To explore this concept, the impact of aminolipids

biosynthesis on colistin tolerance was measured in *A. baumannii* strain ATCC 17978, where

growth rate was not impacted in the *olsB* mutant (**Figure S4F**). Colistin is a cyclic peptide

that directly engages with the negative membrane charge, while the hydrophobic tail forms

pores, leading to bactericidal activity. Colistin is a last resort antibiotic used against multidrug

resistant Gram-negative bacterial infections. Our studies indicate increased colistin

susceptibility in the *olsB*-deficient mutant relative to the wild-type strain (**Figure 4A and**

Figure S5A), suggesting the antibiotic could be more effective against *A. baumannii* when

aminolipid biosynthesis is inhibited. Under high phosphate conditions, the membrane lipid

composition remains typical of standard bacterial membranes, resulting in higher relative

colistin susceptibility (Figure 4C). However, in phosphate limited conditions, lipid turnover

Figure 4: Aminolipids promote colistin tolerance in *A. baumannii*. A. Colistin-dependent killing in wild type and Δ*olsB* mutant strains under phosphate limitation (*n* = 3). Error bars indicate standard deviation. Wild type (WT) strains carrying pOlsB or empty vector were subjected to 5 mg/L colistin exposure over time. CFU/mL were calculated every 0.5 h. **B.** Total lipids were extracted using the Bligh and Dyer method and separated using 2D thin-layer chromatography. Lipids were stained with sulfuric acid (left). Aminolipids were stained using ninhydrin (right). Specific lipids are labelled: PE, phosphatidylethanolamine; PG, phosphatidylglycerol; CL, cardiolipin; LPE, lyso-phosphatidylethanolamine; LL, lysine lipid; OL, ornithine lipid; U3, unknown lipid 3. OL and LL are labelled in red. **C.** Colistin-dependent kill curves (left) and growth rate analysis (right) in wild type expressing empty vector (pEV) or pOlsB under high phosphate conditions (*n* = 3). Error bars indicate standard deviation.

increases, leading to the replacement of phospholipids with non-phosphorylated aminolipids,

which probably contributes to enhanced colistin tolerance. Overexpression of *olsB* in wild-

type *A. baumannii* was sufficient to produce aminolipids during cultivation in excess

phosphate (**Figure 4B**). Analysis demonstrated that *olsB* induction led to increased colistin

tolerance (**Figure 4C and Figure S5B**). Therefore, OlsB-dependent aminolipid biosynthesis

in *A. baumannii* promotes colistin tolerance, a last-line antibiotic for combating multidrug-

resistant Gram-negative bacteria.

PhoR regulates aminolipid biosynthesis in *A. baumannii*

303 Previous work established that aminolipid production generally occurs in response to

304 phosphate limitation (6,14,15,36,37). Consequently, we expected to find a Pho box upstream

305 of the putative *olsB* gene. Using the *E. coli* pho box consensus sequence (38), a putative Pho

306 box was identified that precedes the *olsB* gene in diverse *Acinetobacter* isolates, including

Figure 5: PhoR regulates *olsB* gene expression and aminolipid formation in *A. baumannii* strains.
A. Predicted Pho box sequence alignments from *A. baumannii* strain 17978, 19606, AB5075, AYE, and *A. baylyi olsB* promoters. Black asterisks represent conserved nucleotides in the *E. coli* Pho box consensus (CT*GTCAT*NNNNC*TGTCAT*). **B.** 2D thin-layer chromatography lipid analysis from *A. baumannii* strain ATCC 17978 wild type and mutants grown in minimal media supplemented with 50 µM (limiting) phosphate. **C.** 2D thin-layer chromatography lipid analysis from *A. baumannii* strain AB5075 wild type and the *phoR* Tn*26* mutant in minimal media supplemented with 50 µM (limiting) phosphate. Lipids were stained with sulfuric acid. OL and LL aminolipids are labelled in red.

ATCC 17978 (**Figure 5A**), but not preceding the *olsA* gene. The identified Pho box shows

significant sequence similarity to the *E. coli* consensus (CT*GTCAT*NNNNC*TGTCAT*), with

conserved nucleotides present in multiple *Acinetobacter* strains, including *A. baumannii*

strains ATCC 17978, 19606, AB5075, AYE, and *A. baylyi*. These results are consistent with

our transcriptomics analysis, which also suggested *olsA* gene expression is not responsive to

phosphate concentrations.

Phosphate sensing and diverse responses are regulated by the two-component system,

PhoB/PhoR, in many bacteria (39). When phosphate levels decrease, the sensor kinase, PhoR,

becomes activated, leading to autophosphorylation. Subsequent phosphotransfer to the

cognate response regulator, PhoB results in a conformational change and DNA binding (40)

at target promoters to induce gene expression.

These findings imply that the PhoB/PhoR two-component system regulates OL and

LL biosynthesis in *A. baumannii* through *olsB* expression. To validate this hypothesis, we

generated Δ*phoR* (*A1S_3376*) in *A. baumannii* ATCC 17978 by deleting codons 66 to 409.

Lipid analysis after growth limiting phosphate, showed that the Δ*phoR* mutant accumulates

glycerophospholipids PE, PG, CL, and UPL2, while failing to synthesize OL, LL, and U3,

unlike the wild-type strain (**Figure 5B**). Complementation of the *phoR*-deficient mutant with

the PhoR allele restored OL and LL formation, while aminolipids were absent in the mutant

carrying the empty vector. Expression of *olsB* from a non-native promotor in the Δ*phoR*

mutant restored OL and LL biosynthesis. Additionally, we examined the lipid patterns of the

phoR (*ABUW_0105*)-transposon mutant of *A. baumannii* AB5075 grown under phosphate

limitation (**Figure 5C**). Lipidomic analysis revealed that, unlike the wild-type strain, the

phoR::tn mutant predominantly accumulated phospholipids and was unable to produce OL

Figure 6: Aminolipid biosynthesis is conserved in Gram-negative ESKAPE pathogens. 2D thin-layer chromatography of lipids extracted from wild type *K. pneumoniae* strain KPNIH1, *E. cloacae* strain ATCC 13047, and *P. aeruginosa* strain PAO1 grown in excess (1 mM) or limiting (50 µM) phosphate conditions. Totals lipids were stained with sulfuric acid (left). Aminolipids were stained using ninhydrin (right). Specific lipids are labelled: PE, phosphatidylethanolamine; PG, phosphatidylglycerol; CL, cardiolipin; OL, ornithine lipid; U1, U2, U3, U4, and U5; unknown compounds 1, 2, 3, 4, and 5. **B.** Representative MS1 spectra from shotgun analysis of animolipids isolated from *K. pneumoniae* ($n = 3$) and *E. cloacae* ($n = 3$). The five most enriched lipids are bolded, including two ornithine lipids (OL), labelled in red.

or LLs. Together, these findings suggest that *olsB* expression in *A. baumannii* is mediated by

the *phoR* regulatory gene.

Other Gram-negative ESKAPE pathogens form aminolipids under conditions of

phosphate depletion

In addition to *P. aeruginosa* (14), *A. baumannii* is the second Gram-negative

ESKAPE bacterium where the OL biosynthesis has been described using a OlsBA-dependent

mechanisms. Uniquely, *A. baumannii* also produces LL via the same pathway. These findings

prompted us to investigate if other Gram-negative ESKAPE pathogens, such as *Klebsiella* or

Enterobacter, are also capable of aminolipid biosynthesis under phosphate limitation. TLC

analysis of total lipids from *K. pneumoniae*, *E. cloacae* or *P. aeruginosa* after growth in

minimal medium supplemented with excess phosphates showed production of the canonical

membrane phospholipids PE, PG, and CL (**Figure 6A**). However, when cultivating *P.*

aeruginosa in phosphate limitation (50 μM), a significant decrease in relative phospholipid

levels was observed, and the synthesis of five unknown compounds and OL was induced. *K.*

pneumoniae and *E. cloacae* cultivated in low phosphate concentrations (50 μM), also showed

decreased levels of phospholipid production and accumulation of an unknown lipid. The

unknown lipid exhibited a migration pattern and ninhydrin staining on TLC like OL,

suggesting that aminolipid biosynthesis may be a conserved response to phosphate limiting

growth conditions that could promote tolerance to antibiotics in Gram-negative ESKAPE

pathogens.

To confirm the putative OL identity, total lipids were extracted using the Bligh and

Dyer method after cells were grown in 50 μM phosphate. 1-D TLC (**Figure S6**) on total lipids

stained with ninhydrin showed two prevalent lipids (lane 1). The band corresponding to OLs

was scraped from the plate and extracted. Extracted lipids from three replicates were run on

1-D TLC showing enrichment of the putative OL relative to PE. Negative-ionization mode

MS1 shotgun analysis on the purified lipids showed five enriched lipids in each strain

(**Figure 6B**). MS/MS analysis of the lipids present in *K. pneumoniae* sample revealed two

ornithine lipids (*m/z* 595.50 and 623.53), two PE lipids (*m/z* 662.47 and 702.51), and a *m/z*

consistent with phosphatidylthreonine (*m/z* 760.51) (**Table S4**). Spectra from the *E. cloacae*

lipid analysis included two ornithine lipids (*m/z* 623.54 and 595.50), PE (*m/z* 702.51), a *m/z*

consistent with phosphatidylthreonine (*m/z* 760.51), and an unknown lipid that may be a

modified OL (*m/z* 681.54) (**Table S5**). Together the migration of the lipids via 2-D TLC and

the MS analysis provide strong complementary evidence that growth in limiting phosphate

results in alternative synthesis of OLs.

Discussion

Bacteria have evolved various regulatory mechanisms to sense and adapt to

environmental stress. Here, we show that under phosphate-limiting conditions, *A. baumannii*

produces ornithine and lysine lipids through regulated expression of *olsB*. The aminolipids

alter the lipid bilayer composition to promote tolerance to colistin, an antimicrobial peptide

used to treat Gram-negative bacterial infections. Specifically, aminolipids could reduce the

electrostatic potential for cationic antimicrobial peptides such as colistin to target the cell,

emphasizing changes in bilayer charge as an adaptive mechanism for *A. baumannii* in

challenging environments. Although aminolipids are not directly exposed on the membrane

surface, their accumulation in the inner leaflet of the outer membrane may alter the overall

membrane dynamics, such as lipid packing and membrane fluidity, contributing to enhanced

membrane stability and colistin tolerance. Broadly, cell membrane lipid modifications are

vital for bacterial survival upon exposure to antimicrobial cationic peptides. For example, *V.*

cholerae modifies lipid A with glycine or diglycine residues to resist cationic antimicrobial

peptides (41). Similarly, other Gram-negative bacteria modify their lipid A structures to resist

polymyxins. For instance, *P. aeruginosa* and *A. baumannii* can add positively charged amino

groups, such as ethanolamine to lipid A, reducing to overall negative charge and decreasing

the binding affinity of polymyxin (20). *K. pneumoniae* and *Enterobacter* species have been

shown to add 4-amino-4-deoxy-L-arabinose (L-Ara4N) to lipid A, further neutralizing the

negative charge and conferring resistance (21). Similar strategies are employed by various

bacterial species, such as *P. aeruginosa*, *Rhizobium tropici*, *Staphylococcus aureus*,

Mycobacterium tuberculosis, and *Bacillus subtilis*, which modify phospholipids like PG by

adding amino acids like lysine or alanine, thereby conferring polymyxins resistance

(36,37,42–44).

Membrane lipid remodeling during growth under phosphate limitation is a conserved

strategy across bacteria, involving the substitution of phospholipids with phosphorus-free

lipids alternatives like aminolipids (45). Interestingly, other bacteria, such as *P. aeruginosa*,

also modify their lipid composition in response to phosphate limitation (14). Under these

conditions, *P. aeruginosa* incorporates OL, replacing phospholipids in the membrane. This

remodeling likely ensures that the membrane remains functional and stable despite the

phosphate scarcity, highlighting a similar strategy employed by various bacterial species to

adapt to nutrient stress. These findings further support the idea the replacement of

phospholipids with amino acid-derived lipids, such as ornithine, represents an adaptive

mechanism to phosphate limitation, suggesting that this process may be widely conserved

across Gram-negative bacteria. While the synthesis of these lipids is not directly induced by

the presence of polymyxins, tolerance to polymyxins is an indirect consequence of lipid

membrane remodeling, leading to significant changes in membrane chemical properties. The

alteration could reduce the net negative charge of the membrane, thereby decreasing CAMP

susceptibility.

The *olsB* gene is required for OL biosynthesis, but unlike other OlsB-dependent

pathways, it also induces LL formation in *A. baumannii*, suggesting metabolic plasticity

within the species. Comparing our findings with observations in other bacteria, such as the

soil bacterium *Rhodobacter sphaeroides*, which not only produces OLs but also synthesizes

glutamine lipids (GLs) (46), further highlights the metabolic diversity in aminolipid

biosynthesis. Additionally, recent research has shown that marine bacteria *Ruegeria*

pomeroyi encode two *olsB* paralogs, one responsible for forming OLs and the other for GLs

(30). Although these bacteria inhabit vastly different environments, the similarities suggest

that the ability to use multiple substrates for aminolipid synthesis may be a common survival

strategy in variable environmental conditions.

The transcriptomic analysis uncovered insights into different expression patterns of

the *olsB* and *olsA* genes in response to phosphate limitation. Specifically, overexpression of

the *olsB* gene was observed during phosphate limited growth, indicating a regulatory

mechanism. However, *olsA* expression remained unchanged. The predicted Pho box in the

promoter region of *olsB* suggested that the PhoB/PhoR two-component system regulates *olsB*

under phosphate limitation conditions. The absence of a Pho box-like motif in the promotor

region of *olsA*, in contrast to *olsB*, suggests that *olsA* is regulated by factors distinct from

those controlling *olsB* under phosphate limiting conditions. This was confirmed by RNA

sequencing (**Figure 3A**) and RT-PCR analysis (**Figure S4A**), which both showed that *olsA*

expression remains unchanged in response to phosphate limitation, indicating that its

expression is likely controlled by other regulatory mechanisms or constitutive. Consequently,

olsA is not constitutively expressed under these conditions, but rather its regulation appears

to be independent of phosphate availability. The absence of a similar motif in the promoter

region of *olsA* implies the influence of other regulatory factors on its expression, independent

of phosphate availability. Furthermore, *olsB* overexpression with non-native promoter in *A.*

baumannii during cultivation in excess phosphate resulted in aminolipid formation,

suggesting OlsA-dependent activity occurs after OlsB has formed a lyso-aminolipid. The

differential regulatory events may also suggest that OlsA utilizes other substrates. For

instance, in *Rhodobacter capsulatus*, OlsA functions as a bifunctional enzyme, active in both

OL and phosphatidic acid biosynthesis (47). Additionally, we confirmed the role of the *phoR*

gene as a key regulator in aminolipid biosynthesis in *A. baumannii*, highlighting its fine-

tuned regulation in response to environmental stressors such as phosphate limitation.

Interestingly, this regulatory mechanism shares similarities with species like *Sinorhizobium*

meliloti or *V. cholerae*, where PhoB/PhoR also regulates OLs synthesis under phosphate

limitation conditions (6,15), underscoring the evolutionary conservation of these adaptive

mechanisms across different bacterial taxa.

Finally, the presence of *olsB* and *olsA* orthologues in *A. baumannii* raises questions

about the diversity of aminolipid biosynthesis pathways among pathogens. Interestingly,

bacteria such as *Klebsiella* and *Enterobacter* lack these orthologs, suggesting distinct

biosynthesis pathways compared to *A. baumannii*. This absence prompts further investigation

into alternative pathways or genes involved in aminolipid synthesis and their implications

for antibiotics resistance and environmental adaptation.

Conclusion

The study highlights the impact of phosphate limitation on lipid membrane

composition in *A. baumannii*, resulting in the synthesis of OLs and LLs through regulated

olsB gene expression. Aminolipids can promote tolerance to colistin, an important last-line

antimicrobial. These results also underscore the role for *phoR* gene in regulating aminolipid

synthesis in *A. baumannii* and show that other Gram-negative ESKAPE pathogens produce

aminolipids under phosphate-depleted conditions. Aminolipid biosynthesis is a common

adaptive response to phosphate limitation, which could promote pathogen survival both in

hospital environment and within the host.

Materials and methods

Bacterial strains and growth. All strains and plasmids used in this study are listed in **Table**

S6 in the supplemental material. *E. coli*, *Acinetobacter* strains, *K. pneumoniae*, *E. cloacae* or

P. aeruginosa were initially cultured from frozen stocks on Luria-Bertani (LB Miller) agar

at 37ºC. Isolated colonies were used to inoculate LB Miller broth or minimal medium (Tris

minimal succinate [TMS]) (31); supplemented with different phosphate concentrations at

37ºC. The minimal medium included: Na-succinate 20 mM, NaCl 200 mg/mL, NH_4Cl 450

mg/mL, $CaCl_2$ 200 mg/mL, KCl 200 mg/mL, $MgCl_2$ 450 mg/mL, $FeCl_2$ 10 mg/L, and $MnCl_2$

10 mg/L, with 10 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) buffer

used at pH 7.2. Na_2HPO_4 was then added to achieve a final concentration of 1 mM (excess)

or 50 μM (limiting). All components were dissolved in deionized water and sterilized by

filtration trough 0.22 μm pore-size filters and dissolved deionized H_2O. To generate growth

curves, overnight cultures of *A. baumannii* strains were diluted to an OD_{600} of ~0.05 in 5 mL

of medium, then incubated in glass test tubes at 37ºC for 24 hours. Growth curve data were

analyzed and plotted using GraphPad Prism software.

A. baumannii strain ATCC 17978 has been shown to be a mixture of two variants

(48). Colony PCR using primers specific to the cardiolipin synthase gene (*clsC2*) resulted in

a PCR product, indicating this study was specific to only the *A. baumannii* 17978 UN variant.

A. baumannii strains AB5075 and ATCC 19606 have been shown to produce colonies with

two opacity phenotypes that are phase variable (49). Microscopic examination showed a

majority of opaque forms in strains AB5075 and ATCC 19606 with only 2-3 translucent

colonies out of > 200 colonies found on each plate (*n* = 3 for each strain). The strain ATCC

19606 Δ*olsB* and strain AB5075 *olsB::tn* and *olsA::tn* mutants were also examined, but only

included opaque variants.

Construction of mutant and complementation *A. baumannii* strains. The primers utilized

in this study are listed in **Table S7** in the supplementary material. Genetic mutants of *A.*

baumannii were generated following established protocols (50–53). Briefly, *A. baumannii*

carrying the pMMB67EH^{TetR} plasmid containing the REC$_{Ab}$ coding sequences was diluted

from an overnight culture into LB Miller broth containing 10μg/mL of tetracycline at an

OD$_{600}$ of ~0.05 and incubated for 45 minutes. REC$_{Ab}$ expression was induced by adding 2

mM IPTG, and cells were cultured at 37ºC until they reached mid-log growth phase (OD$_{600}$

of ~0.4). After washing the cells three times in ice-cold 10% glycerol, 10^{10} cells were

electroporated in a 2-mm cuvette at 1.8 mV with 5 μg of a recombineering linear PCR

product. Subsequently, the cells were cultured for 4 hours in 4 mL of LB Miller broth with 2

mM IPTG and then plated on LB agar supplemented with 20 μg/mL of kanamycin. Mutations

were validated using PCR.

To cure isolated mutants of the pMMB67EH^{TetR}::REC$_{Ab}$ plasmid after mutant

isolation, strains were streaked for isolated colonies on LB Miller agar supplemented with 2

mM NiCl$_2$ to select cells that had lost the tetracycline cassette (51–54). Cured insertion

mutants were then electroporated with pMMB67EH^{TetR} carrying the FLP recombinase. Cells

were recovered for 1 hour in 5 mL of LB Miller broth and plated on LB agar containing

10µg/mL of tetracycline and 2 mM IPTG to induce expression of the FLP recombinase.

Excision of the kanamycin cassette was confirmed by PCR.

To complement the *A. baumannii* ATCC 17978 mutants, the coding sequence from

A1S_0889 (*olsB*) was cloned into the KpnI/SalI sites, while the coding sequence from

A1S_3376 (*phoR*) was cloned into the KpnI/BamHI sites in pMMB67EH^{KanR}. Plasmids were

expressed in the respective mutants, and all strains were grown in 30 µg/mL of kanamycin

and 1 mM IPTG for expression.

Analysis of total lipids and aminolipids. Overnight cultures were used to inoculate 10 mL

minimal medium 1 mM phosphate, or 20 mL minimal medium 50 µM phosphate to achieve

an OD$_{600}$ of ~0.05, and then incubated for 24 hours at 37 ºC. After the incubation period,

cells were collected by centrifugation. Lipids were extracted using the Bligh and Dyer (1959)

method (55). The chloroform phase was separated into the individual components on high-

performance TLC silica gel plates. For one-dimensional TLC analysis, the plates were

developed with chloroform-methanol-water (130:50:8 v/v) mixture. For two-dimensional

TLC analysis, a chloroform-methanol-water (140:60:10 v/v) mixture was used in the first

dimension and chloroform-methanol-acetic acid (130:50:20 v/v) mixture was used in the

second dimension. Lipids on TLC were visualized by treating the plates with 10% sulfuric

acid in ethanol at 150ºC (total lipids) or 0.2% ninhydrin in acetone at 100ºC (aminolipids).

Lipid quantifications were done using ImageJ software to detect spots above background.

Each experiment was independently replicated three times, one representative data set was

reported in the quantification, and one representative image was included..

Analysis of ^{32}P-labeled phospholipids. Overnight cultures were diluted to an OD_{600} of ~0.05

in 10 mL LB Miller broth supplemented with 5 µCi/mL ^{32}P ortho-phosphoric acid

(PerkinElmer) and grown until reaching an OD_{600} of ~0.6. After harvesting the cells, lipid

extraction was performed using Bligh and Dyer method (55), followed by analysis using TLC

as previously described. Subsequently, the TLC plates were dried, exposed to a

phosphorimaging screen, and scanned using an Amersham Typhoon laser scanner.

Shotgun Mass Spectrometry: Dried lipid content from Bligh and Dyer extracted TLC bands

of *E. cloacae* WT and *K. pneumoniae* WT biological replicates were solubilized in

chloroform and resuspended in methanol: isopropanol (1:1) at a concentration of 200 µM.

Shotgun analysis was performed in negative-ionization mode on a Thermo Scientific

Orbitrap Fusion Lumos Tribrid mass spectrometer. Nanoelectrospray ionization was

performed using Pd/Au-coated pulled-tip glass capillaries (20 nm i.d.) at a spray voltage of -

1000 V and a capillary temperature of 275°C. Spectra were acquired at a resolution of 50,000

at *m/z* 200, an AGC target between 5E4 and 1E5, and an RF lens of 60%. All MS1 and

MS/MS spectra were an average of 50 scans. The top 5 most abundant species were subjected

to higher energy collisional dissociation (HCD) for identification. MS/MS spectra were

obtained with an isolation width of 1 *m/z* and a normalized collision energy (NCE) ranging

from 22-26%.

Liquid chromatography: Lipids isolated from *Acinetobacter baumannii* were separated

using reversed-phase liquid chromatography (LC) on an Acquity UPLC CSH C18 column

(pore size 130 Å, 1.7 µm particle size, 2.1 mm × 100 mm, Waters) integrated with a Dionex

Ultimate 3000 UHPLC system (Thermo Fisher Scientific). Mobile phase A and B were

comprised of methanol: water: acetonitrile (3:4:3) and isopropanol: water: acetonitrile

(90:2.5:7.5), respectively, each containing 10 mM ammonium formate. Dried lipid content

from TLC bands was extracted via a Bligh and Dyer protocol and resuspended in 50:50

mobile phase (A: B) at a concentration of 100 ng/µL. A 9 µL injection volume was used, and

the column compartment was maintained at a temperature of 50 °C. Aminolipids were

separated at a flow rate of 0.275 mL/min with the following gradient: hold at 10% B (0-1

min), 10-45% B (1-5 min), 45-70% B (5-23 min), 70-95% B (23-24 min), hold at 95% B (24-

29 min), 95-10% B (29-29.5), and hold at 10% B (29.5-35 min).

Untargeted LC-MS with HCD and Targeted LC-MS with UVPD Experiments: An

untargeted negative-ionization mode LC-MS with HCD method was utilized to screen

Acinetobacter baumannii lipid extracts on a Thermo Scientific Orbitrap Fusion Lumos

Tribrid mass spectrometer via heated electrospray ionization. Various source parameters

included a spray voltage of -3800 V, sheath gas setting of 5, aux gas setting of 10, and

capillary temperature of 300 °C. MS1 data was collected at a resolution of 30,000 at *m/z* 200

in the Orbitrap analyzer with a scan range of *m/z* 500-1000, RF lens of 80%, and an AGC

target of 5E5. A data-dependent acquisition method was used where the top 5 most abundant

ions above an intensity threshold of 5E4 were selected for MS/MS analysis with HCD.

Species were isolated using a 1 *m/z* window and subjected to a collision energy corresponding

to NCE of 22%. MS/MS spectra were acquired at a resolution of 30,000 at *m/z* 200 in the

Orbitrap analyzer, AGC target of 1E5, maximum injection time of 250 ms, and 2

microscans/scan.

A targeted positive-ionization mode LC-MS/MS method was performed on a Thermo

Scientific Orbitrap Eclipse Tribrid mass spectrometer equipped with an ArF excimer laser

(Coherent, Inc.) for 193 nm ultraviolet photodissociation (UVPD). The heated electrospray

ionization source was operated at + 3800 V, and the source parameters described in the

untargeted experiments were implemented. MS/MS scans with UVPD were acquired in a

data-dependent manner, with 5 scans collected between each MS1 master scan, and with an

intensity threshold set at 5E4. A targeted mass filter was employed that included the *m/z*

values and start/end retention times for unsaturated aminolipids identified from the negative-

ionization mode LC-MS with HCD run (with a 15-ppm error tolerance). Aminolipids were

isolated with a 1 *m/z* window and activated with 4 laser pulses at 2 mJ/pulse for double bond

localization. MS/MS spectra were acquired at a resolution of 30,000 at *m/z* 200 in the

Orbitrap analyzer, q-value of 0.1, AGC target of 5E5, maximum injection time of 500 ms,

and 5 microscans/scan. Data was manually interpreted using Thermo Xcaliber Qual Browser

and ChemDraw. Precursor and fragment ions were assigned within 10 and 15 ppm mass

error, respectively.

RNA sequencing. Transcriptome sequencing analysis was performed as described

previously, with modification (53). Briefly, total RNA was extracted from *A. baumannii*

ATCC 17978 cultures grown in minimal medium supplemented with either excess (1 mM)

or limiting (50 μM) phosphate at OD_{600} of ~0.5 in triplicate, utilizing the Direct-Zol RNA

miniprep kit (Zymo Research). Genomic DNA contamination was eliminated using the

Turbo DNA-free DNA removal kit (Invitrogen). DNAase-treated RNA samples were then

forwarded to SeqCenter for sequencing on the Illumina NextSeq 550 sequencing.

Subsequently, the CLC Genomic Workbench software (Qiagen) was employed to map the

obtained sequencing data to the *A. baumannii* ATCC 17978 genome annotations and

determine the read per kilobase per million (RPKM) expression values and determine the

weighted-proportions fold changes in expression values between excess or limitation

phosphate conditions. Data were analyzed and plotted using GraphPad Prism software. Data

Accession #: GSE276010.

Quantitative RT-PCR. Relative-abundance quantitative PCR (qPCR) was performed as

previously described In brief, total RNA was extracted from bacterial cultures using the

Direct-Zol RNA Miniprep Kit (Zymo Research), followed by DNase digestion to remove

any genomic DNA. The RNA was then reverse transcribed into cDNA using the Invitrogen

SuperScript III Reverse Transcriptase (Thermo Fisher), with a final RNA concentration of 5

ng/µl. One microliter of the resulting cDNA was used as a template in a 10-µl quantitative

RT-PCR (qRT-PCR) reaction, conducted with Power SYBR Green Master Mix (Applied

Biosystems). qRT-PCR was carried out on a QuantStudio 3 Real-Time PCR System (Thermo

Fisher Scientific). Relative expression levels were calculated using the ΔΔCt method (56),

with normalization of gene targets to the *rpoA* signal.

Microscopy and image analysis. Cells were grown as stated above and fixed with

paraformaldehyde (PFA) and mounted on 1.5% agarose in 1X phosphate-buffered saline

(PBS). Imaging was performed using a Nikon Eclipse Ti-2 wide-field epifluorescence

microscope equipped with a Photometrics Prime 95B camera and a Plan Apo 100x, 1.45

numerical aperture objective lens. Images were captured using NIS Elements software.

Image analysis was conducted using the microbeJ plugin of ImageJ software.

LOS staining and analysis. All cultures were grown in test tubes containing 5 mL of minimal medium with either excess (1 mM) or limiting (50 µM) phosphate at 37°C in a shaker overnight. For the complementation strains, 30 µg/mL kanamycin and 1 mM IPTG were added. The OD_{600} of the overnight cultures was measured and normalized to OD_{600} of ~1. The cells were then centrifuged at 15,000 rpm for 5 minutes. Each pellet was resuspended in 100 µl of 1X Sample Buffer (4X LDS Sample Buffer, 4% β-mercaptoethanol, and water) and boiled in water for 10 minutes. After cooling, proteinase K was added to each sample and mixed. The samples were then incubated in a 55°C water bath overnight. The following day, the samples were boiled in water for 5 minutes and SDS-PAGE was performed. The gel was then fixed and treated according to the protocol in the Pro-Q Emerald 300 Lipopolysaccharide Gel Stain Kit by Thermo Fisher Scientific (P20495).

Colistin susceptibility assays. Overnight cultures were diluted to OD_{600} ~0.150 in minimal medium supplemented with either excess (1 mM) or limiting (50 µM) phosphate containing 5 mg/L colistin. Each culture, comprising 15 mL, was incubated in 125 mL Erlenmeyer flasks at 37ºC with agitation at 250 rpm. Survivors were analyzed at specific time points by serial dilution plating on LB agar.

Funding

The work was supported by funding from the National Institutes of Health (grants R35GM143053 and R01AI168159 to J.M.B and R35GM139658 to J.S.B), and the Robert A. Welch Foundation (F-1155 to J.S.B.). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1.   Raetz CRH, Whitfield C. Lipopolysaccharide endotoxins. *Annu Rev Biochem.*
2002;71:635–700.

2.   Sohlenkamp C, Geiger O. Bacterial membrane lipids: diversity in structures and
pathways. *FEMS Microbiol Rev.* 2016 Jan;40(1):133–59.

3.   Geiger O, González-Silva N, López-Lara IM, Sohlenkamp C. Amino acid-containing
membrane lipids in bacteria. *Prog Lipid Res.* 2010 Jan;49(1):46–60.

4.   Moore EK, Hopmans EC, Rijpstra WIC, Villanueva L, Damsté JSS. Elucidation and
identification of amino acid containing membrane lipids using liquid
chromatography/high-resolution mass spectrometry. *Rapid Commun Mass Spectrom*
RCM. 2016 Mar 30;30(6):739–50.

5.   Vences-Guzmán MÁ, Geiger O, Sohlenkamp C. Ornithine lipids and their structural
modifications: from A to E and beyond. *FEMS Microbiol Lett.* 2012 Oct;335(1):1–10.

6.   Geiger O, Röhrs V, Weissenmayer B, Finan TM, Thomas-Oates JE. The regulator gene
phoB mediates phosphate stress-controlled synthesis of the membrane lipid
diacylglyceryl-N,N,N-trimethylhomoserine in *Rhizobium (Sinorhizobium) meliloti. Mol*
Microbiol. 1999 Apr;32(1):63–73.

7.   Dees C, Shively JM. Localization of quantitation of the ornithine lipid of *Thiobacillus*
thiooxidans. J Bacteriol. 1982 Feb;149(2):798–9.

8.   Palacios-Chaves L, Conde-Álvarez R, Gil-Ramírez Y, Zúñiga-Ripa A, Barquero-Calvo
E, Chacón-Díaz C, et al. *Brucella abortus* ornithine lipids are dispensable outer
membrane components devoid of a marked pathogen-associated molecular pattern. *PloS*
One. 2011 Jan 7;6(1):e16030.

9. Vences-Guzmán MÁ, Guan Z, Ormeño-Orrillo E, González-Silva N, López-Lara IM,

Martínez-Romero E, et al. Hydroxylated ornithine lipids increase stress tolerance in

Rhizobium tropici CIAT899. *Mol Microbiol.* 2011 Mar;79(6):1496–514.

10. Stirrup R, Mausz MA, Silvano E, Murphy A, Guillonneau R, Quareshy M, et al.

Aminolipids elicit functional trade-offs between competitiveness and bacteriophage

attachment in *Ruegeria pomeroyi. ISME J.* 2023 Mar;17(3):315–25.

11. Weissenmayer B, Gao JL, López-Lara IM, Geiger O. Identification of a gene required

for the biosynthesis of ornithine-derived lipids. *Mol Microbiol.* 2002 Aug;45(3):721–33.

12. Gao JL, Weissenmayer B, Taylor AM, Thomas-Oates J, López-Lara IM, Geiger O.

Identification of a gene required for the formation of lyso-ornithine lipid, an intermediate

in the biosynthesis of ornithine-containing lipids. *Mol Microbiol.* 2004 Sep;53(6):1757–

70.

13. Vences-Guzmán MÁ, Guan Z, Escobedo-Hinojosa WI, Bermúdez-Barrientos JR, Geiger

O, Sohlenkamp C. Discovery of a bifunctional acyltransferase responsible for ornithine

lipid synthesis in *Serratia proteamaculans. Environ Microbiol.* 2015 May;17(5):1487–

96.

14. Lewenza S, Falsafi R, Bains M, Rohs P, Stupak J, Sprott GD, et al. The *olsA* gene

mediates the synthesis of an ornithine lipid in *Pseudomonas aeruginosa* during growth

under phosphate-limiting conditions, but is not involved in antimicrobial peptide

susceptibility. *FEMS Microbiol Lett.* 2011 Jul;320(2):95–102.

15. Barbosa LC, Goulart CL, Avellar MM, Bisch PM, von Kruger WMA. Accumulation of

ornithine lipids in *Vibrio cholerae* under phosphate deprivation is dependent on VC0489

(OlsF) and PhoBR system. *Microbiol Read Engl.* 2018 Mar;164(3):395–9.

16. Harding CM, Hennon SW, Feldman MF. Uncovering the mechanisms of *Acinetobacter*

baumannii virulence. *Nat Rev Microbiol.* 2018 Feb;16(2):91–102.

17. Rojas-Jiménez K, Sohlenkamp C, Geiger O, Martínez-Romero E, Werner D, Vinuesa P.

A ClC chloride channel homolog and ornithine-containing membrane lipids of

Rhizobium tropici CIAT899 are involved in symbiotic efficiency and acid tolerance. *Mol*

Plant-Microbe Interact MPMI. 2005 Nov;18(11):1175–85.

18. González-Silva N, López-Lara IM, Reyes-Lamothe R, Taylor AM, Sumpton D, Thomas-

Oates J, et al. The dioxygenase-encoding *olsD* gene from *Burkholderia cenocepacia*

causes the hydroxylation of the amide-linked fatty acyl moiety of ornithine-containing

membrane lipids. *Biochemistry.* 2011 Jul 26;50(29):6396–408.

19. Boll JM, Tucker AT, Klein DR, Beltran AM, Brodbelt JS, Davies BW, et al. Reinforcing

Lipid A Acylation on the Cell Surface of *Acinetobacter baumannii* Promotes Cationic

Antimicrobial Peptide Resistance and Desiccation Survival. *mBio.* 2015;6(3):e00478-

00415.

20. Boll JM, Crofts AA, Peters K, Cattoir V, Vollmer W, Davies BW, et al. A penicillin-

binding protein inhibits selection of colistin-resistant, lipooligosaccharide-deficient

Acinetobacter baumannii. Proc Natl Acad Sci U S A. 2016 11;113(41):E6228–37.

21. Kang KN, Klein DR, Kazi MI, Guérin F, Cattoir V, Brodbelt JS, et al. Colistin

heteroresistance in *Enterobacter cloacae* is regulated by PhoPQ-dependent 4-amino-4-

deoxy-l-arabinose addition to lipid A. *Mol Microbiol.* 2019;111(6):1604–16.

22. Murtha AN, Kazi MI, Schargel RD, Cross T, Fihn C, Cattoir V, et al. High-level

carbapenem tolerance requires antibiotic-induced outer membrane modifications. *PLoS*

Pathog. 2022 Feb;18(2):e1010307.

23. Ernst CM, Peschel A. Broad-spectrum antimicrobial peptide resistance by MprF-
mediated aminoacylation and flipping of phospholipids. *Mol Microbiol.* 2011
Apr;80(2):290–9.

24. Needham BD, Trent MS. Fortifying the barrier: the impact of lipid A remodelling on
bacterial pathogenesis. *Nat Rev Microbiol.* 2013 Jul;11(7):467–81.

25. Henderson JC, Zimmerman SM, Crofts AA, Boll JM, Kuhns LG, Herrera CM, et al. The
Power of Asymmetry: Architecture and Assembly of the Gram-Negative Outer
Membrane Lipid Bilayer. *Annu Rev Microbiol.* 2016 08;70:255–78.

26. de Rudder KE, Thomas-Oates JE, Geiger O. Rhizobium meliloti mutants deficient in
phospholipid N-methyltransferase still contain phosphatidylcholine. *J Bacteriol.* 1997
Nov;179(22):6921–8.

27. Kamischke C, Fan J, Bergeron J, Kulasekara HD, Dalebroux ZD, Burrell A, et al. The
Acinetobacter baumannii Mla system and glycerophospholipid transport to the outer
membrane. *eLife.* 2019 14;8.

28. Raetz CR, Dowhan W. Biosynthesis and function of phospholipids in *Escherichia coli.*
J Biol Chem. 1990 Jan 25;265(3):1235–8.

29. Lopalco P, Stahl J, Annese C, Averhoff B, Corcelli A. Identification of unique cardiolipin
and monolysocardiolipin species in *Acinetobacter baumannii. Sci Rep.* 2017 Jun
7;7(1):2972.

30. Smith AF, Rihtman B, Stirrup R, Silvano E, Mausz MA, Scanlan DJ, et al. Elucidation
of glutamine lipid biosynthesis in marine bacteria reveals its importance under
phosphorus deplete growth in *Rhodobacteraceae. ISME J.* 2019 Jan;13(1):39–49.

31. Jones RA, Shropshire H, Zhao C, Murphy A, Lidbury I, Wei T, et al. Phosphorus stress
induces the synthesis of novel glycolipids in *Pseudomonas aeruginosa* that confer
protection against a last-resort antibiotic. *ISME J.* 2021 Nov;15(11):3303–14.

32. Stankeviciute G, Guan Z, Goldfine H, Klein EA. *Caulobacter crescentus* Adapts to
Phosphate Starvation by Synthesizing Anionic Glycoglycerolipids and a Novel
Glycosphingolipid. *mBio. 2*019 Apr 2;10(2):e00107-19.

33. Escobedo-Hinojosa WI, Vences-Guzmán MÁ, Schubotz F, Sandoval-Calderón M,
Summons RE, López-Lara IM, et al. OlsG (Sinac_1600) Is an Ornithine Lipid N-
Methyltransferase from the Planctomycete *Singulisphaera acidiphila. J Biol Chem.* 2015
Jun 12;290(24):15102–11.

34. Kim SK, Park SJ, Li XH, Choi YS, Im DS, Lee JH. Bacterial ornithine lipid, a surrogate
membrane lipid under phosphate-limiting conditions, plays important roles in bacterial
persistence and interaction with host. *Environ Microbiol.* 2018 Nov;20(11):3992–4008.

35. Gallagher LA, Ramage E, Weiss EJ, Radey M, Hayden HS, Held KG, et al. Resources
for Genetic and Genomic Analysis of Emerging Pathogen *Acinetobacter baumannii. J*
Bacteriol. 2015 Jun 15;197(12):2027–35.

36. Hebecker S, Arendt W, Heinemann IU, Tiefenau JHJ, Nimtz M, Rohde M, et al. Alanyl-
phosphatidylglycerol synthase: mechanism of substrate recognition during tRNA-
dependent lipid modification in *Pseudomonas aeruginosa. Mol Microbiol.* 2011
May;80(4):935–50.

37. Arendt W, Hebecker S, Jäger S, Nimtz M, Moser J. Resistance phenotypes mediated by
aminoacyl-phosphatidylglycerol synthases. *J Bacteriol.* 2012 Mar;194(6):1401–16.

38. Baek JH, Lee SY. Novel gene members in the Pho regulon of *Escherichia coli. FEMS*
Microbiol Lett. 2006 Nov;264(1):104–9.

39. Hsieh YJ, Wanner BL. Global regulation by the seven-component Pi signaling system.

Curr Opin Microbiol. 2010 Apr;13(2):198–203.

40. Gardner SG, McCleary WR. Control of the phoBR Regulon in *Escherichia coli. EcoSal*

Plus. 2019 Sep;8(2).

41. Hankins JV, Madsen JA, Giles DK, Brodbelt JS, Trent MS. Amino acid addition to

Vibrio cholerae LPS establishes a link between surface remodeling in gram-positive and

gram-negative bacteria. *Proc Natl Acad Sci U S A.* 2012 May 29;109(22):8722–7.

42. Sohlenkamp C, Galindo-Lagunas KA, Guan Z, Vinuesa P, Robinson S, Thomas-Oates J,

et al. The lipid lysyl-phosphatidylglycerol is present in membranes of *Rhizobium tropici*

CIAT899 and confers increased resistance to polymyxin B under acidic growth

conditions. *Mol Plant-Microbe Interact MPMI.* 2007 Nov;20(11):1421–30.

43. Klein S, Lorenzo C, Hoffmann S, Walther JM, Storbeck S, Piekarski T, et al. Adaptation

of *Pseudomonas aeruginosa* to various conditions includes tRNA-dependent formation

of alanyl-phosphatidylglycerol. *Mol Microbiol.* 2009 Feb;71(3):551–65.

44. Ernst CM, Staubitz P, Mishra NN, Yang SJ, Hornig G, Kalbacher H, et al. The bacterial

defensin resistance protein MprF consists of separable domains for lipid lysinylation and

antimicrobial peptide repulsion. *PLoS Pathog.* 2009 Nov;5(11):e1000660.

45. Geiger O, Padilla-Gómez J, López-Lara IM. Bacterial Sphingolipids and Sulfonolipids.

In: Geiger O, editor. Biogenesis of Fatty Acids, Lipids and Membranes [Internet]. Cham:

Springer International Publishing; 2019 [cited 2024 Jul 2]. p. 123–37. Available from:

https://doi.org/10.1007/978-3-319-50430-8_12

46. Zhang X, Ferguson-Miller SM, Reid GE. Characterization of ornithine and glutamine

lipids extracted from cell membranes of *Rhodobacter sphaeroides. J Am Soc Mass*

Spectrom. 2009 Feb;20(2):198–212.

47. Aygun-Sunar S, Bilaloglu R, Goldfine H, Daldal F. *Rhodobacter capsulatus* OlsA is a

bifunctional enzyme active in both ornithine lipid and phosphatidic acid biosynthesis. *J*

Bacteriol. 2007 Dec;189(23):8564–74.

48. Wijers CDM, Pham L, Menon S, Boyd KL, Noel HR, Skaar EP, et al. Identification of

Two Variants of *Acinetobacter baumannii* Strain ATCC 17978 with Distinct Genotypes

and Phenotypes. *Infect Immun.* 2021 Nov 16;89(12):e0045421.

49. Tipton KA, Dimitrova D, Rather PN. Phase-Variable Control of Multiple Phenotypes in

Acinetobacter baumannii Strain AB5075. *J Bacteriol.* 2015 Jul 6;197(15):2593–9.

50. Tucker AT, Nowicki EM, Boll JM, Knauf GA, Burdis NC, Trent MS, et al. Defining

gene-phenotype   relationships   in   *Acinetobacter   baumannii*   through   one-step

chromosomal gene inactivation. *mBio.* 2014;5(4):e01313-01314.

51. Kang KN, Kazi MI, Biboy J, Gray J, Bovermann H, Ausman J, et al. Septal Class A

Penicillin-Binding Protein Activity and ld-Transpeptidases Mediate Selection of

Colistin-Resistant Lipooligosaccharide-Deficient *Acinetobacter baumannii. mBio.* 2021

Jan 5;12(1):e02185-20.

52. Kang KN, Boll JM. PBP1A Directly Interacts with the Divisome Complex to Promote

Septal Peptidoglycan Synthesis in *Acinetobacter baumannii. J Bacteriol.* 2022 Dec

20;204(12):e0023922.

53. Islam N, Kazi MI, Kang KN, Biboy J, Gray J, Ahmed F, et al. Peptidoglycan Recycling

Promotes Outer Membrane Integrity and Carbapenem Tolerance in *Acinetobacter*

baumannii. mBio. 2022 Jun 28;13(3):e0100122.

54. Podolsky T, Fong ST, Lee BT. Direct selection of tetracycline-sensitive *Escherichia coli*

cells using nickel salts. *Plasmid.* 1996 Sep;36(2):112–5.

55. Bligh EG, Dyer WJ. A rapid method of total lipid extraction and purification. *Can J*

Biochem Physiol. 1959 Aug;37(8):911–7.

56. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time

quantitative PCR and the 2(-Delta Delta C(T)) *Method.* Methods San Diego Calif. 2001

Dec;25(4):402–8.

Figure 1: Lipid composition of *A. baumannii* strain ATCC 17978 in excess and limiting phosphate concentrations. A. Strains were grown in basal media 2 (BM2) with excess (1mM) or limiting (50 µM) phosphate. Cells were collected, lipids were extracted using the Bligh and Dyer method and separated using 2-dimensional thin-layer chromatography. Total lipids were stained with sulfuric acid (left). Aminolipids were stained using ninhydrin (right). Specific lipids are labelled: PE, phosphatidylethanolamine; PG, phosphatidylglycerol; CL, cardiolipin; LPE, lyso-PE; U1, unknown lipid 1; U2, unknown lipid 2; UPL1, unknown phospholipid 1. Red letters denote aminolipids that provide the focus of the study.

Figure 2: Structural analysis of the lipids produced during phosphate limitation. A. LC-MS trace of U1 lipid extract in negative-ionization mode. **B.** HCD (NCE 22) mass spectrum of *m/z* 621.52 ([M-H]⁻), an ornithine lipid found in the U1 extract**. C.** UVPD (4 pulses at 2 mJ/pulse) mass spectrum of *m/z* 623.53 ([M+H]⁺). **D.** LC-MS trace of U2 lipid extract in negative-ionization mode. **E.** HCD (NCE 22) mass spectrum of *m/z* 635.54 ([M-H]⁻), a lysine lipid found in U2 extract. **F.** UVPD (4 pulses at 2 mJ/pulse) mass spectrum of *m/z* 637.47 ([M+H]⁺). The selected precursor ions are labeled with asterisks in B, C, E, and F.

Figure 3: The *olsB* gene is required for ornithine and lysine lipid formation in phosphate limited growth conditions. A. Volcano plot of differentially regulated genes in excess and limiting phosphate. Lines indicate 3.5-fold cutoffs. **B.** 2D thin-layer chromatography stained with sulfuric acid showing wild type and Δ*olsB* strains grown in phosphate limiting (50 μM) conditions. Specific lipids are labelled: PE, phosphatidylethanolamine; PG, phosphatidylglycerol; CL, cardiolipin; OL, ornithine lipid; LL, lysine lipid; U3, unknown lipid 3; U4, unknown lipid 4. OL and LL aminolipids are labelled in red. **C.** Proposed ornithine lipid (OL) and lysine lipid (LL) biosynthesis pathways in *A. baumannii*.

Figure 4: Aminolipids promote colistin tolerance in *A. baumannii*. A. Colistin-dependent killing in wild type and Δ*olsB* mutant strains under phosphate limitation (*n* = 3). Error bars indicate standard deviation. Wild type (WT) strains carrying pOlsB or empty vector were subjected to 5 mg/L colistin exposure over time. CFU/mL were calculated every 0.5 h. **B.** Total lipids were extracted using the Bligh and Dyer method and separated using 2D thin-layer chromatography. Lipids were stained with sulfuric acid (left). Aminolipids were stained using ninhydrin (right). Specific lipids are labelled: PE, phosphatidylethanolamine; PG, phosphatidylglycerol; CL, cardiolipin; LPE, lyso-phosphatidylethanolamine; LL, lysine lipid; OL, ornithine lipid; U3, unknown lipid 3. OL and LL are labelled in red. **C.** Colistin-dependent kill curves (left) and growth rate analysis (right) in wild type expressing empty vector (pEV) or pOlsB under high phosphate conditions (*n* = 3). Error bars indicate standard deviation.

Figure 5: PhoR regulates *olsB* gene expression and aminolipid formation in *A. baumannii* strains.
A. Predicted Pho box sequence alignments from *A. baumannii* strain 17978, 19606, AB5075, AYE, and *A. baylyi olsB* promoters. Black asterisks represent conserved nucleotides in the *E. coli* Pho box consensus (CT*GTCAT*NNNNC*TGTCAT*). **B.** 2D thin-layer chromatography lipid analysis from *A. baumannii* strain ATCC 17978 wild type and mutants grown in minimal media supplemented with 50 µM (limiting) phosphate. **C.** 2D thin-layer chromatography lipid analysis from *A. baumannii* strain AB5075 wild type and the *phoR* Tn*26* mutant in minimal media supplemented with 50 µM (limiting) phosphate. Lipids were stained with sulfuric acid. OL and LL aminolipids are labelled in red.

Figure 6: Aminolipid biosynthesis is conserved in Gram-negative ESKAPE pathogens. 2D thin-layer chromatography of lipids extracted from wild type *K. pneumoniae* strain KPNIH1, *E. cloacae* strain ATCC 13047, and *P. aeruginosa* strain PAO1 grown in excess (1 mM) or limiting (50 µM) phosphate conditions. Totals lipids were stained with sulfuric acid (left). Aminolipids were stained using ninhydrin (right). Specific lipids are labelled: PE, phosphatidylethanolamine; PG, phosphatidylglycerol; CL, cardiolipin; OL, ornithine lipid; U1, U2, U3, U4, and U5; unknown compounds 1, 2, 3, 4, and 5. **B.** Representative MS1 spectra from shotgun analysis of animolipids isolated from *K. pneumoniae* ($n = 3$) and *E. cloacae* ($n = 3$). The five most enriched lipids are bolded, including two ornithine lipids (OL), labelled in red.

Supplementary Captions

Figure S1: *Escherichia coli* (*Ec*) **and** *Acinetobacter baumannii (Ab)* **lipid composition**

after growth in complex media. A. Strains were grown in complex media (LB broth) in the

presence of ^{32}P-orthopohosphoric acid until mid-logarithmic growth phase. Cells were

collected and total lipids were extracted using the Bligh and Dyer method and separated using

2-dimensional thin-layer chromatography. Labelled lipids include anionic cardiolipin (CL)

and phosphatidylglycerol (PG), aminolipids phosphatidylethanolamine (PE) and lyso-PE

(LPE), and unknown phospholipid 1 (UPL1). **B.** PG, PE, CL and LPE chemical structures.

Head groups are red, the glycerol backbone is blue, and fatty acids are black.

Figure S2: Effect of phosphate availability on growth, cell morphology, and LOS

production. A. Quantification of total lipid in TLC (A) grown in excess or limiting

phosphate concentrations. Lipids are graphed as a percentage of the total. Significance testing

conducted using Student *t* test with two-tailed distribution assuming equal variance. Lines

indicate standard deviation. *$P < 0.05$; ns is not significant. **B.** *A. baumannii* ATCC 17978

was cultured in minimal medium containing either excess (1 mM) or limiting (50 µM)

phosphate for 24 hours. **C.** Phase-contrast images of *A. baumannii* grown under these

conditions, captured during the exponential phase. Scalebar is 10 µm. **D.** Cell length and area

were quantified for each population ($n \geq 150$) using ImageJ software. Each point represents

an individual cell. The experiment was repeated twice, and one representative dataset was

reported. Significance testing conducted using Student *t* test with two-tailed distribution

assuming equal variance. ***$P < 0.001$, ****$P < 0.0001$. **E.** Proteinase K-treated whole-cell

lysate from wild-type *A. baumannii* strains ATCC 17978, ATCC 19606, and AB5075, as

well as from aminolipid-deficient mutants grown under excess (1mM) or limiting (50 µM)

phosphate conditions. The LOS samples were separated using SDS-PAGE and stained using

Pro-Q Emerald 300.

Figure S3: Thin-layer chromatography of *Acinetobacter* aminolipids. A. To produce *A.*

baumannii lipid samples for MS analysis, total lipids were separated using thin-layer

chromatography and scraped from the plate, extracted using the Bligh and Dyer method, and

run alongside lipid controls. Extractions resulted in isolation of U1 & U2, U1, or U2. Lipids

were stained with ninhydrin to visualize aminolipids. Specific lipids are labelled: PE,

phosphatidylethanolamine; U1, unknown lipid 1; U2, unknown lipid 2. **B.** Cells were grown

in minimal medium supplemented with excess (1mM) or limiting (50 µM) phosphate

conditions from indicated *Acinetobacter* strains. Total lipids were extracted from cells grown

in media with limiting phosphate concentrations. Total lipids were spotted on thin-layer

chromatography and separated based on hydrophobicity. Plates were stained with ninhydrin

to visualize aminolipids. Specific lipids are labelled: PE, phosphatidylethanolamine; OL,

ornithine lipid; LL, lysine lipid. **C.** To produce *E. cloacae* and *K. pneumoniae* samples for

MS analysis, total lipids were extracted from cells grown in media with limiting phosphate

concentrations. Total lipids were separated using thin-layer chromatography and scraped

from the plate, extracted using the Bligh and Dyer method, and run alongside lipid controls.

Lipids were stained with ninhydrin to visualize aminolipids. Specific lipids are labelled: PE,

phosphatidylethanolamine; OL; ornithine lipid (predicted).

Figure S4: *olsB* and *olsA* are required for ornithine and lysine lipid biosynthesis in *A.*

***baumannii*. A.** Relative-abundance quantitative PCR (qPCR) of genes after *A. baumannii*

growth in excess or limiting phosphate concentrations (n = 3). Lines indicate standard

deviation. Significance testing was conducted using Student t test with two-tailed distribution

assuming equal variance. *P < 0.05; ns = not significant. **B.** 2D thin-layer chromatography

lipid analysis in ATCC 19606 wild type and the Δ*olsB* mutant strain after growth in limiting

(50 µM) phosphate concentrations. Lipids were stained with sulfuric acid. **C.** Analysis in

AB5075 wild type and transposon (Tn*101*) mutant strains after growth in limiting (50 µM)

phosphate concentrations. Specific lipids are labelled: PE, phosphatidylethanolamine; PG,

phosphatidylglycerol; CL, cardiolipin; OL, ornithine lipid; LL, lysine lipid. OL and LL

aminolipids are labelled in red. **D.** Phase-contrast images of *A. baumannii* strain ATCC

17978 grown under phosphate limiting conditions, captured during exponential phase

growth. Scalebar is 10 µm. **E.** Cell length and area of strain ATCC 17978 were quantified

for each population ($n \geq 150$) using ImageJ software. Each point represents an individual

cell. Significance testing was conducted using Student t test with two-tailed distribution

assuming equal variance. ns = not significant. **F.** Optical density (OD_{600}) measurements of

A. baumannii strains grown in 50 µM phosphate over 24 h.

Figure S5: Aminolipid formation promotes *A. baumannii* tolerance to colistin. Growth

(OD_{600}) of *A. baumannii* ATCC 17978 strains was measured at 37 °C in minimal medium

with limiting phosphate (**A**) or excess phosphate (**B**), and in the presence of colistin at

concentrations of 1, 2.5, or 5 mg/L.

Figure S6: Thin-layer chromatography of *E. cloacae* and *K. pneumoniae* aminolipids.

To produce *E. cloacae* and *K. pneumoniae* samples for MS analysis, total lipids were

extracted from cells grown in media with limiting phosphate concentrations. Total lipids were

separated using thin-layer chromatography and scraped from the plate, extracted using the

Bligh and Dyer method, and run alongside lipid controls. Lipids were stained with ninhydrin

to visualize aminolipids. Specific lipids are labelled: PE, phosphatidylethanolamine; OL;

ornithine lipid (predicted).

Table S1: Identity of aminolipids in unknown lipid 1 biological replicates.

Table S2: Identity of aminolipids in unknown lipid 2 (U2) biological replicates.

Table S3: Differentially regulated genes in excess and limiting phosphate

Table S4: Lipids present in the top five most abundant species from *K. pneumoniae*

biological replicates

Table S5: Lipids present in the top five most abundant species from *E. cloacae* biological

replicates

Table S6: Strains and plasmids used in this study.

Table S7: Primers used in this study.