Title: Mutations in legume genes that influence symbiosis create a complex selective landscape for rhizobial symbionts

Sohini Guha[1], Regina B. Bledsoe[1], Jeremy Sutherland[1], Brendan Epstein[2], Gwendolyn M. Fry[1], Vikram Venugopal[3], Siva Sankari[3], Alejandra Gil Polo[1], Garrett Levin[4], Barney Geddes[4] , Nevin D. Young[5], Peter Tiffin[2], and Liana T. Burghardt[1, *]

[1]The Pennsylvania State University, Department of Plant Science, University Park, Pennsylvania

[2]University of Minnesota, Department of Plant and Microbial Biology, St. Paul, Minnesota

[3]Stowers Institute for Medical Research, Kansas City, MO, USA

[4]North Dakota State University, Department of Department of Microbiological Sciences, Fargo, ND 58102

[5]University of Minnesota, Department of Plant Pathology, St. Paul, Minnesota

[*]Corresponding author (liana.burghardt@psu.edu)

Author for correspondence: Liana T. Burghardt, Department of Plant Science, 109 Tyson Building, University Park, PA 16802, liana.burghardt@psu.edu,1-(814)-863-6168

Running Title: Rhizobial fitness across Medicago mutants

Abstract
In the mutualism between leguminous plants and rhizobia bacteria, rhizobia live inside root
nodules, creating potential for host genes to shape the rhizobial selective environment. Many
host genes that affect symbiosis have been identified; however, the extent to which these genes
affect selection acting on rhizobia is unknown. In this study, we inoculated 18 *Medicago*
truncatula symbiotic mutants (including mutants that alter Nodule Cysteine-Rich (NCR) peptide
production, plant defense, and nodule number regulation) with a mixture of 86 *Sinorhizobium*
meliloti strains. Most mutations resulted in reduced host benefits, but the effects on rhizobial
benefit (i.e., relative strain fitness) varied widely, revealing widespread host-by-strain fitness
interactions. Genome-wide association analyses identified variants on rhizobial replicons pSymA
and pSymB as important in mediating strain fitness responses to host mutations. Whereas most
top variants affected rhizobial fitness with one host mutant (limited effect variants), nine affected
fitness across six or more host mutations. These pervasive variants occurred primarily on
pSymA, the symbiotic replicon, and include *fixL* and some metabolic genes. In contrast to the
limited effect variants, variants with pervasive positive effects on strain fitness when host genes
were mutated tended to adversely affect fitness in wild-type hosts. Competition assays across
Medicago genotypes confirmed a pervasive role for one candidate (malonyl-CoA synthase), and
AlphaFold multimer modelling suggests that many rhizobial top candidates could interact with
host NCR peptides. Our results reveal how host genetic variation affects rhizobial fitness, setting
the stage for improving rhizobial inoculants and breeding legume hosts better adapted to multi-
strain environments.
47

Introduction
Many bacteria live in close association with eukaryotic hosts, and a subset of these bacterial taxa
significantly influence host fitness[1]. If host genetic variation alters selective pressures on
bacteria, that variation can shape bacterial evolution. These genome x genome (GxG)
interactions are an essential prerequisite for co-evolution, and a potential mechanism for
maintaining genetic variation in bacterial and host populations[2]. Most examples of (GxG)
interaction involve pathogens, including a host immune receptor and microbial targets (e.g.,
human HLA and HIV epitopes [3]) or plant R genes [4]. However, examinations of quantitative
pathogen resistance leveraging techniques such as co-GWAS (Genome Wide Association
Studies) [5,6] suggest complex genetic architectures underlie these host-pathogen interactions [7,8].
More recently, studies of (GxG) interactions have expanded to mutualisms [9–12]. These studies
suggest that (GxG) interactions in mutualisms may be equally, if not more, genetically complex
than in parasitism [13–18]. To what extent and in what ways host genes alter bacterial selection has
important implications for bacterial evolution and deploying beneficial microbes in translational
contexts such as agriculture [19,20]. Here, we take a novel approach to measuring (GxG)
interactions by measuring how over a dozen host mutations alter the selective landscape
experienced by scores of strains of mutualistic bacteria.
65
Legume hosts and rhizobium bacteria are a tractable system for studying how individual host
genes alter the selection of their microbial mutualists. Rhizobia inhabit diverse environments,

including soil, the rhizosphere, and root nodules, where they experience selection pressures that
shape their fitness [21]. In nodules, rhizobia convert atmospheric nitrogen into plant assimilable
forms and receive photosynthates for growth [22,23]. Host genetic variation can influence rhizobial
fitness within root nodules [16,24–26] in ways likely shaped by nodule development and subsequent
nitrogen fixation [27]. Decades of work have resulted in extensive collections of mutant lines in
model legumes, including *Lotus* and *Medicago*, which have been exhaustively used to dissect the
functional genetics of nodule formation and function. Most research on these mutants has
focused on their effect on host benefits (e.g., shoot biomass) and investment (e.g., nodule
number/size). Far less attention has been paid to the effect of host mutations on rhizobia [28].
However, doing so is necessary to clarify the potential for host genes to shape the rhizobial
fitness.
79
Measuring the effect of host genes on rhizobia requires considering three aspects of rhizobial
fitness—absolute fitness, relative fitness, and nodule strain diversity. Because most nodules are
initiated by a single rhizobium, rhizobial population sizes in nodules indicate the rewards an
individual bacterium receives from forming a nodule [28,29]. However, measuring population size
alone does not enable an estimate of fitness difference among strains. Assessing strain-level
relative fitness requires comparing strain frequencies before and after exposure to a selective
environment, such as host nodules. A Select and Resequence (S&R) approach leverages whole-
genome shotgun (WGS) sequencing to enable high-throughput tracking of strain frequencies
[25,30]. This method allows for the direct quantification of selection pressures on rhizobia, revealing
how selective an environment is (nodule strain diversity) and which strains are enriched or
depleted compared to the initial community (relative strain fitness) [25,30]. A key advantage of the
S&R method is that, because all strains are already sequenced, GWAS can be used to identify
genetic variants associated with rhizobial relative fitness [16,25,30,31].
93
Work in the legume rhizobium mutualism suggests that host genetic variation can alter these
metrics of rhizobial fitness [16,24,25,30]. For instance, host mutations that increase plant defence
responses (*nad1*) reduce the number of rhizobia released from each nodule, whereas disrupting
genes that cause hosts to form more nodules (*sunn1, sunn4, rdn1*) can increase rhizobia released
at the plant level [32]. Other host genes like *rj2* and *rj4* in soybean restrict the diversity of strains
that form nodules [33,34]. Disruption of the *Sinorhizobium* gene encoding HRRP peptidase,
potentially involved in processing host NCR peptides, can restrict the rhizobial strains in nodules
by modulating host discrimination or defence thresholds [35]. Lastly, disruptions within a host gene
can alter *relative strain fitness*. For instance, members of the NPD gene family in *Medicago*
truncatula differentially influenced which strain frequencies increased in nodules relative to the
inoculated strain community [30]. Taken together, these studies highlight the potential for single
host genes to reshape the selective environment and alter the evolution of rhizobia.
106
Here, we characterize how genetically disrupting host genes alters selection on their nitrogen-
fixing rhizobium symbionts *in planta*. To accomplish this goal, we chose eighteen existing
Medicago truncatula mutant lines with disruptions in the known symbiotic genes. These genes

affect a wide range of host phenotypes and encompass the developmental trajectory of
nodulation. We inoculated these mutants with a mixture of 86 *Sinorhizobium meliloti* strains
(Fig. 1). We studied standard host investment and benefit traits, as well as multiple aspects of
rhizobial fitness (Table 1). We then used a S&R approach to measure the frequencies of rhizobial
strains (relative strain fitness) in nodules [25] and followed up with GWAS, to identify variants in
rhizobium genes that altered strain fitness when host genes were perturbed. We examined the
following questions: 1) Which host symbiosis genes alter rhizobial fitness? 2) Do altered host
genotypes have strain-specific effects on rhizobial fitness (i.e., is there evidence of genome x
genome interactions)? 3) If yes, which rhizobium genes harbour genetic variation underlying the
observed shifts in strain fitness, and are any of those variants selected in multiple host mutations?
And 4) Are any host investment or benefit traits correlated with rhizobial fitness traits? Whereas
few host genotypes altered rhizobial population sizes or nodule strain diversity, many hosts did
shift relative strain fitness. Most rhizobial genes associated with strain fitness shifts were found
to be important within specific hosts, although eighteen variants, (primarily on pSymA) were
important with six or more hosts, suggesting candidate genes that may play fundamental roles in
in-plant rhizobial fitness.
126
127 Materials and methods
128
Host symbiotic mutants: The development of symbiosis in nodules is complex, with each step
regulated by multiple hosts and rhizobium genes [27,36,37]. To encompass these steps, we chose 18
M. truncatula symbiosis mutants (Table S1), which we grouped into three categories: early-
acting, late-acting, and regulation of nodule number (Autoregulation of nodule number; AON) [38].
The early-acting genes (*hcl, ipd3, latd, dmi1-2, dmi2-3,* and *dmi3-1*) affect the initial processes,
such as Nod Factor recognition and nodule organogenesis [39–43]. The late-acting genes (*nad1,*
dnf1, dnf2, dnf4, dnf2/dnf1, dnf5/dnf2, dnf6, dnf7) influence nitrogen fixation [32,44], whereas the
regulation of nodule number genes are in the AON (Autoregulation Of Nodulation) pathway
(*sunn1, sunn4, rdn1*) [45,46]. Whereas these categorizations help set expectations for the impact of
the host mutations, each gene may have additional pleiotropic effects. Together, these mutants
provide a broad view of the effects of host symbiosis genes on their rhizobial partners in multi-
strain inoculations. Seeds for all host mutant lines were obtained from Dr. Dong Wang,
Department of Biochemistry and Molecular Biology, University of Massachusetts, Amherst. The
host mutant lines used here were generated through forward genetic screens and thus may carry
additional background mutations. Our aim is not to assign strict causality to individual host
mutations but rather to examine how alterations in the host genetic landscape broadly influence
rhizobial fitness.
146
Initial inoculum preparation: We created a mixed inoculum using 86 sequenced *Sinorhizobium*
meliloti (henceforth *S. meliloti*) strains, which were part of a collection isolated from soils from
21 locations in Spain and France, using the nodules of *Medicago* plants as a "trap" for capturing
S. meliloti isolates [47]. We grew each strain separately for 72 hours in Tryptone-Yeast extract

(TY) media at 28°C, shaking at 120 rpm, before mixing equal volumes of each culture to create
an initial inoculum with roughly equal proportions of each strain. To inoculate plants, we added
60 ml of the multi-strain mixture to 6 litres of sterile 0.85% NaCl solution and applied 10^7 cells
per pot. This density was chosen to reflect *S. meliloti* populations in field soil [48] and to avoid
triggering quorum-sensing mechanisms [49], which typically occur at higher cell densities. We
measured rhizobial density in the inoculum by plating a dilution series and counting the number
of colonies. We measured the strain frequencies in the inoculum by centrifuging eight 1 ml
aliquots at 18,000 x g for 2 minutes, removing the supernatant, and then freezing the pellet at -
20°C before extracting DNA. Initial communities were sequenced, and frequencies inferred (Fig.
S1) as outlined in the Rhizobial Fitness section below.

Greenhouse experiment to measure multi-strain symbiotic phenotypes: We grew five replicate
pots of each of the 18 *Medicago* mutants plus wildtype A17 (Fig. 1). Before planting, seeds were
scarified by gently abrading with sandpaper, surface sterilized by soaking in a 10% bleach
solution for 90 seconds and rinsed in de-ionized water. Seeds were stratified on wet filter paper
at 4°C in the dark for 2 days, then planted in pots containing steam-sterilized Sunshine Mix #5.
Germinated seedlings were thinned to three plants per pot. All pots were inoculated 48 hours
after seed planting with the initial 86-strain inoculum. Following inoculation, pots were
randomized into five spatial blocks with one representative of each genotype per block. Plants
were watered as needed and fertilized weekly using increasing concentrations of N-free Fårhaeus
medium [50]. Uninoculated plants were included to monitor cross-contamination. Pots were
harvested and traits measured 8 weeks post-inoculation. We measured nodule number per plant,
average area per nodule and total nodulated area per plant as measures of host investment, total
biomass per plant and root-to-shoot ratio as measures of host benefit, and population size of
rhizobia released per plant and per nodule, nodule strain diversity and relative strain fitness in
nodule pools as measures of rhizobial fitness. After 8 weeks, we harvested inoculated plants and
pooled nodules from ~ 3 plants per pot to ensure enough nodules (>100 /pot) to limit stochastic
variation and saturate diversity. None of the uninoculated plants formed nodules. We
photographed nodule pools per replicate to assess nodule number and size using ImageJ. Plant
biomass was measured by weighing the plant roots and shoots after oven-drying at 75 °C for 3
days. Because not all plants survived in every pot, the number of plants occasionally varied. To
control for this variation, all host traits were normalized per plant for analysis.

Rhizobial fitness: We surface-sterilized the nodules from each pot in 10% bleach for 1 minute,
rinsed the nodules in sterile water, and homogenized them for 90 seconds in 4 ml sterile 0.85%
sodium chloride solution using an Omni TH tissue homogenizer. We estimated the number of
rhizobia released from the nodule pools by dilution plating a 500-1000 ul aliquot of the
homogenate and counting the number of colonies formed. We then estimated the ***absolute fitness***
of the rhizobia on both a per-plant and per-nodule level by dividing the total number of colonies
by the number of plants or nodules harvested from each pot and plotting the $\log_{10}$ transformation
of this value. The remaining nodule homogenate was spun at 400 x g for 10 min at 10°C. to
pellet the bacteroids and the plant debris. We collected the supernatant enriched for

undifferentiated rhizobial cells, pelleted them at 18000 x g for 5 min, pipetted off the supernatant
before freezing the pellets at -20 °C until DNA extraction. DNA was extracted using a Qiagen
DNeasy® Plant extraction kit following the manufacturer's protocol, except we used 3 μL of
RNase in the first step and 50 μL of nuclease water for the final elution.
197
Samples were sequenced using Illumina short-read technology at the Huck Genomic Core
Facility at The Pennsylvania State University. Indexed libraries were generated using the
Illumina DNA Prep Kit. 12X libraries were pooled and sequenced on a 150 x 150 paired-end
NextSeq High Output run, resulting in an average read depth of ~75. We trimmed the reads using
TrimGalore! (v0.4.1, www.bioinformatics.babraham.ac.uk/projects/trim_galore/) with default
settings, except with minimum read length: 100 bp; quality threshold: 30 (−-quality); and
minimum adapter match: 3 (−stringency), used bwa mem (v0.7.17) with default settings to align
reads to the *S. meliloti* USDA1106 genome (NCBI BioProject: PRJNA388336) and estimated
strain frequencies using HARP[51]. Strain frequencies were used to 1) estimate **nodule strain**
diversity using Shannon's H index (via the diversity function from the vegan package in R) and
2) calculate **relative strain fitness** of each strain in each replicate of a particular host mutant by
taking the $\log_2$ of the ratio of the final strain frequency in the nodules to the initial frequency in
the inoculum (median initial frequency = 0.0092, 5%-95% quantile: 0.006-0.018). The log
transformation puts increases and decreases in proportional representation on the same scale
(e.g., a fourfold decrease and increase will have a value of −2 and 2, respectively). Strains with
frequency estimates of 0 in the selected communities were assigned a fitness value of −8, less
than the lowest measurable reductions in other strains [25].
215
Statistical analyses: To assess the effects of the host mutants and the symbiotic stage on plant
traits and rhizobial fitness, we used linear models and analysis of variance (ANOVA) for the
univariate traits and redundancy analysis (RDA) for the multivariate trait of relative strain
fitness. All analyses were conducted in the R statistical environment (version 4.3.2). For each
univariate trait, we fit two linear models using the `lm ()` function: one that includes host genotype
and block, and a second that includes the symbiotic disruption stage and block as additive
predictors. We compared variation explained by the two models using the adjusted R^2
(`anova()` function) to infer whether models that included mutant identity explained more
variance than the symbiotic stage alone. Because the genotypes are nested within the stage of
symbiosis, we ran two separate linear models to avoid introducing collinearity. We then
compared the variance explained by each model. To ensure this comparison was not biased by
differences in the number of levels (degrees of freedom) in each predictor, we used the adjusted
R^2, which corrects for model complexity and sample size. This approach allows us to assess the
relative explanatory power of broad functional categories (stage) versus specific genetic
backgrounds (genotypes) on rhizobial community structure. We $\log_{10}$-transformed average
nodule area/plant and total nodule area traits to reduce heteroscedasticity of variance. To assess
pairwise differences between A17 and each mutant for each trait, we used Tukey's Honest
Significant Difference (HSD) tests. Comparisons with an adjusted *P*-value < 0.05 were
considered statistically significant. To investigate how host genotype or symbiotic stage altered

the relative fitness of rhizobial strains (a multivariate trait), we performed an RDA (`rda()`
function of the vegan package, v2.6-4) using relative strain fitness as the response matrix and
host genotype or symbiotic stage as the explanatory variable. The significance of each model
was assessed by extracting the adjusted R^2 and via permutation-based ANOVA (1,000
permutations) using the `anova()` function with `by = "terms"`. To test whether the relative strain
fitness within each mutant differed from A17, we ran the model for each mutant-A17 pairwise
contrast. Predictors with $P < 0.05$ indicated significant changes in strain composition.
242
Genome-wide association study on relative strain fitness shifts in mutants vs. wildtype nodules
The relative strain fitness shift of each strain for each host mutant was calculated by subtracting
the relative strain fitness in A17 from the relative strain fitness in the host mutants. We
conducted 15 GWAS (three mutants formed no nodules) to identify rhizobial variants underlying
host mutation-dependent ***strain fitness shifts*** for each nodulating mutant. To be thorough, we
also conducted a GWAS on strain fitness in each host genotype individually, which highlights
variants associated with strain success in host nodules generally, as well as the effect of the
perturbations. For each of these GWAS, we ran a linear mixed model (LMM) using GEMMA
(version 0.98.3). Analyses were run separately for the three replicons (chromosome, pSymA,
pSymB) to account for the different population structures. We used a published database of
30,076 SNPs representing previously defined linkage-disequilibrium (LD) groups within *S.*
meliloti[16]. SNP filtering was performed with bcftools-1.18 [52] and PLINK 1.9 [53] based on minor
allele frequency (0.05) and missingness (0.2). We selected the top five variant groups for each
mutation-driven fitness shift tested based on their P-value ranks, cross-compared in the other 15
GWAS. To control for multiple tests while enabling comparisons across replicons that differed in
variant group size from ~ 3,000 to 12,000, we adopted a Bonferroni-corrected P-value cutoff of
$P < 10^{-5}$ to approximate a nominal $P < 0.05$ across replicons. This choice of a consistent
intermediate threshold aligns with our goal of examining selection landscapes rather than
identifying individual causal variants. This allowed us to categorise a rhizobial candidate as
'pervasive' if it ranked in the top ten within six or more host mutants or 'limited effect' if it
appeared in five or fewer. We focused on ranks rather than P-values for this comparative
analysis, prioritizing variants with the most evidence, regardless of differences in power or
genetic architecture across host genetic perturbations.
266
Functional validation of malonyl CoA synthase gene through competition assays : To validate
the functional role of a candidate gene identified through GWAS, we mutated the malonyl-CoA
synthase gene within the genetically amenable *S. meliloti* Rm1021 background (mutant strain
named RmND1439). We competed the mutant against the corresponding wild type Rm1021,
across four host genotypes (A17, *dnf3*, *dnf7*, and *sunn4*). The mutant and wild-type strains were
cultured individually and mixed in equal proportions to create the initial inoculum. Strain
representation in the inoculum and post-inoculation nodule pools was quantified by performing a
dilution series and counting colony-forming units (CFU) on selective and non-selective plates.
Competitive fitness was expressed as the $\log_2$ fold change in the proportion of RmND1439

recovered from the nodules relative to its proportion in the initial inoculum. Detailed methods for
strain construction and competition assays are provided in the Supplementary Methods.
278
In-silico prediction of the rhizobial candidate gene functions through protein-protein interaction using
AlphaFold-Multimer and Foldseek: To obtain additional information about the potential for the
protein products of the candidate rhizobial genes to interact with host proteins such as the NCR
peptides directly, we performed in silico protein–protein interaction predictions using
AlphaFold-Multimer [54,55]. The rhizobial candidate proteins were putatively annotated using
Foldseek[56]. To infer candidate gene involvement during symbiosis, we also listed the nodule-
enrichment scores of each rhizobial candidate from the Symbimics database
(https://iant.toulouse.inra.fr/symbimics/) database[57]. These scores quantify the likelihood of the
rhizobial candidate gene involvement during root nodule symbiosis, with higher values
indicating stronger nodule-specific gene expression. Detailed procedures for structural
modelling, interaction scoring, and data filtering are provided in the Supplementary Methods.
290
291 Results
292
Effect of symbiotic mutations on rhizobial population sizes
Community inoculation resulted in nodulation across all host mutants except for *dmi1-2*, *dmi2-3,*
and dmi3-1, consistent with the expectation that these genes are required for nodule formation
(Fig. 2A). For the 15 mutants that formed nodules, we measured the number of rhizobia released
from the nodules and tested if each mutant differed significantly from A17 plants. For instance,
ipd3 releases $\sim 2.1 \times 10^5$ rhizobial cells per nodule and $\sim 2.435 \times 10^7$ per plant, a 10-fold decrease
compared to A17 (Fig. 2B, C). For both, rhizobia released per nodule and plant, symbiosis stage
did not explain nearly as much variation in rhizobia population sizes as individual host mutations
(Table S2). Only two mutants significantly reduced the rhizobial population size at both per plant
and nodule levels: *ipd3 (*per plant P=0.0001; per nodule P=0.04) and *nad1 (*per plant $P = 0.02$;
per nodule P=0.03), and no mutations increased population size (Table S3). Because *ipd3* and
nad1 do not produce significantly different numbers of nodules compared to A17 ($P > 0.05$), the
smaller per-plant rhizobial population size is primarily due to reduced reproduction of rhizobia
within nodules. This makes sense given that *nad1* host mutant is known to evoke a defense-like
response in nodules that kills reproductively active bacteria, and *ipd3* host mutant has aborted
infection threads, the part of the nodule in which reproductively competent rhizobia are found
[43,58].
310
Host symbiotic genes shape relative strain success
The total rhizobial population released from individual nodules and plants does not tell us
whether the effects of the mutations are strain specific. We examined strain-specific effects of
the mutants in two ways. First, we evaluated the *nodule strain diversity,* as measured by
Shannon's H, in each of the 15 nodulating mutants compared to A17. This analysis revealed that
dnf7 was the only mutant with significantly reduced *nodule strain diversity* ($P = 0.009$, Fig. 2C,
Table S3). Second, we compared the relative fitness of each strain in the nodules of each host

mutant to A17 (Fig. 2E). Hierarchical clustering revealed that strain relative fitness did not
cluster solely by symbiotic stage and that the *latd* nodule strain fitness was quite different from
all other hosts, including wildtype. The *latd* host mutant results should be interpreted carefully—
latd uniquely had a single estimate for rhizobial fitness traits, because plants formed so few
nodules that we had to combine nodules across all pots to create a large enough sample for
sequencing. A redundancy analysis with PERMANOVAs reinforced this finding. Whereas the
symbiosis stage explained some variation in relative strain fitness, symbiotic genes had strong
individual effects (Fig. S2, Table S4, S5). Both early-acting (*ipd3* and *nad1)* and late-acting (all
dnf mutants except *dnf4,* which was marginal at $P = 0.06$, and *dnf1*) significantly altered
rhizobial relative fitness in nodules (Fig. 2E). The *dnf* mutants did not cluster together in the
RDA space, reinforcing that defects in nitrogen fixation did not produce uniform effects on
rhizobial enrichment. Some strains consistently exhibited high or low fitness across all host
genotypes (e.g., MAG322 vs. MAG27), but the fitness of most strains depended on host
symbiotic mutations and often changed the rank order of relative strain fitness of the strains
compared to wildtype (Fig. S3). , Strain shifts were often specific to individual host mutants; for
example, MAG726B, MAG701B, and MAG201 had high fitness in *dnf4* but low fitness in *latd*.
The host-dependent changes in strain fitness allow us to use GWAS to identify rhizobial allelic
variants associated with the observed strain fitness shifts caused by each mutation relative to
wildtype.
337
Rhizobial genes underlying strain fitness shifts in response to host mutations were primarily
located on extrachromosomal replicons
To determine genetic variants associated with shifts in strain fitness when the host genes were
disrupted, we performed 16 GWAS, one for the shifts in strain fitness caused by each of the 15
mutants (relative fitness in mutant-relative fitness in A17) and for comparison, one for relative
fitness in A17 (Fig.3, Fig. S4). We found 9 linkage groups on pSymA and 17 linkage groups on
pSymB that were strongly associated with strain fitness shifts ($P < 10^{-5}$) (Table 2, Table S6). In
contrast, linkage groups on the chromosome had weaker effects (none with $P < 10^{-5}$), except for
LD group 838 ($P = 6.38 \times 10^{-6}$), which was associated with increased strain fitness with *dnf6* and
latd. The *P* values of variants across different replicons may differ in part due to replicon-
specific patterns of relatedness among strains. However, the impact of population structure on
GWAS power is not always straightforward to predict. For instance, the chromosome may
exhibit long-range linkage disequilibrium, resulting in large linkage blocks that tag multiple
genes, which can both help or hinder association resolution. Additionally, we ran a second
GWAS to identify the genes underlying strain fitness in each host (Fig. S5, Table S7). As
expected, within the top five candidates, 11% were shared between A17 and the host mutants
(Table S8), the rest were unique for the mutants, indicating host genetic make-up influences
strain fitness (Table S9). Whereas there was some overlap in genetic variants across the two
GWAS (Table S10), the candidates for shifts in strain fitness in response to host genetic
perturbations were largely distinct (only 20 out of 113 variant groups of strain fitness shifts
overlapped with strain fitness) (Table S11). Below, we focus on candidates from the fitness shift

analyses because we are interested in rhizobial responses to changes in the host environment,
rather than general fitness determinants.
361
GWAS identifies rhizobial genes with pervasive and specific effects on strain fitness shifts.
To study the genomic landscape underlying rhizobial fitness shifts in response to host mutations,
we identified the top five variants for each host mutant (in total, 114 variant groups) based on *P*-
ranks (Table S6). Then we compared the *P*-value ranks and effect sizes of these groups across all
the host mutants (Fig. 4, Fig. S5, Table S12). Most of the candidates on the chromosome were
classified as 'limited effect' variant groups because they ranked among the top ten candidates, in
five or fewer host mutations (Table S12). Exceptions included a massive chromosomal variant
group (901) with 144 variants in 65 genes, among the top ten candidates among nine host
mutants. Six variants with pervasive effects were found on pSymB. On pSymB, four small
variant groups were top-ranked in at least three host mutants (ABC transporter and RkpR, a
dihydrodipicolinate synthase family protein, peptidase T, and a hypothetical protein (Fig. S6).
However, the seven pervasive variants on pSymA were most interesting to us because they
spanned similar hosts and had strong statistical support. These candidates included malonyl-CoA
synthase, SGNH/GDSL hydrolase family protein, *fixL,* an oxygen sensor histidine kinase, a
short-chain oxidoreductase, and a transport protein belonging to the MFS (Major Facilitator
Superfamily). We validated one pervasive candidate, the malonyl-CoA synthase, by knocking it
out in *S. meliloti* Rm1021 genetic background (See Supplemental Methods) and competing the
mutant (RmND1439) against the wild type strain across four host genotypes (*sunn4*, *dnf3*, *dnf7*,
and A17). RmND1439 was almost completely absent from nodules, indicating a functional
version of this gene is crucial for strain fitness in all hosts tested (Fig. S7). This result supports
the gene's pervasive effect and confirms that our approach can identify rhizobial genes
influencing strain performance. We further observed that the non-reference variant often
increased strain fitness within the host mutants. In addition, we noted a trend where the pervasive
variants that conferred higher relative fitness in the mutant hosts, tended to have lower fitness in
wildtype A17. For example, the malonyl-COA synthase had an alternate allele associated with
increased strain fitness (positive beta) in several host mutant backgrounds (*ipd3*, *dnf1*, *dnf2*, *dnf6*,
dnf7, *nad1*, *rdn1*, and *sunn1*). In contrast, the same allele was associated with decreased fitness
(negative beta) in the wild-type A17 (Fig. 4, Table S12). This pattern suggests a host-dependent
allelic effect with the allele conferring a fitness advantage in mutant hosts becoming
disadvantageous in the wild-type host.
392
Most limited-effect variant groups (82 of 94) were small (having fewer than three genes). The
majority of these were on pSymB (Table S13). Those with strong statistical support (*P* value<10[-]
[5]) included variants in or nearby genes coding two ABC transporters (*sunn1, ipd3, dnf1/dnf2,*
rdn1), glutamine synthetase (*latd*) methyltransferases (*nad1*), transcriptional regulators (*rdn1*), a
couple of hypothetical proteins (*ipd3, dnf1/dnf2,dnf5/dnf2,dnf2,sunn1*)and genes involved in
purine metabolism (*dnf3*). A rhizobial glutamine synthetase gene, which converts glutamate and
ammonia into glutamine to assimilate nitrogen, was the strongest candidate for variation in strain
fitness shift in the *latd* genotype, which encodes a nitrate transporter [59]. This was unexpected

because rhizobial glutamine synthetases are generally thought to be strongly downregulated in
nodules where host-driven nitrogen assimilation processes dominate [57]. To further investigate the
potential functional relevance of these candidates, we utilized AlphaFold Multimer, coupled with
Foldseek, to predict possible protein–NCR peptide interactions. Host NCR peptides are a logical
starting point for putative plant protein-rhizobial protein interactions because 1) several of our
hosts are NCR mutants [27], 2) NCRs are known to have strain-specific effects [60] and 3) the large
NCR family is derived from antimicrobial peptides, but we only know what a handful of them
do[61–63]. We found that more than half of the top rhizobial candidate genes associated with strain
fitness shifts (20 out of 35) had a high likelihood of physically interacting with one or more NCR
peptides (Table S13, S14). These included rhizobial candidates associated with fitness shifts in
NCR-compromised mutants whose annotations suggest that rhizobia might adjust their
membrane transport, redox balance, and surface defences when host NCR activity changes (e.g.,
xanthine dehydrogenase family protein subunit M, a Teicoplanin resistance protein-like, and an
ABC transporter ATP-binding protein, which could be an NCR efflux protein according to
Foldseek). These findings provide additional support for the biological relevance of these
candidates in mediating rhizobium–host interactions.
417
Correlations of host investment and benefit traits with rhizobial traits
Because mutualistic partners affect each other's fitness, we were curious if any host investment
or benefit traits were correlated with rhizobial fitness. At the level of individuals pots, nodule area,
plant biomass, and root-to-shoot ratio covaried with nodule strain diversity and position on the
first RDA axis describing shifts in strain relative fitness (Fig. 5A). Total nodulated area per plant
covaried with the second RDA axis describing shifts in strain relative fitness. Perhaps
unsurprisingly, the host trait that covaried with the number of CFUs released per plant was the
number of nodules per plant. Across the first PC of trait space, we found the wild-type plants
grouped with the AON mutants, *hcl*, and three *dnfs* (3,4,6). In contrast, the second PC tended to
separate the early-acting, AON, and wildtype plants from the late-acting mutants. Genotypic trait
correlations confirmed these patterns (Fig. 5B). Host mutants that were larger and had lower
root-to-shoot ratios, (i.e., those that benefited more from rhizobial N-fixation), tended to release
more rhizobia (CFUs) from individual nodules, more rhizobia on a whole-plant level, and had
nodule strain communities that resembled those in A17 nodules (Fig. 5B). Most host traits
correlated strongly with strain diversity and position along the first rhizobial relative fitness
RDA axis; The exception was average nodule area/plant, which was most strongly correlated
with host position on the second RDA axis. Details on mutation-specific effects on host traits can
be found in Fig. S6 and Table S3. Our results suggest that symbiotic gene disruptions strongly
shape both partners' fitness in a manner that reinforces mutualism—mutations that reduce
rhizobial fitness also reduce host fitness.
438
439 Discussion
440

By combining a Select-and-Resequence (S&R) experiment with GWAS, we demonstrate that
mutations in known host symbiotic genes create distinct environments that shape rhizobial strain
success and identify rhizobial candidate genes underlying strain fitness shifts in response to host
mutations. By applying a multi-strain inoculation approach here, we have created a more natural
ecological setting to evaluate host-imposed selection on naturally occurring rhizobial genetic
variants. Within this ecological framework, we found rhizobial genetic variants with both
pervasive and limited effects across host mutants, and that symbiotic stage categories provide
little clarity about how rhizobial genes influence strain fitness. We also found that mutations that
increased rhizobial fitness tend to increase host investment and benefit traits, which highlight a
directional effect where host genetic variations reshape selection on rhizobial partners. Our
findings underscore the complexity of mutualism, where both host and rhizobial genomes jointly
determine the outcomes of symbiosis at the community and individual strain levels, adding to
knowledge of how *Medicago* hosts determine rhizobial fitness [16,30].

Host genetic mutations impact rhizobial absolute fitness
Legumes releasing rhizobia from their senescing nodules directly contribute to the soil rhizobial
population [28,48,64]. We found that most mutations did not affect the number of rhizobia released
from each nodule, and those that did, *ipd3* and *nad1*, released fewer rhizobia per nodule. The
reduced number of rhizobia per nodule with these two mutants is not unexpected, given that
disrupting *nad1* leads to overactive plant defence responses in nodules and disrupting *ipd3* traps
rhizobia within infection threads, likely reducing survival [32,43]. Whereas our results suggest that
the effects of host genetic changes on rhizobial fitness may sometimes align with known gene
functions,, we also observed broader positive correlations between rhizobia released per nodule
and average nodule area across mutations. Increasing the number of rhizobia the host supports
represents one pathway to nurture robust populations of rhizobia, and identifying the host allelic
variation that increases this amplification could set the stage for targeted interventions aimed at
bolstering soil rhizobial populations in agricultural fields [65]. However, the host mutant lines used
in this study were generated through forward genetic screens and, therefore, will likely carry
additional background mutations. Thus, our aim is not to assign strict causality to individual
mutations but rather to examine how alterations in the host genetic landscape broadly influence
rhizobial fitness.

Rhizobial fitness shifts depend on host genetics
A large group of NCR peptides is one of the major known host factors for altering strain survival
and growth in nodules [63,66–68]. Although late-acting NCRs can be important determinants of host
strain specificity in *M. truncatula*, our investigation reveals a more complex landscape [27,69]. Our
analyses of two NCR mutations revealed differing effects on strain fitness within nodules. The
dnf7 mutant, which lacks functional NCR169 [62], was the only mutant background where we
observedreduced nodule strain diversity, and it clustered with *ipd3,* which also supported smaller
rhizobial population. In contrast, *dnf4*, which does not produce NCR211 and influences rhizobial
differentiation during bacteroid formation [70], showed shifts in strain relative fitness similar to
dnf3 and *dnf6,* suggesting these uncharacterized mutants may also involve NCR-related

processes. A few rhizobia genes (*bacA, hrrP, yejABEF*) are known to affect the efficacy of
NCRs [35,66,71]; however, none of those genes were top candidates in our GWAS, suggesting
additional mechanisms of host restriction of rhizobia [61,72,73].

Rhizobial variants with pervasive and limited effects across host mutants
Among the ~120 top rhizobial variants, ~18% affected rhizobial fitness in six or more host
mutant genotypes (pervasive genes), and ~82% were associated with 5 or fewer host
mutants(limited effect genes). , Pervasive effect alleles tended to exhibit fitness trade-offs; alleles
linked to higher fitness in host mutants tended to be linked with lower fitness with wild-type
hosts [16]. Many of the GWAS candidates were identified in only one or a few host mutant
backgrounds, with the strongest associations on pSymB. Unlike the pervasive variants, most
limited-effect variants had little to no effect on strain fitness in the wild-type background. Most
candidate genes with pervasive associations were highly expressed in the nitrogen fixation zone
(Zone III) [57]. For example, the *fixL* gene, which regulates the expression of nitrogen fixation
genes in response to low oxygen levels, malonyl-CoA synthase that is induced in the rhizosphere
of alfalfa, responsible for malonate catabolism within root nodules [77], and a NAD(P)H-binding
protein from the conserved NmrA domain family that shows high similarity to a transcriptional
regulator, involved in nitrogen metabolite repression in fungi [78]. These genes represent strong
candidates for uncovering novel mechanisms that influence symbiont performance across diverse
host contexts[79].

Beyond these candidates, most associations appeared host-specific, indicating that each mutated
host background creates a unique challenge for rhizobia reproduction. Notable candidates
include: A Class I SAM-dependent methyltransferase associated with strain fitness in *nad1*
mutants, suggesting the importance of epigenetic modifications for rhizobial survival under host
defences [32] , and *stc4*, encoding stachydrine N-demethylase reductase, which is linked to
enhanced strain fitness in *dnf4* mutants, possibly by protecting differentiating rhizobia [70].
Whereas these genes are candidates for altering strain fitness in some host mutant contexts and
not others, our experiments do not pinpoint the stage at which each host gene exerts the strongest
selective filter. One way to follow-up on these results is with assays comparing rhizobia mutated
in the candidate genes identified in our GWAS on wildtype and mutant hosts, which would allow
one to identify the selective stage—rhizosphere, infection thread, or nodule—where each gene
exerts its influence on strain fitness.

Considerations for interpreting mixed-strain experiments
Final strain fitness in nodules is a composite trait that reflects host genetics and microbial traits
such as rhizosphere colonization, competition for infection, and proliferation within nodules. As
such, shifts in relative strain fitness across host genotypes could also be influenced by
differential competitive abilities among strains before infection of each host genotype. Thus,
whereas our design allows us to detect which host symbiotic genes alter strain fitness, it does not
allow us to disentangle the relative contributions of microbial competition versus host filtering.
This limitation is especially relevant in complex communities, where strain-strain interactions

influence host effects [21]. Whereas data suggest the effects of microbe × microbe interactions on
strain relative fitness in our system are weak [26], we caution that it is important to interpret our
host-driven fitness shifts as emergent outcomes of host × microbe × microbe interactions. An
additional consideration is that our experiment is designed to measure how growing plants of a
single genotype affects the fitness of rhizobial strains. Linking these fine-scale host effects to
long-term dynamics of rhizobial adaptation is an important overarching goal in plant-microbe
interactions that requires incorporating additional ecological and evolutionary contexts, including
the importance of soil selection [80] or the cooccurrence of host genotypes across time and space
[13,81].
534
Conclusion
Despite legumes being used for soil fertility improvement and actively managed for centuries,
biological nitrogen fixation in agriculture remains inconsistent spatially and temporally [83–86].
This challenge has driven interest and investment in developing microbial inoculants to enhance
crop yield, stress tolerance, and disease and pest management [20]. Rhizobia have been included in
microbe-based bioinoculant formulations for centuries [79]. However, most of those strain
selection decisions are dominated by strain performances observed under single-strain assays
(inoculating a single plant with a single strain) in controlled environments. Thus, strains that
show promise as potential bioinoculants under controlled conditions often fail to reach densities
high enough to influence crop yields in realistic agricultural systems [87]. Formulating ways to
sustain rhizobial fitness in agricultural fields is a critical avenue for plants to continue acquiring
nitrogen from their associated rhizobia [88].
547
Advances in host-mediated microbiome engineering have demonstrated that plant genes can be
leveraged to assemble trait-optimized microbial communities, with the selection of beneficial
microbes being dependent on the host genotype [85,89]. The findings in this study suggest a route
towards expanding the search for host genes that alter rhizobial fitness. If natural allelic variants
of these genes also favor colonization by high-performing symbionts, it will enable the
development of "designer plants" with enriched mutualistic microbiomes [90]. Here, we identify
host genes that drive strain-specific selection and show that even functionally similar mutations
can have divergent effects on strain fitness in multi-strain contexts [21]. Whereas our study focused
on loss-of-function mutations, studies on segregating natural genetic variation in *Medicago*
suggest that specific host genes can differentially enrich rhizobia but find limited evidence for
widespread balancing or positive selection [16]. Though we do not expect to find these loss-of-
function mutations segregating in natural populations, this work lays a critical foundation:
identifying the host genes that influence rhizobial fitness. A next step could be to systematically
test the effect of naturally occurring alleles within these genes on rhizobial fitness, to pinpoint
the variants that promote or reduce fitness and the mechanisms by which they do it.
563
Acknowledgments
We thank Prof. Dong Wang of the University of Massachusetts Amherst for generously
providing the *M. truncatula* mutant seeds. We also appreciate the assistance of Lillian Cherry

and Jennifer E. Harris with the tedious task of picking and processing nodules. Prof. Katy Heath
of the University of Illinois Urbana-Champaign and Dr. Mike Grillo (currently Assistant
Professor at Loyola University, Chicago) for providing strains and sequencing data. This work
was supported by the National Science Foundation grants IOS-1856744 to PT, NY, and LTB,
and IOS-2243819, 1724993, 2243817, and 2243819 to PT and LTB. It was completed using
computing resources provided by the Penn State Institute for Computational and Data Science,
and sequencing was performed by the Huck Genomics Core Facility. LTB's work was supported
by the USDA National Institute of Food and Agriculture and Hatch Appropriations under Project
#PEN04760 and Accession #1025611. The authors acknowledge the Huck Institutes' Genomics
Core Facility (RRID:SCR_023645) for providing library prep and sequencing services.

Authorship Credits
LTB, NY, PT idea conception and funding, LTB experimental design, LTB and RGB plant care;
RB, GF, LTB data collection for main experiment; GL, AG, BG constructed the malonyl-CoA
synthase mutant, SG did the competition assays; VV and SS did the Alphafold Multimer and
Foldseek analysis; SG, LTB, RGB, BE, JS, S&R and GWAS analysis; LTB, SG, RB drafted the
manuscript; SG, LTB, RB, JS, BE, PT, NY, GF, BG, GL,VV, AG and SS for the manuscript
revisions.

Competing Interests:
The authors declare no competing interests.

Data Accessibility
All the raw reads from sequencing the rhizobial strains in the initial community and in the
rhizobial genomic DNA extracted from the nodule pools of the individual host mutants have
been deposited in NCBI under the Bioproject PRJNA401437 (Submission ID: SUB14622715).
All raw data and analyses code have been uploaded to the Github
(https://github.com/BurghardtLab/HostMutants2021_C86meliloti)
595

References

1. Friesen, M. L. *et al.* Microbially mediated plant functional traits. *Annu. Rev. Ecol. Evol. Syst.*

42, 23–46 (2011).

2. Percy, D. M. John N. Thompson, The Geographic Mosaic of Coevolution. *Syst. Biol.* **56**,

149–151 (2007).

3. Bartha, I. *et al.* A genome-to-genome analysis of associations between human genetic

variation, HIV-1 sequence diversity, and viral control. *eLife* **2**, e01123 (2013).

4. Jones, J. D. G. & Dangl, J. L. The plant immune system. *Nature* **444**, 323–329 (2006).

5. Amandine, C. *et al.* Unraveling coevolutionary dynamics using ecological genomics. *Trends*

Genet. TIG **38**, 1003–1012 (2022).

6. Demirjian, C., Vailleau, F., Berthomé, R. & Roux, F. Genome-wide association studies in

plant pathosystems: success or failure? *Trends Plant Sci.* **28**, 471–485 (2023).

7. Corwin, J. A. *et al.* The Quantitative Basis of the Arabidopsis Innate Immune System to

Endemic Pathogens Depends on Pathogen Genetics. *PLOS Genet.* **12**, e1005789 (2016).

8. Newberry, E. A., Minsavage, G. V., Holland, A., Jones, J. B. & Potnis, N. Genome-Wide

Association to Study the Host-Specificity Determinants of *Xanthomonas perforans.*

Phytopathology® **113**, 400–412 (2023).

9. Medina, M. & Sachs, J. L. Symbiont genomics, our new tangled bank. *Genomics* **95**, 129–

137 (2010).

10. Clare, E. L., Schiestl, F. P., Leitch, A. R. & Chittka, L. The promise of genomics in the study

of plant-pollinator interactions. *Genome Biol.* **14**, 207 (2013).

11. *Mutualism*. (Oxford University Press, Oxford, United Kingdom, 2015).

12. Batstone, R. T. Genomes within genomes: nested symbiosis and its implications for plant

evolution. *New Phytol.* **234**, 28–34 (2022).

13. Heath, K. D. & Stinchcombe, J. R. Explaining mutualism variation: a new evolutionary

paradox? *Evolution* **68**, 309–317 (2014).

14. Turkarslan, S. *et al.* Synergistic epistasis enhances the co-operativity of mutualistic

interspecies interactions. *ISME J.* **15**, 2233–2247 (2021).

15. Batstone, R. T. *et al.* Genome-wide association studies across environmental and genetic

contexts reveal complex genetic architecture of symbiotic extended phenotypes. *mBio* **13**,

e01823-22 (2022).

16. Epstein, B. *et al.* Combining GWAS and population genomic analyses to characterize

coevolution in a legume-rhizobia symbiosis. *Mol. Ecol.* **32**, 3798–3811 (2023).

17. Carlson, C. & Frederickson, M. E. An emerging view of coevolution in the legume–

rhizobium mutualism. *Mol. Ecol.* **32**, 3793–3797 (2023).

18. Gohar, D. *et al.* Genomic evidence of symbiotic adaptations in fungus-associated bacteria.

iScience **28**, 112253 (2025).

19. Denison, R. F., Kiers, E. T. & West, S. A. Darwinian Agriculture: When Can Humans Find

Solutions Beyond the Reach of Natural Selection? *Q. Rev. Biol.* **78**, 145–168 (2003).

20. Busby, P. E. *et al.* Research priorities for harnessing plant microbiomes in sustainable

agriculture. *PLOS Biol.* **15**, e2001793 (2017).

21. Burghardt, L. T. & diCenzo, G. C. The evolutionary ecology of rhizobia: multiple facets of

competition before, during, and after symbiosis with legumes. *Curr. Opin. Microbiol.* **72**,

102281 (2023).

22. Oldroyd, G. E. D., Murray, J. D., Poole, P. S. & Downie, J. A. The Rules of Engagement in

the Legume-Rhizobial Symbiosis. *Annu. Rev. Genet.* **45**, 119–144 (2011).

23. Poole, P., Ramachandran, V. & Terpolilli, J. Rhizobia: from saprophytes to endosymbionts.

Nat. Rev. Microbiol. **180**, 1691 (2018).

24. Heath, K. D. & Tiffin, P. Context dependence in the coevolution of plant and rhizobial

mutualists. *Proc. R. Soc. B Biol. Sci.* **274**, 1905–1912 (2007).

25. Burghardt, L. T. *et al.* Select and Resequence reveals relative fitness of bacteria in symbiotic

and free-living environments. *Proc. Natl. Acad. Sci.* **115**, 2425–2430 (2018).

26. Burghardt, L. T., Epstein, B., Hoge, M., Trujillo, D. I. & Tiffin, P. Host-associated rhizobial

fitness: dependence on nitrogen, density, community complexity, and legume genotype.

Appl. Environ. Microbiol. **88**, e00526-22 (2022).

27. Roy, S. *et al.* Celebrating 20 Years of Genetic Discoveries in Legume Nodulation and

Symbiotic Nitrogen Fixation. *Plant Cell* **32**, 15–41 (2020).

28. Burghardt, L. T. Evolving together, evolving apart: measuring the fitness of rhizobial

bacteria in and out of symbiosis with leguminous plants. *New Phytol.* **228**, 28–34 (2020).

29. Ratcliff, W. C., Underbakke, K. & Denison, R. F. Measuring the fitness of symbiotic

rhizobia. *Symbiosis* **55**, 85–90 (2011).

30. Burghardt, L. T., Trujillo, D. I., Epstein, B., Tiffin, P. & Young, N. D. A select and

resequence approach reveals strain-specific effects of *Medicago* nodule-specific PLAT-

domain genes. *Plant Physiol.* **182**, 463–471 (2020).

31. Epstein, B. *et al.* Genome-wide association analyses in the model rhizobium *Ensifer meliloti*.

mSphere **3**, e00386-18 (2018).

32. Wang, C. *et al. NODULES WITH ACTIVATED DEFENSE 1* is required for maintenance of

rhizobial endosymbiosis in *Medicago truncatula*. *New Phytol.* **212**, 176–191 (2016).

33. Yang, S., Tang, F., Gao, M., Krishnan, H. B. & Zhu, H. *R* gene-controlled host specificity in

the legume–rhizobia symbiosis. *Proc. Natl. Acad. Sci.* **107**, 18735–18740 (2010).

34. Tang, F., Yang, S., Liu, J. & Zhu, H. *Rj4* , a Gene Controlling Nodulation Specificity in

Soybeans, Encodes a Thaumatin-Like Protein But Not the One Previously Reported. *Plant*

Physiol. **170**, 26–32 (2016).

35. Price, P. A. *et al.* Rhizobial peptidase HrrP cleaves host-encoded signaling peptides and

mediates symbiotic compatibility. *Proc. Natl. Acad. Sci.* **112**, 15244–15249 (2015).

36. Oldroyd, G. E. D. Speak, friend, and enter: signalling systems that promote beneficial

symbiotic associations in plants. *Nat. Rev. Microbiol.* **11**, 252–263 (2013).

37. Wheatley, R. M. *et al.* Lifestyle adaptations of Rhizobium from rhizosphere to symbiosis.

Proc. Natl. Acad. Sci. U. S. A. **117**, 23823–23834 (2020).

38. Ferguson, B. J. *et al.* Legume nodulation: The host controls the party. *Plant Cell Environ.* **42**,

41–51 (2019).

39. Bright, L. J., Liang, Y., Mitchell, D. M. & Harris, J. M. The *LATD* Gene of *Medicago*

truncatula Is Required for Both Nodule and Root Development. *Mol. Plant-Microbe*

Interactions® **18**, 521–532 (2005).

40. Hogg, B. V., Cullimore, J. V., Ranjeva, R. & Bono, J.-J. The *DMI1* and *DMI2* Early

Symbiotic Genes of *Medicago truncatula* Are Required for a High-Affinity Nodulation

Factor-Binding Site Associated to a Particulate Fraction of Roots. *Plant Physiol.* **140**, 365–

373 (2006).

41. Messinese, E. *et al.* A Novel Nuclear Protein Interacts with the Symbiotic DMI3 Calcium-

and Calmodulin-Dependent Protein Kinase of *Medicago truncatula*. *Mol. Plant-Microbe*

Interactions® **20**, 912–921 (2007).

42. Smit, P. *et al. Medicago* LYK3, an Entry Receptor in Rhizobial Nodulation Factor Signaling.

Plant Physiol. **145**, 183–191 (2007).

43. Ovchinnikova, E. *et al.* IPD3 Controls the Formation of Nitrogen-Fixing Symbiosomes in

Pea and *Medicago* Spp. *Mol. Plant-Microbe Interactions®* **24**, 1333–1344 (2011).

44. Starker, C. G., Parra-Colmenares, A. L., Smith, L., Mitra, R. M. & Long, S. R. Nitrogen

Fixation Mutants of *Medicago truncatula* Fail to Support Plant and Bacterial Symbiotic

Gene Expression. *Plant Physiol.* **140**, 671–680 (2006).

45. Schnabel, E., Journet, E.-P., de Carvalho-Niebel, F., Duc, G. & Frugoli, J. The *Medicago*

truncatula SUNN Gene Encodes a CLV1-like Leucine-rich Repeat Receptor Kinase that

Regulates Nodule Number and Root Length. *Plant Mol. Biol.* **58**, 809–822 (2005).

46. Schnabel, E. L. *et al.* The *ROOT DETERMINED NODULATION1* Gene Regulates Nodule

Number in Roots of *Medicago truncatula* and Defines a Highly Conserved, Uncharacterized

Plant Gene Family. *Plant Physiol.* **157**, 328–340 (2011).

47. Riley, A. B., Grillo, M. A., Epstein, B., Tiffin, P. & Heath, K. D. Discordant population

structure among rhizobium divided genomes and their legume hosts. *Mol. Ecol.* **32**, 2646–

2659 (2023).

48. Denison, R. F. & Kiers, E. T. Life histories of symbiotic rhizobia and mycorrhizal fungi.

Curr. Biol. CB **21**, R775-85 (2011).

49. Darch, S. E., West, S. A., Winzer, K. & Diggle, S. P. Density-dependent fitness benefits in

quorum-sensing bacterial populations. *Proc. Natl. Acad. Sci.* **109**, 8259–8263 (2012).

50. Fåhraeus, G. The Infection of Clover Root Hairs by Nodule Bacteria Studied by a Simple

Glass Slide Technique. *Microbiology* **16**, 374–381 (1957).

51. Kessner, D., Turner, T. L. & Novembre, J. Maximum likelihood estimation of frequencies of

known haplotypes from pooled sequence data. *Mol. Biol. Evol.* **30**, 1145–1158 (2013).

52. Li, H. A statistical framework for SNP calling, mutation discovery, association mapping and

population genetical parameter estimation from sequencing data. *Bioinformatics* **27**, 2987–

2993 (2011).

53. Purcell, S. *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-Based

Linkage Analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

54. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,

583–589 (2021).

55. Varadi, M. *et al.* AlphaFold Protein Structure Database in 2024: providing structure coverage

for over 214 million protein sequences. *Nucleic Acids Res.* **52**, D368–D375 (2024).

56. Van Kempen, M. *et al.* Fast and accurate protein structure search with Foldseek. *Nat.*

Biotechnol. **42**, 243–246 (2024).

57. Roux, B. *et al.* An integrated analysis of plant and bacterial gene expression in symbiotic

root nodules using laser-capture microdissection coupled to RNA sequencing. *Plant J.* **77**,

817–837 (2014).

58. Domonkos, Á. *et al.* NAD1 Controls Defense-Like Responses in Medicago truncatula

Symbiotic Nitrogen Fixing Nodules Following Rhizobial Colonization in a BacA-

Independent Manner. *Genes* **8**, 387 (2017).

59. Harris, J. M. & Dickstein, R. Control of root architecture and nodulation by the *LATD/NIP*

transporter. *Plant Signal. Behav.* **5**, 1365–1369 (2010).

60. Wang, Q. *et al.* Nodule-Specific Cysteine-Rich Peptides Negatively Regulate Nitrogen-

Fixing Symbiosis in a Strain-Specific Manner in *Medicago truncatula*. *Mol. Plant-Microbe*

Interactions® **31**, 240–248 (2018).

61. Guefrachi, I. *et al.* Extreme specificity of NCR gene expression in Medicago truncatula.

BMC Genomics **15**, 712 (2014).

62. Horváth, B. *et al.* Loss of the nodule-specific cysteine rich peptide, NCR169, abolishes

symbiotic nitrogen fixation in the Medicago truncatula dnf7 mutant. *Proc. Natl. Acad. Sci.*

U. S. A. **112**, 15232–15237 (2015).

63. Montiel, J. *et al.* Morphotype of bacteroids in different legumes correlates with the number

and type of symbiotic NCR peptides. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 5041–5046 (2017).

64. Brewin, N. J. Development of the Legume Root Nodule. *Annu. Rev. Cell Biol.* **7**, 191–226

(1991).

65. Gopalakrishnan, S. *et al.* Plant growth promoting rhizobia: challenges and opportunities. *3*

Biotech **5**, 355–377 (2015).

66. Haag, A. F. *et al.* Protection of Sinorhizobium against Host Cysteine-Rich Antimicrobial

Peptides Is Critical for Symbiosis. *PLoS Biol.* **9**, e1001169 (2011).

67. Czernic, P. *et al.* Convergent Evolution of Endosymbiont Differentiation in Dalbergioid and

Inverted Repeat-Lacking Clade Legumes Mediated by Nodule-Specific Cysteine-Rich

Peptides. *Plant Physiol.* **169**, 1254–1265 (2015).

68. Mergaert, P. *et al.* Eukaryotic control on bacterial cell cycle and differentiation in the

Rhizobium –legume symbiosis. *Proc. Natl. Acad. Sci.* **103**, 5230–5235 (2006).

69. Walker, L., Lagunas, B. & Gifford, M. L. Determinants of host range specificity in legume-

rhizobia symbiosis. *Front. Microbiol.* **11**, (2020).

70. Kim, M. *et al.* An antimicrobial peptide essential for bacterial survival in the nitrogen-fixing

symbiosis. *Proc. Natl. Acad. Sci.* **112**, 15238–15243 (2015).

71. Nicoud, Q. *et al.* Sinorhizobium meliloti Functions Required for Resistance to Antimicrobial

NCR Peptides and Bacteroid Differentiation. *mBio* **12**, e00895-21 (2021).

72. Burghardt, L. T. *et al.* Transcriptomic basis of genome by genome variation in a legume-

rhizobia mutualism. *Mol. Ecol.* **26**, 6122–6135 (2017).

73. Lima, R. M., Kylarová, S., Mergaert, P. & Kondorosi, É. Unexplored Arsenals of Legume

Peptides With Potential for Their Applications in Medicine and Agriculture. *Front.*

Microbiol. **11**, 1307 (2020).

74. Savolainen, O., Lascoux, M. & Merilä, J. Ecological genomics of local adaptation. *Nat. Rev.*

Genet. **14**, 807–820 (2013).

75. Hoban, S. *et al.* Finding the Genomic Basis of Local Adaptation: Pitfalls, Practical Solutions,

and Future Directions. *Am. Nat.* **188**, 379–397 (2016).

76. Elena, S. F. Local adaptation of plant viruses: lessons from experimental evolution. *Mol.*

Ecol. **26**, 1711–1719 (2017).

77. Karunakaran, R., East, A. K. & Poole, P. S. Malonate Catabolism Does Not Drive N_2

Fixation in Legume Nodules. *Appl. Environ. Microbiol.* **79**, 4496–4498 (2013).

78. Stammers, D. K. The structure of the negative transcriptional regulator NmrA reveals a

structural superfamily which includes the short-chain dehydrogenase/reductases. *EMBO J.*

20, 6619–6626 (2001).

79. Mus, F. *et al.* Symbiotic Nitrogen Fixation and the Challenges to Its Extension to

Nonlegumes. *Appl. Environ. Microbiol.* **82**, 3698–3710 (2016).

80. Burghardt, L. T., Epstein, B. & Tiffin, P. Legacy of prior host and soil selection on rhizobial

fitness in planta. *Evolution* **73**, 2013–2023 (2019).

81. Wippel, K. *et al.* Host preference and invasiveness of commensal bacteria in the Lotus and

Arabidopsis root microbiota. *Nat. Microbiol.* **6**, 1150–1162 (2021).

82. Sugawara, M. *et al.* Variation in bradyrhizobial NopP effector determines symbiotic

incompatibility with Rj2-soybeans via effector-triggered immunity. *Nat. Commun.* **9**, 3139

(2018).

83. Kaminsky, L. M., Trexler, R. V., Malik, R. J., Hockett, K. L. & Bell, T. H. The inherent

conflicts in developing soil microbial inoculants. *Trends Biotechnol.* **37**, 140–151 (2019).

84. Soumare, A. *et al.* Exploiting biological nitrogen fixation: a route towards a sustainable

agriculture. *Plants* **9**, 1011 (2020).

85. Trivedi, P., Batista, B. D., Bazany, K. E. & Singh, B. K. Plant–microbiome interactions

under a changing world: responses, consequences and perspectives. *New Phytol.* **234**, 1951–

1959 (2022).

86. Batstone, R. T., Ibrahim, A. & MacLean, L. T. Microbiomes: Getting to the root of the

rhizobial competition problem in agriculture. *Curr. Biol.* **33**, R777–R780 (2023).

87. Parnell, J. J. *et al.* From the Lab to the Farm: An Industrial Perspective of Plant Beneficial

Microorganisms. *Front. Plant Sci.* **7**, (2016).

88. Porter, S. S. & Sachs, J. L. Agriculture and the disruption of plant–microbial symbiosis.

Trends Ecol. Evol. **35**, 426–439 (2020).

89. Batstone, R. T., O'Brien, A. M., Harrison, T. L. & Frederickson, M. E. Experimental

evolution makes microbes more cooperative with their local host genotype. *Science* **370**,

476–478 (2020).

90. Batstone, R. T., Burghardt, L. T. & Heath, K. D. Phenotypic and genomic signatures of

interspecies cooperation and conflict in naturally occurring isolates of a model plant

symbiont. *Proc. R. Soc. B Biol. Sci.* **289**, 20220477 (2022).

<u>Figure Legends</u>
Fig. 1: Schematic overview of the experiment used to evaluate rhizobial fitness and host
traits across different host mutant backgrounds. A set of 86 *Sinorhizobium* strains was
cultured and combined into a mixed inoculum. This inoculum was applied to *Medicago*
truncatula wild-type (A17) and host mutants with gene disruptions that act at different stages of
symbiosis: Early, Late, and Autoregulation of Nodulation (AON). The number of host mutants
per stage is indicated in parentheses. Host and rhizobial traits were measured at the nodule and
whole-plant scales. The experimental conditions are clarified under 'Experimental Details'. Each
treatment included five pots per host mutant with three plants per pot, grown in randomized
blocks under nitrogen (N)-free conditions for 8 weeks. Traits measured include indicators of
rhizobial benefit, host benefit, and host investment (Refer to Table 1 for more details). Different
coloured nodules indicate nodules formed by different rhizobial strains.
814
Fig. 2: Host symbiotic mutations influence rhizobial fitness. (A) Host mutations altered
nodule numbers in consistent and predictable ways. (B–C) Host genotype also influenced
rhizobial fitness, as measured by $\log_{10}$-transformed colony-forming units per nodule and per
plant (CFU/nodule) (B) and CFU /plant (C). (D) Nodule strain diversity was largely unaffected
by host mutations. Closed triangles indicate a significant increase or decrease relative to wild-
type A17 (based on Tukey-adjusted *P* values). Boxplots are color-coded by the symbiotic stage
disrupted by each host mutation (see Table S1). (E) Heatmap showing the median relative fitness
of each of 86 *S. meliloti* strains across the different host genotypes. Strains enriched relative to
their initial inoculum appear in shades of red; depleted strains appear in shades of blue. Host
genotypes are hierarchically clustered based on similarity in strain fitness profiles. Strains are
ranked by their relative fitness in A17. Mutants in bold differ significantly from A17 (see Table
S5 for RDA paired with the PERMANOVA results).
827
Fig. 3: GWAS of strain fitness shifts in host mutants. (A) The rhizobial strain fitness shifts for
each of the 86 strains, calculated by subtracting individual relative strain fitness measured in A17
from the relative strain fitness in the host mutants. The heatmap is hierarchically based on the
similarity of median 'shift' values across hosts. Shades of orange denote positive fitness shift
compared to A17, whereas shades of purple denote negative fitness shift compared to A17. (B)
Manhattan plot indicating the SNP (single-nucleotide polymorphism) positions, based on S.
meliloti USDA1106 association with the relative strain fitness shifts across the 15 host mutants
and strain fitness in A17. The dotted line indicates the *P*-value threshold used for identifying
candidate genes ($p < 10^{-5}$), and significant associations are color-coded according to host
mutation and A17. The GWAS was run separately on each replicon to account for different
population structures. The 'Δ' indicates 'strain fitness shift' in that host mutant. For comparison,
we also ran the GWAS on A17 relative strain fitness.
840
Fig. 4: Rhizobial candidate genes underlying strain fitness shifts predominantly occur on
pSymA. Heatmaps based on the inverse *P*-value ranks (red-yellow) and β values, which denote
the effect sizes (red-blue) for the top five variant groups on the pSymA across the 15 host

mutants and A17. Rows represent candidate variant groups, and columns represent host mutants.
The host mutants are colour coded based on the stage of disruption they affect. Progressive dark
shades (low P-value ranks) indicate a stronger association between the variant group and strain
fitness shift in that host mutant. Heatmap based on the effect sizes (β values) for the same
candidate variant groups was ordered to match the p-value ranking. Shades of red indicate host
mutants where the corresponding variant has a positive effect on strain fitness shift, and shades
of blue indicate negative effects as compared to A17. Colour intensity corresponds to effect
magnitude. Alternate allele frequencies (i.e., not present in the reference genome USDA1106)
for each candidate are also shown in parentheses next to the gene names. Pervasiveness of a
variant group is defined as the number of times a variant group is among the top 10 candidates
(based on the P value ranks) within a host mutant. A variant group is classified as "pervasive" if
it is in the top 10 across six or more host genotypes. In contrast, a variant is classified as "limited
effect" if it is in the top 10 candidates across five or fewer host genotypes. We have mentioned
the number of times a variant appears in the top 10 under the "Pervasiveness" column in the
figure and have indicated the pervasive genes in bold. (See Fig. S5 for the other replicons.)
859
860
Fig 5: Covariation and correlation structure among host and rhizobial traits.
(A) Principal component analysis (PCA) of host investment traits, benefit traits, and rhizobial
fitness outcomes, conducted at the level of individual pots. Arrows represent the loadings of each
trait on the first two principal components, which together explain 51.2% and 16.7% of the total
variance, respectively. To avoid loss of representation due to missing data, values for two
genotypes (*latd* and *rdn1*) were averaged across replicates prior to PCA, as these genes had
incomplete trait data that would otherwise be excluded by listwise deletion.
(B) All pairwise combinations of genotypic trait correlations (Pearsons). Each ellipse represents
a pairwise Pearson correlation between traits, with ellipse elongation and color saturation
indicating the strength and direction of the correlation (blue = positive, red = negative). Strong
correlations may reflect shared genetic regulation or pleiotropy.

Tables
873
874   **Table 1:** Definitions of the symbiotic traits examined

Symbiotic Traits	Category	Interpretation
CFU* per nodule	Rhizobial Benefit	Rhizobia released per nodule indicates the average individual strain fitness increase from host association
CFU* per plant	Rhizobial Benefit	Rhizobia released per plant indicates the contribution of each host plant to overall rhizobial population size
Nodule strain diversity	Rhizobial Benefit	Shannon diversity index (H) calculated for strains isolate from nodule indicates host selectivity
Strain relative fitness	Rhizobial Benefit	Enrichment/depletion of specific strains relative to frequency in inoculum: $\log_2$(selected freq./initial freq.)
Nodules per plant	Host Investment	Number of successful host attempts to form associations with rhizobia
Average area per nodule	Host Investment	Host investment in each nodule
Total nodule area per plant	Host Investment	Total plant investment in nodules (i.e. nitrogen fixation)
Total biomass per plant	Host Benefit	Proxy for total nitrogen fixed when plants are grown in nitrogen limited conditions
Root-to-shoot ratio	Host Benefit	Lower values indicate lower investment in roots. In a N limited environment we infer that nitrogen needs were met.

875
876   *CFU= Colony Forming Units

877 **Table 2:** Candidate rhizobial genes in small variant groups (three genes or fewer) associated
878 with ($P < 10^{-5}$) variation in rhizobial relative strain fitness. See Table S6 for information on all
879 variant groups.

LD group [1]	Host mutants	Location	Alt. allele freq [2]	Eff. alt. vars in mutants	Eff alt. vars in WT	Variants	Genes ID's [3]
9034	*dnf1,dnf5/dnf2, dnf2, dnf4, dnf7, ipd3, nad1,rdn1 sunn1, sunn4*	pSymA	0.07	+	-	1	CDO30_RS23065:SGNH/GDSL hydrolase family protein
6978.alt [4]	*dnf1, dnf1/dnf2, dnf2, dnf4, dnf7, dnf5/dnf2, ipd3, rdn1, sunn1*	pSymA	0.07	+	none	1	CDO30_RS23160:NAD(P)H-binding protein
5279	*dnf1, dnf2, dnf7, ipd3, nad1,rdn1, sunn1*	pSymA	0.07	+	-	1	CDO30_RS21460:malonyl-CoA synthase
7969	*dnf1, dnf2, dnf6, nad1, sunn4*	pSymA	0.06	+	none	1	CDO30_RS22325:hypothetical protein
6968	*dnf4, dnf7,ipd3, sunn1, sunn4*	pSymA	0.10	+	none	1	CDO30_RS23125:SDR family oxidoreductase (pseudogene)
10122	*dnf1, dnf1/dnf2, dnf2, sunn1*	pSymB	0.07	+	-	1	CDO30_RS27255:hypothetical protein
12516	*dnf1/dnf2, rdn1, sunn1*	pSymB	0.06	+	none	2	CDO30_RS28530:ABC transporter ATP-binding protein; CDO30_RS28525:*RkpR*
10601	*dnf7, ipd3, ltd*	pSymB	0.05	+	none	1	CDO30_RS28740:*pepT*, peptidase T
3770	*rdn1, sunn4*	pSymA	0.14	+	-	1	CDO30_RS18080:*fixL*, oxygen sensor histidine kinase *fixL*
12013	*dnf1/dnf2, ipd3*	pSymB	0.05	+	none	1	CDO30_RS28675:hypothetical protein
10882	*dnf4*	pSymB	0.05	+	none	1	CDO30_RS30150:glycosyltransferase
9671	*dnf3*	pSymB	0.07	+	none	3	CDO30_RS27435:four-carbon acid sugar kinase family protein
10140	*dnf3*	pSymB	0.08	+	none	1	CDO30_RS27435:four-carbon acid sugar kinase family protein
10913	*dnf3*	pSymB	0.05	+	none	2	CDO30_RS27440:xanthine dehydrogenase family protein molybdopterin-binding subunit; CDO30_RS27445:xanthine dehydrogenase family protein subunit M
10915	*dnf3*	pSymB	0.05	+	none	42	CDO30_RS27440:xanthine dehydrogenase family protein molybdopterin-binding subunit; CDO30_RS27445:xanthine dehydrogenase family protein subunit M
8294	*dnf4*	pSymA	0.07	+	none	1	CDO30_RS19605:*stc4*, stachydrine N-demethylase reductase subunit Stc4
8754	*dnf5/dnf2*	pSymB	0.1	+	none	2	CDO30_RS32445:hypothetical protein
9239	*dnf5/dnf2*	pSymB	0.09	+	none	1	CDO30_RS32445:hypothetical protein
10903	*ipd3*	pSymB	0.05	+	none	1	CDO30_RS27310:ABC transporter ATP-binding protein
838	*latd*	chr	0.09	+	none	1	CDO30_RS12270:glutamine synthetase family protein
4107	*nad1*	pSymA	0.11	-	none	1	CDO30_RS22005:amidase

4386	*nad1*	pSymA	0.09	-	none	2	CDO30_RS22060:Atu4866 domain-containing protein;CDO30_RS22055:class I SAM-dependent methyltransferase;
9865	*rdn1*	pSymB	0.07	+	none	7	CDO30_RS32685:four-carbon acid sugar kinase family protein; CDO30_RS32675:NAD(P)-dependent oxidoreductase; CDO30_RS32705:LacI family transcriptional regulator
9926	*rdn1*	pSymB	0.07	+	none	4	CDO30_RS31555:MASE1 domain-containing protein; CDO30_RS32090:TRAP transporter large permease; CDO30_RS32070:class I SAM-dependent methyltransferase
9941	*rdn1*	pSymB	0.07	+	none	1	CDO30_RS25485:pyridoxamine 5'-phosphate oxidase family protein
10728	*rdn1*	pSymB	0.5	+	none	1	CDO30_RS32910:LysR family transcriptional regulator
12841	*sunn1*	pSymB	0.07	+	none	1	CDO30_RS28530:ABC transporter ATP-binding protein

880 **Notes:** [1]LD group numbers were obtained from [1616] , [2]AF= non-reference (alternate) allele frequency, [3]Locus tags
881 are from the *Sinorhizobium meliloti* USDA1106 reference annotation available on MAGE
882 (http://www.genoscope.cns.fr/agc/microscope/home/index.php), [4]A tri allelic SNP

1. Grow 86 *S.meliloti* isolates and mix for inoculum

Experiment Details

- 5 pots/host mutant with 3 plants/pot
- Randomized block design
- N-free fertilizer
- Harvested at 8 weeks

2. Inoculate *Medicago* hosts mutants

Wildtype	Host mutants spanning stage of symbiosis

A17	Early (6)	Late (9)	AON (3)

3. Measure host and rhizobial traits

Rhizobial Benefits	Host Benefits	Host Investments
1. CFU/nodule	5. Total biomass/plant	7. Nodules/plant
2. CFU/plant	6. Root-to-shoot ratio	8. Average area/nodule
3. Nodule isolate diversity		9. Total nodulated area
4. Relative strain fitness		

Per nodule release

Per plant release

Low ⟶ High   Low ⟶ High   Low ⟶ High

Disruption Type:
WT   Early   Late   AON

Pvalue Rank
1   2   3 4 5 9 90

Beta

Pervasiveness

12	CDO30_RS23155:MFS transporter (0.07)
14	CDO30_RS23160:NAD(P)H-binding protein (0.071)
9	CDO30_RS23125:SDR family oxidoreductase (pseudogene)(0.082)
13	CDO30_RS23065:SGNH/GDSL hydrolase family protein(0.075)
8	CDO30_RS22325:hypothetical protein (0.07)
12	CDO30_RS21460:malonyl–CoA synthase (0.07)
9	CDO30_RS18080:*fixL*, oxygen sensor histidine kinase (0.14)
3	CDO30_RS22055:class I SAM–dependent methyltransferase (0.093)
2	CDO30_RS22005:amidase (0.116)
3	CDO30_RS21460:malonyl–CoA synthase (0.07)
	CDO30_RS19190:*nodQ*, bifunctional sulfate adenylyltransferase/adenylyl–sulfate kinase (0.075)
4	CDO30_RS19605:*stc4*, stachydrine N–demethylase reductase subunit Stc4(0.072)
1	CDO30_RS22335:site–specific integrase (0.188)
1	CDO30_RS20200:type II toxin–antitoxin system RelE/ParE family toxin(0.214)
1	CDO30_RS21100:DUF2200 domain–containing protein (0.081)
3	CDO30_RS21085:class I SAM–dependent methyltransferase (0.093)
1	CDO30_RS20205:type II toxin–antitoxin system ParD family antitoxin (0.198)
1	CDO30_RS21175:TRAP transporter small permease (0.081)
1	CDO30_RS18965:*traA*, Ti–type conjugative transfer relaxase (0.267)
2	CDO30_RS21490:S8 family serine peptidase (0.07)
1	CDO30_RS21375:PhzF family phenazine biosynthesis protein (0.198)
1	CDO30_RS21370:cold–shock protein (0.093)
2	CDO30_RS19120:*glmS*, glutamine—fructose–6–phosphate transaminase (isomerizing) (0.372)
1	CDO30_RS24215:CopG family transcriptional regulator (pseudogene)(0.105)
1	CDO30_RS19555:*ccoN*, cytochrome–c oxidase2C cbb3–type subunit I (0.106)
1	CDO30_RS24970:type II toxin–antitoxin system RelE/ParE family toxin (0.128)
1	CDO30_RS22190:aldo/keto reductase(0.314)

* indicates the number of perturbations in which the LD group ranked within the top 10 candidates