Title: Mutations in legume symbiosis genes create a complex selective landscape for natural genetic variation in rhizobia during multi-strain inoculations

Sohini Guha[1], Regina B. Bledsoe[1], Jeremy Sutherland[1], Brendan Epstein[2], Gwendolyn M. Fry[1], Nevin D. Young[3], Peter Tiffin[2], and Liana T. Burghardt[1*]

[1]The Pennsylvania State University, Department of Plant Science University Park, Pennsylvania

[2]University of Minnesota, Department of Plant and Microbial Biology, St. Paul, Minnesota

[3]University of Minnesota, Department of Plant Pathology, St. Paul, Minnesota

[*]corresponding author (liana.burghardt@gmail.com)

Author for correspondence: Liana T. Burghardt, Department of Plant Science, 109 Tyson Building, University Park, PA 16802, liana.burghardt@gmail.com,1-(814)-863-6168

Running Title: Rhizobial fitness across Medicago mutants

Abstract:

In the mutualism between leguminous plants and rhizobia bacteria, rhizobia live inside root

nodules, creating the potential for host genes to shape the rhizobial selective environment.

Single-strain screens have identified many host genes influencing symbiosis. However, it's

unknown whether these genes influence which rhizobial strains colonize and thrive inside

nodules during multi-strain inoculations. In this study, we inoculated 18 *Medicago truncatula*

symbiotic mutants (including mutations that alter NCR peptide production, plant defence, and

nodule number regulation) with a mixture of 86 *Sinorhizobium meliloti* strains. In multi-strain

inoculations, most mutations led to reduced host benefits but widely varying effects on host

investment and rhizobial benefit (i.e., strain relative fitness), revealing widespread host gene by

strain fitness interactions. Genome-wide association studies identify genetic variants on rhizobial

replicons pSymA and pSymB as important in mediating strain fitness responses to host

mutations. While most top variants only affected rhizobial fitness when one host gene was

disrupted, we identified ten variant groups with pervasive effects across six or more host

mutations. These variants occurred primarily on pSymA, the symbiotic replicon, and include *fixL*

and a few metabolic genes. In contrast to the limited-effect variants, variants with pervasive

positive effects in mutants tended to adversely affect strain fitness in wild-type hosts. Our results

reveal how host symbiosis genes perturb the selective landscape and symbiotic outcomes for

rhizobia and set the stage for improving rhizobial inoculants and breeding legume hosts better

adapted for multi-strain environments.

Introduction

The symbiosis between legume plants and rhizobia bacteria involves the formation of specialized

plant organs called root nodules, within which rhizobia convert atmospheric nitrogen into plant

assimilable forms and, in return, receive photosynthates for growth (1,2). This mutualism allows

legumes to grow in nitrogen-poor soil, providing a sustainable alternative to chemical fertilizer

application (3–5). The economic importance of this plant-microbe interaction has motivated

intensive functional genetics research (6). Screening of mutant populations of the annual diploid

legume *Medicago truncatula (M. truncatula)* for mutations in binary symbiotic traits like

nodulation (Nod +/-) or nitrogen fixation (Fix+/-) enabled the discovery of dozens of genes

underlying root nodule symbiosis (6,7). These genes were identified in controlled environments

where hosts were inoculated with a single rhizobia (8,9). However, the effects of at least four

host symbiosis genes on functional nitrogen fixation have been shown to depend on the rhizobial

strain (10–12), suggesting that single-strain results might not allow the inference of the effect of

host genes in natural and agricultural contexts, where a single legume can form nodules with

multiple strains or species of rhizobia (13). By examining the effects of over a dozen well-known

symbiosis genes on host and rhizobial fitness in mixed inoculations, we sought to identify

additional host genes with strain-specific effects and clarify the potential for host genes to shape

the rhizobial selective environment.

Affordable whole genome shotgun sequencing has recently enabled the high-throughput

assessment of rhizobial strain frequencies inside the nodules of *M. truncatula* when multiple

strains compete for nodulation(14–16) . Here, we take advantage of this method to evaluate the

effects of 18 well-studied *M. truncatula* mutants (knockouts or knockdowns) on the host

symbiotic and the rhizobial traits when *M. truncatula* is grown with a mixture of 86

Sinorhizobium meliloti (*S. meliloti*) strains. Besides measuring standard symbiotic phenotypes,

we estimated the relative fitness of each of the rhizobial strains in the community within each

host mutant. We also performed genome-wide associations (GWA) on rhizobial strain fitness

shifts in response to host mutations to identify the rhizobia genes potentially responsible for

rhizobia genome by host genome interactions (Burghardt et al., 2018; Burghardt et al., 2020).

This novel approach allowed us to examine how mutations in host symbiotic genes may perturb

the selection landscape for naturally occurring genetic variation in rhizobia.

The formation and development of symbiosis in nodules is complex, with each step regulated by

multiple host and rhizobial genes (6,18,19). To encompass these steps, we chose 18 host

symbiosis mutants (Table S1). We grouped them into three categories based on their impact on

1) bacterial invasion (Early), 2) nodule organogenesis and nitrogen fixation (Late), or 3)

regulation of nodule number (Autoregulation of nodule number; AON). The early-stage genes

(*hcl, ipd3*, *latd, dmi1-2, dmi2-3,* and *dmi3-1*) affect the initial processes such as Nod Factor

recognition and nodule organogenesis (20–24). The late-stage genes (*nad1, dnf1, dnf2, dnf4,*

dnf2/dnf1, dnf5/dnf2, dnf6, dnf7) influence nitrogen fixation (25), while the AON pathway genes

(*sunn1, sunn4, rdn1*) regulate the overall number of nodules a plant forms (26,27). While these

loose categorizations help set baseline expectations for the impact of the host mutations on the

legume and the rhizobia involved, similar phenotypes can be caused by mutations in different

genetic pathways, and each gene may have additional pleiotropic effects or act in additional

contexts of nodule development when faced with more than one strain. Together, these focal

genotypes provide a broad view of the effects of host symbiosis genes on their rhizobial partners

in multi-strain inoculations.

We measured host benefit and host investment traits commonly used to characterize the

functional effect of host symbiosis genes (Table 1). Host benefit proxies include *total biomass*

per plant and the *root-to-shoot ratio*. Since nitrogen (N) is a growth-limiting resource,

comparing mutant dry *total biomass per plant* to wild type in N-limited conditions is a proxy for

the total nitrogen fixation in nodules and, thus, evidence of overall host benefit from the

symbiosis. Mutations within the genes *ipd3, latd, nad1, dnf1, dnf2, dnf4, dnf2/dnf1, dnf5/dnf2,*

dnf6, dnf7 impair nitrogen fixation and prevent the host from benefiting from the symbiosis, as

evident from reduced biomass (20,24,25,28). A less common metric of host benefit from

rhizobia is the *root-to-shoot ratio*. When nitrogen is not limiting, plants shift to invest more in

above than belowground growth, resulting in lower root-to-shoot ratios (29). Thus, in nitrogen-

limited environments, the benefits of nitrogen fixation by rhizobia should be reflected in lower

root-to-shoot ratios and greater total biomass per plant. In contrast, the host investment traits

nodules per plant, *average area per nodule,* and *total nodule area per plant* reflect how many

costly organs/relationships a legume forms with the rhizobia and the host's average investment

into each nodule or overall investment in nodules, respectively. Disrupting *sunn, sunn4,* or *rdn1*

(AON genes) commonly causes shifts in host investment traits (hypernodulation) (26,27). Here,

we ask if these patterns recur during mixed-strain inoculations where hosts can choose partners.

While researchers often measure the effect of host mutations on host benefits and investment in

symbiosis, it is far less common to measure the effect of host mutations on rhizobia (30). The

benefits rhizobia obtain from hosts can be assessed by estimating the *population size of rhizobia*

released at the individual nodule and whole plant level (31). Host mutations that increase plant

defence responses (*nad1* and *dnf2*) will likely reduce the rhizobia released from each nodule,

while population responses at the host plant level could be compensated for by forming more

nodules (32). Host selectivity is another facet of rhizobial fitness that host genes can affect.

Some host genes can increase the number of strains able to form nodules, causing an increase in

nodule strain diversity, while others may reduce diversity. Lastly, host mutations could affect the

relative strain fitness, or which strains can best compete and thrive in host nodules. To our

knowledge, the only assessment of the effect of host mutations on rhizobial benefits during

mixed strain inoculations has been conducted on the *NPD* gene family (Nodule Plat Domain

encoding genes) of *M. truncatula* where knocking out all 5 NPD genes (*Δnpd1-5*) eliminated

nodulation when inoculated with only *S. meliloti* strain SM1021 but inoculating with a multi-

strain community of *S. meliloti* , partially restored functional nodulation despite nodules

supporting fewer rhizobial strains than WT (17). These results suggest the standard approach of

pairing single hosts to single strains and focusing only on plant traits may miss host by strain

interactions.

Here, we expanded the range of mutations from the *npds* to additional symbiotic genes with

wide-ranging phenotypic effects (Early, Late, and AON) and studied how they influence both

host and rhizobial fitness metrics when inoculated with a mixture of 86 *S. meliloti* strains (Figure

1). In addition to measuring univariate host and rhizobial traits, we used whole genome

sequencing and the select and resequence (S&R) technique to measure the changes in relative

rhizobial strain fitness diversity (Burghardt et al. 2018). Finally, using GWA, we identified

natural genetic variation in rhizobial candidates for host genome x strain genome interaction. We

examined the following key questions: 1) How much of the variation in host and rhizobial traits

can be explained by symbiotic mutation categories (Early, Late, AON)? 2) Which host and

rhizobial symbiotic traits are most sensitive to symbiotic gene mutation in multi-strain

inoculations? 3) Do host symbiosis genes have strain-specific effects on rhizobial fitness (e.g.,

genome x genome interactions)? And 4) How does mutating these host symbiosis genes perturb

the selective landscape for naturally-varying rhizobial genes? Overall, in a multi-strain

inoculation, host symbiosis genes have varied effects on rhizobia and host benefits. Further,

while many host mutations alter selection on rhizobia, the genomic basis of rhizobial fitness

exhibits more overlap on the replicon with the symbiosis genes than the other two replicons.

<u>Materials and methods</u>

Greenhouse experiment to measure multi-strain symbiotic phenotypes

We created a mixed inoculum using 86 sequenced *S. meliloti* strains isolated from soils across

the native range of *M. truncatula* (33). Each strain was grown in liquid tryptone yeast media for

three days to reach similar population sizes (Fig S1) and then mixed in equal proportions. This

community was used to a) inoculate host plants, b) measure rhizobial density via dilution plating,

and c) determine strain frequencies via whole genome sequencing. Seeds from *M. truncatula* WT

and 18 host mutants (Table S1) were surface-sterilized, scarified, cold-stratified, and sown into

pots as described in (14) (see Supplemental Methods for details). Plants were inoculated with

10^6 rhizobia per pot, fertilized with nitrogen-free fertilizer, and harvested 8 weeks post-

inoculation to measure host and rhizobial traits (Table 1).

Nodules were photographed to assess plant investment, and their number and size were measured

using ImageJ. Plant biomass was measured following oven drying of the roots and shoots.

Nodules were pooled, sterilized, and homogenized to estimate rhizobia released via colony

counting. Whole genome sequencing of the rhizobial DNA in the initial inoculum and extracted

from the nodules using the Qiagen Plant DNeasy kit was done to map the resulting reads to

reference genome USDA1106 (34), and calculate the strain frequency using HARP (35).

Calculation of strain diversity and relative fitness

Strain frequencies were used to 1) estimate nodule pool diversity using Shannon diversity (via

the "renyi" function in the vegan package in R) and 2) calculate relative fitness by dividing final

strain frequency by initial frequency (median initial frequency = 0.0092, 5%-95% quantile:

0.006-0.018) and calculating the $\log_2$ of the relative fitness to normalize the distribution. Strains

with zero frequency were assigned a relative fitness of -8, a value less than the lowest

measurable fitness, to allow for statistical analysis (36).

Statistical analyses

Permanova ("ANOVA" function) was used to test for the effect of mutant type (trait ~ type +

block) and host genotypes (trait ~ genotype + block) on traits. We compared the variance

explained by mutant type and host genotype (Table S2) and re-ran models individually on WT

and each type or mutant (genotype)(Tables S3, S4). The fraction of variation explained by

mutation type was compared to that explained by genotype (Table 2). T-tests were used to

compare mutant effects to WT. Correlations between traits were assessed using "corrplot" in R

(Fig S2), and community composition shifts were evaluated using redundancy analysis ("RDA"

function, Table S7, Fig S5).

Genome-wide association study

We conducted a genome-wide association study (GWA) to identify rhizobial variants responsible

for host mutation-dependent strain fitness shifts by subtracting mean strain fitness in mutant

nodules from WT nodules. A linear mixed model (LMM) in GEMMA (v0.98.3) was used. We

used a published database of 30,076 SNPs representing previously defined linkage-

disequilibrium (LD) groups within *S. meliloti* (15). SNP filtering was performed with bcftools

(37) and PLINK (38) based on minor allele frequency (0.05) and missingness (0.2). The

proportion of phenotypic variance explained (PVE) by genetic variants was estimated using the

K-matrix. P-values were calculated via likelihood ratio tests, and top candidate genes were

identified using a P-value cutoff of 10^{-5}. We ranked these variants across 14 host mutations to

categorize them as 'pervasive' if they appeared in six or more perturbations or 'limited effect' if in

five or fewer.

Results

Symbiotic mutation type accounted for only a subset of the trait variation explained by

disrupting specific symbiotic genes. To assess the contributions of symbiotic mutation type

(Early, Late, AON) and the eighteen individual genetic mutations to each trait, we ran two linear

models: one with mutation type as a predictor and the other with genotype. Models with

genotype-specific predictors consistently had higher adjusted R-squared values (R_{adj} ranged from

0.22 to 0.88) than those with only symbiotic mutation types (R_{adj} ranged from 0.08 to 0.63)

(Table 2). Although mutation type explained only a fraction of the phenotypic variance (Table

S2), they explained a significant fraction of variance for all traits except *CFU per nodule* (Table

S2). Notably, 61% of the variation in *total nodule area per plant* was explained by mutation

type, with AON pathway mutations promoting more investment in nodules (Table 2, Table S3).

Mutation type also explained 39% of the variation in the *root-to-shoot ratio* (Table 2), with early

and late-stage disruptors increasing this ratio, indicating host nitrogen limitation (Table S3).

Additionally, they accounted for 40% of the variation in *CFU per plant* (Table 2), with early-

stage mutations reducing and AON mutations promoting the trait (Table S3). While some

variation can be explained at the level of mutation types, the increased explanatory power

provided by including specific genotypes suggests that moving beyond simple symbiotic

mutation types is necessary to explain symbiotic trait variation in mixed inoculations. Thus, next,

we focus on examining specific host mutations.

Some symbiotic traits are more sensitive to genetic mutations in mixed inoculations than

others. To provide a more detailed assessment of the effect of each of the mutated genes, we

studied the effects of each mutation on each of the nine focal traits (Fig 1, Fig S4). Note that host

investment and rhizobial benefits could not be assessed for *dmi1-2*, *dmi2-3, and dmi3-1* mutants

because they were non-nodulating even during multi-strain inoculation. Overall, we found that

host benefit traits (*total biomass per plant* and *root-to-shoot ratio*) tended to be more sensitive to

host symbiotic gene mutations than host investment or rhizobial benefit traits (Fig S4). Similarly,

some genotypic mutations (e.g., *ipd3, nad1,* and *sunn4)* significantly influenced almost all host

and rhizobial traits, while others (e.g., *dnf1*) did not influence any of the traits (Fig S4). Below,

we highlight each trait category with a special focus on rhizobial benefit traits.

Host mutations differentially altered host benefit and investment traits. Perhaps unsurprisingly,

most genetic mutations influenced both biomass (Fig 2A, Fig S4) and the root-to-shoot ratio (Fig

2B, Fig S4). All mutations reduced biomass except *dnf1*, *rdn1,* and *sunn1* (Fig 2A). Similarly,

the *root-to-shoot ratio* increased relative to the wild type in 11 symbiotic mutants, indicating

nitrogen limitation (Fig 2B, Fig S4). In contrast, *sunn4* plants had a low *root-to-shoot ratio,*

presumably due to the very low biomass of these plants. Overall, our data suggest that most

symbiotic mutations identified in single-strain contexts are not strongly strain-dependent

regarding host benefit during multi-strain inoculations, with early- and late-stage mutants

reducing host benefits, but the degree of impairment varied among genes. On the other hand,

host investment traits were differentially affected by the mutants. Two out of the three host

investment traits (*total nodule area per plant* and *nodules per plant)* were higher than wild type

in AON mutants, *dnf4* and *dnf6*, but lower than wild type in six early and late stage mutants (Fig

2C, D; Fig S4). A single mutant (*hcl*) increased *average area per nodule* (Fig 2E, Fig S4).

Importantly, if we had only focused on the commonly used metric of *nodules per plant*, we

would have missed the effect of many early- and late-stage mutants on overall investment during

mixed inoculations. These observations highlight the complexity of symbiotic interactions and

underscore the importance of considering multiple host traits to understand the impacts of host

genetic mutations on plant performance.

The effect of genetic mutations on rhizobial benefits also depended on the trait. Three mutations

(*ipd3, latd,* and *nad1)* reduced both *CFU per plant* and *per nodule* (Fig 3A, B; Fig S4). In

contrast, the *CFU per plant* increased in four hosts (*dnf4*, *dnf6, sunn1,* and *sunn4*). Interestingly,

none of the mutations increased *CFU per nodule* (Table S4, Fig 3B, Fig S4), but rather, the

higher *CFU per plant* was due to forming more nodules. A quarter of genotypic mutations (*ipd3,*

latd, and *dnf7)* reduced the *nodule strain diversity*, while none increased diversity (Fig 3C, Fig

S4, Table S4). The reduced diversity of strains within *latd* mutants comes with the caveat that

latd formed very few nodules, so the observed reduced diversity could be due to a reduction in

the number of nodules rather than a shift in selectivity *per se*. These results suggest that

increased host investment positively influenced rhizobial benefits and that functioning versions

of the selected symbiosis genes tend to expand the range of strains with which hosts form

associations.

Mutations in early and late symbiosis genes, but not AON genes, lead to significant but subtle

changes in strain relative fitness. Interestingly, the eight host mutants affecting strain relative

fitness were all early- and late-stage mutants; none of the three AON genes (*sunn1, sunn4, rdn1*)

mutants significantly affected strain fitness (Fig S4, Fig S5, Table S5, Table S6). Hierarchical

clustering of the strain relative fitness revealed general trends across host genotypes (Fig 4). WT

grouped most closely with *hcl*, while the other early mutations showed varied effects: *latd* was

the most divergent, and *ipd3* clustered with late-stage disruptors *dnf7*, *dnf2*, *dnf1/dnf2*, *dnf5/dnf2,*

and *nad1*. Notably, *dnf3*, *dnf4*, and *dnf6* formed a distinct group of late mutants with the AON

mutants and *dnf1*. Mutants with similar effects on strain fitness also tended to have comparable

above-ground biomass, indicating that host resource allocation to nodules influences strain

fitness differences. While a few strains showed consistently high or low fitness across all

genotypes (e.g., MAG322 vs. MAG27), fitness variability was common across the host mutants.

Hierarchical clustering revealed ~71% of strains to be sensitive to host genotypic mutations.

Some strains strongly responded to specific genetic mutations, for example- MAG322,

MAG386, MAG5, and MAG221 becoming more common with *latd* mutation, and MAG148B

with *hcl* mutation. While these shifts are interesting, the overall pattern is of host mutations

causing slight fitness shifts across many strains. These subtle shifts give us the power to use

GWA to identify rhizobial allelic variants associated with shifts in response to host mutations.

GWA reveals variants on pSymA and pSymB, which are important for rhizobial fitness when

symbiosis genes are disrupted. We performed sixteen GWA, one for each of the shifts in strain

fitness caused by the fifteen mutants (WT minus mutant fitness) and one for WT fitness alone

(Fig 5A), to identify rhizobial variants associated with variation in fitness. The strongest

candidates for genome x genome associations based on P values were found on pSymA and

pSymB (Fig 5A). We found nine rhizobial linkage groups on pSymA and 17 linkage groups on

pSymB that were strongly associated with strain fitness shifts ($P < 10^{-5}$) (Fig 5B, Table 3). In

contrast, linkage groups on the chromosome had weaker effects (no P-values less than 10^{-5}),

except for LD group 838, which was associated with increased strain fitness within *latd*—a host

background with a limited sample size. (Fig 5B*).* Some of these differences in power between

replicons are likely due to the divergent population structure, but the direction of impact is

difficult to predict (e.g., the chromosome has long-distance linkage among variants, creating

large linkage groups tagging many genes).

Mutating host symbiosis genes has widespread effects on the selective landscape for genetic

variation in rhizobia. We were interested to discover if the genetic variants underlying rhizobial

fitness, when the symbiosis pathway was perturbed, had pervasive or limited effects. In other

words, we sought to summarize the genomic selective landscape that host gene mutation creates

for rhizobia, irrespective of arbitrary P-value cut-offs. For each replicon, we identified the five

most strongly associated candidates and then compared the P ranks (Fig 5B and Fig S6) and

effects sizes (Fig S7) of those variant groups across all the host mutations (Table S8 for a

complete list). This resulted in a list of 114 LD groups across the three replicons. To visualize

patterns, we clustered LD groups based on the inverse of P-value ranks (Fig 5B, Fig S6) and on

the effect sizes (beta values) (Fig S7). A vast majority of candidates on the chromosome and

pSymB affected strain fitness in fewer than three host mutants (Table S8, Fig S8). Exceptions on

the chromosome included a massive linkage group (901) with 144 variants in 65 genes. This LD

group was in the top ten rhizobial candidates when nine different host genes were mutated,

spanning all disruption types. On pSymB, four small LD groups were top-ranked in at least four

host mutants (ABC transporter and RKpR, a dihydrodipicolinate synthase family protein, pepT,

peptidase T, and a hypothetical protein) (Fig 5B). However, the strongest signals of shared

genomic selection were on pSymA (Fig S8), where we identified nine linkage groups with

pervasive effects on strain fitness shifts (Fig S6). The effect of these genetic variants on strain

fitness was not confined to a specific symbiotic stage, although the pervasive variants were not

important in two out of the three early-stage mutants (*hcl, latd*) (Fig S6). The non-reference

variant increased strain fitness within the host mutants, and interestingly, only in pervasive genes

did we see evidence of a fitness trade-off of the allele within WT (Fig S7). The pervasive

candidates included genes like malonyl COA synthase, SGNH/GDSL hydrolase family protein,

fixL which was an oxygen sensor histidine kinase, a short chain oxidoreductase, and a transport

protein belonging to the MFS (Major Facilitator Superfamily). Our results indicate that natural

genetic variation in metabolic genes plays a vital role in altering strain competitiveness within

nodules when symbiotic host genes are disrupted—a role challenging to detect using standard

reverse genetics approaches.

Most candidate variants were only important in one or two host symbiotic mutations,

suggesting that each mutation creates unique environments for rhizobia. Most of the limited-

effect LD groups that were small (less than three genes) and showed substantial statistical

support ($p<10^{-5}$) were on pSymB, and many were found with *nad1*, *dnf3*, *dnf4*, *rdn1*, *dnf5/dnf2*,

or *sunn1* hosts (Table 3). These genes included a couple of ABC transporters, each associated

with individual mutants (*sunn1, ipd3, dnf1/dnf2, rdn1*), glutamine synthetase (*latd*), genes

involved in purine metabolism (*dnf3*), methyltransferases (*nad1*), transcriptional regulators

(*rdn1*), and a couple of hypothetical proteins. It was interesting to note that for *latd,* which is

mutated in a nitrate transporter, the gene that is most associated with the observed strain fitness

shift is glutamine synthetase (39). Interestingly, when *dnf3* was disrupted, a LD group containing

45 SNPs in two genes (xanthine dehydrogenase family protein molybdopterin-binding subunit

and xanthine dehydrogenase family protein subunit M), increased strain fitness. Most of these

rhizobial genes did not significantly alter strain fitness in the WT host background, suggesting

this variation may be conditionally neutral. Overall, our results highlight the complexity of host

genome by strain genome interactions underlying rhizobial selection.

<u>Discussion</u>

This study enhances our understanding of how genetic changes in legume hosts affect the

environmental context and fitness of rhizobia within root nodules and the strain-specific nature

of these responses. While single-strain inoculations allow the direct measurement of phenotypic

outcomes, they limit understanding of relative strain responses and host gene effects. Using

multi-strain inoculations, we screened legume symbiotic mutants to find strain-specific effects of

symbiosis genes on host and rhizobial traits -thus exploring potential mechanisms that could

affect the population size and diversity of mutualistic partners (16,17,40,41). This study

expanded on our previous work, revealing the effects of the *NPD* gene family on rhizobial strain

fitness during multi-strain inoculation (17). We showed that host genetic mutations persistently

affect host benefits, with variable effects on host investment and rhizobial benefits. Finally, by

implementing a GWA, we map the selective landscape host gene mutations create for rhizobia

and identify candidate genes underlying the strain fitness shifts in response to each host

mutation. We find rhizobial genetic variants with both pervasive and limited effects across host

mutants and that symbiotic mutation categories provide little clarity about how rhizobial genes

influence strain fitness. Taken together, we present an overview of the selective landscape that

mutating symbiotic host genes creates for rhizobia, which can be leveraged to fine-tune this

mutualism.

As expected, most but not all of our symbiotic mutations showed reduced nitrogen fixation

and/or nodule investment during multi-strain inoculations similar to those observed in prior

single-strain studies. Because we conducted our experiments with N-free fertilizer, we inferred

nitrogen fixation efficacy in a multi-strain context by placing observed mutant biomass and root-

to-shoot ratios on a continuum from no nitrogen fixation (*dmi1-2*, *dmi2-3, and dmi3-1*) to wild-

type levels (*hcl*). Exceptions included *dnf1* and *dnf4,* which exhibited nitrogen fixation during

multi-strain inoculations despite failing to do so in single-strain experiments suggesting that they

can successfully form symbiosis with some strains (42,43). Partial overriding of single strain

phenotypes, in terms of host investment and host benefits, have been previously reported in

Δ*npd1-5* mutants, which are non-fixing with *Sinorhizobium* strain Sm1021, but regain 30%

nitrogen fixation during multi-strain inoculation, highlighting strain-specificity (17). Another

host mutant that did not exhibit the expected phenotype during multi-strain inoculations was the

hcl mutant, which, during single-strain inoculation with the strain Sm2011, formed significantly

fewer nodules, but during community inoculation, accumulated biomass similar to the WT with

comparable nodulation (23). Host mutation-dependent shifts in rhizobial entry mode have been

previously noted with gain of function *snf1* mutants *of Lotus japonicus* (encoding a T265I

calcium calmodulin-dependent kinase CCaMK *),* which cause a shift from complex infection

thread mode to a simpler 'crack entry' mode of rhizobial entry (44). It would be interesting to

probe how rhizobia navigate the *hcl* mutation—indeed, our GWA reveals that all top variants on

all three replicons are unique to the *hcl* mutation. The host mutations that deviate from their

single-strain phenotypes during multi-strain inoculations create additional opportunities for

discovering novel rhizobia genes that ensure successful symbiosis in these hosts, which are

known to be poor or non-fixing with model *Sinorhizobium* strains.

Legumes releasing rhizobia from their senescing nodules into the soil, directly contribute to the

soil rhizobial population (45). Identifying host genes, like the *npd* family that amplify this effect

can lead to targeted soil microbiome interventions (17). A legume can increase the overall

amount of rhizobia released via two routes. First, they can release more rhizobia by increasing

nodule numbers—a pattern identified for two AON genes (*sunn1*, *sunn4)* and two late-stage

genes *(dnf4* and *dnf6)*—suggesting that these genes, when functional, reduce rhizobia released by

limiting nodule formation. This hypothesis seems plausible particularly for *sunn1(***Su**per

Numeric **N**odules1) and *sunn4 (***Su**per **N**umeric **N**odules4*)* which are known to control the

degree of nodulation, preventing the legume from overinvesting when nitrogen needs are met

(26). Alternatively, plants can have comparable nodulation but release more rhizobia per nodule,

as noted previously in *npd2,* which produced fewer nodules than WT during multi-strain

inoculations but released more rhizobia per nodule than WT (17). In contrast, here we found

most mutations had no effect on the number of rhizobia released from each nodule, and *ipd3,*

latd and *nad1* actually reduced the population size released. Thus, functional copies of these

genes support larger populations of rhizobia per nodule—indeed, disrupting *nad1* leads to

overactive plant defence responses in nodules, likely reducing rhizobial survival (32). Overall,

our results suggest that the effects of host genetic mutations on rhizobial fitness are gene-specific

and may differ from wild-type expectations (e.g., bigger nodules release more rhizobia).

Increasing the number of rhizobia the host supports represents one pathway to nurture robust

populations of beneficial microbes, and identifying the host genes that cause this amplification

sets the stage for targeted interventions aimed at bolstering soil rhizobial populations in

agricultural and conservation contexts (46)

Similar to our previous work on the *npd* genes (17), we also found subtle but significant relative

fitness shifts in specific strains within host mutants. For instance, while late-acting NCRs are

often cited as major determinants of host strain specificity in *M. truncatula*, our investigation

reveals a more complex landscape (6,47). Two NCR mutations had differing effects on strain

fitness within nodules. The *dnf7* mutant, lacking functional NCR169 expressed in the N-fixation

zone (48), grouped with the *ipd3* mutant. Surprisingly, *dnf4*, which does not produce NCR211

and influences rhizobial differentiation during bacteroid formation (43), showed similar effects

on strain fitness as *dnf3* and *dnf6*. There are a few ways, by which rhizobia are known to

overcome the inhibitory actions of the NCRs however none of these genes came up as top

candidates in our GWA suggesting additional mechanisms of host restriction of rhizobia. These

initial observations of NCR mutations further motivate exploring the functional roles and

mechanisms of the remaining ~600 NCRs expressed in nodules (49,50), especially since many

NCRs are differentially expressed in hosts depending on which strain inhabits the nodules and

thus could shape the rhizobial selective environment (51).

More broadly, though, our study shows that many other symbiotic genes beyond NCR's,

including AON genes, can significantly alter which rhizobial strain each plant enriches in mixed

inoculations. Mutations in host genes influencing different stages of symbiosis enriched similar

strains. This highlights the complex interplay between host genetics and microbial partners.

Activation of plant defence responses in nodules can also lead to differential strain fitness (32).

Given that *dnf2* and *nad1* both display elevated defence responses in nodules, all three genotypes

with mutations in *dnf2* and *nad1* enriched similar strains (9). The strain fitness pattern in *dnf1*

hosts was most similar to AON mutants, *dnf3, dnf4*, and *dnf6*. Although there are no known

overlapping roles for these mutants, their similarities may arise from functional nitrogen fixation,

allowing the host to reward some strains over others. Lastly, the genetic basis of some symbiosis

genes (e.g., *dnf3* and *dnf6*) remains a mystery-examination of candidate genes from our GWA

could provide insights (6).

Natural variation in symbiosis genes arises and segregates within legume populations, potentially

contributing to variation in selection on rhizobia (52,53). To measure the effect of this host

genetic variation, we explored how mutations in these genes affect rhizobial strain survival and

fitness, specifically asking if the same rhizobial genes were important for rhizobial fitness across

different host mutations or if unique host mutations create distinct selective pressures on

rhizobia. Our GWAS analyses identified rhizobial genes underlying fitness shifts in different

host mutations, revealing genes with both pervasive and limited effects. Variant groups with

widespread effects were primarily located on pSymA and pSymB. Interestingly, alternative

versions of these alleles increased fitness across many host mutations but strongly reduced

fitness in the wild-type. This suggests a potential cost to harbouring these variants, raising the

question of why they persist. These variations may be beneficial in other host accessions or

species (15). We also found many variants important in only one or a few host mutant

backgrounds, with the strongest associations on pSymB and, to some extent, pSymA. Unlike the

pervasive variants, most limited effect variants had little or no effect on strain fitness in WT

backgrounds, a pattern consistent with conditional neutrality (54–56). Together the data suggests

that the alternate versions of the pervasive genes might be maintained through balancing

selection due to host variation, while the unique genes may increase in frequency through neutral

processes and incidentally enhance fitness when specific symbiotic genes are disrupted.

Most candidate genes with pervasive effects are highly expressed in the nitrogen fixation zone

(Zone III), emphasizing the need for downstream validation assays to elucidate their role in

nitrogen fixation (57). For example, the *fixL* gene, an oxygen sensor, regulates oxygen levels

during nitrogen fixation, Malonyl-CoA synthase, induced selectively in the rhizospheres of

alfalfa, is responsible for malonate catabolism within root nodules (58), and a NAD(P)H-binding

protein from the conserved NmrA domain family shows high similarity to a transcriptional

regulator, involved in nitrogen metabolite repression in fungi (59). These genes represent low-

hanging fruit for uncovering novel mechanisms to broaden the host range of symbionts, aiming

to extend nitrogen fixation beyond legumes (60). Beyond these candidates, most genes only

altered strain fitness in specific symbiotic perturbations, indicating that each perturbation creates

a unique challenge for rhizobia survival and reproduction. Further experimentation will uncover

these novel mechanisms of strain survival and elucidate host by rhizobia genome by genome

interactions. Notable candidates include: A Class I SAM-dependent methyltransferase associated

with strain fitness in *nad1* mutants, suggesting the importance of epigenetic modifications for

rhizobial survival under host defences (32) and variants in *stc4*, encoding stachydrine N-

demethylase reductase, enhancing strain fitness in *dnf4* mutants, possibly by protecting

differentiating rhizobia (43). Our results highlight the potential of leveraging natural genetic

variation to explore undiscovered mechanisms in genome-by-genome interactions between hosts

and microbes.

Conclusion:

By identifying how genetic mutations affect symbiosis in multi-strain inoculations, we aimed to

understand the forces shaping rhizobial and legume evolution in ecologically realistic contexts

(13, 30). Measuring the effects of host genetic variation on microbial fitness is challenging but

crucial for understanding the evolution of and maximizing the benefits of root nodule symbiosis.

By re-examining root nodule symbiosis in multi-strain contexts and using a GWA approach for

leveraging natural genetic variation, we identified novel rhizobial genetic factors influencing

strain fitness shifts, laying the groundwork for further investigations of legume genome by

rhizobia genome interactions (30,36). Validating these rhizobia candidates in different contexts

will help fine-tune existing mutualism and potentially engineer new associations between

rhizobia and novel hosts.

Acknowledgments

We thank Prof. Dong Wang of University of Massachusetts, Amherst for generously providing

the *M. truncatula* mutant seeds, Lillian Cherry, Maria Alejandra Gil-Polo, and Jennifer E. Harris

for help with the tedious task of picking and processing nodules. Prof. Katy Heath of University)

of Illinois Urbana- Champaign and Dr. Mike Grillo (currently Assistant Professor in Loyola

University, Chicago) for providing strains and sequencing data. This work was supported by the

National Science Foundation grant IOS-1856744 to PT, NY, and LTB and IOS-2243819 to PT

and LTB and was completed using computing resources provided by Penn State Institute for

Computational and Data Science and sequencing by the Huck Genomics Core Facility. LTB's

work was also supported by HATCH grant no. 1025611. The authors acknowledge the Huck

Institutes' Genomics Core Facility (RRID:SCR_023645) for use the Illumina NextSeq 2000.

Authorship Credits

LTB, NY, PT idea conception and funding, LTB experimental design, LTB, and RGB plant

care; LTB, SG, RB drafted the manuscript; RB, GF, LTB data collection, SG, LTB, RGB, BE,

JS analysis, SG, LTB, RB, JS, BE, PT, NY, GF manuscript revisions

Competing Interests:

The authors declare no competing interests.

Data Accessibility

All the raw reads from sequencing the rhizobial strains in the initial community and in the

rhizobial genomic DNA extracted from the nodule pools of the individual host mutants have

been deposited in NCBI under the Bioproject PRJNA401437 (Submission ID: SUB14622715).

All raw data and analyses code will be uploaded to the Github

(https://github.com/BurghardtLab/HostMutants2021_C86meliloti)

References

1. Oldroyd GED, Murray JD, Poole PS, Downie JA. The Rules of Engagement in the Legume-
Rhizobial Symbiosis. Annu Rev Genet. 2011 Dec 15;45(1):119–44.

2. Poole P, Ramachandran V, Terpolilli J. Rhizobia: from saprophytes to endosymbionts. Nat
Rev Microbiol. 2018 Jan;180:1691.

3. Vance CP. Symbiotic Nitrogen Fixation and Phosphorus Acquisition. Plant Nutrition in a
World of Declining Renewable Resources. Plant Physiol. 2001 Oct 1;127(2):390–7.

4. Udvardi M, Below FE, Castellano MJ, Eagle AJ, Giller KE, Ladha JK, et al. A Research
Road Map for Responsible Use of Agricultural Nitrogen. Front Sustain Food Syst [Internet].
2021 May 31 [cited 2024 Aug 9];5. Available from:
https://www.frontiersin.org/journals/sustainable-food-
systems/articles/10.3389/fsufs.2021.660155/full

5. Jhu MY, Oldroyd GED. Dancing to a different tune, can we switch from chemical to
biological nitrogen fixation for sustainable food security? PLOS Biol. 2023 Mar
14;21(3):e3001982.

6. Roy S, Liu W, Nandety RS, Crook A, Mysore KS, Pislariu CI, et al. Celebrating 20 Years of
Genetic Discoveries in Legume Nodulation and Symbiotic Nitrogen Fixation. Plant Cell.
2020 Jan;32(1):15–41.

7. Barker DG, Bianchi S, Blondon F, Dattée Y, Duc G, Essad S, et al. Medicago truncatula, a
model plant for studying the molecular genetics of theRhizobium-legume symbiosis. Plant
Mol Biol Report. 1990 Feb;8(1):40–9.

8. Mitra RM, Shaw SL, Long SR. Six nonnodulating plant mutants defective for Nod factor-
induced transcriptional changes associated with the legume-rhizobia symbiosis. Proc Natl
Acad Sci. 2004 Jul 6;101(27):10217–22.

9. Domonkos A, Horvath B, Marsh JF, Halasz G, Ayaydin F, Oldroyd GE, et al. The
identification of novel loci required for appropriate nodule development in Medicago
truncatula. BMC Plant Biol. 2013 Dec;13(1):157.

10. Wang Q, Liu J, Li H, Yang S, Körmöczi P, Kereszt A, et al. Nodule-Specific Cysteine-Rich
Peptides Negatively Regulate Nitrogen-Fixing Symbiosis in a Strain-Specific Manner in
Medicago truncatula. Mol Plant-Microbe Interactions®. 2018 Feb;31(2):240–8.

11. Pislariu CI, Sinharoy S, Torres-Jerez I, Nakashima J, Blancaflor EB, Udvardi MK. The
Nodule-Specific PLAT Domain Protein NPD1 Is Required for Nitrogen-Fixing Symbiosis.
Plant Physiol. 2019 Jul;180(3):1480–97.

12. Jardinaud MF, Carrere S, Gourion B, Gamas P. Symbiotic Nodule Development and
Efficiency in the Medicago truncatula Mtefd-1 Mutant Is Highly Dependent on
Sinorhizobium Strains. Plant Cell Physiol. 2023 Feb 16;64(1):27–42.

13. Burghardt LT, diCenzo GC. The evolutionary ecology of rhizobia: multiple facets of
competition before, during, and after symbiosis with legumes. Curr Opin Microbiol. 2023
Apr 1;72:102281.

14. Burghardt LT, Epstein B, Guhlin J, Nelson MS, Taylor MR, Young ND, et al. Select and
resequence reveals relative fitness of bacteria in symbiotic and free-living environments.
Proc Natl Acad Sci. 2018 Mar 6;115(10):2425–30.

15. Epstein B, Burghardt LT, Heath KD, Grillo MA, Kostanecki A, Hämälä T, et al. Combining
GWAS and population genomic analyses to characterize coevolution in a legume☐rhizobia
symbiosis. Mol Ecol. 2022 Jul 21;mec.16602.

16. Batstone RT, Burghardt LT, Heath KD. Phenotypic and genomic signatures of interspecies
cooperation and conflict in naturally occurring isolates of a model plant symbiont. Proc R
Soc B Biol Sci. 2022 Jul 13;289(1978):20220477.

17. Burghardt LT, Trujillo DI, Epstein B, Tiffin P, Young ND. A Select and Resequence
Approach Reveals Strain-Specific Effects of *Medicago* Nodule-Specific PLAT-Domain
Genes. Plant Physiol. 2020 Jan;182(1):463–71.

18. Oldroyd GED. Speak, friend, and enter: signalling systems that promote beneficial symbiotic
associations in plants. Nat Rev Microbiol. 2013 Apr;11(4):252–63.

19. Wheatley RM, Ford BL, Li L, Aroney STN, Knights HE, Ledermann R, et al. Lifestyle
adaptations of Rhizobium from rhizosphere to symbiosis. Proc Natl Acad Sci U S A. 2020
Sep 22;117(38):23823–34.

20. Bright LJ, Liang Y, Mitchell DM, Harris JM. The *LATD* Gene of *Medicago truncatula* Is
Required for Both Nodule and Root Development. Mol Plant-Microbe Interactions®. 2005
Jun;18(6):521–32.

21. Hogg BV, Cullimore JV, Ranjeva R, Bono JJ. The *DMI1* and *DMI2* Early Symbiotic Genes
of *Medicago truncatula* Are Required for a High-Affinity Nodulation Factor-Binding Site
Associated to a Particulate Fraction of Roots. Plant Physiol. 2006 Jan 1;140(1):365–73.

22. Messinese E, Mun JH, Yeun LH, Jayaraman D, Rougé P, Barre A, et al. A Novel Nuclear
Protein Interacts With the Symbiotic DMI3 Calcium- and Calmodulin-Dependent Protein
Kinase of *Medicago truncatula*. Mol Plant-Microbe Interactions®. 2007 Aug;20(8):912–21.

23. Smit P, Limpens E, Geurts R, Fedorova E, Dolgikh E, Gough C, et al. Medicago LYK3, an
Entry Receptor in Rhizobial Nodulation Factor Signaling. Plant Physiol. 2007 Sep
6;145(1):183–91.

24. Ovchinnikova E, Journet EP, Chabaud M, Cosson V, Ratet P, Duc G, et al. IPD3 Controls
the Formation of Nitrogen-Fixing Symbiosomes in Pea and *Medicago* Spp. Mol Plant-
Microbe Interactions®. 2011 Nov;24(11):1333–44.

25. Starker CG, Parra-Colmenares AL, Smith L, Mitra RM, Long SR. Nitrogen Fixation Mutants
of *Medicago truncatula* Fail to Support Plant and Bacterial Symbiotic Gene Expression.
Plant Physiol. 2006 Feb 1;140(2):671–80.

26. Schnabel E, Journet EP, de Carvalho-Niebel F, Duc G, Frugoli J. The Medicago truncatula
SUNN Gene Encodes a CLV1-like Leucine-rich Repeat Receptor Kinase that Regulates
Nodule Number and Root Length. Plant Mol Biol. 2005 Aug;58(6):809–22.

27. Schnabel EL, Kassaw TK, Smith LS, Marsh JF, Oldroyd GE, Long SR, et al. The *ROOT*
DETERMINED NODULATION1 Gene Regulates Nodule Number in Roots of *Medicago*
truncatula and Defines a Highly Conserved, Uncharacterized Plant Gene Family. Plant
Physiol. 2011 Sep 3;157(1):328–40.

28. Domonkos Á, Kovács S, Gombár A, Kiss E, Horváth B, Kováts G, et al. NAD1 Controls
Defense-Like Responses in Medicago truncatula Symbiotic Nitrogen Fixing Nodules
Following Rhizobial Colonization in a BacA-Independent Manner. Genes. 2017 Dec
14;8(12):387.

29. Agren GI. Root□: Shoot Ratios, Optimization and Nitrogen Productivity. Ann Bot. 2003 Oct
8;92(6):795–800.

30. Burghardt LT. Evolving together, evolving apart: measuring the fitness of rhizobial bacteria
in and out of symbiosis with leguminous plants. New Phytol. 2020 Oct;228(1):28–34.

31. Denison RF. Legume Sanctions and the Evolution of Symbiotic Cooperation by Rhizobia.
Am Nat. 2000 Dec;156(6):567–76.

32. Wang C, Yu H, Luo L, Duan L, Cai L, He X, et al. *NODULES WITH ACTIVATED*
DEFENSE 1 is required for maintenance of rhizobial endosymbiosis in *Medicago truncatula*.
New Phytol. 2016 Oct;212(1):176–91.

33. Riley AB, Grillo MA, Epstein B, Tiffin P, Heath KD. Discordant population structure among
rhizobium divided genomes and their legume hosts. Mol Ecol. 2023 May;32(10):2646–59.

34. Nelson M, Guhlin J, Epstein B, Tiffin P, Sadowsky MJ. The complete replicons of 16
Ensifer meliloti strains offer insights into intra- and inter-replicon gene transfer, transposon-
associated loci, and repeat elements. Microb Genomics [Internet]. 2018 May 1 [cited 2024
Sep 24];4(5).

35. Kessner D, Turner TL, Novembre J. Maximum Likelihood Estimation of Frequencies of
Known Haplotypes from Pooled Sequence Data. Mol Biol Evol. 2013 May 1;30(5):1145–58.

36. Burghardt LT, Epstein B, Guhlin J, Nelson MS, Taylor MR, Young ND, et al. Select and
resequence reveals relative fitness of bacteria in symbiotic and free-living environments.
Proc Natl Acad Sci. 2018 Mar 6;115(10):2425–30.

37. Li H. A statistical framework for SNP calling, mutation discovery, association mapping and
population genetical parameter estimation from sequencing data. Bioinformatics. 2011 Nov
1;27(21):2987–93.

38. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, et al. PLINK: A
Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. Am J
Hum Genet. 2007 Sep;81(3):559–75.

39. Harris JM, Dickstein R. Control of root architecture and nodulation by the *LATD/NIP*
transporter. Plant Signal Behav. 2010 Nov;5(11):1365–9.

40. Heath KD, Stinchcombe JR. EXPLAINING MUTUALISM VARIATION: A NEW
EVOLUTIONARY PARADOX?: PERSPECTIVE. Evolution. 2014 Feb;68(2):309–17.

41. Carlson C, Frederickson ME. An emerging view of coevolution in the legume–rhizobium
mutualism. Mol Ecol. 2023;32(14):3793–7.

42. Wang D, Griffitts J, Starker C, Fedorova E, Limpens E, Ivanov S, et al. A Nodule-Specific
Protein Secretory Pathway Required for Nitrogen-Fixing Symbiosis. Science. 2010 Feb
26;327(5969):1126–9.

43. Kim M, Chen Y, Xi J, Waters C, Chen R, Wang D. An antimicrobial peptide essential for
bacterial survival in the nitrogen-fixing symbiosis. Proc Natl Acad Sci. 2015 Dec
8;112(49):15238–43.

44. Madsen LH, Tirichine L, Jurkiewicz A, Sullivan JT, Heckmann AB, Bek AS, et al. The
molecular network governing nodule organogenesis and infection in the model legume Lotus
japonicus. Nat Commun. 2010 Apr 12;1(1):10.

45. Denison RF, Kiers ET. Life histories of symbiotic rhizobia and mycorrhizal fungi. Curr Biol
CB. 2011 Sep;21(18):R775-85.

46. Gopalakrishnan S, Sathya A, Vijayabharathi R, Varshney RK, Gowda CLL, Krishnamurthy
L. Plant growth promoting rhizobia: challenges and opportunities. 3 Biotech. 2015
Aug;5(4):355–77.

47. Walker L, Lagunas B, Gifford ML. Determinants of host range specificity in legume-
rhizobia symbiosis. Front Microbiol [Internet]. 2020 [cited 2023 Jan 10];11. Available from:
https://www.frontiersin.org/articles/10.3389/fmicb.2020.585749

48. Horváth B, Domonkos Á, Kereszt A, Szűcs A, Ábrahám E, Ayaydin F, et al. Loss of the
nodule-specific cysteine rich peptide, NCR169, abolishes symbiotic nitrogen fixation in the
Medicago truncatula dnf7 mutant. Proc Natl Acad Sci U S A. 2015 Dec 8;112(49):15232–7.

49. Guefrachi I, Nagymihaly M, Pislariu CI, Van de Velde W, Ratet P, Mars M, et al. Extreme
specificity of NCR gene expression in Medicago truncatula. BMC Genomics. 2014 Aug
25;15(1):712.

50. Lima RM, Kylarová S, Mergaert P, Kondorosi É. Unexplored Arsenals of Legume Peptides
With Potential for Their Applications in Medicine and Agriculture. Front Microbiol. 2020
Jun 18;11:1307.

51. Burghardt LT, Guhlin J, Chun CL, Liu J, Sadowsky MJ, Stupar RM, et al. Transcriptomic
basis of genome by genome variation in a legume☐rhizobia mutualism. Mol Ecol. 2017
Nov;26(21):6122–35.

52. Stanton-Geddes J, Paape T, Epstein B, Briskine R, Yoder J, Mudge J, et al. Candidate Genes
and Genetic Architecture of Symbiotic and Agronomic Traits Revealed by Whole-Genome,
Sequence-Based Association Genetics in Medicago truncatula. Lukens L, editor. PLoS ONE.
2013 May 31;8(5):e65688.

53. Curtin SJ, Tiffin P, Guhlin J, Trujillo DI, Burghardt LT, Atkins P, et al. Validating Genome-
Wide Association Candidates Controlling Quantitative Variation in Nodulation. Plant
Physiol. 2017 Feb;173(2):921–31.

54. Savolainen O, Lascoux M, Merilä J. Ecological genomics of local adaptation. Nat Rev
Genet. 2013 Nov;14(11):807–20.

55. Hoban S, Kelley JL, Lotterhos KE, Antolin MF, Bradburd G, Lowry DB, et al. Finding the
Genomic Basis of Local Adaptation: Pitfalls, Practical Solutions, and Future Directions. Am
Nat. 2016 Oct 1;188(4):379–97.

56. Elena SF. Local adaptation of plant viruses: lessons from experimental evolution. Mol Ecol.
2017;26(7):1711–9.

57. Roux B, Rodde N, Jardinaud M, Timmers T, Sauviac L, Cottret L, et al. An integrated
analysis of plant and bacterial gene expression in symbiotic root nodules using laser⬜capture
microdissection coupled to RNA sequencing. Plant J. 2014 Mar;77(6):817–37.

58. Karunakaran R, East AK, Poole PS. Malonate Catabolism Does Not Drive N$_2$ Fixation in
Legume Nodules. Appl Environ Microbiol. 2013 Jul 15;79(14):4496–8.

59. Stammers DK. The structure of the negative transcriptional regulator NmrA reveals a
structural superfamily which includes the short-chain dehydrogenase/reductases. EMBO J.
2001 Dec 3;20(23):6619–26.

60. Mus F, Crook MB, Garcia K, Garcia Costas A, Geddes BA, Kouri ED, et al. Symbiotic
Nitrogen Fixation and the Challenges to Its Extension to Nonlegumes. Appl Environ
Microbiol. 2016 Jul;82(13):3698–710.

61. Sugawara M, Takahashi S, Umehara Y, Iwano H, Tsurumaru H, Odake H, et al. Variation in
bradyrhizobial NopP effector determines symbiotic incompatibility with Rj2-soybeans via
effector-triggered immunity. Nat Commun. 2018 Aug 7;9(1):3139.

<u>Figure Legends</u>

Fig.1: Experimental design schematic. We grew 86 sequenced *Sinorhizobium meliloti s*trains

in liquid media and combined them in approximately equal proportions to generate a multi-strain

community. We used this community to inoculate the *M.truncatula* wildtype (A17) and 18

mutant hosts. We measured the traits which represent common proxies for host investment and

benefits. In addition, we measured rhizobial benefits including the population of rhizobia

released per nodule and per plant and applied whole genome shotgun sequencing to determine

the change in the relative fitness of the rhizobial strains prior to and after exposure to the host

mutants.

Fig.2: **Host symbiotic mutations reduce host benefits and differentially impact host**

investments even in mixed inoculations. Hosts benefit traits like (A) total biomass per plant

and (B) root-to-shoot ratio were reduced and increased respectively in almost all the host mutants

except *sunn4*. The host investment traits which include (C) total nodule area per plant, (D)

nodules per plant, and (E) average area per nodule were differentially affected by the host

mutants. Up/down triangles denote when a host mutant significantly increases/decreases a trait

with respect to WT. The boxplots are colour coded based on the stage of symbiosis (disruption

type) to which the individual host mutants belong (See Table S3 for details).

Fig.3: Mutating host symbiosis genes affect rhizobial benefits. Host mutations differentially

influenced rhizobial benefit traits like (A) CFU/plant and (B) Cfu/nodule while (C) nodule

isolate diversity (C) was largely unaffected by the majority of host mutants. Up/down triangles

denote when a host mutant significantly decreases/increases a trait with respect to WT

respectively. The boxplots are colour coded based on the stage of symbiosis (Mutation type) to

which the individual host mutants belong (see Table S3 for details).

Fig.4: Heatmap visualizing variation in strain fitness in pools of wildtype and mutant

nodules. The host mutants (left) are hierarchically clustered based on the similarity of median

strain fitness patterns (Ward D method). The strain fitness are ordered (bottom) by strain fitness

in A17 from high relative fitness (red) to low (blue). Red indicates strain enrichment relative to

the strain frequency in WT, white indicates no fitness change, and blue indicates depletion

relative to WT.

Fig.5: Rhizobial candidate genes underlying strain fitness shifts predominantly occur on

pSymA and pSymB. (A) Manhattan plot indicating SNP(single nucleotide polymorphism)

positions in *S. meliloti* USDA1106 with strong statistical association with strain fitness shifts

across the 14 host mutants when compared to wildtype A17 hosts. The dotted line indicates the P

value threshold used for identifying candidates genes and for clarity only significant associations

are color-coded according to the host gene mutation. GWAs were conducted separately for each

replicon to account for divergent population structure, but are merged here for ease of

comparison. (B) Heatmaps based on hierarchical clustering of the inverse P-value ranks of the

top five variant groups in each of the 14 host mutations. The heatmap shows how the P value

ranks of the variant groups change across the spectrum of the host mutations.

712   <u>Tables</u>
713   Table 1: Definitions of the symbiotic traits examined in this study

Symbiotic Traits	Category	Interpretation
Nodules per plant	Host Investment	Indicates number of successful host attempts to form associations with rhizobia
Average area per nodule	Host Investment	Host investment in each nodule
Total nodule area per plant	Host Investment	Total plant investment in nodules (i.e. nitrogen fixation)
Total biomass per plant	Host Benefit	Proxy for total nitrogen fixed when plants are grown in nitrogen limited conditions
Root-to-shoot ratio	Host Benefit	Lower values indicate lower investment in roots. In a N limited environment we infer that nitrogen needs were met.
CFU* per nodule	Rhizobial Benefit	Rhizobia released per nodule indicates the average individual strain fitness increase from host association
CFU* per plant	Rhizobial Benefit	Rhizobia released per plant indicates the contribution of each host plant to overall rhizobial population size
Nodule strain diversity	Rhizobial Benefit	Shannon diversity index (H) calculated for the nodule pool indicates host selectivity and strength of selection
Strain relative fitness	Rhizobial Benefit	Enrichment/depletion of specific strains in nodule pools relative to other strains

714
715   *CFU= Colony Forming Units

716 Table 2: Comparison of the variance in host and rhizobial traits explained by individual
717 symbiosis genes (Genotype) versus the stage of symbiosis (Type) in a multi-strain inoculation.

Trait	Geno[1]	Type[1]	Geno + Block R_{adj}^2	Type + Block R_{adj}^2
Nodule number per plant	66	42	0.59	0.37
Average area per nodule	61	24	0.50	0.24
Total nodule area per plant	85	61	0.85	0.63
Total biomass per plant per plant	80	25	0.78	0.21
Root-to-shoot ratio	89	39	0.88	0.36
CFU per nodule	61	13	0.54	0.14
CFU per plant	84	40	0.84	0.42
Nodule strain diversity	58	23	0.44	0.19
Strain relative fitness	36	13	0.22	0.09

718 **Notes**: All terms explained more variance than expected by chance ($P_{value}<0.001$) except Type for CFU per nodules
719 ($P_{value} = 0.03$). All results are from linear models except Strain relative fitness, which is from a multivariate RDA
720 analysis and permANOVA. [1]Percentage of Variance Explained, [2]R_{adj} = Adjusted R-squared; Block= Replicate
721

722 Table 3: Candidate rhizobial genes in small linkage groups (LD) (three genes or less) associated
723 with ($P < 10^{-5}$) variation in rhizobial relative strain fitness.

LD group[1]	Host mutants	Alt. allele freq[2]	Eff. alt. vars in mutants	Eff alt. vars in WT	Varia nts	Genes ID's[3]
9034	*dnf1,dnf5/dnf2, dnf2, dnf4, dnf7, ipd3, nad1,rdn1 sunn1, sunn4*	0.07	(+)	(-)	1	CDO30_RS23065:SGNH/GDSL hydrolase family protein
6978.a lt[4]	*dnf1, dnf1/dnf2, dnf2, dnf4, dnf7, dnf5/2, ipd3, rdn1, sunn1*	0.07	(+)	none	1	CDO30_RS23160:NAD(P)H-binding protein
5279	*dnf1, dnf2, dnf7, ipd3, nad1,rdn1, sunn1*	0.07	(+)	(-)	1	CDO30_RS21460:malonyl-CoA synthase
7969	*dnf1, dnf2, dnf6, nad1, sunn4*	0.06	(+)	none	1	CDO30_RS22325:hypothetical protein
6968	*dnf4, dnf7,ipd3, sunn1, sunn4*	0.10	(+)	none	1	CDO30_RS23125:SDR family oxidoreductase (pseudogene)
10122	*dnf1, dnf1/dnf2, dnf2, sunn1*	0.07	(+)	(-)	1	CDO30_RS27255:hypothetical protein
12516	*dnf1dnf2, rdn1, sunn1*	0.06	(+)	none	2	CDO30_RS28530:ABC transporter ATP-binding protein; CDO30_RS28525:*RkpR*
10601	*dnf7, ipd3, ltd*	0.05	(+)	none	1	CDO30_RS28740:*pepT*, peptidase T
3770	*rdn1, sunn4*	0.14	(+)	(-)	1	CDO30_RS18080:*fixL*, oxygen sensor histidine kinase *fixL*
12013	*dnf1dnf2, ipd3*	0.05	(+)	none	1	CDO30_RS28675:hypothetical protein
10882	*dnf4*	0.05	(+)	none	1	CDO30_RS30150:glycosyltransferase
9671	*dnf3*	0.07	(+)	none	3	CDO30_RS27435:four-carbon acid sugar kinase family protein
10140	*dnf3*	0.08	(+)	none	1	CDO30_RS27435:four-carbon acid sugar kinase family protein
10913	*dnf3*	0.05	(+)	none	2	CDO30_RS27440:xanthine dehydrogenase family protein molybdopterin-binding subunit; CDO30_RS27445:xanthine dehydrogenase family protein subunit M
10915	*dnf3*	0.05	(+)	none	42	CDO30_RS27440:xanthine dehydrogenase family protein molybdopterin-binding subunit; CDO30_RS27445:xanthine dehydrogenase family protein subunit M
8294	*dnf4*	0.07	(+)	none	1	CDO30_RS19605:*stc4*, stachydrine N-demethylase reductase subunit Stc4
8754	*dnf5/dnf2*	0.1	(+)	none	2	CDO30_RS32445:hypothetical protein
9239	*dnf5/dnf2*	0.09	(+)	none	1	CDO30_RS32445:hypothetical protein
10903	*ipd3*	0.05	(+)	none	1	CDO30_RS27310:ABC transporter ATP-binding protein
838	*latd*	0.09	(+)	none	1	CDO30_RS12270:glutamine synthetase family protein
4107	*nad1*	0.11	(-)	none	1	CDO30_RS22005:amidase
4386	*nad1*	0.09	(-)	none	2	CDO30_RS22060:Atu4866 domain-containing protein;CDO30_RS22055:class I SAM-dependent methyltransferase;
9865	*rdn1*	0.07	(+)	none	7	CDO30_RS32685:four-carbon acid sugar kinase family protein; CDO30_RS32675:NAD(P)-dependent oxidoreductase; CDO30_RS32705:LacI family transcriptional regulator
9926	*rdn1*	0.07	(+)	none	4	CDO30_RS31555:MASE1 domain-containing protein; CDO30_RS32090:TRAP transporter large permease; CDO30_RS32070:class I SAM-dependent methyltransferase
9941	*rdn1*	0.07	(+)	none	1	CDO30_RS25485:pyridoxamine 5'-phosphate oxidase family protein
10728	*rdn1*	0.5	(+)	none	1	CDO30_RS32910:LysR family transcriptional regulator
12841	*sunn1*	0.07	(+)	none	1	CDO30_RS28530:ABC transporter ATP-binding protein

724 **Notes:** [1]LD group numbers were obtained from (15); [2]AF= non-reference (alternate) allele frequency. [3]Locus tags

725 are from the *Sinorhizobium meliloti* USDA1106 reference annotation available on MAGE

726 (http://www.genoscope.cns.fr/agc/microscope/home/index.php). [4]A tri allelic event

1. Grow 86 *S.meliloti* strains individually

2. Combine the cultures to create a mixed inoculum

3. Inoculate *M.truncatula* genotypes & grow in N-limited conditions

4. Measure nine symbiotic traits after eight weeks

WILDTYPE

A17

EARLY

hcl, ipd3, dmi1-2, dmi2-3, dmi3-1, latd

LATE

dnf1, dnf2, dnf3, dnf4, dnf6, dnf7, dnf1/dnf2, dnf2/ dnf5, nad1

AON

sunn1, sunn4, rdn1

Host Investments

Low ⟶ High

Host Benefits

Low ⟶ High

Rhizobial Benefits

Low ⟶ High

Nodule pool A   Nodule pool B

10^4 rhizobia/nodule 10^6 rhizobia/nodule

10^7 rhizobia release 10^9 rhizobia release

- Nodules per plant
- Average area per nodule
- Total nodule area per plant

- Total biomass per plant
- Root -to-shoot ratio

- Rhizobia released per nodule
- Rhizobia released per plant
- Nodule isolate diversity
- Nodule isolate fitness