Title: Mutations in legume symbiosis genes create a complex selective landscape for rhizobial symbionts

Sohini Guha[1], Regina B. Bledsoe[1], Jeremy Sutherland[1], Brendan Epstein[2], Gwendolyn M. Fry[1], Nevin D. Young[3], Peter Tiffin[2], and Liana T. Burghardt[1][*]

[1]The Pennsylvania State University, Department of Plant Science, University Park, Pennsylvania
[2]University of Minnesota, Department of Plant and Microbial Biology, St. Paul, Minnesota
[3]University of Minnesota, Department of Plant Pathology, St. Paul, Minnesota
[*]Corresponding author (liana.burghardt@gmail.com)

Author for correspondence: Liana T. Burghardt, Department of Plant Science, 109 Tyson Building, University Park, PA 16802, liana.burghardt@gmail.com,1-(814)-863-6168

Running Title: Rhizobial fitness across Medicago mutants

Abstract
In the mutualism between leguminous plants and rhizobia bacteria, rhizobia live inside root
nodules, creating the potential for host genes to shape the rhizobial selective environment. Many
host genes that affect symbiosis have been identified; however, the extent to which these genes
affect selection acting on rhizobia is unknown. In this study, we inoculated 18 *Medicago*
truncatula symbiotic mutants (including mutants that alter NCR peptide production, plant
defence, and nodule number regulation) with a mixture of 86 *Sinorhizobium meliloti* strains.
Most mutations led to reduced host benefits, but effects on rhizobial benefit (i.e., relative strain
fitness) varied widely, revealing widespread host gene by strain fitness interactions. Genome-
wide association analyses identified genetic variants on rhizobial replicons pSymA and pSymB
as important in mediating strain fitness responses to host mutations. While most top variants only
affected rhizobial fitness with one host mutant (limited effect variants), nine affected fitness
across six or more host mutations. These pervasive variants occurred primarily on pSymA, the
symbiotic replicon, and include *fixL* and a few metabolic genes. In contrast to the limited effect
variants, variants with pervasive positive effects on strain fitness when host genes were mutated
tended to adversely affect fitness in wild-type hosts. Our results reveal how host symbiosis genes
affect rhizobial fitness and set the stage for improving rhizobial inoculants and breeding legume
hosts better adapted for multi-strain environments.

Introduction
40
Many bacteria live in close association with eukaryotic hosts, and a subset of these bacterial taxa
significantly influence host fitness[1]. If host genetic variation alters selective pressures on
bacteria, that variation can shape bacterial evolution. These genome x genome interactions are an
essential prerequisite for co-evolution, and a potential mechanism for maintaining genetic
variation in bacterial and host populations[2]. Most examples of genome x genome interaction
involve pathogens, including a host immune receptor and microbial targets (e.g., human HLA
and HIV epitopes [3]) or plant R genes [4]. However, examinations of quantitative pathogen
resistance leveraging techniques such as co-GWAS [5,6] suggest complex genetic architectures
underlie these host-pathogen interactions [7,8]. More recently, studies of genome-by-genome
interactions have expanded to mutualisms [9–12]. These studies suggest that genome-by-genome
interactions in mutualisms may be equally, if not more, genetically complex than in parasitism
[13–18]. To what extent and in what ways host genes alter bacterial selection has important
implications for bacterial evolution and deploying beneficial microbes in translational contexts
such as agriculture [19,20]. Here, we take a novel approach to measuring genome x genome
interactions by measuring how over a dozen host mutations alter the selective landscape
experienced by scores of strains of mutualistic bacteria.
57
Legume hosts and rhizobial bacteria are a tractable system for studying how individual host
genes alter the selection of their microbial mutualists. Rhizobia inhabit diverse environments,
including soil, the rhizosphere, and root nodules, where they experience selection pressures that
shape their fitness [21]. In nodules, rhizobia convert atmospheric nitrogen into plant assimilable
forms and receive photosynthates for growth [22,23]. Host genetic variation can influence rhizobial
fitness within root nodules [16,24–26] in ways likely shaped by nodule development and subsequent
nitrogen fixation [27]. Decades of work have resulted in extensive collections of mutant lines in
model legumes, including *Lotus* and *Medicago*, which have been exhaustively used to dissect the
functional genetics of nodule formation and function. Most research on these mutants has
focused on their effect on host benefits (e.g., shoot biomass) and investment (e.g., nodule
number/size). Far less attention has been paid to the effect of host mutations on rhizobia [28].
However, doing so is necessary to clarify the potential for host genes to shape the rhizobial
fitness.
71
Measuring the effect of host genes on rhizobia requires considering three aspects of rhizobial
fitness—absolute fitness, relative fitness, and nodule strain diversity. Since most nodules are
initiated by a single rhizobium, rhizobial population sizes in nodules indicate the rewards an
individual bacterium receives from forming a nodule [28,29]. However, measuring population size
alone does not enable an estimate of fitness difference among strains. Assessing strain-level
relative fitness requires comparing strain frequencies before and after exposure to a selective
environment, such as host nodules. A Select and Resequence (S&R) approach leverages whole-
genome shotgun (WGS) sequencing to enable high-throughput tracking of strain frequencies
[25,30]. This method allows for the direct quantification of selection pressures on rhizobia, revealing
how selective an environment is (nodule strain diversity) and which strains are enriched or
depleted compared to the initial community (relative strain fitness) [25,30]. A key advantage of the
S&R method is that, since all strains are already sequenced, genome-wide association studies
(GWAS) can be used to identify genetic variants associated with rhizobial relative fitness
[16,25,30,31].
86

Work in the legume rhizobium mutualism suggests that host genetic variation can alter these
metrics of rhizobial fitness [16,24,25,30]. For instance, host mutations that increase plant defence
responses (*nad1*) reduce the number of rhizobia released from each nodule, while disrupting
genes that cause hosts to form more nodules (*sunn1, sunn4, rdn1*) can increase rhizobia released
at the plant level [32]. Other host genes like *rj2* and *rj4* in soybean restrict the diversity of strains
that form nodules [33,34]. Disruption of the *Sinorhizobium* gene encoding HRRP peptidase,
potentially involved in processing host NCR peptides, can restrict the rhizobial strains in nodules
by modulating host discrimination or defence thresholds [35]. Lastly, disruptions within a host
gene can dramatically alter *relative strain fitness.* For instance, members of the NPD gene family
in *Medicago* differentially influenced which strain frequencies increased in nodules relative to
the inoculated strain community [30]. Taken together, these studies highlight the potential for
single host genes to reshape the selective environment and alter the evolution of rhizobia.
99

Here, we characterize how genetically disrupting host symbiosis genes alter selection on their
nitrogen-fixing rhizobial symbionts *in planta*. To accomplish this goal, we chose eighteen
existing *Medicago truncatula* mutant lines disrupted in symbiotic genes. These genes affect a
wide range of host phenotypes and encompass the developmental trajectory of nodulation. We
inoculated these mutants with a mixture of 86 *Sinorhizobium meliloti* strains (Fig. 1). We studied
standard host investment and benefit traits, as well as multiple aspects of rhizobial fitness (Table
1). We then used a Select and Resequence approach to measure the frequencies of rhizobial
strains (relative strain fitness) in nodules [25] and followed up with GWAS, to identify variants in
rhizobial genes that altered strain fitness when host genes were perturbed. We examined the
following questions: 1) Which host symbiosis genes alter rhizobial fitness? 2) Do host symbiosis
genes have strain-specific effects on rhizobial fitness (i.e., is there evidence of genome x genome
interactions)? 3) If yes, which rhizobial genes harbour genetic variation underlying the observed
shifts in strain fitness, and are any of those variants selected in multiple host mutations? And 4)
Are any host investment or benefit traits correlated with rhizobial fitness traits? While few host
mutations altered rhizobial population sizes or nodule strain diversity, many host mutations did
shift relative strain fitness. Most rhizobial genes associated with strain fitness shifts were found
to be important within specific mutants, although eighteen variants, primarily on pSymA were
important with six or more mutations, suggesting candidate genes that may play fundamental
roles in in-plant rhizobial fitness.
119

120   <u>Materials and methods</u>
121

Host symbiotic mutants: The development of symbiosis in nodules is complex, with each step
regulated by multiple hosts and rhizobial genes [27,36,37]. To encompass these steps, we chose 18
host symbiosis mutants (Table S1), which we grouped into three categories: early-acting, late-
acting, and regulation of nodule number (Autoregulation of nodule number; AON) [38]. The early-
acting genes (*hcl, ipd3, latd, dmi1-2, dmi2-3,* and *dmi3-1*) affect the initial processes, such as
Nod Factor recognition and nodule organogenesis [39–43]. The late-acting genes (*nad1, dnf1, dnf2,*
dnf4, dnf2/dnf1, dnf5/dnf2, dnf6, dnf7) influence nitrogen fixation [32,44], while the regulation of
nodule number genes are in the AON (Autoregulation Of Nodulation) pathway (*sunn1, sunn4,*
rdn1) [45,46]. While these categorizations help set expectations for the impact of the host mutations,
each gene may have additional pleiotropic effects. Together, these mutants provide a broad view
of the effects of host symbiosis genes on their rhizobial partners in multi-strain inoculations.

Seeds for all host mutant lines were obtained from Dr. Dong Wang, Department of Biochemistry
and Molecular Biology, University of Massachusetts, Amherst.
135
Initial inoculum preparation: We created a mixed inoculum using 86 sequenced *Sinorhizobium*
meliloti (henceforth *Sinorhizobium*) strains, which were part of a collection isolated from soils
from 21 locations in Spain and France, using the nodules of *Medicago* plants as a "trap" for
capturing *Sinorhizobium* isolates [47]. We grew each strain separately for 72 hours in Tryptone-
Yeast extract (TY) media at 28°C, shaking at 120 rpm, before mixing equal volumes of each
culture to create an initial inoculum with roughly equal proportions of each strain. To inoculate
plants, we added 60 ml of the multi-strain mixture to 6 litres of sterile 0.85% NaCl solution and
applied 10^7 cells per pot. This density was chosen to reflect *Sinorhizobium* populations in field
soil [48] and to avoid triggering quorum sensing mechanisms [49], which typically occur at higher
cell densities. We measured rhizobial density in the inoculum by plating a dilution series and
counting the number of colonies. We measured the strain frequencies in the inoculum by
centrifuging eight 1 ml aliquots at 18,000 x g for 2 minutes, removing the supernatant, and then
freezing the pellet at -20°C before extracting DNA. Initial communities were sequenced, and
frequencies inferred (Fig. S1) as outlined in the Rhizobial Fitness section below.
150
Greenhouse experiment to measure multi-strain symbiotic phenotypes: We grew five replicate
pots of each of the 18 *Medicago* mutants plus wildtype A17 (Fig. 1). Before planting, seeds were
scarified by gently abrading with sandpaper, surface sterilized by soaking in a 10% bleach
solution for 90 seconds, and rinsed in de-ionized water. Seeds were stratified on wet filter paper
at 4°C in the dark for 2 days, then planted in pots containing steam-sterilized Sunshine Mix #5.
Germinated seedlings were thinned to three plants per pot. All pots were inoculated 48 hours
after seed planting with the initial 86-strain inoculum. Following inoculation, pots were
randomized into five spatial blocks with one representative of each genotype per block. Plants
were watered as needed and fertilized weekly using increasing concentrations of N-free Fårhaeus
medium [50]. Uninoculated plants were included to monitor cross-contamination. Pots were
harvested and traits measured 8 weeks post-inoculation. We measured nodule number per plant,
average area per nodule and total nodulated area per plant as measures of host investment, total
biomass per plant and root-to-shoot ratio as measures of host benefit, and population size of
rhizobia released per plant and per nodule, nodule strain diversity and relative strain fitness in
nodule pools as measures of rhizobial fitness. After 8 weeks, we harvested inoculated plants and
pooled nodules from ~ 3 plants per pot to ensure enough nodules (>100 /pot) to limit stochastic
variation and saturate diversity. None of the uninoculated plants formed nodules. We
photographed nodule pools per replicate to assess nodule number and size using ImageJ. Plant
biomass was measured by weighing plant roots and shoots after oven drying at 75 °C for 3 days.
Because not all plants survived in every pot, the number of plants occasionally varied. To control
for this variation, all host traits were normalized per plant for analysis.
172
Rhizobial fitness: We surface sterilized the nodules from each pot in 10% bleach for 1 minute,
rinsed the nodules in sterile water, and homogenized them for 90 seconds in 4 ml sterile 0.85%
sodium chloride solution using an Omni TH tissue homogenizer. We estimated the number of
rhizobia released from the nodule pools by dilution plating a 500-1000 ul aliquot of the
homogenate and counting the number of colonies formed. We then estimated ***absolute fitness*** of
the rhizobia on both a per plant and per nodule level by dividing the total number of colonies by
the number of plants or nodules harvested from each pot and plotting the log10 transformation of
it. The remaining nodule homogenate was spun at 400xg for 10 min at 10°C. to pellet the

bacteroids and the plant debris. We collected the supernatant enriched for undifferentiated
rhizobial cells, pelleted them at 18000xg for 5 min, pipetted off the supernatant before freezing
the pellets at -20 °C until DNA extraction. DNA was extracted using a Qiagen DNeasy® Plant
extraction kit following the manufacturer's protocol, except we used 3 μL of RNase in the first
step and 50 μL of nuclease water for the final elution.
186
Samples were sequenced using Illumina short-read technology at the Huck Genomic Core
Facility at The Pennsylvania State University. Indexed libraries were generated using the
Illumina DNA Prep Kit. 12X libraries were pooled and sequenced on a 150 x 150 paired-end
NextSeq High Output run, resulting in an average read depth of ~75. We trimmed the reads using
TrimGalore! (v0.4.1, www.bioinformatics.babraham.ac.uk/projects/trim_galore/) with default
settings, except with minimum read length: 100 bp; quality threshold: 30 (−-quality); and
minimum adapter match: 3 (−stringency), used bwa mem (v0.7.17) with default settings to align
reads to the *Sinorhizobium (Ensifer) meliloti* USDA1106 genome (NCBI BioProject:
PRJNA388336) and estimated strain frequencies using HARP [51]. Strain frequencies were used
to 1) estimate ***nodule strain diversity*** using Shannon's H index (via the diversity function from
the vegan package in R) and 2) calculate ***relative strain fitness*** of each strain in each replicate of
a particular host mutant by taking the $\log_2$ of the ratio of the final strain frequency in the nodules
to the initial frequency in the inoculum (median initial frequency = 0.0092, 5%-95% quantile:
0.006-0.018). The log transformation puts increases and decreases in proportional representation
on the same scale (e.g., a fourfold decrease and increase will have a value of −2 and 2,
respectively). Strains with frequency estimates of 0 in the selected communities were assigned a
fitness value of −8, less than the lowest measurable reductions in other strains [25].
204
Statistical analyses. To assess the effects of the host mutants and the symbiotic stage on plant
traits and rhizobial fitness, we used linear models and analysis of variance (ANOVA) for the
univariate traits and redundancy analysis (RDA) for the multivariate trait of relative strain
fitness. All analyses were run in the R statistical environment (v4.3.2). For each univariate trait,
we fit two linear models (`lm()` function): one using host genotype and block, and a second using
the symbiotic disruption stage and block as additive predictors. We compared variation
explained by the two models using the adjusted R^2 (`anova()` function) to infer whether models
that included mutant identity explained more variance than the symbiotic stage alone. Because
the genotypes are nested within the stage of symbiosis, we ran two separate linear models to
avoid introducing collinearity. We then compared the variance explained by each model. To
ensure this comparison was not biased by differences in the number of levels (degrees of
freedom) in each predictor, we used the adjusted R^2, which corrects for model complexity and
sample size. This approach allows us to assess the relative explanatory power of broad functional
categories (stage) versus specific genetic backgrounds (genotypes) on rhizobial community
structure. We log10-transformed average nodule area/plant and total nodule area traits to reduce
heteroscedasticity of variance. To assess pairwise differences between A17 and each mutant for
each trait, we used Tukey's Honest Significant Difference (HSD) tests. Comparisons with an
adjusted *p*-value < 0.05 were considered statistically significant. To investigate how host
genotype or symbiotic stage altered the relative fitness of rhizobial strains (a multivariate trait),
we performed an RDA (`rda()` function of the vegan package, v2.6-4) using relative strain fitness
as the response matrix and host genotype or symbiotic stage as the explanatory variable. The
significance of each model was assessed by extracting the adjusted R^2 and via permutation-based
ANOVA (1,000 permutations) using the `anova()` function with `by = "terms"`. To test whether
the relative strain fitness within each mutant differed from A17, we ran the model for each

mutant-A17 pairwise contrast. Predictors with $p < 0.05$ indicated significant changes in strain
composition.
231
Genome-wide association study on relative strain fitness shifts in mutants vs. wildtype nodules
The relative strain fitness shift of each strain for each host mutant was calculated by subtracting
the relative strain fitness in A17 from the relative strain fitness in the host mutants. We
conducted 15 GWAS (three mutants formed no nodules) to identify rhizobial variants underlying
host mutation-dependent ***strain fitness shifts*** for each nodulating mutant. We also conducted a
GWAS to identify genes associated with strain fitness in A17 nodules. For each of these 16
GWAS, we ran a linear mixed model (LMM) in GEMMA (v0.98.3). Analyses were run
separately for the three replicons (chromosome, pSymA, pSymB) to account for the different
population structures. We used a published database of 30,076 SNPs representing previously
defined linkage-disequilibrium (LD) groups within *S. meliloti*[16]. SNP filtering was performed
with bcftools-1.18 [52] and PLINK 1.9 [53] based on minor allele frequency (0.05) and missingness
(0.2). We selected the top five variant groups for each mutation-driven fitness shift tested based
on their *p*-value ranks, cross-compared in the other 15 GWAS. To control for multiple tests
while enabling comparisons across replicons that differed in variant group size from ~ 3,000 to
12,000, we adopted a Bonferroni corrected *p*-value cutoff of $p < 10^5$ to approximate a nominal *p*
< 0.05 across replicons. This choice of a consistent intermediate threshold aligns with our goal of
examining selection landscapes rather than identifying individual causal variants. This allowed
us to categorise a rhizobial candidate as 'pervasive' if it ranked in the top ten within six or more
host mutants or 'limited effect' if it appeared in five or fewer. We focused on ranks rather than *p*-
values for this comparative analysis to focus on variants with the most evidence, regardless of
differences in power or genetic architecture across host genetic perturbations.
253
254 Results
255
Effect of symbiotic mutations on rhizobial population sizes
Community inoculation resulted in nodulation across all host mutants except for *dmi1-2*, *dmi2-3*,
and dmi3-1, consistent with the expectation that these genes are required for nodule formation
(Fig. 2A). For the 15 mutants that formed nodules, we measured the number of rhizobia released
from the nodules and tested if each mutant differed significantly from A17 plants. For instance,
ipd3 releases ~ 2.1×10^5 rhizobial cells per nodule and ~2.435×10^7 per plant, a 10-fold decrease
compared to A17 (Fig. 2B, C). For both, rhizobia released per nodule and plant, symbiosis stage
did not explain nearly as much variation in rhizobia population sizes as individual host mutations
(Table S2). Only two mutants significantly reduced the rhizobial population size at both per
plant and nodule levels: *ipd3 (*per plant $p=0.0001$; per nodule $p=0.04$) and *nad1 (*per plant $p =$
0.02; per nodule $p=0.03$), and no mutations increased population size (Table S3). Because *ipd3*
and *nad1* do not produce significantly different numbers of nodules compared to A17 ($p > 0.05$),
the smaller per-plant rhizobial population size is primarily due to reduced reproduction within
nodules. This makes sense given that *nad1* evokes a defense-like response in nodules that kills
reproductively active bacteria, and *ipd3* causes aborted infection threads, the part of the nodule in
which reproductively competent rhizobia are found [43,54].
272
Host symbiotic genes shape relative strain success
The total rhizobial population released from individual nodules and plants does not tell us
whether the effects of the mutations are strain specific. We examined strain-specific effects of
the mutants in two ways. First, we evaluated the *nodule strain diversity,* as measured by

Shannon's H, in each of the 15 nodulating mutants compared to A17. This analysis revealed that
dnf7 was the only mutant significantly reducing *nodule strain diversity* (p = 0.009, Fig. 2C,
Table S2). Second, we compared the relative fitness of each strain in the nodules of each host
mutant to A17 (Fig. 2E). Hierarchical clustering revealed that strain relative fitness did not
cluster solely by symbiotic stage and that the *latd* nodule strain fitness was quite different from
all other hosts, including wildtype. Note that the *latd* host mutant results should be interpreted
carefully—*latd* uniquely had a single estimate for rhizobial fitness traits, because plants formed
so few nodules that we had to combine nodules across all pots to create a large enough sample
for sequencing. A redundancy analysis with PERMANOVAs reinforced this finding. While the
symbiosis stage explained some variation in relative strain fitness, symbiotic genes had strong
individual effects (Fig. S2, Table S4, S5). Both early-acting (*ipd3* and *nad1*) and late-acting (all
dnf mutants except *dnf4,* which was marginal at p = 0.06, and *dnf1*) significantly altered rhizobial
relative fitness in nodules (Fig. 2E). The *dnf* mutants did not cluster together in the RDA space,
reinforcing that defects in nitrogen fixation did not produce uniform effects on rhizobial
enrichment. Some strains consistently exhibited high or low fitness across all host genotypes
(e.g., MAG322 vs. MAG27), but the fitness of most strains depended on host symbiotic
mutations and often changed the rank order of relative strain fitness of the strains compared to
wildtype (Fig. S3). Importantly, strain shifts were often specific to individual host mutants; for
example, MAG726B, MAG701B, and MAG201 had high fitness in *dnf4* but low fitness in *latd*.
The host-dependent changes in strain fitness allow us to use GWAS to identify rhizobial allelic
variants associated with the observed strain fitness shifts caused by each mutation relative to
wildtype.
299
Rhizobial genes underlying strain fitness shifts in response to host mutations were not on the
main chromosome
To determine genetic variants associated with strain fitness when host genes were disrupted, we
performed 16 GWAS, one for the shifts in strain fitness caused by each of the fifteen mutants
(relative fitness in mutant-relative fitness in A17) and one for relative fitness in A17 (Fig.3, Fig.
S4). We found 9 linkage groups on pSymA and 17 linkage groups on pSymB that were strongly
associated with strain fitness shifts ($p < 10^{-5}$) (Table 2, Table S6). In contrast, linkage groups on
the chromosome had weaker effects (none with $p < 10^{-5}$), except for LD group 838 (p= 6.38 x
10^{-6}), which was associated with increased strain fitness with *dnf6* and *latd*. The *p*-values of
variants across different replicons may differ in part due to replicon-specific patterns of
relatedness among strains. However, the impact of population structure on GWAS power is not
always straightforward to predict. For instance, the chromosome may exhibit long-range linkage
disequilibrium, resulting in large linkage blocks that tag multiple genes, which can both help or
hinder association resolution.
314
GWAS identifies rhizobial genes with pervasive and specific effects on strain fitness shifts.
To study the genomic landscape underlying rhizobial fitness shifts in response to host mutations,
we identified the top five variants for each host mutant (in total 114 variant groups) based on *p*-
ranks (Table S6). Then we compared the *p*-value ranks and effect sizes of these groups across all
the host mutants (Fig. 4, Fig. S5, Table S7). Most of the candidates on the chromosome were
classified as 'limited effect' variant groups since they ranked among the top ten candidates, in
five or fewer host mutations (Table S7). Exceptions included a massive chromosomal variant
group (901) with 144 variants in 65 genes, among the top ten candidates among nine host
mutants. Six variants with pervasive effects were found on pSymB. On pSymB, four small
variant groups were top-ranked in at least three host mutants (ABC transporter and RKpR, a

dihydrodipicolinate synthase family protein, *pepT*, peptidase T, and a hypothetical protein (Fig.
S5). However, the seven pervasive variants on pSymA were most interesting to us because they
spanned similar hosts and had strong statistical support. These candidates included malonyl-CoA
synthase, SGNH/GDSL hydrolase family protein, *fixL*, an oxygen sensor histidine kinase, a
short-chain oxidoreductase, and a transport protein belonging to the MFS (Major Facilitator
Superfamily). Interestingly, the non-reference variant often increased strain fitness within the
host mutants. In addition, we noted a trend where the pervasive variants that conferred higher
relative fitness in the mutant hosts, tended to have lower fitness in wildtype A17. For example,
the malonyl COA synthase had an alternate allele associated with increased strain fitness
(positive beta) in several host mutant backgrounds (*ipd3, dnf1, dnf2, dnf6, dnf7, nad1, rdn1*, and
sunn1). In contrast, the same allele was associated with decreased fitness (negative beta) in the
wild-type A17 (Fig. 4, Table S6). This pattern suggests a host-dependent allelic effect with the
allele conferring a fitness advantage in mutant hosts becoming disadvantageous in the wild-type
host.
339
Only 82 of the 94 limited-effect variant groups were small (having fewer than three genes). The
majority of these were on pSymB (Table S7). Those with strong statistical support ($p<10^{-5}$)
included variants in or nearby genes coding two ABC transporters (*sunn1, ipd3, dnf1/dnf2,*
rdn1), glutamine synthetase (*latd*) methyltransferases (*nad1*), transcriptional regulators (*rdn1*), a
couple of hypothetical proteins (*ipd3, dnf1/dnf2,dnf5/dnf2,dnf2,sunn1)*and genes involved in
purine metabolism (*dnf3*). Interestingly, a rhizobial glutamine synthetase gene, which converts
glutamate and ammonia into glutamine to assimilate nitrogen, was the strongest candidate for
variation in strain fitness shift in hosts with a mutated *latd,* which encodes a nitrate transporter [55].
This was unexpected since rhizobial glutamine synthetases are generally thought to be strongly
downregulated in nodules where host-driven nitrogen assimilation processes dominate [56].
350
Correlations of host investment and benefit traits with rhizobial traits
Because mutualistic partners affect each other's fitness, we were curious if any host investment
or benefit traits were correlated with rhizobial fitness. At the level of individuals pots, nodule area,
plant biomass, and root-to-shoot ratio covaried with nodule strain diversity and position on the
first RDA axis describing shifts in strain relative fitness (Fig. 5A). Total nodulated area per plant
covaried with the second RDA axis describing shifts in strain relative fitness. Perhaps
unsurprisingly, the host trait that covaried with CFUs released per plant was the number of
nodules per plant. Across the first PC of trait space, we found the wild-type plants grouped with
the AON mutants, *hcl*, and three *dnfs* (3,4,6). In contrast, the second PC tended to separate the
early-acting, AON, and wildtype plants from the late-acting mutants. Genotypic trait
correlations confirmed these patterns (Fig. 5B). Host mutants that were larger and had lower
root-to-shoot ratios, i.e., those that benefited more from rhizobial N-fixation, tended to release
more rhizobia (CFUs) from individual nodules, more rhizobia on a whole-plant level, and had
nodule strain communities that resembled those in A17 nodules (Fig. 5B). Interestingly, most
host traits correlated strongly with strain diversity and position along the first rhizobial relative
fitness RDA axis; The exception was average nodule area/plant, which was most strongly
correlated with host position on the second RDA axis. Details on mutation-specific effects on
host traits can be found in Fig. S6 and Table S3. Our results suggest that symbiotic gene
disruptions strongly shape both partners' fitness in directions reinforcing mutualism—mutations
that reduce rhizobial fitness also reduce host fitness.
371
<u>Discussion</u>

By combining a Select-and-Resequence (S&R) experiment with GWAS, we demonstrate that
mutations in host symbiotic genes create distinct environments that shape rhizobial strain success
and identify rhizobial candidate genes underlying strain fitness shifts in response to each host
mutation. We found rhizobial genetic variants with both pervasive and limited effects across host
mutants, and that symbiotic stage categories provide little clarity about how rhizobial genes
influence strain fitness. We also found that mutations that increased rhizobial fitness tend to
increase host investment and benefit traits. Our findings underscore the complexity of
mutualism, where both host and rhizobial genomes jointly determine the outcomes of symbiosis
at the community and individual strain levels, adding to knowledge of how *Medicago* hosts
determine rhizobial fitness [16,30].

Host genetic mutations impact rhizobial absolute fitness
Legumes releasing rhizobia from their senescing nodules directly contribute to the soil rhizobial
population [28,48,57]. We found that most mutations did not affect the number of rhizobia released
from each nodule, and those that did, *ipd3* and *nad1*, released fewer rhizobia per nodule. The
reduced number of rhizobia per nodule with these two mutants is not unexpected, given that
disrupting *nad1* leads to overactive plant defence responses in nodules and disrupting *ipd3* traps
rhizobia within infection threads, likely reducing survival [32,43]. While our results suggest that the
effects of host genetic mutations on rhizobial fitness may sometimes be gene-specific, we did
observe that rhizobia released per nodule and average nodule area were positively correlated
across mutations. Increasing the number of rhizobia the host supports represents one pathway to
nurture robust populations of rhizobia, and identifying the host allelic variation that increases this
amplification could set the stage for targeted interventions aimed at bolstering soil rhizobial
populations in agricultural fields [58].

Rhizobial fitness shifts depend on host genetics
A large group of NCR peptides is one of the major known host factors for altering strain survival
and growth in nodules [59–62]. Although late-acting NCRs can be important determinants of host
strain specificity in *M. truncatula*, our investigation reveals a more complex landscape [27,63]. Our
analyses of two NCR mutations revealed differing effects on strain fitness within nodules. The
dnf7 mutant, which lacks functional NCR169 [64], was the only mutant that reduced nodule strain
diversity, and it clustered with *ipd3,* which reduced rhizobial population size. *dnf4*, which does
not produce NCR211 and influences rhizobial differentiation during bacteroid formation [65],
caused similar shifts in strain relative fitness as *dnf3* and *dnf6,* suggesting these uncharacterized
mutants may also relate to NCRs. A few rhizobia genes (*bacA, hrrP, yejABEF*) are known to
affect the efficacy of NCRs [35,59,66]; however, none of those genes were top candidates in our
GWAS, suggesting additional mechanisms of host restriction of rhizobia [67–69].

Rhizobial variants with pervasive and limited effects across host mutants
Among the ~120 top rhizobial variants, ~18% affected rhizobial fitness in six or more host
mutations (pervasive genes), and ~82% affected rhizobial fitness in 5 or fewer host mutants
(limited effect genes). Interestingly, pervasive effect alleles tended to exhibit fitness trade-offs;
alleles that conferred higher fitness in host mutants tended to confer lower fitness with wild-type
hosts [16]. Many of the GWAS candidates were identified in only one or a few host mutant
backgrounds, with the strongest associations on pSymB. Unlike the pervasive variants, most
limited-effect variants tended to have little or no effect on strain fitness in the wild-type
background. Most candidate genes with pervasive effects were highly expressed in the nitrogen

fixation zone (Zone III) [56]. For example, the *fixL* gene, which regulates the expression of
nitrogen fixation genes in response to low oxygen levels, Malonyl-CoA synthase that is induced
in the rhizosphere of alfalfa, responsible for malonate catabolism within root nodules [73], and a
NAD(P)H-binding protein from the conserved NmrA domain family that shows high similarity
to a transcriptional regulator, involved in nitrogen metabolite repression in fungi [74]. These genes
represent low-hanging fruit for uncovering novel mechanisms that alter the host range of
symbionts [75].
428
Beyond these candidates, most genes only altered strain fitness in specific host mutants,
indicating that each mutation creates a unique challenge for rhizobia reproduction. Notable
candidates include: A Class I SAM-dependent methyltransferase associated with strain fitness in
nad1 mutants, suggesting the importance of epigenetic modifications for rhizobial survival under
host defences [32], and *stc4*, encoding stachydrine N-demethylase reductase, which enhanced
strain fitness in *dnf4* mutants, possibly by protecting differentiating rhizobia [65]. While these
genes are candidates for altering strain fitness in some hosts and not others, our experiments do
not pinpoint at what stage each host gene exerts the strongest selective filter. One way to follow-
up on these results is with assays comparing rhizobia mutated in the candidate genes identified in
our GWAS on wildtype and mutant hosts, which would allow one to identify the selective
stage—rhizosphere, infection thread, or nodule—where each gene exerts its influence on strain
fitness.
441
Considerations for interpreting mixed-strain experiments
Final strain fitness in nodules is a composite trait that reflects host genetics and microbial traits
such as rhizosphere colonization, competition for infection, and proliferation within nodules. As
such, shifts in relative strain fitness across host genotypes could also be influenced by
differential competitive abilities among strains before infection of each host genotype. Thus,
while our design allows us to detect which host symbiotic genes alter strain fitness, it does not
allow us disentangle the relative contributions of microbial competition versus host filtering.
This limitation is especially relevant in complex communities, where strain-strain interactions
influence host effects [21]. While data suggests the effects of microbe × microbe interactions on
strain relative fitness in our system are weak [26], caution that it is important to interpret our host-
driven fitness shifts as emergent outcomes of host × microbe × microbe interactions. An
additional consideration is that our experiment is designed to measure how growing plants of a
single genotype will influence rhizobial strain fitness. Linking these fine-scale host effects to
long-term dynamics of rhizobial adaptation is an important overarching goal in plant-microbe
interactions that requires incorporating additional ecological and evolutionary contexts including
the importance of soil selection [76] or the cooccurrence of host genotypes across time and space
[13,77].
459
460
Conclusion
Despite legumes being used for soil fertility improvement and actively managed for centuries,
biological nitrogen fixation in agriculture remains inconsistent spatially and temporally [79–82].
This challenge has driven interest and investment in developing microbial inoculants to enhance
crop yield, stress tolerance, disease, and pest management [20]. Rhizobia have been included in
microbe-based bioinoculant formulations for centuries [75]. However, most of those strain
selection decisions are dominated by strain performances observed under single-strain assays
(inoculating a single plant with a single strain) in controlled environments. Thus, strains which

show promise as potential bioinoculants under controlled conditions fail to reach densities high
enough to influence crop yield in realistic agricultural systems [83]. Formulating ways to sustain
rhizobial fitness in the agricultural fields is a critical avenue for plants to continue acquiring
nitrogen from their associated rhizobia [84].

Advances in host-mediated microbiome engineering have demonstrated that plant genes can be
leveraged to assemble trait-optimized microbial communities, with the selection of beneficial
microbes being dependent on the host genotype[81,85]. The findings in this study suggest a route
towards expanding the search for host genes that alter rhizobial fitness. If natural allelic variants
of these genes also favor colonization by high-performing symbionts, it will enable the
development of "designer plants" with enriched mutualistic microbiomes [86]. Here, we identify
host genes that drive strain-specific selection and show that even functionally similar mutations
can have divergent effects on strain fitness in multi-strain contexts [21]. While our study focused
on loss-of-function mutations, studies on segregating natural genetic variation in *Medicago*
suggest that specific host genes can differentially enrich rhizobia but find limited evidence for
widespread balancing or positive selection [16]. Though we do not expect to find these loss-of-
function mutations segregating in natural populations, this work lays a critical foundation:
identifying the host genes that influence rhizobial fitness. A next step could be to systematically
test the effect of naturally occurring alleles within these genes on rhizobial fitness, to pinpoint
the variants that promote or reduce fitness and the mechanisms by which they do it.

Acknowledgments
We thank Prof. Dong Wang of University of Massachusetts, Amherst for generously providing
the *M. truncatula* mutant seeds, Lillian Cherry, Maria Alejandra Gil-Polo, and Jennifer E. Harris
for help with the tedious task of picking and processing nodules. Prof. Katy Heath of University
of Illinois Urbana-Champaign and Dr. Mike Grillo (currently Assistant Professor in Loyola
University, Chicago) for providing strains and sequencing data. This work was supported by the
National Science Foundation grant IOS-1856744 to PT, NY, and LTB and IOS-2243819 to PT
and LTB and was completed using computing resources provided by Penn State Institute for
Computational and Data Science and sequencing by the Huck Genomics Core Facility. LTB's
work was also supported by HATCH grant no. 1025611. The authors acknowledge the Huck
Institutes' Genomics Core Facility (RRID:SCR_023645) for providing library prep and
sequencing services.

Authorship Credits
LTB, NY, PT idea conception and funding, LTB experimental design, LTB, and RGB plant
care; LTB, SG, RB drafted the manuscript; RB, GF, LTB data collection, SG, LTB, RGB, BE,
JS analysis, SG, LTB, RB, JS, BE, PT, NY, GF manuscript revisions.

Competing Interests:
The authors declare no competing interests.

Data Accessibility
All the raw reads from sequencing the rhizobial strains in the initial community and in the
rhizobial genomic DNA extracted from the nodule pools of the individual host mutants have
been deposited in NCBI under the Bioproject PRJNA401437 (Submission ID: SUB14622715).
All raw data and analyses code have been uploaded to the Github
(https://github.com/BurghardtLab/HostMutants2021_C86meliloti)

References

1. Friesen, M. L. *et al.* Microbially Mediated Plant Functional Traits. *Annu. Rev. Ecol. Evol. Syst.* **42**, 23–46 (2011).

2. Percy, D. M. John N. Thompson, The Geographic Mosaic of Coevolution. *Syst. Biol.* **56**, 149–151 (2007).

3. Bartha, I. *et al.* A genome-to-genome analysis of associations between human genetic variation, HIV-1 sequence diversity, and viral control. *eLife* **2**, e01123 (2013).

4. Jones, J. D. G. & Dangl, J. L. The plant immune system. *Nature* **444**, 323–329 (2006).

5. Amandine, C. *et al.* Unraveling coevolutionary dynamics using ecological genomics. *Trends Genet.* **38**, 1003–1012 (2022).

6. Demirjian, C., Vailleau, F., Berthomé, R. & Roux, F. Genome-wide association studies in plant pathosystems: success or failure? *Trends Plant Sci.* **28**, 471–485 (2023).

7. Corwin, J. A. *et al.* The Quantitative Basis of the Arabidopsis Innate Immune System to Endemic Pathogens Depends on Pathogen Genetics. *PLOS Genet.* **12**, e1005789 (2016).

8. Newberry, E. A., Minsavage, G. V., Holland, A., Jones, J. B. & Potnis, N. Genome-Wide Association to Study the Host-Specificity Determinants of Xanthomonas perforans. *Phytopathology®* **113**, 400–412 (2023).

9. Medina, M. & Sachs, J. L. Symbiont genomics, our new tangled bank. *Genomics* **95**, 129–137 (2010).

10. Clare, E. L., Schiestl, F. P., Leitch, A. R. & Chittka, L. The promise of genomics in the study of plant-pollinator interactions. *Genome Biol.* **14**, 207 (2013).

11. *Mutualism*. (Oxford University Press, Oxford, United Kingdom, 2015).

12. Batstone, R. T. Genomes within genomes: nested symbiosis and its implications for plant evolution. *New Phytol.* **234**, 28–34 (2022).

13. Heath, K. D. & Stinchcombe, J. R. EXPLAINING MUTUALISM VARIATION: A NEW

EVOLUTIONARY PARADOX?: PERSPECTIVE. *Evolution* **68**, 309–317 (2014).

14. Turkarslan, S. *et al.* Synergistic epistasis enhances the co-operativity of mutualistic

interspecies interactions. *ISME J.* **15**, 2233–2247 (2021).

15. Batstone, R. T. *et al.* Genome-wide association studies across environmental and genetic

contexts reveal complex genetic architecture of symbiotic extended phenotypes. *mBio* **13**,

e01823-22 (2022).

16. Epstein, B. *et al.* Combining GWAS and population genomic analyses to characterize

coevolution in a legume-rhizobia symbiosis. *Mol. Ecol.* mec.16602 (2022)

doi:10.1111/mec.16602.

17. Carlson, C. & Frederickson, M. E. An emerging view of coevolution in the legume–

rhizobium mutualism. *Mol. Ecol.* **32**, 3793–3797 (2023).

18. Gohar, D. *et al.* Genomic evidence of symbiotic adaptations in fungus-associated bacteria.

iScience **28**, 112253 (2025).

19. Denison, R. F., Kiers, E. T. & West, S. A. Darwinian Agriculture: When Can Humans Find

Solutions Beyond the Reach of Natural Selection? *Q. Rev. Biol.* **78**, 145–168 (2003).

20. Busby, P. E. *et al.* Research priorities for harnessing plant microbiomes in sustainable

agriculture. *PLOS Biol.* **15**, e2001793 (2017).

21. Burghardt, L. T. & diCenzo, G. C. The evolutionary ecology of rhizobia: multiple facets of

competition before, during, and after symbiosis with legumes. *Curr. Opin. Microbiol.* **72**,

102281 (2023).

22. Oldroyd, G. E. D., Murray, J. D., Poole, P. S. & Downie, J. A. The Rules of Engagement in

the Legume-Rhizobial Symbiosis. *Annu. Rev. Genet.* **45**, 119–144 (2011).

23. Poole, P., Ramachandran, V. & Terpolilli, J. Rhizobia: from saprophytes to endosymbionts.

Nat. Rev. Microbiol. **180**, 1691 (2018).

24. Heath, K. D. & Tiffin, P. Context dependence in the coevolution of plant and rhizobial

mutualists. *Proc. R. Soc. B Biol. Sci.* **274**, 1905–1912 (2007).

25. Burghardt, L. T. *et al.* Select and resequence reveals relative fitness of bacteria in symbiotic

and free-living environments. *Proc. Natl. Acad. Sci.* **115**, 2425–2430 (2018).

26. Burghardt, L. T., Epstein, B., Hoge, M., Trujillo, D. & Tiffin, P. *Host-Associated Rhizobia*

Fitness: Dependence on Nitrogen, Density, Community Complexity, and Legume Genotype.

http://biorxiv.org/lookup/doi/10.1101/2020.11.20.392183 (2020)

doi:10.1101/2020.11.20.392183.

27. Roy, S. *et al.* Celebrating 20 Years of Genetic Discoveries in Legume Nodulation and

Symbiotic Nitrogen Fixation. *Plant Cell* **32**, 15–41 (2020).

28. Burghardt, L. T. Evolving together, evolving apart: measuring the fitness of rhizobial

bacteria in and out of symbiosis with leguminous plants. *New Phytol.* **228**, 28–34 (2020).

29. Ratcliff, W. C., Underbakke, K. & Denison, R. F. Measuring the fitness of symbiotic

rhizobia. *Symbiosis* **55**, 85–90 (2011).

30. Burghardt, L. T., Trujillo, D. I., Epstein, B., Tiffin, P. & Young, N. D. A Select and

Resequence Approach Reveals Strain-Specific Effects of *Medicago* Nodule-Specific PLAT-

Domain Genes. *Plant Physiol.* **182**, 463–471 (2020).

31. Epstein, B. *et al.* Genome-Wide Association Analyses in the Model Rhizobium *Ensifer*

meliloti. *mSphere* **3**, e00386-18 (2018).

32. Wang, C. *et al.* NODULES WITH ACTIVATED DEFENSE 1 is required for maintenance of

rhizobial endosymbiosis in *Medicago truncatula*. *New Phytol.* **212**, 176–191 (2016).

33. Yang, S., Tang, F., Gao, M., Krishnan, H. B. & Zhu, H. *R* gene-controlled host specificity in

the legume–rhizobia symbiosis. *Proc. Natl. Acad. Sci.* **107**, 18735–18740 (2010).

34. Tang, F., Yang, S., Liu, J. & Zhu, H. *Rj4* , a Gene Controlling Nodulation Specificity in

Soybeans, Encodes a Thaumatin-Like Protein But Not the One Previously Reported. *Plant*

Physiol. **170**, 26–32 (2016).

35. Price, P. A. *et al.* Rhizobial peptidase HrrP cleaves host-encoded signaling peptides and

mediates symbiotic compatibility. *Proc. Natl. Acad. Sci.* **112**, 15244–15249 (2015).

36. Oldroyd, G. E. D. Speak, friend, and enter: signalling systems that promote beneficial

symbiotic associations in plants. *Nat. Rev. Microbiol.* **11**, 252–263 (2013).

37. Wheatley, R. M. *et al.* Lifestyle adaptations of Rhizobium from rhizosphere to symbiosis.

Proc. Natl. Acad. Sci. U. S. A. **117**, 23823–23834 (2020).

38. Ferguson, B. J. *et al.* Legume nodulation: The host controls the party. *Plant Cell Environ.* **42**,

41–51 (2019).

39. Bright, L. J., Liang, Y., Mitchell, D. M. & Harris, J. M. The *LATD* Gene of *Medicago*

truncatula Is Required for Both Nodule and Root Development. *Mol. Plant-Microbe*

Interactions® **18**, 521–532 (2005).

40. Hogg, B. V., Cullimore, J. V., Ranjeva, R. & Bono, J.-J. The *DMI1* and *DMI2* Early

Symbiotic Genes of *Medicago truncatula* Are Required for a High-Affinity Nodulation

Factor-Binding Site Associated to a Particulate Fraction of Roots. *Plant Physiol.* **140**, 365–

373 (2006).

41. Messinese, E. *et al.* A Novel Nuclear Protein Interacts With the Symbiotic DMI3 Calcium-

and Calmodulin-Dependent Protein Kinase of *Medicago truncatula*. *Mol. Plant-Microbe*

Interactions® **20**, 912–921 (2007).

42. Smit, P. *et al.* Medicago LYK3, an Entry Receptor in Rhizobial Nodulation Factor Signaling.

Plant Physiol. **145**, 183–191 (2007).

43. Ovchinnikova, E. *et al.* IPD3 Controls the Formation of Nitrogen-Fixing Symbiosomes in

Pea and *Medicago* Spp. *Mol. Plant-Microbe Interactions®* **24**, 1333–1344 (2011).

44. Starker, C. G., Parra-Colmenares, A. L., Smith, L., Mitra, R. M. & Long, S. R. Nitrogen

Fixation Mutants of *Medicago truncatula* Fail to Support Plant and Bacterial Symbiotic

Gene Expression. *Plant Physiol.* **140**, 671–680 (2006).

45. Schnabel, E., Journet, E.-P., de Carvalho-Niebel, F., Duc, G. & Frugoli, J. The Medicago

truncatula SUNN Gene Encodes a CLV1-like Leucine-rich Repeat Receptor Kinase that

Regulates Nodule Number and Root Length. *Plant Mol. Biol.* **58**, 809–822 (2005).

46. Schnabel, E. L. *et al.* The *ROOT DETERMINED NODULATION1* Gene Regulates Nodule

Number in Roots of *Medicago truncatula* and Defines a Highly Conserved, Uncharacterized

Plant Gene Family. *Plant Physiol.* **157**, 328–340 (2011).

47. Riley, A. B., Grillo, M. A., Epstein, B., Tiffin, P. & Heath, K. D. Discordant population

structure among rhizobium divided genomes and their legume hosts. *Mol. Ecol.* **32**, 2646–

2659 (2023).

48. Denison, R. F. & Kiers, E. T. Life histories of symbiotic rhizobia and mycorrhizal fungi.

Curr. Biol. CB **21**, R775-85 (2011).

49. Darch, S. E., West, S. A., Winzer, K. & Diggle, S. P. Density-dependent fitness benefits in

quorum-sensing bacterial populations. *Proc. Natl. Acad. Sci.* **109**, 8259–8263 (2012).

50. Fåhraeus, G. The Infection of Clover Root Hairs by Nodule Bacteria Studied by a Simple

Glass Slide Technique. *Microbiology* **16**, 374–381 (1957).

51. Kessner, D., Turner, T. L. & Novembre, J. Maximum Likelihood Estimation of Frequencies

of Known Haplotypes from Pooled Sequence Data. *Mol. Biol. Evol.* **30**, 1145–1158 (2013).

52. Li, H. A statistical framework for SNP calling, mutation discovery, association mapping and

population genetical parameter estimation from sequencing data. *Bioinformatics* **27**, 2987–

2993 (2011).

53. Purcell, S. *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-Based

Linkage Analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

54. Domonkos, Á. *et al.* NAD1 Controls Defense-Like Responses in Medicago truncatula

Symbiotic Nitrogen Fixing Nodules Following Rhizobial Colonization in a BacA-

Independent Manner. *Genes* **8**, 387 (2017).

55. Harris, J. M. & Dickstein, R. Control of root architecture and nodulation by the *LATD/NIP*

transporter. *Plant Signal. Behav.* **5**, 1365–1369 (2010).

56. Roux, B. *et al.* An integrated analysis of plant and bacterial gene expression in symbiotic

root nodules using laser-capture microdissection coupled to RNA sequencing. *Plant J.* **77**,

817–837 (2014).

57. Brewin, N. J. Development of the Legume Root Nodule. *Annu. Rev. Cell Biol.* **7**, 191–226

(1991).

58. Gopalakrishnan, S. *et al.* Plant growth promoting rhizobia: challenges and opportunities. *3*

Biotech **5**, 355–377 (2015).

59. Haag, A. F. *et al.* Protection of Sinorhizobium against Host Cysteine-Rich Antimicrobial

Peptides Is Critical for Symbiosis. *PLoS Biol.* **9**, e1001169 (2011).

60. Montiel, J. *et al.* Morphotype of bacteroids in different legumes correlates with the number

and type of symbiotic NCR peptides. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 5041–5046 (2017).

61. Czernic, P. *et al.* Convergent Evolution of Endosymbiont Differentiation in Dalbergioid and

Inverted Repeat-Lacking Clade Legumes Mediated by Nodule-Specific Cysteine-Rich

Peptides. *Plant Physiol.* **169**, 1254–1265 (2015).

62. Mergaert, P. *et al.* Eukaryotic control on bacterial cell cycle and differentiation in the

Rhizobium –legume symbiosis. *Proc. Natl. Acad. Sci.* **103**, 5230–5235 (2006).

63. Walker, L., Lagunas, B. & Gifford, M. L. Determinants of host range specificity in legume-

rhizobia symbiosis. *Front. Microbiol.* **11**, (2020).

64. Horváth, B. *et al.* Loss of the nodule-specific cysteine rich peptide, NCR169, abolishes

symbiotic nitrogen fixation in the Medicago truncatula dnf7 mutant. *Proc. Natl. Acad. Sci.*

U. S. A. **112**, 15232–15237 (2015).

65. Kim, M. *et al.* An antimicrobial peptide essential for bacterial survival in the nitrogen-fixing

symbiosis. *Proc. Natl. Acad. Sci.* **112**, 15238–15243 (2015).

66. Nicoud, Q. *et al.* Sinorhizobium meliloti Functions Required for Resistance to Antimicrobial

NCR Peptides and Bacteroid Differentiation. *mBio* **12**, e00895-21 (2021).

67. Guefrachi, I. *et al.* Extreme specificity of NCR gene expression in Medicago truncatula.

BMC Genomics **15**, 712 (2014).

68. Burghardt, L. T. *et al.* Transcriptomic basis of genome by genome variation in a legume-

rhizobia mutualism. *Mol. Ecol.* **26**, 6122–6135 (2017).

69. Lima, R. M., Kylarová, S., Mergaert, P. & Kondorosi, É. Unexplored Arsenals of Legume

Peptides With Potential for Their Applications in Medicine and Agriculture. *Front.*

Microbiol. **11**, 1307 (2020).

70. Savolainen, O., Lascoux, M. & Merilä, J. Ecological genomics of local adaptation. *Nat. Rev.*

Genet. **14**, 807–820 (2013).

71. Hoban, S. *et al.* Finding the Genomic Basis of Local Adaptation: Pitfalls, Practical Solutions,

and Future Directions. *Am. Nat.* **188**, 379–397 (2016).

72. Elena, S. F. Local adaptation of plant viruses: lessons from experimental evolution. *Mol.*

Ecol. **26**, 1711–1719 (2017).

73. Karunakaran, R., East, A. K. & Poole, P. S. Malonate Catabolism Does Not Drive N$_2$

Fixation in Legume Nodules. *Appl. Environ. Microbiol.* **79**, 4496–4498 (2013).

74. Stammers, D. K. The structure of the negative transcriptional regulator NmrA reveals a

structural superfamily which includes the short-chain dehydrogenase/reductases. *EMBO J.*

20, 6619–6626 (2001).

75. Mus, F. *et al.* Symbiotic Nitrogen Fixation and the Challenges to Its Extension to

Nonlegumes. *Appl. Environ. Microbiol.* **82**, 3698–3710 (2016).

76. Burghardt, L. T., Epstein, B. & Tiffin, P. Legacy of prior host and soil selection on rhizobial

fitness in planta. *Evolution* **73**, 2013–2023 (2019).

77. Wippel, K. *et al.* Host preference and invasiveness of commensal bacteria in the Lotus and

Arabidopsis root microbiota. *Nat. Microbiol.* **6**, 1150–1162 (2021).

78. Sugawara, M. *et al.* Variation in bradyrhizobial NopP effector determines symbiotic

incompatibility with Rj2-soybeans via effector-triggered immunity. *Nat. Commun.* **9**, 3139

(2018).

79. Kaminsky, L. M., Trexler, R. V., Malik, R. J., Hockett, K. L. & Bell, T. H. The Inherent

Conflicts in Developing Soil Microbial Inoculants. *Trends Biotechnol.* **37**, 140–151 (2019).

80. Soumare, A. *et al.* Exploiting Biological Nitrogen Fixation: A Route Towards a Sustainable

Agriculture. *Plants* **9**, 1011 (2020).

81. Trivedi, P., Batista, B. D., Bazany, K. E. & Singh, B. K. Plant–microbiome interactions

under a changing world: responses, consequences and perspectives. *New Phytol.* **234**, 1951–

1959 (2022).

82. Batstone, R. T., Ibrahim, A. & MacLean, L. T. Microbiomes: Getting to the root of the

rhizobial competition problem in agriculture. *Curr. Biol.* **33**, R777–R780 (2023).

83. Parnell, J. J. *et al.* From the Lab to the Farm: An Industrial Perspective of Plant Beneficial

Microorganisms. *Front. Plant Sci.* **7**, (2016).

84. Porter, S. S. & Sachs, J. L. Agriculture and the Disruption of Plant–Microbial Symbiosis.

Trends Ecol. Evol. **35**, 426–439 (2020).

85. Batstone, R. T., O'Brien, A. M., Harrison, T. L. & Frederickson, M. E. Experimental

evolution makes microbes more cooperative with their local host genotype. *Science* **370**,

476–478 (2020).

86. Batstone, R. T., Burghardt, L. T. & Heath, K. D. Phenotypic and genomic signatures of

interspecies cooperation and conflict in naturally occurring isolates of a model plant

symbiont. *Proc. R. Soc. B Biol. Sci.* **289**, 20220477 (2022).

<u>Figure Legends</u>
Fig. 1: Schematic overview of the experiment used to evaluate rhizobial fitness and host traits
across different host mutant backgrounds. A set of 86 *Sinorhizobium* strains was cultured and
combined into a mixed inoculum. This inoculum was applied to *Medicago truncatula* wild-type
(A17) and host mutants with gene disruptions that act at different stages of symbiosis: Early,
Late, and Autoregulation of Nodulation (AON). The number of host mutants per stage is
indicated in parentheses. Host and rhizobial traits were measured at the nodule and whole-plant
scales. The experimental conditions are clarified under 'Experimental Details'. Each treatment
included five pots per host mutant with three plants per pot, grown in randomized blocks under
nitrogen(N)-free conditions for 8 weeks. Traits measured include indicators of rhizobial benefit,
host benefit, and host investment (Refer Table 1 for more details). Different coloured nodules
indicate nodules formed by different rhizobial strains.
729
Fig. 2: Host symbiotic mutations influence rhizobial fitness. (A) Host mutations altered
nodule numbers in consistent and predictable ways. (B–C) Host genotype also influenced
rhizobial fitness, as measured by $\log_{10}$-transformed colony-forming units per nodule and per
plant (CFU/nodule) (B) and CFU /plant (C). (D) Nodule strain diversity was largely unaffected
by host mutations. Closed triangles indicate significant increase or decrease relative to wild-type
A17 (based on Tukey-adjusted *p* values). Boxplots are color-coded by the symbiotic stage
disrupted by each host mutation (see Table S1). (E) Heatmap showing the median relative fitness
of each of 86 *S. meliloti* strains across the different host genotypes. Strains enriched relative to
their initial inoculum appear in shades of red; depleted strains appear in shades of blue. Host
genotypes are hierarchically clustered based on similarity in strain fitness profiles. Strains are
ranked by their relative fitness in A17. Mutants in bold differ significantly from A17 (see Table
S5 for RDA paired with the PERMANOVA results).
742
Fig. 3: GWAS of strain fitness shifts in host mutants. (A) The rhizobial strain fitness shifts
for each of the 86 strains, calculated by subtracting individual relative strain fitness measured in
A17 from the relative strain fitness in the host mutants. The heatmap is hierarchically based on
the similarity of median 'shift' values across hosts. Shades of orange denote positive fitness shift
compared to A17, while shades of purple denote negative fitness shift compared to A17. (B)
Manhattan plot indicating the SNP (single-nucleotide polymorphism) positions, based on S.
meliloti USDA1106 association with the relative strain fitness shifts across the 15 host mutants
and strain fitness in A17. The dotted line indicates the *p* value threshold used for identifying
candidate genes ($p < 10^{-5}$), and significant associations are color-coded according to host
mutation and A17. The GWAS was run separately on each replicon to account for different
population structures. The 'Δ' indicates 'strain fitness shift' in that host mutant. For comparison,
we also ran the GWAS on A17 relative strain fitness.
755
Fig. 4: Rhizobial candidate genes underlying strain fitness shifts predominantly occur on
pSymA. Heatmaps based on the inverse *p*-value ranks (red-yellow) and β values which denote
the effect sizes (red-blue) for the top five variant groups on the pSymA across the 15 host
mutants and A17. Rows represent candidate variant groups, and columns represent host mutants.
The host mutants are colour coded based on the stage of disruption they affect. Progressive dark
shades (low *p*-value ranks) indicate stronger association between the variant group and strain
fitness shift in that host mutant. Heatmap based on the effect sizes (β values) for the same
candidate variant groups was ordered to match the *p*-value ranking. Shades of red indicate host
mutants where the corresponding variant has a positive effect on strain fitness shift, shades of

blue indicate negative effects as compared to A17. Colour intensity corresponds to effect
magnitude. Alternate allele frequencies (i.e., not present in reference genome USDA1106) for
each candidate are also shown in parentheses next to the gene names. Pervasiveness of a variant
group is defined has number of times a variant group is among the top 10 candidates (based on
the *p* value ranks) within a host mutant. A variant group is classified as "pervasive" if it is in the
top10 across 6 or more host genotypes while a variant is classified as "limited effect" if it is in
the top10 candidates across 5 or less host genotypes. We have mentioned the number of times a
variant appears in the top 10 under the "Pervasiveness" column in the figure and have indicated
the pervasive genes in bold. (See Fig. S5 for the other replicons.)

774
775
Fig 5: Covariation and correlation structure among host and rhizobial traits.
(A) Principal component analysis (PCA) of host investment traits, benefit traits, and rhizobial
fitness outcomes, conducted at the level of individual pots. Arrows represent the loadings of each
trait on the first two principal components, which together explain 51.2% and 16.7% of the total
variance, respectively. To avoid loss of representation due to missing data, values for two
genotypes (*latd* and *rdn1*) were averaged across replicates prior to PCA, as these genes had
incomplete trait data that would otherwise be excluded by listwise deletion.
(B) All pairwise combinations of genotypic trait correlations (Pearsons). Each ellipse represents
a pairwise Pearson correlation between traits, with ellipse elongation and color saturation
indicating the strength and direction of the correlation (blue = positive, red = negative). Strong
correlations may reflect shared genetic regulation or pleiotropy.

<u>Tables</u>
788
789 **Table 1:** Definitions of the symbiotic traits examined

Symbiotic Traits	Category	Interpretation
CFU* per nodule	Rhizobial Benefit	Rhizobia released per nodule indicates the average individual strain fitness increase from host association
CFU* per plant	Rhizobial Benefit	Rhizobia released per plant indicates the contribution of each host plant to overall rhizobial population size
Nodule strain diversity	Rhizobial Benefit	Shannon diversity index (H) calculated for strains isolate from nodule indicates host selectivity
Strain relative fitness	Rhizobial Benefit	Enrichment/depletion of specific strains relative to frequency in inoculum: $\log_2$(selected freq./initial freq.)
Nodules per plant	Host Investment	Number of successful host attempts to form associations with rhizobia
Average area per nodule	Host Investment	Host investment in each nodule
Total nodule area per plant	Host Investment	Total plant investment in nodules (i.e. nitrogen fixation)
Total biomass per plant	Host Benefit	Proxy for total nitrogen fixed when plants are grown in nitrogen limited conditions
Root-to-shoot ratio	Host Benefit	Lower values indicate lower investment in roots. In a N limited environment we infer that nitrogen needs were met.

790
791 *CFU= Colony Forming Units

Table 2: Candidate rhizobial genes in small variant groups (three genes or fewer) associated with ($p < 10^{-5}$) variation in rhizobial relative strain fitness. See Table S6 for information on all variant groups.

LD group[1]	Host mutants	Location	Alt. allele freq[2]	Eff. alt. vars in mutants	Eff alt. vars in WT	Variants	Genes ID's[3]
9034	*dnf1,dnf5/dnf2, dnf2, dnf4, dnf7, ipd3, nad1,rdn1 sunn1, sunn4*	pSymA	0.07	+	-	1	CDO30_RS23065:SGNH/GDSL hydrolase family protein
6978.a lt[4]	*dnf1, dnf1/dnf2, dnf2, dnf4, dnf7, dnf5/dnf2, ipd3, rdn1, sunn1*	pSymA	0.07	+	none	1	CDO30_RS23160:NAD(P)H-binding protein
5279	*dnf1, dnf2, dnf7, ipd3, nad1,rdn1, sunn1*	pSymA	0.07	+	-	1	CDO30_RS21460:malonyl-CoA synthase
7969	*dnf1, dnf2, dnf6, nad1, sunn4*	pSymA	0.06	+	none	1	CDO30_RS22325:hypothetical protein
6968	*dnf4, dnf7,ipd3, sunn1, sunn4*	pSymA	0.10	+	none	1	CDO30_RS23125:SDR family oxidoreductase (pseudogene)
10122	*dnf1, dnf1/dnf2, dnf2, sunn1*	pSymB	0.07	+	-	1	CDO30_RS27255:hypothetical protein
12516	*dnf1/dnf2, rdn1, sunn1*	pSymB	0.06	+	none	2	CDO30_RS28530:ABC transporter ATP-binding protein; CDO30_RS28525:*RkpR*
10601	*dnf7, ipd3, ltd*	pSymB	0.05	+	none	1	CDO30_RS28740:*pepT*, peptidase T
3770	*rdn1, sunn4*	pSymA	0.14	+	-	1	CDO30_RS18080:*fixL*, oxygen sensor histidine kinase *fixL*
12013	*dnf1/dnf2, ipd3*	pSymB	0.05	+	none	1	CDO30_RS28675:hypothetical protein
10882	*dnf4*	pSymB	0.05	+	none	1	CDO30_RS30150:glycosyltransferase
9671	*dnf3*	pSymB	0.07	+	none	3	CDO30_RS27435:four-carbon acid sugar kinase family protein
10140	*dnf3*	pSymB	0.08	+	none	1	CDO30_RS27435:four-carbon acid sugar kinase family protein
10913	*dnf3*	pSymB	0.05	+	none	2	CDO30_RS27440:xanthine dehydrogenase family protein molybdopterin-binding subunit; CDO30_RS27445:xanthine dehydrogenase family protein subunit M
10915	*dnf3*	pSymB	0.05	+	none	42	CDO30_RS27440:xanthine dehydrogenase family protein molybdopterin-binding subunit; CDO30_RS27445:xanthine dehydrogenase family protein subunit M
8294	*dnf4*	pSymA	0.07	+	none	1	CDO30_RS19605:*stc4*, stachydrine N-demethylase reductase subunit Stc4
8754	*dnf5/dnf2*	pSymB	0.1	+	none	2	CDO30_RS32445:hypothetical protein
9239	*dnf5/dnf2*	pSymB	0.09	+	none	1	CDO30_RS32445:hypothetical protein
10903	*ipd3*	pSymB	0.05	+	none	1	CDO30_RS27310:ABC transporter ATP-binding protein
838	*latd*	chr	0.09	+	none	1	CDO30_RS12270:glutamine synthetase family protein
4107	*nad1*	pSymA	0.11	-	none	1	CDO30_RS22005:amidase
4386	*nad1*	pSymA	0.09	-	none	2	CDO30_RS22060:Atu4866 domain-containing protein;CDO30_RS22055:class I SAM-dependent methyltransferase;
9865	*rdn1*	pSymB	0.07	+	none	7	CDO30_RS32685:four-carbon acid sugar kinase family protein; CDO30_RS32675:NAD(P)-dependent oxidoreductase; CDO30_RS32705:LacI family transcriptional regulator
9926	*rdn1*	pSymB	0.07	+	none	4	CDO30_RS31555:MASE1 domain-containing protein; CDO30_RS32090:TRAP transporter large permease; CDO30_RS32070:class I SAM-dependent methyltransferase
9941	*rdn1*	pSymB	0.07	+	none	1	CDO30_RS25485:pyridoxamine 5'-phosphate oxidase family protein
10728	*rdn1*	pSymB	0.5	+	none	1	CDO30_RS32910:LysR family transcriptional regulator
12841	*sunn1*	pSymB	0.07	+	none	1	CDO30_RS28530:ABC transporter ATP-binding protein

Notes: [1]LD group numbers were obtained from [1616] , [2]AF= non-reference (alternate) allele frequency, [3]Locus tags are from the *Sinorhizobium meliloti* USDA1106 reference annotation available on MAGE (http://www.genoscope.cns.fr/agc/microscope/home/index.php), [4]A tri allelic SNP

1. Grow 86 *Sinorhizobium* isolates and mix for inoculum

Experiment Details
- 5 pots/host mutant with 3 plants/pot
- Randomized block design
- N-free fertilizer
- Harvested at 8 weeks

2. Inoculate *Medicago* hosts mutants

Wildtype   **Host mutants spanning stage of symbiosis**

| A17 | Early (6) | Late (9) | AON (3) |

3. Measure host and rhizobial traits

Rhizobial Benefits	Host Benefits	Host Investments
1. CFU/nodule	5. Total biomass/plant	7. Nodules/plant
2. CFU/plant	6. Root-to-shoot ratio	8. Average area/nodule
3. Nodule isolate diversity		9. Total nodulated area
4. Relative strain fitness		

Per nodule release

Per plant release

Low ⟶ High   Low ⟶ High   Low ⟶ High

Disruption Type:
WT   Early   Late   AON

Disruption Type:
WT Early Late AON

Pvalue Rank
1 2 3 4 5 9 90

Beta

Pervasiveness

12	CDO30_RS23155:MFS transporter (0.07)
14	CDO30_RS23160:NAD(P)H-binding protein (0.071)
9	CDO30_RS23125:SDR family oxidoreductase (pseudogene)(0.082)
13	CDO30_RS23065:SGNH/GDSL hydrolase family protein(0.075)
8	CDO30_RS22325:hypothetical protein (0.07)
12	CDO30_RS21460:malonyl–CoA synthase (0.07)
9	CDO30_RS18080:*fixL*, oxygen sensor histidine kinase (0.14)
3	CDO30_RS22055:class I SAM–dependent methyltransferase (0.093)
2	CDO30_RS22005:amidase (0.116)
3	CDO30_RS21460:malonyl–CoA synthase (0.07)
1	CDO30_RS19190:*nodQ*, bifunctional sulfate adenylyltransferase/adenylyl–sulfate kinase (0.075)
4	CDO30_RS19605:*stc4*, stachydrine N–demethylase reductase subunit Stc4(0.072)
1	CDO30_RS22335:site–specific integrase (0.188)
1	CDO30_RS20200:type II toxin–antitoxin system RelE/ParE family toxin(0.214)
1	CDO30_RS21100:DUF2200 domain–containing protein (0.081)
3	CDO30_RS21085:class I SAM–dependent methyltransferase (0.093)
1	CDO30_RS20205:type II toxin–antitoxin system ParD family antitoxin (0.198)
1	CDO30_RS21175:TRAP transporter small permease (0.081)
1	CDO30_RS18965:*traA*, Ti–type conjugative transfer relaxase (0.267)
2	CDO30_RS21490:S8 family serine peptidase (0.07)
1	CDO30_RS21375:PhzF family phenazine biosynthesis protein (0.198)
1	CDO30_RS21370:cold–shock protein (0.093)
2	CDO30_RS19120:*glmS*, glutamine–––fructose–6–phosphate transaminase (isomerizing) (0.372)
1	CDO30_RS24215:CopG family transcriptional regulator (pseudogene)(0.105)
1	CDO30_RS19555:*ccoN*, cytochrome–c oxidase2C cbb3–type subunit I (0.106)
1	CDO30_RS24970:type II toxin–antitoxin system RelE/ParE family toxin (0.128)
1	CDO30_RS22190:aldo/keto reductase(0.314)

* indicates the number of perturbations in which the LD group ranked within the top 10 candidates