Apparent timescaling of fossil diversification rates is caused by sampling bias

Bouwe R. Reijenga[1]* & Roger A. Close[1]

[1]Department of Earth Sciences, University of Oxford, Oxford OX1 3AN, UK

*Correspondence: bouwe.reijenga@earth.ox.ac.uk

Key words

speciation, extinction, macroevolution, sampling bias, fossil record, Phanerozoic, age-rate scaling

Highlights

8   • Studies have found that fossil and molecular diversification rates scale with time
9   • Such rate scaling hints at a disconnect between micro- and macroevolution
10   • These trends are absent from the fossil record when controlling for sampling biases
11   • Rate scaling in general may be artefactual, or fossils could show distinct patterns

Summary

Negative scaling relationships between both speciation and extinction rates on the one hand, and the
age or duration of organismal groups on the other, are pervasive, and recovered in both molecular
phylogenetic and fossil time series. The agreement between molecular and fossil data hints at a
universal cause, and potentially to incongruence between micro- and macroevolution. However, the
existence of negative rate scaling in fossil time series has not undergone the same level of scrutiny as
in molecular data. Here, we analyse the marine fossil record across the last ~538.8 Ma of the
Phanerozoic to investigate the presence and strength of negative rate scaling. We find that negative
rate scaling arises under commonly applied age range-based per-capita rates, which do not control
for sampling bias, but are severely reduced or absent when metrics are used that do correct for
sampling. We further show by simulation that even moderately-incomplete sampling of species
occurrences through time may induce rate scaling. We thus conclude that there are no significant
scaling relationships present in these fossil clades, and that any apparent trend is caused by sampling
artefacts and taxonomic practices. If rate scaling in molecular phylogenies is genuine, the absence of
such a relationship in the fossil record will provide a valuable benchmark and constraint on what
processes can cause it.

Results and discussion

The presence of age-rate scaling (ARS) is a long-studied phenomenon in paleobiology[1,2], and bears
close resemblance to the problem of 'spurious' self-correlation when a ratio is plotted against its
denominator across ecology, evolution, geology and statistics[3–7]. For instance, when an evolutionary
rate such as the number of extinctions per-Myr is plotted against time, a negative correlation is
almost inevitable[8]. In the palaeobiological literature, rates of origination and extinction and typically
measured over specified geological intervals such as geological stages. These intervals are irregular in
duration, so, early on, rates were normalised for interval duration to obtain per-Myr rates. However,
this is inaccurate since events are not evenly spread throughout geological stages[2,9], might even be
erroneous if diversification (origination − extinction) happens in pulses[10,11], and propagates error in
the estimation of geological intervals[1,9]. Foote showed that per-Myr rate metrics indeed suffered
from negative correlation between rate and interval duration, leading to the abandonment of
normalisation by interval duration in favour of per-interval rates[1,12].

A recent study by Henao-Diaz et al. has reinvigorated interest in ARS[13], and concluded that
ARS – which might more accurately be called duration-rate scaling, as clades can occur either far
back in time or near the present – was present in both diversification rates estimated from molecular
phylogenetic trees and fossil time series. They compiled a dataset of first and last appearances for 90
orders of mammals, plants, and marine animals from Sepkoski's compendium[14]. Diversification rates
were calculated via Foote's per-capita metric (See STAR Methods) and normalised by interval
duration[15–17]. ARS was subsequently studied between the duration of taxonomic orders and their
mean diversification rates through time. Their findings suggest that rates across different timescales
cannot be compared[18], and if true might reveal differences between micro- and macroevolutionary
processes[19,20]. However, apart from the issue of plotting ratios against their denominator, a key
concern with the use of Foote's metrics is that those equations are biased by sampling and
taxonomic artefacts[9,21,22]. Therefore, the question is whether ARS in fossil diversification rate
analyses is caused by failure to control for sampling biases.

Here, we build on the work of Foote[1] and Henao-Diaz et al.[13] by investigating how different
sampling artefacts might exaggerate ARS, and if they can be addressed using more appropriate
methods. We do note that some palaeobiologists might inherently question the usefulness of
plotting rates against clade duration, and we neither agree nor disagree with this view here. There
are two steps to our analysis: (*i*) We gather occurrence data for marine animal orders from the
Paleobiology Database (*n* = 408 orders after data cleaning; http://www.paleobiodb.org). Foote's
metrics are only capable of using full age-ranges (first and last appearances) of taxa, but fossil
occurrence data containing age, locality and identity for each taxon is now readily available. Fossil
occurrences also inherently contain information about sampling intensity through time. (*ii*) We
compare Foote's per-capita rates to Alroy's second-for-third metrics (See STAR Methods)[21] to
calculate rates of origination and extinction. Second-for-third metrics require occurrences, and use a
sliding-window approach that allows the examination of sampling in small time windows[9,22]. Second-
for-third metrics do not suffer from the biases that age-range metrics such as Foote's per-capita
method suffer from[21,22]. Unless stated otherwise, rates are expressed as the mean genus extinction
and origination per-interval rates of an order and are not normalised for the interval duration (See
STAR Methods). Rates are correlated to the duration of their respective taxonomic order, which is
the time between the first and last appearance.

We find that when per-capita metrics are used, we indeed recover strong negative rate-scaling
between log-transformed estimates of diversification rates and order duration (Figures 1A-1B). These
relationships (origination: β = −0.52; R^2= 0.27; P < 0.001, extinction: β = −0.52; R^2= 0.27; P < 0.001)
are stronger than those that had been reported previously in fossil data (e.g. origination: β = −0.23;
R^2 = 0.15; P < 0.001, and extinction: β = −0.25; R^2 = 0.13; P < 0.01)[13], and are of comparable strength
to those reported for molecular data (e.g. speciation: β = -0.54; R^2 = 0.34 and extinction: β = -0.55; R^2
= 0.16)[13]. However, when we control for incomplete and heterogenous sampling by using second-for-
third metrics, we find only a very weak relationship for origination (β = −0.16; R^2= 0.02; P = 0.014)
and no significant relationship for extinction rate (β = −0.11; R^2= 0.01; P = 0.077; Figures 1C-1D). This
suggests that unequal sampling through time and between clades causes the previously-observed
ARS in the fossil record.

Figure 1. Empirical relationship between log-transformed diversification rates and clade durations.
A and C show mean log-transformed genus-level origination rates (blue), and B and D show mean extinction (red) rates plotted against log
clade duration for 245 marine orders sampled across the Phanerozoic. Rates in A and B are calculated via Foote's per-capita metric, and C
and D are calculated by Alroy's second-for-third metric. Rates are shown as the mean number of events per-interval, per-lineage and not
normalised for interval length. Coloured lines represent ordinary least squares regression fits between log-rate and log-duration, and
shaded areas show 95% confidence intervals. β corresponds to the standardised regression coefficient that is equal to the slope of the
relationship between rate and duration when their variance has been standardised to 1. Results are shown for occurrences binned in
geological stages.

The occurrences that are used for the focal analysis are binned in time intervals based on
geological stages. These stages vary in duration, and if a single origination and extinction pulse were
to happen within a short interval, this might result in ARS. Equally, ARS in molecular phylogenetic
studies tends to be strongest over the shortest time intervals (<10 Myr)[13,23]. To counteract this, we
created composite time-bins with approximately equal duration of ~11 Myr by lumping the same
stage-level intervals[24,25]. Even when using these equal time bins, we still find strong ARS for per-
capita rates (Table S1), although both the variance explained, and the strength of ARS are decreased
substantially for extinction ($\beta = -0.39$; $R^2 = 0.15$; P < 0.001). That a strong relationship remains is not
surprising, because this again shows the strong role of sampling bias. When second-for-third rates
are used, ARS is again reduced. However, even under second-for-third rates, both origination ($\beta =$
-0.18; $R^2 = 0.03$; P = 0.007) and extinction ($\beta = -0.20$; $R^2 = 0.04$; P = 0.003) now show significant
trends, albeit very weak. This is consistent with the idea that when occurrences are binned in longer
time intervals (effectively increasing the minimum age of orders as intervals are aggregated
together), additional survivor and selection bias is applied to the clades in our dataset (see below).
This highlights that heterogenous and incomplete sampling causing ARS in the fossil record arises
independent of timescale and shows an additional role for selection bias.

Sampling biases are thus the likely cause of ARS in fossil time series. Drawing a parallel to
phylogenetic studies, Louca et al.[23] (also see[26]) argued that there are two distinct mechanisms that

lead to a biased sample of species-rich phylogenies that can cause ARS. Mechanism 1 considers that
even if clades share the same intrinsic speciation and extinction rates, clades can show a wide
distribution of clade sizes because of the stochastic variation in rates through time. It is likely that
clades that radiated rapidly at their inception will have survived – the "push of the past"[26,27]. This
survival bias effectively truncates the distribution of clade sizes by excluding species-poor and extinct
clades and can bias towards estimating higher rates, especially in young clades. Mechanism 2
suggests that diversification rates can be truly heterogenous among clades. If high rates are indeed
accurately estimated, the same survival bias may, however, still apply. Additionally, selection bias
may contribute to ARS under both mechanisms by only selecting clades beyond a certain species
diversity-threshold, and taxonomic practices may further exaggerate this if clades are only described
after reaching certain diversity. Diversity thresholds are difficult to avoid in phylogenetic studies,
because extinct clades are not considered, and estimating rates for small clades can be unreliable.
This again disproportionally affects younger clades, because they likely contain fewer species.
Comparable selection and survivor biases could equally affect ARS in the fossil record.

One advantage of using fossil occurrences is that extinct clades are incorporated, which avoids
truncating the possible diversification trajectories a clade can follow. There is also less taxonomic or
researcher-focused bias in selecting clades to study that are above a certain diversity threshold,
because occurrences are described per palaeontological collection and not by specifically sequencing
a particular clade or species. However, this does not mean that estimating rates for small clades
becomes significantly easier, and the clades that are fossilised might equally represent a biased
sample of all clades that ever lived. Therefore, arbitrary diversity-thresholds are still imposed for
methodological reasons, and biased preservation of the most diverse clades cannot be directly ruled
out (see sections below).

We investigate how imposing diversity thresholds might influence ARS by only using orders that
have accumulated ≥10 genera, in line with previous work that has recovered ARS from fossil data[13].
This analysis suggests that there is a minimal impact on observed ARS when we look at occurrences
binned at the stage-level (Table S1). However, when occurrences are binned in equal-time bins, we
do observe a modest but clear decrease in slope and increase in R^2 (Table S1). This is likely due to the
fact that very few young clades with low richness and low rates survive long enough to be sampled,
because the minimum duration has increased, and few remain after filtering. Consequently, there is
a minimal but observable effect of duration- and diversity-threshold selection bias on ARS.

Sampling bias in the fossil record cannot be entirely attributed to diversity-threshold sampling
bias. For species to show up in the fossil record they will need to fossilise, be preserved through time,
and be sampled in the present. The simplest way of thinking about sampling a species is by assigning
it a probability that it will be sampled per time interval – ρ, similar to speciation and extinction
rates[28,29]. Under this assumption, all else being equal, the duration of a species correlates positively
with the chance that it will be sampled least once. For higher-level taxa such as genera and orders,
this also means that the more species they contain, the higher their chance of preservation. We
hypothesise that especially for short duration clades there will be limited opportunities (i.e. in
multiple time bins) to be sampled, which might inflate the number of false originations and
extinctions.

We investigate this hypothesis with a simple heuristic simulation model of species diversification
and sampling. The simulation model is based on a birth-death model with speciation and extinction[22].
Simulations are started with N species (N = 1,000 in Figure 2), and speciation and extinction occur in
pulses at the boundaries between intervals (E = 0.3 in Figure 2). We further assume that the number
of species is constant through time[9,30], so that every species that is lost is immediately replaced by a

new species. The sampling probability, *s*, of species within and between intervals is uniform. The
point of this simulation is not to be exhaustive, nor to pinpoint the precise sampling artefact causing
ARS, but simply to demonstrate whether incomplete sampling could conceivably cause ARS.
Moreover, more complex models might induce their own artefacts leading to ARS by, for instance,
inducing survivorship and selection biases[26,31], which we seek to avoid here.

Figure 2. Diversification rate scaling with clade duration for simulations.
Average extinction (red) and origination (blue) rates are plotted against their clade duration for 1000 repeats. Column 1 shows the
relationships between rates and clade durations when sampling is complete, whereas column 2 shows a scenario of moderate sampling
where each species has 50% chance of being preserved and sampled within a bin. Rows represent the rate metrics used for the empirical
analysis. Row 1 and 3 use Foote's per-capita metrics, and row 2 and 4 use Alroy's second-for-third metrics that correct for sampling bias.
Coloured lines represent non-linear least squares regression fits according to the function $y = ax^b$. 95% confidence intervals are not
shown as they overlap almost completely with the coloured lines. P-values are obtained by comparing the non-linear regression to an
intercept only model. Simulations are shown for a species pool of N = 1000 and an extinction probability of E = 0.3 per interval.

We find that under these simple assumptions, ARS is absent when all species are fully sampled (s
= 1), and diversification appears constant among clades irrespective of clade duration (Figure 2).
However, when we impose lower sampling probabilities (s = 0.5), ARS does manifest, but only under
the per-capita rates. This result is consistent under a wide range of parameter values (Figure S1). This
result confirms that error in the rates for short-duration clades, caused by incomplete sampling and
resulting sampling artefacts, could explain the presence of ARS in the fossil record.

Apart from data driven artefacts, statistical artefacts could cause ARS to arise in fossil data. As
previously mentioned, normalising rates to per-million-year units results in ARS when rates
measured over geological stages of varying durations are compared[1], and, more generally, studies
have attributed ARS to plotting a ratio (events Myr^{-1}) against its denominator $(Myr)^{8,32}$. Our results
(Figure 1) demonstrate that ARS arises in per-capita rates even if they are not normalised for interval
duration. However, it is possible that normalisation could either increase or weaken the strength of
ARS. We find a minimal effect of normalisation on both per-capita and second-for-third rates (Table
S1). These results should not be overinterpreted, because we use rates averaged over time intervals
of irregular duration. The interpretation that steeper slopes equal greater independence between
events and duration, or that this signals the absence of spurious self-correlation consequentially both
hold little weight.

Our results emphasise the importance of controlling for heterogeneous sampling when
calculating rates of origination and extinction from fossil data. This is not a new suggestion, and is
true for many other metrics (e.g. species diversity[33]), but we specifically show that failing to control
for incomplete sampling may induce ARS. With the increased availability of taxonomic occurrence
records, and the development of sophisticated methods for estimating rates (e.g. PyRate[34], fossilised
birth-death range process[35], second-for-third rates[21]), there is little excuse for using tools that have
been shown to be erroneous[11]. Nevertheless, even when a method is chosen that can incorporate
uneven sampling, care should be taken to avoid introducing artefacts relating to plotting rates
against time[5].

Various processes have been argued to cause higher rates at the start of clade radiations, with
diversification slowing down with time[36]. It might therefore be surprising that controlling for uneven
sampling through time significantly reduces ARS in fossil time series. Survivorship bias can cause such
slowdowns and is inherently present in the fossil record[26,31], making it an important null model for
diversification patterns. Assuming constant rates of diversification, young clades will have a higher
risk of extinction compared to older clades as they statistically will harbour more species.
Consequentially, observable clades are likely those that diversified rapidly at their inception – the
"push of the past"[26,27]. If preservation is a per-species probability[28], clades that have experienced
high origination rates are more likely to have entered the fossil record[31]. This leads to ARS, since the
only short-duration clades that can be observed are those that radiated rapidly, whereas older clades
regress towards their mean diversification rates. We find some support for this, because ARS
increases when equal time bins and diversity thresholds are used. However, this effect is minor and
when we correct for the heterogeneity, and incompleteness of sampling through time, ARS all but
disappears. The absence of ARS should be further explored, and we hypothesise that a difference in
taxonomic level causes this. Specifically, genera are the most common unit for fossil diversification
analyses, but in contrast to species, do not go extinct until all their constituent species have gone
extinct (see STAR methods).

In conclusion, we do not find strong support for a relationship between the duration of a clade
and the rates of origination and extinction that it has experienced. Previous studies only found such a
relationship because they used metrics that are susceptible to various sampling biases inherent to

the fossil record – akin to the idea that measuring rates over shorter time scales is more error
prone[37]. Our findings show that biases inherent to the fossil record may cause rate-scaling,
suggesting that artefactual self-correlation[4,8,32] is not always applicable nor that a general cause
necessarily explains rate-scaling. However, the interaction between sampling, survival, and
taxonomic biases provide motivation for further research to uncover the reasons underlying the
different signals present in molecular[23] and fossil data.

Methods
222

Fossil occurrence data assembly
223

We retrieved fossil occurrence data from the Paleobiology Database using the URL API in identical
fashion to previous work[24,38]. We reiterate what has been stated there, but we performed a new
download of the data on the 8th of May 2024. Our analyses focus on marine Animalia. For our API
calls we include all fields for marine Animalia excluding Chordata, Chordata excluding Tetrapoda, and
Tetrapoda. Calls were restricted to marine environments only by the command "envtype=marine".
Separately, lists of occurrences were downloaded for Anthozoa, Brachiopoda, Bryozoa, Conodonta,
Cephalopoda, Crinoidea, Decapoda, Echinoidea, Gastropoda, Graptolithina, marine Bivalvia, and
Trilobita to define taxon sets/clades, and orders were defined by the data field "order".

Diversification rate analyses were conducted at genus level, so our empirical rates pertain to
the origination and extinction of genera. Occurrences were filtered to only those that were assigned
with certainty to at least genus-level using "idqual=genus_certain" in our calls. We only took into
account the latest identification ("idtype=latest") of the occurrence, only valid taxa
("taxon_status=valid"), and removed form taxa (e.g., assigned based on morphological affinity but
not phylogeny) or ichnotaxa (e.g., trace fossils) via "pres=regular". We further filtered the
occurrences to those that had been assigned to a specific order.

Occurrence were binned via two schemes: (i) 100 geological stages, and (ii) 49 composite
equal time bins. The equal time bins each spanning approximately ~11 myr were created by binning
geological stages[39]. We assigned the occurrences to bins based on a majority rule. If a time bin
contained >50% of the geological time range associated with the occurrence, it was assigned to that
bin[39]. Our downstream diversification rate calculations depend on clades ranging over at least 3 time
bins. Any order with fewer bins was removed from the analysis. After applying these sifting criteria,
the two final datasets based on geological and composite time bins consisted of respectively 610,514
and 646,204 occurrences, 408 and 368 orders, a clade duration range of 5.45-520.99 and 31.60-
538.79 Myr and face-value diversity counts ranging from 1 to 1267 and 1 to 1299 genera per order.

Estimating diversification rates
248

The quantification of changing rates of origination and extinction has been one of the central goals of
palaeobiology. A substantial number of metrics have been developed over time, and have been
scrutinised equally as often[1,22,33]. Alroy[15] and Foote[16,17] laid the foundation to modern rate metrics
that quantify how rates change between subsequent time intervals, i.e. geological stages and equal
time bins, whereas later iterations and metrics have built on this framework[9,22,30,40]. Here, we explore
the effect of Foote's[17] per-capita rate metric and Alroy's[40] second-for-third metric on the estimation
of diversification rates and the impact on rate-age-relationships.

Foote's metrics rely on the age ranges of organisms, i.e. their first and last appearance in the
fossil record. Four types of ranges are classified: taxa present both before and after a focal interval,
crossing its base and top (N_{bt}); taxa present before and within the interval but not after, crossing only

its base (N_{bL}); taxa present in and after the interval, crossing only its top; and taxa found only within
the interval (N_{FL}). Summed, these counts are the range-through diversity (N_{bt}). The rates we use here
are Foote's per-capita rates where origination is $\lambda = -ln\left(\frac{N_{bt}}{N_{Ft} + N_{bt}}\right)$, and extinction is $\mu =$
$-ln\left(\frac{N_{bt}}{N_{bL} + N_{bt}}\right)$. Note that Foote's metrics were originally normalised by the interval duration over
which they were measured[17], but this was later reconsidered as discussed below[12]. These range
based metrics are still actively used[41], and are perhaps the best approach to study range based
data[33]. However, range-based metrics have been argued to suffer from edge, Pull of the Recent, and
Signor-Lipps effects, and taxonomic biases. In particular, this means that extinction rates are
increased in intervals preceding mass extinctions (effectively smeared out), rates gradually decline as
the recent is approached, and the importance of rare taxa is overweighted when sampling is good
(i.e. Pull of the Recent) or underweighted when extinction is fast (i.e. Signor-Lipps)[22,40].

Fossil occurrence data has become increasingly available, promoting a way to move beyond
using range-based metrics and focussing on patterns of taxonomic sampling throughout. Alroy's
second-for-third metric has been argued to not suffer from any of the above biases, and relies on a
sliding-window approach, in contrast to full ranges[40]. Our focal time interval is i_0, and we assume a
cohort of taxa that have only been sampled since i_{-1} up to i_2 for extinction and i_{-2} to i_1 for origination.
The following counts are of importance for extinction (origination in brackets if distinct): s_1 = taxa
sampled in i_{-1} and i_0 (i_0 and i_1) but no later (earlier), s_2 = taxa sampled in i_{-1} and i_1 (i_{-1} and i_1) but not in
i_0 or i_2 (i_{-2} and i_0), s_3 = those sampled in i_{-1} and i_2 (i_{-2} and i_1) but not in i_0 or i_1 (i_{-1} and i_0), t_2 = those
sampled in i_{-1} and i_0 (i_0 and i_1) regardless of being sampled later (earlier), and p = i_{-1} and i_1 but not in
i_0. The formula for extinction and origination proportion (not rate) is then calculated as $O = E =$
$(s_1 - s_x)/(t_2 + p)$. If sampling is uniform and given that unextinction cannot occur the counts
should have the expected order $s_1 \geq s_2 \geq s_3$, and s_x would be s_3 (the Gap-Filler equation[22]). However,
in practice they do not and might even show the reverse pattern $s_1 < s_2 < s_3$. To address this Alroy
suggested algorithmically replacing s_x with the second lowest of the counts, e.g. $s_x = s_1$ if $s_2 > s_1 > s_3$.
Conventional rates (e.g. decay coefficients) are then obtained by log-transformation where
$\mu = \log\left(\frac{1}{1-E}\right)$, and $\lambda = \log\left(\frac{1}{1-O}\right)$. Second-for-third rates are obviously heuristic because of the
algorithmic approach, but it incorporates as much information as possible to correct for sample size.

We focus on genus origination and extinction per-interval rates for both Foote's and Alroy's
metrics, and do not normalise for interval duration. That is, we focus on the rate at which species
originate and go extinct per taxon per interval, and do not normalise our rates to be expressed per
million years as was originally done[17]. Raup and Sepkoski, although using percentage extinction and
not true rates, already discussed preferring per-interval metrics in their seminal paper[2]. They
observed that the number of raw extinction counts did not correlate with interval duration as they
are likely clustered in time[9,33]. Foote further explored the consequences of normalisation and found
that if intervals were to experience a single peak in extinction risk, this would dominate the total
extinction profile of short intervals more than for long intervals, effectively causing a negative
correlation between interval duration and extinction rate[1]. We repeat our analysis for time-
normalised rates to understand if negative rate scaling between bins has an effect on average rate
comparisons between clades.

We calculate both Foote's per capita rates and second-for-third rates using the R package
divDyn[42]. Origination and extinction rates were calculated for all orders for both geological stages
and equal time bins, and then averaged per order. In some instances, an average rate of zero is
estimated. Rate estimates of zero likely are caused by sampling error or model misspecification and
make little sense empirically[33] (but see ref[43]). For example, an order must have had a positive

origination rate if it contains more than one species. In our focal analysis, orders with average rate
estimates of zero are removed (i.e. the number of orders is reduced from 408 to 245 for stage-level
bins and from 368 to 220 for equal-time bins). This might bias the analysis towards observing greater
rate-scaling. Regardless, average rates are subsequently correlated across all orders with their
duration using (i) non-linear least squares, and (ii) linear regression by log-transforming both rate and
duration, which would have removed zero estimates.

A simulation model

We investigate if incomplete sampling through time can cause ARS with a simulation model of
species diversification and sampling. The simulation model is based on a prior model used in the
testing of rate metrics[22], and can be abstracted to modelling temporal change in species identities
(i.e. faunal turnover) in a species pool. The species pool has constant species diversity, N, through
time ($N = 10^2$, 10^3, 10^4), which can be justified by the strong correlation between per-interval
extinction rate followed up by high origination in the next interval, across the Phanerozoic marine
fossil record[9,30]. Because we are not interested in measuring diversity, a variable trend might have
sufficed as well. At the end of each interval, species have a probability, E, to go extinct ($E = 0.15$, 0.3,
0.6). Speciation or origination is modelled as the replacement of the extinct species immediately
upon extinction by a new species. Effectively, diversification thus occurs in pulses between intervals.
We simulate 10^3 clades per parameter combination to calculate ARS, with clades experiencing faunal
turnover for a random number of intervals drawn from a uniform distribution ranging from 3 (i.e. the
minimum number of bins required to calculate per-capita rates) to 54 (i.e. roughly the reflecting the
length of the Phanerozoic if each interval is 10 Myr) intervals. The realised time series of faunal
turnover, each seen as a unique clade, are then degraded based on a per-species sampling
probability per interval, s. We model sampling intensity from poor to good ($s = 0.25$, 0.5, 0.75). The
complete and degraded time series are then used to calculate both Foote's per-capita and Alroy's
second-for-third rates and plotted against the number of time intervals to repeat the empirical
analysis (Figures 3, S5, and S6).

Our simulations assume that sampling is uniform through time and across clades. In practice,
sampling within time bins will vary because of variation in preservation (e.g. substrate), sampling
effort, and fossilisation potential of clades (e.g. soft or hard-bodied). Taxonomic biases might further
contribute to error in the estimation of rates by over or under splitting taxa. This might in itself result
from incomplete sampling through time, because infrequently sampled but temporally long-ranged
taxa might be under split[22]. Equally, taxonomic ranks are inherently arbitrary concepts. Especially
when it comes to orders and other higher ranks, it might be difficult to accurately delineate when
they arise, and consequently higher diversity orders might be recognised before low diversity orders.
Therefore, our simulations are by no means exhaustive in terms of the possible ways that sampling
could influence the estimation of diversification rates.

Genus versus species-level diversification

In the main text we discuss how survival bias might cause a "push of the past" under which
clades that radiate rapidly early in their duration are more likely to be observed in the fossil record.
This could lead to a strong negative relationship between diversification rate and clade duration even
when controlled for heterogenous and incomplete sampling, but we do not find this. An explanation
for this absence could be our empirical focus on fossil genera instead of species. This difference in
taxonomic level, often required in fossil analyses, becomes apparent when the survival of a species
(i.e., species duration), is compared to that of a genus. If a species' extinction rate is constant, the
probability of species survival decreases linearly through time[44]. In contrast, if we look at genera,

which are each made up of one or more species, a genus will only go extinct after its last species disappears. With time, as a genus accumulates more species, the survivorship of the genus therefore increases[45–47]. This could be modelled by assuming a small probability of genus formation at each speciation event, and that each genus starts as a single species from which it diversifies. Consequentially, the rate of genus origination would correlate positively with the rate of species formation, albeit stochastic. If a "push of the past" effect occurred on the species-level, a likely identical preservation bias for species and genera would also operate. However, because our results do not reveal ARS, the formation of genera could be largely independent of the formation and extinction of species. We speculate that this might consequently be caused by taxonomic or biological reasons.

From a taxonomic perspective, evolutionary lineages in the fossil record might not be recognised as independent genera until they have shown sufficient morphological differentiation. This might cause different rate-scaling patterns at the level of species (molecular data) and genera (fossil data) to arise, especially if we consider the prevalence of cryptic species in molecular datasets that show little morphological divergence but are reproductively isolated[48]. If most of these cryptic species are unlikely to persist over evolutionary time, which could be the case in some contemporary clades such as sticklebacks[49], we might expect the highest rates in young clades with a substantial proportion of cryptic species. If genera or species are delineated by morphology in the fossil record, such cryptic diversity would only be detectable in molecular datasets.

From a biological perspective, the assumption that a genus has a survival probability, and thus preservation probability, that is a function of the number of species contained could be relaxed. For instance, even if a genus is species-rich, these species might all be extinction-prone because of a property such a specialisation or range size, whereas a genus with a single species might consist of a generalist that is widespread. There are thus two ways in which clades with few genera might have a high survival probability through time. Either each of these genera are widespread, or if the number of species matters, few genera might contain many species purely by chance or taxonomic practice. A push of the past at the level of species may consequentially not be experienced at the level of the genus, either because of stochasticity, the extinction vulnerability of the species within the genus, or by taxonomic relationships not being observable in the fossil record. However, the interaction between these processes is complex and future studies might benefit from extended null models to analyse these patterns.

Quantification and statistical analysis

All statistical analyses are described above (Method details). Statistical analyses and simulations were conducted in R 4.3.1[50].

Acknowledgements

We thank Alex Pigot and David Murrell for discussion and encouragement. Three anonymous reviewers greatly improved the quality of the paper. The Paleobiology Database enabled this work, and contributors are wholeheartedly thanked for their effort. B.R.R. and R.A.C. are funded through the Royal Society by URF\R1\211571 awarded to R.A.C. This is Paleobiology Database publication XXX.

Author contributions

B.R.R. conceptualised the study. R.A.C. curated the data. B.R.R. designed, programmed, and conducted the analysis. B.R.R. wrote the initial draft, which all authors edited and approved.

Declaration of interests

The authors declare no competing interests

References

1. Foote, M. (1994). Temporal variation in extinction risk and temporal scaling of extinction metrics. Paleobiology *20*, 424–444. https://doi.org/10.1017/S0094837300012914.

2. Raup, D.M., and Sepkoski, J.J. (1984). Periodicity of extinctions in the geologic past. Proc. Natl. Acad. Sci. *81*, 801–805. https://doi.org/10.1073/pnas.81.3.801.

3. Jackson, D.A., and Somers, K.M. (1991). The spectre of "spurious" correlations. Oecologia *86*, 147–151. https://doi.org/10.1007/BF00317404.

4. Sadler, P.M. (1981). Sediment Accumulation Rates and the Completeness of Stratigraphic Sections. J. Geol. *89*, 569–584. https://doi.org/10.1086/628623.

5. Pearson, K. Mathematical contributions to the theory of evolution.—On a form of spurious correlation which may arise when indices are used in the measurement of organs. Proc. R. Soc. Lond. *60*, 489–498.

6. Chayes, F. (1949). On Ratio Correlation in Petrography. J. Geol. *57*, 239–254. https://doi.org/10.1086/625606.

7. Gingerich, P. (1983). Rates of Evolution: Effects of Time and Temporal Scaling. Science *222*, 159–161. https://doi.org/10.1126/ science.222.4620.159.

8. De Lisle, S., and Svensson, E.I. (2023). Evolutionary change is remarkably time-independent across scales. Preprint, https://doi.org/10.1101/2023.12.02.569704 https://doi.org/10.1101/2023.12.02.569704.

9. Alroy, J. (2008). Dynamics of origination and extinction in the marine fossil record. Proc. Natl. Acad. Sci. *105*, 11536–11542. https://doi.org/10.1073/pnas.0802597105.

10. Foote, M. (2005). Pulsed origination and extinction in the marine realm. Paleobiology *31*, 6–20.

11. Foote, M. (2007). Extinction and Quiescence in Marine Animal Genera. Paleobiology *33*, 261–272.

12. Foote, M. (2003). Origination and Extinction through the Phanerozoic: A New Approach. J. Geol. *111*, 125–148. https://doi.org/10.1086/345841.

13. Henao Diaz, L.F., Harmon, L.J., Sugawara, M.T.C., Miller, E.T., and Pennell, M.W. (2019). Macroevolutionary diversification rates show time dependency. Proc. Natl. Acad. Sci. *116*, 7403–7408. https://doi.org/10.1073/pnas.1818058116.

14. Sepkoski, J.J. (1978). A kinetic model of Phanerozoic taxonomic diversity I. Analysis of marine orders. Paleobiology *4*, 223–251. https://doi.org/10.1017/S0094837300005972.

15. Alroy, J. (1996). Constant extinction, constrained diversification, and uncoordinated stasis in North American Mammals. Palaeogeogr. Palaeoclimatol. Palaeoecol. *127*, 285–311. https://doi.org/10.1016/S0031-0182(96)00100-9.

16. Foote, M. (1999). Morphological Diversity in the Evolutionary Radiation of Paleozoic and Post-Paleozoic Crinoids. Paleobiology *25*, 1–115.

17. Foote, M. (2000). Origination and extinction components of taxonomic diversity: general problems. Paleobiology *26*, 74–102.

18. Ho, S.Y.W., Lanfear, R., Bromham, L., Phillips, M.J., Soubrier, J., Rodrigo, A.G., and Cooper, A. (2011). Time-dependent rates of molecular evolution. Mol. Ecol. *20*, 3087–3101. https://doi.org/10.1111/j.1365-294X.2011.05178.x.

19. Harmon, L.J., Pennell, M.W., Henao-Diaz, L.F., Rolland, J., Sipley, B.N., and Uyeda, J.C. (2021). Causes and Consequences of Apparent Timescaling Across All Estimated Evolutionary Rates. Annu. Rev. Ecol. Evol. Syst. *52*, 587–609. https://doi.org/10.1146/annurev-ecolsys-011921-023644.

432   20.   Rolland, J., Henao-Diaz, L.F., Doebeli, M., Germain, R., Harmon, L.J., Knowles, L.L., Liow, L.H., Mank, J.E., Machac, A.,
433   Otto, S.P., et al. (2023). Conceptual and empirical bridges between micro- and macroevolution. Nat. Ecol. Evol. *7*,
434   1181–1193. https://doi.org/10.1038/s41559-023-02116-7.

435   21.   Alroy, J. (2015). A more precise speciation and extinction rate estimator. Paleobiology *41*, 633–639.
436   https://doi.org/10.1017/pab.2015.26.

437   22.   Alroy, J. (2014). Accurate and precise estimates of origination and extinction rates. Paleobiology *40*, 374–397.
438   https://doi.org/10.1666/13036.

439   23.   Louca, S., Henao-Diaz, L.F., and Pennell, M. (2022). The scaling of diversification rates with age is likely explained by
440   sampling bias. Evolution *76*, 1625–1637. https://doi.org/10.1111/evo.14515.

441   24.   Close, R.A., Benson, R.B.J., Saupe, E.E., Clapham, M.E., and Butler, R.J. (2020). The spatial structure of Phanerozoic
442   marine animal diversity. Science *368*, 420–424. https://doi.org/10.1126/science.aay8309.

443   25.   Alroy, J. (2010). The Shifting Balance of Diversity Among Major Marine Animal Groups. Science *329*, 1191–1194.
444   https://doi.org/10.1126/science.1189910.

445   26.   Budd, G.E., and Mann, R.P. (2018). History is written by the victors: The effect of the push of the past on the fossil
446   record. Evolution *72*, 2276–2291. https://doi.org/10.1111/evo.13593.

447   27.   Nee, S., Holmes, E.C., May, R.M., and Harvey, P.H. (1994). Extinction rates can be estimated from molecular
448   phylogenies. Philos. Trans. R. Soc. Lond. B. Biol. Sci. *344*, 77–82. https://doi.org/10.1098/rstb.1994.0054.

449   28.   Foote, M., and Raup, D.M. (1996). Fossil preservation and the stratigraphic ranges of taxa. Paleobiology *22*, 121–140.
450   https://doi.org/10.1017/S0094837300016134.

451   29.   Foote, M. (1997). Estimating taxonomic durations and preservation probability. Paleobiology *23*, 278–300.
452   https://doi.org/10.1017/S0094837300019692.

453   30.   Foote, M. (2023). Diversity-Dependent Diversification in the History of Marine Animals. Am. Nat. *201*, 680–693.
454   https://doi.org/10.1086/723626.

455   31.   Budd, G.E., and Mann, R.P. (2020). The dynamics of stem and crown groups. Sci. Adv. *6*.

456   32.   Sheets, H.D., and Mitchell, C.E. (2001). Uncorrelated change produces the apparent dependence of evolutionary rate
457   on interval. Paleobiology *27*, 429–445. https://doi.org/10.1666/0094-8373(2001)027<0429:UCPTAD>2.0.CO;2.

458   33.   Alroy, J. (2010). Fair Sampling of Taxonomic Richness and Unbiased Estimation of Origination and Extinction Rates.
459   Paleontol. Soc. Pap. *16*, 55–80. https://doi.org/10.1017/s1089332600001819.

460   34.   Silvestro, D., Salamin, N., Antonelli, A., and Meyer, X. (2019). Improved estimation of macroevolutionary rates from
461   fossil data using a Bayesian framework. Paleobiology *45*, 546–570. https://doi.org/10.1017/pab.2019.23.

462   35.   Stadler, T., Gavryushkina, A., Warnock, R.C.M., Drummond, A.J., and Heath, T.A. (2018). The fossilized birth-death
463   model for the analysis of stratigraphic range data under different speciation modes. J. Theor. Biol. *447*, 41–55.
464   https://doi.org/10.1016/j.jtbi.2018.03.005.

465   36.   Moen, D., and Morlon, H. (2014). Why does diversification slow down? Trends Ecol. Evol. *29*, 190–197.
466   https://doi.org/10.1016/j.tree.2014.01.010.

467   37.   O'Meara, B.C., and Beaulieu, J.M. (2024). Noise leads to the perceived increase in evolutionary rates over short time
468   scales. PLOS Comput. Biol. *20*, e1012458. https://doi.org/10.1371/journal.pcbi.1012458.

469   38.   Peters, S.E., and McClennen, M. (2016). The Paleobiology Database application programming interface. Paleobiology
470   *42*, 1–7. https://doi.org/10.1017/pab.2015.39.

471   39.   Close, R.A., Benson, R.B.J., Alroy, J., Carrano, M.T., Cleary, T.J., Dunne, E.M., Mannion, P.D., Uhen, M.D., and Butler,
472   R.J. (2020). The apparent exponential radiation of Phanerozoic land vertebrates is an artefact of spatial sampling
473   biases. Proc. R. Soc. B Biol. Sci. *287*. https://doi.org/10.1098/rspb.2020.0372.

40. Alroy, J. (2015). A more precise speciation and extinction rate estimator. Paleobiology *41*, 633–639.
https://doi.org/10.1017/pab.2015.26.

41. Foote, M., Edie, S.M., and Jablonski, D. (2024). Ecological structure of diversity-dependent diversification in
Phanerozoic marine bivalves. Biol. Lett. *20*, 20230475. https://doi.org/10.1098/rsbl.2023.0475.

42. Kocsis, Á.T., Reddin, C.J., Alroy, J., and Kiessling, W. (2019). The r package divDyn for quantifying diversity dynamics
using fossil sampling data. Methods Ecol. Evol. *10*, 735–743. https://doi.org/10.1111/2041-210X.13161.

43. Foote, M. (2007). Extinction and Quiescence in Marine Animal Genera. Paleobiology *33*, 261–272.

44. Van Valen, L. (1973). A new evolutionary law. Evol. Theory *1*, 1–30.

45. Raup, D.M. (1975). Taxonomic survivorship curves and Van Valen's Law. Paleobiology *1*, 82–96.
https://doi.org/10.1017/S0094837300002220.

46. Raup, D.M. (1978). Cohort analysis of generic survivorship. Paleobiology *4*, 1–15.
https://doi.org/10.1017/S0094837300005649.

47. Raup, D.M. (1991). A kill curve for Phanerozoic marine species. Paleobiology *17*, 37–48.
https://doi.org/10.1017/S0094837300010332.

48. Struck, T.H., Feder, J.L., Bendiksby, M., Birkeland, S., Cerca, J., Gusarov, V.I., Kistenich, S., Larsson, K.-H., Liow, L.H.,
Nowak, M.D., et al. (2018). Finding Evolutionary Processes Hidden in Cryptic Species. Trends Ecol. Evol. *33*, 153–163.
https://doi.org/10.1016/j.tree.2017.11.007.

49. Rabosky, D.L. (2016). Reproductive isolation and the causes of speciation rate variation in nature. Biol. J. Linn. Soc.
118, 13–25.

50. R Core team (2024). R: A language and environment for statistical computing. (R Foundation for Statistical Computing).

494