A critical affinity window for IgSF proteins DIP-α and Dpr10 is required for proper

motor neuron arborization

Davys H. Lopez[1,2], Kevin D. Rostam[2], Sumaira Zamurrad[2], Shuwa Xu[3], Richard S.

Mann[2,4,5]

[1]Department of Genetics and Development, Columbia University Medical Center,

[2]Zuckerman Mind Brain and Behavior Institute, Columbia University, [3]Division of Biology

and Biological Engineering, California Institute of Technology, [4]Department of Systems

Biology, Columbia University Medical Center, [5]Department of Biochemistry and

Molecular Biophysics, Columbia University.

Abstract

For flies to walk properly, motor neurons (MNs) from the ventral nerve cord (VNC) need to reach the correct muscle, and arborize appropriately during development. The canonical view of how this is achieved is that cell surface proteins are expressed pre- and post-synaptically that bind to each other like molecular "lock-and-keys" that guide neurons to their targets. The binding affinities of these molecules can vary by more than 100-fold. In the fly leg neuromuscular system, three MNs express *DIP-α* and their target muscles express its cognate partner, *dpr10*, both of which encode members of the Immunoglobulin superfamily (IgSF). Although, both of these molecules are necessary for the maintenance of MN-muscle contacts, the role that specific affinities play in this process has not been examined. Here we use knock-in mutations into *DIP-α* and *dpr10* that either decrease or increase the affinity between these two proteins. Compared to control animals, decreasing the affinity results in phenotypes similar to *DIP-α* or *dpr10 n*ull animals, where MN axons fail to maintain contacts with their muscle targets and retract their filopodia, resulting in stunted and/or branchless axons. We also find that the three *DIP-α*-expressing motor neurons behave differently to the loss of affinity. Surprisingly, if the affinity increases past a certain threshold, a similar branchless phenotype is observed in adult legs. Live imaging during pupal development shows that MN filopodia are unable to productively engage their muscle targets and behavioral assays suggest that these defects lead to locomotor deficits. These data suggest that CAM affinities are tuned to a specific range to achieve proper neuronal morphology.

Introduction

Animal movement requires developing motor neurons (MNs) to arborize and synapse onto the correct muscles. For axons to establish the precise set of synapses required for neural circuit formation, it is helpful to conceptualize this process into a series of broad steps: axon guidance or pathfinding, in which axons navigate close to their cellular targets[1–4], synaptic specificity, where short-range interactions establish synaptic choice[2,5,6], branching and arborization[7,8], and synaptic plasticity, which is an activity-dependent process that can modify preexisting synapses[9,10]. In all these steps, cell adhesion molecules (CAMs) have been implicated as playing critical roles[2,5]. For example, CAMs that bind hetero- and homophilically such as the *beat-la* and *sidestep* have been implemented during axon pathfinding[3,4,11,12], molecules such as the L1 protein subfamily are needed for synaptic specificity[5,13–15], Dscam1 and Protocadherins are needed for self-avoidance[16–18] and β1 integrins and contactins are used for synaptic plasticity[5,9,10,17,19–21]

In Drosophila, the DIPs (Dpr-interacting proteins) and Dprs (Defective proboscis response) are a subset of CAMs that are also members of the Ig domain Superfamily (IgSF) proteins. The "Dpr-ome" interaction network was initially discovered using an *in vitro* interactome assay that detected interactions between extracellular protein domains[22,23]. This interactome and its subsequent analysis revealed that there are 9 Dprs, each with two Ig domains, that bind selectively to DIPs, which have 21 members, each with three Ig domains[22,24,25]

DIPs and dprs are expressed by a subset of neurons, suggesting a role in synaptic specificity. For instance, photoreceptors R7 and R8 and lamina neurons L1-5

in the fly visual system express a subset of Dprs, with their axonal targets in the medulla

expressing matching DIPs [24]. In the larval nervous system Dprs are more broadly

expressed than DIPs in both MNs and SNs, and each neuron expresses a unique

combination of Dpr and DIP genes[26]. This is also the case in the adult leg

neuromuscular system where *dprs* are more widely expressed than the *DIPs*, in the

majority of adult leg MNs and leg sensory neurons[27]. Furthermore, in the mushroom

body, 16 of 21 *dpr* members were shown to be dynamically expressed in developing

γ-KCs. Together, this indicates that DIPs and Dprs expressed by different neurons are

likely to be involved in many stages of neural development[28].

The DIPs and Dprs have also been shown to play an important role in

establishing neuronal morphologies and constructing neural circuits. For example,

dpr11 and *DIP-γ* mutant animals have defects in presynaptic terminal development in

the larval neuromuscular system and in the visual system both are required for the

photoreceptor neuron, yR7, to target the correct M6 layer[25]. Furthermore, *DIP-γ* mutant

flies also have a reduction in Dm8 neurons compared to controls[25]. *DIP-ε* has also been

shown to have a critical role in establishing wildtype *fruP1* neuronal patterns, in both a

cell-autonomous and non-cell-autonomous manner[29]. And, Dpr12/DIP-δ transneuronal

interactions are required for γ-Kenyon cells to find their proper targets in the mushroom

body[28]. In the optic lobe, DIP-α heterophilic interactions with Dpr6 and 10 regulate

arborization within layers, synapse number, layer specificity, and cell survival of

amacrine-like Dm4 and Dm12 neurons[30].

In the larval motor system, *DIP-α* is expressed by only two motor neurons (MNs)

in each hemisegment, ISNb/d-1s and ISN-1s. In *DIP-α* mutants, one branch of the ISN-

1s MN fails to innervate muscle 4 (m4)[31]. In addition, in *DIP-α* mutants, the axons that

fail to innervate m4 now innervate muscle m1[31]. Interestingly, during development ISN-

1s still projects to m4 but fails to be stabilized. DIP-α cognate partners, Dpr6 and Dpr10,

are expressed in 1s and in muscles, and knockdown of *dpr10* in the muscle, or in *dpr10*

null animals, there is 100% loss of 1s boutons on m4 similar to the *DIP-α* null. *dpr6*

hemizygous mutants have no obvious defect in ISN-1s[31].

DIP-α also has a role in the adult leg motor system. Using live imaging, it has

been shown that the leg neuromuscular system develops through sequential rounds of

stereotyped MN branching and defasciculation followed by muscle targeting,

arborization, and synapse formation[27]. Using gene trap reporters into *DIP-α*[27,32], it was

shown that it is expressed by three MNs that target the long tendon muscle in the femur,

αFe-ltm; the long tendon muscle in the tibia, αTi-ltm; and the tarsal depressor muscle in

the tibia, αTi-tadm (Figure 1C). In *DIP-α* null mutants, the terminal axon branches of

αFe-ltm are absent in almost all samples, while those of the αTi-ltm are missing in

approximately 70% and those of the αTi-tadm are rarely missing. *dpr10* is expressed

broadly by all muscles and in *dpr10* null animals the three DIP-α-expressing MNs

display the same phenotype as *DIP-α* null animals[27]. To differentiate between whether

axons fail to send out filopodia or whether they send out filopodia but are not able to

arborize, live imaging was used to visualize developing *DIP-α* null axons[27]. These

videos showed that, compared to controls with an intact *DIP-α* locus, axons send out

filopodia to the correct muscle target but fail to arborize and eventually retract, resulting

in adults with no axon branches[27].

The binding affinities of all interacting DIP/Dpr pairs have been measured using

biophysical methods. K_Ds for homophilic and heterophilic binding for the DIP/Dpr pairs

vary by more than 100-fold, ranging from <2 to >200 µM[24,33]. Interestingly, DIP-α and

Dpr10, the binding pair examined here, have one of the highest affinity interactions in

the DIP-Dpr interactome, with a K_D of 1.4 µM[24,33]. Motivated by these observations, a

series of mutations were designed based on X-ray crystal structures that include the

DIP-α–Dpr10 binding interface that either decrease or increase binding affinity, and

were knocked into the *DIP-α* and *dpr10* loci[33]. In the optic lobe, it was shown that

reducing the affinity of either *DIP-α* and *dpr10* causes loss-of-function phenotypes such

as synaptic targeting and cell survival defects, and that the severity of these phenotypes

scales with the magnitude of affinity reduction[33]. Interestingly, although increasing

affinity did not alter targeting, it did affect cell survival in this system. These results

indicate that affinity are tuned to achieve proper cell survival and synaptic

connections[33]. In contrast to the visual system, no affect on leg MN survival has been

observed in *DIP-α* and *dpr10* mutants. This may be because in the visual system,

neurons are produced in excess and those that do not form functional synapses are

eliminated via apoptosis[34]. In contrast, the wiring of the leg motor neuron to muscle

innervation is hardwired and every individual leg motor has a unique target and

morphology[35,36].

Here, we take advantage of the precision of the leg motor neuromuscular system

to examine the role of affinity between DIP-α and Dpr10. We find that decreasing or

increasing affinity does not affect cell survival nor does it cause mistargeting, but rather

has a role in stabilizing MN axon arborization. Using a quantitative neuron tracing

analysis[37] we find that the three DIP-α-expressing MNs (αMNs) have different affinity

requirements. Furthermore, we find that increasing affinity does not result in more

branches but causes aberrant neuronal morphology. Furthermore, we show that

changing the affinity affects proper locomotion. Together, these results reveal that

affinities between interacting CAMs have been tuned to achieve proper neuronal

development.

Results

Leg MNs arborize through all of pupal development

We used time lapse imaging to obtain a better description for how αMNs

coordinate their branching as they contact nascent groups of muscle precursors.

Previously, we live imaged αMNs until about 45 hours After Pupal Formation (hrs

APF)[27]. However, to extend this analysis to axon morphology changes that occur

between mid-pupa and adult we improved our live imaging protocol, allowing us to

visualize development later in pupal stages. We used *DIP-α-T2A-Gal4>20XUAS-*

6XGFP to visualize the three αMNs together with *Mef2-QF>QUAS-10XmCherry* to

visualize muscles as they develop from ~20-108 hrs APF, which encompasses almost

all of pupal development (Figure 1A and Videos 1 and S1). Because we can only

visualize structures close to the coverslip, we are limited to visualizing MNs and

muscles in the tibia, which is close to the surface of the pupa.

As early as 20 hrs APF αTI-MNs have already targeted nascent clusters of

myoblasts within the tibia, as previously described[27]. Interestingly, by visualizing the

developing myoblasts with myr::tdTomato, we find that the myoblasts are organized into

interconnected clusters that will become future muscle groups i.e. the tarsal depressor

muscles (tadm) and the Tibia long tendon muscle (Ti-ltm) (Video 2 and Figure S4).

These clusters then split and become discrete muscle groups. Between 25-55 hrs APF,

the myoblasts will fuse and become a syncytial multifiber muscle (Figure 1B). By 20-60

hrs APF, the videos show single mobile myoblasts that end up joining and fusing with a

particular muscle fiber (Figure 1B). At ~50 hrs APF αTi-MNs have reached and

stabilized a nascent axon terminus with a limited number of branches on their targeted

muscle. However, from ~50 hrs APF and onward, this initial nascent axon terminus will

continue to give rise to new branches and elongate its existing branches until a few

hours before the animal ecloses (Figure 1B). During this later phase, the movement of

the αMN filopodia will slow and take on a bead-on-string morphology, indicating a

transition from a growth cone to synapse formation (Video 1). Based on these

observations, there are at least two different phases of axon morphogenesis, an initial

phase where the αTi-MNs choose and stabilize muscle fibers as they are developing,

and a second phase when αTi-MNs limit their mobility but continue to arborize while

forming synapses. This indicates that MN arborization occurs for much longer than

previously noted[27].

172   Reducing DIP-α affinity to Dpr10 impedes arborization

Reducing the affinity between DIP-α and Dpr10 causes cell death of Dm4 and

Dm12 neurons in the optic lobe and mistargeting of Dm12 from medulla layers M3 to

M8[33]. Using the same *DIP-α* and *dpr10* alleles used in that study we examined the final

extent of leg MN arborization. These alleles alter the affinity between DIP-α and Dpr10

in graded fashion over a wide range, do not affect binding specificity, and do not alter

homophilic binding between DIP-α and itself. We used the following *DIP-α* mutants:

K81Q (K_D = 31.8 µM, *DIP-α*$^{-20F}$; superscripts indicate the direction and fold change (F)

in affinity relative to wild type), K81Q G74S (K_D = 68.0 µM, *DIP-α*$^{-50F}$) (Figure 2A) and

dpr10 mutants: V144K (K_D = 11.3 µM, *dpr10*$^{-8F}$), Q138D (K_D = 27.6 µM, *dpr10*$^{-20F}$), and

G99D, Q142E, V144K (K_D = 50.0 µM, *dpr10*$^{-40F}$) (Figure 3A). Although *dpr10*, but not

dpr6, is necessary for the terminal branching of the αMNs[27], all *dpr10* mutant

chromosomes are also mutant for *dpr6*. The αMNs were visualized using a previously

characterized MiMIC line converted to T2A-Gal4, and repeated using a transcriptional

enhancer isolated from *DIP-α* called A8 fused to Gal4 (*A8-Gal4*)[27] (Figure S1). To

acquire quantitative information regarding branch number and branch length, we traced

and quantified both parameters using the semi-automatic tracing software on FIJI[37], and

compared control animals with wildtype alleles to animals carrying either one copy of

the mutant *DIP-α* allele and one copy of the *DIP-α-T2A-Gal4* (which is a strong

hypomorph[27]) or two copies of the *DIP-α* allele with *A8-Gal4* (Figure S1)[37]. We also

examined animals with null alleles as a negative control.

Compared to controls, adult animals with *DIP-α* decreased affinity alleles (*DIP-α* $^{-}$

^{20F} and *DIP-α* $^{-50F}$) show phenotypes similar to *DIP-α* null animals, where αMN axons fail

to arborize onto their muscle target and have no terminal branches (Figure 2B).

Unfortunately, using *DIP-α-T2A-Gal4,* we are unable to visualize the αFe-ltm, because

even in *DIP-α-T2A-Gal4/+* animals αFe-ltm fails to arborize ~50% of the time[27].

However, using *A8-Gal4* to directly compare all three αMNs we find that the three αMNs

have different expressivities in the absence of *DIP-α* or *dpr10*: αFe-ltm completely fails

to arborize, αTi-ltm terminal branches are reduced or lost in about ~70-80% of animals,

and αTi-tadm is rarely affected[27]. In addition to these observations with null alleles, we

also find there is a graded affinity requirement for each of the αMNs. For instance, we

find that for αFe-ltm there is less arborization for the *DIP-α*$^{-50F}$ allele compared to the

DIP-α^{-20F} allele, suggesting that the affinity required to stabilize the branch is within this

range (Figure S1). αTi-ltm also fails to arborize although it seems that the expressivity is

not as strong as in the αFe-ltm (Figure S1B and C). Finally, although the αTi-tadm rarely

shows a complete loss of arborization, both branch number and branch length are

reduced (Figure 2D and Figure S2B). Together, these data reveal that the three αMNs

all require different minimal affinities to properly arborize onto their target muscles.

Reducing Dpr10 affinity to DIP-α impedes arborization

Considering that we see *DIP-α*$^{-20F}$ flies lose the terminal branches of αFe-ltm, we

wondered whether we would see a similar loss with the *dpr10*$^{-8F}$ allele. As with the *DIP-*

α alleles*,* we examined the *dpr10* alleles using both *DIP-α-T2A-GAL4* and *A8-Gal4* as

readouts of αMN morphology. Using the *A8-Gal4* as a readout, we find that for *dpr10*$^{-8F}$,

αFe-ltm branches are not completely absent but there is a decrease in branch number

and branch length, while the -20F allele of *dpr10* results in the complete loss of

branches in this MN (Figure S1D and F). Furthermore, we find that for αTi-ltm the

complete absence to arborize occurs at a lower frequency compared to αFe-ltm for all

dpr10 alleles, indicating that the αFe-ltm has a higher affinity requirement than the αTi-

ltm (Figure S1). However, using *DIP-α-T2A-Gal4*, αTi-ltm fails to arborize in most *dpr10*$^{-}$

[8F] animals. This discrepancy may be explained because *DIP-α-T2A-Gal4* is a strong

hypomorph of *DIP-α* [27], possibly resulting in lower levels of DIP-α and a compromised

interaction between αMNs and muscles. For αTi-tadm there is a slight decrease in

branch number and length in both readouts and in both cases the αTi-tadm arborization

phenotype is less strong than with αTi-ltm (Figure 2C and D, Figure S1D and F).

Together, these results suggest that the affinity requirement for the arborization

of αFe-ltm is higher than the αTi-ltm, consistent with the differences observed in

expressivity for *DIP-α* and *dpr10* null alleles. αTi-tadm is rarely fully lost in either *DIP-α*

or *dpr10* null alleles indicating that it relies on additional mechanisms to achieve its

proper branch number and length. Finally, these results indicate that varying the affinity

from the postsynaptic target has the same effects as varying the affinity from the

presynaptic αMN.

Increasing affinity also impedes αMN arborization

Given the loss of arborization when the DIP-α::Dpr10 affinity was reduced, we

considered the possibility that increasing the affinity might result in more and/or longer

branches. To address this question we used the following *DIP-α* and *dpr10* mutants:

DIP-α^{G74A} (K_D = 0.90 μM, *DIP-α*$^{+2F}$) and *dpr10*^{Q142M} (K_D = 0.19 μM, *dpr10*$^{+10F}$). In

addition, we also analyzed the combination of *DIP-α*$^{+2F}$ and *dpr10*$^{+10F}$, which results in

an increased affinity of +15F (*DIP-α & dpr10*$^{+15F}$; K_D = 0.10 μM) (Figure 4A). We find

that in *DIP-α*$^{+2F}$ animals, αMN branch number and length is not affected (Figure 4C).

Surprisingly, in *dpr10*$^{+10F}$ flies, αTi-ltm and αTi-tadm branch number and branch length

were on average shorter than in control flies (Figure 4C and D). In *DIP-α & dpr10*$^{+15F}$

flies, some αTi-ltm and αTi-tadm axons completely fail to arborize. Together, these

results show that increasing affinity, like reducing affinity, also adversely affects terminal

branch arborization. We note however that using *A8-Gal4* as the readout, the lack of

arborization phenotype is not as strong as in the experiments done with the *DIP-α-T2A-*

Gal4, even in a sensitized background with only one functional copy of *DIP-α* (*DIP-α &*

dpr10^{+15}/DIP-α^{Null}). Nevertheless, there is a reduction in branch number and branch

length for both the αFe-ltm and αTi-ltm in this genotype (Figure S2). Together, these

experiments indicate that increasing the affinity between DIP-α and Dpr10 only has

strong effects when the affinity is above a certain threshold, and that the penetrance of

this effect depends on the status of the *DIP-α* locus (see Discussion).

To determine if αMNs with reduced arborization are able to make synapses, we

expressed the marker Bruchpilot (Brp-short::mStraw[38]) in *DIP-α & dpr10^{+15F}* and control

flies. Although we were not able get single bouton resolution of Brp puncta, we are

nevertheless able to see that in *DIP-α & dpr10^{+15F}* flies, αTi-ltm and αTi-tadm have Brp

puncta, suggesting that they assemble synapses (Figure S3).

Increasing affinity prevents the stabilization of developing αMNs

It is possible that *DIP-α & dpr10^{+15F}* αMNs do not stabilize on their muscle targets

either because 1) they never send out tertiary branches or 2) they send out branches

but they are not capable of stabilizing on their targets. To test this, we took time lapse

videos and compared control to *DIP-α & dpr10^{+15F}* flies, imaged from ~25-80 hrs APF,

the time window when αMNs form their secondary and terminal branches and stabilize

on their muscle targets[27] (Video 2). In control animals, at ~25 hrs APF both αTi-ltm and

αTi-tadm have already reached their muscle targets. Between 35-55 hrs APF the axons

will branch and extend filopodia in the general area of their muscle targets and then at

~60 hrs APF some filopodia will stabilize. In *DIP-α & dpr10*$^{+15F}$ animals, both αTi-ltm and

αTi-tadm reach their muscle targets successfully at 24 hrs APF. Very few differences

can be seen in the next 24 hours. However, at 60 hrs APF the tertiary branches that

have extended fail to stabilize and instead retract resulting in the phenotype that we see

in the adult. Interestingly, this is similar to what has been described for *DIP-α* nulls

(Figure 5A). These results indicate that αMNs can initially "recognize" the correct

muscle target and send out filopodia to the correct myoblast cluster but fail to maintain

the interaction long enough for the axons to fully arborize.

Changing affinity leads to behavioral defects

The three αMNs directly target the Fe-ltm, Ti-ltm, and Ti-tadm muscles[27]. It is

plausible that locomotion defects might occur when the terminal branches in these MNs

fail to properly arborize. Previously our lab established a free walking behavioral assay

that reports a dozen different walking parameters[39] (Figure 5B). We compared both

DIP-α^{Null} and *DIP-α & dpr10*$^{+15F}$ to control animals because these alleles are highly

penetrant. Both *DIP-α*^{Null} and *DIP-α & dpr10*$^{+15F}$ flies have a decreased overall velocity

and median walking velocity. Furthermore, *DIP-α & dpr10*$^{+15F}$ flies also have a reduced

absolute angular velocity compared to controls (Figure 5C). Although, it is possible that

these effects are the result of other, non-MN *DIP-α*-expressing neurons elsewhere in

the nervous system, they demonstrate that changing the affinity between DIP-α::Dpr10

adversely affects locomotion.

Discussion

Here we use single neuron morphology measurements to understand the importance of affinity between two CAMs that are required to maintain contacts between MNs and muscles in the Drosophila leg neuromuscular system[37]. We address this question using a series of knock in alleles of both *DIP-α* and *dpr10* that change the affinity of these proteins, as measured *in vitro*[33]. We show that decreasing the affinity past a certain threshold leads to phenotypes similar to the null phenotype[27]. Surprisingly, we also find that increasing the affinity past a certain threshold also reduces the amount of axonal arborization. We also show via live imaging that axons in flies with high affinity alleles are able to extend filopodia to the correct muscle target but ultimately fail to stabilize and arborize. Together, these data show that a specific affinity window for these interacting CAMs is needed for proper arborization.

The loss of branching in the high affinity mutants was initially surprising. Our lab has recently shown that *dpr10* is also expressed by αMNs and that removing MN-expressed *dpr10* leads to abnormally long branches in both the αTi-ltm and αTi-tadm[40]. Therefore, a possible mechanism that contributes to the phenotype seen in the *DIP-α & dpr10^{+15F}* mutant is that the higher affinity interaction between DIP-α and Dpr10 is occurring in *cis* (in MNs), which reduces DIP-α's ability to interact with Dpr10 in *trans* in muscle cells. This could explain why the filopodia still target the correct muscles but are not able to "anchor" long enough to generate stable terminal branches. Dpr10 is also expressed by other MNs in addition to the αMNs. Therefore, another explanation is that DIP-α in αMNs is interacting with Dpr10 present in other MNs. However, this seems less

likely as we would expect the filopodia to become less mobile in the *DIP-α & dpr10*$^{+15F}$

mutant due to more and stronger interneuronal contacts.

Another interesting observation is that, like the time lapse videos of *DIP-α null*[27],

the live imaging videos of *DIP-α & dpr10*$^{+15F}$ mutants reveal that filopodia still extend out

to their correct muscle target but unable to make stable contacts with their target and

then retract (Figure 5A and Video 2). This reinforces the idea that DIP-α::Dpr10 *trans*

interactions are needed to stabilize terminal branches, and not for axon guidance or

synaptic specificity. Further, this suggests that there must be additional mechanisms

that help MN axons recognize their targets. We propose that the proper DIP-α::Dpr10

affinity range is needed to stabilize the contact long enough to recruit more DIP-α::dpr10

trans interactions and keep the axon arborizing.

It is not clear why the phenotype is stronger in *DIP-α & dpr10*$^{+15F}$*/DIP-α-T2A-Gal4*

animals compared to *DIP-α & dpr10*$^{+15F}$ */DIP-α*^{Null} animals using *A8-Gal4*, when

theoretically the amount of DIP-α should be the same in both genotypes. It is possible

that a truncated DIP-α protein generated by the *DIP-α-T2A-Gal4* allele is further

sensitizing the phenotype. One way to test this is to overexpress Dpr10 in αMNs and/or

in muscles in *DIP-α & dpr10*$^{+15F}$*/DIP-α*^{Null} flies and see whether we obtain similar

phenotypes to those seen in *DIP-α & dpr10*$^{+15F}$*/DIP-α-T2A-Gal4* flies. This would also

allow us to determine whether the phenotype is due to differences in levels in the MNs

or the muscles. We have shown that overexpressing *dpr10* in muscle does not have a

phenotype but that overexpressing *dpr10* and *dpr6* in αMNs leads to *cis*-inhibition and a

reduction in aborization[40]. Therefore, it would be interesting to see if *DIP-α & dpr10*$^{+15F}$

/*DIP-α* ^{Null}; *A8-Gal4>>UAS-dpr10* flies have the same phenotype and is indeed a result

of *cis* inhibition.

Furthermore, it is interesting that different neural systems utilize the same

molecules for different purposes. As noted in the Introduction, reducing DIP-α::Dpr10

affinity in the visual system leads to cell survival and synaptic targeting defects. Here,

DIP-α::Dpr10 is not required for cell survival nor synapse formation. We propose that

DIP-α and Dpr10 interact with different molecules in different contexts to achieve a

variety of functions. These differences may also be a consequence of the more hard-

wired nature of how MNs are born compared to the more plastic way in which an excess

of neurons are born in the visual system, leading some to apoptose.

Methods

Dissection and microscopy

For adult leg dissection, flies were first immersed in 70% Ethanol for ~1 min and

rinsed in 0.3% Triton-X in PBS 3X. Abdomen and heads were then removed and legs

attached to thorax were fixed overnight at 4°C in 4% paraformaldehyde in 0.3% Triton-X

in PBS. The next day, legs and thorax were washed five times in 0.3% Triton-X in PBS

then stored in 80% VECTASHIELD (in PBS) overnight at 4°C. Legs were then removed

from the thorax and placed onto a glass slide in a drop of VECTASHIELD mounting

medium (Vector Labs). 18X18mm coverslips were then placed on top with dental wax

on the corners to prevent the coverslip from crushing the legs. 0.5 µm-thick sections in

the Z axis were imaged using a Zeiss LSM 800 Confocal Laser Scanning Microscope.

Motor neuron axon branch quantification and statistical analysis

To quantify leg motor neurons only one T1 leg from each animal was imaged and their neurons traced using the Simple Neuron Tracer from ImageJ[37]. For each neuron, with the exception of the αTi-tadm, the tracing began at the first bifurcation point. For the αTi-tadm, only the most distal branch was traced due to presence of collateral branches which would have potential misrepresented the terminal branch number and branch length. Both total terminal branch number and the sum of cable length values were separately plotted and analyzed using the GraphPad Prism 9.0 software. For all graphs statistical significance was determined using an unpaired nonparametric two tailed Mann-Whitney test. Error bars represent mean with 95% confidence intervals. ns= no statistical difference. *$p < 0.05$ **$p < 0.01$ ***$p < 0.001$ ****$p < 0.0001$.

Live Imaging of MN development in pupa

Pupae were first staged by picking late (wandering) L3 larva and placing them in a separate vial with a moist tissue, kept at 25°C and then picked 24 hours later. A small window on the left ventral side of the pupal case was made using forceps to expose just the T1 leg. Individual pupae were placed on a glass slide, surrounded by two layers of filter paper dampened with distilled water. A 5 ul drop of distilled water was placed at the center of a glass coverslip (N-1.5) and placed exactly over the exposed T1 leg. Petroleum jelly surrounding the filter paper was used to seal the space between the coverslip and the glass slide to retain humidity. Samples were imaged on a Zeiss LSM

800 microscope, 25X objective, with a 10 min interval between each z-stack series.

Videos were generated using the FIJI software[41] at five frames per second.

Fly arena experiments

The fly arena setup was constructed by a previous member of the lab and described in

Howard et al, 2009[39]. Briefly, the polycarbonate plastic arena sits in an aluminum plate,

which maintains a level surface and is covered with a circular acrylic disc that has a

small hole for mouth pipetting in flies. The inside of the lid is freshly coated with a thin

layer of fluon (Amazon, B00UJH12A) to prevent flies from walking on the ceiling. Videos

were recorded with a Point Grey Blackfly Mono USB3 camera fitted with a Tamron ½"

F/1.2 ER C-mount lens (B&H photo) was mounted above the arena and connected to a

computer running Ubuntu 20.04 LTS. A Kodax 3x3" 89B Opaque IR filter (B&H photo)

was placed in front of the camera detector to allow for detection of IR but not visible

light. Backlighting was provided by a plate of LEDs arranged in parallel rows underneath

the arena. An acrylic diffuser was placed in front of the camera detector to allow for

detection of IR but not visible light. Infrared LEDs were acquired from

ledlightsworld.com (SMD3528-300). The IR lights were set at 100% brightness.

Data Acquisition: For recordings, 3-5 day old mated females were kept in a

humidity and temperature controlled incubator with a 12 hr day/night. All behavior

videos were recorded between the morning activity peak time (9am-12pm). Prior to the

recording, the arena was leveled, the lid cleaned and layered with fluon to prevent the

flies from walking on the lid. Each recorded video had 10 flies. Videos were taken on

multiple days and pooled for the final dataset (N = 30). For every recording session, flies

were mouth pipetted into the arena through the small hole in the lid and the lid was slid

to move the hole out of the camera view. A blackout curtain was used to cover the

arena to protect from any contaminating light. All videos were recorded at a rate of 30

frames per second and stored in a compressed "fly format" using software written by

Andrew Straw at the University of Frieburg. Videos were tracked using the Matlab

based FlyTracker software from the Caltech vision lab[42]. Before tracking, pixel to mm

conversion was calibrated using the inbuilt GUI of the tracker software and a single

calibration file was generated for all the videos recorded on the same day. Background

model and thresholds were adjusted to provide optimal recognition of animals and were

not standardized between recording sessions. All analysis on data from arena

experiments was performed in MATLAB using scripts written by Dr. Clare E. Howard

and modified by Dr. Sumaira Zamurrad.

Behavioral classifiers:

<u>Walk:</u> Walking frames were defined using a dual threshold Schmitt trigger filter

(original script written by Irene Kim – Dickinson lab). Speed thresholds were set at

1 and 2.5 mm/s, and time thresholds were .1 s. Walking frames were also specified

to be those in which the fly was not already engaged in a jump.

<u>Stop:</u> Stop frames were classified as any frames where animals were not performing

walking or jumping behaviors.

424
425
426   **Parameters:**

427
<u>Walk Frequency</u>: the percent of frames classified as walking during the recording
period.

<u>Overall Velocity</u>: the median of all velocities over the recording period.

<u>Median Walking</u> Velocity: the median of velocities during all frames when the animal is
classified as walking.

<u>Maximum Walking Velocity</u>: the maximum velocity an animal reaches during
walking.

<u>Angular velocity</u>: the median value of angular velocity. This parameter takes into
account directionality of turning.

<u>Absolute Angular Velocity</u>: the median of the absolute value of angular velocities.
This parameter does not take into account directionality of turning.

<u>Walking bout duration</u>: the length of each bout was calculated, and the median
of all bout lengths was taken for each animal.

<u>Stop bout number and duration</u>: calculated as for walking bouts.

For each animal, the value of each parameter was defined as described above for a

ten-minute walking period in the dark. Comparisons with control were drawn with

Kruskall-Wallis (for non-normally distributed data) analysis.

Acknowledgements

This work was funded by a grant from NINDS, NS070644, to R.S.M. We thank Drs.

Honig, Shapiro, Zinn, and Zipursky for sharing data and ideas in the early stages of this

work and Dr. L. Venkatasubramanian for training and advice in the early phase of this

project.

Author contributions

D.L. executed all of these experiments, including the statistical analyses, with help from

K.R. S.Z. helped with the behavioral analysis. S.X. provided the knock-in alleles and

additional information regarding these chromosomes. D.L. wrote the initial draft. R.S.M.

conceptualized the study, provided feedback, and edited the manuscript.

Figure legends

Figure 1. Overview of fly leg MN arborization.

A. Life cycle of *Drosophila*. Imaging was done during most of pupal development, from 20 hours (hrs) APF and onwards.

B. Drawings and individual frames from videos of *DIP-α-T2A-Gal4>>20X6XGFP, Mef2-QF>>10XQUAS-6XmCherry* flies. First row shows αMNs in green and muscles in magenta. Sequence shows αMNs developing in the context of myoblast fusion. Second row shows developing αMNs. At early time points, αMNs are more mobile but from 60 hrs APF onwards axon termini begin to stabilize yet continue to mature until eclosion. Third row shows stills from the video (see Video 1).

C. Adult fly (left) and leg (right). αMNs, visualized here with *A8-Gal4, 20X6XUAS-GFP*, are born in the VNC and extend to arborize on muscles in the tibia and femur. Ti-ltm= Tibia long tendon muscle, tadm-tarsal depressor muscle, talm-tarsal levator muscle

Figure 2. Analysis of *DIP-α* alleles with reduced affinity.

A. Affinity range of *DIP-α* mutants in log scale.

B. Representative images of *yw/DIP-α-T2A-Gal4>>20XUAS-6XGFP* (control) and *DIP-α^{null}/DIP-α-T2A-Gal4>>20XUAS-6XGFP* tibia leg segments.

C. Schematic of how MN morphologies are quantified.

D. Quantification of terminal branch number of *DIP-α* mutant alleles over *DIP-α-T2A-*

Gal4>>20XUAS-6XGFP. αTi-ltm: control, n=10; *DIP-α^{-20F}* n=10; *DIP-α^{-50F}*, n=10; *DIP-*

α^{Null} n=10; αTi-tadm: control, n=12; *DIP-α^{-20F}* n=10; *DIP-α^{-50F}* n=11; *DIP-α^{Null}* n=11.

E. Quantification of total branch length of control and *DIP-α* mutant alleles over *DIP-α-*

T2A-Gal4>>20XUAS-6XGFP. αTi-ltm: control, n=10; *DIP-α^{-20F}* n=10; *DIP-α^{-50F}* n=10;

DIP-α^{Null} n=10; αTi-tadm: control, n=12; *DIP-α^{-20F}* n=10; *DIP-α^{-50F}* n=11; *DIP-α^{Null}* n=10.

For all graphs statistical significance was determined using an unpaired nonparametric

two tailed Mann-Whitney test. Error bars represent mean with 95% confidence intervals.

ns= no statistical difference. *p<0.05 **p<0.01 ***p<0.001 ****p<0.0001.

Figure 3. Analysis of *dpr10* alleles with reduced affinity.

A. Affinity range of *dpr10* mutants in log scale.

B. Representative images of *yw/DIP-α-T2A-Gal4>>20XUAS-6XGFP* (control) and

yw/DIP-α-T2A-Gal4;dpr6$^-$dpr10$^-$//>>20XUAS-6XGFP tibia leg segments.

C. Quantification of terminal branch number of *dpr10* mutants, *yw/DIP-α-T2A-Gal4;*

dpr6$^-$dpr10$^{affinity\ mutant}$//>>20XUAS-6XGFP. αTi-ltm: control, n=10; *dpr10^{-8F}* n=10; *dpr10$^-$*

^{20F} n=10; *dpr10^{-40F}* n=10; *dpr10^{Null}* n=9; αTi-tadm: control, n=12; *dpr10^{-8F}* n=8; *dpr10^{-20F}*

n=11; *dpr10^{-40F}* n=11; *dpr10^{Null}* n=10**.**

D. Quantification of total branch length for control and *yw/DIP-α-T2A-Gal4; dpr6$^-$*

dpr10$^{affinity\ mutant}$//>>20XUAS-6XGFP .αTi-ltm: control, n=10; *dpr10^{-8F}* n=10; *dpr10^{-20F}*

n=10; *dpr10^{-40F}* n=10; *dpr10^{Null}* n=9; αTi-tadm: control, n=12; *dpr10^{-8F}* n=8; *dpr10^{-20F}*

n=11; *dpr10^{-40F}* n=11; *dpr10^{Null}* n=10. The control samples in Figure 2D and E were the

same because all experiments were done at the same time. For all graphs statistical

significance was determined using an unpaired nonparametric two tailed Mann-Whitney

test. Error bars represent mean with 95% confidence intervals. ns= no statistical

difference. *p<0.05 **p<0.01 ***p<0.001 ****p<0.0001.

Figure 4. Analysis of increased affinity genotypes.

A. Affinity range of increased affinity mutants in log scale.

B. Representative images of *yw/DIP-α-T2A-Gal4>>20XUAS-6XGFP* (control) and *DIP-*

α^{G74A}/DIP-α-T2A-Gal4;dpr6⁻dpr10^{Q142M}//>>20XUAS-6XGFP (+15F) tibia segments.

C. Quantification of terminal branch number of *dpr10* mutants, *DIP-α^{G74A}/DIP-α-T2A-*

Gal4//>>20XUAS-6XGFP(+2F);   yw/DIP-α-T2A-Gal4;   dpr6⁻dpr10^{Q142M}//>>20XUAS-

6XGFP(+10F);   and   DIP-α^{G74A}/DIP-α-T2A-Gal4;   dpr6⁻dpr10^{Q142M}//>>20XUAS-

6XGFP(+15F). αTi-ltm: control, n=10; *DIP-α^{+2F}* n=10; *dpr10^{+10F}* n=10; *DIP-α^{+2F}* &

dpr10^{+10F} n=10; αTi-tadm: control, n=12; *DIP-α^{+2F}* n=12; *dpr10^{+10F}* n=12; *DIP-α^{+2F}* &

dpr10^{+10F} n=12.

D. Quantification of total branch length in control and *DIP-α^{G74A}/DIP-α-T2A-*

Gal4//>>20XUAS-6XGFP(+2F);   yw/DIP-α-T2A-Gal4;   dpr6⁻dpr10^{Q142M}//>>20XUAS-

6XGFP(+10F); and DIP-α^{G74A}/DIP-α-T2A-Gal4; dpr6⁻dpr10^{Q142M}//>>20XUAS-6XGFP

(+15F) flies. αTi-ltm: control, n=10; *DIP-α^{+2F}* n=10; *dpr10^{+10F}* n=10; *DIP-α^{+2F}* & *dpr10^{+10F}*

n=10; αTi-tadm: control, n=12; *DIP-α $^{+2F}$* n=12; *dpr10^{+10F}* n=12 *DIP-α $^{+2F}$* & *dpr10^{+10F}*

n=12. The control samples for Figure 2D and E were the same because all experiments

were done at the same time. For all graphs statistical significance was determined using

an unpaired nonparametric two tailed Mann-Whitney test. Error bars represent mean

with 95% confidence intervals. ns= no statistical difference. *$p<0.05$ **$p<0.01$

***$p<0.001$ ****$p<0.0001$.

Figure 5. Live imaging and locomotion analysis.

A. Still images of a timelapse video (see Video 2) of *yw/DIP-α-T2A-Gal4>>20XUAS-*

6XGFP pupa (top row) and *DIP-α^{G74A}/DIP-α-T2A-Gal4;dpr6⁻dpr10^{Q142M}//>>20XUAS-*

6XGFP (+15F) (bottom row) flies. Yellow arrowheads point to small filopodia that

aberrantly extend but ultimately retract in *DIP-α^{G74A}/DIP-α-T2A-Gal4; dpr6⁻dpr10^{Q142M}*

MNs.

B. Free walking arena schematic.

C. Walking parameters measured in control (*yw, 20XUAS-6XGFP*); *DIP-α^{Null}, 20XUAS-*

6XGFP, and +15F gain-of-affinity *DIP-α^{G74A}//20XUAS-6XGFP;dpr6⁻dpr10^{Q142M}(+15)//*

genotypes. Control n=10; *DIP-α^{Null}* n=10; *DIP-α & dpr10^{+15F}* n=10. *$p<0.05$

Supplementary Figure Legends

Figure S1. Analysis of decreased affinity mutants using *A8-Gal4* as readout.

A. Representative femur and tibia images of *yw, A8-Gal4>>20XUAS-6XGFP* (control)

and *DIP-α^{Null}, A8-Gal4>>20XUAS-6XGFP* femur and tibia leg segments.

B. Quantification of terminal branch number of *DIP-α* mutants; αFe-ltm: control n=8;

DIP-α^{-20F}, A8-Gal4 n=10; *DIP-α^{-50F}, A8-Gal4* n=10; *DIP-α^{Null}, A8-Gal4* n=10; αTi-ltm:

control n=9; *DIP-α^{-20F}, A8-Gal4* n=9; *DIP-α^{-50F}, A8-Gal4* n=10; *DIP-α^{Null}, A8-Gal4* n=8;

αTi-tadm: control n=9; *DIP-α^{-20F}, A8-Gal4* n=10; *DIP-α^{-50F}, A8-Gal4* n=9; *DIP-α^{Null}, A8-*

Gal4 n=10.

C. Quantification of total branch length of control and *DIP-α* mutants. αFe-ltm: control

n=8; *DIP-α^{-20F}, A8-Gal4* n=10; *DIP-α^{-50F}, A8-Gal4* n=10; *DIP-α^{Null}, A8-Gal4* n=10; αTi-

ltm: control n=9; *DIP-α^{-20F}, A8-Gal4* n=9; *DIP-α^{-50F}, A8-Gal4* n=10; *DIP-α^{Null}, A8-Gal4*

n=9; αTi-tadm: control n=9; *DIP-α^{-20F}, A8-Gal4* n=10; *DIP-α^{-50F}, A8-Gal4* n=9; *DIP-α^{Null},*

A8-Gal4 n=10.

D. Representative images of *yw, A8-Gal4>>20XUAS-6XGFP* (control) and *dpr6$^-$,dpr10$^-$*

,A8-Gal4>>20XUAS-6XGFP femur and tibia leg segments.

E. Quantification of terminal branch number of *dpr10* mutants. αFe-ltm: control n= 11;

dpr6$^-$,dpr10^{-8F}, A8-Gal4 n= 9*; dpr6$^-$,dpr10^{-20F}, A8-Gal4* n=10; *dpr6$^-$,dpr10^{-40F}, A8-Gal4*

n=11; *dpr6$^-$, dpr10^{Null}, A8-Gal4* n= 12; αTi-ltm: Control n= 11; *dpr6$^-$,dpr10^{-8F}, A8-Gal4* n=

9; *dpr6$^-$,dpr10^{-20F}, A8-Gal4* n=10; *dpr6$^-$,dpr10^{-40F}, A8-Gal4* n=11; *dpr6$^-$, dpr10^{Null}, A8-*

Gal4 n= 12; αTi-tadm: Control n= 11; *dpr6$^-$,dpr10^{-8F}, A8-Gal4* n= 9; *dpr6$^-$,dpr10^{-20F}, A8-*

Gal4 n=10; *dpr6$^-$,dpr10^{-40F}, A8-Gal4* n=10; *dpr6$^-$, dpr10^{Null}, A8-Gal4* n=12.

F. Quantification of total branch length of *dpr10* mutants. αFe-ltm: control n=11; *dpr6$^-$*

,dpr10^{-8F}, A8-Gal4 n=9; *dpr6$^-$,dpr10^{-20F}, A8-Gal4* n=10; *dpr6$^-$,dpr10^{-40F}, A8-Gal4* n=11;

dpr6$^-$, dpr10^{Null}, A8-Gal4 n=12; αTi-ltm: control n=11; *dpr6$^-$,dpr10^{-8F}, A8-Gal4* n=9; *dpr6$^-$*

,dpr10^{-20F}, A8-Gal4 n=10; *dpr6$^-$,dpr10^{-40F}, A8-Gal4* n=11; *dpr6$^-$, dpr10^{Null}, A8-Gal4* n=12;

αTi-tadm: control n=11; *dpr6$^-$,dpr10^{-8F}, A8-Gal4* n=9; *dpr6$^-$,dpr10^{-20F}, A8-Gal4* n=10;

dpr6$^-$,dpr10^{-40F}, A8-Gal4 n=11; *dpr6$^-$, dpr10^{Null}, A8-Gal4* n=12. For all graphs statistical

significance was determined using an unpaired nonparametric two tailed Mann-Whitney

test. Error bars represent mean with 95% confidence intervals. ns= no statistical

difference. *p<0.05 **p<0.01 ***p<0.001 ****p<0.0001. All quantifications done with

20XUAS-6XGFP.

Figure S2. Analysis of increased affinity mutants with A8-Gal4 as the readout.

A. Representative images of control, *yw, DIP-α^{Null}, A8-Gal4, dpr6 dpr10^{Q142M}*

heterozygous>>20XUAS-6XGFP and *DIP-α^{G74A}, DIP-α^{Null}, A8-Gal4, dpr6 dpr10^{Q142M}*

>>20XUAS-6XGFP homozygous femur and tibia leg segments.

B. Quantification of terminal branch number of control, *yw, DIP-α^{Null}, A8-Gal4, dpr6-*

dpr10^{Q142M} heterozygous>>*20X-6XGFP* and *DIP-α^{G74A}, DIP-α^{Null}, A8-Gal4, dpr6-*

dpr10^{Q142M} homozygous genotypes. αFe-ltm: control n=11; *DIP-α^{+2}, DIP-α^{Null}, dpr6-*

dpr10^{+10} n=13; αTi-ltm: control n=10; *DIP-α^{+2}, DIP-α^{Null}, dpr6-dpr10^{+10}* n=13; αTi-tadm:

control n=11; *DIP-α^{+2}, DIP-α^{Null}, dpr6-dpr10^{+10}* n=13.

C. Quantification of total branch length of control *yw, DIP-α^{Null}, A8-Gal4, dpr6-*

dpr10^{Q142M} heterozygous>>*20X-6XGFP* and *DIP-α^{G74A}/ DIP-α^{Null}, A8-Gal4, dpr6-*

dpr10^{Q142M} homozygous genotypes. αFe-ltm: control n=11, *DIP-α^{+2}/DIP-α^{Null}, dpr6-*

dpr10^{+10} n=13; αTi-ltm: control n=10; *DIP-α^{+2}/ DIP-α^{Null}, dpr6-dpr10^{+10}* n=14; αTi-tadm:

control n=11; *DIP-α^{+2}/ DIP-α^{Null}, dpr6-dpr10^{+10}* n=13. For all graphs statistical

significance was determined using an unpaired nonparametric two tailed Mann-Whitney

test. Error bars represent mean with 95% confidence intervals. ns= no statistical

difference. *p<0.05 **p<0.01 ***p<0.001 ****p<0.0001. All quantifications were done

with *20XUAS-6XGFP.*

Figure S3. Truncated axon termini have boutons

A. Representative images of tibia leg segments from control, *yw, DIP-α-T2A-Gal4, dpr6-dpr10^{Q142M} heterozygous>>20XUAS-6XGFP, UAS-brp-short::mStraw* and *DIP-α^{G74A}, DIP-α-T2A-Gal4, dpr6-dpr10^{Q142M} homozygous>>20XUAS-6XGFP, UAS-brp-short::mStraw* flies.

B. High magnification images of boxed regions in **A** for the control. **B:** αTi-ltm. **B':** αTi-tadm.

C. High magnification images of boxed regions in **A** for the +15F genotype. **C:** αTi-ltm. **C':** αTi-tadm.

Figure S4. **Leg myoblast fusion during pupal development.**

Still images of Mef2 expressing leg myoblasts during pupal development. Arrowhead points to Ti-ltm connecting to femur muscles. Arrows point to two femur muscle groups segregating during development.

Video legends

Video 1:

https://www.dropbox.com/scl/fi/idapupdlq2gkcb033upzg/Video_1.mov?rlkey=fjwstxhhs4h4wepbkzbxhrlqt&dl=0

DIP-α-T2A-Gal4>>20XUAS-6XGFP,Mef2-QUAS>>10XQUAS-6xmCherry.

Green=αMNs; Magenta=myoblasts

Video 2:

https://www.dropbox.com/scl/fi/9wfaszkqwem6eyw33rnzc/Video_2.mov?rlkey=ln4xzlqn9

t7ttf5ile6vtdg3g&dl=0

Left: yw/DIP-α-T2A-Gal4>>20XUAS-6XGFP; Right: DIP-α^{+2}/DIP-α-T2A-

Gal4;dpr6,dpr10^{+10} (+15)>>20XUAS-X6GFP.

Video S1:

https://www.dropbox.com/scl/fi/t1hjvkovek7n1xsteoj4z/Video_S1.mov?rlkey=zotp832dd

ptw9i3aarqq8x058&dl=0

VGlut-LexA>>13XLexAop-myr::GFP, Mef2-Gal4>10XUAS-myr::tdTomato

Literature cited

1. Chiba, A. Early development of the drosophila neuromuscular junction: A model for studying neuronal networks in development. *Int. Rev. Neurobiol.* **43**, 1–24 (1999).

2. Moreland, T. & Poulain, F. E. To Stick or Not to Stick: The Multiple Roles of Cell Adhesion Molecules in Neural Circuit Assembly. *Front. Neurosci.* **16**, 1–16 (2022).

3. Stoeckli, E. T. Understanding axon guidance: Are we nearly there yet? *Dev.* **145**, (2018).

4. Van Vactor, D., Sink, H., Fambrough, D., Tsoo, R. & Goodman, C. S. Genes that control neuromuscular specificity in Drosophila. *Cell* **73**, 1137–1153 (1993).

5. Sanes, J. R. & Zipursky, S. L. Synaptic Specificity, Recognition Molecules, and Assembly of Neural Circuits. *Cell* **181**, 536–556 (2020).

6. Tan, L. *et al.* Ig Superfamily Ligand and Receptor Pairs Expressed in Synaptic Partners in Drosophila Accession Numbers GSE68235 Tan et al. *Cell* **163**, (2015).

7. Grueber, W. B. & Sagasti, A. Self-avoidance and Tiling□: Mechanisms. 1–16 (2010).

8. Gibson, D. A. & Ma, L. Developmental regulation of axon branching in the vertebrate nervous system. *Development* **138**, 183–195 (2011).

9. Park, Y. K. & Goda, Y. Integrins in synapse regulation. *Nat. Rev. Neurosci.* **17**, 745–756 (2016).

10. Rohrbough, J., Grotewiel, M. S., Davis, R. L. & Broadie, K. Integrin-mediated

regulation of synaptic morphology, transmission, and plasticity. *J. Neurosci.* **20**, 662   6868–6878 (2000).

11. Sink, H. *et al. sidestep Encodes a Target-Derived Attractant Essential for Motor* 664   *Axon Guidance in Drosophila. Cell* **105**, (2001).

12. Siebert, M., Banovic, D., Goellner, B. & Aberle, H. Drosophila motor axons 666   recognize and follow a Sidestep-labeled substrate pathway to reach their target 667   fields. doi:10.1101/gad.520509

13. Ango, F. *et al.* Ankyrin-based subcellular gradient of neurofascin, an 669   immunoglobulin family protein, directs GABAergic innervation at Purkinje axon 670   initial segment. *Cell* **119**, 257–272 (2004).

14. Telley, L. *et al.* Dual Function of NRP1 in Axon Guidance and Subcellular Target 672   Recognition in Cerebellum. *Neuron* **91**, 1276–1291 (2016).

15. Ashrafi, S. *et al.* Neuronal Ig/Caspr Recognition Promotes the Formation of 674   Axoaxonic Synapses in Mouse Spinal Cord. *Neuron* **81**, 120–129 (2014).

16. Lemieux, M., Thiry, L., Laflamme, O. D. & Bretzner, F. Role of dscam in the 676   development of neural control of movement and locomotion. *Int. J. Mol. Sci.* **22**, 677   1–26 (2021).

17. Tadros, W. *et al.* Dscam Proteins Direct Dendritic Targeting through Adhesion. 679   *Neuron* **89**, 480–493 (2016).

18. Lefebvre, J. L., Kostadinov, D., Chen, W. V., Maniatis, T. & Sanes, J. R. 681   Protocadherins mediate dendritic self-avoidance in the mammalian nervous 682   system. *Nature* **488**, 517–521 (2012).

19. Denda, S. & Reichardt, L. F. Studies on Integrins in the Nervous System.

Methods Enzymol. **426**, 203–221 (2007).

20.   Kinold, J. C., Pfarr, C. & Aberle, H. Sidestep-induced neuromuscular miswiring

causes severe locomotion defects in Drosophila larvae. (2018).

doi:10.1242/dev.163279

21.   Shimoda, Y. & Watanabe, K. Contactins: Emerging key roles in the development

and function of the nervous system. *Cell Adhes. Migr.* **3**, 64–70 (2009).

22.   Özkan, E. *et al.* XAn extracellular interactome of immunoglobulin and LRR

proteins reveals receptor-ligand networks. *Cell* **154**, 228 (2013).

23.   Zinn, K. & Özkan, E. Neural immunoglobulin superfamily interaction networks.

Curr. Opin. Neurobiol. **45**, 99–105 (2017).

24.   Cosmanescu, F. *et al.* Neuron-Subtype-Specific Expression, Interaction Affinities,

and Specificity Determinants of DIP/Dpr Cell Recognition Proteins. *Neuron* **100**,

1385-1400.e6 (2018).

25.   Carrillo, R. A. *et al.* Control of Synaptic Connectivity by a Network of Drosophila

IgSF Cell Surface Proteins. *Cell* **163**, 1770–1782 (2015).

26.   Wang, Y. *et al.* Systematic expression profiling of Dpr and DIP genes reveals cell

surface codes in Drosophila larval motor and sensory neurons. *Dev.* **49**, (2022).

27.   Venkatasubramanian, L. *et al.* Stereotyped terminal axon branching of leg motor

neurons mediated by IgSF proteins DIP-α and Dpr10. *Elife* **8**, 1–27 (2019).

28.   Bornstein, B. *et al.* Transneuronal Dpr12/DIP-δ interactions facilitate

compartmentalized dopaminergic innervation of Drosophila mushroom body

axons . *EMBO J.* **40**, 1–21 (2021).

29.   Brovero, S. G. *et al.* Investigation of drosophila fruitless neurons that express

dpr/dip cell adhesion molecules. *Elife* **10**, 1–69 (2021).

30. Xu, S. *et al.* Interactions between the Ig-Superfamily Proteins DIP-α and Dpr6/10

Regulate Assembly of Neural Circuits. *Neuron* **100**, 1369-1384.e6 (2018).

31. Ashley, J. *et al.* Transsynaptic interactions between IgSF proteins DIP-α and

Dpr10 are required for motor neuron targeting specificity. *Elife* **8**, 1–24 (2019).

32. Venken, K. J. T. *et al.* MiMIC: A highly versatile transposon insertion resource for

engineering Drosophila melanogaster genes. *Nat. Methods* **8**, 737–747 (2011).

33. Xu, S. *et al.* Affinity requirements for control of synaptic targeting and neuronal

cell survival by heterophilic IgSF cell adhesion molecules. *Cell Rep.* **39**, 110618

(2022).

34. Courgeon, M. & Desplan, C. Coordination between stochastic and deterministic

specification in the Drosophila visual system. *Science (80-.).* **366**, (2019).

35. Enriquez, J. *et al.* Specification of Individual Adult Motor Neuron Morphologies by

Combinatorial Transcription Factor Codes. *Neuron* **86**, 955–970 (2015).

36. Guan, W. *et al.* Post-transcriptional regulation of transcription factor codes in

immature neurons drives neuronal diversity. *Cell Rep.* **39**, (2022).

37. Arshadi, C., Günther, U., Eddison, M., Harrington, K. I. S. & Ferreira, T. A. SNT: a

unifying toolbox for quantification of neuronal anatomy. *Nat. Methods* **18**, 374–377

(2021).

38. Mosca, T. J. & Luo, L. Synaptic organization of the Drosophila antennal lobe and

its regulation by the Teneurins. *Elife* **3**, e03726 (2014).

39. Howard, C. E. *et al.* Serotonergic Modulation of Walking in Drosophila. *Curr. Biol.*

29, 4218-4230.e8 (2019).

40.   Morano, N. C. *et al.* Cis inhibition of co-expressed DIPs and Dprs shapes neural

731   development. *BioRxiv* (2024).

732   41.   Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years

733   of image analysis. *Nat. Methods* **9**, 671–675 (2012).

734   42.   Eyjolfsdottir, E. *et al.* Detecting social actions of fruit flies. *Lect. Notes Comput.*

735   *Sci. (including Subser. Lect. Notes Artif. Intell. Lect. Notes Bioinformatics)* **8690**

736   **LNCS**, 772–787 (2014).

738
739
740

A.

Day 1   Day 2-5   Day 5-9   Adult

B.

20 hrs. APF   45 hrs. APF   60 hrs. APF   85 hrs. APF

αMNs Muscles

VNC   Myoblast fusion

αMNs

αTi-ltm   2° Bundles Form   MNs Mature
αTi-tadm

αMNs Muscles

Ti-ltm   Ti-ltm   Ti-ltm   Ti-ltm
tadm   tadm   talm   tadm   talm   tadm   talm

C.

VNC   DIP-α-A8-GAL4   Cuticle
T3   T2   T1
Femur   Femur   Tibia   Coxa   Trochanter   Tibia   150 μm

A.

DIP-α^{WT} DIP-α^{-50F} DIP-α^{NULL}

10^{-7} 10^{-6} 10^{-5} 10^{-4} 0

DIP-α^{-20F}

Affinity (μM)

B.

Control   DIP-α^{Null}

DIP-α-T2A-Gal4 | Cuticle

Tibia

αTi-ltm

αTi-tadm

50 μm

C.

ℓ_3

b_2 b_3

ℓ_2

b_1

b_4

ℓ_1 ℓ_4

$b_1 + b_2 + b_3 + b_4 + b_{n...} =$ Branch No.

$\ell_1 + \ell_2 + \ell_3 + \ell_4 + \ell_{n...} =$ Branch Length

D.

αTi-ltm   αTi-tadm

Branch No.

***   ***

Control α^{-20F} α^{-50F} α^{Null} Control α^{-20F} α^{-50F} α^{Null}

E.

αTi-ltm   αTi-tadm

Branch Length (μm)

***   ***   **   ***

Control α^{-20F} α^{-50F} α^{Null} Control α^{-20F} α^{-50F} α^{Null}

A. dpr10^WT dpr10^-20F dpr10^NULL

Affinity (μM)

10^-7 10^-6 10^-5 10^-4 0

dpr10^-8F dpr10^-40F

B.

Control | dpr10^Null

Tibia

DIP-α-T2A-Gal4 Cuticle
αTi-ltm
αTi-tadm

50 μm

C.

Branch No.

αTi-ltm | αTi-tadm

Control 10^-8F 10^-20F 10^-40F 10^Null

D.

Branch Length (μm)

αTi-ltm | αTi-tadm

Control 10^-8F 10^-20F 10^-40F 10^Null

A.

DIP-α^{+2F}

DIP-α^{+2F}::dpr10^{+10}(+15)

DIP-α::dpr10^{WT}

10^{-7} 10^{-6} 10^{-5} 10^{-4} 0

dpr10^{+10F}

Affinity (μM)

B.

Control   DIP-α+^{2F},dpr10^{+10F}(+15F)

Tibia

DIP-α-T2A-Gal4   Cuticle

αTi-ltm

αTi-tadm

50 μm

C. Branch No.

αTi-ltm   αTi-tadm

ns   *   *   ns   ****

Control   α^{+2F} 10^{+10F} α^{+2F} & 10^{+10F} (+15F)   Control   α^{+2F} 10^{+10F} α^{+2F} & 10^{+10F} (+15F)

D. Branch Length (μm)

αTi-ltm   αTi-tadm

ns   **   *   ns   ns   ***

Control   α^{+2F} 10^{+10F} α^{+2F} & 10^{+10F} (+15F)   Control   α^{+2F} 10^{+10F} α^{+2F} & 10^{+10F} (+15F)

Day 1   **Day 2-5**   **Day 5-9**   **Adult**

A.

	25 hrs. apf	35 hrs. apf	45 hrs. apf	55 hrs. apf	65 hrs. apf
Control					
DIP-α+²ᶠ & dpr10⁺¹⁰ᶠ(+15F)					

αTi-Itm
αTi-Tadm
50 μm

B.

C.

z-score

'Walk Frequency' 'Overall Velocity' 'Median Walking Velocity' 'Maximum Walking Velocity' 'Angular Velocity' 'Absolute Angular Velocity'

- Control
- DIP-α^{Null}
- DIP-α dpr10^{+15F}

ns ns * * * * ns ns ns ns * ns