Effects of parental care on skin microbial community composition in poison frogs

Marie-Therese Fischer[1,6*], Katherine S. Xue[1], Elizabeth K. Costello[2], Mai Dvorak[1], Gaëlle Raboisson[1], Anna Robaczewska[2], Stephanie N. Caty[1], David A. Relman[2,3,4], Lauren A. O'Connell[1,5]

[1] Department of Biology, Stanford University, Stanford, CA 94305, USA
[2] Department of Medicine, Stanford University School of Medicine, Stanford, CA 94305, USA
[3] Department of Microbiology & Immunology, Stanford University School of Medicine, Stanford, CA 94305, USA
[4] Infectious Diseases Section, Veterans Affairs Palo Alto Health Care System, Palo Alto, CA 94304, USA
[5] Wu Tsai Institute for Neuroscience, Stanford University, Stanford, CA 94305, USA
[6] Division of Microbial Ecology, University of Vienna, 1030 Vienna, Austria

Keywords: microbiome, vertical transmission, tadpole transport

*To whom correspondence should be addressed:
Marie-Therese Fischer
Department of Biology
Stanford University
371 Jane Stanford Way
Stanford, CA 94305
fi5cher@stanford.edu

Word count: Abstract: 330, Main text: 17437

Abstract

Parent-offspring interactions constitute the first contact of many newborns with their environment, priming community assembly of microbes through priority effects. Early exposure to microbes can have lasting influences on the assembly and functionality of the host's microbiota, leaving a life-long imprint on host health and disease. Studies of the role played by parental care in microbial acquisition have primarily focused on humans and hosts with agricultural relevance. Anuran vertebrates offer the opportunity to examine microbial community composition across life stages as a function of parental investment. In this study, we investigate vertical transmission of microbiota during parental care in a poison frog (Family *Dendrobatidae*), where fathers transport their offspring piggyback-style from terrestrial clutches to aquatic nurseries. We found that substantial bacterial colonization of the embryo begins after hatching from the vitelline envelope, emphasizing its potential role as microbial barrier during early development. Using a laboratory cross-foster experiment, we demonstrated that poison frogs performing tadpole transport serve as a source of skin microbes for tadpoles on their back. To study how transport impacts the microbial skin communities of tadpoles in an ecologically relevant setting, we sampled frogs and tadpoles of sympatric species that do or do not exhibit tadpole transport in their natural habitat. We found more diverse microbial communities associated with tadpoles of transporting species compared to a non-transporting frog. However, we detected no difference in the degree of similarity between adult and tadpole skin microbiotas, based on whether the frog species exhibits transporting behavior or not. Using a field experiment, we confirmed that tadpole transport can result in the persistent colonization of tadpoles by isolated microbial taxa associated with the caregiver's skin, albeit often at low prevalence. This is the first study to describe vertical transmission of skin microbes in anuran amphibians, showing that offspring transport may serve as a mechanism for transmission of parental skin microbes. Overall, these findings provide a foundation for further research on how vertical transmission in this order impacts host-associated microbiota and physiology.

Introduction

Host-associated microbial communities contribute to health and facilitate ecological adaptations by playing critical roles in host development, nutrition, pathogen exclusion, and immune response (Belkaid & Hand, 2014; Donald & Finlay, 2023; Gensollen et al., 2016; Mazmanian et al., 2005; Reynolds & Bettini, 2023; Sampson & Mazmanian, 2015; Von Frieling et al., 2018). Across host species, early exposure to microbes can exert lasting influences on the assembly and functionality of the host's microbiota (Al Nabhani & Eberl, 2020; Powell et al., 2014; Warne et al., 2017; Zepeda Mendoza et al., 2018). Timing of microbial contact is particularly critical in newborns and can prime community assembly through priority

effects, where species that arrive early impact establishment of later colonizers (Debray et al., 2022;
Fukami, 2015; D. Sprockett et al., 2018). In many species, the transmission of host-adapted microbes is
facilitated by parental care strategies, such as feeding, grooming and direct skin contact (for review see e.g.
S. Wang et al., 2020 for humans, Klug & Bonsall, 2014 for animals; see also Blyton et al., 2022; D. W.
Chen & Garud, 2022; Pascoe et al., 2017; D. D. Sprockett et al., 2020; Sylvain & Derome, 2017). To date,
most studies investigating effects of exposure to microbiota during parental care center on humans and
livestock, while vertical transmission remains largely unexplored in other classes of vertebrates (but see
Kouete et al., 2023). Therefore, investigating early-life microbial colonization in other species that have
dedicated parenting strategies offers opportunities to understand microbe-host interactions in a wider
variety of ecologically relevant contexts.

Acquisition of host-adapted skin microbes is especially important in amphibians, which rely on
healthy skin for physiological processes such as respiration, osmoregulation, immune response and barrier
function (Çömden et al., 2023; Duellman & Trueb, 1986; Varga et al., 2019). The amphibian skin
microbiota also serves to exclude pathogens, as the skin microbiome can confer resistance to infections by
the deadly chytrid fungus *Batrachochytrium* (Alexiev et al., 2023; Harris et al., 2006, 2009; Loudon et al.,
2014; Vredenburg et al., 2011; Woodhams et al., 2007), which decimates amphibian populations worldwide
(Daszak et al., 2003; Luedtke et al., 2023; Wake & Vredenburg, 2008). Many studies have characterized
microbe-host interactions in adult frogs in the context of chytrid infections, but few have focused on
microbiomes associated with tadpoles (Fontaine et al., 2022; Santos et al., 2023; Warne et al., 2017, 2019;
Weinfurther et al., 2023). Those studies that have looked at tadpoles have primarily characterized the
changes in bacterial communities across development and metamorphosis (discussed e.g. in Hughey et al.,
2017; Kueneman et al., 2014; Prest et al., 2018; Warne et al., 2017, 2019). Furthermore, most of this
research has focused on temperate-region, pond-spawning species, where opportunities to investigate
effects of parental care on community assembly are limited. These species generally lay many eggs at once
into bodies of water to avoid desiccation and larvae develop without further parental contact or care. In the
tropics, a warm and humid climate has favored the progression from aquatic to terrestrial reproduction,
where clutches are laid on land and parents then care for their offspring (McDiarmid & Altig, 1999; Wells,
2007). Neotropical frogs show more diverse reproductive strategies where parents construct foam nests,
attend clutches, defend eggs against predators, and guard, transport or feed their tadpoles (reviewed e.g. in
Schulte et al., 2020, see also Crump, 2015; Delia et al., 2013; Requena et al., 2009; Warkentin, 1995).
Priority effects influence community assembly of frog embryos (K. R. Jones et al., 2023, 2024) but previous
studies have found no direct evidence for vertical transmission of microbes during egg attendance (Hughey
et al., 2017). However, this form of offspring care does not involve direct contact between the parents and
offspring as developing tadpoles in the clutch are surrounded by several layers of gelatinous jelly and a

double vitelline envelope before they hatch (see Altig & McDiarmid, 2007 and Méndez-Tepepa et al., 2023

for a review). While the vitelline layer may serve as an antimicrobial barrier in chickens (Guyot et al., 2016;

Mann, 2008), studies of its role in microbial colonization of anuran embryos are currently lacking.

Parental care involving skin-to-skin contact has developed multiple times independently in

Neotropical dendrobatoid frogs (Superfamily *Dendrobatoidea*), commonly referred to as poison frogs

(Grant et al., 2017; Weygoldt, 1987). Closely related species in this clade display a variety of parenting

behaviors ranging from egg attendance and transport of hatched tadpoles to provisioning larvae with

unfertilized eggs (Crump, 1974; Ringler et al., 2023; Wells, 2007). The most widespread form of care

beyond clutch attendance is tadpole transport, where parents shuttle hatched tadpoles piggy-back style from

terrestrial clutches to water pools (Furness & Capellini, 2019). Transport can last multiple hours to days,

where tadpoles adhere tightly to the backs of their caregivers (Pašukonis et al., 2019; Pröhl & Berke, 2001;

Ringler et al., 2013, p. 201; Wells, 1980). Therefore, poison frogs provide the opportunity to evaluate the

effects of parental care on offspring skin microbial community assembly. The adaptive value of parental

care in this clade has received considerable attention (reviewed in Schulte et al., 2020; Summers & Tumulty,

2014), but how reproductive strategies involving skin-to-skin contact impact microbial colonization of

tadpoles and whether these effects persist to later developmental stages has, to our knowledge, not been

studied in any anuran to date.

The variable poison frog *Ranitomeya variabilis (Rv)* (Zimmermann & Zimmermann, 1988) is

particularly well-suited to study vertical transmission of microbes during tadpole transport. This small

species is diurnal, commonly found across South America and readily reproduces in captivity. Males of this

species often assist their larvae with hatching (Brown et al., 2008) and carry tadpoles on their back for up

to 48 hours (Lötters et al., 2007). The cannibalistic tadpoles grow up in individual pools where they feed

on algae and leaf debris until they complete their development after about five months. In the natural study

population located in the reserve 'Les Nouragues' in French Guiana, these frogs live on rocky outcrops, use

bromeliads as a resource for reproduction and shuttle tadpoles from the oviposition site in arboreal plants

to water reservoirs in leaf axils of terrestrial bromeliads (Poelman et al., 2013; Poelman & Dicke, 2007,

2008; Sarthou, 2001). This species is sympatric with other anuran species with differing parenting

strategies: the poison frog *Allobates femoralis* (*Af*) shuttles tadpoles to water but transporting periods are

shorter and tadpoles live in groups. The leptodactylid frog *Leptodactylus longirostris* (*Ll*) deposits eggs in

ephemeral rock pools and does not exhibit parental care. These cohabiting species are potentially useful for

determining if reproductive strategies impact patterns of microbial diversity of tadpoles.

In this study, we combine laboratory experiments with sampling of anuran populations in the wild

to examine vertical transmission of microbes during tadpole transport and address implications for

community structure across life stages. In the laboratory, we investigated microbial colonization of hatched
and unhatched *Rv* embryos and then tested the hypothesis that transporting frogs serve as a source for
bacteria on tadpole skins using cross-foster experiments. We complemented these experiments with a
comparative field study examining microbiomes associated with tadpoles and adults of three anuran species
with varying parenting behaviors to examine how shuttling affects community composition on the skin of
tadpoles in their native habitat. To our knowledge, this is the first study to document and characterize
vertical transmission of microbes during parental care in any anuran.

Results

Microbial colonization of a poison frog embryo occurs after hatching

Clarifying the timing of initial microbial colonization is essential for determining whether and how
priority effects mediate community assembly of vertically transmitted microbes in early life, or whether
these microbes arrive into an already established microbial landscape. We used non-sterile frogs of our
captive laboratory colony to determine the timing of microbial colonization of poison frog embryos.
Embryos within the vitelline membrane exhibit a characteristic C-shaped posture due to the membrane's
restriction on their movement and adopt a straight posture after hatching (Fig. S1A). First, we qualitatively
examined bacterial presence in *Rv* egg jellies and embryos across tadpole developmental stages using a
broad range PCR targeting the 16S rRNA gene. We manually separated embryos in the vitelline envelope
from the surrounding jelly and processed each for DNA extraction (Fig 1A). Bacteria were detected in egg
jellies across all developmental stages (formation of the dorsal lip to hatchling) but were not detected in
embryos prior to hatching from the vitelline membrane (N = 12) (Fig. S1B). After hatching, bacteria were
detected in tadpoles that remained in the jelly and in tadpoles that were transported by their caregiver (Fig.
S1B). Next, we confirmed differences in bacterial load between jelly and embryo by examining variations
in 16S rRNA gene copy numbers using a droplet digital PCR (ddPCR) approach suited for low biomass
samples (Abellan-Schneyder et al., 2021). We found that unhatched embryos contained on average 2942
copies of the 16S rRNA gene per µl (min = 742, max = 11584, std = ±3086) while jellies contained on
average 6128 times as many (min = 457417, max = 66115451, mean = 18029604 ±18586471; Kruskal
Wallis (KW): chi-squared = 18.6667, df = 2, p <0.00001) (Fig. 1B, Fig. S1C, Table S1). 16S rRNA gene
copy concentrations were on average 8310 times higher in hatched relative to unhatched embryos (min =
737340, max = 48159335, mean = 24448337 ±33532414; KW: p = 0.0195), but did not differ significantly
from concentrations detected in the jelly (KW: p = 0.5) (Fig. S1C, Table S1).

Figure 1. **Microbes colonize a poison frog embryo after hatching and are vertically transmitted during tadpole transport.** (**A**) Embryos in the vitelline envelope (dashed line) were manually separated from jelly using tweezers. The vitelline envelope (dashed line) containing the embryo was transferred to sterile water and opened to free the embryo. The embryo was washed in fresh sterile water before homogenization. (**B**) 16S rRNA gene copy number variations across developmental stages of embryos and jelly. DNA was isolated from whole embryos and jellies of different developmental stages and tested for bacterial presence using droplet digital PCR (ddPCR). Rounded mean copy numbers/µl/stage are displayed. (**C**) After hatching, siblings of a clutch were either (1) not transported (middle arrow), (2) transported by their biological parent (upper arrow), or (3) transported by a foster poison frog of a different species (*Oophaga. sylvatica*) (lower arrow). (**D**) We performed 16S rRNA gene specific amplicon sequencing targeting the V4 region on swabs from the transporting frogs and the skins of the transported tadpoles and used Sourcetracker to identify the sources of taxa that had been acquired by tadpoles. The function was trained on communities of adult *Rv* and *Os* that had served as caregivers. Source proportions of both species (Os: orange dots and Rv: blue dots) were determined for each tadpole (N = 24), resulting in 2 data points per tadpole. Source proportions (*Os*: orange dots, *Rv*: blue dots) were determined in tadpoles transported by *Os* (orange boxes), *Rv* (blue boxes) and non-transported tadpoles (grey boxes) and compared with a Kruskal-Wallis test.

Evidence for vertical transmission of microbiome during parental care in laboratory-reared poison frogs

We used a cross-foster design to test the hypothesis that frogs carrying tadpoles on their backs serve as a source of microbes for communities on tadpole skin. Siblings of the same clutch (N = 6 clutches from 2 *Rv* pairs) were randomly assigned to three groups: (1) not carried (N = 6 tadpoles), (2) carried on the back of their biological parent for six hours (N = 9 tadpoles), or (3) carried by a heterospecific poison frog (*Oophaga sylvatica, Os*) for six hours (N = 9 tadpoles) (Fig. 1C). For this design, sequence-based surveys of amplified 16S rRNA genes were used to assess the composition of skin-associated microbial communities from tadpoles and their adult caregivers (i.e., the frogs carrying the tadpoles, typically referred

to as "transporting" frogs). We found that bacterial community composition on caregiver skin was primarily
shaped by host species (Rv versus Os; Adonis: Bray-Curtis: $F_{1, 12}$ = 5.6538, R^2 = 0.33949, p = 0.001;
unweighted Unifrac: $F_{1, 12}$ = 2.0078, R^2 = 0.15435, p = 0.013), allowing us to distinguish between these
potential sources of tadpole-colonizing microbes (Fig. S2A). Indeed, we found that after six hours of being
carried on the back, bacterial community composition on Rv tadpole skin was influenced by caregiving
species (Rv versus Os; Adonis: Bray-Curtis: $F_{1, 12}$ = 1.75, R^2 = 0.099, p = 0.017; Unifrac: $F_{1, 12}$ = 1.67, R^2 =
0.095, p = 0.015). Despite their distinct compositions, we observed substantial overlap between
communities of tadpoles carried by Os and Rv (Fig. S2B), possibly reflecting similarities in tadpoles' skin
communities arising from a shared clutch environment after hatching until transport.

We used 'Sourcetracker' (Knights et al., 2011) to determine the relative contribution of the
transporting frogs as a source of bacteria detected in tadpoles' skin after six hours of transport. Tadpoles of
a clutch that were transported by Os always shared a higher proportion of their communities with this
heterospecific caregiver than siblings that were not transported (KW: chi^2 = 14.4696, df = 2, p = 0.0003;
difference in proportion: median = 39.33, min = 0.872 %, max = 68.478 %) or siblings transported by the
biological parent (KW: p = 0.015; percentage of higher proportion: median = 37.07 %, min = 0.874 %,
max = 72.97 %) (Fig. 1D, Table S2). Similarly, all but two transported tadpoles shared a larger proportion
of their communities with the transporting species than with the non-transporting species (p = 0.001) (Table
S2). Transported tadpoles (N = 18) generally shared higher proportions of their skin communities with the
transporting species than did non-transported tadpoles (N = 6) (Kruskal Wallis chi^2 = 19.4029, df = 2, p =
0.0001) (Fig. S2D). Non-transported tadpoles were colonized by between 5 and 57 Amplicon Sequence
Variants (ASVs) (median = 12). We conducted a separate analysis defining non-transported tadpoles as
potential microbe source to represent community proportions that were acquired in the clutch before
caregiver contact (Fig. S2C). We found that transported tadpoles shared between 0 and 46.99 % (median =
13%) of their communities with non-transported tadpoles. Sourcetracker detected higher proportions of
clutch-acquired microbes in tadpoles transported by Os (pairwise Wilcoxon test, p = 0.02). This likely
reflects the higher degree of similarity between the source clutch and Rv, as the clutch remained in the Rv
tank during development, making it harder to distinguish between these sources. Overall, our results suggest
that microbes are vertically transmitted during tadpole transport in a poison frog.

Tadpoles of the poison frogs *Ranitomeya variabilis* and *Allobates femoralis* host more diverse skin
communities than tadpoles of the leptodactylid frog *Leptodactylus longirostris*.

To investigate how vertical transmission of microbes during tadpole transport affects microbial
skin communities of tadpoles in an ecologically relevant context, we studied the skin microbiome
composition and diversity of Rv tadpoles and adults in a natural population. To assess the broader relevance

of our findings to frog species that transport their tadpoles, we also sampled two sympatric species with

differing reproductive strategies: *Allobates femoralis (Af)* and *Leptodactylus longirostris* (*Ll*) (Fig. 2A). In

contrast to *Rv* that transports its cannibalistic tadpoles to individual pools in bromeliads, *Af* lives in forest

leaf litter and shuttles multiple non-cannibalistic tadpoles to the same pool where they grow up

gregariously. *Ll* is a nocturnal leptodactylid frog inhabiting the rock savanna that does not exhibit tadpole

transport but deposits eggs directly in ephemeral rock pools where the larvae grow up together (Fig. S3).

We compared bacterial communities from 137 skin swabs of adult frogs, tadpoles, and the aquatic

environment belonging to the tadpoles of these three species. We sampled 44 adults and 21 tadpoles (Gosner

stages 29 - 41, categorized as "medium" and "large") of the species *Rv*, ten adults and 14 tadpoles (Gosner

stages 34 - 41, categorized as "medium" and "large") of the species *Ll*, and ten adult and eight tadpoles

(Gosner stage 25 - 26, categorized as "small") of the species *Af* (Table S3). Additionally, we monitored the

growth of 184 *Rv* tadpoles in separate bromeliad pools for up to 82 days. Out of 45 tadpoles that were

monitored for over 50 days, 13 (three "small" and ten "medium") tadpoles did not advance in their sizing

category, reflecting that the complete larval developmental in this population spans multiple months.

Cannibalistic *Rv* tadpoles grow up in isolated pools that were sampled as single replicates when collecting

the tadpole. Due to their development in groups, *Af* and *Ll* tadpoles were collected from fewer water bodies

and represent a narrower developmental window (stages 34 to 41 for *Ll* and stages 25 and 26 for *Af*). If

multiple tadpoles were collected from the same aquatic environment (*Af* and *Ll* tadpoles), we sampled

triplicates of the respective environment.

For each species, the aquatic environment of the tadpoles displayed a higher average phylum-level

diversity than the inhabiting tadpoles or adults (Table S3). As previously reported for Neotropical frogs

(Hughey et al., 2017; Kueneman et al., 2016), ASVs belonging to Phylum Proteobacteria dominated the

microbial community across all species and life stages (mean relative abundance: 70.9%), followed by

Bacteroidota (10.3%), Actinobacteriota (4.1%), Firmicutes (4%), Verrucomicrobiota (2.3%) and

Planctomycetota (1.1%) (Fig. S4). The abundance of the frog pathogen *Batrachochytrium dendrobatidis*

(*Bd*) in the rock savanna was low, with detection in two samples of *Rv* adults and one *Rv* tadpole, but no

positive *Ll* sample. We detected *Bd* on one adult *Af* individual collected in primary forest but not on tadpoles

of this species collected in the same area. In all species, tadpole communities contained fewer taxa with

known *Batrachochytrium*-growth inhibiting function than did environmental samples or adults (Table S3).

We found no significant differences in the observed ASV richness, diversity (Shannon), or

evenness between tadpoles and adults of the poison frogs *Rv* and *Af*. In contrast, *Ll* tadpoles were

significantly less diverse than *Ll* adults (Fig 2B, Table S4). We followed up on this finding by comparing

the diversity of tadpole-associated communities across species and found higher microbial diversity and

evenness in *Rv* and *Af* tadpoles compared to *Ll* tadpoles (Fig. 2C). Differences in community diversity of tadpoles were not a reflection of habitat diversity: microbial communities associated with *Rv* aquatic environments were less diverse and even compared to those in *Af* and *Ll* habitats (Fig. 2D). These variations may therefore reflect differences in life history traits among the three species.

Figure 2. Tadpoles skin microbiome is shaped by their environment and is more diverse in *Rv* and *Af* compared to *Ll* tadpoles. **(A)** We compared the skin microbiome of three anuran species: two species of poison frogs inhabiting different habitats that transport their offspring (*Rv* and *Af*) and a leptodactylid frog *(Ll)* that deposits its eggs in water without transporting the tadpoles. **(B)** Alpha diversity measures (observed ASV richness, Shannon diversity and evenness) for tadpoles (T) and adults (A) of each species were compared. Differences were determined with an ANOVA or Kruskal Wallis test, significance (p < 0.01) is indicated by **. **(C)** Comparison of ASV richness, Shannon diversity, and evenness of communities associated with poison frog tadpoles (*Af* or *Rv*) and non-poison frog species (*Ll*). **(D)** Comparison of Shannon diversity and evenness of communities associated with the aquatic habitats of *Af*, *Rv,* and *Ll*. Bars in boxplots represent median values. The dataset was separately rarefied to the lowest read depth of each comparison.

<u>Tadpole transport is not associated with higher degree of similarity between adult and tadpole skin microbiotas.</u>

To assess how microbes transmitted during parental care shape communities associated with tadpoles, we tested the hypothesis that tadpoles of the transporting species *Rv* and *Af* would have communities more similar to conspecific adults compared to tadpoles of the non-transporting species *Ll*. We first evaluated variation in bacterial communities associated with adults, tadpoles and their microenvironment by using a principal coordinate analysis (PCoA) to construct ordinations based on Bray-Curtis dissimilarities to explore factors that influence skin community composition (Fig. 3A). Tadpoles of all species clustered separately from adults, each other, and their aquatic environment (PERMANOVA on genus level, df = 8, R^2 = 0.52909, F = 18.117, p adj = 0.001) (Fig. 3A, B, Table S5). Consistent with findings in pond-spawning anurans (Kueneman et al., 2014, 2016; Prest et al., 2018), skin communities varied significantly across life stages in all species (PERMANOVA on genus level: species: df = 5, R^2 = 0.29674, F = 11.139, p = 0.001, life stage: df = 2, R^2 = 0.23148, F = 20.331, p = 0.001). Parental care also explained some of the observed variance (PERMANOVA on genus level: df = 2, R^2 = 0.07513, F = 5.4831, p = 0.001). To further examine patterns of similarities between life stages, we compared the core communities of adults, tadpoles and their aquatic environments across a range of prevalence and abundance cutoffs (prevalence 100% or 75%; relative abundance no cutoff, $\geq$ 0.1% (low) and $\geq$ 1% (high) (Table S6). Independent of parenting strategy and species, tadpoles shared core communities of their microbiome with their environments, but not with adult individuals. Adult poison frogs never shared core taxa with the aquatic environment of their tadpoles (Fig. 3C, Table S6), even though adults were commonly found in the nurseries. In contrast, the aquatic environment contributed substantially to the skin community of adult *Ll*, which shared high- and low-abundance ($\geq$ 0.1%) core genera with the pond water, but not with the pond-inhabiting tadpoles (Fig. 3C, Table S7).

Figure 3. Species, life stage and parental care affect clustering of microbial communities. (**A**) In a Principal Coordinate Analysis constructed with Bray-Curtis distances (axis 1 and 2 on the left, axes 2 and 3 on the right) tadpoles cluster significantly differently from each other, adults, and their aquatic environment. Significances were determined with a PERMANOVA followed by a pairwise Adonis posthoc test. Points in ordination plots represent the communities of each sample, circles represent confidence ellipses. (**B**) Principal Coordinate Analysis constructed with Bray-Curtis and Unifrac distances for adults, tadpoles, and aquatic environment of each species. (**C**) Number of core species (prevalence > 75%, relative abundance > 0.1%) shared between adults, tadpoles and the respective aquatic environment of each species.

To assess if microbes transmitted during tadpole transport constitute a stable part of skin communities associated with tadpoles, we compared community distances between skin communities of tadpoles and adults of the three frog species. We predicted that tadpoles and adults might have more similar skin microbiomes in the transporting species *Rv* and *Af* than in the non-transporting species *Ll*. We found no difference in community distances between tadpoles and adults in *Af* and *Ll* (iterated beta regression averaged over 999 random subsamples of similar sample size (IBR): Unifrac: $p = 0.3$). Communities associated with *Rv* tadpoles were more dissimilar from adult communities than microbiota of *Ll* tadpoles and adults (IBR: Unifrac: $p = 0.008$). Taken together, our results indicate that microbes acquired during tadpole transport do not create lasting similarities in composition between tadpoles and adults of wild populations.

After 4 weeks, transported *Ranitomeya variabilis* tadpoles retain isolated members of their caretaker's core community.

To test how offspring transport in *Rv* affected the community composition in tadpoles, we conducted a field experiment where we reared transported (N=10) and non-transported (N=10) tadpoles in artificial cups. Transported tadpoles were collected from the backs of their caregiver (N= 8), while we bypassed parental care in the non-transported group by transferring eggs to cups before the tadpoles hatched (Fig. 4A). Two out of ten non-transported tadpoles died before sampling. Richness and evenness of tadpole communities did not differ significantly between experimental conditions (Anova: Observed ASVs: df = 1, F = 2.7234, p.adj = 0.3552; Shannon: df = 1, F = 1.2067, p.adj = 0.8646; evenness: df = 1, F = 0.0009, p.adj = 2.9313), though PCoA on Unifrac distances revealed differences in microbiome composition attributable to the presence or absence of transport (Fig. 4B) (PERMANOVA: df = 1, R^2 = 0.11759, F = 2.1321, p = 0.002). Compared to transported tadpoles, non-transported tadpoles showed less overlap of genera in their microbial community with their aquatic environment (Fig. 4C). Specifically, the genera *Pelomonas, Rhodomicrobium* (Proteobacteria), *Mycobacterium* (Actinobacteriota) and *Candidatus Koribacter* (Acidobacteriota) primarily colonized transported tadpoles, while the genus *Limnohabitans* (Proteobacteria) was found on non-transported tadpoles. *Cetobacterium* and *Burkholderia-Caballeronia-Paraburkholderia* were found to colonize tadpoles from both experimental conditions, but at different abundances (ANCOMBC 2, p < 0.05) (Table S8).

Figure 4. Tadpole transport influences community structure. **(A)** Tadpoles collected from the back of their caregiver ("transported") and reared in artificial cups for one month were compared to six-week-old tadpoles that hatched from eggs transferred to artificial cups and did not experience transport by adult frogs ("non-transported"). **(B)** Principal Coordinate Analysis constructed using unweighted Unifrac distances, transported tadpoles cluster significantly differently from non-transported tadpoles. Significances were determined with a PERMANOVA followed by a pairwise Adonis post hoc test. **(C)** Venn diagram comparing unrarefied ASVs agglomerated on a genus level between transported tadpoles, non-transported tadpoles, and the aquatic environment. **(D)** Bubble diagram depicting the presence (circle) or absence (dot) of 10 ASVs shared between parents and transported tadpoles as well as their possible source ("transporting parent ", "nursery water", "refill water" or "untested"). Non-transported tadpoles that died prior to sampling are indicated by 'X'.

We further tested the hypothesis that related tadpole-adult pairs with a history of direct contact would have more similar communities than unrelated tadpole-adult pairs without such history. One month after transport, we found that skin communities of tadpoles were not more similar to the transporting parent than to unrelated adults of the population (iterated beta regression averaged over 999 random subsamples of similar sample size (N=10) (IBR): Unifrac: p = 0.27), confirming our results from the population sampling. However, a Sourcetracker analysis on unrarefied data revealed notable relative contributions of the parental skin community to the community on four transported tadpoles (tadpoles 1, 2, 6, 8) (min = 0% to max = 17%, median = 0%, IQR = 10.5) (Table S9). We followed up on this finding by directly identifying

taxa shared between caregivers and the respective transported offspring without imposing an abundance cutoff. We found that 8 out of 10 tadpoles shared one or two ASVs with the frog that had transported them in relative abundances ranging from 0.02 to 19.3 % (Fig. 4D) (Table S9). Shared taxa belong to the genera *Akkermansia, Elsteraceae, Aquitalea, Methylocella, Methylobacterium-Methylorubrum, Staphylococcus, Pseudomonas* and *Acinetobacter*. Seven out of ten shared microbes constituted members of the ten most abundant ASVs found on the respective transporting frog (Table S9). Absolute abundances of shared ASVs likely originating from the parental source pool (as identified by Sourcetracker) after one month of growth ranged from 7804 to 172326 sequence copies per tadpole (Table S10). Finally, we determined the success of shared ASVs in colonizing transported and non-transported experimental tadpoles. Even though at least six (min: 6, avg: 6.38, max: 7) of the ten ASVs were consistently present in the aquatic environment of non-transported experimental tadpoles, only one (family *Enterobacteriaceae*) successfully colonized their skin (Fig 4D, not transported tadpoles) (Table S11).

Discussion

In this study we aimed to understand the main factors influencing the establishment of microbial communities associated with anuran tadpoles receiving parental care. To better understand sources contributing pioneering taxa, we first aimed to determine the ontogenetic window in which anuran embryos are colonized. Next, we tested if a specialized anuran parenting strategy, tadpole transport, transmits early colonizers. Then, we sampled natural populations of frogs to investigate if parental signatures persist into later ontogenetic stages in an ecologically relevant context. We found that substantial microbial colonization of the poison frog *Rv* occurs after embryos hatch from the vitelline envelope. Furthermore, we demonstrated through cross-fostering experiments that transporting frogs serve as a source of skin microbes for tadpoles. Next, we sampled sympatric species in a Neotropical habitat and revealed that tadpoles of poison frogs showed more diverse microbial communities than tadpoles of leptodactylid frogs. Finally, a field experiment in the wild population indicated that microbial communities of *Rv* tadpoles after four weeks of growth were not more similar to transporting caregivers than to unrelated adults, although some tadpoles did maintain isolated ASV that were shared with the transporting adults. Together, our results demonstrate that microbial colonization of a poison frog tadpole begins at an ontogenetic stage immediately preceding parental contact, that parental care in poison frogs facilitates the transfer of microbes to newly hatched tadpoles, and that signatures of these parentally transmitted microbes can persist, albeit at low prevalences, through the first month of tadpole development. This study shows that tadpole transport may serve as a mechanism to transmit host-adapted microbes, thereby filling a knowledge gap in our understanding of the function of parental care in anurans.

Early microbial colonizers have been shown to shape community structure of newborns in insects
(E. W. Jones et al., 2022), mammals (Blyton et al., 2022; Kaur et al., 2022; Sprockett et al., 2018), fish (
Almany, 2004), birds (Chen et al., 2020) and anurans (K. R. Jones et al., 2023, 2024) via priority effects
(see Debray et al., 2022; Fukami, 2015 for reviews). For example, inoculation of embryos has been shown
to increase the relative abundance of beneficial taxa such as *Janthinobacterium lividum* in tadpoles (Jones
et al., 2024), whereas efforts to introduce the same organism into established adult communities have not
led to long-term persistence (Bletz, 2013; Woodhams et al., 2016). Thus, determining critical
developmental periods for microbial colonization of host tissue is crucial to assess which sources contribute
these pioneering taxa. We detected microbes in embryos of all stages but found the abundance to be
considerably lower than in the surrounding jelly. Additionally, we found that substantial colonization of *Rv*
tadpoles happens after hatching from the vitelline membrane, at an ontogenetic stage immediately
preceding skin-to-skin contact with the caregiver. These results underscore that the opportunistic
colonization of the anuran embryo is much more restricted than that of the surrounding jelly (Hayes et al.,
2009; Kueneman et al., 2014; Walke et al., 2011), which harbors stable microbial assemblages (Hughey et
al., 2017; Walke et al., 2011). Such restrictions might result from host-led selection as observed in marine
mammals (Switzer et al., 2023) and/or a barrier function of the vitelline membrane that limits microbial
access to the developing embryo. For example, the latter was described for the avian vitelline membrane,
where it is mediated by antimicrobial proteins (Guyot et al., 2016; Mann, 2008; Mine & Kovacs-Nolan,
2006). The substantial microbial colonization that occurred after hatching suggests that the anuran vitelline
envelope plays an important role in restricting microbial access to poison frog embryos until they are
sufficiently developed to be transported by a caregiver. Those bacteria that are present in unhatched tadpoles
might have been acquired prior to hatching, such as in the case of the amphibian symbiont *Oophila*
amblystomatis (Kerney et al., 2011; Kim et al., 2014)*,* which is acquired by oviductal transmission and
grows on the inside of individual salamander and frog vitelline membranes. Alternatively, the vitelline
membrane may function as a selective barrier, permitting the passage of certain taxa, with hosts acting as
additional filter to select suitable colonizers. Future studies should therefore address the presence of
antimicrobial proteins, biofilms, and immune components in the anuran embryo and vitelline membrane to
shed light on the mechanisms that regulate microbial colonization of anuran embryos.

Across species, newborns might acquire bacteria not only through transfer from environmental
source pools and other hosts (e.g. arthropods: Douglas, 2018; Mushegian et al., 2018, amphibians: Rebollar
et al., 2016, mammals: Bik et al., 2016) but also through vertical transmission during parent-offspring
interactions (reptiles: Bunker et al., 2021; fish: Sylvain & Derome, 2017; humans: Wang et al., 2020).
Modes of parental inheritance involve indirect transmission, like contact smearing onto the egg surface

during or after oviposition, as found in European firebugs (Salem et al., 2015), and direct social acquisition,

as observed in social bees (Kwong et al., 2017; Leftwich et al., 2020). Previous studies in anurans have

investigated vertical transmission of microbes during parental care in the context of indirect transmission

during egg attendance without considering less widespread parenting strategies involving direct parental

contact. These studies did not find evidence for parental inoculation, as hatched tadpoles supported bacterial

communities very similar to adult frogs but independent of caregiver identity (Hughey et al., 2017). In

contrast, we investigated vertical transmission in an anuran system where parenting involves direct skin-

to-skin contact between tadpoles and their caregiver. We found that both transported and non-transported

tadpoles in our cross-foster experiment shared microbial ASVs with each other, indicative of acquisition of

microbes from the jelly upon hatching as demonstrated for other anuran clades (Hughey et al., 2017; Jones

et al., 2024; Warne et al., 2017). After six hours of contact, transported tadpoles -but not their siblings-

shared more microbes with the surrogate heterospecific frog than with the parental species, suggesting that

microbes can transmit vertically during parental care in poison frogs. Thus, in this clade both clutch

environment and transporting caregivers can serve as source pool for hatched tadpoles.

Most poison frogs transport tadpoles on their backs, but the mechanism of adherence remains

unclear. Similar to natural conditions, tadpoles that are experimentally placed onto a caregiver's back also

gradually adhere to the dorsal skin, where they remain firmly attached for several hours as the adult

navigates dense terrain. Although transport durations were standardized, species-specific factors- such as

microbial density at the contact site, microbial taxa identity, and skin physiology such as moisture -might

influence microbial transmission between the transporting frog and the tadpole. While these differences

may have contributed to varying transmission efficacies observed between the two frog species in our

experiment, none of these factors should compromise the correct microbial source assignment. We thus

conclude that transporting frogs serve as a source of microbiota for transported tadpoles. However, further

studies on species-specific physiological traits and adherence mechanisms are needed to clarify what

modulates the efficacy of microbial transmission during transport, both under experimental and natural

conditions. For example, evidence for duration-dependent microbial transmission is seen in humans, where

recent research highlights a cumulative contribution of the paternal microbiome to the assembly of infants'

microbial communities (Dubois et al., 2024). In anurans, the time that hatched tadpoles spend in contact

with jelly-associated microbes or in skin-to-skin contact with parents might therefore affect the success and

stochasticity of colonizers that are transmitted during parental care. These transport logistics vary between

and within anuran species (Altig & McDiarmid, 2007; McDiarmid & Altig, 1999) and are closely linked to

trade-off decisions of the caregivers, which often need to balance territory defense, mating, and caring for

offspring (Pašukonis et al., 2019; Ringler et al., 2013; Wells, 2007). For instance, *Rv* males often assist

tadpoles with hatching by tearing the capsule apart with their rear legs to initiate transport (Brown et al., 2008), but have also been observed to pick-up free-swimming tadpoles (Schulte & Mayer, 2017). Likewise, *Af* tadpoles have been observed to remain in the jelly for several days before being transported (Peignier et al., 2022; Ringler et al., 2013). Parentally transmitted microbes might thus constitute pioneering taxa in an unoccupied niche if caregivers assist with hatching or encounter established assemblages as in the latter scenario. Prolonged parental care has been shown to enhance the generational transmission of symbiotic microorganisms in burying beetles (e.g. Körner et al., 2023) and thus, duration of the skin-to-skin contact during tadpoles transport might influence microbial acquisition in a time-dependent manner. How duration of tadpole transport, which ranges from a few hours to several days depending on the species (Lötters et al., 2007; McDiarmid & Altig, 1999; Pašukonis et al., 2019), affects stochasticity of transmitted microbes remains unexplored. Collectively, our findings suggest that early life microbes are introduced to the poison frog host in a mixed transmission mode, where community members can be sourced environmentally from the egg jelly and socially from a caregiving frog. Recent evidence from Kouete et al. documented the first case of vertical transmission in amphibians, observed in subterranean caecilians that protect and feed their offspring (Kouete et al., 2023). To our knowledge, ours is the first study that extends evidence for vertical transmission during parental care to the order of anurans.

To investigate if parental signatures persist into later ontogenetic stages, we tested the hypothesis that a history of direct contact would result in more similar skin communities between tadpoles and caregivers relative to unrelated adults. Based on previous findings which documented that priority inoculation of tadpoles increased the relative abundance of some beneficial inoculates (Jones et al., 2024), we hypothesized that vertical transmission manifests in persistent associations with parentally acquired microbes. Similar to Hughey et al., we found that vertical transmission did not lead to a higher degree of similarity between caregiver and tadpole skin microbiota relative to unrelated adult-tadpole pairs. Likewise, communities associated with adults and tadpoles of transporting species were no more similar than those of non-transporting species. While poison frog tadpoles do acquire caregiver-specific microbes during transport, most of these microbes do not persist on the tadpoles' skin long-term. This pattern can likely be attributed to the capacity of tadpole skin- and gut microbiota to flexibly adapt to environmental changes (Emerson & Woodley, 2024; Santos et al., 2023; Scarberry et al., 2024). It may also reflect the limited compatibility of skin microbiota from terrestrial adults with aquatic habitats or tadpole skin, which differs structurally from that of adults (Faszewski et al., 2008). As a result, many transmitted microbes are probably outcompeted by microbial taxa continuously supplied by the aquatic environment. Interestingly, microbial communities of the non-transporting *Ll* were more similar to their adult counterparts than those of poison frogs. This pattern might reflect differences in life history among the species. Adult *Ll* commonly inhabit the rock pools where their tadpoles develop, but adults of the two poison frog species visit tadpole nurseries

only sporadically for deposition. These differences in habitat use may result in adult *Ll* hosting skin
microbiota that are better adapted to aquatic environments as compared to *Rv* and *Af*. Additionally, their
presence in the tadpoles' habitat could make *Ll* a more consistent source of microbiota for developing
tadpoles. Transient shifts in the composition of tadpole-associated communities followed by later
convergence have also been observed in wood frogs experimentally inoculated with bullfrog gut microbes
(Warne et al., 2019). Notably, despite this compositional convergence, effects of early microbial disruption
on host physiology and disease susceptibility persisted- an outcome also documented in fruit flies (Cox et
al., 2014). Therefore, it is possible that vertically acquired microbes shape communities of poison frogs in
different ways than by gaining abundance priority on tadpole skin. For example, they may durably imprint
the immune system (Fallet et al., 2022; Gensollen et al., 2016) and alter host metabolic phenotypes (Cox et
al., 2014; Sommer & Bäckhed, 2013). Additionally, isolated ASVs contributing to central genera of
communities on adults persisted on some tadpoles in varying abundance. There is clear evidence that low-
abundant keystone taxa, despite their rarity, can act as drivers of compositional changes (Han & Vaishnava,
2023). For instance, low abundance ASVs drive compositional changes in the hindgut of subterranean
termites after dietary alterations (Benjamino et al., 2018) and confer resistance to Salmonella-induced
colitis in mice (Herp et al., 2019). Moreover, despite their abundance in the water of all tadpole nurseries,
most of the ASVs that tadpoles shared with caregivers failed to colonize non-transported hosts. These
results suggest that vertical transmission may be a more effective mode of transmission of host-adapted
microbes compared to horizontal transmission from the environment. Whether vertically transmitted
microbes shape the anuran immune response and if retaining certain ASVs allows them to become dominant
community members after metamorphosis remains to be investigated.

We found that the higher diversity of communities associated with tadpoles of species that exhibit
offspring transport was not linked to the diversity found in tadpoles' aquatic habitat but might instead be
associated with other life history traits. Although skin alkaloids are found on many adult poison frogs
including *Rv* and influence skin community composition (Caty et al., 2024), they are not present on adult
Af or on tadpoles of any species included in this study (Villanueva et al., 2022) and are thus unlikely to
cause the observed differences in diversity. It is possible that oviposition strategy (*Af* and *Rv* lay terrestrial
egg clutches whereas *Ll* lays eggs directly in water) or physiological characteristics like skin pH or
abundance of antimicrobial proteins (Conlon, 2011; Faszewski et al., 2008; Kueneman et al., 2014)
contributed to increased diversity in *Rv* and *Af* relative to *Ll* tadpoles. Alternatively, transport might have
been co-opted in an evolutionary context to increase microbial diversity in tadpoles and shape immune
recognition and health in the anuran host in a manner comparable to humans (Al Nabhani & Eberl, 2020;
Donald & Finlay, 2023; Gensollen et al., 2016). While increased microbial diversity is not inherently

advantageous, it has been associated with beneficial outcomes such as improved immune function, lower disease risk, and enhanced fitness in multiple other vertebrate systems. For example, higher diversity of gut microbial communities is associated with enhanced exploratory behavior in songbirds (Florkowski & Yorzinski, 2023) and improved pathogen resilience in fish (de Bruijn et al., 2018). Previous research in frogs has linked higher diversity of tadpole skin communities to enhanced parasite resistance later in life (Knutie et al., 2017; Warne et al., 2019). However, increased microbial diversity is not a known outcome of vertical transmission, and further studies across a broader range of transporting and non-transporting species are needed to assess the role of transport in shaping diversity of tadpole-associated microbial communities. Moreover, the low chytrid infection-rate in the studied natural populations limits the ability to draw conclusions about differences in disease susceptibility within and between species or life stages in our dataset.

Elucidating the key factors that select for parental care and account for the diversity of care strategies across various species remains a challenge in the field of evolutionary and behavioral ecology (Balshine, 2012; Clutton-Brock & Scott, 1991; Gross, 2005; Royle et al., 2012). Microbiota management is documented to actively drive, rather than merely accompany sociality and parenting behavior in other organisms like carrion feeding insects (Biedermann & Rohlfs, 2017; Körner et al., 2023) and has recently been proposed as a factor driving the evolution of paternal care in vertebrates (Gurevich, 2020). Even though commensal skin microbiota undoubtably play a pivotal role in the health and pathogen defense of amphibians, it remains poorly understood how complex parental care strategies in this class aid microbial transmission. This is especially surprising as amphibians have recently been identified as the most threatened class of vertebrates (Daszak et al., 2003; Luedtke et al., 2023; Wake & Vredenburg, 2008) due to their susceptibility to globalization-related threats like climate change and disease-spread. Our observation that anuran parental care transmits microbes fills a knowledge gap about the function of parental care in this class and highlights the need for further studies across more anuran species with contrasting reproductive strategies.

Methods

Captive-bred animals

Captive-bred *Rv* of our laboratory colony are kept non-sterile and reproduce year-long. Males attend the terrestrial clutch of 3-5 eggs and transport larva to water when they reach Gosner stage 24-25 where they hatch from the vitelline membrane. *Rv* were housed in pairs and provided with horizontally mounted film canisters as egg deposition sites, and film canisters filled with water (treated with reverse osmosis R/O Rx, Josh's Frogs, Owosso, MI) for tadpole deposition (Goolsby et al., 2023).

 <u>Microbial colonization of embryos and egg jelly</u>

We used individually packed sterile transfer pipets (Samco™, Thermo Scientific, 7.7 mL, cat # 202-1SPK)
with a cut tip to move eggs and embryos. All tools were treated with 8.25% Sodium Hypochlorite
(CloroxPro) and wiped down with 70% Ethanol before use and between tissues and tadpoles. The
experimenter always wore gloves that were changed between individuals and disinfected with 70% Ethanol
before handling frogs.

Triplicates of five different developmental stages (Gosner, 1960) before (stages 10, 19, 23, 24) and after
hatching from the vitelline membrane (stages 24, 25) were collected from 3 different breeding pairs. We
manually separated the jelly from the egg sac with autoclaved forceps, a procedure commonly done in
preparation for microinjections. The embryo in the vitelline membrane was transferred to a small petri dish
with sterile, filtered deionized water and opened using forceps. Tadpoles were transferred to a new petri
dish with clean water twice to remove transient bacteria from the skin and were then euthanized and
homogenized. Organic matter was removed from the jelly, and both jelly and tadpole tissues were flash
frozen and stored at -80 °C until DNA isolation. Isolation of DNA was performed as detailed below, and 1
µL of isolated DNA was used for PCR amplification of the full length 16S rRNA region (~1500 bp) with
1 µL of 10 µM forward 27F (5'-AGAGTTTGATCMTGGCTCAG-3') and reverse primer 1492R (5'-
TACGGYTACCTTGTTAYGACTT-3') (Fredriksson et al., 2013), 12.5 µL of OneTaq Hot Start Quick
Load polymerase (New England Biolabs) and 9 µL water. We followed an amplification protocol with
denaturation for 30 s at 94 °C followed by 30 cycles of denaturation for 15 s at 94 °C, annealing for 30 s at
58 °C and extension for 2 min at 68 °C and a final extension period of 5 min at 68 °C. To determine the
presence or absence of amplification, we visualized 5 µL of each PCR product on a 1% Agarose gel (90mV
for 90 min) with 2 µL of 1kb GeneRuler for sizing.

Quantitative analysis of 16S rRNA copy numbers with droplet digital PCR (ddPCR)

Quantification of prokaryotic concentration for each sample was determined by ddPCR for all samples
(Doyle et al., 2025). Sample DNA was serially diluted 1:10 in sterile nuclease-free water to dilute the
concentration of any PCR inhibitors using a liquid handler (Agilent, Agilent Velocity 11 Vprep). 16S rRNA
concentrations from embryonic tissue are reported from undiluted DNA as its concentration was
approaching the limit of detection in diluted reactions. Universal 16S rRNA primers (331F/797R) and 16S
rRNA HPLC-purified FAM probes were used, as previously described (Langenfeld et al., 2021). Primer-
probe mixture was created by mixing forward primer, reverse primer, and probe 1:1:1 for a total volume of
0.264uL per reaction. Each ddPCR reaction was composed of 11uL of ddPCR Supermix for probes (Bio-

Rad, 1863024), 0.264uL of primer-probe mixture, 4.736uL of sterile nuclease-free water and 6uL of sample, for a total of 22uL/reaction. As a positive control, equal amounts of NIST Microbial Pathogen DNA Standards for Detection and Identification (NIST, RM8376) components A through R were combined in equal amounts. Each ddPCR plate included a positive control, NIST mixture, and negative control, sterile nuclease-free water, in quadruplicate. ddPCR reactions were performed with the QX200 AutoDG Droplet Digital PCR system (Bio-Rad). PCR amplification was performed with the Bio-Rad T100 thermocycler using the following program: 95°C for 10 minutes, 40 cycles of 95°C for 30 seconds and 56°C for 1 minute and 72°C for 2 minutes, followed by 1 cycle of 4°C for 5 minutes and 95°C for 5 minutes with a ramp speed of 2°C/second at each step. Amplified reactions were quantified using a ddPCR reader. Thresholds were set for each ddPCR reaction based on the negative and positive control rain plots generated by the QX Manager Software, 2.1 Standard Edition (Bio-Rad). All reactions had greater than 15,000 accepted droplets. 16S rRNA copies per droplet were calculated by taking the natural logarithm of the ratio of accepted droplets to negative droplets for each reaction. Values were corrected for droplet volume and ddPCR reaction volume to calculate total 16S rRNA copies per reaction. We further calculate and report the total 16S rRNA copies per μL by accounting for the sample ddPCR volume, dilution factor, and DNA extraction volume. Differences in copy number concentrations detected in jellies, hatched and unhatched tadpoles were determined using a Kruskal Wallis test with p-values adjusted for multiple comparisons using Benjamini Hochberg correction (R Core Team, 2023) and followed by a Dunn test.

Vertical transmission of bacteria

Cross-fostering tadpoles onto non-parental frogs has been used previously to study navigation in poison frogs (Pašukonis et al., 2017). According to our experience, successful adherence to both parent and heterospecific frogs depends on the developmental readiness of tadpoles, which must have retracted their gills and be capable of hatching from the vitelline envelope through vigorous movement. Another factor influencing cross-fostering success is the docility of the frog during initial attachment, as erratic movements easily dislodge tadpoles before adherence is established. *Rv* are small, jumpy frogs that are easily stressed by handling, making experimental fostering of tadpoles- even their own- impractical. Therefore, we favored an experimental design where tadpoles initiate natural transport, and parental frogs pick them up with a 100% success rate. We chose the poison frog *Os* as foster frog because adults are docile, parental care in this species involves transporting tadpoles, and skin microbial communities differ from *Rv*- a critical prerequisite for our Sourcetracker analysis. The use of the docile *Os* as the foster species enabled a 100% cross-fostering success rate, with no notable differences in adherence strength of tadpoles after six hours.

Siblings of one clutch were randomly assigned to one of three treatments: (1) no transport, (2) transport by

the biological caregiver, or (3) transport by the heterospecific surrogate frog *Os*. Tanks of *Rv* breeding pairs

were monitored for clutches of at least three eggs daily. Embryos assigned to group one or three were placed

into a petri dish in a separate tank without access to adults when tadpoles reached a developmental stage

close to hatching (Gosner stage 22) to bypass transport by the biological father. The remaining eggs in the

parental tank were monitored with security cameras (Wyze v3, Goolsby et al., 2023) that notify the user of

any movement in the canister. After tadpole pickup by the biological father, all water canisters for tadpole

deposition were removed from the tank to standardize transport time to 6 hours for both groups. Siblings

of group three were transferred to a new petri dish, hatched using a sterile brush and directly transferred to

the back of a surrogate frog (as described in Pašukonis et al., 2017) that was caught in a fresh plastic bag

and rinsed with sterile water. The surrogate frog was placed into a plastic containment (Sterilite 16428012)

with moist, autoclaved paper towels for the duration of the experiment. After 6 hours, the transporting

biological father and the surrogate frog were caught in a fresh plastic bag. Each tadpole was removed from

the back and washed twice to remove transient bacteria by transferring them to new petri dishes with 10 ml

sterile water. Non-transported tadpoles were removed from the jelly and washed in a similar way. Each frog

was rinsed with sterile water, and a skin swab of their back was collected using a sterile Puritan applicator

(Puritan Medical Products, 25-206 1PD BT) (Caty et al., 2024). All tadpoles were euthanized, their skin

was collected, and skin and applicators were stored at -20 °C until DNA extraction. We chose to sample

whole skin instead of swabs to detect small amounts of transferred bacteria on and in the skin.

<u>Study species, reproductive strategies and life history</u>

Oophaga sylvatica (Os) (Funkhouser, 1956; CITES Appendix II, IUCN Conservation status: Near

Threatened) is a large, diurnal poison frog (Family *Dendrobatidae*) inhabiting lowland and submontane

rainforests in Colombia and Ecuador. While male *Os* care for the clutch of up to seven eggs, females of

this species transport 1-2 tadpoles at a time to water-filled leaf axils where tadpoles complete their

development (Pašukonis et al., 2022; Silverstone, 1973; Summers, 1992). Notably, females return

regularly to these deposition sites to provision their offspring with unfertilized eggs.

Ranitomeya variabilis (Rv) (Zimmermann & Zimmermann 1988; CITES Appendix II, IUCN Conservation

status: Least Concern) is a small diurnal poison frog (Family *Dendrobatidae*) with male uniparental care

that lives largely arboreal and uses bromeliads as a resource for reproduction. Adult frogs defend territories,

are polygamous and typically lay clutches of 3-4 eggs into small arboreal bromeliads (*Catopsis*

berteroniana) that are abundant on Clusia trees in the rock savanna. Male frogs transport hatched tadpoles

to individual water bodies that form in the leaf axils of the large terrestrial tank bromeliads (*Aechmea*

aquilega). The cannibalistic tadpoles grow up in individual pools of ~80 ml (Poelman et al., 2013) where

they feed on algae and leaf debris until they complete their development after about 3 - 5 months (Poelman
& Dicke, 2007).

Allobates femoralis (Af) (Boulenger, 1884; CITES Appendix II, IUCN Conservation status: Least Concern)
are polygamous diurnal poison frogs (Family *Aromobatidae*) that inhabit the understory of primary tropical
forests. Females are attracted by advertisement calls of the territorial males and deposit clutches of 11-25
eggs in the leaf litter (Fischer et al., 2020; Stückler et al., 2019; Weygoldt, 1980). Fathers care for their
offspring by attending clutches and shuttle hatched tadpoles to water bodies where the social tadpoles grow
up sharing a pool (Lescure, 1976; Ringler et al., 2013, 2018).

Leptodactylus longirostris (Ll) (Boulenger, 1882; IUCN Conservation status: Least Concern) are medium
sized nocturnal leptodactylid frogs (Family *Leptodactylidae*) that inhabit the rock outcrops. They shelter in
rock crevices and ground covering vegetation during the day and dwell in seasonal rock pools on the
Inselberg plateau during nights. They breed in the rainy season and lay their eggs in ephemeral rock pools.
After clutch deposition, they do not care further for their tadpoles, which collectively grow up in the
deposition pond and feed on algae and plant debris.

Field site and natural populations

The natural study site of *Rv* and *Ll* is situated on top of the mountain 'Inselberg' in vicinity to the Centre
Nationale de la Recherche Scientifique managed research station (4°5' N, 52°41' W) within the Nature
Reserve ´Les Nouragues´ in French Guiana (Bongers et al., 2011). Patches of Clusia trees are separated by
bare granite rocks and exposed to extreme environmental conditions (e.g. temperature oscillations between
18°C - 75°C (Sarthou, 2001). *Rv* and *Ll* (IUCN Conservation status: Least Concern) inhabit the granite
outcrop of the study site, *Af* inhabit forest floors of primary terra-firme forests surrounding the campsites
'Inselberg' and 'Saut Pararé' (4°02'N/52°41'W) in the Nouragues reserve.

Sampling of frogs in the field

Frogs were caught in fresh plastic bags upon encounter, rinsed with sterile water to remove transient
bacteria and soil particles and then swabbed with 20 strokes each on the dorsal and lateral sides as well as
on each leg and between the toes using Sterile Polyester Applicators (Puritan Medical Products, 25-206
1PD BT). Due to limitations in the availability of dry ice at the remote study site, applicators were directly
transferred into 800 µL of buffer CD1 (a lysis buffer containing chaotropic salts) of the Qiagen PowerSoil
Pro extraction kit and frozen until DNA extraction. Previous studies have found no difference in OTU
richness and evenness between native frozen and lysis buffer-stored swabs (Hallmaier-Wacker et al., 2018).

    Field sampling of tadpoles and their environment

To avoid disturbing microbial communities of tadpoles and their habitats before sampling, free-swimming

tadpoles were visually categorized as "small" (with a body length of less than 6 mm, typically Gosner stages

25 and 26), "medium" (with a body length exceeding 6 mm but no visible dorsal pattern, stages 26 - 33),

"large" (visible yellow dorsal pattern, stages 34 - 41) or "metamorph" (front limbs visible, stages 42 - 46).

Tadpoles of stages 29 – 41 (Gosner) were collected from bromeliad pools for swabbing when they at least

doubled the body size of freshly hatched tadpoles, at the earliest four weeks after deposition. The water and

sediment of the nursery was sampled prior to tadpole extraction by submerging a sterile applicator tip into

the leaf axil pool and moving it over all plant material contained in the water for 10 seconds. *Rv* tadpoles

were extracted from bromeliad leaf axils with a custom designed vacuum extractor consisting of an inlet

hose wide enough to allow the passage of a late-stage tadpole and a suction hose with air filter, both tightly

connected to the lid of a one-liter bottle. The inlet hose was submerged into the bromeliad pool while

applying suction to the second hose to create a vacuum that transferred the pool content into the collector.

After each use, all parts of the collector were cleaned with soapy water and rinsed, followed by a sodium

hypochlorite disinfection for 1 hour and a rinse in 70% Ethanol as described for the dissection tools. After

collection, tadpoles were transferred to a petri dish with sterile water twice, using individual sterile transfer

pipettes with a cut tip. For swabbing, tadpoles were collected in the bulb of a new sterile pipette. The water

was discarded, and the tadpole was swabbed on all body parts with a sterile applicator tip by moving the

tip over its body for 10 seconds. We further processed swabs in the same way as adult swabs. *Af* tadpoles

of stage 25 - 26 (Gosner) were collected from an artificial pool in the vicinity of camp 'Saut Parare'. *Ll*

tadpoles of Goser stage 34 - 41 were collected from three natural pools in the study plot 'Inselberg'. We

sampled triplicates of each pool containing multiple tadpoles. All tadpoles were processed for swabbing in

a similar way as *Rv* tadpoles.

Field experiment to compare transported and non-transported *Rv* tadpoles

To directly compare parental frogs with transported and non-transported tadpoles while reducing predation

and variation in the quality of their nurseries, we reared ten tadpoles that were collected from the back of

eight caregivers and ten tadpoles that hatched from eggs without experiencing transport in polypropylene

cups (USP #77876, 60 ml). Each cup was punctured at the 50 ml water level, covered with a sterilized

mosquito net secured with a rubber band to minimize predation, and attached to the stem of a Clusia tree.

All tadpoles received water collected from bromeliads as their initial aquatic environment with Clusia

leaves as shelter and food. Cups were exposed to rain and during dry periods, rainwater was collected and

mixed with leaf litter to refill cups. Triplicates of water samples were obtained using the procedure

described for bromeliad water. We caught transporting frogs upon encounter, transferred tadpoles from
their backs to cups, and swabbed the parenting frog as described above. To avoid transport, eggs from 5
clutches were transferred to cups before tadpoles hatched. Tadpoles were sampled after growing for at least
26 days (min: 26, max: 77), when all tadpoles of a group exceeded ~5 mm in body size, as smaller
individuals exhibited increased mortality after swabbing in previous lab experiments. After swabbing,
tadpoles were measured and photo documented.

DNA extraction and 16S rRNA gene sequencing

The Qiagen PowerSoil Pro Kit was used to extract DNA from all swabs and tissues. The protocol was
adapted for use with swabs as described previously (Caty, Alvarez-Buylla, et al., 2024). DNA
concentrations were quantified using a Qubit. Samples were pooled by volume and the V4 region of the
16S rRNA gene was amplified using 515F (GTGYCAGCMGCCGCGGTAA) and 806R
(GGACTACNVGGGTWTCTAAT) primers (Bletz et al., 2017) and barcoded using standard Illumina
unique dual indices (UDIs). We performed two separate sequencing runs: Laboratory collected samples
were sequenced in a 2x300nt paired-end configuration on an Illumina MiSeq v3 run, field collected samples
in a 2x250nt paired-end configuration on a NovaSeq 6000 SP Flowcell (Roy J Carver Biotechnology
Center, University of Illinois).

Sequence processing and bioinformatics

We annotated in-line barcodes based on the first 7 bases of each sequencing read (umi-tools, Smith et al.,
2017), split out reads that matched each known barcode combination (grep, GNU Project, 1998), trimmed
the remaining primer sequences from the sequencing reads (cutadapt, Martin, 2011) counted the number of
sequencing reads in each file and removed files (including negative controls) with low read numbers (<100
reads) from the dataset. We processed the remaining reads with the R Divisive Amplicon Denoising
Algorithm package "dada2" (version 1.28.0) (Callahan et al., 2016). Taxonomy was assigned using the
Silva 138 database (Ref NR99) (Quast et al., 2012). The count table, taxonomy table, and sample-associated
data were integrated into a phyloseq object using the R package "phyloseq" (version 1.44.0) (McMurdie &
Holmes, 2013) for further data analysis. Any ASVs detected in control sequencing reactions without DNA
as well as ASVs with a phylum designation that was "NA", Eukaryotic, belonging to the family of
Mitochondria or the class of Chloroplasts were removed from the datasets. Samples with very low reads
(clearly below the first quartile) were excluded from each group.

 ITS sequence processing and bioinformatics

Amplification targeted the second internal transcribed spacer (ITS2) region of the fungal ribosomal cistron

using primers ITS3 and ITS4 (White et al., 1990). Full primer sequences included Illumina Nextera

transposase adapters (underlined) and were as follows: ITS3-5'-TCG TCG GCA GCG TCA GAT GTG

TAT AA GAG ACA GGC ATC GAT GAA GAA CGC AGC-3' and ITS4-5'-GTC TCG TGG GCT CGG

AGA TGT GTA TAA GAG ACA GTC CTC CGC TTA TTG ATA TGC-3'. Reaction mixes contained 1x

AccuStart II PCR SuperMix (Quantabio), 0.4 µM forward and reverse primers, and 2 µl of template DNA.

The thermal cycling regime consisted of an initial denaturation and enzyme activation step at 94°C for 3

min, followed by 30 cycles of 94°C for 45 sec, 50°C for 60 sec, and 72°C for 90 sec, with a final extension

step at 72° for 10 min. Ten additional PCR cycles were used to tag diluted amplicons with unique dual

indices (Illumina; cycling regime as above but with annealing at 54°C). Amplicons were pooled, cleaned

using AMPure XP beads (Beckman Coulter), and sequenced on an Illumina MiSeq instrument (2 x 300 nt)

at the DNA Services Lab, Roy J. Carver Biotechnology Center, University of Illinois at Urbana-Champaign.

Reads processing was consistent with the DADA2 ITS workflow

(https://benjjneb.github.io/dada2/ITS_workflow.html). Raw reads were trimmed of non-biological

sequence using cutadapt (Martin, 2011). DADA2 was used to filter and truncate the reads, infer amplicon

sequence variants (ASVs), and remove chimeras (Callahan et al., 2016). Taxonomy was assigned using

DADA2's implementation of the RDP naïve Bayesian classifier (Q. Wang et al., 2007) against a UNITE

reference database (version 9.0; Abarenkov et al., 2023). The ASV count table, taxonomy table, and sample-

associated data were integrated into a single object in R using the phyloseq package (McMurdie & Holmes,

2013).

Microbiota composition analyses

All statistical analyses were performed in R (v 4.3.2) (R Core Team, 2023), and "ggplot2" (v 3.4.4)

(Wickham, 2016) was used for visualizations if not stated otherwise. Illustrations of all species were created

in Adobe Illustrator (v 28.3) (Adobe Inc., 2024) and all figures were compiled in InkScape (v 1.1) (Inkscape

Project, 2021).

Microbiota composition analyses of captive frogs and source tracking

First, we calculated beta diversity between skin communities of adult frogs that served as transporting frogs

to determine if communities are distinct enough to distinguish them as microbial source communities. We

applied Principal Coordinates Analysis (PCoA) using the package "phyloseq" to visualize variation in the

genus-agglomerated dataset based on two dissimilarity indices: Bray-Curtis, a widely used measure

considering relative abundances and unweighted Unifrac, a presence-absence measure of ASVs based on

phylogenetic distances. Data were converted to relative abundances before calculating Bray-Curtis or rarefied using a rarefaction over 300 iterations (function phyloseq_mult_raref_avg in the package 'metagmisc' (Mikryukov, 2023) before calculating Unifrac distances. We confirmed equal dispersion between compared groups using the betadisper function of the package "vegan" (v2.6.4) (Oksanen et al., 2022) followed by a Tukey post-hoc test. Variation was analyzed with a PERMANOVA (function 'adonis' within the package "vegan") using Bray-Curtis and unweighted Unifrac dissimilarities as responses and species or transporting species as predictors. Significant PERMANOVAs were followed by pairwise multilevel post-hoc comparisons between groups with the function pairwise.adonis (package "pairwiseAdonis", (Arbizu, 2017). We applied the same method to compare if communities between tadpoles that had been transported by a homospecific frog (*Rv*) or a heterospecific frog (*Os*) differed after six hours of tadpole transport. We further used 'Sourcetracker' (Knights et al., 2011) to determine whether transporting frogs served as a source for bacteria detected on tadpole skin after six hours of transport. This Bayesian approach was developed to estimate proportion of contaminants in a given community that come from possible source environments and is commonly used to determine if parental communities serve as source for offspring communities (Kouete et al., 2023; Murphy et al., 2023; Switzer et al., 2023). We performed two sets of analysis to determine the source proportions of adult *Rv* and *Os* in tadpole communities. First, we defined transporting frogs of both species as possible microbial sources and all tadpoles including non-transported controls as sinks. Then, we defined transporting frogs and non-transported control tadpoles as possible sources to visualize the proportion of bacteria acquired in the clutch prior to transport. For both approaches, we rarefied the source dataset of transporting frogs to decrease the influence of high-coverage source samples on the analysis. For the first approach, we plotted the source proportions of *Rv* and *Os* communities for non-transported tadpoles, tadpoles transported by *Rv*, and tadpoles transported by *Os*, and compared between the experimental conditions for each source (*Rv* or *Os*) separately using a Kruskal Wallis test, with p-values adjusted for multiple comparisons using Benjamini Hochberg correction (R Core Team, 2023). Additionally, differences in community proportions shared with the transporting species ('transporting frog') or the non-transporting species ('non-transporting frog') were determined for non-transported and transported tadpoles and compared with the same analysis. For the second approach, we determined the source proportions that transported tadpoles had acquired from the clutch, from *Rv* adults, and from *Os* adults, and we determined differences between tadpoles transported by *Rv* or by *Os* using a paired Wilcoxon test with Benjamini Hochberg correction (R Core Team, 2023).

To assess training quality, we evaluated model self-assignment using source samples. We selected the model trained on a dataset rarefied to the read depth of the adult frog sample with the lowest read count (48162 reads), as it showed the best overall self-assignment performance, whereas models trained on

datasets rarefied to the lowest overall read depth performed worse. Unlike studies using technical replicates,

our source samples represent distinct biological individuals and sampling timepoints, where natural

microbiome variability is expected within each source category. Consequently, we considered self-

assignment rates above 70% acceptable. All source samples were correctly assigned to their respective

categories (*Rv*, *Os*, or control), but with varying proportions of reads assigned as 'Unknown'. Adult frog

sources were reliably self-identified with high confidence (*Os*: 97.2% median, IQR = 1.4; *Rv*: 76.3%

median, IQR = 38.1). Adult *Rv* frogs displayed a higher proportion of 'Unknown' assignments compared to

Os, reflecting greater biological variability among individuals and/or a higher proportion of rare taxa not

well captured in the training set. The control tadpole source showed lower self-assignment accuracy

(median = 30.5%, IQR = 17.1), as expected given the low microbial biomass of these samples, which

resulted in low read depth. Low read-depth limits the information available to inform the iterative updating

steps in Gibbs sampling and reduces confidence in source assignments. We therefore verified the robustness

of our results by performing the second Sourcetracker analysis as described above, training the model only

on adult sources and assigning all tadpoles, including low-biomass controls, as sinks (as described above).

Self-assignment rates for the second training set varied (*Os*: 79.2% median, IQR = 29; *Rv*: 96.6% median,

IQR = 3.7), while results remained consistent across analyses, supporting the reliability of our findings.

Microbiota composition analyses of wild populations

We calculated mean and standard deviation across all samples per group for the number of phyla and

families from a rarefied dataset to compare groups. The total number of unique phyla and families for each

group was calculated from the unrarefied dataset to include low abundance taxa. We plotted the relative

abundance of taxa agglomerated at the phylum level with an abundance greater than 2%. In addition, we

identified the most abundant genera (N = 10) for each group based on overall mean relative abundances.

The number of *Batrachochytrium*-inhibiting taxa was calculated by blasting the rarefied dataset against the

AmphiBac-Database-2023.2 (Woodhams et al., 2015, updated by Bletz 2023); Antifungal database:

AmphibBac_InhibitoryStrict_2023.2; https://github.com/AmphiBac/AmphiBac-Database) and calculating

the average across all samples of a group.

All alpha diversity analyses were conducted with datasets rarefied to 90% of the read number of the sample

with the fewest reads in each comparison and visualized with boxplots. We compared richness, Shannon

index and evenness between different life stages of the same species, as well as between tadpoles from

different species and their environments. Differences in alpha diversity measures between groups were

determined using an ANOVA (for normally distributed, homoscedastic data) followed by a Tukey post-hoc

test or Kruskal-Wallis (for non-normally distributed data) followed by a Dunn test. P-values were adjusted

for multiple comparisons using Benjamini-Hochberg corrections.

We determined beta diversity across frog species, life stages, and aquatic environments using two

commonly used dissimilarity measurements as described for the captive-bred dataset. ASVs found in only

one frog were removed from the full, non-rarefied dataset. Variation was analyzed with a PERMANOVA

(function adonis within the package "vegan"), using Bray-Curtis and unweighted Unifrac dissimilarities as

responses and species, life stage, species- life stage interaction and parenting behavior as predictors.

Significant PERMANOVAs were followed by pairwise multilevel post-hoc comparisons between groups

with the function pairwise.adonis (package "pairwiseAdonis", Martinez Arbizu, 2020).

As previously suggested (Neu et al., 2021), we determined genus level agglomerated taxa between adults,

tadpoles and aquatic environments from unrarefied data converted to relative abundances across different

prevalence (prev) and abundance (abd) cutoffs. This approach avoids using arbitrary thresholds to define a

taxonomic 'core'. The following cutoffs were evaluated using the function "core_members" of the package

"microbiome" (v1.22.0, Lahti & Shetty, 2012, 2023): (1) prev 100% without abd cutoff, (2) prev >75%

with abd >1%, (3) prev >75% with abd >0.1% and (4) prev >75% without abd cutoff. We listed the most

abundant genera for each species (*Rv*, *Af* and *Ll*) and life stage (adult or tadpoles) with respect to their

presence or absence in the core of the aquatic environment of the respective species and used the package

"eulerr" (Larson, 2022) to visualize overlaps of genera with prevalence over 75% and relative abundance

over 0.1% with a Venn diagram.

To test whether community distances between tadpoles and adults are smaller in poison frogs than in a

species without parental care, we performed a beta regression with an underlying logit function (R package

"betareg", Zeileis et al., 2021) to analyze unweighted Unifrac distances of a rarefied dataset. This approach

is suited to model predictions on our bounded distance dataset with high variance. ASVs were agglomerated

at the genus level prior to calculating distances between tadpoles and adults of different species, while

distances between related and unrelated pairs were calculated on a ASV level. To account for differences

in samples size between groups, we performed iterated analyses with random subsamples of equal sample

size (N = 8 for distances between adults and tadpoles between species and N = 10 for related vs. not-related

pairs of *Rv*) and report averaged p-values (following the approach of Hughey et al., 2017).

Microbiota composition analyses of wild experimental tadpoles

We determined and illustrated alpha and beta diversities of microbial communities on transported and non-

transported tadpoles and significant differences between the two groups as described for wild populations.

Variation was analyzed with a PERMANOVA (function adonis within the package "vegan"), using
unweighted Unifrac dissimilarity measures calculated from a rarefied dataset. Due to the limited sample
size and low biomass of some tadpoles, we chose to work with relative ASV abundances rather than
rarefaction to determine disparities and overlaps between the groups and their aquatic environment,
following previous examples (McMurdie & Holmes, 2013; Prest et al., 2018). Venn diagrams were created
without prevalence or abundance cutoffs and include all low abundance taxa. Differentially abundant
genera between transported and non-transported tadpoles were determined with the package
"ANCOMBC2" (version 2.2.2) (Lin, 2023).

We used 'Sourcetracker' (Knights et al., 2011) on unrarefied ASVs to determine proportions of the
communities of adult caregivers ("source") that can be detected on tadpoles that were collected from their
back and then reared in artificial cups for one month ("sink"). For this approach, we defined the
communities of each caregiver as source and transported tadpoles as sinks and reported minimum,
maximum and median source proportions detected among transported tadpoles.

ASVs shared between the transported tadpole and its transporting caregiver were identified using the
function "common_taxa" of the package "phylosmith" (Smith, 2023). We then evaluated the presence and
relative abundance of these shared taxa in transported and non-transported tadpoles, in their respective
aquatic cup environments, and in the water used to refill their cups. We converted relative abundances to
binary presence-absence information for illustration in a bubble chart and determined the ten taxa with the
highest relative abundance for each transporting caregiver individually from the unrarefied dataset.

Quantitative analysis of 16S rRNA gene copy numbers with digital PCR (dPCR)

Absolute abundances were estimated for ASVs that were shared between tadpoles after a one-month growth
period and their respective caregivers, and for which Sourcetracker analysis identified the caregiver as a
likely source of microbiota. We followed the quantitative sequencing framework described by Barlow et
al., 2020, measuring total microbial load via digital PCR (dPCR) with the same universal 16S rRNA primers
used to amplify the V4 region in our sequencing dataset. Absolute 16S rRNA gene copy numbers obtained
from dPCR were then multiplied by the relative abundances from our amplicon sequencing dataset to
calculate ASV-specific scaled absolute abundances. All dPCR reactions were carried out on a QIAcuity
Digital PCR System (Qiagen) using Nanoplates with a 8.5K partition configuration, using the following
cycling program: 95°C for 2 minutes, 40 cycles of 95°C for 30 seconds and 52°C for 30 seconds and 72°C
for 1 minute, followed by 1 cycle of 40°C for 5 minutes. Reactions were prepared using the QIAcuity
EvaGreen PCR Kit (Qiagen, Cat. No. 250111) with 2 µL of DNA template per reaction, following the

manufacturer's protocol, and included a negative no-template control and a cleaned and sequenced PCR
product as positive control. Samples were measured in triplicates and serial dilutions were performed to
ensure accurate quantification. Data were processed with the QIAcuity Software Suite (v3.1.0.0). The
threshold was set based on the negative and positive controls in 1D scatterplots. We report mean copy
numbers per microliter with standard deviations, correcting for template input, dPCR reaction volume, and
dilution factor. Mean gene copy numbers per tadpole were additionally calculated by accounting for the
DNA extraction (elution) volume.

Permits

Approval of the field monitoring by the scientific institution of the authors:

The monitoring project in the Nouragues was reviewed by the Administrative Panel on Laboratory Animal
Care (APLAC) at Stanford University and approved under the APLAC-protocol 33691 (Protocol Director
LAO). The APLAC committee is Stanford's Institutional Animal Care and Use Committee (IACUC). It is
appointed by the University Vice Provost and Dean of Research.

Approval from local field site:

Non-invasive field experiments included in this publication were approved by the scientific committee of
the Nouragues Reserve. The field station in the Nouragues Reserve is managed by the CNRS (Centre
National de la Recherche Scientifique), is part of the USR 3456 LEEISA (Laboratoire Ecologie, Evolution
Interactions des Systèmes Amazoniens, founded in 2016) and located in the nature reserve 'Les Nouragues',
which is operated by the Office national des forêts (ONF) Guyane. Additional tissue collections were
approved by the Direction Générale des Territoires et de la Mer (DGTM approval number # R03-2022-08-
10-00001).

Data availability

899 Data files and data analysis scripts used to generate the results are available on DataDryad (pending
900 acceptance).

Acknowledgements

We would like to thank Ami S. Bhatt for her support in facilitating the contributions of Mai Dvorak to this
work. We are grateful to the staff of the CNRS Guyane, the Nouragues Ecological Research Station and

the ONF, especially Patrick Chatelet and Jennifer Devillechabrol, for logistic and moral support in the field.
We thank Daniel Shaykevich for reviewing all versions of the manuscript, Camilo Rodríguez-Lopez for
statistical consultation and Philippe Gaucher for endless knowledge, discussions and guidance in the world
of amphibians. This research was conducted at Stanford University, which is located on the ancestral and
unceded land of the Muwekma Ohlone tribe. Our field work was conducted in the Nature Reserve 'Les
Nouragues' in French Guiana, that was founded on land ancestrally inhabited by the Amerindien Nouragues
('Norak') tribe.

Declaration of Interests

912   The authors declare no competing interests.

Author Contributions

Conceptualization: MTF, SC

Data Curation: MTF, KSX, EKC, MD, GR, AR

Formal Analysis: MTF, KSX, EKC, MD

Investigation: MTF

Visualization: MTF, LAO

Methodology: MTF

Writing: MTF

Review and Editing: KSX, EKC, DAR, LAO

Project Administration: MTF, LAO

Supervision: KSX, EKC, SC, DAR, LAO

Resources: DAR, LAO

Funding Acquisition: MTF, LAO

Funding

927   This research was funded with grants from National Institutes of Health (DP2HD102042) and the
928   McKnight Foundation to LAO. MTF is supported by an Erwin Schrödinger fellowship from the FWF (J-
929   4526B). LAO is a New York Stem Cell Foundation – Robertson Investigator. DAR was supported by the
930   Thomas C. and Joan M. Merigan Endowment at Stanford University and by the Good Ventures Fund.

References

Abarenkov, K., Zirk, A., Piirmann, T., Pöhönen, R., Ivanov, F., Nilsson, R. H., & Kõljalg, U. (2023). *UNITE general FASTA release for eukaryotes* [Application/gzip]. UNITE Community. https://doi.org/10.15156/BIO/2938069

Abellan-Schneyder, I., Schusser, A. J., & Neuhaus, K. (2021). ddPCR allows 16S rRNA gene amplicon sequencing of very small DNA amounts from low-biomass samples. *BMC Microbiology*, *21*(1), 349. https://doi.org/10.1186/s12866-021-02391-z

Adobe Inc. (2024). *Adobe Illustrator, Version 28.3*. https://www.adobe.com/products/illustrator.html

Al Nabhani, Z., & Eberl, G. (2020). Imprinting of the immune system by the microbiota early in life. *Mucosal Immunology*, *13*(2), 183–189. https://doi.org/10.1038/s41385-020-0257-y

Alexiev, A., Chen, M. Y., Korpita, T., Weier, A. M., & McKenzie, V. J. (2023). Together or Alone: Evaluating the Pathogen Inhibition Potential of Bacterial Cocktails against an Amphibian Pathogen. *Microbiology Spectrum*, *11*(2), e01518-22. https://doi.org/10.1128/spectrum.01518-22

Almany, G. R. (2004). Priority Effects in Coral Reef Fish Communities of the Great Barrier Reef. *Ecology*, *85*(10), 2872–2880. https://doi.org/10.1890/03-3166

Altig, R., & McDiarmid, R. W. (2007). MORPHOLOGICAL DIVERSITY AND EVOLUTION OF EGG AND CLUTCH STRUCTURE IN AMPHIBIANS. *Herpetological Monographs*, *21*(1), 1. https://doi.org/10.1655/06-005.1

Arbizu, P. M. (2017). *pairwiseAdonis: Pairwise Multilevel Comparison using Adonis*.

Balshine, S. (2012). Patterns of parental care in vertebrates. In N. J. Royle & P. T. Smiseth (Eds.), *The Evolution of Parental Care* (pp. 62–80). Oxford University Press. https://doi.org/10.1093/acprof:oso/9780199692576.003.0004

Barlow, J. T., Bogatyrev, S. R., & Ismagilov, R. F. (2020). A quantitative sequencing framework for absolute abundance measurements of mucosal and lumenal microbial communities. *Nature Communications*, *11*(1), 2590. https://doi.org/10.1038/s41467-020-16224-6

Belkaid, Y., & Hand, T. W. (2014). Role of the Microbiota in Immunity and Inflammation. *Cell*, *157*(1), 121–141. https://doi.org/10.1016/j.cell.2014.03.011

Benjamino, J., Lincoln, S., Srivastava, R., & Graf, J. (2018). Low-abundant bacteria drive compositional

changes in the gut microbiota after dietary alteration. *Microbiome*, *6*(1), 86.

https://doi.org/10.1186/s40168-018-0469-5

Biedermann, P. H., & Rohlfs, M. (2017). Evolutionary feedbacks between insect sociality and microbial

management. *Current Opinion in Insect Science*, *22*, 92–100. https://doi.org/10.1016/j.cois.2017.06.003

Bik, E. M., Costello, E. K., Switzer, A. D., Callahan, B. J., Holmes, S. P., Wells, R. S., Carlin, K. P.,

Jensen, E. D., Venn-Watson, S., & Relman, D. A. (2016). Marine mammals harbor unique microbiotas

shaped by and yet distinct from the sea. *Nature Communications*, *7*(1), 10516.

https://doi.org/10.1038/ncomms10516

Bletz, M. C., Myers, J., Woodhams, D. C., Rabemananjara, F. C. E., Rakotonirina, A., Weldon, C.,

Edmonds, D., Vences, M., & Harris, R. N. (2017). Estimating Herd Immunity to Amphibian

Chytridiomycosis in Madagascar Based on the Defensive Function of Amphibian Skin Bacteria. *Frontiers*

in Microbiology, *8*. https://doi.org/10.3389/fmicb.2017.01751

Blyton, M. D. J., Soo, R. M., Hugenholtz, P., & Moore, B. D. (2022). Maternal inheritance of the koala

gut microbiome and its compositional and functional maturation during juvenile development.

Environmental Microbiology, *24*(1), 475–493. https://doi.org/10.1111/1462-2920.15858

Bongers, F., Charles-Dominique, P., Forger, P., & Théry, M. (2011). *Nouragues: Dynamics and Plant-*

Animal Interactions in a Neotropical Rainforest. Springer Netherlands.

Brown, J. L., Morales, V., & Summers, K. (2008). Divergence in parental care, habitat selection and

larval life history between two species of Peruvian poison frogs: An experimental analysis. *Journal of*

Evolutionary Biology, *21*, 1534–1543. https://doi.org/10.1111/j.1420-9101.2008.01609.x

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016).

DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods*, *13*, 581–583.

https://doi.org/10.1038/nmeth.3869

Caty, S. N., Alvarez-Buylla, A., Vasek, C., Tapia, E. E., Martin, N. A., McLaughlin, T., Weber, P. K.,

Mayali, X., Coloma, L. A., Morris, M. M., & O'Connell, L. A. (2024). *A toxic environment selects for*

specialist microbiome in poison frogs (p. 2024.01.10.574901). bioRxiv.

https://doi.org/10.1101/2024.01.10.574901

Caty, S. N., Vasek, C., Fischer, M.-T., Meneses, A. M., Shaykevich, D. A., & O'Connell, L. A. (2024).

Isolation of microbes from the skin of terrestrial frogs. https://www.protocols.io/view/isolation-of-

microbes-from-the-skin-of-terrestrial-c3icykaw

Chen, C.-Y., Chen, C.-K., Chen, Y.-Y., Fang, A., Shaw, G. T.-W., Hung, C.-M., & Wang, D. (2020).

Maternal gut microbes shape the early-life assembly of gut microbiota in passerine chicks via nests.

Microbiome, *8*(1), 129. https://doi.org/10.1186/s40168-020-00896-9

Chen, D. W., & Garud, N. R. (2022). Rapid evolution and strain turnover in the infant gut microbiome.

Genome Research, *32*(6), 1124–1136. https://doi.org/10.1101/gr.276306.121

Clutton-Brock, T. H., & Scott, D. (1991). *The Evolution of Parental Care* (Vol. 64). Princeton University

Press. https://doi.org/10.2307/j.ctvs32ssj

Çömden, E., Yenmiş, M., & Çakır, B. (2023). The Complex Bridge between Aquatic and Terrestrial Life:

Skin Changes during Development of Amphibians. *Journal of Developmental Biology*, *11*(1), 6.

https://doi.org/10.3390/jdb11010006

Conlon, J. M. (2011). The contribution of skin antimicrobial peptides to the system of innate immunity in

anurans. *Cell and Tissue Research*, *343*(1), 201–212. https://doi.org/10.1007/s00441-010-1014-4

Cox, L. M., Yamanishi, S., Sohn, J., Alekseyenko, A. V., Leung, J. M., Cho, I., Kim, S. G., Li, H., Gao,

Z., Mahana, D., Zárate Rodriguez, J. G., Rogers, A. B., Robine, N., Loke, P., & Blaser, M. J. (2014).

Altering the intestinal microbiota during a critical developmental window has lasting metabolic

consequences. *Cell*, *158*(4), 705–721. https://doi.org/10.1016/j.cell.2014.05.052

Crump, M. L. (1974). Reproductive Strategies in a Tropical Anuran Community. *Misc Publ Mus Nat Hist*

Univ Kansas, *61*, 1–68.

Crump, M. L. (2015). Anuran Reproductive Modes: Evolving Perspectives. *Journal of Herpetology*,

49(1), 1–16. https://doi.org/10.1670/14-097

Daszak, P., Cunningham, A. A., & Hyatt, A. D. (2003). Infectious disease and amphibian population

declines. *Diversity and Distributions*, *9*(2), 141–150. https://doi.org/10.1046/j.1472-4642.2003.00016.x

de Bruijn, I., Liu, Y., Wiegertjes, G. F., & Raaijmakers, J. M. (2018). Exploring fish microbial

communities to mitigate emerging diseases in aquaculture. *FEMS Microbiology Ecology*, *94*(1), fix161.

https://doi.org/10.1093/femsec/fix161

Debray, R., Herbert, R. A., Jaffe, A. L., Crits-Christoph, A., Power, M. E., & Koskella, B. (2022).

Priority effects in microbiome assembly. *Nature Reviews Microbiology*, *20*(2), 109–121.

https://doi.org/10.1038/s41579-021-00604-w

Delia, J. R. J., Ramírez-Bautista, A., & Summers, K. (2013). Parents adjust care in response to weather

conditions and egg dehydration in a Neotropical glassfrog. *Behavioral Ecology and Sociobiology*, *67*(4),

557–569. https://doi.org/10.1007/s00265-013-1475-z

Donald, K., & Finlay, B. B. (2023). Early-life interactions between the microbiota and immune system:

Impact on immune system development and atopic disease. *Nature Reviews Immunology*, *23*(11), 735–

748. https://doi.org/10.1038/s41577-023-00874-w

Doyle, B., Reynolds, G. Z. M., Dvorak, M., Maghini, D. G., Natarajan, A., & Bhatt, A. S. (2025).

Absolute quantification of prokaryotes in the microbiome by 16S rRNA qPCR or ddPCR. *Nature*

Protocols, 1–36. https://doi.org/10.1038/s41596-025-01165-5

Dubois, L., Valles-Colomer, M., Ponsero, A., Helve, O., Andersson, S., Kolho, K.-L., Asnicar, F.,

Korpela, K., Salonen, A., Segata, N., & Vos, W. M. de. (2024). Paternal and induced gut microbiota

seeding complement mother-to-infant transmission. *Cell Host & Microbe*, *32*(6), 1011-1024.e4.

https://doi.org/10.1016/j.chom.2024.05.004

Duellman, W. E., & Trueb, L. (1986). *Biology of Amphibians*. John Hopkins University Press, Baltimore,

London.

Emerson, K. J., & Woodley, S. K. (2024). Something in the water: Aquatic microbial communities

influence the larval amphibian gut microbiota, neurodevelopment and behaviour. *Proceedings of the*

Royal Society B: Biological Sciences, *291*(2017), 20232850. https://doi.org/10.1098/rspb.2023.2850

Fallet, M., Montagnani, C., Petton, B., Dantan, L., De Lorgeril, J., Comarmond, S., Chaparro, C., Toulza,

E., Boitard, S., Escoubas, J.-M., Vergnes, A., Le Grand, J., Bulla, I., Gueguen, Y., Vidal-Dupiol, J.,

Grunau, C., Mitta, G., & Cosseau, C. (2022). Early life microbial exposures shape the Crassostrea gigas

immune system for lifelong and intergenerational disease protection. *Microbiome*, *10*(1), 85.

https://doi.org/10.1186/s40168-022-01280-5

Faszewski, E. E., Tyrell, A., Guin, S., & Kaltenbach, J. C. (2008). Metamorphic changes in localization

of sugars in skin of the leopard frog, *Rana pipiens*. *Journal of Morphology*, *269*(8), 998–1007.

https://doi.org/10.1002/jmor.10639

Fischer, M.-T., Ringler, M., Ringler, E., & Pašukonis, A. (2020). Reproductive behavior drives female

space use in a sedentary Neotropical frog. *PeerJ*, *8*, e8920. https://doi.org/10.7717/peerj.8920

Florkowski, M. R., & Yorzinski, J. L. (2023). Gut microbiome diversity and composition is associated

with exploratory behavior in a wild-caught songbird. *Animal Microbiome*, *5*(1), 8.

https://doi.org/10.1186/s42523-023-00227-x

Fontaine, S. S., Mineo, P. M., & Kohl, K. D. (2022). Experimental manipulation of microbiota reduces

host thermal tolerance and fitness under heat stress in a vertebrate ectotherm. *Nature Ecology &*

Evolution. https://doi.org/10.1038/s41559-022-01686-2

Fredriksson, N. J., Hermansson, M., & Wilén, B.-M. (2013). The Choice of PCR Primers Has Great

Impact on Assessments of Bacterial Community Diversity and Dynamics in a Wastewater Treatment

Plant. *PLoS ONE*, *8*(10), e76431. https://doi.org/10.1371/journal.pone.0076431

Fukami, T. (2015). Historical Contingency in Community Assembly: Integrating Niches, Species Pools,

and Priority Effects. *Annual Review of Ecology, Evolution, and Systematics*, *46*(1), 1–23.

https://doi.org/10.1146/annurev-ecolsys-110411-160340

Furness, A. I., & Capellini, I. (2019). The evolution of parental care diversity in amphibians. *Nature*

Communications, *10*(1), 4709. https://doi.org/10.1038/s41467-019-12608-5

Gensollen, T., Iyer, S. S., Kasper, D. L., & Blumberg, R. S. (2016). How colonization by microbiota in

early life shapes the immune system. *Science*, *352*(6285), 539–544.

https://doi.org/10.1126/science.aad9378

GNU Project. (1998). *Grep 2.5*.

Goolsby, B. C., Fischer, M.-T., Pareja-Mejia, D., Lewis, A. R., Raboisson, G., & O'Connell, L. A.

(2023). *Home security cameras as a tool for behavior observations and science equity* (p.

2023.04.17.537238). bioRxiv. https://doi.org/10.1101/2023.04.17.537238

Grant, T., Rada, M., Anganoy-Criollo, M., Batista, A., Dias, P. H., Jeckel, A. M., Machado, D. J., &

Rueda-Almonacid, J. V. (2017). Phylogenetic Systematics of Dart-Poison Frogs and Their Relatives

Revisited (Anura: Dendrobatoidea). *South American Journal of Herpetology*, *12*(SI 1).

https://doi.org/10.2994/SAJH-D-17-00017.1

Gross, M. R. (2005). The Evolution of Parental Care. *The Quarterly Review of Biology*, *80*(1), 37–45.

https://doi.org/10.1086/431023

Guyot, N., Labas, V., Harichaux, G., Chessé, M., Poirier, J.-C., Nys, Y., & Réhault-Godbert, S. (2016).

Proteomic analysis of egg white heparin-binding proteins: Towards the identification of natural

antibacterial molecules. *Scientific Reports*, *6*, 27974. https://doi.org/10.1038/srep27974

Hallmaier-Wacker, L. K., Lueert, S., Roos, C., & Knauf, S. (2018). The impact of storage buffer, DNA

extraction method, and polymerase on microbial analysis. *Scientific Reports*, *8*(1), 6292.

https://doi.org/10.1038/s41598-018-24573-y

Han, G., & Vaishnava, S. (2023). Microbial underdogs: Exploring the significance of low-abundance

commensals in host-microbe interactions. *Experimental & Molecular Medicine*, *55*(12), 2498–2507.

https://doi.org/10.1038/s12276-023-01120-y

Harris, R. N., Brucker, R. M., Walke, J. B., Becker, M. H., Schwantes, C. R., Flaherty, D. C., Lam, B. A.,

Woodhams, D. C., Briggs, C. J., Vredenburg, V. T., & Minbiole, K. P. C. (2009). Skin microbes on frogs

prevent morbidity and mortality caused by a lethal skin fungus. *The ISME Journal*, *3*(7), 818–824.

https://doi.org/10.1038/ismej.2009.27

Harris, R. N., James, T. Y., Lauer, A., Simon, M. A., & Patel, A. (2006). Amphibian Pathogen

Batrachochytrium dendrobatidis Is Inhibited by the Cutaneous Bacteria of Amphibian Species.

EcoHealth, *3*(1), 53–56. https://doi.org/10.1007/s10393-005-0009-1

Hayes, R. A., Crossland, M. R., Hagman, M., Capon, R. J., & Shine, R. (2009). Ontogenetic variation in

the chemical defenses of cane toads (Bufo marinus): Toxin profiles and effects on predators. *Journal of*

Chemical Ecology, *35*(4), 391–399. https://doi.org/10.1007/s10886-009-9608-6

Herp, S., Brugiroux, S., Garzetti, D., Ring, D., Jochum, L. M., Beutler, M., Eberl, C., Hussain, S., Walter,

S., Gerlach, R. G., Ruscheweyh, H. J., Huson, D., Sellin, M. E., Slack, E., Hanson, B., Loy, A., Baines, J.

F., Rausch, P., Basic, M., … Stecher, B. (2019). Mucispirillum schaedleri Antagonizes Salmonella

Virulence to Protect Mice against Colitis. *Cell Host & Microbe*, *25*(5), 681-694.e8.

https://doi.org/10.1016/j.chom.2019.03.004

Hughey, M. C., Delia, J., & Belden, L. K. (2017). Diversity and stability of egg-bacterial assemblages:

The role of paternal care in the glassfrog *Hyalinobatrachium colymbiphyllum*. *Biotropica*, *49*(6), 792–

802. https://doi.org/10.1111/btp.12461

Inkscape Project. (2021). *Inkscape, Version 1.1*. https://inkscape.org

Jones, E. W., Carlson, J. M., Sivak, D. A., & Ludington, W. B. (2022). Stochastic microbiome assembly

depends on context. *Proceedings of the National Academy of Sciences*, *119*(7), e2115877119.

https://doi.org/10.1073/pnas.2115877119

Jones, K. R., Belden, L. K., & Hughey, M. C. (2024). Priority effects alter microbiome composition and

increase abundance of probiotic taxa in treefrog tadpoles. *Applied and Environmental Microbiology*,

90(6), e00619-24. https://doi.org/10.1128/aem.00619-24

Jones, K. R., Hughey, M. C., & Belden, L. K. (2023). Colonization order of bacterial isolates on treefrog

embryos impacts microbiome structure in tadpoles. *Proceedings of the Royal Society B: Biological*

Sciences, *290*(1995), 20230308. https://doi.org/10.1098/rspb.2023.0308

Kaur, H., Ali, S. A., & Yan, F. (2022). Interactions between the gut microbiota-derived functional factors

and intestinal epithelial cells – implication in the microbiota-host mutualism. *Frontiers in Immunology*,

13, 1006081. https://doi.org/10.3389/fimmu.2022.1006081

Klug, H., & Bonsall, M. B. (2014). What are the benefits of parental care? The importance of parental

effects on developmental rate. *Ecology and Evolution*, *4*(12), 2330–2351.

https://doi.org/10.1002/ece3.1083

Knights, D., Kuczynski, J., Charlson, E. S., Zaneveld, J., Mozer, M. C., Collman, R. G., Bushman, F. D.,

Knight, R., & Kelley, S. T. (2011). Bayesian community-wide culture-independent microbial source

tracking. *Nature Methods*, *8*(9), 761–763. https://doi.org/10.1038/nmeth.1650

Körner, M., Steiger, S., & Shukla, S. P. (2023). Microbial management as a driver of parental care and

family aggregations in carrion feeding insects. *Frontiers in Ecology and Evolution*, *11*, 1252876.

https://doi.org/10.3389/fevo.2023.1252876

Kouete, M., Bletz, M., Labumbard, B., Woodhams, D., & Blackburn, D. (2023). Parental care contributes

to vertical transmission of microbes in a skin-feeding and direct-developing caecilian. *Animal*

Microbiome, *5*. https://doi.org/10.1186/s42523-023-00243-x

Kueneman, J. G., Parfrey, L. W., Woodhams, D. C., Archer, H. M., Knight, R., & McKenzie, V. J.

(2014). The amphibian skin-associated microbiome across species, space and life history stages.

Molecular Ecology, *23*(6), 1238–1250. https://doi.org/10.1111/mec.12510

Kwong, W. K., Medina, L. A., Koch, H., Sing, K.-W., Soh, E. J. Y., Ascher, J. S., Jaffé, R., & Moran, N.

A. (2017). Dynamic microbiome evolution in social bees. *Science Advances*, *3*(3), e1600513.

https://doi.org/10.1126/sciadv.1600513

Lahti, L., & Shetty, S. (2012). *Microbiome R package*.

Lahti, L., & Shetty, S. (2023). *microbiome: Microbiome Analytics*.

https://doi.org/10.18129/B9.bioc.microbiome

Langenfeld, K., Chin, K., Roy, A., Wigginton, K., & Duhaime, M. B. (2021). Comparison of

ultrafiltration and iron chloride flocculation in the preparation of aquatic viromes from contrasting sample

types. *PeerJ*, *9*, e11111. https://doi.org/10.7717/peerj.11111

Larson, E. J. (2022). *eulerr: Area-Proportional Euler and Venn Diagrams with Ellipses*. https://cran.r-

project.org/package=eulerr

Leftwich, P. T., Edgington, M. P., & Chapman, T. (2020). Transmission efficiency drives host–microbe

associations. *Proceedings of the Royal Society B: Biological Sciences*, *287*(1934), 20200820.

https://doi.org/10.1098/rspb.2020.0820

Lescure, J. (1976). Etude des têtards de deux Phyllobates (Dendrobatidae). *Bull Soc Zool Fr*, *101*, 299–

306.

Lin, H. (2023). *ANCOMBC: Microbiome differential abudance and correlation analyses with bias

correction*. https://doi.org/10.18129/B9.bioc.ANCOMBC

Lötters, S., Jungfer, K.-H., Henkel, F. W., & Schmidt, W. (n.d.). *Poison frogs- Biology, Species and

Captive Husbandry*.

Loudon, A. H., Holland, J. A., Umile, T. P., Burzynski, E. A., Minbiole, K. P. C., & Harris, R. N. (2014).

Interactions between amphibians' symbiotic bacteria cause the production of emergent anti-fungal

metabolites. *Frontiers in Microbiology*, *5*. https://doi.org/10.3389/fmicb.2014.00441

Luedtke, J. A., Chanson, J., Neam, K., Hobin, L., Maciel, A. O., Catenazzi, A., Borzée, A., Hamidy, A.,

Aowphol, A., Jean, A., Sosa-Bartuano, Á., Fong G., A., De Silva, A., Fouquet, A., Angulo, A., Kidov, A.

A., Muñoz Saravia, A., Diesmos, A. C., Tominaga, A., … Stuart, S. N. (2023). Ongoing declines for the

world's amphibians in the face of emerging threats. *Nature*, *622*(7982), 308–314.

https://doi.org/10.1038/s41586-023-06578-4

Mann, K. (2008). Proteomic analysis of the chicken egg vitelline membrane. *PROTEOMICS*, *8*(11),

2322–2332. https://doi.org/10.1002/pmic.200800032

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet. Journal, *17*(1), 10–12.

Mazmanian, S. K., Liu, C. H., Tzianabos, A. O., & Kasper, D. L. (2005). An Immunomodulatory

Molecule of Symbiotic Bacteria Directs Maturation of the Host Immune System. *Cell*, *122*(1), 107–118.

https://doi.org/10.1016/j.cell.2005.05.007

McDiarmid, R. W., & Altig, R. (Eds.). (1999). *Tadpoles- the biology of anuran larvae*. University of

Chicago Press.

McMurdie, P. J., & Holmes, S. (2013). phyloseq: An R package for reproducible interactive analysis and

graphics of microbiome census data. *PLoS ONE*, *8*(4), e61217.

Méndez-Tepepa, M., Morales-Cruz, C., García-Nieto, E., & Anaya-Hernández, A. (2023). A review of

the reproductive system in anuran amphibians. *Zoological Letters*, *9*(1), 3.

https://doi.org/10.1186/s40851-023-00201-0

Mikryukov, V. (2023). *metagMisc: Miscellaneous functions for metagenomic analysis*.

Mine, Y., & Kovacs-Nolan, J. (2006). New insights in biologically active proteins and peptides derived

from hen egg. *World's Poultry Science Journal*, *62*(1), 87–96. https://doi.org/10.1079/WPS200586

Murphy, K. M., Le, S. M., Wilson, A. E., & Warner, D. A. (2023). The Microbiome as a Maternal Effect:

A Systematic Review on Vertical Transmission of Microbiota. *Integrative And Comparative Biology*,

63(3), 597–609. https://doi.org/10.1093/icb/icad031

Neu, A. T., Allen, E. E., & Roy, K. (2021). Defining and quantifying the core microbiome: Challenges

and prospects. *Proceedings of the National Academy of Sciences*, *118*(51), e2104429118.

https://doi.org/10.1073/pnas.2104429118

Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B.,

Solymos, P., Stevens, M. H. H., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker, B., Borcard,

D., Carvalho, G., Chirico, M., Caceres, M. D., Durand, S., … Weedon, J. (2022). *vegan: Community*

Ecology Package. https://github.com/vegandevs/vegan

Pascoe, E. L., Hauffe, H. C., Marchesi, J. R., & Perkins, S. E. (2017). Network analysis of gut microbiota

literature: An overview of the research landscape in non-human animal studies. *The ISME Journal*,

11(12), 2644–2651. https://doi.org/10.1038/ismej.2017.133

Pašukonis, A., Beck, K. B., Fischer, M.-T., Weinlein, S., Stückler, S., & Ringler, E. (2017). Induced

parental care in a poison frog: A tadpole cross-fostering experiment. *The Journal of Experimental*

Biology, *220*, 3949–3954. https://doi.org/10.1242/jeb.165126

Pašukonis, A., Loretto, M.-C., & Rojas, B. (2019). How far do tadpoles travel in the rainforest? Parent-

assisted dispersal in poison frogs. *Evolutionary Ecology*, *0123456789*. https://doi.org/10.1007/s10682-

019-09994-z

Pašukonis, A., Serrano-Rojas, S. J., Fischer, M.-T., Loretto, M.-C., Shaykevich, D. A., Rojas, B., Ringler,

M., Roland, A. B., Marcillo-Lara, A., Ringler, E., Rodríguez, C., Coloma, L. A., & O'Connell, L. A.

(2022). Contrasting parental roles shape sex differences in poison frog space use but not navigational

performance. *eLife*, *11*, e80483. https://doi.org/10.7554/eLife.80483

Peignier, M., Furdi, B., Bégué, L., & Ringler, E. (2022). <p>A comparative table for staging anuran

embryos from terrestrial clutches based on the Brilliant-thighed Poison Frog, Allobates

femoralis (Anura: Dendrobatidae)</p>: Staging anuran embryos from terrestrial clutches.

Herpetology Notes, *15*, 723–727.

Poelman, E. H., & Dicke, M. (2007). Offering offspring as food to cannibals: Oviposition strategies of

Amazonian poison frogs (Dendrobates ventrimaculatus). *Evolutionary Ecology*, *21*, 215–227.

https://doi.org/10.1007/s10682-006-9000-8

Poelman, E. H., & Dicke, M. (2008). Space Use of Amazonian Poison Frogs: Testing the Reproductive

Resource Defense Hypothesis. *Journal of Herpetology*, *42*(2), 270–278. https://doi.org/10.1670/07-

1031.1

Poelman, E. H., van Wijngaarden, R. P. A., & Raaijmakers, C. E. (2013). Amazon poison frogs

(Ranitomeya amazonica) use different phytotelm characteristics to determine their suitability for egg and

tadpole deposition. *Evolutionary Ecology*, *27*(4), 661–674. https://doi.org/10.1007/s10682-013-9633-3

Powell, J. E., Martinson, V. G., Urban-Mead, K., & Moran, N. A. (2014). Routes of Acquisition of the

Gut Microbiota of the Honey Bee Apis mellifera. *Applied and Environmental Microbiology*, *80*(23),

7378–7387. https://doi.org/10.1128/AEM.01861-14

Prest, T. L., Kimball, A. K., Kueneman, J. G., & McKenzie, V. J. (2018). Host-associated bacterial

community succession during amphibian development. *Molecular Ecology*, *27*(8), 1992–2006.

https://doi.org/10.1111/mec.14507

Pröhl, H., & Berke, O. (2001). Spatial distributions of male and female strawberry poison frogs and their

relation to female reproductive resources. *Oecologia*, *129*, 534–542.

https://doi.org/10.1007/s004420100751

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., & Glöckner, F. O. (2012). The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. *Nucleic Acids Research*, *41*(D1), D590–D596. https://doi.org/10.1093/nar/gks1219

R Core Team. (2023). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing. https://www.R-project.org/

Rebollar, E. A., Simonetti, S. J., Shoemaker, W. R., & Harris, R. N. (2016). Direct and Indirect Horizontal Transmission of the Antifungal Probiotic Bacterium Janthinobacterium lividum on Green Frog (Lithobates clamitans) Tadpoles. *Applied and Environmental Microbiology*, *82*(8), 2457–2466. https://doi.org/10.1128/AEM.04147-15

Requena, G. S., Buzatto, B. A., Munguía-Steyer, R., & Machado, G. (2009). Efficiency of uniparental male and female care against egg predators in two closely related syntopic harvestmen. *Animal Behaviour*, *78*(5), 1169–1176. https://doi.org/10.1016/j.anbehav.2009.07.035

Reynolds, H. M., & Bettini, M. L. (2023). Early-life microbiota-immune homeostasis. *Frontiers in Immunology*, *14*, 1266876. https://doi.org/10.3389/fimmu.2023.1266876

Ringler, E., Pasukonis, A., Hödl, W., Ringler, M., Pašukonis, A., Hödl, W., & Ringler, M. (2013). Tadpole transport logistics in a Neotropical poison frog: Indications for strategic planning and adaptive plasticity in anuran parental care. *Frontiers in Zoology*, *10*(1), 67. https://doi.org/10.1186/1742-9994-10-67

Ringler, E., Rojas, B., Stynoski, J. L., & Schulte, L. M. (2023). What Amphibians Can Teach Us About the Evolution of Parental Care. *Annual Review of Ecology, Evolution, and Systematics*, *54*(1), 43–62. https://doi.org/10.1146/annurev-ecolsys-102221-050519

Ringler, E., Szipl, G., Harrigan, R. J., Bartl-Binder, P., Mangione, R., & Ringler, M. (2018). Hierarchical decision-making balances current and future reproductive success. *Molecular Ecology*, *27*, 2289–2301. https://doi.org/10.1111/mec.14583

Royle, N. J., Smiseth, P. T., & Kölliker, M. (Eds.). (2012). *The evolution of parental care* (First edition). Oxford University Press.

Salem, H., Florez, L., Gerardo, N., & Kaltenpoth, M. (2015). An out-of-body experience: The extracellular dimension for the transmission of mutualistic bacteria in insects. *Proceedings of the Royal Society B: Biological Sciences*, *282*(1804). https://doi.org/10.1098/rspb.2014.2957

Sampson, T. R., & Mazmanian, S. K. (2015). Control of Brain Development, Function, and Behavior by

the Microbiome. *Cell Host & Microbe*, *17*(5), 565–576. https://doi.org/10.1016/j.chom.2015.04.011

Santos, B., Martins, F. M. S., Sabino-Pinto, J., Licata, F., & Crottini, A. (2023). Skin and gut

microbiomes of tadpoles vary differently with host and water environment: A short-term experiment

using 16S metabarcoding. *Scientific Reports*, *13*(1), 16321. https://doi.org/10.1038/s41598-023-43340-2

Sarthou. (2001). Plant communities on a granitic outcrop. In F. Bongers, P. Charles-Dominique, P. M.

Forget, & M. Théry (Eds.), *Nouragues: Dynamics and plant-animal interactions in a neotropical*

rainforest. (p. pp 65-78). Kluwer Academic Publisher.

Scarberry, N., McAdams, Z. L., Benson, D., Herrick, J., Moore, B. C., & Ericsson, A. C. (2024).

Longitudinal characterization of the captive adult and tadpole Wyoming toad (Anaxyrus baxteri)

microbiome. *Frontiers in Amphibian and Reptile Science*, *1*, 1329897.

https://doi.org/10.3389/famrs.2023.1329897

Schulte, L. M., & Mayer, M. (2017). Poison frog tadpoles seek parental transportation to escape their

cannibalistic siblings. *Journal of Zoology*, *303*(2), 83–89. https://doi.org/10.1111/jzo.12472

Schulte, L. M., Ringler, E., Rojas, B., & Stynoski, J. L. (2020). Developments in Amphibian Parental

Care Research: History, Present Advances, and Future Perspectives. *Herpetological Monographs*, *34*(1),

71. https://doi.org/10.1655/HERPMONOGRAPHS-D-19-00002.1

Silverstone, P. A. (1973). Observations on the Behavior and Ecology of a Colombian Poison-Arrow Frog,

the Kõkoé-Pá (Dendrobates histrionicus Berthold). *Herpetologica*, *29*(4), 295–301.

Smith, S. (2023). *phylosmith: Functions to help analyze data as phyloseq objects*. https://schuyler-

smith.github.io/phylosmith/

Sommer, F., & Bäckhed, F. (2013). The gut microbiota—Masters of host development and physiology.

Nature Reviews Microbiology, *11*(4), 227–238. https://doi.org/10.1038/nrmicro2974

Sprockett, D. D., Martin, M., Costello, E. K., Burns, A. R., Holmes, S. P., Gurven, M. D., & Relman, D.

A. (2020). Microbiota assembly, structure, and dynamics among Tsimane horticulturalists of the Bolivian

Amazon. *Nature Communications*, *11*(1), 3772. https://doi.org/10.1038/s41467-020-17541-6

Sprockett, D., Fukami, T., & Relman, D. A. (2018). Role of priority effects in the early-life assembly of

the gut microbiota. *Nature Reviews Gastroenterology & Hepatology*, *15*(4), 197–205.

https://doi.org/10.1038/nrgastro.2017.173

Stückler, S., Ringler, M., Pašukonis, A., Weinlein, S., Hödl, W., & Ringler, E. (2019). Spatio-temporal

Characteristics of the prolonged Courtship in the Brilliant-thighed Poison Frog, Allobates femoralis.

Herpetologica, Accepted.

Summers, K. (1992). Mating strategies in two species of dart-poison frogs: A comparative study. *Animal*

Behaviour, *43*, 907–919. https://doi.org/10.1016/S0003-3472(06)80004-7

Summers, K., & Tumulty, J. (2014). Parental Care, Sexual Selection, and Mating Systems in Neotropical

Poison Frogs. In *Sexual Selection* (pp. 289–320). Elsevier. https://doi.org/10.1016/B978-0-12-416028-

6.00011-6

Switzer, A. D., Callahan, B. J., Costello, E. K., Bik, E. M., Fontaine, C., Gulland, F. M. D., & Relman, D.

A. (2023). Rookery through rehabilitation: Microbial community assembly in newborn harbour seals after

maternal separation. *Environmental Microbiology*, *25*(11), 2182–2202. https://doi.org/10.1111/1462-

2920.16444

Sylvain, F.-É., & Derome, N. (2017). Vertically and horizontally transmitted microbial symbionts shape

the gut microbiota ontogenesis of a skin-mucus feeding discus fish progeny. *Scientific Reports*, *7*(1),

5263. https://doi.org/10.1038/s41598-017-05662-w

Varga, J. F. A., Bui-Marinos, M. P., & Katzenback, B. A. (2019). Frog Skin Innate Immune Defences:

Sensing and Surviving Pathogens. *Frontiers in Immunology*, *9*, 3128.

https://doi.org/10.3389/fimmu.2018.03128

Villanueva, E. D., Brooks, O. L., Bolton, S. K., Savastano, N., Schulte, L. M., & Saporito, R. A. (2022).

Maternal Provisioning of Alkaloid Defenses are Present in Obligate but not Facultative Egg Feeding

Dendrobatids. *Journal of Chemical Ecology*, *48*(11–12), 900–909. https://doi.org/10.1007/s10886-022-

01394-y

Von Frieling, J., Fink, C., Hamm, J., Klischies, K., Forster, M., Bosch, T. C. G., Roeder, T., Rosenstiel,

P., & Sommer, F. (2018). Grow With the Challenge – Microbial Effects on Epithelial Proliferation,

Carcinogenesis, and Cancer Therapy. *Frontiers in Microbiology*, *9*, 2020.

https://doi.org/10.3389/fmicb.2018.02020

Vredenburg, V., Briggs, C. J., & Harris, R. (2011). Host pathogen dynamics of amphibian

chytridiomycosis: The role of the skin microbiome in health and disease. *Fungal Diseases: An Emerging*

Threat to Human, Animal, and Plant Health, 342–355.

Wake, D. B., & Vredenburg, V. T. (2008). Are we in the midst of the sixth mass extinction? A view from

the world of amphibians. *Proceedings of the National Academy of Sciences*, *105*(supplement_1), 11466–

11473. https://doi.org/10.1073/pnas.0801921105

Walke, J. B., Harris, R. N., Reinert, L. K., Rollins-Smith, L. A., & Woodhams, D. C. (2011). Social

Immunity in Amphibians: Evidence for Vertical Transmission of Innate Defenses: Vertical Transmission

of Amphibian Immunity. *Biotropica*, *43*(4), 396–400. https://doi.org/10.1111/j.1744-7429.2011.00787.x

Wang, Q., Garrity, G. M., Tiedje, J. M., & Cole, J. R. (2007). Naive Bayesian classifier for rapid

assignment of rRNA sequences into the new bacterial taxonomy. *Applied and Environmental*

Microbiology, *73*(16), 5261–5267. https://doi.org/10.1128/AEM.00062-07

Wang, S., Ryan, C. A., Boyaval, P., Dempsey, E. M., Ross, R. P., & Stanton, C. (2020). Maternal Vertical

Transmission Affecting Early-life Microbiota Development. *Trends in Microbiology*, *28*(1), 28–45.

https://doi.org/10.1016/j.tim.2019.07.010

Warkentin, K. M. (1995). Adaptive plasticity in hatching age: A response to predation risk trade-offs.

Proceedings of the National Academy of Sciences, *92*(8), 3507–3510.

https://doi.org/10.1073/pnas.92.8.3507

Warne, R. W., Kirschman, L., & Zeglin, L. (2017). Manipulation of Gut Microbiota Reveals Shifting

Community Structure Shaped by Host Developmental Windows in Amphibian Larvae. *Integrative and*

Comparative Biology, *57*(4), 786–794. https://doi.org/10.1093/icb/icx100

Warne, R. W., Kirschman, L., & Zeglin, L. (2019). Manipulation of gut microbiota during critical

developmental windows affects host physiological performance and disease susceptibility across

ontogeny. *Journal of Animal Ecology*, *88*(6), 845–856. https://doi.org/10.1111/1365-2656.12973

Weinfurther, K. D., Stuckert, A. M. M., Muscarella, M. E., Peralta, A. L., & Summers, K. (2023).

Evidence for a Parabasalian Gut Symbiote in Egg-Feeding Poison Frog Tadpoles in Peru. *Evolutionary*

Biology, *50*(2), 239–248. https://doi.org/10.1007/s11692-023-09602-7

Wells, K. D. (1980). Social behavior and communication of a dendrobatid frog (Colostethus trinitatis).

Herpetologica, *36*, 189–199.

Wells, K. D. (2007). *The Ecology and Behavior of Amphibians*. The University of Chicago Press,

Chicago and London.

Weygoldt, P. (1980). Zur Fortpflanzungsbiologie von Phyllobates femoralis (Boulenger) im Terrarium.

Salamandra, *16*(4), 215–226.

Weygoldt, P. (1987). Evolution of parental care in dart poison frogs (Amphibia: Anura: Dendrobatidae).

Journal of Zoological Systematics and Evolutionary Research, *25*(1), 51–67.

https://doi.org/10.1111/j.1439-0469.1987.tb00913.x

White, T., Bruns, T., Lee, S., Taylor, J., Innis, M., Gelfand, D., & Sninsky, J. (1990). Amplification and

Direct Sequencing of Fungal Ribosomal RNA Genes for Phylogenetics. In *Pcr Protocols: A Guide to*

Methods and Applications, (Vol. 31, pp. 315–322).

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

https://ggplot2.tidyverse.org

Woodhams, D. C., Alford, R. A., Antwis, R. E., Archer, H., Becker, M. H., Belden, L. K., Bell, S. C.,

Bletz, M., Daskin, J. H., Davis, L. R., Flechas, S. V., Lauer, A., Gonzalez, A., Harris, R. N., Holden, W.

M., Hughey, M. C., Ibáñez, R., Knight, R., Kueneman, J., … McKenzie, V. (2015). Antifungal isolates

database of amphibian skin-associated bacteria and function against emerging fungal pathogens:

Ecological Archives E096-059. *Ecology*, *96*(2), 595–595. https://doi.org/10.1890/14-1837.1

Woodhams, D. C., Vredenburg, V. T., Simon, M.-A., Billheimer, D., Shakhtour, B., Shyr, Y., Briggs, C.

J., Rollins-Smith, L. A., & Harris, R. N. (2007). Symbiotic bacteria contribute to innate immune defenses

of the threatened mountain yellow-legged frog, Rana muscosa. *Biological Conservation*, *138*(3–4), 390–

398. https://doi.org/10.1016/j.biocon.2007.05.004

Zeileis, A., Cribari-Neto, F., Gruen, B., & Kosmidis, I. (2021). *betareg: Beta Regression*.

https://CRAN.R-project.org/package=betareg

Zepeda Mendoza, M. L., Xiong, Z., Escalera-Zamudio, M., Runge, A. K., Thézé, J., Streicker, D., Frank,

H. K., Loza-Rubio, E., Liu, S., Ryder, O. A., Samaniego Castruita, J. A., Katzourakis, A., Pacheco, G.,

Taboada, B., Löber, U., Pybus, O. G., Li, Y., Rojas-Anaya, E., Bohmann, K., … Gilbert, M. P. T. (2018).

Hologenomic adaptations underlying the evolution of sanguivory in the common vampire bat. *Nature*

Ecology & Evolution, *2*(4), 659–668. https://doi.org/10.1038/s41559-018-0476-8

1358