Demographic responses to climatic changes during the Final Palaeolithic in Europe

Isabell Schmidt[1]*, Birgit Gehlen[1], Katja Winkler[2], Alvaro Arrizabalaga[3], Nico Arts[4,5], Nuno Bicho[6], Philippe Crombé[7], Berit Valentin Eriksen[8], Sonja B. Grimm[9], Katarina Kapustka[10], Mathieu Langlais[11], Ludovic Mevel[12], Nicolas Naudinot[13], Zdeňka Nerudová[14], Marcel Niekus[4,15], Marco Peresani[16,17], Felix Riede[18], Florian Sauer[1], Werner Schön[19], Iwona Sobkowiak-Tabaka[20], Hans Vandendriessche[7], Mara-Julia Weber[8], Annabell Zander[21], Andreas Zimmermann[1], Andreas Maier[1]

[1] Institute of Prehistoric Archaeology, University of Cologne, Cologne, Germany.

[2] Heinrich Schliemann-Institute of Ancient Studies, Seminar for Prehistoric Archaeology, University of Rostock, Germany.

[3] University of the Basque Country, Faculty of Arts, Vitoria-Gasteiz, Spain.

[4] Faculty of Archaeology, Leiden University, The Netherlands.

[5] Research bureau Old Land, Eindhoven, The Netherlands.

[6] Interdisciplinary Center for Archaeology and the Evolution of Human Behaviour, Universidade do Algarve, Faro, Portugal.

[7] Department of Archaeology, Prehistory Research Unit, Ghent University, Ghent, Belgium.

[8] Museum for Archaeology, Schleswig-Holstein State Museums Foundation Schloss Gottorf, Schleswig, Germany.

[9] Leibniz-Zentrum für Archäologie, Schleswig, Germany.

[10] Institute of Archaeology of the Czech Academy of Sciences, Prague, Czech Republic.

[11] CNRS Laboratoire PACEA - UMR 5199, Université de Bordeaux, France.

[12] CNRS - UMR 8068 TEMPS MSH MONDES, Nanterre, France.

[13] Muséum national d'Histoire naturelle – CNRS HNHP UMR 7194, Paris, France.

[14] Centre for Cultural Anthropology, Moravian Museum, Brno, Czech Republic.

[15] Stichting STONE/Foundation for Stone Age Research in the Netherlands, Groningen, The Netherlands.

[16] Department of Humanities, Prehistoric and Anthropological Sciences Unit, University of Ferrara, Ferrara,
Italy.

[17] Institute of Environmental Geology and Geoengineering, National Council of Research, Milano, Italy.

[18] Department of Archaeology and Heritage Studies, Aarhus University, Denmark.

[19] Department of Old World and Asian Studies, University Erlangen-Nürnberg, Germany.

[20] Faculty of Archaeology, Adam Mickiewicz University, Poznań, Poland.

[21] Department of Archaeology, University of York, York, United Kingdom

* Corresponding author

E-mail: isabell.schmidt@uni-koeln.de (IS)

Abstract

The European Final Palaeolithic witnessed marked changes in almost all societal domains. Despite a rich body of evidence, our knowledge of palaeodemographic processes and regional population dynamics still needs to be improved. In this study, we present regionally differentiated estimates of absolute numbers and population densities for the Greenland Interstadial 1d-a (GI-1d-a; 14-12.7 ka BP) and the Greenland Stadial 1 (GS-1; 12.7-11.6 ka BP) for Southern, Western, Northern and Central Europe. The data were obtained by applying the Cologne Protocol, a geostatistical approach for estimating prehistoric population size and density, to a newly compiled dataset of Final Palaeolithic sites. On a large spatio-temporal scale, we observe a shift of the main areas of human occupation from the Franco-Cantabrian region, which was intensely occupied during most phases of the preceding Upper Palaeolithic, to regions north of the Alps. At smaller scales, we observe divergent regional trends in the Final Palaeolithic meta-population: during GI 1d-a, a decreasing population in southwestern Europe and an increasing population in north-eastern Central Europe. For the first time since the dispersal of anatomically modern humans into Europe, we see that Central Europe becomes the dominant demographic growth area. Subsequently, the climatic cooling of GS-1 coincides with a pronounced population decline in most parts of the study area. An apparent increase in population density occurs only in north-eastern Central Europe and north-eastern Italy. Our estimates suggest that the total population was reduced by half. Similar results, with a relationship between decreasing temperatures and decreasing populations, have already been observed for the late phase of the Gravettian, when populations were reduced to only one third of those estimated for the early phase. Yet, in contrast to the collapse of local populations during the late Gravettian, the increase in population densities in Central Europe during GS-1 indicates population movements eastwards, possibly in response to deteriorating climatic conditions, particularly in western regions during the Younger Dryas.

Keywords: Allerød (≈ GI-1d-a), Younger Dryas (≈ GS-1), technocomplex, Late Glacial ecology, demography, human population movement

Introduction

The Late Glacial, comprising Greenland Interstadial 1 (GI-1) and Stadial 1 (GS-1), was a period

when the impact of rapidly changing environmental conditions drove high-magnitude and high-amplitude

ecosystem shifts. The extensive dispersal of shrub and tree species into northern Western and Central Europe

alongside the associated fauna during GI-1 contributed to creating a European habitat with particularly

pronounced regional differences. Although the so-called 'mammoth steppe' [1] of preceding periods was

also never a uniform habitat [2], this emerging, regionally diverse ecotonal patchwork presumably had

consequences for most fundamental properties of hunter-gatherer societies, such as subsistence strategies,

social networks, and – prominently – the structure and distribution of the underlying meta-population.

In Europe, these changes are culturally associated with the transition from the Late Upper to the

Final Palaeolithic period (e.g., [3–6]). Subsistence strategies change from hunting migratory game to a

preference for more stationary, thermophilous game. Settlement and land use patterns are marked by sites

with functionally differentiated areas, such as Le Closeau or Reichwalde [7,8] rather than large

agglomerative sites, such as Gönnersdorf [9–11]. Changes in lithic technology, particularly in core reduction

strategies, signalled a less normative use of lithic raw material of varying quality, alongside a regionalisation

in lithic projectile morphology [3,12–18]. Visual representations are no longer found in caves but begin to

appear frequently on pebbles. In addition, abstract representations become much more dominant than

figurative ones. From a demographic perspective, these profound changes went hand in hand with expanding

hunter-gatherer groups into the northern European lowlands. From the extensive dispersal of anatomically

modern humans into Europe during the Aurignacian up to the repopulation of Central Europe after the Last

Glacial Maximum, the meta-population in Europe had undergone substantial fluctuations regarding absolute

numbers and densities of people as well as social interconnectedness of sub-populations and spatial

distributions of genetic haplotypes [19–23]. In its broadest outline, however, the large-scale distribution

pattern of hunter-gatherers remained remarkably stable, with one demographic centre in the Franco-

Cantabrian region and another in eastern Central Europe, covering Lower Austria, Moravia, southern

Poland, and northern Hungary. In the Final Palaeolithic, this pattern changed drastically.

This change occurred between 15 and 14 ka BP, but there is no generally accepted date or event

marking the end of the Late Upper and the beginning of the Final Palaeolithic in Western and Central

Europe. Particularly problematic for a chronological delimitation is a plateau in the calibration curve during

this period that causes wide calibration age ranges [24,25] and, thus, reduces chronological resolution.

Further complicating matters are regionally differentiated environmental responses to the climatic shift at

the transition from the Late Pleniglacial to GI-1e, which appears rather abrupt in the Greenland ice core

records and may have been related to a substantial meltwater discharge into the North Atlantic [26] and

tipping points in sea-ice cover [27]. Atmospheric warming over Europe would have started well before, and

the associated ecosystem change likely was more gradual [28,29]. There is growing evidence that such an

early start and comparatively slow change of ecosystems did not only take place south of the Alps [30,31],

but also in Central Europe [32,33]. Variable response periods of the vegetation in diverse parts of Europe

(e.g., [33–36]) further indicate regionally distinct ecosystem reactions differentiated by both leads and lags.

Such differences are not limited to warming events but also occurred during the cooling of GS-1 ($\approx$ Younger

Dryas) and subsequent warming when de- and increases in seasonal temperature and precipitation varied

markedly between different areas throughout Europe [37–41]. An earlier and more gradual ecosystem

change prior to GI-1e also has consequences for our understanding of the timing of human population

dispersal into Northern Europe. It is in accord with an apparently earlier northward expansion of hunter-

gatherers of the Creswellian and Hamburgian [42–44].

From a palaeogenetic point of view, the limited data of this period indicates a major turnover

between 15 and 14 ka BP, during which the so-called 'Oberkassel cluster' became the dominant haplogroup

in Central Europe [23]. Thus, while it is difficult to pinpoint the start of the Final Palaeolithic, it is undisputed

that archaeological phenomena after 14 ka BP (post-GI-1e) are attributed to the Final Palaeolithic

[3,18,45,46]. Therefore, we chose this date as the older boundary of our study, while the younger boundary

coincides with the end of GS-1 and the end of the Pleistocene at 11.6 ka BP.

Here, we present regionally differentiated estimates of absolute numbers and densities of people for

two phases of the Final Palaeolithic according to the Cologne Protocol [47]. A previous study considered

the Final Palaeolithic as a whole [21], but limited data granularity has hitherto precluded analyses at a finer

resolution. Through the involvement of regional experts from different countries, we compiled an extensive

database [48], allowing for a chronological subdivision of the assemblages into two phases. The older phase,

dates between 14 and 12.7 ka BP and thus corresponds to GI-1d-a or the Older Dryas and the Allerød *sensu*

Iversen [49], while the younger phase dates to between 12.7 and 11.6 ka BP and thus corresponds to GS-1

or the Younger Dryas (**Fig 1**). Based on our estimates, we address how the human meta-population

responded demographically to environmental change associated with the climatic cooling from GI-1d-a to

GS-1 at a subcontinental and regional spatial scale. We compare these results with the demographic response

during the Gravettian [50], that was similarly linked to major cooling, and discuss similarities and

differences, as well as implications for social networks and behavioural adjustments.

Fig 1. Chronological framework of this study. The time-bins defined in the upper part of the graph indicate the different typo-chronological resolutions of the assemblages that were taken into account for the database and analysis.

Material and Methods

The database underlying this study comprises archaeological sites from Southern, Western, Northern and Central Europe. It has been compiled with the regional expertise of the participating authors and is available online as an open-access resource [48]. To benchmark this detailed recording of the evidence across the study area, unpublished (and thus not verifiable) material and single finds were not considered. Some differences, however, turned out to be inevitable due to data accessibility (publication, database management of heritage authorities) or regionally varying properties of the record (e.g., chronological resolution). In sum, the database considers material from 1788 sites (**Fig 2**, excavations and surface collections). 1548 assemblages can either be directly dated and/or typologically or stratigraphically assigned to one of the two phases. GI-1d-a (14-12.7 ka BP) comprises 988 assemblages and GS-1 (12.7-11.6 ka BP),

comprises 560 assemblages. Material from assemblages that cover larger time-bins (see **Fig 1**) contain 237

assemblages assigned to the Final Palaeolithic in general, 131 assemblages assigned to the Final Palaeolithic

and the Preboreal in general, and 75 assemblages are recorded separately because their chronological

placement relates to the final GS-1 and the succeeding Preboreal phase. Frequent generic attribution is most

problematic for evidence from Central Europe.

Fig 2. Distribution of Final Palaeolithic sites in the study area. Green dots: GI-1d-a; blue dots: GS-1.

Filled circles: dataset B (chronologically subdivided sites); filled and open circles: dataset A (all sites; see

text and **Fig 1**. Base map details: see supporting information).

Submerged coastal areas and so-called Doggerland provided large, formerly inhabitable

landmasses. Evidence of human occupation from these areas is extremely scarce and, therefore, virtually

absent in the dataset. Examples of long-distance raw material transport, e.g., of Helgoland flint [51–53],

speak to human presence across these now submerged areas. If diagnostic archaeological artefacts are

recovered from below current sea level, they often lack geospatial information. Our calculations of the

estimates, regional comparisons and subsequent interpretations of the results therefore rest upon the

available evidence from the current landmass, as mapped in **Fig 2**.

On the North European Plain and adjacent areas, GI-1d-a mainly comprises sites attributed to the

Federmesser-Gruppen or *Federmesser* culture (e.g., [54–59]). The beginning of this phase overlaps slightly

with the Havelte phase of the Hamburgian [42,46,56,60,61] and potentially with the latest Creswellian sites

[54,62,63]. Given that most sites attributed to these industries predate GI-1d-a (see also [64]), Havelte and

Creswellian assemblages were consistently excluded from the dataset. The disappearance of these 'cultures'

may in itself be indicative of a demographic break at this important tipping-point transition [44,56,65] as

indicated by results from the British Isles [66]. During GS-1, assemblages assigned to the Ahrensburgian

and Swiderian occur in the Central and Northern lowlands [58,67–69], at time persisting into the subsequent

Preboreal period (e.g., [70]). The position of the Bromme culture in this framework, situated at the transition

from Allerød to Younger Dryas, is subject to discussion [56,71,72]. A high number of surface collections

characterises the archaeological evidence on the northern Plains in general.

In the Benelux, Ahrensburgian and Epi-Ahrensburgian assemblages [73–77] replace the

Federmesser-Gruppen during GS-1, with most dates placing the Ahrenbsurgian notably late into GS-1 or

even into the Holocene [78]. Long Blade industries [79,80] and Belloisian/Long Blade assemblages [80–

84] emerged in the UK and the Paris Basin, respectively. Assemblages with so-called 'Riesenklingen' (giant

blades) also occur in the Netherlands [76], north-western Germany (e.g., [85]) and Denmark [86], where

they are assigned to the younger half of GS-1 and occasionally seem to diffuse into the Preboreal [87].

In France, the term 'Azilian' is used in competition with 'Federmesser-Gruppen'. The latter is more

commonly used in northern France, particularly in the Somme valley. This distinction has more to do with

taxonomy and the history of research [88] than with any archaeological reality. However, differences cannot

be overlooked typologically and technologically, especially when comparing certain sites (e.g., Le Closeau

vs Saleux), although it is not yet possible to explain the exact meaning of this variability. This taxonomic

question also applies to the 'Belloisian', 'Laborian' and 'Epi-Ahrensbourgian' entities, which are sometimes

supposed to be part of the same technical trend ('FBBT'; [89–92]). The situation is, of course, different for

assemblages related to the different phases of the Epigravettian period, and particularly to the most recent

ones. They reflect the probable spread of these traditions beyond southeastern France - at least along the

Rhone corridor and certainly as far as the Jura range [93–96] - during the GS-1 period. In the actual state of

knowledge, we can highlight offsets between their chronometric boundaries and typological changes

[83,90,97–99].

Along the Atlantic façade of the Iberian Peninsula, regional chronologies differ in the assignment

of taxonomic units to certain periods [100–103]. In Portugal, the Magdalenian is seen as ending before the

Younger Dryas. The following phase is frequently called Epipalaeolithic [102,104] or Early Mesolithic

[103]. Either way, the lithic assemblages appear very similar in the two phases in both technology and

typology. In Cantabrian Spain and Pyrenees, Azilian is the most used term.

In Italy, complexes dated to the Late Glacial interstadial and the Younger Dryas are framed within

the late and terminal Epigravettian, although differences exist between regions in chronology and

terminology [105,106]. Indeed, from the end of the Last Glacial Maximum up to the Pleistocene/Holocene

boundary, the Epigravettian modified its technological features, the typology of tools and of weapons, as

related to the use of bow and arrow, particularly in the latest phases [107]. The shortening of certain tools,

such as frontal end-scrapers [108], and the spread of backed knives occur alongside with the appearance of

geometric microliths and the increasing use of the microburin blow technique. These modifications seem to

be shared at larger scale in Eastern and Mediterranean Europe [12].

The greatest difficulties for a chronological subdivision were encountered along the northern range

of the Alps, particularly in southern Germany and the Czech Republic. In contrast to many other areas, there

is no discernible shift in typology between GI-1d-a and GS-1 in south-eastern Germany, and assemblages

of both phases are attributed to the *Federmesser-Gruppen* [109]. The absence of chronometric data further

complicates matters since most assemblages are surface collections and organic preservation is poor. In the

Czech Republic, *Federmesser-Gruppen* are only present in the north-western part of Bohemia towards the

end of GS-1 [110,111]. In contrast, assemblages of GI-1d-a and the early GS-1 and in other parts of the

country are at times assigned to the Epimagdalenian or Final Palaeolithic. However, many sites are surface

collections and can therefore not be securely associated with either of the two phases [112,113]. In the

sandstone regions, stratified and radiometrically dated sites are known [33,114,115], which may help refine

the chronology of the Final Palaeolithic in this region. Better dated sites and assemblages occur in

southernmost Germany and Switzerland [116–121].

Alongside the database on Final Palaeolithic sites and assemblages, we collected data on lithic raw

material transport from the literature. Where possible, the information was divided into the two

chronological phases (**Fig 1**), too. If a site was known to contain assemblages from both periods but raw

material data are presented in the literature generically as 'Final Palaeolithic', these data were assigned to

both periods equally. If a site was known to be only generically assigned to the Final Palaeolithic, the raw

material data was recorded but not considered for the phase-specific calculation of the demographic

estimates. This accounts especially for sites from the area of SE Germany and the Czech Republic (e.g.,

[109,110]). We also excluded raw materials documented by single pieces and only considered raw materials

representing at least 1% of an assemblage (see also [47]). In a second step, we determined the Raw Material

Catchment Areas (RMCA) for each assemblage by adding a 5 km radius buffer around the archaeological

site and constructing a convex hull around the site-buffer and the related raw material sources. We excluded

RMCAs which only comprised a single source-to-site distance, as well as RMCAs which were smaller than

500 km^2 since they turned out to be exceptional outliers among the overall RMCA sizes (these assemblages

are from Grotte du le Bichon and Blot for GI-1d-a and Gahlen-Schermbeck, Bad Breisig and Babí Pec for

GS-1). Eventually, for GI-1d-a, 53 out of 108 RMCAs and for GS-1, 39 out of 97 RMCAs were used to

calculate demographic estimates (**S1 and S2 Table**). The scarcity or even lack of available raw material

data in some Core Areas (e.g., Great Britain, Denmark, and northern Spain) required transferring RMCA-

sizes from adjacent areas.

The calculation of population estimates follows the Cologne Protocol [47]. In brief, it statistically

divides a certain map section, called Total Area of Calculation, into areas of high hunter-gatherer activities,

called Core Areas, and areas of little or no activities. The Core Areas are delimited by calculating areas with

equal site densities, encircled by the so-called optimally describing isoline. To account for differences in

site density within Europe, the optimally describing isolines have been modelled individually for three

regions (for details, see **S1 and S2 Figs**). The Total Area of Calculation for the current study is 2.6 million

km^2 of Southern, Western, Northern, and Central Europe, following the current coastlines. To estimate the

number of hunter-gatherer groups per Core Area, the size of each optimally describing isoline (in km^2) is

divided by the quartiles (Q1, Q2, Q3) of the related RMCA. The results are then multiplied with an average

group size (42.5) derived from ethno-historic cases selected based on socio-economic considerations (see

[47,122]), providing the estimate of the absolute number of people. Population densities are then calculated

for each Core Area by dividing the respective numbers of people by the size of the Core Area, and for the

Total Area of Calculation by dividing the number of all people by the size of the selected map section.

To account for the fact that a chronological subdivision of the dataset was more feasible in some

regions and less so in others, we decided to calculate for each phase two demographic estimates based on

different datasets: the so-called 'dataset A' includes all sites, subdivided assemblages as well as sites that

are only generically assigned to the Final Palaeolithic period or the time-bins indicated in **Fig 1**. Information

about human presence recorded from sites with low chronological resolution is thus retained in the

modelling, which may then, however, lead to overestimations for a given phase at regional scales. The

second demographic estimate, based on 'dataset B', in contrast, considers only assemblages with a clear

chronological subdivision assigned to them, attributable either to GI-1d-a or to GS-1, or sites that can be

clearly assigned to both phases. Due to the overall smaller sample size of 'dataset B', it is expected to

produce lower estimates – likely underestimating the number of people – than the ones derived from dataset

A, but it provides the best available high chronological control. All estimates are given as a mean estimate

and, based on our calculation protocol, with a minima and maxima estimate (related to Q1 and Q3 of the

RMCA).

292   Results

In the following, we will discuss the results of the estimates from dataset A and dataset B for each

chronological phase, first considering dataset A with all sites, and then the smaller but chronologically

highly resolved dataset B.

For the estimates from dataset A, all Core Areas of GI-1d-a (**Fig 3**) enclose an area of approximately

635,000 km^2 and contain 90% of the sites (1220 out of 1356), while those of GS-1 (**Fig 4**), in contrast,

enclose only about 481,000 km^2 and contain 82% of the sites (825 out of 1003; for further details, see **S1**

Fig). The mean estimate for the total population for GI-1d-a amounts to roughly 8100 people at any given

time, with minima and maxima ranging from 4800 to 14,300 (**Table 1**). For the GS-1, the mean estimate

for the total population drops to 4250 people living at the same time, with minima and maxima ranging from

2200 to 8800 people (**Table 2**). This decrease in population size is accompanied by a decrease in population

density within the Core Areas from a mean value of 1.3 people per 100 km² (p/100km²) in GI-1d-a (min.

0.8, max. 2.2) to 0.9 p/100km² in GS-1 (min. 0.5, max. 1.8). In the regional results, the Czech Republic and

SE Germany are combined by the Core Area and have, with around 2500 people, the largest and with 2.2

p/100 km² also the densest population in GI-1d-a. However, they also show the strongest decline to only

460 people and 0.9 p/100 km² in GS-1. The second highest estimates for GI-1d-a are found in northern Spain

and the French Pyrenees with 1200 people, closely followed by Poland and NE Germany with 1100 people.

Despite these similar numbers, the density varies considerably between 1.7 p/100 km² in the former and 0.8

p/100 km² in the latter areas. For the total area of calculation ('study area' in **Fig 2**), the estimated population

density drops from 0.003 in GI-1d-a to 0.002 p/km² in GS-1.

Fig 3. Site distribution and modelled Core Areas for GI-1d-a. The data from dataset A (thick lines, empty and filled dots) and dataset B (shaded areas, filled dots only) are shown. For details on areal data and population estimates see **Table 1** (data A) and **S3 Table** (data B). Reconstructions of ice sheets, river channels, and coastline (>45°N) correspond to GI-1c-a conditions (see supporting information and [3] and references therein).

Fig 4. Site distribution and modelled Core Areas for GS-1. The data from dataset A (thick lines, empty and filled dots) and dataset B (shaded areas, filled dots only) are shown. For details on areal data and population estimates see **Table 2** (data A) and **S4 Table** (data B). Reconstructions of ice sheets, river channels, and coastline (>45°N) correspond to GS-1 conditions (see [3] and references therein).

329   **Table 1. Regionally distinguished demographic estimates for GI-1d-a based on dataset A (all sites).**

Core Area region	ODI (km²)	Q	RMCA (km²)	N_{raw}	N_{groups}	N_{people}	$D_{population}$
		1	2670		*19.4*	*826*	*0.016*
Great Britain	51,900	2	5117	17	**10.1**	**431**	**0.008**
		3	5896		*8.8*	*374*	*0.007*
		1			*7.5*	*317*	*0.016*
N France	19,940	2			**3.9**	**166**	**0.008**
		3			*3.4*	*144*	*0.007*
		1			*35.2*	*1,496*	*0.016*
Benelux & NW Germany	94,010	2			**18.4**	**781**	**0.008**
		3			*15.9*	*678*	*0.007*
		1			*10.4*	*441*	*0.016*
S Scandinavia	27,700	2			**5.4**	**230**	**0.008**
		3			*4.7*	*200*	*0.007*
		1			*50.0*	*2,125*	*0.016*
Poland & NE Germany	133,500	2			**26.1**	**1,109**	**0.008**
		3			*22.6*	*962*	*0.007*
		1	1057		*109.7*	*4,661*	*0.040*
Czech Rep. & SE Germany	115,900	2	1948	10	**59.5**	**2,529**	**0.022**
		3	2626		*44.1*	*1,876*	*0.016*
		1	2604		*12.9*	*547*	*0.016*
Switzerland & SW Germany	33,540	2	3550	19	**9.4**	**402**	**0.012**
		3	8662		*3.9*	*165*	*0.005*
		1			*5.8*	*246*	*0.016*
Italy	15,050	2			**4.2**	**180**	**0.012**
		3			*1.7*	*74*	*0.005*
		1			*9.8*	*415*	*0.016*
SE France	25,440	2			**7.2**	**305**	**0.012**
		3			*2.9*	*125*	*0.005*
		1	1578		*24.9*	*1,058*	*0.027*
SW France	39,290	2	2522	7	**15.6**	**662**	**0.017**
		3	20730		*1.9*	*81*	*0.002*
		1			*44.2*	*1,878*	*0.027*
Spain & French Pyrenees	69,720	2			**27.6**	**1,175**	**0.017**
		3			*3.4*	*143*	*0.002*
		1			*5.9*	*252*	*0.027*
Portugal	9,351	2			**3.7**	**158**	**0.017**
		3			*0.5*	*19*	*0.002*
		1			*336*	*14,262*	*0.022*
Sum Core Area	635,341	2		53	191	8,126	0.013
		3			*114*	*4,839*	*0.008*

		Q					$D_{population}$
		1					*0.006*
Total Area of Calculation	**2,600,000**	2					**0.003**
		3					*0.001*

330 Thin horizontal lines indicate regions with transfer of RMCA data (see **S3 Fig**). ODI = optimally describing

331 isoline in km²; Q = Quartile of RMCA [1 = lower, 2 = median, 3 = upper]; RMCA (km²) = raw material

332 catchment area in km²; N_{raw} = number of RMCAs; N_{groups} = number of groups; N_{people} = number of people;

333 $D_{population}$ = population density within Core Areas as p/km², and the Total Area of Calculation (bottom rows).

335 **Table 2. Regionally distinguished demographic estimates for GS-1 based on dataset A.**

Core Area region	ODI (km²)	Q	RMCA (km²)	N_{raw}	N_{groups}	N_{people}	$D_{population}$
		1	2934		*8.8*	*374*	*0.014*
Great Britain	25,830	2	5187	4	**5.0**	**212**	**0.008**
		3	7462		*3.5*	*147*	*0.006*
		1			*4.8*	*202*	*0.014*
N France	13,950	2			**2.7**	**114**	**0.008**
		3			*1.9*	*79*	*0.006*
		1			*17.9*	*761*	*0.014*
Benelux & NW Germany	52,500	2			**10.1**	**430**	**0.008**
		3			*7.0*	*299*	*0.006*
		1			*0.5*	*20*	*0.014*
S Scandinavia	1,350	2			**0.3**	**11**	**0.008**
		3			*0.2*	*8*	*0.006*
		1			*50.8*	*2,159*	*0.014*
Poland & NE Germany	149,000	2			**28.7**	**1,221**	**0.008**
		3			*20.0*	*849*	*0.006*
		1	970		*55.3*	*2,352*	*0.044*
Czech Rep. & SE Germany	53,670	2	4959	11	**10.8**	**460**	**0.009**
		3	14826		*3.6*	*154*	*0.003*
		1	3160		*7.3*	*310*	*0.013*
Switzerland & SW Germany	23,050	2	4336	14	**5.3**	**226**	**0.010**
		3	9988		*2.3*	*98*	*0.004*
		1			*10.3*	*437*	*0.013*
Italy	32,500	2			**7.5**	**319**	**0.010**
		3			*3.3*	*138*	*0.004*
		1			*6.2*	*262*	*0.013*

SE France	19,480	2			**4.5**	**191**	**0.010**
		3			*2.0*	*83*	*0.004*
		1	2420		*10.2*	*434*	*0.018*
SW France	24,740	2	4337	10	**5.7**	**242**	**0.010**
		3	13094		*1.9*	*80*	*0.003*
		1			*31.3*	*1329*	*0.018*
Spain & French Pyrenees	75,710	2			**17.5**	**742**	**0.010**
		3			*5,8*	*246*	*0.003*
		1			*3.6*	*154*	*0.018*
Portugal	8,780	2			**2.0**	**86**	**0.010**
		3			*0.7*	*28*	*0.003*
		1			*207*	*8794*	*0.018*
Sum Core Area	**480,560**	2		39	**100**	**4254**	**0.009**
		3			*52*	*2209*	*0.005*
		1					*0.003*
Total Area of Calculation	**2,600,000**	2					**0.002**
		3					*0.001*

Thin horizontal lines indicate regions with transfer of RMCA data (see **S4 Fig**). For an explanation on

abbreviations see Table 1.

For the estimates from dataset B (see **S3 and S4 Tables**), the size of all Core Areas of GI-1d-a (**Fig**

3: shaded areas) is reduced to 489,000 km² (85% of the sites; 842 out of 988), and those of GS-1 to

392,000km² (**Fig 4**: shaded areas; 74% of the sites; 415 out of 560). The mean estimate for the total

population for GI-1d-a sums up to only roughly 5700 people (min. 3100, max. 9900) and to 3500 people

(min. 1900, max. 6200) for GS-1. The mean density of GI-1d-a, 1.2 p/100 km² (min. 0.6, max. 2.0), in

contrast, is only slightly lower than for dataset A, and remains at 0.9 p/100 km² for GS-1 (min. 0.5, max.

1.6). The largest regional population for dataset B estimates during GI-1d-a is found in northern Spain and

the French Pyrenees with 1100 people, followed by Benelux and NW Germany with 820 people. With 770

people, NE Germany and Poland display estimates similar to those of the Czech Republic and SE Germany.

The largest differences between estimates from datasets A and B are observed during GI-1d-a

especially for the Czech Republic and SE Germany, but also for the area of Poland and NE Germany and

for Switzerland and SW Germany (**Fig 5, for details, see S5 Fig**). For GS-1, differences are observable for

the Czech Republic and SE Germany, where dataset A estimates around 460 people and dataset B around

80 people (**S6 Fig**). The density for this area, however, remains at 2.2 p/100 km² for GI-1d-a and 0.9 p/100

km² for GS-1, fully in line with the dataset A estimates. For the total area of calculation the estimated

population density drops in dataset B from 0.002 in GI-1d-a to 0.001 p/km² in GS-1.

Fig 5. Comparison of demographic estimates from GI-1d-a and GS-1. Depicted are the mean estimates

from dataset A (bright fill) and dataset B (dark fill).

A consistent signal in our palaeodemographic reconstructions is a strong decline in the meta-

population from GI-1d-a to GS-1 (**Fig 5**). In Portugal, SE France, and S Scandinavia, likewise Czech

Republic and SE Germany, the regional population may have dropped below a hundred people. Notable

exceptions from this trend are Poland, NE Germany, and Italy (especially the northeastern area, compare

Figs 4 and **5**). Moreover, in all estimates, the former region hosts the largest regional population of around

1200 during GS-1, thus being almost twice as large as the second highest regional population (data A = 740;

data B = 590) located in (northern) Spain and the French Pyrenees. In Switzerland and SW Germany, the

estimates from dataset B result in a relatively stable population (data A = 260, data B = 230).

Discussion

Large-scale distribution patterns of Palaeolithic sites are influenced by several confounding factors

that lie outside the decision-making processes of past hunter-gatherers. Many other parameters can influence

the likelihood of site discovery, such as the geomorphological development of a landscape as shown, for

instance, in the Great Adriatic Po-Plain [123] and other coastal areas [124]. Modern land use (e.g., forestry,

vineyards, ploughed fields, quarries) or specific taphonomic settings (presence of caves and rock shelters or

wetlands; loess or coversand deposition) can also play an important and sometimes even dominant role

[125]. Other factors are the research histories of the different areas, in some cases linked to the proximity

of universities and other institutes (research-driven versus development-led archaeological excavations)

with strong prehistoric research traditions, the accessibility of collected data (such as regional or national

databases, or grey literature) and their revision by experts (cf. [67,109,126,127]). Finally, differences in

artefact size or the recognisability of iconic artefacts, such as the tanged points of the Ahrensburgian and

Swiderian, can skew the observed patterning, either by making them harder to find (cf. [73,128]), or by

skewing site numbers towards those characterised by particularly large artefacts (cf.[129]). These biases

need to be discussed and necessitate caution but should allow research to address large-scale questions such

as population dynamics in general [15].

The distribution of Final Palaeolithic sites in the study area (**Fig 2**) is notably uneven but comparable

between GI-1d-a and GS-1 (**Figs 3 and 4**). Generally – and in sharp contrast to all periods of the preceding

Upper Palaeolithic – the main area of the human population is not located in the Franco-Cantabrian region

but in regions north of the Alps. Given that modern land use will not affect Upper and Final Palaeolithic

sites differently and given further that the Late Glacial warming made higher-latitude areas newly accessible

391 for Final Palaeolithic hunter-gatherers, it seems highly unlikely that this large-scale spatial pattern is

392 exclusively the result of research biases [125]. At smaller scales, however, the effects of research biases

393 might be more pronounced. For our dataset, this is particularly relevant for a conspicuous void that stretches

394 in the south-west to north-east direction from southwestern France up to the Upper Rhine Valley, separating

395 the investigated area into a south-western and north-eastern part. Final Palaeolithic sites are scarce in this

396 void, but it is unclear whether this reflects prehistoric reality or results from multiple biases listed above.

397 The latter view is supported by the absence of strong research traditions and academic structures in this

398 region compares with, for instance, the southwest or the northeast of France and a relative scarcity of

399 contexts favouring preservation and recognition of Upper and Final Palaeolithic sites. Difficulties in locating

400 raw material sourcing sites and a general scarcity of typological markers (e.g., at Le Closeau) might

401 additionally hamper the attribution of sites to the Final Palaeolithic in some areas. Also, modern population

402 densities and industrial activity are either low or very high in some of the western areas of Germany – some

403 sites are surely waiting to be discovered while others have long since been destroyed. The comparably long

404 and strong archaeological research tradition here, however, speaks in favour of the existence of a void, at

405 least in Germany (i.e. Hesse and western Thuringia). A comparison of the extent of the void with Core Areas

406 calculated for the immediately preceding period of the Final Magdalenian (**Fig 6,** [122]), shows that the

407 void widens considerably in the Final Palaeolithic towards the southwest and northeast. Core Areas are

408 modelled in the void, at least for the French area, and also during earlier periods of the Upper Palaeolithic

409 record, the early Gravettian and Solutrean technocomplexes [50,130]. It is difficult to see why potential

410 biases would particularly affect Final Palaeolithic sites, leaving those of the Upper Palaeolithic in place.

411 Also, differences in material culture on either side of the void (e.g. occurrences of painted cobbles), a lack

412 of evidence for objects that were transported across it, and hints from palaeogenetic data warrant a closer

413 examination of this observation.

Fig 6. Comparison of Core Areas from the Final Magdalenian (ca. 16-14 ka BP, orange areas, after [122]) to Core Areas of the GI-1d-a (green areas). For details on background map, see Fig 3.

In order to assess biases by typological visibility, we had a look at the number of radiocarbon-dated sites in both phases based on the entries in the Radiocarbon Palaeolithic Europe Database v.27 ([131]; we are aware of a later version but used v.27 when conducting this study). The analysis shows similar trends to our demographic estimates. For the entire map section and in absolute numbers, there is a decline of more than 50 % in radiocarbon dated sites from GI-1d-a to GS-1 (**S5 Table**). Regionally differentiated and in

relative terms (**S7 Fig**), we see a decline in all areas, but it is specifically pronounced in Central Europe. A

fundamental difference, however, is our increasing estimates for NE Italy, Poland, and NE Germany (**Fig**

5). It is thus possible that our estimates for these regions are positively skewed due to greater typological

recognisability of assemblages produced during GS-1. The estimated demographic increase, however, can

also be the result of behavioural responses to contemporaneous climate change, such as population

movements towards the east.

A comparison to other palaeodemographic studies might also reveal potential biases. Yet, in

contrast to periods of the Upper Palaeolithic, such a comparison is difficult for the Final Palaeolithic

estimates because of divergent chronological frameworks. For instance, the youngest period in the study by

Bocquet-Appel et al. [132], is termed 'Late Glacial' and comprises the interval roughly between 19.1 and

13.3 ka BP. Tallavaara et al. [20] provide estimates in 1 ka steps, but the last two steps are centred at 14 and

13 ka BP, respectively, therefore essentially missing the Younger Dryas. While the former period is too

large for a meaningful comparison, the latter provides two estimates at the fringes for our first phase and

none for the second phase of GS-1. The chronologically matching study by French and Collins for

southwestern France [19] also sees a decline in demographic signatures from GI-1d-a to GS-1. A revision

of the Final Magdalenian assemblages in this area resulted in a reattribution to the Laborian, chronologically

placing the sites to the GS-1 [133]. These findings and very recent research developments on the Laborian

could change the demographic picture here. Rather more synoptically on a European scale, Ordonez and

Riede [134] reconstruct forager population densities with reference to ethnographic data and climatic

limiting factors as derived from high-resolution climate data. Their study similarly documents a population

downturn during GS-1.

A large-scale shift to the east as a potential reaction to the GS-1 cooling

At a large spatial scale, a ~25% decrease in the size of the Core Areas from GI-1d-a to GS-1 is

observable in most parts of the investigated area and across both datasets A (**Fig 7**) and B (**S8 Fig**). At the

same time, Core Areas in western Britain, Brittany, and Denmark disappeared, whereas new Core Areas

formed in eastern Poland, particularly in Masuria. Core Areas in Iberia and southwestern France shrank but

remained rather stable in their geographic location. Notably, in the rest of the investigated area, Core Areas

showed a pronounced geographic shift towards the east during GS-1. Given that it seems unlikely that this

large-scale shift is exclusively caused by biases (see discussion above), the question arises as to whether it

can be seen as a reaction to changing climatic conditions. Here, a comparison to the developments during

the Gravettian is instructive. Demographic studies, which compared the internal shift from an early phase

(33-29 ka BP) to a late phase (29-25 ka BP) of the Gravettian is the only other instance since the arrival of

anatomically modern humans in Europe where we see a pronounced demographic decline coinciding with

climatic cooling [50]. During the late phase of the Gravettian, the estimates indicate a reduction across all

regions, suggesting that no or only little population movement occurred over larger distances. This may

indicate that a behavioural response to environmental change in terms of mobility did not happen or that

population movement was not a sufficiently effective risk mitigation strategy, given that the climatic

changes affected all of Europe [135]. Instead, it seems that the regional populations north of 50° latitude

collapsed entirely, leaving western Central Europe largely depopulated until the Late Magdalenian [50,136].

This is in accord with palaeogenetic findings, indicating a continuity in western and southwestern Europe

(Fournol cluster) and a disappearance of traits associated with the Věstonice cluster in Central and Southern

Europe [23]. Interestingly – and in stark contrast to the Gravettian – the estimates for the Final Palaeolithic

show a regionally more differentiated demographic pattern. In the following, we discuss the environmental

change of the Younger Dryas in more detail, focusing on regional differences that might have contributed

to the observed pattern.

Fig 7. Comparison of Final Palaeolithic Core Areas (based on data A) from GI-1d-a to GS-1. For a
comparison of data B Core Areas, see **S8 Fig**. Grey diamonds (after [137]) indicate documented ash
(=tephra, microtephra or heavy minerals) from the Laacher See eruption, dated to 13 ky BP [138].

The Younger Dryas (GS-1) is a phenomenon driven by processes in the North Atlantic and with
partly varying effects on the ecosystems in different parts of Europe, often with a west-east-trending gradient
(e.g., [34–37,41,109,139,140]). In northern Germany, western Belgium [141] and Poland [142], for
instance, the climate of GS-1 drove the return of open landscapes, while the vegetation in Switzerland and

eastern France seems to have been affected only weakly [143–145]. In a general comparison to the GI-1d-

a, there is a sharp decline in winter temperatures with much colder temperatures between September and

May, with strongly increased snow depth in north-western Europe during May, particularly in western

Britain, Brittany, and Central France and a lesser extent in northern Central Europe [41].

As a consequence of this marked cooling, the thawing of the permafrost, ice-covered lakes, and

other bodies of water was delayed. The period with stable warm summer temperatures was considerably

shorter, restricted to about three months (June to August) in eastern Europe and to only two months (July to

August) in north-western Europe (e.g., [40,146,147]). This led to a considerable decline in the length of the

growing season, again with a pronounced west-east trend. While the regions along the Atlantic coast of

France and Britain experienced a shortening of more than a month, Eastern Britain, Central France, Iberia,

and Italy saw a reduction of 10-20 days, while northern Central Europe and the Balkans saw a decrease of

between 2 and 10 days [41]. Shifts in the phenological patterns of the growing season can have considerable

influence on the direction of migrations of herbivores [148] and might, over the course of centuries also

influence the large-scale distribution of hunter-gatherer populations [22,149]. Interestingly, isotope analyses

on reindeer teeth from the Hamburgian and Ahrensburgian layer at Stellmoor (northern Germany) suggest

that reindeer herds generally migrated in an east-west direction on the North European Plain during the

Younger Dryas [150], although these patterns could have been more complex in detail [151], cf. [38].

Another highly relevant phenomenon is a general shift towards dryer conditions during the Younger Dryas,

which probably led to the lowering of the groundwater table and, thus, reduced water availability in some

areas [152,153]. This phenomenon is particularly well documented in the Scheldt basin in north-western

Belgium. Here, a lowering of the groundwater table probably led to a disappearance of numerous shallow

lakes and ponds in the interior of the Lower Scheldt basin, while the neighbouring Meuse basin seems to

have been less affected [140]. In addition, the lowlands of the North European Plain were affected by

intensified winds forming cover sand dunes [154,155]. Lower groundwater tables, sand cover in which

water seeps away easily and longer periods with ice-cover of the remaining water bodies might have posed

challenges to animals and humans alike. Yet, the cover increased the in situ preservation of sites but also

reduced their archaeological visibility [156].

In sum, the Younger Dryas has precipitated a swath of largely unfavourable environmental changes

for humans. These changes were more pronounced closer to the Atlantic seaboard and less so towards the

east. This is in line with our observation of an eastward shift of the Core Areas in north-western and Central

Europe. While populations declined in most areas, there is a clear increase in NE Germany and Poland as

well as stable estimates for Switzerland and SW Germany in the scenario based on dataset A. At the same

time, the estimates for SE Germany and the Czech Republic show the sharpest decline of all regions in both

scenarios, dataset A and B. In S Scandinavia, i.e. Denmark, the population dropped almost to zero, with

only a few camps appearing in the East. Considering the overall decline of regional populations, independent

population growth in NE Italy, NE Germany, and Poland seems unlikely. Therefore, these regional

population dynamics may be seen as first evidence of large-scale population movement as a response to

climatic cooling. If the population shift represents a prehistoric reality, then the source of the movement into

NE Germany and Poland would have mainly been SE Germany and the Czech Republic, with some people

moving towards Switzerland but the majority moving towards Poland. The sink areas of Poland, NE and

NW Germany also could have had an influx from Denmark. NE Italy might have received an influx from

SE France or western Italy, although there is a lack of archaeological evidence for contact in this direction

[157]. Possible source areas are also regions at the margins or outside the investigated area, such as central-

southern Italy, the western Balkans, or now submerged coastal areas.

Finally, environmental events beyond climate trends may also have played a role in shaping human

responses at this time. The spatial congruence between the void area across Central Europe and the most

massive ash fallout of the cataclysmic Laacher See eruption that occurred at 13,006±9 cal BP [138] in the

Eifel volcanic zone is notable, especially to the south-east and north-east of the eruptive centre (**Fig 7**). The

eruption occurred at the tail end of the cool GI-1b which likely stressed contemporaneous forager

populations. In Belgium and Central Germany, several sites evidence stratigraphic relations where

Federmesser-Gruppen occupation is capped by Laacher See fallout, terminating these occupations

[158,159]. The short- and medium-term environmental impacts of the eruption were varied and, depending

on local fallout thickness, likely considerable [160–164]. These may have depressed population in the

affected regions already prior to the onset of GS-1 and may have resulted in sustained legacies of landscape

avoidance that lasted into GS-1.

Demographic dynamics, minimal viable population, and social networks

Depending on whether one favours the mean estimate of the total population during GI-1d-a from

dataset A (8100, also including generically assigned sites) or the one from dataset B (5700, using only

chronologically subdivided sites) as more reliable, there is either an increase (dataset A) or decrease (dataset

B) after the Final Magdalenian, which produced a mean estimate of 7300 (min. 4500, max. 10200) people

[122]. However, some northern regions were frequently occupied for the first time after the Last Glacial

Maximum (**Fig 6**), suggesting a stable metapopulation if not an increase. Less ambiguous is the subsequent

decline in GS-1 to 4300 and 3500 people for the estimates based on datasets A and B, respectively. In both

scenarios, the population is reduced by roughly half. This marked decline raises questions about the viability

of the remaining population and the consequences for regional as well as long-distance social networks.

Defining what constitutes minimum viable populations in humans is difficult. A lower threshold of

~1500 individuals, comprising around 770 individuals in the reproductive age cohort, has been suggested

[165–167]. While the effects of drift-related inbreeding can be avoided at even lower numbers of 100 to 500

people [166,168], such minuscule populations become highly sensitive to negative stochastic effects,

especially in high-risk environments [169]. Despite being relatively low, our estimates are in the range of

viable populations, including, notably, the estimates for GS-1, which are still high compared to those of

most other periods of the Upper Palaeolithic [130,170]. This difference is particularly striking regarding the

Gravettian, where mean estimated population sizes dropped from 2800 people during the early phase to only

1000 during the late phase [50].

The estimated population decline likely also affected the size, structure, and connectivity of social

networks. On the one hand, a reduction in the number of people who can maintain social contacts is likely

to lead to distortions in the networks. Here, networks at regional scales are usually maintained by more

people than networks at larger scales, which are thus more prone to disturbances. On the other hand, actively

increasing the extent and connectivity of networks can be a way to mitigate environmental risk and

population decline [171] – as long as the increasing physical distances can be bridged (cf. [172–174]). Social

networks exist at different spatial scales, and besides palaeogenetic studies [23], they usually become

archaeologically visible through different classes of objects circulating within them that have characteristic

transport distances and patterns [175]. For networks at a regional scale, lithic raw materials are probably the

most important source of information [176]. The materials usually transported in economically significant

numbers and over distances well below 300 km are here considered likely to be acquired by embedded

procurement [177,178] and thus by members of the same regional group that also discarded them (but see

[179]). In the available raw material data, there is indeed an increase in the size of the catchments during

the Final Palaeolithic, very pronounced in all areas of the study region except for the NW European area

(see RMCA median values in **Tables 1 and 2; S1 and S2 Tables**).

For large-scale networks (connecting two or more regional networks) to become visible, transport

patterns of objects with a presumably different acquisition mode are better suited. Ideally, these objects are

not procured directly by the group that discards them but acquired via exchange between groups. Here, sub-

recent and fossil mollusc shells seem to be highly informative. In contrast to lithic raw materials, they are

regularly transported over more than 600 km and during the Late Upper Palaeolithic, they trace a network

that spans between the Atlantic and Mediterranean coast, the Paris Basin up the Rhine-Meuse Area and

across to the Swabian Jura [180–183]. They thus also span across the area of the central void in the Final

Palaeolithic dataset (**Fig 7**). For the Final Palaeolithic itself, however, information on transported mollusc

shells is extremely rare. There are singular instances of long-distance transports of lithic raw materials (e.g.

obsidian found in Poland), yet there is no available evidence that systematically reflects social networks at

that scale. The absence of evidence for regular long-distance transports of objects in contrast to the Late

Upper Palaeolithic may signal reduced contact at that scale. By this token, the extensive void in the centre

of the investigated area might also be an expression of reduced social connectivity between hunter-gatherers

in southwestern France and Iberia on the one hand and in Northern and Central Europe on the other. The

latter area can possibly be further divided by mid-scale networks through the distribution of raw materials

such as red Helgoland flint [52] or the use of amber [184,185] or by differentiated symbolic use [97,185].

If there had been decreasing contact between the south-western and north-eastern parts of the investigated

area during GI-1d-a, the marked population decrease in GS-1 probably would have additionally distorted

the networks.

Recent palaeogenetic evidence suggests a differential development in both regions during the Final

Palaeolithic. There appears to be a probably south-north directed movement of people during GI-1d-a,

passing west of the Alps from Italy into Central Europe and the British Isles, whereas the population in

Iberia shows more pronounced genetic continuity with several admixture events [23,66]. Whether this

suggested south-north dispersal is consilient with the proposed northward dispersal of backed points from

south-eastern France along the Rhône and Saône axis and alongside the disappearance of reindeer, as often

emphasised by Thévenin [45,186,187], is beyond the scope of this paper. However, these proposals have

mainly been based on typological analyses carried out on assemblages whose archaeostratigraphic reliability

is difficult to prove. Furthermore, given the available data, the Azilian traditions appear after the

Magdalenian, and some can be considered as a final episode of these traditions [3,90,188]. Other, more

recent traditions of the Azilian (GI-1) are part of their extension and of a common evolution of technical

traditions on a European scale. The situation is perhaps more complex at the very end of the Palaeolithic

(GS-1) with the existence of a certain duality of technical traditions in France. The presence of assemblages

with characteristics close to those known from the Laborian and Epigravettian periods has been proposed

[83,93,96,189]. However, the chronological framework of these phenomena remains to be defined and the

techno-economic data needs to be improved. At the moment, identifying this large empty area with very

little to no evidence being assignable to the Final Palaeolithic calls for future research.

Conclusion

During the GI-1d-a, the higher-latitude areas of Europe – parts of which were previously unoccupied

or only ephemerally visited by anatomically modern humans – became more regularly and frequently visited

by contemporaneous hunter-gatherer groups. At this time, the western and central European lowlands

became a hotspot of population dynamics for the first time. The climatic cooling during GS-1 (Younger

Dryas) led to a marked population decline but also different regional responses. Populations in the Iberian

Peninsula and southwestern France shrank while staying in the same area, a response comparable to the one

observed during the Gravettian. Populations in Northern and Central Europe, in contrast, reacted with a

pronounced shift towards the east. While populations declined in the western part, they increased in the

eastern part. We argue this was not because of population growth but because of population movement from

west to east. The same phenomenon can be observed for the area of south-eastern France up to north-eastern

Italy, where the former region also showed a population decline, while the latter saw a population increase,

again possibly due to population influx. The shift of populations towards the east was most probably a

reaction to deteriorating environmental conditions brought about by the Younger Dryas, including longer

winters, higher snow cover and a strongly reduced growing season. These changes affected the western parts

of Europe particularly hard, especially the Atlantic coast, and became more and more buffered towards the

east.

The question thus arises as to why the two sub-populations in the southwestern part and the rest of

the investigated area reacted so differently. Answers to this question are not readily evident. Maybe the two

sub-populations had already become largely decoupled during the Allerød, as might be indicated by the

central void area. Maybe hunter-gatherers in the topographically rather uniform lowlands had a different

attitude towards and aptitude for long-distance movements or a more permeable territorial structure. The

possibility that some behavioural or technological innovation – using poison for efficiently killing prey,

snowshoes, skis, travois or the like – transformed their ability to move longer distances at lower costs should

not be discounted. Future studies are needed to illuminate these aspects. To further refine our

palaeodemographic estimates, a higher chronological resolution of the dataset will likely reveal currently

invisible patterns, which in combination with high-resolution climate models and palaeoenvironmental

studies will allow for a more differentiated view of the biocultural processes that occurred at the end of the

Pleistocene. In-depth studies of material culture change and a better understanding of transport patterns of

raw materials and objects of far-distance transports could provide data on the size and connectivity of social

networks and therefore offer additional details to the coarse picture and potentially identify non-

environmental explanations for the observed patterns of population dynamics.

Acknowledgements

The authors would like to thank Andrea Darida, Martin Müller and Nina Avci for their help in

compiling the database and Dennis Batz for editing the references. All authors revised and added new data

on sites and assemblages for regions of their expertise. Felix Riede, Katja Winkler, Florian Sauer and Birgit

Gehlen provided regional databases at an early stage of the project, that were incorporated after adjustment

to the specific requirements. All geospatial calculations were conducted using MapInfo 8.5, maps were

created using QGis 3.34. The German Research Foundation funded the research of Isabell Schmidt –

Project-ID 57444011 – SFB 806 'Our Way to Europe'. Andreas Maier's contribution is made as part of the

project 'HESCOR', funded from the programme 'Profilbildung 2022', an initiative of the Ministry of

Culture and Science of the State of Northrhine Westphalia (HESCOR PB22-081). Research contributed to

this paper by Berit Valentin Eriksen, Mara-Julia Weber and Sonja B. Grimm was funded by the DFG,

German Research Foundation, project number 290391021-SFB 1266, 'Scales of Transformation'. Research

of Werner Schön in southwest Bavaria (Allgäu) was funded by the DFG, German Research Foundation,

project numbers 386654307 and 57444011. Felix Riede's contribution is made as part of CLIOARCH, an

ERC Consolidator Grant project, and has received funding from the European Research Council (ERC)

under the European Union's Horizon 2020 research and innovation programme (grant agreement

No. 817564). Annabell Zander's contribution is made as part of her British Academy Postdoctoral

Fellowship at the University of York. MN would like to thank Dion Stoop (Noordelijk Archeologisch Depot,

Nuis), Huub Scholte Lubberink (RAAP Oost, Zutphen) and Haije Veenstra (RAAP Noord, Drachten). The

authors are solely responsible for the content of this publication.

References

1.   Guthrie RD. Origin and causes of the mammoth steppe: a story of cloud cover, woolly mammal
tooth pits, buckles, and inside-out Beringia. Quat Sci Rev. 2001;20: 549–574. doi:10.1016/S0277-
3791(00)00099-8

2.   Drucker DG. The Isotopic Ecology of the Mammoth Steppe. Annu Rev Earth Planet Sci. 2022;50:
395–418. doi:10.1146/annurev-earth-100821-081832

3.   Grimm SB. Resilience and reorganisation of social systems during the Weichselian lateglacial in
North-West Europe: an evaluation of the archaeological, climatic, and environmental record. Mainz,
Regensburg: Verlag des Römisch-Germanischen Zentralmuseums, Schnell & Steiner; 2019.

4.   Jacquier J, Mevel L, Bodu P. Less simple than expected. New technological and functional insights
on the Late Azilian site of Le Closeau (France). Quartär. 2021;68: 181–214.
doi:10.7485/QU.2021.68.94302

5.   Rozoy J-G. Les Derniers Chasseurs: L'Épipaléolithique en France et en Belgique: Essai de Synthèse.
Charleville: Champenoise; 1978.

6.   Terberger T, Eriksen BV, editors. Hunters in a changing world: environment and archaeology of the
Pleistocene - Holocene Transition (ca. 11000-9000 B.C.) in Northern Central Europe; workshop of
the U.I.S.P.P.-Commission XXXII at Greifswald in September 2002. Rahden/Westfalen: Verlag
Marie Leidorf; 2004.

7.   Vollbrecht J. Spätpaläolithische Besiedlungsspuren aus Reichwalde. Dresden: Theiss; 2005.

8.   Bodu P, Debout G, Bignon O. Variabilité des habitudes tardiglaciaires dans le Bassin parisien:
l'organisation spatiale et sociale de l'Azilien ancien du Closeau. Bull Soc Préhist Fr. 2006;103: 711–
728. doi:10.3406/bspf.2006.13501

9. Bolus M. Die Siedlungsbefunde des späteiszeitlichen Fundplatzes Niederbieber (Stadt Neuwied): Ausgrabungen 1981–1988. Bonn: Habelt; 1992.

10. Bosinski G. Upper and Final Palaeolithic settlement patterns in the Rhineland, West Germany. In: Dibble HL, Montet-White A, editors. Upper Pleistocene Prehistory of Western Eurasia. Philadelphia: The University Museum, University of Pennsylvania; 1988. pp. 375–86.

11. Terberger T. Die Siedlungsbefunde des Magdalénien-Fundplatzes Gönnersdorf: Konzentrationen III und IV. Stuttgart: Steiner; 1997.

12. Fasser N, Visentin D, Duches R, Peresani M, Fontana F. Lithic projectile technology in the western Late Epigravettian: The case study of north-eastern Italy. Quat Int. 2024;694: 70–90. doi:10.1016/j.quaint.2024.04.006

13. Fat Cheung C. L'Azilien pyrénéen parmi les sociétés du Tardiglaciaire ouest-européen: apport de l'étude des industries lithiques. Archéologie et Préhistoire. Ph.D., Université Toulouse le Mirail - Toulouse II. 2015.

14. Fischer A. Pioneers in deglaciated landscapes: the expansion and adaptation of late Palaeolithic societies in Southern Scandinavia. In: Barton N, Roberts AJ, Roe DA, editors. The late Glacial in north-west Europe: human adaptation and environmental change at the end of the Pleistocene. London: Council for British Archaeology; 1991. pp. 100–121.

15. Riede F, Matzig DN, Biard M, Crombé P, De Pablo JF-L, Fontana F, et al. A quantitative analysis of Final Palaeolithic/earliest Mesolithic cultural taxonomy and evolution in Europe. Peresani M, editor. PLoS ONE. 2024;19: e0299512. doi:10.1371/journal.pone.0299512

16. Valentin B. Les groupes humains et leurs traditions au Tardiglaciaire dans le Bassin parisien. Apports de la technologie lithique comparée. Ph.D., Université Panthéon-Sorbonne. 1995.

17. Valentin B. L'usage de la pierre tendre pour la débitage des lames. Circonstances de quelques innovations au cours du Tardiglaciaire dans le Bassin parisien. In: Pion G, editor. Le paléolithique supérieur récent: nouvelles données sur le peuplement et l'environnement Actes de la Table ronde de Chambéry. Paris: Société Préhistorique Française; 2000. pp. 253–260.

18. Valentin B, Fosse G, Billard C. Aspects et rythmes de l'azilianisation dans le Bassin parisien: caractérisation de l'industrie lithique recueillie au Cornet (locus 33) à Ambenay, Eure. Gallia Préhist. 2004;46: 171–209. doi:10.3406/galip.2004.2042

19. French JC, Collins C. Upper Palaeolithic population histories of Southwestern France: a comparison of the demographic signatures of 14C date distributions and archaeological site counts. J Archaeol Sci. 2015;55: 122–134. doi:10.1016/j.jas.2015.01.001

20. Tallavaara M, Luoto M, Korhonen N, Järvinen H, Seppä H. Human population dynamics in Europe over the Last Glacial Maximum. Proc Natl Acad Sci. 2015;112: 8232–8237. doi:10.1073/pnas.1503784112

21. Schmidt I, Gehlen B, Zimmermann A. Population estimates for the Final Palaeolithic (14,000 to 11,600 years cal. BP) of Europe - challenging evidence and methodological limitations. In: Mevel L, Weber M-J, Maier A, editors. En Mouvement / On the Move / In Bewegung. Paris: Société Préhistorique Française; 2021. pp. 221–237.

22. Maier A, Ludwig P, Zimmermann A, Schmidt I. The Sunny Side of the Ice Age: Solar Insolation as a Potential Long-Term Pacemaker for Demographic Developments in Europe Between 43 and 15 ka Ago. PaleoAnthropology. 2022;1: 35–51. doi:10.48738/2022.ISS1.100

23. Posth C, Yu H, Ghalichi A, Rougier H, Crevecoeur I, Huang Y, et al. Palaeogenomics of Upper Palaeolithic to Neolithic European hunter-gatherers. Nature. 2023;615: 117–126. doi:10.1038/s41586-023-05726-0

24. Adolphi F, Muscheler R, Friedrich M, Güttler D, Wacker L, Talamo S, et al. Radiocarbon calibration uncertainties during the last deglaciation: Insights from new floating tree-ring chronologies. Quat Sci Rev. 2017;170: 98–108. doi:10.1016/j.quascirev.2017.06.026

25. Reimer PJ, Austin WEN, Bard E, Bayliss A, Blackwell PG, Bronk Ramsey C, et al. The IntCal20 Northern Hemisphere Radiocarbon Age Calibration Curve (0–55 cal kBP). Radiocarbon. 2020;62: 725–757. doi:10.1017/RDC.2020.41

26. Deschamps P, Durand N, Bard E, Hamelin B, Camoin G, Thomas AL, et al. Ice-sheet collapse and sea-level rise at the Bølling warming 14,600 years ago. Nature. 2012;483: 559–564. doi:10.1038/nature10902

27. Rasmussen TL, Thomsen E, Moros M. North Atlantic warming during Dansgaard-Oeschger events synchronous with Antarctic warming and out-of-phase with Greenland climate. Sci Rep. 2016;6: 20535. doi:10.1038/srep20535

28. Naughton F, Costas S, Gomes SD, Desprat S, Rodrigues T, Goñi MS, et al. Coupled ocean and atmospheric changes during Greenland stadial 1 in southwestern Europe. Quat Sci Rev. 2019;212: 108–120. doi:10.1016/j.quascirev.2019.03.033

29. Landais A, Capron E, Masson-Delmotte V, Toucanne S, Rhodes R, Popp T, et al. Ice core evidence for decoupling between midlatitude atmospheric water cycle and Greenland temperature during the last deglaciation. Clim Past. 2018;14: 1405–1415. doi:10.5194/cp-14-1405-2018

30. Vescovi E, Ravazzi C, Arpenti E, Finsinger W, Pini R, Valsecchi V, et al. Interactions between climate and vegetation during the Lateglacial period as recorded by lake and mire sediment archives in Northern Italy and Southern Switzerland. Quat Sci Rev. 2007;26: 1650–1669. doi:10.1016/j.quascirev.2007.03.005

31. Samartin S, Heiri O, Lotter AF, Tinner W. Climate warming and vegetation response after Heinrich event 1 (16 700–16 000 cal yr BP) in Europe south of the Alps. Climate of the Past. 2012;8: 1913–1927. doi:10.5194/cp-8-1913-2012

32. Lemanik A, Baca M, Wertz K, Socha P, Popović D, Tomek T, et al. The impact of major warming at 14.7 ka on environmental changes and activity of Final Palaeolithic hunters at a local scale (Orawa-Nowy Targ Basin, Western Carpathians, Poland). Archaeol Anthropol Sci. 2020;12: 66. doi:10.1007/s12520-020-01020-6

33. Reade H, Grimm SB, Tripp JA, Neruda P, Nerudová Z, Roblíčková M, et al. Magdalenian and Epimagdalenian chronology and palaeoenvironments at Kůlna Cave, Moravia, Czech Republic. Archaeol Anthropol Sci. 2021;13: 4. doi:10.1007/s12520-020-01254-4

34. Hoek WZ. Vegetation response to the ∼14.7 and ∼11.5 ka cal. BP climate transitions: is vegetation lagging climate? Glob Planet Change. 2001;30: 103–115. doi:10.1016/S0921-8181(01)00081-9

35. Ammann B, Van Leeuwen JFN, Van Der Knaap WO, Lischke H, Heiri O, Tinner W. Vegetation responses to rapid warming and to minor climatic fluctuations during the Late-Glacial Interstadial (GI-1) at Gerzensee (Switzerland). Palaeogeogr Palaeoclimatol Palaeoecol. 2013;391: 40–59. doi:10.1016/j.palaeo.2012.07.010

36. Krüger S, Mortensen MF, Dörfler W. Sequence completed – palynological investigations on Lateglacial/Early Holocene environmental changes recorded in sequentially laminated lacustrine sediments of the Nahe palaeolake in Schleswig-Holstein, Germany. Rev Palaeobot Palyn. 2020;280: 104271. doi:10.1016/j.revpalbo.2020.104271

37. Rea BR, Pellitero R, Spagnolo M, Hughes P, Ivy-Ochs S, Renssen H, et al. Atmospheric circulation over Europe during the Younger Dryas. Sci Adv. 2020;6: eaba4844. doi:10.1126/sciadv.aba4844

38. Krüger S. Of birches, smoke and reindeer dung – Tracing human-environmental interactions palynologically in sediments from the Nahe palaeolake. J Archaeol Sci Rep. 2020;32: 102370. doi:10.1016/j.jasrep.2020.102370

39. Lane CS, Brauer A, Blockley SPE, Dulski P. Volcanic ash reveals time-transgressive abrupt climate change during the Younger Dryas. Geology. 2013;41: 1251–1254. doi:10.1130/G34867.1

40. Schenk F, Väliranta M, Muschitiello F, Tarasov L, Heikkilä M, Björck S, et al. Warm summers during the Younger Dryas cold reversal. Nat Commun. 2018;9: 1634. doi:10.1038/s41467-018-04071-5

41. Schenk F, Wohlfarth B. The imprint of hemispheric-scale climate transitions on the European climate during the last deglaciation (15.5 ka to 9 ka BP). 2019. Report No.: AB 2019-01. Available: https://www.skb.com/publication/2492488/

42. Grimm SB, Weber M-J. The chronological framework of the Hamburgian in the light of old and new 14C dates. Quartär. 2008;55: 17–40. doi:10.7485/QU55_02

43. Jacobi RM, Higham TFG. The early Lateglacial re-colonization of Britain: new radiocarbon evidence from Gough's Cave, southwest England. Quat Sci Rev. 2009;28: 1895–1913. doi:10.1016/j.quascirev.2009.03.006

44. Pedersen JB, Assmann JJ, Normand S, Karger DN, Riede F. Climate Niche Modeling Reveals the Fate of Pioneering Late Pleistocene Populations in Northern Europe. Current Anthropology. 2023;64: 599–608. doi:10.1086/726700

45. Thévenin A. L'"Azilien" et les cultures à pointes à dos courbe: esquisse géographique et chronologique. Bulletin de la Société préhistorique française. 1997;94: 393–411.

46. Holzkämper J, Maier A, Richter J. „Dark Ages" illuminated – Rietberg and related assemblages possibly reducing the hiatus between the Upper and Late Palaeolithic in Westphalia. Quartär. 2013; 115–136. doi:10.7485/QU60_6

47. Schmidt I, Hilpert J, Kretschmer I, Peters R, Broich M, Schiesberg S, et al. Approaching prehistoric
demography: proxies, scales and scope of the Cologne Protocol in European contexts. Phil Trans R
Soc B. 2021;376: 20190714. doi:10.1098/rstb.2019.0714

48. Schmidt I, Gehlen B, Winkler K, Arrizabalaga A, Arts N, Bicho N, et al. Database for the Final
Palaeolithic archaeological record in western, southern and central Europe. [object Object]; 2024.
doi:10.5281/ZENODO.13132357

49. Iversen J. The late-glacial flora of Denmark and its relation to climate and soil. Danmarks
Geologiske Undersøgelse II Række. 1954;80: 87–119.

50. Maier A, Zimmermann A. Populations headed south? The Gravettian from a palaeodemographic
point of view. Antiquity. 2017;91: 573–588. doi:10.15184/aqy.2017.37

51. Beuker J, Niekus JLT. Het gebruik van rode Helgoland-vuursteen in het laat-paleolithicum. Nieuwe
Drentse Volksalmanak. 2003;120: 112–120.

52. Terberger T, Schneider J, Fries JE. The site of Edewecht 82-East, district of Ammerland, and the
role of Red Heligoland flint in the Late Glacial. Praehist Zeitschr. 2020;95: 1–16. doi:10.1515/pz-
2020-0006

53. Baales M, Heuschen W, Kehl M, Manz A, Nolde N, Riemenschneider D, et al. Western visitors at
the Blätterhöhle (city of Hagen, southern Westphalia) during the Younger Dryas? A new final
palaeolithic assemblage type in western Germany. PLoS ONE. 2023;18: e0284479.
doi:10.1371/journal.pone.0284479

54. De Bie M, Caspar J-P. Rekem: A Federmesser camp on the Meuse River bank: Archeologie in
Vlaanderen 3. Leuven: Leuven University Press; 2000.

55. Grimm SB, Weber M-J, Mevel L, Sobkowiak-Tabaka I, editors. From the Atlantic to Beyond the
Bug River: Finding and Defining the Federmesser-Gruppen / Azilian: Proceedings of Session A5b
(Commission »The Final Palaeolithic of Northern Eurasia«) of the XVIIth UISPP Congress: 2014
Sep 1-7; Burgos: Spain. Heidelberg: Propylaeum; 2020. doi:10.11588/PROPYLAEUM.575

56. Grimm SB, Eriksen BV, Krüger S, Reuter T, Wild M, Weber M-J. Late Glacial occupation of
northern Germany and adjacent areas: Revisiting the archives. In: Gaudzinski-Windheuser S, Jöris
O, editors. The Beef behind all possible pasts The tandem-Festschrift in honour of Elaine Turner and
Martin Street. Heidelberg: Propylaeum; 2021. pp. 433–457.

57. Sobkowiak-Tabaka I, Diachenko A. Approaching Late Paleolithic Seasonal Mobility on the North
European Plain. In: Sobkowiak-Tabaka I, Diachenko A, Wiśniewski A, editors. Quantifying Stone
Age Mobility: Scales and Parameters. Cham: Springer International Publishing; 2022. pp. 97–128.
doi:10.1007/978-3-030-94368-4

58. Sobkowiak-Tabaka I. Społeczności późnego paleolitu w dorzeczu Odry. Poznań: Instytut
Archeologii i Etnologii Polskiej Akademii Nauk; 2011.

59. Sobkowiak-Tabaka I. Living on the edge, or how resilient people settled the North. In: Djindjian F,
editor. Les sociétés humaines face aux changements climatiques. Oxford: Archaeopress
Archaeology; 2022. pp. 297–319.

60.   Weber M-J, Clausen I, Housley RA, Miller CE, Riede F, Usinger H. New information on the
Havelte Group site Ahrenshöft LA 58 D (Nordfriesland, Germany) - Preliminary results of the 2008
fieldwork. Quartär. 2010;57: 7–24. doi:10.7485/QU57_01

61.   Holzkämper J, Maier A. Neue Ergebnisse zur jungpaläolithischen Besiedlung Nordrhein-Westfalens.
25 Jahre Archäologie im Rheinland 1987 - 2011. Stuttgart: Theiss; 2012. pp. 58–60.

62.   Barton RNE. Regional and chronological patterns in lithic raw material behaviour during the late
glacial and some implications for the British Later Upper Palaeolithic. Lithics. 2010;31: 33–43.

63.   Jacobi R, Higham T. The Later Upper Palaeolithic Recolonisation of Britain: New Results from
AMS Radiocarbon Dating. The Ancient Human Occupation of Britain. Amsterdam: Elsevier; 2011.
pp. 223–247. doi:10.1016/B978-0-444-53597-9.00012-1

64.   Pedersen JB, Maier A, Riede F. A punctuated model for the colonisation of the Late Glacial margins
of northern Europe by Hamburgian hunter-gatherers. Quartär. 2018;65: 85–104.
doi:10.7485/QU65_4

65.   Brovkin V, Brook E, Williams JW, Bathiany S, Lenton TM, Barton M, et al. Past abrupt changes,
tipping points and cascading impacts in the Earth system. Nat Geosci. 2021;14: 550–558.
doi:10.1038/s41561-021-00790-5

66.   Charlton S, Brace S, Hajdinjak M, Kearney R, Booth T, Reade H, et al. Dual ancestries and
ecologies of the Late Glacial Palaeolithic in Britain. Nat Ecol Evol. 2022;6: 1658–1668.
doi:10.1038/s41559-022-01883-z

67.   Winkler K. Ahrensburgien und Swiderien im mittleren Oderraum: Technologische und typologische
Untersuchungen an Silexartefakten der Jüngeren Dryaszeit. Kiel: Wachholtz Verlag; 2019.

68.   Stefański D, Wilczyński J. Extralocal raw materials in the Swiderian Culture: Case study of
Kraków-Bieżanów sites. Anthropologie. 2012;50: 427–442.

69.   Weber M-J, Grimm SB, Baales M. Between warm and cold: Impact of the Younger Dryas on human
behavior in Central Europe. Quat Int. 2011;242: 277–301. doi:10.1016/j.quaint.2010.12.002

70.   Grimm SB, Groß D, Gerken K, Weber M-J. On the onset of the Early Mesolithic on the North
German Plain. In: Zander A, Gehlen B, Gramsch B, editors. From the Early Preboreal to the
Subboreal period – Current Mesolithic research in Europe: studies in honour of Bernhard Gramsch.
Kerpen-Loogh: Welt und Erde Verlag; 2020. pp. 15–37.

71.   Eriksen BV. Late Palaeolithic settlement in Denmark – how do we read the record? In: Kobusiewicz
M, Kozlowski JK, editors. Post-pleniglacial re-colonisation of the great European Lowland: Papers
presented at the Conference organised by the International Union of Prehistoric and Protohistoric
Sciences, Commission 8, held at the Jagellonian University, Kraków, in June 1998. Kraków: Polska
Akademia Umiejętności; 1999. pp. 157–73.

72.   Riede F. The Laacher See-eruption (12,920 BP) and material culture change at the end of the
Allerød in Northern Europe. J Archaeol Sci. 2008;35: 591–599. doi:10.1016/j.jas.2007.05.007

73.   Crombé P, Deeben J, Van Strydonck M. Hunting in a changing environment: The transition from the
Younger Dryas to the (Pre)boreal in Belgium and the Southern Netherlands. In: Jaubert J, Fourment

N, Depaepe P, editors. Transitions, ruptures et continuité en préhistoire: actes du XXVII Congrès préhistorique de France: Bordeaux-Les Eyzies: 31 mai-5 juin 2010 Volume 2: Paléolithique et Mésolithique. Paris: Société Préhistorique Française; 2014. pp. 583–604.

74. Crombé P, Sergant J, Verbrugge A, De Graeve A, Cherretté B, Mikkelsen J, et al. A sealed flint knapping site from the Younger Dryas in the Scheldt valley (Belgium): Bridging the gap in human occupation at the Pleistocene–Holocene transition in W Europe. J Archaeol Sci. 2014;50: 420–439. doi:10.1016/j.jas.2014.07.021

75. Vermeersch PM. La transition Ahrensbourgien-Mésolithique ancien en Campine belge et dans le sud sableux des Pays-Bas. In: Fagnart J-P, Thévenin A, Ducrocq T, Souffi B, Coudrec P, editors. Le début du Mésolithique en Europe du Nord-Ouest, Mémorie XLV de la Societé préhistorique française. Paris: Societé Préhistorique Française; 2008. pp. 11–29.

76. Niekus MJ, Boekschoten GR, Deeben JHC. A Late Preboreal site from Zwolle, province of Overijssel, and some remarks on the Ahrensburgian in the Netherlands. In: Eriksen BV, Rensink E, Harris S, editors. The Final Palaeolithic of Northern Eurasia: proceedings of the Amersfoort, Schleswig and Burgos UISPP commission meetings. Kiel: Ludwig; 2019. pp. 51–79.

77. Vandendriessche H. Flintknapping from the Late Glacial to the Early Holocene: the Belgian Scheldt Valley sites of Ruien and Kerkhove. Leiden: Sidestone Press; 2022.

78. Crombé P, Pironneau C, Robert P, Van Der Sloot P, Boudin M, De Groote I, et al. Human response to the Younger Dryas along the southern North Sea basin, Northwest Europe. Sci Rep. 2024;14: 18074. doi:10.1038/s41598-024-68686-z

79. Barton RNE. Long Blade Technology in Southern Britain. In: Bonsall C, editor. The Mesolithic in Europe: papers presented at the third international symposium, Edinburgh, 1985. Edinburgh: John Donald; 1989. pp. 264–271.

80. Barton RNE, Roberts AJ. The transition from the Younger Dryas to the Pre-Boreal in southern Britain: some new perspectives on the spatial patterning and chronology of long blade sites. In: Montoya C, Fagnart J-P, Locht J-L, editors. Préhistoire de l'Europe du nord-ouest: mobilités, climats et identités culturelles Volume 2: Paléolithique supérieur ancien, Paléolithique final – Mésolithique, Congrès préhistorique de France: Compte Rendu de la 28ème session. Paris: Société Préhistorique Française; 2019. pp. 381–389.

81. Fagnart J-P. La fin des temps glaciaires dans le nord de la France: approches archéologiques et environnementales des occupations humaines du Tardiglaciaire. Paris: Societé Préhistorique Française; 1997.

82. Valentin B. Jalons pour une Paléohistoire des derniers chasseurs (XIVe-VIe millénaire avant J.-C.). Paris: Publications de la Sorbonne; 2008.

83. Naudinot N, Fagnart J-P, Langlais M, Mevel L, Valentin B. Les dernières sociétés du Tardiglaciaire et des tout débuts de l'Holocène en France: Bilan d'une trentaine d'années de recherche. Gallia Préhist. 2019; 5–45. doi:10.4000/galliap.1394

84. Weber M-J, Valentin B, Bodu P. It's all about volume - New observations on specific aspects of core reduction at the Belloisian site Donnemarie-Dontilly (Seine-et-Marne, France). In: Eriksen BV,

Rensink E, Harris S, editors. The Final Palaeolithic of Northern Eurasia: proceedings of the
Amersfoort, Schleswig and Burgos UISPP commission meetings. Kiel: Ludwig; 2019. pp. 91–104.

85. Street M, Baales M, Gehlen B, Heinen M, Heuschen W, Orschiedt J, et al. Archaeology across the
Pleistocene-Holocene boundary in western Germany: Human responses to rapid environmental
change. In: Montoya C, Fagnart J-P, Locht J-L, editors. Préhistoire de l'Europe du nord-ouest:
mobilités, climats et identités culturelles Volume 2: Paléolithique supérieur ancien, Paléolithique
final – Mésolithique, Congrès préhistorique de France: Compte Rendu de la 28ème session. Paris:
Société Préhistorique Française; 2019. pp. 491–510.

86. Mevel L, Weber M-J, Berg-Hansen IM. Towards a reconstruction of the techno-economic variability
of the Ahrensburgian in Schleswig-Holstein. In: Montoya C, Fagnart J-P, Locht J-L, editors.
Préhistoire de l'Europe du nord-ouest: mobilités, climats et identités culturelles Volume 2:
Paléolithique supérieur ancien, Paléolithique final – Mésolithique, Congrès préhistorique de France:
Compte Rendu de la 28ème session. Paris: Société Préhistorique Française; 2019. pp. 433–453.

87. Sørensen M, Lübke H, Groß D. The Early Mesolithic in Southern Scandinavia and Northern
Germany. In: Milner N, Conneller C, Taylor B, editors. Star Carr: a persistent place in a changing
world. Heslington, York: White Rose University Press; 2018. pp. 305–329.

88. Bodu P, Valentin B. Groupes à Federmesser ou Aziliens dans le sud et l'ouest du Bassin parisien.
Propositions pour un nouveau modèle d'évolution. Bull Soc Préhist Fr. 1997;94: 341–348.
doi:10.3406/bspf.1997.10698

89. Naudinot N, Jacquier J. Socio-economic organization of Final Paleolithic societies: New
perspectives from an aggregation site in Western France. J Anthropol Archaeol. 2014;35: 177–189.
doi:10.1016/j.jaa.2014.05.004

90. Langlais M, Chevallier A, Fat Cheung C, Jacquier J, Marquebielle B, Naudinot N. The Pleistocene-
Holocene transition in Southwestern France: A focus on the Laborian. Quat Int. 2020;564: 37–47.
doi:10.1016/j.quaint.2019.09.045

91. Gibaud A, Delvigne V, Langlais M, Bracco J-P. Frontière de préhistorique·s ou frontière de
préhistorien·ne·s : quid des litho-espaces durant la transition Pléistocène-Holocène en France
centrale? Gallia Préhist. 2023; 75–102. doi:10.4000/galliap.3790

92. Naudinot N. Entre steppes et forêts: Chronique atlantique des derniers chasseurs-cueilleurs
paléolithiques. Éditions Totem; in press.

93. Fornage-Bontemps S. Le niveau A4 de Rochedane, l'est de la France et la question des influences
épigravettiennes à la fin du tardiglaciaire. Besançon: Presses universitaires de Franche-Comté; 2017.

94. Fornage-Bontemps S. Quand les idées franchissent les montagnes. L'Est de la France et la question
de la diffusion des influences épigravettiennes au nord des Alpes entre l'Allerød et la fin du Dryas
recent. In: Naudinot N, Meignen L, Binder D, Querre G, editors. Les systèmes de mobilité de la
préhistoire au Moyen Age : XXXVe rencontres internationales d'archéologie et d'histoire
d'Antibes : actes des rencontres, 14-16 octobre 2014. Nice: APDCA; 2016. pp. 337–352.

95. Mevel L, Fornage-Bontemps C, Bereiziat G. Au carrefour des influences culturelles ? Les industries
lithiques de la fin du Tardiglaciaire entre Alpes du nord et Jura, 11 500-9 500 CalBC. In: Langlais
M, Naudinot N, Peresani M, editors. Les groupes culturels de la transition Pléistocène - Holocène

entre Atlantique et Adriatique: actes de la séance de la Société préhistorique française Bordeaux 24-
25 mai 2012. Paris: Société Préhistorique Française; 2014. pp. 45–81.

96. Mevel L, Pion G, Fornage-Bontemps S. Changements techniques et géographie culturelle à
l'extrême fin du Paléolithique dans les Alpes du nord françaises. Les stratigraphies de l'abri de La
Fru (Savoie) revisitées. In: Jaubert J, Fourment N, Depaepe P, editors. Transitions, ruptures et
continuité en préhistoire: actes du XXVII Congrès préhistorique de France: Bordeaux-Les Eyzies:
31 mai-5 juin 2010 Volume 1. Paris: Société Préhistorique Française; 2014. pp. 527–546.

97. Naudinot N, Tomasso A, Messager E, Finsinger W, Ruffaldi P, Langlais M. Between Atlantic and
Mediterranean: Changes in technology during the Late Glacial in Western Europe and the climate
hypothesis. Quat Int. 2017;428: 33–49. doi:10.1016/j.quaint.2016.01.056

98. Tomasso A, Fat Cheung C, Fornage-Bontemps S, Langlais M, Naudinot N. Winter is coming: What
happened in western European mountains between 12.9 and 12.6 ka cal. BP (beginning of the GS1).
Quat Int. 2018;465: 210–221. doi:10.1016/j.quaint.2017.12.020

99. Langlais M, Fat Cheung C, Laroulandie V, Lefebvre A, Marquebielle B, Pétillon J-M. Dix mille ans
de peuplement humain sur le versant nord des Pyrénées (21000-11000 cal BP) entre unification et
régionalisation culturelles. Munibe, Antropologia-Arkeologia. 2020;71: 59–71.
doi:10.21630/maa.2020.71.18

100. Aura JE, Jordá JF, Montes L, Utrilla P. Human responses to Younger Dryas in the Ebro valley and
Mediterranean watershed (Eastern Spain). Quat Int. 2011;242: 348–359.
doi:10.1016/j.quaint.2011.01.023

101. Bicho N, Haws J. The Magdalenian in central and southern Portugal: Human ecology at the end of
the Pleistocene. Quat Int. 2012;272–273: 6–16. doi:10.1016/j.quaint.2012.02.055

102. Bicho N, Haws J, Almeida F. Hunter–gatherer adaptations and the Younger Dryas in central and
southern Portugal. Quat Int. 2011;242: 336–347. doi:10.1016/j.quaint.2010.12.025

103. Gameiro C, Aubry T, Almeida F, Dimuccio L, Gabriel S, Gaspar R, et al. Archaeology of the
Pleistocene-Holocene transition in Portugal: Synthesis and prospects. Quat Int. 2020;564: 113–137.
doi:10.1016/j.quaint.2020.03.018

104. Bicho N, Cascalheira J, Marreiros J, Gonçalves C, Pereira T, Dias R. Chronology of the Mesolithic
occupation of the Muge valley, central Portugal: The case of Cabeço da Amoreira. Quat Int.
2013;308–309: 130–139. doi:10.1016/j.quaint.2012.10.049

105. Naudinot N, Tomasso A, Tozzi C, Peresani M. Changes in mobility patterns as a factor of 14 C date
density variation in the Late Epigravettian of Northern Italy and Southeastern France. J Archaeol
Sci. 2014;52: 578–590. doi:10.1016/j.jas.2014.05.021

106. Ruiz-Redondo A, Vukosavljević N, Tomasso A, Peresani M, Davies W, Vander Linden M. Mid and
Late Upper Palaeolithic in the Adriatic Basin: Chronology, transitions and human adaptations to a
changing landscape. Quat Sci Rev. 2022;276: 107319. doi:10.1016/j.quascirev.2021.107319

107. Duches R, Peresani M, Pasetti P. Success of a flexible behavior: Considerations on the manufacture
of Late Epigravettian lithic projectile implements according to experimental tests. Archaeol
Anthropol Sci. 2018;10: 1617–1643. doi:10.1007/s12520-017-0473-x

108. Peresani M, Tomio C, Dalmeri G. Les grattoirs épigravettiens et leur « raccourcissement » durant le Tardiglaciaire en Italie. Reflets d'un changement dans l'économie du débitage. In: Langlais M, Naudinot N, Peresani M, editors. Les groupes culturels de la transition Pléistocène - Holocène entre Atlantique et Adriatique: actes de la séance de la Société préhistorique française Bordeaux 24-25 mai 2012. Paris: Société Préhistorique Française; 2014. pp. 205–220.

109. Sauer F. Late Palaeolithic Land Use Patterns in Bavaria. Ph.D., Friedrich-Alexander-Universität. 2018.

110. Vencl S. Prehistory of Bohemia 1: The Palaeolithic and Mesolithic. Praha: Archeologický ústav AV CR; 2013.

111. Mlejnek O, Záhorák V. Předběžná zpráva o výzkumu mezolitické a pozdně paleolitické lokality u Uherska ve východních Čechách. Excavation of the Mesolithic and Late Palaeolithic site near Uhersko in Eastern Bohemia: preliminary report. Ročenka. 2018; 50-66.

112. Eigner J, Prekop F. Tašovice 1 a 2: dvě pozdně paleolitické a mezolitické lokality na Karlovarsku = Tašovice 1 and 2: two late palaeolithic and mesolithic sites in the Karlovy Vary region. Praha: Národní muzeum; 2021.

113. Kapustka K, Eigner J, Parkman M, Řezáč M, Přichystal A, Pokorný P, et al. Pozdně paleolitické a mezolitické osídlení Šumavy:možnosti výzkumu, datování a interpretaceLate Palaeolithic and Mesolithic Settlement of Šumava:The Possibilities of Research, Dating and Interpretation. Pam Arch. 2020; 5–59. doi:10.35686/PA2020.1

114. Svoboda J, Pokorný P, Horáček I, Sázelová S, Abraham V, Divišová M, et al. Late Glacial and Holocene sequences in rockshelters and adjacent wetlands of Northern Bohemia, Czech Republic: Correlation of environmental and archaeological records. Quat Int. 2018;465: 234–250. doi:10.1016/j.quaint.2017.05.009

115. Šída P, Pokorný P. Mesolithic of the Northern Bohemia III: evolution of prehistoric landscape in Bohemian Paradise: vegetation, fauna, humans. Brno: Archeologický ústav AV ČR; 2020.

116. Jochim MA, Kind C-J, Kleinmann A, Merkt J, Stephan E. Eine spätpaläolithische Fundstelle am Ufer des Federsees: Bad Buchau-Kappel, Flurstück Gemeindebeunden. Fundber Baden-Württemberg. 2015;35: 37–134. doi:10.11588/FBBW.2015.0.44521

117. Hornauer-Jahnke TK, Noack ES. Der Übergang vom Jung- zum Spätpaläolithikum in Südwestdeutschland: Der Zigeunerfels. In: Baales M, Pasda C, editors. "All der holden Hügel ist keiner mir fremd .": Festschrift zum 65 Geburtstag von Claus-Joachim Kind. Bonn: Habelt; 2019. pp. 315–336.

118. Schön W, Affolter J, Gehlen B, Guggenmos A, Schröppel J, Guggenmos S. Siedlungsrelikte eiszeitlicher Menschen im Allgäu. In: Batzel K, editor. Natur und Mensch während der Eiszeit im Allgäu. Aichach-Friedberg: Likias-Verlag; 2024.

119. Schön W, Affolter J, Gehlen B, Gulisano G, Guggenmos A, Guggenmos S, et al. Mesolithische Besiedlung und Rohstoffversorgung im Allgäu. In: Uthmeier T, Maier A, editors. Stone Age: studying technologies of non-analogous environments and glacial ecosystems: papers in honor of Jürgen Richter. Bonn: Habelt; 2024. pp. 399–422.

120. Nielsen EH. Paläolithikum und Mesolithikum in der Zentralschweiz: Mensch und Umwelt zwischen 17 000 und 5 500 v. Chr. Luzern: Kantonaler Lehrmittelverlag; 2009.

121. Sedlmeier J. Die letzten Wildbeuter der Eiszeit: neue Forschungen zum Spätpaläolithikum im Kanton Basel-Landschaft. Basel: Schwabe Verlag; 2015.

122. Kretschmer I. Demographische Untersuchungen zu Bevölkerungsdichten, Mobilität und Landnutzungsmustern im späten Jungpaläolithikum. Rahden/Westf: Verlag Marie Leidorf; 2015.

123. Peresani M, Monegato G, Ravazzi C, Bertola S, Margaritora D, Breda M, et al. Hunter-gatherers across the great Adriatic-Po region during the Last Glacial Maximum: Environmental and cultural dynamics. Quat Int. 2021;581–582: 128–163. doi:10.1016/j.quaint.2020.10.007

124. Niekus MJ, Cohen KM, Johansen L, Stapert D. Opgezogen, opgespoten en opgeraapt: vuurstenen en hardstenen artefacten van de Zandmotor en hun sedimentaire context. Amersfoort: Rijksdienst voor het Cultureel Erfgoed; 2023.

125. Boemke BJ, Maier A, Schmidt I, Römer W, Lehmkuhl F. Testing the representativity of Palaeolithic site distribution: The role of sampling bias in the European Upper and Final Palaeolithic record. Quat Sci Rev. 2023;316: 108220. doi:10.1016/j.quascirev.2023.108220

126. Baales M. Umwelt und Jagdökonomie der Ahrensburger Rentierjäger im Mittelgebirge. Bonn: Dr. Rudolf Habelt GmbH; 1996.

127. Sauer F, Riede F. A Critical Reassessment of Cultural Taxonomies in the Central European Late Palaeolithic. J Archaeol Method Theory. 2019;26: 155–184. doi:10.1007/s10816-018-9368-0

128. Vaquero M, Romagnoli F. Searching for Lazy People: the Significance of Expedient Behavior in the Interpretation of Paleolithic Assemblages. J Archaeol Method Theory. 2018;25: 334–367. doi:10.1007/s10816-017-9339-x

129. Riede F. The 'Bromme Problem' – Notes on Understanding the Federmessergruppen and Bromme Culture Occupation in Southern Scandinavia during the Allerød and Early Younger Dryas Chronozones. In: Sørensen M, Buck Pedersen K, editors. Problems in Palaeolithic and Mesolithic research. Copenhagen: University of Copenhagen & Museum of Southeast Denmark; 2017. pp. 61–85.

130. Maier A, Lehmkuhl F, Ludwig P, Melles M, Schmidt I, Shao Y, et al. Demographic estimates of hunter–gatherers during the Last Glacial Maximum in Europe against the background of palaeoenvironmental data. Quat Int. 2016;425: 49–61. doi:10.1016/j.quaint.2016.04.009

131. Vermeersch PM. Radiocarbon Palaeolithic Europe database: A regularly updated dataset of the radiometric data regarding the Palaeolithic of Europe, Siberia included. Data in Brief. 2020;31: 105793. doi:10.1016/j.dib.2020.105793

132. Bocquet-Appel J-P, Demars P-Y, Noiret L, Dobrowsky D. Estimates of Upper Palaeolithic meta-population size in Europe from archaeological data. J Archaeol Sci. 2005;32: 1656–1668. doi:10.1016/j.jas.2005.05.006

133. Langlais M, Fat Cheung C. Le site laborien d'Auberoche (Le Change, Dordogne), collections Daleau et Daniel: Anciennes fouilles, nouvelle attribution. Bull Soc Préhist Fr. 2019;116: 155–158.

134. Ordonez A, Riede F. Changes in limiting factors for forager population dynamics in Europe across
the last glacial-interglacial transition. Nat Commun. 2022;13: 5140. doi:10.1038/s41467-022-32750-
x

135. Halstead P, O'Shea J, editors. Bad Year Economics: Cultural Responses to Risk and Uncertainty.
Cambridge: Cambridge University Press; 1989. doi:10.1017/CBO9780511521218

136. Miller R. Mapping the expansion of the Northwest Magdalenian. Quat Int. 2012;272–273: 209–230.
doi:10.1016/j.quaint.2012.05.034

137. Riede F. LSTbase - a collation of Laacher See tephra occurrences. Zenodo; 2024.
doi:10.5281/ZENODO.13732535

138. Reinig F, Wacker L, Jöris O, Oppenheimer C, Guidobaldi G, Nievergelt D, et al. Precise date for the
Laacher See eruption synchronizes the Younger Dryas. Nature. 2021;595: 66–69.
doi:10.1038/s41586-021-03608-x

139. Theuerkauf M, Joosten H. Younger Dryas cold stage vegetation patterns of central Europe - climate,
soil and relief controls. Boreas. 2012;41: 391–407. doi:10.1111/j.1502-3885.2011.00240.x

140. Crombé P. The impact of environmental changes on the human occupation of the Scheldt basin
during the Younger Dryas. In: Montoya C, Fagnart J-P, Locht J-L, editors. Préhistoire de l'Europe
du nord-ouest: mobilités, climats et identités culturelles Volume 2: Paléolithique supérieur ancien,
Paléolithique final – Mésolithique, Congrès préhistorique de France: Compte Rendu de la 28ème
session. Paris: Société Préhistorique Française; 2019. pp. 407–416.

141. Bos JAA, Gelorini V, Van Der Meeren T, Peleman J, Court-Picon M, Demiddele H, et al. The
Younger Dryas and Preboreal landscape in the Moervaart area (northwestern Belgium) and the
apparent decrease in human occupation. Veget Hist Archaeobot. 2018;27: 697–715.
doi:10.1007/s00334-017-0666-3

142. Latałowa M. Late Glacial 14,000 –10,000 14C yr BP (ca. 15,500 (16,000)-11,500 cal yr BP). In:
Ralska-Jasiewiczowa M, Latałowa M, Wasylikowa K, Tobolski K, Madeyska E, Wright jr. HE, et
al., editors. Late glacial and holocene history of vegetation in Poland based on isopollen maps.
Kraków: W. Szafer Institute of Botany; 2004. pp. 385–391.

143. Ammann B, Birks HJB, Brooks SJ, Eicher U, Von Grafenstein U, Hofmann W, et al. Quantification
of biotic responses to rapid climatic changes around the Younger Dryas — a synthesis. Palaeogeogr
Palaeoclimatol Palaeoecol. 2000;159: 313–347. doi:10.1016/S0031-0182(00)00092-4

144. Magny M, Aalbersberg G, Bégeot C, Benoitruffaldi P, Bossuet G, Disnar J, et al. Environmental and
climatic changes in the Jura mountains (eastern France) during the Lateglacial–Holocene transition:
a multi-proxy record from Lake Lautrey. Quat Sci Rev. 2006;25: 414–445.
doi:10.1016/j.quascirev.2005.02.005

145. Cupillard C, Magny M, Bocherens H, Bridault A, Bégeot C, Bichet V, et al. Changes in ecosystems,
climate and societies in the Jura Mountains between 40 and 8 ka cal BP. Quat Int. 2015;378: 40–72.
doi:10.1016/j.quaint.2014.05.032

146. Isarin RFB, Bohncke SJP. Mean July Temperatures during the Younger Dryas in Northwestern and Central Europe as Inferred from Climate Indicator Plant Species. Quat Res. 1999;51: 158–173. doi:10.1006/qres.1998.2023

147. Van Asch N, Kloos ME, Heiri O, De Klerk P, Hoek WZ. The younger dryas cooling in northeast Germany: summer temperature and environmental changes in the Friedländer Groβe Wiese region. J Quat Sci. 2012;27: 531–543. doi:10.1002/jqs.2547

148. Merkle JA, Monteith KL, Aikens EO, Hayes MM, Hersey KR, Middleton AD, et al. Large herbivores surf waves of green-up during spring. Proc R Soc B. 2016;283: 20160456. doi:10.1098/rspb.2016.0456

149. Maier A, Tharandt L, Linsel F, Krakov V, Ludwig P. Where the Grass is Greener — Large-Scale Phenological Patterns and Their Explanatory Potential for the Distribution of Paleolithic Hunter-Gatherers in Europe. J Archaeol Method Theory. 2023 [cited 9 Feb 2024]. doi:10.1007/s10816-023-09628-3

150. Price TD, Meiggs D, Weber M-J, Pike-Tay A. The migration of Late Pleistocene reindeer: isotopic evidence from northern Europe. Archaeol Anthropol Sci. 2017;9: 371–394. doi:10.1007/s12520-015-0290-z

151. Rivals F, Drucker DG, Weber M-J, Audouze F, Enloe JG. Dietary traits and habitats of the reindeer (Rangifer tarandus) during the Late Glacial of Northern Europe. Archaeol Anthropol Sci. 2020;12: 98. doi:10.1007/s12520-020-01052-y

152. Usinger H. Ein weit verbreiteter Hiatus in spätglazialen Seesedimenten: mögliche Ursache für Fehlinterpretationen von Pollendiagrammen und Hinweis auf klimatisch verursachte Seespiegelbewegungen. Eiszeitalter und Gegenwart. 1981;31: 91–107.

153. Renssen H, Goosse H, Roche DM, Seppä H. The global hydroclimate response during the Younger Dryas event. Quat Sci Rev. 2018;193: 84–97. doi:10.1016/j.quascirev.2018.05.033

154. Kolstrup E. Lateglacial older and younger coversand in northwest Europe: chronology and relation to climate and vegetation. Boreas. 2007;36: 65–75. doi:10.1111/j.1502-3885.2007.tb01181.x

155. Brauer A, Haug GH, Dulski P, Sigman DM, Negendank JFW. An abrupt wind shift in western Europe at the onset of the Younger Dryas cold period. Nat Geosci. 2008;1: 520–523. doi:10.1038/ngeo263

156. Clausen I, Hartz S. Fundplätze des Spätglazials am Sorgetal bei Alt-Duvenstedt, Kreis Rendsburg-Eckernförde. Offa. 1988;45: 17–35.

157. Mussi M, Peresani M. Human settlement of Italy during the Younger Dryas. Quat Int. 2011;242: 360–370. doi:10.1016/j.quaint.2011.03.008

158. Riede F. Changes in mid- and far-field human landscape use following the Laacher See eruption (c. 13,000 BP). Quat Int. 2016;394: 37–50. doi:10.1016/j.quaint.2014.07.008

159. Riede F. Splendid isolation: the eruption of the Laacher See Volcano and Southern Scandinavian Late Glacial hunter-gatherers. Aarhus: Aarhus University Press; 2017.

160. Riede F, Bazely O. Testing the 'Laacher See hypothesis': a health hazard perspective. Journal of Archaeological Science. 2009;36: 675–683. doi:10.1016/j.jas.2008.10.013

161. Riede F, Kierdorf U. The eruption of the Laacher See volcano (~13,000 years BP) and possible fluoride poisoning amongst contemporaneous wildlife and human foragers—Outline of a hypothesis and the way to test it. Intl J of Osteoarchaeology. 2020;30: 855–871. doi:10.1002/oa.2916

162. Riede F, Wheeler JM. Testing the 'Laacher See hypothesis': tephra as dental abrasive. Journal of Archaeological Science. 2009;36: 2384–2391. doi:10.1016/j.jas.2009.06.020

163. Niemeier U, Riede F, Timmreck C. Simulation of ash clouds after a Laacher See-type eruption. Clim Past. 2021;17: 633–652. doi:10.5194/cp-17-633-2021

164. Andreasen LS, Cornér J, Abbott PM, Sinclair VA, Riede F, Timmreck C. Changes in Northern Hemisphere extra-tropical cyclone frequency following volcanic eruptions. Environ Res: Climate. 2024;3: 025002. doi:10.1088/2752-5295/ad2c0e

165. Hey J. On the Number of New World Founders: A Population Genetic Portrait of the Peopling of the Americas. PLoS Biol. 2005;3: e193. doi:10.1371/journal.pbio.0030193

166. Smith CM. Estimation of a genetically viable population for multigenerational interstellar voyaging: Review and data for project Hyperion. Acta Astron. 2014;97: 16–29. doi:10.1016/j.actaastro.2013.12.013

167. Bradshaw CJA, Ulm S, Williams AN, Bird MI, Roberts RG, Jacobs Z, et al. Minimum founding populations for the first peopling of Sahul. Nat Ecol Evol. 2019;3: 1057–1063. doi:10.1038/s41559-019-0902-6

168. Traill LW, Brook BW, Frankham RR, Bradshaw CJA. Pragmatic population viability targets in a rapidly changing world. Biol Conserv. 2010;143: 28–34. doi:10.1016/j.biocon.2009.09.001

169. Eller E, Hawks J, Relethford JH. Local Extinction and Recolonization, Species Effective Population Size, and Modern Human Origins. Hum Biol. 2004;76: 689–709. doi:10.3378/027.081.0623

170. Schmidt I, Zimmermann A. Population dynamics and socio-spatial organization of the Aurignacian: Scalable quantitative demographic data for western and central Europe. PLoS ONE. 2019;14: e0211562. doi:10.1371/journal.pone.0211562

171. Whallon R. Social networks and information: Non-"utilitarian" mobility among hunter-gatherers. J Anthropol Archaeol. 2006;25: 259–270. doi:10.1016/j.jaa.2005.11.004

172. Mandryk CAS. Hunter-Gatherer Social Costs and the Nonviability of Submarginal Environments. J Anthropol Res. 1993;49: 39–71. doi:10.1086/jar.49.1.3630629

173. Riede F. Climate Change, Demography and Social Relations: An Alternative View of the Late Palaeolithic Pioneer Colonization of Southern Scandinavia. In: McCartan S, Schulting R, Warren G, Woodman P, editors. Mesolithic Horizons Papers Presented at the Seventh International Conference on the Mesolithic in Europe. Oxford: Oxbow; 2009. pp. 3–10.

174. Riede F. Success and Failure during the Late Glacial Pioneer Human Re-Colonisation of Southern
Scandinavia. In: Riede F, Tallavaara M, editors. Lateglacial and postglacial pioneers in Northern
Europe. Oxford: Archaeopress; 2014. pp. 33–52.

175. Maier A, Sauer F, Bergsvik KA. Transport Patterns as Heuristic Testing Variables for the Social
Coherence of Taxonomic Units at Different Spatial Scales. J Paleo Arch. 2022;5: 8.
doi:10.1007/s41982-022-00120-0

176. Delvigne V, Lafarge A, Fernandes P, Pesesse D, Angevin R, Bindon P, et al. Quels territoires en
préhistoire? Une analyse par réseaux de lieux pour penser l'espace au Paléolithique supérieur. In:
Mevel L, Weber M-J, Maier A, editors. En Mouvement / On the Move / In Bewegung. Paris: Société
Préhistorique Française; 2021. pp. 27–69.

177. Floss H. Rohmaterialversorgung im Paläolithikum des Mittelrheingebietes. Bonn: Habelt; 1994.

178. Binford LR. Organization and Formation Processes: Looking at Curated Technologies. J Anthropol
Res. 1979;35: 255–273. doi:10.1086/jar.35.3.3629902

179. Tomasso A, Porraz G. Hunter-gatherer mobility and embedded raw-material procurement strategies
in the mediterranean Upper Paleolithic. Evol Anthropol. 2016;25: 164–174. doi:10.1002/evan.21488

180. Eriksen BV. Fossil mollusks and exotic raw materials in late glacial and early postglacial find
contexts – a complement to lithic studies. In: Fisher LE, Eriksen BV, editors. Lithic raw material
economies in late Glacial and early Postglacial Europe. Oxford: Archaeopress; 2002. pp. 27–52.

181. Álvarez-Fernández E. Magdalenian personal ornaments on the move: a review of the current
evidence in Central Europe. Zephyrus. 2009;63: 45–59.

182. Schwendler RH. Diversity in social organization across Magdalenian Western Europe ca. 17–12,000
BP. Quat Int. 2012;272–273: 333–353. doi:10.1016/j.quaint.2012.03.054

183. Maier A. The Central European Magdalenian: Regional Diversity and Internal Variability.
Dordrecht: Springer; 2015. doi:10.1007/978-94-017-7206-8

184. Burdukiewicz JM. Further research on amber in Palaeolithic archaeology. In: Burdukiewicz JM,
Cyrek K, Dyczek P, Szymczak K, editors. Understanding the past: papers offered to Stefan K
Kozłowski. Warsaw: Center for Research on the Antiquity of Southeastern Europe, University of
Warsaw; 2009. pp. 69–74.

185. Petersen PV. Zigzag lines and other protective patterns in Palaeolithic and Mesolithic art. Quat Int.
2021;573: 66–74. doi:10.1016/j.quaint.2019.09.029

186. Thévenin A. Géographie et cultures au Tardiglaciaire- L'impact de l'axe Rhône Saône. In: Pion G,
editor. Le paléolithique supérieur récent: nouvelles données sur le peuplement et l'environnement
Actes de la Table ronde de Chambéry. Paris: Société préhistorique française; 2000. pp. 67–79.

187. Thévenin A. L'espace culturel «Méditerranéen» ses expansions de la fin du Tardiglaciaire au début
du Postglaciaire. In: Barbaza M, Jaubert J, editors. Territoires, déplacements, mobilité, échanges
pendant la Préhistoire: terres et hommes du Sud Actes du 126e Congrès national des sociétés
historiques et scientifiques, Toulouse, 2001. Paris: Comité des travaux historiques et scientifiques;
2005. pp. 429–451.

188. Mevel L. Les premières sociétés aziliennes : nouvelle lecture de la genèse du phénomène
d'azilianisation dans les Alpes du Nord à partir des deux niveaux d'occupation de l'abri de La Fru
(Saint-Christophe-la-Grotte, Savoie). Bull Soc Préhist Fr. 2013;110: 657–689.
doi:10.3406/bspf.2013.14319

189. Jacquier J, Langlais M, Naudinot N. Late Laborian trapezoids: Function and origin of the first
transverse projectile tips of Western Europe prehistory. Quat Int. 2020;564: 48–60.
doi:10.1016/j.quaint.2020.01.009

1230

1231

Supporting information

Information on Maps

Tables

S1 Table. Sites with data on Raw Material Catchment Areas (RMCA) assigned to GI-1d-a.

S2 Table. Sites with data on Raw Material Catchment Areas (RMCA) assigned to GS-1.

S3 Table. Regionally distinguished demographic estimates for GI-1d-a based on dataset B (i.e., only directly dated or specifically attributable to GI-1d-a).

S4 Table. Regionally distinguished demographic estimates for GS-1 based on dataset B (i.e., only directly dated or specifically attributable to GS-1).

S5 Table. Number of radiocarbon-dated sites divided by GI-1d-a and GS-1.

Figures

S1 Fig. Maps (left) and diagrams (right) from the definition of the Core Areas based on the sites dated to Greenland Interstadial (GI) 1d-a.

S2 Fig. Maps (left) and diagrams (right) from the definition of the Core Areas based on the sites dated to Greenland Stadial (GS) 1.

S3 Fig. Map of all Raw Material Catchment Areas (RMCA) from sites dated to GI-1d-a that were collected from the literature (n = 108).

S4 Fig. Map of all Raw Material Catchment Areas (RMCA) from sites dated to GS-1 that were collected from the literature (n = 97).

S5 Fig. GI-1d-a: Comparison of estimates of people for dataset A (left bar) and B (right bar) per region.

S6 Fig. GS-1: Comparison of estimates of people for dataset A (left bar) and B (right bar) per region.

S7 Fig. Diachronic tendencies in the frequency of regionally differentiated radiocarbon dates per region.

1255   S8 Fig. Comparison of Final Palaeolithic Core Areas (data B) from GI-1d-a (green areas) to GS-1 (blue

1256   areas).

Supporting information References

1258