Title:

Attentional Information Routing in The Human Brain

Authors: Kianoush Banaie Boroujeni[1*], Randolph F. Helfrich[2], Ian C. Fiebelkorn[3], Nicole

Bentley[4], Jack J. Lin[5], Robert T. Knight[6], Sabine Kastner[1,7]

Affiliations:

[1]Princeton Neuroscience Institute, Princeton University, Princeton, NJ 08544
[2]Hertie Institute for Clinical Brain Research, Center for Neurology, University Medical Center Tübingen, Tübingen,
Germany.
[3]Department of Neuroscience and Del Monte Institute for Neuroscience, University of Rochester, Rochester, NY
14627, USA
[4]Department of Neurosurgery, University of Alabama at Birmingham, Birmingham, Alabama, USA
[5]Department of Neurology, University of California Davis, Davis, CA, USA
[6]Departments of Psychology and Neuroscience, University of California Berkeley, Berkeley, CA, USA
[7]Department of Psychology, Princeton University, Princeton, NJ 08544
17
18

*Corresponding author: Kianoush Banaie Boroujeni. Email: kb9590@princeton.edu

Abstract

Brain-wide communication supports behaviors that require coordination between sensory

and associative regions. However, how large-scale brain networks route sensory information at

fast timescales to guide upcoming actions remains unclear. Using spiking neural networks and

human intracranial electrophysiology during spatial attention tasks, where participants detected a

target at cued locations, we show that high-frequency activity bursts (HFAb) serve as information-

carrying events, facilitating fast and long-range communications. HFAbs emerged as bouts of

neural population spiking and were coordinated brain-wide through low-frequency rhythms. At

the network-level, HFAb coordination identified distinct cue- and target-activated subnetworks.

HFAbs following the cue onset in cue-subnetworks predicted successful target detection and

preceded the information in target-subnetworks following target onset. Our findings suggest

HFAbs as a neural mechanism for fast brain-wide information routing that supports attentional

performance.

Introduction

Prioritizing information from the external environment to guide ongoing behavior and upcoming

actions requires coordination of neural activity on fast timescales across large-scale networks

distributed across distant brain areas (*1−7*). This fast coordination allows information to be routed

selectively from sensory to higher level executive brain networks (*8−12*). However, the

mechanisms by which these fast neural dynamics emerge and enable information routing within

large-scale brain networks are not well understood.

Investigating these questions at the network-level in the human brain is challenging due to spatial

or temporal constraints of the techniques that have been employed. Studies examining network-

level interactions have largely been based on connectivity maps derived from functional magnetic

resonance imaging (fMRI) which has insufficient temporal resolution to capture fast routing

dynamics   (*13−17*).   Studies   using   high   temporal   resolution   methods   such   as

electroencephalography (EEG) and magnetoencephalography (MEG), on the other hand, have

limitations in spatial precision for measuring network dynamics (*18*, *19*). Lastly, single unit

recordings provide both fine temporal and spatial signals but lack the broad coverage for

addressing questions of brain-wide communication (*7*, *20*). In addition to these technical

challenges, cortex-wide neural dynamics have been mainly evaluated via spectral coherence

between brain regions in low-frequency bands or coupling of high frequency activity to low

frequency rhythms, which provides only an indirect measure of information routing and also

cannot capture fast network interactions (*21–27*).

High frequency broadband activity (HFB) in human intracranial electroencephalography (iEEG),

offers a unique opportunity to address these limitations by providing spatially localized and

temporally precise data that can be instrumental to understanding fast cortical dynamics (*28*, *29*).

These high-frequency activities can reflect different neural mechanisms depending on their

recording source, including aggregated spiking activity of nearby neurons, neural post-synaptic

activity, brain state shifting into spiking states, and spike current leakage to local field potentials

(LFPs) (*30–33*). HFB has been widely reported to correlate with different cognitive functions,

including attentional processes (*24*, *28*, *29*, *34–37*). In addition, HFBs show long-range phase

synchronization with different frequency bands in the human brain, making them interesting

candidates for studying fast and brain-wide communications (*22*, *24*, *38–40*). While previous

research has highlighted the significant engagement of HFB in cognitive processes and its potential

involvement in interareal communication through coupling to low-frequency rhythms, it remains

unclear how HFB dynamics can inform about brain-wide communication and information routing

at fast timescales.

Here, we established a novel method to investigate fast information routing within large-scale

brain networks. We first detected transient high-frequency activity bursts (HFAbs) in human iEEG

data from epilepsy patients performing spatial attention tasks. HFAbs were brain-wide

synchronized and linked to slow rhythms (4-25 Hz) and were found to occur more frequently

during sensory and perceptual processing. HFAbs during cue processing and their local decoupling

from low-frequency dynamics were associated with successful target detection. Network-level

synchronization of HFAbs, identified functionally specialized brain subnetworks that activated in

response to cues and targets. These subnetworks showed distinct topographical and temporal

organization, with activity in cue-subnetworks preceding that in target-subnetworks. Using

computational modeling, we further showed that HFAbs may reflect state transitions in neural

populations into a coherent spiking activity state, characterized by bouts of elevated neuronal

firing, which suggests HFAbs can serve as neural population-level signatures of information-

encoding events. Our results demonstrate a novel mechanism for fast neural coordination in large-

scale brain networks that supports attentional processes and facilitates information routing between

functionally specialized brain subnetworks.

Results

HFA bursts predict behavioral accuracy

To examine neural mechanisms of fast information routing we used human iEEG data recorded

from epilepsy patients to quantify high frequency activity dynamics. We analyzed data from two

spatial attention tasks in which patients were cued either exogenously or endogenously to a spatial

location to detect visual targets (**Fig. 1A**, **S1A**). Throughout the paper, the main figures mostly

present results of experiment 1, while the results of experiment 2 are presented as supplementary

information to indicate reproducibility and generalization across attention tasks (for brain heatmap

plots, we combined both experiments to improve 3D rendering coverage).

In experiment 1, patients performed a spatial attention task as described in (*41*). Each trial

started with the presentation of two vertical or horizontal bar stimuli. Patients were instructed to

fixate their gaze at the center of the display. A transient spatial cue appeared at the end of one bar,

exogenously cueing an upcoming target location. Following a delay period, a target (i.e.,

luminance changes at perceptual threshold) appeared at the cued location or infrequently at equally

distant non-cued locations. Patients were required to respond if they detected the target (**Fig.**

1A). In experiment 2, patients were endogenously cued to a hemifield and expected to report a

target if it appeared in the cued hemifield (**Fig. S1A**, see **Methods,** and for more details see (*4*)).

First, we used an adaptive method (*7*) and detected reliable bursts of high frequency activity at

each electrode (HFAbs, 65-115 Hz, intermittent high amplitude oscillatory events lasting more

than 2.5 frequency cycles; **Fig. 1B**). The frequency band was selected based on average spectral

peaks observed across subjects (91.2 ± 20.9 Hz, n = 12, see **Methods** for further details). We then

calculated the HFAb density (number of HFAb events per unit of time) for each electrode to

examine their evoked response by different task epochs. HFAbs showed higher density averaged

across all electrodes in response to both cue and target (**Fig. 1B, S1B, S2A**). The HFAb activation

to targets was in most subjects negatively correlated with that to the cue, suggesting distinct groups

of electrodes processing cues and targets (4 of n = 7, Spearman correlation, p<0.05, R = -0.05 ±

0.07, mean ± standard error across all 7 participants, **Fig. 1D**). The topography of burst responses

to cues was largely confined to occipital and superior parietal regions (**Fig. 1E**), whereas those to

target responses were more widely distributed across different brain areas, including parietal,

motor, pre-motor, and lateral prefrontal cortices (**Fig. 1F**).

At the population level, HFAb responses to cues tended to be stronger in correct than in incorrect

trials, while the population-level burst responses to targets were significantly stronger for correct

detections (random permutation test. P<0.05, **Fig. 1G**, **S1C**). Cue responses (the average

normalized HFAb density within 500 ms following the cue-onset) were stronger in parietal areas,

whereas HFAb target-evoked responses were found to be stronger in motor/premotor areas for

correct, but not for missed trials (**Fig. 1H**, also see **Fig. S1E, S2C** for more individual examples).

Overall, cue responses were non-zero for correct hit and reject trials (Wilcoxon signed-rank test,

P<0.001), and they were also different in correct as compared to missed and false alarm trials

(Kruskal Wallis test, P=0.001, Dunn's test, P=0.008 and P=0.01, respectively; **Fig. 1I**). Target

responses were significantly different between correct hit and other outcome conditions (Kruskal

Wallis test, P<0.001, Dunn's test, P<0.001, **Fig. 1I**). We used Generalized Linear Mixed Effect

(GLME) models to examine the relationship between cue and target processing and outcome on a

trial-by-trial basis (see **Methods**). There was a main effect of outcome on both cue response (t=2.8,

P=0.005) and target response (GLME, t=15.2, P<0.001). Further, there was a main effect of cue

validity on the target response (GLME, t=4.7, P<0.001), and a main effect of the cue laterality on

the cue response (GLME, t=5.1, P<0.001). No significant main effect was observed for laterality

on target response (**Fig. S3A**, P=0.13), or validity on cue response (**Fig. S3B**, P=0.32, See **Fig.**

S3C,D for the effect of laterality and cue validity on cue and target response in experiment 2, as

well as **Fig. S3E,F** for the temporal profiles of HFAbs around cue and target onsets in both

experiments).

Next, we examined whether the burst density of electrodes that showed significant activation in

response to cues (Cue+ electrodes, see **Methods**) predicted whether participants would

successfully complete a trial (correct hits and correct rejections). We trained a classifier on the

burst density using a sliding window of 350 ms around the cue onset. In Cue+ electrodes and

within 500 ms after the cue onset, the burst density predicted higher than the baseline level and the

chance level if the trial would be executed correctly (binomial test, FDR corrected for dependent

samples, P<0.05, **Fig. 1L, S3G**). This prediction was consistent across both experiments (**Fig.**

S3H) and was not dependent on an individual subject (**Fig. S3I**).

These findings indicate that HFAbs occur more frequently in response to sensory cues and targets,

while exhibiting distinct spatial profiles across the brain. The cue response was predominantly

observed in occipital and parietal regions and predicted performance accuracy following the cue-

onset on a trial-by-trial basis. These results suggest a potential role for HFAbs in encoding and

routing spatially relevant sensory information for subsequent target detection.

HFA bursts are synchronized in cortical networks

High frequency activity dynamics in brain networks are shown to be organized by low-frequency

rhythms, involving in different cognitive processes (*22*). Thus, we examined if HFAbs are

coordinated with slower rhythms of local population activity, and if behavioral variables affect

these local coordination patterns. For each electrode, we extracted the LFP around the HFAb

centers and measured the HFAb-triggered LFP average, as well as phase locking value (PLV) of

HFAbs to the low frequency dynamics of LFP. The HFAb-triggered LFP showed consistent

evoked potentials with low frequency side-lobes across all subjects (**Fig. 2A**). HFAbs triggered

spectral peaks (**Fig. 2B**, top) and phase locking peaks (**Fig. 2B**, bottom) predominantly in the theta

frequency band (4-8 Hz), as well as in alpha and beta frequency bands (8-14 Hz and 15-25 Hz,

respectively). The phase locking of HFAbs to the local low-frequency field dynamics was evident

in most electrodes and across all subjects (**Fig. 2C,** 9 ± 0.24 Hz**,** significant peaks are shown with

black dots; see **Fig. 2D, S4C** for examples of individual electrodes showing phase distributions of

HFAbs locked to one or more frequency bands, also see **Fig. S4A,B** for a topographic

representation of the frequency of phase locking peaks at the group average and individual subject

levels).

We next examined whether sensory cues or target processing affected HFAb phase locking to

the local population. Overall, HFAbs showed a transient decrease in their local phase locking to

low-frequency rhythms relative to the baseline following cue and target onsets (**Fig. 2E**). This

decrease in phase locking value is referred to as *decoupling*. HFAbs decoupled from the theta/alpha

and beta frequency bands after the cue onset and after the target onset (**Fig. 2F, S4D,**

randomization test, $P < 0.05$, FDR corrected for dependent samples). This decoupling was not

induced by event-triggered potentials (**Fig. S4E**, see **Methods**). Also HFAb coupling strength

changed only in correct trials, but not in incorrect trials (**Fig. 2G,H,** the number of bursts was

controlled across trial conditions, see **Methods**, see **Fig. S4F** for individual examples). The

decoupling was consistent across most participants in both tasks at the network level (**Fig. 2I,**

S4G,H, 4 and 8 of the $n = 12$ subjects showed significant decoupling at the network level after the

cue and target onset, respectively; Wilcoxon rank-sum test, $P<0.05$).

We then examined whether alterations in HFAb coupling to low frequencies were related to HFAb

activation profiles in response to cues and targets. We calculated correlations between coupling

ratios and burst density for each subject (a coupling ratio measured changes in HFAb coupling to

low frequencies after cues or targets relative to the baseline within 1000 ms before the cue and

target onset, see **Methods**). The coupling ratio at target and cue onsets was negatively correlated

with burst density for most participants (**Fig. 2J, S4I**, 8 of $n = 12$ subjects showed a significant

negative correlation, Spearman correlation, $P<0.05$). Burst density and coupling ratio were

correlated within the same event, indicating that coupling was not dependent upon burst activation

to the other events (**Fig. 2J**, Fisher test, $P<0.001$). Overall, the dynamic coupling of HFAbs to

local low-frequency activity suggests that while these HFAbs are primarily organized by local

dynamics, they can decouple from their local network when receiving and processing sensory

information as well as during perceptual processing and decision making.

HFAbs were coordinated brain-wide and their baseline synchronization identified

functionally specialized subnetworks

We found that HFAbs evoked by cue and target demonstrated different topography. In addition,

the processing of cues and targets involved the decoupling of HFAbs from local low-frequency

dynamics. These observations suggest the possibility that brain networks may coordinate at a large

scale when processing these events, rather than processing them locally. Thus, we asked whether

the brain-wide network could be organized into distinct functionally specialized subnetworks, each

characterized by unique high frequency coordination patterns. To avoid stimulus-driven

coordination in the network, we only considered burst events outside of the cue/delay and

target/response periods. To examine this coordination, we measured the power of high frequency

broadband activity (65-175 Hz, HFB, to capture broader spectral contents) in one electrode aligned

to the center of HFAb events in another electrode (HFAb-triggered HFB). HFAb-triggered HFB

between most channels revealed activity coordination in high frequency bands (**Fig. 3A**). The

HFAb coordination between electrodes was organized through a low frequency rhythm showing

spectral peaks mostly in theta band (4-8 Hz) among the majority of electrodes (**Fig. 3B**, top) and

consistent across all subjects (**Fig. 3B**, bottom, similar patterns were observed in experiment 2,

Fig. S5A,B). HFAb coordination strength (the sharpness of HFAb-triggered HFB) was inversely

related to distance between electrodes, with closer sites showing stronger coordination (**Fig. 3C**,

S5E, Spearman correlation between coordination kurtosis and distance between electrodes, R = -

0.25, P<0.001).

To extract prominent temporal features of the high frequency coordination in the network, we

reduced dimensionality of HFAb-triggered HFB by Principal Component Analysis (PCA),

considering each electrode pair as a single dimension. All subjects showed zero-lag synchronized

and near-symmetric distributions of high frequency activity in their first component, explaining

on average more than 20% of the total variance (**Fig. 3D, Fig. S5D,E**). We projected HFAb

coordination between electrode pairs onto their PC space and calculated the scores for all electrode

pairs on the synchronized PC (**Fig. S5F**, PC1). Using a correlation-based K-means clustering

technique, we identified electrodes that were consistently grouped together based on their score on

this synchronized component (**Fig. S5F,** see **Methods**). While the clustering algorithm generates

most stable clusters for each cluster number (K = 2-8, see **Fig. S6A** and **S7A** for individual

examples), we defined a series of accuracy metrics to find a balance between the highest number

of reliable clusters and the highest degree of accuracy in each subject (**Fig. S6B,C** and **Fig. S7B,C,**

see **Methods**). An example of clustering for one subject with eight optimal clusters is shown (**Fig.**

3E, also see **Fig. S8** for topographical organizations of clusters in different subjects).

Next, we investigated whether these clusters were functionally specialized. In most subjects,

clusters showed distinct activation profiles following cue and target onsets (**Fig. 3G,** see **Fig. S9**

for an example of activation profiles for different cluster numbers, also see **Fig. S6D** and **Fig. S7D**

for additional examples). We labeled clusters that were significantly activated by cue and target

events as *cue* and *target-subnetworks*, respectively (**Fig. 3**h; Wilcoxon signed-rank test, P<0.05;

FDR corrected for dependent sample). Across both tasks, 6 out of =12 subjects showed stable cue-

subnetworks, and 10 showed stable target networks, respectively. Cue-subnetworks were

predominantly located in occipital and parietal cortices, while target-subnetworks were more

widely distributed across different brain areas including parietal, motor, premotor, and lateral

prefrontal cortices (**Fig. 3H**).

We examined if stable cue and target-subnetworks could predict trial outcomes. A classifier was

trained on the HFAb density of cue and target-subnetworks in sliding windows around the cue

onset to predict trial outcomes. The HFAb density within 98-374 ms after sensory cues predicted

successful detection of upcoming targets better than both baseline and chance level (binomial test,

P<0.001, FDR corrected for dependent samples). This prediction was only true for the cue-

subnetwork (**Fig. 3I**) and was not driven by a single subject (see **Fig. S10** for more details on the

classifier results in both experiments). As a control analysis, we re-referenced the datasets from

common average referencing to local composite referencing (LCR, each electrode was referenced

to its nearest neighbors, see **Methods**), which did not alter any of the main results.

Together, these findings suggest that large-scale brain networks can be divided into subnetworks

with distinct HFAb temporal dynamics. These subnetworks are further characterized with their

unique responses to sensory cues and targets, with the cue-activated subnetworks containing

information predictive of performance accuracy following cue onsets.

Cue-subnetworks precede the information flow triggered by targets within target-

subnetworks

Our observation that cue-subnetworks responses predict performance accuracy led to a key

question: whether cue-subnetworks routed information to target-subnetworks during target

processing. To address this question, we examined the information flow between cue and target-

subnetworks. We only considered subjects that showed both stable cue and target-subnetworks (3

subjects in each experiment). We used two different approaches. First, we quantified whether

HFAbs exhibited any temporally ordered activity pattern during the target-to-response period

between cue and target-subnetworks. When we averaged the time-lags between cue and target-

subnetworks over different cluster numbers, we observed that the HFAbs in occipital, posterior

parietal, and prefrontal areas led over motor/premotor areas during target processing (**Fig. 4A**, see

Fig. S11A for more examples). Individual subjects showed stronger HFB in target-subnetworks

within 150 ms after HFAbs in cue-subnetworks (**Fig. 4B, Fig. S11B,** Wilcoxon rank-sum test,

$p<0.05$, the opposite pattern was observed in the other direction). On average, HFAbs in cue-

subnetworks led the activity of target-subnetworks while target-subnetworks lagged cue-

subnetworks during the target processing period (within 150 ms around the HFAb-triggered HFB,

Wilcoxon rank-sum test, $p<0.001$, **Fig. 4C).** For comparison, no lead-lag patterns were observed

between cue and target-subnetworks during the cue-to-target interval **(**$P=0.44$, n.s., **Fig. S11C).**

We quantified the peak of these lead-lag relationships for individual subjects. The HFB in target-

subnetworks showed an activity peak following HFAbs in cue-subnetworks, while the peak of

HFB in cue-subnetworks preceded HFAbs in target-subnetworks (**Fig. 4D**, Wilcoxon rank-sum

test, $p<0.05$, 16 ± 7 ms; $n = 6$**).**

Next, we used delayed mutual information (DMI) to further quantify how information was

directionally coupled between cue and target-subnetworks. DMI can inform about (i) when cue

and target-subnetworks shared the most information relative to target onset, and (ii) the specific

time-lags at which these two subnetworks showed maximum inter-predictability (see **Fig. 4E** for

an individual example).

The DMI between cue and target-subnetworks showed a peak up to 500 ms after target onset (**Fig.**

4F, 275 ± 22 ms; n = 6). Time-lag distributions of the maximum DMI between cue and target-

subnetworks showed that cue-subnetworks preceded the information in target-subnetworks during

the target processing period (Wilcoxon signed-rank test, P<0.001; 67 ± 16ms, n = 6, **Fig. 4F**). This

pattern was consistent for individual subjects (**Fig. 4E,G**, see **Fig. S11D** for more examples) as

well as on an average group level (**Fig. 4F**). The information precession in cue-subnetworks over

target-subnetworks was only evident after the target onset (no significant time-lag was observed

around the cue onset, showing this effect is not a reflection of visual hierarchy activation **Fig.**

S11E), and limited to trials with a valid sensory cue (the target appeared as instructed by the visual

cue, **Fig. 4G**). Overall, these results indicate that information processing in cue-subnetworks

precedes that in target-subnetworks when the cue poses valid information about the target.

Computational modeling of HFAbs through spiking neural networks

We used spiking neural networks to investigate potential neural mechanisms underlying HFAbs

in iEEG signals. We simulated two interconnected networks of spiking neurons (*42*), each

consisting of 1000 neurons, including an average of ~80% excitatory pyramidal neurons and ~20%

inhibitory interneurons with different subtypes (**Fig. 5A, S12A,** see **Methods**). Each network was

fed by external input currents primarily to the excitatory population.

We implemented two recording sites on top of each network measuring electrical field dynamics

of postsynaptic and transmembrane potentials from all neurons in that network (**Fig. 5A**). A non-

ohmic filter was used to attenuate higher frequencies in the iEEG signal (**Fig. 5B**, **S12B,C** see

Methods). The recorded iEEG signal showed low-frequency spectral peaks reflecting the

resonance frequency of each network (*7*) and high frequency spectral peaks in the range of 65-115

Hz (**Fig. 5C**). Similar to our experimental results, we detected HFAbs as transient oscillatory

events of high amplitude 65-115Hz at each recording site (**Fig. 5D**).

To investigate the neural mechanisms underlying these HFAbs, we calculated the spike density for

the population of neurons, and the HFAb density (number of bursts per unit of time) from each

recording site. Burst density was significantly correlated with spike density as compared to bursts

with random timings (randomization test, P<0.05, **Fig. 5D**). This was further quantified by feeding

the network different levels of external current coherence (**Fig. S12C**). Increasing the input

coherence to the network showed a linear increase in correlation between the HFAb density and

the spike density (Spearman correlation, R=0.87, P<0.001, **Fig. 5E**). This observation is consistent

with our experimental findings that HFAbs are evoked in response to cues and targets, suggesting

that they may reflect large-scale information-carrying events emerging from transiently elevated

excitation in local networks of neurons.

Next, we asked how HFAbs coordinate with slower rhythms locally. We measured HFAb-

triggered LFPs at each recording site, which showed an average evoked response with low-

frequency sidelobes (**Fig. 5F**). The HFAb-triggered LFP spectrum showed peaks in low

frequencies as well as the burst frequency band (**Fig. 5G**). The cross-spectral correlation was

dependent on synaptic time constants, external input strength, neural connectivity strength, and

coherent input consistency (**Fig. 5G**). In addition to the spectral correlation between HFAbs and

the LFP, HFAbs also showed phase locking to different frequencies in the theta, alpha and beta

frequency bands (4-25 Hz, **Fig. 5H**). These results are consistent with the prominent local cross-

spectral coupling observed in our experimental data, suggesting local neural network mechanisms

underlying the coupling of high- and low-frequency dynamics.

We also tested for possible network mechanisms underlying the decoupling following cue and

target processing. We modeled this scenario through feedforward networks and measured the time-

resolved PLV for both networks after feeding one with a brief input pulse (~50% of the population

simultaneously received in-phase inputs simulating an external stimuli; **Fig. 5I, S12E,F**). The

external impulse desynchronized both networks, resulting in a transient decoupling of HFAb from

low-frequency activity as compared to the baseline (Wilcoxon rank-sum test, P<0.001, **Fig. 5I,**

S12G,H). This confirmed that desynchronization was not due to lower burst rate (**Fig. S12I**).

Lastly, we examined how HFAbs might be coordinated between distributed networks of neural

populations. Similar to our experimental data, we measured HFAb-triggered HFB by aligning HFB

power in one network to HFAb times in the other network. HFAbs showed different rhythmically

organized coordination patterns between networks depending on inter-network connectivity and

external inputs. Reciprocally connected networks and networks with correlated external

inputs showed synchronized high frequency activity (**Fig. 5J**), while feedforward

networks showed a lead-lag pattern within 150 ms following HFAbs (Wilcoxon rank-sum test,

P<0.001, **Fig. 5K**). In addition to the internally generated rhythmic structure of HFAbs between

networks, external inputs may also contribute to these rhythmic coordination patterns. A rhythmic

input to either or both networks entrained its rhythm in the coordination between the two networks

(Spearman correlation, R = 0.83, P < 0.001, **Fig. 5L,M**).

In sum, our modeling results suggest that HFAbs reflect a state transition in neural populations

into a coherent spiking activity state. By default, these HFAbs are phase locked to low frequency

activity, but decouple when transient inputs interfere with internally generated network

states. HFAbs are coordinated across interconnected networks or networks with correlated external

drives.

Discussion

Here we present a novel framework for understanding the mechanisms underlying large-scale information routing in brain networks at fast timescales. We identified brain subnetworks characterized by similar high-frequency activity coordination patterns, which tracked the encoding and transfer of spatially relevant sensory cue information within large-scale brain networks. Using computational modeling, we showed that HFAbs can emerge as bouts of elevated excitatory drive to local networks, which may serve as fast communication units across the brain. These burst events were synchronized across distributed brain networks in iEEG data, and their network-level synchronization patterns revealed functionally specialized subnetworks for processing cues and targets. HFAbs in cue-subnetworks predicted successful target detections on a trial-by-trial basis and preceded the activity of target-subnetworks during target processing. These findings demonstrate, to our knowledge, for the first time that high-frequency activities can be involved in both forming a brain-wide representation of sensory inputs and facilitating the communication of these sensory signals to associative brain networks for detecting upcoming targets and decision-making processes.

High-frequency activities (HFB or high gamma) have been widely observed in human intracranial recordings and are associated with a wide range of cognitive and circuit functions (*22, 24, 26, 28, 29, 37, 38*). Our results align with these previous studies, showing prevalent high-frequency activity in human iEEG signals. However, instead of continuous measures, we characterized these activities as discrete burst events that may reflect coherent bouts of excitation in neural populations, as previously shown in monkey electrophysiology and modeling studies as an important mechanism in facilitating information routing between distant brain areas (*7, 43, 44*).

This discrete description of high-frequency dynamics is also supported by previous research in

animal models emphasizing their circuit-level origins and their essential roles in long range

communications (*43*, *45–49*). In addition, while cross-frequency coupling have typically been

observed between high-frequency amplitude and low-frequency phase in theta and alpha bands

(*21*, *22*, *24*, *41*), our discrete quantification of these high-frequencies enabled the study to explore

their interactions with a broader frequency range (including low and high beta bands, 15-35 Hz).

This unique characterization of high-frequency activities opens new avenues for understanding

fast and long-range information routing and network-level communication mechanisms in

different cognitive processes.

One of the main findings of our study was revealing the synchronized inter-regional temporal

structure of high-frequency coordination through HFAbs. We found that the most prominent

pattern of high frequency dynamics in the brain network was zero-lag amplitude synchronization

(**Fig. 3D**). This long-range synchronization of HFAbs is consistent with previous studies showing

phase locking of high frequency dynamics in brain networks (*38–40*). Zero-lag synchronization

patterns in cortical networks are thought to enhance large-scale information processing and

facilitate representational states of sensory information in the brain (*2*, *50–52*). While our results

cannot directly identify the origins of these synchronized interactions, two different scenarios

might be involved. One potential mechanism relies on subcortical areas (*53–57*), particularly the

higher-order thalamic nuclei (*53*, *58–61*). Our modeling suggested that long-range synchronization

might result from correlated inputs to two distinct networks of neurons (**Fig. 5L**). Such correlated

inputs, if originating from subcortical structures such as higher-order thalamic nuclei, can facilitate

the long-range synchronizations between separate or weakly connected cortical areas (*53, 62*).

Another mechanism depends on long-range intracortical interactions. We found that cortical

synchronization was largely between areas with similar activation profiles during cognitive

functions and spatially neighboring regions (**Fig. 3, S8**). Additionally, our modeling suggested that

reciprocally connected networks with similar connectivity strength can result in synchronized

dynamics (**Fig. 5J**). This zero-lag synchronization induced by reciprocal connections has been

observed experimentally in long-range neuronal interactions (*50, 51*) as well as in network

modeling (*63*). Recent findings in mice support this idea by showing that deep layer 6b neurons,

which are recipient of long-range cortical projection neurons (*64*), are involved in brain state

shifting into a spiking state, and strong high gamma activity in brain networks (*33*). While our

modeling and empirical results can support either of these mechanisms, future studies need to

explore the extent to which cortico-cortical versus thalamocortical interactions drive and influence

this high frequency synchronization in large-scale cortical networks.

We further showed that network-level baseline synchronization fingerprints of high-frequency

dynamics outside of task events can be used to identify distinct functional brain subnetworks

through a novel clustering framework. While the baseline state did not contain cue or target

processing periods, it could identify subnetworks with unique activation patterns in response to

those behavioral events (**Fig. 3**). Our findings are consistent with several fMRI studies suggesting

functional brain networks are continuously active, even during resting states (*65, 66*). However,

using methods with high spatial and temporal resolution we can understand both variabilities

within each subnetwork, as well as their dynamic interactions, which may not readily be resolved

by imaging techniques. Our results revealed that brain subnetworks, identified by their network

synchronization, exhibit lead-lag interactions during behavioral events that involved information

routing. Specifically, cue-subnetworks lagged target-subnetworks during target presentation when

spatial cues provided relevant information about target locations (**Fig. 4**). These observations are

particularly unique as they bridge the gap between studies focusing on local and cross-regional

mechanisms of fast information routing (*7*, *26*, *43*, *46*, *48*, *58*, *67*) and those examining

communication in large-scale brain networks beyond pairwise interactions (*17*, *68*, *69*).

Lastly, another key and rare finding of our study is the coupling dynamics of HFAbs and local

low-frequency activities. We found that the phase-locking of HFAbs to low-frequency dynamics

can occur with random external input patterns in the modeling, and this is a widespread

phenomenon in iEEG signals, consistent with previous studies (*22*, *70*, *71*). However, our findings

further suggest that HFAbs transiently decouple from the phase of local low-frequency activities

in response to cue and target events (**Fig. 2**). Our modeling supports this observation, showing that

transient inputs, whether directly fed into a network (e.g., a visually evoked response from

subcortical structures) or indirectly through a feedforward structure, can interrupt the default state

coupling of HFAbs to low frequencies (**Fig. 5I**). Network interferences, characterized by brief and

strong responses, can temporarily induce local cross-frequency decoupling possibly by increasing

response heterogeneity and disrupting the timing of inhibitory and excitatory neuron activation

(*72*, *73*). This desynchronization can facilitate cognitive processes like selective attention,

perception, and memory retrieval by enriching represented information and suppressing internally

regulated activity states (*72*, *74–78*). We observed that HFAb decoupling from low-frequency

dynamics, accompanied by increased HFAb density following cue and target events, resembling a

desynchronized up-state in cortical networks. Importantly, this decoupling was more pronounced

in correct trials during both cue and target processing, further supporting the role of

desynchronization in facilitating accurate cognitive performance (**Fig. 2H**). Future studies need to

investigate the origins of these coupling dynamics, their implications in other cognitive domains,

and their causal contributions to local and brain-wide information processing.

In summary, our study provides a systematic approach to understanding large-scale cortical

communications, showing that HFAbs act as fast communication units in the brain, supporting

long-range information processing, facilitating attentional information routing, and identifying

distinct and functionally specialized brain subnetworks.

Acknowledgments

This work was supported by the C.V. Starr Fellowship (KBB), German Research Foundation,

Emmy Noether Program (DFG HE8329/2-1; RFH), NINDS (R01NS021135; RTK), and NEI

(2R01EY017699; SK), NIMH (2R01MH064043, P50MH109429; SK). The funders had no role

in study design, data collection and analysis, the decision to publish, or the preparation of this

manuscript.

455
456
Author Contributions

Conceptualization, K.B.B. and S.K.; Methodology, K.B.B.; Formal Analysis, K.B.B.;

Investigation, R.F.H., and J.J.L.; Software, R.F.H. and I.C.F.; Visualization, K.B.B; Writing –

Original Draft, K.B.B.; Writing – Review & Editing, K.B.B., R.F.H., N.B., R.T.K., and S.K..;

Supervision, S.K.

Data and code accessibility

Custom programming code for analysis and modeling is available upon publication and will be

deposited in an online directory.

Competing Interest: The authors declare no competing interests.

470
Fig. 1. HFAb activation profile predicts behavioral outcome on a trial-by-trial basis. (A) A
schematic of experiment 1 task outline. Two bars appear on the screen. A transient cue informs
the subject of the most probable location of an upcoming target. A target at perceptual threshold
contrast change appears at a cued or infrequently at an equidistant uncued location after a variable
interval. Subjects report the change detection. **(B)** An example trial of detected transient high-
frequency bursts (HFAbs, shown in red) around the cue onset. The brain shows localization of
electrodes across all subjects. **(C)** Normalized HFAb density profiles for individual subjects and
averaged across participants aligned to the cue (left) and target onset (right). **(D)** Correlation of
HFAb responses evoked by cue and target onset (each point represents one electrode, each line
represents regression for one subject, the orange line shows regression across all electrodes). **(E,
F)** Group average heatmap 3D rendering of HFAb responses to **(E)** cues and **(F)** targets. **(G)**
Activation profiles of HFAbs grouped by trial outcome. The dashed lines indicate the significance
level (P<0.05, permutation test). **(H)** An individual subject example of HFAb responses to cue and

target within 500 ms and 700 ms after the cue and target onset, respectively, for correct and
incorrect trials (black circles indicate electrode locations). (**I**) HFAb responses to cues (left) and
targets (right) grouped by trial outcome. (**J**) HFAb responses to targets for valid and invalid cue
conditions. (**K**) HFAb responses to cues ipsi- and contra-lateral to electrodes. (Each line represents
one subject, with thick lines showing group average, horizontal lines showing significant
differences between groups, and asterisks denoting non-zero responses). (**L**) Classifier accuracy
in predicting trial outcome based on HFAb density around the cue onset for cue responsive
electrodes and cue unresponsive electrodes (error bars indicate standard error of the mean across
all realizations and cross validations). Red lines indicate significantly higher prediction accuracy
than baseline and chance levels ($P<0.05$, binomial test, FDR corrected for dependent samples).
494
495

496
Fig. 2. HFAbs dynamically phase lock to low frequency LFPs and decouple transiently after
cue and target onsets. (**A**) HFAb-triggered LFP averages across all electrodes for individual
subjects (gray lines) and across all individuals (purple line). (**B**) HFAb-triggered LFP spectrum
(top) and phase locking values (PLV, bottom) across all electrodes (each line represents one
subject). (**C**) PLVs for individual electrodes (each row). Black circles indicate the maximum peak
of PLV for each electrode, and white dashed lines delineate results from individual subjects. (**D**)
Example of phase-frequency distribution of HFAbs for an electrode with theta phase locking (top)
and beta phase locking (bottom). (**E**) Time-resolved PLV analysis averaged across all subjects
aligned to cue (left) and target (right) onsets. (**F**) The proportion of time points where phase locking
dropped significantly below the baseline averaged across all electrodes (P<0.05, random
permutation test; thick lines indicate segments significantly different from the chance level,
P<0.05, binomial test, FDR corrected for dependent samples; see **Methods**). (**G**) An individual
brain heatmap example showing coupling ratio between HFAb and low frequency (4-25 Hz) LFP
after cue (top) and target onsets (bottom) for correct and incorrect trials. (**H**) Similar to F,
proportion of time points with significant decoupling from low frequency LFPs (4-25 Hz) after
cue and target onsets for correct and incorrect trials. (**I**) Average cue-target coupling ratios across
electrodes for individual subjects. Each line shows the median ± standard error, thick lines indicate
significant non-zero coupling ratios (P<0.05, Wilcoxon test). (**J**) Correlation between burst density
within 500 ms after the target onset and coupling ratios for cue (purple) and target (green)
processing. Circles indicate individual subjects, and filled circles indicate significant correlations.
The horizontal line indicates significant differences between the two conditions (P<0.001,
Wilcoxon test).

519
Fig. 3. Long-range coordination of HFAbs identifies functionally specialized subnetworks.
520
(A) High frequency coordination was measured by calculating HFAb-triggered HFB between
electrode pairs for HFAbs outside of cue/delay and target/response periods (top shows an electrode
pair example, and bottom shows HFAb-triggered HFB averaged over all electrode pairs for
individual subjects). **(B)** Normalized PSD over HFAb-triggered HFB for all electrodes (each
electrode relative to the rest of the network; white dashed lines are the border between subjects).
(C) HFAb-triggered HFB for electrodes distanced in four quantiles, from green (short) to red
(long). **(D)** The first two principal components of HFAb-triggered HFB for individual subjects.
(E) A pair-wise grouping probability matrix for an individual example with an optimal number of
eight clusters. Each element indicates the probability of two electrodes being grouped together
across all random samples (white lines indicate cluster borders, clusters are ordered from top to
bottom by their stability, see **Methods**). **(F)** Cluster topography for the K = 8 number of clusters
for the individual subject shown in E. **(G)** Normalized burst densities around cue and target onsets
averaged across electrodes within each cluster for the same subject as in E and F. (shaded error
bars indicate the standard error of the mean; significant activations are indicated by thicker lines,
$P<0.05$, Wilcoxon test). **(H)** Topographical organization of electrodes in clusters that show
significant activation in response to cues and targets across all subjects (i.e., "cue-subnetworks"
and "target-subnetworks"). **(I)** Classifier accuracy based on HFAb density around cue onset for

cue-subnetworks (purple) and target-subnetworks (green). Error bars indicate the standard error of
the mean across all realizations and cross-validations. Red lines indicate significantly higher
prediction accuracy than the baseline and chance level (binomial test, P<0.05, FDR corrected for
dependent samples).

543
Fig. 4. Cue-subnetworks precede target-subnetworks during target processing. (A) An
individual visualization example of the lead-lag patterns between electrodes in cue and target-
subnetworks (across all cluster numbers) during the target processing period. The lead and lag
temporal patterns are shown in a color gradient from red to blue. **(B)** Example of HFAb-triggered
HFB for HFAbs during target processing, when HFB is measured in target-subnetworks relative
to HFAbs in cue-subnetworks (purple) vs HFB is measured in cue-subnetworks relative to HFAbs
in target-subnetworks (green). Red and blue shades denote significant difference between
directions denoting a cue lead and a target lag respectively (Wilcoxon rank-sum test, P<0.05) **(C)**
Group-level average of HFAb-triggered HFB, similar to B. **(D)** HFAb-triggered HFB median peak
for electrode pairs corresponding to the same groups as in C (the Wilcoxon test was used to
determine whether there was a difference in the distribution of peak positions between the two
directions. Each row indicates the results for one individual. Error bars represent the standard error
of the mean). **(E)** Heatmaps of normalized DMI values averaged across all members of the cue
and target-subnetworks. The left panel shows DMI when the target-subnetwork is fixed, and the
cue-subnetwork is shifted. The right panel shows DMI when the cue-subnetwork is fixed, and the
target-subnetwork is shifted. The white line indicates the time-lag distribution of DMI peaks across
electrode pairs. The cross signs indicate the median of DMI peak time-lags ± standard error. **(F)**
Group average of DMI across all individuals, similar to E. **(G)** Comparisons of DMI peak
distributions between the two opposite directions (green for shifting the cue-subnetwork and
purple for shifting the target-subnetwork) for trials with valid (right) and invalid cues (left). Each
individual is shown by a cross sign, which refers to the median ± standard error for DMI peaks
across all electrode pairs between the two subnetworks.

567
Fig. 5. Computational modeling of iEEG signals. (A) Two interconnected cubic networks of
point source spiking neurons were simulated (bottom panels show raster plots for neurons in each
network). A recording disk was implemented on top of each network to measure neural activity.
(**B**) Example of simulated iEEG signals. (**C**) The power spectral density (PSD) of iEEG signals.
(**D**) A single trial example of spike density and raster (top) detected HFAbs (middle, HFAbs are
shown in red), and HFAb density (bottom). (**E**) HFAb density and spike density show linearly
increasing correlations as input coherence increases. (**F**) HFAb-triggered LFP. (**G**) Cycle-balanced
power spectrum triggered around HFAbs when the input to the network has high (red) or low
coherence (cyan). (**H**) HFAb phase distribution at different frequencies. (**I**) The PLV of HFAbs to
low frequencies after a transient input (thick black line) is fed to the network. (**J,K**) HFAb-
triggered HFB between (**J**) reciprocally connected and (**K**) directional feedforward networks
(connections from N1 to N2, see **Fig. S12E** for further details) with varying connectivity strengths
(purple and green traces show HFAb-triggered HFB when bursts are extracted from N1 and N2,
respectively). (**L**) HFAb-triggered HFB between disconnected networks with shared rhythmic
inputs. (**M**) PSD over the HFAb-triggered HFB for rhythmic inputs of varying frequencies.

References

1.   Y. B. Saalmann, I. N. Pigarev, T. R. Vidyasagar, Neural Mechanisms of Visual Attention:
How Top-Down Feedback Highlights Relevant Locations. *Science* **316**, 1612–1615
(2007).

2.   P. Fries, Neuronal Gamma-Band Synchronization as a Fundamental Process in Cortical
Computation. *Annual Review of Neuroscience* **32**, 209–224 (2009).

3.   G. G. Gregoriou, S. J. Gotts, H. Zhou, R. Desimone, High-Frequency, Long-Range
Coupling Between Prefrontal and Visual Cortex During Attention. *Science* **324**, 1207–
1210 (2009).

4.   S. M. Szczepanski, N. E. Crone, R. A. Kuperman, K. I. Auguste, J. Parvizi, R. T. Knight,
Dynamic Changes in Phase-Amplitude Coupling Facilitate Spatial Attention Control in
Fronto-Parietal Cortex. *PLOS Biology* **12**, e1001936 (2014).

5.   T. J. Buschman, S. Kastner, From Behavior to Neural Dynamics: An Integrated Theory of
Attention. *Neuron* **88**, 127–144 (2015).

6.   I. C. Fiebelkorn, S. Kastner, Functional Specialization in the Attention Network. *Annu Rev*
Psychol **71**, 221–249 (2020).

7.   K. Banaie Boroujeni, T. Womelsdorf, Routing states transition during oscillatory bursts
and attentional selection. *Neuron*, doi: 10.1016/j.neuron.2023.06.012 (2023).

8.   T. Womelsdorf, P. Fries, Neuronal coherence during selective attentional processing and
sensory–motor integration. *Journal of Physiology-Paris* **100**, 182–193 (2006).

9.   S. M. Doesburg, A. B. Roggeveen, K. Kitajo, L. M. Ward, Large-Scale Gamma-Band
Phase Synchronization and Selective Attention. *Cerebral Cortex* **18**, 386–396 (2008).

10.   P. Lakatos, G. Karmos, A. D. Mehta, I. Ulbert, C. E. Schroeder, Entrainment of Neuronal
Oscillations as a Mechanism of Attentional Selection. *Science* **320**, 110–113 (2008).

11.   M. Siegel, T. H. Donner, R. Oostenveld, P. Fries, A. K. Engel, Neuronal Synchronization
along the Dorsal Visual Pathway Reflects the Focus of Spatial Attention. *Neuron* **60**, 709–
719 (2008).

12.   J. Hirvonen, S. Monto, S. H. Wang, J. M. Palva, S. Palva, Dynamic large-scale network
synchronization from perception to action. *Network Neuroscience* **2**, 442–463 (2018).

13.   J. Fan, B. D. McCandliss, J. Fossella, J. I. Flombaum, M. I. Posner, The activation of
attentional networks. *NeuroImage* **26**, 471–479 (2005).

14.   R. T. Canolty, K. Ganguly, S. W. Kennerley, C. F. Cadieu, K. Koepsell, J. D. Wallis, J. M.
Carmena, Oscillatory phase coupling coordinates anatomically dispersed functional cell
assemblies. *PNAS* **107**, 17356–17361 (2010).

15.   U. Braun, A. Schäfer, H. Walter, S. Erk, N. Romanczuk-Seiferth, L. Haddad, J. I.
618   Schweiger, O. Grimm, A. Heinz, H. Tost, A. Meyer-Lindenberg, D. S. Bassett, Dynamic
619   reconfiguration of frontal brain networks during executive cognition in humans.
620   *Proceedings of the National Academy of Sciences* **112**, 11678–11683 (2015).

621   16.   D. S. Bassett, O. Sporns, Network neuroscience. *Nat Neurosci* **20**, 353–364 (2017).

622   17.   A. Avena-Koenigsberger, B. Misic, O. Sporns, Communication dynamics in complex
623   brain networks. *Nat Rev Neurosci* **19**, 17–33 (2018).

624   18.   M. J. Brookes, J. R. Hale, J. M. Zumer, C. M. Stevenson, S. T. Francis, G. R. Barnes, J. P.
625   Owen, P. G. Morris, S. S. Nagarajan, Measuring functional connectivity using MEG:
626   Methodology and comparison with fcMRI. *NeuroImage* **56**, 1082–1104 (2011).

627   19.   R. Hari, L. Parkkonen, The brain timewise: how timing shapes and supports brain
628   function. *Philosophical Transactions of the Royal Society B: Biological Sciences* **370**,
629   20140170 (2015).

630   20.   M. J. Higley, J. A. Cardin, Spatiotemporal dynamics in large-scale cortical networks.
631   *Current Opinion in Neurobiology* **77**, 102627 (2022).

632   21.   O. Jensen, L. L. Colgin, Cross-frequency coupling between neuronal oscillations. *Trends*
633   *in Cognitive Sciences* **11**, 267–269 (2007).

634   22.   R. T. Canolty, R. T. Knight, The functional role of cross-frequency coupling. *Trends in*
635   *Cognitive Sciences* **14**, 506–515 (2010).

636   23.   H. Jiang, A. Bahramisharif, M. A. J. van Gerven, O. Jensen, Measuring directionality
637   between neuronal oscillations of different frequencies. *NeuroImage* **118**, 359–367 (2015).

638   24.   B. Voytek, A. S. Kayser, D. Badre, D. Fegen, E. F. Chang, N. E. Crone, J. Parvizi, R. T.
639   Knight, M. D'Esposito, Oscillatory dynamics coordinating human frontal networks in
640   support of goal maintenance. *Nat Neurosci* **18**, 1318–1324 (2015).

641   25.   M. Bonnefond, S. Kastner, O. Jensen, Communication between Brain Areas Based on
642   Nested Oscillations. *eNeuro* **4**, ENEURO.0153-16.2017 (2017).

643   26.   J. W. Y. Kam, J. J. Lin, A.-K. Solbakk, T. Endestad, P. G. Larsson, R. T. Knight, Default
644   network and frontoparietal control network theta connectivity supports internal attention.
645   *Nat Hum Behav* **3**, 1263–1270 (2019).

646   27.   E. L. Johnson, J. J. Lin, D. King-Stephens, P. B. Weber, K. D. Laxer, I. Saez, F. Girgis, M.
647   D'Esposito, R. T. Knight, D. Badre, A rapid theta network mechanism for flexible
648   information encoding. *Nat Commun* **14**, 2872 (2023).

649   28.   J.-P. Lachaux, N. Axmacher, F. Mormann, E. Halgren, N. E. Crone, High-frequency
650   neural activity and human cognition: past, present and possible future of intracranial EEG
651   research. *Prog Neurobiol* **98**, 279–301 (2012).

652   29.   J. Parvizi, S. Kastner, Human Intracranial EEG: Promises and Limitations. *Nat Neurosci*
653   **21**, 474–483 (2018).

654   30.   S. Ray, J. H. R. Maunsell, Different Origins of Gamma Rhythm and High-Gamma
655   Activity in Macaque Visual Cortex. *PLOS Biology* **9**, e1000610 (2011).

656   31.   K. Banaie Boroujeni, P. Tiesinga, T. Womelsdorf, Adaptive spike-artifact removal from
657   local field potentials uncovers prominent beta and gamma band neuronal synchronization.
658   *Journal of Neuroscience Methods* **330**, 108485 (2020).

659   32.   M. Leszczyński, A. Barczak, Y. Kajikawa, I. Ulbert, A. Y. Falchier, I. Tal, S. Haegens, L.
660   Melloni, R. T. Knight, C. E. Schroeder, Dissociation of broadband high-frequency activity
661   and neuronal firing in the neocortex. *Science Advances* **6**, eabb0977 (2020).

662   33.   T. A. Zolnik, A. Bronec, A. Ross, M. Staab, R. N. S. Sachdev, Z. Molnár, B. J. Eickholt,
663   M. E. Larkum, Layer 6b controls brain state via apical dendrites and the higher-order
664   thalamocortical system. *Neuron* **112**, 805-820.e4 (2024).

665   34.   S. Ray, E. Niebur, S. S. Hsiao, A. Sinai, N. E. Crone, High-frequency gamma activity (80-
666   150 Hz) is increased in human cortex during selective attention. *Clin Neurophysiol* **119**,
667   116–133 (2008).

668   35.   M. Haller, J. Case, N. E. Crone, E. F. Chang, D. King-Stephens, K. D. Laxer, P. B. Weber,
669   J. Parvizi, R. T. Knight, A. Y. Shestyuk, Persistent neuronal activity in human prefrontal
670   cortex links perception and action. *Nat Hum Behav* **2**, 80–91 (2018).

671   36.   A. B. Martin, X. Yang, Y. B. Saalmann, L. Wang, A. Shestyuk, J. J. Lin, J. Parvizi, R. T.
672   Knight, S. Kastner, Temporal Dynamics and Response Modulation across the Human
673   Visual System in a Spatial Attention Task: An ECoG Study. *J. Neurosci.* **39**, 333–352
674   (2019).

675   37.   C. Jacques, J. Jonas, S. Colnat-Coulbois, L. Maillard, B. Rossion, Low and high frequency
676   intracranial neural signals match in the human associative cortex. *eLife* **11**, e76544 (2022).

677   38.   G. Arnulfo, S. H. Wang, V. Myrov, B. Toselli, J. Hirvonen, M. M. Fato, L. Nobili, F.
678   Cardinale, A. Rubino, A. Zhigalov, S. Palva, J. M. Palva, Long-range phase
679   synchronization of high-frequency oscillations in human cortex. *Nat Commun* **11**, 5363
680   (2020).

681   39.   N. Williams, S. H. Wang, G. Arnulfo, L. Nobili, S. Palva, J. M. Palva, Modules in
682   connectomes of phase-synchronization comprise anatomically contiguous, functionally
683   related regions. *NeuroImage* **272**, 120036 (2023).

684   40.   J. C. Garrett, I. A. Verzhbinsky, E. Kaestner, C. Carlson, W. K. Doyle, O. Devinsky, T.
685   Thesen, E. Halgren, Binding of cortical functional modules by synchronous high-
686   frequency oscillations. *Nat Hum Behav*, 1–15 (2024).

41. R. F. Helfrich, I. C. Fiebelkorn, S. M. Szczepanski, J. J. Lin, J. Parvizi, R. T. Knight, S.
Kastner, Neural Mechanisms of Sustained Attention Are Rhythmic. *Neuron* **99**, 854-
865.e5 (2018).

42. E. M. Izhikevich, Which model to use for cortical spiking neurons? *IEEE Transactions on*
Neural Networks **15**, 1063–1070 (2004).

43. A. Palmigiano, T. Geisel, F. Wolf, D. Battaglia, Flexible information routing by transient
synchrony. *Nat Neurosci* **20**, 1014–1022 (2017).

44. M. Lundqvist, E. K. Miller, J. Nordmark, J. Liljefors, P. Herman, Beta: bursts of
cognition. *Trends in Cognitive Sciences* **28**, 662–676 (2024).

45. L. L. Colgin, T. Denninger, M. Fyhn, T. Hafting, T. Bonnevie, O. Jensen, M.-B. Moser, E.
I. Moser, Frequency of gamma oscillations routes flow of information in the hippocampus.
Nature **462**, 353–357 (2009).

46. T. Akam, D. M. Kullmann, Oscillations and Filtering Networks Support Flexible Routing
of Information. *Neuron* **67**, 308–320 (2010).

47. J. Yamamoto, J. Suh, D. Takeuchi, S. Tonegawa, Successful Execution of Working
Memory Linked to Synchronized High-Frequency Gamma Oscillations. *Cell* **157**, 845–
857 (2014).

48. G. Hahn, A. Ponce-Alvarez, G. Deco, A. Aertsen, A. Kumar, Portraits of communication
in neuronal networks. *Nat Rev Neurosci* **20**, 117–127 (2019).

49. A. Fernandez-Ruiz, A. Sirota, V. Lopes-dos-Santos, D. Dupret, Over and above
frequency: Gamma oscillations as units of neural circuit operations. *Neuron* **111**, 936–953
(2023).

50. P. König, A. K. Engel, W. Singer, Relation between oscillatory activity and long-range
synchronization in cat visual cortex. *Proceedings of the National Academy of Sciences* **92**,
290–294 (1995).

51. P. R. Roelfsema, A. K. Engel, P. König, W. Singer, Visuomotor integration is associated
with zero time-lag synchronization among cortical areas. *Nature* **385**, 157–161 (1997).

52. P. J. Uhlhaas, G. Pipa, B. Lima, L. Melloni, S. Neuenschwander, D. Nikolić, W. Singer,
Neural Synchrony in Cortical Networks: History, Concept and Current Status. *Front*
Integr Neurosci **3**, 17 (2009).

53. L. L. Gollo, C. Mirasso, A. E. P. Villa, Dynamic control for synchronization of separated
cortical areas through thalamic relay. *NeuroImage* **52**, 947–955 (2010).

54. L. L. Gollo, C. R. Mirasso, M. Atienza, M. Crespo-Garcia, J. L. Cantero, Theta Band
Zero-Lag Long-Range Cortical Synchronization via Hippocampal Dynamical Relaying.
PLOS ONE **6**, e17756 (2011).

55. Y. B. Saalmann, S. Kastner, Cognitive and Perceptual Functions of the Visual Thalamus.
Neuron **71**, 209–223 (2011).

56. N. K. Logothetis, O. Eschenko, Y. Murayama, M. Augath, T. Steudel, H. C. Evrard, M.
Besserve, A. Oeltermann, Hippocampal–cortical interaction during periods of subcortical
silence. *Nature* **491**, 547–553 (2012).

57. S. J. Sara, S. Bouret, Orienting and Reorienting: The Locus Coeruleus Mediates Cognition
through Arousal. *Neuron* **76**, 130–141 (2012).

58. Y. B. Saalmann, M. A. Pinsk, L. Wang, X. Li, S. Kastner, The Pulvinar Regulates
Information Transmission Between Cortical Areas Based on Attention Demands. *Science*,
doi: 10.1126/science.1223082 (2012).

59. S. Vijayan, N. J. Kopell, Thalamic model of awake alpha oscillations and implications for
stimulus processing. *Proceedings of the National Academy of Sciences* **109**, 18553–18558
(2012).

60. I. C. Fiebelkorn, M. A. Pinsk, S. Kastner, The mediodorsal pulvinar coordinates the
macaque fronto-parietal network during rhythmic spatial attention. *Nat Commun* **10**, 215
(2019).

61. S. Kastner, I. C. Fiebelkorn, M. K. Eradath, Dynamic pulvino-cortical interactions in the
primate attention network. *Current Opinion in Neurobiology* **65**, 10–19 (2020).

62. T. Menara, Y. Qin, D. S. Bassett, F. Pasqualetti, Relay Interactions Enable Remote
Synchronization in Networks of Phase Oscillators. *IEEE Control Systems Letters* **6**, 500–
505 (2022).

63. L. L. Gollo, C. Mirasso, O. Sporns, M. Breakspear, Mechanisms of Zero-Lag
Synchronization in Cortical Motifs. *PLOS Computational Biology* **10**, e1003548 (2014).

64. T. A. Zolnik, J. Ledderose, M. Toumazou, T. Trimbuch, T. Oram, C. Rosenmund, B. J.
Eickholt, R. N. S. Sachdev, M. E. Larkum, Layer 6b Is Driven by Intracortical Long-
Range Projection Neurons. *Cell Reports* **30**, 3492-3505.e5 (2020).

65. S. M. Smith, P. T. Fox, K. L. Miller, D. C. Glahn, P. M. Fox, C. E. Mackay, N. Filippini,
K. E. Watkins, R. Toro, A. R. Laird, C. F. Beckmann, Correspondence of the brain's
functional architecture during activation and rest. *Proceedings of the National Academy of*
Sciences **106**, 13040–13045 (2009).

66. A. R. Laird, S. B. Eickhoff, C. Rottschy, D. Bzdok, K. L. Ray, P. T. Fox, Networks of task
co-activations. *NeuroImage* **80**, 505–514 (2013).

67. X.-J. Wang, G. R. Yang, A disinhibitory circuit motif and flexible information routing in
the brain. *Current Opinion in Neurobiology* **49**, 75–83 (2018).

68.   F. Battiston, G. Cencetti, I. Iacopini, V. Latora, M. Lucas, A. Patania, J.-G. Young, G. Petri, Networks beyond pairwise interactions: Structure and dynamics. *Physics Reports* **874**, 1–92 (2020).

69.   S. L. Bressler, V. Menon, Large-scale brain networks in cognition: emerging methods and principles. *Trends in Cognitive Sciences* **14**, 277–290 (2010).

70.   O. Jensen, B. Gips, T. O. Bergmann, M. Bonnefond, Temporal coding organized by coupled alpha and gamma oscillations prioritize visual processing. *Trends in Neurosciences* **37**, 357–369 (2014).

71.   J. I. Chapeton, R. Haque, J. H. Wittig, S. K. Inati, K. A. Zaghloul, Large-Scale Communication in the Human Brain Is Rhythmically Modulated through Alpha Coherence. *Current Biology* **29**, 2801-2811.e5 (2019).

72.   K. D. Harris, A. Thiele, Cortical state and attention. *Nat Rev Neurosci* **12**, 509–523 (2011).

73.   R. Gast, S. A. Solla, A. Kennedy, Neural heterogeneity controls computations in spiking neural networks. *Proceedings of the National Academy of Sciences* **121**, e2311885121 (2024).

74.   S. Hanslmayr, T. Staudigl, M.-C. Fellner, Oscillatory power decreases and long-term memory: the information via desynchronization hypothesis. *Front. Hum. Neurosci.* **6** (2012).

75.   S. Hanslmayr, B. P. Staresina, H. Bowman, Oscillations and Episodic Memory: Addressing the Synchronization/Desynchronization Conundrum. *Trends in Neurosciences* **39**, 16–25 (2016).

76.   C. B. Beaman, S. L. Eagleman, V. Dragoi, Sensory coding accuracy and perceptual performance are improved during the desynchronized cortical state. *Nat Commun* **8**, 1308 (2017).

77.   A. Nandy, J. J. Nassi, M. P. Jadi, J. Reynolds, Optogenetically induced low-frequency correlations impair perception. *eLife* **8**, e35123 (2019).

78.   N. Weisz, N. G. Kraft, G. Demarchi, Auditory cortical alpha/beta desynchronization prioritizes the representation of memory items during a retention period. *eLife* **9**, e55508 (2020).

79.   R. Egly, J. Driver, R. D. Rafal, Shifting visual attention between objects and locations: Evidence from normal and parietal lesion subjects. *Journal of Experimental Psychology: General* **123**, 161–177 (1994).

80.   I. C. Fiebelkorn, Y. B. Saalmann, S. Kastner, Rhythmic Sampling within and between Objects despite Sustained Attention at a Cued Location. *Current Biology* **23**, 2553–2558 (2013).

81. B. Boashash, "Estimating and Interpreting the Instantaneous Frequency of a Signal – Part
1: Fundamentals" in *Proceedings of the Ieee* (1992), pp. 520–538.

82. M. R. H. Hill, I. Fried, C. Koch, Quantification and classification of neuronal responses in
kernel-smoothed peristimulus time histograms. *Journal of Neurophysiology* **113**, 1260–
1274 (2015).

83. O. J. Dunn, Multiple Comparisons among Means. *Journal of the American Statistical
Association* **56**, 52–64 (1961).

84. Y. Benjamini, D. Yekutieli, False Discovery Rate–Adjusted Multiple Confidence Intervals
for Selected Parameters. *Journal of the American Statistical Association* **100**, 71–81
(2005).

85. N. I. Fisher, *Statistical Analysis of Circular Data* (Cambridge University Press, 1995).

86. G. W. Milligan, M. C. Cooper, An examination of procedures for determining the number
of clusters in a data set. *Psychometrika* **50**, 159–179 (1985).

87. A. K. Jain, R. C. Dubes, *Algorithms for Clustering Data* (Prentice-Hall, Inc., USA, 1988).

88. M. Halkidi, Y. Batistakis, M. Vazirgiannis, On Clustering Validation Techniques. *Journal
of Intelligent Information Systems* **17**, 107–145 (2001).

89. L. Vendramin, R. J. G. B. Campello, E. R. Hruschka, Relative clustering validity criteria:
A comparative overview. *Statistical Analysis and Data Mining: The ASA Data Science
Journal* **3**, 209–235 (2010).

90. D. Freedman, P. Diaconis, On the histogram as a density estimator:L2 theory. *Z.
Wahrscheinlichkeitstheorie verw Gebiete* **57**, 453–476 (1981).

91. Y. N. Billeh, B. Cai, S. L. Gratiy, K. Dai, R. Iyer, N. W. Gouwens, R. Abbasi-Asl, X. Jia,
J. H. Siegle, S. R. Olsen, C. Koch, S. Mihalas, A. Arkhipov, Systematic Integration of
Structural and Functional Data into Multi-scale Models of Mouse Primary Visual Cortex.
Neuron **106**, 388-403.e18 (2020).

92. T. C. Potjans, M. Diesmann, The Cell-Type Specific Cortical Microcircuit: Relating
Structure and Activity in a Full-Scale Spiking Network Model. *Cerebral Cortex* **24**, 785–
806 (2014).

93. J. Barral, A. D Reyes, Synaptic scaling rule preserves excitatory–inhibitory balance and
salient neuronal network dynamics. *Nat Neurosci* **19**, 1690–1696 (2016).

94. E. M. Izhikevich, Simple model of spiking neurons. *IEEE Transactions on Neural
Networks* **14**, 1569–1572 (2003).

95.   C. K. Pfeffer, M. Xue, M. He, Z. J. Huang, M. Scanziani, Inhibition of inhibition in visual
cortex: the logic of connections between molecularly distinct interneurons. *Nat Neurosci*
16, 1068–1076 (2013).

96.   M. M. Karnani, J. Jackson, I. Ayzenshtat, J. Tucciarone, K. Manoocheri, W. G. Snider, R.
Yuste, Cooperative Subnetworks of Molecularly Similar Interneurons in Mouse
Neocortex. *Neuron* **90**, 86–100 (2016).

97.   M. Medalla, J. P. Gilman, J.-Y. Wang, J. I. Luebke, Strength and Diversity of Inhibitory
Signaling Differentiates Primate Anterior Cingulate from Lateral Prefrontal Cortex. *J.*
Neurosci. **37**, 4717–4734 (2017).

98.   J. P. Gilman, M. Medalla, J. I. Luebke, Area-Specific Features of Pyramidal Neurons—a
Comparative Study in Mouse and Rhesus Monkey. *Cerebral Cortex* **27**, 2078–2094
(2017).

99.   M. Medalla, J. I. Luebke, Diversity of Glutamatergic Synaptic Strength in Lateral
Prefrontal versus Primary Visual Cortices in the Rhesus Monkey. *J. Neurosci.* **35**, 112–
127 (2015).

100.   J. M. Amatrudo, C. M. Weaver, J. L. Crimins, P. R. Hof, D. L. Rosene, J. I. Luebke,
Influence of Highly Distinctive Structural Properties on the Excitability of Pyramidal
Neurons in Monkey Visual and Prefrontal Cortices. *J. Neurosci.* **32**, 13644–13660 (2012).

101.   A. Roxin, N. Brunel, D. Hansel, G. Mongillo, C. van Vreeswijk, On the Distribution of
Firing Rates in Networks of Cortical Neurons. *J. Neurosci.* **31**, 16217–16226 (2011).

102.   J. Milstein, F. Mormann, I. Fried, C. Koch, Neuronal Shot Noise and Brownian 1/f2
Behavior in the Local Field Potential. *PLOS ONE* **4**, e4338 (2009).

103.   C. Bédard, H. Kröger, A. Destexhe, Modeling Extracellular Field Potentials and the
Frequency-Filtering Properties of Extracellular Space. *Biophysical Journal* **86**, 1829–1842
(2004).

850