1

Both density- and frequency-dependent effects determine plant growth in a dune

heath ecosystem

Christian Damgaard [ID][✉][1], Mathias Emil Kaae[1], Jacob Weiner[2], and Jesper Leth Bak[1]

[1]Department of Ecoscience, Aarhus University, 8000 Aarhus C, Denmark

[2]Department of Plant and Environmental Sciences, University of Copenhagen, Frederiksberg,

Denmark

Correspondence: Christian Damgaard (cfd@ecos.au.dk)

Abstract

We tested the hypothesis that both density- and frequency-dependent interactions play important roles in determining plant growth in a dune heath ecosystem at several levels of available nitrogen. Plant growth was measured using the pin-point method in a five-block experiment with four nitrogen levels. To maximize statistical power, we used only three taxonomic groups: *Calluna vulgaris*, *Avenella flexuosa* (the two most dominant species), and all other vascular plant species together. The results show that both Lotka-Volterra type interspecific competition and frequency-dependency play significant roles in determining the growth of the species in the community. Significant interspecific density-dependent competition was observed in 4 out of the 6 possible cases. Nitrogen addition increased the competitive effect of *C. vulgaris* on the growth of the other species. Both *C. vulgaris* and *A. flexuosa* showed frequency-dependent positive feedback dynamics on growth when they were relatively dominant at the plot scale, and this effect increased with added nitrogen. In plots with added nitrogen, there was a beneficial effect of being relatively rare on the growth of the group of other species. The study highlights the importance of the combined effects of density and frequency dependency in determining plant growth.

keywords: plant-plant interactions; plant-soil feedback; frequency-dependence; unmeasured variables; measurement uncertainty; hierarchical modelling

Introduction

To develop predictive models of ecosystems in changing environments, it is important to understand the mechanisms that govern plant community dynamics and quantify their relative importance. Historically, the most important mechanisms for understanding plant community dynamics are competition for resources, e.g. light, water and soil nutrients that are necessary for plant growth (Goldberg et al. 1990). Species coexistence has primarily been examined with density-dependent Lotka-Volterra type models, where population growth is affected by intra- and interspecific population densities, and investigating the conditions for invasion when a species is rare (Adler et al. 2018; Barabás et al. 2018; Chesson 1994; 2000). This body of theory has been referred to as "modern coexistence theory". But plant communities may also be regulated by frequency-dependent mechanisms, where plant growth is regulated by the frequency of the species within the community. There has been increasing awareness that frequency-dependent mechanisms may play important roles in the dynamics of plant communities (Bever et al. 1997; Chisholm and Fung 2020; Connell et al. 1984; Suding et al. 2024). Such frequency-dependency may be caused by numerous mechanisms, such as species-specific soil-plant interactions that hinder local establishment and growth of conspecific dominant plant species, or because relatively low population sizes of natural enemies of rare species may give them an advantage (Heinen et al. 2020; Mazzoleni et al. 2015a; Mazzoleni et al. 2015b; van der Putten et al. 2013), or dominant plant species may be favored by positive feedback mechanisms, e.g. by resource monopolizing strategies (van der Putten et al. 2013). It is surprising that many studies demonstrate the importance of frequency dependence for coexistence, but it is completely ignored in "modern coexistence theory". It has also been argued that higher-order

interactions among species are important (Singh and Baruah 2020), and that intraspecific

genetic variation may play a role in the outcome of species interactions (Ehlers et al. 2016).

Generally, the relative importance of different mechanisms in regulating plant community

dynamics in different ecosystems is still an open question (e.g. De Long et al. 2023).

Dune heathland ecosystems are natural or semi-natural habitats situated on leached,

nutrient-poor, sandy soils. The vegetation is adapted to withstand the harsh, windy

environment and is dominated by dwarf shrubs, e.g. *Calluna vulgaris* (heather), *Empetrum*

nigrum (crowberry), *Vaccinium uliginosum* (bog bilberry), and graminoids, especially

Avenella flexuosa (wavy hair-grass), *Carex arenaria* (sand sedge), *Corynephorus canescens*

(grey hair-grass), and *Festuca ovina (*sheep fescue). Bryophytes and lichens are also

common. Dwarf shrubs and grasses often coexist in a dynamic patchy spatial configuration.

The cyclical successional processes leading to such a patchy distribution of dwarf shrubs and

grasses was described by Watt (1947), a key feature of which is the life cycle of *Calluna*

vulgaris plants, which senescence when they are 30–40 years old (Gimingham 1988; Watt

1955).

One of the essential abiotic factors that regulate plant growth and community dynamics

is the level of available nitrogen (Craine 2009). Nitrogen deposition from the combustion of

fossil fuels and animal husbandry is known to play an important role in current heathland

ecosystems (Alonso et al. 2001; Bähring et al. 2017; Damgaard 2019a; 2019b; Jones and

Power 2012; Southon et al. 2013). If nitrogen input is sufficiently high, especially after the

vegetation is opened by burning or beetle infestation, grass species may outcompete ericoid

dwarf shrubs and the dwarf shrub vegetation may be replaced by acid grassland vegetation

or, in the absence of management, encroachment by trees (Alonso et al. 2001; Heil and

Bobbink 1993; van Voorn et al. 2016). Under increased nitrogen availability, *A. flexuosa*

outcompetes *C. vulgaris*, which is adapted to nitrogen-limited environments in which its

slowly decaying litter is decomposed by ericoid mycorrhiza and the nitrogen in its litter is

absorbed by its roots (Diemont and Heil 1984; Johansson 2001; Smith and Read 2008; Talbot

et al. 2008). The positive feedback between heather and ericoid mycorrhiza may lead to

frequency-dependency in nitrogen limited environments, where the competitive growth of

C. vulgaris is accelerated when it is dominant (Aerts et al. 1990).

The role of competition in determining plant growth and therefore community dynamics

is investigated empirically by modelling the observed net negative plant-plant interactions

("competition in the broad sense" sensu Weiner 1993) using Lotka-Volterra-type

competition models (e.g Adler et al. 2018; Damgaard 1998; 2005; Law and Dieckmann 2000).

Although the use of net effects in the modelling of species interaction has been criticized, it

is still the best currently available tool we have for describing inter- and intra-specific

interactions in communities. Hypothesizing that frequency dependence plays an important

role in plant community dynamics, Damgaard and Weiner (2021) added frequency

dependent competition to a Lotka-Volterra-type competition model. Their model was fitted

to empirical growth data using a hierarchical Bayesian framework, in which the possible role

of unmeasured variables that influence individual plant performance and measurement

errors was accounted for (Fig. 1). The inclusion of frequency dependence improved the

model fit significantly.

Here, we test the hypotheses that (1) both density- and frequency-dependent

competition play important roles in determining plant growth in a dune heath ecosystem,

and (2) that the relative importance of these different mechanisms depends on the amount

of available nitrogen. Growth models in which the effect of density- and frequency-

dependent competition are parameterized and fit to pin-point cover data sampled in both

early and late summer over a three-year period. The competitive growth of *C. vulgaris* and *A.*

flexuosa, and all other higher plants grouped together are then modeled as a function of the

cover in early summer (Damgaard and Weiner 2021).

Material and Methods

Competitive growth data

At a dune heath site, Vust Hede (Fig. 2), with relatively low background nitrogen deposition

(9 kg N ha^{-1} year^{-1}), five experimental blocks were established in 2018 (Kaae et al. 2024).

Each block had four treatment plots with added nitrogen (0, 5, 10, 25 kg N ha^{-1} year^{-1}). At the

sites, there were natural populations of roe deer, red deer and fallow deer in an estimated

combined density of 9.95 ha^{-1} day^{-1} (Appendix A), and herbivory by the deer population was

shown to affect plant species composition (Kaae et al. 2024). In the summer of 2020,

approximately 70-80% of the heather plants were infested by the heather beetle (*Lochmeae*

suturalis)*,* leading to discoloring of leaves and reduction in growth (Kaae et al. 2024).

To study the species interactions on plant growth, a 0.5 m x 0.5 m quadrat was

permanently outlined within each plot. In the quadrat, plant abundance was measured non-

destructively by the "pin-point" (also called "point intercept") method (Kent and Coker 1992)

using a pin-point frame with the same dimension as the quadrat and with 25 pin-positions

regularly placed at distances of 10 cm. At each position, a pin with a diameter of 0.5 mm was

passed vertically through the vegetation, and the number of contacts to each species was

recorded. The sampling was performed in early (June) and late summer (August to October)

in the three years 2019, 2020, and 2021 (Kaae et al. 2024).

The pin-point method provides estimates of two important plant ecological variables:

plant cover and vertical density, which are well suited for studying species interactions in

natural and semi-natural herbaceous perennial plant communities, where it is difficult to

distinguish individual genets (Damgaard 2011; Damgaard et al. 2009). The cover of a specific

plant species is defined as the proportion of pins in the grid that touch the species, thus,

plant cover measures the cover of the plant species when it is projected onto the two-

dimensional ground surface. The vertical density is defined as the number of times a pin

touches a specific species, and this measure is correlated with plant biomass (Jonasson 1983;

1988; Kaae et al. 2024).

In the subsequent analysis, the observed plant taxa were grouped into *C. vulgaris*, *A.*

flexuosa grass and an aggregated group of all other vascular plant species.

The pinpoint data is available in an electronic supplement (Appendix B)

Model

The effect of species interactions on plant growth was analyzed in a hierarchical model (Fig.

1c) by modelling how the vertical density of *C. vulgaris*, *A. flexuosa* and the other species at

the end of the growing season depended on the cover at the beginning of the growing

season. The underlying assumption is that the measure of vertical density at the end of the

growing season as a function of the cover at the beginning of the growing season may be

used as a measure of population growth (Damgaard 2011; Damgaard et al. 2009).

We used a modified version of the model in Damgaard and Weiner (2021),

$$f_i(\boldsymbol{x}) = (a_i + an_i\, n)x_i^{b_i}\, Exp\left(\sum_{j=1, j\neq i}^{S} -(c_{ij} + cn_{ij}n)\, x_j^{d_j}\right) Exp\left(-(e_i + en_i\, n)\frac{x_i}{\sum_{j=1}^{S} x_j}\right) \quad (1),$$

where $f_i(\boldsymbol{x})$ is the expected vertical density of species *i* at the end of the growing season,

and x_i is the cover of species *i* at the beginning of the growing season. The first term models

the intraspecific density dependence (i.e. if the species were in a monospecific population).

The second term models the effect of interspecific density dependence (as in the Lotka-

Volterra models), and the third term models the effect of frequency-dependency within the

plant community. Thus, the density-dependent and frequency-dependent effects are

partitioned and estimated in separate terms. More specifically, the parameters a_i, an_i and

b_i model the expected vertical density as a function of cover and nitrogen (n) in the absence

of inter-specific density-dependent and frequency-dependent Effects, while c_{ij}, cn_{ij} and d_j

model the inter-specific interactions of species j on the vertical density of species i. If

$\left(c_{ij} + cn_{ij}\, n\right) > 0$, there is evidence of competitive inter-specific interactions, otherwise

there is evidence of facilitative inter-specific interactions. The parameters e_i and en_i model

the frequency-dependent effect of species i. If $(e_i + en_i\, n) = 0$, then there is no evidence of

frequency-dependence, and the population growth model (1) simplifies to a Lotka-Volterra

competition model. If $(e_i + en_i\, n) > 0$, then rare species have an advantage over dominant

species and, conversely, if $(e_i + en_i\, n) < 0$, then dominant species have an advantage over

rare species. The parameters b_i and d_j model possible additional non-linearity in the effect

of cover on vertical density with a domain in the interval (0.5, 2).

The measurement errors of cover and vertical density have previously been assumed to

be binomially distributed and generalized-Poisson distributed, respectively (Damgaard et al.

2014). However, since we wanted to model the covariance between the early and later

growth stages to account for the possible effect of unmeasured variables (Rinella et al.

2020), both distributions are approximated by standard normal distributions, where (i) the

observed pin-point cover measure, y, in a pin-point frame with n pin-positions and an

expected cover q is transformed to $z_1(y; q) = \frac{y - n\, q}{\sqrt{n\, q(1-q)}}$, and (ii) the observed pin-point

vertical density measure, vd, with an expected value λ and with a species-specific scale

parameter υ_i is transformed to $z_2(vd; \lambda, \nu_i) = \frac{vd - \lambda}{\sqrt{\lambda}\ \nu_i}$. The measurement error of the

standardized cover and vertical density measures is consequently modelled as

$(z_1(y; q), z_2(vd; \lambda, \nu_i)) \sim N_2 \left((0,0), \begin{pmatrix} 1 & \rho_i \\ \rho_i & 1 \end{pmatrix} \right)$ (2),

where ρ_i is a species-specific correlation coefficient in the domain (-1, 1; (Damgaard and

Weiner 2021). If $\rho_i = 0$, then the measurement errors of the standardized cover and vertical

density measures are assumed to be uncorrelated, i.e. the competition model, not additional

unmeasured variables, accounts for the covariance between the early and later growth

stages.

The structural uncertainty of the competitive growth model was assumed to be normally

distributed and modelled as $vd_i \sim N(f_i(\boldsymbol{x}), \sigma_i)$, where σ_i is the species-specific process error

(see Table 1).

Estimation

We treat the three years of competitive growth data from each of the plots as independent

events because i) the within-plots among-year covariation is expected to be minimal, since

we model the plant growth of each year, and ii) considering plot as a random effect is not

compatible with modelling covariance between early and late plant growth.

The joint Bayesian posterior probability distribution of the parameters in the model was

calculated using Markov Chain Monte Carlo (MCMC; Metropolis-Hastings) simulations of

900,000 iterations with a burn-in period of 800,000 iterations and normal candidate

distributions (Carlin and Louis 1996). The prior probability distributions of all parameters and

latent variables were assumed to be uniformly distributed within their specified domains

under the additional constraints that $0 < a_i < 500, -5 < c_{ij} < 5, -5 < e_i < 5, 0.5 < \nu_i <$

8, and $-0.5 < \rho_i < 0.8$, except for σ_i, which was assumed to be inverse gamma distributed

with parameters $(0.1, 0.1)$.

Plots of the deviance and trace plots of parameters were inspected to check the fitting

and mixing properties of the sampling procedure. Residual plots of the vertical density for

each species were inspected to check the fitting properties of the different models. The

statistical inferences were assessed using the calculated credible intervals, i.e. 95%

percentiles of the marginal posterior distribution of the parameters and the probabilities

that the marginal posterior distributions of the parameters were either above or below zero.

All calculations were done using Mathematica (Wolfram 2019). The software code and all

results are given in the electronic supplement (Appendix B).

Prediction of relative vertical density

To investigate the estimated effects of both density- and frequency-dependent competition

on plant growth (vertical density) at different cover values of the three species and nitrogen

levels, the mean relative predicted vertical density was calculated by inserting samples from

the joint posterior distribution of the model parameters into equation 1 and dividing by the

cover of the species, $f_i(\boldsymbol{x})/x_i$.

Results

The MCMC iterations of the plant growth model converged with acceptable mixing

properties, although there was sizeable covariation among parameters (Appendix C). Based

on the residual plots (Fig. S1), the fit of the model was judged to be acceptable. A summary

of the marginal posterior probability distributions of all parameters is shown in Table 2.

The vertical density in the absence of interspecific species interaction was modelled by

the parameters a_i, an_i and b_i. The vertical density was an increasing function of added

nitrogen, only for all the group of other vascular plant species (Table 2, an_3).

The competitive effect of *A. flexuosa* and all other vascular species on within-season

growth of *C. vulgaris* was relatively large without added nitrogen (Table 2, c_{12}, c_{13}), but

there was no significant effect of the nitrogen treatment (Table 2, cn_{12}, cn_{13}). Similarly, the

competitive effect of all other vascular species on *A. flexuosa* was relatively large without

added nitrogen (Table 2, c_{23}), and there was no significant effect of the nitrogen level (Table

2, cn_{23}). The competitive effect of *C. vulgaris* on all other vascular species was significantly

positive without added nitrogen (Table 2, c_{31}), and this effect increased significantly with

the level of added nitrogen (Table 2, cn_{31}). There was no significant competitive effects of *A.*

flexuosa on the growth of all other vascular species (Table 2, c_{32}, cn_{32}), or of *C. vulgaris* on

the growth of *A. flexuosa* (Table 2, c_{21}, cn_{21}).

There were significant frequency-dependent effects. For *C. vulgaris* and *A. flexuosa*, there

were significant positive effects of being relatively dominant at the plot scale (Table 2, Fig. 3,

e_1, e_2), and this effect increased significantly with added nitrogen (Table 2, Fig. 3, en_1, en_2).

On the other hand, all other vascular species were significantly positively affected when

relatively rare in the plots (Table 2, e_3, en_3).

The mean relative predicted vertical density of both *C. vulgaris* and *A. flexuosa* increased

with cover (Fig. 4), indicating that the frequency-dependent positive feedback mechanisms

are important for the plant community dynamics in heathland ecosystems. The predicted

increases were relatively independent of the amount of added nitrogen (results not shown).

For *C. vulgaris* and *A. flexuosa*, there was no significant effect of unmeasured

environmental variables (Table 2, ρ_1, ρ_2) , i.e. the covariance between the early and later

growth stages is assumed to be accounted for by the competition model (eq. 1) and not by

additional unmeasured variables. However, for all other vascular species there was an

indication that unmeasured environmental variables had a significant positive effect on the

observed covariance between the early and later growth stages that was not accounted for

by the species interaction model (Table 2, ρ_3).

Discussion

In the dune heath ecosystem studied, we have presented evidence that both density- and

frequency-dependent interspecific competition play significant roles in determining plant

growth in the plant community studied. Significant interspecific density-dependent

competition was observed in 4 out of the 6 possible cases of three species interactions. We

expected that nitrogen addition would alter the species interactions, but this was only

demonstrated in the effect of *C. vulgaris* on the growth of all other vascular species, where

nitrogen addition increased the competitive effect of *C. vulgaris*. It has been suggested that

C. vulgaris may retain a competitive advantage with increased deposition of nitrogen

because of the vertical structure of its leaves, the mature dense vegetation and the fact that

leaves remain on the plant during winter (Aerts et al. 1990; Aerts and Heil 1993). It is

possible that the effects of nitrogen addition occur only after several years, after which the

level of available nitrogen has reached a new and higher steady-state level. It will be

interesting to follow the effect of nitrogen on plant species interactions in the coming years.

C. vulgaris and *A. flexuosa* showed positive feedback dynamics on growth when they

were relatively dominant at the plot scale, whereas the growth of all other vascular species

was favored when they were relatively rare. The positive feedback dynamics of *C. vulgaris*

could be explained by the resource monopolizing strategy mediated by ericoid mycorrhiza

(Diemont and Heil 1984; Johansson 2001; Smith and Read 2008; Talbot et al. 2008), although

the estimated positive feedback was found to increase with added nitrogen for both species,

which is somewhat surprising in light of the resource monopolizing hypothesis. On the other

hand, there was a beneficial effect of being relatively rare on the growth of all other vascular

species in plots when nitrogen was added. This advantage of being rare could be due to the

absence of negative interactions with soil pathogens or other natural enemies (e.g. van der

Putten et al. 2013). This hypothesis is consistent with the finding of positive covariation

between early and late season growth (Rinella et al. 2020). It has been shown that soil

inoculation significantly affects heathland plant communities (van der Bij et al. 2017; Wubs

et al. 2016) and that the composition of fungal communities depends on the composition of

the plant community (Radujković et al. 2020), suggesting that plant–soil biotic interactions

are especially important in heathland ecosystems. However, since the aggregated group

"other plant species" is made up of many species, the estimated effects will be the net result

of different species effects, and the effects of individual species is unknown. It would require

a larger experimental set-up to determine the density- and frequency-dependent

competition effects of each individual species.

For the two dominant species and the group of remaining species, the results show

significant frequency-dependent effects with positive effects of either being relatively

dominant or relatively rare.

Implicitly, it may be assumed that biomass at the end of the growing season translates to

cover at the start of the next growing season in a species-specific way, so that the observed

plant growth patterns may be used to predict community dynamics (Damgaard 2011;

Damgaard et al. 2009). The demonstrated positive effects of being relatively dominant for *C.*

vulgaris and *A. flexuosa* at the plot scale are corroborated by previous studies, where it was

concluded that the more locally dominant of the two species will outcompete the other

(Damgaard et al. 2009). The observed advantage of being dominant is also in agreement

with divergent long-term successional patterns on dry heath plots, where plots either

became dominated by grass or dwarf shrubs (Ransijn et al. 2015).

The observed widespread infestation of the heather beetle in 2020 had adverse effects

on *C. vulgaris* cover and growth. Such infestations may play an important role in

transforming a dwarf shrub-dominated heath to a grass-dominated heath. However, our

results suggest that a single heather beetle outbreak does not have long-term effects on the

growth and competitive ability of heather at this dune heath site (Kaae et al. 2024). It

appears that *C. vulgaris* is able to recover relatively quickly after infestation events, although

recurrent outbreaks could result in a transformation to a grass-dominated heath.

As previously demonstrated (Damgaard and Weiner 2021), there is compelling evidence

that the standard use of the Lotka-Volterra model in studies of species interactions in plant

communities is inadequate, and that most species interaction studies would be improved by

including frequency-dependency. It has also been demonstrated that if the effects of

measurement errors are ignored, model predictions may be biased (Carroll et al. 2006;

Damgaard 2020; Damgaard and Weiner 2021; Detto et al. 2019). It is therefore also

generally advisable to consider the possible effect of measurement errors when modelling

plant species interactions.

Traditional vegetation sampling in relatively small plots is not ideal for studying patch

dynamic mechanisms, such as positive feedback mechanisms, and cyclical vegetation

processes with ageing, which occur at a larger spatial scale. More research on the combined

effects of density- and frequency-dependent competition at larger spatial scales is needed.

New drone-based sampling methods of plant abundance that operate at a larger spatial

scales (e.g. Damgaard 2021) may provide much needed information for the study of plant

patch dynamics.

In conclusion, including frequency- as well as density-dependent effects, and unmeasured

variables in models of plant community dynamics could result in important improvements in

our understanding of plant growth, plant community dynamics and coexistence theory.

Modern statistical modelling methods make this easy to do.

Acknowledgements

323   We are grateful for funding from Aage V. Jensens Fonde and the Danish Long-term

324   Ecosystem Research Network (LTER-DK).

References

Adler PB, Smull D, Beard KH, Choi RT, Furniss T, Kulmatiski A, Meiners JM, Tredennick AT, Veblen KE. 2018. Competition and coexistence in plant communities: Intraspecific competition is stronger than interspecific competition. Ecol Lett. 21(9):1319-1329.

Aerts R, Berendse F, Caluwe Hd, Schmitz M. 1990. Competition in heathland along an experimental gradient of nutrient availability. Oikos. 57:310-318.

Aerts R, Heil GW. 1993. Heathlands. Patterns and processes in a changing environment.: Kluwer Academic Publishers.

Alonso I, Hartley SE, Thurlow M. 2001. Competition between heather and grasses on scottish moorlands: Interacting effects of nutrient enrichment and grazing regime. Journal of Vegetation Science. 12(2):249-260.

Barabás G, D'Andrea R, Stump SM. 2018. Chesson's coexistence theory. Ecological Monographs. 88(3):277-303.

Bever JD, Westover KM, Antonovics J. 1997. Incorporating the soil community into plant population dynamics: The utility of the feedback approach. J Ecol. 85(5):561-573.

Bähring A, Fichtner A, Ibe K, Schütze G, Temperton VM, von Oheimb G, Härdtle W. 2017. Ecosystem functions as indicators for heathland responses to nitrogen fertilisation. Ecological Indicators. 72:185-193.

Carlin BP, Louis TA. 1996. Bayes and empirical bayes methods for data analysis. London: Chapman & Hall.

Carroll RJ, Ruppert D, Stefanski LA, Crainiceanu C. 2006. Measurement error in nonlinear models: A modern perspective. Boca Raton: CRC.

Chesson P. 1994. Multispecies competition in variable environments. Theoretical Population Biology. 45(3):227-276.

Chesson P. 2000. Mechanisms of maintenance of species diversity. Annual Review of Ecology and Systematics. 31:343-366.

Chisholm RA, Fung T. 2020. Janzen-connell effects are a weak impediment to competitive exclusion. Am Nat. 196(5):649-661.

Connell JH, Tracey JG, Webb LJ. 1984. Compensatory recruitment, growth, and mortality as factors maintaining rain forest tree diversity. Ecological Monographs. 54(2):141-164.

Craine JM. 2009. Resource strategies of wild plants. Princeton University Press.

Damgaard C. 1998. Plant competition experiments: Testing hypotheses and estimating the probability of coexistence. Ecology. 79:1760-1767.

Damgaard C. 2005. Evolutionary ecology of plant-plant interactions - an empirical modelling approach. Aarhus, Denmark: Aarhus University Press.

Damgaard C. 2011. Measuring competition in plant communities where it is difficult to distinguish individual plants. Computational Ecology and Software. 1:125-137.

Damgaard C. 2019a. Is the degree of spatial aggregation of calluna vulgaris regulated by nitrogen deposition or precipitation? Acta Oecologica. 97:1-5.

Damgaard C. 2019b. Spatio-temporal structural equation modeling in a hierarchical bayesian framework: What controls wet heathland vegetation? Ecosystems. 22:152-164.

Damgaard C. 2020. Measurement uncertainty in ecological and environmental models. Trends Ecol Evol. 35:871-873.

Damgaard C. 2021. Integrating hierarchical statistical models and machine-learning algorithms for ground-truthing drone images of the vegetation: Taxonomy, abundance and population ecological models. Remote Sensing. 13(6).

Damgaard C, Riis-Nielsen T, Schmidt IK. 2009. Estimating plant competition coefficients

and predicting community dynamics from non-destructive pin-point data: A case study
with *calluna vulgaris* and *deschampsia flexuosa*. Plant Ecol. 201:687–697.
Damgaard C, Strandberg B, Mathiassen SK, Kudsk P. 2014. The effect of glyphosate on the
growth and competitive effect of perennial grass species in semi-natural grasslands.
Journal of environmental science and health Part B, Pesticides, food contaminants, and
agricultural wastes. 49(12):897-908.
Damgaard C, Weiner J. 2021. The need for alternative plant species interaction models.
Journal of Plant Ecology. 14:771-780.
De Long JR, Heinen R, Heinze J, Morriën E, Png GK, Sapsford SJ, Teste FP, Fry EL. 2023.
Plant-soil feedback: Incorporating untested influential drivers and reconciling
terminology. Plant and Soil. 485(1):7-43.
Detto M, Visser MD, Wright SJ, Pacala SW. 2019. Bias in the detection of negative density
dependence in plant communities. Ecol Lett. 22(11):1923-1939.
Diemont WH, Heil GW. 1984. Some long-term observations on cyclical and serial processes
in dutch heathlands. Biol Conserv. 30(3):283-290.
Ehlers BK, Damgaard CF, Laroche F. 2016. Intraspecific genetic variation and species
coexistence in plant communities. Biology letters. 12(1):20150853.
Gimingham CH. 1988. A reappraisal of cyclical processes in calluna heath. Vegetatio.
77(1/3):61-64.
Goldberg DE, Grace J, Tilman D. 1990. Components of resource competition in plant
communities. In: Grace J, Tilman D, editors. Perspective on plant competition. San
Diego: Academic press. p. 27-49.
Heil GW, Bobbink R. 1993. "Calluna", a simulation model for evaluation of impacts of
atmospheric nitrogen deposition on dry heathlands. Ecological Modelling. 68:161-
182.
Heinen R, Hannula SE, De Long JR, Huberty M, Jongen R, Kielak A, Steinauer K, Zhu F,
Bezemer TM. 2020. Plant community composition steers grassland vegetation via soil
legacy effects. Ecol Lett. 23(6):973-982.
Johansson M. 2001. Fungal associations of danish calluna vulgaris roots with special
reference to ericoid mycorrhiza. Plant and Soil. 231(2):225-232.
Jonasson S. 1983. The point intercept method for non-destructive estimation of biomass.
Phytocoenologia. 11:385-388.
Jonasson S. 1988. Evaluation of the point intercept method for the estimation of plant
biomass. Oikos. 52:101-106.
Jones AG, Power SA. 2012. Field-scale evaluation of effects of nitrogen deposition on the
functioning of heathland ecosystems. J Ecol. 100(2):331-342.
Kent M, Coker P. 1992. Vegetation description and analysis. London: Belhaven Press.
Kaae ME, Hu F, Bak JL, Strandberg MT, Damgaard CF. 2024. How do nitrogen deposition,
mowing, and deer grazing drive vegetation changes on dune heaths?
Ecologies.5(1):116-131. doi:10.3390/ecologies5010008.
Law R, Dieckmann U. 2000. A dynamical system for neighborhoods in plant communities.
Ecology. 81:2137-2148.
Mazzoleni S, Bonanomi G, Incerti G, Chiusano ML, Termolino P, Mingo A, Senatore M,
Giannino F, Cartenì F, Rietkerk M et al. 2015a. Inhibitory and toxic effects of
extracellular self-DNA in litter: A mechanism for negative plant–soil feedbacks? New
Phytologist. 205(3):1195-1210.
Mazzoleni S, Cartenì F, Bonanomi G, Senatore M, Termolino P, Giannino F, Incerti G,
Rietkerk M, Lanzotti V, Chiusano ML. 2015b. Inhibitory effects of extracellular self-
DNA: A general biological process? New Phytologist. 206(1):127-132.

Radujković D, van Diggelen R, Bobbink R, Weijters M, Harris J, Pawlett M, Vicca S,
Verbruggen E. 2020. Initial soil community drives heathland fungal community
trajectory over multiple years through altered plant–soil interactions. New Phytologist.
225(5):2140-2151.
Ransijn J, Kepfer-Rojas S, Verheyen K, Riis-Nielsen T, Schmidt IK. 2015. Hints for
alternative stable states from long-term vegetation dynamics in an unmanaged
heathland. Journal of Vegetation Science. 26(2):254-266.
Rinella MJ, Strong DJ, Vermeire LT. 2020. Omitted variable bias in studies of plant
interactions. Ecology. 101(6):e03020.
Singh P, Baruah G. 2020. Higher order interactions and species coexistence. Theoretical
Ecology. 14:71–83.
Smith SE, Read DJ. 2008. Mycorrhizal symbiosis. New York: Academic Press.
Southon GE, Field C, Caporn SJM, Britton AJ, Power SA. 2013. Nitrogen deposition reduces
plant diversity and alters ecosystem functioning: Field-scale evidence from a
nationwide survey of uk heathlands. PLOS ONE. 8(4):e59031.
Suding KN, Collins CG, Hallett LM, Larios L, Brigham LM, Dudney J, Farrer EC, Larson JE,
Shackelford N, Spasojevic MJ. 2024. Biodiversity in changing environments: An
external-driver internal-topology framework to guide intervention. Ecology.
105(8):e4322.
Talbot JM, Allison SD, Treseder KK. 2008. Decomposers in disguise: Mycorrhizal fungi as
regulators of soil c dynamics in ecosystems under global change. Functional Ecology.
22(6):955-963.
van der Bij AU, Pawlett M, Harris JA, Ritz K, van Diggelen R. 2017. Soil microbial
community assembly precedes vegetation development after drastic techniques to
mitigate effects of nitrogen deposition. Biol Conserv. 212:476-483.
van der Putten WH, Bardgett RD, Bever JD, Bezemer TM, Casper BB, Fukami T, Kardol P,
Klironomos JN, Kulmatiski A, Schweitzer JA et al. 2013. Plant–soil feedbacks: The
past, the present and future challenges. J Ecol. 101(2):265-276.
van Voorn GAK, Kooi BW, Bregt AK. 2016. Over-shading is critical for inducing a regime
shift from heathland to grassland under nitrogen enrichment. Ecol Complex. 27:74-83.
Watt AS. 1947. Pattern and process in the plant community. J Ecol. 35:1-22.
Watt AS. 1955. Bracken versus heather, a study in plant sociology. J Ecol. 43(2):490-506.
Weiner J. 1993. Competition among plants. Treballs de La Societat Catalana de Biologia.
44:99–109.
Wolfram S. 2019. Mathematica. 12.0.0.0 ed. Champaign, USA: Wolfram Research, Inc.
Wubs ERJ, van der Putten WH, Bosch M, Bezemer TM. 2016. Soil inoculation steers
restoration of terrestrial ecosystems. Nature Plants. 2(8):16107.
460
461

462
463 **Tables**
464

465 Table 1. Parameter explanations.

Parameter	Description
a_i	Vertical density of species *i* as a function of cover in the absence inter-specific density-dependent and frequency-dependent effects, and without added nitrogen
an_i	The effect of added nitrogen on a_i
b_i	Non-linear effect of cover on plant growth
c_{ij}	Competition coefficients, inter-specific density-dependent interaction effect of species *j* on the vertical density of species *i*.
cn_{ij}	The effect of added nitrogen on c_{ij}
d_i	Non-linear effect of cover on density-dependent inter-specific interaction effect
e_i	the frequency-dependent effect of species *i* on plant growth.
en_i	The effect of added nitrogen on e_i
σ_i	Structural uncertainty of the competitive growth model
v_i	Scale parameter of the measurement error of vertical density
ρ_1	Unexplained correlation coefficient between cover and vertical density

Table 2. Percentiles of the marginal posterior probability distribution of the model parameters. The subscripts indicate the taxa: 1 - *Calluna vulgaris*, 2 - *Avenella flexuosa*, and 3 - all other vascular plant species. The probability that the marginal distribution is above zero is given in the last column, where significant values are indicated in bold.

Parameter	2.5%	50%	97.5%	P(X > 0)
a_1	60.354	66.953	72.629	-
a_2	12.062	16.665	23.747	-
a_3	37.188	62.952	68.653	-
an_1	-1.248	-0.588	0.351	0.208
an_2	-0.641	-0.405	0.697	0.100
an_3	0.228	0.961	1.910	**0.993**
b_1	1.541	1.820	1.987	-
b_2	0.939	1.459	1.941	-
b_3	1.635	1.916	1.996	-
c_{12}	2.086	3.127	4.295	**>0.999**
c_{13}	0.817	1.385	1.817	**>0.999**
c_{21}	-1.036	0.114	1.691	0.541
c_{23}	0.596	1.490	2.892	**>0.999**
c_{31}	0.222	0.651	1.568	**>0.999**
c_{32}	-0.719	0.117	0.872	0.636
cn_{12}	-0.118	0.008	0.228	0.548
cn_{13}	-0.006	0.022	0.079	0.877
cn_{21}	-0.256	-0.058	0.086	0.344
cn_{23}	-0.074	0.065	0.136	0.709
cn_{31}	0.118	0.236	0.346	**>0.999**
cn_{32}	-0.067	0.005	0.178	0.532
d_1	0.826	1.511	1.962	-
d_2	1.157	1.656	1.986	-
d_3	1.552	1.823	1.988	-
e_1	-0.260	-0.202	-0.133	**<0.001**
e_2	-0.663	-0.467	0.013	**0.027**
e_3	0.859	1.282	1.466	**>0.999**
en_1	-0.124	-0.104	-0.095	**<0.001**
en_2	-0.315	-0.250	-0.176	**<0.001**
en_3	-0.054	-0.005	0.006	0.244
σ_1	0.060	0.092	0.138	-
σ_2	0.036	0.080	0.145	-
σ_3	0.042	0.333	1.059	-
v_1	7.437	7.889	7.996	-
v_2	7.679	7.944	7.998	-
v_3	7.796	7.960	7.999	-
ρ_1	-0.313	-0.048	0.283	0.372
ρ_2	-0.406	-0.062	0.341	0.390
ρ_3	0.671	0.773	0.799	**>0.999**

Figures

Fig. 1. Flowcharts illustrating components of the model. (a) The effect of an unmeasured

variable (z_u) on individual plant performance at both an early (x_i) and later growth stage

(y_i) of plant species i. The competitive growth process is modelled by the function, f_i, which

depends on x_i and possibly on some measured environmental variables (z_m). (b) The effect

of the unmeasured variable affecting both earlier and later growth stages is modelled as the

part of the covariance between the early and later growth stages that is not explained by the

independent factors in the competition model. (c) Hierarchical model in which true but

unknown factors affecting plant performance are modelled by latent variables (squares) and

measurement errors are separated from process errors. Data are denoted by circles.

Fig. 2. Map showing the location of Vust Hede (57°7'23.412" N, 9°0'42.443" E). The

rectangles are the blocks (1-5) with experimental manipulations (Kaae et al. 2024).

492   Fig. 3. Marginal posterior distributions of the parameters e_i and en_i that model frequency-

493   dependent growth. If $(e_i + en_i\, n) > 0$, then rare species have an advantage over dominant

494   species and, conversely, if $(e_i + en_i\, n) < 0$, then dominant species have an advantage over

495   rare species.

497
498
499
500
Fig 4. The predicted relative vertical density of *Calluna vulgaris* (purple) and *Avenella*

flexuosa (blue) as a function of its cover without added nitrogen. The sum of the cover of the

two species is set to one, i.e., only these two species are assumed to be locally present.

Electronic Supplements

The electronic supplements are located at https://osf.io/n6s9w/

Appendix A: Deer densities at Vust heath

Appendix B: Pin-point data (Excel file)

Appendix C: Software code (Mathematica notebook, get free viewer at

https://www.wolfram.com/player/)

Fig. S1: Residual plots of vertical density

513