The epitranscriptomic m6A RNA modification modulates synaptic function in ageing and in a mouse model of synucleinopathy

Avika Chopra[1], Mary Xylaki[1,2], Fanzheng Yin[1], Ricardo Castro-Hernández[3], Madiha Merghani[1], Valentina Grande[1], Brit Mollenhauer[2,4], André Fischer[3,5,6], and Tiago F. Outeiro[1,7,8,9*]

1. University Medical Center Göttingen, Department of Experimental Neurodegeneration, Center for Biostructural Imaging of Neurodegeneration, 37073, Göttingen, Germany.

2. Department of Neurology, University Medical Centre Göttingen, 37073, Göttingen, Germany.

3. Department for Systems Medicine and Epigenetics, German Center for Neurodegenerative Diseases (DZNE), Göttingen, Germany

4. Paracelsus Elena Klinik, Kassel, Germany

5. Department of Psychiatry and Psychotherapy, University Medical Center Göttingen, Göttingen, Germany

6. Cluster of Excellence "Multiscale Bioimaging: from Molecular Machines to Networks of Excitable Cells" (MBExC), University of Göttingen, Germany

7. Max Planck Institute for Multidisciplinary Sciences, 37075 Göttingen, Germany.

8. Translational and Clinical Research Institute, Faculty of Medical Sciences, Newcastle University, Framlington Place, Newcastle Upon Tyne, NE2 4HH, UK.

9. Scientific employee with an honorary contract at Deutsches Zentrum für Neurodegenerative Erkrankungen (DZNE), 37075, Göttingen, Germany.

24   ***Correspondence to:** Tiago F. Outeiro

25   Department of Experimental Neurodegeneration, Centre for Biostructural Imaging of

26   Neurodegeneration, University Medical Centre Göttingen, Göttingen, Germany

27   Email: tiago.outeiro@med.uni-goettingen.de

Abstract

N6-methyladenosine (m6A) is the most abundant and conserved transcriptional modification in eukaryotic RNA, regulating RNA fate. While the functions of m6A in the development of the mammalian brain have been extensively studied, its roles in synaptic plasticity, cognitive decline, motor function, or other brain circuits remain underexplored. To date, the role of this modification in Parkinson's disease (PD) and other synucleinopathies has been largely unknown. Here, we investigated the m6A epitranscriptome in a mouse model of synucleinopathy. We performed m6A RNA immunoprecipitation sequencing (meRIP-seq) to obtain the m6A epitranscriptome of the midbrain in young (3 mo) and aged (15 mo) A30P-aSyn transgenic mice (aSyn Tg) and C57BL6 control wild type (Wt) mice. We observed hypermethylation of synaptic genes in 3 mo aSyn Tg mice compared to age-matched Wt mice. This methylation was reduced during ageing, with synaptic genes becoming increasingly hypomethylated. Using immunofluorescence imaging alongside biochemical analysis, we further investigated the expression of m6A regulatory enzymes — writer, N6-Adenosine-Methyltransferase Complex Catalytic Subunit (METTL3); reader, YTH N6-methyladenosine RNA-binding protein (YTHDF1); and eraser, fat mass and obesity-associated protein (FTO) — in the cortex, striatum, hippocampus, and cerebellum of Wt and aSyn Tg mice, as well as in primary cortical neuronal cultures. We observed that the levels of METTL3, YTHDF1 and FTO were similar between Wt and aSyn Tg mice. Interestingly, the writer protein METTL3 was found in both the nucleus and in the post-synaptic compartment in neuronal cultures. Our findings suggest that alterations in the regulation of m6A RNA methylation may be associated with

50   neurodegeneration and ageing and that this level of epitranscriptomic regulation plays a

51   significant role at the synapse.

53   **Keywords:** epigenetics, m6A, Parkinson's disease, ageing, METTL3

Introduction

Epigenetics is a process that acts as a bridge between genotype and phenotype, changing the outcome of a gene without modifying the DNA sequence. It encompasses various types of alterations such as methylation of DNA, histone modifications, and chromatin remodeling [1,2]. The field that studies posttranscriptional changes of RNA bases is known as epitranscriptomics and aims to identify modifications in the transcriptome and unravel their regulatory roles [3]. N^6-methyladenine (m6A) modification is the most prevalent mRNA modification occurring at a frequency of 0.15 - 0.6% of all adenosines [4–6] and is found in different types of RNAs including protein coding, messenger RNA (mRNA) [7,8]. This modification is being heavily studied due to its prevalence and importance in eukaryotes, where it regulates various cellular functions [8–12]. Methylation marks are mostly added co-transcriptionally to specific consensus motif sequences and are mostly found close to the 3' un-translated region (3'-UTR) at the stop codon [4,13].

The m6A modification machinery consists of three groups of enzymes, namely methyltransferases, demethylases and detectors of this modification, also termed writers, erasers, and readers, respectively. The modification can be added onto the RNA by methyltransferases including methyltransferase-like 3 (METTL3), methyltransferase-like 14 (METTL14), and adapter protein Wilms tumour 1-associated protein (WTAP) [5,14–19]. METTL3 was the first discovered methyltransferase and was described as a key component in the methylation of transcripts involved in physiological processes such as embryonic development and brain development [20–23]. The catalytically active part of METTL3 forms a heterodimer with METTL14, which provides the structure to install the m6A modification onto RNA [5,24–27]. Apart from these, the m6A methyltransferase complex comprises other

proteins, including RNA binding proteins (RBP) like RBP15, which recruit this machinery to

the target mRNA to facilitate methylation [15,28]. Demethylases, for instance, fat mass and

obesity-associated protein (FTO) and alkB homologue 5 (ALKBH5) remove the m6A

modification marks making this a dynamic process [29,30]. FTO was the first eraser protein

identified [29,31] and it carries out oxidative demethylation where m6A is converted in

several steps to N^6-hydroxymethyl adenosine (hm6A), then to N^6-formyladenosine (f6A) and

lastly to adenine [31]. Additionally, reader proteins like the YT521-B homology (YTH)

domain family of proteins including YTHDF1, YTHDF2/3, insulin-like growth factor 2 mRNA-

binding proteins (IGF2BP1), heterogeneous nuclear ribonucleoprotein (HNRNPA2B1)

recognise the modification and modulate alternative splicing and processing of target

transcripts [8,32]. The ones located in the cytoplasm include YTHDF1, YTHDF2, and YTHDF3.

The main function of these proteins is unclear; however, studies indicate different roles of

these enzymes in a cell. While YTHDF1 promotes the translation of mRNA thereby

promoting its protein expression, YTHDF2 promotes the degradation of mRNA, and YTHDF3

promotes both translation and degradation of mRNA [33,34].

m6A methylation has been associated with ageing, and downregulation in the expression of

FTO and an overall increase in RNA m6A modification was found in human follicular fluid

(FF) and granulosa cells (GCs) from aged donors and murine ovaries from old mice [35].

Another recent study has also highlighted the relationship between elevated m6A levels and

brain ageing in both mice and humans [36], although a significant downregulation of m6A

methylation in ageing mice and in postmortem brain tissue from Alzheimer's disease

patients has also been reported [37].

Strikingly, ageing is also a major risk factor for the onset and progression of

neurodegenerative disorders such as Parkinson's disease (PD) and related

synucleinopathies, disorders associated with the accumulation of aggregated alpha-

synuclein (aSyn) protein in the brain. Despite the growing interest in the role of m6A

methylation in ageing, its role in PD is still unclear and remains controversial. While a recent

study indicated that the expression levels of m6A are significantly reduced in cellular models

of PD and in the striatum of a rodent model of PD [38], a different study showed that the

m6A methyl transferases were significantly decreased while the demethylases were

increased in the substantia nigra and in the striatum of mouse models of PD [39]. Ageing

also affects multiple homeostatic pathways including mitochondrial function and protein

degradation pathways, ultimately leading to cell death in the substantia nigra [40–42].

In this study, we investigated the m6A RNA modification profile in the context of ageing and

neurodegeneration in a mouse model of synucleinopathy. We performed methylated RNA

immunoprecipitation sequencing (meRIP-seq) to obtain the m6A epitranscriptome of the

midbrain of C57BL6 (Wt) control and A30P aSyn PD (aSyn Tg) mice at 3 months (3 mo) and

15 months (15 mo) of age. We observed an increase in the m6A methylation in the genes

involved in the molecular and biological processes at the synapse. Several genes involved in

synaptic processes were hypermethylated in 3 mo aSyn Tg vs. Wt mice, while the

methylation marks were reduced in 15 mo aSyn Tg vs. age-matched Wt mice. Using

immunofluorescence imaging of primary cortical neurons, we observed that the localisation

of the writer METTL3 enzyme is not only limited to the nucleus but is also present at the

post-synapse. Our findings suggest that m6A RNA methylation is possibly playing a role in

neurodegeneration and ageing, and is involved in regulating synaptic function.

Results

m6A alterations during ageing in Wt mice and in a transgenic model of synucleinopathy

We began by examining the distribution of differentially expressed transcripts and transcripts with m6A methylation in the healthy ageing brain of Wt mice. We further extended the analysis to aSyn Tg mice, in order to compare the effects of ageing on m6A modification and assess this change in the context of synucleinopathy. We extracted and dissected the midbrains from five 3 mo and five 15 mo Wt and an equal number of aSyn Tg mice, as this brain region is particularly relevant in the context of PD and related synucleinopathies. Methylated RNA immunoprecipitation sequencing (meRIP-seq) was performed to determine the midbrain-specific epitranscriptomic landscape in 3 mo and 15 mo mice (Figure 1A, B). First, we compared the differential expression of genes due to ageing in 15 mo vs 3 mo mice in both Wt and aSyn Tg animals. We observed a stronger effect in aSyn Tg mice where 270 transcripts were differentially expressed when compared with Wt mice, where only 79 transcripts were differentially expressed. Most differentially expressed transcripts in aSyn Tg mice were downregulated (Supplementary Figure S1C). However, when we compared the differentially methylated transcripts, we found similar numbers between Wt (420 genes) and Tg (424 genes) mice (Figure 1A).

On the other hand, the effect of aSyn A30P overexpression was assessed by a comparison between aSyn Tg and Wt mice at 3 mo and 15 mo. We did not identify major differences in the number of differentially expressed transcripts which were 128 at 3 mo and 142 at 15 mo. Similarly, the differentially methylated transcripts showed no significant changes

between Wt and aSyn Tg at 3 mo (384 genes) and 15 mo (353 genes) (Figure 1B). While we

did not see a significant difference in the differential methylation, we found that the

prevalence of hyper- and hypomethylated transcripts was reversed (Figure 1B).

As a consequence of ageing, the methylation levels of the differentially methylated m6A

transcripts in 15 mo vs. 3 mo Wt mice we found 418 hypermethylated and only 2

hypomethylated transcripts. On the contrary, 171 genes were hypermethylated and 244

transcripts were hypomethylated in 15 mo vs. 3 mo aSyn Tg mice (Figure 1D). A transcript

can carry multiple m6A methylation marks and some of these marks likely increase while

others decrease simultaneously. Such transcripts are referred to as mixed transcripts. We

found only 9 such mixed transcripts (Figure 1D, left column) in aSyn Tg mice during ageing.

To identify the effect of aSyn expression on the levels of the m6A modification, we

compared 3 mo aSyn Tg mice with age-matched Wt mice. We found 380 hypermethylated

transcripts and only 4 transcripts were hypomethylated, and no mixed transcripts. On the

other hand, when comparing 15 mo aSyn Tg vs. Wt mice, a reverse trend was observed,

with 108 hypermethylated, 240 hypomethylated, and only 5 mixed transcripts (Figure 1D,

right column).

We found 6135 methylated transcripts in common between all groups (Figure 1E). The 3 mo

Wt mice displayed 266 unique methylated transcripts and the 15 mo Wt mice 255. These

numbers were lower for the aSyn Tg group where the 3 mo mice showed 165 unique

methylated transcripts and the 15 mo mice showed 189. At 15 months, both genotypes

showed 124 commonly methylated transcripts, suggesting the occurrence of conserved

processes in normal ageing and synucleinopathies. The aSyn Tg mice displayed 53

methylated transcripts correlated with aSyn expression and conserved with ageing.

The overall distribution of m6A methylation peaks on mRNA did not display any striking

anomalies in any of the groups. As expected, the majority of the m6A modifications were

localised close to the 3' UTR on mRNA (Figure 1F), as previously reported for ageing-related

genes [36].

m6A levels are reduced in synaptic transcripts upon ageing

To investigate the cellular processes and the subcellular regions where the differentially

methylated transcripts were identified in 15 mo aSyn Tg vs. Wt mice, we performed gene

ontology (GO) term analysis. We found that synaptic function was the biological process

most represented. Our analysis revealed that transcripts with m6A methylation marks have

strong enrichment for biological processes such as receptor internalization, excitatory post-

synaptic potential, axonal transport, and dendritic spine morphogenesis. The identified

transcripts were also involved in cellular processes including neuron projection cytoplasm

and synaptic vesicle membrane (Figure 2A). Previous data from our group has also shown

that while the average dendritic spine number was not significantly different, a strong

difference was observed in the abundance of different types of spines in aSyn Tg neurons

when compared with Wt neurons [67].

We also observed that when we compared the differentially enriched genes in aSyn Tg 15

mo vs. 3 mo, we saw a strong enrichment of pathways involved in the immune system. The

genes with highest enrichment score were involved in innate immune system, adaptive

immune system, immunoglobulin binding, phagocytic vesicle, defense response, innate

immune response, and regulation of immune system processes among others

(Supplementary Figure S2).

We used the SynGO software [64] (https://syngoportal.org/), an experimentally annotated

database for synaptic location and functional GO, to specifically check the m6A methylation

of synaptic transcripts. These differentially methylated genes are primarily involved in

synaptic organisation, metabolism, transport, signaling, and both pre-and post-synaptic

processes. Based on meRIP and RNA sequencing analysis, sunburst plots of Wt and aSyn Tg

mice were generated (Figure 2B). Our findings suggest that the differentially methylated

transcripts that are involved at the synapse and in synaptic function were majorly

hypermethylated in 3 mo aSyn Tg mice compared to 3 mo Wt mice (Figure 2A). In 3 mo

mice, the hypermethylated transcripts were concentrated in synapse organisation and at

the pre-synapse (Figure 2B). Conversely, at 15 months of age, the majority of the

differentially methylated transcripts were hypomethylated when aSyn Tg mice are

compared with Wt mice, which were largely involved in signalling, pre- and post-synapse,

and synapse organisation (Figure 2C).

In summary, we observed that the differentially methylated genes, both hyper- and

hypomethylated genes in 15 mo aSyn Tg vs. Wt mice were mostly hypomethylated and

involved in synaptic processes, especially at the pre-and post-synapse. However, at 3 mo,

most of the differentially methylated genes were hypermethylated.

208 **The levels of m6A regulatory proteins do not change in different brain regions in Wt and**

209 **aSyn Tg mice**

To validate the meRIP-Seq results and to evaluate if the overall levels of the m6A

modification are affected in aSyn Tg vs. Wt mice, we performed an m6A ELISA, for total RNA

isolated from cortical and midbrain brain tissue of young and aged, Wt and aSyn Tg mice.

We found that overall, the levels of m6A modification marks were higher in the cortex

compared to the midbrain. In cortical brain tissue, we observed that the m6A marks in the

aSyn Tg mice do not display the same pattern as in Wt animals (Figure 3A). A significant

reduction in m6A marks was observed in 12 mo aSyn Tg mice compared to 12 mo Wt mice

(Wt; mean = 0.032 ± 0.018 OD, aSyn Tg; mean = 0.011 ± 0.002 OD, adjusted p-value =

0.045). In the younger animals, however not significant, a slight reduction in m6A marks was

observed in 3 mo aSyn Tg were compared with 3 mo Wt mice, indicative of an effect that

probably starts at a young age and progresses with ageing (Figure 3A). In the midbrain, we

did not observe significant differences in the m6A marks between the groups (Figure 3A).

This is consistent with the meRIP-Seq data where we observed that the total number of

differentially methylated transcripts was not altered among the groups (Figure 1A, B). This

suggests that despite the observed hypo- and hyper-methylation in certain transcripts

(Figure 1D) the overall m6A levels remain in an equilibrium in the midbrain.

Next, we asked if the observed changes in the m6A marks in the cortex correlate with

altered levels of the three m6A regulatory proteins in the cortical brain tissue. To do so, we

performed immunoblotting analysis to assess the expression of m6A regulators — METTL3,

YTHDF1, and FTO. We used three different age groups – 3 mo, 6mo and 12 mo from the Wt

and aSyn Tg mice for our analysis. We observed no significant differences in the expression

levels of these m6A regulatory proteins neither within genotypes (Tg vs. Wt) nor across the

different age groups (Figure 3B). Nonetheless, we observed a trend towards a reduction in

the levels of METTL3 and FTO in Wt mice during ageing (Figure 3B).

Considering no difference in the protein levels of m6A regulators, we sought to identify if

the sub-regional and/or subcellular localisation of the three m6A regulatory proteins is

altered. For this, we assessed the levels of METTL3, YTHDF1 and FTO by

immunofluorescence intensity analysis in four different regions of the mouse brain, namely

cortex, striatum, hippocampus, and cerebellum. MAP2 co-staining was used only to identify

the neurons present in the areas of the brain. All these proteins were present in our

selected areas of interest, but their compartmentalization varied across different regions of

the neurons.

The m6A writer protein, METTL3 was observed as small puncta within the nucleus in the

cortex, striatum, hippocampus, and cerebellum in both aged Wt and aSyn Tg brain (Figure

4A). We found no significant differences in protein levels between aSyn Tg vs. Wt mice

based on the fluorescence intensity (Figure 4B).

The m6A reader protein, YTHDF1, on the other hand, localised only in the soma of neurons

in all four areas of interest, with a similar expression pattern in Wt and aSyn Tg mouse brain

slices. (Figure 4A). No significant difference in the fluorescence intensity of the protein was

observed in 15 mo aSyn Tg mice when they were compared with Wt mice (Figure 4B).

Finally, the m6A eraser, FTO, was detected in the nucleus and the soma of cells in the

cortex, striatum, hippocampus, and cerebellum. FTO was also localized in the dendrites of

neurons in the cortex and cerebellum in both Wt and aSyn Tg mouse brain slices (Figure 4A).

After analysis of the expression of this protein in the soma, no significant differences were

observed in the different regions of either aSyn Tg or Wt mice (Figure 4B).

Our findings show that the m6A regulatory proteins – METTL3, YTHDF1 and FTO are present

in the cortex, striatum, hippocampus, and the cerebellum in the mouse brain. In addition,

we found that the levels of METTL3, YTHDF1 and FTO do not change in cortical brain tissue.

METTL3 is reduced in the aSyn Tg mouse post-synapse

Based on our bioinformatics analysis, we observed that the majority of m6a genes were

synaptic, so we asked if/and to what degree the m6A regulators are present at synapses

acting as local regulators since the overall regulator levels remain unchanged. We used

primary cortical neurons that allow higher imaging potential and performed co-staining of

the three m6A regulatory proteins – METTL3, YTHDF1 and FTO alongside Syn1, pre-synaptic

marker and PSD95, post-synaptic marker to study the localisation of these proteins at the

synapse. Previous studies have indicated reduced synaptic activity in neurons derived from

sporadic PD patients when compared with controls [65,66].

Two experimenters performed the colocalisation analysis, and similar results were obtained.

METTL3 was observed to be present in the nucleus as nuclear speckles and it significantly

co-localized with PSD95, indicating its presence at the post-synapse in both Wt (PCC = 0.810

± 0.039) and aSyn Tg (PCC = 0.748 ± 0.075) primary neurons. In contrast, METTL3 did not co-

localize with synaptotagmin1 in Wt (PCC: 0.489 ± 0.086) or aSyn Tg (PCC: 0.476 ± 0.050)

primary neurons, indicating no noticeable localisation at the pre-synapse (Figure 5B).

Similarly, we investigated the compartmentalization of YTHDF1, which did not overlap with

either synaptotagmin1 (PCC: Wt = 0.382 ± 0.090, aSyn Tg = 0.294 ± 0.065) or PSD95 (PCC:

Wt = 0.310 ± 0.088, aSyn Tg = 0.280 ± 0.069) (Figure 5B). The m6A eraser protein, FTO, was

identified within the cell nucleus and in some cases, in the soma of primary neurons. No

significant co-localisation of FTO was observed at the pre- (PCC, Wt = 0.468 ± 0.109, aSyn Tg

= 0.488 ± 0.095) or post-synapse (PCC, Wt = 0.366 ± 0.063, aSyn Tg = 0.463 ± 0.081) (Figure

5B).

Overall, we observed that the expression of METTL3 was reduced at the post-synapse in

aSyn Tg primary neurons suggesting a possible loss of METTL3 at the synapse.

Primary neurons from aSyn Tg mice have a greater number of synapses than Wt animals

Since we observed a reduction in the localisation of METTL3 at the synapse in aSyn Tg

primary neurons, we wondered whether this has any influence on the synapse morphology

and number, or vice versa. For this, we used primary cortical neurons and performed

proximity analysis of the pre-and post-synaptic markers Syn1 and PSD95 using the IMARIS

software Imaris (RRID: SCR_007370) and the data were analysed blindly to avoid bias. We

found that the number of synapses in aSyn Tg primary neurons was significantly increased

when compared with Wt neurons (Wt; mean = 2607 ± 763.1, aSyn Tg; mean = 3152 ± 1187,

p-value = 0.044) (Figure 6).

Discussion

The involvement of epitranscriptomics in PD is still not well understood. In this study, our

objective was to begin to unravel the role of m6A RNA methylation in the A30P aSyn Tg

mouse model of PD/synucleinopathy, in disease and physiological ageing.

Our findings revealed that 99.5% of the transcripts were hypermethylated in 15 mo vs. 3 mo

Wt mice suggesting that ageing might contribute to a significant increase in m6A

methylation. This observation aligns with a recent study that has demonstrated an increase

in m6A methylation during ageing in both mice and humans [36]. This indicates that the

accumulation of m6A methylation marks might be a general feature of ageing in the brain,

potentially influencing various biological processes. It has also been shown that these

epigenetic changes arise mainly in young and aged mice, while during the adolescent stage

these genes are stabilised [68]. In contrast, when comparing 15 mo aSyn Tg mice with their

3 mo counterpart littermates, more transcripts were hypomethylated. This decrease in

methylation with age in aSyn Tg mice might suggest that the loss of methylation might be

related to aSyn expression/pathology. There is a possibility that the overexpressed aSyn is

somehow interacting with the m6A regulators and inhibiting methylation. Previous studies

have shown that the 6-hydroxydopamine rat PD model has significantly reduced m6A levels

in the striatal brain region and in the cellular model of PD [69]. Similarly, a reduction in the

total levels of m6A was also reported in the MPTP-treated PD mouse model and in MPP+-

treated MN9D cells [70].

The effect of aSyn expression was more directly studied in the comparison between 3 mo

aSyn Tg and Wt mice. The increase in methylation suggests that aSyn expression may be

linked to enhanced methylation activity in early life. However, in 15 mo aSyn Tg vs. Wt mice,

we observed more hypomethylated genes, suggesting that A30P aSyn expression, and likely

pathology, might interfere with the writing or erasing of methylation marks during ageing.

Importantly, our results are in accordance with a recent publication suggesting that the

levels of m6A are significantly lower in PD patients [71].

Interestingly, based on GO term analysis, we found that differentially methylated transcripts

were predominantly associated with synaptic function and that methylation was lost with

ageing. Hypermethylation of transcripts involved in the synaptic processes might lead to an

inhibition of mRNA translation and thereafter reduced functionality of the synapse. This is

particularly relevant as more than half of the causative genes and risk factors in PD have

been identified to function at the synapse [72] and this change in methylation pattern might

indirectly affect the functioning of the synapse. Considering that different RNA-based

epigenetic mechanisms occur locally at the synapse, this underscores their potential

importance in synaptic regulation [74]. Additionally, GO term analysis of the differentially

expressed transcripts showed a strong enrichment in immune response-related pathways.

Notably, m6A is implicated in the regulation of inflammation under physiological and

pathological conditions, known as "inflammatory ageing", or inflammaging. This is a chronic

inflammatory state arising during ageing and it is characterised by an increase in pro-

inflammatory factors such as tumor necrosis factor (TNF-α), interleukins (IL-1β and IL-6), and

C-reactive protein (CRP), to name just a few [73].

The assessment of total m6A methylation marks showed a significant reduction in the cortex

of 12 mo aSyn Tg mice when compared with age-matched Wt mice. This suggests that the

overall m6A methylation decreases in the ageing brain. This finding is consistent with recent

studies that reported a predominant reduction in m6A methylation in the frontal cortex and

cingulate cortex from PD patients [71,75]. In vitro studies have also reported a significant

reduction in m6A levels in the cellular PD models [70]. Despite this, we observed similar

expression levels of m6A regulatory proteins. However, our findings are consistent with

those from a recent study reporting that the levels of YTHDF1 and FTO mRNA remain

unchanged in PD. However, a reduction in the levels of METTL3 mRNA in PD patients was

also reported in that study, in contrast to what we observed [71]. Some studies have also

reported up-regulation of ALKBH5, IGF2BP2 in the substantia nigra while, YTHDF1 and

fragile X mRNA (FMR1) were downregulated. In the striatum, an up-regulation of FMR1 and

Cbl proto-oncogene like 1 (CBLL1) and down-regulation of METTL3, RBM15 and IGF2BP3

was observed [76].

Although we found no significant differences in the levels of m6A regulatory proteins, we

observed that the three proteins were present in the cortex, striatum, hippocampus, and

cerebellum. Our findings are consistent with those from previous studies reporting that the

m6A-methyltransferase complex, including METTL3, is predominantly found in the nucleus,

but that it can also be localised in the cytoplasm [77–79]. As methylation generally occurs

co-transcriptionally in the nucleus, while reading occurs largely at the cytosol, YTHDF1 is

primarily cytosolic [80]. FTO, on the other hand, is localised both in the nucleus and in the

cytosol [81]. These observations suggest that an inhibitory factor affecting their function

and aSyn aggregation is potentially playing a role. Previous studies have shown that aSyn

binds to histone proteins, which may alter the protein surface due to post-translation

modifications of histone tails, altering their affinity for DNA and other histones [82]. Another

study showed that aSyn expression increases histone-H3 lysine-9 (H3K9) methylation,

suggesting a possible interaction of aSyn with epigenetic writer proteins [83]. This could

constitute a mechanism through which aSyn affects gene methylation, ultimately affecting

synaptic functioning [84,85].

Given that most of the hypermethylated genes were synaptic, we assessed the presence of

these regulators at the synapse. Our findings suggest the presence of METTL3 at the post-

synapse, indicating that the enzyme might be locally active at the synapse. The dynamic

nature of m6A methylation may be localised to different regions within a cell, depending on

the temporal requirements of the respective genes [86,87]. In another study, m6A

regulatory proteins were found localised adjacent to the synapse and in the dendrites, also

suggesting the sub-cellular localisation of m6A methylated mRNA which might lead to

synaptic modifications [88,89]. Our findings revealed that the expression of METTL3 is

reduced at the post-synapse in aSyn Tg DIV14 neurons, suggesting a possible loss of METTL3

at the post-synapse. Due to this reduction at an early stage, the nucleus might be sending

hypermethylated transcripts to the dendrites to compensate for this loss. However, this may

be lost during ageing. Based on our bioinformatics analysis, we observed that methylated

transcripts associated with synaptic function were reduced in aSyn Tg mice with ageing. This

suggests that a reduction in the m6A methylation in synaptic genes may be directly linked

with the reduced METTL3 localisation at the synapse in aSyn Tg neurons.

Conclusion

In conclusion, our study highlights the critical role of m6A RNA methylation in the

progression of PD and its implications in ageing, particularly focusing on the A30P aSyn

mouse model. Our findings indicate that aSyn expression/pathology may influence RNA

methylation patterns, leading to reduced levels of m6A in mice. The differentially

methylated transcripts were predominantly synaptic, indicating that m6A modifications play

a crucial role in synapse physiology and potentially contribute to the synaptic dysfunction

observed in PD [90]. Although we observed no significant changes in the levels of m6A

regulatory proteins, the reduction in METTL3 at the synapse in aSyn Tg mice informs on the

importance of subcellular compartmentalization of these proteins in the regulation of RNA

methylation. Therefore, our research provides new insights into the epigenetic mechanisms

underlying PD and underscores the need for further exploration of m6A RNA methylation as

a potential therapeutic target for PD and other related synucleinopathies.

Materials and methods

Resource availability

Lead contact for reagent and resource sharing

Further information and requests for resources and reagents should be directed to and will

be fulfilled by the lead contact, Tiago F. Outeiro (tiago.outeiro@med.uni-goettingen.de).

Experimental model and subject details

Animals

Five male homozygous [A30P] aSyn transgenic (aSyn Tg) mice (C57BL/6J-Tg(Th-

SNCA*A30P*A53T) 39Eric/J, Strain #:008239, RRID: IMSR_JAX:008239) and five wild type

(Wt) littermates, 3 mo, 6mo, 12 mo and 15 mo were included in this study for RNA

sequencing and analysis. Additional animals used for each experiment are mentioned in the

respective section. Animals were housed in standard cages with 12h light/12h dark cycle

and ad libitum access to food and water in the animal facility of the University Medical

Centre Göttingen (Göttingen, Germany). Animal procedures were performed in accordance

with the European Community (Directive 2010/63/EU); and institutional and national

guidelines under the Lower Saxony State Office for Consumer Protection and Food Safety

(LAVES) (license number 19.3213). Animals were anesthetised with carbon dioxide (CO_2) and

then sacrificed using cervical dislocation. For tissue processing, the brain was surgically

extracted from the skull and was further dissected to isolate the cortex and midbrain. The

tissue was snap-frozen in liquid nitrogen and preserved at -80°C until use.

For the preparation of brain slices, the animals were perfused with 1x phosphate buffer

saline (PBS) followed by 4% paraformaldehyde (PFA). The whole brain was dissected out of

the skull and placed overnight in 4% PFA at 4°C. The following day, the brain was washed

once with 1x PBS and then placed in 15% saccharose and 30% saccharose with 0.1% sodium

azide (NaN_3). The brains were placed at 4°C until used. Sagittal cryo-sections of 30 μm

thickness were cryo-sectioned using Leica CM3050 S cryostat for each mouse brain. The

sections were placed in 1x PBS with NaN_3 until they were used for immunohistochemistry

(IHC) protocol.

Neuronal cultures

Primary cortical neuronal cultures were prepared from both aSyn Tg and Wt mice as previously described (Saal et al., 2015). Cells were seeded on coverslips coated with poly-L-ornithine (0.1 mg/mL in borate buffer; PLO) (Sigma-Aldrich, MO, USA) or culture plates (Corning, Merck, Darmstadt, Germany) for immunocytochemistry and maintained in neuronal cell culture medium (Neurobasal medium, Gibco Invitrogen, CA, USA), supplemented with 1% penicillin-streptomycin (PAN Biotech, Aidenbach, Germany), 0.25% GlutaMAX (Gibco Invitrogen, CA, USA), and 2% B27 (B-27™ Plus Supplement (50X), Gibco, #A3582801). The cells were maintained at 37°C with 5% CO_2, and one-third of the medium was replaced with fresh media every 3-4 days.

Immunocytochemistry

Primary neurons were first washed with 1x PBS (PAN Biotech, Aidenbach, Germany) and fixed with 4% of PFA for 20 min at room temperature (RT). Cells were first permeabilized with 0.1% Triton X-100 (Sigma-Aldrich, MO, USA) for 10 min and then blocked with 1.5% normal goat serum (NGS) in 1x PBS (Sigma-Aldrich, MO, USA) for 1h at RT. The cells were then incubated with the primary antibodies diluted in blocking solution (FTO (mouse, Abcam – ab92821, 1:1000), METTL3 (rabbit, Cell Signalling - #51104, 1:1000), YTHDF1 (rabbit, Proteintech 17479-1-AP, 1:1000), MAP2 (rabbit, Proteintech – #17490-1-AP, 1:1000), postsynaptic density protein 95 (PSD95) (mouse, Synaptic systems 124011, 1:1000), PSD95(D74D3) (rabbit, Cell Signalling - #3409, 1:1000), synaptotagmin 1 (Syn1) (mouse, Synaptic systems - #105311, 1:1000) synaptotagmin1 (rabbit, Synaptic systems – 105102/3,

1:1000)) overnight at 4°C, with constant shaking at a speed of 40-50 rpm. The following day,

cells were washed three times for 10 min each with 1x PBS (PAN Biotech, Aidenbach,

Germany) and then incubated with fluorescently conjugated secondary antibodies (Alexa

Fluor 488 donkey anti-mouse IgG – Invitrogen, Alexa Fluor 488 donkey anti-rabbit IgG -

Invitrogen, Alexa Fluor 555 donkey anti-mouse IgG – Invitrogen, Alexa Fluor 555 donkey

anti-rabbit IgG – Invitrogen, Alexa fluor 633 goat anti-chicken IgG - Invitrogen) for 2h at RT,

on a shaker at 40-50 rpm. Lastly, nuclei were counter-stained with DAPI (Carl Roth,

Karlsruhe, Germany) and mounted on glass slides (Epredia™ SuperFrostPlus™ Adhesion)

with mounting media (Invitrogen™ Fluoromount-G™) for microscopy.

Immunohistochemistry

The free-floating tissue was carefully transferred from PBS and $NaNH_3$ to 1x tris-buffered

saline (TBS) in a new 24-well plate with immunohistochemistry (IHC) grids. The tissue was

washed three times for 5 min each with 1x TBS to remove the remnants of NaN_3. Following

this, tissue antigen retrieval was performed using citrate buffer (10 mM citrate, 0.05%

Tween 20, pH 6.0) at 90°C for 30 min in a water bath. The tissue samples were allowed to

cool down until they reached RT. Tissue sections were washed thrice for 10 min each with

1x TBS and then incubated with 0.3% Triton X-100 for 20 min. Without washing, the tissue

slices were incubated directly in a blocking solution (5% BSA, and 1% NGS in 1x TBS) for 2h

at RT. The tissue sections were then incubated for 48h with antibodies diluted 1:200 times

(FTO (mouse, Abcam – ab92821, 1:1000), METTL3 (rabbit, Cell Signalling - #51104, 1:1000),

YTHDF1 (rabbit, Proteintech #17479-1-AP, 1:1000), MAP2 (rabbit, Proteintech – #17490-1-

AP, 1:1000)) at 4°C, on a shaker. The following day, tissue sections were washed thrice for

10 min each with 1x TBS at RT. Afterwards, the tissue sections were incubated in secondary

antibody (1:400 dilution) (Alexa Fluor 488 donkey anti-mouse IgG – Invitrogen, Alexa Fluor

488 donkey anti-rabbit IgG - Invitrogen, Alexa Fluor 555 donkey anti-mouse IgG –

Invitrogen, Alexa Fluor 555 donkey anti-rabbit IgG – Invitrogen, Alexa fluor 633 goat anti-

chicken IgG - Invitrogen) overnight at 4°C on the shaker. The following day, the tissue was

washed thrice for 10 min each with 1x TBS to remove the unbound secondary antibody. The

tissue sections were incubated for 5 min in DAPI (Carl Roth, Karlsruhe, Germany) followed

by a 10 min wash with 1x TBS. The tissue sections were then mounted onto glass slides

(Epredia™ SuperFrostPlus™ Adhesion) and left to dry in the dark at RT. 8-10µl of mounting

media (Invitrogen™ Fluoromount - G™) was applied as a drop onto each section on the glass

slide and a glass coverslip (Avantor, 631-0147, 0.16-0.19mm thickness, #1.5) was carefully

placed on the tissue to avoid any air bubbles.

Western blots

Cortical brain tissue was extracted from three adult mice per age group (age group 2 mo, 6

mo and 12 mo) from each Wt and Tg mice. The tissue from adult mice was lysed using

homogenisation beads (1.4 mm diameter ceramic beads, 91-PCS-CK14B, PEQLAB

Biotechnologie, Germany) in RIPA buffer (50 mM Tris, pH 8.0, 0.15M NaCl, 0.1% SDS, 1.0%

NP-40, 0.5% Na-Deoxycholate, 2 mM EDTA) supplemented with protease and phosphatase

inhibitors cocktail (cOmplete™, Mini Protease Inhibitor Cocktail, Roche). The

homogenisation was performed in Precellys 24 tissue homogenizer (Bertin Instruments) at

65 000 × g for 2 × 30 sec runs with a 30 sec break in between. Protein concentrations in the

lysates were determined by the BCA protein assay (Pierce Biotechnology, #23225) following

the manufacturer's protocol. The spectrophotometric recordings were performed using a

plate reader (Tecan, Infinite® M200 pro) at 562 nm. Volume for 20 µg of protein per sample

was calculated, to which Laemmli sample buffer (250 mM Tris-HCl pH 6.8, 10% SDS, 1.25%

Bromophenol Blue, 5% ß-mercaptoethanol, 50% glycerol) was added (1:5 dilution of

Laemmli: protein). The proteins were denatured for 5 min at 95°C, with 600 rpm in a

thermal shaker. The samples were loaded on a 12% SDS-PAGE electrophoresis gel and

separated at 120V for approximately 1h. The proteins were then transferred to PVDF

membranes (iBlot2 Transfer Stack, PVDF, Life, #IB24002X3) using iBlot2 (Invitrogen, CA,

USA) using the P0 protocol for 7 min. Membranes were blocked with 5% BSA (Sigma Aldrich,

#AG9418) in TBS (pH 8) with 0.05% Tween-20 (TBS-T) and then incubated with the

appropriate primary antibody (FTO (mouse, Abcam – ab92821, 1:5000), METTL3 (rabbit, Cell

Signalling - #51104, 1:1000), YTHDF1 (rabbit, Proteintech #17479-1-AP, 1:2500)) overnight in

5% BSA (Sigma-Aldrich, MO, USA) in TBS-T at 4°C. After three washes with TBS-T,

membranes were incubated for 2h with horseradish peroxidase (HRP) conjugated secondary

antibodies (Cytiva sheep anti-mouse IgG, 1:8000 dilution, Cytiva donkey anti-rabbit IgG,

1:8000 dilution). Following incubation, membranes were washed three times with TBS-T for

10 min each and developed in a chemiluminescence system (Fusion FX Vilber Lourmat,

Vilber, France) using chemiluminescent HRP substrate (Millipore, MA, USA). Intensities of

specific bands were normalised to a protein loading control.

RNA extraction

For brain tissue, TRIzol reagent (Invitrogen, CA, USA) and ceramic beads (1.4 mm diameter

ceramic beads, 91-PCS-CK14B, PEQLAB Biotechnologie, Germany) were added in vials with

frozen tissue and homogenisation was performed with the Precellys 24 tissue homogenizer

(Bertin Instruments) at 65 000 g for 2 x 30 sec runs with a 30 sec break in between. RNA

isolation was performed with TRIzol reagent according to the manufacturer's instructions.

Briefly, 0.5 ml of TRIzol was added to the tissue samples and incubated for 5 min before

homogenisation. After homogenisation, 0.1 ml of chloroform was added to each tube. The

tubes were shaken vigorously and incubated at RT for 2-3 min. The tubes were centrifuged

at 14 000g, for 10min at 4°C. The top aqueous layer was carefully transferred to a fresh

tube. Total RNA was then precipitated using Clean and Concentrator 5, Zymo research kit

(Zymo Research, #R1014). Following RNA extraction and precipitation, RNA was dissolved in

RNAse-free water and concentration and quality were estimated with NanoDrop 1000

spectrophotometer (Thermo Fisher Scientific, MA, USA) by measuring absorption at 260 nm

and the ratios 260/280 nm and 230/280nm respectively.

m6A fluorescence ELISA

m6A RNA methylation ELISA (m6A RNA methylation, Abcam, #ab233491) was used for the

detection of m6A RNA methylation in the cortical brain and midbrain tissue from 3 mo and

12 mo/15 mo Wt and aSyn Tg mice. For this experiment, we used cortical mouse brain

tissue from Wt mice each at 3 mo and 12 mo (six 3 mo and six 12 mo), and aSyn Tg mice (six

3 mo and five 12 mo). Midbrain tissue was used from Wt mice (three 3 mo and four 15 mo)

and aSyn Tg mice (five 3 mo and four 15 mo). This protocol ensures that the total m6A

methylation is measured in both mRNAs and small RNAs. The protocol was performed

according to the manufacturer's protocol. Briefly, total RNA isolated from the tissue was

added to bind to the assay wells. The wells were then washed to remove the unbound RNA

and diluted capture antibody was added to the wells. Unbound capture antibody was

washed out with multiple washing steps. Diluted detector antibody was then added along

with fluoro-enhancer. The wells were washed again and incubated with fluoro developer

mixture to initiate the fluorescence reaction. The fluorescence was then measured at

590nm using plate reader (Tecan, Infinite® M200 pro).

MeRiP

Total RNA was treated with DNase to remove any DNA remnants and further, 5-10 μg of

total RNA was depleted for rRNA using the RiboMinus Eukaryote Ribosomal Removal Kit

(Invitrogen). The large RNA fraction (>200 bp) was fragmented to a fragment size of

approximately 100-120 nt by RNA Fragmentation Reagent at 70°C for 5 min (#AM8740,

Thermo Fisher Scientific, Waltham, MA). 5% of this RNA sample was kept as inputs and the

rest was subjected to immunoprecipitation (IP). 300 μg of fragmented RNA was used for

each IP. 3 μg of anti-m6A antibody was incubated with the RNA IP buffer (0.2 M Tris-HCl pH

7.5, 0.5 M NaCl and 2% (vol/vol) Igepal CA-630) for 2h at 4°C with constant rotation. RNA

antibody mixes were then crosslinked twice with 0.15 J/cm2 of UV light (254 nm) in a UVP

crosslinker (Analitik Jena). Antibody-RNA conjugates were incubated with 30 μl of Protein

A/G beads overnight (ON) at 4°C with rotation in 500 μl IP buffer supplemented with 200

units. The beads with immunoprecipitated RNA were washed with IP buffer five times, and

further washed with low-salt (50 mM Tris pH 7.4, 50 mM NaCl, 1mM EDTA, 1, 0.1% NP-40,

0.1% SDS) and high-salt (same as low-salt but with 500mM NaCl) buffers at 4°C with rotation

to remove any nonspecific binding. RNA was eluted by incubating in 150 µl PKD buffer

(Qiagen) with 10 µl of Proteinase K (Millipore) for 1 hour at 37°C with agitation. Eluted RNA

was cleaned before proceeding to library preparation.

Library Preparation and Sequencing

Samples were prepared for sequencing using the SMARTer Stranded Total RNA-Seq Kit v2 —

Pico Input Mammalian (Takara) according to the manufacturer's protocol. All the RNA

obtained from the IP samples was used for library preparation, for input samples, 2 ng was

used. The libraries were amplified for a total of 12 cycles for input and 16 cycles for IP

samples. RNA-Seq samples were prepared with the TruSeq Stranded mRNA Library Prep Kit

(Illumina) according to the manufacturer's instructions for which, 50 ng of rRNA-depleted

RNA from each of the inputs was used. Prepared libraries were sequenced in a Hiseq 2000

System (Illumina) for 50 cycles in single-end reads. Additional metadata is also available via

the GEO database (GSE275432).

Bioinformatic Analysis of meRIP-seq and RNA-seq

Raw reads were processed and demultiplexed using bcl2fastq (v2.20.2), and low-quality

reads were filtered out with Cutadapt v1.11.0 [43]. Filtered reads were mapped to the

mouse (mm10) genome using the STAR aligner v2.5.2b [44]. The resulting bam files were

sorted and indexed, and the unmapped reads were removed using SAMtools v1.9.0 [45].

Methylation sites were determined using MeTPeak v1.0.0 [46], and differential methylation

(hypo- and hyper-methylated regions) was assessed with ExomePeak v2.16.0 [47] using 3

mo samples as control and 15 mo as treatment for the first comparison and for the second

comparison, Wt samples were used as controls while aSyn Tg were used as treatment. An

adjusted P value (padj, also termed as FDR [false discovery rate]) cutoff of 0.05 and FC cutoff

of 1.2 or 1.5 were used as indicated in the text. Only consistently significantly, differentially

methylated peaks were used, unless indicated.

For RNA-seq analyses, read counts were obtained with subread's featureCounts v1.5.1 [48]

from the bam files of input samples. Differential gene expression was determined by

DESeq2 v3.5.12 [49] using normalised read counts and correcting for covariates detected by

RUVseq v1.16.1 [50]. Cutoffs of padj ≤ 0.05, FC ≥ 1.2, and BaseMean ≥ 50 were applied to

the results (Supplementary Figure S1A, B). Background expressed genes were determined

for each region as those genes with a BaseMean > 50 in the corresponding input sample.

For visualization, bam files of both IP and input samples were collapsed for PCR duplicates

using SAMtools, and IP samples were normalized to their corresponding inputs and to their

library size using deeptools v3.2.1 [51] bamCompare. The resulting normalized tracks were

visualized in the IGV Browser 2.9.2 [52].

GO Analysis

GO term enrichment analyses were performed using the App ClueGO v2.5.3 [53] in

Cytoscape 3.7.2 [53], with GO Term Fusion enabled to collapse terms containing very similar

gene lists and using a custom background corresponding to expressed genes in the

corresponding species as obtained from RNA-seq results of the corresponding input samples

of the meRIP experiments. GO term tables for biological processes, cellular components,

pathways, and KEGG were produced and are labelled accordingly in the figures. The

resulting enriched GO terms were visualized with a custom script using ggplot2 v3.3.5 [54]

playing the adjusted p-value (padj) for the GO term, the number of genes from the list that

belong to said term, and the percentage of the total genes in the GO term that are present

in the list. Synaptic GO enrichment analyses were performed with SynGO (v1.1,

syngoportal.org) [55].

Additional Bioinformatics Packages and Tools

Scripts and analysis pipelines were written in R (3.5.2) [56]. Peak annotation was performed

with Homer v4.10.4 [57] and Annotatr v1.8.0 [58]. Guitar plots were produced with the

Guitar v1.20.1 [59] R package. Volcano plots were generated with plot.ly/orca v4.9.4.1 [60]

(Supplementary Figure S1C). Area-proportional Venn diagrams were produced with BioVenn

(www.biovenn.nl), and multiple list comparisons performed with Intervene / UpSet

(asntech.shinyapps.io/intervene/). Mouse/human homologues were determined by their

annotation in NCBI's HomoloGene database using the HomoloGene (v1.4.68.19.3.27) R

package. Odds ratios and p values to determine significance in overlapped datasets were

calculated with the GeneOverlap R package v1.18.0 [61]. De novo motif analyses were

performed with Homer's findMotifsGenome, and motifs containing the DRACH consensus

sequence out of the top 10 most significant are dipcasplayed. KEGG pathway enrichment

was produced with KEGG Mapper (www.genome.jp/kegg/mapper/) [62]. Microscopy

images were pre-processed with Fiji, and quantification was automated in Cell Profiler

(cellprofiler.org) [63]. Graphs, heat maps, and statistical analyses were performed on

GraphPad Prism version 9.3.1 for Mac.

Microscopy

Images of the immunofluorescence-stained neurons (three experiments in total, with at

least 10 regions of interest acquired per experiment) and brain tissue were acquired using a

confocal microscope ZEISS LSM900. The images for the neuronal cell culture were acquired

as a z-stack using 63x magnification with Airyscan mode. Brain slices were imaged at 40x

magnification, with tiles in confocal mode.

Statistical analysis

The cell culture images for m6A regulators and their colocalisation at the synapse were

analysed using FIJI with the plugin JaCoP, while the synapse counting was performed using

IMARIS. IMARIS analysis was performed following the protocol from the Queensland Brain

Institute. The protein signals were further classified inside the dendritic or outside using

membranes based on the MAP2 signals. The signals were then divided into four groups,

representing PSD95 inside the neuron, PSD95 outside the neuron, Synaptotagmin1 (Syn1)

inside the neuron and Syn1 outside the neuron. When the signal for PSD95 inside the

neuron was at < 500µm from the signal for Syn1, it was considered a synapse. Colocalisation

of synaptic markers with m6A regulators was estimated using Pearson's correlation

coefficient (PCC) estimated using the JaCoP plugin in ImageJ, FIJI. All IHC images were also

analysed for their expression levels using FIJI software. Data analysis was performed using

the GraphPad software version 9 for Windows (GraphPad Software, La Jolla California

USA, https://www.graphpad.com). For group comparisons, one-way ANOVA with Tukey's

test was used, while comparisons of two groups of means were done with an unpaired

student's t-test. All data are expressed as mean ± SD. Differences are considered significant

with $p < 0.05$ (*$p < 0.05$, **<0.01, ***<0.001).

Declarations

Ethics approval

Animal procedures were performed in accordance with the European Community (Directive

2010/63/EU); and institutional and national guidelines under the Lower Saxony State Office

for Consumer Protection and Food Safety (LAVES) (license number 19.3213).

Consent for publication

All authors approved the publication of the article.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the

corresponding author upon reasonable request. All raw data files for meRIPseq are available

on the GEO database (GSE275432).

Competing interests

The authors declare that they have no competing interests.

Funding

TFO and AF are supported by the Deutsche Forschungsgemeinschaft (DFG, German

Research Foundation) under Germany's Excellence Strategy - EXC 2067/1- 390729940), and

by SFB1286 (Projects B6 and B8). AF is supported by the Niedersächsisches Vorab:

Forschungskooperation Niedersachsen – Israel (ZN3460).

Authors' contributions

AC performed, analysed, and interpreted the data from cell cultures, ELISA, IHC, and

western blot, and interpreted the meRIP results. MM and VG performed experiments in

animals. MX interpreted the data for meRIP analysis and was a major contribution in writing

the manuscript. YF performed and analysed ICC data. RCH performed the meRIP and

subsequent data analysis. AC and TO wrote the manuscript. All authors read and approved

the final manuscript.

Figure Legends

Figure 1. m6A modification landscape in ageing and aSyn Tg mice. (A) Bar plots representing the variations in the differentially expressed genes and differentially methylated genes in the midbrain from aSyn Tg mice than Wt mice when 3mo mice are compared with 15mo applying equal cutoffs for FC and adjusted P value (FC > 1.2 and padj ≤ 0.05). Left: with age, right: with aSyn (B) Bar plots representing the effects of aSyn pathology on differential expression and differential methylation in 3mo and 15mo Wt vs. aSyn Tg mice. (C) Overview of protocols utilized in this project. (D) Pie charts representing the overview of the distribution of hypomethylated, hypermethylated and mixed transcripts as an effect of aSyn pathology and ageing. (E) Venn diagram representing the number of differentially methylated genes involved in young and old Wt and aSyn Tg mice. It also provides a summary of the overlapping genes across different genotypes. (F) Guitar plots representing the distribution of m6A methylation on mRNA and lncRNA. CDS stands for coding sequence, and UTR stands for untranslated region.

Figure 2. Gene ontology (GO) analysis for 3 mo and 15 mo Wt vs. aSyn Tg mice. (A) GO analysis for the biological processes and cellular processes in 15 mo aSyn Tg vs. Wt mice, FDR = 1.5. (B) Sunburst plots representing GO analysis for the genes involved in the processes at the synapse in aSyn Tg vs. Wt mice at 3 mo. Graphs created with syngoportal.org. (C) Sunburst plots representing GO analysis for the genes involved in the processes at the synapse in aSyn Tg vs. Wt mice aged mice at 15 mo. Graphs created with syngoportal.org.

Figure 3. Levels of m6A RNA and m6A regulators in mouse brain tissue. (A) Bar plots

representing the levels of m6A modification on total RNA using ELISA from cortical and

midbrain regions of Wt and aSyn Tg mice. A significant reduction in the levels of m6A was

observed in the aged, 12 mo aSyn Tg mouse cortex when compared with 12 mo Wt mice (*,

ajd P value = 0.045). No significant differences were observed in the midbrain region of the

brain. (B) Biochemical analysis using western blot comparing the expression levels of

METTL3, YTHDF1 and FTO in the cortical tissue from 3 mo, 6mo and 12 mo Wt vs. aSyn Tg

mice. Scatter plot showing no significant difference in the expression level when comparing

Wt with aSyn Tg mice.

Figure 4. Distribution of m6A regulatory proteins METTL3, YTHDF1 and FTO in the cortex,

striatum, hippocampus and cerebellum of mouse brain. (A) Representative images of

mouse brain tissue slices (30µm) fixed, and stained with MAP2, METTL3, YTHDF1 and FTO.

The images are acquired using a confocal microscope, 40x magnification with tiles, and scale

bar = 50µm. White arrows point to the neurons with the protein expression. (B) Bar plots

showing the comparison between the mean intensity of METTL3, YTHDF1 and FTO in 15mo

Wt vs. aSyn Tg mice.

Figure 5. METTL3 is localised at the post-synapse in Wt and aSyn Tg DIV14 primary

neurons. (A) Representative images of DIV14 primary neurons, fixed and stained with

Histone H1, MAP2, PSD95, Synaptotagmin 1, METTL3, YTHDF1, and FTO in Wt and aSyn Tg

neurons. Images are acquired using airy-scan at 63x magnification, scale bar = 10μm. (B) Box

and whisker plot representing the colocalisation of m6A regulators in Wt and aSyn Tg D-14

primary neurons at the pre-and post-synapse. METTL3 colocalises with the post-synapse,

(Pearson's correlation coefficient, Wt = 0.810, aSyn Tg = 0.748). A significant difference (*,

adj p value = 0.018) was observed between METTL3 at the post-synapse in Wt neurons

when compared with aSyn Tg neurons. (C) Box and whisker plot representing the

colocalisation of m6A regulators at the pre-and post-synapse in Wt and aSyn Tg primary

neurons. METTL3 colocalisation is significantly reduced in aSyn Tg neurons when compared

with Wt at the post-synapse

Figure 6. The number of synapses is increased in aSyn Tg primary neurons. (A)

Representative immunofluorescent images of DIV14 primary neurons stained with pre-

(Syn1, magenta) and post-synaptic (PSD95, green) markers and MAP2 in red. Acquired at

63x magnification, airy scan mode. Scale bar = 10μm. (B) Scatter plot for synapse counting

for Wt and aSyn Tg mouse models with a significant reduction in the number of synapses (*,

adj p value = 0.044) in aSyn Tg primary neurons when compared with Wt.

Supplementary Figures

Figure S1. Gene expression analysis in aSyn Tg and Wt mice. (A) Principle component

analysis (PCA) plot of total Wt vs. aSyn Tg mice (PC1 = 40.5%, PC2 = 19.2%). (B) PCA plot of

aSyn Tg – 3 mo and 15 mo mice, showing a clear segregation (PC1 = 49.8%, PC2 = 23.6%). (C)

Volcano plot showing that the expression of most of the genes is down regulated in aSyn Tg

mice, due to ageing (FDR < 0.05, abs(logFC) >1), aSyn Tg 15 mo vs. 3mo.

Figure S2. Differentially expressed gene pathways in aSyn Tg 15 mo vs. 3 mo. GO term

analysis comparing the enrichment of pathways, molecular function, cellular components

and biological processes of differentially methylated genes. Most pathway enrichment was

observed in immune system and immune response linked pathways including –

immunoglobulin binding, phagocytic vesicle, defense response and innate and adaptive

immune response, FC > 1.2, FDR = 1.5.

References

1. Allis D. No Title. Epigenetics, second edition [Internet]. 2015. Available from: chrome-

extension://efaidnbmnnnibpcajpcglclefindmkaj/https://www.cshlpress.com/pdf/sample/2014/epige

netics2/EPIFM.pdf

2. Goldberg AD, Allis CD, Bernstein E. Epigenetics: A Landscape Takes Shape. Cell. 2007;128:635–8.

3. Saletore Y, Meyer K, Korlach J, Vilfan ID, Jaffrey S, Mason CE. The birth of the Epitranscriptome:

deciphering the function of RNA modifications. Genome Biol [Internet]. 2012;13:175. Available from:

http://genomebiology.biomedcentral.com/articles/10.1186/gb-2012-13-10-175

4. Dominissini D, Moshitch-Moshkovitz S, Schwartz S, Salmon-Divon M, Ungar L, Osenberg S, et al.

Topology of the human and mouse m6A RNA methylomes revealed by m6A-seq. Nature [Internet].

2012;485:201–6. Available from: https://www.nature.com/articles/nature11112

5. Liu J, Yue Y, Han D, Wang X, Fu Y, Zhang L, et al. A METTL3–METTL14 complex mediates

mammalian nuclear RNA N6-adenosine methylation. Nat Chem Biol [Internet]. 2014;10:93–5.

Available from: https://www.nature.com/articles/nchembio.1432

6. Ke S, Alemu EA, Mertens C, Gantman EC, Fak JJ, Mele A, et al. A majority of m 6 A residues are in

the last exons, allowing the potential for 3⬜ UTR regulation. Genes Dev [Internet]. 2015;29:2037–53.

Available from: http://genesdev.cshlp.org/lookup/doi/10.1101/gad.269415.115

7. Liu H, Begik O, Lucas MC, Ramirez JM, Mason CE, Wiener D, et al. Accurate detection of m6A RNA

modifications in native RNA sequences. Nat Commun [Internet]. 2019;10:4079. Available from:

https://www.nature.com/articles/s41467-019-11713-9

8. Shi H, Wei J, He C. Where, When, and How: Context-Dependent Functions of RNA Methylation

Writers, Readers, and Erasers. Mol Cell [Internet]. 2019;74:640–50. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S109727651930317X

9. Yang Y, Hsu PJ, Chen Y-S, Yang Y-G. Dynamic transcriptomic m6A decoration: writers, erasers,

readers and functions in RNA metabolism. Cell Res [Internet]. 2018;28:616–24. Available from:

https://www.nature.com/articles/s41422-018-0040-8

10. Zhao BS, Wang X, Beadell A V., Lu Z, Shi H, Kuuspalu A, et al. m6A-dependent maternal mRNA

clearance facilitates zebrafish maternal-to-zygotic transition. Nature [Internet]. 2017;542:475–8.

Available from: https://www.nature.com/articles/nature21355

11. Dinescu S, Ignat S, Lazar AD, Constantin C, Neagu M, Costache M. Epitranscriptomic Signatures in

lncRNAs and Their Possible Roles in Cancer. Genes (Basel) [Internet]. 2019;10:52. Available from:

https://www.mdpi.com/2073-4425/10/1/52

12. Licht K, Jantsch MF. Rapid and dynamic transcriptome regulation by RNA editing and RNA

modifications. Journal of Cell Biology [Internet]. 2016;213:15–22. Available from:

https://rupress.org/jcb/article/213/1/15/38525/Rapid-and-dynamic-transcriptome-regulation-by-

RNA

13. Dominissini D, Moshitch-Moshkovitz S, Salmon-Divon M, Amariglio N, Rechavi G. Transcriptome-

wide mapping of N6-methyladenosine by m6A-seq based on immunocapturing and massively

parallel sequencing. Nat Protoc [Internet]. 2013;8:176–89. Available from:

https://www.nature.com/articles/nprot.2012.148

14. Bokar JA, Shambaugh ME, Polayes D, Matera AG, Rottman FM. Purification and cDNA cloning of

the AdoMet-binding subunit of the human mRNA (N6-adenosine)-methyltransferase. RNA [Internet].

1997;3:1233–47. Available from: http://www.ncbi.nlm.nih.gov/pubmed/9409616

15. Schwartz S, Mumbach MR, Jovanovic M, Wang T, Maciag K, Bushkin GG, et al. Perturbation of

m6A Writers Reveals Two Distinct Classes of mRNA Methylation at Internal and 5⬚ Sites. Cell Rep

[Internet]. 2014;8:284–96. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S2211124714004422

16. Pendleton KE, Chen B, Liu K, Hunter O V., Xie Y, Tu BP, et al. The U6 snRNA m 6 A

Methyltransferase METTL16 Regulates SAM Synthetase Intron Retention. Cell [Internet].

2017;169:824-835.e14. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0092867417305305

17. Patil DP, Chen C-K, Pickering BF, Chow A, Jackson C, Guttman M, et al. m6A RNA methylation

promotes XIST-mediated transcriptional repression. Nature [Internet]. 2016;537:369–73. Available

from: https://www.nature.com/articles/nature19342

18. Wen J, Lv R, Ma H, Shen H, He C, Wang J, et al. Zc3h13 Regulates Nuclear RNA m6A Methylation

and Mouse Embryonic Stem Cell Self-Renewal. Mol Cell [Internet]. 2018;69:1028-1038.e6. Available

from: https://linkinghub.elsevier.com/retrieve/pii/S1097276518301345

19. Agarwala SD, Blitzblau HG, Hochwagen A, Fink GR. RNA Methylation by the MIS Complex

Regulates a Cell Fate Decision in Yeast. PLoS Genet. 2012;8:e1002732.

20. Li Q, Li X, Tang H, Jiang B, Dou Y, Gorospe M, et al. NSUN2-Mediated m5C Methylation and

METTL3/METTL14-Mediated m6A Methylation Cooperatively Enhance p21 Translation. J Cell

Biochem [Internet]. 2017;118:2587–98. Available from:

https://onlinelibrary.wiley.com/doi/10.1002/jcb.25957

21. Xu K, Yang Y, Feng G-H, Sun B-F, Chen J-Q, Li Y-F, et al. Mettl3-mediated m6A regulates

spermatogonial differentiation and meiosis initiation. Cell Res [Internet]. 2017;27:1100–14. Available

from: https://www.nature.com/articles/cr2017100

22. Aguilo F, Zhang F, Sancho A, Fidalgo M, Di Cecilia S, Vashisht A, et al. Coordination of m 6 A

mRNA Methylation and Gene Transcription by ZFP217 Regulates Pluripotency and Reprogramming.

Cell Stem Cell [Internet]. 2015;17:689–704. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S1934590915004129

23. Hess ME, Hess S, Meyer KD, Verhagen LAW, Koch L, Brönneke HS, et al. The fat mass and obesity

associated gene (Fto) regulates activity of the dopaminergic midbrain circuitry. Nat Neurosci.

2013;16:1042–8.

24. Ping X-L, Sun B-F, Wang L, Xiao W, Yang X, Wang W-J, et al. Mammalian WTAP is a regulatory

subunit of the RNA N6-methyladenosine methyltransferase. Cell Res [Internet]. 2014;24:177–89.

Available from: https://www.nature.com/articles/cr20143

25. Wang X, Lu Z, Gomez A, Hon GC, Yue Y, Han D, et al. N6-methyladenosine-dependent regulation

of messenger RNA stability. Nature [Internet]. 2014;505:117–20. Available from:

https://www.nature.com/articles/nature12730

26. Śledź P, Jinek M. Structural insights into the molecular mechanism of the m6A writer complex.

Elife [Internet]. 2016;5. Available from: https://elifesciences.org/articles/18434

27. Wang X, Feng J, Xue Y, Guan Z, Zhang D, Liu Z, et al. Structural basis of N6-adenosine methylation

by the METTL3–METTL14 complex. Nature. 2016;534:575–8.

28. Moindrot B, Cerase A, Coker H, Masui O, Grijzenhout A, Pintacuda G, et al. A Pooled shRNA

Screen Identifies Rbm15, Spen, and Wtap as Factors Required for Xist RNA-Mediated Silencing. Cell

Rep [Internet]. 2015;12:562–72. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S2211124715006798

29. Jia G, Fu Y, Zhao X, Dai Q, Zheng G, Yang Y, et al. N6-Methyladenosine in nuclear RNA is a major

substrate of the obesity-associated FTO. Nat Chem Biol [Internet]. 2011;7:885–7. Available from:

https://www.nature.com/articles/nchembio.687

30. Zheng G, Dahl JA, Niu Y, Fedorcsak P, Huang C-M, Li CJ, et al. ALKBH5 Is a Mammalian RNA

Demethylase that Impacts RNA Metabolism and Mouse Fertility. Mol Cell [Internet]. 2013;49:18–29.

Available from: https://linkinghub.elsevier.com/retrieve/pii/S1097276512008921

31. Fu Y, Jia G, Pang X, Wang RN, Wang X, Li CJ, et al. FTO-mediated formation of N6-

hydroxymethyladenosine and N6-formyladenosine in mammalian RNA. Nat Commun [Internet].

2013;4:1798. Available from: https://www.nature.com/articles/ncomms2822

32. Alarcón CR, Lee H, Goodarzi H, Halberg N, Tavazoie SF. N6-methyladenosine marks primary

microRNAs for processing. Nature [Internet]. 2015;519:482–5. Available from:

https://www.nature.com/articles/nature14281

33. Zaccara S, Jaffrey SR. A Unified Model for the Function of YTHDF Proteins in Regulating m6A-

Modified mRNA. Cell [Internet]. 2020;181:1582-1595.e18. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0092867420305742

34. Chen J, Wei X, Yi X, Jiang D-S. RNA Modification by m6A Methylation in Cardiovascular Disease.

Ghose J, editor. Oxid Med Cell Longev [Internet]. 2021;2021:1–13. Available from:

https://www.hindawi.com/journals/omcl/2021/8813909/

35. Jiang Z, Wang Y, Li Z, Dai Z, He Y, Chu K, et al. The m6A mRNA demethylase FTO in granulosa cells

retards FOS-dependent ovarian aging. Cell Death Dis [Internet]. 2021;12:744. Available from:

https://www.nature.com/articles/s41419-021-04016-9

36. Shafik AM, Zhang F, Guo Z, Dai Q, Pajdzik K, Li Y, et al. N6-methyladenosine dynamics in

neurodevelopment and aging, and its potential role in Alzheimer's disease. Genome Biol.

2021;22:17.

37. Castro-Hernández R, Berulava T, Metelova M, Epple R, Peña Centeno T, Richter J, et al.

Conserved reduction of m ^{6} A RNA modifications during aging and neurodegeneration is linked to

changes in synaptic transcripts. Proceedings of the National Academy of Sciences. 2023;120.

38. Chen YG, Chen R, Ahmad S, Verma R, Kasturi SP, Amaya L, et al. N6-Methyladenosine

Modification Controls Circular RNA Immunity. Mol Cell [Internet]. 2019;76:96-109.e9. Available

from: https://linkinghub.elsevier.com/retrieve/pii/S1097276519305465

39. Yu Z, Huang L, Xia Y, Cheng S, Yang C, Chen C, et al. Analysis of m6A modification regulators in

the substantia nigra and striatum of MPTP-induced Parkinson's disease mice. Neurosci Lett

[Internet]. 2022;791:136907. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0304394022004682

40. Schapira AHV, Cooper JM, Dexter D, Jenner P, Clark JB, Marsden CD. MITOCHONDRIAL COMPLEX

I DEFICIENCY IN PARKINSON'S DISEASE. The Lancet [Internet]. 1989;333:1269. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0140673689923660

41. Krishnan KJ, Reeve AK, Samuels DC, Chinnery PF, Blackwood JK, Taylor RW, et al. What causes

mitochondrial DNA deletions in human cells? Nat Genet [Internet]. 2008;40:275–9. Available from:

https://www.nature.com/articles/ng.f.94

42. Pickrell AM, Pinto M, Hida A, Moraes CT. Striatal Dysfunctions Associated with Mitochondrial

DNA Damage in Dopaminergic Neurons in a Mouse Model of Parkinson's Disease. The Journal of

Neuroscience [Internet]. 2011;31:17649–58. Available from:

https://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.4871-11.2011

43. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet J [Internet]. 2011;17:10. Available from:

http://journal.embnet.org/index.php/embnetjournal/article/view/200

44. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-

seq aligner. Bioinformatics [Internet]. 2013;29:15–21. Available from:

https://academic.oup.com/bioinformatics/article/29/1/15/272537

45. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map

format and SAMtools. Bioinformatics [Internet]. 2009;25:2078–9. Available from:

https://academic.oup.com/bioinformatics/article/25/16/2078/204688

46. Cui X, Meng J, Zhang S, Chen Y, Huang Y. A novel algorithm for calling mRNA m 6 A peaks by

modeling biological variances in MeRIP-seq data. Bioinformatics [Internet]. 2016;32:i378–85.

Available from: https://academic.oup.com/bioinformatics/article/32/12/i378/2289056

47. Meng J, Cui X, Rao MK, Chen Y, Huang Y. Exome-based analysis for RNA epigenome sequencing

data. Bioinformatics [Internet]. 2013;29:1565–7. Available from:

https://academic.oup.com/bioinformatics/article/29/12/1565/291595

48. Liao Y, Smyth GK, Shi W. The Subread aligner: fast, accurate and scalable read mapping by seed-

and-vote. Nucleic Acids Res [Internet]. 2013;41:e108–e108. Available from:

https://academic.oup.com/nar/article/41/10/e108/1075719

49. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq

data with DESeq2. Genome Biol [Internet]. 2014;15:550. Available from:

http://genomebiology.biomedcentral.com/articles/10.1186/s13059-014-0550-8

50. Risso D, Ngai J, Speed TP, Dudoit S. Normalization of RNA-seq data using factor analysis of

control genes or samples. Nat Biotechnol [Internet]. 2014;32:896–902. Available from:

https://www.nature.com/articles/nbt.2931

51. Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, et al. deepTools2: a next

generation web server for deep-sequencing data analysis. Nucleic Acids Res [Internet].

2016;44:W160–5. Available from: https://academic.oup.com/nar/article-

lookup/doi/10.1093/nar/gkw257

52. Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, et al. Integrative

genomics viewer. Nat Biotechnol [Internet]. 2011;29:24–6. Available from:

https://www.nature.com/articles/nbt.1754

53. Bindea G, Mlecnik B, Hackl H, Charoentong P, Tosolini M, Kirilovsky A, et al. ClueGO: a Cytoscape

plug-in to decipher functionally grouped gene ontology and pathway annotation networks.

Bioinformatics. 2009;25:1091–3.

54. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: A Software

Environment for Integrated Models of Biomolecular Interaction Networks. Genome Res [Internet].

2003;13:2498–504. Available from: http://genome.cshlp.org/lookup/doi/10.1101/gr.1239303

55. Wickham H. Toolbox. ggplot2. New York, NY: Springer New York; 2009. p. 65–90.

56. Koopmans F, van Nierop P, Andres-Alonso M, Byrnes A, Cijsouw T, Coba MP, et al. SynGO: An

Evidence-Based, Expert-Curated Knowledge Base for the Synapse. Neuron. 2019;103:217-234.e4.

57. R Development Core Team. R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing [Internet]. 2015. Available from: http://www.r-project.org

58. Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, et al. Simple Combinations of Lineage-

Determining Transcription Factors Prime cis-Regulatory Elements Required for Macrophage and B

Cell Identities. Mol Cell [Internet]. 2010;38:576–89. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S1097276510003667

59. Cavalcante RG, Sartor MA. annotatr: genomic regions in context. Valencia A, editor.

Bioinformatics [Internet]. 2017;33:2381–3. Available from:

https://academic.oup.com/bioinformatics/article/33/15/2381/3092365

60. Cui X, Wei Z, Zhang L, Liu H, Sun L, Zhang S-W, et al. Guitar: An R/Bioconductor Package for Gene

Annotation Guided Transcriptomic Analysis of RNA-Related Genomic Features. Biomed Res Int

[Internet]. 2016;2016:1–8. Available from: http://www.hindawi.com/journals/bmri/2016/8367534/

61. Sievert C. Interactive Web-Based Data Visualization with R, plotly, and shiny. Chapman and

Hall/CRC; 2020.

62. L Shen. C. Sievert, Interactive Web-Based Data Visualization with R, Plotly, and Shiny (Champ

Hall, 2019). 2014.

63. Kanehisa M, Sato Y, Kawashima M. <scp>KEGG</scp> mapping tools for uncovering hidden

features in biological data. Protein Science. 2022;31:47–53.

64. Jones TR, Kang IH, Wheeler DB, Lindquist RA, Papallo A, Sabatini DM, et al. CellProfiler Analyst:

data exploration and analysis software for complex image-based screens. BMC Bioinformatics

[Internet]. 2008;9:482. Available from:

https://bmcbioinformatics.biomedcentral.com/articles/10.1186/1471-2105-9-482

65. Xylaki M, Paiva I, Al-Azzani M, Gerhardt E, Jain G, Islam MR, et al. miR-101a-3p Impairs Synaptic

Plasticity and Contributes to Synucleinopathy. J Parkinsons Dis. 2023;13:179–96.

66. Koopmans F, van Nierop P, Andres-Alonso M, Byrnes A, Cijsouw T, Coba MP, et al. SynGO: An

Evidence-Based, Expert-Curated Knowledge Base for the Synapse. Neuron. 2019;103:217-234.e4.

67. Stern S, Lau S, Manole A, Rosh I, Percia MM, Ben Ezer R, et al. Reduced synaptic activity and

dysregulated extracellular matrix pathways in midbrain neurons from Parkinson's disease patients.

NPJ Parkinsons Dis. 2022;8:103.

68. Rosh I, Tripathi U, Hussein Y, Rike WA, Djamus J, Shklyar B, et al. Synaptic dysfunction and

extracellular matrix dysregulation in dopaminergic neurons from sporadic and E326K-GBA1

Parkinson's disease patients. NPJ Parkinsons Dis. 2024;10:38.

69. Licatalosi DD, Mele A, Fak JJ, Ule J, Kayikci M, Chi SW, et al. HITS-CLIP yields genome-wide

insights into brain alternative RNA processing. Nature. 2008;456:464–9.

70. Chen X, Yu C, Guo M, Zheng X, Ali S, Huang H, et al. Down-Regulation of m6A mRNA Methylation

Is Involved in Dopaminergic Neuronal Death. ACS Chem Neurosci. 2019;10:2355–63.

71. Geng Y, Long X, Zhang Y, Wang Y, You G, Guo W, et al. FTO-targeted siRNA delivery by MSC-

derived exosomes synergistically alleviates dopaminergic neuronal death in Parkinson's disease via

m6A-dependent regulation of ATM mRNA. J Transl Med. 2023;21:652.

72. He H, Zhang Q, Liao J, Lei J, Luo M, Huang J, et al. METTL14 is decreased and

regulates m^{6}A modification of α-synuclein in Parkinson's disease. J Neurochem.

2023;166:609–22.

73. Soukup S, Vanhauwaert R, Verstreken P. Parkinson's disease: convergence on synaptic

homeostasis. EMBO J. 2018;37.

74. Liu-Yesucevitz L, Bassell GJ, Gitler AD, Hart AC, Klann E, Richter JD, et al. Local RNA Translation at

the Synapse and in Disease: Figure 1. The Journal of Neuroscience. 2011;31:16086–93.

75. Sun J, Cheng B, Su Y, Li M, Ma S, Zhang Y, et al. The Potential Role of m6A RNA Methylation in the

Aging Process and Aging-Associated Diseases. Front Genet. 2022;13.

76. Martinez De La Cruz B, Gell C, Markus R, Macdonald I, Fray R, Knight HM. m^{6}A mRNA

methylation in human brain is disrupted in Lewy body disorders. Neuropathol Appl Neurobiol.

2023;49.

77. Yu Z, Huang L, Xia Y, Cheng S, Yang C, Chen C, et al. Analysis of m6A modification regulators in

the substantia nigra and striatum of MPTP-induced Parkinson's disease mice. Neurosci Lett.

2022;791:136907.

78. Harper JE, Miceli SM, Roberts RJ, Manley JL. Sequence specificity of the human mRNA N6-

adenosine methylase *in vitro*. Nucleic Acids Res. 1990;18:5735–41.

79. Bokar JA, Shambaugh ME, Polayes D, Matera AG, Rottman FM. Purification and cDNA cloning of

the AdoMet-binding subunit of the human mRNA (N6-adenosine)-methyltransferase. RNA.

1997;3:1233–47.

80. Ping X-L, Sun B-F, Wang L, Xiao W, Yang X, Wang W-J, et al. Mammalian WTAP is a regulatory

subunit of the RNA N6-methyladenosine methyltransferase. Cell Res. 2014;24:177–89.

81. Zou Z, Sepich-Poore C, Zhou X, Wei J, He C. The mechanism underlying redundant functions of

the YTHDF proteins. Genome Biol. 2023;24:17.

82. Gulati P, Avezov E, Ma M, Antrobus R, Lehner P, O'Rahilly S, et al. Fat mass and obesity-related

(FTO) shuttles between the nucleus and cytoplasm. Biosci Rep. 2014;34.

83. Goers J, Manning-Bog AB, McCormack AL, Millett IS, Doniach S, Di Monte DA, et al. Nuclear

Localization of α-Synuclein and Its Interaction with Histones. Biochemistry. 2003;42:8465–71.

84. Sugeno N, Jäckel S, Voigt A, Wassouf Z, Schulze-Hentrich J, Kahle PJ. α-Synuclein enhances

histone H3 lysine-9 dimethylation and H3K9me2-dependent transcriptional responses. Sci Rep.

2016;6:36328.

85. Surguchov A. α-Synuclein and Mechanisms of Epigenetic Regulation. Brain Sci. 2023;13:150.

86. Jowaed A, Schmitt I, Kaut O, Wüllner U. Methylation Regulates Alpha-Synuclein Expression and Is

Decreased in Parkinson's Disease Patients' Brains. The Journal of Neuroscience. 2010;30:6355–9.

87. Merkurjev D, Hong W-T, Iida K, Oomoto I, Goldie BJ, Yamaguti H, et al. Synaptic N6-

methyladenosine (m6A) epitranscriptome reveals functional partitioning of localized transcripts. Nat

Neurosci. 2018;21:1004–14.

88. Flamand MN, Meyer KD. The epitranscriptome and synaptic plasticity. Curr Opin Neurobiol.

2019;59:41–8.

89. Madugalle SU, Meyer K, Wang DO, Bredy TW. RNA N6-Methyladenosine and the Regulation of

RNA Localization and Function in the Brain. Trends Neurosci. 2020;43:1011–23.

90. Flamand MN, Meyer KD. m6A and YTHDF proteins contribute to the localization of select

neuronal mRNAs. Nucleic Acids Res. 2022;50:4464–83.

91. Picconi B, Piccoli G, Calabresi P. Synaptic Dysfunction in Parkinson's Disease. 2012. p. 553–72.

1012

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.