The epitranscriptomic m6A RNA modification modulates synaptic function in ageing and in a mouse model of synucleinopathy

Avika Chopra[1], Mary Xylaki[1,2], Fanzheng Yin[1], Ricardo Castro-Hernández[3], Madiha Merghani[1], Valentina Grande[1], Brit Mollenhauer[2,4], André Fischer[3,5,6], and Tiago F. Outeiro[1,7,8,9*]

1. University Medical Center Göttingen, Department of Experimental Neurodegeneration, Center for Biostructural Imaging of Neurodegeneration, 37073, Göttingen, Germany.

2. Department of Neurology, University Medical Centre Göttingen, 37073, Göttingen, Germany.

3. Department for Systems Medicine and Epigenetics, German Center for Neurodegenerative Diseases (DZNE), Göttingen, Germany

4. Paracelsus Elena Klinik, Kassel, Germany

5. Department of Psychiatry and Psychotherapy, University Medical Center Göttingen, Göttingen, Germany

6. Cluster of Excellence "Multiscale Bioimaging: from Molecular Machines to Networks of Excitable Cells" (MBExC), University of Göttingen, Germany

7. Max Planck Institute for Multidisciplinary Sciences, 37075 Göttingen, Germany.

8. Translational and Clinical Research Institute, Faculty of Medical Sciences, Newcastle University, Framlington Place, Newcastle Upon Tyne, NE2 4HH, UK.

9. Scientific employee with an honorary contract at Deutsches Zentrum für Neurodegenerative Erkrankungen (DZNE), 37075, Göttingen, Germany.

25   ***Correspondence to:** Tiago F. Outeiro

26   Department of Experimental Neurodegeneration, Centre for Biostructural Imaging of

27   Neurodegeneration, University Medical Centre Göttingen, Göttingen, Germany

28   Email: tiago.outeiro@med.uni-goettingen.de

Abstract

N6-methyladenosine (m6A) is the most abundant and conserved transcriptional modification in eukaryotic RNA, regulating RNA fate. While the functions of m6A in the development of the mammalian brain have been extensively studied, its roles in synaptic plasticity, cognitive decline, motor function, or other brain circuits remain underexplored. To date, the role of this modification in Parkinson's disease (PD) and other synucleinopathies has been largely unknown. Here, we investigated the m6A epitranscriptome in a mouse model of synucleinopathy. We performed m6A RNA immunoprecipitation sequencing (meRIP-seq) to obtain the m6A epitranscriptome of the midbrain in young (3 mo) and aged (15 mo) A30P-aSyn transgenic mice (aSyn Tg) and C57BL6 control wild type (Wt) mice. We observed hypermethylation of synaptic genes in 3 mo aSyn Tg mice compared to age-matched Wt mice. This methylation was reduced during ageing, with synaptic genes becoming increasingly hypomethylated. Using immunofluorescence imaging alongside biochemical analysis, we further investigated the expression of m6A regulatory enzymes — writer, N6-Adenosine-Methyltransferase Complex Catalytic Subunit (METTL3); reader, YTH N6-methyladenosine RNA-binding protein (YTHDF1); and eraser, fat mass and obesity-associated protein (FTO) — in the cortex, striatum, hippocampus, and cerebellum of Wt and aSyn Tg mice, as well as in primary cortical neuronal cultures. We observed that the levels of METTL3, YTHDF1 and FTO were similar between Wt and aSyn Tg mice. Interestingly, the writer protein METTL3 was found in both the nucleus and in the post-synaptic compartment in neuronal cultures. Our findings suggest that alterations in the regulation of m6A RNA methylation may be associated with neurodegeneration and ageing and that this level of epitranscriptomic regulation plays a significant role at the synapse.

53   **Keywords:** epigenetics, m6A, Parkinson's disease, ageing, METTL3

Introduction

Epigenetics is a process that acts as a bridge between genotype and phenotype, changing the outcome of a gene without modifying the DNA sequence. It encompasses various types of alterations such as methylation of DNA, histone modifications, and chromatin remodeling [1,2]. The field that studies posttranscriptional changes of RNA bases is known as epitranscriptomics and aims to identify modifications in the transcriptome and unravel their regulatory roles [3]. N^6-methyladenine (m6A) modification is the most prevalent mRNA modification occurring at a frequency of 0.15 - 0.6% of all adenosines [4–6] and is found in different types of RNAs including protein coding, messenger RNA (mRNA) [7,8]. This modification is being heavily studied due to its prevalence and importance in eukaryotes, where it regulates various cellular functions [8–12]. Methylation marks are mostly added co-transcriptionally to specific consensus motif sequences and are mostly found close to the 3' un-translated region (3'-UTR) at the stop codon [4,13].

The m6A modification machinery consists of three groups of enzymes, namely methyltransferases, demethylases and detectors of this modification, also termed writers, erasers, and readers, respectively. The modification can be added onto the RNA by methyltransferases including methyltransferase-like 3 (METTL3), methyltransferase-like 14 (METTL14), and adapter protein Wilms tumour 1-associated protein (WTAP) [5,14–19]. METTL3 was the first discovered methyltransferase and was described as a key component in the methylation of transcripts involved in physiological processes such as embryonic development and brain development [20–23]. The catalytically active part of METTL3 forms a heterodimer with METTL14, which provides the structure to install the m6A modification onto RNA [5,24–27]. Apart from these, the m6A methyltransferase complex comprises other proteins, including RNA binding proteins (RBP) like RBP15, which recruit this machinery to

the target mRNA to facilitate methylation [15,28]. Demethylases, for instance, fat mass and

obesity-associated protein (FTO) and alkB homologue 5 (ALKBH5) remove the m6A

modification marks making this a dynamic process [29,30]. FTO was the first eraser protein

identified [29,31] and it carries out oxidative demethylation where m6A is converted in

several steps to N^6-hydroxymethyl adenosine (hm6A), then to N^6-formyladenosine (f6A) and

lastly to adenine [31]. Additionally, reader proteins like the YT521-B homology (YTH)

domain family of proteins including YTHDF1, YTHDF2/3, insulin-like growth factor 2

mRNA-binding proteins (IGF2BP1), heterogeneous nuclear ribonucleoprotein

(HNRNPA2B1) recognise the modification and modulate alternative splicing and processing

of target transcripts [8,32]. The ones located in the cytoplasm include YTHDF1, YTHDF2,

and YTHDF3. The main function of these proteins is unclear; however, studies indicate

different roles of these enzymes in a cell. While YTHDF1 promotes the translation of mRNA

thereby promoting its protein expression, YTHDF2 promotes the degradation of mRNA, and

YTHDF3 promotes both translation and degradation of mRNA [33,34].

m6A methylation has been associated with ageing, and downregulation in the expression of

FTO and an overall increase in RNA m6A modification was found in human follicular fluid

(FF) and granulosa cells (GCs) from aged donors and murine ovaries from old mice [35].

Another recent study has also highlighted the relationship between elevated m6A levels and

brain ageing in both mice and humans [36], although a significant downregulation of m6A

methylation in ageing mice and in postmortem brain tissue from Alzheimer's disease patients

has also been reported [37].

Strikingly, ageing is also a major risk factor for the onset and progression of

neurodegenerative disorders such as Parkinson's disease (PD) and related synucleinopathies,

disorders associated with the accumulation of aggregated alpha-synuclein (aSyn) protein in

the brain. Despite the growing interest in the role of m6A methylation in ageing, its role in

PD is still unclear and remains controversial. While a recent study indicated that the

expression levels of m6A are significantly reduced in cellular models of PD and in the

striatum of a rodent model of PD [38], a different study showed that the m6A methyl

transferases were significantly decreased while the demethylases were increased in the

substantia nigra and in the striatum of mouse models of PD [39]. Ageing also affects multiple

homeostatic pathways including mitochondrial function and protein degradation pathways,

ultimately leading to cell death in the substantia nigra [40–42].

In this study, we investigated the m6A RNA modification profile in the context of ageing and

neurodegeneration in a mouse model of synucleinopathy. We performed methylated RNA

immunoprecipitation sequencing (meRIP-seq) to obtain the m6A epitranscriptome of the

midbrain of C57BL6 (Wt) control and A30P aSyn PD (aSyn Tg) mice at 3 months (3 mo)

and 15 months (15 mo) of age. We observed an increase in the m6A methylation in the genes

involved in the molecular and biological processes at the synapse. Several genes involved in

synaptic processes were hypermethylated in 3 mo aSyn Tg vs. Wt mice, while the

methylation marks were reduced in 15 mo aSyn Tg vs. age-matched Wt mice. Using

immunofluorescence imaging of primary cortical neurons, we observed that the localisation

of the writer METTL3 enzyme is not only limited to the nucleus but is also present at the

post-synapse. Our findings suggest that m6A RNA methylation is possibly playing a role in

neurodegeneration and ageing, and is involved in regulating synaptic function.

Results

m6A alterations during ageing in Wt mice and in a transgenic model of synucleinopathy

We began by examining the distribution of differentially expressed transcripts and transcripts

with m6A methylation in the healthy ageing brain of Wt mice. We further extended the

analysis to aSyn Tg mice, in order to compare the effects of ageing on m6A modification and

assess this change in the context of synucleinopathy. We extracted and dissected the

midbrains from five 3 mo and five 15 mo Wt and an equal number of aSyn Tg mice, as this

brain region is particularly relevant in the context of PD and related synucleinopathies.

Methylated RNA immunoprecipitation sequencing (meRIP-seq) was performed to determine

the midbrain-specific epitranscriptomic landscape in 3 mo and 15 mo mice (Figure 1A, B).

First, we compared the differential expression of genes due to ageing in 15 mo vs 3 mo mice

in both Wt and aSyn Tg animals. We observed a stronger effect in aSyn Tg mice where 270

transcripts were differentially expressed when compared with Wt mice, where only 79

transcripts were differentially expressed. Most differentially expressed transcripts in aSyn Tg

mice were downregulated (Supplementary Figure S1C). However, when we compared the

differentially methylated transcripts, we found similar numbers between Wt (420 genes) and

Tg (424 genes) mice (Figure 1A).

On the other hand, the effect of aSyn A30P overexpression was assessed by a comparison

between aSyn Tg and Wt mice at 3 mo and 15 mo. We did not identify major differences in

the number of differentially expressed transcripts which were 128 at 3 mo and 142 at 15 mo.

Similarly, the differentially methylated transcripts showed no significant changes between Wt

and aSyn Tg at 3 mo (384 genes) and 15 mo (353 genes) (Figure 1B). While we did not see a

significant difference in the differential methylation, we found that the prevalence of hyper-

and hypomethylated transcripts was reversed (Figure 1B).

As a consequence of ageing, the methylation levels of the differentially methylated m6A

transcripts in 15 mo vs. 3 mo Wt mice we found 418 hypermethylated and only 2

hypomethylated transcripts. On the contrary, 171 genes were hypermethylated and 244

transcripts were hypomethylated in 15 mo vs. 3 mo aSyn Tg mice (Figure 1D). A transcript

can carry multiple m6A methylation marks and some of these marks likely increase while

others decrease simultaneously. Such transcripts are referred to as mixed transcripts. We

found only 9 such mixed transcripts (Figure 1D, left column) in aSyn Tg mice during ageing.

To identify the effect of aSyn expression on the levels of the m6A modification, we

compared 3 mo aSyn Tg mice with age-matched Wt mice. We found 380 hypermethylated

transcripts and only 4 transcripts were hypomethylated, and no mixed transcripts. On the

other hand, when comparing 15 mo aSyn Tg vs. Wt mice, a reverse trend was observed, with

108 hypermethylated, 240 hypomethylated, and only 5 mixed transcripts (Figure 1D, right

column).

We found 6135 methylated transcripts in common between all groups (Figure 1E). The 3 mo

Wt mice displayed 266 unique methylated transcripts and the 15 mo Wt mice 255. These

numbers were lower for the aSyn Tg group where the 3 mo mice showed 165 unique

methylated transcripts and the 15 mo mice showed 189. At 15 months, both genotypes

showed 124 commonly methylated transcripts, suggesting the occurrence of conserved

processes in normal ageing and synucleinopathies. The aSyn Tg mice displayed 53

methylated transcripts correlated with aSyn expression and conserved with ageing.

The overall distribution of m6A methylation peaks on mRNA did not display any striking

anomalies in any of the groups. As expected, the majority of the m6A modifications were

localised close to the 3' UTR on mRNA (Figure 1F), as previously reported for ageing-

related genes [36].

m6A levels are reduced in synaptic transcripts upon ageing

To investigate the cellular processes and the subcellular regions where the differentially methylated transcripts were identified in 15 mo aSyn Tg vs. Wt mice, we performed gene ontology (GO) term analysis. We found that synaptic function was the biological process most represented. Our analysis revealed that transcripts with m6A methylation marks have strong enrichment for biological processes such as receptor internalization, excitatory post-synaptic potential, axonal transport, and dendritic spine morphogenesis. The identified transcripts were also involved in cellular processes including neuron projection cytoplasm and synaptic vesicle membrane (Figure 2A). Previous data from our group has also shown that while the average dendritic spine number was not significantly different, a strong difference was observed in the abundance of different types of spines in aSyn Tg neurons when compared with Wt neurons [67].

We also observed that when we compared the differentially enriched genes in aSyn Tg 15 mo vs. 3 mo, we saw a strong enrichment of pathways involved in the immune system. The genes with highest enrichment score were involved in innate immune system, adaptive immune system, immunoglobulin binding, phagocytic vesicle, defense response, innate immune response, and regulation of immune system processes among others (Supplementary Figure S2).

We used the SynGO software [64] (https://syngoportal.org/), an experimentally annotated database for synaptic location and functional GO, to specifically check the m6A methylation of synaptic transcripts. These differentially methylated genes are primarily involved in synaptic organisation, metabolism, transport, signaling, and both pre-and post-synaptic processes. Based on meRIP and RNA sequencing analysis, sunburst plots of Wt and aSyn Tg mice were generated (Figure 2B). Our findings suggest that the differentially methylated

transcripts that are involved at the synapse and in synaptic function were majorly

hypermethylated in 3 mo aSyn Tg mice compared to 3 mo Wt mice (Figure 2A). In 3 mo

mice, the hypermethylated transcripts were concentrated in synapse organisation and at the

pre-synapse (Figure 2B). Conversely, at 15 months of age, the majority of the differentially

methylated transcripts were hypomethylated when aSyn Tg mice are compared with Wt mice,

which were largely involved in signalling, pre- and post-synapse, and synapse organisation

(Figure 2C).

In summary, we observed that the differentially methylated genes, both hyper- and

hypomethylated genes in 15 mo aSyn Tg vs. Wt mice were mostly hypomethylated and

involved in synaptic processes, especially at the pre-and post-synapse. However, at 3 mo,

most of the differentially methylated genes were hypermethylated.

208   The levels of m6A regulatory proteins do not change in different brain regions in Wt

209   and aSyn Tg mice

To validate the meRIP-Seq results and to evaluate if the overall levels of the m6A

modification are affected in aSyn Tg vs. Wt mice, we performed an m6A ELISA, for total

RNA isolated from cortical and midbrain brain tissue of young and aged, Wt and aSyn Tg

mice. We found that overall, the levels of m6A modification marks were higher in the cortex

compared to the midbrain. In cortical brain tissue, we observed that the m6A marks in the

aSyn Tg mice do not display the same pattern as in Wt animals (Figure 3A). A significant

reduction in m6A marks was observed in 12 mo aSyn Tg mice compared to 12 mo Wt mice

(Wt; mean = 0.032 ± 0.018 OD, aSyn Tg; mean = 0.011 ± 0.002 OD, adjusted p-value =

0.045). In the younger animals, however not significant, a slight reduction in m6A marks was

observed in 3 mo aSyn Tg were compared with 3 mo Wt mice, indicative of an effect that

probably starts at a young age and progresses with ageing (Figure 3A). In the midbrain, we

did not observe significant differences in the m6A marks between the groups (Figure 3A).

This is consistent with the meRIP-Seq data where we observed that the total number of

differentially methylated transcripts was not altered among the groups (Figure 1A, B). This

suggests that despite the observed hypo- and hyper-methylation in certain transcripts (Figure

1D) the overall m6A levels remain in an equilibrium in the midbrain.

Next, we asked if the observed changes in the m6A marks in the cortex correlate with altered

levels of the three m6A regulatory proteins in the cortical brain tissue. To do so, we

performed immunoblotting analysis to assess the expression of m6A regulators — METTL3,

YTHDF1, and FTO. We used three different age groups – 3 mo, 6mo and 12 mo from the Wt

and aSyn Tg mice for our analysis. We observed no significant differences in the expression

levels of these m6A regulatory proteins neither within genotypes (Tg vs. Wt) nor across the

different age groups (Figure 3B). Nonetheless, we observed a trend towards a reduction in the

levels of METTL3 and FTO in Wt mice during ageing (Figure 3B).

Considering no difference in the protein levels of m6A regulators, we sought to identify if the

sub-regional and/or subcellular localisation of the three m6A regulatory proteins is altered.

For this, we assessed the levels of METTL3, YTHDF1 and FTO by immunofluorescence

intensity analysis in four different regions of the mouse brain, namely cortex, striatum,

hippocampus, and cerebellum. MAP2 co-staining was used only to identify the neurons

present in the areas of the brain. All these proteins were present in our selected areas of

interest, but their compartmentalization varied across different regions of the neurons.

The m6A writer protein, METTL3 was observed as small puncta within the nucleus in the

cortex, striatum, hippocampus, and cerebellum in both aged Wt and aSyn Tg brain (Figure

4A). We found no significant differences in protein levels between aSyn Tg vs. Wt mice

based on the fluorescence intensity (Figure 4B).

The m6A reader protein, YTHDF1, on the other hand, localised only in the soma of neurons

in all four areas of interest, with a similar expression pattern in Wt and aSyn Tg mouse brain

slices. (Figure 4A). No significant difference in the fluorescence intensity of the protein was

observed in 15 mo aSyn Tg mice when they were compared with Wt mice (Figure 4B).

Finally, the m6A eraser, FTO, was detected in the nucleus and the soma of cells in the cortex,

striatum, hippocampus, and cerebellum. FTO was also localized in the dendrites of neurons in

the cortex and cerebellum in both Wt and aSyn Tg mouse brain slices (Figure 4A). After

analysis of the expression of this protein in the soma, no significant differences were

observed in the different regions of either aSyn Tg or Wt mice (Figure 4B).

Our findings show that the m6A regulatory proteins – METTL3, YTHDF1 and FTO are

present in the cortex, striatum, hippocampus, and the cerebellum in the mouse brain. In

addition, we found that the levels of METTL3, YTHDF1 and FTO do not change in cortical

brain tissue.

METTL3 is reduced in the aSyn Tg mouse post-synapse

Based on our bioinformatics analysis, we observed that the majority of m6a genes were

synaptic, so we asked if/and to what degree the m6A regulators are present at synapses acting

as local regulators since the overall regulator levels remain unchanged. We used primary

cortical neurons that allow higher imaging potential and performed co-staining of the three

m6A regulatory proteins – METTL3, YTHDF1 and FTO alongside Syn1, pre-synaptic

marker and PSD95, post-synaptic marker to study the localisation of these proteins at the

synapse. Previous studies have indicated reduced synaptic activity in neurons derived from

sporadic PD patients when compared with controls [65,66].

Two experimenters performed the colocalisation analysis, and similar results were obtained.

METTL3 was observed to be present in the nucleus as nuclear speckles and it significantly

co-localized with PSD95, indicating its presence at the post-synapse in both Wt (PCC =

0.810 ± 0.039) and aSyn Tg (PCC = 0.748 ± 0.075) primary neurons. In contrast, METTL3

did not co-localize with synaptotagmin1 in Wt (PCC: 0.489 ± 0.086) or aSyn Tg (PCC: 0.476

± 0.050) primary neurons, indicating no noticeable localisation at the pre-synapse (Figure

5B). Similarly, we investigated the compartmentalization of YTHDF1, which did not overlap

with either synaptotagmin1 (PCC: Wt = 0.382 ± 0.090, aSyn Tg = 0.294 ± 0.065) or PSD95

(PCC: Wt = 0.310 ± 0.088, aSyn Tg = 0.280 ± 0.069) (Figure 5B). The m6A eraser protein,

FTO, was identified within the cell nucleus and in some cases, in the soma of primary

neurons. No significant co-localisation of FTO was observed at the pre- (PCC, Wt = $0.468 \pm$

0.109, aSyn Tg = 0.488 ± 0.095) or post-synapse (PCC, Wt = 0.366 ± 0.063, aSyn Tg = 0.463

± 0.081) (Figure 5B).

Overall, we observed that the expression of METTL3 was reduced at the post-synapse in

aSyn Tg primary neurons suggesting a possible loss of METTL3 at the synapse.

Primary neurons from aSyn Tg mice have a greater number of synapses than Wt

animals

Since we observed a reduction in the localisation of METTL3 at the synapse in aSyn Tg

primary neurons, we wondered whether this has any influence on the synapse morphology

and number, or vice versa. For this, we used primary cortical neurons and performed proximity analysis of the pre-and post-synaptic markers Syn1 and PSD95 using the IMARIS software Imaris (RRID: SCR_007370) and the data were analysed blindly to avoid bias. We found that the number of synapses in aSyn Tg primary neurons was significantly increased when compared with Wt neurons (Wt; mean = 2607 ± 763.1, aSyn Tg; mean = 3152 ± 1187, p-value = 0.044) (Figure 6).

Discussion

The involvement of epitranscriptomics in PD is still not well understood. In this study, our objective was to begin to unravel the role of m6A RNA methylation in the A30P aSyn Tg mouse model of PD/synucleinopathy, in disease and physiological ageing.

Our findings revealed that 99.5% of the transcripts were hypermethylated in 15 mo vs. 3 mo Wt mice suggesting that ageing might contribute to a significant increase in m6A methylation. This observation aligns with a recent study that has demonstrated an increase in m6A methylation during ageing in both mice and humans [36]. This indicates that the accumulation of m6A methylation marks might be a general feature of ageing in the brain, potentially influencing various biological processes. It has also been shown that these epigenetic changes arise mainly in young and aged mice, while during the adolescent stage these genes are stabilised [68]. In contrast, when comparing 15 mo aSyn Tg mice with their 3 mo counterpart littermates, more transcripts were hypomethylated. This decrease in methylation with age in aSyn Tg mice might suggest that the loss of methylation might be related to aSyn expression/pathology. There is a possibility that the overexpressed aSyn is somehow interacting with the m6A regulators and inhibiting methylation. Previous studies

have shown that the 6-hydroxydopamine rat PD model has significantly reduced m6A levels
in the striatal brain region and in the cellular model of PD [69]. Similarly, a reduction in the
total levels of m6A was also reported in the MPTP-treated PD mouse model and in MPP+-
treated MN9D cells [70].

The effect of aSyn expression was more directly studied in the comparison between 3 mo
aSyn Tg and Wt mice. The increase in methylation suggests that aSyn expression may be
linked to enhanced methylation activity in early life. However, in 15 mo aSyn Tg vs. Wt
mice, we observed more hypomethylated genes, suggesting that A30P aSyn expression, and
likely pathology, might interfere with the writing or erasing of methylation marks during
ageing. Importantly, our results are in accordance with a recent publication suggesting that
the levels of m6A are significantly lower in PD patients [71].

Interestingly, based on GO term analysis, we found that differentially methylated transcripts
were predominantly associated with synaptic function and that methylation was lost with
ageing. Hypermethylation of transcripts involved in the synaptic processes might lead to an
inhibition of mRNA translation and thereafter reduced functionality of the synapse. This is
particularly relevant as more than half of the causative genes and risk factors in PD have been
identified to function at the synapse [72] and this change in methylation pattern might
indirectly affect the functioning of the synapse. Considering that different RNA-based
epigenetic mechanisms occur locally at the synapse, this underscores their potential
importance in synaptic regulation [74]. Additionally, GO term analysis of the differentially
expressed transcripts showed a strong enrichment in immune response-related pathways.
Notably, m6A is implicated in the regulation of inflammation under physiological and
pathological conditions, known as "inflammatory ageing", or inflammaging. This is a chronic
inflammatory state arising during ageing and it is characterised by an increase in pro-

inflammatory factors such as tumor necrosis factor (TNF-α), interleukins (IL-1β and IL-6),

and C-reactive protein (CRP), to name just a few [73].

The assessment of total m6A methylation marks showed a significant reduction in the cortex

of 12 mo aSyn Tg mice when compared with age-matched Wt mice. This suggests that the

overall m6A methylation decreases in the ageing brain. This finding is consistent with recent

studies that reported a predominant reduction in m6A methylation in the frontal cortex and

cingulate cortex from PD patients [71,75]. In vitro studies have also reported a significant

reduction in m6A levels in the cellular PD models [70]. Despite this, we observed similar

expression levels of m6A regulatory proteins. However, our findings are consistent with

those from a recent study reporting that the levels of YTHDF1 and FTO mRNA remain

unchanged in PD. However, a reduction in the levels of METTL3 mRNA in PD patients was

also reported in that study, in contrast to what we observed [71]. Some studies have also

reported up-regulation of ALKBH5, IGF2BP2 in the substantia nigra while, YTHDF1 and

fragile X mRNA (FMR1) were downregulated. In the striatum, an up-regulation of FMR1

and Cbl proto-oncogene like 1 (CBLL1) and down-regulation of METTL3, RBM15 and

IGF2BP3 was observed [76].

Although we found no significant differences in the levels of m6A regulatory proteins, we

observed that the three proteins were present in the cortex, striatum, hippocampus, and

cerebellum. Our findings are consistent with those from previous studies reporting that the

m6A-methyltransferase complex, including METTL3, is predominantly found in the nucleus,

but that it can also be localised in the cytoplasm [77–79]. As methylation generally occurs co-

transcriptionally in the nucleus, while reading occurs largely at the cytosol, YTHDF1 is

primarily cytosolic [80]. FTO, on the other hand, is localised both in the nucleus and in the

cytosol [81]. These observations suggest that an inhibitory factor affecting their function and

aSyn aggregation is potentially playing a role. Previous studies have shown that aSyn binds

to histone proteins, which may alter the protein surface due to post-translation modifications

of histone tails, altering their affinity for DNA and other histones [82]. Another study showed

that aSyn expression increases histone-H3 lysine-9 (H3K9) methylation, suggesting a

possible interaction of aSyn with epigenetic writer proteins [83]. This could constitute a

mechanism through which aSyn affects gene methylation, ultimately affecting synaptic

functioning [84,85].

Given that most of the hypermethylated genes were synaptic, we assessed the presence of

these regulators at the synapse. Our findings suggest the presence of METTL3 at the post-

synapse, indicating that the enzyme might be locally active at the synapse. The dynamic

nature of m6A methylation may be localised to different regions within a cell, depending on

the temporal requirements of the respective genes [86,87]. In another study, m6A regulatory

proteins were found localised adjacent to the synapse and in the dendrites, also suggesting the

sub-cellular localisation of m6A methylated mRNA which might lead to synaptic

modifications [88,89]. Our findings revealed that the expression of METTL3 is reduced at the

post-synapse in aSyn Tg DIV14 neurons, suggesting a possible loss of METTL3 at the post-

synapse. Due to this reduction at an early stage, the nucleus might be sending

hypermethylated transcripts to the dendrites to compensate for this loss. However, this may

be lost during ageing. Based on our bioinformatics analysis, we observed that methylated

transcripts associated with synaptic function were reduced in aSyn Tg mice with ageing. This

suggests that a reduction in the m6A methylation in synaptic genes may be directly linked

with the reduced METTL3 localisation at the synapse in aSyn Tg neurons.

Conclusion

In conclusion, our study highlights the critical role of m6A RNA methylation in the progression of PD and its implications in ageing, particularly focusing on the A30P aSyn mouse model. Our findings indicate that aSyn expression/pathology may influence RNA methylation patterns, leading to reduced levels of m6A in mice. The differentially methylated transcripts were predominantly synaptic, indicating that m6A modifications play a crucial role in synapse physiology and potentially contribute to the synaptic dysfunction observed in PD [90]. Although we observed no significant changes in the levels of m6A regulatory proteins, the reduction in METTL3 at the synapse in aSyn Tg mice informs on the importance of subcellular compartmentalization of these proteins in the regulation of RNA methylation. Therefore, our research provides new insights into the epigenetic mechanisms underlying PD and underscores the need for further exploration of m6A RNA methylation as a potential therapeutic target for PD and other related synucleinopathies.

Materials and methods

Resource availability

Lead contact for reagent and resource sharing

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Tiago F. Outeiro (tiago.outeiro@med.uni-goettingen.de).

Experimental model and subject details

Animals

Five male homozygous [A30P] aSyn transgenic (aSyn Tg) mice (C57BL/6J-Tg(Th-SNCA*A30P*A53T) 39Eric/J, Strain #:008239, RRID: IMSR_JAX:008239) and five wild type (Wt) littermates, 3 mo, 6mo, 12 mo and 15 mo were included in this study for RNA sequencing and analysis. Additional animals used for each experiment are mentioned in the respective section. Animals were housed in standard cages with 12h light/12h dark cycle and ad libitum access to food and water in the animal facility of the University Medical Centre Göttingen (Göttingen, Germany). Animal procedures were performed in accordance with the European Community (Directive 2010/63/EU); and institutional and national guidelines under the Lower Saxony State Office for Consumer Protection and Food Safety (LAVES) (license number 19.3213). Animals were anesthetised with carbon dioxide (CO_2) and then sacrificed using cervical dislocation. For tissue processing, the brain was surgically extracted from the skull and was further dissected to isolate the cortex and midbrain. The tissue was snap-frozen in liquid nitrogen and preserved at -80°C until use.

For the preparation of brain slices, the animals were perfused with 1x phosphate buffer saline (PBS) followed by 4% paraformaldehyde (PFA). The whole brain was dissected out of the skull and placed overnight in 4% PFA at 4°C. The following day, the brain was washed once with 1x PBS and then placed in 15% saccharose and 30% saccharose with 0.1% sodium azide (NaN_3). The brains were placed at 4°C until used. Sagittal cryo-sections of 30 µm thickness were cryo-sectioned using Leica CM3050 S cryostat for each mouse brain. The sections were placed in 1x PBS with NaN_3 until they were used for immunohistochemistry (IHC) protocol.

Neuronal cultures

Primary cortical neuronal cultures were prepared from both aSyn Tg and Wt mice as

previously described (Saal et al., 2015). Cells were seeded on coverslips coated with poly-L-

ornithine (0.1 mg/mL in borate buffer; PLO) (Sigma-Aldrich, MO, USA) or culture plates

(Corning, Merck, Darmstadt, Germany) for immunocytochemistry and maintained in

neuronal cell culture medium (Neurobasal medium, Gibco Invitrogen, CA, USA),

supplemented with 1% penicillin-streptomycin (PAN Biotech, Aidenbach, Germany), 0.25%

GlutaMAX (Gibco Invitrogen, CA, USA), and 2% B27 (B-27™ Plus Supplement (50X),

Gibco, #A3582801). The cells were maintained at 37°C with 5% CO_2, and one-third of the

medium was replaced with fresh media every 3-4 days.

Immunocytochemistry

Primary neurons were first washed with 1x PBS (PAN Biotech, Aidenbach, Germany) and

fixed with 4% of PFA for 20 min at room temperature (RT). Cells were first permeabilized

with 0.1% Triton X-100 (Sigma-Aldrich, MO, USA) for 10 min and then blocked with 1.5%

normal goat serum (NGS) in 1x PBS (Sigma-Aldrich, MO, USA) for 1h at RT. The cells

were then incubated with the primary antibodies diluted in blocking solution (FTO (mouse,

Abcam – ab92821, 1:1000), METTL3 (rabbit, Cell Signalling - #51104, 1:1000), YTHDF1

(rabbit, Proteintech 17479-1-AP, 1:1000), MAP2 (rabbit, Proteintech – #17490-1-AP,

1:1000), postsynaptic density protein 95 (PSD95) (mouse, Synaptic systems 124011, 1:1000),

PSD95(D74D3) (rabbit, Cell Signalling - #3409, 1:1000), synaptotagmin 1 (Syn1) (mouse,

Synaptic systems - #105311, 1:1000) synaptotagmin1 (rabbit, Synaptic systems – 105102/3,

1:1000)) overnight at 4°C, with constant shaking at a speed of 40-50 rpm. The following day,

cells were washed three times for 10 min each with 1x PBS (PAN Biotech, Aidenbach,

Germany) and then incubated with fluorescently conjugated secondary antibodies (Alexa

Fluor 488 donkey anti-mouse IgG – Invitrogen, Alexa Fluor 488 donkey anti-rabbit IgG -

Invitrogen, Alexa Fluor 555 donkey anti-mouse IgG – Invitrogen, Alexa Fluor 555 donkey

anti-rabbit IgG – Invitrogen, Alexa fluor 633 goat anti-chicken IgG - Invitrogen) for 2h at

RT, on a shaker at 40-50 rpm. Lastly, nuclei were counter-stained with DAPI (Carl Roth,

Karlsruhe, Germany) and mounted on glass slides (Epredia™ SuperFrostPlus™ Adhesion)

with mounting media (Invitrogen™ Fluoromount-G™) for microscopy.

Immunohistochemistry

The free-floating tissue was carefully transferred from PBS and $NaNH_3$ to 1x tris-buffered

saline (TBS) in a new 24-well plate with immunohistochemistry (IHC) grids. The tissue was

washed three times for 5 min each with 1x TBS to remove the remnants of NaN_3. Following

this, tissue antigen retrieval was performed using citrate buffer (10 mM citrate, 0.05% Tween

20, pH 6.0) at 90°C for 30 min in a water bath. The tissue samples were allowed to cool

down until they reached RT. Tissue sections were washed thrice for 10 min each with 1x

TBS and then incubated with 0.3% Triton X-100 for 20 min. Without washing, the tissue

slices were incubated directly in a blocking solution (5% BSA, and 1% NGS in 1x TBS) for

2h at RT. The tissue sections were then incubated for 48h with antibodies diluted 1:200 times

(FTO (mouse, Abcam – ab92821, 1:1000), METTL3 (rabbit, Cell Signalling - #51104,

1:1000), YTHDF1 (rabbit, Proteintech #17479-1-AP, 1:1000), MAP2 (rabbit, Proteintech –

#17490-1-AP, 1:1000)) at 4°C, on a shaker. The following day, tissue sections were washed

thrice for 10 min each with 1x TBS at RT. Afterwards, the tissue sections were incubated in

secondary antibody (1:400 dilution) (Alexa Fluor 488 donkey anti-mouse IgG – Invitrogen,

Alexa Fluor 488 donkey anti-rabbit IgG - Invitrogen, Alexa Fluor 555 donkey anti-mouse

IgG – Invitrogen, Alexa Fluor 555 donkey anti-rabbit IgG – Invitrogen, Alexa fluor 633 goat

anti-chicken IgG - Invitrogen) overnight at 4°C on the shaker. The following day, the tissue

was washed thrice for 10 min each with 1x TBS to remove the unbound secondary antibody.

The tissue sections were incubated for 5 min in DAPI (Carl Roth, Karlsruhe, Germany)

followed by a 10 min wash with 1x TBS. The tissue sections were then mounted onto glass

slides (Epredia™ SuperFrostPlus™ Adhesion) and left to dry in the dark at RT. 8-10µl of

mounting media (Invitrogen™ Fluoromount - G™) was applied as a drop onto each section

on the glass slide and a glass coverslip (Avantor, 631-0147, 0.16-0.19mm thickness, #1.5)

was carefully placed on the tissue to avoid any air bubbles.

Western blots

Cortical brain tissue was extracted from three adult mice per age group (age group 2 mo, 6

mo and 12 mo) from each Wt and Tg mice. The tissue from adult mice was lysed using

homogenisation beads (1.4 mm diameter ceramic beads, 91-PCS-CK14B, PEQLAB

Biotechnologie, Germany) in RIPA buffer (50 mM Tris, pH 8.0, 0.15M NaCl, 0.1% SDS,

1.0% NP-40, 0.5% Na-Deoxycholate, 2 mM EDTA) supplemented with protease and

phosphatase inhibitors cocktail (cOmplete™, Mini Protease Inhibitor Cocktail, Roche). The

homogenisation was performed in Precellys 24 tissue homogenizer (Bertin Instruments) at 65

000 $\times$ g for 2 $\times$ 30 sec runs with a 30 sec break in between. Protein concentrations in the

lysates were determined by the BCA protein assay (Pierce Biotechnology, #23225) following

the manufacturer's protocol. The spectrophotometric recordings were performed using a plate

reader (Tecan, Infinite® M200 pro) at 562 nm. Volume for 20 µg of protein per sample was

calculated, to which Laemmli sample buffer (250 mM Tris-HCl pH 6.8, 10% SDS, 1.25%

Bromophenol Blue, 5% ß-mercaptoethanol, 50% glycerol) was added (1:5 dilution of

Laemmli: protein). The proteins were denatured for 5 min at 95°C, with 600 rpm in a thermal

shaker. The samples were loaded on a 12% SDS-PAGE electrophoresis gel and separated at

120V for approximately 1h. The proteins were then transferred to PVDF membranes (iBlot2

Transfer Stack, PVDF, Life, #IB24002X3) using iBlot2 (Invitrogen, CA, USA) using the P0

protocol for 7 min. Membranes were blocked with 5% BSA (Sigma Aldrich, #AG9418) in

TBS (pH 8) with 0.05% Tween-20 (TBS-T) and then incubated with the appropriate primary

antibody (FTO (mouse, Abcam – ab92821, 1:5000), METTL3 (rabbit, Cell Signalling -

#51104, 1:1000), YTHDF1 (rabbit, Proteintech #17479-1-AP, 1:2500)) overnight in 5% BSA

(Sigma-Aldrich, MO, USA) in TBS-T at 4°C. After three washes with TBS-T, membranes

were incubated for 2h with horseradish peroxidase (HRP) conjugated secondary antibodies

(Cytiva sheep anti-mouse IgG, 1:8000 dilution, Cytiva donkey anti-rabbit IgG, 1:8000

dilution). Following incubation, membranes were washed three times with TBS-T for 10 min

each and developed in a chemiluminescence system (Fusion FX Vilber Lourmat, Vilber,

France) using chemiluminescent HRP substrate (Millipore, MA, USA). Intensities of specific

bands were normalised to a protein loading control.

RNA extraction

For brain tissue, TRIzol reagent (Invitrogen, CA, USA) and ceramic beads (1.4 mm diameter

ceramic beads, 91-PCS-CK14B, PEQLAB Biotechnologie, Germany) were added in vials

with frozen tissue and homogenisation was performed with the Precellys 24 tissue

homogenizer (Bertin Instruments) at 65 000 g for 2 x 30 sec runs with a 30 sec break in

between. RNA isolation was performed with TRIzol reagent according to the manufacturer's

instructions. Briefly, 0.5 ml of TRIzol was added to the tissue samples and incubated for 5

min before homogenisation. After homogenisation, 0.1 ml of chloroform was added to each

tube. The tubes were shaken vigorously and incubated at RT for 2-3 min. The tubes were

centrifuged at 14 000g, for 10min at 4°C. The top aqueous layer was carefully transferred to a

fresh tube. Total RNA was then precipitated using Clean and Concentrator 5, Zymo research

kit (Zymo Research, #R1014). Following RNA extraction and precipitation, RNA was

dissolved in RNAse-free water and concentration and quality were estimated with NanoDrop

1000 spectrophotometer (Thermo Fisher Scientific, MA, USA) by measuring absorption at

260 nm and the ratios 260/280 nm and 230/280nm respectively.

m6A fluorescence ELISA

m6A RNA methylation ELISA (m6A RNA methylation, Abcam, #ab233491) was used for

the detection of m6A RNA methylation in the cortical brain and midbrain tissue from 3 mo

and 12 mo/15 mo Wt and aSyn Tg mice. For this experiment, we used cortical mouse brain

tissue from Wt mice each at 3 mo and 12 mo (six 3 mo and six 12 mo), and aSyn Tg mice

(six 3 mo and five 12 mo). Midbrain tissue was used from Wt mice (three 3 mo and four 15

mo) and aSyn Tg mice (five 3 mo and four 15 mo). This protocol ensures that the total m6A

methylation is measured in both mRNAs and small RNAs. The protocol was performed

according to the manufacturer's protocol. Briefly, total RNA isolated from the tissue was

added to bind to the assay wells. The wells were then washed to remove the unbound RNA

and diluted capture antibody was added to the wells. Unbound capture antibody was washed

out with multiple washing steps. Diluted detector antibody was then added along with fluoro-

enhancer. The wells were washed again and incubated with fluoro developer mixture to

initiate the fluorescence reaction. The fluorescence was then measured at 590nm using plate

reader (Tecan, Infinite® M200 pro).

MeRiP

Total RNA was treated with DNase to remove any DNA remnants and further, 5-10 µg of total RNA was depleted for rRNA using the RiboMinus Eukaryote Ribosomal Removal Kit (Invitrogen). The large RNA fraction (>200 bp) was fragmented to a fragment size of approximately 100-120 nt by RNA Fragmentation Reagent at 70°C for 5 min (#AM8740, Thermo Fisher Scientific, Waltham, MA). 5% of this RNA sample was kept as inputs and the rest was subjected to immunoprecipitation (IP). 300 µg of fragmented RNA was used for each IP. 3 µg of anti-m6A antibody was incubated with the RNA IP buffer (0.2 M Tris-HCl pH 7.5, 0.5 M NaCl and 2% (vol/vol) Igepal CA-630) for 2h at 4°C with constant rotation. RNA antibody mixes were then crosslinked twice with 0.15 J/cm2 of UV light (254 nm) in a UVP crosslinker (Analitik Jena). Antibody-RNA conjugates were incubated with 30 µl of Protein A/G beads overnight (ON) at 4°C with rotation in 500 µl IP buffer supplemented with 200 units. The beads with immunoprecipitated RNA were washed with IP buffer five times, and further washed with low-salt (50 mM Tris pH 7.4, 50 mM NaCl, 1mM EDTA, 1, 0.1% NP-40, 0.1% SDS) and high-salt (same as low-salt but with 500mM NaCl) buffers at 4°C with rotation to remove any nonspecific binding. RNA was eluted by incubating in 150 µl PKD buffer (Qiagen) with 10 µl of Proteinase K (Millipore) for 1 hour at 37°C with agitation. Eluted RNA was cleaned before proceeding to library preparation.

Library Preparation and Sequencing

Samples were prepared for sequencing using the SMARTer Stranded Total RNA-Seq Kit v2 — Pico Input Mammalian (Takara) according to the manufacturer's protocol. All the RNA obtained from the IP samples was used for library preparation, for input samples, 2 ng was

used. The libraries were amplified for a total of 12 cycles for input and 16 cycles for IP

samples. RNA-Seq samples were prepared with the TruSeq Stranded mRNA Library Prep Kit

(Illumina) according to the manufacturer's instructions for which, 50 ng of rRNA-depleted

RNA from each of the inputs was used. Prepared libraries were sequenced in a Hiseq 2000

System (Illumina) for 50 cycles in single-end reads. Additional metadata is also available via

the GEO database (GSE275432).

Bioinformatic Analysis of meRIP-seq and RNA-seq

Raw reads were processed and demultiplexed using bcl2fastq (v2.20.2), and low-quality

reads were filtered out with Cutadapt v1.11.0 [43]. Filtered reads were mapped to the mouse

(mm10) genome using the STAR aligner v2.5.2b [44]. The resulting bam files were sorted

and indexed, and the unmapped reads were removed using SAMtools v1.9.0 [45].

Methylation sites were determined using MeTPeak v1.0.0 [46], and differential methylation

(hypo- and hyper-methylated regions) was assessed with ExomePeak v2.16.0 [47] using 3 mo

samples as control and 15 mo as treatment for the first comparison and for the second

comparison, Wt samples were used as controls while aSyn Tg were used as treatment. An

adjusted P value (padj, also termed as FDR [false discovery rate]) cutoff of 0.05 and FC

cutoff of 1.2 or 1.5 were used as indicated in the text. Only consistently significantly,

differentially methylated peaks were used, unless indicated.

For RNA-seq analyses, read counts were obtained with subread's featureCounts v1.5.1 [48]

from the bam files of input samples. Differential gene expression was determined by DESeq2

v3.5.12 [49] using normalised read counts and correcting for covariates detected by RUVseq

v1.16.1 [50]. Cutoffs of padj $\leq$ 0.05, FC $\geq$ 1.2, and BaseMean $\geq$ 50 were applied to the results

(Supplementary Figure S1A, B). Background expressed genes were determined for each

region as those genes with a BaseMean > 50 in the corresponding input sample.

For visualization, bam files of both IP and input samples were collapsed for PCR duplicates

using SAMtools, and IP samples were normalized to their corresponding inputs and to their

library size using deeptools v3.2.1 [51] bamCompare. The resulting normalized tracks were

visualized in the IGV Browser 2.9.2 [52].

GO Analysis

GO term enrichment analyses were performed using the App ClueGO v2.5.3 [53] in

Cytoscape 3.7.2 [53], with GO Term Fusion enabled to collapse terms containing very similar

gene lists and using a custom background corresponding to expressed genes in the

corresponding species as obtained from RNA-seq results of the corresponding input samples

of the meRIP experiments. GO term tables for biological processes, cellular components,

pathways, and KEGG were produced and are labelled accordingly in the figures. The

resulting enriched GO terms were visualized with a custom script using ggplot2 v3.3.5 [54]

playing the adjusted p-value (padj) for the GO term, the number of genes from the list that

belong to said term, and the percentage of the total genes in the GO term that are present in

the list. Synaptic GO enrichment analyses were performed with SynGO (v1.1,

syngoportal.org) [55].

Additional Bioinformatics Packages and Tools

Scripts and analysis pipelines were written in R (3.5.2) [56]. Peak annotation was performed

with Homer v4.10.4 [57] and Annotatr v1.8.0 [58]. Guitar plots were produced with the

Guitar v1.20.1 [59] R package. Volcano plots were generated with plot.ly/orca v4.9.4.1 [60]

(Supplementary Figure S1C). Area-proportional Venn diagrams were produced with

BioVenn (www.biovenn.nl), and multiple list comparisons performed with Intervene / UpSet

(asntech.shinyapps.io/intervene/). Mouse/human homologues were determined by their

annotation in NCBI's HomoloGene database using the HomoloGene (v1.4.68.19.3.27) R

package. Odds ratios and p values to determine significance in overlapped datasets were

calculated with the GeneOverlap R package v1.18.0 [61]. De novo motif analyses were

performed with Homer's findMotifsGenome, and motifs containing the DRACH consensus

sequence out of the top 10 most significant are dipcasplayed. KEGG pathway enrichment was

produced with KEGG Mapper (www.genome.jp/kegg/mapper/) [62]. Microscopy images

were pre-processed with Fiji, and quantification was automated in Cell Profiler

(cellprofiler.org) [63]. Graphs, heat maps, and statistical analyses were performed on

GraphPad Prism version 9.3.1 for Mac.

Microscopy

Images of the immunofluorescence-stained neurons (three experiments in total, with at least

10 regions of interest acquired per experiment) and brain tissue were acquired using a

confocal microscope ZEISS LSM900. The images for the neuronal cell culture were acquired

as a z-stack using 63x magnification with Airyscan mode. Brain slices were imaged at 40x

magnification, with tiles in confocal mode.

635 **Statistical analysis**

636 The cell culture images for m6A regulators and their colocalisation at the synapse were

637 analysed using FIJI with the plugin JaCoP, while the synapse counting was performed using

638 IMARIS. IMARIS analysis was performed following the protocol from the Queensland Brain

639 Institute. The protein signals were further classified inside the dendritic or outside using

640 membranes based on the MAP2 signals. The signals were then divided into four groups,

641 representing PSD95 inside the neuron, PSD95 outside the neuron, Synaptotagmin1 (Syn1)

642 inside the neuron and Syn1 outside the neuron. When the signal for PSD95 inside the neuron

643 was at $< 500\mu m$ from the signal for Syn1, it was considered a synapse. Colocalisation of

644 synaptic markers with m6A regulators was estimated using Pearson's correlation coefficient

645 (PCC) estimated using the JaCoP plugin in ImageJ, FIJI. All IHC images were also analysed

646 for their expression levels using FIJI software. Data analysis was performed using the

647 GraphPad software version 9 for Windows (GraphPad Software, La Jolla California

648 USA, https://www.graphpad.com). For group comparisons, one-way ANOVA with Tukey's

649 test was used, while comparisons of two groups of means were done with an unpaired

650 student's t-test. All data are expressed as mean $\pm$ SD. Differences are considered significant

651 with $p < 0.05$ (*$p < 0.05$, **<0.01, ***<0.001).

652

Declarations

653

654 *Ethics approval*

Animal procedures were performed in accordance with the European Community (Directive

2010/63/EU); and institutional and national guidelines under the Lower Saxony State Office

for Consumer Protection and Food Safety (LAVES) (license number 19.3213).

Consent for publication

All authors approved the publication of the article.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the

corresponding author upon reasonable request. All raw data files for meRIPseq are available

on the GEO database (GSE275432).

Competing interests

The authors declare that they have no competing interests.

Funding

TFO and AF are supported by the Deutsche Forschungsgemeinschaft (DFG, German

Research Foundation) under Germany's Excellence Strategy - EXC 2067/1- 390729940), and

by SFB1286 (Projects B6 and B8). AF is supported was supported by the Niedersächsisches

Vorab: Forschungskooperation Niedersachsen – Israel (ZN3460).

Authors' contributions

AC performed, analysed, and interpreted the data from cell cultures, ELISA, IHC, and

western blot, and interpreted the meRIP results. MM and VG performed experiments in

animals. MX interpreted the data for meRIP analysis and was a major contribution in writing

the manuscript. YF performed and analysed ICC data. RCH performed the meRIP and

subsequent data analysis. AC and TO wrote the manuscript. All authors read and approved

the final manuscript.

 Figure Legends

Figure 1. m6A modification landscape in ageing and aSyn Tg mice. (A) Bar plots

representing the variations in the differentially expressed genes and differentially methylated

genes in the midbrain from aSyn Tg mice than Wt mice when 3mo mice are compared with

15mo applying equal cutoffs for FC and adjusted P value (FC > 1.2 and padj ≤ 0.05). Left:

with age, right: with aSyn (B) Bar plots representing the effects of aSyn pathology on

differential expression and differential methylation in 3mo and 15mo Wt vs. aSyn Tg mice.

(C) Overview of protocols utilized in this project. (D) Pie charts representing the overview of

the distribution of hypomethylated, hypermethylated and mixed transcripts as an effect of

aSyn pathology and ageing. (E) Venn diagram representing the number of differentially

methylated genes involved in young and old Wt and aSyn Tg mice. It also provides a

summary of the overlapping genes across different genotypes. (F) Guitar plots representing

the distribution of m6A methylation on mRNA and lncRNA. CDS stands for coding

sequence, and UTR stands for untranslated region.

Figure 2. Gene ontology (GO) analysis for 3 mo and 15 mo Wt vs. aSyn Tg mice. (A) GO

analysis for the biological processes and cellular processes in 15 mo aSyn Tg vs. Wt mice,

FDR = 1.5. (B) Sunburst plots representing GO analysis for the genes involved in the

processes at the synapse in aSyn Tg vs. Wt mice at 3 mo. Graphs created with

syngoportal.org. (C) Sunburst plots representing GO analysis for the genes involved in the

processes at the synapse in aSyn Tg vs. Wt mice aged mice at 15 mo. Graphs created with

syngoportal.org.

Figure 3. Levels of m6A RNA and m6A regulators in mouse brain tissue. (A) Bar plots

representing the levels of m6A modification on total RNA using ELISA from cortical and

midbrain regions of Wt and aSyn Tg mice. A significant reduction in the levels of m6A was

observed in the aged, 12 mo aSyn Tg mouse cortex when compared with 12 mo Wt mice (*,

ajd P value = 0.045). No significant differences were observed in the midbrain region of the

brain. (B) Biochemical analysis using western blot comparing the expression levels of

METTL3, YTHDF1 and FTO in the cortical tissue from 3 mo, 6mo and 12 mo Wt vs. aSyn

Tg mice. Scatter plot showing no significant difference in the expression level when

comparing Wt with aSyn Tg mice.

Figure 4. Distribution of m6A regulatory proteins METTL3, YTHDF1 and FTO in the

cortex, striatum, hippocampus and cerebellum of mouse brain. (A) Representative

images of mouse brain tissue slices (30µm) fixed, and stained with MAP2, METTL3,

YTHDF1 and FTO. The images are acquired using a confocal microscope, 40x magnification

with tiles, and scale bar = 50µm. White arrows point to the neurons with the protein

expression. (B) Bar plots showing the comparison between the mean intensity of METTL3,

YTHDF1 and FTO in 15mo Wt vs. aSyn Tg mice.

Figure 5. METTL3 is localised at the post-synapse in Wt and aSyn Tg DIV14 primary

neurons. (A) Representative images of DIV14 primary neurons, fixed and stained with

Histone H1, MAP2, PSD95, Synaptotagmin 1, METTL3, YTHDF1, and FTO in Wt and aSyn

Tg neurons. Images are acquired using airy-scan at 63x magnification, scale bar = 10µm. (B)

Box and whisker plot representing the colocalisation of m6A regulators in Wt and aSyn Tg

D-14 primary neurons at the pre-and post-synapse. METTL3 colocalises with the post-

synapse, (Pearson's correlation coefficient, Wt = 0.810, aSyn Tg = 0.748). A significant

difference (*, adj p value = 0.018) was observed between METTL3 at the post-synapse in Wt

neurons when compared with aSyn Tg neurons. (C) Box and whisker plot representing the

colocalisation of m6A regulators at the pre-and post-synapse in Wt and aSyn Tg primary

neurons. METTL3 colocalisation is significantly reduced in aSyn Tg neurons when compared

with Wt at the post-synapse

Figure 6. The number of synapses is increased in aSyn Tg primary neurons. (A)

Representative immunofluorescent images of DIV14 primary neurons stained with pre-

(Syn1, magenta) and post-synaptic (PSD95, green) markers and MAP2 in red. Acquired at

63x magnification, airy scan mode. Scale bar = 10μm. (B) Scatter plot for synapse counting

for Wt and aSyn Tg mouse models with a significant reduction in the number of synapses (*,

adj p value = 0.044) in aSyn Tg primary neurons when compared with Wt.

 Supplementary Figures

Figure S1. Gene expression analysis in aSyn Tg and Wt mice. (A) Principle component

analysis (PCA) plot of total Wt vs. aSyn Tg mice (PC1 = 40.5%, PC2 = 19.2%). (B) PCA

plot of aSyn Tg – 3 mo and 15 mo mice, showing a clear segregation (PC1 = 49.8%, PC2 =

23.6%). (C) Volcano plot showing that the expression of most of the genes is down regulated

in aSyn Tg mice, due to ageing (FDR < 0.05, abs(logFC) >1), aSyn Tg 15 mo vs. 3mo.

Figure S2. Differentially expressed gene pathways in aSyn Tg 15 mo vs. 3 mo. GO term

analysis comparing the enrichment of pathways, molecular function, cellular components and

biological processes of differentially methylated genes. Most pathway enrichment was

observed in immune system and immune response linked pathways including –

immunoglobulin binding, phagocytic vesicle, defense response and innate and adaptive

immune response, FC > 1.2, FDR = 1.5.

References

1. Allis D. No Title. Epigenetics, second edition [Internet]. 2015. Available from: chrome-extension://efaidnbmnnnibpcajpcglclefindmkaj/https://www.cshlpress.com/pdf/sample/2014/epigenetics2/EPIFM.pdf

2. Goldberg AD, Allis CD, Bernstein E. Epigenetics: A Landscape Takes Shape. Cell. 2007;128:635–8.

3. Saletore Y, Meyer K, Korlach J, Vilfan ID, Jaffrey S, Mason CE. The birth of the Epitranscriptome: deciphering the function of RNA modifications. Genome Biol [Internet]. 2012;13:175. Available from: http://genomebiology.biomedcentral.com/articles/10.1186/gb-2012-13-10-175

4. Dominissini D, Moshitch-Moshkovitz S, Schwartz S, Salmon-Divon M, Ungar L, Osenberg S, et al. Topology of the human and mouse m6A RNA methylomes revealed by m6A-seq. Nature [Internet]. 2012;485:201–6. Available from: https://www.nature.com/articles/nature11112

5. Liu J, Yue Y, Han D, Wang X, Fu Y, Zhang L, et al. A METTL3–METTL14 complex mediates mammalian nuclear RNA N6-adenosine methylation. Nat Chem Biol [Internet]. 2014;10:93–5. Available from: https://www.nature.com/articles/nchembio.1432

6. Ke S, Alemu EA, Mertens C, Gantman EC, Fak JJ, Mele A, et al. A majority of m 6 A residues are in the last exons, allowing the potential for 3′ UTR regulation. Genes Dev [Internet]. 2015;29:2037–53. Available from: http://genesdev.cshlp.org/lookup/doi/10.1101/gad.269415.115

7. Liu H, Begik O, Lucas MC, Ramirez JM, Mason CE, Wiener D, et al. Accurate detection of m6A RNA modifications in native RNA sequences. Nat Commun [Internet]. 2019;10:4079. Available from: https://www.nature.com/articles/s41467-019-11713-9

8. Shi H, Wei J, He C. Where, When, and How: Context-Dependent Functions of RNA Methylation

Writers, Readers, and Erasers. Mol Cell [Internet]. 2019;74:640–50. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S109727651930317X

9. Yang Y, Hsu PJ, Chen Y-S, Yang Y-G. Dynamic transcriptomic m6A decoration: writers, erasers,

readers and functions in RNA metabolism. Cell Res [Internet]. 2018;28:616–24. Available from:

https://www.nature.com/articles/s41422-018-0040-8

10. Zhao BS, Wang X, Beadell A V., Lu Z, Shi H, Kuuspalu A, et al. m6A-dependent maternal

mRNA clearance facilitates zebrafish maternal-to-zygotic transition. Nature [Internet]. 2017;542:475–

8. Available from: https://www.nature.com/articles/nature21355

11. Dinescu S, Ignat S, Lazar AD, Constantin C, Neagu M, Costache M. Epitranscriptomic Signatures

in lncRNAs and Their Possible Roles in Cancer. Genes (Basel) [Internet]. 2019;10:52. Available

from: https://www.mdpi.com/2073-4425/10/1/52

12. Licht K, Jantsch MF. Rapid and dynamic transcriptome regulation by RNA editing and RNA

modifications. Journal of Cell Biology [Internet]. 2016;213:15–22. Available from:

https://rupress.org/jcb/article/213/1/15/38525/Rapid-and-dynamic-transcriptome-regulation-by-RNA

13. Dominissini D, Moshitch-Moshkovitz S, Salmon-Divon M, Amariglio N, Rechavi G.

Transcriptome-wide mapping of N6-methyladenosine by m6A-seq based on immunocapturing and

massively parallel sequencing. Nat Protoc [Internet]. 2013;8:176–89. Available from:

https://www.nature.com/articles/nprot.2012.148

14. Bokar JA, Shambaugh ME, Polayes D, Matera AG, Rottman FM. Purification and cDNA cloning

of the AdoMet-binding subunit of the human mRNA (N6-adenosine)-methyltransferase. RNA

[Internet]. 1997;3:1233–47. Available from: http://www.ncbi.nlm.nih.gov/pubmed/9409616

15. Schwartz S, Mumbach MR, Jovanovic M, Wang T, Maciag K, Bushkin GG, et al. Perturbation of

m6A Writers Reveals Two Distinct Classes of mRNA Methylation at Internal and 5′ Sites. Cell Rep

[Internet]. 2014;8:284–96. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S2211124714004422

16. Pendleton KE, Chen B, Liu K, Hunter O V., Xie Y, Tu BP, et al. The U6 snRNA m 6 A

Methyltransferase METTL16 Regulates SAM Synthetase Intron Retention. Cell [Internet].

2017;169:824-835.e14. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0092867417305305

17. Patil DP, Chen C-K, Pickering BF, Chow A, Jackson C, Guttman M, et al. m6A RNA methylation

promotes XIST-mediated transcriptional repression. Nature [Internet]. 2016;537:369–73. Available

from: https://www.nature.com/articles/nature19342

18. Wen J, Lv R, Ma H, Shen H, He C, Wang J, et al. Zc3h13 Regulates Nuclear RNA m6A

Methylation and Mouse Embryonic Stem Cell Self-Renewal. Mol Cell [Internet]. 2018;69:1028-

1038.e6. Available from: https://linkinghub.elsevier.com/retrieve/pii/S1097276518301345

19. Agarwala SD, Blitzblau HG, Hochwagen A, Fink GR. RNA Methylation by the MIS Complex

Regulates a Cell Fate Decision in Yeast. PLoS Genet. 2012;8:e1002732.

20. Li Q, Li X, Tang H, Jiang B, Dou Y, Gorospe M, et al. NSUN2-Mediated m5C Methylation and

METTL3/METTL14-Mediated m6A Methylation Cooperatively Enhance p21 Translation. J Cell

Biochem [Internet]. 2017;118:2587–98. Available from:

https://onlinelibrary.wiley.com/doi/10.1002/jcb.25957

21. Xu K, Yang Y, Feng G-H, Sun B-F, Chen J-Q, Li Y-F, et al. Mettl3-mediated m6A regulates

spermatogonial differentiation and meiosis initiation. Cell Res [Internet]. 2017;27:1100–14. Available

from: https://www.nature.com/articles/cr2017100

22. Aguilo F, Zhang F, Sancho A, Fidalgo M, Di Cecilia S, Vashisht A, et al. Coordination of m 6 A

mRNA Methylation and Gene Transcription by ZFP217 Regulates Pluripotency and Reprogramming.

Cell Stem Cell [Internet]. 2015;17:689–704. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S1934590915004129

23. Hess ME, Hess S, Meyer KD, Verhagen LAW, Koch L, Brönneke HS, et al. The fat mass and

obesity associated gene (Fto) regulates activity of the dopaminergic midbrain circuitry. Nat Neurosci.

2013;16:1042–8.

24. Ping X-L, Sun B-F, Wang L, Xiao W, Yang X, Wang W-J, et al. Mammalian WTAP is a

regulatory subunit of the RNA N6-methyladenosine methyltransferase. Cell Res [Internet].

2014;24:177–89. Available from: https://www.nature.com/articles/cr20143

25. Wang X, Lu Z, Gomez A, Hon GC, Yue Y, Han D, et al. N6-methyladenosine-dependent

regulation of messenger RNA stability. Nature [Internet]. 2014;505:117–20. Available from:

https://www.nature.com/articles/nature12730

26. Śledź P, Jinek M. Structural insights into the molecular mechanism of the m6A writer complex.

Elife [Internet]. 2016;5. Available from: https://elifesciences.org/articles/18434

27. Wang X, Feng J, Xue Y, Guan Z, Zhang D, Liu Z, et al. Structural basis of N6-adenosine

methylation by the METTL3–METTL14 complex. Nature. 2016;534:575–8.

28. Moindrot B, Cerase A, Coker H, Masui O, Grijzenhout A, Pintacuda G, et al. A Pooled shRNA

Screen Identifies Rbm15, Spen, and Wtap as Factors Required for Xist RNA-Mediated Silencing. Cell

Rep [Internet]. 2015;12:562–72. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S2211124715006798

29. Jia G, Fu Y, Zhao X, Dai Q, Zheng G, Yang Y, et al. N6-Methyladenosine in nuclear RNA is a

major substrate of the obesity-associated FTO. Nat Chem Biol [Internet]. 2011;7:885–7. Available

from: https://www.nature.com/articles/nchembio.687

30. Zheng G, Dahl JA, Niu Y, Fedorcsak P, Huang C-M, Li CJ, et al. ALKBH5 Is a Mammalian RNA

Demethylase that Impacts RNA Metabolism and Mouse Fertility. Mol Cell [Internet]. 2013;49:18–29.

Available from: https://linkinghub.elsevier.com/retrieve/pii/S1097276512008921

31. Fu Y, Jia G, Pang X, Wang RN, Wang X, Li CJ, et al. FTO-mediated formation of N6-

hydroxymethyladenosine and N6-formyladenosine in mammalian RNA. Nat Commun [Internet].

2013;4:1798. Available from: https://www.nature.com/articles/ncomms2822

32. Alarcón CR, Lee H, Goodarzi H, Halberg N, Tavazoie SF. N6-methyladenosine marks primary

microRNAs for processing. Nature [Internet]. 2015;519:482–5. Available from:

https://www.nature.com/articles/nature14281

33. Zaccara S, Jaffrey SR. A Unified Model for the Function of YTHDF Proteins in Regulating m6A-

Modified mRNA. Cell [Internet]. 2020;181:1582-1595.e18. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0092867420305742

34. Chen J, Wei X, Yi X, Jiang D-S. RNA Modification by m6A Methylation in Cardiovascular

Disease. Ghose J, editor. Oxid Med Cell Longev [Internet]. 2021;2021:1–13. Available from:

https://www.hindawi.com/journals/omcl/2021/8813909/

35. Jiang Z, Wang Y, Li Z, Dai Z, He Y, Chu K, et al. The m6A mRNA demethylase FTO in

granulosa cells retards FOS-dependent ovarian aging. Cell Death Dis [Internet]. 2021;12:744.

Available from: https://www.nature.com/articles/s41419-021-04016-9

36. Shafik AM, Zhang F, Guo Z, Dai Q, Pajdzik K, Li Y, et al. N6-methyladenosine dynamics in

neurodevelopment and aging, and its potential role in Alzheimer's disease. Genome Biol. 2021;22:17.

37. Castro-Hernández R, Berulava T, Metelova M, Epple R, Peña Centeno T, Richter J, et al.

Conserved reduction of m^{6}A RNA modifications during aging and neurodegeneration is linked to

changes in synaptic transcripts. Proceedings of the National Academy of Sciences. 2023;120.

38. Chen YG, Chen R, Ahmad S, Verma R, Kasturi SP, Amaya L, et al. N6-Methyladenosine

Modification Controls Circular RNA Immunity. Mol Cell [Internet]. 2019;76:96-109.e9. Available

from: https://linkinghub.elsevier.com/retrieve/pii/S1097276519305465

39. Yu Z, Huang L, Xia Y, Cheng S, Yang C, Chen C, et al. Analysis of m6A modification regulators

in the substantia nigra and striatum of MPTP-induced Parkinson's disease mice. Neurosci Lett

[Internet]. 2022;791:136907. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0304394022004682

40. Schapira AHV, Cooper JM, Dexter D, Jenner P, Clark JB, Marsden CD. MITOCHONDRIAL

COMPLEX I DEFICIENCY IN PARKINSON'S DISEASE. The Lancet [Internet]. 1989;333:1269.

Available from: https://linkinghub.elsevier.com/retrieve/pii/S0140673689923660

41. Krishnan KJ, Reeve AK, Samuels DC, Chinnery PF, Blackwood JK, Taylor RW, et al. What

causes mitochondrial DNA deletions in human cells? Nat Genet [Internet]. 2008;40:275–9. Available

from: https://www.nature.com/articles/ng.f.94

42. Pickrell AM, Pinto M, Hida A, Moraes CT. Striatal Dysfunctions Associated with Mitochondrial

DNA Damage in Dopaminergic Neurons in a Mouse Model of Parkinson's Disease. The Journal of

Neuroscience [Internet]. 2011;31:17649–58. Available from:

https://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.4871-11.2011

43. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet

J [Internet]. 2011;17:10. Available from:

http://journal.embnet.org/index.php/embnetjournal/article/view/200

44. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal

RNA-seq aligner. Bioinformatics [Internet]. 2013;29:15–21. Available from:

https://academic.oup.com/bioinformatics/article/29/1/15/272537

45. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map

format and SAMtools. Bioinformatics [Internet]. 2009;25:2078–9. Available from:

https://academic.oup.com/bioinformatics/article/25/16/2078/204688

46. Cui X, Meng J, Zhang S, Chen Y, Huang Y. A novel algorithm for calling mRNA m 6 A peaks by

modeling biological variances in MeRIP-seq data. Bioinformatics [Internet]. 2016;32:i378–85.

Available from: https://academic.oup.com/bioinformatics/article/32/12/i378/2289056

47. Meng J, Cui X, Rao MK, Chen Y, Huang Y. Exome-based analysis for RNA epigenome

sequencing data. Bioinformatics [Internet]. 2013;29:1565–7. Available from:

https://academic.oup.com/bioinformatics/article/29/12/1565/291595

48. Liao Y, Smyth GK, Shi W. The Subread aligner: fast, accurate and scalable read mapping by

seed-and-vote. Nucleic Acids Res [Internet]. 2013;41:e108–e108. Available from:

https://academic.oup.com/nar/article/41/10/e108/1075719

49. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq

data with DESeq2. Genome Biol [Internet]. 2014;15:550. Available from:

http://genomebiology.biomedcentral.com/articles/10.1186/s13059-014-0550-8

50. Risso D, Ngai J, Speed TP, Dudoit S. Normalization of RNA-seq data using factor analysis of

control genes or samples. Nat Biotechnol [Internet]. 2014;32:896–902. Available from:

https://www.nature.com/articles/nbt.2931

51. Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, et al. deepTools2: a next

generation web server for deep-sequencing data analysis. Nucleic Acids Res [Internet].

2016;44:W160–5. Available from: https://academic.oup.com/nar/article-

lookup/doi/10.1093/nar/gkw257

52. Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, et al. Integrative

genomics viewer. Nat Biotechnol [Internet]. 2011;29:24–6. Available from:

https://www.nature.com/articles/nbt.1754

53. Bindea G, Mlecnik B, Hackl H, Charoentong P, Tosolini M, Kirilovsky A, et al. ClueGO: a

Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks.

Bioinformatics. 2009;25:1091–3.

54. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: A Software

Environment for Integrated Models of Biomolecular Interaction Networks. Genome Res [Internet].

2003;13:2498–504. Available from: http://genome.cshlp.org/lookup/doi/10.1101/gr.1239303

55. Wickham H. Toolbox. ggplot2. New York, NY: Springer New York; 2009. p. 65–90.

56. Koopmans F, van Nierop P, Andres-Alonso M, Byrnes A, Cijsouw T, Coba MP, et al. SynGO: An

Evidence-Based, Expert-Curated Knowledge Base for the Synapse. Neuron. 2019;103:217-234.e4.

57. R Development Core Team. R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing [Internet]. 2015. Available from: http://www.r-project.org

58. Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, et al. Simple Combinations of

Lineage-Determining Transcription Factors Prime cis-Regulatory Elements Required for Macrophage

and B Cell Identities. Mol Cell [Internet]. 2010;38:576–89. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S1097276510003667

59. Cavalcante RG, Sartor MA. annotatr: genomic regions in context. Valencia A, editor.

Bioinformatics [Internet]. 2017;33:2381–3. Available from:

https://academic.oup.com/bioinformatics/article/33/15/2381/3092365

60. Cui X, Wei Z, Zhang L, Liu H, Sun L, Zhang S-W, et al. Guitar: An R/Bioconductor Package for

Gene Annotation Guided Transcriptomic Analysis of RNA-Related Genomic Features. Biomed Res

Int [Internet]. 2016;2016:1–8. Available from: http://www.hindawi.com/journals/bmri/2016/8367534/

61. Sievert C. Interactive Web-Based Data Visualization with R, plotly, and shiny. Chapman and

Hall/CRC; 2020.

62. L Shen. C. Sievert, Interactive Web-Based Data Visualization with R, Plotly, and Shiny (Champ

Hall, 2019). 2014.

63. Kanehisa M, Sato Y, Kawashima M. <scp>KEGG</scp> mapping tools for uncovering hidden

features in biological data. Protein Science. 2022;31:47–53.

64. Jones TR, Kang IH, Wheeler DB, Lindquist RA, Papallo A, Sabatini DM, et al. CellProfiler

Analyst: data exploration and analysis software for complex image-based screens. BMC

Bioinformatics [Internet]. 2008;9:482. Available from:

https://bmcbioinformatics.biomedcentral.com/articles/10.1186/1471-2105-9-482

65. Xylaki M, Paiva I, Al-Azzani M, Gerhardt E, Jain G, Islam MR, et al. miR-101a-3p Impairs

Synaptic Plasticity and Contributes to Synucleinopathy. J Parkinsons Dis. 2023;13:179–96.

66. Koopmans F, van Nierop P, Andres-Alonso M, Byrnes A, Cijsouw T, Coba MP, et al. SynGO: An

Evidence-Based, Expert-Curated Knowledge Base for the Synapse. Neuron. 2019;103:217-234.e4.

67. Stern S, Lau S, Manole A, Rosh I, Percia MM, Ben Ezer R, et al. Reduced synaptic activity and

dysregulated extracellular matrix pathways in midbrain neurons from Parkinson's disease patients.

NPJ Parkinsons Dis. 2022;8:103.

68. Rosh I, Tripathi U, Hussein Y, Rike WA, Djamus J, Shklyar B, et al. Synaptic dysfunction and

extracellular matrix dysregulation in dopaminergic neurons from sporadic and E326K-GBA1

Parkinson's disease patients. NPJ Parkinsons Dis. 2024;10:38.

69. Licatalosi DD, Mele A, Fak JJ, Ule J, Kayikci M, Chi SW, et al. HITS-CLIP yields genome-wide

insights into brain alternative RNA processing. Nature. 2008;456:464–9.

70. Chen X, Yu C, Guo M, Zheng X, Ali S, Huang H, et al. Down-Regulation of m6A mRNA

Methylation Is Involved in Dopaminergic Neuronal Death. ACS Chem Neurosci. 2019;10:2355–63.

71. Geng Y, Long X, Zhang Y, Wang Y, You G, Guo W, et al. FTO-targeted siRNA delivery by

MSC-derived exosomes synergistically alleviates dopaminergic neuronal death in Parkinson's disease

via m6A-dependent regulation of ATM mRNA. J Transl Med. 2023;21:652.

72. He H, Zhang Q, Liao J, Lei J, Luo M, Huang J, et al. <scp>METTL14</scp> is decreased and

regulates <scp> m ^{6} A </scp> modification of α☐synuclein in Parkinson's disease. J Neurochem.

2023;166:609–22.

73. Soukup S, Vanhauwaert R, Verstreken P. Parkinson's disease: convergence on synaptic

homeostasis. EMBO J. 2018;37.

74. Liu-Yesucevitz L, Bassell GJ, Gitler AD, Hart AC, Klann E, Richter JD, et al. Local RNA

Translation at the Synapse and in Disease: Figure 1. The Journal of Neuroscience. 2011;31:16086–93.

75. Sun J, Cheng B, Su Y, Li M, Ma S, Zhang Y, et al. The Potential Role of m6A RNA Methylation

in the Aging Process and Aging-Associated Diseases. Front Genet. 2022;13.

76. Martinez De La Cruz B, Gell C, Markus R, Macdonald I, Fray R, Knight HM. m ^{6} A mRNA

methylation in human brain is disrupted in Lewy body disorders. Neuropathol Appl Neurobiol.

2023;49.

77. Yu Z, Huang L, Xia Y, Cheng S, Yang C, Chen C, et al. Analysis of m6A modification regulators

in the substantia nigra and striatum of MPTP-induced Parkinson's disease mice. Neurosci Lett.

2022;791:136907.

78. Harper JE, Miceli SM, Roberts RJ, Manley JL. Sequence specificity of the human mRNA N6-

adenosine methylase *in vitro*. Nucleic Acids Res. 1990;18:5735–41.

79. Bokar JA, Shambaugh ME, Polayes D, Matera AG, Rottman FM. Purification and cDNA cloning

of the AdoMet-binding subunit of the human mRNA (N6-adenosine)-methyltransferase. RNA.

1997;3:1233–47.

80. Ping X-L, Sun B-F, Wang L, Xiao W, Yang X, Wang W-J, et al. Mammalian WTAP is a

regulatory subunit of the RNA N6-methyladenosine methyltransferase. Cell Res. 2014;24:177–89.

81. Zou Z, Sepich-Poore C, Zhou X, Wei J, He C. The mechanism underlying redundant functions of

the YTHDF proteins. Genome Biol. 2023;24:17.

82. Gulati P, Avezov E, Ma M, Antrobus R, Lehner P, O'Rahilly S, et al. Fat mass and obesity-related

(FTO) shuttles between the nucleus and cytoplasm. Biosci Rep. 2014;34.

83. Goers J, Manning-Bog AB, McCormack AL, Millett IS, Doniach S, Di Monte DA, et al. Nuclear

Localization of α-Synuclein and Its Interaction with Histones. Biochemistry. 2003;42:8465–71.

84. Sugeno N, Jäckel S, Voigt A, Wassouf Z, Schulze-Hentrich J, Kahle PJ. α-Synuclein enhances

histone H3 lysine-9 dimethylation and H3K9me2-dependent transcriptional responses. Sci Rep.

2016;6:36328.

85. Surguchov A. α-Synuclein and Mechanisms of Epigenetic Regulation. Brain Sci. 2023;13:150.

86. Jowaed A, Schmitt I, Kaut O, Wüllner U. Methylation Regulates Alpha-Synuclein Expression and

Is Decreased in Parkinson's Disease Patients' Brains. The Journal of Neuroscience. 2010;30:6355–9.

87. Merkurjev D, Hong W-T, Iida K, Oomoto I, Goldie BJ, Yamaguti H, et al. Synaptic N6-

methyladenosine (m6A) epitranscriptome reveals functional partitioning of localized transcripts. Nat

Neurosci. 2018;21:1004–14.

88. Flamand MN, Meyer KD. The epitranscriptome and synaptic plasticity. Curr Opin Neurobiol.

2019;59:41–8.

89. Madugalle SU, Meyer K, Wang DO, Bredy TW. RNA N6-Methyladenosine and the Regulation of

RNA Localization and Function in the Brain. Trends Neurosci. 2020;43:1011–23.

90. Flamand MN, Meyer KD. m6A and YTHDF proteins contribute to the localization of select

neuronal mRNAs. Nucleic Acids Res. 2022;50:4464–83.

91. Picconi B, Piccoli G, Calabresi P. Synaptic Dysfunction in Parkinson's Disease. 2012. p. 553–72.

1012