A GAF domain mediates inositol pyrophosphate substrate channeling in PPIP5K phosphatases. (89

characters)

Pierre Raia[1], Kitaik Lee[1,2], Simon M. Bartsch[3,4], Felix Rico-Resendiz[1], Dorothea Fiedler[3,4], Michael

Hothorn[1,#]

[1]Structural Biology Laboratory, Department of Plant Sciences, University of Geneva, 30 Quai E. Ansermet,

1211 Geneva, Switzerland.

[2]Present address: Center for Structural Biology, Center for Cancer Research, National Cancer Institute (NCI),

Frederick, MD 21702-1201, U.S.A.

[3]Department of Chemical Biology, Leibniz-Forschungsinstitut für Molekulare Pharmakologie, Robert-

Rössle-Straße 10, 13125, Berlin, Germany.

[4]Institute of Chemistry, Humboldt-Universität zu Berlin, Brook-Taylor-Straße 2, 12489 Berlin, Germany

[#]Corresponding author michael.hothorn@unige.ch

ORCiDs

Pierre Raia https://orcid.org/0000-0003-3469-1922

Kitaik Lee https://orcid.org/0000-0002-7761-5586

Simon M. Bartsch https://orcid.org/0000-0002-2215-7901

Felix Rico-Resendiz https://orcid.org/0000-0003-0260-7298

Dorothea Fiedler https://orcid.org/0000-0002-0798-946X

Michael Hothorn https://orcid.org/0000-0002-3597-5698

Keywords

Inositol pyrophosphate, inositol pyrophosphate phosphatase, histidine acid phosphatase, phytase, enzyme

mechanism, GAF domain, phosphate homeostasis, *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*,

Arabidopsis thaliana.

Abstract

Inositol pyrophosphates are highly phosphorylated nutrient messengers. The final step of their biosynthesis is catalyzed by diphosphoinositol pentakisphosphate kinase (PPIP5K) enzymes, which are conserved among fungi, plants, and animals. PPIP5Ks contain an N-terminal kinase domain that generates the active messenger 1,5-InsP$_8$ and a C-terminal phosphatase domain that participates in PP-InsP catabolism. The balance between kinase and phosphatase activities controls the cellular levels and signaling capacity of 1,5-InsP$_8$. Here, we present crystal structures of the apo and substrate-bound Vip1 phosphatase domain from S. *cerevisiae* (ScVip1^{PD}). ScVip1^{PD} is a phytase-like inositol 1-pyrophosphate phosphatase with two conserved histidine phosphatase catalytic motifs. The enzyme has a strong preference for 1,5-InsP$_8$ and is inhibited by inorganic phosphate. ScVip1^{PD} has an α-helical insertion domain stabilized by a structural Zn^{2+} binding site, and a unique GAF domain that exists in an open and closed state, allowing channeling of the 1,5-InsP$_8$ substrate to the active site. Mutations that alter the active site, that restrict the movement of the GAF domain or that modify the charge of the substrate channel significantly inhibit the activity of the yeast enzyme *in vitro*, and the function of the Arabidopsis PPIP5K VIH2 *in planta*. Structural analyses of full-length PPIP5Ks suggest that the kinase and phosphatase are independent enzymatic modules. Taken together, our work reveals the structure, enzymatic mechanism and regulation of eukaryotic PPIP5K phosphatases. (221 words)

Introduction

Inositol pyrophosphates (PP-InsPs) are eukaryotic nutrient messengers (Shears, 2018). These intracellular signaling molecules consist of a densely phosphorylated six-carbon *myo*-inositol ring with pyrophosphate groups attached, for example, at position 1 (1-InsP$_7$), position 5 (5-InsP$_7$), or both positions (1,5-InsP$_8$). PP-InsPs regulate cell signaling by a variety of mechanisms, including mediating protein-protein interactions, allosterically regulating enzyme or transporter activities, or post-translationally modifying proteins (Gu *et al*, 2024; Nguyen Trung *et al*, 2022).

A central role of PP-InsPs in cellular inorganic phosphate (Pi) homeostasis is conserved among fungi (Wild *et al*, 2016; Chabert *et al*, 2023; Guan *et al*, 2023), protozoa (Cordeiro *et al*, 2017), algae (Couso *et al*, 2016), plants (Stevenson-Paulik *et al*, 2005; Zhu *et al*, 2019; Dong *et al*, 2019; Ried *et al*, 2021; Riemer *et al*, 2021; Zhou *et al*, 2021; Guan *et al*, 2022), and animals (Gu *et al*, 2017b; Li *et al*, 2020; Wang *et al*, 2020; Li *et al*, 2024). Pi homeostasis is primarily controlled by 1,5-InsP$_8$ binding to PP-InsP sensing SPX receptor proteins (Wild *et al*, 2016; Dong *et al*, 2019; Ried *et al*, 2021; Li *et al*, 2020; Zhou *et al*, 2021; Guan *et al*, 2022; Chabert *et al*, 2023). 1,5-InsP$_8$ levels are low under Pi starvation conditions, whereas 1,5-InsP$_8$ accumulates under Pi-sufficient growth conditions or upon Pi resupply (Riemer *et al*, 2021; Chabert *et al*, 2023). As cellular Pi homeostasis and other essential cellular processes appear to be mechanistically linked to 1,5-InsP$_8$ levels, understanding the regulation of 1,5-InsP$_8$ biosynthesis and catabolism is of fundamental importance.

PP-InsPs are generated from InsP$_6$ through a series of phosphorylation steps catalyzed by different small molecule kinases. In yeast, the inositol hexakiphosphate kinase Kcs1 (Saiardi *et al*, 1999; Dubois *et al*, 2002) and in plants, the inositol 1,3,4-trisphosphate 5/6-kinases ITPK1 and 2 (Laha *et al*, 2019), preferentially transfer a second phosphate to position 5 of InsP$_6$ or 1-InsP$_7$ to generate 5-InsP$_7$ and 1,5-InsP$_8$, respectively. The diphosphoinositol pentakisphosphate kinases (PPIP5Ks), known as Vip1 in *S. cerevisiae*, Asp1 in *Schizosaccharomyces pombe* (Fridy *et al*, 2007; Mulugu *et al*, 2007; Dollins *et al*, 2020), PPIP5K in amoeba and in animals (Gokhale *et al*, 2011; Wang *et al*, 2012; Weaver *et al*, 2013; Gu *et al*, 2017b; Nair *et al*, 2018; Desfougères *et al*, 2022) or VIP/VIH in plants (Desai *et al*, 2014; Laha *et al*, 2015; Couso *et al*, 2016; Zhu *et al*, 2019; Shukla *et al*, 2021; Laurent *et al*, 2024), preferentially transfer a phosphate to position 1, to generate 1-InsP$_7$ and 1,5-InsP$_8$, respectively. Consistent with the function of 1,5-InsP$_8$ as messenger in Pi homeostasis in Arabidopsis, *vih1/vih2* loss-of-function mutants have severely reduced 1,5-InsP$_8$ levels and display constitutive Pi starvation responses (Zhu *et al*, 2019; Dong *et al*, 2019; Ried *et al*, 2021).

In addition to the conserved Pi homeostasis-related phenotypes, loss-of-function of PPIP5K enzymes results in altered growth, cell and vacuolar morphology, chromosome segregation, host colonization and DNA damage-induced autophagy (Mulugu *et al*, 2007; Pascual-Ortiz *et al*, 2018; Dollins *et al*, 2020; Gu *et al*, 2017a). PPIP5K mutants are associated with hearing loss in humans and mice (Yousaf *et al*, 2018). In human cell lines, depletion of PPIP5K activity induced broad changes in cell metabolism (Gu *et al*, 2021). In plants, PPIP5K activity has been linked to jasmonic acid and salicylic acid hormone signaling, and to cell wall architecture (Laha *et al*, 2015; Shukla *et al*, 2021; Gulabani *et al*, 2022; Laurent *et al*, 2024).

There are several PP-InsP catabolic enzymes, including a family of histidine acid phosphatases
located in C-terminal half of PPIP5Ks (Mulugu *et al*, 2007). Thus, PPIP5Ks are dual-function enzymes that
contain two distinct domains with opposing enzymatic activities: An N-terminal kinase domain that
preferentially generates 1,5-InsP$_8$ from 5-InsP$_7$ (Gokhale *et al*, 2011; Dollins *et al*, 2020; Benjamin *et al*,
2022a), and a C-terminal phosphatase domain with inositol 1-pyrophosphate phosphatase activity (Pöhlmann
et al, 2014; Wang *et al*, 2015a; Nair *et al*, 2018; Dollins *et al*, 2020; Benjamin *et al*, 2022a). The relative PP-
InsP kinase and phosphatase activities in PPIP5K are thought to modulate the cellular levels of 1,5-InsP$_8$ (Gu
et al, 2017b; Zhu *et al*, 2019; Gu *et al*, 2024).

PPIP5K loss-of-function mutants show altered PP-InsP pools in various eukaryotes (Mulugu *et al*,
2007; Pascual-Ortiz *et al*, 2018; Dollins *et al*, 2020; Chabert *et al*, 2023; Gu *et al*, 2016, 2017a, 2021; Laha
et al, 2015; Couso *et al*, 2016; Zhu *et al*, 2019; Riemer *et al*, 2021; Laurent *et al*, 2024). PPIP5K knockout
mutants have reduced 1-InsP$_7$ and 1,5-InsP$_8$ levels, while the 5-InsP$_7$ substrate of the PPIP5K kinase domain
accumulates (Qiu *et al*, 2020; Chabert *et al*, 2023; Riemer *et al*, 2021; Laurent *et al*, 2024). 1,5-InsP$_8$ pools
can be restored to wild-type like levels in genetic rescue experiments with phosphatase-dead but not with
kinase-dead versions of the corresponding PPIP5K enzyme (Gu *et al*, 2017a; Pascual-Ortiz *et al*, 2018; Zhu
et al, 2019; Dollins *et al*, 2020; Gu *et al*, 2021).

How regulation of PPIP5Ks controls 1,5-InsP$_8$ pools in response to changes in Pi nutrient availability
is not well understood. Currently, it is known that the ATP/ADP ratio decreases in response to Pi starvation,
which in turn reduces 1,5-InsP$_8$ synthesis by the kinase domains of human PPIP5K1/2 or plant VIH1/2 (Gu
et al, 2017b; Zhu *et al*, 2019). Pi itself has been reported as an inhibitor of the phosphatase domain of human
and fungal PPIP5Ks (Gu *et al*, 2017b; Zhu *et al*, 2019; Benjamin *et al*, 2022a).

The structure and enzymatic mechanism of the human and fungal PPIP5K kinase domains have been
well established (Wang *et al*, 2012, 2014; Benjamin *et al*, 2022b). However, currently there is no structural
information on the PPIP5K phosphatase domain and its enzymatic mechanism and regulation remain to be
elucidated. Based on sequence comparison, the PPIP5K phosphatase domain has been assigned to the
histidine acid phosphatase superfamily (Mulugu *et al*, 2007), specifically to the branch 2, which includes the
well-characterized *Escherichia coli* and fungal phytases, as well as glucose-1-phosphate, and acid
phosphatases (Rigden, 2008). The PPIP5K phosphatase domain contains the catalytic RHxxR motif, which is
conserved among histidine acid phosphatases (Mulugu *et al*, 2007). However, the lack of sequence
conservation in the substrate-binding domains within the superfamily precludes predictions about substrate
recognition and specificity. A "cryptic" pleckstrin homology (PH) domain located near the conserved RHxxR
motif has been hypothesized to play a critical role in substrate binding (Gokhale *et al*, 2011). PH domains,
typically found in signaling proteins, bind phospholipids and molecules derived from phospholipid
headgroups, including inositol phosphate and PP-InsPs (Scheffzek & Welti, 2012). We now report the
structure of a PPIP5K phosphatase domain and characterize its substrate binding and enzymatic mechanism,
its regulation by nutrients and its structural organization within the full-length enzyme.

Results

Overall structure of the ScVip1 kinase and phosphatase domains

We first attempted to characterize the structure and catalytic mechanism of the histidine acid phosphatase domain (PD) from plant PPIP5Ks (Desai *et al*, 2014; Laha *et al*, 2015; Zhu *et al*, 2019). However, we were unable to express sufficient amounts of the isolated phosphatase domains from either *Arabidopsis thaliana* VIH1/VIH2 or from *Marchantia polymorpha* VIP1 (Laurent *et al*, 2024) (Fig EV1A). Therefore, we expressed and purified the phosphatase domains from *S. cerevisiae* Vip1 (ScVip1^{PD}, residues 536-1107) as well as its N-terminal kinase domain (ScVip1^{KD}, residues 186-522) in baculovirus-infected insect cells (see Methods).

The structure of the ScVip1^{KD} was solved by molecular replacement to a resolution of ~1.2 Å (Table 1). Apo crystals of the ScVip1^{PD} initially diffracted to ~3.1 Å. The structure was solved by molecular replacement - single wavelength anomalous diffraction (MR-SAD) on a platinum derivative (Table 1). In this structure, a large disordered loop (residues 851-918) was located and subsequently replaced by a short Gly-Ser-Ser-Gly linker. Crystallization of ScVip1$^{PDΔ848-918}$ yielded a second crystal form diffracting at 3.4 Å resolution (Table 1).

The structure of ScVip1^{KD} in complex with adenosine diphosphate revealed the typical ATP-grasp fold and substrate binding sites previously seen in other PPIP5K structures (Wang *et al*, 2012; Benjamin *et al*, 2022b) (Fig 1A,B). ScVip1^{KD} shares significant sequence and structural homology with HsPPIP5K2 and with the kinase domain of *S. pombe* Asp1 (Figs EV2, EV3). The C-terminal α11 helix of the ScVip1 kinase domain (residues 498-520) is well defined in our structure (Figs 1B, EV2).

The two apo structures of ScVip1^{PD} revealed a conserved histidine acid phosphatase core with two insertion domains (Fig 1C). The N-terminus of ScVip1^{PD} (residue 536) and the most C-terminal α9 helix (residues 1079-1094) are well defined by electron density (Fig 1C). Based on these domain boundaries, we hypothesize that the ScVip1 kinase and phosphatase domains are surrounded by large, unstructured N- and C-terminal extensions, and are connected by a flexible ~15 amino acid linker (Figs 1A, EV2, see below). Plant and human PPIP5Ks contain sequence-diverse linkers of similar length between their kinase and phosphatase domains (Fig EV2). A conserved metal binding site occupied by a Zn^{2+} ion is formed by His651 and His1064 from the histidine acid phosphatase core, and by Cys793 and His836 from the α-helical insertion domain (Figs 1C, EV2, see below).

A PPIP5K unique GAF domain

Next, we performed structural homology searches with ScVip1^{PD} as a search model using the DALI web server (http://ekhidna2.biocenter.helsinki.fi/dali/) (Holm & Rosenström, 2010). We found that ScVip1^{PD} shares the core phosphatase domain fold and the α-helical insertion domain with bacterial, plant and human histidine acid phosphatases (Figs 2A, EV4). The relative orientation of the α-helical insertion domain is conserved among histidine acid phosphatases with very different substrate preferences (Fig 2A), including

phytases that hydrolyze $InsP_6$ (Fig 2B). We located a second insertion domain specific to PPIP5Ks in a
position normally occupied by the much smaller β1-insert in other histidine acid phosphatases (Fig 2A).
Analysis with DALI revealed a weak structural homology of this second insertion domain with bacterial and
human GAF (Aravind & Ponting, 1997) or PAS (Ponting & Aravind, 1997) domains involved in light,
metabolite and cyclic nucleotide sensing (Ho *et al*, 2000; Henry & Crosson, 2011) (Figs 2C, EV4). The GAF
domain is conserved among fungal, plant and animal PPIP5Ks (Fig EV2). The central β-sheet of the
ScVip1^{PD} GAF domain is reminiscent of the inositol phosphate / inositol polyphosphate binding surface
previously identified in PH domains, but its topology is different (Fig 2C) (Milburn *et al*, 2003; Wang *et al*,
2015b).

ScVip1^{PD} specifically hydrolyses 1,5-InsP$_8$

We identified 1,5-$InsP_8$ as the preferred substrate for the isolated ScVip1 phosphatase domain in
quantitative nuclear magnetic resonance (NMR) phosphatase assays using ^{13}C-labeled substrates (Figs 3A,
EV5). ScVip1^{PD} is a specific inositol pyrophosphate phosphatase, with a high preference for the 1-position of
PP-InsPs (Figs 3A, EV5), as previously reported (Dollins *et al*, 2020). In malachite green-based phosphatase
assays, we derived similar K_M values for 1,5-$InsP_8$ and 1-$InsP_7$, but the catalytic efficiency of ScVip1^{PD} for
1,5-$InsP_8$ is ~18fold higher when compared to 1-$InsP_7$ (Fig 3B). Notably, 1,5-$InsP_8$ is the reaction product of
the N-terminal kinase domain of ScVip1 (Mulugu *et al*, 2007).

A sequence motif central to catalysis in histidine acid phosphatases is located in the active site cleft
sandwiched between the phosphatase core and the α-helical insertion domain (Fig 3C). The motif with the
consensus sequence Arg-His-x-x-Arg (RHxxR, where x can be any amino acid) is structurally conserved
between ScVip1^{PD} and bacterial phytases (Fig 3C). Mutation of either Arg547, His548 or Arg551 to alanine
greatly reduces 1,5-$InsP_8$ hydrolysis in NMR- and malachite green-based enzyme assays, respectively (Figs
3D,E, EV5). This is consistent with the established function of the two arginine residues in $InsP_6$ substrate
binding in bacterial phytases (Lim *et al*, 2000), and the essential role of the nucleophilic acceptor histidine in
catalysis (Ostanin *et al*, 1992). Asp304 in *E. coli* phytase is the proton donor involved in the degradation of
the phosphohistidine intermediate (Ostanin & Van Etten, 1993). In our ScVip1^{PD} structures, a glutamate
residue (Glu991) occupies the position of the phytase proton donor aspartate (Fig 3C). Mutation of Glu911 to
alanine activates the enzyme (Figs 3D,E, EV5, see below). Similarly, deletion of the large disordered loop in
the α-helical insertion domain (ScVip1$^{PDΔ848-918}$) moderately activates the engineered enzyme *in vitro* (Fig
3D).

In conclusion, the isolated phosphatase domain of ScVip1 is an inositol 1-pyrophosphate
phosphatase with a substrate preference for 1,5-$InsP_8$ and with an active site similar to $InsP_6$ metabolizing
phytases.

 Inhibition of ScVip1^{PD} by Pi

Gu et al. previously reported that low millimolar concentrations of Pi inhibited the inositol 1-
pyrophosphate phosphatase activity of human PPIP5K1 and PPIP5K2 (Gu *et al*, 2017b). Addition of 10 mM
Pi inhibited the phosphatase activity of ScVip1 (Zhu *et al*, 2019). Similar results were reported for full-length
Asp1 (Benjamin *et al*, 2022a). We found that ScVip1^{PD} is inhibited by Pi in a concentration-dependent
manner, as inferred from NMR-based assays (Fig 4A). As previously reported for SpAsp1 (Benjamin *et al*,
2022a), sulfate was less effective than Pi in inhibiting the ScVip1^{PD} phosphatase activity (Fig 4B). Nitrate
had no detectable effect on the hydrolysis of 1,5-InsP$_8$ (Fig 4C). We conclude that Pi at concentrations found
in the yeast cytosol (Auesukaree *et al*, 2004), directly inhibits the enzymatic activity of the isolated ScVip1
phosphatase domain.

GAF domain movements control enzyme activity

All four molecules in the asymmetric unit of our ScVip1^{PD} crystals display the GAF domain in an
"open" conformation (Figs 5A, EV6, Table 1). In ScVip1$^{PDΔ848-918}$ crystals two molecules in the asymmetric
unit also adopt the "open" conformation (Figs 5A, EV6, Table 1). Crystal packing interactions between two
adjacent GAF domains in our ScVip1$^{PDΔ848-918}$ crystals are mediated in part by a second Zn^{2+} binding site. The
Zn^{2+} ion is coordinated along a pseudo two-fold axis involving residues His573 and Glu575 from the GAF
domain (Fig EV6). Given that ScVip1^{PD} behaves as a monomer in solution (Fig EV7) and that the histidine
and glutamate residues are not conserved among other PPIP5Ks (Fig EV2), the second Zn^{2+} coordination site
is likely a crystallization artifact (Fig EV6).

Next, we replaced the catalytic loop residues Arg547, His548 and Arg551 in ScVip1$^{PD\ Δ848-918}$ with
alanine and obtained crystals in complex with the substrate 1,5-InsP$_8$ diffracting at ~2.4 Å resolution
(ScVip1$^{PD\ Δ848-918\ RHR-AAA}$, Table 1). Notably, the GAF domain adopts a "closed" conformation in the substrate-
bound structure (Fig. 5A). We hypothesized that the movements of the GAF domain are enabled by a hinge
region in close proximity to the catalytic loop (RHxxR motif) of the enzyme (Fig 5A,B). Consequently,
mutation of the hinge residues Pro553, Gly646 and Gly647 (P553V-G646V-G647A) inactivated ScVip1^{PD} in
NMR- and malachite green-based phosphatase assays, respectively (Fig 5B-D). This suggests that movement
of the GAF domain is part of the enzymatic cycle of ScVip1^{PD}. Notably, the catalytic loop (Arg547, His548,
Arg551) does not appear to be involved in the domain movement when comparing the open and closed
conformation structures (Fig 5A). The conserved Zn^{2+} binding site in ScVip1^{PD} positions His836 to form a
salt bridge with Asp550, apparently stabilizing the position of the catalytic loop (Fig 5B). Mutation of the
Zn^{2+} binding site residues His651 or Cys793 to alanine reduces the phosphatase activity in our enzyme
assays, suggesting that the Zn^{2+} binding site is involved in the structural stabilization of the enzyme and,
importantly, of its catalytic loop (Fig 5B-E). Consistent with this, mutation of Zn^{2+} coordinating residues
reduces the structural stability of ScVip1^{PD}, whereas excess of ZnCl$_2$ further stabilizes the enzyme in thermal
shift assays *in vitro* (Fig 5E, EV7).

The GAF domain is involved in substrate binding and channeling

In our ScVip1$^{PD \ \Delta848-918 \ RHR-AAA}$ – 1,5-InsP$_8$ complex structure, we found a domain-swapped

crystallographic dimer stabilized by an intermolecular disulfide bridge (Fig EV8). One molecule of 1,5-InsP$_8$

is bound to each monomer, either on the surface of the enzyme involving the GAF domain or near the

catalytic center (Fig 6A, EV8). The GAF domain adopts a closed conformation in both monomers (Fig 6A).

In both binding sites, all phosphate groups of 1,5-InsP$_8$ are well defined by electron density (Fig 6B), with

the 1-β-phosphate facing away from the active site (Fig 6C). Conserved basic residues from the GAF domain

and the α-insertion domain contribute to the formation of the outer substrate binding surface (Fig 6C). A

large network of lysine residues from the GAF domain and from the base of the α-insertion domain forms the

second substrate binding surface, which is located in close proximity to the RHxxR catalytic loop (Fig 6C).

Most of the basic residues contributing to the outer and the inner 1,5-InsP$_8$ binding surfaces map to the

central β-sheet of the GAF domain (Fig 6C,D). Comparison of the outer and inner substrate binding sites in

ScVip1^{PD} suggests that the 1,5-InsP$_8$ substrate may undergo a ~ 90º rotation upon entering the active site (Fig

6D).

We next compared the open conformation of apo ScVip1^{PD} with the substrate-bound structures (Fig

7A). We found that the conformational changes strongly alter the surface charge distribution of the enzyme,

with the GAF domain opening and closing on a large and well conserved basic surface area surrounding the

substrate binding sites and formed by the GAF and α-insertion domains (Fig 7A,B). By mapping the

positions of the 1,5-InsP$_8$ molecules bound to the outer and inner substrate binding sites, we uncovered a

potential substrate binding channel (Fig 7C). We hypothesize that the 1,5-InsP$_8$ substrate first targets the

outer substrate binding site of ScVip1^{PD} in its open conformation. Substrate binding to this outer site may

trigger closure of the GAF domain, resulting in rotation of the substrate and its translocation to the active site

(Fig 7C). In its closed conformation, the PPIP5K GAF domain contributes to the formation of a substrate

channel that may facilitate transport of 1,5-InsP$_8$ to the catalytic center of the enzyme as well as the exit of

the 5-InsP$_7$ product from the enzyme (Fig 7C).

Mutation of highly conserved residues from the outer substrate binding site (Lys558, Lys605,

Lys732, Lys817 to alanine) (Fig 7D) inactivated the enzyme (Fig 7E,F), suggesting that 1,5-InsP$_8$ is indeed

initially bound to the surface of the enzyme before entering the active site. We next mutated residues that

contribute to the PPIP5K substrate channel. Mutation of Lys817, Asp819 and Ser821 originating from the α-

helical insertion domain (Fig 7D) moderately reduced the 1,5-InsP$_8$ phosphohydrolase activity of ScVip1^{PD} in

both NMR- and malachite green-based assays (Fig 7E,F). In contrast, mutation of 1,5-InsP$_8$ binding residues

from the GAF domain (Glu576, Lys623, Gln625, Lys627) inactivated the enzyme (Fig 7D-F). Mutations

close to the active site of the enzyme (Lys554, Lys557, Lys644) also reduced the activity of ScVip1^{PD} (Fig

7D-F), but in addition structurally destabilized the enzyme (Fig EV7). We therefore generated an additional

single Lys554 to alanine mutant, which is structurally intact (Fig EV7) but which also showed very low

enzyme activity in our assays (Fig 7D-F). Taken together, our structure-function analyses suggest that

PPIP5K phosphatase domains contain two 1,5-InsP$_8$ binding sites that contribute to substrate channeling into
the active site.

An active site glutamate mediates substrate specificity of ScVip1^{PD}

As described above, the canonical phytase proton donor aspartate is replaced by Glu991 in ScVip1,
which adopts different conformations in our apo and substrate-bound crystal structures (Figs 3C, 8A,B).
Replacement of Glu991 with alanine activates ScVip1^{PD} (Fig 3D,E). In addition, we observed that the
ScVip1^{PD} Glu991 to Ala mutant readily accepts 1-InsP$_7$ as a substrate in both NMR and malachite green-
based phosphatase assays, but remains unable to hydrolyze InsP$_6$ (Fig 8C,D). It has been previously reported
that mutation of Asp304 in AppA strongly alters the stereospecificity of the *E. coli* enzyme (Acquistapace *et*
al, 2022) Notably, multiple inositol polyphosphate phosphatases that lack strong stereospecificity for InsP$_6$
are characterized by an alanine residue at this position (Fig 8A,B) (Stentz *et al*, 2014). We conclude that
Glu991, which is conserved among PPIP5Ks from different species (Fig EV2) is involved in the selection of
the 1,5-InsP$_8$ substrate. The inositol ring occupies a different position in our ScVip1^{PD} – 1,5-InsP$_8$ complex
structure when compared to known phytases (Fig 8B). We speculate that this is an adaptation to the
hydrolysis of a pyrophosphate substrate. The long and uniquely shaped substrate channel may play an
additional role in the selection of 1,5-InsP$_8$ (Fig 7C).

Mutation of key motifs inhibit the function of the VIH2 phosphatase domain in planta

Returning to our original plan to characterize the catalytic mechanism of plant PPIP5K phosphatases
(see above), we next mapped all residues and sequence motifs characterized in ScVip1^{PD} to AtVIH2 from
Arabidopsis thaliana, which shares 25% sequence identity with ScVip1^{PD} (Figs 9A, EV2). We have
previously reported that expression of a phosphatase-dead variant of full-length AtVIH2 (AtVIH2$^{R372A-H373A}$)
in Col-0 wild-type background reduced rosette size and intracellular Pi levels (Zhu *et al*, 2019). We used
these phenotypes to perform structure-function studies with the AtVIH2 phosphatase domain. Constitutive
expression of wild-type AtVIH2 had no detectable effect on either rosette area or cellular Pi levels (Fig 9B-
E). Expression of a kinase-dead version of the enzyme (AtVIH2$^{K219A-D292A}$) resulted in hyperaccumulation of
cellular Pi, as previously reported (Fig 9B-E) (Zhu *et al*, 2019). Three different mutant combinations
targeting the catalytic core of the phosphatase in Arabidopsis VIH2 (compare Fig 3C-E) were all associated
with significantly reduced rosette areas and cellular Pi levels (highlighted in blue in Fig 9A-E). Mutations
targeting the conserved structural Zn^{2+} binding site (compare Fig 5B-E) were also associated with reduced
rosette areas and cellular Pi levels (highlighted in purple in Fig 9A-E). A mutant in the phosphatase hinge
region showed less severe growth phenotypes and only slightly reduced Pi levels (shown in red in Fig 9A-E),
whereas the corresponding mutations in ScVip1^{PD} strongly affected the activity and structural stability of the
enzyme *in vitro* (Fig 5A,C-E). Consistent with the moderate effect of mutations in the α-helical insert
contributing to the outer 1,5-InsP$_8$ binding site (Fig 7D-F), the corresponding AtVIH2$^{R722S-D724A-T726A}$ mutant
looks similar to the AtVIH2 wild-type expressing control (Fig 9A-E). Importantly, mutations targeting either

9 9

the outer or inner GAF domain binding surfaces (compare Fig 7D-F) both resulted in a severe reduction in

rosette area and in low Pi levels (shown in yellow in Fig 9A-E). In conclusion, our mutational analysis of the

AtVIH2 phosphatase domain confirms the importance of the catalytic center of the enzyme and highlights

the critical contributions of the outer and inner 1,5-InsP$_8$ binding sites in the PPIP5K-specific GAF domain.

The ScVip1 kinase and phosphatase domains are independent enzymatic modules

We expressed and purified full-length SpAsp1 from *E. coli* (Fig 10A), as previously described

(Benjamin *et al*, 2022a), applied the apo enzyme to carbon-coated grids and performed negative-stain

electron microscopy (Fig 10B, see Methods). We calculated a low-resolution envelope from two-dimensional

projections of single particles and docked the experimental structures of the sequence-related ScVip1 kinase

and phosphatase domains (Figs 10C, EV2). Full-length SpAsp1 adopts a L-shaped conformation (Fig 10C).

The N-terminal kinase and the C-terminal phosphatase domains are perpendicular to each other (Fig 10C).

The exact orientation of the kinase domain is somewhat ambiguous but is reasonably constrained by the size

of the 16 amino acid linker (Fig 10C). In our model, the catalytic centers of the kinase and phosphatase

modules are far apart, at least in the conformation adopted by the apo enzyme on the electron microscope

grid (Fig 10C). Consistent with this more open conformation of PPIP5K, the inositol 1-pyrophosphate

phosphatase activity of ScVip1[KD-PD] (residues 186-1107) is comparable to that of ScVip1[PD] (Fig 10D). Next,

we performed hydrogen/deuterium exchange mass spectrometry experiments on ScVip1[KD-PD]. Addition of

1,5-InsP$_8$ to the apo enzyme protects regions from the α-helical insertion domain and the GAF domain (Figs

10E, EV9), further supporting our substrate binding model for ScVip1[PD] (Fig 6C), and the proposed

dynamics of the GAF domain (Figs 5A, 7A-C). Addition of an ATP substrate analog reveals additional

protected regions in the vicinity of the active site of the kinase domain (Figs 10E, EV9). No changes

consistent with the formation of new kinase – phosphatase domain interfaces are observed upon addition of

either the phosphatase or kinase substrate (Figs 10E, EV9). Taken together, these experiments suggest that

the PPIP5K kinase and phosphatase modules are enzymatically and structurally independent entities.

Discussion

PPIP5Ks play a central role in inositol pyrophosphate metabolism and signaling (Gu *et al*, 2024).

The enzyme activity, substrate specificity, 3-dimensional structure and catalytic mechanism of the N-

terminal kinase domain of PPIP5K are well established (Gokhale *et al*, 2011; Wang *et al*, 2012; Weaver *et

al*, 2013; An *et al*, 2019; Benjamin *et al*, 2022a, 2022b), but the C-terminal phosphatase domain has only

been partially characterized biochemically (Pöhlmann *et al*, 2014; Wang *et al*, 2015a; Nair *et al*, 2018;

Dollins *et al*, 2020; Benjamin *et al*, 2022a), and has resisted structural analysis (Gu *et al*, 2024). We were

able to overcome this problem by focusing on the phosphatase domain of ScVip1, by using a eukaryotic

expression system, by fine-mapping the phosphatase domain boundaries and by engineering of a large

unstructured loop (see Methods, Table 1). The structure of ScVip1[PD] confirms its evolutionary relationship

with bacterial and animal histidine acid phosphatases (Mulugu *et al*, 2007). We found that the active site of

ScVip1^{PD} is very similar to that of InsP$_6$ hydrolyzing phytases, with both enzymes sharing the structurally conserved RHxxR catalytic motif (Fig 3C-E) (Mulugu *et al*, 2007; Rigden, 2008).

A second catalytic histidine (His993 in ScVip1, His807 in SpAsp1) was originally defined by sequence comparison with *E. coli* AppA (Fridy *et al*, 2007; Mulugu *et al*, 2007), where the corresponding His303 is involved in InsP$_6$ substrate binding and in catalysis (Ostanin *et al*, 1992; Lim *et al*, 2000). Structural comparisons of EcAppA and ScVip1^{PD} show that this motif has been incorrectly assigned. His303 in the bacterial phytase corresponds to a poorly conserved lysine residue (Lys990) in ScVip1 (Figs EV10, EV2). His993 in ScVip1 or His807 in SpAsp1 are not part of the active site but rather map to an α-helical segment buried in the phosphatase domain (Fig EV10). Thus, substitution of SpAsp1 H807 with alanine may disrupt the structural integrity of the enzyme rather than affect its catalytic activity, providing an alternative explanation for the reduced enzyme activities and associated phenotypes reported for this mutant variant (Pöhlmann *et al*, 2014; Wang *et al*, 2015a; Pascual-Ortiz *et al*, 2018). Our *in vitro* (Fig 3) and *in vivo* (Fig 9) analyses of the catalytic center of the enzyme now provide a reliable set of mutant combinations for engineering phosphatase-dead PPIP5Ks for genetic rescue experiments in various eukaryotes.

A conserved aspartate (Asp304) has been previously reported as the proton donor in *E. coli* phytase (Ostanin & Van Etten, 1993) and mutation of Asp304 strongly alters the substrate stereospecificity of this enzyme (Acquistapace *et al*, 2022). In ScVip1^{PD} this aspartate is replaced by glutamate (Glu991), which is conserved among PPIP5Ks from different species, and which adopts different conformations in our apo and substrate-bound structures (Fig 8A,B). The ScVip1^{E991A} mutant exhibits an overall increase in enzyme activity and reduced stereospecificity (Fig 8C,D). We hypothesize that both Asp304 in *E. coli* phytase and Glu991 in ScVip1^{PD} influence the size and charge distribution of the substrate binding site, with mutations increasing the processivity of the enzyme at the expense of substrate selectivity.

Mapping previously identified SpAsp1 missense mutations from a genetic suppressor screen onto our ScVip1^{PD} structure revealed that the His686 (His836 in ScVip1^{PD}) to Tyr and Cys643 (Cys793 in ScVip1^{PD}) to Tyr mutants target the structural Zn^{2+} binding site in PPIP5K (Fig 1C). This further highlights the contribution of the Zn^{2+} binding site to the structural integrity of the enzyme (Fig EV7) (Garg *et al*, 2020).

In the case of ScVip1^{PD} , a K_M of ~ 5 μM has been reported for 1-InsP$_7$ (Dollins *et al*, 2020). Using saturating substrate concentrations, we have derived K_M values of ~70 μM and 45 μM for 1-InsP$_7$ and 1,5-InsP$_8$, respectively, and 1,5-InsP$_8$ is the preferred substrate of ScVip1^{PD} *in vitro* (Fig 3A,B). For comparison, the PP-InsP phosphatase Siw14 from yeast has a K_M for 5-InsP$_7$ of ~ 10 μM (Wang *et al*, 2018a). The cytosolic concentration of 1,5-InsP$_8$ in *S. cerevisiae* has been recently estimated at ~ 0.3 μM (Chabert *et al*, 2023). However, the local concentration in proximity to the N-terminal Vip1 kinase domain may be higher.

Our structures reveal that the insertion GAF domain is a distinctive structural feature of PPIP5K phosphatases (Fig 2A). In phytases, the β1-insert maps to the position of the GAF domain. It has been previously demonstrated that this β1-insert undergoes conformational changes upon InsP$_6$ binding (Fig 2) (Lim *et al*, 2000). Our structures reveal that the PPIP5K GAF domain undergoes much larger conformational

changes and plays a pivotal role in substrate binding and in channeling of the substrate to the active site (Figs
5-7). This is further highlighted by the growth phenotypes of mutant versions of AtVIH2 targeting the GAF
domain (Fig 9). Despite the low degree of sequence conservation, substrate binding, catalysis and regulation
appear to be well conserved among PPIP5Ks from different organisms (Fig 9).

In our 1,5-InsP$_8$ complex structure, one substrate molecule is bound to the outer GAF domain, the
second molecule maps close to the active site (Fig 6A-C). It is noteworthy that a PP-InsP "capture site"
outside the active site has also been identified in the PPIP5K kinase domain (Wang *et al*, 2014).
Orthophosphate is a known inhibitor of human and yeast PPIP5K phosphatase domains, thereby regulating
1,5-InsP$_8$ levels and contributing to phosphate homeostasis (Gu *et al*, 2017b; Zhu *et al*, 2019; Benjamin *et al*,
2022a). We speculate that the inhibitory effect of orthophosphate in ScVip1^{PD} (Fig 4A) may be caused by the
binding of this metabolite to the highly basic substrate channel, where it may block access of the 1,5-InsP$_8$
substrate to the outer substrate binding surface (Fig 7A,C). We attempted to further characterize this
intriguing regulatory mechanism, but ScVip1^{PD} crystals grown in the presence of Pi, sulfate or tungstate did
not diffract.

The low-resolution envelope of full-length SpAsp1 and our hydrogen/deuterium exchange mass
spectrometry experiments collectively indicate that the PPIP5K kinase and phosphatase domains share only a
small interface. The active sites of the kinase and phosphatase domains are situated at a considerable distance
from one another, which is incompatible with the hypothesis of direct substrate channeling (Fig 10C,E). A
similar L-shaped arrangement of kinase and phosphatase domains has been previously observed in the
crystal structure of fructose-2,6-bisphosphate kinase phosphatase (Lee *et al*, 2003). Consistent with this very
open architecture of the enzyme, the phosphatase activity of full-length ScVip1 is not drastically different
from that of isolated ScVip1^{PD} (Fig 10D). However, this appears not to be the case for the kinase activity in
human PPIP5K2, where the isolated kinase domain appeared much more active compared to the full-length
enzyme (Yousaf *et al*, 2018). The domain boundaries of the PPIP5K kinase and phosphatase domains
suggest the presence of large unstructured N- and C-terminal loops, as well as the occurence of a large
disordered linker region within the α-helical insertion domain the phosphatase module (Figs 1A-C, EV2).
These sequence-diverse linker regions may represent protein interaction sites (Machkalyan *et al*, 2016;
Yousaf *et al*, 2018) that could alter the architecture and regulation of the full-length enzyme. Moreover, the
linkers are found phosphorylated in yeast (Fig EV2), suggesting that additionals regulatory mechanisms of
PPIP5K remain to be elucidated.

Methods
Expression construct cloning
The kinase (residues 186-522, ScVip1^{KD}), phosphatase (residues 536-1107, ScVip1^{PD}), loop-truncated
phosphatase (residues 536-1107, Δ848-918, ScVip1PD$^{Δ848-918}$), and kinase-phosphatase tandem domains
(residues 186-1107, ScVip1^{KD-PD}) from *Saccharomyces cerevisiae* Vip1 were cloned from synthetic genes
(Twist Bioscience) codon-optimized for expression in *Trichoplusia ni* (Tnao) cells (Invitrogen) in vector

pBB3, providing tobacco etch virus protease (TEV) cleavable N-terminal 10xHis and twin StrepII affinity
tags. Point mutations in the ScVip1 phosphatase domain were generated by site-directed mutagenesis and
subcloned in pBB3. For crystallization optimization, the ScVip1$^{PD\ \Delta848-918}$ construct was engineered by
replacing the flexible loop with a short Gly-Ser-Ser-Gly linker. The ScVip1$^{PD\ \Delta848-918}$ construct was further
optimized for co-crystallization with 1,5-InsP$_8$ by replacing Arg547, His548 and Arg551 in the catalytic
center with alanines. Full-length Asp1 from *Schizosaccaromyces pombe* was cloned from a synthetic gene
codon-optimized for expression in *Escherichia coli* into the pMH-HSSumo vector, providing N-terminal
small ubiquitin-like modifier fusion protein (Sumo) containing 7xHis and StrepII affinity tags, as described
(Hothorn *et al*, 2011; Benjamin *et al*, 2022a).

Protein expression and purification
ScVip1^{KD}, ScVip1^{PD}, ScVip1$^{PD\ \Delta848-918}$, ScVip1PD $^{\Delta848-918\ RHR-AAA}$, ScVip1^{KD-PD}, and their point-mutated
versions were expressed in Tnao cells (Invitrogen) using recombinant baculoviruses (Hashimoto *et al*, 2010).
Typically, 10 mL of recombinant P3 baculovirus was used to infect 200 mL of Tnao insect cells growing in
SF900 media (Gibco) at a cell density of approximately 2.0×10^6 cells/mL. The cells were incubated for 72 h
at 28°C with shaking at 110 rpm. Cells were harvested by centrifugation at 1,000 g for 20 minutes at 4°C
when cell viability was 80–85% and GFP fluorescence indicated >50% infection rate. The cell pellets were
flash-frozen in liquid N$_2$ and stored at -80°C in lysis buffer containing 50 mM Bis-Tris HCl (pH 7.5), 500
mM NaCl, protease inhibitor cocktail tablets (cOmplete EDTA-free; Roche), and DNase I (Roche). For
protein purification, cells were thawed and lysed by sonication (Branson Sonifier DS450). The lysate was
clarified by ultracentrifugation at 55,000 g for 60 min at 4°C and filtered through 0.45 µm (Durapore Merck)
filters. The supernatant was loaded onto a Ni^{2+} affinity column (HisTrap HP 5 mL, Cytiva), washed with 50
mM Bis-Tris HCl (pH 7.5), 500 mM NaCl, 20 mM imidazole, 0.5 mM EDTA, and eluted with wash buffer
supplemented with 300 mM imidazole (pH 8.0) directly onto a Strep-Tactin XT 4Flow 5 mL column (IBA).
The final elution from the Strep-Tactin column was performed using BXT buffer (IBA) supplemented with
NaCl to reach a final concentration of 500 mM. The eluted fractions were incubated with TEV protease for
16 h at 4°C to remove the 10xHis and twin StrepII affinity tags. The cleaved protein was separated from the
tags by a second Ni^{2+} affinity chromatography step. Further purification was achieved by size-exclusion
chromatrography on a HiLoad Superdex 200 16/600 pg column (Cytiva) equilibrated in 20 mM HEPES (pH
7.5) and 150 mM NaCl. The peak fractions were concentrated to 5-15 mg/mL and immediately used for
crystallization experiments or flash-frozen in liquid nitrogen and stored at -80°C for enzymatic assays.
The expression and purification protocol for full-length SpAsp1 was derived from the work of
Benjamin et al. (Benjamin *et al*, 2022a). Protein expression was done in *E. coli* BL21 (DE3) RIL cells
grown in Terrific Broth containing 50 µg/mL kanamycin, 34 µg/mL chloramphenicol, and 1% (v/v) ethanol.
When the culture reached an OD$_{600nm}$ of 0.8, protein expression was induced with 0.5 mM isopropyl β-D-1-
thiogalactopyranoside (IPTG) at 18°C for 16 h. Cells were harvested by centrifugation and resuspended in a
buffer containing 50 mM Tris-HCl (pH 8.0), 500 mM NaCl, and 10% glycerol, supplemented with a

cOmplete protease inhibitor cocktail tablet, DNase I, and 0.5 mg/mL lysozyme (Roth). Cell lysis was

achieved by sonication. The lysate was then clarified by centrifugation at 46,500 g for 1 h, and the

supernatant was loaded onto a Ni^{2+} affinity column (HisTrap HP 5 mL, Cytiva) equilibrated in 50 mM Tris-

HCl (pH 8.0), 500 mM NaCl, and 10% (v/v) glycerol. After a washing step with the same buffer, SpAsp1

was eluted with buffer supplemented with 250 mM imidazole (pH 8.0). The 7xHis-StrepII-SUMO tag was

cleaved by treatment with Ulp1 protease during an overnight incubation at 4°C (Malakhov *et al*, 2004). The

tag-free SpAsp1 protein was separated from the fusion tag by a second Ni^{2+} affinity chromatography step.

SpAsp1 was then diluted into 50 mM Tris-HCl (pH 8.0) and 10% (v/v) glycerol to reach a final NaCl

concentration of 50 mM and loaded onto a HiTrap Heparin HP 5 mL column (Cytiva) equilibrated with 50

mM Tris-HCl (pH 8.0), 50 mM NaCl, and 10% (v/v) glycerol. SpAsp1 was eluted against a gradient of 50

mM Tris-HCl (pH 8.0), 1 M NaCl, and 10% (v/v) glycerol. Finally, SpAsp1 was applied to a HiLoad

Superdex 200 16/600 pg column (Cytiva) equilibrated in 20 mM HEPES (pH 7.5) and 150 mM NaCl. The

peak fractions containing SpAsp1 were concentrated and immediately used for negative stain electron

microscopy.

Crystallization and data collection

ScVip1[KD] (8.0 mg/mL in 20 mM Hepes pH 7.5, 150 mM NaCl) was incubated with 5 mM ADP and

20 mM $MgCl_2$ for 1 h on ice prior to crystallization. Crystals developed in hanging drops composed of 1.0

µL of protein solution and 1.0 µL of crystallization buffer (23 % [v/v] PEG 3,350, 0.1 M citric acid / BIS-

Tris propane pH 6.4) suspended over 1.0 mL of the latter as reservoir solution. A complete dataset to 1.18 Å

resolution was collected at beam line X06DA of the Swiss Light Source, Villigen, Switzerland (Table 1).

Triclinic crystals of apo ScVip1[PD] developed in hanging drops composed of 1.5 µL of protein solution (15

mg/mL in 20 mM Hepes pH 7.4, 150 mM NaCl) and 1.5 µL of crystallization buffer (20 % [v/v] PEG 3,350,

0.2 M NH_4NO_3). Crystals were derivatized for 18 h in crystallization buffer supplemented with 0.1 mM

K_2PtCl_4 and subsequently snap-frozen in liquid N_2 after serial transfer in crystallization buffer supplemented

with 20 % (v/v) glycerol as cryoprotectant. A single anomalous diffraction (SAD) dataset at the 'white line'

L3 edge was collected at beam line X06DA (Table 1). Native crystals of ScVip1[PD] developed in 20% v/v

(+/-)-2-Methyl-2,4-pentanediol, 0.1 M Tris pH 8.0. Crystals were cryoprotected by addition of glycerol to a

final concentration of 20% (v/v) and a complete dataset with apparent orthorhombic symmetry was collected

at beam line X06DA to 3.05 Å resolution (Table 1). Crystals of the engineered ScVip1[PDΔ848-918] construct (10

mg/mL in 20 mM Hepes pH7.5, 150 mM NaCl, 0.5 mM $(CH_3CO_2)_2Zn$) developed in 2 µL drops containing

10 % (w/v) polyvinylpyrrolidone K15, 5 mM $CoCl_2$, 0.1M Tris pH 7.0 as crystallization buffer and 0.1 µL

additive solution (5mM 13:0 Lyso PC aka 1-tridecanoyl-2-hydroxy-sn-glycero-3-phosphocholine) using

microseeding protocols. Crystals were cryoprotected by serial transfer into crystallization buffer containing

glyerol to a final concentration of 30% (v/v) and snap frozen in liquid N_2. Data collection at beam line

X06DA yielded a complete dataset at 3.4 Å resolution (Table 1). The ScVip1[PD Δ848-918 RHR-AAA] – 1,5-InsP[8]

complex crystallized in sitting drops containing 0.25 µL protein solution (12 mg/mL in 20 mM Hepes 7.5,

150 mM NaCl, 2.5 mM 1,5-InsP$_8$) and 0.25 μL of crystallization buffer (Morpheus II screen condition H2,

32.5% v/v Precipitant Mix 6: 25% w/v PEG 4000, 40% w/v 1,2,6-Hexanetriol; 0.04M Polyamines: 0.01M

Spermine   tetrahydrochloride,   0.01M   Spermidine   trihydrochloride,   0.01M   1,4-Diaminobutane

dihydrochloride, 0.01M DL-Ornithine monohydrochloride; 0.1M Buffer System 4: pH6.5 MOPSO, Bis-Tris)

(Gorrec, 2009). Crystals were cryoprotected by serial transfer in crystallization buffer supplemented with

ethylene glycol to a final concentration of 25 % (v/v) and diffracted up to 2.36 Å resolution (Table 1). Data

processing was done with XDS (version June 30, 2023) (Kabsch, 1993).

Crystallographic structure solution and refinement

The structure of ScVip1^{KD} was solved by molecular replacement method using the program PHASER

(McCoy *et al*, 2007) and the isolated kinase domain of HsPPIP5K2 as search model (https://www.rcsb.org/

PDB-ID 3t9a). The solution comprises a monomer in the asymmetric unit. The structure was completed in

alternating cycles of manual model building in COOT (Emsley & Cowtan, 2004) and restrained refinement

with anisotropic atomic displacement parameters as implemented in PHENIX.REFINE (Afonine *et al*, 2012).

The structure of ScVip1^{PD} was solved by molecular replacement combined with single anomalous diffraction

(MR-SAD) using a low redundancy SAD dataset collected from a K$_2$PtCl$_2$-derivatized crystal and a fragment

of the *Francisella tularensis* histidine acid phosphatase core domain (PDB-ID 3it3) in the program

PHASER-EP (Read & McCoy, 2011). The resulting solution comprised four molecules in the asymmetric

unit, four Pt^{2+} and four Zn^{2+} ions (Table 1). The starting phases and model were used for non-crystallographic

symmetry (NCS) density modification in PHENIX.RESOLVE (Terwilliger, 2003) and the resulting 3.5 Å

electron density map was sufficient to build a nearly complete poly-alanine model that was used to determine

the structure of ScVip1^{PD} apo by molecular replacement. ScVip1^{PD} apo crystals had an apparent *C* 2 2 2

symmetry, however no packing solution could be identified with PHASER (McCoy *et al*, 2007) and analysis

with PHENIX.XTRIAGE suggested an abnormal distribution of intensities. The structure was subsequently

solved in space group *P* 1 and input into the program ZANUDA (Lebedev & Isupov, 2014) for space group

validation. ZANUDA returned a solution in space group *P* 2$_1$, with four molecules in the asymmetric unit

(Table 1). The a and c axes are almost identical, with a twin fraction of ~0.5, similar to previous reports (Ban

et al, 1999; Wittmann & Rudolph, 2007). The structure was completed in alternating cycles of manual model

building and restrained NCS refinement with twin operator l, -k, h in PHENIX.REFINE (Afonine *et al*,

2012). A large, apparently disordered loop (residues 851-918) was identified in this structure and

subsequently replaced by a short Gly-Ser-Ser-Gly linker. The structures of ScVip1$^{PDΔ848-918}$ and ScVip1$^{PD\ Δ848-}$

$^{918\ RHR-AAA}$ were solved by molecular replacement using the ScVip1 apo structure as search model and refined

with PHENIX.REFINE (Afonine *et al*, 2012) (Table 1). The stereochemistry of the refined models was

assessed with PHENIX.MOLPROBITY (Davis *et al*, 2007). Structural diagrams were generated with

CHIMERA (Pettersen *et al*, 2004).

NMR-based phosphatase assay

ScVip1 phosphatase assays were performed as previously described (Zhu *et al*, 2019). Reactions contained 100 µM of the respective [$^{13}C_6$]-labeled PP-InsP in 20 mM HEPES pH 7.0, 150 mM NaCl, 0.2 mg/mL BSA and D_2O to a total volume of 600 µL. Reactions were supplemented with sodium phosphate, sodium sulfate or sodium nitrate, as indicated. Reaction mixtures were pre-incubated at 37°C and the reaction was started by adding the respective amount of enzyme (ScVip1^{PD} (WT): 10 nM for 1-InsP$_7$ and 1,5-InsP$_8$, 700 nM for 5-InsP$_7$; 1.5 µM R547A-H548A-R551A; 1.25 µM R547A; 1.5 µM H548A; 1.5 µM R551A; 10 nM Δ848-918; 1.5 µM K554A; E991A: 2 µM for InsP$_6$, 20 nM for 1-InsP$_7$, 10 nM for 1,5-InsP$_8$; 1 µM P553V-G646V-G647A; 10 nM C793A; 10 nM H651A; 350 nM K558A-K605A-K732A-K817A; 1 µM K554A-K556A-K644A; 10 nM K817S-D819A-S821A; 50 nM E576A-K623A-Q625S-K627A). Turnover of PP-InsPs was monitored continuously at 37°C using a nuclear magnetic resonance spectroscopy (NMR) pseudo-2D spin-echo difference experiment (Harmel *et al*, 2019). Individual NMR spectra were recorded every 86 s. The relative intensity changes of the C2 peaks of the respective PP-InsPs as a function of reaction time were used for quantification (Harmel *et al*, 2019; Zhu *et al*, 2019). Data analysis was carried out with GraphPad PRISM 5. Trend lines were added to the raw data, following either a linear regression model or the first derivative of the equation of a one phase decay model, normalized by the enzyme's mass concentration.

In order to determine the substrate preference of ScVip1^{PD}, reaction mixtures containing 20 mM HEPES pH 7.0, 150 mM NaCl, 0.2 mg/mL BSA and 100 µM of the respective [$^{13}C_6$]-labeled (PP-)InsP were prepared and pre-incubated at 37°C. ScVip1^{PD} was added subsequently (100 nM for 1-InsP$_7$ and 1,5-InsP$_8$, 1 µM for InsP$_6$ and 5-InsP$_7$) and the reactions were incubated at 37°C for 3 h. As a control, H_2O was added instead of the enzyme. Reactions were quenched by heating to 95°C for 5 min. The samples were centrifuged at 15,000 g for 5 min and the supernatant was analyzed by a ^{1}H-^{13}C-BIRD-HMQC 2D NMR as described (Harmel *et al*, 2019).

Malachite green-based phosphatase assay

The phosphatase activity of ScVip1^{PD} was determined using the Malachite Green Phosphate Assay Kit (Sigma-Aldrich). This assay detects the release of inorganic phosphate catalyzed by ScVip1PD from substrates such as 1,5-InsP$_8$, 1-InsP$_7$, 5-InsP$_7$ or InsP$_6$ during the phosphatase reaction (Ames, 1966; Baykov *et al*, 1988). For the assay, 100 nM of ScVip1^{PD} was incubated at 37°C for 2,4, and 8 min in 100 µL reaction mixtures containing 20 mM Hepes (pH 7.5), 150 mM NaCl and varying concentrations of substrates (1,5-InsP$_8$ ranging from 8 µM to 700 µM; 1-InsP$_7$ ranging from 1 µM to 2000 µM, unless indicated otherwise). The reaction was quenched by mixing 40 µL of the reaction mixture with 10 µL of malachite green reagent. The color formation was measured after 25 min on a plate reader (Tecan Sparc) at 620 nm. When precipitation of the reagent occurred at high phosphate concentrations (>100 µM), the reaction mixture was 2-8 times diluted in buffer prior quenching. The quantity of released inorganic phosphatase was estimated using the standard curve of OD$^{620\ nm}$ versus phosphate standard concentrations. Blanks OD$_{620\ nm}$ were

measured without enzyme at each time point to discard effect of non-enzymatic hydrolysis of PP-InsPs or the

eventual presence of inorganic phosphate in the buffer.

Thermal shift assay

Thermal shift assays were conducted with 25 to 100 µM of wild-type or mutant ScVip1^{PD} in 20 mM

Hepes (pH 7.5), 150 mM NaCl, and 8fold dilution of SYPRO Orange dye (Thermo Fisher Scientific).

Protein samples were heated with an increasing gradient of 0.05°C/s from 25 to 99°C and melting curves

were recorded using a QuantStudio 5 Real-Time PCR System system (Thermo Fisher Scientific).

Analytical size-exclusion chromatography

Gel filtration experiments were performed using a Superdex 200 Increase 10/300 GL column (GE

Healthcare) pre-equilibrated in 20 mM Hepes pH 7.5, 150 mM NaCl. 200 µl of the respective protein was

loaded sequentially onto the column, and elution at 0.75 ml/min was monitored by ultraviolet absorbance at

280 nm. Peak fractions were analyzed by SDS–PAGE gel electrophoresis.

Molecular cloning and generation of stable transgenic A. thaliana lines

The Golden Gate system was used to generate plasmids for Arabidopsis transformation (Binder *et al*,

2014). Mutant versions of the AtVIH2 coding sequence were synthesized (Twist bio-science, San

Francisco,CA) (Table EV1). The different mutants were then cloned into the level one (L1) vector (Table

EV1). To construct the binary vector L1 vectors containing the AtUBI10 promoter (proUBI10), the

corresponding AtVIH2 coding sequence , a FLAG tag, the Nos terminator and Fast Red as a fluorescent

marker for the seed coat (Table EV1). The plasmids were transformed into *A. tumefaciens* strain GV3101.

Flower dipping was then used to transform 5-week-old plants (Col-0) (Clough & Bent, 1998). T1 plants were

selected by red fluorescence using a Nikon SMZ18 stereomicroscope equipped with an RFP-B filter. Lines

with single T-DNA insertions were selected by the segregation analysis. All experiments were performed

using stable T3 generation lines (Table EV2).

Plant material, seed sterilization, and plant growth conditions

Seeds were sterilized using the chloride gas protocol. Seeds were then sown on ½MS media

containing 1% (w/v) sucrose, MES (0.5 g/L), pH 5.7 and agar (8g/L). 120 mm square Petri dishes were

incubated for 48h at 4°C in the dark and transferred to a growth chamber at 22 °C with a 16h/8h light/dark

cycle under a fluorescent light source. 7d old seedlings were transferred to soil in 8 cm pots and grown as

indicated.

Image acquisition and rosette area quantification

To evaluate the rosette phenotypes of the different T3 lines, 16 plants (in 4 pots) were analyzed. After

2 weeks in soil, top images were recorded and the rosette area was manually segmented in Fiji (Schindelin *et*

al, 2012). Specifically, images were converted to 8 bit followed by manual selection of individual plants.

Contrast was adjusted, and the rosette was selected using the threshold. The scale (240 px to 1.5 cm) was set

before quantification. Finally, the area of the segmented image was determined using the Particle Quantify

function as implemented in Fiji.

Determination of cellular Pi concentration

For Pi quantification 3-weeks-old plants were analyzed (8 independent plants per sample). One

cotyledonous leaf per plant was harvested, the fresh weight was recorded and the leaf was transferred to an

Eppendorf tube (1.5 ml) containing 500µl miliQ H_2O. The tubes were then frozen in liquid nitrogen and

boiled at 80 ºC for 5 min. This step was repeated two times. Measurement was performed using a

colorimetric molybdate assay (Ames, 1966). The master solution contains contained 72 µL of ammonium

molybdate solution (0.0044 % [w/v] of ammonium molybdate tetrahydrate, 0.23 % [v/v] of 18 M H_2SO_4), 16

µL of 10 % (w/v) acetic acid and 12 µL of miliQ H_2O. For each reaction, 20 µl of sample were added in

triplicate to a 96-well plate and incubated with 100 µl of working solution for 1 h at 37 ºC. Absorbance was

measured at 820 nm in a Spark plate reader (Tecan). The concentration was determined according to the

standard curve with a concentration range of 1, 0.5, 0.2, 0.1 and 0 mM Na_2HPO_4.

Western blotting

Proteins were extracted from 3-weeks-old plants. About 100 mg of sample were collected in 2 ml

Eppendorf tubes and frozen in liquid nitrogen. The tissue was homogenized in a tissue lyzer (MM400,

Retsch). Then, 300 µl of 1x PBS buffer supplemented with the plant-specific protease inhibitor cocktail

(PE0230, Merck) was added to each tube and incubated for 30 min at 4 ºC in an orbital shaker (Intelli-

Mixer™ RM-2M, ELMI). The samples were then centrifuged at 10,000 g for 20 min at 4 ºC. Then, 150 µl of

the supernatant was transferred in a new tube containing 30 µl of SDS loading buffer and incubated at 95 ºC

for 5 min. A 10% BIS-Tris acrylamide SDS-PAGE was run with the MES running buffer. Proteins were

transferred to a 0.2 µ PVDF membrane (IB34002, Thermo Scientific™) using a semi-dry protocol (iBlot3,

Thermo Scientific). Membranes were blotted with TBS-Tween (0.1%)-milk (5%) for 1 h at room

temperature. For FLAG detection, membranes were incubated with a monoclonal Anti-FLAG-M2-

Peroxidase (HRP) antibody (A8592, Merck) using a dilution of 1:5000. Detection was performed using

SuperSignal West Femto Maximum Sensitivity Substrate (34095, Thermo Scientific).

 Electron microscopy and image processing

The full-length SpAsp1 envelope was analyzed using negative-stain electron microscopy. A total of 5

µl of monomeric peak fractions from a gel filtration run (concentration 0.01 mg per mL) were applied to

glow-discharged, carbon-coated copper grids (400 mesh, Electron Microscopy Sciences). After incubation

for 1 min, excess liquid was blotted off. The grids were then sequentially passed through one drop of buffer

(20 mM Hepes, pH 7.5, and 150 mM NaCl) and two drops of 2% (w/v) uranyl acetate solution, with the

sample incubated in the last drop for 1 min before blotting and air drying. A total of 1,565 micrographs were

recorded at the DCI-Geneva (cryoGEnic) electron microscopy platform using a Talos L120C microscope

operating at 120 kV and equipped with a Falcon II direct electron detector. Data acquisition was automated

using EPU software (Thermo Fisher Scientific) with a pixel size of 1.531 Å and a total electron dose of 52

e-/Å². Data processing was performed using CryoSPARC (version 4.1) (Punjani *et al*, 2017). Initially,

contrast transfer function (CTF) parameters were estimated using patch-based CTF estimation. Micrographs

with a CTF fit better than 10 Å resolution were selected, resulting in 1,497 micrographs. Manual picking of

177 SpAsp1 particles, which were extracted in boxes of 250 × 250 pixels, were used to train the blob picker

tuner. After this process, 239 exposures were rejected due to containing either too few or excessively high

numbers of picked particles, leaving a particle stack of 270,221 particles (averaging 220 particles per

micrograph) with defocus values between 0.3 and 1.3 µm. Two rounds of reference-free 2D class averaging

were employed to further refine the dataset and remove the stain artifacts, resulting in 107,485 particles. An

ab initio reconstruction with six classes was followed by 2D classification and non-uniform refinement for

each class. The best 3D reconstruction, representing full-length SpAsp1, comprised 11,837 particles.

Hydrogen/deuterium exchange mass spectrometry

HDX-MS experiments were performed at the UniGe Protein Platform (University of Geneva,

Switzerland) following an established protocol with minimal modifications (Wang *et al*, 2018b). Details of

reaction conditions and all data are presented in Fig EV10. HDX reactions were performed in 50 µl volumes

with a final protein concentration of 2.1 µM of ScVip1 and a 100-fold molar excess of AMP-PNP and / or

PCP-IP$_8$ (Hostachy *et al*, 2021). Briefly, 107 picomoles of protein were pre-incubated with ligands for 1 h on

ice in a final volume of 7 µl. The deuterium on-exchange reaction was initiated by adding 43 µl of D$_2$O

exchange buffer (10 mM Tris pH 8 / 150 mM NaCl in D$_2$O) to the protein-ligand mixture. Reactions were

carried-out at room temperature for 2 incubation times (30 s, 300 s) and terminated by the sequential addition

of 20 µl of ice-cold quench buffer (4 M Gdn-HCl / 1 M NaCl / 0.1 M NaH$_2$PO$_4$ pH 2.5 / 1 % (v/v) formic

acid). Samples were immediately frozen in liquid nitrogen and stored at -80 °C for up to two weeks. All

experiments were repeated in triplicate. To quantify deuterium uptake into the protein, samples were thawed

and injected in a UPLC system immersed in ice with 0.1 % FA as liquid phase. The protein was digested via

two immobilized pepsin columns (Thermo Scientific #23131), and peptides were collected onto a VanGuard

precolumn trap (Waters). The trap was then eluted, and peptides were separated with a C18, 300Å, 1.7 µm

particle size Fortis Bio 100 x 2.1 mm column over a gradient of 8 – 30 % buffer C over 20 min at 150 µl/

min (buffer B: 0.1 % [v/v] formic acid; buffer C: 100 % acetonitrile). Mass spectra were acquired on an
Orbitrap Velos Pro (Thermo), for ions from 400 to 2200 m/z using an electrospray ionization source operated
at 300 °C, 5 kV of ion spray voltage. Peptides were identified by data-dependent acquisition of a non-
deuterated sample after MS/ MS and data were analyzed by Mascot. All peptides analysed are shown in Fig
EV10. Deuterium incorporation levels were quantified using HD examiner software version 3.3 (Sierra
Analytics), and quality of every peptide was checked manually. Results are presented as percentage of
maximal deuteration compared to theoretical maximal deuteration level. Changes in deuteration level
between two states were considered significant if >7% and >0.5 Da and $p < 0.02$ (unpaired t-test) for a single
deuteration time (Dataset EV1).

Data availability

Crystallographic coordinates and associated structure factors have been deposited with the Protein
Data Bank (http://rcsb.org) doi:10.2210/pdb9gr8/pdb (ScVip1^{KD} – ADP, PDB-ID 8gr8),
doi:10.2210/pdb9grh/pdb (ScVip1^{PD} – apo, PDB-ID 9grh), doi:10.2210/pdb9grn/pdb (ScVip1$^{PDΔ848-918}$ – apo,
PDB-ID 9grn) and doi:10.2210/pdb9gro/pdb (ScVip1$^{PD \ Δ848-918 \ RHR-AAA}$ – 1,5-InsP$_8$, PDB-ID 9gro).

Acknowlegements

We thank P. Rieu and A. Cargenato for critical reading of the manuscript, Oscar Vadas from the
Protein Purification and Characterisation Platform at the Faculty of Medicine, University of Geneva for
performing the different HDX-MS experiments. This work was supported by Sinergia grant CRSII5_209412
from the Swiss National Science Foundation (to D.F. and M.H.) and by Howard Hughes Medical Institute
International Research Scholar Award 55008733 (to M.H.).

Disclosure and competing interests

The authors declare no competing interests.

References

Acquistapace IM, Thompson EJ, Kühn I, Bedford MR, Brearley CA & Hemmings AM (2022) Insights to the
    Structural Basis for the Stereospecificity of the Escherichia coli Phytase, AppA. *IJMS* 23: 6346

Afonine PV, Grosse-Kunstleve RW, Echols N, Headd JJ, Moriarty NW, Mustyakimov M, Terwilliger TC,
    Urzhumtsev A, Zwart PH & Adams PD (2012) Towards automated crystallographic structure
    refinement with phenix.refine. *Acta Cryst D* 68: 352–367

Ames B (1966) Assay of inorganic phosphate, total phosphate and phosphatases. *Methods Enzymol* 8: 115–
    118

An Y, Jessen HJ, Wang H, Shears SB & Kireev D (2019) Dynamics of Substrate Processing by PPIP5K2, a
    Versatile Catalytic Machine. *Structure* 27: 1022-1028.e2

Aravind L & Ponting CP (1997) The GAF domain: an evolutionary link between diverse phototransducing
    proteins. *Trends Biochem Sci* 22: 458–459

Auesukaree C, Homma T, Tochio H, Shirakawa M, Kaneko Y & Harashima S (2004) Intracellular phosphate serves as a signal for the regulation of the PHO pathway in Saccharomyces cerevisiae. *J Biol Chem* 279: 17289–17294

Ban N, Nissen P, Hansen J, Capel M, Moore PB & Steitz TA (1999) Placement of protein and RNA structures into a 5 A-resolution map of the 50S ribosomal subunit. *Nature* 400: 841–847

Baykov AA, Evtushenko OA & Avaeva SM (1988) A malachite green procedure for orthophosphate determination and its use in alkaline phosphatase-based enzyme immunoassay. *Anal Biochem* 171: 266–270

Benjamin B, Garg A, Jork N, Jessen HJ, Schwer B & Shuman S (2022a) Activities and Structure-Function Analysis of Fission Yeast Inositol Pyrophosphate (IPP) Kinase-Pyrophosphatase Asp1 and Its Impact on Regulation of *pho1* Gene Expression. *mBio* 13: e01034-22

Benjamin B, Goldgur Y, Jork N, Jessen HJ, Schwer B & Shuman S (2022b) Structures of Fission Yeast Inositol Pyrophosphate Kinase Asp1 in Ligand-Free, Substrate-Bound, and Product-Bound States. *mBio* 13: e03087-22

Binder A, Lambert J, Morbitzer R, Popp C, Ott T, Lahaye T & Parniske M (2014) A modular plasmid assembly kit for multigene expression, gene silencing and silencing rescue in plants. *PLoS One* 9: e88218

Chabert V, Kim G-D, Qiu D, Liu G, Michaillat Mayer L, Jamsheer K M, Jessen HJ & Mayer A (2023) Inositol pyrophosphate dynamics reveals control of the yeast phosphate starvation program through 1,5-IP8 and the SPX domain of Pho81. *Elife* 12: RP87956

Clough SJ & Bent AF (1998) Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. *The Plant Journal* 16: 735–743

Cordeiro CD, Saiardi A & Docampo R (2017) The inositol pyrophosphate synthesis pathway in Trypanosoma brucei is linked to polyphosphate synthesis in acidocalcisomes. *Mol Microbiol* 106: 319–333

Couso I, Evans BS, Li J, Liu Y, Ma F, Diamond S, Allen DK & Umen JG (2016) Synergism between Inositol Polyphosphates and TOR Kinase Signaling in Nutrient Sensing, Growth Control, and Lipid Metabolism in Chlamydomonas. *Plant Cell* 28: 2026–2042

Davis IW, Leaver-Fay A, Chen VB, Block JN, Kapral GJ, Wang X, Murray LW, Arendall WB, Snoeyink J, Richardson JS, *et al* (2007) MolProbity: all-atom contacts and structure validation for proteins and nucleic acids. *Nucleic Acids Res* 35: W375-383

Desai M, Rangarajan P, Donahue JL, Williams SP, Land ES, Mandal MK, Phillippy BQ, Perera IY, Raboy V & Gillaspy GE (2014) Two inositol hexakisphosphate kinases drive inositol pyrophosphate synthesis in plants. *Plant J* 80: 642–653

Desfougères Y, Portela-Torres P, Qiu D, Livermore TM, Harmel RK, Borghi F, Jessen HJ, Fiedler D & Saiardi A (2022) The inositol pyrophosphate metabolism of Dictyostelium discoideum does not regulate inorganic polyphosphate (polyP) synthesis. *Advances in Biological Regulation* 83: 100835

Dollins DE, Bai W, Fridy PC, Otto JC, Neubauer JL, Gattis SG, Mehta KPM & York JD (2020) Vip1 is a kinase and pyrophosphatase switch that regulates inositol diphosphate signaling. *Proceedings of the National Academy of Sciences* 117: 9356–9364

Dong J, Ma G, Sui L, Wei M, Satheesh V, Zhang R, Ge S, Li J, Zhang T-E, Wittwer C, *et al* (2019) Inositol Pyrophosphate InsP8 Acts as an Intracellular Phosphate Signal in Arabidopsis. *Molecular Plant*

Dubois E, Scherens B, Vierendeels F, Ho MMW, Messenguy F & Shears SB (2002) In Saccharomyces cerevisiae, the Inositol Polyphosphate Kinase Activity of Kcs1p Is Required for Resistance to Salt Stress, Cell Wall Integrity, and Vacuolar Morphogenesis. *Journal of Biological Chemistry* 277: 23755–23763

Dunnett CW (1955) A Multiple Comparison Procedure for Comparing Several Treatments with a Control. *Journal of the American Statistical Association* 50: 1096–1121

Emsley P & Cowtan K (2004) Coot: model-building tools for molecular graphics. *Acta Cryst D* 60: 2126–2132

Fridy PC, Otto JC, Dollins DE & York JD (2007) Cloning and characterization of two human VIP1-like inositol hexakisphosphate and diphosphoinositol pentakisphosphate kinases. *J Biol Chem* 282: 30754–30762

Garg A, Shuman S & Schwer B (2020) A genetic screen for suppressors of hyper-repression of the fission yeast PHO regulon by Pol2 CTD mutation T4A implicates inositol 1-pyrophosphates as agonists of precocious lncRNA transcription termination. *Nucleic Acids Res* 48: 10739–10752

Gokhale NA, Zaremba A & Shears SB (2011) Receptor-dependent compartmentalization of PPIP5K1, a kinase with a cryptic polyphosphoinositide binding domain. *Biochemical Journal* 434: 415–426

Gorrec F (2009) The MORPHEUS protein crystallization screen. *J Appl Crystallogr* 42: 1035–1042

Gu C, Li X, Zong G, Wang H & Shears SB (2024) IP8: A quantitatively minor inositol pyrophosphate signaling molecule that punches above its weight. *Advances in Biological Regulation* 91: 101002

Gu C, Liu J, Liu X, Zhang H, Luo J, Wang H, Locasale JW & Shears SB (2021) Metabolic supervision by PPIP5K, an inositol pyrophosphate kinase/phosphatase, controls proliferation of the HCT116 tumor cell line. *Proc Natl Acad Sci USA* 118: e2020187118

Gu C, Nguyen H-N, Ganini D, Chen Z, Jessen HJ, Gu Z, Wang H & Shears SB (2017a) KO of 5-InsP7 kinase activity transforms the HCT116 colon cancer cell line into a hypermetabolic, growth-inhibited phenotype. *Proceedings of the National Academy of Sciences* 114: 11968–11973

Gu C, Nguyen H-N, Hofer A, Jessen HJ, Dai X, Wang H & Shears SB (2017b) The Significance of the Bifunctional Kinase/Phosphatase Activities of Diphosphoinositol Pentakisphosphate Kinases (PPIP5Ks) for Coupling Inositol Pyrophosphate Cell Signaling to Cellular Phosphate Homeostasis. *J Biol Chem* 292: 4544–4555

Gu C, Wilson MSC, Jessen HJ, Saiardi A & Shears SB (2016) Inositol Pyrophosphate Profiling of Two HCT116 Cell Lines Uncovers Variation in InsP8 Levels. *PLOS ONE* 11: e0165286

Guan Z, Chen J, Liu R, Chen Y, Xing Q, Du Z, Cheng M, Hu J, Zhang W, Mei W, *et al* (2023) The cytoplasmic synthesis and coupled membrane translocation of eukaryotic polyphosphate by signal-activated VTC complex. *Nat Commun* 14: 718

Guan Z, Zhang Q, Zhang Z, Zuo J, Chen J, Liu R, Savarin J, Broger L, Cheng P, Wang Q, *et al* (2022) Mechanistic insights into the regulation of plant phosphate homeostasis by the rice SPX2 - PHR2 complex. *Nat Commun* 13: 1581

Gulabani H, Goswami K, Walia Y, Roy A, Noor JJ, Ingole KD, Kasera M, Laha D, Giehl RFH, Schaaf G, *et al* (2022) Arabidopsis inositol polyphosphate kinases IPK1 and ITPK1 modulate crosstalk between SA-dependent immunity and phosphate-starvation responses. *Plant Cell Rep* 41: 347–363

Hamada K, Kato M, Shimizu T, Ihara K, Mizuno T & Hakoshima T (2005) Crystal structure of the protein histidine phosphatase SixA in the multistep His-Asp phosphorelay. *Genes to Cells* 10: 1–11

Harmel RK, Puschmann R, Nguyen Trung M, Saiardi A, Schmieder P & Fiedler D (2019) Harnessing 13C-labeled myo-inositol to interrogate inositol phosphate messengers by NMR. *Chem Sci* 10: 5267–5274

Hashimoto Y, Zhang S & Blissard GW (2010) Ao38, a new cell line from eggs of the black witch moth, Ascalapha odorata (Lepidoptera: Noctuidae), is permissive for AcMNPV infection and produces high levels of recombinant proteins. *BMC Biotechnol* 10: 50

Henry JT & Crosson S (2011) Ligand-Binding PAS Domains in a Genomic, Cellular, and Structural Context. *Annual Review of Microbiology* 65: 261–286

Ho YJ, Burden LM & Hurley JH (2000) Structure of the GAF domain, a ubiquitous signaling motif and a new class of cyclic GMP receptor. *The EMBO Journal* 19: 5288–5299

Holm L & Rosenström P (2010) Dali server: conservation mapping in 3D. *Nucl Acids Res* 38: W545–W549

Hostachy S, Utesch T, Franke K, Dornan GL, Furkert D, Türkaydin B, Haucke V, Sun H & Fiedler D (2021) Dissecting the activation of insulin degrading enzyme by inositol pyrophosphates and their bisphosphonate analogs. *Chem Sci* 12: 10696–10702

Hothorn M, Dabi T & Chory J (2011) Structural basis for cytokinin recognition by Arabidopsis thaliana histidine kinase 4. *Nat Chem Biol* 7: 766–768

Kabsch W (1993) Automatic processing of rotation diffraction data from crystals of initially unknown symmetry land cell constants. *Journal of Applied Crystallography* 26: 795–800

Laha D, Johnen P, Azevedo C, Dynowski M, Weiß M, Capolicchio S, Mao H, Iven T, Steenbergen M, Freyer M, *et al* (2015) VIH2 Regulates the Synthesis of Inositol Pyrophosphate InsP8 and Jasmonate-Dependent Defenses in Arabidopsis. *Plant Cell* 27: 1082–1097

Laha D, Parvin N, Hofer A, Giehl RFH, Fernandez-Rebollo N, von Wirén N, Saiardi A, Jessen HJ & Schaaf G (2019) Arabidopsis ITPK1 and ITPK2 Have an Evolutionarily Conserved Phytic Acid Kinase Activity. *ACS Chem Biol* 14: 2127–2133

Landau M, Mayrose I, Rosenberg Y, Glaser F, Martz E, Pupko T & Ben-Tal N (2005) ConSurf 2005: the projection of evolutionary conservation scores of residues on protein structures. *Nucleic Acids Res* 33: W299-302

Laurent F, Bartsch SM, Shukla A, Rico-Resendiz F, Couto D, Fuchs C, Nicolet J, Loubéry S, Jessen HJ, Fiedler D, *et al* (2024) Inositol pyrophosphate catabolism by three families of phosphatases controls plant growth and development. 2024.04.30.591831 doi:10.1101/2024.04.30.591831 [PREPRINT]

Lebedev AA & Isupov MN (2014) Space-group and origin ambiguity in macromolecular structures with pseudo-symmetry and its treatment with the program Zanuda. *Acta Cryst D* 70: 2430–2443

Lee Y-H, Li Y, Uyeda K & Hasemann CA (2003) Tissue-specific Structure/Function Differentiation of the Liver Isoform of 6-Phosphofructo-2-kinase/Fructose-2,6-bisphosphatase. *Journal of Biological Chemistry* 278: 523–530

Li X, Gu C, Hostachy S, Sahu S, Wittwer C, Jessen HJ, Fiedler D, Wang H & Shears SB (2020) Control of XPR1-dependent cellular phosphate efflux by InsP8 is an exemplar for functionally-exclusive inositol pyrophosphate signaling. *PNAS* 117: 3568–3574

Li X, Kirkpatrick RB, Wang X, Tucker CJ, Shukla A, Jessen HJ, Wang H, Shears SB & Gu C (2024) Homeostatic coordination of cellular phosphate uptake and efflux requires an organelle-based receptor for the inositol pyrophosphate IP8. *Cell Rep* 43: 114316

Lim D, Golovan S, Forsberg CW & Jia Z (2000) Crystal structures of Escherichia coli phytase and its complex with phytate. *Nat Struct Biol* 7: 108–113

Machkalyan G, Trieu P, Pétrin D, Hébert TE & Miller GJ (2016) PPIP5K1 interacts with the exocyst complex through a C-terminal intrinsically disordered domain and regulates cell motility. *Cellular Signalling* 28: 401–411

Malakhov MP, Mattern MR, Malakhova OA, Drinker M, Weeks SD & Butt TR (2004) SUMO fusions and SUMO-specific protease for efficient expression and purification of proteins. *J Struct Funct Genomics* 5: 75–86

McCoy AJ, Grosse-Kunstleve RW, Adams PD, Winn MD, Storoni LC & Read RJ (2007) Phaser crystallographic software. *J Appl Crystallogr* 40: 658–674

Milburn CC, Deak M, Kelly SM, Price NC, Alessi DR & Van Aalten DMF (2003) Binding of phosphatidylinositol 3,4,5-trisphosphate to the pleckstrin homology domain of protein kinase B induces a conformational change. *Biochemical Journal* 375: 531–538

Mulugu S, Bai W, Fridy PC, Bastidas RJ, Otto JC, Dollins DE, Haystead TA, Ribeiro AA & York JD (2007) A Conserved Family of Enzymes That Phosphorylate Inositol Hexakisphosphate. *Science* 316: 106–109

Nair VS, Gu C, Janoshazi AK, Jessen HJ, Wang H & Shears SB (2018) Inositol pyrophosphate synthesis by diphosphoinositol pentakisphosphate kinase-1 is regulated by phosphatidylinositol(4,5)bisphosphate. *Bioscience Reports* 38: BSR20171549

Nguyen Trung M, Furkert D & Fiedler D (2022) Versatile signaling mechanisms of inositol pyrophosphates. *Current Opinion in Chemical Biology* 70: 102177

Ortlund E, LaCount MW & Lebioda L (2003) Crystal Structures of Human Prostatic Acid Phosphatase in Complex with a Phosphate Ion and α-Benzylaminobenzylphosphonic Acid Update the Mechanistic Picture and Offer New Insights into Inhibitor Design , . *Biochemistry* 42: 383–389

Ostanin K, Harms EH, Stevis PE, Kuciel R, Zhou MM & Van Etten RL (1992) Overexpression, site-directed mutagenesis, and mechanism of Escherichia coli acid phosphatase. *J Biol Chem* 267: 22830–22836

Ostanin K & Van Etten RL (1993) Asp304 of Escherichia coli acid phosphatase is involved in leaving group protonation. *J Biol Chem* 268: 20778–20784

Pascual-Ortiz M, Saiardi A, Walla E, Jakopec V, Künzel NA, Span I, Vangala A & Fleig U (2018) Asp1 Bifunctional Activity Modulates Spindle Function via Controlling Cellular Inositol Pyrophosphate Levels in Schizosaccharomyces pombe. *Mol Cell Biol* 38: pii: e00047-18

Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC & Ferrin TE (2004) UCSF Chimera--a visualization system for exploratory research and analysis. *J Comput Chem* 25: 1605–1612

Pöhlmann J, Risse C, Seidel C, Pohlmann T, Jakopec V, Walla E, Ramrath P, Takeshita N, Baumann S, Feldbrügge M, *et al* (2014) The Vip1 inositol polyphosphate kinase family regulates polarized growth and modulates the microtubule cytoskeleton in fungi. *PLoS Genet* 10: e1004586

Ponting CP & Aravind L (1997) PAS: a multifunctional domain family comes to light. *Curr Biol* 7: R674-677

Punjani A, Rubinstein JL, Fleet DJ & Brubaker MA (2017) cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nat Methods* 14: 290–296

Qiu D, Wilson MS, Eisenbeis VB, Harmel RK, Riemer E, Haas TM, Wittwer C, Jork N, Gu C, Shears SB, *et al* (2020) Analysis of inositol phosphate metabolism by capillary electrophoresis electrospray ionization mass spectrometry. *Nat Commun* 11: 6035

Read RJ & McCoy AJ (2011) Using SAD data in Phaser. *Acta Crystallogr D Biol Crystallogr* 67: 338–344

Ried MK, Wild R, Zhu J, Pipercevic J, Sturm K, Broger L, Harmel RK, Abriata LA, Hothorn LA, Fiedler D, *et al* (2021) Inositol pyrophosphates promote the interaction of SPX domains with the coiled-coil motif of PHR transcription factors to regulate plant phosphate homeostasis. *Nature Communications* 12: 384

Riemer E, Qiu D, Laha D, Harmel RK, Gaugler P, Gaugler V, Frei M, Hajirezaei M-R, Laha NP, Krusenbaum L, *et al* (2021) ITPK1 is an InsP6/ADP phosphotransferase that controls phosphate signaling in Arabidopsis. *Molecular Plant* 14: 1864–1880

Rigden DJ (2008) The histidine phosphatase superfamily: structure and function. *Biochemical Journal* 409: 333–348

Saiardi A, Erdjument-Bromage H, Snowman AM, Tempst P & Snyder SH (1999) Synthesis of diphosphoinositol pentakisphosphate by a newly identified family of higher inositol polyphosphate kinases. *Current Biology* 9: 1323–1326

Scheffzek K & Welti S (2012) Pleckstrin homology (PH) like domains – versatile modules in protein–protein interaction platforms. *FEBS Letters* 586: 2662–2673

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, *et al* (2012) Fiji: an open-source platform for biological-image analysis. *Nat Methods* 9: 676–682

Shears SB (2018) Intimate connections: Inositol pyrophosphates at the interface of metabolic regulation and cell signaling. *Journal Cellular Physiology* 233: 1897–1912

Shukla A, Kaur M, Kanwar S, Kaur G, Sharma S, Ganguli S, Kumari V, Mazumder K, Pandey P, Rouached H, *et al* (2021) Wheat inositol pyrophosphate kinase TaVIH2-3B modulates cell-wall composition and drought tolerance in Arabidopsis. *BMC Biol* 19: 261

Stentz R, Osborne S, Horn N, Li AWH, Hautefort I, Bongaerts R, Rouyer M, Bailey P, Shears SB, Hemmings AM, *et al* (2014) A Bacterial Homolog of a Eukaryotic Inositol Phosphate Signaling Enzyme Mediates Cross-kingdom Dialog in the Mammalian Gut. *Cell Reports* 6: 646–656

Stevenson-Paulik J, Bastidas RJ, Chiou S-T, Frye RA & York JD (2005) Generation of phytate-free seeds in *Arabidopsis* through disruption of inositol polyphosphate kinases. *Proc Natl Acad Sci USA* 102: 12612–12617

Terwilliger TC (2003) SOLVE and RESOLVE: automated structure solution and density modification. *Meth Enzymol* 374: 22–37

Varadi M, Bertoni D, Magana P, Paramval U, Pidruchna I, Radhakrishnan M, Tsenkov M, Nair S, Mirdita M, Yeo J, *et al* (2024) AlphaFold Protein Structure Database in 2024: providing structure coverage for over 214 million protein sequences. *Nucleic Acids Research* 52: D368–D375

Wang H, Falck JR, Hall TMT & Shears SB (2012) Structural basis for an inositol pyrophosphate kinase surmounting phosphate crowding. *Nat Chem Biol* 8: 111–116

Wang H, Godage HY, Riley AM, Weaver JD, Shears SB & Potter BVL (2014) Synthetic Inositol Phosphate Analogs Reveal that PPIP5K2 Has a Surface-Mounted Substrate Capture Site that Is a Target for Drug Discovery. *Chemistry & Biology* 21: 689–699

Wang H, Gu C, Rolfes RJ, Jessen HJ & Shears SB (2018a) Structural and biochemical characterization of Siw14: A protein-tyrosine phosphatase fold that metabolizes inositol pyrophosphates. *Journal of Biological Chemistry* 293: 6905–6914

Wang H, Lo W-T, Vujičić Žagar A, Gulluni F, Lehmann M, Scapozza L, Haucke V & Vadas O (2018b) Autoregulation of Class II Alpha PI3K Activity by Its Lipid-Binding PX-C2 Domain Module. *Mol Cell* 71: 343-351.e4

Wang H, Nair VS, Holland AA, Capolicchio S, Jessen HJ, Johnson MK & Shears SB (2015a) Asp1 from *Schizosaccharomyces pombe* Binds a [2Fe-2S] $^{2+}$ Cluster Which Inhibits Inositol Pyrophosphate 1-Phosphatase Activity. *Biochemistry* 54: 6462–6474

Wang Q, Vogan EM, Nocka LM, Rosen CE, Zorn JA, Harrison SC & Kuriyan J (2015b) Autoinhibition of Bruton's tyrosine kinase (Btk) and activation by soluble inositol hexakisphosphate. *Elife* 4: e06074

Wang Z, Jork N, Bittner T, Wang H, Jessen HJ & Shears SB (2020) Rapid stimulation of cellular Pi uptake by the inositol pyrophosphate InsP $_8$ induced by its photothermal release from lipid nanocarriers using a near infra-red light-emitting diode. *Chem Sci* 11: 10265–10278

Weaver JD, Wang H & Shears SB (2013) The kinetic properties of a human PPIP5K reveal that its kinase activities are protected against the consequences of a deteriorating cellular bioenergetic environment. *Bioscience Reports* 33: e00022

Wild R, Gerasimaite R, Jung J-Y, Truffault V, Pavlovic I, Schmidt A, Saiardi A, Jessen HJ, Poirier Y, Hothorn M, *et al* (2016) Control of eukaryotic phosphate homeostasis by inositol polyphosphate sensor domains. *Science* 352: 986–990

Wittmann JG & Rudolph MG (2007) Pseudo-merohedral twinning in monoclinic crystals of human orotidine-5'-monophosphate decarboxylase. *Acta Crystallogr D Biol Crystallogr* 63: 744–749

Yousaf R, Gu C, Ahmed ZM, Khan SN, Friedman TB, Riazuddin S, Shears SB & Riazuddin S (2018) Mutations in Diphosphoinositol-Pentakisphosphate Kinase PPIP5K2 are associated with hearing loss in human and mouse. *PLoS Genet* 14: e1007297

Zhou J, Hu Q, Xiao X, Yao D, Ge S, Ye J, Li H, Cai R, Liu R, Meng F, *et al* (2021) Mechanism of phosphate sensing and signaling revealed by rice SPX1-PHR2 complex structure. *Nat Commun* 12: 7040

Zhu J, Lau K, Puschmann R, Harmel RK, Zhang Y, Pries V, Gaugler P, Broger L, Dutta AK, Jessen HJ, *et al* (2019) Two bifunctional inositol pyrophosphate kinases/phosphatases control plant phosphate homeostasis. *eLife* 8: e43582

  Figure legends

Figure 1 PPIP5K enzymes contain an unusual histidine acid phosphate domain.

(**A**) Schematic diagram of the *S. cerevisiae* Vip1 kinase (orange, $ScVip1^{KD}$) and phosphatase (blue, $ScVip1^{PD}$)

domains (https://www.uniprot.org/ Uniprot ID Q06685), highlighting structured regions as blocks (residues

186-520 and 536-1094) and putative linker regions as solid lines (residues 1-185, 521-535, and 1095-1146).

(**B**) Ribbon diagram of the $ScVip1^{KD}$ structure bound to ADP (in bonds representation) and including a omit

2 ($F_o - F_c$) electron density map contoured at 1 σ (blue mesh). (**C**) Ribbon diagram of $ScVip1^{PD\ \Delta848-918}$, with

the histidine acid phosphatase core and α-helical insertion domain shown in light blue, and the PPIP5K-

specific GAF domain in dark blue, respectively. The position of the engineered GSSG linker is indicated

with a dotted line. The inlet provides a view of the zinc binding site, with the Zn^{2+} ion shown as a sphere (in

gray) and the coordinating histidine and cysteine residues depicted in bonds representation.

Figure 2 A unique GAF domain in PPIP5Ks replaces the canonical β1-insert in histidine acid

phosphatases.

(**A**) Structural comparison of $ScVip1^{PD\ \Delta848-918}$ (ribbon diagram, in blue) with the *E. coli* phytase AppA

(https://www.rcsb.org/ PDB-ID 7Z2S, yellow to red) (Acquistapace *et al*, 2022), human prostatic acid

phosphatase (PDB-ID 1ND6, light- to dark-green) (Ortlund *et al*, 2003), human liver fructose-2,6-

bisphosphatase (PDB-ID 1K6M, purple and dark gray) (Lee *et al*, 2003) and the sensor histidine kinase

phosphatase SixA from *E. coli* (PDB-ID 1UJC, pink and light gray) (Hamada *et al*, 2005). (**B**) Structural

superposition of $ScVip1^{PD\ \Delta848-918}$ and EcAppA (colors as in panel A, root mean square deviation r.m.s.d. is ~

1.1 Å comparing 105 corresponding C_α atoms from the histidine acid phosphatase core). The Zn^{2+} binding

site in $ScVip1^{PD}$ is not conserved among other histidine acid phosphatases. (**C**) Structural comparison of the

GAF domain in $ScVip1^{PD}$ and structurally related GAF (PDB-ID 1F5M, r.m.s.d. is ~0.9 Å comparing 26

corresponding C_α atoms in the β-sheet) (Ho *et al*, 2000) and inositol polyphosphate-binding PH domains

(PDB-IDs 4Y94, 1UNQ) (Wang *et al*, 2015b) (Milburn *et al*, 2003). Note that GAF and PH domains share

similar sized β-sheets with different topologies.

Figure 3 $ScVip1^{PD}$ specifically hydrolyzes 1,5-InsP$_8$ using a conserved phytase active site.

(**A**) Table summary of the initial enzyme velocities (v_0) for $ScVip1^{PD}$ versus 1,5-InsP$_8$, 1-InsP$_7$, or 5-InsP7

derived from an NMR-based phosphatase assay. Pseudo-2D NMR spectra of $ScVip1^{PD}$ incubated with

different substrates (blue traces) are shown below, overlaid with the respective reaction products 5-InsP$_7$, or

InsP$_6$ (red traces). (**B**) Initial enzyme velocities (v_0) for $ScVip1^{PD}$ – catalyzed hydrolysis of 1,5-InsP$_8$ and 5-

InsP$_7$ as a function of inositol pyrophosphate concentration and using a malachite green-based phosphatase

assay. The Michaelis-Menten equation was fitted to the data using non-linear regression. Extracted kinetic

parameters are summarized in the table as means ± S.E.M. (n=2). (**C**) Structural superposition of the active

sites of $ScVip1^{PD\ \Delta848-918}$ (in bond representation, in dark blue) and EcAppA (in orange, PDP-ID 1DKL,

r.m.s.d. is ~1.0 Å comparing 67 corresponding C_α atoms from the histidine acid phosphatase core). In
EcAppA, His17 is the catalytic residue, Asp304 is the proton-donor, and Arg16, Arg20, and Arg92 are
involved in $InsP_6$ substrate binding. (**D-E**) Initial velocities of 1,5-$InsP_8$ hydrolysis catalyzed by wild-type
and mutant versions of $ScVip1^{PD}$, measured by NMR-based (**D**) or malachite green-based (**E**) phosphatase
assays.

Figure 4 $ScVip1^{PD}$ phosphatase activity is inhibited by inorganic phosphate and sulfate.
(**A-B**) NMR time course experiments for $ScVip1^{PD}$-catalyzed hydrolysis of 100 µM 1,5-$InsP_8$ to 5-$InsP_7$ in
the presence of increasing concentrations of (**A**) PO_4^{2-}, (**B**) SO_4^{2-}, or (**C**) at a constant concentration of 25
mM $NaNO_3$. The measured initial velocity of 980 nmol min^{-1} mg^{-1} is similar to the initial velocity in absence
of $NaNO_3$ (1100 nmol min^{-1} mg^{-1}, compare Fig3A).

Figure 5 $ScVip1^{PD}$ is regulated by GAF domain movements and stabilized by a zinc-binding site.
(**A**) Structural superposition of apo $ScVip1^{PD\ \Delta848-918}$ reveals the GAF insertion domain in either its open
(chain A, dark blue ribbon diagram) or closed (chain D, in light blue) conformation. The RHxxR and hinge
regions are shown alongside (in bonds representation) (**B**) Details of the $ScVip1^{PD\ \Delta848-918}$ active site: Asp550
stabilize the catalytic loop by engaging in a salt bridge (green dotted line) with the Zn^{2+} ion coordinating
His836. The active site and Zn^{2+} coordinating residues are depicted in bonds representation, the Zn^{2+} ion is
shown as a gray sphere. (**C-D**) Initial velocities of 1,5-$InsP_8$ hydrolysis catalyzed by wild-type or mutant
version of $ScVip1^{PD}$, measured by NMR (**C**) or malachite green (**D**) phosphatase assays. (**E**) Table
summarizing the thermal melting profiles of wild-type or mutant $ScVip1^{PD}$. The ΔTm (in ºC) represent the
difference between the melting temperature of a designated $ScVip1^{PD}$ mutant protein and the wild-type
control.

Figure 6 The GAF domain of $ScVip1^{PD}$ binds 1,5-$InsP_8$ in two capture sites.
(**A**) Ribbon diagrams of $ScVip1^{PD\ \Delta848-918\ RHR-AAA}$ bound to 1,5-$InsP_8$ (in bonds representation). 1,5-$InsP_8$ binds
either at the surface of $ScVip1^{PD\ \Delta848-918\ RHR-AAA}$ (chain A, top) or close to the active site (chain B, bottom). (**B**)
Detailed view of the 1,5-$InsP_8$ molecules shown in two orientations and including a 2 ($F_o - F_c$) electron
density map (blue mesh) contoured at 2.5 σ in chain A (top) or at 1 σ in chain B (bottom), respectively. "1-β-
P" and "5-β-P" designate the positions of the β-phosphate of the pyrophosphate group attached to the
position 1 or 5 of the *myo*-inositol ring. (**C**) Detailed view of the 1,5-$InsP_8$ binding sites. The residues
interacting with 1,5-$InsP_8$ are shown in bonds representation, hydrogen bonds are indicted by dotted lines (in
green). (**D**) Ribbon diagram of the isolated GAF domain from the $ScVip1^{PD\ \Delta848-918\ RHR-AAA}$ structure, depicting
the relative positions of the outer (chain A) and inner (chain B, modeled) 1,5-$InsP_8$ substrate (in bonds
representation) binding sites.

Figure 7 A GAF domain-induced substrate channel transports 1,5-InsP$_8$ to the active site.

(**A**) Electrostatic surface charge distribution of the apo ScVip1$^{PD \; \Delta848\text{-}918}$ (top) and 1,5-InsP$_8$-bound ScVip1^{PD}

$^{\Delta848\text{-}918 \; RHR\text{-}AAA}$ (bottom) structures. (**B**) Evolutionary conservation (from cyan, variable to pink, invariant) of

surface residues mapped onto the apo ScVip1$^{PD \; \Delta848\text{-}918}$ (top) and 1,5-InsP$_8$-bound ScVip1$^{PD \; \Delta848\text{-}918 \; RHR\text{-}AAA}$

(bottom) structures. Views as in (**A**). Residue conservation was derived from the multiple sequence

alignment shown in Fig EV2 and analyzed with Consurf (Landau *et al*, 2005) (**C**) Cutaway front view of the

apo ScVip1$^{PD \; \Delta848\text{-}918}$ surface modeled with the outer and inner 1,5-InsP$_8$ molecules (top). Same view for the

1,5-InsP$_8$-bound ScVip1$^{PD \; \Delta848\text{-}918 \; RHR\text{-}AAA}$ surface modeled with the outer and inner 1,5-InsP$_8$ molecules

(bottom). The closed GAF domain forms the substrate channel. (**D**) Groups of mutated residues involved in

1,5-InsP$_8$ channeling (in bonds representation) mapped onto the ScVip1$^{PD \; \Delta848\text{-}918 \; RHR\text{-}AAA}$ structure (shown as

molecular surface). The active site of ScVip1^{PD} is indicated with a red asterisk. (**E-F**) Initial velocities of 1,5-

InsP$_8$ hydrolysis for wild-type or mutant versions of ScVip1^{PD}, measured by NMR-based (**E**) or malachite

green-based (**F**) phosphatase assays.

Figure 8 A conserved glutamate controls substrate sterospecificity in ScVip1^{PD}.

(**A**) Structure-based sequence alignment of the presumed catalytic motifs in ScVip1^{PD}, *E. coli* AppA and the

multiple inositol polyphosphate phosphatase MINPP from *Bacteroides thetaiotaomicron*. The RHxxR motif

is conserved among all enzymes, while the proton donor aspartate in EcAppA is replaced by Glu991 in

ScVip1^{PD} and by an alanine residue in MINPP. (**B**) Structural comparison of the ScVip1^{PD}, EcAppA (PDB-ID

7Z2S) (Acquistapace *et al*, 2022) and MINPP (PDB-ID 4FDU) (Stentz *et al*, 2014) active sites. Glu991 (in

bonds representation, colored from cyan to blue) adopts different conformation in our structures. Asp304

controlling substrate stereospecificity in EcAppA (Acquistapace *et al*, 2022), and Ala324 in the highly

promiscuous MINPP enzyme are highlighted in blue. Initial velocities of 1,5-InsP$_8$, 1-InsP$_7$ and 5-InsP$_7$

hydrolysis for wild-type or Glu991Ala versions of ScVip1^{PD}, measured by NMR-based (**C**) or malachite

green-based (**D**) phosphatase assays.

Figure 9 Structure-function analysis of the AtVIH2 phosphatase domain.

(**A**) Ribbon diagram of the ScVip1PD$^{\Delta848\text{-}918}$ structure, with the histidine phosphatase core in white, the α-

helical insert in light gray, and the GAF domain in dark gray. Residues involved in phosphatase catalysis,

Zn^{2+} ion coordination, GAF domain movement, and 1,5-InsP$_8$ binding are shown in bond representation.

Corresponding residues in AtVIH2 are also indicated, illustrating the conserved functional regions. (**B**)

Rosette phenotype of three weeks old Col-0 wild-type plants and lines expressing full-length Flag-tagged

versions of AtVIH2 under the control of the UBI10 promoter. (**C**) Quantification of the rosette area. Box

plots show 16 plants per genotype. Multiple comparisons of the genotypes vs. Col-0 were performed

according to Dunnett (Dunnett, 1955) as implemented in GraphPad prism v10.3.0 (*** $p < 0.001$, ** $p <$

0.005, * $p < 0.01$). (**D**) Quantification of cellular Pi levels. For each genotype, 8 individual plants were

measured using 3 technical replicates. The estimated Pi concentration was normalized by fresh weight. (**E**)

Western blot of Flag-tagged AtVIH2. The theoretical molecular mass of AtVIH2 is ~118 kDa (indicated by a

black arrow; 1:Col-0, 2:AtVIH2^{WT},3:AtVIH2$^{K219A-D292A}$, 4:AtVIH2$^{R372A-H373A}$, 5:AtVIH2$^{R372A-H373A-R376A}$,

6:AtVIH2$^{R372A-H373A-R376A-E882A}$, 7:AtVIH2^{H741A}, 8:AtVIH2^{C696A}, 9:AtVIH2$^{H505A-H741A-H975A}$, 10:AtVIH2$^{P378V-G500V-}$

^{G501A}, 11:AtVIH2$^{R722S-D724A-T726A}$, 12:AtVIH2$^{K383A-K447A-K598A-R722A}$, 13:AtVIH2$^{K379A-K381A-K498A}$.

Figure 10 Arrangement of the the kinase and phosphatase domains in a full-length PPIP5K.

(**A**) Coomassie blue-stained SDS-PAGE of purified SpAsp1. (**B**) A representative micrograph of the

negatively stained SpAsp1 used for image processing. (**C**) Low resolution envelope of SpAsp1 determined

by negative stain electron microscopy (left). Individual crystal structures of ScVip1KD (orange ribbon

diagram) and ScVip1$^{PD\ \Delta848-918}$ (blue ribbon diagram) were docked into the L-shaped negative stain map of

SpAsp1 (right). The linkers truncated in the crystal structures of ScVip1 are potentially visible in the SpAsp1

envelope (black dashed lines). (**D**) Initial velocities of 1,5-InsP$_8$ hydrolysis catalyzed by ScVip1^{PD} or

ScVip1^{KD-PD}, measured by NMR-based phosphatase assays. (**E**) Hydrogen-deuterium exchange data mapped

on the ScVip1^{KD-PD} model. The protected regions in ScVip1 after incubation with non-hydrolysable 1,5-InsP$_8$

(PCP-IP$_8$) are shown in light green, the protected regions after incubation with non-hydrolysable 1,5-InsP$_8$

(PCP-IP$_8$) and ATP (AMPPNP) analogs, respectively are shown in dark green.

Figure EV1 Expression of AtVIH1^{PD}, AtVIH2^{PD} and MpVIP1^{PD} phosphatase domains in baculovirus-

infected Tnao insect cells.

Western blot analysis showing the expression level and solubility of different PPIP5K phosphatase domain

(PD) constructs from different plant species. The constructs analyzed are *Arabidopsis thaliana* (*At*) VIH1^{PD}

(residues 351-1010, Uniprot ID Q84WW3), VIH2^{PD} (residues 351-1046, Uniprot ID F4J8C6), and

Marchantia polymorpha (*Mp)* VIP1^{PD} (residues 382-1041 Uniprot ID Mp8g06840).

Figure EV2 Multiple sequence alignment of PPIP5K phosphatase domains.

Multiple-sequence alignment of S. *cerevisiae* Vip1 (Uniprot ID Q06685) with *Schizosaccharomyces pombe*

Asp1 (Uniprot ID O74429), *Arabidopsis thaliana* VIH1 and VIH2 (Uniprot ID Q84WW3 and F4J8C6,

respectively), *Marchantia polyorpha* VIP1 (Uniprot ID Mp8g06840), and *Homo sapiens* PPIP5K1 and

PPIP5K2 (Q6PFW1 and O43314, respectively). Secondary structure elements are colored according to

different domains as indicated on top. Functionally important conserved residues are highlighted. Dashed

lines indicate regions of the protein truncated in the engineered constructs of ScVip1 kinase domain (residues

186-522) and ScVip1 phosphatase domain (residues 536-1107, Δ848-918) used for crystallization.

Figure EV3 ScVip1 harbors a structurally conserved PPIP5K kinase domain.

Structural comparison of different PPIP5K kinase domain structures. Shown are C$_\alpha$ traces of the

Saccharomyces cerevisiae Vip1 kinase domain (in orange, bound to ADP in bonds representation), *Homo*

sapiens PPIP5K2 (PDB id: 3T9E, in green and bound to ADP and 5-InsP$_7$) (Wang *et al*, 2012) and

Schizosaccharomyces pombe Asp1 (PDB id: 8E1H, in purple) (Benjamin *et al*, 2022b). The kinase structures

align with a r.m.s.d of ~1.5 Å comparing 310 C$_\alpha$ atoms.

Figure EV4 ScVip1^{PD} is structurally related to histidine acid phosphatases and contains a GAF

domain.

Top twenty hits from the DALI web server (http://ekhidna2.biocenter.helsinki.fi/dali/) (Holm & Rosenström,

2010) using the histidine acid phosphatase core containing the α-helical insertion domain (residues 536-553;

647-1094) (**A**), or the isolated GAF domain in ScVip1^{PD} (residues 554-646) (**B**) as search model against the

Protein Data Bank.

Figure EV5 NMR-based enzyme assays – raw data.

NMR time course experiments of PP-InsP hydrolysis catalyzed by ScVip1^{PD} and respective mutant constructs

using 100µM of [^{13}C$_6$]-labeled PP-InsP substrates. Concentrations of enzyme were varied based on the

respective construct and substrate. PP-InsP turnover was measured using a pseudo-2D spin-echo difference

nuclear magnetic resonance spectroscopy experiment. Quantification was based on the relative intensities of

the C2 signals of the respective (PP-)InsP species.

Figure EV6 Arrangement of GAF domains in the ScVip1^{PD} and ScVip1$^{PD\Delta848-918}$ apo structures.

(**A**) Structural superposition of chains A (shown as C$_\alpha$ traces) from the ScVip1^{PD} (in blue) and ScVip1$^{PD\Delta848-918}$

(in cyan) apo crystal forms reveal the GAF domain in an open conformation (r.m.s.d is ~0.5 Å comparing

457 corresponding C$_\alpha$ atoms). The asymmetric unit of ScVip1^{PD} apo contains four molecules, all of which

present the GAF domain in the open conformation. (**B**) ScVip1$^{PD\Delta848-918}$ crystals also contain four molecules

in the asymmetric unit with chains A and B in the open and chains C and D in the closed conformation,

respectively. Shown are ribbon diagrams, colors as in Fig 1B. (**C**) A second Zn^{2+} binding site in ScVip1$^{PD\Delta848-918}$

is formed along a pseudo 2-fold axis involved two neighboring GAF domains. This second binding site

likely represent a crystallization artifact. The crystallization condition contained an excess of (CH$_3$CO$_2$)$_2$Zn

and the residues involved in Zn^{2+} binding are not conserved among other PPIP5Ks (His573, Glu575,

compare Fig EV2).

Figure EV7 Structural integrity profiling of *Sc*Vip1^{PD} mutant proteins.

(**A**) Coomassie blue-stained SDS-PAGE analysis of wild-type and mutant versions of ScVip1^{PD} (theoretical

molecular weight is ~65.5 kDa). (**B**) Analytical size-exclusion chromatography of wild-type and mutant

versions of ScVip1PD. Shown are A$_{280nm}$ absorption traces as a function of column elution volume (in mL).

The red arrow indicates the characteristic elution volume of ScVip1PD, which behaves as a monomer in

solution. (**C**) Melting temperatures of wild-type (blue traces, N=5) and mutant versions of ScVip1^{PD} derived

from thermal shift assays (N.D. not determined). Note that addition of divalent metal chelators such as EDTA

reduces the thermal stability of the enzyme. Addition of $ZnCl_2$ but not $MgCl_2$ increases ScVip1^{PD} stability,

likely by targeting the structural Zn^{2+} binding site.

Figure EV8 An intermolecular disulfide bridge stabilizes the crystallographic dimer in the *Sc*Vip1^{PD}

$^{\Delta848-918\ RHR-AAA}$ – 1,5-InsP$_8$ complex structure.

Ribbon diagram of a crystallographic domain-swapped dimer in the asymmetric unit of our ScVip1$^{PD\ \Delta848-918}$

$^{RHR-AAA}$ crystals (chain A is shown in blue, chain B in green, 1,5-InsP$_8$ molecules are depicted in bonds

representation). The inlet provides a detailed view of the Zn^{2+} binding site, with the Zn^{2+} ions shown as a

sphere (in gray) and the coordinating residues depicted in bonds representation. The view includes a 2 (F_o –

F_c) electron density map contoured at 1 σ (blue mesh). Cys793 is involved in an intermolecular disulfide

bridge between ScVip1$^{PD\ \Delta848-918\ RHR-AAA}$ chain A (in blue) and chain B (in green). In the crystal structure of the

apo ScVip1$^{PD\ \Delta848-918}$ (compare Fig1C), Cys793 replaces Glu796 and coordinates the Zn^{2+} ion.

EV9 Summary of the HDX-MS experiments

(**A**) Experimental details of the HDX analyses. (**B**) D_2O incorporation into ScVip1^{KD-PD}. (**C**) HDX differences

comparing apo ScVip1^{KD-PD}, ScVip1^{KD-PD} in complex with 1,5-InsP$_8$ and ScVip1^{KD-PD} in the presence of 1,5-

InsP$_8$ and a non-hydrolyzable ATP analog (blue, protection; red, deprotection).

Figure EV10 The "HD" catalytic motif presents in phytases is not present in ScVip1^{PD}.

Detailed views of the active sites of *E. coli* phytase AppA (in orange, PDP-ID 7Z2S) (Acquistapace *et al*,

2022) bounds to the substrate analogue InsS$_6$, *S. cerevisiae* Vip1$^{PD\ \Delta848-918}$, and the Alphafold model of *S.*

pombe Asp1^{PD} (in purple, https://alphafold.ebi.ac.uk/entry/O74429) (Varadi *et al*, 2024). The InsP$_6$ and 1,5-

InsP$_8$ molecules are depicted in bonds representation, and the side chains of residues belonging to the

catalytic motifs 1 and 2 are depicted bonds representation. Residues His303 and Asp304 forming the HD

motif in AppA correspond to Lys990 and Glu991 in ScVip1^{PD} and not to His993 and Ile994, as previously

thought (Fridy *et al*, 2007; Mulugu *et al*, 2007; Pöhlmann *et al*, 2014; Wang *et al*, 2015a; Pascual-Ortiz *et al*,

2018).

Table EV1 Synthetic gene and primer sequences used in this study

Table EV2 Overview of transgenic Arabidopsis lines used in this study

Dataset EV1 Hydrogen/deuterium exchange mass spectrometry – raw data and experimental settings

Table 1 – Crystallographic data collection and refinement statistics.

	ScVip1[KD] ADP	ScVip1[PD] Pt MR-SAD	ScVip1[PD] apo	ScVip1[PDΔ848-918] apo	ScVip1[PD Δ848-918] RHR-AAA 1,5-InsP$_8$
PDB-ID	9GR8		9GRH	9GRN	9GRO
Data collection					
Wavelength	1.000036	1.071846	0.999993	1.000031	1.000000
Space group	$P\,2_1$	$P\,1$	$P\,2_1$	$C\,2$	$P\,3_1\,2\,1$
Cell dimensions					
a, b, c (Å)	40.7, 83.2, 51.8	84.5, 84.7, 101.1	84.4, 194.9, 84.4	249.5, 90.5, 200.9	114.6, 114.6, 172.2
α, β, γ (°)	90, 94.32, 90	86.6, 74.1, 86.6	90, 112.2, 90	90, 126.91, 90	90, 90, 120
Resolution (Å)	43.86 – 1.18 (1.25 – 1.18)	48.55 – 3.50 (3.71 – 3.50)	48.7 – 3.05 (3.23 – 3.05)	45.26 – 3.40 (3.60 – 3.40)	49.68 – 2.36 (2.51 – 2.36)
R_{meas}[#]	0.104 (2.02)	0.106 (1.04)	0.158 (1.71)	0.46 (2.70)	0.110 (2.94)
CC(1/2)[#]	1.0 (0.55)	1.0 (0.40)	1.0 (0.48)	0.98 (0.30)	1.0 (0.52)
$I/\sigma\,I$[#]	13.3 (1.2)	7.2 (1.0)	10.1 (1.2)	5.2 (1.0)	19.6 (1.0)
Completeness (%)[#]	97.5 (93.4)	95.2 (95.3)	99.7 (99.8)	99.0 (98.5)	99.9 (99.6)
Redundancy[#]	6.7 (6.2)	1.8 (1.9)	7.0 (7.2)	6.9 (7.2)	20.4 (19.7)
Wilson B-factor[#]	18.3	111.3	83.8	72.2	73.2
Phasing					
No. sites (Pt/Zn)		4/4			
Figure of merit[*]		0.64			
Refinement					
Resolution (Å)	43.86 – 1.18		48.7 – 3.05	45.26 – 3.50	49.68 – 2.36
No. reflections	215,320		48,029	49,444	54,125
Twin operator / fraction[+]			l, -k, h / 0.5		
R_{work} / R_{free}[+]	0.15 / 0.18		0.267 / 0.31	0.24 / 0.29	0.22 / 0.26
No. atoms					
protein	2,845		15,722	15,461	7,819
ligand	47		4	255	241
solvent	397			46	51
Res. B-factors[+]					
protein	19.8		120.1	101.1	91.1
ligand	28.3		113.1	113.9	110.3
solvent	34.7			57.3	74.3
R.m.s deviations[$]					
bond lengths (Å)	0.021		0.002	0.006	0.003
bond angles (°)	1.81		0.52	1.0	0.53
Ramachandran plot[$]:					
most favored regions (%)	98.2		95.8	96.7	96.7
outliers (%)	0		0.5	0.1	0.0
MolProbity score[$]	1.21		1.84	2.18	1.30

912 [#]as defined in XDS (Kabsch, 1993)
913 [*]as defined in PHASER-EP (Read & McCoy, 2011)
914 [+]as defined in phenix.refine (Afonine *et al*, 2012)
915 [$]as defined in Molprobity (Davis et al, 2007)

Figure 1. PPIP5K enzymes contain an unusual histidine acid phosphate domain.
(**A**) Schematic diagram of the *S. cerevisiae* Vip1 kinase (orange, ScVip1[KD]) and phosphatase (blue, ScVip1[PD]) domains (https://www.uniprot.org/ Uniprot ID Q06685), highlighting structured regions as blocks (residues 186-520 and 536-1094) and putative linker regions as solid lines (residues 1-185, 521-535, and 1095-1146). (**B**) Ribbon diagram of the ScVip1[KD] structure bound to ADP (in bonds representation) and including a omit 2 ($F_o − F_c$) electron density map contoured at 1 σ (blue mesh). (**C**) Ribbon diagram of ScVip1[PD Δ848-918], with the histidine acid phosphatase core and α-helical insertion domain shown in light blue, and the PPIP5K-specific GAF domain in dark blue, respectively. The position of the engineered GSSG linker is indicated with a dotted line. The inlet provides a view of the zinc binding site, with the Zn^{2+} ion shown as a sphere (in gray) and the coordinating histidine and cysteine residues depicted in bonds representation.

Figure 2. A unique GAF domain in PPIP5Ks replaces the canonical β1-insert in histidine acid phosphatases.
(**A**) Structural comparison of ScVip1$^{PD\,\Delta848-918}$ (ribbon diagram, in blue) with the *E. coli* phytase AppA (https://www.rcsb.org/ PDB-ID 7Z2S, yellow to red) (Acquistapace *et al*, 2022), human prostatic acid phosphatase (PDB-ID 1ND6, light- to dark-green) (Ortlund *et al*, 2003), human liver fructose-2,6-bisphosphatase (PDB-ID 1K6M, purple and dark gray) (Lee *et al*, 2003) and the sensor histidine kinase phosphatase SixA from *E. coli* (PDB-ID 1UJC, pink and light gray) (Hamada *et al*, 2005). (**B**) Structural superposition of ScVip1$^{PD\,\Delta848-918}$ and EcAppA (colors as in panel A, root mean square deviation r.m.s.d. is ~ 1.1 Å comparing 105 corresponding C$_\alpha$ atoms from the histidine acid phosphatase core). The Zn^{2+} binding site in ScVip1^{PD} is not conserved among other histidine acid phosphatases. (**C**) Structural comparison of the GAF domain in ScVip1^{PD} and structurally related GAF (PDB-ID 1F5M, r.m.s.d. is ~0.9 Å comparing 26 corresponding C$_\alpha$ atoms in the β-sheet) (Ho *et al*, 2000) and inositol polyphosphate-binding PH domains (PDB-IDs 4Y94, 1UNQ) (Wang *et al*, 2015b) (Milburn *et al*, 2003). Note that GAF and PH domains share similar sized β-sheets with different topologies.

Reaction	v_0 (nmol min^{-1} mg^{-1})
1,5-InsP$_8$ → 5-InsP$_7$	1100
1-InsP$_7$ → InsP$_6$	230
5-InsP$_7$ → InsP$_6$	0

Substrate	Vmax (nmol min^{-1} mg^{-1})	K_M (µM)	k_{cat} (s^{-1})	k_{cat}/K_M (10^6M^{-1}s^{-1})
1,5-InsP$_8$	1570 ± 100	44 ± 12	1.4 ± 0.08	0.032
1-InsP$_7$	147 ± 13	72 ± 28	0.13 ± 0.01	0.0018

Residue mutations	v_0 (nmol min^{-1} mg^{-1})
WT	1100
ΔG848-S918	2000
R547A-H548A-R551A	30
R547A	2
H548A	0
R551A	10
E991A	2200

Figure 3. ScVip1^{PD} specifically hydrolyzes 1,5-InsP$_8$ using a conserved phytase active site.
(**A**) Table summary of the initial enzyme velocities (v $_0$) for ScVip1^{PD} versus 1,5-InsP$_8$, 1-InsP$_7$, or 5-InsP$_7$ derived from an NMR-based phosphatase assay. Pseudo-2D NMR spectra of ScVip1^{PD} incubated with different substrates (blue traces) are shown below, overlaid with the respective reaction products 5-InsP$_7$, or InsP$_6$ (red traces). (**B**) Initial enzyme velocities (v $_0$) for ScVip1^{PD} – catalyzed hydrolysis of 1,5-InsP$_8$ and 5-InsP$_7$ as a function of inositol pyrophosphate concentration and using a malachite green-based phosphatase assay. The Michaelis-Menten equation was fitted to the data using non-linear regression. Extracted kinetic parameters are summarized in the table as means ± S.E.M. (n=2). (**C**) Structural superposition of the active sites of ScVip1^{PD} $^{Δ848-918}$ (in bond representation, in dark blue) and EcAppA (in orange, PDP-ID 1DKL, r.m.s.d. is ~1.0 Å comparing 67 corresponding C$_α$ atoms from the histidine acid phosphatase core). In EcAppA, His17 is the catalytic residue, Asp304 is the proton-donor, and Arg16, Arg20, and Arg92 are involved in InsP$_6$ substrate binding. (**D-E**) Initial velocities of 1,5-InsP$_8$ hydrolysis catalyzed by wild-type and mutant versions of ScVip1^{PD}, measured by NMR-based (**D**) or malachite green-based (**E**) phosphatase assays.

Figure 4. ScVip1[PD] phosphatase activity is inhibited by inorganic phosphate and sulfate.
(**A-B**) NMR time course experiments for ScVip1[PD]-catalyzed hydrolysis of 100 µM 1,5-InsP$_8$ to 5-InsP$_7$ in the presence of increasing concentrations of (**A**) PO$_4$$^{2-}$, (**B**) SO$_4$$^{2-}$, or (**C**) at a constant concentration of 25 mM NaNO$_3$. The measured initial velocity of 980 nmol min^{-1} mg^{-1} is similar to the initial velocity in absence of NaNO$_3$ (1100 nmol min^{-1} mg^{-1}, compare Fig3A).

Figure 5. ScVip1[PD] is regulated by GAF domain movements and stabilized by a zinc-binding site.
(**A**) Structural superposition of apo ScVip1[PD] [Δ848-918] reveals the GAF insertion domain in either its open (chain A, dark blue ribbon diagram) or closed (chain D, in light blue) conformation. The RHxxR and hinge regions are shown alongside (in bonds representation) (**B**) Details of the ScVip1[PD] [Δ848-918] active site: Asp550 stabilize the catalytic loop by engaging in a salt bridge (green dotted line) with the Zn^{2+} ion coordinating His836. The active site and Zn^{2+} coordinating residues are depicted in bonds representation, the Zn^{2+} ion is shown as a gray sphere. (**C-D**) Initial velocities of 1,5-InsP8 hydrolysis catalyzed by wild-type or mutant version of ScVip1[PD], measured by NMR (**C**) or malachite green (**D**) phosphatase assays. (**E**) Table summarizing the thermal melting profiles of wild-type or mutant ScVip1[PD]. The ΔTm (in ºC) represent the difference between the melting temperature of a designated ScVip1[PD] mutant protein and the wild-type control.

Figure 6. The GAF domain of ScVip1^{PD} binds 1,5-InsP$_8$ in two capture sites.
(**A**) Ribbon diagrams of ScVip1$^{PD \ \Delta848-918 \ RHR-AAA}$ bound to 1,5-InsP$_8$ (in bonds representation). 1,5-InsP$_8$ binds either at the surface of ScVip1$^{PD \ \Delta848-918 \ RHR-AAA}$ (chain A, top) or close to the active site (chain B, bottom). (**B**) Detailed view of the 1,5-InsP$_8$ molecules shown in two orientations and including a 2 ($F_o - F_c$) electron density map (blue mesh) contoured at 2.5 σ in chain A (top) or at 1 σ in chain B (bottom), respectively. "1-β-P" and "5-β-P" designate the positions of the β-phosphate of the pyrophosphate group attached to the position 1 or 5 of the *myo*-inositol ring. (**C**) Detailed view of the 1,5-InsP$_8$ binding sites. The residues interacting with 1,5-InsP$_8$ are shown in bonds representation, hydrogen bonds are indicted by dotted lines (in green). (**D**) Ribbon diagram of the isolated GAF domain from the ScVip1$^{PD \ \Delta848-918 \ RHR-AAA}$ structure, depicting the relative positions of the outer (chain A) and inner (chain B, modeled) 1,5-InsP$_8$ substrate (in bonds representation) binding sites.

Figure 7. A GAF domain-induced substrate channel transports 1,5-InsP$_8$ to the active site.

(**A**) Electrostatic surface charge distribution of the apo ScVip1$^{PD\ \Delta848-918}$ (top) and 1,5-InsP$_8$-bound ScVip1$^{PD\ \Delta848-918\ RHR-AAA}$ (bottom) structures. (**B**) Evolutionary conservation (from cyan, variable to pink, invariant) of surface residues mapped onto the apo ScVip1$^{PD\ \Delta848-918}$ (top) and 1,5-InsP$_8$-bound ScVip1$^{PD\ \Delta848-918\ RHR-AAA}$ (bottom) structures. Views as in (A). Residue conservation was derived from the multiple sequence alignment shown in Fig EV2 and analyzed with Consurf (Landau *et al*, 2005) (**C**) Cutaway front view of the apo ScVip1$^{PD\ \Delta848-918}$ surface modeled with the outer and inner 1,5-InsP$_8$ molecules (top). Same view for the 1,5-InsP$_8$-bound ScVip1$^{PD\ \Delta848-918\ RHR-AAA}$ surface modeled with the outer and inner 1,5-InsP$_8$ molecules (bottom). The closed GAF domain forms the substrate channel. (**D**) Groups of mutated residues involved in 1,5-InsP$_8$ channeling (in bonds representation) mapped onto the ScVip1$^{PD\ \Delta848-918\ RHR-AAA}$ structure (shown as molecular surface). The active site of ScVip1^{PD} is indicated with a red asterisk. (**E-F**) Initial velocities of 1,5-InsP$_8$ hydrolysis for wild-type or mutant versions of ScVip1^{PD}, measured by NMR-based (**E**) or malachite green-based (**F**) phosphatase assays.

Figure 8. A conserved glutamate controls substrate sterospecificity in ScVip1[PD].

(**A**) Structure-based sequence alignment of the presumed catalytic motifs in ScVip1[PD], *E. coli* AppA and the multiple inositol polyphosphate phosphatase MINPP from *Bacteroides thetaiotaomicron*. The RHxxR motif is conserved among all enzymes, while the proton donor aspartate in *E. coli* AppA is replaced by Glu991 in ScVip1[PD] and by an alanine residue in MINPP. (**B**) Structural comparison of the ScVip1[PD], AppA (PDB-ID 7Z2S) (Acquistapace *et al*, 2022) and MINPP (PDB-ID 4FDU) (Stentz *et al*, 2014) active sites. Glu991 (in bonds representation, colored from cyan to blue) adopts different conformation in our structures. Asp304 controlling substrate stereospecificity in AppA (Acquistapace *et al*, 2022), and Ala324 in the highly promiscuous MINPP enzyme are highlighted in purple. Initial velocities of 1,5-InsP$_8$, 1-InsP$_7$ and 5-InsP$_7$ hydrolysis for wild-type or Glu991Ala versions of ScVip1[PD], measured by NMR-based (**C**) or malachite green-based (**D**) phosphatase assays.

A

Hinge of the GAF domain
Sc G646-G647 / At G500-G501
Sc P553 / At P378

Zinc binding site
Sc H836 / At H741
Sc C793 / At C696
Sc H1064 / At H975
Sc H651 / At H505

1,5-InsP₈ binding site GAF domain
Sc K605 / At K447
Sc K558 / At K383
Sc K644 / At K498
Sc K556 / At K381
Sc K554 / At K379
Sc K605 / At K598

1,5-InsP₈ binding site α-helical insert
Sc K817 / At R722
Sc D819 / At D724
Sc S821 / At T726

Catalytic site
Sc E991 / At E882
Sc R551 / At R376
Sc R547 / At R372
Sc H548 / At H373

Figure 9. Structure-function analysis of the AtVIH2 phosphatase domain.
(**A**) Ribbon diagram of the ScVip1PD$^{\Delta848\text{-}918}$ structure, with the histidine phosphatase core in white, the α-helical insert in light gray, and the GAF domain in dark gray. Residues involved in phosphatase catalysis, Zn^{2+} ion coordination, GAF domain movement, and 1,5-InsP₈ binding are shown in bond representation. Corresponding residues in AtVIH2 are also indicated, illustrating the conserved functional regions. (**B**) Rosette phenotype of three weeks old Col-0 wild-type plants and lines expressing full-length Flag-tagged versions of AtVIH2 under the control of the UBI10 promoter. (**C**) Quantification of the rosette area. Box plots show 16 plants per genotype. Multiple comparisons of the genotypes vs. Col-0 were performed according to Dunnett (Dunnett, 1955) as implemented in GraphPad prism v10.3.0 (*** p ≤ 0.001, ** p ≤ 0.005, * p ≤ 0.01). (**D**) Quantification of cellular Pi levels. For each genotype, 8 individual plants were measured using 3 technical replicates. The estimated Pi concentration was normalized by fresh weight. (**E**) Western blot of Flag-tagged AtVIH2. The theoretical molecular mass of AtVIH2 is ~118 kDa (indicated by a black arrow; 1:Col-0, 2:AtVIH2^{WT},3:AtVIH2$^{K219A\text{-}D292A}$, 4:AtVIH2$^{R372A\text{-}H373A}$, 5:AtVIH2$^{R372A\text{-}H373A\text{-}R376A}$, 6:AtVIH2$^{R372A\text{-}H373A\text{-}R376A\text{-}E882A}$, 7:AtVIH2^{H741A}, 8:AtVIH2^{C696A}, 9:AtVIH2$^{H505A\text{-}H741A\text{-}H975A}$, 10:AtVIH2$^{P378V\text{-}G500V\text{-}G501A}$, 11:AtVIH2$^{R722S\text{-}D724A\text{-}T726A}$, 12:AtVIH2$^{K383A\text{-}K447A\text{-}K598A\text{-}R722A}$, 13:AtVIH2$^{K379A\text{-}K381A\text{-}K498A}$.

Figure 10. Arrangement of the the kinase and phosphatase domains in a full-length PPIP5K.
(**A**) Coomassie blue-stained SDS-PAGE of purified SpAsp1. (**B**) A representative micrograph of the negatively stained SpAsp1 used for image processing. (**C**) Low resolution envelope of SpAsp1 determined by negative stain electron microscopy (left). Individual crystal structures of ScVip1KD (orange ribbon diagram) and ScVip1$^{PD\ \Delta848\text{-}918}$ (blue ribbon diagram) were docked into the L-shaped negative stain map of SpAsp1 (right). The linkers truncated in the crystal structures of ScVip1 are potentially visible in the SpAsp1 envelope (black dashed lines). (**D**) Initial velocities of 1,5-InsP$_8$ hydrolysis catalyzed by ScVip1^{PD} or ScVip1$^{KD\text{-}PD}$, measured by NMR-based phosphatase assays. (**E**) Hydrogen-deuterium exchange data mapped on the ScVip1$^{KD\text{-}PD}$ model. The protected regions in ScVip1 after incubation with non-hydrolysable 1,5-InsP$_8$ (PCP-IP$_8$) are shown in light green, the protected regions after incubation with non-hydrolysable 1,5-InsP$_8$ (PCP-IP$_8$) and ATP (AMPPNP) analogs, respectively are shown in dark green.

T.E: total extract; I: insoluble; S: soluble
Anti-his$_6$-peroxidase from mouse IgG$_1$

Figure EV1 Expression of AtVIH1[PD], AtVIH2[PD] and MpVIP1[PD] phosphatase domains in baculovirus-infected Tnao insect cells.
Western blot analysis showing the expression level and solubility of different PPIP5K phosphatase domain (PD) constructs from different plant species. The constructs analyzed are *Arabidopsis thaliana* (*At)* VIH1[PD] (residues 351-1010, Uniprot ID Q84WW3), VIH2[PD] (residues 351-1046, Uniprot ID F4J8C6), and *Marchantia polymorpha* (*Mp)* VIP1[PD] (residues 382-1041 Uniprot ID Mp8g06840).

Figure EV2 Multiple sequence alignment of PPIP5K phosphatase domains.
Multiple-sequence alignment of S. *cerevisiae* Vip1 (Uniprot ID Q06685) with *Schizosaccharomyces pombe* Asp1 (Uniprot ID O74429), *Arabidopsis thaliana* VIH1 and VIH2 (Uniprot ID Q84WW3 and F4J8C6, respectively), *Marchantia polyorpha* VIP1 (Uniprot ID Mp8g06840), and *Homo sapiens* PPIP5K1 and PPIP5K2 (Q6PFW1 and O43314, respectively). Secondary structure elements are colored according to different domains as indicated on top. Functionally important conserved residues are highlighted. Dashed lines indicate regions of the protein truncated in the engineered constructs of ScVip1 kinase domain (residues 186-522) and ScVip1 phosphatase domain (residues 536-1107, Δ848-918) used for crystallization.

Figure EV3 ScVip1 harbors a structurally conserved PPIP5K kinase domain.
Structural comparison of different PPIP5K kinase domain structures. Shown are C_α traces of the *Saccharomyces cerevisiae* Vip1 kinase domain (in orange, bound to ADP in bonds representation), *Homo sapiens* PPIP5K2 (PDB id: 3T9E, in green and bound to ADP and 5-InsP$_7$) (Wang *et al*, 2012) and *Schizosaccharomyces pombe* Asp1 (PDB id: 8E1H, in purple) (Benjamin *et al*, 2022b). The kinase structures align with a r.m.s.d of ~1.5 Å comparing 310 C_α atoms.

A

PDB id	Z	Description
1ND6	21.5	Prostatic acid phosphatase (*H. sa*)
3IT2	21.4	Histidine acid phosphatase (*F. tu*)
…	…	…
1NT4	20.2	Periplasmic glucose-1 phosphatase (*E. co*)
1DKL	20.0	Phytase (*E. co*)
…	…	…
4JOC	18.8	Lysophosphatidic acid phosphatase type 6 (*H. sa*)
5CX4	18.8	Inositol polyphosphate phosphatase (*B. th*)
…	…	…
3EOZ	12.8	Phosphoglycerate mutase (*P. fa*)
3DCY	12.5	TIGAR (TP53 induced glycolysis and apoptosis regulator) (*H. sa*)
1UJC	12.4	SixA histidine phosphatase (*E. co*)
5XD2	12.3	Nudix MutT1 (*M. sm*)
3E9E	12.3	TIGAR fructose-2,6- and fructose-1,6-bisphosphatase (*D. re*)
3O0T	12.1	Ser/Thr phosphatase PGAM5 (phosphoglycerate mutase 5) (*H. sa*)
5UM0	11.9	2,3-bisphosphoglycerate-dependent phosphoglycerate mutase (*N. go*)
…	…	…
5ZR2	11.4	Phosphoserine phosphatase (*E. hi*)
1EBB	11.3	Cofactor-dependent phosphoglycerate mutase homolog PhoE (or YhfR) (*B. st*)
…	…	…
6NRU	10.9	Alpha-ribazole phosphatase (*S. fl*)
…	…	…
3D6A	9.9	2H-phosphatase domain of Sts-2 (*M. mu*)
…	…	…
1K6M	8.2	Liver PFKFB (6-phosphofructo-2-kinase/Fructose-2,6-bisphosphatase) (*H. sa*)
…	…	…
3PGM	7.6	Phosphoglycerate mutase (*S.ce*)
…	…	…

α-helical insert

Histidine phosphatase fold

B

PDB id	Z	Description
6FWS	5.9	Arch domain of the ATP-dependent DNA helicase DinG (*E. co*)
7zsa	5.8	RNA polymerase II subunit RPB1 (*S. ce*)
7O4K	5.5	Arch domain of the transcription factor IIH helicase subunit (*S. ce*)
…	…	…
3R5A	4.6	Tetrahydrodipicolinate N-succinyletransferase (*P. ae*)
…	…	…
5N0L	4.3	GAF domain from the nutrient sensor CodY (*C. di*)
3IBJ	4.3	GAF domain from the phosphodiesterase 2A (*H. sa*)
4JGO	4.2	PAS-like domain from the sporulation kinase D (*B. su*)
3A1J	4.2	Rad9 subunit from the Rad9-Hus1-Rad1 DNA sliding clamp (*H. sa*)
…	…	…
3EEH	4.0	Putative light and redox sensing histidine kinase (*H. ma*)
7JTK	4.0	GAF domain from RSP2 subunit of the flagellar radial spoke 1 (*C. re*)
…	…	…
8W7P	3.9	PAS domain from a sensor histidine kinase (*P. sp. FPU-7*)
…	…	…
5IJJ	3.8	SPX domain from a glycerophosphodiester phosphodiesterase 1 (*T. th*)
…	…	…
4MCW	3.8	GAF domain from a cyclic-di-GMP phosphodiesterase (*P. ma. EX-H1*)
…	…	…
5W10	3.8	GAF domain from a cAMP-dependent diguanylate cyclase (*L. in*)
…	…	…
6HMJ	3.6	PAS domain from a photoreceptor PAL (*N. mu. DSM 44233*)
…	…	…
8UFI	3.6	GAF domain from a phosphodiesterase 6 (*B. ta*)
8HXX	3.6	Transcriptional regulatory protein SIN3 (*S. ce*)
…	…	…

GAF-like domain

Figure EV4 ScVip1^PD is structurally related to histidine acid phosphatases and contains a GAF domain.
Top twenty hits from the DALI web server (http://ekhidna2.biocenter.helsinki.fi/dali/) (Holm & Rosenström, 2010) using the histidine acid phosphatase core containing the α-helical insertion domain (residues 536-553; 647-1094) (**A**), or the isolated GAF domain in ScVip1^PD (residues 554-646) (**B**) as search model against the Protein Data Bank.

Figure EV5 NMR-based enzyme assays – raw data.

NMR time course experiments of PP-InsP hydrolysis catalyzed by ScVip1-PD and respective mutant constructs using 100µM of [$^{13}C_6$]-labeled PP-InsP substrates. Concentrations of enzyme were varied based on the respective construct and substrate. PP-InsP turnover was measured using a pseudo-2D spin-echo difference nuclear magnetic resonance spectroscopy experiment. Quantification was based on the relative intensities of the C2 signals of the respective (PP-)InsP species.

Figure EV6 Arrangement of GAF domains in the ScVip1[PD] and ScVip1[PDΔ848-918] apo structures.

(**A**) Structural superposition of chains A (shown as C_α traces) from the ScVip1[PD] (in blue) and ScVip1[PDΔ848-918] (in cyan) apo crystal forms reveal the GAF domain in an open conformation (r.m.s.d is ~0.5 Å comparing 457 corresponding C_α atoms). The asymmetric unit of ScVip1[PD] apo contains four molecules, all of which present the GAF domain in the open conformation. (**B**) ScVip1[PDΔ848-918] crystals also contain four molecules in the asymmetric unit with chains A and B in the open and chains C and D in the closed conformation, respectively. Shown are ribbon diagrams, colors as in Fig 1B. (**C**) A second Zn^{2+} binding site in ScVip1[PDΔ848-918] is formed along a pseudo 2-fold axis involved two neighboring GAF domains. This second binding site likely represent a crystallization artifact. The crystallization condition contained an excess of $(CH_3CO_2)_2Zn$ and the residues involved in Zn^{2+} binding are not conserved among other PPIP5Ks (His573, Glu575, compare Fig EV2).

Figure EV7 Structural integrity profiling of *Sc*Vip1^{PD} mutant proteins.

(**A**) Coomassie blue-stained SDS-PAGE analysis of wild-type and mutant versions of ScVip1^{PD} (theoretical molecular weight is ~65.5 kDa). (**B**) Analytical size-exclusion chromatography of wild-type and mutant versions of ScVip1PD. Shown are A$_{280nm}$ absorption traces as a function of column elution volume (in mL). The red arrow indicates the characteristic elution volume of ScVip1PD, which behaves as a monomer in solution. (**C**) Melting temperatures of wild-type (blue traces, N=5) and mutant versions of ScVip1^{PD} derived from thermal shift assays (N.D. not determined). Note that addition of divalent metal chelators such as EDTA reduces the thermal stability of the enzyme. Addition of ZnCl$_2$ but not MgCl$_2$ increases ScVip1^{PD} stability, likely by targeting the structural Zn^{2+} binding site.

Figure EV8 An intermolecular disulfide bridge stabilizes the crystallographic dimer in the *Sc*Vip1^{PD Δ848-918 RHR-AAA} – 1,5-InsP₈ complex structure.

Ribbon diagram of a crystallographic domain-swapped dimer in the asymmetric unit of our ScVip1^{PD Δ848-918 RHR-AAA} crystals (chain A is shown in blue, chain B in green, 1,5-InsP$_8$ molecules are depicted in bonds representation). The inlet provides a detailed view of the Zn^{2+} binding site, with the Zn^{2+} ions shown as a sphere (in gray) and the coordinating residues depicted in bonds representation. The view includes a 2 $(F_o − F_c)$ electron density map contoured at 1 σ (blue mesh). Cys793 is involved in an intermolecular disulfide bridge between ScVip1^{PD Δ848-918 RHR-AAA} chain A (in blue) and chain B (in green). In the crystal structure of the apo ScVip1^{PD Δ848-918} (compare Fig1C), Cys793 replaces Glu796 and coordinates the Zn^{2+} ion.

Description	ScVip1[KD-PD]	ScVip1[KD-PD] + 1,5-InsP8	ScVip1[KD-PD] + AMPPNP + 1,5-InsP8
Reaction volume	50 µl	50 µl	50 µl
% D_2O in the reaction	77,0%	77,0%	77,0%
Temperature	22 °C	22 °C	22 °C
D_2O incubation times (sec)	30 ; 300	30 ; 300	30 ; 300
Control sample	Non-deuterated (ND) ScVip1		
Quench buffer (20 µl)	4M Gdn-HCl, 1M NaCl, 0.1M NaH_2PO_4 (pH 2.5), 1% Formic Acid (FA)		
Number of peptides anayzed	121	121	121
Sequence coverage	97,0%	97,0%	97,0%
Replicates	3 ; 3	3 ; 3	2 ; 2
Criteria for HDX rate difference	Difference of HDX level at a given timepoint is > 7 % and > 0.5 Da and p values of student t-test is < 0.02		
Protein amount (nmol)	0,107	0,107	0,107
Ligand amount (nmol)	-	10,71	10.71 + 10.71

EV9 Summary of the HDX-MS experiments

(**A**) Experimental details of the HDX analyses. (**B**) D_2O incorporation into ScVip1[KD-PD]. (**C**) HDX differences comparing apo ScVip1[KD-PD], ScVip1[KD-PD] in complex with 1,5-InsP8 and ScVip1[KD-PD] in the presence of 1,5-InsP8 and a non-hydrolyzable ATP analog (blue, protection; red, deprotection).

Figure EV10 The "HD" catalytic motif presents in phytases is not present in ScVip1[PD].

Detailed views of the active sites of *E. coli* phytase AppA (in orange, PDP-ID 7Z2S) (Acquistapace *et al*, 2022) bounds to the substrate analogue InsS$_6$, *S. cerevisiae* Vip1[PD] [Δ848-918], and the Alphafold model of *S. pombe* Asp1[PD] (in purple, https://alphafold.ebi.ac.uk/entry/O74429) (Varadi *et al*, 2024). The InsS$_6$ and 1,5-InsP$_8$ molecules are depicted in bonds representation, and the side chains of residues belonging to the catalytic motifs 1 and 2 are depicted bonds representation. Residues His303 and Asp304 forming the HD motif in AppA correspond to Lys990 and Glu991 in ScVip1[PD] and not to His993 and Ile994, as previously thought (Fridy *et al*, 2007; Mulugu *et al*, 2007; Pöhlmann *et al*, 2014; Wang *et al*, 2015a; Pascual-Ortiz *et al*, 2018).