Using correlative and mechanistic species distribution models to predict vector-borne disease risk for the current and future environmental and climatic change: a case study of West Nile Virus in the UK.

Withers, A. J.[1], Croft, S.[1], Budgey, R.[1], Warren, D.[1] and Johnson, N.[2]

[1] National Wildlife Management Centre, Animal and Plant Health Agency, Sand Hutton, YO41 1LZ

[2] Virology Department, Animal and Plant Health Agency, Surrey, KT15 3NB

- **corresponding author:** amy.parkes-withers@apha.gov.uk

- **data availability statement:** All data underlying this work are held in publicly accessible repositories and can be downloaded for use in research under licence. Outputs can be made available under license by contacting the corresponding author.
- **funding statement:** This work is funded by EXSE0566: Development of improved tools and approaches for control of zoonotic arthropod-borne diseases of animals and their vectors Funded by the Department for Environment, Food and Rural Affairs and the Scottish and Welsh Governments and was co-funded by the European Union's Horizon Europe Project 101136346 EUPAHW.
- **conflict of interest disclosure:** The authors declare no conflicts of interest.
- **ethics approval statement:** n/a
- **patient consent statement:** n/a
- **permission to reproduce material from other sources:** n/a
- **clinical trial registration:** n/a

Abstract

Globally, vector-borne diseases have significant impacts on both animal and human health, and these are predicted to increase with the effects of climate change. Understanding the drivers of such diseases can help inform surveillance and control measures to minimise risks both now and in the future. In this study, we illustrate a generalised approach for assessing disease risk combining species distribution models of vector and wildlife hosts with data on livestock and human populations using the potential emergence of West Nile Virus (WNV) in the UK as a case study. Currently absent in the UK, WNV is an orthoflavivirus with a natural transmission cycle between *Culex* mosquitos (*Cx. pipiens* and *Cx. modestus*) and birds. It can spread into non-target hosts (e.g., equids, humans) via mosquito bites where it can cause febrile disease with encephalitis and mortality in severe cases. We compared six correlative species distribution models and selected the most appropriate for each vector based on a selection of performance measures and compared this to mechanistic species distribution models and known distributions. We then combined these with correlative species distribution models of representative avian hosts, equines, and human population data to predict risk of WNV occurrence. Our findings highlighted areas at greater risk of WNV due to higher habitat suitability for both avian hosts and vectors, and considered how this risk could change by 2100 under a best-case Shared Socioeconomic Pathway (SSP1) and worst-case (SSP5) future climate scenario. Generally, WNV risk in the future was found to increase in south-eastern UK and decrease further north. Overall, this paper presents how current and future vector distributions can be modelled and combined with projected host distributions to predict areas at greater risk of novel diseases. This is important for policy decision making and contingency preparedness to enable adaptation to changing environments and the resulting shifts in vector-borne diseases that are predicted to occur.

Introduction

Vector-borne diseases have a significant impact upon global animal and human health, accounting for around 17% of all infectious diseases in humans, leading to over 700,000 deaths a year worldwide (World Health Organisation, 2020). Additionally, impacts on the livestock industry can result in devastating economic losses, as well as contributing to food shortages due to limited productivity (Faburay, 2015). Most insect vector species, including mosquitos, ticks, and biting midges, are exotherms meaning that their lifecycle and geographic ranges are highly influenced by the climate and environment. Consequently, the risk from the diseases they carry is expected to be significantly affected by future climate change and urbanisation, with range shifts predicted to occur globally (Endo and Amarasekare, 2022, Rocklöv and Dubrow, 2020, Noll et al., 2023). Understanding these potential range shifts is important for surveillance so mitigation measures can be developed to reduce impact on both human and animal health. For the UK, the risk of vector-borne disease is predicted to increase with the potential for increased globalisation and warming temperatures leading to a higher number of invasive mosquito species introductions and favourable conditions that could support year-round vector survival, leading to outbreaks of diseases such as dengue, chikungunya, and West Nile fever (Medlock and Leach, 2015, Townroe and Callaghan, 2014).

West Nile Virus (WNV) is an orthoflavivirus with a natural transmission cycle between mosquitos and birds. On rare occasions, mosquitos will spread WNV into non-target, dead-end hosts, such as humans and equids. Whilst WNV is usually asymptomatic in these non-target hosts, it can become symptomatic (e.g., headache, fever, rash) in around 20% of cases, with 1 in 150 human cases becoming neuroinvasive (e.g., acute aseptic meningitis, encephalitis) which can occasionally lead to severe illness or death (Pradier et al., 2012, World Health Organisation, 2017). In recent years, outbreaks of WNV have occurred in Europe in areas where the disease has not been present historically (e.g., France, Germany), and predictions suggest WNV could reach the UK (Vogels et al., 2017, Constant et al., 2022, Bessell et al., 2016). The most likely route of WNV introduction to the UK is via migrating birds from endemic regions of mainland Europe (Bessell et al., 2016). Some birds migrate to the UK from areas in Africa where WNV is present, however, the time taken for this migration to be completed means that viremia will have declined and transmission will no longer be possible when the birds arrive at their final destination (Pradier et al., 2012).

As well as significant effects on human and animal health, WNV can have severe economic impacts. For example, predictions of the economic impact in Belgium were around €4,500 per human case that led to hospitalisation which could be thousands during an outbreak (Humblet et al., 2016). As well as additional costs to the equestrian sector of around €17 million to vaccinate horses

in at-risk areas to prevent further economic damage (Humblet et al., 2016). Therefore,

understanding the distribution of WNV vectors and hosts in the UK can reveal where the risk is

higher, this is important for contingency planning and mitigation to reduce the impact of an

outbreak by enabling measures to be put in place to protect human and animal health, and

consequently reduce economic stress (Medlock and Leach, 2015, Medlock et al., 2018, Gandy et al.,

2022, Lawson et al., 2022).

There are two primary vectors of WNV in the UK, these are *Culex pipiens* f. *pipiens* (hereafter *Cx.*

pipiens) and *Culex modestus*. *Cx. pipiens* has a widespread distribution, and mostly feeds on birds

but has been known to feed on mammals so could potentially spread WNV into non-target hosts

(e.g., humans, equids). C*x. modestus* is an invasive species in the UK that was first discovered in the

south of England (Portsmouth) in the 1940's, then rediscovered in the north Kent marshes in

southeast England in 2003 where it had established (Golding et al., 2012, Vaux et al., 2015). Since

then, multiple other records have occurred across the UK, but the distribution of *Cx. modestus* is

currently much more limited than that of *Cx. pipiens*. *Cx. modestus* is of interest as a key bridge

vector for WNV transmission and host spillover as it is known to bite both mammals and birds,

especially in areas where frequent overlap of these potential blood meals occurs (Rudolf et al., 2021,

Hernández-Triana et al., 2020, Soto and Delang, 2023).

When modelling the potential range of insect distributions there are two main approaches;

correlative and mechanistic (Koch, 2021). Correlative species distribution models have been widely

applied to predict distributions across all taxa by relating occurrence data to a set of environmental

variables to determine suitability and then using the suitability model to predict species distributions

across the whole study area (Koch, 2021, Elith et al., 2010, Hijmans and Elith, 2013, Valavi et al.,

2021). There are a broad range of correlative species distribution models available, such as Random

Forest (RF), Boosted Regression Tree (BRT), Support Vector Machine (SVM), Neural Network (NET)

and MaxEnt (MAX). Each of these approaches have previously been used to produce species

distribution models for a large range of taxa, including mosquitos, and each can outperform the

other depending on the species and presence or absence data available (European Centre for

Disease Prevention and Control, 2019, Beeman et al., 2021, Simons et al., 2019, Furlong et al., 2022,

Senay et al., 2013, Senay and Worner, 2019, Richman et al., 2018, Gorris et al., 2021). It is rarely

clear which individual model is the most appropriate. Ensemble approaches have been used to

remedy this issue and can be beneficial, allowing multiple outputs to be combined into a single

consensus model. However, determining how to combine different models is difficult and many

approaches have been suggested (Simons et al., 2019, European Centre for Disease Prevention and

Control, 2019, Beeman et al., 2021).

The correlative approaches discussed above each require presence-absence data (i.e., known absences) or presence-background (i.e., characterise where a species is less likely to be found based on known presences). Presence-absence data is often unavailable for species distribution modelling as this requires a consistent sampling effort over a widespread area to ensure that absences are true absences and not due to sampling bias (Elith et al., 2006, VanDerWal et al., 2009, Lobo et al., 2010, Iturbide et al., 2015, Croft and Smith, 2019). Therefore, presence-background is the most common approach for species distribution modelling, however, this requires artificially generating pseudo-absence or background points to represent areas where the species is absent. It can be difficult to determine background data for species with limited presence records, especially if it is an invasive species as they may not have reached all suitable locations within the area yet (Elith et al., 2010, Giovanelli et al., 2010). Whilst this is true of many species it requires extra consideration for insects as host and range expansions can occur rapidly in some species due to short life-cycles and multiple generations each year, meaning that the predicted distribution of insects may be underestimated if background points are created in areas where survival is possible (Koch, 2021). Conversely, mechanistic models do not rely on sampling data as they are based on *a priori* knowledge of a species and its response to climatic variables, including factors such as seasonal population growth, environmental stress, and survival constraints (Koch, 2021, Sutherst and Maywald, 1985, Kriticos et al., 2015). This means that they are usually limited to plants and insects where dependencies are more climatic so laboratory studies to measure tolerances (e.g., temperature, drought) can inform the models (Sutherst and Maywald, 1985, Poutsma et al., 2008, Taylor and Kumar, 2012, Khormi and Kumar, 2014, Kriticos et al., 2015, Early et al., 2022). A common mechanistic modelling approach is CLIMEX, which predicts a species' potential range based on biological constraints (Sutherland et al., 2021, Kriticos et al., 2015). Previously, CLIMEX modelling has successfully been used to determine potential distributions of invasive mosquitos including *Aedes aegypti* and *Aedes atropalpus* (Khormi and Kumar, 2014, Scholte et al., 2009), and it can be beneficial for species with limited presence-absence data as it does not rely on sampling data (Sutherst and Maywald, 1985, Taylor and Kumar, 2012, Khormi and Kumar, 2014, Kriticos et al., 2015, Early et al., 2022).

Here, an approach of producing disease risk maps based on species distribution modelling of vector and host habitat suitability is presented, using WNV emergence in the UK as a case study. Correlative species distribution models are used to predict the range of each vector (*Cx. pipiens* and *Cx. modestus*), and a mechanistic species distribution model was also produced for each vector to further validate the correlative approaches. These models were then used to predict distributions across the UK for the current climate as well as future predictions to the year 2100 for two shared socioeconomic pathways; SSP1 (climate protection measures and development compatible with

target of 2°C) and SSP5 (worst case scenario without climate measures and mitigations) (Fick and

Hijmans, 2017). These individual vector maps were then combined to create an overall vector risk

map for all climate scenarios. Following this, correlative species distribution models were produced

of representative avian hosts for corvid and non-corvid Passeriformes. Once all current and future

suitability predictions had been produced for vectors and hosts, these were then combined with

horse or human density data to produce overall WNV risk maps for the current climate (based on

horse or human density) and future environment (based on human density).

Methods

Model extent

158   Both vector and avian host models were conducted at a global extent, with a spatial-

159   resolution of 30 arc-seconds (0.00833º), which equates to about 1km^2 at the equator. The models

160   were then used to predict distribution across the whole UK at the same resolution 0.00833º.

Presence and absence data

Vector presence

Global species presence data for all confirmed records of *Culex pipiens* and *Culex modestus*

were downloaded from the National Biodiversity Network (NBN) Atlas and the Global Biodiversity

Information Facility (GBIF) for all time up to the download date of 7[th] November 2022 (NBN Atlas,

2022, GBIF Org User, 2022). An additional presence record for *Cx. modestus* was included from a

recent paper reporting the first recording of the species in Finland (Culverwell and Vapalahti, 2023).

For each species, all presences were combined and cleaned in R (v. 4.3.3) to remove duplicated

records and invalid points (e.g., over the sea) (CoordinateCleaner::*clean_coordinates*, v3.0.1, Zizka et

al. (2019)). Presence points were then thinned based on environmental conditions prior to modelling

to further reduce sampling bias (flexsdm::*occ*_filt, v1.3.4, Velazco et al. (2022)) (Castellanos et al.,

2019).

Host presence

A wide variety of avian hosts can be susceptible to and spread WNV, however, the Passerine

order are generally considered the most susceptible (Ziegler et al., 2022, Montecino-Latorre and

Barker, 2018, Durand et al., 2010, Bergmann et al., 2023, Komar et al., 2003, Centers for Disease

Control and Prevention, 2021). Here, we considered common species susceptible to WNV frequently

observed around areas that may be suitable for mosquito vectors and humans or horses to act as

representative avian hosts (Brugman et al., 2017). Selected avian host species were aggregated into

two groups, corvids consisting of the common raven (*Corvus corax*), rook (*Corvus frugilegus*) and

carrion crow (*Corvus corone*) and non-corvids comprising of the house sparrow (*Passer domesticus*), blackbird (*Turdus merula*) and barn swallow (*Hirundo rustica*).

Bird presence records for our selected species were taken from GBIF for April to October from 2013 to 2023 on 4th June 2024 for all six avian host species included in these models (GBIF. Org User, 2024, GBIF.Org User, 2024d, GBIF.Org User, 2024e, GBIF.Org User, 2024b, GBIF.Org User, 2024a, GBIF.Org User, 2024c). GBIF was used for presences as it includes eBird observations, alongside many other submitters, thereby maximising the number of presence records available. These presences were cleaned using clean_coordinates in R (v.4.4.0) to remove duplicated records and invalid points (e.g., over the sea) (CoordinateCleaner::*clean_coordinates*, v3.0.1, Zizka et al. (2019)). Presence points were thinned based on environmental conditions prior to modelling to further reduce sampling bias (flexsdm::*occ_*filt, v1.3.4, Velazco et al. (2022)) (Castellanos et al., 2019).

Predictor variable selection

Twenty-one environmental predictor variables were selected for vectors and hosts for the current and future environmental conditions based on two Shared Socioeconomic Pathways (SSP), SSP126 and SSP585 (Table 1). For vectors and hosts a global extent was used, with a resolution of 0.00833, 0.00833 (i.e., about $1km^2$). To reduce multicollinearity in models Pearson correlation coefficient was used to remove climatic variables with a >0.7 correlation coefficient irrespective of presence or absence points (i.e., correlation across all predictors) to minimise the effects of spatial autocorrelation (*vifcor,* usdm, (Naimi et al., 2014)), and a dendrogram of clusters of correlates is available in Supplementary Information (SI) 1. Following this, the remaining variables included were bio2 (mean diurnal range), bio8 (mean temperature of the wettest quarter), bio9 (mean temperature of the driest quarter), bio14 (precipitation of the driest month), bio15 (precipitation seasonality), bio18 (precipitation of the warmest quarter), bio19 (precipitation of the coldest quarter), land use and human population density.

206 Table 1: The predictor variables used in modelling vectors and hosts. Each predictor variable was
207 downloaded for the current time and for two Shared-Socioeconomic Pathways (SSP) in the future
208 (up to 2100), these were SSP1 and SSP5. Those variables in italics were maintained in the model
209 following variable selection based on correlation.

Variable	Resolution	Source
Bioclimatic variables based on UK-ESM1-0-LL: • BIO1 = Annual Mean Temperature • *BIO2 = Mean Diurnal Range* • BIO3 = Isothermality • BIO4 = Temperature Seasonality • BIO5 = Max Temperature of Warmest Month • BIO6 = Min Temperature of Coldest Month • BIO7 = Temperature Annual Range • *BIO8 = Mean Temperature of Wettest Quarter* • *BIO9 = Mean Temperature of Driest Quarter* • BIO10 = Mean Temperature of Warmest Quarter • BIO11 = Mean Temperature of Coldest Quarter • BIO12 = Annual Precipitation • BIO13 = Precipitation of Wettest Month • *BIO14 = Precipitation of Driest Month* • *BIO15 = Precipitation Seasonality* • BIO16 = Precipitation of Wettest Quarter • BIO17 = Precipitation of Driest Quarter • *BIO18 = Precipitation of Warmest Quarter* • *BIO19 = Precipitation of Coldest Quarter*	0.00833, 0.00833 (approximately 1km^2)	(Fick and Hijmans, 2017)
Land use (7 land types): Forest, Grassland, Barren, Cropland, Urban, Water, Permanent snow and ice	1km^2 (re-projected for modelling to 0.00833, 0.00833)	(Chen et al., 2022)
Human population density	1km^2 (re-projected for modelling to 0.00833, 0.00833)	(Gao, 2020, Gao, 2017)

Modelling approaches

For vectors, seven different modelling approaches were used, six of these were correlative

approaches and one was a mechanistic approach used to determine suitability based on the

biological constraints of *Cx. pipiens* and *Cx. modestus*. For hosts, four correlative approaches were

used as CLIMEX is most appropriate for species strongly limited by climate due to factors such as

temperature and moisture, whereas avian distribution is influenced strongly by additional factors.

Mechanistic modelling was carried out using CLIMEX (version 4, Kriticos et al. (2015)). All correlative

species distribution modelling and the production of figures was carried out in R (v 4.4.0).

 Correlative species distribution models for vectors and hosts

 Vectors

Based on the presence data available following filtering, a model calibration area was

determined based on a 100km buffer around presence points for *Cx. pipiens* and *Cx. modestus*, this

radius was selected following trials of models using different buffers (5km, 50km, and 100km) and

comparing performance across several different performance measures (full details are provided in

Supplementary Information (SI) 2). Within this calibration area we randomly allocate 4716 and 144

background points (dismo::*randomPoints,* v.1.3-14, Hijmans et al. (2023)) for *Cx. pipiens* and *Cx.*

modestus respectively to act as pseudo-absences. This method has the fewest assumptions about

the presence data. The number of background points was selected as three-times the number of

presences available as large numbers of absences have been shown to not be beneficial for species

with small presence datasets, with fewer absences leading to increased accuracy in machine learning

models (Liu et al., 2019, Barbet-Massin et al., 2012, Thuiller et al., 2024).

Prior to model fitting, the presence and background points datasets were partitioned into two

datasets, one for training and one for testing. For vectors, this was based on the spatial

autocorrelation of points based on Moran's I, environmental similarity, and the number of points

available to ensure a balanced division of points between the training and testing datasets

(flexsdm::*part_sband,* v1.3.4, Velazco et al. (2022)) (Roberts et al., 2017, Santini et al., 2021, Velazco

et al., 2022). Using a structured approach to partition a large database into training and testing

datasets can help to overcome the sampling and spatial bias that can later affect performance

measures and the evaluation of models in partitioning is carried out randomly (Valavi et al., 2022,

Elith et al., 2006, Valavi et al., 2021, Roberts et al., 2017, Elith et al., 2010). Following data

partitioning based on spatial autocorrelation, 1319 presence and 3064 background points were in

the training dataset and 253 presence and 1531 background points were in the testing datasets for

Culex pipiens. For *Cx. modestus,* 27 presence and 72 background points were in the training dataset

and 21 presence and 70 background points were in the testing datasets.

The correlative modelling approaches used were Random Forest (RF, flexsdm::*tune_raf, v1.3.4,*

Velazco et al. (2022)), Maxent (MAX, flexsdm::*tune_max, v1.3.4, Velazco et al. (2022)*), Generalized

Boosted Regression (GBM, flexsdm::*tune_gbm, v1.3.4, Velazco et al. (2022)*), Neural Network (NET,

flexsdm::*tune_net, v1.3.4, Velazco et al. (2022)*), Support Vector Machine (SVM, flexsdm::*tune_svm,*

v1.3.4, Velazco et al. (2022)) and an Ensemble model based on all five individual models

(flexsdm::fit_ensemble*, v1.3.4, Velazco et al. (2022)).*

All individual correlative models were fit using the *tune* functions in the *flexsdm* package, with

selections based on three different thresholds, which were when sensitivity and specificity were

equal (*equal_sens_spec*), when the sum of sensitivity and specificity was highest (*max_sens_spec*)

and when the Sorensen threshold was highest (*max_sorensen*). A number of performance metrics

were calculated during model fitting and considered for model evaluation, as well as visual

comparison to known presence locations (European Centre for Disease Prevention and Control and

European Food Safety Authority, 2022, GBIF Org User, 2022). A number of measures were used as

there is often variation in how each model is scored, and determining suitability on a single measure

can lead to poor model selection due to inherent, undetectable biases, such as model overfitting and

sampling bias in the data used to build the models (Velazco et al., 2022, Valavi et al., 2022, Valavi et

al., 2021, Hijmans and Elith, 2013, Roberts et al., 2017). The performance metrics used were True

positive rate (TPR), True negative rate (TNR), Sorensen, Jaccard, F-measure of presence-background

(FPB), Omission rate (OR), True skill statistic (TSS), Area under curve (AUC), Boyce index and Inverse

mean absolute error (IMAE) (Velazco et al., 2022, Valavi et al., 2022, Valavi et al., 2021, Hijmans and

Elith, 2013, Roberts et al., 2017). The TPR and TNR are the proportion of observed presences (TPR)

or absences (TNR) in the test data that are correctly predicted by the model, and are sometimes

referred to as sensitivity and specificity respectively. Several performance metrics are based on the

TPR and TNR alongside the falsely classified presences (False Positive Rate; FPR) and absences (False

Negative Rate; FNR) that give an indication of model performance, including the Sorensen and

Jaccard similarity indices, FPB, OR and TSS. The Sorensen and Jaccard similarity indices are scored on

a scale of 0 to 1, where 0 is a very poor similarity and 1 high similarity and thus indicates a better

performing model. They are calculated very similarly, however, Sorensen assigns more importance

to true positives by weighting them higher than true negatives in the index. The FPB is on a scale of 0

to 2, where a higher score is indicative of better model performance based on combining the TPR,

TNR, FPR and FNR. The OR is the proportion of presences correctly assigned by the model. The TSS is

calculated by combining the TPR and TNR to give an overall indication of performance, and is also on

a scale of 0 to 1 where the closer to 1 the score is, the better the model is. The AUC is widely used in

species distribution modelling to compare the TPR and FPR to assess model performance on a scale

of 0 to 1, where 0.5 suggests the model is no better than random and 0.75 is generally considered

good. The Boyce index measures how predictions made using the test dataset differ from a random

distribution of the observed presences, and is scored on a scale of -1 to 1, where 1 indicates a good

model and 0 suggests the model is no different to random. Negative values suggest that the

predictions are frequently incorrect (i.e., presences located in areas predicted to be unsuitable). The

IMAE considers the total number of correctly assigned presences and absences in relation to the

total number of observations, and it is also given on a scale of 0 to 1, where 1 reflects a model

performing well with no errors. Full details on how these performance measures are calculated are

available in the *flexsdm* R package (Velazco et al., 2022).

Hosts

Using the presence data available following filtering for each individual species, a model

calibration area was determined based on a convex polygon around presence points to represent

the accessible area as this has previously been shown to improve model performance and worked

well for other bird species distribution models (Rojas-Soto et al., 2024, Palacio et al., 2021, Rousseau

and Betts, 2022). Background points were randomly allocated within the defined model calibration

area (dismo::*randomPoints,* v.1.3-14, Hijmans et al. (2023)) for each species to act as pseudo-

absences as this method has the fewest assumptions about the presence data. Due to the large

number of presence data available, an equal number of background points was created (Liu et al.,

2019, Barbet-Massin et al., 2012, Thuiller et al., 2024). Any datapoints outside the extent of

environmental predictors were excluded from the dataset. Prior to model fitting, the species

presence and background data were partitioned into training and testing datasets. Following the

same approach as for vectors a structured approach was used to partition data into training and

testing datasets, this was based on spatial autocorrelation of points based on Morans I,

environmental similarity, and the number of points available to ensure even division of points

between the training and testing datasets (flexsdm::*part_sband,* v1.3.4, Velazco et al. (2022)).

(Roberts et al., 2017, Santini et al., 2021, Velazco et al., 2022). Individual species data were then

combined for modelling into corvid (common raven, rook, carrion crow) or non-corvid (house

sparrow, blackbird, barn swallow) representative groups. In the corvid group there were 33,132

(22,994 and 10,138 in the training and testing datasets respectively) presences and 32,002 (19,476

and 12,526 in the training and testing datasets respectively) background points. In the non-corvid

group there were 43,085 (25,944 and 17,141 in the training and testing datasets respectively)

presences and 42,273 (22,108 and 20,165 in the training and testing datasets respectively)

background points.

Three individual correlative models were fitted using the *tune* functions in the *flexsdm*

package, with selections using a threshold when the sum of sensitivity and specificity was highest

(*max_sens_spec*). The correlative modelling approaches used were Random Forest (RF,

flexsdm::*tune_raf,* v1.3.4, Velazco et al. (2022)), Generalized Boosted Regression (GBM,

flexsdm::*tune_gbm,* v1.3.4, Velazco et al. (2022)), and Neural Network (NET, flexsdm::*tune_net,*

v1.3.4, Velazco et al. (2022)). An Ensemble model based on the previous three individual models was

also fit (*fit_ensemble, flexsdm)* (Velazco et al., 2022)*.* Several performance metrics were calculated

during model fitting and considered for model evaluation, these were True positive rate (TPR), True

negative rate (TNR), Sorensen, Jaccard, F-measure (FPB), Omission rate (OR), True skill statistic (TSS),

Area under curve (AUC), Boyce index and Inverse mean absolute error (IMAE) (Velazco et al., 2022,

Valavi et al., 2022, Valavi et al., 2021, Hijmans and Elith, 2013, Roberts et al., 2017). These

performance metrics have been described in detail in the vector section previously.

Mechanistic species distribution models for vectors

To further validate our choice of species distribution models of vectors the mechanistic

model CLIMEX was used, to ensure that areas predicted to be of higher suitability for *Culex*

mosquitos using the correlative approaches were consistent with the known eco-physiological

tolerance of *Cx. pipiens* and *Cx. modestus*. The mechanistic model was developed using the CLIMEX

Compare Locations Genetic Algorithm (single species) function (v. 4) (Sutherst and Maywald, 1985,

Kriticos et al., 2015). This was used to determine the most suitable parameter values based on the

global presence datasets available alongside the literature and expert opinion to ensure the model

was as reliable as possible (Kriticos et al., 2015, Sutherst and Maywald, 1985). Sensitivity analysis

was carried out using the Compare Locations with Sensitivity Analysis function in CLIMEX, which

showed that the most influential parameters for the ecoclimatic index calculation were soil moisture

(*limiting low moisture, lower optimal moisture*), temperature (*lower optimal temperature)* and

diapause (*diapause induction daylength, diapause termination temperature*) (Supplementary

Information (SI) 3). As insufficient data was available to accurately parametrise for soil moisture

constraints on both species the soil moisture index was removed from the model as incorrectly fitted

parameters can greatly influence the suitability indices if the model is highly sensitive to them

(Taylor and Kumar, 2012). The parameter values used in the CLIMEX model are shown in Table 2,

and parameters not shown were not included in fitting the model. CLIMEX outputs were converted

to a raster in ArcGIS Pro v. 3.0.1 (Esri, 2022) using the Point to Raster function, with the Ecoclimatic

Index (EI) as the value field, and a cell size (i.e. resolution) of 0.167 degrees (approximately 1.85 x

1.85km). Validation of CLIMEX models has previously been carried out by using a presence dataset

to determine how many known presence points fall within areas predicted to be suitable by the

CLIMEX model (Taylor and Kumar, 2012). This approach of validation was used here, with

environmental suitability categories created based on the ecoclimatic index with a score of 0 to 1

considered unsuitable, 1 to 10 as marginally suitable, 10 to 20 as favourable and >20 as very

favourable. This followed the environmental classification approach used for the CLIMEX model for

the mosquito *Aedes aegyptii* (Khormi and Kumar, 2014).

Table 2: Parameter values used for CLIMEX modelling of *Cx. pipiens* and *Cx. modestus*.

Parameter	*Cx. pipiens*		*Cx. modestus*	
	Value	Reason/Reference	Value	Reason/Reference
Temperature index				
DV0 Limiting low temperature	7	(Ewing et al., 2016)	10	Informed by Genetic Algorithm
DV1 Lower optimal temperature	15	(Ewing et al., 2016)	15	Based on *Cx. pipiens*
DV2 Upper optimal temperature	25	(Ewing et al., 2016)	25	Based on *Cx. pipiens*
DV3 Limiting high temperature	29	Informed by Genetic Algorithm	29.3	Informed by Genetic Algorithm
Diapause index				
DPD0 Diapause induction daylength	13.5	(Ewing et al., 2016)	13.5	Based on *Cx. pipiens*
DPT0 Diapause induction temperature	15	(Ewing et al., 2016)	15	Based on *Cx. pipiens*
DPT1 Diapause termination temperature	6.5	Informed by Genetic Algorithm	6.5	Based on *Cx. pipiens*
DPD Diapause development days	-100	Expert advice	-100	Based on *Cx. pipiens*
DPSW (0 for Winter, 1 for Summer)	0	Winter diapause	0	Winter diapause
Day degree accumulation above DV0				
DV0 Limiting low temperature	7	(Ewing et al., 2016)	10	Informed by Genetic Algorithm
DV3 Limiting high temperature	29	Informed by Genetic Algorithm	29.3	Informed by Genetic Algorithm
MTS Model timestep	7	Default	7	Temperate Default
Day degree accumulation above DVCS				
DVCS Cold stress threshold temperature	-15	Informed by Genetic Algorithm	-12.5	Informed by Genetic Algorithm
MTS Model timestep	7	Default	7	Default
Day degree accumulation above DVHS				
DVHS Heat stress threshold temperature	34	(Ewing et al., 2016)	27.7	Informed by Genetic Algorithm
MTS Model timestep	7	Default	7	Default
Day degrees per generation				
PDD Degree-days per generation	83	Informed by Genetic Algorithm	356	Informed by Genetic Algorithm

Model predictions

Based on the global model, predictions of species distributions for the two *Culex* species
across the UK were created (flexsdm:: *sdm_predict,* v1.3.4, Velazco et al. (2022)). For the MaxEnt
binary predictions the threshold for suitability was based on the maximum sensitivity-specificity
threshold (max_sens_spec), and for all other models continuous predictions were made.

To consider model transferability, predictions to regions outside the extent of the training
data, were assessed using Mahalanobis distance and the shape method, with distance calculations
based on the environmental distance between the training data (post-variable selection), and the
area being projected into (flexsdm::*extra_eval,* v1.3.4, Velazco et al. (2022)) (Velazco et al., 2022,
Velazco et al., 2023). The higher the Shape value, the greater the environmental distance is from the
training data to the projection area (i.e., the more novel), indicating that areas with high Shape
values have less reliable predictions than those with low Shape values (Velazco et al., 2023). Model
transferability was measured both for the current and future environmental conditions under SSP1
and SSP5 for both *Cx. pipiens* and *Cx. modestus*. To further consider model extrapolation, partial
dependence plots for the variables were created for each of the individual correlative models to
determine how each model handled predictions to novel environmental conditions within the UK
(flexsdm*::p_pdp*, v1.3.4, Velazco et al. (2022)).

Future predictions were made using the best performing model for *Cx. pipiens* and *Cx.*
modestus considering the known distributions, the mechanistic CLIMEX model and the different
performance scores. Future predictions were produced for the year 2100 based on the SSP126 and
SSP585 (flexsdm:: *sdm_predict,* v1.3.4, Velazco et al. (2022)). Combined vector risk maps were then
produced by multiplying the habitat suitability scores of cells for both *Cx. pipiens* and *Cx. modestus*
for the current and future time periods to show overall vector risk based on habitat suitability up to
the year 2100 for SSP1 and SSP5.

To produce a map of WNV risk the non-/corvid host species distribution maps were
combined into one host risk map by summing the suitability score of all cells overlayed to give an
overall host suitability score that included both representative corvid and non-corvid hosts. This host
distribution model was then combined with the vector risk map by multiplying the host and vector
maps together to produce an overall WNV risk map that considered both vectors and hosts. To
further describe WNV risk in terms of human and equine health, this process was repeated to
include layers representative of human population density both in the current environment and into
the future for SSP1 and SPP5 in the year 2100 (Gao, 2020, Gao, 2017) and the current horse
population density in the UK (Gilbert et al., 2022).

Results

Transferability of *Culex* species distribution models to areas and time-periods not represented by the presence data

Overall, the environmental conditions covered by the *Cx. pipiens* presence data represent those found across much of the UK, with only areas in northern Scotland (e.g., north-west Scotland, Shetland Islands) and very urban areas of England (e.g., London and Birmingham) showing as less represented (Figure 1a). Compared to *Cx. pipiens,* there was greater variation between the training data and predicted data for *Cx. modestus*, however many of the environmental conditions of the UK were covered by the data available, with London and north-west Scotland most poorly represented (Figure 1d). Similar patterns of environmental representation continued in future predictions for both species, with SSP5 showing more extrapolation was needed from the training data compared to SSP1 (Figure 1b-c for *Cx. pipiens* and Figure 1e-f for *Cx. modestus*). Partial dependence plots were created for each of the individual five correlative models to determine how each of the models' handled data within the novel areas for both *Cx. pipiens* and *Cx. modestus* (Supplementary Information (SI) 4).

Figure 1a-f: The Shape values for extrapolation of the study area based on the presence data available for the *Culex* species distribution models. The lower the shape value the more similar the regions are, thus the higher values represent environmental conditions poorly represented by the presence data available. Extrapolation values are shown for *Culex pipiens* for the a) current and b) future (year 2100) environment under scenario SSP1 and c) SSP5, and for *Culex modestus* for the d) current and e) future (year 2100) environment under scenario SSP1 and f) and SSP5.

Correlative species distribution models for vectors in the current climate and environmental conditions

Culex pipiens

The overall pattern of the individual *Cx. pipiens* distribution models was similar across the UK, with areas of higher suitability around the north-east Scotland, south-east, central, and north-west in England and coastal Wales (Figure 2a-f). In all models the areas with lowest suitability for *Cx. pipiens* were middle-east England and northern Scotland. However, it was the areas of lower suitability that had higher variability with GBM and SVM models predicting wider spread unsuitability compared to the RF, MaxEnt and Net models (Figure 2a-f).

Model performance can be measured in many ways and, based on the different performance metrics, the most consistently high scoring model for *Cx. pipiens* was the Ensemble model, with the Sorensen index, Jaccard, FPB, TSS and AUC all being the highest, with the remaining scores being highest for either the GBM (TNR, OR, IMAE) or RF (TPR, Boyce) models. Considering this, the *Cx. pipiens* model that performed best when predicting habitat suitability in the UK was the mean weighted Ensemble model (Table 3). This model reflects the known distribution of *Cx. pipiens* in the UK, with higher suitability in the south compared to the north (European Centre for Disease Prevention and Control and European Food Safety Authority, 2023), and that predicted by the CLIMEX model which is discussed further in the mechanistic species distribution model section.

Culex modestus

Overall, the pattern of suitability across the UK for *Cx. modestus* was relatively similar for the RF, GBM and Net models, showing predominantly unsuitable habitat, apart from areas of southern and central England, particularly in the south-east (Figure 3a-f). The MaxEnt and SVM models showed widespread suitability across the UK, though most of these areas were scored as relatively low suitability compared to the other models (Figure 3a-f).

The best performing model for *Cx. modestus* was the Net model, which scored highest in the TPR, Sorensen, Jaccard, FPB, TSS and AUC performance measures. Both the RF and Ensemble Mean Threshold models also scored highly for certain metrics (TNR/OR and Boyce/IMAE, respectively). Considering this, the Net model for *Cx. modestus* was taken to be the most reliable prediction of habitat suitability in the UK (Table 3). The pattern of suitability for the Net model aligns with the known presence data for this species, which is predominantly associated with the North Kent marshes in the south-east in the UK (Golding et al., 2012) and with that predicted by the CLIMEX model which is discussed in the mechanistic species distribution model section.

Table 3: Model performance scores and standard deviation (sd) of the Culex models. The performance metrics are True positive rate (TPR), True negative rate (TNR), Sorensen, Jaccard, F-measure (FPB), Omission rate (OR), True skill statistic (TSS), Area under curve (AUC), Boyce index and Inverse mean absolute error (IMAE). Higher values indicate better model performance.

	score range	TPR 0 to 1		TNR 0 to 1		Sorensen 0 to 1		Jaccard 0 to 1		FPB 0 to 2		OR 0 to 1		TSS -1 to 1		AUC 0 to 1		Boyce -1 to 1		IMAE 0 to 1	
		mean	sd	mean	sd	mean	sd	mean	sd	mean	sd	mean	sd	mean	sd	mean	sd	mean	sd	mean	sd
										Culex pipiens											
SVM		0.70	0.10	0.71	0.05	0.50	0.21	0.35	0.19	0.70	0.38	0.30	0.10	0.42	0.14	0.75	0.10	0.70	0.18	0.69	0.01
Neural Network		0.63	0.19	0.76	0.09	0.51	0.16	0.35	0.15	0.69	0.30	0.37	0.19	0.39	0.10	0.73	0.08	0.87	0.04	0.71	0.01
GBM		0.60	0.20	0.80	0.06	0.51	0.19	0.35	0.17	0.71	0.34	0.40	0.20	0.40	0.14	0.75	0.09	0.79	0.24	0.71	0.03
Random Forest		0.73	0.05	0.67	0.16	0.49	0.21	0.34	0.18	0.67	0.37	0.27	0.05	0.40	0.12	0.74	0.06	0.98	0.03	0.69	0.01
MaxEnt		0.72	0.08	0.67	0.03	0.49	0.16	0.33	0.14	0.67	0.28	0.28	0.08	0.40	0.05	0.74	0.06	0.96	0.06	0.60	0.00
Ensemble mean weighted		0.65	0.21	0.77	0.10	0.52	0.17	0.36	0.15	0.72	0.30	0.35	0.21	0.42	0.11	0.76	0.08	0.95	0.02	0.68	0.01
										Culex modestus											
SVM		0.68	0.12	0.66	0.11	0.5	0.02	0.33	0.02	0.67	0.04	0.32	0.12	0.34	0.01	0.66	0.02	0.89	0.07	0.64	0.02
Neural Network		0.79	0.17	0.61	0.16	0.53	0.03	0.36	0.03	0.72	0.06	0.21	0.17	0.39	0.01	0.68	0.08	0.93	0.08	0.67	0.02
GBM		0.58	0.19	0.76	0.16	0.5	0	0.33	0	0.67	0	0.42	0.19	0.34	0.03	0.63	0.03	0.95	0.06	0.71	0.04
Random Forest		0.32	0.18	0.93	0.04	0.4	0.17	0.26	0.13	0.52	0.26	0.68	0.18	0.25	0.14	0.56	0.08	0.72	0.15	0.67	0
MaxEnt		0.63	0.21	0.7	0.09	0.5	0.1	0.33	0.09	0.67	0.18	0.37	0.21	0.33	0.12	0.61	0.08	0.73	0.05	0.54	0.1
Ensemble mean threshold		0.66	0.27	0.7	0.18	0.52	0.07	0.35	0.07	0.7	0.13	0.34	0.27	0.36	0.08	0.67	0.05	0.96	0.02	0.72	0.05

Figure 2a-f: Species distribution models for *Culex pipiens* using a) MaxEnt, b) Random Forest, c) Generalized Boosted Regression, d) Neural Network, e) Support Vector Machine and f) A mean weighted Ensemble model of a – e, that was weighted based on model performance.

450 Figure 3a-f: Species distribution models for *Culex modestus* using a) MaxEnt, b) Random Forest, c)

451 Generalized Boosted Regression, d) Neural Network, e) Support Vector Machine and f) A mean

452 threshold Ensemble model of a – e based on cells with suitability above the maximum sensitivity and

453 specificity threshold.

Mechanistic species distribution models for vectors in the current climate and environmental conditions

Culex pipiens

Based on the CLIMEX model, Southern England is most suitable for *Cx. pipiens,* with a higher

ecoclimatic index compared to central and northern regions. Overall, much of the UK is suitable for

Cx. pipiens as an ecoclimatic index greater than 1 indicates that some growth could occur. The only

region calculated to be unsuitable is an area of higher elevation in central-north Scotland (Figure 4a).

Validation of the CLIMEX model to known presences was able to use 655 *Cx. pipiens* presence points

that fell within the study area, and these were assigned to categories based on the ecoclimatic index

score of the model. Of these, most presence points fell into areas considered to be very favourable

(N = 646), with the remaining presences located in favourable (N = 8) or marginally suitable (N = 1)

areas. No presences were recorded in areas considered unsuitable based on the ecoclimatic index.

Culex modestus

The CLIMEX model for *Cx. modestus* predicted that the south-east of England had the

highest ecoclimatic index in the UK, whereas much of northern and central Scotland was unsuitable

(ecoclimatic index was very low or 0 suggesting that no population growth could occur; Figure 4b).

Validation using known presence records showed that all 6 *Cx. modestus* records for the UK were in

areas that were very favourable based on the ecoclimatic index, however, the limited sample size

does reduce the reliability of validation for this species. Therefore, as has previously been done

when modelling with CLIMEX (Early et al., 2022), visual comparisons to known and predicted *Cx.*

modestus distributions were also considered and these did align with the output of the CLIMEX

model (Golding et al., 2012, European Centre for Disease Prevention and Control, 2019).

Figure 4a, b: The ecoclimatic index of the UK for a) *Cx. pipiens* and b) *Cx. modestus*. An ecoclimatic
index greater than 1 indicates some survival can occur, with the higher the number the more
suitable the area is climatically for the species.

West Nile Virus case study; current and future predictions of risk in the UK

Risk based on *Culex* vector distribution

Considering future projections for both *Cx. pipiens* and *Cx. modestus*, there are many areas

across the UK that see an increase or decrease in suitability for both SSP1 and SSP5 scenarios.

Increases in suitability are predominantly in areas currently identified as highly suitable, such as

across much of eastern England (Figure 5b-e). In the year 2100, the overall pattern of change for the

SSP5 scenario is very similar to that for SSP1, however the changes are predicted to be greater,

reflecting the increased change in the environment with the SSP5 scenario (Figure 5b-e).

Figure 5a-e: Habitat suitability for *Culex spp.* (*Cx. pipiens* and *Cx. modestus*) in a) the current climate
and in the year 2100 under b) SSP1 and c) SSP5 scenarios, and the change in suitability between the
current and future climates for scenario d) SPP1 and e) SSP5.

Risk based on *Culex* and representative avian host distribution

To get an improved understanding of WNV risk, a species distribution map of the representative

corvid and non-corvid hosts was also created (Figure 6a-c). The results here focus on the species

distribution modelling of vectors as the primary focus of this study; for full results of the species

distribution models for the avian hosts see Supplementary Information (SI) 5. Similar to the vector

distribution predictions, when considering the future environment some areas showed an increase

in suitability in south-east England and north-west Scotland whereas others had reduced suitability

such as in the north-east and central England. Overall there was much more widespread suitability

across the UK for host species, including areas of high suitability in northern Scotland which was

largely unsuitable for both vectors (Figure 5a-e, Figure 6a-c).

Figure 6a-c: Habitat suitability for the combined, representative host species (corvid and non-corvid)
in a) the current climate and the change in suitability by year 2100 under scenario b) SSP1 and c)
SSP5. For full details of predictions for host suitability see Supplementary Information (SI) 5.

By combining vector and host distribution maps a WNV risk map was produced to show areas of

high suitability for both vectors and hosts, as these areas are likely to have increased viral

transmission and a higher likelihood of the virus establishing. The WNV risk map showed that the

greatest risk is in England, particularly urbanised areas in the south-east, north-east and north-west

(Figure 7). Predicting into 2100 revealed that many of the areas currently deemed to be at risk will

have an increased risk under both SSP1 and SSP5 scenarios and, as expected based on the host and

vector maps, this change will be greater under SSP5 (Figure 7).

Figure 7a-e: The WNV risk in the UK calculated by considering both vector (*Cx. pipiens*, *Cx. modestus*)
and host (corvid and non-corvid) species distribution maps to infer overall risk. The figure shows a)
the current risk, b) predicted risk in 2100 under scenario SSP1, c) the predicted risk in 2100 under
scenario SSP5, and the change in risk by 2100 under scenario d) SSP1 and e) SSP5.

Risk based on *Culex* vectors, representative avian hosts and non-target human and equine
hosts

To further define risk in terms of human and equine health, risk maps based on vector and

host distributions were overlaid with maps reflecting human and horse densities in the UK. As

expected, the greatest risk to horses and humans coincided with areas of high suitability for vector

and host populations, such as the south-east and north-west of England (Figure 8a-d). Only human

density estimates are available for the future, and these revealed a very similar pattern of risk under

both SSP1 and SSP5, with at-risk areas expanding following the predicted spread of urbanised areas

with higher population densities (Figure 8a-d).

528 Figure 8a-d: The WNV risk in the UK calculated by considering both vector (*Cx. pipiens*, *Cx.*
529 *modestus*), target reservoir host (corvid and non-corvid) and dead-end host species distribution
530 maps to infer overall risk. The figure shows a) the risk to UK horses in the current climate, b) the risk
531 to the human population in the current climate, c) the risk to the human population in 2100 under
532 scenario SSP1 and, D) the risk to the human population in 2100 under scenario SSP5.

Discussion

The focus of this study was to determine a general framework of using species distribution

modelling of vectors and hosts to highlight areas at risk of vector-borne diseases in the UK. Here, it is

proposed that correlative species distribution models can be used to predict vector distributions,

and that mechanistic modelling approaches can help to validate correlative models. These vector

species distribution models can then be merged with correlative target host species distribution

models to assess risk from disease, with further refinements of disease risk predictions by then

combining with data for dead-end hosts such as humans, equines, or livestock.

Implications for disease risk in the UK

Using this approach, a case study on WNV was carried out and a risk map produced, showing that hotspots of WNV risk exist across the UK, with the greatest risk being in the south-east and north-west England. When projecting this risk into the future under SSP1 and SSP5 environmental scenarios some areas have an increased risk, including much of eastern England and hotspots in the north-west. This risk was also considered in relation to both human and horse population densities and, as expected, this showed disease risk hotspots located in densely populated areas where both vectors and hosts were likely to be present. To further improve the risk map for horses, improved horse population density estimate maps for the UK will be required as this is currently an area with limited data availability, with current predictions likely to be underestimated. However, the current map is likely to reflect the overall distribution of horses and thus show the potential risk to horses within the UK. As such, these risk maps will be important for targeting surveillance so any disease incursion or outbreak can be detected quickly, enabling a rapid response that aims to control the disease prior to spread and establishment.

Whilst this study has focussed on WNV, there are also other pathogens spread by *Cx. pipiens* and *Cx. modestus*, such as Usutu virus, avian malaria (*Plasmodium spp.*) and heartworm (*Difilaria immitis*) (Soto and Delang, 2023, Farajollahi et al., 2011). The vector risk maps presented in this study will also represent areas of higher risk from these diseases in the UK and could be refined further using the approach described here to combine vector and relevant non-/target host species distribution models. Using this approach could improve our understanding of the risk of diseases spread by *Culex* vectors across the UK. Furthermore, this methodology could also be tailored for other vectors, countries or regions to determine disease risk based on the likelihood of vector presence elsewhere.

Correlative and mechanistic approaches to species distribution modelling of *Culex* vectors in the UK

Whilst avian species distribution modelling was carried out to represent WNV host availability across the UK, carefully selected species were used to create representative host models. This is due to *Culex* species not showing a strong host preference for specific avian species and many different bird species being competent hosts for WNV (Brugman et al., 2013, Brugman et al., 2017, Centers for Disease Control and Prevention, 2021, Ziegler et al., 2022, Bergmann et al., 2023, Bessell et al., 2016). Therefore, vectors may be a key limiting factor in the spread of WNV and have higher habitat specificity so were the primary focus on this study and much of this discussion will focus on the species distribution modelling of the vectors.

The species distribution models of *Cx. pipiens* and *Cx. modestus* highlighted that both

species had areas of high suitability across the UK, with these areas often overlapping, highlighting

similarities in the species environmental needs. The transferability of the *Culex* data used to train the

models is important to consider as the WNV risk models are predominantly driven by vector habitat

suitability both in the current and future maps. Here, reviewing extrapolation of presence data with

the Shape method proposed by Velazco *et al.* (2023) showed that the ability to predict to current

and future climate in the UK was good for *Cx. pipiens* but more limited for *Cx. modestus*. This is to be

expected as *Cx. modestus* is an invasive species in the UK that may not have reached its full range,

and data availability is much more limited for this species (Golding et al., 2012, Soto and Delang,

2023, Elith et al., 2010, Venette, 2017). This further highlights the importance in improving current

knowledge of vector distributions, especially for species that have very limited data, such as *Cx.*

modestus.

Selecting which species distribution models are most appropriate can be a challenge, as each

model can produce markedly different results and cross-validation scoring approaches can vary

considerably as well as being strongly affected by biases in the data (Elith et al., 2006, Elith et al.,

2010, Hijmans and Elith, 2013, Roberts et al., 2017, Valavi et al., 2022). Throughout the modelling

process used in this study, steps were taken to ensure that biases were minimised. For example,

some performance scores can be artificially inflated by how data is partitioned into testing and

training datasets for cross-validation (Velazco et al., 2022, Roberts et al., 2017). To overcome this

host and vector data were partitioned based on environmental conditions in the model calibration

area. This was particularly important to consider for vector data as it is more limited than the avian

host data, with strong spatial biases in presence locations due to insect sampling methods typically

relying on trapping which can have biases due to trap attractiveness and locations (Koch, 2021,

Petrovskii et al., 2012).

The model that performed best was taken forward for both vectors and hosts. For *Cx.*

pipiens this was the mean-weighted Ensemble model whereas for *Cx. modestus* this was the Neural

Network model, highlighting how model performance can vary for different species. Species-specific

variations in model performance have been recorded previously for mosquito species in Europe

(European Centre for Disease Prevention and Control, 2019), as well as across 225 other species

(Valavi et al., 2022)*.* These differences in model suitability and scoring can be due to a number of

reasons, such as variation in sampling effort, balances between presence and absences and species

characteristics (e.g., species niches) (Valavi et al., 2022). Whilst performance metrics were used in

this study, visual comparisons to known distributions, previous predictions, and to the CLIMEX

mechanical model were also considered to aid model selection (European Centre for Disease

Prevention and Control, 2019, European Centre for Disease Prevention and Control and European

Food Safety Authority, 2023, Golding et al., 2012). This was important as it allowed consideration of

biological limits which can reflect geographical survival ranges of vector species.

There was more data available on *Cx. pipiens* to parametrise the CLIMEX model, and the

genetic algorithm was used to further inform some parameters based on known presence data

(Ewing et al., 2016, Ewing et al., 2017, Ewing et al., 2019). By comparing the CLIMEX output to

presence data available to inform on model performance, most presences were in areas considered

to be highly suitable for *Cx. pipiens* and no presences were in areas predicted to be unsuitable,

indicating the CLIMEX model performed well. Parameterising the CLIMEX model for *Cx. modestus*

was more challenging than for *Cx. pipiens* as data was much more limited (Soto and Delang, 2023).

Moreover, *Cx. modestus* has more specific habitat requirements for different life-stages, with larvae

typically recorded in permanent water bodies only (Soto and Delang, 2023). As such, land cover is

not considered in the CLIMEX modelling approach. Modelling using CLIMEX for species with limited

distributions has been carried out based on the upper and lower limits of the observed distribution

(Khormi and Kumar, 2014), or by matching to the known distribution of a species (Poutsma et al.,

2008, Venette, 2017). However, for invasive species with limited distributions and poor data

availability these approaches could lead to errors in estimating species distribution, as both rely on

absences in distribution data being true absences and may not reflect the biological limits of a

species (Venette, 2017). Therefore, to overcome these challenges the *Cx. modestus* CLIMEX model

relies on the genetic algorithm results, supplementing missing parameters with those used for *Cx.*

pipiens as it is a closely related species that also has winter diapause (Soto and Delang, 2023). The

resulting *Cx. modestus* model performed well in model validation, with all presence data available in

areas considered to be very favourable and visual comparisons showing alignment with the known

distribution of this species (Golding et al., 2012, European Centre for Disease Prevention and

Control, 2019). This would suggest that the model is informative, however, to further improve the

CLIMEX model, extensive research to refine model parameterisation would be beneficial for *Cx.*

modestus, especially those parameters that show high levels of sensitivity (i.e. temperature limits,

soil moisture), which can lead to strong variations in suitability predictions based on ecoclimatic

indices (Taylor and Kumar, 2012).

Comparisons of the mechanistic and correlative models showed similarities for *Cx.*

modestus, with both approaches predicting higher suitability in the south-east and very low

suitability in northern-Scotland. Similarly, the *Cx. pipiens* mechanistic model closely reflects the

output of the correlative models, with both showing southern England and the western coastal areas

of Wales as more suitable than more northern areas. The CLIMEX model does not highlight urban

areas as more suitable for *Cx. pipiens* however the correlative models can identify this, which is expected as *Cx. pipiens* is commonly associated with urbanisation (Ruybal et al., 2016, Krol et al., 2024, Townroe and Callaghan, 2014). This is likely due to land use being included in the correlative models whereas the CLIMEX model is based solely on biological limits and climate information. Therefore, for vectors with strong associations to land use types correlative models may be more suitable, with CLIMEX used to support where the climatic conditions are suitable for vector survival and reproduction, with correlation between the two modelling approaches indicative of strong models.

Conclusion

By combining both vector and host distributions it is possible to assess potential risks from novel diseases in the UK. This can highlight where a pathogen is most likely to establish due to increased environmental suitability for both vectors and hosts, both presently and in the future. This study has highlighted that the risk of WNV risk establishment is expected to increase in some areas whilst decreasing in others, leading to areas of high and low risk across the UK in the future. Overall, a novel method, combining correlative and mechanistic vector species distribution models, correlative host species distribution models and non-target host data to consider disease risk is presented. It is hoped that this approach will be transferable to other diseases and geographic regions in the future.

References

BARBET-MASSIN, M., JIGUET, F., ALBERT, C. H. & THUILLER, W. 2012. Selecting pseudo-absences for species distribution models: how, where and how many? *Methods in Ecology and Evolution,* 3**,** 327-338. *doi:* https://doi.org/10.1111/j.2041-210X.2011.00172.x

BEEMAN, S. P., MORRISON, A. M., UNNASCH, T. R. & UNNASCH, R. S. 2021. Ensemble ecological niche modeling of West Nile virus probability in Florida. *PLOS ONE,* 16**,** e0256868. *doi:* 10.1371/journal.pone.0256868

BERGMANN, F., SCHMOOCK-WELLHAUSEN, M., FAST, C., HOLICKI, C., MICHEL, F., WYSOCKI, P., SADEGHI, B., GROSCHUP, M. & ZIEGLER, U. 2023. Longitudinal Study of the Occurrence of Usutu Virus and West Nile Virus Infections in Birds in a Zoological Garden in Northern Germany. *Pathogens,* 12**,** 753. *doi:* 10.3390/pathogens12060753

BESSELL, P. R., ROBINSON, R. A., GOLDING, N., SEARLE, K. R., HANDEL, I. G., BODEN, L. A., PURSE, B. V. & BRONSVOORT, B. M. D. C. 2016. Quantifying the Risk of Introduction of West Nile Virus into Great Britain by Migrating Passerine Birds. *Transboundary and Emerging Diseases,* 63**,** e347-e359. *doi:* https://doi.org/10.1111/tbed.12310

BRUGMAN, V. A., HERNÁNDEZ-TRIANA, L. M., ENGLAND, M. E., MEDLOCK, J. M., MERTENS, P. P. C., LOGAN, J. G., WILSON, A. J., FOOKS, A. R., JOHNSON, N. & CARPENTER, S. 2017. Blood-feeding patterns of native mosquitoes and insights into their potential role as pathogen vectors in the Thames estuary region of the United Kingdom. *Parasites & Vectors,* 10**,** 163. *doi:* 10.1186/s13071-017-2098-4

BRUGMAN, V. A., HORTON, D. L., PHIPPS, L. P., JOHNSON, N., COOK, A. J. C., FOOKS, A. R. & BREED,
A. C. 2013. Epidemiological perspectives on West Nile virus surveillance in wild birds in Great
Britain. *Epidemiology & Infection,* 141**,** 1134-1142. *doi:* 10.1017/S095026881200177X
CASTELLANOS, A. A., HUNTLEY, J. W., VOELKER, G. & LAWING, A. M. 2019. Environmental filtering
improves ecological niche models across multiple scales. *Methods in Ecology and Evolution,*
10**,** 481-492. *doi:* https://doi.org/10.1111/2041-210X.13142
CENTERS FOR DISEASE CONTROL AND PREVENTION 2021. Species of dead birds in which West Nile
virus has been detected, United States 1999-2016. Available from:
https://www.cdc.gov/westnile/dead-birds/index.html
CHEN, G., LI, X. & LIU, X. 2022. Global land projection based on plant functional types with a 1-km
resolution under socio-climatic scenarios. *Scientific Data,* 9**,** 125. *doi:* 10.1038/s41597-022-
01208-6
CONSTANT, O., GIL, P., BARTHELEMY, J., BOLLORÉ, K., FOULONGNE, V., DESMETZ, C., LEBLOND, A.,
DESJARDINS, I., PRADIER, S., JOULIÉ, A., SANDOZ, A., AMARAL, R., BOISSEAU, M.,
RAKOTOARIVONY, I., BALDET, T., MARIE, A., FRANCES, B., REBOUL SALZE, F., TINTO, B., VAN
DE PERRE, P., SALINAS, S., BECK, C., LECOLLINET, S., GUTIERREZ, S. & SIMONIN, Y. 2022. One
Health surveillance of West Nile and Usutu viruses: a repeated cross-sectional study
exploring seroprevalence and endemicity in Southern France, 2016 to 2020.
Eurosurveillance, 27**,** 2200068. *doi:* doi:https://doi.org/10.2807/1560-
7917.ES.2022.27.25.2200068
CROFT, S. & SMITH, G. C. 2019. Structuring the unstructured: estimating species-specific absence
from multi-species presence data to inform pseudo-absence selection in species distribution
models. *bioRxiv***,** 656629. *doi:* 10.1101/656629
CULVERWELL, C. L. & VAPALAHTI, O. 2023. First record of *Culex modestus* in Finland. *Journal of the*
European Mosquito Control Association, 41**,** 63-66. *doi:*
https://doi.org/10.52004/JEMCA2023.0003
DAVIES, N. B. 1976. Food, Flocking and Territorial Behaviour of the Pied Wagtail (*Motacilla alba*
yarrellii Gould) in Winter. *Journal of Animal Ecology,* 45**,** 235-253. *doi:* 10.2307/3777
DURAND, B., BALANÇA, G., BALDET, T. & CHEVALIER, V. 2010. A metapopulation model to simulate
West Nile virus circulation in Western Africa, Southern Europe and the Mediterranean basin.
Veterinary research, 41**,** 32. *doi:* 10.1051/vetres/2010004
EARLY, R., RWOMUSHANA, I., CHIPABIKA, G. & DAY, R. 2022. Comparing, evaluating and combining
statistical species distribution models and CLIMEX to forecast the distributions of emerging
crop pests. *Pest Management Science,* 78**,** 671-683. *doi:* https://doi.org/10.1002/ps.6677
ELITH, J., H. GRAHAM, C., P. ANDERSON, R., DUDÍK, M., FERRIER, S., GUISAN, A., J. HIJMANS, R.,
HUETTMANN, F., R. LEATHWICK, J., LEHMANN, A., LI, J., G. LOHMANN, L., A. LOISELLE, B.,
MANION, G., MORITZ, C., NAKAMURA, M., NAKAZAWA, Y., MCC. M. OVERTON, J.,
TOWNSEND PETERSON, A., J. PHILLIPS, S., RICHARDSON, K., SCACHETTI-PEREIRA, R., E.
SCHAPIRE, R., SOBERÓN, J., WILLIAMS, S., S. WISZ, M. & E. ZIMMERMANN, N. 2006. Novel
methods improve prediction of species' distributions from occurrence data. *Ecography,* 29**,**
129-151. *doi:* https://doi.org/10.1111/j.2006.0906-7590.04596.x
ELITH, J., KEARNEY, M. & PHILLIPS, S. 2010. The art of modelling range-shifting species. *Methods in*
Ecology and Evolution, 1**,** 330-342. *doi:* https://doi.org/10.1111/j.2041-210X.2010.00036.x
ENDO, A. & AMARASEKARE, P. 2022. Predicting the Spread of Vector-Borne Diseases in a Warming
World. *Frontiers in Ecology and Evolution,* 10. *doi:* 10.3389/fevo.2022.758277
ESRI, I. 2022. ArcGIS Pro version 3.0.1. . Available from: https://www.esri.com/en-
us/arcgis/products/arcgis-pro/overview
EUROPEAN CENTRE FOR DISEASE PREVENTION AND CONTROL 2019. A spatial modelling method for
vector surveillance. Available from: https://www.ecdc.europa.eu/en/publications-
data/spatial-modelling-method-vector-surveillance

EUROPEAN CENTRE FOR DISEASE PREVENTION AND CONTROL AND EUROPEAN FOOD SAFETY AUTHORITY. 2022. *Mosquito maps* [Online]. Stockholm. Available: https://ecdc.europa.eu/en/disease-vectors/surveillance-and-disease-data/mosquito-maps [Accessed 05/10/2022].

EUROPEAN CENTRE FOR DISEASE PREVENTION AND CONTROL AND EUROPEAN FOOD SAFETY AUTHORITY. 2023. *Culex pipiens group - current known distribution: October 2023* [Online]. Available: https://www.ecdc.europa.eu/en/publications-data/culex-pipiens-group-current-known-distribution-october-2023 [Accessed 28/06/2024].

EWING, D. A., COBBOLD, C. A., PURSE, B. V., NUNN, M. A. & WHITE, S. M. 2016. Modelling the effect of temperature on the seasonal population dynamics of temperate mosquitoes. *Journal of Theoretical Biology,* 400**,** 65-79. *doi:* https://doi.org/10.1016/j.jtbi.2016.04.008

EWING, D. A., PURSE, B. V., COBBOLD, C. A., SCHÄFER, S. M. & WHITE, S. M. 2017. Seasonal abundance data of all *Culex pipiens* life stages at a UK field site. NERC Environmental Information Data Centre. *doi:*

EWING, D. A., PURSE, B. V., COBBOLD, C. A., SCHÄFER, S. M. & WHITE, S. M. 2019. Uncovering mechanisms behind mosquito seasonality by integrating mathematical models and daily empirical population data: *Culex pipiens* in the UK. *Parasites & Vectors,* 12**,** 74. *doi:* 10.1186/s13071-019-3321-2

FABURAY, B. 2015. The case for a 'one health' approach to combating vector-borne diseases. *Infection Ecology & Epidemiology,* 5**,** 28132. *doi:* 10.3402/iee.v5.28132

FARAJOLLAHI, A., FONSECA, D. M., KRAMER, L. D. & MARM KILPATRICK, A. 2011. "Bird biting" mosquitoes and human disease: A review of the role of *Culex pipiens* complex mosquitoes in epidemiology. *Infection, Genetics and Evolution,* 11**,** 1577-1585. *doi:* https://doi.org/10.1016/j.meegid.2011.08.013

FICK, S. E. & HIJMANS, R. J. 2017. WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology,* 37**,** 4302-4315. *doi:* https://doi.org/10.1002/joc.5086

FURLONG, M., ADAMU, A., HICKSON, R. I., HORWOOD, P., GOLCHIN, M., HOSKINS, A. & RUSSELL, T. 2022. Estimating the Distribution of Japanese Encephalitis Vectors in Australia Using Ecological Niche Modelling. *Tropical Medicine and Infectious Disease,* 7. *doi:* 10.3390/tropicalmed7120393

GANDY, S. L., HANSFORD, K. M. & MEDLOCK, J. M. 2022. Possible expansion of *Ixodes ricinus* in the United Kingdom identified through the Tick Surveillance Scheme between 2013 and 2020. *Medical and Veterinary Entomology,* n/a. *doi:* https://doi.org/10.1111/mve.12612

GAO, J. 2017. Downscaling Global Spatial Population Projections from 1/8-degree to 1-km Grid Cells. Available from: https://doi.org/10.5065/D60Z721H

GAO, J. 2020. Global 1-km Downscaled Population Base Year and Projection Grids Based on the Shared Socioeconomic Pathways, Revision 01. Palisades, New York: NASA Socioeconomic Data and Applications Center (SEDAC). *doi:* 10.7927/q7z9-9r69

GBIF ORG USER 2022. Occurrence Download: *Culex*. The Global Biodiversity Information Facility. *doi:* 10.15468/DL.FQC9T3

GBIF. ORG USER 2024. Occurrence Download - *Hirundo rustica*. The Global Biodiversity Information Facility. *doi:* 10.15468/DL.E8TZ2B

GBIF.ORG USER 2024a. Occurrence Download - *Corvus corone*. The Global Biodiversity Information Facility. *doi:* 10.15468/DL.Q7TAWD

GBIF.ORG USER 2024b. Occurrence Download - *Turdus merula* The Global Biodiversity Information Facility. *doi:* 10.15468/DL.7MYK3H

GBIF.ORG USER 2024c. Occurrence Download - *Corvus corax*. The Global Biodiversity Information Facility. *doi:* 10.15468/DL.QZ6KFF

GBIF.ORG USER 2024d. Occurrence Download - *Corvus frugilegus*. The Global Biodiversity Information Facility. *doi:* 10.15468/DL.DQH9EQ

GBIF.ORG USER 2024e. Occurrence Download - *Passer domesticus*. The Global Biodiversity
Information Facility. *doi:* 10.15468/DL.YWR2JF
GILBERT, M., CINARDI, G., DA RE, D., WINT, W. G. R., WISSER, D. & ROBINSON, T. P. 2022. Global
horses distribution in 2015 (5 minutes of arc). V1 ed.: Harvard Dataverse. *doi:*
doi:10.7910/DVN/JJGCTX
GIOVANELLI, J., SIQUEIRA, M., HADDAD, C. & ALEXANDRINO, J. 2010. Modeling a spatially restricted
distribution in the Neotropics: How the size of calibration area affects the performance of
five presence-only methods. *Ecological Modelling,* 221**,** 215-224. *doi:*
10.1016/j.ecolmodel.2009.10.009
GOLDING, N., NUNN, M. A., MEDLOCK, J. M., PURSE, B. V., VAUX, A. G. C. & SCHÄFER, S. M. 2012.
West Nile virus vector *Culex modestus* established in southern England. *Parasites & Vectors,*
5**,** 32. *doi:* 10.1186/1756-3305-5-32
GORRIS, M. E., BARTLOW, A. W., TEMPLE, S. D., ROMERO-ALVAREZ, D., SHUTT, D. P., FAIR, J. M.,
KAUFELD, K. A., DEL VALLE, S. Y. & MANORE, C. A. 2021. Updated distribution maps of
predominant *Culex* mosquitoes across the Americas. *Parasites & Vectors,* 14**,** 547. *doi:*
10.1186/s13071-021-05051-3
HERNÁNDEZ-TRIANA, L. M., BRUGMAN, V. A., PRAMUAL, P., BARRERO, E., NIKOLOVA, N. I., RUIZ-
ARRONDO, I., KAISER, A., KRÜGER, A., LUMLEY, S., OSÓRIO, H. C., IGNJATOVIĆ-ĆUPINA, A.,
PETRIĆ, D., LAURE SETIER-RIO, M., BØDKER, R. & JOHNSON, N. 2020. Genetic diversity and
population structure of *Culex modestus* across Europe: does recent appearance in the
United Kingdom reveal a tendency for geographical spread? *Medical and Veterinary*
Entomology, 34**,** 86-96. *doi:* https://doi.org/10.1111/mve.12412
HIJMANS, R. & ELITH, J. 2013. *Species Distribution Modeling with R,* 692-705, Elsevier Inc.*doi:*
10.1016/B978-0-12-384719-5.00318-X
HIJMANS, R. J., PHILLIPS, S., LEATHWICK, J. & ELITH, J. 2023. Dismo: Species Distribution Modeling.
1.3-14 ed. Available from: https://github.com/rspatial/dismo.
HUMBLET, M. F., VANDEPUTTE S FAU - FECHER-BOURGEOIS, F., FECHER-BOURGEOIS F FAU -
LÉONARD, P., LÉONARD P FAU - GOSSET, C., GOSSET C FAU - BALENGHIEN, T., BALENGHIEN T
FAU - DURAND, B., DURAND B FAU - SAEGERMAN, C. & SAEGERMAN, C. 2016. Estimating the
economic impact of a possible equine and human epidemic of West Nile virus infection in
Belgium. *Eurosurveillance* 21. *doi:*
ITURBIDE, M., BEDIA, J., HERRERA, S., DEL HIERRO, O., PINTO, M. & GUTIÉRREZ, J. M. 2015. A
framework for species distribution modelling with improved pseudo-absence generation.
Ecological Modelling, 312**,** 166-174. *doi:* https://doi.org/10.1016/j.ecolmodel.2015.05.018
KHORMI, H. M. & KUMAR, L. 2014. Climate change and the potential global distribution of *Aedes*
aegypti: spatial modelling using geographical information system and CLIMEX. *Geospatial*
Health, 8**,** 405-415. *doi:* 10.4081/gh.2014.29
KOCH, F. H. 2021. Considerations regarding species distribution models for forest insects.
Agricultural and Forest Entomology, 23**,** 393-399. *doi:* https://doi.org/10.1111/afe.12458
KOMAR, N., LANGEVIN, S., HINTEN, S., NEMETH, N., EDWARDS, E., HETTLER, D., DAVIS, B., BOWEN,
R. & BUNNING, M. 2003. Experimental Infection of North American Birds with the New York
1999 Strain of West Nile Virus. *Emerg Infect Dis. 9(3):311-322,* 9. *doi:*
KRITICOS, D. J., MAYWALD, G. F., YONOW, T., ZURCHER, E. J., HERRMANN, N. I. & SUTHERST, R. W.
2015. CLIMEX Version 4: Exploring the effects of climate on plants, animals and diseases.
CSIRO. Available from: https://www.hearne.software/Software/CLIMEX-DYMEX/Editions
KROL, L., LANGEZAAL, M., BUDIDARMA, L., WASSENAAR, D., DIDASKALOU, E. A., TRIMBOS, K.,
DELLAR, M., VAN BODEGOM, P. M., GEERLING, G. W. & SCHRAMA, M. 2024. Distribution of
Culex pipiens life stages across urban green and grey spaces in Leiden, The Netherlands.
Parasites & Vectors, 17**,** 37. *doi:* 10.1186/s13071-024-06120-z
LAWSON, B., ROBINSON, R. A., BRISCOE, A. G., CUNNINGHAM, A. A., FOOKS, A. R., HEAVER, J. P.,
HERNÁNDEZ-TRIANA, L. M., JOHN, S. K., JOHNSON, N., JOHNSTON, C., LEAN, F. Z. X.,

MACGREGOR, S. K., MASTERS, N. J., MCCRACKEN, F., MCELHINNEY, L. M., MEDLOCK, J. M., PEARCE-KELLY, P., SEILERN-MOY, K., SPIRO, S., VAUX, A. G. C. & FOLLY, A. J. 2022. Combining host and vector data informs emergence and potential impact of an Usutu virus outbreak in UK wild birds. *Scientific Reports,* 12**,** 10298. *doi:* 10.1038/s41598-022-13258-2

LIU, C., NEWELL, G. & WHITE, M. 2019. The effect of sample size on the accuracy of species distribution models: considering both presences and pseudo-absences or background sites. *Ecography,* 42**,** 535-548. *doi:* https://doi.org/10.1111/ecog.03188

LOBO, J. M., JIMÉNEZ-VALVERDE, A. & HORTAL, J. 2010. The uncertain nature of absences and their importance in species distribution modelling. *Ecography,* 33**,** 103-114. *doi:* https://doi.org/10.1111/j.1600-0587.2009.06039.x

MEDLOCK, J. M., HANSFORD, K. M., VAUX, A. G. C., CULL, B., GILLINGHAM, E. & LEACH, S. 2018. Assessment of the Public Health Threats Posed by Vector-Borne Disease in the United Kingdom (UK). *International Journal of Environmental Research and Public Health,* 15. *doi:* 10.3390/ijerph15102145

MEDLOCK, J. M. & LEACH, S. A. 2015. Effect of climate change on vector-borne disease risk in the UK. *The Lancet Infectious Diseases,* 15**,** 721-730. *doi:* https://doi.org/10.1016/S1473-3099(15)70091-5

MONTECINO-LATORRE, D. & BARKER, C. M. 2018. Overwintering of West Nile virus in a bird community with a communal crow roost. *Scientific Reports,* 8**,** 6088. *doi:* 10.1038/s41598-018-24133-4

NAIMI, B., HAMM, N. A. S., GROEN, T. A., SKIDMORE, A. K. & TOXOPEUS, A. G. 2014. Where is positional uncertainty a problem for species distribution modelling? *Ecography,* 37**,** 191-203. *doi:* https://doi.org/10.1111/j.1600-0587.2013.00205.x

NBN ATLAS. 2022. *Occurance Records - Culicidae* [Online]. Available: https://records.nbnatlas.org/occurrences/search?q=lsid:NBNSYS0000040182&fq=occurrenc e_status:present&nbn_loading=true#tab_mapView [Accessed 07/11/2022].

NOLL, M., WALL, R., MAKEPEACE, B. L., NEWBURY, H., ADASZEK, L., BØDKER, R., ESTRADA-PEÑA, A., GUILLOT, J., DA FONSECA, I. P., PROBST, J., OVERGAAUW, P., STRUBE, C., ZAKHAM, F., ZANET, S. & ROSE VINEER, H. 2023. Predicting the distribution of *Ixodes ricinus* and *Dermacentor reticulatus* in Europe: a comparison of climate niche modelling approaches. *Parasites & Vectors,* 16**,** 384. *doi:* 10.1186/s13071-023-05959-y

PALACIO, R. D., NEGRET, P. J., VELÁSQUEZ-TIBATÁ, J. & JACOBSON, A. P. 2021. A data-driven geospatial workflow to map species distributions for conservation assessments. *Diversity and Distributions,* 27**,** 2559-2570. *doi:* https://doi.org/10.1111/ddi.13424

PETROVSKII, S., BEARUP, D., AHMED, D. A. & BLACKSHAW, R. 2012. Estimating insect population density from trap counts. *Ecological Complexity,* 10**,** 69-82. *doi:* https://doi.org/10.1016/j.ecocom.2011.10.002

POUTSMA, J., LOOMANS, A. J. M., AUKEMA, B. & HEIJERMAN, T. 2008. Predicting the potential geographical distribution of the harlequin ladybird, *Harmonia axyridis*, using the CLIMEX model. *BioControl,* 53**,** 103-125. *doi:* 10.1007/s10526-007-9140-y

PRADIER, S., LECOLLINET, S. & LEBLOND, A. 2012. West Nile virus epidemiology and factors triggering change in its distribution in Europe. *Scientific & Technical Review,* 31**,** 829-844. *doi:* dx.doi.org/10.20506/rst.31.3.2167

RICHMAN, R., DIALLO, D., DIALLO, M., SALL, A. A., FAYE, O., DIAGNE, C. T., DIA, I., WEAVER, S. C., HANLEY, K. A. & BUENEMANN, M. 2018. Ecological niche modeling of *Aedes* mosquito vectors of chikungunya virus in southeastern Senegal. *Parasites & Vectors,* 11**,** 255. *doi:* 10.1186/s13071-018-2832-6

ROBERTS, D. R., BAHN, V., CIUTI, S., BOYCE, M. S., ELITH, J., GUILLERA-ARROITA, G., HAUENSTEIN, S., LAHOZ-MONFORT, J. J., SCHRÖDER, B., THUILLER, W., WARTON, D. I., WINTLE, B. A., HARTIG, F. & DORMANN, C. F. 2017. Cross-validation strategies for data with temporal, spatial,

882   hierarchical, or phylogenetic structure. *Ecography,* 40**,** 913-929. *doi:*
883   https://doi.org/10.1111/ecog.02881

884 ROCKLÖV, J. & DUBROW, R. 2020. Climate change: an enduring challenge for vector-borne disease
885   prevention and control. *Nature Immunology,* 21**,** 479-483. *doi:* 10.1038/s41590-020-0648-y

886 ROJAS-SOTO, O., FORERO-RODRÍGUEZ, J. S., GALINDO-CRUZ, A., MOTA-VARGAS, C., PARRA-HENAO,
887   K. D., PEÑA-PENICHE, A., PIÑA-TORRES, J., ROJAS-HERRERA, K., SÁNCHEZ-RODRÍGUEZ, J. D.,
888   TORO-CARDONA, F. A. & TRINIDAD-DOMÍNGUEZ, C. D. 2024. Calibration areas in ecological
889   niche and species distribution modelling: Unravelling approaches and concepts. *Journal of*
890   *Biogeography,* n/a. *doi:* https://doi.org/10.1111/jbi.14834

891 ROUSSEAU, J. S. & BETTS, M. G. 2022. Factors influencing transferability in species distribution
892   models. *Ecography,* 2022**,** e06060. *doi:* https://doi.org/10.1111/ecog.06060

893 RUDOLF, I., ŠIKUTOVÁ, S., ŠEBESTA, O., MENDEL, J., MALENOVSKÝ, I., KAMPEN, H., MEDLOCK, J. &
894   SCHAFFNER, F. 2021. Overwintering of *Culex modestus* and Other Mosquito Species in a
895   Reedbed Ecosystem, Including Arbovirus Findings. *Journal of the American Mosquito Control*
896   *Association,* 36, 257-260. *doi:* 10.2987/20-6949.1

897 RUYBAL, J. E., KRAMER, L. D. & KILPATRICK, A. M. 2016. Geographic variation in the response of
898   *Culex pipiens* life history traits to temperature. *Parasites & Vectors,* 9**,** 116. *doi:*
899   10.1186/s13071-016-1402-z

900 SANTINI, L., BENÍTEZ-LÓPEZ, A., MAIORANO, L., ČENGIĆ, M. & HUIJBREGTS, M. A. J. 2021. Assessing
901   the reliability of species distribution projections in climate change research. *Diversity and*
902   *Distributions,* 27**,** 1035-1050. *doi:* https://doi.org/10.1111/ddi.13252

903 SCHOLTE, E. J., DEN HARTOG, W., BRAKS, M., REUSKEN, C., DIK, M. & HESSELS, A. 2009. First report
904   of a North American invasive mosquito species *Ochlerotatus atropalpus* (Coquillett) in the
905   Netherlands, 2009. *Eurosurveillance,* 14**,** 19400. *doi:*
906   doi:https://doi.org/10.2807/ese.14.45.19400-en

907 SENAY, S. D. & WORNER, S. P. 2019. Multi-Scenario Species Distribution Modeling. *Insects,* 10. *doi:*
908   10.3390/insects10030065

909 SENAY, S. D., WORNER, S. P. & IKEDA, T. 2013. Novel Three-Step Pseudo-Absence Selection
910   Technique for Improved Species Distribution Modelling. *PLOS ONE,* 8**,** e71218. *doi:*
911   10.1371/journal.pone.0071218

912 SIMONS, R. R. L., CROFT, S., REES, E., TEARNE, O., ARNOLD, M. E. & JOHNSON, N. 2019. Using species
913   distribution models to predict potential hot-spots for Rift Valley Fever establishment in the
914   United Kingdom. *PLOS ONE,* 14**,** e0225250. *doi:* 10.1371/journal.pone.0225250

915 SOTO, A. & DELANG, L. 2023. *Culex modestus*: the overlooked mosquito vector. *Parasites & Vectors,*
916   16**,** 373. *doi:* 10.1186/s13071-023-05997-6

917 SUTHERLAND, L. N., POWELL, G. S. & BYBEE, S. M. 2021. Validating species distribution models to
918   illuminate coastal fireflies in the South Pacific (Coleoptera: Lampyridae). *Scientific Reports,*
919   11**,** 17397. *doi:* 10.1038/s41598-021-96534-x

920 SUTHERST, R. W. & MAYWALD, G. F. 1985. A computerised system for matching climates in ecology.
921   *Agriculture, Ecosystems & Environment,* 13**,** 281-299. *doi:* https://doi.org/10.1016/0167-
922   8809(85)90016-7

923 TAYLOR, S. & KUMAR, L. 2012. Sensitivity Analysis of CLIMEX Parameters in Modelling Potential
924   Distribution of *Lantana camara L*. *PLOS ONE,* 7**,** e40969. *doi:* 10.1371/journal.pone.0040969

925 THUILLER, W., GEORGES, D., GUEGUEN, M., ENGLER, R., BRIENER, F. & LAFOUR, B. 2024. biomod2:
926   Ensemble platform for species distribution modelling. 4.2-6-1 ed. Available from:
927   https://biomodhub.github.io/biomod2/

928 TOWNROE, S. & CALLAGHAN, A. 2014. British Container Breeding Mosquitoes: The Impact of
929   Urbanisation and Climate Change on Community Composition and Phenology. *PLOS ONE,* 9**,**
930   e95325. *doi:* 10.1371/journal.pone.0095325

VALAVI, R., ELITH, J., LAHOZ-MONFORT, J. J. & GUILLERA-ARROITA, G. 2021. Modelling species
presence-only data with random forests. *Ecography,* 44**,** 1731-1742. *doi:*
https://doi.org/10.1111/ecog.05615
VALAVI, R., GUILLERA-ARROITA, G., LAHOZ-MONFORT, J. J. & ELITH, J. 2022. Predictive performance
of presence-only species distribution models: a benchmark study with reproducible code.
Ecological Monographs, 92**,** e01486. *doi:* https://doi.org/10.1002/ecm.1486
VANDERWAL, J., SHOO, L. P., GRAHAM, C. & WILLIAMS, S. E. 2009. Selecting pseudo-absence data
for presence-only distribution modeling: How far should you stray from what you know?
Ecological Modelling, 220**,** 589-594. *doi:* https://doi.org/10.1016/j.ecolmodel.2008.11.010
VAUX, A. G. C., GIBSON, G., HERNANDEZ-TRIANA, L. M., CHEKE, R. A., MCCRACKEN, F., JEFFRIES, C. L.,
HORTON, D. L., SPRINGATE, S., JOHNSON, N., FOOKS, A. R., LEACH, S. & MEDLOCK, J. M.
2015. Enhanced West Nile virus surveillance in the North Kent marshes, UK. *Parasites &*
Vectors, 8**,** 91. *doi:* 10.1186/s13071-015-0705-9
VELAZCO, S. J. E., ROSE, M. B., DE ANDRADE, A. F. A., MINOLI, I. & FRANKLIN, J. 2022. flexsdm: An r
package for supporting a comprehensive and flexible species distribution modelling
workflow. *Methods in Ecology and Evolution,* 13**,** 1661-1669. *doi:*
https://doi.org/10.1111/2041-210X.13874
VELAZCO, S. J. E., ROSE, M. B., DE MARCO JR, P., REGAN, H. M. & FRANKLIN, J. 2023. How far can I
extrapolate my species distribution model? Exploring shape, a novel method. *Ecography,*
2024. *doi:* https://doi.org/10.1111/ecog.06992
VENETTE, R. C. 2017. Climate Analyses to Assess Risks from Invasive Forest Insects: Simple Matching
to Advanced Models. *Current Forestry Reports,* 3**,** 255-268. *doi:* 10.1007/s40725-017-0061-4
VOGELS, C. B. F., HARTEMINK, N. & KOENRAADT, C. J. M. 2017. Modelling West Nile virus
transmission risk in Europe: effect of temperature and mosquito biotypes on the basic
reproduction number. *Scientific Reports,* 7**,** 5022. *doi:* 10.1038/s41598-017-05185-4
WORLD HEALTH ORGANISATION. 2020. *Vector-borne diseases* [Online]. Available:
https://www.who.int/news-room/fact-sheets/detail/vector-borne-
diseases#:~:text=Vector%2Dborne%20diseases%20account%20for,infection%20transmitted
%20by%20Anopheline%20mosquitoes. [Accessed 08/03/2024].
WORLD HEALTH ORGANISATION. 2017. *West Nile virus* [Online]. Available:
https://www.who.int/news-room/fact-sheets/detail/west-nile-virus. [Accessed 02/09/2024].
ZIEGLER, U., BERGMANN, F., FISCHER, D., MÜLLER, K., HOLICKI, C. M., SADEGHI, B., SIEG, M., KELLER,
M., SCHWEHN, R., REUSCHEL, M., FISCHER, L., KRONE, O., RINDER, M., SCHÜTTE, K.,
SCHMIDT, V., EIDEN, M., FAST, C., GÜNTHER, A., GLOBIG, A., CONRATHS, F. J., STAUBACH, C.,
BRANDES, F., LIERZ, M., KORBEL, R., VAHLENKAMP, T. W. & GROSCHUP, M. H. 2022. Spread
of West Nile Virus and Usutu Virus in the German Bird Population, 2019-2020.
Microorganisms [Online], 10.
ZIZKA, A., SILVESTRO, D., ANDERMANN, T., AZEVEDO, J., DUARTE RITTER, C., EDLER, D., FAROOQ, H.,
HERDEAN, A., ARIZA, M., SCHARN, R., SVANTESSON, S., WENGSTRÖM, N., ZIZKA, V. &
ANTONELLI, A. 2019. CoordinateCleaner: Standardized cleaning of occurrence records from
biological collection databases. *Methods in Ecology and Evolution,* 10**,** 744-751. *doi:*
https://doi.org/10.1111/2041-210X.13152