A gene-based model of fitness and its implications for genetic variation:

Linkage disequilibrium

Parul Johri[1,2,3,*] and Brian Charlesworth[4,*]

[1]Department of Biology, University of North Carolina at Chapel Hill, NC 27599, United
States
[2]Department of Genetics, University of North Carolina at Chapel Hill, NC 27599, United
States
[3]Integrative Program for Biological and Genome Sciences, University of North Carolina at
Chapel Hill, NC 27599, United States
[4]Institute of Ecology and Evolution, School of Biological Sciences, University of Edinburgh
EH9 3FL, United Kingdom
13
14
15 *correspondence may be addressed to these authors: pjohri@unc.edu;
16 brian.charlesworth@ed.ac.uk
17

19 **Keywords:** complementation, dominance, epistasis, fitness models, linkage disequilibrium

21 **Running title:** Fitness models and linkage disequilibrium

Abstract

23
A widely used model of the effects of mutations on fitness (the "sites" model) assumes that

heterozygous recessive or partially recessive deleterious mutations at different sites in a gene

complement each other, similarly to mutations in different genes. However, the general lack

of complementation between major effect allelic mutations suggests an alternative possibility,

which we term the "gene" model. This assumes that a pair of heterozygous deleterious

mutations in *trans* behave effectively as homozygotes, so that the fitnesses of *trans*

heterozygotes are lower than those of *cis* heterozygotes. We examine the properties of the

two different models, using both analytical and simulation methods. We show that the gene

model predicts positive linkage disequilibrium (LD) between deleterious variants within the

coding sequence, under conditions when the sites model predicts zero or slightly negative

LD. We also show that focussing on rare variants when examining patterns of LD, especially

with Lewontin's D' measure, is likely to produce misleading results with respect to

inferences concerning the causes of the sign of LD. Synergistic epistasis between pairs of

mutations was also modeled; it is less likely to produce negative LD under the gene model

than the sites model. The theoretical results are discussed in relation to patterns of LD in

natural populations of several species.

Introduction

There is much interest in the effects of deleterious mutations on properties of populations

such as their genetic load, inbreeding depression, and variance in fitness, tracing back to the

pioneering work of Haldane (1937), Muller (1950), Morton et al. (1956) and Crow (1958),

and to more recent studies of the genetics of *Drosophila* fitness components (reviewed by

Crow 1993; Charlesworth 2015). In addition, there have been a number of recent theoretical

studies of measures of linkage disequilibrium (LD) among deleterious mutations (e.g., Garcia

and Lohmueller 2021; Roze 2021; Good 2022; Ragsdale 2022; Lyulina et al. 2004),

stimulated by population genomic studies where the sign of LD among rare, putatively

deleterious variants has been used to make inferences concerning features such as epistatic

fitness interactions and Hill-Robertson interference (Pool et al. 2012; Sohail et al. 2017;

Sandler et al. 2021; Garcia and Lohmueller 2021; Ragsdale 2022; Stolyarova et al. 2022).

The theoretical predictions are mostly based on population genetics models of diploid

organisms that tacitly assume that nucleotide sites subject to mutation and selection can be

treated as independent of each other with respect to their fitness effects, and of whether they

are in the same or different genes (reviewed by Kyriazis et al. 2023). Models of this type

were first introduced by Haldane (1937), who assumed that fitnesses were multiplicative

across loci. Consider individuals that are heterozygous at two different nucleotide sites.

Assume that the fitnesses of the two types of double heterozygotes are the same, regardless of

whether or not the two mutations are carried on the same haplotype (*cis*), or on the maternal

and paternal haplotypes (*trans*) – the "sites" model (see Figure 1A). With the same selection

coefficients against homozygotes (*s*) and dominance coefficients (*h*) at each site, the fitnesses

of these individuals are both $(1 - hs)^2$ if effects are multiplicative across sites and $1 - 2hs$ if

effects are additive. If selection is sufficiently weak that $s^2 << s$, as is assumed here, the

additive and multiplicative models are nearly equivalent. For mathematical convenience, we

will use the additive model from now on.

In contrast, if the gene is treated as a single entity that is subject to mutation, and there

is no allelic complementation between mutations within the same gene (more accurately, a

cistron as defined by Benzer [1957]), the fitness of an individual that is heterozygous for two

mutations depends on whether or not they are in *cis* or *trans* (Figure 1B) – the "gene model".

The fact that double heterozygotes with mutations in *cis* under the gene model have higher

fitnesses than those with mutations in *trans* implies an intragenic position effect with respect

to fitness when $h < \frac{1}{2}$. Other things being equal, this is likely to favor positive linkage

disequilibrium (LD), if the sign of the coefficient of linkage disequilibrium (*D*) is defined as

positive when the frequencies of haplotypes carrying pairs of deleterious mutant alleles and

pairs of wild-type alleles are in excess of random expectation. This is because the lower

fitness of *trans* compared with *cis* double heterozygotes when $h < \frac{1}{2}$ means that + + and – –

haplotypes will be transmitted more frequently to the next generation relative to + – and – +

haplotypes than when the two genotypes have the same fitness.

80   Position effects of this kind have rarely been included in population genetics models,

with the exception of a study of two-locus polymorphisms by Nordborg et al. (1995), and the

models of mutation and selection at multiple sites within a gene by Clark (1998), Thornton et

al. (2013) and Sanjak et al. (2015), and at two sites by Ragsdale (2022), who all used

somewhat different fitness models from ours. Such gene-based fitness models are likely to

result in patterns of LD between deleterious mutations that are different from those expected

under the sites model (Ragsdale 2022).

87   The purpose of the present paper is to examine the consequences of the differences

between the gene and sites models for patterns of linkage disequilibrium (LD) among

variants at sites under selection, using a model of a single coding sequence subject to

deleterious mutations. We do not attempt to present a completely general model of a multi-

site system, but instead analyze a simple model that brings out the main differences between

the two different representations of mutational effects on fitness. An analytical treatment of a

model with two sites under selection in an infinite, randomly mating population is given first,

followed by stochastic simulations of a multi-site sequence that is intended to provide an

approximate description of a coding sequence. We show that the difference between the two

models has significant consequences for the patterns of LD among such mutations, with the

gene model leading to positive LD among pairs of deleterious variants under conditions

where the additive model is associated with zero or slightly negative LD. Interpretations of

such patterns in natural populations need to take into account the possibility that the gene

model is likely to be a more accurate representation of the dominance properties of

deleterious mutations. We also show that focusing on rare variants, as advocated by Garcia

and Lohmueller (2021) and Good (2022), is likely to produce estimates of positive LD

among deleterious variants in agreement with Good (2022); the opposite finding of Garcia

and Lohmueller (2021) is due to their use of Lewontin's D' statistic (Lewontin 1964), which

is biased towards positive values when rare variants are used. The consequences of the

differences between the two models for the genetic load, inbreeding load, and genetic

variance in fitness are described in the companion paper (Johri and Charlesworth 2025).

Methods

Multi-locus simulations

Forward simulations were performed using SLiM 4.0 (Haller and Messer 2023), with a

Wright-Fisher population of $N = 1000$ diploid individuals, a mutation rate of $u = 0.5 \times 10^{-5}$

per nucleotide site/generation, and a rate of crossing over of 10^{-5} per nucleotide

site/generation. A gene with 1000 selected sites was simulated, assuming a dinucleotide

model, *i.e*, there were two alleles at each site: a wildtype (denoted by +) and a selectively

deleterious mutant (denoted by –) allele. Note that the two alleles were equally likely to

mutate to each other, i.e., the + allele could mutate to the – allele at rate u and vice versa.

This model contrasts with similar previous studies (Garcia and Lohmueller 2021; Good 2022;

Ragsdale 2022), where mutation was unidirectional from + to –. This procedure has the

advantage that it allows a true stationary distribution of allele frequencies to be reached

(Wright 1931), in contrast to models with one-way mutation, where all sites will eventually

become fixed for mutant alleles. It thus provides a more realistic description of evolution

resulting from mutations involving base substitutions. With sufficiently strong selection ($N_e s$

$\gg 1$), where N_e is the effective population size and s is the selective disadvantage to a mutant

homozygote (see Figure 1), there is little effect of back-mutation.

Simulations were run either assuming the same selection coefficient at each site or a

distribution of fitness effects (DFE). In the first case, the selective disadvantage of mutant

homozygotes scaled by twice the population size was fixed at $\gamma = 2Ns = 2$, 20, 100, or

1000. When simulating a distribution of fitness effects (DFE), a gamma distribution was

assumed for most of the results presented here, with shape parameter $\alpha = 0.3$ and mean

scaled selection coefficients of $\bar{\gamma} = 2N\bar{s} = 2$, 20, 100, or 1000. In this case, the homozygous

selective disadvantage of a – allele at a particular site remained constant throughout a given

simulation, although it varied across simulation replicates. The dominance coefficient (h) of

the mutant allele (see Figure 1) was fixed at 0.0, 0.2, or 0.5. The fitness of a diploid

individual was calculated using either the gene or the sites model, as described below. All

sites (polymorphic and fixed) were included when calculating the fitnesses of individuals.

Simulations were run for $10N$ generations from an initial population of N identical diploid

individuals with no variation. For each scenario, 1000 replicates were simulated, and used to

determine the means, standard deviations and standard errors across replicates.

Relating the simulations to Drosophila melanogaster *populations*

These simulation parameters can be placed in the context of the *D. melanogaster* genome, for

which there are on average about 5 exons per gene with a mean length of ~300 bp (Misra et

al. 2002). As only about two-thirds of coding sites are nonsynonymous, there are expected to

be ~1000 (= $300 \times 5 \times 2/3$) selected sites per gene, corresponding to our simulations. Intron

lengths, and hence gene lengths, vary across the genome, with a mean and median length of 5

kb and 2 kb respectively; we have not included introns or synonymous sites in our

simulations. Assuming $N_e = 10^6$ for *D. melanogaster* and a sex-averaged rate of crossing over

of 10^{-8} per site/generation (Johri et al. 2020), $N_e G r \approx 20 - 50$ over a genic region, where G

is the mean length of a gene. With our simulation parameters, the 1000 selected sites have

$N_e G r = 10$ when rescaled to a population size of 10^6, and therefore approximately represents

a genic region of *D. melanogaster*. Silent site polymorphism data imply that $N_e u \approx 0.003 -$

0.005 in *D. melanogaster* East African populations (Johri et al. 2020), consistent with the

$N u = 0.005$ used in our simulations. This rescaling uses the principle that, when all

evolutionary forces are weak, the outcome of evolution is determined by the products of the

effective population size and the magnitudes of the deterministic forces (e.g., Ewens 2004,

Chap. 5).

Because our simulations were computationally very expensive, it was not feasible to

perform them with a large population size. Rescaling by a large scaling factor in forward-in-

time simulations can affect population genetic statistics (Dabi and Schrider 2024; Ferrari et

al. 2025); rescaling has, however, been found to produce accurate results in the absence of

recombination (Kaiser and Charlesworth 2010). As we only simulated short regions of 1000

selected sites, rescaling is likely to produce accurate results in our case as well, since the

frequency of crossing over between the most widely separated sites is nearly proportional to

the distance between them, so that the recombination rates can be scaled by population size

when using a fixed rate of crossing over per basepair. This would not be the case for large

distances between sites, for which double crossovers cause a non-linear relation between

distance and recombination rate. This means that multiplication of the rate of crossing over

per basepair by a given factor does not result in a proportional change in the frequency of

recombination. To test the effects of rescaling, additional simulations were conducted with a

larger population size of 5000 diploid individuals, while maintaining the same values of Nu

($u = 2 \times 10^{-6}$ per site/generation) and Nr ($r = 2 \times 10^{-6}$ per site/generation) as the

simulations above. Selection coefficients followed a gamma distribution with an expected

value of $2N\bar{s} = 100$ and a shape parameter of 0.3.

Simulations of human populations

A Wright-Fisher population of $N = 1000$ diploid individuals was simulated with a mutation

rate $u = 2.5 \times 10^{-7}$ per nucleotide site/generation and crossing over rate (r) of 2×10^{-7} per

nucleotide site/generation. This resulted in $Nu = 0.00025$ and $Nr = 0.0002$ in our simulations,

which are consistent with the estimates of these parameters for human populations: N_e of

20,000, mutation rate of 1.25×10^{-8} per site/generation (Kong et al. 2012), and

recombination rate of 1.0×10^{-8} per site/generation (Altshuler et al. 2005). Assuming that

there are 8 exons per gene on average (Johri et al. 2021), that each exon is about 350 bp, and

that 70% of all coding sites in humans are nonsynonymous (Kryukov et al. 2007), there

should be about 1960 selected sites in a gene. We therefore simulated 2000 selected sites. We

again ignored introns; because genes tend to be coldspots for recombination in many species

(e.g., Myers et al. 2010; Joseph et al. 2024), the presence of introns should not greatly affect

recombination rates within genes. The DFE was assumed to be a gamma distribution with

shape parameter $\alpha = 0.23$ and $\bar{\gamma} = 850$, as estimated by Eyre-Walker et al. (2006) and Kim et

al. (2017).

Modeling the fitnesses of individuals

As mentioned in the Introduction, two different models of main effects on fitness (the sites

and gene models) are used to calculate the fitnesses of diploid genotypes. The sites model

assumes that heterozygous mutations at all sites complement each other fully. The gene

model assumes a complete absence of complementation between heterozygous mutations

present in *trans*. The difference in fitness between *cis*- and *trans*-heterozygotes can be

regarded mathematically as a form of epistasis, although its biological basis is different from

the type of epistasis usually considered in population genetics models.

In both cases, additivity of the main effects across sites was assumed in most of our

numerical work, as was also done by Good (2022) and Ragsdale (2022), in contrast to the

multiplicative fitness model that is often used in multi-locus studies. The difference between

additive and multiplicative fitness models involves second-order terms in the selection

coefficients. For a single gene with a modest number of segregating sites, as simulated here,

there should be little difference between additive and multiplicative models, unless selection

is much stronger than is generally assumed here. As just described, our simulations use

rescaling of the deterministic parameters, so that their products with N_e match those for much

larger natural populations. The second-order terms in the selection coefficients involved in

multiplicative fitness models makes such rescaling problematical; this difficulty is avoided

when the additive model used.

In addition to the main effects of each site, a simple model of pairwise epistasis of

the standard type was used, in which a single parameter (ϵ) describes a pairwise interaction

effect on fitness. To avoid confusion with the effect of non-complementation under the gene

model, we will refer to this as either antagonistic or synergistic epistasis. The total number of

sites within a gene is denoted by G. The parameter ϵ is positive if there is synergistic

epistasis, so that double mutants reduce fitness by more than is expected from the effects of

each mutation on its own, and negative if there is antagonistic (diminishing returns) epistasis,

where double mutants behave in the opposite fashion (Crow 1970; Kondrashov 1988).

Analytical results are derived in sections 1-3 of the Appendix and in Supplementary

File S1 for the case of $G = 2$, with the same selection coefficient s at each site. In general, in

addition to a possible difference in fitness between *cis*- and *trans*-heterozygotes, there are

three possible epistatic coefficients that measure departure from additivity in a symmetric

fitness model, denoted by the coefficients e_1, e_2 and e_3 in Table 1. In order to avoid multiple

epistatic parameters, we assume that the magnitude of such epistasis scales linearly with the

sizes of the selective effects at each site with a constant of proportionality ϵ, so that the

fitness of a mutant double homozygote is $1 - 2s - 2\epsilon s$. The fitnesses of all possible genotypes

under the sites model are defined in the upper section of Table 1, where the assumption of

linear scaling and complete complementation of mutations at different sites in double

heterozygotes gives $e_3 = 2\epsilon s$, $e_1 = 2\epsilon h s$ and $e_2 = \epsilon(1 + h)s$. For the gene model, a double

heterozygote with mutations in *cis* has fitness $1 - 2hs - 2\epsilon hs$, whereas the absence of

complementation in a *trans* double heterozygote implies that its fitness is $1 - s$.

It is less obvious how to assign the fitness of a genotype for which one site is

homozygous for a mutation and the other is heterozygous, e.g., $+ -/- -$. The representation in

Table 1 assigns a fitness reduction of $(1 + k)(1 + \epsilon)s$ to this genotype. For most of our

numerical results, we assume that a lack of complementation implies that the effects of each

mutation on the fitness of this genotype combine additively, with the heterozygous site

contributing a fitness reduction of ½s and the homozygous site a reduction of s, so that $k =$

0.5. This giving a net additive effect of 1.5s and an epistatic effect of 1.5ϵs. With $e_3 = 2\epsilon s$, we

thus have $e_1 = \epsilon hs$ and $e_2 = 1.5\epsilon s$.

An extreme alternative is to assume that $k = h$, as for the sites model, so that the single

heterozygous site has the same effect as if the other site were wild-type. There are no data

available to distinguish between these alternatives. It is shown in section 3 of the Appendix

and section 5 of Supplementary File S1 that the second alternative is likely to give a

substantially larger positive LD measure for pairs of deleterious mutations, so that our use of

the additive assumption is conservative from the point of view of predicting positive LD.

Modeling fitness with multiple sites and no epistasis

For describing fitness (w) when there are G selected sites within a gene, as in the simulations,

consider the two haplotypes (1 and 2) of a diploid individual, where $w = 1$ when all sites are

wild-type. With the same selection coefficient at all sites, fitness for the gene model is

calculated as follows.

If there is at least one – allele in 1 or 2, but not both, we have:

254
$$w = 1 - T_1 hs \qquad (1a)$$

256 where T_1 is the total number of – alleles over all G sites.

257   If there is at least one – allele in both 1 and 2, we have

259
$$w = 1 - 0.5 T_2 s \qquad (1b)$$

261 where T_2 is the total number of – alleles present.

262   In the case of a distribution of fitness effects (DFE), with variable selection

263 coefficient s_i at each site, these equations become:

265
$$w = 1 - h \sum_{i=1}^{T_1} s_i \qquad (2a)$$

267
$$w = 1 - 0.5 \sum_{i=1}^{T_2} s_i \qquad (2b)$$

269   For the sites model with a fixed selection coefficient, we have:

271
$$w = 1 - T_3 hs - T_4 s \qquad (3)$$

273   where T_3 is the total number of heterozygous (+/–) sites and T_4 is the total number of mutant

274   homozygous (–/–) sites. With a DFE, Equation (3) becomes:

276
$$w = 1 - h \sum_{i=1}^{T_3} s_i - \sum_{j=1}^{T_4} s_j \qquad (4)$$

The gene model with multiple sites

For the gene model, an extension of the analytical treatment of the two sites model to

multiple sites subject to the same selection regime (in the absence of epistasis) is described in

section 4 of the Appendix. This is based on the assumptions that LD is sufficiently weak that

the distribution of numbers of mutations per haplotype follows a Poisson distribution, that the

number of selected sites G is large, and that selection is sufficiently strong that the

equilibrium frequencies of deleterious mutations at each site are very small. Since significant

levels of synergistic or antagonistic epistasis generate LD, we have not included epistasis in

this approach. We consider only the gene model, since LD for the sites model without

epistasis is expected to be negligible.

Modeling epistasis with multiple sites

All epistatic effects are assumed to be between pairs of sites that are either homozygous or

heterozygous for mutant alleles, *i.e*, there are no higher-order epistatic effects. Each pairwise

interaction involves the sum of a joint measure of the fitness effects of the alleles at the pair

of sites in question. The simplest model, which applies to both the sites and gene models, is

to assume that the epistatic contribution from a pair of homozygous sites i and j is $\epsilon(s_i + s_j)$;

this is summed over all pairs of homozygous sites, and the net fitness reduction of the

genotype where all sites are homozygotes for mutations is given by this term and the sum of

the s_i over all homozygous mutant sites.

For the sites model, a heterozygous site i can be considered as contributing a main

effect of hs_i to the fitness reduction of a genotype, and ϵhs_i to each of the interactions with

other sites. Its interaction with a homozygous site j thus contributes $\epsilon(hs_i + s_j)$ and its

interaction with another heterozygous site k contributes $\epsilon h(s_i + s_k)$. Similarly, a pair of

302 homozygous sites i and j contributes a fitness reduction of $\epsilon(s_i + s_j)$. The net epistatic

303 contribution is obtained by summing over all pairs of sites, and the net fitness reduction is

304 given by this sum plus the sum of the fitness reductions from each heterozygous and

305 homozygous mutant site. Note that for both the sites and the gene model, if the fitness of a

306 diploid individual was found to be negative in the simulations, it was changed to a value of

307 zero.

308   For the gene model, a heterozygous individual whose mutations are all in *cis* would

309 have an epistatic component given by the sum over all pairs of i and j of $\epsilon h(s_i + s_j)$. The net

310 fitness reduction is the sum of this quantity plus the sum of the hs_i over all the heterozygous

311 sites. For an individual who is homozygous for a mutation, or heterozygous in *trans,* at more

312 than one site, the sum of the epistatic contributions over all pairs of sites is obtained by

313 weighting ϵs_i for heterozygous sites by ½ instead of by h, and ϵs_i for homozygous sites by 1.

314 The net fitness reduction is given by the sum of this quantity and the sum of the main effects

315 across sites, weighting s_i at homozygous sites by 1 and heterozygous sites by ½. This is

316 slightly different from the symmetric two-site model of Table 1, where a heterozygote for

317 two mutations in *trans* behaves like a homozygote.

319 **Estimating the linkage disequilibrium statistics from the simulations**

320 Summaries of linkage disequilibrium (LD) between sites from simulated data were calculated

321 using the mean of the coefficient of linkage disequilibrium D across all pairs of segregating

322 sites:

323 $$D = p_{ab} - q_a q_b \qquad (5)$$

325 and the two normalized measures of LD:

326 $$\bar{\sigma}_d = \frac{\bar{D}}{\sqrt{\overline{p_a p_b q_a q_b}}} \qquad (6)$$

328 $$\bar{\sigma}_d^1 = \frac{\bar{D}}{\overline{p_a p_b q_a q_b}} \qquad (7)$$

330 where q_a is the selected (deleterious) or minor allele frequency at the first site, q_b is the

331 deleterious/minor allele frequency at the second site, $p_a = 1 - q_a$, $p_b = 1 - q_b$, and p_{ab} is

332 the frequency of the haplotype "*a b*" which corresponds to the presence of either the

333 deleterious or the minor alleles at both sites. The overbars represent the mean of the

respective quantities over all SNP pairs and simulation replicates. Here, the mean of D and

the mean of the product $p_a p_b q_a q_b$ was calculated across all SNPs for each simulated

replicate. For each replicate, a single value of $\bar{\sigma}_d$ and $\bar{\sigma}_d^1$ was obtained by taking the ratio of

the mean D and the mean product of allele frequencies. The means and SEs reported in tables

and figures are those across the 1000 replicate simulations. The fluctuations across replicates

in the values of the three statistics ($\bar{D}$, $\bar{\sigma}_d$ and $\bar{\sigma}_d^1$) will thus be somewhat different in

magnitude, so there is no guarantee that their overall means will have the same sign.

However, this problem did not greatly affect our results, because it only happened when $\bar{D}$

was not significantly different from 0.

$\bar{\sigma}_d$ corresponds approximately to the mean correlation coefficient between the two

loci (Ohta and Kimura 1969); the denominator of the statistic $\bar{\sigma}_d^1$, used by Good (2022), and

Ragsdale (2022) greatly magnifies the apparent magnitude of LD compared with $\bar{\sigma}_d$.

However, the denominator of $\bar{\sigma}_d^1$ (the mean crossproduct of the four allele frequencies at a

pair of segregating sites, rather than its square root) means that its range is not confined to

(0,1), in contrast to the correlation coefficient.

Note that in our case, a *deleterious* allele may not be equivalent to a *derived* allele,

since back mutations from – to + are possible. The distinction is important, since many

empirical and theoretical studies of LD have relied on classifying variants as derived versus

ancestral; neither of these are necessarily equivalent to *minor* alleles (for a discussion, see

Potapova and Kondrashov 2023) We have used combinations of deleterious alleles as the

basis for calculations of the direction of D; with sufficiently strong selection ($N_e s > 4$), there

will be little difference between these two classifications, since deleterious alleles are almost

always derived. We also show simulation results for LD statistics where combinations of

minor alleles were used as the criterion for positive LD, as has often been done in empirical

studies.

Two-locus simulations

We also investigated the effect of weighting LD statistics toward low frequency derived

variants, a procedure suggested by Garcia and Lohmueller (2021) and Good (2022). We used

the approach of Good (2022) for simulating a two-site haploid model without recombination.

This procedure assumes that a mutation from A_1 to A_2 at the first site (A) is present at a

frequency q, whose probability distribution is given by the standard formula for drift and

selection (Equations S1.18 of section 6 of Supplementary File S1). A second mutation, from

B_1 to B_2, was introduced at the second site, either in *cis* with A_1 (case 1, probability $1 - q$) or

in *cis* with A_2 (case 2, probability q). The fitness model for determining the post-selection

frequencies of the haplotypes in a given generation assumed that the relative fitnesses of

A_1B_1, A_1B_2, A_2B_1 and A_2B_2 are 1, $1 - s$, $1 - s$ and $1 - 2s(1 + \epsilon)$, respectively, where ϵ is a

measure of epistasis of the same form as that used in the two-site diploid model. As described

in section 6 of Supplementary File S1, multinomial sampling of N haplotypes was then

carried out each generation, until one of the two mutations was fixed or lost, and the mean

values over all generations and replicates of D, D^2 and the allele frequency cross-product $P =$

$p_A q_A p_B q_B$ (where subscripts A and B denote sites A and B) were obtained. The means of D, D^2

and P, denoted by $\bar{D}$, $\overline{D^2}$ and $\bar{P}$, were found by dividing the respective sums of these

statistics by the sum of the durations of the replicate simulations.

Data availability

The scripts used to determine the properties of two-site equilibrium populations, perform all

the simulations, and calculate statistics are provided at

https://github.com/paruljohri/Gene_vs_sites_model/tree/main.

Results

Deterministic results for two selected sites

Here we examine the properties of equilibria under mutation-selection balance at two

selected sites within a coding sequence. The fitness matrices in Table 1 are used for this

purpose. The mutation rate from + to − at both sites is u. The symmetry of the fitness models

means that the equilibrium allele frequencies are the same at both sites, so that the system can

be completely described by the frequency q of the − allele at each site and the coefficient of

linkage disequilibrium D between them. The correlation coefficient between the two selected

sites is then given by $R = D/q(1 - q)$. An algebraic treatment is given in sections 1-3 of the

Appendix and sections 1-5 of Supplementary File S1. We focus on the case when the

recombination frequency, r, is zero. This gives insights into the behavior of the system with

low frequencies of recombination, such as those between pairs of sites within a gene. The

complexity of the equilibrium equations, even when $r = 0$, means that exact results for the

equilibrium haplotype frequencies can only be obtained numerically, as described in section 1

of the Appendix.

399   It is, however, possible to obtain approximate analytical results for the sign and

400 magnitude of D and R at equilibrium without full solutions for the equilibrium haplotype

401 frequencies. As might be expected from results in the earlier literature (e.g., Felsenstein 1965;

402 Kimura and Maruyama 1966; Kondrashov 1988; Charlesworth 1990; Shnol and Kondrashov

403 1993; Roze 2021), a multiplicative version of the sites model with no synergistic or

404 antagonistic epistasis results in equilibria with $\widehat{D} = 0$ (where the carat indicates equilibrium).

405 In contrast, the additive version results in equilibria with negative values of $\widehat{D}$, reflecting the

406 weak synergistic epistasis on the multiplicative scale that is associated with additive fitnesses

407 (see sections 2 and 3 of File S1). Explicit approximate expressions for the equilibrium values

408 of $\widehat{D}$ and $\widehat{R}$ for the additive sites model with no epistasis and a low frequency of

409 recombination are given by Equations (A9a), (A9b) and (A9c); the effect of epistasis on $\widehat{R}$ is

410 described by Equation (A9d). Equations (A9) show that, if the ratio of recombination rate to

411 selection coefficient (r/s) is $<< \hat{q}$ or $<< h$, recombination has only a small effect on the

412 magnitude of D and R.

413   In contrast, for the additive gene model with no synergistic epistasis, Equation (A10)

414 shows that $\widehat{D}$ and $\widehat{R}$ are positive when h is sufficiently small relative to ½ and $k \leq$ ½, selection

415 is sufficiently weak, and $\hat{q}$ is sufficiently large. $\widehat{D}$ and $\widehat{R}$ become slightly negative when these

416 conditions are violated, even in the absence of synergistic epistasis, but are always positive

417 for the corresponding multiplicative model when $h \leq k \leq$ ½ (Equation A11). Equations (A10)

418 and (A11) show that a smaller value of k is associated with larger LD values; in the absence

419 of synergistic epistasis, the values of $\widehat{D}$ and $\widehat{R}$ with $k = h <$ ½ are approximately twice those

420 with $h < k =$ ½ , reflecting the weaker selection against – – haplotypes in the former case (see

421 Equation A3b). Unsurprisingly, with sufficiently strong synergistic epistasis, $\widehat{D}$ and $\widehat{R}$

422 become negative (and vice-versa for antagonistic epistasis).

423   These conclusions are borne out by the results of iterating the exact recursion

424 relations (Equations A1 and A2), which are shown in Figure 2. The upper panels in Figure 2

425 show the results for the sites model, and the lower panels those for the gene model with $k =$

426 ½. In both cases, $u = 0.0001$ and $s = 0.01$. These are clearly not realistic values of the

427 mutation and selection parameters but produce relatively large values of $\widehat{R}$, enabling the gene

428 and sites models to be compared more convincingly than when $\widehat{R}$ is very small. The left-hand

429 panels plot $\widehat{R}$ (in percent) against the epistasis coefficient, $\epsilon,$ for three different dominance

430 coefficients, h. The right-hand panels plot $\widehat{R}$ against h for three different values of ϵ. Since

431 antagonistic epistasis ($\epsilon < 0$) is well known to induce positive LD (Kondrashov 1988;

Charlesworth 1990; Shnol and Kondrashov 1993), we have focused here on synergistic

epistasis ($\epsilon > 0$), as well as the zero epistasis case. As is shown below, the general properties

of the deterministic case mirror those of the multi-site simulations, although the results are

quantitatively very different.

Several points of interest emerge. As expected from the theoretical results, the sites

model shows consistently negative but very small values of $\hat{R}$, which are very close to the

negative of one-half of the mutation rate ($- 5 \times 10^{-5}$) when $\epsilon = 0$, independently of h (the

difference from 0 is too small to be detectable in the figure, but can be seen in the results of

the exact iterations provided in Supplementary File S2). The dominance coefficient has larger

effects under the sites model, the larger ϵ; $\hat{R}$ declines almost linearly with ϵ, becoming

negative when ϵ is sufficiently large. In contrast, for the gene model with no epistasis, $\hat{R}$

remains positive until h becomes close to ½, decreasing monotonically as h increases. With

synergistic epistasis, the gene model is associated with larger values of $\hat{R}$ compared with the

sites model when $h < $ ½, and the difference is more marked, the smaller h, with small h being

associated with larger magnitudes of $\hat{R}$. In addition, with the strongest value of epistasis ($\epsilon =$

0.32), $\hat{R}$ first declines with h and then increases again as it approaches ½ but remains

negative. The approximations describe the general patterns quite well. In the absence of

epistasis, $\hat{R}$ for the multiplicative gene model is always positive for $h < 0.5$, with a maximal

value at $h = 0$, when the very small value of the denominator in Equation (A11) overcomes

the fact that the numerator is proportional to $\hat{q}^2$ and is thus very small.

Sets of results for other values of u and s are shown in Supplementary File S2. The

general features described above can be seen in these cases. However, as might be expected,

the smaller the mean deleterious allele frequency, the smaller the differences between the

gene and sites models, and the smaller the magnitudes of $\hat{D}$ and $\hat{R}$. While $\hat{D}$ and $\hat{R}$ are still

positive for the gene model in the absence of synergistic epistasis and sufficiently small hs,

under the sites model they become negative at much smaller ϵ and $\hat{q}$ values. Thus, in order to

determine whether there are any biologically important consequences of the features just

described, it is necessary to determine what happens with multiple selected sites, where larger

effects are to be expected.

Deterministic results for the gene model with multiple sites

As described in section 4 of the Appendix, the results from the two-site gene model without

epistasis can be extended to an arbitrary number of selected sites, all subject to the same

selection regime, using the assumptions described in the above section *The gene model with*

multiple sites. Equations (A19) provide approximate expressions for $\hat{D}$ and $\hat{R}$ for the additive

gene model without epistasis, given a value of $\hat{q}$, on the assumption that $\hat{q} \ll 1$ and that

number of selected sites (G) is large. Equations (A21) show how to determine $\hat{q}$ for given

values of G, u, s and h. The results were derived for the case $k = \frac{1}{2}$, since this was used in the

simulations. The conclusion that positive LD between pairs of deleterious mutations is

expected under the gene model, provided that $h < \frac{1}{2}$, is confirmed by this analysis. In contrast

to the two-site model, where $\hat{D}$ and $\hat{R}$ depend on $\hat{q}^3$ and $\hat{q}^2$ respectively (see Equation A11),

$\hat{D}$ is proportional to $\hat{q}^2 \mu e^{-\mu}\left(\frac{1}{2} - h\right)$ and $\hat{R}$ to $\hat{q} \mu e^{-\mu}\left(\frac{1}{2} - h\right)$ where $\mu = G\hat{q}$ is the mean

number of deleterious mutations per haplotype; this reflects the cumulative effects of LD

across multiple sites. The next section presents comparisons of the predictions of Equations

(A19) with LD statistics obtained in the simulations.

Simulation results for multiple sites

The analytical results presented above were supplemented with forward simulations of a

single gene with multiple selected sites, with an equal probability of mutations between the +

and – alleles and vice versa (see the section *Methods*, *Multi-locus simulations*). We first

describe the results with no synergistic or antagonistic epistasis.

Linkage disequilibrium with no epistasis

As discussed in the section *Deterministic results for two selected sites*, deleterious alleles are

more likely to be associated on the same haplotypes in the gene model than in the sites

model, resulting in larger positive mean D values for the gene model with suitable parameter

values, as can be seen from the results in Figure 2. Table 2 shows the results of simulations

without epistasis in which s was fixed, and LD statistics were calculated by assigning positive

values to cases when $++$ and $--$ haplotypes were in excess of random expectation (entries

under "selected allele"). In these examples, the gene model results in higher mean LD than

the sites model for closely linked sites whenever there are significantly non-zero values. The

difference between the two models increases with increasing recessivity and strength of

selection, with mean positive values of $\overline{D}$ and $\overline{\sigma}_d$ for the gene model versus negative values

with the sites model when $h < 0.5$ and $\gamma \geq 20$, except when $h = 0$ and $\gamma = 100$, when small but

significantly positive values of $\overline{D}$ and $\overline{\sigma}_d$ were found with the sites model, consistent with the

results of Roze (2021) for small h values. For $\gamma = 2$, however, negative values were found

with both models, consistent with the operation of Hill-Robertson interference (HRI) effects

generated by the interaction between drift and selection with multiple linked selected sites

(Hill and Robertson 1966; Garcia and Lohmueller 2021; Roze 2021), which are not allowed

for in the two-site deterministic model.

It is of interest to examine whether these results are consistent with the theoretical

predictions for the multi-site gene model outlined in the previous section. The simulation

results for $\gamma = 20$ and $\gamma = 100$ were used, since the results for $\gamma = 2$ must be heavily influenced

by drift, whereas the theoretical predictions are purely deterministic. A complication in

making this comparison is that the LD statistics for the simulations are presented for pairs of

segregating sites, since $D = 0$ for invariant sites and $\bar{\sigma}_d$ is defined only for segregating sites.

As expected from general theory on strongly selected deleterious mutations (e.g.

Charlesworth and Jain 2014), the proportion of sites that are segregating at a given time with

$\gamma = 20$ or 100 is very small (see Table S5 in File S3).

The deterministic results for mean D are expected to correspond to the expected value

of D over all pairs of sites. For this reason, the predictions for $\bar{D}$ in Table 2 were obtained by

dividing the predictions of $\hat{D}$ from Equation (A19a) by the squares of the mean proportions

of segregating sites for the appropriate values of γ and h, obtained from Table S5. The values

of $\hat{q}$ were obtained from Table 3 of Johri and Charlesworth (2025); these agree closely with

the predictions from Equations (A21) (see Johri and Charlesworth (2025).

The predicted value of $\bar{\sigma}_d$ for a given parameter set can be obtained by dividing the

predicted value of $\bar{D}$ by one half of the nucleotide site diversity at selected sites (π),

normalized by the proportion of segregating sites. The values of π for the parameters in

Table 2 were obtained as described by Jain and Charlesworth (2014), equating the selection

coefficient in their formulae to $s[1 + (\mu - 1)e^{-\mu}(1 - 2h)]$ in Equation (A20b). Overall,

the agreement between observed and predicted values is remarkably good, considering the

approximations involved, suggesting that the multi-site gene model provides an accurate

description of the population genetics of LD when selection is strong and G is large.

Similar results to those in Table 2 were found with a gamma distribution of selection

coefficients, with the gene model yielding positive values of $\bar{D}$ and $\bar{\sigma}_d$ with $h < 0.5$ and $\bar{\gamma} \geq$

100, as opposed to slightly negative (but not significantly non-zero) values for the sites model

(Figure 3 and Figure S1 in Supplementary File S3). The theoretical results for the two-sites

model predict that $\bar{D}$ and $\bar{\sigma}_d$ values are expected to be slightly negative for the sites model.

Consistent with this prediction, we observed small negative values when $h < ½$. These effects
could also be caused by HRI.

To test this possibility, we analyzed the results with a DFE for variants separated by a
greater-than-average distance (800-100 base pairs; Table S1 in Supplementary File S3). The
$\bar{D}$ values for selected variants under the sites model then approach zero, consistent with
reduced effects of HRI with higher frequencies of recombination, whereas the deterministic
theory suggests only a weak effect of recombination with the low values of r/s for pairs of
sites within a gene (see the previous section *Deterministic results for two selected sites*). In
contrast, for the gene model under a gamma distribution with $\bar{\gamma} = 1000$, $\bar{D}$ and $\bar{\sigma}_d$ are still
substantially greater than zero, suggesting significantly positive LD even at larger distances
(Table S1 in Supplementary File S3). In addition, the magnitude of negative LD is greatest
with $h = 0.5$, whereas the deterministic sites model predicts zero LD (see Figure 2).

Furthermore, as discussed in the previous section, the multiplicative fitness sites
model is not expected to generate any LD, in contrast to the additive model. When fitnesses
were assigned to be multiplicative in our multi-locus simulations, $\bar{D}$ and $\bar{\sigma}_d$ values were
sometimes significantly negative when sites were less than 100 bases apart, especially with h
$= 0$ (Table S2 in Supplementary File S3). For alleles between 800-1000 bases apart, $\bar{D}$ and $\bar{\sigma}_d$
values were close to zero in all cases (Table S2 in Supplementary File S3). This suggests that
HRI was probably partly responsible for the slightly negative values observed in our multi-
site simulations of the sites model.

Interestingly, when an excess frequency of haplotypes with pairs of minor alleles was
used as the criterion for positive D, there is not much difference in $\bar{D}$ and $\bar{\sigma}_d$ between the
gene and sites models when selection is weak, with mainly positive values for both (Table 2
and Figure 3). The sign of $\bar{D}$ is expected to be positive for minor alleles in cases where the
overall value of $\bar{D}$ for derived alleles is zero. This comes about because there is a reasonable
chance that, when derived variants in a pairwise comparison on different haplotypes, one of
the derived alleles at one site in a pairwise comparison is present at a frequency greater than
0.5 and is regarded as a major allele while the derived allele at the other site is regarded as a
minor allele, so that D for the minor allele combination is positive. Taking the average over
all possible situations, this results in a bias toward positive LD among minor alleles
(Potapova and Kondrashov 2023). However, with stronger selection against – alleles and $h <$
½, the gene model yields slightly lower values of $\bar{D}$ and $\bar{\sigma}_d$, possibly due to the lower allele

frequencies under the gene model compared with the sites model (see Johri and Charlesworth

2025).

564   Our forward-in-time simulations were conducted with small populations (1000

diploid individuals) that are a factor of 1000 smaller than the value of N_e commonly used for

Drosophila populations, so that the mutation rate, recombination rate, and selection

coefficients were a factor of 1000 larger than the natural population values (see the Methods

section). To check if rescaling biased our results, we conducted simulations with a much

larger population size (5000 individuals). LD statistics obtained using minor allele

frequencies were entirely unaffected by rescaling (Table S3 in Supplementary File S3).

rescaled population. The differences between the two N values in the means of the LD

statistics for pairs of deleterious variants were within the limits of sampling error, so that

there is no evidence that rescaling has a major systematic effect on these statistics.

574   In contrast to the simulations with *Drosophila*-like parameters, which we have

discussed above, simulations of human-like populations yielded high variances between

replicates and somewhat unclear patterns for the LD statistics (Table 3). Overall, however,

LD between selected sites is higher with the gene model than the sites model when $h < ½$. In

this case, we show $\bar{\sigma}_d^1$ as well as $\bar{\sigma}_d$, for comparison with the analyses of human data by

Ragsdale (2022) (see *Discussion, Patterns of linkage disequilibrium in population genetic*

data). The difference in LD between the two models is much clearer when using $\bar{\sigma}_d$ or $\bar{\sigma}_d^1$ as

opposed to $\bar{D}$, because $\bar{\sigma}_d$ and $\bar{\sigma}_d^1$ (partially) correct for differences in allele frequencies.

LD with epistasis

Similarly to the analytical results discussed earlier, there is a substantial decrease in $\bar{D}$ and $\bar{\sigma}_d$

(using the selected/derived allele criterion) with increasing strength of synergistic epistasis

for both models when h is close to 0.0, but only small changes when h is 0.2 or 0.5 (Figures 4

and Figure S2 in Supplementary File S3). $\bar{D}$ and $\bar{\sigma}_d$ between selected alleles remain

significantly higher with the gene model than the sites model (for which both statistics

become negative) with large ϵ. There seems to be no significant difference between the gene

and sites model with these parameters when $h = 0.2$ and ϵ is very small, consistent with their

large SEs relative to the means when $\bar{\gamma} = 100$ (see Figure 3 for the results with no epistasis,

and Figure S3 in Supplementary File S3 for a comparison of the distribution of the statistics

for the results in Figure 3 and Figure 4); $\bar{D}$ and $\bar{\sigma}_d$ are both not significantly different from

zero in this case. Importantly, however, with $h = 0.2$ and $\epsilon > 0$, $\bar{\sigma}_d$ is negative with the sites

model, but is close to zero with the gene model, so that the two models display differences in

LD even with epistasis.

597   When the same LD statistics are calculated with minor/major alleles as the sign

criterion, there is a substantial decrease in LD with an increase in the epistasis coefficient;

however, the values always remain positive and there is no significant difference in $\bar{\sigma}_d$

between the two fitness models (Figure 4), although $\bar{D}$ obtained using minor alleles is

consistently and slightly smaller for the gene than the sites model, possibly due to lower

mean allele frequencies at equilibrium (see Table 3 of Johri and Charlesworth 2025). In

summary, if synergistic epistasis were prevalent amongst deleterious alleles, LD between

selected/derived alleles will probably still be higher and positive with the gene model than

the sites model when $h < \frac{1}{2}$, whereas no difference is expected between LD statistics

calculated using minor alleles as the sign criterion.

607   With much lower rates of crossing over (0.1 × the standard rate used here), when

mutations are fully recessive, LD calculated using minor alleles is significantly higher with

the sites model than the gene model (Figures S4 and S5 in Supplementary File S3). The

difference between the two models is largest when there is no epistasis, but persists even

when the epistasis coefficient is 0.32. The predicted differences in LD between the gene and

the sites models using the derived and minor allele sign criteria can thus be entirely opposite,

depending on the degree of dominance. HRI in the absence of synergistic epistasis seems to

produce only a weak tendency towards negative LD when $h = \frac{1}{2}$, and for the sites but not the

gene model when $h = 0$ or 0.2.

Effects on LD statistics of weighting towards low allele frequencies

We now consider the potential effects of preferentially using rare variants on estimates of

LD, a procedure proposed by Garcia and Lohmueller (2021) and Good (2022). Use of such a

procedure could produce substantially different patterns from those described above. Garcia

and Lohmueller (2021) suggested that the effects of HRI on LD between deleterious

mutations would be more easily detected in samples from populations if doubleton variants

were used for estimating LD. This procedure might help to correct for the effect of allele

frequency on LD statistics; in addition, because rare variants are more likely to be deleterious

than common ones, larger negative D values might be expected for nonsynonymous than

synonymous variants when conditioning on doubletons. In their multi-locus simulations, they

indeed found negative LD between derived variants at sites subject to purifying selection,

both with $h = 0$ and $h = ½$, with LD being measured by Lewontin's D' statistic (Lewontin

1964). In contrast, Good (2022) proposed that multiplying allele frequencies by a function

that gives greater weights to low frequency variants should lead to positive expected D

values, for both neutral and deleterious variants. He validated this expectation by a

combination of simulations and analytical formulae, using a model of a haploid population;

this is equivalent to a diploid population with $h = ½$ in the absence of epistasis. However,

while the simulations performed by Garcia and Lohmueller (2021) involved multiple selected

sites, those of Good (2022) involved only two sites.

In order to reconcile these seemingly contradictory findings, we used the approach

described in the section *Methods: Two-locus simulations* and section 6 of Supplementary File

S1. This enables the means of D, D^2 and the crossproduct P of the four allele frequencies at

sites segregating for two mutations ($\bar{D}$, $\overline{D^2}$ and $\bar{P}$) to be found, yielding estimates of the

standardized LD statistics $\bar{\sigma}_d = \bar{D}/\sqrt{\bar{P}}$, $\bar{\sigma}_d^1 = \bar{D}/\bar{P}$ and $\sigma_d^2 = \overline{D^2}/\bar{P}$. σ_d^2 corresponds

approximately to the mean squared correlation coefficient between the two loci (Ohta and

Kimura 1969); the other two statistics are defined by Equations (6) and (7). These statistics

were estimated either with or without weighting. The weighting factor f is 0 if no weighting

is applied, and $f > 1$ if low frequency mutant alleles receive preferential weights (f is the

inverse of the factor used by Good (2002)).

For neutral variants, the unweighted case gave values of $\bar{D}$ and $\bar{\sigma}_d$ that were non-

significantly negative and close to zero: $\bar{D} = 0.75 \times 10^{-4} \pm 5.14 \times 10^{-4}$ and $\bar{\sigma}_d = 0.00144 \pm$

0.00713. The value of σ_d^2 was close to the theoretical value of 0.45 for the case of no

recombination under the infinite sites model (Ohta and Kimura 1971). In contrast, $\bar{\sigma}_d$ and $\bar{\sigma}_d^1$

with weighting ($f = 10$)were 0.0107 and 0.915, respectively. The values when doubletons

were sampled were $\bar{\sigma}_d = 0.133$ and $\sigma_d^1 = 13.5$.

Some representative results for $\bar{\sigma}_d$ with selection are shown in Figure 5, for two

different strengths of selection ($\gamma = 2Ns = 10$ and $\gamma = 2Ns = 50$) and varying amounts of

synergistic epistasis. With no epistasis and no weighting, $\bar{D}$ and $\bar{\sigma}_d$ were slightly negative, but

became increasingly negative as ϵ increased. The case of $\gamma = 50$ without epistasis gave no

evidence for significantly non-zero LD ($\bar{D} = -5.35 \times 10^{-7} \pm 4.16 \times 10^{-7}$; $\bar{\sigma}_d = -8.88 \times 10^{-5} \pm$

6.93×10^{-5}), implying that HRI effects with two sites are not sufficiently strong to overcome

any tendency for positive LD to be produced by purifying selection. Weighting causes the LD

statistics to take larger values, which are positive with large weighting factors towards rare

alleles ($f = 100$; see the details in section 7 of Supplementary File S1) even for the largest

value of ϵ (0.64). They remain positive over the whole range of ϵ if doubletons are sampled.

Further results are shown in Supplementary Files S4 and S5.

These results show that focussing on mutations at low frequencies results in positive

LD between derived variants, which can persist even if there is strong synergistic epistasis,

especially if doubletons are sampled. A heuristic argument that explains this effect is

provided in section 7 of Supplementary File S1. These considerations suggest that the use of

LD statistics estimated from low frequency variants is not helpful in diagnosing HRI effects

caused by purifying selection. On the contrary, it is likely to conceal the effects of HRI or

synergistic epistasis on LD between derived variants.

This raises the question of why Garcia and Lohmueller (2021) found consistently

negative LD in their multi-locus simulations of deleterious mutations when sampling

doubletons. It turns out that this was because they applied the D' statistic (Lewontin 1964) to

estimates of D for derived variants. D' is defined as $D/|D_{max}|$, where $|D_{max}|$ is the

maximum value of the absolute value of D conditioned on the allele frequencies. The basis

for this conclusion is described in section 5 of the Appendix and section 8 of Supplementary

File S1, where it is shown that purifying selection can cause D' at selected sites to be lower

than at neutral sites, reaching negative values even if their frequencies are similar and if

doubletons are used.

To avoid these sources of bias, one should use unweighted values of D or $\bar{\sigma}_d$ and

avoid using D'. It is, however, currently unclear how nonequilibrium scenarios like

population size changes might affect the signs of these statistics and whether unweighted

values of D would be the best to use in all cases.

684   Discussion

685   The role of complementation

A major finding of this work is the effect on LD statistics of the distinction between the gene

and sites models, which we discuss in detail below. In the accompanying paper (Johri and

Charlesworth 2025), we discuss the implications for population genetics statistics relevant to

the genetic load and inbreeding load caused by deleterious mutations. Our results raise the

question of how widely the gene model might apply to real organisms. The distinction

between the two models is based on the well-established finding of classical and molecular

genetics that two recessive loss-of-function mutations affecting the amino-acid sequence of

the same polypeptide usually fail to complement each other when present in the same cell in

694 *trans* in a double heterozygote, due to the fact that both copies of the polypeptide are non-

695 functional. In contrast, the same mutations present in *cis* in a double heterozygote result in

696 the cell containing one functional and one non-functional polypeptide, with near wild-type

697 performance (see Figure 1). This finding is of great historical importance, since it underlies

698 the recognition of the gene as a coding sequence whose sites could be separated by

699 recombination, and to Seymour Benzer's concept of the cistron as the functional genetic unit

700 (reviewed by Benzer 1957; Pontercorvo 1958; Hawley and Gilliland 2006).

701   If mutations within the same coding sequence fail to complement each other when in

702 *trans*, the assumptions of the gene model are satisfied. The sites model must usually apply to

703 mutations affecting different coding sequences, which necessarily complement each other

704 unless there are unusual types of epistasis between them, e.g., due to interactions between

705 mutations in different members of a multi-protein complex. Examples of such a failure of

706 complementation between non-allelic loss-of-function mutations are reviewed by Hawley and

707 Gilliland (2006) and Navarro-Quiles et al. (2024); these appear to be very rare compared with

708 complementation.

709   It should be noted, however, that there are exceptions to a lack of allelic

710 complementation (reviewed by Fincham et al. 1979, Chap. 4; Hawley and Gilliland 2006). In

711 such cases, loss-of-function mutations affecting the same polypeptide complement each

712 other, so that *trans* double heterozygotes show wild-type or near wild-type performance. In

713 many cases, some mutations within the same gene complement each other, whereas others do

714 not, allowing the construction of complementation maps (Fincham et al. 1979, Chap. 4).

715 Some mutations within the same gene may thus obey the gene model and others the sites

716 model. Allelic complementation has been best studied in fungi like *Neurospora*, and often

717 involves cases where multimers of the same polypeptide form part of the same protein

718 molecule (Fincham et al. 1979, pp. 348-353). In other cases, spatial interactions between

719 mutations in distinct functional subunits of the same polypeptide are involved (Fincham et al.

720 1979, p. 352; Eng et al. 2015). It is often found that interallelic complementation is

721 incomplete, when phenotypes such as enzyme activity are measured (Fincham et al. 1979, pp.

722 348-351), so that the distinction between the two models is somewhat blurred. Similarly,

723 while Chandler et al. (2017) found multiple instances of hypomorphic alleles that

724 complement each other phenotypically, the extent of complementation between two alleles

725 was dependent on the genetic background. For simplicity, we have not attempted to include

726 any of these complexities in our models.

It is, of course, less clear what happens to the heterozygous fitness effects of more

minor effect mutations in the same coding sequence when configured in *cis* or *trans*, and

there appears to be little relevant data. The simplest assumption is that the effects parallel

those of loss-of-function mutations and are recessive or partially recessive, but are additive in

their effects when in *trans*, as we have assumed here when setting k in Table 1 to ½. There

are, of course, many other possibilities, including the extreme alternative of setting $k = h$ to

represent the fitness effect of a heterozygous effect of a mutation at one site on a background

of homozygosity for a mutation at another site.

The "gene-based model" of Thornton et al. (2013) and Sanjak et al. (2015) is similar

to ours, in that it assumes that a *trans* heterozygote with n mutations in a gene has the same

fitness as a homozygote with n mutations. They did not, however, consider the position

effect, *i.e*, the effects of the difference in fitness between *trans* and *cis* heterozygotes. Clark

(1998) evaluated the effects of allelic complementation or non-complementation between

pairs of deleterious alleles on allele frequencies at a multiallelic locus, but did not examine

patterns of LD. Ragsdale (2022, Table S1) describes two versions of a two-site model. His

"simple dominance" model is identical with our sites model (allowing for the difference in

notation). His "gene-based" model is similar to our gene model, in that it assigns different

fitnesses to *cis* and *trans* double heterozygotes. However, it differs in several other respects,

e.g., by assigning the same fitness to single homozygotes, single heterozygotes, double

homozygotes, and *cis* double heterozygotes.

The situation with respect to mutations affecting the same non-coding regulatory

sequence such as a UTR, an enhancer or an insulator is somewhat unclear, although it seems

likely that mutations within the same cis-regulatory sequence should show similar *cis*/*trans*

behavior to mutations in the same polypeptide, fitting the gene model, while mutations in

different cis-regulators of the same gene should complement each other, and hence fit the

sites model. The phenomenon of transvection involves complementation between mutations

in different cis-regulatory sequences of the same gene when the two homologous

chromosomes or regions of homologs are paired in a cell nucleus, but failure of

complementation when they are unpaired (Lewis 1954; Duncan 2002; Mellert and Truman

2012; Galouzis and Prud'homme 2021); the normal situation in this case is for them to fit the

sites model. However, a recent study of hypomorphic regulatory mutations in two *D.*

melanogaster genes showed a predominance of failure to complement (Chandler et al. 2017).

The role of epistasis

The extent to which deleterious mutations with minor effects on fitness have epistatic effects rather than obeying an additive or multiplicative model for their combined effects, and the typical nature of such interactions, is still an unsettled issue with respect to both theoretical (Diaz-Colunga et al. 2023) and empirical (Bakerlee et al. 2022) questions. Attempts to characterize epistasis for fitness components among deleterious mutations in a variety of organisms have used analyses of the change in the means over generations of mutation accumulation lines or populations subject to inbreeding. These experiments have yielded somewhat conflicting results, although overall there seems to be a predominance of weak synergistic epistasis (reviewed by Charlesworth 1998; Fry 2004; de Visser et al. 2011; García Dominguez et al. 2019). A problem with interpreting these results is that data on fitness, or its components such as viability or fecundity, are inherently noisy and there may be statistical biases (due to regression towards the mean) that lead to an underestimation of synergistic epistasis among deleterious mutations (Berger and Postma 2014).

Microbial systems have allowed estimates of the fitness consequences of combining deletions of two or more genes, providing extensive information about epistasis among drastically deleterious mutations, using haploid genomes (e.g., Costanzo et al. 2016; Kuzmin et al. 2018). New technologies that allow the introduction of multiple mutations into the same coding sequence have opened a path towards characterizing epistasis between mutations at the intragenic level. This can be done either at the level of a specific biochemical function, such as resistance to an antibiotic (e.g., Bershtein et al. 2006), fluorescence intensity of the green fluorescent protein (e.g., Sarkisyan et al. 2016; Somermeyer et al. 2022), or by measurements of the fitness effects of mutant strains in microbial competition experiments (e.g., Bank et al. 2015, 2016; Puchta et al. 2016; Domingo et al. 2018; Bakerlee et al. 2022; Chen et al. 2023; Johnson et al. 2023; Papkou et al. 2023). Interestingly, although the stability of proteins is nearly linearly reduced by increasing numbers of mutations (De Pristo et al. 2005; Bershtein et al. 2006), the effect of mutations on function or fitness in the experiments just cited seems to show a predominance of synergistic epistasis among deleterious mutations, despite complexities such as the dependence of patterns of interaction between the same mutations in different environments (Bank et al. 2016) or between-species differences in fitness landscapes (Somermeyer et al. 2022).

There appear to be no reliable estimates of the parameters involved in quantifying epistasis (such as our ϵ) in organisms such as *Drosophila* or humans, making it difficult to evaluate the relevance of our model of epistasis to patterns of LD in natural populations. It is, however, possible that synergistic epistasis among deleterious mutations could reduce the

positive LD expected under the gene model to near zero, unless mutations are highly

recessive (see Figure 4). The extent to which the sign of LD between deleterious variants is

affected by epistasis versus HRI also remains an open question, but our simulations suggest

that synergistic epistasis is more likely to produce detectable negative LD than HRI in

genomic regions where recombination rates are not unusually low.

Linkage disequilibrium: theory

It has generally been accepted that multiplicative fitness models of multi-locus mutation and

selection in randomly mating, discrete generation populations predict an absence of linkage

disequilibrium when equilibrium has been reached in an infinite population (Haldane 1937;

Kimura and Maruyama 1966). This is confirmed by our analytical results for the two-site

version of the sites model with multiplicative fitnesses (section 2 of Supplementary File S1).

Previous theoretical studies have found that synergistic epistasis on a multiplicative scale

results in negative LD when a positive sign of D is assigned to haplotypes with pairs of

mutant or wild-type alleles, whereas antagonistic epistasis causes positive LD (Kondrashov

1982, 1988; Charlesworth 1990; Shnol and Kondrashov 1993). Furthermore, in discrete

generation finite populations, there is a general expectation that negative LD will be

generated by Hill-Robertson interference in the absence of epistasis on the multiplicative

fitness scale, provided that linkage is sufficiently tight and mutations are not highly recessive

(Hill and Robertson 1966; Felsenstein 1974; Garcia and Lohmueller 2021; Roze 2021),

although the opposite pattern can arise with sufficiently small values of h (Roze 2021;

Ragsdale 2002).

Since an absence of epistasis on a linear scale produces weakly synergistic epistasis

on a multiplicative scale, it might be expected that our deterministic two-site model would

result in small negative values of the two measures of LD, D and R, when the epistatic

parameter ϵ is equal to zero. The analytical results presented here confirm this expectation for

the sites model (see section 2 of the Appendix and Supplementary File S1, sections 2 and 3,

but the properties of the gene model are significantly different (section 3 of the Appendix and

section 4 of File S1). Both the analytical results and results of exact iterations

(Supplementary File S2) show that the additive sites model exhibits negative values of R that

are very close to one-half the mutation rate when $\epsilon = 0$, and much larger negative values

when $\epsilon > 0$.

In contrast, the correlation coefficient R is often positive under the additive gene

model when $h < \frac{1}{2}$, for sufficiently small values of ϵ. This reflects the fact that the higher

fitness of *cis* compared with *trans* double heterozygotes (see Table 1) produces an excess of

+ + and – – haplotypes compared with + – and – + haplotypes, as was mentioned in the

Introduction. However, the fitness of the – –/+ – and – –/– + genotypes, $1 - (1 + k)s$ in the

absence of epistasis, also influences the magnitude of LD, through the contribution of these

genotypes to the marginal fitness of the – – haplotype; a reduction in this marginal fitness

causes a reduction in the frequency of the – – haplotype, diminishing the value of D and R.

For this reason, our numerical results that assume $h \leq k = \frac{1}{2}$ are conservative with respect to

the magnitude of positive LD among pairs of deleterious mutations. In the absence of a

cis/*trans* effect, a modification of the sites model, with the fitnesses of – –/+ – and – –/– +

becoming $1 - (1 + k)s$ rather than $1 - (1 + h)s$, results in negative LD with $h < k \leq \frac{1}{2}$ (see

section 5 of Supplementary File S1). This model generates positive LD only under the

biologically implausible condition $k < h$, which would imply that a homozygote for – has a

higher fitness than a heterozygote. This finding emphasizes the fact that positive LD under

the deterministic two-site gene model arises from the *cis*/*trans* effect, unless epistasis is

antagonistic rather than synergistic.

Similarly, the finding of positive LD in simulations of the gene model without

epistasis, at least with $h < \frac{1}{2}$ and sufficiently strong selection (see *Simulation results for*

multiple sites; *linkage disequilibrium with no epistasis*), makes sense given the intragenic

position effect on fitness. The results in Table 2 show that the approximations for LD derived

in section 4 of the Appendix for the multi-site gene model (Equations A19 and A21) agree

well with the simulation results with $k = \frac{1}{2}$ when the selection regime is the same for all sites

and selection is strong relative to drift, so that this approach provides useful insights into

when to expect positive LD between variants within the same gene.

In the simulations of multiple mutations within the same gene, diploid genotypes with

cis and *trans* configurations of mutations are likely to be relatively abundant in a randomly

mating population, unless selection is very intense. For example, with a scaled selection

coefficient of $\gamma = 20$ and $h = 0.2$, the mean frequency of a deleterious mutation at a given site

under the gene model with $k = \frac{1}{2}$ is approximately 0.001 (see Table 3 of Johri and

Charlesworth 2025). The mean number of mutations per haplotype with 1000 selected sites in

a coding sequence is thus approximately 1. Since mean D values are small in this case (Table

2), a Poisson distribution should give a good approximation to the distribution of the number

of mutations per haplotype. This implies a frequency of approximately 0.13 of individuals

with two mutations in *cis* over a wild-type haplotype, 0.14 with two mutations in *trans,* and

0.06 with three or more mutations in *cis* over a wild-type haplotype; the frequencies of wild-

type homozygotes and heterozygotes for single mutations are only 0.14 and 0.28,

respectively. The frequency of individuals that are homozygous at one or more sites is

approximately 0.001 in this example, so they will have a negligible effect on the haplotype

dynamics compared with the heterozygous carriers of mutations, implying that the value of k

can play only a small role when the number of selected sites is large. Positive LD under the

multi-site gene model can thus arise from the *cis/trans* effect.

As mentioned above, positive LD has also been found for the sites model in recent

theoretical studies that included finite population effects. Roze (2021) studied a diploid finite

population model with two loci under a multiplicative version of the sites model, assuming γ

$= 2N_e s \gg 1$ but $s \ll 1$. He found small positive mean D with $h < 0.25$ in both his

approximate analytical treatment and in multi-locus simulations (see his Figure 1). Similarly,

the analysis of the additive sites model for a two-locus system in a finite population by

Ragsdale (2022) found positive mean values of D and σ_d^1 for sufficiently small values of h,

provided that γ was sufficiently large. We similarly found small positive mean values of D

and σ_d for the sites model when $\gamma=100$ and $h=0.0$ (Table 2). Ragsdale's model of an

intragenic position effect also generated positive mean LD. No such effect is expected in an

infinite population (see Figure 2).

However, for very small values of the recombination rate and with $h \ll \frac{1}{2}$, negative

LD was found for the sites model by both Roze (2021) and Ragsdale (2022), reflecting the

build-up of associative overdominance caused by the interaction between drift and selection,

such that a predominance of + − and − + haplotypes is generated when linkage is close (Zhao

and Charlesworth 2016; Becher et al. 2020; Gilbert et al. 2020; Abu-Awad and Waller 2023).

The interactions between selection and drift when there is dominance are thus rather

complex.

Our multi-locus simulations of the sites model with multiplicative rather than linear

fitnesses failed to generate significantly positive LD (Table S2 in Supplementary File S3),

when deleterious versus beneficial alleles were used for the sign criterion. The absence of

positive LD in our simulations may be due to the stronger effects of HRI in multi-site

compared with two-site models. The existence of positive LD among derived or selected

mutations discussed here should not be confused with the much more frequently observed

positive D when an excess frequency of haplotypes with pairs of minor alleles is used as the

sign criterion for D (e.g., Figure 3). As mentioned in the section *Simulation results for*

multiple sites: linkage disequilibrium with no epistasis, this effect is expected even for

neutral variants (Potapova and Kondrashov 2023).

Patterns of linkage disequilibrium in population genetic data

Previous theoretical and simulation studies have suggested that, while HRI and synergistic

epistasis can create negative LD between selected variants, processes such as antagonistic

epistasis, admixture and population bottlenecks could also result in positive LD (Sohail et al.

2017; Sandler et al. 2021; Garcia and Lohmueller 2021; Ragsdale 2022). Several recent

studies have evaluated the sign of LD among variants in coding sequences in natural

populations of *D. melanogaster*, humans, *Capsella grandiflora* and *Schizophyllum commune*

(Pool et al. 2012; Sohail et al. 2017; Sandler et al. 2021; Garcia and Lohmueller 2021;

Ragsdale 2022; Stolyarova et al. 2022). Garcia and Lohmueller (2021) found a larger

magnitude of negative LD among doubleton nonsynonymous variants compared with

synonymous variants in humans when using Lewontin's D' statistic (Lewontin 1964). As

described in the subsection *Effects of weighting towards low frequencies on LD statistics*, D'

is biased towards negative values when low frequency variants are used, and differences in

D' between neutral and selected sites do not necessarily reflect the effects of epistasis or HRI

alone.

Sohail et al. (2017) and Sandler et al. (2021) used estimates of LD based on the excess or

deficiency over the Poisson expectation of the variance in the numbers of minor alleles over a

set of defined genomic regions, which is proportional to the sum of pairwise D values over

the regions in question, whereas Stolyarova et al. (2022) used estimates of the correlation

coefficients between pairs of variants, polarizing with respect to minor versus major alleles.

Both of these methods are likely to give higher magnitudes of the LD statistic than when

combinations of derived alleles are used for determining the sign of LD (Potapova and

Kondrashov 2023). Ragsdale (2022) polarized LD using the ancestral/derived criterion; he

employed the σ_d^1 measure, whose denominator involves the cross products of the four variant

frequencies at a pair of diallelic sites (see Equation 7). This will tend to give *higher*

magnitudes for sites with lower frequencies, for a given correlation coefficient.

These methodological differences across studies make it hard to compare the results, but

nevertheless there are some patterns in common. Notably, in most cases the magnitude of

signed LD between synonymous sites was found to be significantly higher than that between

nonsynonymous variants, with the exception of the analysis of the hypervariable mushroom

species *S. commune* by Stolyarova et al. (2022), where the opposite pattern was observed –

see their Appendix 2, Figure 1.

There seems to be a preponderance of slightly negative values of LD between loss-of-

function (LOF) mutations (presumed to be the most harmful), but positive values between

pairs of synonymous and nonsynonymous variants. Because these patterns applied to loosely

linked variants in humans and *Drosophila*, and the extent of under-dispersion of loosely

linked loss-of-function alleles was found to be much higher than explained by simulations

that included the demographic history of humans, Sohail *et al.* (2017) proposed that

synergistic epistasis was responsible.

A possible explanation for a predominance of negative signed LD between loosely

linked LOF mutations is that these are kept at very low frequencies in the population. This

means that a finite sample from the population has a high probability of not containing the –

– haplotype at a pair of diallelic sites. This automatically generates $D < 0$, even if D is zero in

the population as a whole, as would be expected for most pairs of sites that are not tightly

linked. With two sites with $D = 0$ in the population and a minor allele frequency q at each

site, a sample of size k has a probability of approximately $[1 - \exp(-kq)]^2$ of segregating at

both sites, and a probability of approximately kq^2 of containing the – – haplotype, giving a

net probability of $1 - kq^2/[1 - \exp(-kq)]^2$ that a segregating sample contains no – –

haplotypes. With $k = 1000$ and $q = 0.001$, this probability is equal to 0.9975 for 10000

replicate pairs. But with $q = 0.01$, it is only 0.9. Despite the fact that the expected value of D

is zero in both cases, with $q = 0.001$ a finite set of pairs of segregating sites is likely to have a

high probability of containing none or only a few – – haplotypes, resulting in a large

preponderance of cases with $D < 0$. This is illustrated in section of Supplementary File S1. It

is thus possible that observations of negative LD between loosely linked or independent LOF

mutations, such as those shown in Figure S6 of Sohail et al. (2017) are the result of this

sampling effect. It follows that it cannot safely be inferred that synergistic epistasis is the

cause of this pattern unless a very large number of pairs of variants is analysed; however, our

analytical and simulation results confirm that it does indeed generate negative LD for the

sites model but not necessarily for the gene model (see Figures 2 - 4). As noted by Ragsdale

(2022, p.10) only a few hundred pairs of within-gene LOF mutations were present in his

human population dataset.

Our modeling work adds a new perspective on findings of positive LD among closely

linked pairs of putatively selected variants. If fitness is governed by the gene model, then LD

between variants at non-synonymous sites within the same gene should be relatively larger

than under the sites model, and the sign should often be positive, provided the effects of

sampling just discussed do not overwhelm the effects of selection. However, as we have

discussed in the section *The role of complementation*, the gene model is unlikely to apply

between different genes but is likely to be applicable within a gene. Our models thus make

the following predictions about signed LD patterns in population genomics data. (*a*) LD

between derived or deleterious alleles at nonsynonymous sites within genes should be

positive or nearly zero (assuming that the population is panmictic and at equilibrium). (*b*) LD

between derived or deleterious nonsynonymous variants in the same gene should be higher

than between variants in different genes. (*c*) There should be higher LD between derived

nonsynonymous vs derived synonymous variants within the same gene.

There are, however, several caveats that need to be borne in mind when using these

findings to interpret data. First, the sign of LD statistics when calculated using minor/major

alleles (as opposed to using derived/ancestral) is almost always positive, unless the minor

alleles are very rare (Potapova and Kondrashov 2023). This applies to neutral sites (Potapova

and Kondrashov 2023) and to deleterious variants that obey the sites model as well as the

gene model – see Figure 3. It is thus not surprising that the studies which polarize their

variants by the minor/major allele criterion (Pool *et al*. 2012; Sohail et al. 2017; Sandler et al.

2021; Stolyarova et al. 2022) often found positive LD. Second, weighting towards low allele

frequencies (Good 2022) also tends to promote positive LD, as described in the subsection

Effects of weighting towards low frequencies on LD statistics, and should thus be avoided if

the sign of LD is used for inferences about selection. It follows that patterns of signed LD

using this criterion do not necessarily imply effects of admixture, non-equilibrium

demography, or antagonistic/complementary epistasis, which have all been proposed as

possible explanations in these previous studies.

This leaves the study of humans by Ragsdale (2022), who found that LD between

derived missense variants was found to be significantly positive within the same domain of a

gene, as opposed to near-zero values between SNPs in different domains, even after matching

the distance between SNPs found within compared between domains. This observation could

be explained by a gene model operating at the level of domains within genes, perhaps

because there is higher probability of complete failure of complementation within domains

versus across the entire gene. Ragsdale's estimates of $\bar{\sigma}_d^1$ from the Yoruba population (which

is closest to having a stable size) between missense variants within genes is ~0.02 (his Figure

8). As the average length of a gene in humans is quite large (~ 60 kb), it is difficult to

compare this estimate to that obtained in our simulations. Other estimates of $\bar{\sigma}_d^1$ within 100

base pairs provided by Ragsdale (2002) range from 0.25-0.3 (within the same gene; his

Figure S17), ~0.5 (within the same domain; his Figure S18), and ~0.2 (outside domains; his

Figure S19). While estimates of $\bar{\sigma}_d^1$ within 100 bases in our simulations of human-like

populations are within this range (Table S4 in Supplementary File S3), the large SEs make it

difficult to make any reliable conclusion. In general, however, both demographic processes

(like admixture and bottlenecks) as well as selective processes such as positive epistasis can

result in positive LD between derived nonsynonymous variants, and joint modeling of these

processes in a specific population would be required to distinguish between these scenarios,

taking into account important features such as recombination hotspots, which strongly

influence the rates of recombination within genes in taxa such as mammals and flowering

plant (e.g., Myers et al. 2010; Choi et al. 2013; Baudat et al. 2014; Sandler et al. 2021; Joseph

et al. 2024).

There are multiple reasons why within-gene LD should be larger in magnitude, and

more likely to be positive, than between-gene LD: (a) On average, there is a much higher

frequency of recombination between pairs of sites within different genes than within the same

gene (b) Fitness follows the gene model within genes but not between genes (c) There might

be antagonistic epistasis within genes but not between genes. While it might be possible to

differentiate between (a) and (b) by comparing the patterns of LD between species with

compact genes with little spacing and/or low recombination rates between genes with

patterns in species with high recombination rates between genes, it will be harder to

distinguish between (b) and (c).

In summary, it will probably be quite difficult to distinguish among the different

possible explanations for patterns of signed LD from population genomic data, but it is

essential to use the ancestral/derived or beneficial/deleterious criterion rather than the

minor/major allele criterion for the sign of LD. The problem is further complicated by the

fact that, if average gene lengths and rates of recombination are sufficiently large, even

patterns of LD within genes may largely reflect genome-wide patterns generated by

demographic processes rather than the effects of selection. Analyses of large-scale

resequencing data in the light of population genetic models of the type investigated here,

using suitable choices of species with respect to their genomic properties and demographic

histories, offer the best way to approach this problem.

1027

Evolution of the recombination rate within genes

Previous models of recombination in the presence of selection against recurrent deleterious mutations have shown that higher rates of recombination are likely to evolve if there is synergistic epistasis among mutations at different sites, and reduced rates to evolve if there is antagonistic epistasis (Feldman et al. 1980; Kondrashov 1982, 1988; Charlesworth 1990; Otto and Feldman 1997; Roze 2021). As we have shown, under the gene model, pairs of deleterious mutations result in a smaller fitness cost in *cis* than when they are in *trans*, promoting positive linkage disequilibrium. This suggests that recombination modifiers that increase the recombination rate within genes are likely to be disfavored, whereas modifiers that reduce them will be favored. This is consistent with the fact that recombination hotspots are often found in promoter regions or transcription start sites upstream of genes, and that genes are often relatively "cold" with respect to recombination rates (Myers et al. 2010; Joseph et al. 2024). Further modeling of this possible consequence of the *cis-trans* effect for the evolution of recombination rates would seem to be desirable.

Acknowledgments

We thank Kirk Lohmueller, Denis Roze and three anonymous reviewers for their helpful comments on an earlier version of the manuscript. This research was conducted using computational resources provided by ITS Research Computing at the University of North Carolina at Chapel Hill. PJ was funded by the National Institute of General Medical Sciences of the National Institutes of Health under the award number R35GM154969.

Literature Cited

Abu-Awad D, Waller DM. 2023. Conditions for maintaining and eroding pseudo-overdominance and its contribution to inbreeding depression. Peer Comm. J. 3:e8.

Altshuler D, Donnelly P, The International HapMap Consortium. 2005. A haplotype map of the human genome. Nature. 437:1299–1320.

Bakerlee CW, Nguyen Ba AN, Shulgina Y, Rojas Echenique JI, Desai MM. 2022. Idiosyncratic epistasis leads to global fitness-correlated trends. Science 376:630-635.

Bank C, Hietpas RT, Jensen JD, Bolon DN. 2015. A systematic survey of an intragenic epistatic landscape. Mol. Biol. Evol. 31:229-238.

Bank C, Matuszewski S, Hietpas RT, Jensen JD. 2016. On the (un)predictability of a large intragenic fitness landscape. Proc. Natl. Acad. Sci. USA 113:14085-14090.

Baudat F, Imai Y, De Massy B. 2014. Meiotic recombination in mammals: localization and regulation. Nat. Rev. Genet. 14:794-806.

Becher H, Jackson B, Charlesworth B. 2020. Patterns of genetic variability in genomic regions with low rates of recombination. Curr. Biol. 30:94-100.

Benzer S. 1957. The elementary units of heredity. In: McElroy WD, Glass B, editors. The Chemical Basis of Heredity. Baltimore, MD: Johns Hopkins Press. p. 70-93.

Berger D, Postma E. 2014. Biased estimates of diminishing-returns epistasis? Empirical evidence revisited. Genetics 198:1417-1420.

Bershtein S, Segal M, Bekerman S, Tokuriki N, Tawfik DS. 2006. Robustness–epistasis link shapes the fitness landscape of a randomly drifting protein. Nature 444:929-931.

Bürger, R. 2000. The mathematical theory of selection, recombination and mutation. Chichester UK, John Wiley & Sons.

Chandler CH, Chari S, Kowalaki A, Choi L, Tack D, DeNieu M, et al. 2017. Complex effects of genetic background on expressivity, complementation, and ordering of allelic effects. PloS Genet. 13:e1007075.

Charlesworth B. 1990. Mutation-selection balance and the evolutionary advantage of sex and recombination. Genet Res. 55 :199-221.

Charlesworth B. 1998. The effect of synergistic epistasis on the inbreeding load. Genet. Res. 71:85-89.

Charlesworth B. 2015. Causes of natural variation in fitness: evidence from studies of Drosophila populations. Proc. Natl. Acad. Sci. USA 12:1662-1669.

Charlesworth B, Jain K. 2014. Purifying selection, drift, and reversible mutation with arbitrarily high mutation rates. Genetics 198:1587-1602.

Chen Y, Hu R, LI K, Zhang Y, Fu L, Zhang J, Si T. 2023. Deep mutational scanning of an oxygen-Independent fluorescent protein CreiLOV for comprehensive profiling of mutational and epistatic effects. ACS Synth. Biol. 12:1461-1473.

Choi K, Zhao X, Kelly KA, Venn O, Higgins JD, Yelina NE, Hardcastle TJ, Ziolkowski PA, Copenhaver GP, Franklin FCH, et al. 2013. Arabidopsis meiotic crossover hot spots overlap with H2A.Z nucleosomes at gene promoters. Nat Genet. 45(11):1327–1336.

Clark AG. 1998. Mutation-selection balance with multiple alleles. Genetica 102:41–47.

Costanzo M, VanderSluis B, Koch EN, Baryshnikova A, Pons C, Tan G, et al. 2016. A global genetic interaction network maps a wiring diagram of cellular function. Science 353:aaf1420-1421-1414.

Crow JF. 1958. Some possibilities for measuring selection intensities in man. Hum. Biol. 30:1-13.

Crow JF. 1970. Genetic loads and the cost of natural selection. In: Kojima K, editor. Mathematical Topics in Population Genetics. Berlin: Springer-Verlag. p. 128-177.

Crow JF. 1993. Mutation, mean fitness, and genetic load. Oxf. Surv. Evol. Biol. 9:3-42.

Dabi A, Schrider DR. 2025. Population size rescaling significantly biases outcomes of forward-in-time population genetic simulations. Genetics 229:iyae180.

De Pristo MA, Weinreich DM, Hartl DL. 2005. Missense meandering in sequence space: a biophysical view of protein evolution. Nature Rev. Genet. 6:678-687.

Domingo J, Diss G, Lehner B. 2018. Pairwise and higher-order genetic interactions during the evolution of a tRNA. Nature 558:117-121.

De Visser JAGM, Cooper TF, Elena SF. 2011. The causes of epistasis. Proc. R. Soc. B. 278:3617-3624.

Diaz-Colunga J, Skwara A, Gowda K, Diaz-Uriarte R, Tikhonov M, Bajic D, Sanchez A. 2023. Global epistasis on fitness landscapes. Phil. Trans. R. Soc. B. 378:20220053.

Duncan IW. 2002. Transvection effects in Drosophila. Annu. Rev. Genet. 36:521-556.

Eng T, Guacci V, Koshland D. 2015. Interallelic complementation provides functional evidence for cohesin–cohesin interactions on DNA. Mol. Biol. Cell 26:4224-4235.

Ewens WJ. 2004. Mathematical Population Genetics. 1. Theoretical Introduction. New York: Springer.

Eyre-Walker A, Woolfit M, Phelps T. 2006. The distribution of fitness effects of new deleterious amino acid mutations in humans. Genetics. 173:891–900.

Feldman MW, Christiansen FB, Brooks LD. 1980. Evolution of recombination in a constant environment. Proc Natl Acad Sci U S A. 77:4838–4841.

Felsenstein J. 1965. The effect of linkage on directional selection. Genetics 52:349-63.

Felsenstein J. 1974. The evolutionary advantage of recombination. Genetics 78:737-756.

Ferrari T, Feng S, Zhang X, Mooney J. 2025. Parameter scaling in population genetics simulations may produce unintended background selection: Considerations for scaled simulation design. Genome Biol. Evol. 17:evaf097.

Fincham JRS, Day PR, Radford A. 1979. Fungal Genetics. Berkeley, CA: University of California Press.

Fry JD. 2004. On the rate and linearity of viability declines in Drosophila mutation-accumulation experiments. Genetics 166:797-806.

Galouzis CC, Prud'homme B. 2021. Transvection regulates the sex-biased expression of a fly X-linked gene. Science 371:396-400.

Garcia JA, Lohmueller KE. 2021. Negative linkage disequilibrium between amino acid changing variants reveals interference among deleterious mutations in the human genome. PloS Genet. 17:e1009676.

García-Dominguez S, García C, Quesada H, Caballero A. 2019. Accelerated inbreeding depression suggests synergistic epistasis for deleterious mutations in *Drosophila melanogaster*. Heredity 123:709-722.

Gilbert KJ, Pouyet F, Excoffier L, Peischl S. 2020. Transition from background selection to associative overdominance promotes diversity in regions of low recombination. Curr. Biol. 30:101-107.

Good BH. 2022. Linkage disequilibrium between rare mutations. Genetics 220:iyac004.

Haldane JBS. 1937. The effect of variation on fitness. Am. Nat. 71:337-349.

Haldane JBS. 1927. A mathematical theory of natural and artificial selection. Part V. Selection and mutation. Proc. Camb. Phil. Soc. 23:838-844.

Haller, BC, Messer, PW. 2023 SLiM 4: Multispecies eco-evolutionary modeling. Am. Nat. 201: E127–E139.

Hawley RS, Gilliland WD. 2006. Sometimes the result Is not the answer: The truths and the lies that come from using the complementation test. Genetics 174:5-15.

Hill WG, Robertson A. 1966. The effect of linkage on limits to artificial selection. Genet. Res. 8:269-294.

Johri P, Charlesworth B. 2025. A gene-based model of fitness and its implications for genetic variation: Genetic and inbreeding loads. (*companion paper*)

Johri P, Charlesworth B, Jensen JD. 2020. Toward an evolutionarily appropriate null model: Jointly inferring demography and purifying selection. Genetics 215:173-192.

Johri P, Riall K, Becher H, Excoffier L, Charlesworth B, Jensen JD. 2021. The impact of purifying and background selection on the inference of population history: problems and prospects. Mol Biol Evol. 38:2986–3003.

Johnson MS, Reddy G, Desai MM. 2023. Epistasis and evolution: recent advances and an outlook for prediction. BMC Biol. 21:120.

Joseph J, Prentout D, Laverré A, Duret L. 2024. High prevalence of PRDM9-independent recombination hotspots in placental mammals. Proc. Natl. Acad. Sci. USA 121:e2401973121.

Kaiser VB, Charlesworth B. 2010. Muller's ratchet and the degeneration of the *Drosophila miranda* neo-Y chromosome. Genetics 185:339-348.

Karlin S. 1975. General two-locus selection models: some objectives, results and interpretations. Theor. Pop. Biol. 7:364-398.

1216

1217 Kim BY, Huber CD, Lohmueller KE. 2017. Inference of the distribution of selection coefficients for
1218　　 new nonsynonymous mutations using large samples. Genetics. 206:345–361.

1219

1220 Kimura M, Maruyama T. 1966. The mutational load with epistatic gene interactions in fitness.
1221　　 Genetics 54:1303-1312.

1222

1223 Kondrashov AS. 1982. Selection against harmful mutations in large sexual and asexual populations.
1224　　 Genet. Res 44:325-332.

1225

1226 Kondrashov AS. 1988. Deleterious mutations and the evolution of sexual reproduction. Nature
1227　　 336:435-440.

1228

1229 Kong A, Frigge ML, Masson G, Besenbacher S, Sulem P, Magnusson G, Gudjonsson SA,
1230　　 Sigurdsson A, Jonasdottir Aslaug, Jonasdottir Adalbjorg, et al. 2012. Rate of *de novo* mutations
1231　　 and the importance of father's age to disease risk. Nature. 488(7412):471–475.

1232

1233 Kryukov GV, Pennachio LA, Sunyaev S. 2007. Most rare missense alleles are deleterious in humans:
1234　　 implications for complex disease and association studies. Am. J. Hum. Genet. 80:727-739.

1235

1236 Kuzmin E, VanderSluis B, Wang W, Tan G, Despande R, Cheng Y, et al. 2018. Systematic analysis
1237　　 of complex genetic interactions. Science 360:eaao1729.

1238

1239 Kyriazis CC, Robinson JA, Lohmueller KE. 2023. Using computational simulations to model
1240　　 deleterious variation and genetic load in natural populations. Am. Nat. 302:737-752.

1241

1242 Lewis EB. 1954. The theory and application of a new method of detecting chromosomal
1243　　 rearrangements in *Drosophila melanogaste*r. Am. Nat. 88:225-239.

1244

1245 Lewontin RC. 1964. The interaction of selection and linkage. I. General considerations, heterotic
1246　　 models. Genetics 49:49-67.

1247

1248 Lyulina AS, Liu Z, Good BH. 2024. Linkage equilibrium between rare mutations. Genetics
1249　　 228:iyae145.

1250

1251 Mellert DJ, Truman JW. 2012. Transvection is common throughout the Drosophila genome.
1252　　 Genetics 191:1121-1141.

1253

1254 Misra S, Crosby MA, Mungall CJ, Matthews BB, Campbell KS, et al. 2002. Annotation of the
1255　　 *Drosophila melanogaster* euchromatic genome: a systematic review. Gen. Biol. 3:Research0083.

1256

1257 Morton NE, Crow JF, Muller HJ. 1956. An estimate of the mutational damage in man from data on
1258　　 consanguineous marriages. Proc. Natl. Acad. Sci. USA 42:855-863.

1259

1260 Muller HJ. 1950. Our load of mutations. Am. J. Human Genet. 2:111-176.

1261

1262 Myers S, Bottolo L, Freeman C, McVean G, Donnelly G. 2010. A fine-scale map of recombination
1263　　 rates and hotspots across the human genome. Science 310:321-324.

1264

Navarro-Quiles C, Lup SD, Munoz-Nortes T, Candela H, Micol JL. 2024. The genetic and molecular basis of haploinsufficiency in flowering plants. Trends. Plant Sci. 29:72-95.

Nordborg M, Franklin IR, Feldman MW. 1995. Effects of cis-trans viability selection on some two-locus models. Theor. Pop. Biol. 47:365-392.

Ohta T, Kimura M. 1969. Linkage disequilibrium due to random genetic drift. Genet. Res. 13:47-55.

Ohta T, Kimura M. 1971. Linkage disequilibrium between two segregating nucleotide sites under steady flux of mutations in a finite population. Genetics 68:571-580.

Otto SP, Feldman MW. 1997. Deleterious mutations, variable epistatic interactions, and the evolution of recombination. Theor Popul Biol. 51:134–147.

Papkou A, Garcia-Pastor L, Escudero JA, Wagner A. 2023. A rugged yet easily navigable fitness landscape. Science 382:eadh3860.

Pontecorvo G. 1958. Trends in Genetic Analysis. New York, NY: Columbia University Press.

Pool JE, Corbett-Detig RB, Sugino RP, Stevens KA, Cardeno CM, Crepeau MW, et al. 2012. Population genomics of sub-Saharan *Drosophila melanogaster*: African diversity and non-African admixture. PloS Genet 8:e1003080.

Potapova NA, Kondrashov AS. 2023. Positive association between alleles at selectively neutral loci. Biorxiv 2023.06.14.544962:1-12.

Puchta O, Czeke B, Czaja H, D. T, Sanguinetti G, Kudla G. 2016. Network of epistatic interactions within a yeast snoRNA. Science 352:840-844.

Ragsdale AP. 2022. Local fitness and epistatic effects lead to distinct patterns of linkage disequilibrium in protein-coding genes. Genetics 221:iyac097.

Roze D. 2021. A simple expression for the strength of selection on recombination generated by interference among mutations. Proc. Natl. Acad. Sci. USA 118:e2022805118.

Sandler G, Wright SI, Agrawal AF. 2021. Patterns and causes of signed linkage disequilibria in flies and plants. Mol. Biol. Evol. 38:4310-4321.

Sanjak JS, Long AD, Thornton KR. 2015. A model of compound heterozygous, loss-of-function alleles Is broadly consistent with observations from complex-disease GWAS datasets. PloS Genet. 13:e1006573.

Sarkisyan KS, Bolotin DA, Meer MV, Usmanova DR, Mishin AS, Sharanov GV, et. al. 2016. Local fitness landscape of the green fluorescent protein. Nature 533:397-401.

Shnol EE, Kondrashov AS. 1993. Effect of selection on the phenotypic variance. Genetics 134:995-996.

1313 Sohail M, Vakhrusheva OA, Sul JH, Pulit SA, Francioli LC, Genome of the Netherlands
1314   Consortium, et al. 2017. Negative selection in humans and fruit flies involves synergistic
1315   epistasis. Science 356:539-542.
1316
1317 Somermeyer LG, Fleiss A, Mishin AS, Bozhanova NG, Igolkina AA, Meiler J, Pujol M-EA,
1318   Putintseva EV, Sarkisyan KS, Kondrashov FA. 2022. Heterogeneity of the GFP fitness landscape
1319   and data-driven protein design. Elife 11:e75842.
1320
1321 Stolyarova AV, Neretina TV, Zvyagina EA, Fedotova AV, Kondrashov AS, Bazykin GA. 2022.
1322   Complex fitness landscape shapes variation in a hyperpolymorphic species. Elife 11:e76073.
1323
1324 Thornton KR, Foran AJ, Long AD. 2013. Properties and modeling of GWAS when complex disease
1325   risk is due to non-complementing, deleterious mutations in genes of large effect. PloS Genet.
1326   9:e1003258.
1327
1328 Wright S. 1931. Evolution in Mendelian populations. Genetics 16:97-159.
1329
1330 Zhao L, Charlesworth B. 2016. Resolving the conflict between associative overdominance and
1331   background selection. Genetics 203:1315-1334.
1332
1333

Appendix

1. Deterministic treatment of the sites and gene models: recursion equations for the

haplotype frequencies

The full recursion relations for a general fitness matrix and mutation model are given below; these

differ slightly in form from those usually used for models of mutation and selection, where the

changes in haplotype frequencies due to selection and mutation are added to each other (e.g.,

Haldane [1937]). This is because the small equilibrium values of D resulting from the interaction

between mutation and selection are very sensitive to approximations, and incorrect signs of D can

result if approximations are made prematurely.

Assume that there are two diallelic loci, A and B, with alleles A_1, A_2 and B_1, B_2, with

subscript 1 corresponding to + in Table 1 and subscript 2 to –. Let the frequencies of the four

haplotypes A_1B_1, A_1B_2, A_2B_1 and A_2B_2 be x_1, x_2, x_3 and x_4, respectively. The corresponding allele

frequencies are p_1, q_1 (A_1 and A_2) and p_2, q_2 (B_1 and B_2). The coefficient of linkage disequilibrium

is $D = x_1 x_4 - x_2 x_3$. Assume that A_2 and B_2 are both deleterious and are kept at such low frequencies

that mutations can be treated as unidirectional, with mutation rates u_1 and u_2 for A_1 to A_2 and B_1 to

B_2, respectively. Let w_{ij} be the fitness of individuals formed by a pair of haplotypes (i and j) with

frequencies x_i and x_j. Under the gene model, the fitnesses of the two double heterozygotes (w_{14} and

w_{23}) are unequal, in contrast to the usual assumptions of two-locus models (e.g., Karlin 1975). The

marginal fitness of haplotype is $w_i = \sum_j x_j w_{ij}$ and the population mean fitness is $\bar{w} = \sum_i w_i$.

Recombination occurs between the two loci with frequency r per generation. Random mating is

assumed.

Assume that the order of events within a generation is selection, mutation and

recombination. The post-selection frequency of haplotype i is $x_i^* = x_i w_i / \overline{w}$. Mutation changes the

haplotype frequencies to new values x_i^{**} as follows:

1358

$$x_1^{**} = x_1^*(1 - u_1)(1 - u_2), \qquad x_2^{**} = (x_2^* + x_1^* u_2)(1 - u_1) \qquad (A1a)$$

1359

$$x_3^{**} = (x_3^* + x_1^* u_1)(1 - u_2), \qquad x_4^{**} = x_4^* + x_1^* u_1 u_2 + x_2^* u_1 + x_3^* u_2 \qquad (A1b)$$

1360

1361

In the absence of recombination, these are the haplotype frequencies in the new generation.

If recombination occurs post-mutation, the haplotype frequencies in the new generation are given

by the following expressions, where the triple asterisks indicate haplotype frequencies after

mutation without any selection within the generation in question (*i.e*, the frequencies given by

Equations (A1) without the asterisks on the right-hand side); the effects of selection on the double

heterozygotes are captured by the terms in w_{14} and w_{23}:

1368

$$x_1' = x_1^{**} - r(x_1^{***} x_4^{***} w_{14} - x_2^{***} x_3^{***} w_{23}) \overline{w}^{-1} \qquad (A2a)$$

1369

$$x_2' = x_2^{**} + r(x_1^{***} x_4^{***} w_{14} - x_2^{***} x_3^{***} w_{23}) \overline{w}^{-1} \qquad (A2b)$$

1370

$$x_3' = x_3^{**} + r(x_1^{***} x_4^{***} w_{14} - x_2^{***} x_3^{***} w_{23}) \overline{w}^{-1} \qquad (A2c)$$

1371

$$x_4' = x_4^{**} - r(x_1^{***} x_4^{***} w_{14} - x_2^{***} x_3^{***} w_{23}) \overline{w}^{-1} \qquad (A2d)$$

1372

1373

These recursion equations can be used to obtain the equilibrium values for the system,

as well as the approach to equilibrium. Given the symmetry of the fitness matrix in Table 1,

for the system studied here we can write $x_2 = x_3 = x = q(1 - q) - D$; $x_4 = y = q^2 + D$; $x_1 = z = 1$

$- 2x - y$; $q = x + y$. It follows that only the recursions for x and y need to be considered. Let

the mutation rate per locus from $+$ to $-$ be u. If $u \ll hs$, the frequency q of the $-$ allele at each

locus is $\ll 1$ and reverse mutations can be ignored. The coefficient of linkage disequilibrium

is denoted by $D = yz - x^2$. Together with the fitness values for each genotype, these variables

can be used to determine the recursion relations for the frequencies of the four haplotypes, x_1

$(+ +)$, $x_2 (- +)$, $x_3 (+ -)$ and $x_4 (- -)$.

Denote the marginal fitnesses of the $- +$ and $+ -$ haplotypes by w_x and that of the $- -$

haplotype by w_y; the marginal fitness of the $+ +$ haplotype is w_z. From Table 1, we obtain the

following expressions for the marginal fitnesses under the sites model:

1386

$$w_x = 1 - [h + q + 2h\epsilon x + (1 + h)\epsilon y]s \qquad (A3a)$$

$$w_y = 1 - 2[h + (1 - h)q + h\epsilon + (1 - h)\epsilon q]s \qquad (A3b)$$

$$w_z = 1 - 2[hq + h\epsilon y]s \qquad (A3c)$$

The equivalent expressions for the gene model are as follows:

$$w_x = 1 - [h + 2(1 - h)x + (1 + k - h)y + (1 + k)\epsilon y]s \qquad (A3d)$$

$$w_y = 1 - 2[h + (1 + k - 2h)x + (1 - h)y](1 + \epsilon)s \qquad (A3e)$$

$$w_z = 1 - 2hs(q + \epsilon y) \qquad (A3f)$$

For both models, the mean fitness of the population is given by:

$$\overline{w} = 2xw_x + yw_y + zw_z \qquad (A3g)$$

If these expressions are substituted into Equations (A1), using specified values of the mutation rate, recombination rates and selection parameters, a numerical solution to the equilibrium version of the recursions could be found by an iterative method. It is, however, simpler to run the recursions to close to equilibrium, thereby obtaining estimates of the equilibrium values of x and y and all derived quantities of interest. The initial starting point of the recursions was arbitrarily chosen to have $D = 0$, with q given by the solution of the linear equation that approximates the cubic equation describing mutation-selection equilibrium at a single locus when q is sufficiently small but non-negligible (Bürger 2000, p.100):

$$\hat{q} \approx \frac{u}{(u + hs)} \qquad (A4)$$

2. Properties of equilibria with respect to LD under the additive and multiplicative sites models

As shown in part 1 of Supplementary File S1, when $r = 0$, a formula for the equilibrium value of D, $\hat{D}$, can be obtained for a general symmetric fitness model (Equation S1.3). Neglecting the small terms in the mutation rate in this expression, we have:

$$\hat{D}[\overline{w}^2 - w_x^2] \approx -\lambda \hat{y}\hat{z} \qquad (A5a)$$

1419 where

$$\lambda = w_x^2 - w_y w_z \qquad \text{(A5b)}$$

1420

1421

1422   The analysis in section 2 of Supplementary File S1 shows that $\widehat{D} = 0$ under the
1423 multiplicative sites model with no epistasis (Equation S1.6b).

1424   Equations (S1.7) and (S1.8) in section 3 of Supplementary File S1 show that, for the
1425 additive sites model with no epistasis, we have:

1426

$$\lambda = \{h^2 + 2h(1 - 2h)q + [1 - 4h(1 - h)]q^2\}s^2 \qquad \text{(A5c)}$$

1428

1429   We can approximate q for an equilibrium population by equating it to the equilibrium
1430 solution $\hat{q}$ given by Equation (A4). We can also approximate the haplotype frequencies by
1431 neglecting the small terms involving $\widehat{D}$, so that $\hat{x} \approx \hat{q}(1 - \hat{q}), \hat{y} \approx \hat{q}^2, \hat{z} = (1 - \hat{q})^2$. Ignoring
1432 terms in $\hat{q}^2$, Equation (S10b) then gives the following approximation:

1433

$$\overline{w}^2 - w_x^2 \approx 2[h + (1 - 4h)\hat{q}]s \qquad \text{(A6)}$$

1435

1436 Equations (A5c) and (A.6) yield the following approximation for $\widehat{D}$:

1437

$$\widehat{D} \approx -\tfrac{1}{2}s\hat{q}^2(1 - \hat{q})^2\{h^2 + 2h(1 - 2h)\hat{q} + [1 - 4h(1 - h)]\hat{q}^2\}[h + (1 - 4h)\hat{q}]^{-1}$$

1439 $$\text{(A7)}$$

1440

1441   This expression can be extended to include the effect of recombination when r, D and
1442 s are sufficiently small that their second-order terms can be neglected. In this case, Equations
1443 (A2) imply that the change in D caused by recombination is approximately $-rD$. If this term
1444 is added to the right-hand side of the recursion equation for D (Equation S1.3), and $\overline{w}^2 D$ is
1445 approximated by D, Equation (A5a) becomes:

1446

$$\widehat{D}[\overline{w}^2 - w_x^2] \approx -(\lambda\hat{y}\hat{z} + r\widehat{D}) \qquad \text{(A8)}$$

1448

1449 This yields the following approximate expression for $\widehat{D}$:

1450

1451 $$\widehat{D} \approx -\tfrac{1}{2}s\hat{q}^2(1 - \hat{q})^2\{h^2 + 2h(1 - 2h)\hat{q} + [1 - 4h(1 - h)]\hat{q}^2\}\{h[1 - 2\hat{q}(1 - 2\hat{q})] + \hat{q}(1 - 2\hat{q}) + r/s\}^{-1}$$

1452 $\qquad$ (A9a)

1453 This equation can be converted into an expression for the equilibrium value of the correlation

1454 coefficient, $\hat{R}$, by dividing by $\hat{q}(1 - \hat{q})$:

1455

1456 $\qquad \hat{R} \approx -\frac{1}{2}s\hat{q}(1 - \hat{q})\{h^2 + 2h(1 - 2h)\hat{q} + [1 - 4h(1 - h)]\hat{q}^2\}[h + (1 - 4h)\hat{q} + r/s\}^{-1}$

1457 $\qquad$ (A9b)

1458 If second-order terms in $\hat{q}$ are neglected, this expression simplifies to:

1459

1460 $\qquad \hat{R} \approx -\frac{1}{2}s\hat{q}[h^2 + 2h(1 - 2h)\hat{q}][h + (1 - 4h)\hat{q} + r/s]^{-1}$ (A9c)

1461

$\qquad$ The magnitudes of $\hat{D}$ and $\hat{R}$ under the additive sites model are, however, very small in

the absence of epistasis, of the order of the mutation rate. This can be seen as follows.

Provided h is bounded away from 0, and $u << hs$, we have $\hat{q} \approx \frac{u}{hs} \ll 1$ (Haldane 1937). The

leading term in braces in the numerator of Equation (A9b) is h^2, and the leading term in

braces in the denominator is h, so that the magnitude of the ratio of numerator to denominator

in Equation (A9b) is approximately $\frac{1}{2}hs\hat{q} \approx \frac{1}{2}u$. If $h = 0$, $\hat{q} \approx \sqrt{u/s}$, so that the term in braces

in the numerator is approximately u/s and the term in braces in the denominator is

approximately $\hat{q}$. This cancels the $\hat{q}$ term in the numerator, so that (assuming $\hat{q} \ll 1$), the

magnitude of the ratio of numerator to denominator is approximately $\frac{1}{2}u$, as before. It follows

that, provided that $\hat{q} \ll 1$, $\hat{R}$ should be approximately equal to the negative of one-half of the

mutation rate, and independent of the dominance coefficient and strength of selection.

Provided that the epistasis coefficient ϵ is of the same order as s or greater, much larger

negative values of $\hat{D}$ and $\hat{R}$ are expected with synergistic epistasis ($\epsilon > 0$), and positive

values are expected with antagonistic epistasis ($\epsilon < 0$) – see Equation (A9c).

$\qquad$ Equations (A9) must be modified if epistasis is present, which introduces extra terms

of the order of ϵs. Use of Equations (A3) leads to the following expression for the additive

sites model when $\epsilon > s$:

1479

1480 $\qquad \hat{R} \approx -\epsilon\hat{q}(1 - \hat{q})[h + (1 - 3h)\hat{q}][h + (1 - 4h)\hat{q} + 2h\hat{q}\epsilon + r/s]^{-1}$ (A9d)

1481

1482 **3. Properties of equilibria with respect to LD under the additive and multiplicative gene**

1483 **models**

1484   The same type of approach can be applied to the additive and multiplicative gene models, as

1485   described in section 4 of Supplementary File S1. Equation (S1.16c) yields the following

1486   approximation for $\hat{R}$:

1487

1488 $$\hat{R} \approx \tfrac{1}{2}\hat{q}(1-\hat{q})\{2\hat{q}[(1-h-k)-2h(1-2h)s]-h^2 s - 2\,[h+(1+k-2h)\hat{q}]\epsilon\}[h+2(1-3h)\hat{q}+r/s]^{-1}$$

1489   (A10)

1490   Similarly, for the multiplicative gene model without synergistic or antagonistic

1491   epistasis, Equation (S1.16d) yields:

1492

1493 $$\hat{R} \approx \hat{q}^2(1-\hat{q})\,\{(1-h-k)+(k-h)s\}[h+2(1-3h\,)\hat{q}+r/s]^{-1} \quad \text{(A11)}$$

1494

1495   For $h \leq \frac{1}{2}$, and $h \leq k \leq \frac{1}{2}$, $\hat{R} > 0$.

1496

4. Extension to a large number of selected sites

It is possible to derive an approximation for the LD measures considered above in the case of

a large number of sites subject to the same selection regime. It is assumed that Equation (A8)

applies to a given pair of sites, provided that expressions for w_x, w_y for such a pair of sites, as

well as for $\bar{w}$, can be obtained. The equilibrium allele frequency at each site, $\hat{q}$, is treated

initially as a given. For the purpose of comparisons with the simulation results, we consider

only the additive gene model with $k = \frac{1}{2}$. The effects of LD on the distribution of the number

of deleterious mutations per haplotype are ignored, so that the probability that a haplotype

carries i mutations (P_i) is assumed to follow a Poisson distribution with mean $\mu = G\hat{q}$, where

$\hat{q}$ is the equilibrium frequency of mutant alleles at a given site ($G \gg 1$):

1507

$$P_i = \frac{\mu^i}{i!}\exp(-\mu) \qquad \text{(A12)}$$

Consider first the net marginal fitness of a haplotype that lacks mutations at a pair of

randomly chosen sites, w_z. Such a haplotype may carry 0, 1, 2, ... $G - 2$ mutations located at

other sites. Ignoring LD, the probability of i mutations at other sites is P_i; the marginal fitness

of such a haplotype is denoted by w_{zi}. Taking into account the other haplotypes that are

encountered to form diploid individuals, and using the fitness function for the gene model

with $k = \frac{1}{2}$, we have:

$$w_{z0} = 1 - hs\sum_{j=0}^{\infty} jP_j = 1 - \mu hs \qquad \text{(A13}a\text{)}$$

1516

1517 For $i \geq 1$, we have:

$$w_{zi} = 1 - iP_0hs - \frac{s}{2}\sum_{j=1}^{\infty}(j+i)P_j$$

$$= 1 - ie^{-\mu}hs - \frac{s}{2}[\mu + i(1-e^{-\mu})] \qquad (A13b)$$

1520 This yields:

$$w_z = \sum_{i=0}^{\infty} w_{zi}P_i = 1 - 2\mu e^{-\mu}hs - \frac{s}{2}\sum_{i=1}^{\infty}[\mu + i(1-e^{-\mu})]P_i$$

$$w_z = 1 - [\mu - (1-2h)\mu e^{-\mu}]s \qquad (A13c)$$

1525 Next, consider the marginal fitness of a haplotype with a mutation at one of the two chosen
1526 sites. As before, the probability that such a haplotype carries i mutations at other sites is P_i
1527 and the marginal fitness of such a haplotype is:

$$w_{xi} = 1 - (i+1)P_0hs - \frac{s}{2}\sum_{j=1}^{\infty}(j+i+1)P_j \quad (i = 0, 1, \ldots)$$

$$= 1 - (i+1)e^{-\mu}hs - [\mu + (i+1)(1-e^{-\mu})]\frac{s}{2} \qquad (A14a)$$

1531 so that:

$$w_x = \sum_{i=0}^{\infty} w_{xi}P_i = 1 - (\mu+1)e^{-\mu}hs - [\mu + (\mu+1)(1-e^{-\mu})]\frac{s}{2} \qquad (A14b)$$

1534 The same procedure can be carried out for the marginal fitness of a haplotype with
1535 mutations at both chosen sites, w_y. The fitness of such a haplotype that carries i mutations at
1536 other sites is:

$$w_{yi} = 1 - (i+2)P_0hs - \frac{s}{2}\sum_{j=1}^{\infty}(j+i+2)P_j \quad (i = 0, 1, \ldots)$$

$$= 1 - (i+2)e^{-\mu}hs - [\mu + (i+2)(1-e^{-\mu})]\frac{s}{2} \qquad (A15a)$$

1539 and:

$$w_y = \sum_{i=0}^{\infty} w_{yi}P_i = 1 - (\mu+2)e^{-\mu}hs - [\mu + (\mu+2)(1-e^{-\mu})]\frac{s}{2} \qquad (A15b)$$

1542 If second-order terms in s are neglected, we have:

$$w_y w_z - w_x^2 \approx -2\mu e^{-\mu}hs - \mu(1-e^{-\mu})s - (\mu+2)e^{-\mu}hs - [\mu + (\mu+2)(1-e^{-\mu})]\frac{s}{2}$$

$$+ 2(\mu+1)e^{-\mu}hs + [\mu + (\mu+1)(1-e^{-\mu})]s$$

$$= \mu e^{-\mu}\left(\frac{1}{2} - h\right)s \qquad (A16)$$

1548 Furthermore, if $\hat{q} \ll 1$ and $s \ll 1$, we have:

1549

$$\overline{w} \approx w_z = 1 - [(2h-1)\mu e^{-\mu} + \mu]s \qquad (A17)$$

1551

1552 Using this expression together with Equation (A14b), we obtain:

1553

1554 $$\overline{w}^2 - w_x^2 \approx -4\mu e^{-\mu}hs - 2\mu(1-e^{-\mu})s + 2(\mu+1)e^{-\mu}hs + [\mu + (\mu+1)(1-e^{-\mu})]s$$

1555 $$= [(\mu-1)e^{-\mu}(1-2h)+1]s \qquad (A18)$$

1556

1557 Substituting from Equations (A16) and (A18) into Equation (A8) yields the expressions:

1558

1559 $$\widehat{D} \approx \hat{q}^2 \mu e^{-\mu}\left(\tfrac{1}{2}-h\right)[(\mu-1)e^{-\mu}(1-2h)+1]^{-1} \qquad (A19a)$$

1560

1561 $$\widehat{R} \approx \hat{q}\mu e^{-\mu}\left(\tfrac{1}{2}-h\right)[(\mu-1)e^{-\mu}(1-2h)+1]^{-1} \qquad (A19b)$$

1562

1563 Equations (A13c) and (A14b) can be used to obtain an equation for the value of μ and

1564 hence $\hat{q} = \mu/G$. Provided that $q << 1$, the change in q due to selection at a given site is given

1565 by:

1566 $$\Delta q_s \approx q(w_x - \overline{w}) \approx q(w_x - w_z) \qquad (A20a)$$

1567

1568 Substituting from Equations (A13c) and (A14b) and simplifying, the following expression is

1569 obtained:

1570 $$\Delta q_s \approx -\tfrac{1}{2}sq[1 + (\mu-1)e^{-\mu}(1-2h)] \qquad (A20b)$$

1571

1572 At mutation-selection equilibrium, this must balance the change in q due to mutation, which

1573 is approximately equal to the mutation rate u. The equilibrium value of μ is then determined

1574 by the following transcendental equation, which can be solved numerically by Newton-

1575 Raphson iteration:

1576

1577 $$\frac{2Gu}{s} \approx \hat{\mu}[1 + (\hat{\mu}-1)e^{-\hat{\mu}}(1-2h)] \qquad (A21a)$$

1578

With semi-dominance ($h = \tfrac{1}{2}$), this equation is equivalent to the standard equation

$\hat{q} = 2u/s$ (Haldane 1927), as expected from the equivalence of the sites and gene models in

this case. For smaller values of h, $\hat{q}$ is reduced below this value, as seen in the two-site model

described above. Importantly, for $h = 0$, $\hat{q}$ does not obey the standard single locus result for a

completely recessive mutation, $\hat{q} = \sqrt{u/s}$ (Haldane 1927), but instead takes much larger

values.

But for sufficiently small values of G, $\hat{\mu}$ is $\ll 1$, and Equation (A21a) can be

approximated by:

1587
$$\frac{Gu}{s} \approx \hat{\mu}[h + \hat{\mu}(1 - 2h)] \qquad\qquad (A21b)$$

1588

1589 When $h = 0$, this yields the standard equation for complete recessivity. When $\hat{\mu} \ll h$, this

1590 reduces to the standard Haldane equation $\hat{q} = u/(hs)$.

1591

1592 **5. The effect of using Lewontin's D' statistic on the sign of LD**

1593 Let the allele frequencies at the two sites in question, A and B, be p_A, q_A, p_B and q_B, where p

1594 and q refer to ancestral and derived variants, respectively. If D is defined as positive for

1595 associations between derived variants, we have:

1596

1597
$$|D_{max}| = \min(p_A\,p_B, q_A\,q_B\,), \quad D < 0 \qquad\qquad (A22a)$$

1598

1599
$$|D_{max}| = \min(p_A\,q_B, q_A\,p_B\,), \quad D > 0 \qquad\qquad (A22b)$$

1600

First, consider the situation if unweighted haplotype frequencies from the whole

population are used for the LD statistics. As described in the section *Methods: two-locus*

simulations, the LD statistics were calculated after a second mutation at site B, was

introduced at a site segregating for a mutation at site A, either in *cis* with the ancestral A

variant (case 1) or in *cis* with the derived variant (case 2). For case 1 in the neutral model

described above, $D < 0$ for all generations, and $D' = -q_A q_B/q_A\,q_B = -1$ if $q_A\,q_B < p_A\,p_B$

or $D' = -q_A q_B/p_A\,p_B > -1$ if $q_A\,q_B > p_A\,p_B$. Given that q_A and q_B are the frequencies of

derived variants, whose probability distribution is skewed towards low frequencies, the large

majority of cases will involve the first situation, and the expectation of D' for case 1 will be –

1 or close to – 1.

For case 2, $D' = p_A q_B/p_A q_B = 1$ if $p_A\,q_B < q_A\,p_B$ and $D' = p_A q_B/q_A p_B > 1$ if

$p_A\,q_B > q_A\,p_B$. Again, the majority of cases will satisfy the first condition, since q_B is likely

to be much smaller than q_A, due to B$_2$ arriving in the population as the second mutation. The

expectation of D' for case 2 will thus be 1 or close to 1. Case 1 has a higher probability of

occurring than case 2 (their probabilities are proportional to the integrals between $1/N_H$ and 1

$- 1/ N_H$ of pq^{-1} and 1, respectively), so that the overall expectation of D' is negative. If

purifying selection is acting, the probability of case 1 is increased over case 2, due to the

increased skew of the probability distribution of allele frequencies towards low values, so the

magnitude of D' will increase with the strength of selection.

The properties of D' for doubleton variants are described in section 8 of

Supplementary File S1.

1622

1623

1624

1625

1626

1627

1628

1629   **TABLES**

1630
1631

1632   **Table 1. Models of the fitness effects of two sites with a fixed selection coefficient**

1633

1634   The "sites" model

1635

	++	+ -	- +	- -
+ +	1	$1 - hs$	$1 - hs$	$1 - 2hs - e_1$
+ -	$1 - hs$	$1 - s$	$1 - 2hs - e_1$	$1 - s - hs - e_2$
- +	$1 - hs$	$1 - 2hs - e_1$	$1 - s$	$1 - s - hs - e_2$
- -	$1 - 2hs - e_1$	$1 - s - hs - e_2$	$1 - s - hs - e_2$	$1 - 2s - e_3$

1636
1637 $e_1 = 2\epsilon hs$; $e_2 = \epsilon(1 + h)s$; $e_3 = 2\epsilon s$.

1638
1639   The "gene" model

1640

	+ +	+ -	- +	- -
+ +	1	$1 - hs$	$1 - hs$	$1 - 2hs - e_1$
+ -	$1 - hs$	$1 - s$	$1 - s$	$1 - (1+k)s - e_2$
- +	$1 - hs$	$1 - s$	$1 - s$	$1 - (1+k)s - e_2$
- -	$1 - 2hs - e_1$	$1 - (1+k)s - e_2$	$1 - (1+k)s - e_2$	$1 - 2s - e_3$

1641
1642
1643 $e_1 = 2\epsilon hs$; $e_2 = (1 + k)\epsilon s$; $e_3 = 2\epsilon s$.

1644

1645

1646
1647

Table 2. Statistical summaries of linkage disequilibrium between selected alleles for the gene
and sites models, with fixed scaled selection coefficient $\gamma = 2Ns$. 1000 selected sites were
simulated, with $Nu = 0.005$ and $Nr = 0.01$, where N is the population size ($N = 1000$), and u
and r are the mutation and recombination rate per site/generation respectively. $\bar{D}$ and $\bar{\sigma}_d$
were calculated for alleles that were less than 100 sites apart. The sign of the LD statistic $\bar{\sigma}_d$
was assigned either by giving a positive sign to cases with excesses over random
combinations of pairs of selectively deleterious alleles, or to cases with excesses of
combinations of minor alleles. The means and standard errors (SEs) for 1000 replicate
simulations are shown. The predicted values of $\bar{D}$ for the gene model with $\gamma = 20$ and 100 are
also shown; these were obtained as described in the text; the corresponding values for the
sites model are expected to be close to zero.
1659

selected allele		h=0.0		h=0.2		h=0.5	
		gene	sites	gene	sites	gene	sites
$\gamma = 2$							
$\bar{D}$	mean	-0.00021	-0.00032	-0.00016	-0.00026	-0.00016	-0.00021
	SE	0.00004	0.00004	0.00004	0.00004	0.00004	0.00004
$\bar{\sigma}_d$	mean	-0.00241	-0.00280	-0.00184	-0.00247	-0.00175	-0.00238
	SE	0.00054	0.00036	0.00056	0.00047	0.00051	0.00055
$\gamma = 20$							
$\bar{D}$	mean	0.00019	-0.00010	0.00008	-0.00006	-0.00006	-0.00002
	SE	0.00002	0.00003	0.00002	0.00002	0.00002	0.00002
	predicted	0.00015	0	0.00009	0	0	0
$\bar{\sigma}_d$	mean	0.00733	-0.00080	0.00347	-0.00052	-0.00153	-0.00022
	SE	0.00070	0.00045	0.00061	0.00056	0.00053	0.00057
	predicted	0.0052	0	0.00313	0	0	0
$\gamma = 100$							
$\bar{D}$	mean	0.00008	0.00011	0.00004	-0.00003	-0.00001	0.00003
	SE	0.00001	0.00001	0.00001	0.00001	0.00002	0.00002
	predicted	0.00004	0	0.00002	0	0	0
$\bar{\sigma}_d$	mean	0.00447	0.00361	0.00247	-0.00126	0.00005	0.00211
	SE	0.00083	0.00043	0.00087	0.00065	0.00142	0.00161
	predicted	0.00393	0	0.00157	0	0	0

minor allele							
$\gamma = 2$							
$\bar{D}$	mean	0.00288	0.00593	0.00305	0.00473	0.00289	0.00289
	SE	0.00010	0.00014	0.00009	0.00012	0.00009	0.00009
$\bar{\sigma}_d$	mean	0.03604	0.05513	0.03831	0.05033	0.03637	0.03644
	SE	0.00111	0.00120	0.00113	0.00121	0.00103	0.00109
$\gamma = 20$							
$\bar{D}$	mean	0.00019	0.00038	0.00009	0.00002	-0.00005	-0.00002
	SE	0.00002	0.00004	0.00002	0.00002	0.00002	0.00002
$\bar{\sigma}_d$	mean	0.00737	0.00618	0.00360	0.00136	-0.00121	-0.00013
	SE	0.00070	0.00056	0.00062	0.00058	0.00053	0.00057
$\gamma = 100$							
$\bar{D}$	mean	0.00008	0.00012	0.00004	-0.00003	-0.00001	0.00003
	SE	0.00001	0.00001	0.00001	0.00001	0.00002	0.00002
$\bar{\sigma}_d$	mean	0.00451	0.00375	0.00251	-0.00119	0.00015	0.00211
	SE	0.00083	0.00043	0.00088	0.00066	0.00142	0.00161

1660

1661

1662

Table 3: Statistical summaries of linkage disequilibrium under the gene and sites models in

human-like populations, using a gamma distribution of scaled selection coefficients with

shape parameter 0.23 and mean $\bar{\gamma} = 2N\bar{s}$. 2000 selected sites were simulated, with $Nu =$

0.00025 and $Nr = 0.0002$, where N is the population size ($N = 1000$), u is the mutation rate

per site/generation, and r is the rate of crossing over between adjacent sites. The signs of

$\bar{D}$ and $\bar{\sigma}_d$ were assigned either by giving a positive sign to cases with excesses over random

combinations of pairs of selectively deleterious alleles, or to cases with excesses of

combinations of minor alleles. The means and standard errors (SEs) for 1000 replicate

simulations are shown.

1672

selected alleles		h=0.0		h=0.2		h=0.5	
		gene	sites	gene	sites	gene	sites
$\bar{\gamma} = 850$							
$\bar{D}$	mean	0.00015	0.00031	0.00101	0.00030	-0.00096	-0.00105
	SE	0.00045	0.00035	0.00062	0.00066	0.00063	0.00074
$\bar{\sigma}_d$	mean	0.01130	0.00650	0.01587	0.00345	-0.00706	-0.00345
	SE	0.00472	0.00353	0.00591	0.00547	0.00525	0.00600
$\bar{\sigma}_d^1$	mean	0.46777	0.25534	0.61505	0.14143	-0.12457	0.20649
	SE	0.17992	0.08900	0.21642	0.11249	0.09229	0.20202
minor alleles							
$\bar{\gamma} = 850$							
$\bar{D}$	mean	0.00123	0.00137	0.00319	0.00319	0.00244	0.00285
	SE	0.00046	0.00036	0.00063	0.00065	0.00071	0.00078
$\bar{\sigma}_d$	mean	0.01924	0.02183	0.03972	0.03556	0.02356	0.03175
	SE	0.00473	0.00373	0.00601	0.00548	0.00560	0.00625
$\bar{\sigma}_d^1$	mean	0.54411	0.51751	0.94606	0.55386	0.20582	0.60968
	SE	0.17966	0.09122	0.21699	0.11278	0.09449	0.20269

1673

1674

1675

1676

1677
1678

1679 **FIGURES**

Mutations within a single gene **Mutations across multiple genes**

Additive <u>sites</u> model Additive <u>sites</u> model

fitness $(w) = 1\text{-}2hs$

fitness $(w) = 1\text{-}2hs$

Additive <u>gene</u> model

★ – allele (deleterious)

fitness $(w) = 1\text{-}s$

1680

1681

1682   **Figure 1:** A schematic of the additive gene versus additive sites models. Note that the

1683   additive sites model is approximately equivalent to the standard multiplicative fitness model.

1684

1685

1686
1687

1688
1689
Figure 2. Linkage disequilibrium (LD) as a function of the synergistic epistasis coefficient

(ϵ; left-hand panels) and dominance coefficient (h; right hand panels) for the two-sites

deterministic equilibrium model, with $u = 0.0001$, $s = 0.01$, and $r = 0$. The upper panels are

the results for the additive sites model; the lower panels are for the additive gene model with

$k = ½$. LD is measured by the correlation coefficient (R) multiplied by 100. The solid curves

are the exact solutions to the recursion equations for D (Equations S1.2 of Supplementary

File S1). The dashed curves are the analytic approximations for R; see sections 2-3 of the

Appendix, Equations (A.9) and (A.10). The horizontal lines in the lower panels indicate zero

values of

R.

1700

1701

1702

1703

Figure 3: Statistical summaries of linkage disequilibrium as measured by $\bar{\sigma}_d$ under the gene

and sites models for a gamma distribution of scaled selection coefficients with shape

parameter 0.3 and a mean of $\bar{\gamma} = 2N\bar{s}$. 1000 selected sites were simulated, with $Nu = 0.005$

and $Nr = 0.01$, where N is the population size ($N = 1000$), u is the mutation rate per

site/generation, and r is the rate of crossing over between adjacent sites. $\bar{\sigma}_d$ was calculated for

alleles that were less than 100 basepairs apart. The sign of $\bar{\sigma}_d$ was assigned either by giving a

positive sign to cases with excesses over random combinations of pairs of selectively

deleterious alleles, or to cases with excesses of combinations of minor alleles (see the

Methods section). The means and standard errors (SEs) for 1000 replicate simulations are

shown. Similar results for $\bar{D}$ are shown in Figure S1 of Supplementary File S3.

1714

1715

1716

1717

1718

1719

1720

Figure 4: Statistical summaries of the $\bar{\sigma}_d$ measure of linkage disequilibrium between all pairs of sites segregating for deleterious mutations for gene and sites model with epistasis, for closely linked pairs of alleles (1-100 basepairs apart). The simulation parameters were the same as in Figure 3. The sign of the LD statistic $\bar{\sigma}_d$ was assigned either by giving a positive sign to cases with excesses over random combinations of pairs of selectively deleterious alleles or to cases with excesses of combinations of minor alleles. The error bars represent the SEs across replicates. Similar results for $\bar{D}$ are shown in Figure S2 of Supplementary File S3.

Figure 5. Mean linkage disequilibrium (LD) as a function of the epistasis coefficient (ϵ), from stochastic simulations of the two-site haploid model with $r = 0$. LD is measured by the $\bar{\sigma}_d$ statistic described in the text (multiplied by 100). The results for 10^6 replicate simulations with two different values of $\gamma = 2Ns$ are shown, with $N = 1000$. The dashed curve with filled markers shows the population values of LD with no weighting towards low allele frequencies; the full curve with open markers is for doubleton variants in a sample of size 100. The dashed curve with open markers shows population values with a weight of 10 toward low frequency variants; the full curve with filled markers is for a weight of 100. See the main text for further details.