A gene-based model of fitness and its implications for genetic variation: linkage disequilibrium

Parul Johri[1,2,3,*] and Brian Charlesworth[4,*]

[1]Department of Biology, University of North Carolina at Chapel Hill, NC 27599, United States
[2]Department of Genetics, University of North Carolina at Chapel Hill, NC 27599, United States
[3]Integrative Program for Biological and Genome Sciences, University of North Carolina at Chapel Hill, NC 27599, United States
[4]Institute of Ecology and Evolution, School of Biological Sciences, University of Edinburgh EH9 3FL, United Kingdom

*correspondence may be addressed to these authors: pjohri@unc.edu;
brian.charlesworth@ed.ac.uk

Keywords: complementation, dominance, epistasis, fitness models, linkage disequilibrium

Running title: Fitness models and linkage disequilibrium

Abstract

A widely used model of the effects of mutations on fitness (the "sites" model) assumes that heterozygous recessive or partially recessive deleterious mutations at different sites in a gene complement each other, similarly to mutations in different genes. However, the general lack of complementation between major effect allelic mutations suggests an alternative possibility, which we term the "gene" model. This assumes that a pair of heterozygous deleterious mutations in *trans* behave effectively as homozygotes, so that the fitnesses of *trans* heterozygotes are lower than those of *cis* heterozygotes. We examine the properties of the two different models, using both analytical and simulation methods. We show that the gene model predicts positive linkage disequilibrium (LD) between deleterious variants within the coding sequence, under conditions when the sites model predicts zero or slightly negative LD. We also show that focussing on rare variants when examining patterns of LD, especially with Lewontin's D' measure, is likely to produce misleading results with respect to inferences concerning the causes of the sign of LD. Synergistic epistasis between pairs of

37 mutations was also modeled; it is less likely to produce negative LD under the gene model

38 than the sites model. The theoretical results are discussed in relation to patterns of LD in

39 natural populations of several species.

Introduction

There is much interest in the effects of deleterious mutations on properties of populations

such as their genetic load, inbreeding depression, and variance in fitness, tracing back to the

pioneering work of Haldane (1937), Muller (1950), Morton et al. (1956) and Crow (1958),

and to more recent studies of the genetics of *Drosophila* fitness components (reviewed by

Crow 1993; Charlesworth 2015). In addition, there have been a number of recent theoretical

studies of measures of linkage disequilibrium (LD) among deleterious mutations (e.g., Garcia

and Lohmueller 2021; Roze 2021; Good 2022; Ragsdale 2022), stimulated by population

genomic studies where the sign of LD among rare, putatively deleterious variants has been

used to make inferences concerning features such as epistatic fitness interactions and Hill-

Robertson interference (Pool et al. 2012; Sohail et al. 2017; Sandler et al. 2021; Garcia and

Lohmueller 2021; Ragsdale 2022; Stolyarova et al. 2022).

The theoretical predictions are mostly based on population genetics models of diploid

organisms that tacitly assume that nucleotide sites subject to mutation and selection can be

treated as independent of each other with respect to their fitness effects, and of whether they

are in the same or different genes (reviewed by Kyriazis et al. 2023). Models of this type

were first introduced by Haldane (1937), who assumed that fitnesses were multiplicative

across loci. Consider individuals that are heterozygous at two different nucleotide sites.

Assume that the fitnesses of the two types of double heterozygotes are the same, regardless of

whether or not the two mutations are carried on the same haplotype (*cis*), or on the maternal

and paternal haplotypes (*trans*) – the "sites" model (see Figure 1A). With the same selection

coefficients against homozygotes (s) and dominance coefficients (h) at each site, the fitnesses

of these individuals are both $(1 - hs)^2$ if effects are multiplicative across sites and $1 - 2hs$ if

effects are additive. If selection is sufficiently weak that $s^2 << s$, as is assumed here, the

additive and multiplicative models are nearly equivalent. For mathematical convenience, we

will use the additive model from now on.

In contrast, if the gene is treated as a single entity that is subject to mutation, and there

is no allelic complementation between mutations within the same gene (more accurately, a

cistron as defined by Benzer [1957]), the fitness of an individual that is heterozygous for two

mutations depends on whether or not they are in *cis* or *trans* (Figure 1B) – the "gene model".

The fact that double heterozygotes with mutations in *cis* under the gene model have higher

fitnesses than those with mutations in *trans* implies an intragenic position effect with respect

to fitness when $h < ½$. This is more likely to favor positive linkage disequilibrium (LD) than

the sites model, if the sign of the coefficient of linkage disequilibrium (D) is defined as

positive when the frequencies of haplotypes carrying pairs of deleterious mutant alleles and

pairs of wild-type alleles are in excess of random expectation. This is because the lower

fitness of *trans* compared with *cis* double heterozygotes when $h < ½$ means that $+ +$ and $- -$

haplotypes will be transmitted more frequently to the next generation relative to $+ -$ and $- +$

haplotypes than when the two genotypes have the same fitness.

Position effects of this kind have rarely been included in population genetics models,

with the exception of a study of two-locus polymorphisms by Nordborg et al. (1995), and the

models of mutation and selection at multiple sites within a gene by Clark (1998), Thornton et

al. (2013) and Sanjak et al. (2015), and at two sites by Ragsdale (2022), who all used

somewhat different fitness models from ours. Such gene-based fitness models are likely to

result in patterns of LD between deleterious mutations that are different from those expected

under the sites model (Ragsdale 2022).

The purpose of the present paper is to examine the consequences of the differences

between the gene and sites models for patterns of linkage disequilibrium (LD) among

variants at sites under selection, using a model of a single coding sequence subject to

deleterious mutations. We do not attempt to present a completely general model of a multi-

site system, but instead analyze a simple model that brings out the main differences between

the two different representations of mutational effects on fitness. An analytical treatment of a

model with two sites under selection in an infinite, randomly mating population is given first,

followed by stochastic simulations of a multi-site sequence that is intended to provide an

approximate description of a coding sequence. We show that the difference between the two

models has significant consequences for the patterns of LD among such mutations, with the

gene model leading to positive LD among pairs of deleterious variants under conditions

where the additive model is associated with zero or slightly negative LD. Interpretations of

such patterns in natural populations need to take into account the possibility that the gene

model is likely to be a more accurate representation of the dominance properties of

deleterious mutations. We also show that focusing on rare variants, as advocated by Garcia

and Lohmueller (2021) and Good (2022), is likely to produce estimates of positive LD

among deleterious variants in agreement with Good (2022); the opposite finding of Garcia

and Lohmueller (2021) is due to their use of Lewontin's D' statistic (Lewontin 1964), which

is biased towards positive values when rare variants are used. The consequences of the

differences between the two models for the genetic load, inbreeding load, and genetic

variance in fitness are described in the companion paper (Johri and Charlesworth 2025).

Methods

Multi-locus simulations

Forward simulations were performed using SLiM 4.0 (Haller and Messer 2023), with a

Wright-Fisher population of N =1000 diploid individuals, a mutation rate of $u = 0.5 \times 10^{-5}$

per nucleotide site/generation, and a rate of crossing over of 10^{-5} per nucleotide

site/generation. A gene with 1000 selected sites was simulated, assuming a dinucleotide

model, *i.e*, there were two alleles at each site: a wildtype (denoted by +) and a selectively

deleterious mutant (denoted by –) allele. Note that the two alleles were equally likely to

mutate to each other, i.e., the + allele could mutate to the – allele at rate u and vice versa.

This model contrasts with similar previous studies (Garcia and Lohmueller 2021; Good 2022;

Ragsdale 2022), where mutation was unidirectional from + to –. This procedure has the

advantage that it allows a true stationary distribution of allele frequencies to be reached

(Wright 1931), in contrast to models with one-way mutation, where all sites will eventually

become fixed for mutant alleles. It thus provides a more realistic description of evolution

resulting from mutations involving base substitutions.

Simulations were run either assuming the same selection coefficient at each site or a

distribution of fitness effects (DFE). In the first case, the selective disadvantage of mutant

homozygotes scaled by twice the population size was fixed at $\gamma = 2Ns = 2$, 20, 100, or

1000. When simulating a distribution of fitness effects (DFE), a gamma distribution was

assumed for most of the results presented here, with shape parameter $\alpha = 0.3$ and mean

scaled selection coefficients of $\bar{\gamma} = 2N\bar{s} = 2$, 20, 100, or 1000. The homozygous selective

disadvantage of a – allele at a particular site remained constant throughout a given

simulation, although it varied across simulation replicates. The dominance coefficient (h) of

the mutant allele was fixed at 0.0, 0.2, or 0.5. The fitness of a diploid individual was

calculated using either the gene or the sites model, as described below. All sites (polymorphic

and fixed) were included when calculating the fitnesses of individuals. Simulations were run

for $10N$ generations from an initial population of N identical diploid individuals with no

variation. For each scenario, 1000 replicates were simulated, and used to determine the

means, standard deviations and standard errors across replicates.

Relating the simulations to Drosophila melanogaster *populations*

These simulation parameters can be placed in the context of the *D. melanogaster* genome, for

which there are on average about 5 exons per gene with a mean length of ~300 bp (Misra et

al. 2002). As only about two-thirds of coding sites are nonsynonymous, there are expected to

be ~1000 (= $300 \times 5 \times 2/3$) selected sites per gene, corresponding to our simulations. Intron

lengths, and hence gene lengths, vary across the genome, with a mean and median length of 5

kb and 2 kb respectively; we have not included introns or synonymous sites in our

simulations. Assuming $N_e = 10^6$ for *D. melanogaster* and a sex-averaged rate of crossing over

of 10^{-8} per site/generation (Johri et al. 2020), $NGr \approx 20 - 50$ over a genic region, where G

is the mean length of a gene. With our simulation parameters, the 1000 selected sites have

$NGr = 10$ when rescaled to a population size of 10^6, and therefore approximately represents

a genic region of *D. melanogaster*. Silent site polymorphism data imply that $Nu \approx 0.003 -$

0.005 in *D. melanogaster* East African populations (Johri et al. 2020), consistent with the

$Nu = 0.005$ used in our simulations. This rescaling uses the principle that, when all

evolutionary forces are weak, the outcome of evolution is determined by the products of the

effective population size and the magnitudes of the deterministic forces (e.g., Ewens 2004,

Chap. 5).

Because our simulations were computationally very expensive, it was not feasible to

perform them with a large population size. Rescaling by a large scaling factor in forward-in-

time simulations can affect population genetic statistics (Dabi and Schrider 2024; Ferrari et

al. 2024); rescaling has, however, been found to produce accurate results in the absence of

recombination (Kaiser and Charlesworth 2010). As we only simulated short regions of 1000

selected sites, rescaling is likely to produce accurate results in our case as well, since the

frequency of crossing over between the most widely separated sites is nearly proportional to

the distance between them, so that the recombination rates can be scaled by population size

when using a fixed rate of crossing over per basepair. This would not be the case for large

distances between sites, for which double crossovers cause a non-linear relation between

distance and recombination rate. This means that multiplication of the rate of crossing over

per basepair by a given factor does not result in a proportional change in the frequency of

recombination. To test the effects of rescaling, additional simulations were conducted with a

larger population size of 5000 diploid individuals, while maintaining the same values of Nu

($u = 2 \times 10^{-6}$ per site/generation) and Nr ($r = 2 \times 10^{-6}$ per site/generation) as the

simulations above. Selection coefficients followed a gamma distribution with an expected

value of $2N\bar{s} = 100$ and a shape parameter of 0.3.

Simulations of human populations

A Wright-Fisher population of $N = 1000$ diploid individuals was simulated with a mutation

rate $u = 2.5 \times 10^{-7}$ per nucleotide site/generation and crossing over rate (r) of 2×10^{-7} per

nucleotide site/generation. This resulted in $Nu = 0.00025$ and $Nr = 0.0002$ in our simulations,

which are consistent with the estimates of these parameters for human populations: N_e of

20,000, mutation rate of 1.25×10^{-8} per site/generation (Kong et al. 2012), and

recombination rate of 1.0×10^{-8} per site/generation (Altshuler et al. 2005). Assuming that

there are 8 exons per gene on average (Johri et al. 2021), that each exon is about 350 bp, and

that 70% of all coding sites in humans are nonsynonymous (Kryukov et al. 2007), there

should be about 1960 selected sites in a gene. We therefore simulated 2000 selected sites. We

again ignored introns; since genes tend to be coldspots for recombination in many species

(e.g., Myers et al. 2010; Joseph et al. 2024), the presence of introns should not greatly affect

recombination rates within genes. The DFE was assumed to be a gamma distribution with

shape parameter $\alpha = 0.23$ and $\bar{\gamma} = 850$, as estimated by Eyre-Walker et al. (2006) and Kim et

al. (2017).

Modeling the fitnesses of individuals

As mentioned in the Introduction, two different models of main effects on fitness (the sites

and gene models) are used to calculate the fitnesses of diploid genotypes. The sites model

assumes that heterozygous mutations at all sites complement each other fully. The gene

model assumes a complete absence of complementation between heterozygous mutations

present in *trans*. The difference in fitness between *cis*- and *trans*-heterozygotes can be

regarded mathematically as a form of epistasis, although its biological basis is different from

the type of epistasis usually considered in population genetics models.

In both cases, additivity of the main effects across sites was assumed in most of our

numerical work, as was also done by Good (2022) and Ragsdale (2022), in contrast to the

multiplicative fitness model that is often used in multi-locus studies. The difference between

additive and multiplicative fitness models involves second-order terms in the selection

coefficients. For a single gene with a modest number of segregating sites, as simulated here,

there should be little difference between additive and multiplicative models, unless selection

is much stronger than is generally assumed here. As just described, our simulations use

rescaling of the deterministic parameters, so that their products with N_e match those for much

larger natural populations. The second-order terms in the selection coefficients involved in

multiplicative fitness models makes such rescaling problematical; this difficulty is avoided

when the additive model used.

In addition to the main effects of each site, a simple model of pairwise epistasis of

the standard type was used, in which a single parameter (ϵ) describes a pairwise interaction

effect on fitness. To avoid confusion with the effect of non-complementation under the gene

model, we will refer to this as either antagonistic or synergistic epistasis. The total number of

sites within a gene is denoted by G. The parameter ϵ is positive if there is synergistic

epistasis, so that double mutants reduce fitness by more than is expected from the effects of

each mutation on its own, and negative if there is antagonistic (diminishing returns) epistasis,

where double mutants behave in the opposite fashion (Crow 1970; Kondrashov 1988).

Analytical results are derived in the Appendix and Supplementary File S1 for the case

of $G = 2$, with the same selection coefficient s at each site. In general, in addition to a

possible difference in fitness between *cis*- and *trans*-heterozygotes, there are three possible

epistatic coefficients that measure departure from additivity in a symmetric fitness model,

denoted by the coefficients e_1, e_2 and e_3 in Table 1. In order to avoid multiple epistatic

parameters, we assume that the magnitude of such epistasis scales linearly with the sizes of

the selective effects at each site with a constant of proportionality ϵ, so that the fitness of a

mutant double homozygote is $1 - 2s - 2\epsilon s$. The fitnesses of all possible genotypes under the

sites model are defined in the upper section of Table 1, where the assumption of linear

scaling and complete complementation of mutations at different sites in double heterozygotes

gives $e_3 = 2\epsilon s$, $e_1 = 2\epsilon h s$ and $e_2 = \epsilon(1 + h)s$. For the gene model, a double heterozygote with

mutations in *cis* has fitness $1 - 2hs - 2\epsilon hs$, whereas the absence of complementation in a

trans double heterozygote implies that its fitness is $1 - s$.

It is less obvious how to assign the fitness of a genotype for which one site is

homozygous for a mutation and the other is heterozygous, e.g., $+ -/- -$. The representation in

Table 1 assigns a fitness reduction of $(1 + k)(1 + \epsilon)s$ to this genotype. For most of our

numerical results, we assume that a lack of complementation implies that the effects of each

mutation on the fitness of this genotype combine additively, with the heterozygous site

contributing a fitness reduction of ½s and the homozygous site a reduction of s, so that $k =$
0.5. This giving a net additive effect of 1.5s and an epistatic effect of 1.5ϵs. With $e_3 = 2\epsilon s$, we
thus have $e_1 = \epsilon h s$ and $e_2 = 1.5\epsilon s$.

An extreme alternative is to assume that $k = h$, as for the sites model, so that the single
heterozygous site has the same effect as if the other site were wild-type. There are no data
available to distinguish between these alternatives. It is shown in the Appendix that the
second alternative is likely to give a substantially larger positive LD measure for pairs of
deleterious mutations, so that our use of the additive assumption is conservative from the
point of view of predicting positive LD.

Modeling fitness with multiple sites and no epistasis
For describing fitness (w) when there are G selected sites within a gene, as in the simulations,
consider the two haplotypes (1 and 2) of a diploid individual, where $w = 1$ when all sites are
wild-type. With the same selection coefficient at all sites, fitness for the gene model is
calculated as follows.

If there is at least one – allele in 1 or 2, but not both, we have:

$$w = 1 - T_1 hs \qquad (1a)$$

where T_1 is the total number of – alleles over all G sites.

If there is at least one – allele in both 1 and 2, we have

$$w = 1 - 0.5T_2 s \qquad (1b)$$

where T_2 is the total number of – alleles present.

In the case of a distribution of fitness effects (DFE), with variable selection
coefficient s_i at each site, these equations become:

$$w = 1 - h\sum_{i=1}^{T_1} s_i \qquad (2a)$$

$$w = 1 - 0.5\sum_{i=1}^{T_2} s_i \qquad (2b)$$

For the sites model with a fixed selection coefficient, we have:

$$w = 1 - T_3 hs - T_4 s \qquad (3)$$

271 where T_3 is the total number of heterozygous (+/−) sites and T_4 is the total number of mutant

272 homozygous (−/−) sites. With a DFE, Equation (3) becomes:

$$w = 1 - h\sum_{i=1}^{T_3} s_i - \sum_{j=1}^{T_4} s_j \qquad (4)$$

The gene model with multiple sites

For the gene model, an extension of the analytical treatment of the two sites model to

multiple sites subject to the same selection regime (in the absence of epistasis) is described in

the final section of the Appendix. This is based on the assumptions that LD is sufficiently

weak that the distribution of numbers of mutations per haplotype follows a Poisson

distribution, that the number of selected sites G is large, and that selection is sufficiently

strong that the equilibrium frequencies of deleterious mutations at each site are very small.

Since significant levels of synergistic or antagonistic epistasis generate LD, we have not

included epistasis in this approach. We consider only the gene model, since LD for the sites

model without epistasis is expected to be negligible.

Modeling epistasis with multiple sites

All epistatic effects are assumed to be between pairs of sites that are either homozygous or

heterozygous for mutant alleles, *i.e*, there are no higher-order epistatic effects. Each pairwise

interaction involves the sum of a joint measure of the fitness effects of the alleles at the pair

of sites in question. The simplest model, which applies to both the sites and gene models, is

to assume that the epistatic contribution from a pair of homozygous sites i and j is $\epsilon(s_i + s_j)$;

this is summed over all pairs of homozygous sites, and the net fitness reduction of the

genotype where all sites are homozygotes for mutations is given by this term and the sum of

the s_i over all homozygous mutant sites.

For the sites model, a heterozygous site i can be considered as contributing a main

effect of hs_i to the fitness reduction of a genotype, and ϵhs_i to each of the interactions with

other sites. Its interaction with a homozygous site j thus contributes $\epsilon(hs_i + s_j)$ and its

interaction with another heterozygous site k contributes $\epsilon h(s_i + s_k)$. Similarly, a pair of

homozygous sites i and j contributes a fitness reduction of $\epsilon(s_i + s_j)$. The net epistatic

contribution is obtained by summing over all pairs of sites, and the net fitness reduction is

given by this sum plus the sum of the fitness reductions from each heterozygous and

homozygous mutant site. Note that for both the sites and the gene model, if the fitness of a

diploid individual was found to be negative in the simulations, it was changed to a value of

zero.

For the gene model, a heterozygous individual whose mutations are all in *cis* would

have an epistatic component given by the sum over all pairs of i and j of $\epsilon h(s_i + s_j)$. The net

fitness reduction is the sum of this quantity plus the sum of the hs_i over all the heterozygous

sites. For an individual who is homozygous for a mutation, or heterozygous in *trans,* at more

than one site, the sum of the epistatic contributions over all pairs of sites is obtained by

weighting ϵs_i for heterozygous sites by ½ instead of by h, and ϵs_i for homozygous sites by 1.

The net fitness reduction is given by the sum of this quantity and the sum of the main effects

across sites, weighting s_i at homozygous sites by 1 and heterozygous sites by ½. This is

slightly different from the symmetric two-site model of Table 1, where a heterozygote for

two mutations in *trans* behaves like a homozygote.

Estimating the linkage disequilibrium statistics from the simulations

Summaries of linkage disequilibrium (LD) between sites from simulated data were calculated

using the mean of the coefficient of linkage disequilibrium D across all pairs of segregating

sites:

321
$$D = p_{ab} - q_a q_b \tag{5}$$

323   and the two normalized measures of LD:

324
$$\bar{\sigma}_d = \frac{\bar{D}}{\sqrt{\overline{p_a p_b q_a q_b}}} \tag{6}$$

326
$$\bar{\sigma}_d^1 = \frac{\bar{D}}{\overline{p_a p_b q_a q_b}} \tag{7}$$

where q_a is the selected (deleterious) or minor allele frequency at the first site, q_b is the

deleterious/minor allele frequency at the second site, $p_a = 1 - q_a$, $p_b = 1 - q_b$, and p_{ab} is

the frequency of the haplotype "*a b*" which corresponds to the presence of either the

deleterious or the minor alleles at both sites. The overbars represent the mean of the

respective quantities over all SNP pairs and simulation replicates. Here, the mean of D and

the mean of the product $p_a p_b q_a q_b$ was calculated across all SNPs for each simulated

replicate. For each replicate, a single value of $\bar{\sigma}_d$ and $\bar{\sigma}_d^1$ was obtained by taking the ratio of

the mean D and the mean product of allele frequencies. The means and SEs reported in tables

and figures are those across the 1000 replicate simulations. The fluctuations across replicates

in the values of the three statistics ($\bar{D}$, $\bar{\sigma}_d$ and $\bar{\sigma}_d^1$) will thus be somewhat different in

magnitude, so there is no guarantee that their overall means will have the same sign.

However, this problem did not greatly affect our results, because it only happened when $\bar{D}$

was not significantly different from 0.

$\bar{\sigma}_d$ corresponds approximately to the mean correlation coefficient between the two

loci (Ohta and Kimura 1969); the denominator of the statistic $\bar{\sigma}_d^1$, used by Good (2022), and

Ragsdale (2022) greatly magnifies the apparent magnitude of LD compared with $\bar{\sigma}_d$.

However, the denominator of $\bar{\sigma}_d^1$ (the mean crossproduct of the four allele frequencies at a

pair of segregating sites, rather than its square root) means that its range is not confined to

(0,1), in contrast to the correlation coefficient.

Note that in our case, a derived allele may not always be equivalent to a *deleterious*

allele, since back mutations from – to + are possible. We have used the *selected* alleles as the

basis for calculations of the direction of D; with sufficiently strong selection ($Ns > 4$), there

will be little difference between these two classifications, since deleterious alleles are almost

always derived.

Two-locus simulations

We also investigated the effect of weighting LD statistics toward low frequency derived

variants, a procedure suggested by Garcia and Lohmueller (2021) and Good (2022). We used

the approach of Good (2022) for simulating a two-site haploid model without recombination.

This procedure assumes that a mutation from A_1 to A_2 at the first site (A) is present at a

frequency q, whose probability distribution is given by the standard formula for drift and

selection (Equations S1.17 of section 5 of Supplementary File S1). A second mutation, from

B_1 to B_2, was introduced at the second site, either in *cis* with A_1 (case 1, probability $1 - q$) or

in *cis* with A_2 (case 2, probability q). The fitness model for determining the post-selection

frequencies of the haplotypes in a given generation assumed that the relative fitnesses of

A_1B_1, A_1B_2, A_2B_1 and A_2B_2 are 1, $1 - s$, $1 - s$ and $1 - 2s(1 + \epsilon)$, respectively, where ϵ is a

measure of epistasis of the same form as that used in the two-site diploid model. As described

in section 2 of Supplementary File S1, multinomial sampling of N haplotypes was then

carried out each generation, until one of the two mutations was fixed or lost, and the mean

values over all generations and replicates of D, D^2 and the allele frequency cross-product $P =$

$p_{A}q_{A}p_{B}q_{B}$ (where subscripts A and B denote sites A and B) were obtained. The means of D, D^2

and P, denoted by $\overline{D}$, $\overline{D^2}$ and $\overline{P}$, were found by dividing the respective sums of these

statistics by the sum of the durations of the replicate simulations.

Data availability

The scripts used to determine the properties of two-site equilibrium populations, perform all

the simulations, and calculate statistics are provided at

https://github.com/paruljohri/Gene_vs_sites_model/tree/main.

Results

Deterministic results for two selected sites

Here we examine the properties of equilibria under mutation-selection balance at two

selected sites within a coding sequence. The fitness matrices in Table 1 are used for this

purpose. The mutation rate from $+$ to $-$ at both sites is u. The symmetry of the fitness models

means that the equilibrium allele frequencies are the same at both sites, so that the system can

be completely described by the frequency q of the $-$ allele at each site and the coefficient of

linkage disequilibrium D between them. The correlation coefficient between the two selected

sites is then given by $R = D/q(1-q)$. An algebraic treatment is given in the Appendix and

sections 1-4 of Supplementary File S1. We focus on the case when the recombination

frequency, r, is zero. This gives insights into the behavior of the system with low frequencies

of recombination, such as those between pairs of sites within a gene. The complexity of the

equilibrium equations, even when $r = 0$, means that exact results for the equilibrium

haplotype frequencies can only be obtained numerically, as described in the Appendix.

It is, however, possible to obtain approximate analytical results for the sign of D at

equilibrium without full solutions for the equilibrium haplotype frequencies. As might be

expected from results in the earlier literature (e.g., Felsenstein 1965; Kimura and Maruyama

1966; Kondrashov 1988; Charlesworth 1990; Shnol and Kondrashov 1993; Roze 2021), a

multiplicative version of the sites model with no synergistic or antagonistic epistasis results

in equilibria with $\widehat{D} = 0$ (where the carat indicates equilibrium). In contrast, the additive

version results in equilibria with negative values of $\widehat{D}$, reflecting the weak synergistic

epistasis on the multiplicative scale that is associated with additive fitnesses (see sections 2

and 3 of File S1). Explicit approximate expressions for the equilibrium values of $\widehat{D}$ and $\widehat{R}$ for

the additive sites model with no epistasis and a low frequency of recombination are given by

Equations (A9*a*), (A9*b*) and (A9*c*); the effect of epistasis on $\widehat{R}$ is described by Equation

(A9*d*). Equations (A9) show that, if the ratio of recombination rate to selection coefficient

(r/s) is $<< \hat{q}$ or $<< h$, recombination has only a small effect on the magnitude of D and R.

The magnitudes of $\widehat{D}$ and $\widehat{R}$ under the additive sites are, however, very small in the

absence of epistasis, of the order of the mutation rate. This can be seen as follows. Provided h

is bounded away from 0, and $u << hs$, we have $\hat{q} \approx \frac{u}{hs} \ll 1$ (Haldane 1937). The leading term

in braces in the numerator of Equation (A9*b*) is h^2, and the leading term in braces in the

denominator is h, so that the magnitude of the ratio of numerator to denominator in Equation

(A9*b*) is approximately $\frac{1}{2}hs\hat{q} \approx \frac{1}{2}u$. If $h = 0$, $\hat{q} \approx \sqrt{u/s}$, so that the term in braces in the

numerator is approximately u/s and the term in braces in the denominator is approximately

$\hat{q}$. This cancels the $\hat{q}$ term in the numerator, so that (assuming $\hat{q} \ll 1$), the magnitude of the

ratio of numerator to denominator is approximately $\frac{1}{2}u$, as before. It follows that, provided

that $\hat{q} \ll 1$, $\widehat{R}$ should be approximately equal to the negative of one-half of the mutation rate,

and independent of the dominance coefficient and strength of selection. Provided that the

epistasis coefficient ϵ is of the same order as s or greater, much larger negative values of $\widehat{D}$

and $\widehat{R}$ are expected with synergistic epistasis ($\epsilon > 0$), and positive values are expected with

antagonistic epistasis ($\epsilon < 0$) – see Equation (A9*c*).

In contrast, for the additive gene model with no synergistic epistasis, Equation (A10)

shows that $\widehat{D}$ and $\widehat{R}$ are positive when h is sufficiently small relative to ½ and $k \leq$ ½, selection

is sufficiently weak, and $\hat{q}$ is sufficiently large. $\widehat{D}$ and $\widehat{R}$ become slightly negative when these

conditions are violated, even in the absence of synergistic epistasis, but are always positive

for the corresponding multiplicative model when $h \leq k \leq$ ½ (Equation A11). Equations (A10)

and (A11) show that a smaller value of k is associated with larger LD values; in the absence

of synergistic epistasis, the values of $\widehat{D}$ and $\widehat{R}$ with $k = h <$ ½ are approximately twice those

with $h < k =$ ½ , reflecting the weaker selection against $--$ haplotypes in the former case (see

Equation A3*b*). Unsurprisingly, with sufficiently strong synergistic epistasis, $\widehat{D}$ and $\widehat{R}$

become negative (and vice-versa for antagonistic epistasis).

These conclusions are borne out by the results of iterating the exact recursion

relations (Equations A1 and A2), which are shown in Figure 2. The upper panels in Figure 2

show the results for the sites model, and the lower panels those for the gene model with $k =$

½. In both cases, $u = 0.0001$ and $s = 0.01$These are clearly not realistic values of the mutation

and selection parameters but produce relatively large values of $\hat{R}$, enabling the gene and sites

models to be compared more convincingly than when $\hat{R}$ is very small. The left-hand panels

plot $\hat{R}$ (in percent) against the epistasis coefficient, ϵ, for three different dominance

coefficients, h. The right-hand panels plot $\hat{R}$ against h for three different values of ϵ. Since

antagonistic epistasis ($\epsilon < 0$) is well known to induce positive LD (Kondrashov 1988;

Charlesworth 1990; Shnol and Kondrashov 1993), we have focused here on synergistic

epistasis ($\epsilon > 0$), as well as the zero epistasis case. As is shown below, the general properties

of the deterministic case mirror those of the multi-site simulations, although the results are

quantitatively very different.

Several points of interest emerge. As expected from the theoretical results, the sites

model shows consistently negative but very small values of $\hat{R}$, which are very close to the

negative of one-half of the mutation rate ($- 5 \times 10^{-5}$) when $\epsilon = 0$, independently of h. The

dominance coefficient has larger effects under the sites model, the larger ϵ; $\hat{R}$ declines almost

linearly with ϵ, becoming negative when ϵ is sufficiently large. In contrast, for the gene

model with no epistasis, $\hat{R}$ remains positive until h becomes close to ½, decreasing

monotonically as h increases. With synergistic epistasis, the gene model is associated with

larger values of $\hat{R}$ compared with the sites model when $h < ½$, and the difference is more

marked, the smaller h, with small h being associated with larger magnitudes of $\hat{R}$. In addition,

with the strongest value of epistasis ($\epsilon = 0.32$), $\hat{R}$ first declines with h and then increases

again as it approaches ½ but remains negative. The approximations describe the general

patterns quite well, with the exception of the sites model with strong synergistic epistasis and

small h, where the approximation incorrectly predicts an increase in $\hat{R}$ as h approaches zero.

Sets of results for other values of u and s are shown in Supplementary File S2. The

general features described above can be seen in these cases. However, as might be expected,

the smaller the mean deleterious allele frequency, the smaller the differences between the

gene and sites models, and the smaller the magnitudes of $\hat{D}$ and $\hat{R}$. While $\hat{D}$ and $\hat{R}$ are still

positive for the gene model in the absence of synergistic epistasis and sufficiently small hs,

under the sites model they become negative at much smaller ϵ and $\hat{q}$ values. Thus, in order to

determine whether there are any biologically important consequences of the features just

described, it is necessary to use computer simulations to examine what happens with multiple

selected sites, where larger effects are to be expected.

Deterministic results for the gene model with multiple sites

As described in the last section of the Appendix, the results from the two-site gene model

without epistasis can be extended to an arbitrary number of selected sites, all subject to the

same selection regime, using the assumptions described in the above section *The gene model*

with multiple sites. Equations (A19) provide approximate expressions for $\widehat{D}$ and $\widehat{R}$ for the

additive gene model without epistasis, given a value of $\hat{q}$, on the assumption that $\hat{q} \ll 1$ and

that number of selected sites (G) is large. Equations (A21) show how to determine $\hat{q}$ for

given values of G, *u*, *s* and *h*. The results were derived for the case $k = \frac{1}{2}$, since this was used

in the simulations. The conclusion that positive LD between pairs of deleterious mutations is

expected, provided that $h < \frac{1}{2}$, is confirmed by this analysis. In contrast to the two-site

model, where $\widehat{D}$ and $\widehat{R}$ depend on $\hat{q}^3$ and $\hat{q}^2$ respectively (see Equations A11), $\widehat{D}$ is

proportional to $\hat{q}^2 \mu e^{-\mu}(\frac{1}{2} - h)$ and $\widehat{R}$ to $\hat{q}\mu e^{-\mu}(\frac{1}{2} - h)$ where $\mu = G\hat{q}$ is the mean number of

deleterious mutations per haplotype; this reflects the cumulative effects of LD across multiple

sites. The next section presents comparisons of the predictions of Equation (A19a) with the

simulation results for LD.

Simulation results for multiple sites

The analytical results presented above were supplemented with forward simulations of a

single gene with multiple selected sites, with an equal probability of mutations between the +

and – alleles and vice versa (see the section *Methods*, *Multi-locus simulations*). We first

describe the results with no synergistic or antagonistic epistasis.

Linkage disequilibrium with no epistasis

As discussed in the section *Deterministic results for two selected sites*, deleterious alleles are

more likely to be associated on the same haplotypes in the gene model than in the sites

model, resulting in larger positive mean *D* values for the gene model with suitable parameter

values, as can be seen from the results in Figure 2. Table 2 shows the results of simulations

without epistasis in which *s* was fixed, and LD statistics were calculated by assigning positive

values to cases when + + and – – haplotypes were in excess of random expectation (entries

under "selected alleles"). In these examples, the gene model results in higher mean LD than

the sites model for closely linked sites whenever there are significantly non-zero values. The

difference between the two models increases with increasing recessivity and strength of

selection, with mean positive values of $\overline{D}$ and $\bar{\sigma}_d$ for the gene model versus negative values

with the sites model when $h < 0.5$ and $\gamma \geq 20$. For $\gamma = 2$, however, negative values were found

with both models, consistent with the operation of Hill-Robertson interference (HRI) effects

generated by the interaction between drift and selection with multiple linked selected sites

(Hill and Robertson 1966; Garcia and Lohmueller 2021; Roze 2021), which are not allowed

for in the two-site deterministic model.

It is of interest to examine whether these results are consistent with the theoretical

predictions for the multi-site gene model outlined in the previous section. The simulation

results for $\gamma = 20$ and $\gamma = 100$ were used, since the results for $\gamma = 2$ must be heavily influenced

by drift, whereas the theoretical predictions are purely deterministic. A complication in

making this comparison is that the LD statistics for the simulations are presented for pairs of

segregating sites, since $D = 0$ for invariant sites and $\bar{\sigma}_d$ is defined only for segregating sites.

As expected from general theory on strongly selected deleterious mutations (e.g.

Charlesworth and Jain 2014), the proportion of sites that are segregating at a given time with

$\gamma = 20$ or 100 is very small (see Table S5 in File S3). The deterministic results for mean D are

expected to correspond to the expected value of D over all pairs of sites. For this reason, the

predictions for $\bar{D}$ in Table 2 were obtained by dividing the predictions of $\hat{D}$ from Equation

(A19a) by the squares of the mean proportions of segregating sites for the appropriate values

of γ and h, obtained from Table S5. The values of $\hat{q}$ were obtained from Table 3 of Johri and

Charlesworth (2025); these agree closely with the predictions from Equations (A21) (see

Johri and Charlesworth (2025). The predicted value of $\bar{\sigma}_d$ for a given parameter set can be

obtained by dividing the predicted value of $\bar{D}$ by one half of the nucleotide site diversity at

selected sites (π), normalized by the proportion of segregating sites. The values of π for the

parameters in Table 2 were obtained as described by Jain and Charlesworth (2014), equating

the selection coefficient in their formulae to $s[1 + (\mu - 1)e^{-\mu}(1 - 2h]$ in Equation (A20b).

Overall, the agreement between observed and predicted values is remarkably good,

considering the approximations involved, suggesting that the multi-site gene model provides

an accurate description of the population genetics of LD when selection is strong and G is

large.

Similar results were found with a gamma distribution of selection coefficients, with

the gene model yielding positive values of $\bar{D}$ and $\bar{\sigma}_d$ with $h < 0.5$ and $\bar{\gamma} \geq 100$, as opposed to

slightly negative (but not significantly non-zero) values for the sites model (Figure 3 and

Figure S1 in Supplementary File S3). The theoretical results for the two-sites model predict

that $\bar{D}$ and $\bar{\sigma}_d$ values are expected to be slightly negative for the sites model. Consistent with

this prediction, we observed small negative values when $h < ½$. These effects could also be

caused by HRI.

To test this possibility, we analyzed the results with a DFE for variants separated by a

greater-than-average distance (800-100 base pairs; Table S1 in Supplementary File S3). The

$\overline{D}$ values for selected variants under the sites model then approach zero, consistent with

reduced effects of HRI with higher frequencies of recombination, whereas the deterministic

theory suggests only a weak effect of recombination with the low values of r/s for pairs of

sites within a gene (see the previous section). In contrast, for the gene model under a gamma

distribution with $\overline{\gamma} = 1000$, $\overline{D}$ and $\overline{\sigma}_d$ are still substantially greater than zero, suggesting

significantly positive LD even at larger distances (Table S1 in Supplementary File S3). In

addition, the magnitude of negative LD is greatest with $h = 0.5$, the opposite of what the

deterministic site model predicts (see Figure 2). Furthermore, as discussed in the previous

section, the multiplicative fitness sites model is not expected to generate any LD, in contrast

to the additive model. When fitnesses were assigned to be multiplicative in our multi-locus

simulations, $\overline{D}$ and $\overline{\sigma}_d$ values were sometimes significantly negative when sites were less than

100 bases apart, especially with $h = 0$ (Table S2 in Supplementary File S3). For alleles

between 800-1000 bases apart, $\overline{D}$ and $\overline{\sigma}_d$ values were close to zero in all cases (Table S2 in

Supplementary File S3). This suggests that HRI was probably partly responsible for the

slightly negative values observed in our multi-site simulations of the sites model.

Interestingly, when an excess frequency of haplotypes with pairs of minor alleles was

used as the criterion for positive D, there is not much difference in $\overline{D}$ and $\overline{\sigma}_d$ between the

gene and sites models when selection is weak, with mainly positive values for both (Table 2

and Figure 3). However, with stronger selection against – alleles and $h < ½$, the gene model

yields slightly lower values of $\overline{D}$ and $\overline{\sigma}_d$. The sign of $\overline{D}$ is expected to be positive for minor

alleles in cases where the overall value of $\overline{D}$ for derived alleles is zero; this is because, in

cases where two derived alleles are in negative LD, and one is at a frequency greater than 0.5

while the other is at a frequency lower than 0.5, LD between their corresponding minor

alleles is positive (Potapova and Kondrashov 2023). The difference between the gene and

sites models probably reflects the lower allele frequencies with the gene model. It should also

be noted that we estimated $\overline{D}$ with respect to *selected* alleles, which were not necessarily

derived (due to back mutations). However, as was pointed out in the section *Estimating the*

linkage disequilibrium statistics from the simulations, this should only affect our results when

selection is weak (i.e., when $\gamma \leq 4$); selected alleles are equivalent to derived alleles in most

other cases.

Our forward-in-time simulations were conducted with small populations (1000

diploid individuals) that are a factor of 1000 smaller than the value of N_e commonly used for

Drosophila populations, so that the mutation rate, recombination rate, and selection

coefficients were a factor of 1000 larger than the natural population values (see the Methods

section). To check if rescaling biased our results, we conducted simulations with a much

larger population size (5000 individuals). LD statistics obtained using minor allele

frequencies were entirely unaffected by rescaling (Table S3 in Supplementary File S3).

rescaled population. The differences between the two N values in the means of the LD

statistics for pairs of deleterious variants were within the limits of sampling error, so that

there is no evidence that rescaling has a major systematic effect on these statistics.

In contrast to the simulations with *Drosophila*-like parameters, which we have

discussed above, simulations of human-like populations yielded high variances between

replicates and somewhat unclear patterns for the LD statistics (Table 3). Overall, however,

LD between selected sites is higher with the gene model than the sites model when $h < \frac{1}{2}$. In

this case, we show $\bar{\sigma}_d^1$ as well as $\bar{\sigma}_d$, for comparison with the analyses of human data by

Ragsdale (2022) (see *Discussion, Patterns of linkage disequilibrium in population genetic*

data). The difference in LD between the two models is much clearer when using $\bar{\sigma}_d$ or $\bar{\sigma}_d^1$ as

opposed to $\bar{D}$, because $\bar{\sigma}_d$ and $\bar{\sigma}_d^1$ (partially) correct for differences in allele frequencies.

LD with epistasis

Similarly to the analytical results discussed earlier, there is a substantial decrease in $\bar{D}$ and $\bar{\sigma}_d$

(using the selected/derived allele criterion) with increasing strength of synergistic epistasis

for both models when h is close to 0.0, but only small changes when h is 0.2 or 0.5 (Figures 4

and Figure S2 in Supplementary File S3). $\bar{D}$ and $\bar{\sigma}_d$ between selected alleles remain

significantly higher with the gene model than the sites model (for which both statistics

become negative) with large ϵ. There seems to be no significant difference between the gene

and sites model with these parameters when $h = 0.2$ and ϵ is very small, consistent with their

large SEs relative to the means when $\bar{\gamma} = 100$ (see Figure 3 for the results with no epistasis,

and Figure S3 in Supplementary File S3 for a comparison of the distribution of the statistics

for the results in Figure 3 and Figure 4); $\bar{D}$ and $\bar{\sigma}_d$ are both not significantly different from

zero in this case. Importantly, however, with $h = 0.2$ and $\epsilon > 0$, $\bar{\sigma}_d$ is negative with the sites

model, but is close to zero with the gene model, so that the two models display differences in

LD even with epistasis.

When the same LD statistics are calculated with minor/major alleles as the sign

criterion, there is a substantial decrease in LD with an increase in the epistasis coefficient;

however, the values always remain positive and there is no significant difference in $\bar{\sigma}_d$

between the two fitness models (Figure 4), although $\bar{D}$ obtained using minor alleles is

consistently and slightly smaller for the gene than the sites model, possibly due to lower

mean allele frequencies at equilibrium (see Table 3 of Johri and Charlesworth 2025). In

summary, if synergistic epistasis were prevalent amongst deleterious alleles, LD between

selected/derived alleles will probably still be higher and positive with the gene model than

the sites model when $h < \frac{1}{2}$, whereas no difference is expected between LD statistics

calculated using minor alleles as the sign criterion.

With much lower rates of crossing over ($0.1 \times$ the standard rate used here), when

mutations are fully recessive, LD calculated using minor alleles is significantly higher with

the sites model than the gene model (Figures S4 and S5 in Supplementary File S3). The

difference between the two models is largest when there is no epistasis, but persists even

when the epistasis coefficient is 0.32. The predicted differences in LD between the gene and

the sites models using the derived and minor allele sign criteria can thus be entirely opposite,

depending on the degree of dominance. HRI in the absence of synergistic epistasis seems to

produce only a weak tendency towards negative LD when $h = \frac{1}{2}$, and for the sites but not the

gene model when $h = 0$ or 0.2.

Effects on LD statistics of weighting towards low allele frequencies

Garcia and Lohmueller (2021) suggested that the effects of HRI on LD between deleterious

mutations would be more easily detected in samples from populations if doubleton variants

were used for estimating LD. This procedure might help to correct for the effect of allele

frequency on LD statistics; in addition, because rare variants are more likely to be deleterious

than common ones, larger negative D values might be expected for nonsynonymous than

synonymous variants when conditioning on doubletons. In their multi-locus simulations, they

indeed found negative LD between derived variants at sites subject to purifying selection,

both with $h = 0$ and $h = \frac{1}{2}$, with LD being measured by Lewontin's D' statistic (Lewontin

1964). In contrast, Good (2022) proposed that multiplying allele frequencies by a function

that gives greater weights to low frequency variants should lead to positive expected D

values, for both neutral and deleterious variants. He validated this expectation by a

combination of simulations and analytical formulae, using a model of a haploid population;

this is equivalent to a diploid population with $h = ½$ in the absence of epistasis. However,

while the simulations performed by Garcia and Lohmueller (2021) involved multiple selected

sites, those of Good (2022) involved only two sites.

In order to reconcile these seemingly contradictory findings, we used the approach

described in the section *Methods: Two-locus simulations*. This enables the means of D, D^2

and the crossproduct P of the four allele frequencies at sites segregating for two mutations

($\bar{D}$, $\overline{D^2}$ and $\bar{P}$) to be found, yielding estimates of the standardized LD statistics $\bar{\sigma}_d = \bar{D}/\sqrt{\bar{P}}$,

$\bar{\sigma}_d^1 = \bar{D}/\bar{P}$ and $\sigma_d^2 = \overline{D^2}/\bar{P}$. σ_d^2 corresponds approximately to the mean squared correlation

coefficient between the two loci (Ohta and Kimura 1969); the other two statistics are defined

by Equations (6) and (7). These statistics were estimated either with or without weighting; in

the latter case, the population frequencies of haplotypes carrying the derived, deleterious

variants at each site (frequencies q_A and q_B, respectively) were weighted by $\exp(-fq_A)$ and the

frequencies of haplotypes carrying the deleterious allele (B$_2$) at the second site by $\exp(-fq_B)$.

The factor f is 0 if no weighting is applied, and $f > 1$ if low frequency mutant alleles receive

preferential weights (f is the inverse of the factor used by Good [2002)]). In addition, the

effect of calculating LD statistics from variants present only at a pre-set frequency in a

sample of k haplotypes (Garcia and Lohmueller 2021) was determined by trinomial sampling

of k haplotypes each generation, recording statistics only for samples where the numbers of

alleles at each site in the sample matched the pre-set allele frequency (usually 2 out of k).

For neutral variants, the unweighted case gave values of $\bar{D}$ and $\bar{\sigma}_d$ that were non-

significantly negative and close to zero: $\bar{D} = 0.75 \times 10^{-4} \pm 5.14 \times 10^{-4}$ and $\bar{\sigma}_d = 0.00144 \pm$

0.00713 over 5 repeated simulations of 10^6 introductions of B$_2$. The value of σ_d^2 was close to

the theoretical value of 0.45 for the case of no recombination under the infinite sites model

(Ohta and Kimura 1971). In contrast, with $f = 10$ and a single set of 10^6 introductions of B$_2$,

$\bar{\sigma}_d$ and $\bar{\sigma}_d^1$ were 0.0107 and 0.915, respectively. The values when doubletons were sampled

were $\bar{\sigma}_d = 0.133$ and $\sigma_d^1 = 13.5$.

Some representative results for $\bar{\sigma}_d$ with selection (one set of 10^6 introductions of B$_2$; N

$= 1000$) are shown in Figure 5, for two different strengths of selection ($\gamma = 2Ns = 10$ and $\gamma =$

$2Ns = 50$) and varying amounts of synergistic epistasis. With no epistasis and no weighting,

$\bar{D}$ and $\bar{\sigma}_d$ were slightly negative, but became increasingly negative as ϵ increased. 20

repetitions of the case of $\gamma = 50$ without epistasis gave no evidence for significantly non-zero

LD ($\overline{D} = -5.35 \times 10^{-7} \pm 4.16 \times 10^{-7}$; $\overline{\sigma}_d = -8.88 \times 10^{-5} \pm 6.93 \times 10^{-5}$), implying that HRI

effects with two sites are not sufficiently strong to overcome any tendency for positive LD to

be produced by purifying selection. Weighting causes the LD statistics to take larger values,

which are positive with $f = 100$ even for the largest value of ϵ (0.64). They remain positive

over the whole range of ϵ if doubletons are sampled. Further results are shown in

Supplementary Files S4 and S5.

These results show that focussing on mutations at low frequencies results in positive

LD between derived variants, which can persist even if there is strong synergistic epistasis,

especially if doubletons are sampled. A heuristic argument that explains this effect is

provided in section 6 of Supplementary File S1. These considerations suggest that the use of

LD statistics estimated from low frequency variants is not helpful in diagnosing HRI effects

caused by purifying selection. On the contrary, it is likely to conceal the effects of HRI or

synergistic epistasis on LD between derived variants.

This raises the question of why Garcia and Lohmueller (2021) found consistently

negative LD in their multi-locus simulations of deleterious mutations when sampling

doubletons. It turns out that this was because they applied the D' statistic (Lewontin 1964) to

estimates of D for derived variants. D' is defined as $D/|D_{max}|$, where $|D_{max}|$ is the

maximum value of the absolute value of D conditioned on the allele frequencies. Using our

notation, when defining D as positive for associations between derived variants we have:

$$|D_{max}| = \min(p_A\, p_B, q_A\, q_B\,), \quad D < 0 \qquad\qquad (8a)$$

$$|D_{max}| = \min(p_A\, q_B, q_A\, p_B\,), \quad D > 0 \qquad\qquad (8b)$$

First, consider the situation if unweighted haplotype frequencies from the whole

population are used for the LD statistics. As described in the section *Methods: two-locus*

simulations, the LD statistics were calculated after a second mutation at site B, was

introduced at a site segregating for a mutation at site A, either in *cis* with the ancestral A

variant (case 1) or in *cis* with the derived variant (case 2). For case 1 in the neutral model

described above, $D < 0$ for all generations, and $D' = -q_A q_B/q_A\, q_B = -1$ if $q_A\, q_B < p_A\, p_B$

or $D' = -q_A q_B/p_A\, p_B > -1$ if $q_A\, q_B > p_A\, p_B$. Given that q_A and q_B are the frequencies of

derived variants, whose probability distribution is skewed towards low frequencies, the large

majority of cases will involve the first situation, and the expectation of D' for case 1 will be –

1 or close to – 1. For case 2, $D' = p_A q_B / p_A q_B = 1$ if $p_A q_B < q_A p_B$ and $D' = p_A q_B / q_A p_B$

> 1 if $p_A q_B > q_A p_B$. Again, the majority of cases will satisfy the first condition, since q_B is

likely to be much smaller than q_A, due to B_2 arriving in the population as the second

mutation. The expectation of D' for case 2 will thus be 1 or close to 1. Case 1 has a higher

probability of occurring than case 2 (their probabilities are proportional to the integrals

between $1/N_H$ and $1 – 1/N_H$ of pq^{-1} and 1, respectively), so that the overall expectation of D'

is negative. If purifying selection is acting, the probability of case 1 is increased over case 2,

due to the increased skew of the probability distribution of allele frequencies towards low

values, so the magnitude of D' will increase with the strength of selection.

This shows that unweighted estimates of mean D' that take into account the sign of D

are biased towards negative values. It seems likely that this will also apply to estimates based

on doubleton frequencies, although the expectation of D itself in this case is biased towards

positivity unless there is strong synergistic epistasis (see Figure 5). This expectation was

tested by using the simulations described above to estimate mean D' from doubletons. For the

cases of neutrality, $\gamma = 10$ and $\gamma = 50$ (with no epistasis), the mean D' values for cases 1 and 2

were – 1 and 1, respectively; the overall mean D' values were – 0.705, – 0.857 and – 0.922,

respectively.

Consistent with the findings of Garcia and Lohmueller (2021), neutral variants have a

smaller magnitude than selected variants of negative LD as measured by D'. However, this

effect is entirely due to the fact that the probability of case 2 increases as selection becomes

stronger. In other words, purifying selection itself can bias D' values at selected alleles to be

lower than those of neutral alleles, even if their frequencies are similar, so that this effect

should be taken into account when testing for HRI effects. To avoid it, one should use

unweighted values of D or $\bar{\sigma}_d$. It is, however, currently unclear how nonequilibrium scenarios

like population size changes might affect the signs of these statistics and whether unweighted

values of D would be the best to use in all cases.

Discussion

The role of complementation

A major finding of this work is the effect on LD statistics of the distinction between the gene

and sites models, which we discuss in detail below. In the accompanying paper (Johri and

Charlesworth 2025), we discuss the implications for population genetics statistics relevant to

the genetic load and inbreeding load caused by deleterious mutations. This raises the question

of how widely the gene model might apply to real organisms. The distinction between the

two models is based on the well-established finding of classical and molecular genetics that

two recessive loss-of-function mutations affecting the amino-acid sequence of the same

polypeptide usually fail to complement each other when present in the same cell in *trans* in a

double heterozygote, due to the fact that both copies of the polypeptide are non-functional. In

contrast, the same mutations present in *cis* in a double heterozygote result in the cell

containing one functional and one non-functional polypeptide, with near wild-type

performance (see Figure 1). This finding is of great historical importance, since it underlies

the recognition of the gene as a coding sequence whose sites could be separated by

recombination, and to Seymour Benzer's concept of the cistron as the functional genetic unit

(reviewed by Benzer 1957; Pontercorvo 1958; Hawley and Gilliland 2006).

If mutations within the same coding sequence fail to complement each other when in

trans, the assumptions of the gene model are satisfied. The sites model must usually apply to

mutations affecting different coding sequences, which necessarily complement each other

unless there are unusual types of epistasis between them, e.g., due to interactions between

mutations in different members of a multi-protein complex. Examples of such a failure of

complementation between non-allelic loss of function mutations are reviewed by Hawley and

Gilliland (2006) and Navarro-Quiles et al. (2024); these appear to be very rare compared with

complementation.

It should be noted, however, that there are exceptions to a lack of allelic

complementation (reviewed by Fincham et al. 1979, Chap. 4; Hawley and Gilliland 2006). In

such cases, loss-of-function mutations affecting the same polypeptide complement each

other, so that *trans* double heterozygotes show wild-type or near wild-type performance. In

many cases, some mutations within the same gene complement each other, whereas others do

not, allowing the construction of complementation maps (Fincham et al. 1979, Chap. 4).

Some mutations within the same gene may thus obey the gene model and others the sites

model. Allelic complementation has been best studied in fungi like *Neurospora*, and often

involves cases where multimers of the same polypeptide form part of the same protein

molecule (Fincham et al. 1979, pp. 348-353). In other cases, spatial interactions between

mutations in distinct functional subunits of the same polypeptide are involved (Fincham et al.

1979, p. 352; Eng et al. 2015). It is often found that interallelic complementation is

incomplete, when phenotypes such as enzyme activity are measured (Fincham et al. 1979, pp.

348-351), so that the distinction between the two models is somewhat blurred. For simplicity,

we have not attempted to include any of these complexities in our models.

It is, of course, less clear what happens to the heterozygous fitness effects of more minor effect mutations in the same coding sequence when configured in *cis* or *trans*, and there appears to be little relevant data. The simplest assumption is that the effects parallel those of loss-of-function mutations and are recessive or partially recessive, but are additive in their effects when in *trans*, as we have assumed here when setting k in Table 1 to ½. There are, of course, many other possibilities, including the extreme alternative of setting $k = h$ to represent the fitness effect of a heterozygous effect of a mutation at one site on a background of homozygosity for a mutation at another site. As shown in the section *Deterministic results for two selected sites*, the case of $k = ½$ is conservative with respect to the magnitude of positive LD among pairs of deleterious mutations, as a larger k results in a lower marginal fitness of $--$ haplotypes.

In the simulations of multiple mutations within the same gene, diploid genotypes with both *cis* and *trans* configurations of mutations are likely to be relatively abundant in a randomly mating population, unless selection is very intense. For example, with a scaled selection coefficient of $\gamma = 20$ and $h = 0.2$, the mean frequency of a deleterious mutation at a given site under the gene model with $k = ½$ is approximately 0.001 (see Table 3 of Johri and Charlesworth 2025). The mean number of mutations per haplotype with 1000 selected sites in a coding sequence is thus approximately 1. Since mean D values are small in this case (Table 2), a Poisson distribution should give a good approximation to the distribution of the number of mutations per haplotype. This implies a frequency of approximately 0.13 of individuals with two mutations in *cis* over a wild-type haplotype, 0.14 with two mutations in *trans*, and 0.06 with three or more mutations in *cis* over a wild-type haplotype; the frequencies of wild-type homozygotes and heterozygotes for single mutations are only 0.14 and 0.28, respectively. The frequency of individuals that are homozygous at one or more sites is approximately 0.001 in this example, so they will have a negligible effect on the haplotype dynamics compared with the heterozygous carriers of mutations, implying that the value of k can play only a small role when the number of selected sites is large.

The "gene-based model" of Thornton et al. (2013) and Sanjak et al. (2015) is similar to ours, in that it assumes that a *trans* heterozygote with n mutations in a gene has the same fitness as a homozygote with n mutations. They did not, however, consider the position effect, *i.e*, the effects of the difference in fitness between *trans* and *cis* heterozygotes. Clark (1998) evaluated the effects of allelic complementation or non-complementation between pairs of deleterious alleles on allele frequencies at a multiallelic locus, but did not examine

patterns of LD. Ragsdale (2022, Table S1) describes two versions of a two-site model. His
"simple dominance" model is identical with our sites model (allowing for the difference in
notation). His "gene-based" model is similar to our gene model, in that it assigns different
fitnesses to *cis* and *trans* double heterozygotes. However, it differs in several other respects,
e.g., by assigning the same fitness to single homozygotes, single heterozygotes, double
homozygotes, and *cis* double heterozygotes.

The situation with respect to mutations affecting the same non-coding regulatory
sequence such as a UTR, an enhancer or an insulator is somewhat unclear, although it seems
likely that mutations within the same cis-regulatory sequence should show similar *cis*/*trans*
behavior to mutations in the same polypeptide, fitting the gene model, while mutations in
different cis-regulators of the same gene should complement each other, and hence fit the
sites model. The phenomenon of transvection involves complementation between mutations
in different cis-regulatory sequences of the same gene when the two homologous
chromosomes or regions of homologs are paired in a cell nucleus, but failure of
complementation when they are unpaired (Lewis 1954; Duncan 2002; Mellert and Truman
2012; Galouzis and Prud'homme 2021); the normal situation is thus for them to fit the sites
model.

The role of epistasis
The extent to which deleterious mutations with minor effects on fitness have epistatic effects
rather than obeying an additive or multiplicative model for their combined effects, and the
typical nature of such interactions, is still an unsettled issue with respect to both theoretical
(Diaz-Colunga et al. 2023) and empirical (Bakerlee et al. 2022) questions. Attempts to
characterize epistasis for fitness components among deleterious mutations in a variety of
organisms have used analyses of the change in the means over generations of mutation
accumulation lines or populations subject to inbreeding. These experiments have yielded
somewhat conflicting results, although overall there seems to be a predominance of weak
synergistic epistasis (reviewed by Charlesworth 1998; Fry 2004; de Visser et al. 2011; García
Dominguez et al. 2019). A problem with interpreting these results is that data on fitness, or
its components such as viability or fecundity, are inherently noisy and there may be statistical
biases (due to regression towards the mean) that lead to an underestimation of synergistic
epistasis among deleterious mutations (Berger and Postma 2014).
Microbial systems have allowed estimates of the fitness consequences of combining
deletions of two or more genes, providing extensive information about epistasis among

drastically deleterious mutations, using haploid genomes (e.g., Costanzo et al. 2016; Kuzmin

et al. 2018). New technologies that allow the introduction of multiple mutations into the same

coding sequence have opened a path towards characterizing epistasis between mutations at

the intragenic level. This can be done either at the level of a specific biochemical function,

such as resistance to an antibiotic (e.g., Bershtein et al. 2006), fluorescence intensity of the

green fluorescent protein (e.g., Sarkisyan et al. 2016; Somermeyer et al. 2022), or by

measurements of the fitness effects of mutant strains in microbial competition experiments

(e.g., Bank et al. 2015, 2016; Puchta et al. 2016; Domingo et al. 2018; Bakerlee et al. 2022;

Chen et al. 2023; Papkou et al. 2023). Interestingly, although the stability of proteins is nearly

linearly reduced by increasing numbers of mutations (De Pristo et al. 2005; Bershtein et al.

2006), the effect of mutations on function or fitness in the experiments just cited seems to

show a predominance of synergistic epistasis among deleterious mutations, despite

complexities such as the dependence of patterns of interaction between the same mutations in

different environments (Bank et al. 2016) or between-species differences in fitness

landscapes (Somermeyer et al. 2022). However, there appear to be no reliable estimates of

the parameters involved in quantifying epistasis (such as our ϵ), making it difficult to

evaluate the relevance of our model of epistasis. The extent to which the sign of LD between

deleterious variants is affected by epistasis versus HRI remains an open question, but our

simulations suggest that synergistic epistasis is more likely to produce detectable negative

LD than HRI in genomic regions where recombination rates are not unusually low.

Linkage disequilibrium: theory

It has generally been accepted that multiplicative fitness models of multi-locus mutation and

selection in randomly mating, discrete generation populations predict an absence of linkage

disequilibrium when equilibrium has been reached in an infinite population (Haldane 1937;

Kimura and Maruyama 1966). This is confirmed by our analytical results for the two-site

version of the sites model with multiplicative fitnesses (section 2 of Supplementary File S1),

as well as numerical evaluations of the resulting equilibria (Figure 2 and Supplementary File

S2). Previous theoretical studies have found that synergistic epistasis on a multiplicative

scale results in negative LD when a positive sign of D is assigned to haplotypes with pairs of

mutant or wild-type alleles, whereas antagonistic epistasis causes positive LD (Kondrashov

1982, 1988; Charlesworth 1990; Shnol and Kondrashov 1993). Furthermore, in discrete

generation finite populations, there is a general expectation that negative LD will be

generated by Hill-Robertson interference in the absence of epistasis on the multiplicative

fitness scale, provided that linkage is sufficiently tight (Hill and Robertson 1966; Felsenstein

1974; Garcia and Lohmueller 2021; Roze 2021).

Since an absence of epistasis on a linear scale produces weakly synergistic epistasis

on a multiplicative scale, it might be expected that our deterministic two-site model would

result in small negative values of the two measures of LD, D and R, when the epistatic

parameter ϵ is equal to zero. The analytical results presented here confirm this expectation for

the sites model (see the Appendix and Supplementary File S1, sections 2 and 3, but the

properties of the gene model are significantly different (Appendix and section 4 of File S1).

The analytical results and Figure 2 show that the additive sites model exhibits negative values

of R that are very close to one-half the mutation rate when $\epsilon = 0$, and much larger negative

values when $\epsilon > 0$. However, the correlation coefficient R is often positive under the additive

gene model when $h < ½$, for sufficiently small values of ϵ.

The intragenic position effect that characterizes the gene model is expected to favor

positive LD, especially when the k parameter of Table 1 is less than one-half. As noted

earlier, a smaller value of k results in weaker selection against – – haplotypes, enhancing the

value of D. With $h = k = ½$ and no synergistic or antagonistic epistasis, the additive sites and

additive gene models have identical properties. Note, however, that setting $k = h$ and equating

the fitness of *cis* and *trans* double heterozygotes to $1 – 2hs$ or $(1 – 2hs)^2$ reduces to the sites

model, so that a *cis-trans* effect is a necessary condition for positive LD under the

assumptions made here, in the absence of antagonistic epistasis.

Similarly, the finding of positive LD in simulations of the gene model without

epistasis, at least with $h < ½$ and when selection is sufficiently strong (see *Simulation results*

for multiple sites; *linkage disequilibrium with no epistasis*), makes sense given the intragenic

position effect on fitness. The results in Table 2 show that the approximations for LD derived

in the Appendix for the multi-site gene model (Equations A19 and A21) agree well with the

simulation results with $k = ½$ when the selection regime is the same for all sites and selection

is strong relative to drift, so that this approach provides useful insights into when positive LD

between variants within the same gene is expected under the gene model.

Positive LD has also been found for the sites model in recent theoretical studies that

included finite population effects. Roze (2021) studied a diploid finite population model with

two loci under a multiplicative version of the sites model, assuming $\gamma = 2N_e s \gg 1$ but $s \ll 1$.

He found small positive mean D and σ_d values with $h < 0.25$ in both his approximate

analytical treatment and in multi-locus simulations (see his Figure 1). Similarly, the analysis

of the additive sites model for a two-locus system in a finite population by Ragsdale (2022) found positive mean values of D and σ_d^1 for sufficiently small values of h, provided that γ was sufficiently large. We similarly observe small positive values of D and σ_d for the sites model when $\gamma=100$ and $h=0.0$ (Table 2). Ragsdale's model of an intragenic position effect also generated positive mean LD. No such effect is expected in an infinite population (see Figure 2).

It is probable that this positive mean LD under the sites model without epistasis reflects the fact that, with a high degree of recessivity, $+ +$ haplotypes behave as dominant with respect to fitness. Their rate of change in frequency per generation due to selection is thus approximately $2sq^2$, since the marginal fitness of $+ +$ is 1 and the mean fitness of the population is approximately $1 - 2sq^2$ when $h = 0$ (see Equations A3). In contrast, the rate of change of $- -$ is approximately $- sq^3$. In the case of complete recessivity, an increase in the frequency of $+ +$ caused by drift is thus unopposed by selection, and an increase in the frequency of $- -$ is only weakly opposed, so that the resulting chance net increase in D is likely to persist, especially as the accompanying decrease in the frequency of $+ -$ and $- +$ haplotypes is unopposed by selection.

However, for very small values of the recombination rate and $h << \frac{1}{2}$, negative LD was found for the sites model by both Roze (2021) and Ragsdale (2022), reflecting the build-up of associative overdominance caused by the interaction between drift and selection, such that a predominance of $+ -$ and $- +$ haplotypes is generated when linkage is close (Zhao and Charlesworth 2016; Becher et al. 2020; Gilbert et al. 2020; Abu-Awad and Waller 2023). The interactions between selection and drift when there is dominance are thus rather complex.

Our multi-locus simulations of the sites model with multiplicative rather than linear fitnesses failed to generate significantly positive LD (Table S2 in Supplementary File S3), when deleterious versus beneficial alleles were used for the sign criterion. The absence of positive LD in our simulations may be due to the stronger effects of HRI in multi-site compared with two-site models. The existence of positive LD among derived or selected mutations discussed here should not be confused with the much more frequently observed positive D when an excess frequency of haplotypes with pairs of minor alleles is used as the sign criterion (e.g., Figure 3). As explained in the section *Simulation results for multiple sites: linkage disequilibrium with no epistasis*, this effect is expected even for neutral variants (Potapova and Kondrashov 2023).

929 **Patterns of linkage disequilibrium in population genetic data**

930 Previous theoretical and simulation studies have suggested that, while HRI and synergistic

931 epistasis create negative LD between selected variants, processes such as antagonistic

932 epistasis, admixture, and population bottlenecks could also result in positive LD (Sohail et al.

933 2017; Sandler et al. 2021; Garcia and Lohmueller 2021; Ragsdale 2022). Several recent

934 studies have obtained estimates of signed LD among variants in coding sequences in natural

935 populations of *D. melanogaster*, humans, *Capsella grandiflora* and *Schizophyllum commune*

936 (Pool et al. 2012; Sohail et al. 2017; Sandler et al. 2021; Garcia and Lohmueller 2021;

937 Ragsdale 2022; Stolyarova et al. 2022). Garcia and Lohmueller (2021) found a larger

938 magnitude of negative LD among doubleton nonsynonymous variants compared with

939 synonymous variants in humans when using Lewontin's D' statistic (Lewontin 1964). As

940 described in the subsection *Effects of weighting towards low frequencies on LD statistics, D'*

941 is biased towards negative values when low frequency variants are used, and differences in

942 D' between neutral and selected sites do not necessarily reflect the effects of epistasis or HRI

943 alone. Corrections for this bias are thus needed to disentangle the relative contribution of the

944 causes of signed LD at selected sites, if this method is to be used.

945 Sohail et al. (2017) and Sandler et al. (2021) used estimates of LD based on the excess or

946 deficiency over the Poisson expectation of the variance in the numbers of minor alleles over a

947 set of defined genomic regions, which is proportional to the sum of pairwise D values over

948 the regions in question, whereas Stolyarova et al. (2022) used estimates of the correlation

949 coefficients between pairs of variants, polarizing with respect to minor versus major alleles.

950 Both of these methods are likely to give higher magnitudes of the LD statistic for variants at

951 higher frequencies. Ragsdale (2022) polarized LD using the ancestral/derived criterion; he

952 employed the σ_d^1 measure, whose denominator involves the cross products of the four variant

953 frequencies at a pair of diallelic sites (see Equation 7). This will tend to give *higher* values

954 for sites with lower frequencies, for a given correlation coefficient.

955 These methodological differences across studies make it hard to compare the results, but

956 nevertheless there are some patterns in common. Notably, in most cases signed LD between

957 synonymous sites was found to be significantly higher than that between nonsynonymous

958 variants, with the exception of the analysis of the hypervariable mushroom species *S.*

959 *commune* by Stolyarova et al. (2022), where the opposite pattern was observed – see their

960 Appendix 2, Figure 1.

There seems to be a preponderance of slightly negative values of LD between loss-of-

function (LOF) mutations (presumed to be the most harmful), but positive values between

pairs of synonymous and nonsynonymous variants. Because these patterns applied to loosely

linked variants in humans and *Drosophila*, and the extent of under-dispersion of loosely

linked loss-of-function alleles was found to be much higher than explained by simulations

that included the demographic history of humans, Sohail *et al.* (2017) proposed that

synergistic epistasis was responsible.

A possible explanation for a predominance of negative signed LD between loosely

linked LOF mutations is that these are kept at very low frequencies in the population. This

means that a finite sample from the population has a high probability of not containing the $-$

$-$ haplotype at a pair of diallelic sites. This automatically generates $D < 0$, even if D is zero in

the population as a whole, as would be expected for most pairs of sites that are not tightly

linked. With two sites with $D = 0$ in the population and a minor allele frequency q at each

site, a sample of size k has a probability of approximately $[1 - \exp(-kq)]^2$ of segregating at

both sites, and a probability of approximately kq^2 of containing the $-$ $-$ haplotype, giving a

net probability of $1 - kq^2/[1 - \exp(-kq)]^2$ that a segregating sample contains no $-$ $-$

haplotypes. With $k = 1000$ and $q = 0.001$, this probability is equal to 0.9975 for 10000

replicate pairs. But with $q = 0.01$, it is only 0.9. Despite the fact that the expected value of D

is zero in both cases, with $q = 0.001$ a finite set of pairs of segregating sites is likely to have a

high probability of containing none or only a few $-$ $-$ haplotypes, resulting in a large

preponderance of cases with $D < 0$. This is illustrated in section 7 of Supplementary File S1.

It is thus possible that observations of negative LD between loosely linked or independent

LOF mutations, such as those shown in Figure S6 of Sohail et al. (2017) are the result of this

sampling effect. It follows that it cannot safely be inferred that synergistic epistasis is the

cause of this pattern unless a very large number of pairs of variants is analysed; our analytical

and simulation results confirm that it does indeed generate negative LD for the sites model

but not necessarily for the gene model (see Figures 2 - 4).

Our modeling work adds a new perspective on findings of positive LD among closely

linked pairs of putatively selected variants. If fitness is governed by the gene model, then LD

between variants at non-synonymous sites within the same gene should be relatively larger

than under the sites model, and the sign should often be positive, provided the effects of

sampling just discussed do not overwhelm the effects of selection. However, as we have

discussed in the section *The role of complementation*, the gene model is unlikely to apply

between different genes but is likely to be applicable within a gene. Our models thus make

the following predictions about signed LD patterns in population genomics data. (*a*) LD between derived or deleterious alleles at nonsynonymous sites within genes should be positive or nearly zero (assuming that the population is panmictic and at equilibrium). (*b*) LD between derived or deleterious nonsynonymous variants in the same gene should be higher than between variants in different genes. (*c*) There should be higher LD between derived nonsynonymous vs derived synonymous variants within the same gene.

There are, however, several caveats that need to be borne in mind when using these findings to interpret data. First, the sign of LD statistics when calculated using minor/major alleles (as opposed to using derived/ancestral) is almost always positive, unless the minor alleles are very rare (Potapova and Kondrashov 2023). This applies to neutral sites (Potapova and Kondrashov 2023) and to deleterious variants that obey the sites model as well as the gene model – see Figure 3. It is thus not surprising that the studies which polarize their variants by the minor/major allele criterion (Pool *et al*. 2012; Sohail et al. 2017; Sandler et al. 2021; Stolyarova et al. 2022) often found positive LD. Second, weighting towards low allele frequencies (Good 2022) also tends to promote positive LD, as described in the subsection *Effects of weighting towards low frequencies on LD statistics*, and should thus be avoided if the sign of LD is used for inferences about selection. It follows that patterns of signed LD using this criterion do not necessarily imply effects of admixture, non-equilibrium demography, or antagonistic/complementary epistasis, which have all been proposed as possible explanations in these previous studies.

This leaves the study of humans by Ragsdale (2022), who found that LD between derived missense variants was found to be significantly positive within the same domain of a gene, as opposed to near-zero values between SNPs in different domains, even after matching the distance between SNPs found within compared between domains. This observation could be explained by a gene model operating at the level of domains within genes, perhaps because there is higher probability of complete failure of complementation within domains versus across the entire gene. Ragsdale's estimates of $\bar{\sigma}_d^1$ from the Yoruba population (which is closest to having a stable size) between missense variants within genes is ~0.02 (his Figure 8). As the average length of a gene in humans is quite large (~ 60 kb), it is difficult to compare this estimate to that obtained in our simulations. Other estimates of $\bar{\sigma}_d^1$ within 100 base pairs provided by Ragsdale (2002) range from 0.25-0.3 (within the same gene; his Figure S17), ~0.5 (within the same domain; his Figure S18), and ~0.2 (outside domains; his Figure S19). While estimates of $\bar{\sigma}_d^1$ within 100 bases in our simulations of human-like

populations are within this range (Table S4 in Supplementary File S3), the large SEs make it

difficult to make any reliable conclusion. In general, however, both demographic processes

(like admixture and bottlenecks) as well as selective processes such as positive epistasis can

result in positive LD between derived nonsynonymous variants, and joint modeling of these

processes in a specific population would be required to distinguish between these scenarios,

taking into account important features such as recombination hotspots, which strongly

influence the rates of recombination within genes in taxa such as mammals and flowering

plant (e.g., Myers et al. 2010; Choi et al. 2013; Baudat et al. 2014; Sandler et al. 2021; Joseph

et al. 2024).

There are multiple reasons why within-gene LD could be higher than between-gene

LD: (a) On average, there is a much higher frequency of recombination between pairs of sites

within different genes than within the same gene (b) Fitness follows the gene model within

genes but not between genes (c) There might be antagonistic epistasis within genes but not

between genes. While it might be possible to differentiate between (a) and (b) by comparing

the patterns of LD between species with compact genes with little spacing and/or low

recombination rates between genes with patterns in species with high recombination rates

between genes, it will be harder to distinguish between (b) and (c).

In summary, it will probably be quite difficult to distinguish among the different

possible explanations for patterns of signed LD from population genomic data, but it is

essential to use the ancestral/derived or beneficial/deleterious rather than the minor/major

allele criterion for the sign of LD. The problem is further complicated by the fact that, if

average gene lengths and rates of recombination are sufficiently large, even patterns of LD

within genes may mostly reflect genome-wide patterns generated by demographic processes

rather than the effects of selection. Analyses of large-scale resequencing data in the light of

population genetic models of the type investigated here, using suitable choices of species

with respect to their genomic properties and demographic histories, offer the best way to

approach this problem.

1055

Evolution of the recombination rate within genes

Previous models of recombination in the presence of selection against recurrent deleterious

mutations have shown that higher rates of recombination are likely to evolve if there is

synergistic epistasis among mutations at different sites, and reduced rates to evolve if there is

antagonistic epistasis (Feldman et al. 1980; Kondrashov 1982, 1988; Charlesworth 1990;

Otto and Feldman 1997; Roze 2021). As we have shown, under the gene model, pairs of

deleterious mutations result in a smaller fitness cost in *cis* than when they are in *trans*,

promoting positive linkage disequilibrium. This suggests that recombination modifiers that

increase the recombination rate within genes are likely to be disfavored, whereas modifiers

that reduce them will be favored. This is consistent with the fact that recombination hotspots

are often found in promoter regions or transcription start sites upstream of genes, and that

genes are often relatively "cold" with respect to recombination rates (Myers et al. 2010;

Joseph et al. 2024). Further modeling of this possible consequence of the *cis-trans* effect for

the evolution of recombination rates would seem to be desirable.

1070

Acknowledgments

1072 We thank Kirk Lohmueller, Denis Roze and three anonymous reviewers for their helpful

1073 comments on an earlier version of the manuscript. This research was conducted using

1074 computational resources provided by ITS Research Computing at the University of North

1075 Carolina at Chapel Hill. PJ was funded by the National Institute of General Medical Sciences

1076 of the National Institutes of Health under the award number R35GM154969.

1077

Literature Cited

1079 Abu-Awad D, Waller DM. 2023. Conditions for maintaining and eroding pseudo-overdominance
1080 and its contribution to inbreeding depression. Peer Comm. J. 3:e8.
1081

1082 Altshuler D, Donnelly P, The International HapMap Consortium. 2005. A haplotype map of the
1083 human genome. Nature. 437:1299–1320.
1084

1085 Bakerlee CW, Nguyen Ba AN, Shulgina Y, Rojas Echenique JI, Desai MM. 2022. Idiosyncratic
1086 epistasis leads to global fitness-correlated trends. Science 376:630-635.
1087

1088 Bank C, Hietpas RT, Jensen JD, Bolon DN. 2015. A systematic survey of an intragenic epistatic
1089 landscape. Mol. Biol. Evol. 31:229-238.
1090

1091 Bank C, Matuszewski S, Hietpas RT, Jensen JD. 2016. On the (un)predictability of a large intragenic
1092 fitness landscape. Proc. Natl. Acad. Sci. USA 113:14085-14090.
1093

1094 Baudat F, Imai Y, De Massy B. 2014. Meiotic recombination in mammals: localization and
1095 regulation. Nat. Rev. Genet. 14:794-806.
1096

1097 Becher H, Jackson B, Charlesworth B. 2020. Patterns of genetic variability in genomic regions with
1098 low rates of recombination. Curr. Biol. 30:94-100.
1099

1100 Benzer S. 1957. The elementary units of heredity. In: McElroy WD, Glass B, editors. The Chemical
1101 Basis of Heredity. Baltimore, MD: Johns Hopkins Press. p. 70-93.
1102

1103   Berger D, Postma E. 2014. Biased estimates of diminishing-returns epistasis? Empirical evidence
1104   revisited. Genetics 198:1417-1420.
1105

1106   Bershtein S, Segal M, Bekerman S, Tokuriki N, Tawfik DS. 2006. Robustness–epistasis link shapes
1107   the fitness landscape of a randomly drifting protein. Nature 444:929-931.
1108

1109   Bürger, R. 2000. The mathematical theory of selection, recombination and mutation. Chichester UK,
1110   John Wiley & Sons.
1111

1112   Charlesworth B. 1990. Mutation-selection balance and the evolutionary advantage of sex and
1113   recombination. Genet Res. 55 :199-221.
1114

1115   Charlesworth B. 1998. The effect of synergistic epistasis on the inbreeding load. Genet. Res. 71:85-
1116   89.
1117

1118   Charlesworth B. 2015. Causes of natural variation in fitness: evidence from studies of Drosophila
1119   populations. Proc. Natl. Acad. Sci. USA 12:1662-1669.
1120

1121   Charlesworth B, Jain K. 2014. Purifying selection, drift, and reversible mutation with arbitrarily high
1122   mutation rates. Genetics 198:1587-1602.
1123   Chen Y, Hu R, LI K, Zhang Y, Fu L, Zhang J, Si T. 2023. Deep mutational scanning of an oxygen-
1124   Independent fluorescent protein CreiLOV for comprehensive profiling of mutational and epistatic
1125   effects. ACS Synth. Biol. 12:1461-1473.
1126

1127   Choi K, Zhao X, Kelly KA, Venn O, Higgins JD, Yelina NE, Hardcastle TJ, Ziolkowski PA,
1128   Copenhaver GP, Franklin FCH, et al. 2013. Arabidopsis meiotic crossover hot spots overlap with
1129   H2A.Z nucleosomes at gene promoters. Nat Genet. 45(11):1327–1336.
1130

1131   Clark AG. 1998. Mutation-selection balance with multiple alleles. Genetica 102:41–47.
1132

1133   Costanzo M, VanderSluis B, Koch EN, Baryshnikova A, Pons C, Tan G, et al. 2016. A global
1134   genetic interaction network maps a wiring diagram of cellular function. Science 353:aaf1420-
1135   1421-1414.
1136

1137   Crow JF. 1958. Some possibilities for measuring selection intensities in man. Hum. Biol. 30:1-13.
1138

1139   Crow JF. 1970. Genetic loads and the cost of natural selection. In: Kojima K, editor. Mathematical
1140   Topics in Population Genetics. Berlin: Springer-Verlag. p. 128-177.
1141

1142   Crow JF. 1993. Mutation, mean fitness, and genetic load. Oxf. Surv. Evol. Biol. 9:3-42.
1143

1144   Dabi A, Schrider DR. 2025. Population size rescaling significantly biases outcomes of forward-in-
1145   time population genetic simulations. Genetics 229:iyae180.
1146

1147   De Pristo MA, Weinreich DM, Hartl DL. 2005. Missense meandering in sequence space: a
1148   biophysical view of protein evolution. Nature Rev. Genet. 6:678-687.
1149

1150   Domingo J, Diss G, Lehner B. 2018. Pairwise and higher-order genetic interactions during the
1151   evolution of a tRNA. Nature 558:117-121.
1152

De Visser JAGM, Cooper TF, Elena SF. 2011. The causes of epistasis. Proc. R. Soc. B. 278:3617-3624.

Diaz-Colunga J, Skwara A, Gowda K, Diaz-Uriarte R, Tikhonov M, Bajic D, Sanchez A. 2023. Global epistasis on fitness landscapes. Phil. Trans. R. Soc. B. 378:20220053.

Duncan IW. 2002. Transvection effects in Drosophila. Annu. Rev. Genet. 36:521-556.

Eng T, Guacci V, Koshland D. 2015. Interallelic complementation provides functional evidence for cohesin–cohesin interactions on DNA. Mol. Biol. Cell 26:4224-4235.

Ewens WJ. 2004. Mathematical Population Genetics. 1. Theoretical Introduction. New York: Springer.

Eyre-Walker A, Woolfit M, Phelps T. 2006. The distribution of fitness effects of new deleterious amino acid mutations in humans. Genetics. 173:891–900.

Feldman MW, Christiansen FB, Brooks LD. 1980. Evolution of recombination in a constant environment. Proc Natl Acad Sci U S A. 77:4838–4841.

Felsenstein J. 1965. The effect of linkage on directional selection. Genetics 52:349-63.

Felsenstein J. 1974. The evolutionary advantage of recombination. Genetics 78:737-756.

Ferrari T, Feng S, Zhang X, Mooney J. 2024. Towards simulation optimization: An examination of the impact of scaling on coalescent and forward simulations. bioRxiv 2024.04.27.591463.

Fincham JRS, Day PR, Radford A. 1979. Fungal Genetics. Berkeley, CA: University of California Press.

Fry JD. 2004. On the rate and linearity of viability declines in Drosophila mutation-accumulation experiments. Genetics 166:797-806.

Galouzis CC, Prud'homme B. 2021. Transvection regulates the sex-biased expression of a fly X-linked gene. Science 371:396-400.

Garcia JA, Lohmueller KE. 2021. Negative linkage disequilibrium between amino acid changing variants reveals interference among deleterious mutations in the human genome. PloS Genet. 17:e1009676.

García-Dominguez S, García C, Quesada H, Caballero A. 2019. Accelerated inbreeding depression suggests synergistic epistasis for deleterious mutations in *Drosophila melanogaster*. Heredity 123:709-722.

Gilbert KJ, Pouyet F, Excoffier L, Peischl S. 2020. Transition from background selection to associative overdominance promotes diversity in regions of low recombination. Curr. Biol. 30:101-107.

Good BH. 2022. Linkage disequilibrium between rare mutations. Genetics 220:iyac004.

Haldane JBS. 1937. The effect of variation on fitness. Am. Nat. 71:337-349.

Haldane JBS. 1927. A mathematical theory of natural and artificial selection. Part V. Selection and mutation. Proc. Camb. Phil. Soc. 23:838-844.

Haller, BC, Messer, PW. 2023 SLiM 4: Multispecies eco-evolutionary modeling. Am. Nat. 201: E127–E139.

Hawley RS, Gilliland WD. 2006. Sometimes the result Is not the answer: The truths and the lies that come from using the complementation test. Genetics 174:5-15.

Hill WG, Robertson A. 1966. The effect of linkage on limits to artificial selection. Genet. Res. 8:269-294.

Johri P, Charlesworth B. 2025. A gene-based model of fitness and its implications for genetic variation: Genetic and inbreeding loads. (*companion paper*)

Johri P, Charlesworth B, Jensen JD. 2020. Toward an evolutionarily appropriate null model: Jointly inferring demography and purifying selection. Genetics 215:173-192.

Johri P, Riall K, Becher H, Excoffier L, Charlesworth B, Jensen JD. 2021. The impact of purifying and background selection on the inference of population history: problems and prospects. Mol Biol Evol. 38(7):2986–3003.

Joseph J, Prentout D, Laverré A, Duret L. 2024. High prevalence of PRDM9-independent recombination hotspots in placental mammals. Proc. Natl. Acad. Sci. USA 121:e2401973121.

Kaiser VB, Charlesworth B. 2010. Muller's ratchet and the degeneration of the *Drosophila miranda* neo-Y chromosome. Genetics 185:339-348.

Karlin S. 1975. General two-locus selection models: some objectives, results and interpretations. Theor. Pop. Biol. 7:364-398.

Kim BY, Huber CD, Lohmueller KE. 2017. Inference of the distribution of selection coefficients for new nonsynonymous mutations using large samples. Genetics. 206:345–361.

Kimura M, Maruyama T. 1966. The mutational load with epistatic gene interactions in fitness. Genetics 54:1303-1312.

Kondrashov AS. 1982. Selection against harmful mutations in large sexual and asexual populations. Genet. Res 44:325-332.

Kondrashov AS. 1988. Deleterious mutations and the evolution of sexual reproduction. Nature 336:435-440.

Kong A, Frigge ML, Masson G, Besenbacher S, Sulem P, Magnusson G, Gudjonsson SA, Sigurdsson A, Jonasdottir Aslaug, Jonasdottir Adalbjorg, et al. 2012. Rate of *de novo* mutations and the importance of father's age to disease risk. Nature. 488(7412):471–475.

Kryukov GV, Pennachio LA, Sunyaev S. 2007. Most rare missense alleles are deleterious in humans: implications for complex disease and association studies. Am. J. Hum. Genet. 80:727-739.

Kuzmin E, VanderSluis B, Wang W, Tan G, Despande R, Cheng Y, et al. 2018. Systematic analysis of complex genetic interactions. Science 360:eaao1729.

Kyriazis CC, Robinson JA, Lohmueller KE. 2023. Using computational simulations to model deleterious variation and genetic load in natural populations. Am. Nat. 302:737-752.

Lewis EB. 1954. The theory and application of a new method of detecting chromosomal rearrangements in *Drosophila melanogaste*r. Am. Nat. 88:225-239.

Lewontin RC. 1964. The interaction of selection and linkage. I. General considerations, heterotic models. Genetics 49:49-67.

Mellert DJ, Truman JW. 2012. Transvection Is common throughout the Drosophila genome. Genetics 191:1121-1141.

Misra S, Crosby MA, Mungall CJ, Matthews BB, Campbell KS, et al. 2002. Annotation of the *Drosophila melanogaster* euchromatic genome: a systematic review. Gen. Biol. 3:Research0083.

Morton NE, Crow JF, Muller HJ. 1956. An estimate of the mutational damage in man from data on consanguineous marriages. Proc. Natl. Acad. Sci. USA 42:855-863.

Muller HJ. 1950. Our load of mutations. Am. J. Human Genet. 2:111-176.

Myers S, Bottolo L, Freeman C, McVean G, Donnelly G. 2010. A fine-scale map of recombination rates and hotspots across the human genome. Science 310:321-324.

Navarro-Quiles C, Lup SD, Munoz-Nortes T, Candela H, Micol JL. 2024. The genetic and molecular basis of haploinsufficiency in flowering plants. Trends. Plant Sci. 29:72-95.

Nordborg M, Franklin IR, Feldman MW. 1995. Effects of cis-trans viability selection on some two-locus models. Theor. Pop. Biol. 47:365-392.

Ohta T, Kimura M. 1969. Linkage disequilibrium due to random genetic drift. Genet. Res. 13:47-55.

Ohta T, Kimura M. 1971. Linkage disequilibrium between two segregating nucleotide sites under steady flux of mutations in a finite population. Genetics 68:571-580.

Otto SP, Feldman MW. 1997. Deleterious mutations, variable epistatic interactions, and the evolution of recombination. Theor Popul Biol. 51:134–147.

Papkou A, Garcia-Pastor L, Escudero JA, Wagner A. 2023. A rugged yet easily navigable fitness landscape. Science 382:eadh3860.

Pontecorvo G. 1958. Trends in Genetic Analysis. New York, NY: Columbia University Press.

Pool JE, Corbett-Detig RB, Sugino RP, Stevens KA, Cardeno CM, Crepeau MW, et al. 2012. Population genomics of sub-Saharan *Drosophila melanogaster*: African diversity and non-African admixture. PloS Genet 8:e1003080.

Potapova NA, Kondrashov AS. 2023. Positive association between alleles at selectively neutral loci. Biorxiv 2023.06.14.544962:1-12.

Puchta O, Czeke B, Czaja H, D. T, Sanguinetti G, Kudla G. 2016. Network of epistatic interactions within a yeast snoRNA. Science 352:840-844.

Ragsdale AP. 2022. Local fitness and epistatic effects lead to distinct patterns of linkage disequilibrium in protein-coding genes. Genetics 221:iyac097.

Roze D. 2021. A simple expression for the strength of selection on recombination generated by interference among mutations. Proc. Natl. Acad. Sci. USA 118:e2022805118.

Sandler G, Wright SI, Agrawal AF. 2021. Patterns and causes of signed linkage disequilibria in flies and plants. Mol. Biol. Evol. 38:4310-4321.

Sanjak JS, Long AD, Thornton KR. 2015. A model of compound heterozygous, loss-of-function alleles Is broadly consistent with observations from complex-disease GWAS datasets. PloS Genet. 13:e1006573.

Sarkisyan KS, Bolotin DA, Meer MV, Usmanova DR, Mishin AS, Sharanov GV, et. al. 2016. Local fitness landscape of the green fluorescent protein. Nature 533:397-401.

Shnol EE, Kondrashov AS. 1993. Effect of selection on the phenotypic variance. Genetics 134:995-996.

Sohail M, Vakhrusheva OA, Sul JH, Pulit SA, Francioli LC, Genome of the Netherlands Consortium, et al. 2017. Negative selection in humans and fruit flies involves synergistic epistasis. Science 356:539-542.

Somermeyer LG, Fleiss A, Mishin AS, Bozhanova NG, Igolkina AA, Meiler J, Pujol M-EA, Putintseva EV, Sarkisyan KS, Kondrashov FA. 2022. Heterogeneity of the GFP fitness landscape and data-driven protein design. Elife 11:e75842.

Stolyarova AV, Neretina TV, Zvyagina EA, Fedotova AV, Kondrashov AS, Bazykin GA. 2022. Complex fitness landscape shapes variation in a hyperpolymorphic species. Elife 11:e76073.

Thornton KR, Foran AJ, Long AD. 2013. Properties and modeling of GWAS when complex disease risk is due to non-complementing, deleterious mutations in genes of large effect. PloS Genet. 9:e1003258.

Wright S. 1931. Evolution in Mendelian populations. Genetics 16:97-159.

Zhao L, Charlesworth B. 2016. Resolving the conflict between associative overdominance and background selection. Genetics 203:1315-1334.

Appendix

Deterministic treatment of the sites and gene models: recursion equations for the

haplotype frequencies

The full recursion relations for a general fitness matrix and mutation model are given below; these

differ slightly in form from those usually used for models of mutation and selection, where the

changes in haplotype frequencies due to selection and mutation are added to each other (e.g.,

Haldane [1937]). This is because the small equilibrium values of D resulting from the interaction

between mutation and selection are very sensitive to approximations, and incorrect signs of D can

result if approximations are made prematurely.

Assume that there are two diallelic loci, A and B, with alleles A_1, A_2 and B_1, B_2, with

subscript 1 corresponding to + in Table 1 and subscript 2 to –. Let the frequencies of the four

haplotypes A_1B_1, A_1B_2, A_2B_1 and A_2B_2 be x_1, x_2, x_3 and x_4, respectively. The corresponding allele

frequencies are p_1, q_1 (A_1 and A_2) and p_2, q_2 (B_1 and B_2). The coefficient of linkage disequilibrium

is $D = x_1x_4 - x_2 x_3$. Assume that A_2 and B_2 are both deleterious and are kept at such low frequencies

that mutations can be treated as unidirectional, with mutation rates u_1 and u_2 for A_1 to A_2 and B_1 to

B_2, respectively. Let w_{ij} be the fitness of individuals formed by a pair of haplotypes (i and j) with

frequencies x_i and x_j. Under the gene model, the fitnesses of the two double heterozygotes (w_{14} and

w_{23}) are unequal, in contrast to the usual assumptions of two-locus models (e.g., Karlin 1975). The

marginal fitness of haplotype is $w_i = \sum_j x_j w_{ij}$ and the population mean fitness is $\overline{w} = \sum_i w_i$.

Recombination occurs between the two loci with frequency r per generation. Random mating is

assumed.

Assume that the order of events within a generation is selection, mutation and

recombination. The post-selection frequency of haplotype i is $x_i^* = x_i w_i / \overline{w}$. Mutation changes the

haplotype frequencies to new values x_i^{**} as follows:

1374

1375 $$x_1^{**} = x_1^*(1 - u_1)(1 - u_2), \quad x_2^{**} = (x_2^* + x_1^* u_2)(1 - u_1) \qquad (A1a)$$

1376 $$x_3^{**} = (x_3^* + x_1^* u_1)(1 - u_2), \quad x_4^{**} = x_4^* + x_1^* u_1 u_2 + x_2^* u_1 + x_3^* u_2 \qquad (A1b)$$

1377

In the absence of recombination, these are the haplotype frequencies in the new generation.

If recombination occurs post-mutation, the haplotype frequencies in the new generation are given

by the following expressions, where the triple asterisks indicate haplotype frequencies after

mutation without any selection within the generation in question (*i.e*, the frequencies given by

Equations (A1) without the asterisks on the right-hand side); the effects of selection on the double

heterozygotes are captured by the terms in w_{14} and w_{23}:

1384

$$x_1' = x_1^{**} - r(x_1^{***}x_4^{***}w_{14} - x_2^{***}x_3^{***}w_{23})\bar{w}^{-1} \qquad (A2a)$$

$$x_2' = x_2^{**} + r(x_1^{***}x_4^{***}w_{14} - x_2^{***}x_3^{***}w_{23})\bar{w}^{-1} \qquad (A2b)$$

$$x_3' = x_3^{**} + r(x_1^{***}x_4^{***}w_{14} - x_2^{***}x_3^{***}w_{23})\bar{w}^{-1} \qquad (A2c)$$

$$x_4' = x_4^{**} - r(x_1^{***}x_4^{***}w_{14} - x_2^{***}x_3^{***}w_{23})\bar{w}^{-1} \qquad (A2d)$$

1389

These recursion equations can be used to obtain the equilibrium values for the system,

as well as the approach to equilibrium. Given the symmetry of the fitness matrix in Table 1,

for the system studied here we can write $x_2 = x_3 = x = q(1-q) - D$; $x_4 = y = q^2 + D$; $x_1 = z = 1$

$- 2x - y$; $q = x + y$. It follows that only the recursions for x and y need to be considered. Let

the mutation rate per locus from $+$ to $-$ be u. If $u << hs$, the frequency q of the $-$ allele at each

locus is $<< 1$ and reverse mutations can be ignored. The coefficient of linkage disequilibrium

is denoted by $D = yz - x^2$. Together with the fitness values for each genotype, these variables

can be used to determine the recursion relations for the frequencies of the four haplotypes, x_1

$(+\,+)$, x_2 $(-\,+)$, x_3 $(+\,-)$ and x_4 $(-\,-)$.

Denote the marginal fitnesses of the $-\,+$ and $+\,-$ haplotypes by w_x and that of the $-\,-$

haplotype by w_y; the marginal fitness of the $+\,+$ haplotype is w_z. From Table 1, we obtain the

following expressions for the marginal fitnesses under the sites model:

1402

$$w_x = 1- [h + q + 2h\epsilon x + (1+h)\epsilon y]s \qquad (A3a)$$

$$w_y = 1- 2[h + (1-h)q)] + h\epsilon + (1-h)\epsilon q]s \qquad (A3b)$$

$$w_z = 1- 2[hq + h\epsilon y]s \qquad (A3c)$$

1406

1407   The equivalent expressions for the gene model are as follows:

1408

$$w_x = 1- [h + 2(1-h)x + (1+k-h)y + (1+k)\epsilon y]s \qquad (A3d)$$

$$w_y = 1- 2[h + (1+k-2h)x + (1-h)y](1+\epsilon)s \qquad (A3e)$$

$$w_z = 1- 2hs(q + \epsilon y) \qquad (A3f)$$

1412

1413 For both models, the mean fitness of the population is given by:

1414

$$\bar{w} = 2xw_x + yw_y + zw_z \qquad\qquad (A3g)$$

If these expressions are substituted into Equations (A1), using specified values of the mutation rate, recombination rates and selection parameters, a numerical solution to the equilibrium version of the recursions could be found by an iterative method. It is, however, simpler to run the recursions to close to equilibrium, thereby obtaining estimates of the equilibrium values of x and y and all derived quantities of interest. The initial starting point of the recursions was arbitrarily chosen to have $D = 0$, with q given by the solution of the linear equation that approximates the cubic equation describing mutation-selection equilibrium at a single locus when q is sufficiently small but non-negligible (Bürger 2000, p.100):

$$\hat{q} \approx \frac{u}{(u+hs)} \qquad\qquad (A4)$$

Properties of equilibria with respect to LD under the additive and multiplicative sites models

As shown in part 1 of Supplementary File S1, when $r = 0$, a formula for the equilibrium value of D, $\hat{D}$, can be obtained for a general symmetric fitness model (Equation S1.3). Neglecting the small terms in the mutation rate in this expression, we have:

$$\hat{D}[\bar{w}^2 - w_x^2] \approx -\lambda \hat{y} \hat{z} \qquad\qquad (A5a)$$

where

$$\lambda = w_x^2 - w_y w_z \qquad\qquad (A5b)$$

The analysis in section 2 of Supplementary File S1 shows that $\hat{D} = 0$ under the multiplicative sites model with no epistasis (Equation S1.6b).

Equations (S1.7) and (S1.8) in section 3 of Supplementary File S1 show that, for the additive sites model with no epistasis, we have:

$$\lambda = \{h^2 + 2h(1 - 2h)q + [1 - 4h(1 - h)]q^2\}s^2 \qquad\qquad (A5c)$$

We can approximate q for an equilibrium population by equating it to the equilibrium solution $\hat{q}$ given by Equation (A4). We can also approximate the haplotype frequencies by

1447 neglecting the small terms involving $\widehat{D}$, so that $\hat{x} \approx \hat{q}(1 - \hat{q}), \hat{y} \approx \hat{q}^2, \hat{z} = (1 - \hat{q})^2$. Equation

1448 (S10b) then gives the following approximation:

1449

1450 $$\overline{w}^2 - w_x^2 \approx 2\{h[1 - 2\hat{q}(1 - 2\hat{q})] + \hat{q}(1 - 2\hat{q})]s \qquad (A6)$$

1451

1452 Equations (A5c) and (A.6) yield the following approximation for $\widehat{D}$:

1453

1454 $$\widehat{D} \approx -\tfrac{1}{2}s\hat{q}^2(1 - \hat{q})^2\{h^2 + 2h(1 - h)\hat{q} + [1 - 4h(1 - h)]\hat{q}^2\}\{h[1 - 2\hat{q}(1 - 2\hat{q})] + \hat{q}(1 - 2\hat{q})\}^{-1}$$

1455 $$(A7)$$

1456

1457 This expression can be extended to include the effect of recombination when r, D and

1458 s are sufficiently small that their second-order terms can be neglected. In this case, Equations

1459 (A2) imply that the change in D caused by recombination is approximately $- rD$. If this term

1460 is added to the right-hand side of the recursion equation for D (Equation S1.3), and $\overline{w}^2 D$ is

1461 approximated by D, Equation (A5a) becomes:

1462

1463 $$\widehat{D}[\overline{w}^2 - w_x^2] \approx -(\lambda \hat{y}\hat{z} + r\widehat{D}) \qquad (A8)$$

1464

1465 This yields the following approximate expression for $\widehat{D}$:

1466

1467 $$\widehat{D} \approx -\tfrac{1}{2}s\hat{q}^2(1 - \hat{q})^2\{h^2 + 2h(1 - h)\hat{q} + [1 - 4h(1 - h)]\hat{q}^2\}\{h[1 - 2\hat{q}(1 - 2\hat{q})] + \hat{q}(1 - 2\hat{q}) + r/s\}^{-1}$$

1468 $$(A9a)$$

1469 This equation can be converted into an expression for the equilibrium value of the correlation

1470 coefficient, $\hat{R}$, by dividing by $\hat{q}(1 - \hat{q})$:

1471

1472 $$\hat{R} \approx -\tfrac{1}{2}s\hat{q}(1 - \hat{q})\{h^2 + 2h(1 - h)\hat{q} + [1 - 4h(1 - h)]\hat{q}^2\}\{h[1 - 2\hat{q}(1 - 2\hat{q})] + \hat{q}(1 - 2\hat{q}) + r/s\}^{-1}$$

1473 $$(A9b)$$

1474 If second-order terms in $\hat{q}$ are neglected, this expression simplifies to:

1475

1476 $$\hat{R} \approx -\tfrac{1}{2}s\hat{q}[h^2 + 2h(1 - h)\hat{q}][h(1 - 2\hat{q}) + \hat{q} + r/s]^{-1} \qquad (A9c)$$

1477

1478   These expressions must be modified if epistasis is present, which introduces extra

1479 terms of the order of ϵs. Use of Equations (A3) leads to the following expression for the

1480 additive sites model when $\epsilon > s$:

1481

1482 $$\hat{R} \approx -\tfrac{1}{2}\epsilon \hat{q}\,(1 - \hat{q})[h + (1 - h)\hat{q}]\{h[1 - 4\hat{q}(1 - \hat{q})] + \hat{q}(1 + 2\epsilon) + r/s\}^{-1} \quad (A9d)$$

1483

1484 **Properties of equilibria with respect to LD under the additive and multiplicative gene**

1485 **models**

1486 The same type of approach can be applied to the additive and multiplicative gene models, as

1487 described in section 4 of Supplementary File S1. Equation (S1.16c) yields the following

1488 approximation for $\hat{R}$:

1489

1490 $$\hat{R} \approx \hat{q}(1 - \hat{q})\{\hat{q}[(1 - h - k) - 2h(1 - 2h)s] - h^2 s - [h + (1 + k - 2h)\hat{q}]\epsilon\}[h + 2(1 - 2h)\hat{q} + r/s]^{-1}$$

1491 $$(A10)$$

1492   Similarly, for the multiplicative gene model without synergistic or antagonistic

1493 epistasis, Equation (S1.16d) yields:

1494

1495 $$\hat{R} \approx \hat{q}^2(1 - \hat{q})\{(1 - h - k) + [k - h(2 - 3h)]s\}[h + 2(1 - 2h)\hat{q} + r/s]^{-1} \quad (A11)$$

1496

1497 For $h \leq \tfrac{1}{2}$, and $h \leq k \leq \tfrac{1}{2}$, $\hat{R} > 0$.

1498

Extension to a large number of selected sites

It is possible to derive an approximation for the LD measures considered above in the case of

a large number of sites subject to the same selection regime. It is assumed that Equation (A8)

applies to a given pair of sites, provided that expressions for w_x, w_y for such a pair of sites, as

well as for $\overline{w}$, can be obtained. The equilibrium allele frequency at each site, $\hat{q}$, is treated

initially as a given. For the purpose of comparisons with the simulation results, we consider

only the additive gene model with $k = \tfrac{1}{2}$. The effects of LD on the distribution of the number

of deleterious mutations per haplotype are ignored, so that the probability that a haplotype

carries i mutations (P_i) is assumed to follow a Poisson distribution with mean $\mu = G\hat{q}$, where

$\hat{q}$ is the equilibrium frequency of mutant alleles at a given site ($G \gg 1$):

1509

$$P_i = \frac{\mu^i}{i!} \exp(-\mu) \qquad (A12)$$

Consider first the net marginal fitness of a haplotype that lacks mutations at a pair of

randomly chosen sites, w_z. Such a haplotype may carry 0, 1, 2, ... $G-2$ mutations located at

other sites. Ignoring LD, the probability of i mutations at other sites is P_i; the marginal fitness

of such a haplotype is denoted by w_{zi}. Taking into account the other haplotypes that are

encountered to form diploid individuals, and using the fitness function for the gene model

with $k = \frac{1}{2}$, we have:

1517
$$w_{z0} = 1 - hs \sum_{j=0}^{\infty} jP_j = 1 - \mu hs \qquad (A13a)$$

1518

1519   For $i \geq 1$, we have:

1520
$$w_{zi} = 1 - iP_0 hs - \frac{s}{2}\sum_{j=1}^{\infty}(j + i)P_j$$

1521
$$= 1 - ie^{-\mu}hs - \frac{s}{2}[\mu + i(1 - e^{-\mu})] \qquad (A13b)$$

1522   This yields:

1523
$$w_z = \sum_{i=0}^{\infty} w_{zi}P_i = 1 - 2\mu e^{-\mu}hs - \frac{s}{2}\sum_{i=1}^{\infty}[\mu + i(1 - e^{-\mu})]P_i$$

1524

1525
$$w_z = 1 - [\mu - (1 - 2h)\mu e^{-\mu}]s \qquad (A13c)$$

1526

1527   Next, consider the marginal fitness of a haplotype with a mutation at one of the two chosen

1528   sites. As before, the probability that such a haplotype carries i mutations at other sites is P_i

1529   and the marginal fitness of such a haplotype is:

1530

1531
$$w_{xi} = 1 - (i + 1)P_0 hs - \frac{s}{2}\sum_{j=1}^{\infty}(j + i + 1)P_j \quad (i = 0, 1, ...)$$

1532
$$= 1 - (i + 1)e^{-\mu}hs - [\mu + (i + 1)(1 - e^{-\mu})]\frac{s}{2} \qquad (A14a)$$

1533   so that:

1534
$$w_x = \sum_{i=0}^{\infty} w_{xi}P_i = 1 - (\mu + 1)e^{-\mu}hs - [\mu + (\mu + 1)(1 - e^{-\mu})]\frac{s}{2} \qquad (A14b)$$

1535

1536   The same procedure can be carried out for the marginal fitness of a haplotype with

1537   mutations at both chosen sites, w_y. The fitness of such a haplotype that carries i mutations at

1538   other sites is:

1539
$$w_{yi} = 1 - (i + 2)P_0 hs - \frac{s}{2}\sum_{j=1}^{\infty}(j + i + 2)P_j \quad (i = 0, 1, ...)$$

1540
$$= 1 - (i + 2)e^{-\mu}hs - [\mu + (i + 2)(1 - e^{-\mu})]\frac{s}{2} \qquad (A15a)$$

1541   and:

1542
$$w_y = \sum_{i=0}^{\infty} w_{yi}P_i = 1 - (\mu + 2)e^{-\mu}hs - [\mu + (\mu + 2)(1 - e^{-\mu})]\frac{s}{2} \qquad (A15b)$$

1543

1544   If second-order terms in s are neglected, we have:

1545

1546 $w_y w_z - w_x^2 \approx -2\mu e^{-\mu} hs - \mu(1 - e^{-\mu})s - (\mu + 2)e^{-\mu} hs - [\mu + (\mu + 2)(1 - e^{-\mu})]\frac{s}{2}$

1547 $\qquad\qquad + 2(\mu + 1)e^{-\mu} hs + [\mu + (\mu + 1)(1 - e^{-\mu})]s$

1548 $\qquad\qquad = \mu e^{-\mu}\left(\frac{1}{2} - h\right)s$ $\qquad\qquad\qquad\qquad$ (A16)

1549

1550   Furthermore, if $\hat{q} \ll 1$ and $s \ll 1$, we have:

1551

1552 $$\bar{w} \approx w_z = 1 - [(2h - 1)\mu e^{-\mu} + \mu]s \qquad\qquad (A17)$$

1553

1554   Using this expression together with Equation (A14b), we obtain:

1555

1556 $\bar{w}^2 - w_x^2 \approx -4\mu e^{-\mu} hs - 2\mu(1 - e^{-\mu})s + 2(\mu + 1)e^{-\mu} hs + [\mu + (\mu + 1)(1 - e^{-\mu})]s$

1557 $\qquad\qquad = [(\mu - 1)e^{-\mu}(1 - 2h) + 1]s$ $\qquad\qquad\qquad$ (A18)

1558

1559   Substituting from Equations (A16) and (A18) into Equation (A8) yields the expression:

1560

1561 $$\hat{D} \approx \hat{q}^2 \mu e^{-\mu}\left(\frac{1}{2} - h\right)[(\mu - 1)e^{-\mu}(1 - 2h) + 1]^{-1} \qquad (A19)$$

1562

1563   Equations (A13c) and (A14b) can be used to obtain an equation for the value of μ and

1564   hence $\hat{q} = \mu/G$. Provided that $q \ll 1$, the change in q due to selection at a given site is given

1565   by:

1566 $$\Delta q_s \approx q(w_x - \bar{w}) \approx q(w_x - w_z) \qquad\qquad (A20a)$$

1567

1568   Substituting from Equations (A13c) and (A14b) and simplifying, the following expression is

1569   obtained:

1570 $$\Delta q_s \approx -\tfrac{1}{2}sq[1 + (\mu - 1)e^{-\mu}(1 - 2h] \qquad\qquad (A20b)$$

1571

1572   At mutation-selection equilibrium, this must balance the change in q due to mutation, which

1573   is approximately equal to the mutation rate u. The equilibrium value of μ is then determined

1574   by the following transcendental equation, which can be solved numerically by Newton-

1575   Raphson iteration:

1576

1577
$$\frac{2Gu}{s} \approx \hat{\mu}[1 + (\hat{\mu} - 1)e^{-\hat{\mu}}(1 - 2h] \qquad (A21a)$$

1578

With semi-dominance ($h = \frac{1}{2}$), this equation is equivalent to the standard equation

$\hat{q} = 2u/s$ (Haldane 1927), as expected from the equivalence of the sites and gene models in

this case. For smaller values of h, $\hat{q}$ is reduced below this value, as seen in the two-site model

described above. Importantly, for $h = 0$, $\hat{q}$ does not obey the standard single locus result for a

completely recessive mutation, $\hat{q} = \sqrt{u/s}$ (Haldane 1927), but instead takes much larger

values.

But for sufficiently small values of G, $\hat{\mu}$ is $<< 1$, and Equation (A21a) can be

approximated by:

1587
$$\frac{Gu}{s} \approx \hat{\mu}[h + \hat{\mu}(1 - 2h)] \qquad (A21b)$$

1588

1589   When $h = 0$, this yields the standard equation for complete recessivity. When $\hat{\mu} \ll h$, this

1590   reduces to the standard Haldane equation $\hat{q} = u/(hs)$.

1591

1592

1593

1594

1595 **TABLES**

1596
1597

1598 **Table 1. Models of the fitness effects of two sites with a fixed selection coefficient**

1599

1600 The "sites" model

1601

	++	+ -	- +	- -
+ +	1	$1 - hs$	$1 - hs$	$1 - 2hs - e_1$
+ -	$1 - hs$	$1 - s$	$1 - 2hs - e_1$	$1 - s - hs - e_2$
- +	$1 - hs$	$1 - 2hs - e_1$	$1 - s$	$1 - s - hs - e_2$
- -	$1 - 2hs - e_1$	$1 - s - hs - e_2$	$1 - s - hs - e_2$	$1 - 2s - e_3$

1602

1603 $e_1 = 2\epsilon hs$; $e_2 = \epsilon(1 + h)s$; $e_3 = 2\epsilon s$.

1604

1605 The "gene" model

1606

	+ +	+ -	- +	- -
+ +	1	$1 - hs$	$1 - hs$	$1 - 2hs - e_1$
+ -	$1 - hs$	$1 - s$	$1 - s$	$1 - (1+k)s - e_2$
- +	$1 - hs$	$1 - s$	$1 - s$	$1 - (1+k)s - e_2$
- -	$1 - 2hs - e_1$	$1 - (1+k)s - e_2$	$1 - (1+k)s - e_2$	$1 - 2s - e_3$

1607
1608

1609 $e_1 = 2\epsilon hs$; $e_2 = (1 + k)\epsilon s$; $e_3 = 2\epsilon s$.

1610

1611

1612
1613

Table 2. Statistical summaries of linkage disequilibrium between selected alleles for the gene

and sites models, with fixed scaled selection coefficient $\gamma = 2Ns$. 1000 selected sites were

simulated, with $Nu = 0.005$ and $Nr = 0.01$, where N is the population size ($N = 1000$), and u

and r are the mutation and recombination rate per site/generation respectively. $\bar{D}$ and $\bar{\sigma}_d$

were calculated for alleles that were less than 100 sites apart. The sign of the LD statistic $\bar{\sigma}_d$

was assigned either by giving a positive sign to cases with excesses over random

combinations of pairs of selectively deleterious alleles, or to cases with excesses of

combinations of minor alleles. The means and standard errors (SEs) for 1000 replicate

simulations are shown. The predicted values of $\bar{D}$ for the gene model with $\gamma = 20$ and 100 are

also shown; these were obtained as described in the text; the corresponding values for the

sites model are expected to be close to zero.

1625

selected allele		$h=0.0$		$h=0.2$		$h=0.5$	
		gene	sites	gene	sites	gene	sites
$\gamma = 2$							
$\bar{D}$	mean	-0.00021	-0.00032	-0.00016	-0.00026	-0.00016	-0.00021
	SE	0.00004	0.00004	0.00004	0.00004	0.00004	0.00004
$\bar{\sigma}_d$	mean	-0.00241	-0.00280	-0.00184	-0.00247	-0.00175	-0.00238
	SE	0.00054	0.00036	0.00056	0.00047	0.00051	0.00055
$\gamma = 20$							
$\bar{D}$	mean	0.00019	-0.00010	0.00008	-0.00006	-0.00006	-0.00002
	SE	0.00002	0.00003	0.00002	0.00002	0.00002	0.00002
	predicted	0.00015	0	0.00009	0	0	0
$\bar{\sigma}_d$	mean	0.00733	-0.00080	0.00347	-0.00052	-0.00153	-0.00022
	SE	0.00070	0.00045	0.00061	0.00056	0.00053	0.00057
	predicted	0.0052	0	0.00313	0	0	0
$\gamma = 100$							
$\bar{D}$	mean	0.00008	0.00011	0.00004	-0.00003	-0.00001	0.00003
	SE	0.00001	0.00001	0.00001	0.00001	0.00002	0.00002
	predicted	0.00004	0	0.00002	0	0	0
$\bar{\sigma}_d$	mean	0.00447	0.00361	0.00247	-0.00126	0.00005	0.00211
	SE	0.00083	0.00043	0.00087	0.00065	0.00142	0.00161
	predicted	0.00393	0	0.00157	0	0	0

minor allele							
$\gamma = 2$							
$\bar{D}$	mean	0.00288	0.00593	0.00305	0.00473	0.00289	0.00289
	SE	0.00010	0.00014	0.00009	0.00012	0.00009	0.00009
$\bar{\sigma}_d$	mean	0.03604	0.05513	0.03831	0.05033	0.03637	0.03644
	SE	0.00111	0.00120	0.00113	0.00121	0.00103	0.00109
$\gamma = 20$							
$\bar{D}$	mean	0.00019	0.00038	0.00009	0.00002	-0.00005	-0.00002
	SE	0.00002	0.00004	0.00002	0.00002	0.00002	0.00002
$\bar{\sigma}_d$	mean	0.00737	0.00618	0.00360	0.00136	-0.00121	-0.00013
	SE	0.00070	0.00056	0.00062	0.00058	0.00053	0.00057
$\gamma = 100$							
$\bar{D}$	mean	0.00008	0.00012	0.00004	-0.00003	-0.00001	0.00003
	SE	0.00001	0.00001	0.00001	0.00001	0.00002	0.00002
$\bar{\sigma}_d$	mean	0.00451	0.00375	0.00251	-0.00119	0.00015	0.00211
	SE	0.00083	0.00043	0.00088	0.00066	0.00142	0.00161

1626

1627

1628

Table 3: Statistical summaries of linkage disequilibrium under the gene and sites models in

human-like populations, using a gamma distribution of scaled selection coefficients with

shape parameter 0.23 and mean $\bar{\gamma} = 2N\bar{s}$. 2000 selected sites were simulated, with $Nu =$

0.00025 and $Nr = 0.0002$, where N is the population size ($N = 1000$), u is the mutation rate

per site/generation, and r is the rate of crossing over between adjacent sites. The signs of

$\bar{D}$ and $\bar{\sigma}_d$ were assigned either by giving a positive sign to cases with excesses over random

combinations of pairs of selectively deleterious alleles, or to cases with excesses of

combinations of minor alleles. The means and standard errors (SEs) for 1000 replicate

simulations are shown.

1638

selected alleles		h=0.0		h=0.2		h=0.5	
		gene	sites	gene	sites	gene	sites
$\bar{\gamma} = 850$							
$\bar{D}$	mean	0.00015	0.00031	0.00101	0.00030	-0.00096	-0.00105
	SE	0.00045	0.00035	0.00062	0.00066	0.00063	0.00074
$\bar{\sigma}_d$	mean	0.01130	0.00650	0.01587	0.00345	-0.00706	-0.00345
	SE	0.00472	0.00353	0.00591	0.00547	0.00525	0.00600
$\bar{\sigma}_d^1$	mean	0.46777	0.25534	0.61505	0.14143	-0.12457	0.20649
	SE	0.17992	0.08900	0.21642	0.11249	0.09229	0.20202
minor alleles							
$\bar{\gamma} = 850$							
$\bar{D}$	mean	0.00123	0.00137	0.00319	0.00319	0.00244	0.00285
	SE	0.00046	0.00036	0.00063	0.00065	0.00071	0.00078
$\bar{\sigma}_d$	mean	0.01924	0.02183	0.03972	0.03556	0.02356	0.03175
	SE	0.00473	0.00373	0.00601	0.00548	0.00560	0.00625
$\bar{\sigma}_d^1$	mean	0.54411	0.51751	0.94606	0.55386	0.20582	0.60968
	SE	0.17966	0.09122	0.21699	0.11278	0.09449	0.20269

1639

1640

1641

1642

1643
1644

1645 **FIGURES**

1646
1647

1648 **Figure 1:** A schematic of the additive gene versus additive sites models. Note that the

1649 additive sites model is approximately equivalent to the standard multiplicative fitness model.

1650

1651

1652

1653

1654

1655

Figure 2. Linkage disequilibrium (LD) as a function of the synergistic epistasis coefficient

(ϵ; left-hand panels) and dominance coefficient (h; right hand panels) for the deterministic

equilibrium model with two sites, with $u = 0.0001$, $s = 0.01$, and $r = 0$. The upper panels are

the results for the additive sites model; the lower panels are for the additive gene model with

$k = ½$. LD is measured by the correlation coefficient (R) multiplied by 100. The solid curves

are the exact solutions to the recursion equations for D (Equations S1.2 of Supplementary

File S1). The dashed curves are the analytic approximations for R; see the Appendix,

Equations (A.9) and (A.10). The horizontal lines in the lower panels indicate zero values of

R.

1665

1666

1667

Figure 3: Statistical summaries of linkage disequilibrium as measured by $\bar{\sigma}_d$ under the gene and sites models for a gamma distribution of scaled selection coefficients with shape parameter 0.3 and a mean of $\bar{\gamma} = 2N\bar{s}$. 1000 selected sites were simulated, with $Nu = 0.005$ and $Nr = 0.01$, where N is the population size ($N = 1000$), u is the mutation rate per site/generation, and r is the rate of crossing over between adjacent sites. $\bar{\sigma}_d$ was calculated for alleles that were less than 100 basepairs apart. The sign of $\bar{\sigma}_d$ was assigned either by giving a positive sign to cases with excesses over random combinations of pairs of selectively deleterious alleles, or to cases with excesses of combinations of minor alleles (see the Methods section). The means and standard errors (SEs) for 1000 replicate simulations are shown. Similar results for $\bar{D}$ are shown in Figure S1 in Supplementary File S3.

Figure 4: Statistical summaries of the $\bar{\sigma}_d$ measure of linkage disequilibrium between all pairs of sites segregating for deleterious mutations for gene and sites model with epistasis, for closely linked pairs of alleles (1-100 basepairs apart). The simulation parameters were the same as in Figure 3. The sign of the LD statistic $\bar{\sigma}_d$ was assigned either by giving a positive sign to cases with excesses over random combinations of pairs of selectively deleterious alleles or to cases with excesses of combinations of minor alleles. The error bars represent the SEs across replicates. Similar results for $\bar{D}$ are shown in Figure S2 in Supplementary File S3.

1688

Figure 5. Mean linkage disequilibrium (LD) as a function of the epistasis coefficient (ϵ),

from stochastic simulations of the two-site haploid model with $r = 0$. LD is measured by the

$\bar{\sigma}_d$ statistic described in the text (multiplied by 100). The results for 10^6 replicate simulations

with two different values of $\gamma = 2Ns$ are shown, with $N = 1000$. The dashed curve with filled

markers shows the population values of LD with no weighting towards low allele

frequencies; the full curve with open markers is for doubleton variants in a sample of size

100. The dashed curve with open markers shows population values with a weight of 10

toward low frequency variants; the full curve with filled markers is for a weight of 100. See

the main text for further details.

1698

1699

1700

1701