1 A structural MRI marker predicts individual differences in impulsivity and classifies

2 patients with behavioral-variant frontotemporal dementia from matched controls

4 Valérie Godefroy[1], Anaïs Durand[2], Marie-Christine Simon[4], Bernd Weber[5], Joseph Kable[6], Caryn

5 Lerman[7], Fredrik Bergström[8,9], Richard Levy[2], Bénédicte Batrancourt[2], Liane Schmidt, Hilke

6 Plassmann[2,3]* and Leonie Koban[1]*

**Author affiliations:**

(1) Université Claude Bernard Lyon 1, CNRS, INSERM, Centre de Recherche en Neurosciences de

Lyon CRNL U1028 UMR5292, F-69500, Bron, France ;

(2) Paris Brain Institute (ICM), INSERM U 1127, CNRS UMR 7225, Sorbonne University, Paris,

France UMR 7225, Sorbonne University, Paris, France;

(3) Marketing Area, INSEAD, Fontainebleau, France,

(4) Institute for Nutrition Science, University of Bonn, Bonn, Germany;

(5) Center for Economics and Neuroscience, University of Bonn, Bonn, Germany;

(6) Department of Psychology, University of Pennsylvania, Philadelphia, PA, USA;

(7) USC Norris Comprehensive Cancer Center

(8) Faculty of Psychology and Educational Sciences, University of Coimbra, Portugal

(9) Department of Psychology, University of Gothenburg, Sweden

*shared senior authorship

**Short title: MRI-BASED PREDICTION OF IMPULSIVITY**

**Correspondence to:** Valérie Godefroy, PhD

Université Claude Bernard Lyon 1, CNRS, INSERM, Centre de Recherche en Neurosciences de Lyon

CRNL U1028 UMR5292

CH Le Vinatier, Bât. 462, Neurocampus 95 Bd Pinel - 69675 BRON CEDEX, France

E-mail: vlrgodefroy@gmail.com

And :Leonie Koban, PhD, CRNL, CH Le Vinatier, Bât. 462, Neurocampus 95 Bd Pinel - 69675 BRON

CEDEX, France ; email: leonie.koban@cnrs.fr

**Acknowledgments**

This study was funded by an ANR Tremplin-ERC grant to HP, a Sorbonne Emergence Grant to HP

and LK, and an ERC Starting Grant to LK. Study 2 was funded by National Cancer Institute Grants

R01-CA-170297 to J.W.K. and C.L. and R35-CA-197461 to C.L. Study 3 was funded by grant ANR-

10-IAIHU-06 from the program 'Investissements d'avenir', by grant FRM DEQ20150331725 from the

foundation 'Fondation pour la recherche médicale', and by the ENEDIS company. FB was supported

by Fundação para a Ciência e Tecnologia (CEECIND/03661/2017). We thank all the participants and

organizers of the three studies mentioned in this paper as well as all the students who contributed to

help data collection.

**Data availability statement**

Deidentified data for the three studies are available at https://figshare.com/s/a090ef2531a7d7b9a2e4
and upon request to the first author.

**ABSTRACT**

Impulsivity and higher preference for sooner over later rewards (i.e., delay discounting) are transdiagnostic markers of many psychiatric and neurodegenerative disorders. Yet, their neurobiological basis is still debated. Here, we aimed at 1) identifying a structural MRI signature of delay discounting in healthy adults, and 2) validating it in patients with behavioral variant frontotemporal dementia (bvFTD)—a neurodegenerative disease characterized by high impulsivity. We used a machine-learning algorithm to predict individual differences in delay discounting rates based on whole-brain grey matter density maps in healthy male adults (Study 1, N=117). This resulted in a cross-validated prediction-outcome correlation of $r$=0.35 ($p$=0.0028). We tested the validity of this brain signature in an independent sample of 166 healthy adults (Study 2) and its clinical relevance in 24 bvFTD patients and 18 matched controls (Study 3). In Study 2, responses of the brain signature did not correlate significantly with discounting rates, but in both Studies 1 and 2, they correlated with psychometric measures of trait urgency—a measure of impulsivity. In Study 3, brain-based predictions correlated with discounting rates, separated bvFTD patients from controls with 81% accuracy, and were associated with the severity of disinhibition among patients. Our results suggest a new structural brain pattern—the Structural Impulsivity Signature (SIS)—which predicts individual differences in impulsivity from whole-brain structure, albeit with small-to-moderate effect sizes. It provides a new brain target that can be tested in future studies to assess its diagnostic value in bvFTD and other neurodegenerative and psychiatric conditions characterized by high impulsivity.

**Keywords:** brain signature; machine-learning; dementia; decision-making; delay discounting; intertemporal choice; prediction

## BACKGROUND

Impulsivity is the tendency to act in a rush and to seek immediate rewards without consideration of potentially negative long-term consequences [1]. Trait impulsivity varies substantially within the general population, with high impulsivity being a hallmark of many psychiatric and neurological conditions [2]. Despite the many negative consequences of high impulsivity for health and life in general [3,4], its neurobiological correlates are still unclear, and it is unknown whether individual differences in impulsivity can be reliably predicted based on structural brain features [5–7]. Neurobiological measures of impulsivity could help to understand the mechanisms and disentangle the heterogeneity of symptoms related to maladaptive behavior and decision-making. Brain signatures of impulsivity could also constitute new targets for diagnosis and treatment. They might aid in the diagnosis and monitoring of conditions such as behavioral variant frontotemporal dementia (bvFTD)—a neurodegenerative disorder characterized by frontal and temporal brain atrophy, with high impulsivity and inappropriate behaviors as core symptoms [8]. In this study, we aimed at developing a structural brain signature of individual differences in impulsivity, and tested whether it could accurately classify patients with bvFTD from matched healthy controls.

The idea that any psychological construct would depend on only one or a few isolated brain regions has been more and more challenged. A new paradigm of "brain signatures" (or "neuromarkers") promoting a multivariate *brain patterns* view has therefore emerged, to complement the traditional univariate brain mapping approach examining brain regions independently [9]. Brain signatures are predictive models of mental events or of individual variables (such as impulsivity) that take into account distributed information across multiple brain systems [10]. Brain signatures using structural data are increasingly used in the field of translational neuroimaging, especially for applications in patients with neurodegenerative conditions [11]. One of the greatest advantages of these predictive models which predict behavior from brain features is that they can be tested across studies, labs and populations to challenge their generalizability. We used this brain signature approach to identify a network of spatially distributed structural features associated with impulsivity, as measured by delay discounting. The present study applies the "component process" framework of brain signatures [11]. Instead of predicting a given heterogenous condition such as bvFTD, we aimed at identifying a predictive model of a key symptom (i.e., impulsivity), which is a common factor across

different diseases. This framework is also suited to the purpose of predicting a specific patient's

clinical profile in a perspective of personalized medicine.

Several arguments support the idea that delay discounting—how much people prefer smaller

sooner over larger later rewards—is a reliable measure of stable individual differences in a specific

facet of impulsivity (that is the urgency to get short-term rather than long-term reward). Individual

differences in delay discounting are relatively stable over time and show significant genetic heritability

[12–14]. Delay discounting moreover constitutes a potential transdiagnostic marker of conditions with high

impulsivity since it has been found to be altered across multiple psychiatric [15] and neurodegenerative

conditions [16]. Recent studies have therefore started to investigate the neurobiological basis of

individual differences in delay discounting [7,14,17–23]. However, less is known about how these candidate

brain markers of delay discounting are expressed in psychiatric and neurological conditions

characterized by increased impulsivity.

Characterized by multiple impulsivity-related symptoms, bvFTD is a good example to

demonstrate the clinical potential (in particular for diagnosis) of a structural brain signature of delay

discounting. BvFTD is the most common clinical variant of syndromes associated with predominant

degeneration of the prefrontal and temporal regions as well as the basal ganglia. It is characterized by

significant changes in personality and behavior including disinhibition (socially inappropriate and

generally impulsive behaviors), as well as executive function deficits [8]. Brain regions known to be

related to delay discounting such as the orbitofrontal cortex (OFC), ventromedial prefrontal cortex

(vmPFC) and ventral striatum [24–26] are often affected in bvFTD [27,28]. Relatedly, most studies found an

alteration of delay discounting in bvFTD patients compared to controls [16,29–32].

Here, we first trained and cross-validated a structural MRI-based brain signature in a healthy

adult population (Study 1, N=117) using LASSO-PCR (least absolute shrinkage and selection

operator-principal component regression)—an established machine-learning algorithm [33,34] —to

predict individual differences in delay discounting rates from subjects' grey matter maps (N=117).

Brain markers of individual differences need to be tested in different and completely independent

samples and studies to establish their robustness and generalizability [10]. Thus, in Study 2, we tested

the replicability of the brain signature in a second independent sample of healthy adults (N=166). In

Study 3, we tested the validity of the structural brain signature in a clinical population of patients with

behavioral variant frontotemporal dementia, who often show high impulsivity and were shown to be

steeper discounters (N = 42, including 24 bvFTD patients and 18 matched controls) [35]. If a consistent

pattern of grey matter density across the brain can reliably predict delay discounting and more

generally impulsivity, then the brain-predicted discounting should be higher in bvFTD patients than in

controls and should be related to the level of clinically assessed impulsivity in patients. In addition to

testing the generalizability of the brain signature developed in Study 1, we analyzed the topographical

distribution of the most important structural alterations contributing to differences of brain-predicted

delay discounting.

## MATERIALS AND METHODS

### Participants

The research reported here complies with all relevant ethical regulations. The study protocols were approved by the institutional review board of Bonn University's Medical School (Study 1), by the University of Pennsylvania Institutional Review Board (Study 2), and by the French Ethics Committee "Comité de Protection des Personnes Sud Méditerranée I" (Study 3).

*Study 1*

In Study 1, participants were recruited in the context of a seven-week dietary intervention study (https://osf.io/rj8sw/?view_only=af9cba7f84064e61b29757f768a8d3bf) at the University of Bonn in Germany. In this study, only male participants were recruited, with the following inclusion criteria: age between 20 and 60 years, right-handedness, non-smoker, no excessive drug or alcohol use in the past year, no psychiatric or neurological disease, body mass index (BMI) between 20 and 34, no other chronic illness or medication, following a typical Western diet without dietary restrictions, and no MRI exclusion criteria (e.g., large tattoos, metal in the body,). For the present purpose, we used only the behavioral and structural MRI data collected during a baseline session before the dietary intervention. N=117 participants were tested for the baseline session of Study 1. However, four participants were excluded from the present analyses due to being outliers on grey matter density maps (three participants) and due to very incoherent choices at the intertemporal choice task (one participant). Thus, the data of a total of 113 participants was used for the analyses.

*Study 2*

In Study 2, participants were recruited in the context of a ten-week cognitive training study (registered at clinicaltrials.gov as Clinical trial reg. no. NCT01252966) at the University of Pennsylvania, USA. Individuals between 18 and 35 years of age who reported home computer and internet access were recruited. Exclusion criteria were: an IQ score of <90 on Shipley Institute of Living Scale, self-reported history of neurological, psychiatric, or addictive disorders (excluding nicotine), positive breath alcohol reading (>0.01), color blindness, left-handedness, and claustrophobia. Here, we focused on behavioral and structural MRI data collected during the baseline session before the cognitive training. In Study 2, N=166 participants (mean age=24.5, 59% male) were included in the baseline session and all were included in our data analyses.

*Study 3*

For Study 3, participants were recruited in the context of a clinical study at the Paris Brain Institute,

France (clinicaltrials.gov: NCT03272230). This study was designed to investigate the behavioral

correlates and neural bases of neuropsychiatric symptoms associated with behavioral variant

frontotemporal dementia (bvFTD). BvFTD patients were recruited in two tertiary referral centers, at the

Pitié-Salpe☐trière Hospital and the Lariboisière Fernand-Widal Hospital, in Paris. Patients were

diagnosed according to the International Consensus Diagnostic Criteria [8]. To be included, bvFTD

patients had to present a Mini-Mental State Evaluation (MMSE) score of at least 20. Healthy controls

(HC) were recruited by an online announcement. Inclusion criteria included a MMSE score of at least

27 and matching the demographic characteristics of the bvFTD group. In total, 24 bvFTD patients

(mean age=66.6, 66.6% male) and 18 controls matched to patients for age and sex (mean age=62.6,

44.4% male) were recruited in this clinical study (see Supplementary table 1). Data of all participants

were used for our analyses.

**Intertemporal choice tasks**

*Study 1*

During the intertemporal choice (ITC) task performed in an MRI scanner, participants in Study 1 were

presented with 108 trials offering a choice between a smaller sooner (SS) reward option and a larger

later (LL) reward option [14]. Participants were informed that one of their choices could be paid out at the

end of the experiment, which made their choices non-hypothetical and incentive-compatible. The two

options were displayed on the left or right of the screen (position randomized) for 4 seconds.

Participants used their left or right index finger to press the response key corresponding to their choice

(left index for left option or right index for right option). The option chosen by the participant was then

highlighted by a yellow frame which remained on the screen until the end of the 4 second trial. Trials

were presented in randomized order (see Koban et al., 2023 for further details on the trial structure).

*Study 2*

During the ITC performed in an MRI scanner, participants had to make 120 choices between the same

smaller immediate reward ($20 today) and a varying larger reward available after a longer delay (e.g.,

$40 in a month) [36]. Participants were informed that one of their choices could be paid out at the end of

the experiment, which made their choices non-hypothetical and incentive-compatible. Each trial

started with the presentation of the amount and delay of the larger later option. Once subjects had

made their choice, a checkmark on the screen indicated if the larger later option was chosen and a "X"

indicated that the immediate option was chosen for 1 s. Subjects had 4 s to make their choice.

*Study 3*

In Study 3, participants performed two ITC tasks on a computer screen, one using monetary rewards

(from 8 to 35 euros) and one using food rewards (from 8 to 35 chocolates) in randomized order [35]. In

this study, using these two tasks allowed us to test the validity of our brain signature for the prediction

of discounting of several types of reward, and thus to investigate generalizability across reward

domains. Each of these tasks included 32 choices between SS and LL options. Participants were

instructed that one of their 32 choices could be randomly selected and the option that they had chosen

would be given to them. Thus, like in Study 1 and 2, participants' choices were non-hypothetical and

incentive-compatible. For each trial, participants could indicate their choice by pressing either a blue

key on the keyboard with their right-hand index to select the option on the left or a yellow key with their

right-hand middle finger to select the option on the right. Once the choice had been made, a message

on the screen indicated which option had been chosen. Trials were presented in randomized order.

**Other measures of impulsivity traits and symptoms**

*Study 1*

In Study 1, along with choice data collected from the ITC task, we used self-report data from the

Impulsive Behavior Short Scale–8 (I-8), which measures the psychological construct of trait impulsivity

according to the Urgency, lack of Premeditation, lack of Perseverance, and Sensation seeking (UPPS)

model with four subscales comprising two items each [37]. We predicted that the trait of *urgency—*

*defined as the tendency to act rashly in an emotional context* (e.g., "I sometimes do things to cheer

myself up that I later regret")— would be closest to brain-predicted delay discounting, as both urgency

and delay discounting are supposed to measure a tendency to prefer most immediate rewards at the

expense of potential long-term gains.

*Study 2*

In Study 2, we used data from the UPPS-P Impulsive Behavior Scale, which measures trait impulsivity

according to the UPPS model with five subscales: positive urgency, negative urgency, lack of

premeditation, lack of perseverance, and sensation seeking [38]. Paralleling Study 1, we predicted that

urgency would be the most closely related to brain-based predictions. We used the average of the

subscales of positive urgency (rash actions taken in response to positive emotional states) and

negative urgency (rash actions taken in response to negative emotional states) to test this hypothesis.

*Study 3*

This clinical study did not include a trait measure of impulsivity such as the UPPS scale. However,

clinical measures of core symptoms of bvFTD were available, in particular for two symptoms closely

related to impulsivity: inhibition deficit and dysexecutive syndrome (i.e., dysfunction in executive

functions). In another recent investigation of the same sample, we found that these two bvFTD

symptoms are related to higher discounting rates of both money and food [35]. We further used the

Hayling Sentence Completion Test (HSCT) [39] considered as an objective measure of inhibition deficit,

and the Frontal Assessment Battery (FAB) [40] as a measure of executive functions (lower scores

indicating worse executive functions). In the HSCT, participants are asked to complete 15 sentences

using the appropriate word, as fast as possible (automatic condition, part A), and 15 sentences using a

completely unrelated word (inhibition condition, part B). We used the Hayling error score (number of

errors in part B) as a measure of the difficulty to inhibit a prepotent response, as in Flanagan et al. [41].

**MRI data acquisition and preprocessing**

*Study 1*

Brain imaging data for Study 1 were acquired using a Siemens Trio 3T scanner. Structural images

were acquired using a T1 weighted MPRAGE sequence with the following parameters: TR 1660 ms;

TE 2.54 ms; FoV 256 mm; 208 slices; slice thickness 0.80 mm; TI 850 ms; flip angle 9°; voxel size 0.8

mm isomorphic; total acquisition time 6:32 min. T1 images were preprocessed for Voxel Based

Morphometry (VBM) analyses with SPM 12. We used the SPM module "Segment" for segmentation,

bias correction and rigid alignment of T1 images. These images were then used as input into the

DARTEL SPM module to create a customized DARTEL template and individual 'flow fields' for each

subject. DARTEL determines the nonlinear deformations for warping all grey and white matter images

so that they match each other. Finally, the SPM module "Normalise to MNI space" generated spatially

normalized grey matter images using the deformations estimated in the previous step and images

were spatially smoothed with a 6 mm Gaussian FWHM kernel. Among the obtained grey matter

images, three outliers (based on Mahalanobis distance of individual grey matter density maps with

Bonferroni correction) were detected and excluded from further analyses.

*Study 2*

Brain imaging data for Study 2 were acquired using a Siemens Trio 3T scanner (with a 32-channel

head coil). Structural images were acquired using a T1 weighted MPRAGE sequence with the

following parameters: TR 1630 ms; TE 3.11 ms; FOV 192x256; 160 slices; slice thickness 1 mm; TI

1100 ms; flip angle 15°; voxel size 0.9375 × 0.9375 × 1.000 mm; total acquisition time 4:35 min. We

used existing data preprocessed by Kable and colleagues [36]. T1 images were preprocessed for VBM

analyses using the default preprocessing pipeline of the Computational Anatomy Toolbox (CAT12) for

SPM12. T1-weighted images underwent spatial adaptive non-local means (SANLM) denoising filter,

were bias corrected, and affine-registered, followed by standard SPM unified tissue segmentation into

grey matter, white matter, and cerebral spinal fluid. The grey matter volume images were spatially

registered to a common template using Geodesic Shooting, resampled to 1.5 mm3, and spatially

smoothed with an 8 mm Gaussian FWHM kernel.

*Study 3*

Brain imaging data for Study 3 were acquired using a Siemens Prisma whole-body 3T scanner (with a

12-channel head coil). Structural images were acquired using a T1 weighted MPRAGE sequence with

the following parameters: TR 2400 ms; TE 2.17 ms; FOV 224 mm; 256 slices; slice thickness 0.70

mm; TI 1000 ms; flip angle 8°; voxel size 0.7 mm isomorphic; total acquisition time 7:38 min. T1

images were preprocessed for Voxel Based Morphometry (VBM) analyses using SPM 12, following

the same steps as in Study 1.

**Data analyses**

The analyses detailed in the following subsections aimed to: (1) develop and validate a structural brain

signature predicting delay discounting in a healthy population (Study 1); (2) test the validity of

predictions of this structural brain signature as measures of impulsivity in independent studies

involving different types of populations, including healthy (Study 2) and clinical samples (Study 3). All

analyses were performed using R Studio (1.2.1335) and Matlab (R2017b). The global analytic

approach is summarized in Figure 1A. The specific analyses conducted in each study to check the

validity of brain-based predictions are detailed in Figure 1B.

## A. General analytic approach

## B. Delay discounting paradigms and prediction validity assessment

**Figure 1. Methodological approach for the development and validation of a structural brain signature of impulsivity. A)** Grey matter density (GMD) maps from healthy participants of Study 1 were used for the prediction of delay discounting (log(k)) by LASSO-PCR with 10-fold cross-validation. In each fold, the classifier was trained on 90% of the data and tested on the remaining 10% hold-out data to evaluate its predictive accuracy. The predictive whole-brain pattern obtained from Study 1 was then tested in two independent samples, on the data of participants of Study 2 (healthy participants), and Study 3 (patients with neurodegenerative dementia and matched healthy control participants). The brain pattern was applied to the grey matter density maps of each study's participants to evaluate the validity of its predictions in different types of population. **B)** Several tests were performed in each of the four studies to assess the validity of the structural signature trained and cross-validated in

Study 1. In Study 1 (the training and cross-validation sample), permutation tests on different metrics
(MSE, RMSE, MAE) and in particular on the correlation between predicted and actual log(k) were
used to investigate the predictive accuracy of the developed brain pattern; the validity of predictions
was also assessed through testing their correlation with out-of-sample log(k) measured several weeks
later and with a self-report measure of the urgency component of impulsivity trait (subscale of
Impulsive Behavior Short Scale). Study 2 and 3 served as independent test samples to further validate
and generalize the structural signature developed in Study 1. In Study 2, we tested whether brain-
based predictions correlated with the actual log(k)'s computed in the sample and with self-reported
urgency trait (mean of positive and negative urgency subscales of UPPS- Impulsive Behavior Scale).
In Study 3, which involved patients with behavioral variant frontotemporal dementia (bvFTD) matched
with healthy controls, we tested: 1) correlations between the brain pattern predictions and observed
delay discounting for two types of stimuli (money and food) across patients and controls; 2) the ability
of brain-based predictions to distinguish patients from controls; 3) correlations between measures of
impulsivity symptoms (inhibition and executive deficits) and brain-based predictions among patients.

312
313　*Computation of discount rates*

314　In all three studies, the individual discounting rate (k) was estimated by fitting logistic regressions to

315　the individual choice data, with the assumption that the subjective value (SV) of the choice options

316　followed hyperbolic discounting, as follows:

317
$$SV = \frac{A}{1 + kD}$$

where A is the amount of the option, D is the delay until the receipt of the reward (for immediate

choice, $D = 0$), and k is a discounting rate parameter that varies across subjects. Higher values of k

indicate greater discounting and thus higher preference for sooner rewards. In Study 1, we used

logistic regressions (as described in Wileyto et al., 2004) to estimate the individual parameter k from

the participant's answers in the ICT task at baseline and we used the log(k) values as the parameter to

be predicted. Individual k's were log-transformed in all studies to obtain non-skewed distributions of

discounting parameters. In Study 2, we also used the log(k) values at baseline (see [36]). In Study 3, we

used the log(k) values calculated in bvFTD patients matched with controls for both monetary and food

rewards (see [35]).

*LASSO-PCR, training and cross-validation of the brain pattern predicting log(k) in Study 1*

We used a regression-based standard machine learning algorithm, LASSO-PCR (least absolute

shrinkage and selection operator-principal component regression) [34], to train a classifier to predict

log(k) from the individual whole brain grey matter density (GMD) maps. LASSO-PCR uses principal

components analysis (PCA) to reduce the dimensionality of the data and LASSO regression to predict

the outcome (log(k)) from the extracted component scores. The components identified by the PCA

correspond to groups of brain regions that covary with each other in terms of grey matter density. The

LASSO algorithm fits a regularized regression model predicting log(k) from the identified components.

This algorithm iteratively shrinks the regression weights towards zero, thus selecting a subset of

predictors and reducing the contribution of unstable components. LASSO-PCR is suited to make

predictions from thousands of voxels across the whole-brain, in particular because it solves the issue

of multicollinearity between voxels and brain regions (see [43,44]). Moreover, it is possible to reconstruct

voxel weights across the brain (from voxel loadings on PCA components and LASSO regression

coefficients of components), yielding predictive brain maps that are easier to interpret than component

weights. To assess the accuracy of this predictive modeling from GMD maps, we used a 10-fold cross-

validation process. The brain classifier was trained on 90% of the data and tested on the remaining

10% with 10 iterations, so that each participant was used for training the model in nine folds and for

testing the accuracy of its prediction in the remaining fold. Ten-fold cross-validation is within the range

of typically recommended folds (between 5 and 10) and allowed for a large training sample size at

each iteration [45,46]. Default regularization parameters were used for all machine-learning analyses to

avoid overfitting of the model to the data. We used four metrics to assess the accuracy of the model

predictions: the mean squared error (MSE) of prediction, the root mean squared error (RMSE), the

mean absolute error (MAE), and the correlation between the model predictions (from the 10 hold-out

test samples) and observed log(k)'s (prediction-outcome correlation).

*Test of the validity of predicted log(k) in Study 1*

To test the reliability of the predictions, we used permutation tests assessing the statistical significance

of the accuracy metrics (MSE, RMSE, MAE and prediction-outcome correlation). More precisely, 5000

iterations of randomly permuting the log(k) values were used to generate null distributions of these

four metrics and thus to assess the probability of: (MSE < actual MSE), (RMSE < actual RMSE),

(Mean abs. error < actual Mean abs. error) and of (prediction-outcome correlation < actual prediction-

outcome correlation) under the null hypothesis. To further confirm the validity of out-of-sample

predictions of log(k), we performed correlation tests between the predicted log(k) and: (1) calculated

log(k) values for the ITC task performed seven weeks later (at the end of the dietary intervention); (2)

the urgency trait subscale of the Impulsive Behavior Short Scale–8 (I-8). Since we had directional

hypotheses, we used one-tailed correlation tests for all correlations between predicted and observed

log(k).

*Predictions of the brain pattern in an independent sample of healthy participants in Study 2*

To assess the predictions of the brain classifier developed in Study 1 in participants of Study 2, we

calculated the dot product between the predictive weight map and the grey matter density map of each

participant of Study 2. The dot product (computed as a linear combination of the participant's voxel

grey matter density multiplied by voxel weight across the brain), plus the classifier's intercept, provides

a pattern response and thereby a predicted value of log(k) for each participant. This allowed us to test

the correlations between the predicted log(k) values and: (1) the actual log(k) values computed in the

sample; (2) the average of positive and negative urgency measures from the UPPS-P Impulsive

Behavior Scale.

*Predictions of the brain pattern in patients with neurodegenerative dementia in Study 3*

To assess the predictions of the brain classifier developed in Study 1 in participants of Study 3, we

calculated again the dot product as a measure of pattern response and thereby a predicted value of

log(k) for each participant of Study 3. This allowed us to test: (1) the correlation between the predicted

log(k) and the actual log(k) values (for both monetary and food rewards) across the whole sample

(bvFTD patients and matched controls); (2) whether predicted log(k) values could accurately

discriminate between bvFTD patients and controls, using a single-interval test (thresholded for optimal

overall accuracy). Further, we explored whether the predicted log(k)'s were related to the severity of

inhibition deficit (measured by Hayling error score) and of dysexecutive syndrome (i.e., lower FAB total

score) among bvFTD patients.

*Bootstrapping and thresholding of the predictive brain pattern obtained in Study 1*

We used a bootstrapping analysis to detect the brain regions that were the most robust contributors to

predict log(k). Sampling with replacement from the initial sample of Study 1 participants generated

5,000 samples. The LASSO-PCR algorithm yielded a predictive brain pattern (voxel weights across

the brain) from the data (paired GMD map – log(k) outcome) in each of these 5,000 samples. For each

voxel weight in the whole-brain pattern, the probability of being different from 0 (either above or below

0) could be estimated across the 5,000 samples. Thus, two-tailed, uncorrected p-values were

calculated for each voxel across the whole brain and false discovery rate (FDR) correction was used

to correct for multiple comparisons. Bootstrapped weights were thresholded at q=0.05 FDR-corrected

across the whole weight map, as well as at p=0.05 uncorrected for display.

*Spatial distribution of weights in the predictive brain pattern obtained in Study 1*

To further characterize the spatial distribution of regions predicting log(k) and their link to different

functional networks, we investigated the similarity between the predictive brain pattern (resulting from

the LASSO-PCR procedure) and term-based meta-analytic images [47] representing functional networks

that have been previously hypothesized [48] to contribute to temporal discounting, namely brain areas

related to valuation, executive control and memory/prospection. We calculated the spatial correlation

coefficients (Pearson's r) between the brain pattern (map of weights) and each of the meta-analytic

maps (thresholded meta-analytic uniformity maps from Neurosynth) corresponding to the following list

of terms: "value", "reward", "emotion", "affect", "executive", "conflict", "cognitive control", "attention",

"planning", "imagery", "memory", "episodic memory". These spatial correlations provide descriptive

insight into the importance of the contribution of GMD within specific functional networks to predict

individual differences in delay discounting [14,49].

**RESULTS**

**Development and cross-validation of a structural brain signature predicting delay discounting**

**in healthy adults (Study 1)**

*Individual differences in impulsivity*

On average, participants had a fitted log(k) parameter of -5.94 (median log(k)=-5.49, corresponding to

k=0.0041). Discounting rates were characterized by substantial individual differences (SD=2.00), with

log(k) ranging from -11.92 to -2.16. These individual differences were very stable over a 7-week period

as reported previously [14]. On the I-8 subscale of urgency trait, participants' average scores varied

between 1 and 5 (mean=2.72; median=2.5; SD=0.84). Log(k) showed a trend for a weak positive

correlation with the urgency trait (R=0.17, p=0.06, 95%-CI= [-0.009, 0.35]).

*Cross-validated predictions of delay discounting - Validity of predicted log(k) in healthy participants*

The 10-fold cross-validation procedure revealed a significant accuracy of the brain-based prediction

(see Figure 2A and 2B and Supplementary figure 1): the predictions had a mean squared error of 3.45

(permutation test: p=0.0026), a root mean squared error of 1.86 (permutation test: p=0.0026), a mean

absolute error for predicted log(k) of 1.46 (permutation test: p=0.0022), and a cross-validated

prediction-outcome correlation of R=0.35 (permutation test: p=0.0028) (Figure 2C).

Further, supporting the reliability and conceptual validity of the brain-predicted log(k)'s, we found that

brain-based predictions at baseline significantly correlated with (out-of-sample) log(k)'s computed from

the ITC task performed seven weeks later (R=0.34, p<0.001, 95%-CI= [0.18, 1]) (Figure 2D). This

suggests that a relatively stable part of the between-person variability in delay discounting was

explained by individual differences in brain structure. Moreover, higher brain-predicted log(k) values

were associated with higher self-reported urgency (R=0.20, p=0.037, 95%-CI= [0.01, 0.37]) (Figure

2E).

Like the actual measures of log(k) (see [14]), brain-based predictions of log(k) did not significantly

correlate with age (R=-0.11, p=0.24, 95%-CI= [-0.29, 0.07]), education (R=-0.15, p=0.10, 95%-CI= [-

0.33, 0.03]), income (R=-0.12, p=0.21, 95%-CI= [-0.30, 0.07]), BMI (R= -0.04, p=0.66, 95%-CI= [-0.22,

0.14]), and percentage of body fat (R= -0.13, p=0.18, 95%-CI= [-0.31, 0.06]) (see more details in

Supplementary figure 2).

**Performance of the Structural Impulsivity Signature in a second independent sample of healthy**

**participants (Study 2)**

Study 2 tests the predictions of the Structural Impulsivity Signature (SIS) in a second MRI dataset of

healthy participants, that has used a different protocol, scanner, different preprocessing pipeline, in a

socio-demographically different participant population.

*Individual differences in impulsivity*

The mean log(k) parameter in Study 2 was -4.09 (median log(k)=-3.94, corresponding to a k of 0.019).

Individual differences in the discounting parameter were less variable (SD=0.98) as compared to

Study 1, with log(k) ranging from -7.08 to -2.12. Participants had average urgency trait scores (means

of positive and negative urgency) varying between 1.00 and 3.01 (mean=1.76; median=1.68;

SD=0.48). In Study 2, log(k) had a trend for a negative correlation with urgency (R=-0.14, p=0.06,

95%-CI= [-0.29, 0.008]). Therefore, in Study 2, the discounting rate does not seem to be related to

individual differences in impulsivity.

*Brain-based predictions of impulsivity - Validity of predicted log(k) in a second independent sample of*

*healthy participants*

For each participant in Study 2, we calculated the predicted individual log(k) as the dot-product

between the weight map developed in Study 1 and the individual GMD map. We then tested whether

predicted log(k) correlated with observed individual log(k) and with the impulsivity trait of urgency

(UPPS subscales). While we did not find a significant link between predicted and observed log(k) in

Study 2 (R=0.06, p=0.21, 95%-CI= [-0.07, 1]), predicted log(k) was positively associated with urgency

scales (R=0.15, p=0.047, 95%-CI= [0.002, 0.30], see Figure 2F), as in Study 1. Thus, the results of

Study 2 partially validate the developed structural brain signature as a brain signature of impulsivity.

**Figure 2. Predictive validity of the structural brain pattern in Study 1 and Study 2. A)** Mean squared error (MSE) of prediction and significance obtained by permutation test (5,000 samples – N=113 males). **B)** Mean absolute error (MAE) of prediction and significance obtained by permutation test (5,000 samples – N=113 males). **C)** Correlation between predicted log(k) and actual log(k) in Study 1 and significance of prediction-outcome correlation obtained by permutation test (5,000 samples – N=113 males). **D)** Test of the parametric correlation between predicted log(k) and actual log(k) assessed 7 weeks later in Study 1 (R=0.34, p<0.001, 95%-CI= [0.15, 0.50]). **E)** Test of the parametric correlation between predicted log(k) and self-reported urgency (subscale of I-8 Impulsive Behavior Short Scale) in Study 1 (R=0.20, p=0.037, 95%-CI= [0.01, 0.37]). **F)** Test of the parametric correlation between predicted log(k) and self-reported urgency (mean of positive and negative urgency subscales of UPPS-P Impulsive Behavior Scale) in Study 2 (R=0.15, p=0.047, 95%-CI= [0.002, 0.30]).

**Validation of the structural brain signature in a clinical sample of bvFTD patients and matched controls (Study 3)**

Our last analysis step aimed at further testing the generalizability of the SIS by evaluating its validity in a patient population that is characterized by impulsivity. Study 3 employed a distinct protocol from Studies 1 and 2 (different ITC task, different MRI scanner and parameters), and in a different, older population including dementia patients with substantial structural atrophy. This further allowed us to investigate the clinical relevance of the SIS (1) for classifying patients with bvFTD differently from matched control participants and (2) for predicting the core symptoms of disinhibition and executive deficits in patients with bvFTD [8].

*Differences of impulsivity between bvFTD patients and healthy controls*

In line with the core symptoms of this disorder, bvFTD patients presented significantly higher delay discounting (i.e. more impatient or impulsive choices) compared to controls, for both money rewards and food rewards (see [35]). They also showed higher inhibition deficit (Hayling-error score; $t=5.71$, $p<0.001$, Cohen's $d=1.60$, 95%-CI=[0.87, 2.33]) and lower executive performances (FAB score; $t=-7.31$, $p<0.001$, Cohen's $d=-2.00$, 95%-CI=[-1.23, -2.77]) compared to controls (see Supplementary table 1).

*Brain-based predictions of impulsivity – Validity of predicted log(k) in bvFTD patients*

To investigate the predictive validity of our classifier in Study 3, we first tested whether predicted log(k)'s (obtained from the brain pattern applied to each participant's grey matter density map) were correlated with actual log(k)'s calculated in this study across the whole sample (patients and controls). This analysis showed that the predicted log(k) values were positively correlated with actual log(k) values, for both monetary rewards (R=0.30, p=0.03, 95%-CI= [0.03, 1], mean absolute error of 2.08) and for food rewards (R=0.40, p=0.006, 95%-CI= [0.15, 1], mean absolute error of 2.65) (see Figure 3.A and 3.B).

We next tested whether the SIS predictions could distinguish bvFTD patients from controls. As expected, we found that brain-predicted log(k) was significantly higher in bvFTD patients than in controls (t=3.60, p= 0.0009, Cohen's d=1.09, 95%-CI=[0.41, 1.76] – see Figure 3.C). Notably, brain-

predicted log(k) significantly predicted whether a grey matter density map was from a bvFTD patient or

from a control participant, with a classification accuracy of 81 % (p= 0.002, sensitivity = 87.5%,

specificity = 72.2%, - see Figure 3.D). Interestingly, the actual log(k)'s calculated for monetary and

food rewards in this sample revealed slightly lower predictive accuracies and especially lower

specificities: 73.7 % accuracy for monetary rewards (p= 0.07, sensitivity = 100%, specificity = 37.5%,)

and 76.3 % for food rewards (p= 0.01, sensitivity = 100%, specificity = 47.1%).

We next investigated the relationship between brain-predicted log(k) and clinical measures of bvFTD

core symptoms of disinhibition and executive deficits. Across both the patient and control groups,

higher predicted log(k) was associated with higher inhibition deficit (higher Hayling-error score;

R=0.55, p=0.0002, 95%-CI= [0.30, 0.74]) and higher executive troubles (lower FAB score; R=-0.56,

p=0.0001, 95%-CI= [-0.74, -0.30]). More interestingly, even within the group of bvFTD patients, higher

predicted log(k) was associated with higher inhibition deficit (higher Hayling-error score; R=0.52,

p=0.01, 95%-CI= [0.14, 0.77]) and higher executive troubles (lower FAB score; R=-0.43, p=0.04, 95%-

CI= [-0.71, -0.03]) (see Figure 3.E and 3.F). Further, we checked that predicted log(k) was still

significantly related to lack of inhibition (i.e., higher Hayling-error scores; B=8.63, p=0.02, 95%-CI=

[1.51, 15.7]) within bvFTD patients even after controlling for executive function deficit; this added result

showed that the relationship between brain-based predictions and disinhibition symptom was not only

due to shared variance with the severity of dysexecutive syndrome. Together, these findings show that

the SIS significantly and accurately classified bvFTD patients from matched controls, and that it

tracked the severity of key symptoms in these patients.

**Figure 3. Predictive validity of the structural brain pattern in Study 3. (A)** Parametric correlation between predicted log(k) and actual log(k) assessed with monetary rewards in Study 3 (R=0.30, p=0.07, 95%-CI= [-0.02, 0.57]). Patients are represented as squares in darker blue and controls as circles in lighter blue. (**B**) Parametric correlation between predicted log(k) and actual log(k) assessed with food rewards in Study 3 (R=0.45, p=0.01, 95%-CI= [0.1, 0.64]). Patients are represented as squares in darker blue and controls as circles in lighter blue. (**C**) As expected, predicted log(k) was higher in bvFTD patients (N=24) than in controls (N=18) (t=3.60, p=0.0009, Cohen's d=1.09, 95%-CI=[0.41, 1.76]). (**D**) ROC curve showing the performance of the brain-based prediction of log(k) in classification of bvFTD patients versus healthy controls (single interval test thresholded for optimal accuracy: accuracy=81 %, p= 0.002, AUC = 0.80, sensitivity = 87.5%, specificity = 72.2%). (**E**) Higher predicted log(k) was related to greater inhibition deficits (Hayling-error score) in bvFTD patients (R=0.52, p=0.01, 95%-CI= [0.14, 0.77]). (**F)** Higher predicted log(k) was related to more impaired executive functions (as measured with the FAB score) in bvFTD patients (R=-0.43, p=0.04, 95%-CI= [-0.71, -0.03]).

**Spatial distribution of weights in the structural brain signature (Study 1)**

*Thresholded pattern of structural brain signature*

Bootstrapping results revealed the positive and negative weights that most strongly contributed to GMD-based prediction of individual differences in delay discounting. At a threshold of q=0.05 FDR-corrected, we found two clusters in which grey matter density positively contributed to discounting differences (which means that higher grey matter density was associated with higher impatience); these clusters were in the left lateral parietal cortex (supramarginal gyrus) and left lateral occipital cortex (superior division). At a threshold of p=0.001 uncorrected, we found additional clusters contributing positive weights, especially in regions of the valuation system [50] such as the right orbitofrontal cortex (OFC), ventromedial prefrontal cortex (vmPFC) and right ventral striatum.

At q=0.05 FDR-corrected, there was one cluster in the posterior cingulate cortex (PCC) and adjacent lingual gyrus (including retrosplenial cortex) in which grey matter density contributed negatively to discounting differences (i.e., in which lower grey matter density was associated with higher impatience). At a threshold of p=0.001 uncorrected, other important regions contributing negative weights were found in the left hippocampus, the right anterior insulae (AI), dorsal anterior cingulate cortex (ACC), and amygdalae. For display purposes, the bootstrapped weight map is displayed in Figure 3A at a more comprehensive threshold (p=0.05 uncorrected, see also Supplementary table 2).

*Similarity of structural brain signature to meta-analytic maps*

When comparing the predictive map of log(k) with meta-analytic uniformity maps [47], we observed that the highest similarities (spatial correlation r's > 0.1 in absolute value) were with the "Emotions", "Affect", "Conflict" and "Imagery" meta-analytic maps (Figure 4B). These spatial correlations were all negative, indicating that greater grey matter density in areas related to emotions, affect, conflict processing, and imagery contributes to predicting lower delay discounting or more 'patient' decision-making (or conversely, lower grey matter density in these areas predicts higher discounting and more impulsive decision-making). The "Emotions", "Affect", "Conflict" and "Imagery" meta-analytic maps correspond to overlapping functional networks (see Figure 4.B). Among the most overlapping regions between these four networks (in red), the AI and dorsal ACC, corresponding to robust negative weights in the brain pattern, are known to be major hubs of the salience network [51].

**Study 1: Spatial distribution of weights in the structural brain pattern predicting delay discounting**

**A. Thresholded weights of the brain pattern after bootstrapping**

*amygdalae*   *Right OFC / vmPFC*   *Right VS*   *Right AI*

*Right lingual gyrus*   *Left hippocampus*   *Dorsal ACC*   *Left lateral occipital cortex*

*Left lateral parietal cortex*   Z-values   -0.005 ▮ 0.005

**B. Spatial similarity of the brain pattern with meta-analytic maps**

Overlap between "Emotions", "Affect", "Conflict" and "Imagery" meta-analytic maps

Value / Reward / Emotions / Affect / Executive / Conflict / Cognitive control / Attention / Planning / Imagery / Memory / Episodic memory

-0.15  -0.10  -0.05  0.00  0.05  0.10  0.15
Spatial Correlation (Pearson's r)

Intensity of overlap: 1  2  3  4

580
581
**Figure 4. Spatial organization of the structural brain pattern developed in Study 1. A)** Whole-
brain weight map thresholded at p=0.05 (uncorrected for multiple comparisons across the brain)
resulting from a bootstrapping procedure (5,000 samples); negative weights (contributing to lower
discounting with higher grey matter density) are shown in blue. Positive weights (contributing to higher
discounting with higher grey matter density) are shown in orange. The three framed clusters
correspond to the three clusters in which peaks are significant at q=0.05 FDR-corrected. Regions
indicated in italics are some of the main regions significant at p=0.001, uncorrected (OFC: orbitofrontal
cortex; vmPFC: ventromedial prefrontal cortex; VS: ventral striatum; AI: anterior insula; ACC: anterior
cingulate cortex). **B)** On the left, spatial correlations of the unthresholded delay discounting brain
pattern with thresholded meta-analytic uniformity maps from Neurosynth (http://www.neurosynth.org).
As in [14], we selected meta-analytic maps corresponding to three types of functions assumed to be
involved in delay discounting: 1/ valuation and emotion processing; 2/ executive control; 3/ memory
and prospection. Spatial correlations are descriptive and indicate the extent of spatial similarities
between the structural brain pattern and the functional networks of interest [47]. Highest correlations (or

| similarities) were observed with the "Emotions", "Affect", "Conflict", and "Imagery" meta-analytic maps,
| and were all negative, meaning that higher grey matter density in these functional regions is
| associated with lower discounting. On the right, we show the spatial distribution and overlap between
| the four meta-analytic maps found to be the most negatively correlated with the structural brain pattern
| (from 1, corresponding to non-overlapping regions from only one map, to 4, corresponding to regions
| of overlap between the 4 maps).

**604 Spatial distribution of brain regions contributing to higher predicted log(k) in bvFTD (Study 3)**

To identify the main brain regions which contributed to differentiate bvFTD patients from controls on

the brain-predicted log(k), we contrasted bvFTD patients versus controls in terms of voxel-wise pattern

expression of the predictive map of log(k). To this end, for each bvFTD patient and each control

participant, we computed an 'importance map' as the unsummed matrix dot product between the

predictive structural weight map and the individual grey matter density map. Since higher resulting dot

product contributes to higher predicted discounting, the importance map shows which brain regions

contributed to increase (or decrease) predicted discounting in each individual. We performed a t-test

contrasting bvFTD patients and controls (bvFTD > controls) on the resulting importance maps, with a

family-wise error (FWE) correction applied to p-values to correct for multiple comparisons across the

brain (see Figure 5.C). This contrast shows the regions in which structural atrophy contributed

positively to higher predicted discounting in bvFTD than in controls (regions in red). These included

the OFC, anterior insulae, dorsal ACC, striatum, thalamus, amygdalae, hippocampus, and middle

temporal regions. These regions corresponded to areas combining the presence of negative weights

in the predictive brain pattern (i.e., voxels for which higher GMD predicts lower discounting and more

patient decision-making, shown in Figure 5.B) and the presence of significant grey matter atrophy due

to bvFTD pathology (see atrophy pattern in Figure 5.A). Thus, the contrast shown in Figure 5.C also

maps the regions in which the SIS is the most similar to bvFTD atrophy pattern.

**Figure 5. Spatial distribution of regions contributing to higher predicted discounting in bvFTD in Study 3.** We computed an importance map as the unsummed matrix dot product between the Structural Impulsivity Signature (SIS) (developed in Study 1) and the individual grey matter density map of each Study 3 participant. Since higher resulting dot product contributes to higher predicted

discounting, the importance map shows how brain regions contribute to increased (or decreased)
predicted discounting in each individual. We performed a t-test contrasting bvFTD patients and
controls (bvFTD > controls) on the resulting importance maps, to show in particular the regions in
which the contribution to higher discounting was significantly higher in bvFTD than in controls. Within
regions showing atrophy in bvFTD (see 6.A), those corresponding to negative (/positive) weights in the
whole-brain predictive pattern (see 6.B) contributed to increase (/decrease) discounting in bvFTD (see
6.C). (**A**) VBM–derived grey matter atrophy map of bvFTD patients contrasted with matched controls
(bvFTD<Controls), FWE-corrected and thresholded at p < 0.05. (**B**) Unthresholded whole-brain weight
map of the structural brain pattern developed in Study 1 and used in Study 2 to predict delay
discounting in bvFTD patients (N=24) and matched controls (N=18). Negative weights (contributing to
lower discounting with higher grey matter density) are in blue and positive weights (contributing to
higher discounting with higher grey matter density) are in orange. (**C**) Contrast between bvFTD
patients and controls (bvFTD>Controls)) on the importance map, FWE-corrected and thresholded at p
< 0.05; this map shows regions contributing to increase discounting in bvFTD patients (compared to
controls) in red and regions contributing to decrease discounting in bvFTD patients (compared to
controls) in blue, the balance being in favor of a global increase in predicted discounting in bvFTD
patients.

**DISCUSSION**

Impulsive and maladaptive decision-making is a transversal feature of many mental disorders, especially prominent in behavioral-variant frontotemporal dementia (bvFTD). Yet, its relationship with individual brain characteristics, in particular brain structure, is still debated. Here, we used a machine learning technique to develop a brain signature (i.e., a multi-variate brain model) of individual differences in delay discounting—a facet of impulsivity—based on whole-brain grey matter density patterns. We performed out-of-sample cross-validation in a first sample of 117 healthy adults (Study 1) used for brain signature development. We further tested the generalizability of this brain signature developed in Study 1 in two independent studies: a second sample of 166 healthy adults (Study 2) and a clinical study including 24 bvFTD patients and 18 matched controls (Study 3). Individual differences of whole-brain grey matter density reliably predicted individual differences in discounting rates in the first sample of healthy adults but not in the second independent sample. However, the brain signature predicted individual differences of urgency (a subcomponent of impulsivity according to the UPPS model) with small-to-moderate effect sizes in both the first and the second samples of healthy adults. Most importantly, in the clinical study, we found that this structural signature of impulsivity (SIS) separated bvFTD patients from controls with 81% accuracy and that it significantly predicted not only individual differences in delay discounting across participants but also inhibition deficit (objectively assessed from the Hayling test), even within the group of bvFTD patients. Thus, the SIS might be more closely and reliably related to the broader concepts of impulsivity, urgency, and inhibition deficits rather than to specifically delay discounting, which may be more driven by cultural and educational factors than trait urgency. In sum, our results suggest that: 1) it is possible to predict individual differences in impulsivity from whole-brain structure and 2) this novel brain signature is sensitive to the structural atrophy that is characteristic of bvFTD, making it a novel candidate neuromarker for improving bvFTD diagnosis.

The identification of the SIS advances our knowledge of the neurobiology underlying individual differences in impulsivity. Higher discounting (i.e., greater impulsivity) was associated with higher grey matter density in clusters of the lateral parietal and occipital cortex as well as in regions of the OFC, vmPFC, ventral striatum, lateral PFC, precentral gyrus, and precuneus. Functional activation of these regions during intertemporal choices and in response to rewards has previously been shown to predict

higher discounting [14,52]. The SIS obtained from Study 1 also revealed regions in which greater grey

matter density contributes to lower individual impulsivity. Among the strongest negative contributors,

we found clusters corresponding to hub regions of the salience network (anterior insulae, dorsal ACC,

amygdalae). Dorsal ACC and anterior insula were also consistently found as significant regions

predicting delay discounting from whole-brain functional MRI [14]. These regions are associated with the

processing of emotionally significant internal and external stimuli [51,53,54] and awareness of present and

future affective states [55]; they are also supposed to be involved in switching between large-scale

networks to facilitate access to attention and working memory resources in the presence of a salient

event [56]. These areas are also known to be involved in cognitive conflict processing [57,58] and previous

studies have shown their response to difficult choices (characterized by choice conflicts between

options) during delay discounting [59]. Thus, our results suggest that more impulsive individuals might

be those for whom lower affective, attentional, and conflict processing would lead to more impulsive

decision-making, favouring immediately rewarding options over long-term consequences of behavior.

The SIS has the potential to contribute to the early diagnosis of conditions characterized by

high impulsivity, such as bvFTD. Brain signatures can in particular help the diagnosis of conditions

involving brain lesions that are sometimes difficult to detect by mere visual inspection of MRI scans,

especially at early stages of the disease. In addition, brain signatures can constitute neuroimaging

markers with diagnostic value that can be used across different samples and populations [11,60]. The SIS

may contribute to the diagnosis of bvFTD by complementing other brain models able to detect bvFTD.

A few previous studies successfully trained structural MRI classifiers for the specific purpose of

distinguishing FTD patients from controls (e.g., [61–63]). These bvFTD classifiers have shown their

accuracy to detect patients with clear structural brain damage but their ability to distinguish individuals

at risk of developing FTD due to genetic mutations is likely to be limited to the period just before

symptom onset [64,65]. Under the hypothesis of a continuum of marked impulsivity in presymptomatic

individuals and patients [16], the SIS might serve the early prediction and monitoring of bvFTD before

symptom onset. Impulsive behaviors may be present in an attenuated form long before clinical

diagnosis and hard to detect with traditional clinical methods. A neuromarker predicting impulsivity

may be sensitive to specific brain modifications that appear very early in individuals predisposed to

FTD (possibly as neurodevelopmental lesions [66]) and would thus allow to enhance the monitoring of

clinical signs of these subtle behavioral changes. Future tests of this brain signature in

presymptomatic populations will allow to evaluate these potential clinical applications.

As it predicts nearly 30% of the variance of inhibition deficit among bvFTD patients, the SIS

may be sensitive to lesions in a structural network underlying the core bvFTD symptom of disinhibition.

In addition to its potential contribution to the early detection of presymptomatic individuals, this brain

signature may thus aid differential diagnosis and provide insight into the neuropsychological profiles of

patients. The SIS may for instance help to distinguish bvFTD from other neurodegenerative or

neuropsychiatric conditions with different core symptoms. The differential diagnosis of Alzheimer's

disease and bvFTD can in particular be challenging. Using neuromarkers such as the SIS in cases of

diagnostic uncertainty potentially impacting the choice of treatment could therefore be highly valuable

[67] and should be an avenue for future studies. Moreover, the SIS could become a useful tool to

disentangle the phenotypic heterogeneity within bvFTD population [68]. The characterization of different

clinical and behavioral profiles within the bvFTD spectrum could help to better understand the

pathology, and to better adapt treatments according to patients' specific needs.

Despite holding promises for future clinical applications, we note that our results also point at

challenges in generalizing the brain signature to other independent samples of healthy adults. We

were successful at predicting delay discounting from whole-brain grey matter in a first rather

homogenous sample of healthy adults (male participants, controlled experimental conditions) showing

significant variability in terms of impulsivity and a positive correlation between the discounting rate and

urgency. In a second independent sample of healthy adults with lower variance of impulsivity and a

slightly negative correlation between the discounting rate and urgency, we could not replicate the

association with measured discounting rates but found evidence of the conceptual validity (i.e., a link

with the urgency trait) of brain-based predictions. This suggests that the variance captured by the SIS

developed in the first sample is more reliably related to individual differences in urgency than to

individual differences in discounting. The fact that urgency was slightly negatively correlated with the

discounting rate in the second healthy sample questions the idea that delay discounting necessarily

captures individual differences in impulsivity. These two constructs overlap but are not equivalent and

previous studies have already reported an absence of link between delay discounting and some

psychometric measures of impulsivity (e.g., [69,70]). Discounting rate is also a state-dependent variable

[71] and depends on situational factors such as cultural and social context [72]. In addition, the links

between personality and discounting rates may depend on participants' cognitive abilities [73]. Therefore, association between delay discounting and other measures of trait impulsivity may vary according to samples and studies. A promising approach for future studies would therefore be to predict latent variables that underlie different observed variables related to the same concept of impulsivity (instead of only one observed variable such as the discounting rate), which might achieve better performance in terms of replicability and generalizability [9]. Although multivariate brain signatures can be replicable with moderate sample sizes [74], future studies aiming to develop brain signatures of impulsivity could also benefit from using larger and more diverse samples [75]. More generally, we note that our results suggest a relatively small contribution of interindividual variability in brain structure to interindividual variability in impulsivity among healthy adults. Effect sizes of associations between predicted and observed impulsivity are however in line with those reported for most brain signatures of behavioral individual differences using structural features [9]. Moreover, like variability in brain structure, variability in genotype accounts for a rather small part of the variance of impulsivity [76]. The magnitude of associations between brain structure and behaviors may be limited in the general population but these associations might be more salient within populations with a marked variability of both brain and behavior such as patients with neurodegenerative conditions.

In conclusion, our results advance our knowledge of the association between impulsivity and brain structure in healthy adults and in patients with bvFTD. They also point at inherent challenges in developing replicable and generalizable brain signatures of individual differences based on brain structure. By identifying a structural network associated with individual differences in discounting rates, our results provide insight into the potential neurobiological bases of trait impulsivity (and in particular its urgency component). The good performance of the SIS among patients with bvFTD suggests a possible continuum of brain-impulsivity relationship across healthy and clinical conditions. Most noteworthy, the SIS separates bvFTD patients from controls with high accuracy, pointing at the potential clinical value for the diagnosis of bvFTD, in particular for the purpose of stratifying this heterogenous condition. MRI can be instrumental to confirm an FTD diagnosis [67] and the SIS only requires a preprocessed T1-weighted scan to reach a prediction. It holds promise as a phenotypic marker in patients with neurodegenerative or psychiatric conditions associated with high impulsivity. Future studies could test its clinical potential and whether this brain signature could be used in a real-life patient workflow.

**REFERENCES**

1. Chamberlain, S. R. & Sahakian, B. J. The neuropsychiatry of impulsivity. *Curr. Opin. Psychiatry* **20**, 255–261 (2007).

2. Chamberlain, S. R., Stochl, J., Redden, S. A. & Grant, J. E. Latent traits of impulsivity and compulsivity: toward dimensional psychiatry. *Psychol. Med.* **48**, 810–821 (2018).

3. Goodwin, B. C., Browne, M., Hing, N. & Russell, A. M. Applying a revised two-factor model of impulsivity to predict health behaviour and well-being. *Personal. Individ. Differ.* **111**, 250–255 (2017).

4. Rogers, M. M., Kelley, K. & McKinney, C. Trait impulsivity and health risk behaviors: A latent profile analysis. *Personal. Individ. Differ.* **171**, 110511 (2021).

5. Bjork, J. M., Momenan, R. & Hommer, D. W. Delay discounting correlates with proportional lateral frontal cortex volumes. *Biol. Psychiatry* **65**, 710–713 (2009).

6. MacNiven, K. H., Leong, J. K. & Knutson, B. Medial forebrain bundle structure is linked to human impulsivity. *Sci. Adv.* **6**, eaba4788 (2020).

7. Pehlivanova, M. *et al.* Diminished cortical thickness is associated with impulsive choice in adolescence. *J. Neurosci.* **38**, 2471–2481 (2018).

8. Rascovsky, K. *et al.* Sensitivity of revised diagnostic criteria for the behavioural variant of frontotemporal dementia. *Brain* **134**, 2456–2477 (2011).

9. Genon, S., Eickhoff, S. B. & Kharabian, S. Linking interindividual variability in brain structure to behaviour. *Nat. Rev. Neurosci.* **23**, 307–318 (2022).

10. Kragel, P. A., Koban, L., Barrett, L. F. & Wager, T. D. Representation, Pattern Information, and Brain Signatures: From Neurons to Neuroimaging. *Neuron* **99**, 257–273 (2018).

11. Woo, C.-W., Chang, L. J., Lindquist, M. A. & Wager, T. D. Building better biomarkers: brain models in translational neuroimaging. *Nat. Neurosci.* **20**, 365–377 (2017).

12. Anokhin, A. P., Grant, J. D., Mulligan, R. C. & Heath, A. C. The Genetics of Impulsivity: Evidence for the Heritability of Delay Discounting. *Biol. Psychiatry* **77**, 887–894 (2015).

13. Kirby, K. N. One-year temporal stability of delay-discount rates. *Psychon. Bull. Rev.* **16**, 457–462 (2009).

14. Koban, L. *et al.* An fMRI-based brain marker of individual differences in delay discounting. *J.*

*Neurosci.* (2023).

15. Amlung, M. *et al.* Delay Discounting as a Transdiagnostic Process in Psychiatric Disorders: A

Meta-analysis. *JAMA Psychiatry* **76**, 1176 (2019).

16. Godefroy, V. *et al.* Altered delay discounting in neurodegeneration: insight into the underlying

mechanisms and perspectives for clinical applications. *Neurosci. Biobehav. Rev.* 105048 (2023).

17. Ballard, K. & Knutson, B. Dissociable neural representations of future reward magnitude and

delay during temporal discounting. *Neuroimage* **45**, 143–150 (2009).

18. Bergström, F., Lerman, C. & Kable, J. W. Reduced cortical complexity in ventromedial prefrontal

cortex is associated with a greater preference for risky and immediate rewards. *Imaging Neurosci.*

**2**, 1–14 (2024).

19. Cooper, N., Kable, J. W., Kim, B. K. & Zauberman, G. Brain activity in valuation regions while

thinking about the future predicts individual discount rates. *J. Neurosci.* **33**, 13150–13156 (2013).

20. Hare, T. A., Hakimi, S. & Rangel, A. Activity in dlPFC and its effective connectivity to vmPFC are

associated with temporal discounting. *Front. Neurosci.* **8**, 50 (2014).

21. Lebreton, M. *et al.* A critical role for the hippocampus in the valuation of imagined outcomes.

*PLoS Biol.* **11**, e1001684 (2013).

22. Li, N. *et al.* Resting-State Functional Connectivity Predicts Impulsivity in Economic Decision-

Making. *J. Neurosci.* **33**, 4886–4895 (2013).

23. van den Bos, W., Rodriguez, C. A., Schweitzer, J. B. & McClure, S. M. Connectivity strength of

dissociable striatal tracts predict individual differences in temporal discounting. *J. Neurosci.* **34**,

10298–10310 (2014).

24. Bartra, O., McGuire, J. T. & Kable, J. W. The valuation system: a coordinate-based meta-analysis

of BOLD fMRI experiments examining neural correlates of subjective value. *Neuroimage* **76**, 412–

427 (2013).

25. Levy, D. J. & Glimcher, P. W. The root of all value: a neural common currency for choice. *Curr.*

*Opin. Neurobiol.* **22**, 1027–1038 (2012).

26. Rangel, A. & Clithero, J. A. Value normalization in decision making: theory and evidence. *Curr.*

*Opin. Neurobiol.* **22**, 970–981 (2012).

27. Chare, L. *et al.* New criteria for frontotemporal dementia syndromes: clinical and pathological

diagnostic implications. *J. Neurol. Neurosurg. Psychiatry* **85**, 865–870 (2014).

28. Karageorgiou, E. & Miller, B. Frontotemporal Lobar Degeneration: A Clinical Approach. *Semin.*

*Neurol.* **34**, 189–201 (2014).

29. Beagle, A. J. *et al.* Amount and delay insensitivity during intertemporal choice in three

neurodegenerative diseases reflects dorsomedial prefrontal atrophy. *Cortex J. Devoted Study*

*Nerv. Syst. Behav.* **124**, 54–65 (2020).

30. Bertoux, M., O'Callaghan, C., Flanagan, E., Hodges, J. R. & Hornberger, M. Fronto-striatal

atrophy in behavioral variant frontotemporal dementia and Alzheimer's disease. *Front. Neurol.* **6**,

147 (2015).

31. Chiong, W. *et al.* Neuroeconomic dissociation of semantic dementia and behavioural variant

frontotemporal dementia. *Brain J. Neurol.* **139**, 578–587 (2016).

32. Manuel, A. L. *et al.* Interactions between decision-making and emotion in behavioral-variant

frontotemporal dementia and Alzheimer's disease. *Soc. Cogn. Affect. Neurosci.* **15**, 681–694

(2020).

33. Han, D., Qu, L., Sun, L. & Sun, Y. Variable selection for a mark-specific additive hazards model

using the adaptive LASSO. *Stat. Methods Med. Res.* **30**, 2017–2031 (2021).

34. Tibshirani, R. Regression shrinkage and selection via the lasso. *J. R. Stat. Soc. Ser. B Methodol.*

**58**, 267–288 (1996).

35. Godefroy, V. *et al.* High delay discounting relates to core symptoms and to pulvinar atrophy in

frontotemporal dementia. Preprint at https://doi.org/10.1101/2024.05.16.594526 (2024).

36. Kable, J. W. *et al.* No Effect of Commercial Cognitive Training on Brain Activity, Choice Behavior,

or Cognitive Performance. *J. Neurosci.* **37**, 7390–7402 (2017).

37. Groskurth, K., Nießen, D., Rammstedt, B. & Lechner, C. M. The impulsive behavior short scale–8

(I-8): A comprehensive validation of the English-language adaptation. *Plos One* **17**, e0273801

(2022).

38. Whiteside, S. P. & Lynam, D. R. The five factor model and impulsivity: Using a structural model of

personality to understand impulsivity. *Personal. Individ. Differ.* **30**, 669–689 (2001).

39. Burgess, P. W. & Shallice, T. The hayling and brixton tests. (1997).

40. Dubois, B., Slachevsky, A., Litvan, I. & Pillon, B. The FAB: a frontal assessment battery at

bedside. *Neurology* **55**, 1621–1626 (2000).

41. Flanagan, E. C. *et al.* False recognition in behavioral variant frontotemporal dementia and

Alzheimer's disease—disinhibition or amnesia? *Front. Aging Neurosci.* **8**, 177 (2016).

42. Wileyto, E. P., Audrain-McGovern, J., Epstein, L. H. & Lerman, C. Using logistic regression to

estimate delay-discounting functions. *Behav. Res. Methods Instrum. Comput.* **36**, 41–51 (2004).

43. Wager, T. D., Atlas, L. Y., Leotti, L. A. & Rilling, J. K. Predicting Individual Differences in Placebo

Analgesia: Contributions of Brain Activity during Anticipation and Pain Experience. *J. Neurosci.*

**31**, 439–452 (2011).

44. Wager, T. D. *et al.* An fMRI-Based Neurologic Signature of Physical Pain. *N. Engl. J. Med.* **368**,

1388–1397 (2013).

45. Poldrack, R. A., Huckins, G. & Varoquaux, G. Establishment of best practices for evidence for

prediction: a review. *JAMA Psychiatry* **77**, 534–540 (2020).

46. Scheinost, D. *et al.* Ten simple rules for predictive modeling of individual differences in

neuroimaging. *NeuroImage* **193**, 35–45 (2019).

47. Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C. & Wager, T. D. Large-scale

automated synthesis of human functional neuroimaging data. *Nat. Methods* **8**, 665–670 (2011).

48. Peters, J. & Büchel, C. The neural mechanisms of inter-temporal decision-making: understanding

variability. *Trends Cogn. Sci.* **15**, 227–239 (2011).

49. Koban, L., Jepma, M., López-Solà, M. & Wager, T. D. Different brain networks mediate the effects

of social and conditioned expectations on pain. *Nat. Commun.* **10**, 1–13 (2019).

50. Lempert, K. M., Steinglass, J. E., Pinto, A., Kable, J. W. & Simpson, H. B. Can delay discounting

deliver on the promise of RDoC? *Psychol. Med.* **49**, 190–199 (2019).

51. Menon, V. Salience Network. in *Brain Mapping* 597–611 (Elsevier, 2015). doi:10.1016/B978-0-12-

397025-1.00052-X.

52. Hariri, A. R. *et al.* Preference for immediate over delayed rewards is associated with magnitude of

ventral striatal activity. *J. Neurosci.* **26**, 13213–13217 (2006).

53. Seeley, W. W. The salience network: a neural system for perceiving and responding to

homeostatic demands. *J. Neurosci.* **39**, 9878–9882 (2019).

54. Seeley, W. W., Zhou, J. & Kim, E.-J. Frontotemporal dementia: what can the behavioral variant

teach us about human brain organization? *The Neuroscientist* **18**, 373–385 (2012).

55. Craig, A. D., How do you feel — now? The anterior insula and human awareness. *Nat. Rev.*

*Neurosci.* **10**, 59–70 (2009).

56.   Menon, V. & Uddin, L. Q. Saliency, switching, attention and control: a network model of insula

function. *Brain Struct. Funct.* **214**, 655–667 (2010).

57.   Botvinick, M. M., Braver, T. S., Barch, D. M., Carter, C. S. & Cohen, J. D. Conflict monitoring and

cognitive control. *Psychol. Rev.* **108**, 624 (2001).

58.   Van Veen, V., Cohen, J. D., Botvinick, M. M., Stenger, V. A. & Carter, C. S. Anterior cingulate

cortex, conflict monitoring, and levels of processing. *Neuroimage* **14**, 1302–1308 (2001).

59.   Hoffman, W. F. *et al.* Cortical activation during delay discounting in abstinent methamphetamine

dependent individuals. *Psychopharmacology (Berl.)* **201**, 183–193 (2008).

60.   Haller, S., Lovblad, K.-O., Giannakopoulos, P. & Van De Ville, D. Multivariate Pattern Recognition

for Diagnosis and Prognosis in Clinical Neuroimaging: State of the Art, Current Challenges and

Future Trends. *Brain Topogr.* **27**, 329–337 (2014).

61.   Davatzikos, C., Resnick, S. M., Wu, X., Parmpi, P. & Clark, C. M. Individual patient diagnosis of

AD and FTD via high-dimensional pattern classification of MRI. *NeuroImage* **41**, 1220–1227

(2008).

62.   Gonzalez-Gomez, R., Ibañez, A. & Moguilner, S. Multiclass characterization of frontotemporal

dementia variants via multimodal brain network computational inference. *Netw. Neurosci.* 1–29

(2022).

63.   Moguilner, S. *et al.* Visual deep learning of unprocessed neuroimaging characterises dementia

subtypes and generalises across non-stereotypic samples. *EBioMedicine* **90**, (2023).

64.   Feis, R. A. *et al.* Single-subject classification of presymptomatic frontotemporal dementia mutation

carriers using multimodal MRI. *NeuroImage Clin.* **20**, 188–196 (2018).

65.   Feis, R. A. *et al.* A multimodal MRI-based classification signature emerges just prior to symptom

onset in frontotemporal dementia mutation carriers. *J. Neurol. Neurosurg. Psychiatry* **90**, 1207–

1214 (2019).

66.   Lee, S. E. *et al.* Network degeneration and dysfunction in presymptomatic C9ORF72 expansion

carriers. *NeuroImage Clin.* **14**, 286–297 (2017).

67.   Sheikh-Bahaei, N., Sajjadi, S. A. & Pierce, A. L. Current Role for Biomarkers in Clinical Diagnosis

of Alzheimer Disease and Frontotemporal Dementia. *Curr. Treat. Options Neurol.* **19**, 46 (2017).

68.   O'Connor, C. M. *et al.* Behavioral-variant frontotemporal dementia: Distinct phenotypes with

unique functional profiles. *Neurology* **89**, 570–577 (2017).

69. Lane, S. D., Cherek, D. R., Rhoades, H. M., Pietras, C. J. & Tcheremissine, O. V. Relationships

Among Laboratory and Psychometric Measures of Impulsivity: Implications in Substance Abuse

and Dependence. *Addict. Disord. Their Treat.* **2**, 33 (2003).

70. Reynolds, B., Ortengren, A., Richards, J. B. & de Wit, H. Dimensions of impulsive behavior:

Personality and behavioral measures. *Personal. Individ. Differ.* **40**, 305–315 (2006).

71. Odum, A. L. & Baumann, A. A. L. Delay discounting: State and trait variable. in *Impulsivity: The*

*behavioral and neurological science of discounting* 39–65 (American Psychological Association,

Washington, DC, US, 2010). doi:10.1037/12069-002.

72. Schwenke, D., Wehner, P. & Scherbaum, S. Effects of individual and dyadic decision-making and

normative reference on delay discounting decisions. *Cogn. Res. Princ. Implic.* **7**, 71 (2022).

73. Hirsh, J. B., Morisano, D. & Peterson, J. B. Delay discounting: Interactions between personality

and cognitive ability. *J. Res. Personal.* **42**, 1646–1650 (2008).

74. Spisak, T., Bingel, U. & Wager, T. Replicable multivariate BWAS with moderate sample sizes.

*bioRxiv* 2022–06 (2022).

75. Marek, S. *et al.* Reproducible brain-wide association studies require thousands of individuals.

*Nature* **603**, 654–660 (2022).

76. Sanchez-Roige, S. *et al.* Genome-wide association study of delay discounting in 23,217 adult

research participants of European ancestry. *Nat. Neurosci.* **21**, 16–18 (2018).

# A. General analytic approach

Development of a brain signature
predicting log(k) (Study 1)

Resulting whole-brain pattern
after voxel weight reconstruction

Predictors:
GMD maps

LASSO-PCR
with 10-fold cross-
validation

Training on:
90% of participants

Test on hold-out:
10% of participants

X10

Component
reduction

LASSO
regression

Outcome:
log(k)

Prediction validity
assessment
(permutation tests)

Application to new subjects in
independent samples (Studies 2 & 3)

GMD maps

Dot
product

Brain-predicted log(k)

Prediction validity
assessment

# B. Delay discounting paradigms and prediction validity assessment

## STUDY 1

N = 117 healthy adults (men)

Please choose:

10 €
today

22 €
in 2 weeks

Money delay discounting task

Actual log(k)
Urgency trait
(I-8)

Predicts ?

Brain-
predicted
log(k)

## STUDY 2

N = 166 healthy adults

Accept or reject vs baseline offer
(20 $ today)

42 $ in 35 days?

Money delay discounting task

Actual log(k)
Urgency trait
(UPPS)

Predicts ?

Brain-
predicted
log(k)

## STUDY 3

N = 24 bvFTD patients & 18 matched controls

Would you rather have:

11 €
today

44 €
in 14 days

Money delay discounting task

Would you rather have:

11 chocolates
today

44 chocolates
in 14 days

Food delay discounting task

Actual log(k) money - Actual log(k) food
Patient vs Control
Patient's inhibition deficit & executive deficit

Predicts ?

Brain-
predicted
log(k)

# Validity of brain-based predictions of impulsivity in healthy participants

## Study 1 – Prediction of delay discounting

**A.** *MSE* = 3.45, *p* = 0.0026

**B.** *MAE* = 1.46, *p* = 0.0022

**C.** *R* = 0.35, *p* = 0.0028

**D.** *R* = 0.34, *p* = 0.0003

## Study 1 – Prediction of urgency

**E.** *R* = 0.2, *p* = 0.037

## Study 2 – Prediction of urgency

**F.** *R* = 0.15, *p* = 0.047

**Validity of brain-based predictions of impulsivity in patients with bvFTD matched with controls (Study 3)**

# Study 1: Spatial distribution of weights in the structural brain pattern predicting delay discounting

## A. Thresholded weights of the brain pattern after bootstrapping

amygdalae

Right OFC / vmPFC

Right VS

Right AI

Right lingual gyrus

Left hippocampus

Dorsal ACC

Left lateral occipital cortex

Left lateral parietal cortex

Z-values

-0.005 ———— 0.005

## B. Spatial similarity of the brain pattern with meta-analytic maps

Value
Reward
Emotions
Affect
Executive
Conflict
Cognitive control
Attention
Planning
Imagery
Memory
Episodic memory

-0.15  -0.10  -0.05  0.00  0.05  0.10  0.15

Spatial Correlation (Pearson's r)

Overlap between "Emotions", "Affect", "Conflict" and "Imagery" meta-analytic maps

1   2   3   4

Intensity of overlap

# Study 3: Spatial distribution of brain regions contributing to higher predicted delay discounting in bvFTD patients

*Lower grey matter density in bvFTD*

*Negative weight*     *Positive weight*

*Contribution to higher discounting*     *Contribution to lower discounting*

Atrophy pattern of bvFTD (5.A)    **X**    Brain pattern predicting discounting (5.B)    **=**    Importance map = pattern of contribution to alter discounting in bvFTD (5.C)

**A. Thresholded *T*-map of bvFTD vs Controls contrast (bvFTD<Controls) on whole-brain grey matter density**

*T*-values   5.54 — 8.93

**B. Whole-brain weight map of the structural brain pattern predicting delay discounting**

*β*-values   -0.005 — 0.005

**C. Thresholded *T*-map of bvFTD vs Controls contrast (bvFTD>Controls) on whole-brain importance map**

*T*-values   -9 — 9