Target journal: Frontiers in Drug Discovery

Special issue: Transport of Peptides to the CNS

Full title: Sympathetic innervation of interscapular brown adipose tissue is not a

predominant mediator of OT-elicited reductions of body weight gain and adiposity in male

diet-induced obese rats

Abbreviated title: Contribution of SNS to Oxytocin-Elicited Weight Loss

Melise M. Edwards[1], Ha K. Nguyen[1], Andrew D. Dodson[1], Adam J. Herbertson[1],

Mackenzie K. Honeycutt[1], Jared D. Slattery[1], June R. Rambousek[1], Edison Tsui[1], Tami

Wolden-Hanson[1], Tomasz Wietecha[2,4], James L. Graham[5], Geronimo P. Tapia[7,8], Carl

Sikkema[1,6], Kevin D. O'Brien[3,4], Thomas O. Mundinger[2], Elaine R. Peskind[1,6], Vitaly Ryu[9],

Peter J. Havel[5,6], Arshad M. Khan[7,8], Gerald J. Taborsky, Jr.[1,2], and James E. Blevins[1,2]

[1]VA Puget Sound Health Care System, Office of Research and Development Medical

Research Service, Department of Veterans Affairs Medical Center, Seattle, WA 98108,

USA

[2]Division of Metabolism, Endocrinology and Nutrition, Department of Medicine, University

of Washington School of Medicine, Seattle, WA 98195, USA

[3]Division of Cardiology, Department of Medicine, University of Washington School of

Medicine, Seattle, WA 98195, USA

[4]UW Medicine Diabetes Institute, University of Washington School of Medicine, Seattle,

WA 98109

[5]Department of Nutrition, University of California, Davis, CA 95616, USA

[6]Department of Molecular Biosciences, School of Veterinary Medicine, University of

California, Davis, CA 95616, USA

[6]Department of Psychiatry and Behavioral Sciences, University of Washington School of

Medicine, Seattle, Washington

[7]UTEP Systems Neuroscience Laboratory, Department of Biological Sciences, The
University of Texas at El Paso, El Paso, TX 79968, USA

[8]Border Biomedical Research Center, The University of Texas at El Paso, El Paso, TX
79968, USA

[9]Icahn School of Medicine at Mount Sinai, New York, NY, USA

Corresponding author's address:

J. E. Blevins, Ph.D.

VA Puget Sound Health Care System

Research-151

1660 South Columbian Way

Seattle, WA 98108

Key words: Obesity, brown adipose tissue, white adipose tissue, oxytocin, sympathetic

nervous system

This material was based upon work supported by the Office of Research and
Development, Medical Research Service, Department of Veterans Affairs (VA) and the
VA Puget Sound Health Care System Rodent Metabolic Phenotyping Core and the

Cellular and Molecular Imaging Core of the Diabetes Research Center at the University
of Washington and supported by National Institutes of Health (NIH) grant P30DK017047.
This work was also supported by the VA Merit Review Award 5 I01BX004102, from the
United States (U.S.) Department of Veterans Affairs Biomedical Laboratory Research and
Development Service and NIH 5R01DK115976 grant to James Blevins. PJH's research
program also received research support during the project period from NIH grants DK-
095980, HL-091333, HL-107256 and a multi-campus grant from the University of
California Office of the President. Work in the UTEP Systems Neuroscience Laboratory
is supported by NIH Grant SC1GM127251 awarded to AMK. GPT is a Dodson Research
Excellence Funds recipient.
58
59 The contents do not represent the views of the U.S. Department of Veterans Affairs or
60 the United States Government.
61

ACKNOWLEDGMENTS
The authors thank the technical support of Nishi Ivanov, Hailey Chadwick, Alex Vu and
Ana R. Arvizu Morales. In addition, the authors are appreciative of the efforts by Drs.
Michael Schwartz and Dianne Lattemann for providing feedback throughout the course
of these studies. Disclosures: JEB had a financial interest in OXT Therapeutics, Inc., a
company developing highly specific and stable analogs of oxytocin to treat obesity and
metabolic disease. The author's interests were reviewed and are managed by their local
institutions in accordance with their conflict-of-interest policies. The other authors have
nothing to report.
71

Abstract

Recent studies indicate that central administration of oxytocin (OT) reduces body weight (BW) in high fat diet-induced obese (DIO) rodents by reducing energy intake and increasing energy expenditure (EE). Previous studies in our lab have shown that administration of OT into the fourth ventricle (4V; hindbrain) elicits weight loss and stimulates interscapular brown adipose tissue temperature (T_{IBAT}) in DIO rats. We hypothesized that OT-elicited stimulation of sympathetic nervous system (SNS) activation of IBAT contributes to its ability to activate BAT and reduce BW in DIO rats. To test this, we determined the effect of disrupting SNS activation of IBAT on OT-elicited stimulation of T_{IBAT} and reduction of BW in DIO rats. We first confirmed that bilateral surgical SNS denervation to IBAT was successful based on having achieved ≥ 60% reduction in IBAT norepinephrine (NE) content from DIO rats. NE content was selectively reduced in IBAT by 94.7 ± 2.7, 96.8 ± 1.8 and 85.9 ± 6.1% ($P<0.05$) at 1, 6 and 7-weeks post-denervation, respectively, and was unchanged in liver or inguinal white adipose tissue. We then measured the impact of bilateral surgical SNS denervation to IBAT on the ability of acute 4V OT (1, 5 µg) to stimulate T_{IBAT} in DIO rats. We found that the high dose of 4V OT (5 µg) stimulated T_{IBAT} similarly between sham and denervated rats (P=NS) and that the effects of 4V OT to stimulate T_{IBAT} did not require beta-3 adrenergic receptor signaling. We subsequently measured the effect of bilateral surgical denervation of IBAT on the effect of chronic 4V OT (16 nmol/day) or vehicle infusion to reduce BW, adiposity and energy intake in DIO rats. Chronic 4V OT reduced BW gain by −7.2 ± 9.6 g and −14.1 ± 8.8 g in sham and denervated rats ($P<0.05$ vs vehicle treatment), respectively, and this effect was similar between groups (P=NS). These effects were associated with reductions

105   in adiposity and energy intake ($P<0.05$). Collectively, these findings support the

106   hypothesis that sympathetic innervation of IBAT is not required for central OT to increase

107   BAT thermogenesis and reduce BW gain and adiposity in male DIO rats.

113
114
115

Introduction

While the pleiotropic hormone, oxytocin (OT), is largely associated with male and female sexual and reproductive behavior (i.e. uterine contraction, milk ejection reflex, and penile erection) [1; 2] and social behaviors (i.e. parent-infant bonds, pair bonds, and increased trust) [3; 4], questions remain as far as how OT functions in the regulation of body weight (BW) [5; 6; 7; 8]. Although suppression of food intake is thought to contribute to OT-elicited weight loss, pair-feeding studies in mice and rats suggest that OT's effects on weight loss cannot be completely explained by its ability to reduce food intake [9; 10; 11].

In addition to OT's well-characterized effects on food intake, previous studies have shown that OT increases other metabolic functions that control BW, namely energy expenditure (EE) [12; 13; 14; 15]. While brown adipose tissue thermogenesis (BAT) is important in the control of EE (see [16; 17] for review), it is unclear if OT's effects on EE stem from 1) non-shivering BAT thermogenesis, 2) non-shivering and shivering thermogenesis in skeletal muscle [18], 3) spontaneous physical activity-induced thermogenesis [19], 4) white adipose tissue thermogenesis or 5) hormonal mediators (e.g. leptin [20], secretin [21; 22], irisin [23], fibroblast growth factor-21 [24] or thyroid hormone [25] (see [26; 27] for review). We have found that acute CNS injections of OT increase interscapular BAT temperature (T_{IBAT}) (functional readout of BAT thermogenesis) in mice and rats [28; 29]. Moreover, the timing of OT-elicited elevations of T_{IBAT} coincide with the onset of OT-elicited reductions of BW in diet-induced obese (DIO) rats [28]. Sutton and colleagues reported that chemogenetic activation of OT neurons within the paraventricular nucleus (PVN) stimulates both subcutaneous BAT

temperature and EE in *Oxytocin-Ires-Cre* mice [30]. In addition, Yuan et al found that

systemic administration of OT increases thermogenic gene expression in IBAT and

stimulates BAT differentiation *in vitro* in DIO mice [31]. In contrast, genetic or

pharmacologically-induced deficits in OT signaling are associated with reductions in BAT

thermogenesis [32; 33; 34; 35], EE [14; 15; 32; 36] and obesity [15; 36; 37; 38] in mice.

More recently, we found that SNS innervation of IBAT is not a predominant mediator of

OT-elicited reductions of BW and adiposity in male diet-induced obese mice [29]. One

outstanding question is whether OT-elicited elevation of IBAT thermogenesis and weight

loss require increased sympathetic nervous system (SNS) outflow to IBAT in DIO rats

and whether the effect of OT on BAT thermogenesis may involve beta-3 adrenergic

receptors (β3-AR) and hindbrain OTR. Here, we sought to clarify the role of SNS outflow

to IBAT in mediating the effects of hindbrain OTR stimulation on both weight loss and

BAT thermogenesis in a DIO rat model. Furthermore, we measured gross motor activity

to test the extent to which increases in spontaneous physical activity may contribute to

the effects of acute 4V OT on BAT thermogenesis.

We hypothesized that OT-elicited stimulation of sympathetic nervous system (SNS)

activation of IBAT contributes to its ability to activate BAT and reduce BW in DIO rats. We

initially verified the success of the IBAT denervation procedure by measuring IBAT

norepinephrine (NE) content at 1-week post-denervation in lean rats. We then determined

whether these effects can be replicated in DIO rats at 1, 6 and 7-weeks post-

sham/denervation. To assess the role of SNS innervation of BAT in contributing to OT-

elicited increases in BAT thermogenesis, we determined the impact of bilateral surgical

SNS denervation to IBAT on the ability of acute 4V OT (1, 5 μg) to stimulate T_{IBAT} (as

functional measure of BAT thermogenesis) in DIO rats. To assess the role of SNS

innervation of BAT in contributing to OT-elicited weight loss, we subsequently measured

the effect of bilateral surgical denervation of IBAT on the effect of chronic 4V OT (16

nmol/day over 29 days) or vehicle infusion to reduce BW, adiposity and energy intake in

DIO rats. We subsequently examined whether these effects were associated with a

decrease adipocyte size and energy intake. In addition, we measured the temporal profile

of acute 4V OT on BAT temperature, core temperature and gross motor activity to

determine if spontaneous physical activity might have contributed to the effects of 4V OT

on BAT thermogenesis. Lastly, we determined the extent to which 4V OT activates an

early marker of neuronal activation, phosphorylated extracellular signal-regulated kinases

1 and 2-immunoreactivity (pERK1/2), within the hindbrain NTS in DIO rats.

Methods

Animals

Adult male Long-Evans rats (~ 326–932 g BW) were initially obtained from Envigo

(Indianapolis, IN) and maintained for at least 4 months on a high fat diet (HFD) prior to

study onset. All animals were housed individually in Plexiglas cages in a temperature-

controlled room (22±2°C) under a 12:12-h light-dark cycle. All rats were maintained on a

6 a.m./6 p.m. light cycle except for those in **Study 7A** [(7 a.m./7 p.m. light cycle (telemetry

studies) or 1 a.m./1 p.m. reversed light cycle (food intake studies)] and **Study 7B** (1 a.m./1

p.m. reversed light cycle). Rats had *ad libitum* access to water and a HFD providing 60%

kcal from fat (Research Diets, D12492, New Brunswick, NJ). By design, rats used in

Study 1 and **Supplemental Study 1** were maintained on a low-fat chow diet (13% kcal

from fat; diet no. 5001, LabDiet, St. Louis, MO). The research protocols were approved

both by the Institutional Animal Care and Use Committee of the Veterans Affairs Puget

Sound Health Care System (VAPSHCS) and the University of Washington in accordance

with NIH's *Guide for the Care and Use of Laboratory Animals* (NAS, 2011) [39] .

Drug Preparation

Fresh solutions of OT acetate salt (Bachem Americas, Inc., Torrance, CA) were prepared

the day of each experiment. OT was solubilized in sterile water (**Studies 5 and 6**). Fresh

solutions of OT acetate salt (Bachem Americas, Inc., Torrance, CA) were solubilized in

sterile water, loaded into Alzet® minipumps (model 2004; DURECT Corporation,

Cupertino, CA) and subsequently primed in sterile 0.9% saline at 37º C for approximately

40 hours prior to minipump implantation based on manufacturer's recommended

instructions (**Study 6**). The β3-AR agonist**,** CL 316243 (Tocris/Bio-Techne Corporation,

Minneapolis, MN)**,** was solubilized in sterile water each day of each experiment (**Study**

3; Supplemental Study 2). The β3-AR antagonist, SR 59230A, was dissolved in 100%

dimethyl sulfoxide (DMSO) each day of each experiment (**Study 5**).

SNS denervation procedure

A dissecting microscope (Leica M60/M80; Leica Microsystems, Buffalo Grove, IL) was

used throughout the procedure. A 1" midline incision was made in the skin dorsally at the

level of the thorax and continued rostrally to the base of the skull. Connective tissue was

blunt-dissected away from the adipose tissue with care to avoid cutting the large

thoracodorsal artery that is located medially to both pads. Both left and right fat pads were

separated from the midline. Each fat pad was lifted up and the intercostal nerve bundles were located below. Once the nerves were located, a sharp-point forceps was used to pull the nerve bundles straight up while using a 45 degree scissors to cut and remove 3-5 mm of nerves. The interscapular incision was closed with 4-0 non-absorbable monofilament Ethilon (nylon) sutures or with standard metal wound clips. Nerves were similarly identified but not cut for sham-operated animals. Rats were treated pre-operatively with the analgesic ketoprofen (2 mg/kg; Fort Dodge Animal Health, Overland Park, KS) prior to the completion of the denervation or sham procedure. This procedure was combined with transponder implantations for studies that involved IBAT temperature measurements in response to acute 4V injections (**Study 4**). Animals were allowed to recover for approximately 5–7 days prior to implantation of 4V cannulas. Note that surgical denervation was used in place of chemical denervation because chemical denervation can result in SNS terminal recovery within IBAT [40; 41] within only 10 days post-chemical denervation [40].

4V cannulations for acute injections

Animals were implanted with a cannula (P1 Technologies, Roanoke, VA) that was directed towards the 4V as previously described [42; 43; 44]. Briefly, rats under isoflurane anesthesia were placed in a stereotaxic apparatus [Digital Lab Standard Stereotaxic, Rat, (Item 51900), Stoelting Co., Wood Dale, IL] with the incisor bar positioned 3.3 mm below the interaural line. A 26-gauge cannula (P1 Technologies) was stereotaxically positioned into the 4V [–3.5 mm caudal to the interaural line; 1.4 mm lateral to the midline, and 6.2

mm ventral to the skull surface [45]] and secured to the surface of the skull with dental

cement and stainless-steel screws.

4V cannulations for chronic infusions

Briefly, rats were implanted with a cannula within the 4V with a side port that was

connected to an osmotic minipump (model 2004, DURECT Corporation, Cupertino, CA)

as previously described [28; 46]. Rats under isoflurane anesthesia were placed in a

stereotaxic apparatus [Digital Lab Standard Stereotaxic, Rat, (Item 51900) Stoelting Co.]

with the incisor bar positioned 3.3 mm below the interaural line. A 30-gauge cannula (P1

Technologies) was stereotaxically positioned into the 4V [–3.5 mm caudal to the interaural

line; 1.4 mm lateral to the midline, and 7.2 mm ventral to the skull surface [45]] and

secured to the surface of the skull with dental cement and stainless steel screws. A 2.4"

piece of plastic Tygon^{TM} Microbore Tubing (0.020" × 0.060" OD; Cole-Parmer) was

tunneled subcutaneously along the midline of the back and connected to the 21-gauge

sidearm osmotic minipump-cannula assembly. A stainless steel 22-gauge pin plug

(Instech Laboratories, Inc., Plymouth Meeting, PA) was temporarily inserted at the end of

the tubing during a two week postoperative recovery period, after which it was replaced

by an osmotic minipump (DURECT Corporation) containing saline or OT. Rats were

treated with the analgesic ketoprofen (2 mg/kg; Fort Dodge Animal Health) and the

antibiotic enrofloxacin (5 mg/kg; Bayer Healthcare LLC., Animal Health Division,

Shawnee Mission, KS) at the completion of the 4V cannulations and were allowed to

recover at least 10 days prior to implantation of osmotic minipumps.

Implantation of implantable telemetry devices (PDT-4000 E-Mitter) into abdominal cavity

Animals were anesthetized with isoflurane and subsequently underwent a sham surgery (no implantation) or received implantations of a sterile PDT-4000 E-Mitter (23 mm long × 8 mm wide; Starr Life Sciences Company) into the intraperitoneal cavity. The abdominal opening was closed with 4-0 Vicryl absorbable suture and the skin was closed with 4-0 monofilament nonabsorbable suture. Vetbond glue was used to seal the wound and bind any tissue together between the sutures. Sutures were removed within two weeks after the PDT-4000 E-Mitter implantation. All PDT-4000 E-Mitters were confirmed to have remained within the abdominal cavity at the conclusion of the study.

Implantation of temperature transponders underneath IBAT

Animals were anesthetized with isoflurane and had the dorsal surface along the upper midline of the back shaved and the area was scrubbed with 70% ethanol followed by betadine swabs. A one-inch incision was made at the midline of the interscapular area. The temperature transponder (14 mm long/2 mm wide) (HTEC IPTT-300; Bio Medic Data Systems, Inc., Seaford, DE) was implanted underneath the left IBAT pad as previously described [28; 47; 48] and secured in place by suturing it to the brown fat pad with sterile silk suture. The interscapular incision was closed with Nylon sutures (5-0), which were removed in awake animals 10–14 days after surgery.

Acute IP or 4V injections and measurements of T_{IBAT}

On an experimental day, animals received either IP (CL 316243 or saline vehicle; 0.1 ml/kg injection volume) or 4V injections (OT or saline vehicle; 1 μL injection volume, 4V)

290 during the early part of the light cycle following 4 hours of food deprivation. Injections

291 were completed in a crossover design over approximately 7-day (CL 316243) or 48-h

292 (OT) intervals such that each animal served as its own control. Animals remained without

293 access to food for an additional 4 h (**Studies 3 & 4**) during the T_{IBAT} measurements. A

294 handheld reader (DAS-8007-IUS Reader System; Bio Medic Data Systems, Inc.) was

295 used to collect measurements of T_{IBAT}. Rats underwent all treatments in a randomized

296 order separated by at least 7 or 8 days between treatments.

298 *Changes of Core Temperature and Gross Motor Activity.* Telemetry recordings of core

299 temperature and gross motor activity were measured from each rat in the home cage

300 immediately prior to injections and for a 4-h period after injections. Core temperature and

301 gross motor activity were recorded every 5 min.

303 *Body Composition*

304 Determinations of lean body mass and fat mass were made on un-anesthetized rats by

305 quantitative magnetic resonance using an EchoMRI 4-in-1-700^{TM} instrument (Echo

306 Medical Systems, Houston, TX) at the VAPSHCS Rodent Metabolic Phenotyping Core.

307 Measurements were taken prior to 4V cannulations and minipump implantations as well

308 as at the end of the infusion period.

310 *Tissue collection for NE content measurements*

311 Rats were euthanized by rapid conscious decapitation at 1 week (**Study 1**), 6 weeks

312 (**Study 2**), 7 weeks (**Study 2, Study 5**), 9 weeks (**Study 6**) or 10–11 weeks (**Study 4**)

post-sham or denervation procedure. Trunk blood and tissues (IBAT, EWAT, IWAT, liver

and/or pancreas) were collected from 4-h fasted rats. Tissue was rapidly removed,

wrapped in foil and frozen in liquid N_2. Samples were stored frozen at −80°C until analysis.

Note that anesthesia was not used when collecting tissue for NE content as it can cause

the release of NE from SNS terminals within the tissue [41].

Norepinephrine (NE) content measurements (biochemical confirmation of IBAT

denervation procedure)

NE content was measured in IBAT, EWAT, IWAT, liver and/or pancreas using previously

established techniques [49]. Successful denervation was noted by $\geq 60\%$ reduction in

IBAT NE content as previously noted [50]. Experimental animals that did not meet this

criterion were excluded from the data analysis.

Study Protocols

Study 1A: Determine the success of the surgical denervation of only the superficial

nerves at 1-week post-sham or denervation in lean rats by measuring NE content.

Rats underwent sham or bilateral SNS denervation procedures of only the superficial

nerves that innervate IBAT and, to prevent the confound of anesthesia on NE content,

animals were euthanized by rapid conscious decapitation at 1-week post-sham or

denervation procedure.

Study 1B: Determine the success of the surgical denervation procedure at 1-week post-sham or denervation in lean rats by measuring NE content.

Rats underwent sham or bilateral SNS denervation procedures of both superficial and more ventrally located intercostal nerves that innervate IBAT and, to prevent the confound of anesthesia on NE content, animals were euthanized by rapid conscious decapitation at 1-week post-sham or denervation procedure.

Study 2A: Determine the success of the surgical denervation procedure at 1, 6 and 7-weeks post-sham or denervation in DIO rats by measuring NE content.

Rats were fed *ad libitum* and maintained on HFD for approximately 5 months prior to underdoing sham or SNS denervation procedures. In addition to weekly BW measurements, body composition measurements were obtained at baseline and at 6 and 7-weeks post-denervation/sham procedures. Rats were euthanized by rapid conscious decapitation 1, 6 and 7-weeks post-sham or denervation procedure.

Study 3: Determine if surgical denervation of IBAT changes the ability of the β-3R agonist, CL 316243, to increase T_{IBAT} in DIO rats.

Rats were fed *ad libitum* and maintained on HFD for approximately 4.25 months prior to underdoing sham or SNS denervation procedures and implantation of temperature transponders underneath the left IBAT depot. Rats were allowed to recover for at least 1 week during which time they were adapted to a daily 4-h fast, handling and mock injections. On an experimental day, 4-h fasted rats received Cl 316243 (0.1 or 1 mg/kg) or vehicle (sterile water) during the early part of the light cycle in a crossover design at

approximately 7-day intervals such that each animal served as its own control (approximately 1–3 weeks post-sham or denervation procedures). T_{IBAT} was measured at baseline (–2 h; 9:00 a.m.), immediately prior to IP injections (0 h; 9:45–10:00 a.m.), and at 0.25, 0.5, 0.75, 1, 1.25, 1.5, 2, 3, 4, and 24-h post-injection (10:00 a.m.). Food intake and BW were measured daily. This dose range was based on doses of CL 316243 found to be effective at reducing food intake and weight gain in rats [51; 52]. Animals were euthanized by rapid conscious decapitation at 13 weeks post-sham or denervation procedure.

Study 4: Determine the extent to which OT-induced activation of sympathetic outflow to IBAT contributes to its ability to increase T_{IBAT} in DIO rats.

Rats from **Study 3** were implanted with 4V cannulas approximately 1 month following sham/denervation procedures and implantation of temperature transponders underneath the left IBAT depot. Rats were allowed to recover for at least 2 weeks during which time they were adapted to a daily 4-h fast, handling and mock injections. On an experimental day, 4-h fasted rats received OT (1 or 5 μg/μl) or vehicle during the early part of the light cycle in order to maximize the effects of OT [15; 52] during a time when circulating NE levels [53] and IBAT catecholamine levels are lower [54]. Injections were completed in a crossover design at approximately 48-h intervals such that each animal served as its own control (approximately 7-weeks post-sham or denervation procedures). T_{IBAT} was measured at baseline (–2 h; 9:00 a.m.), immediately prior to 4V injections (0 h; 9:45–10:00 a.m.), and at 0.25, 0.5, 0.75, 1, 1.25, 1.5, 2, 3, 4, and 24-h post-injection (10:00

a.m.). Food intake and BW were measured daily. This dose range was based on doses

of 4V OT found to be effective at stimulating T_{IBAT} in DIO rats in previous studies [28].

Study 5: Determine the extent to which OT-induced activation of sympathetic

outflow to IBAT requires activation of β3-AR to increase T_{IBAT} in DIO rats

Rats were fed *ad libitum* and maintained on HFD for approximately 4 months prior to

being implanted with temperature transponders underneath the left IBAT depot. Rats

were subsequently implanted with 4V cannulas approximately 1 month following

implantation of temperature transponders. Rats were allowed to recover for 4 weeks

during which time they were adapted to a daily 4-h fast, handling and mock injections. On

an experimental day, 4-h fasted rats received the β3-AR antagonist, SR 59230A (0.5

mg/kg, IP) or vehicle (DMSO) approximately 20 min prior to 4V administration of either

OT (5 μg/μl) or vehicle. Injections occurred during the early part of the light cycle in order

to maximize the effects of OT [15; 52] during a time when circulating NE levels [53] and

IBAT catecholamine levels are lower [54]. Injections were completed in a crossover

design at approximately 7-day intervals such that each animal served as its own control.

T_{IBAT} was measured at baseline (-2 h; 9:00 a.m.), immediately prior to 4V injections (0 h;

9:45-10:00 a.m.), and at 0.25, 0.5, 0.75, 1, 1.25, 1.5, 2, 3, 4, and 24-h post-injection (10:00

a.m.). Food intake and BW were measured daily. This dose of SR 59230A was based on

other studies [55; 56; 57] and doses that we found to be subthreshold for reducing IBAT

temperature in our pilot studies (data not shown) and able to block the effects of the β3-

AR agonist, CL 316243, on T_{IBAT} in DIO rats (**Supplemental Figure 1**).

Study 6A: Determine the extent to which OT-induced activation of sympathetic outflow to IBAT contributes to its ability to elicit weight loss in DIO rats.

Rats were fed *ad libitum* and maintained on HFD for at least 4.5 months prior to receiving 4V cannulas and minipumps to infuse vehicle or OT (16 nmol/day) over 29 days as previously described [28]. This dose was selected based on a dose of 4V OT found to be effective at reducing BW in DIO rats [28]. Daily food intake and BW were also tracked for 29 days. Animals were euthanized by rapid conscious decapitation at 7 weeks post-sham or denervation procedure. Trunk blood and tissues (IBAT, EWAT, IWAT, liver and pancreas) were collected from 4-h fasted rats and tissues were subsequently analyzed for IBAT NE content to confirm success of denervation procedure relative to sham-operated animals and other tissues (EWAT, IWAT, liver and pancreas).

Study 6B: Determine the success of the surgical denervation procedure at 6-8 weeks post-sham or denervation in DIO rats by measuring thermogenic gene expression.

Rats from **Study 6A** were used for these studies. All rats underwent sham or bilateral surgical SNS denervation procedures prior to being implanted with 4V cannulas and minipumps. Animals were euthanized by rapid conscious decapitation at approximately 6–8 weeks post-sham or denervation procedure. Tissues were collected following a 4-h fast.

Study 6C: Determine the extent to which OT-induced activation of sympathetic outflow to IBAT impacts thermogenic gene expression in IBAT and EWAT in DIO mice.

Rats from **Study 6A** were used for these studies. All rats underwent sham or bilateral surgical SNS denervation procedures prior to being implanted with 4V cannulas and minipumps. Animals were euthanized by rapid conscious decapitation at approximately 6–8 weeks post-sham or denervation procedure. Tissues were collected following a 4-h fast.

Study 7A: Determine the extent to which 4V OT increases gross motor activity, core temperature, T$_{IBAT}$ and food intake in DIO rats.

Telemetry measurements:

Rats were fed *ad libitum* and maintained on HFD for approximately 6 months prior to being implanted with temperature transponders underneath IBAT, intra-abdominal telemetry devices and 4V cannulas as previously described. Rats were implanted temperature transponders approximately 4 days prior to being implanted with intra-abdominal telemetry devices. Animals were allowed up to 3 weeks post-op recovery prior to being implanted with 4V cannulas. Rats were allowed to recover for at least 2 weeks during which time they were adapted to a daily 4-h fast, handling and mock injections. Like **Study 4**, on an experimental day, 4-h fasted rats received OT (1 or 5 μg/μl) or vehicle during the early part of the light cycle. Injections were completed in a crossover design at approximately 48-h intervals such that each animal served as its own control. T$_{IBAT}$ was measured at baseline (-2 h; 9:00 a.m.), immediately prior to 4V injections (0 h; 9:45-10:00

a.m.), and at 0.25, 0.5, 0.75, 1, 1.25, 1.5, 2, 3, 4, and 24-h post-injection (10:00 a.m.).

Food intake and BW were measured daily.

Food intake measurements:

Animals were treated identically as described above with the exception that animals were

adapted to a 1 a.m. (lights on)/1 p.m. (lights off) light cycle and animals received acute

4V injections at the end of the light cycle (12:40-1 p.m.). Food was returned at the onset

of the dark cycle. Food was measured at 0.5, 1, 2, 3, 4 and 24-h post-injection.

Study 7B: Determine the extent to which 4V OT activates an early marker of

neuronal activation, phosphorylated extracellular signal-regulated kinases 1 and 2-

immunoreactivity (pERK1/2), within the hindbrain NTS in DIO rats.

Rats from **Study 7A** were used for this study. Following the conclusion of **Study 7A**, 4-h

fasted rats received OT (5 μg/μl) or vehicle during the late part of the light cycle. Animals

remained fasted for an additional 0.25 or 0.5-h post-injection, at which time animals were

euthanized with an IP overdose of ketamine cocktail and transcardially perfused with PBS

followed by 4% paraformaldehyde in 0.1 M PBS. Brains were removed, stored overnight

in fresh fixative at 4°C and subsequently transferred to 0.1 M PBS containing 25%

sucrose for 48 h. Brains were then frozen by submersion for 20–30 s in isopentane chilled

with dry ice.

Tissue Processing and Immunohistochemistry

Perfusion-fixed frozen brain specimens were shipped overnight in dry ice to The University of Texas at El Paso for histological analysis, but arrived thawed due to an unforeseen shipping delay beyond our control. Therefore, they were prepared to be re-frozen by placing them into a 20% sucrose cryoprotectant solution and stored at 4°C before processing. Phospho-ERK1/2 labeling in the dorsal vagal complex did not appear to be adversely affected in our peroxidase-based reactions and was found in the same cellular compartments previously noted by others [58; 59].

Brains were coronally-cut into two tissue blocks, and the hindbrain portion was re-frozen with its rostral face down onto an ice-cold solid brass freezing stage (3.25 × 7.25 × 1.5 in; Brain Research Laboratories, Newton, MA, USA; cat #3488-RJ) modified to fit a modern 'Naples' object-holder and carrier of a sliding microtome (Reichert; Vienna, Austria; Nr. 15 156). Dry ice pellets in the stage troughs were used to keep the stage at freezing temperatures. The frozen blocks were cut into 30 µm-thick coronal sections, collected as six series into 24-well tissue culture plates filled with cryoprotectant (50% 0.05 M sodium phosphate buffer, 30% ethylene glycol, 20% glycerol; pH 7.4 at room temperature), and stored at −20°C before undergoing further processing.

Using Tris-buffered saline (TBS) (0.05 M; pH 7.4 at room temperature), sections were washed of cryoprotectant (5 washes for 5 minutes each; 5 × 5) and were incubated in 0.007% phenylhydrazine solution for 20 minutes at room temperature. This was followed by 5 × 5 TBS washes, then an incubation in blocking solution consisting of TBS containing 2% (v/v) normal donkey serum (catalog #S30-100ML; EMD-Millipore, Burlington, MA) and 0.1% (v/v) Triton X-100 for ~1 hour at room temperature. The tissue

was then reacted with an antibody solution consisting of blocking solution containing a

primary antibody targeting the phosphorylated forms of ERKs 1 and 2 (pERK1/2; EC

2.7.11.24) (1:100 dilution; catalog #9101S; Cell Signaling Technology, Danvers, MA;

RRID:AB_331646) for ~12 hours at 4°C. This was followed by a 5×5 wash period, then

reaction of the sections with an affinity-purified donkey anti-rabbit IgG secondary antibody

(1:500 dilution; catalog #711-065-152; Jackson ImmunoResearch, West Grove, PA;

RRID:AB_2340593) for ~5 hours at room temperature. The sections underwent another

5×5 wash before incubation in an avidin-biotin complex (ABC) reagent (VECTASTAIN®

ABC-HRP Kit, catalog #PK-4000; Vector Laboratories, Newark, CA; RRID:AB_2336818)

for ~1 hour at room temperature. The tissue was rinsed via a 5×5 TBS wash before

being subjected to a Ni^{2+}-enhanced 3,3'-diaminobenzidene peroxidase reaction to

visualize antibody labeling.

Following the reaction, the sections underwent a final 5×5 wash before being

mounted onto gelatin subbed slides. Once air-dried, the mounted sections were

dehydrated in ascending concentrations of ethanol (50%, 70%, 95%, $3 \times 100\%$; 3 min

each), and cleared in xylenes (2×12 min). As individual slides were removed from the

staining jar, they were immediately coated with 5 drops of dibutyl phthalate xylenes (DPX)

mountant for histology (cat #06522-100ML, EMD-Millipore, Burlington, MA), coverslipped,

and laid to dry.

Widefield Microscopy

Using an Olympus BX63 light microscope with an X–Y–Z motorized stage and an

attached DP74 color CMOS camera (cooled, 20.8 MP pixel-shift, 60 fps; Olympus

America, Inc., Waltham, MA), photomicrographs were captured under bright-field and

dark-field illumination. Image acquisition was processed using cellSense™ Dimension

software installed on a Hewlett Packard PC workstation. Widefield mosaic

photomicrographs of the sections were captured using a ×20 magnification objective lens

(Olympus UPlanSApo, N.A. 0.50, FN 26.5 mm) and 15% overlap of stitched tiles. Each

image was exported as full-resolution uncompressed TIFF-formatted files.

Anatomical observations for determining the plane of section

To assess the recruitment of ERK1/2 in the NTS of OT- and vehicle-treated

animals, we relied on the *Brain Maps 4.0 (BM4.0)* open-access atlas [60]

(RRID:SCR_017314) to localize sections to a particular region of the dorsal vagal

complex (*BM4.0* atlas level 70) to have a stable landmark for comparing immunoreactive

signal. This was performed by referencing the region delineations represented in *BM4.0*

templates of caudal hindbrain levels, relying mostly on the morphology of the area

postrema (AP). The AP is represented at two *BM4.0* atlas levels, levels 69 (Bregma =

−13.76 mm) and 70 (Bregma = −14.16 mm), respectively. At level 69, the dorsal AP

boundaries extend laterally, and the nucleus has a more flattened dorsal border when

compared to more caudal representations. Ventral to the NTS, the DMX is a large, blimp-

shaped cell group that loses height and decreases in cellular area traveling in the

posterior direction. At level 70, the AP takes on a rounder shape as it reduces in size

proceeding towards its caudal pole nestled in the dorsal midline. The cytoarchitectural

features of these regions surrounding the NTS are reliable landmarks for the

determination of a rostral or caudal AP-containing section, and were used to select the

sections that best corresponded to atlas level 70 for analysis here.

Regional Parcellations and Cell Tabulation

Once the sections that most closely adhered to *BM4.0* atlas level 70 region

boundaries were identified, the photomicrographs were imported into Adobe Illustrator

CC (AI) (Version 28.6) and annotated manually. This was performed by superimposing

the bright-field and dark-field section images over the *BM4.0* digital template. In a

separate layer, the reviewer used the *Pencil Tool* to parcellate the region boundaries of

the immunolabeled section, relying on the cytoarchitectural features to guide their

tracings. The myeloarchitecture, that is more easily visualized under dark-field viewing

conditions, was used to trace important white matter areas, such as the solitary tract. If a

region was displaced due to damage of the tissue, then its inferred borders were

represented.

Another layer was created to mark the locations of pERK1/2-immunoreactive cell

profiles using the *Ellipse Tool*, in which the reviewer placed a colored ellipse over each

immunolabeled perikaryon. Considering that the borders of the ventral NTS and dorsal

DMX do not have a clear-cut boundary but rather contain an intermixing of cells, the

morphology of each annotated cell profile was assessed for its consistency with the

smaller neuronal size of the NTS or larger perikaryon of the motor DMX, so as to only

consider activation of the NTS here.

Atlas nomenclature

We used the standardized nomenclature system provided by Swanson (2018) [56]

to describe and name regions of the hindbrain for which we mapped activation patterns.

This approach is being used by us as part of our longstanding effort to encourage

investigators to place their data into a common spatial model so that they are interrelated

with other datasets mapped in the same space [61; 62]. This also requires a common

nomenclature. First devised in a comprehensive reference work for human brain [63], the

nomenclature system includes an *(author, date)* attribution as *part of the name of the*

structure, to help readers know the provenance of where the term was first described as

defined in the atlas. For those terms where such information was not traced, the

assignation "(>1840)" is used instead to note that it was in use sometime after the

development of cell theory. We applied this nomenclature to those regions of the

hindbrain we examined for the early activation marker, phosphorylated p44/42 MAP

Kinases (pERK1/2). These regions include the *area postrema (>1840)* (AP), *nucleus of*

solitary tract (>1840) (NTS), and *dorsal motor nucleus of vagus nerve (>1840)* (DMX).

Blood collection

Trunk blood was collected from 4-h (**Studies 1 & 2**, **Study 6A–C**), 4.25 or 4.5 h

(**Study 7B**) or 6-h (**Studies 4 & 5**) fasted rats within a 2-h window towards the beginning

of the light cycle (10:00 a.m.–12:00 p.m.) as previously described in DIO CD® IGS and

Long-Evans rats and C57BL/6J mice [28; 43] with the exception of **Study 7B**, in which

blood was collected towards the end of the light cycle. Treatment groups were

counterbalanced at time of euthanasia to avoid time of day bias. Blood samples [up to 3

mL] were collected immediately prior to transcardial perfusion by cardiac puncture in

chilled K2 EDTA Microtainer Tubes (Becton-Dickinson, Franklin Lakes, NJ). Whole blood

was centrifuged at 6,000 rpm for 1.5 min at 4°C; plasma was removed, aliquoted and

stored at −80°C for subsequent analysis.

Plasma hormone measurements

Plasma leptin and insulin were measured using electrochemiluminescence detection

[Meso Scale Discovery (MSD®), Rockville, MD] using established procedures [28; 64].

Intra-assay coefficient of variation (CV) for leptin was 5.9% and 1.6% for insulin. The

range of detectability for the leptin assay is 0.137–100 ng/mL and 0.069–50 ng/mL for

insulin. Plasma fibroblast growth factor-21 (FGF-21) (R&D Systems, Minneapolis, MN)

and irisin (AdipoGen, San Diego, CA) levels were determined by ELISA. The intra-assay

CV for FGF-21 and irisin were 2.2% and 9.1%, respectively; the ranges of detectability

were 31.3–2,000 pg/mL (FGF-21) and 0.078–5 μg/mL (irisin). Plasma adiponectin was

also measured using electrochemiluminescence detection [Millipore Sigma (Burlington,

MA)] using established procedures [28; 64]. The intra-assay CV for adiponectin was

3.7%. The range of detectability for the adiponectin assay is 2.8–178 ng/mL. The data

were normalized to historical values using a pooled plasma quality control sample that

was assayed in each plate.

Blood glucose and lipid measurements

Blood was collected for glucose measurements by tail vein nick in 4-h or 6-h fasted

rats and measured with a glucometer using the AlphaTRAK 2 blood glucose monitoring

system (Abbott Laboratories, Abbott Park, IL) [28; 65]. Total cholesterol (TC) [Fisher

Diagnostics (Middletown, VA)] and free fatty acids (FFAs) [Wako Chemicals USA, Inc.,

Richmond, VA)] were measured using an enzymatic-based kits. Intra-assay CVs for TC

and FFAs were 1.4 and 2.3%, respectively. These assay procedures have been validated

for rodents [66].

Adipose tissue processing for adipocyte size

Adipose tissue depots were collected at the end of the infusion period in DIO rats from

Studies 6A–C (IBAT, EWAT and IWAT). IBAT, IWAT, and EWAT depots were dissected

and placed in 4% paraformaldehyde-PBS for 24 h and then placed in 70% ethanol (EtOH)

prior to paraffin embedding. Sections (5 μm-thick) were obtained using a rotary

microtome, slide-mounted using a floatation water bath (37°C) and baked for 30 min at

60°C to give approximately 15 or 16 slides/fat depot with two sections/slide.

Adipocyte size analysis

Adipocyte size analysis was performed on deparaffinized and digitized IWAT and EWAT

sections. The average cell area from two randomized photomicrographs was determined

using the built-in particle counting method of ImageJ software (National Institutes of

Health, Bethesda, MD). Slides were visualized using bright field on an Olympus BX51

microscope (Olympus Corporation of the Americas; Center Valley, PA) and photographed

using a Canon EOS 5D SR DSLR (Canon U.S.A., Inc., Melville, NY) camera at ×100

magnification. Values for each tissue within a treatment were averaged to obtain the

mean of the treatment group.

Tissue collection for quantitative real-time PCR (qPCR)

Tissue (IBAT, IWAT and EWAT) was collected from a subset of 4-h (**Study 6**) or 6-h fasted rats (**Study 4**). IBAT, IWAT, and EWAT were collected within a 2-h window towards the start of the light cycle (10:00 a.m.–12:00 p.m.) as previously described in DIO CD® IGS/Long-Evans rats and C57BL/6J mice [28; 43; 52]. Tissue was rapidly removed, wrapped in foil and frozen in liquid N_2. Samples were stored frozen at −80°C until analysis.

qPCR

RNA extracted from samples of IBAT and IWAT (**Studies 4–7**) were analyzed using the RNeasy Lipid Mini Kit (Qiagen Sciences Inc, Germantown, MD) followed by reverse transcription into cDNA using a high-capacity cDNA archive kit (Applied Biosystems, Foster City, CA). Quantitative analysis for relative levels of mRNA in the RNA extracts was measured in duplicate by qPCR on an Applied Biosystems 7500 Real-Time PCR system (Thermo Fisher Scientific, Waltham, MA) and normalized to the cycle threshold value of Nono mRNA in each sample. The TaqMan® probes used in the study were Thermo Fisher Scientific Gene Expression Assay probes. The probe for rat *Nono* (Rn01418995_g1), uncoupling protein-1 (UCP-1) (*Ucp1*; catalog no. Rn00562126_m1), uncoupling protein-2 (UCP-2) (*Ucp2*; catalog no. Mm00627599_m1), uncoupling protein-3 (UCP-3) (*Ucp3*; catalog no. Rn00565874_m1), β1-adrenergic receptor (β1-AR) (*Adrb1*; catalog no. Rn00824536_s1), β2-adrenergic receptor (β2-AR) (*Adrb2*; catalog no. Rn005600650_s1), β3-adrenergic receptor (β3-AR) (*Adrb3*; catalog no. Rn01478698_g1), alpha 2-adrenergic receptor (alpha 2-AR) (*Adra2a*; catalog no. Rn00562488_s1)], type 2 deiodinase (D2) (*Dio2*; catalog no. Rn00581867_m1), PR

domain containing 16 (*Prdm16*; catalog no. Rn01516224_m1), cytochrome c oxidase

subunit 8b (*Cox8b*; catalog no. Rn00562884_m1), G-protein coupled receptor 120

(*Gpr120*; catalog no. Rn01759772_m1), bone morphogenetic protein 8b (*bmp8b*; catalog

no. Rn01516089_gH), cell death-inducing DNA fragmentation factor alpha-like effector A

(*Cidea*; catalog no. Rn04181355_m1), peroxisome proliferator-activated receptor gamma

coactivator 1 alpha (*Ppargc1a*; catalog no. Rn00580241_m1) and cannabinoid receptor

1(*Cnr1*; catalog no. Mm01212171_s1) were acquired from Thermo Fisher Scientific.

Relative amounts of target mRNA were determined using the Comparative C_T or $2^{-\Delta\Delta C_T}$

method [67] following adjustment for the housekeeping gene, Nono. Specific mRNA

levels of all genes of interest were normalized to the cycle threshold value of *Nono* mRNA

in each sample and expressed as changes normalized to controls (vehicle/sham

treatment).

Statistical Analyses

All results are expressed as means ± SE. Comparisons between multiple groups involving

between-subjects designs were made using one-way ANOVA, as appropriate, followed

by a post-hoc Fisher's least significant difference test. Comparisons involving within-

subjects designs were made using a one-way repeated-measures ANOVA followed by a

post-hoc Fisher's least significant difference test. Analyses were performed using the

statistical program SYSTAT (Systat Software, Point Richmond, CA). Differences were

considered significant at $P < 0.05$, 2-tailed.

Results

Study 1A: Determine the success of the surgical denervation of only the superficial nerves at 1-week post-sham or denervation in lean rats by measuring NE content.

The goal of this study was to determine the success of lesioning only the more superficial intercostal nerves that innervate IBAT in lean rats by confirming a reduction of NE content that was specific to IBAT relative to other tissues (IWAT and liver). Lesions of only the more superficial nerves resulted in a tendency for IBAT NE content to be reduced in denervated rats relative to sham-operated control rats (80.3 ± 6.0%; P=0.118) (data not shown). NE content remained unchanged in IWAT or liver from denervated rats relative to sham rats (P=NS). There were also no significant differences in BWs between sham and denervation groups at 1 -week post-denervation or sham surgery (P=NS; data not shown).

Study 1B: Determine the success of the surgical denervation procedure at 1-week post-sham or denervation in lean rats by measuring NE content.

The goal of this study was to verify the success of the SNS denervation of both superficial and more ventrally located intercostal nerves that innervate IBAT by confirming a reduction of NE content that was specific to IBAT relative to other tissues (IWAT, liver and pancreas). IBAT NE content was reduced in denervated rats by 91.5±5.4% in denervated rats relative to sham-operated control rats [(F(1,8) = 57.941, P=0.000) (**Figure 1**). In contrast, NE content was unchanged in IWAT, liver or pancreas in denervated rats relative to sham rats (P=NS).

There were no significant differences in BWs between sham and denervation groups at

1, 6 or 7-weeks post-denervation or sham surgery (P=NS; data not shown).

Study 2: Determine the success of the surgical denervation procedure at 1, 6 and

7-weeks post-sham or denervation in DIO rats by measuring NE content.

The goal of this study was to 1) verify the success of the SNS denervation procedure in

DIO rats by confirming a reduction of NE content that was specific to IBAT relative to

other tissues (IWAT, liver and pancreas) and 2) confirm that these changes would persist

for extended periods of time out to 7 weeks. By design, DIO rats were obese as

determined by both BW (695 ± 13.9 g) and adiposity (230.1 ± 10.1 g fat mass; 32.7 ±

0.9% adiposity) after maintenance on the HFD for approximately 5 months prior to

sham/denervation procedures. Sham and denervation groups were matched for BW, fat

mass and lean mass such that there was no difference in baseline BW, lean mass or fat

mass between groups prior to surgery (P=NS).

IBAT NE content was reduced by 94.7 ± 2.7, 96.8 ± 1.8 and 85.9±6.1% at 1- [($F_{(1,8)}$ =

123.847, P=0.000)], 6- [($F_{(1,8)}$ = 28.121, P=0.001)] and 7-weeks [($F_{(1,8)}$ = 17.081,

P<0.0003)] post-denervation (**Figure 2**) relative to IBAT NE content from sham operated

rats. In contrast, NE content was unchanged in IWAT or liver in denervated rats relative

to sham rats (P=NS) at 1, 6 or 7-weeks post-denervation. Similarly, there was also

difference in NE content in pancreas at 6 or 7-weeks post-denervation relative to sham

rats. However, there was a reduction of NE content in pancreas at 1-week post-

denervation relative to sham rats (P<0.05).

There were no significant differences in BWs between sham and denervation groups at

1, 6 or 7-week post-surgery (P=NS; data not shown). Similarly, there was no difference

in fat mass or lean mass between groups at 6 or 7-weeks post-denervation or sham

surgeries (P=NS; data not shown). These findings are in agreement with others who have

reported no difference in BW [68; 69; 70; 71; 72] or fat mass [69; 71] in hamsters or mice

following bilateral surgical or chemical denervation of IBAT.

Study 3:　Determine if surgical denervation of IBAT changes the ability of the β3-

AR agonist, CL 316243, to increase T_{IBAT} in DIO rats

The goal of this study was to confirm there was no functional defect in the ability of IBAT

to respond to direct β3-AR stimulation because of the denervation procedure relative to

sham operated animals. By design, DIO rats were obese as determined by both BW

(684.1 ± 11.5 g) and adiposity (226.1 ± 8.1 g fat mass; 33 ± 0.8% adiposity) after

maintenance on the HFD for approximately four months prior to sham/denervation

procedures. IBAT NE content was reduced in denervated rats by 87.5 ± 3.2% in

denervated rats relative to sham-operated control rats [(F(1,12) = 67.209, P=0.000). In

contrast, NE content was unchanged in IWAT, EWAT, liver or pancreas in denervated

rats relative to sham rats (P=NS). There was no significant difference in BW between

sham and denervation groups at the end of the study (P=NS; data not shown).

In sham rats, CL 316243 (1 mg/kg) increased T_{IBAT} at 0.25, 0.5, 0.75, 1, 1.25, 1.5, 1.75,

2, 3 and 4-h post-injection. The lowest dose (0.1 mg/kg) also stimulated T_{IBAT} at 0.25, 0.5,

0.75, 1, 1.25, 1.5, 1.75, 2, 3 and 4-h post-injection. CL 316243 also stimulated T_{IBAT} at 24-

h post-injection at both doses ($P<0.05$; **Figure 3A**). Similar findings were apparent when

measuring change in T_{IBAT} relative to baseline T_{IBAT} (**Figure 3B**).

Similarly, in denervated rats, CL 316243 (1 mg/kg) increased T_{IBAT} at 0.25, 0.5, 0.75, 1,

1.25, 1.5, 1.75, 2, 3 and 4-h post-injection. The lowest dose (0.1 mg/kg) also stimulated

T_{IBAT} at 0.25, 0.5, 0.75, 1, 1.25, 1.5, 1.75, 2, 3 and 4-h post-injection. CL 316243 also

stimulated T_{IBAT} at 24-h post-injection at the high dose ($P<0.05$) and tended to stimulate

24-h T_{IBAT} at the low dose as well ($P=0.152$; **Figure 3D**). Similar findings were apparent

when measuring change in T_{IBAT} relative to baseline T_{IBAT} (**Figure 3E**). Importantly, there

was no difference in the T_{IBAT} response to CL 316243 (0.1 or 1 mg/kg) when the data

were averaged over the 1-h or 4-h post-injection period between sham and denervated

rats (P=NS). Overall, these findings indicate that IBAT denervation did not result in a

change in the ability of CL 316243 to increase BAT thermogenesis (surrogate measure

of EE) in DIO mice relative to sham operated rats.

Energy intake

In sham rats, CL 316243 reduced daily energy intake at both 0.1 and 1 mg/kg by 49 and

58.3% (P<0.05). Similarly, in denervated rats, CL 316243 also reduced daily energy

intake at both 0.1 and 1 mg/kg ($P<0.05$) by 38 and 64.5% relative to vehicle (**Figure 3C**).

Body Weight (BW)

While CL 316243 did not impact BW in either group the high dose reduced BW gain in

the denervated rats ($P<0.05$; **Figure 3F**).

Importantly, there was no difference in the effectiveness of CL 316243 (0.1 or 1 mg/kg)

to reduce food intake or weight gain between sham and denervated rats (*P*=NS). Overall,

these findings indicate that IBAT denervation did not result in a significant change in the

ability of CL 316243 to reduce food intake or BW gain in DIO rats relative to sham

operated rats.

Study 4: Determine the extent to which OT-induced activation of sympathetic

outflow to IBAT contributes to its ability to increase T_{IBAT} in DIO rats.

After having confirmed there was no functional defect in the ability of IBAT to respond to

direct β3-AR stimulation (**Study 3**), the goal of this study was to determine if OT-elicited

elevation of T_{IBAT} requires intact SNS outflow to IBAT. DIO rats from **Study 3** were

subsequently used in this study. Animals weighed approximately 631.6±10.3 g at the start

of the studies.

In sham rats, OT (5 μg) increased T_{IBAT} at 0.75, 1, 1.25, 1.5, and 1.75-h post-injection.

The high dose also tended to stimulate T_{IBAT} at 0.25 (0.05<*P*<0.1) and 2-h (*P*=0.052) post-

injection. The lowest dose (1 μg) also stimulated T_{IBAT} at 1.75-h post-injection and tended

to stimulate T_{IBAT} at 0.75-h post-injection (0.05<*P*<0.1; **Figure 4A**). Similar findings were

apparent when measuring change in T_{IBAT} relative to baseline T_{IBAT} (**Figure 4B**).

In denervated rats, OT (5 μg) increased T_{IBAT} at 1.25-h post-injection (*P*<0.05) and also

tended to stimulate T_{IBAT} at 0.75 and 1-h post-injection (0.05<*P*<0.1). The lowest dose (1

μg) tended to stimulate T_{IBAT} at 1.25-h post-injection (0.05<*P*<0.1; **Figure 4C**). Similar

findings were apparent when measuring change in T_{IBAT} relative to baseline T_{IBAT} (**Figure**

4D) as well as when examining these effects in lean rats (**Supplemental Figure 1**).

Importantly, there was no difference in the T_{IBAT} response to 4V OT (5 μg) at 1.25-h post-

injection between sham and denervated rats (*P*=NS). Overall, these findings indicate that

IBAT denervation did not result in a significant change in the ability of 4V OT to increase

BAT thermogenesis in denervated rats relative to sham operated rats.

Study 5: Determine the extent to which OT-induced activation of sympathetic

outflow to IBAT requires activation of β3-AR to increase T_{IBAT} in DIO rats.

The goal of this study was to determine whether hindbrain administration of OT required

activation of β3-AR to stimulate BAT thermogenesis. We initially identified a dose of the

β3-AR antagonist, SR 59230A, that was sufficient to block the effects of the β3-AR

agonist, CL316243, to stimulate T_{IBAT} at 0.5-h and 0.75-h post-CL316243 treatment

(P=NS vehicle vs CL316243; **Supplemental Figure 2**). This dose was then used to

address whether 4V OT stimulated T_{IBAT} through a β3-AR driven mechanism.

4V OT (in the absence of the β3-AR antagonist) increased T_{IBAT} at 0.75, 1, 1.25, 1.75, 2,

and 4-h post-injection (*P*<0.05) and tended to stimulate T_{IBAT} at 90, and 105-min post-

injection (0.05<*P*<0.1). Following β3-AR antagonist pre-treatment, 4V OT stimulated T_{IBAT}

at 1.25-h post-injection (*P*=0.033) and tended to stimulate T_{IBAT} at 0.75 (*P*=0.071), 1

(*P*=0.084) and 1.5-h (*P*=0.058) post-injection (**Figure 5**). 4V OT also appeared to

stimulate T_{IBAT} at 1.75 (*P*=0.118) and 2-h post-injection (*P*=0.127).

We also found that 4V OT (5 μg) was effective (n=11; data not shown) at stimulating T_{IBAT}

in lean rats even in the presence of the β3-AR antagonist, SR 59230A, that was given at

2- (1 mg/kg) and 10-fold higher (5 mg/kg) doses approximately 12 minutes prior to 4V OT

(n=8; data not shown). Note that in contrast to the study design in the DIO rats, the timing

of the drug treatments in the lean rats occurred toward the end of the light cycle.

Overall, these findings suggest that OT does not require activation of β3-AR to stimulate

BAT thermogenesis.

Study 6A: Determine the extent to which OT-induced activation of sympathetic

outflow to IBAT contributes to its ability to impact BW in DIO rats.

The goal of this study was to determine if OT-elicited weight loss requires intact SNS

outflow to IBAT. By design, DIO rats were obese as determined by both BW (679.6 ± 9.2

g) and adiposity (209.8 ± 6.4 g fat mass; 30.7 ± 0.7% adiposity) after maintenance on the

HFD for at least 4.5 months prior to sham/denervation procedures. One subset (N=4) was

maintained on the HFD for approximately 6 months prior to sham/denervation procedures

and animals were divided equally among treatment groups. IBAT NE content was

reduced in denervated rats by 90.2 ± 2.6% in denervated rats relative to sham-operated

control rats [(F(1,35) = 47.925, P=0.000)]. In contrast, NE content was unchanged in

EWAT, liver or pancreas in denervated rats relative to sham rats (P=NS). In contrast,

there was an increase in NE content in IWAT in denervated rats relative to sham rats

[(F(1,35) = 6.421, P=0.016)]. There was no significant difference in BW between sham

and denervation groups at the end of the study (P=NS; data not shown).

As expected, in sham rats, 4V vehicle resulted in 5.9±1.9% weight gain relative to vehicle

pre-treatment [(F(1,9) = 9.610, *P*=0.013)]. In contrast, 4V OT failed to reduce BW relative

to OT pre-treatment (*P*=NS) (**Figure 6A**) but it reduced weight gain (**Figure 6B**)

throughout the 29-day infusion period. OT treatment reduced weight gain on day 3 and

7-29 (*P*<0.05) and tended to reduce weight gain on day 6 (*P*=0.088). By the end of the

infusion period (infusion day 29), OT had reduced weight gain by -7.2 ± 9.6 g relative to

vehicle treated animals 49.1 ± 10.0 g (*P*<0.05). OT reduced relative fat mass (pre- vs

post-intervention) (**Figure 6C**; *P*<0.05), fat mass and relative lean mass (pre- vs post-

intervention) but had no effect on total lean body mass (*P*=NS). These effects that were

mediated, at least in part, by a modest reduction of energy intake that was evident during

weeks 2 and 4 of the treatment period (**Figure 6D**; *P*<0.05). OT also tended to reduce

energy intake at week 3 of the treatment period (*P*=0.061).

In denervated rats, 4V vehicle resulted in an expected 5.9±1.9% weight gain relative to

vehicle pre-treatment [(F(1,7) = 9.039, *P*=0.020)]. In contrast, 4V OT tended to BW by 2.9

± 1.4% relative to OT pre-treatment [(F(1,8) = 3.166, *P*=0.113)]. (**Figure 56**) and reduced

weight gain (**Figure 6B**) throughout the 29-day infusion period. OT treatment reduced

weight gain on days 3–29 (*P*<0.05). By the end of the infusion period (infusion day 29),

OT had reduced weight gain by −14.1 ± 8.8 g relative to vehicle treated animals 44 ± 13.4

g (*P*<0.05). OT reduced relative fat mass (pre- vs post-intervention) (**Figure 6C**; *P*<0.05)

and fat mass but had no effect on total lean body mass (*P*=NS). These effects that were

mediated, at least in part, by a modest reduction of energy intake that persisted during

weeks 2 and 3 of the treatment period (**Figure 6D**; *P*<0.05). OT also tended to reduce

energy intake during week 1 of the treatment period (P=0.091). Similar effects were also

observed in rats with more pronounced obesity (**Supplemental Figure 3**).

Importantly, there was no difference in the effectiveness of 4V OT to reduce weight gain

at the end of the infusion period (day 29), energy intake, and relative fat mass or fat mass

between sham and denervated rats (P=NS). Based on these collective findings, we

conclude that SNS innervation of IBAT is not a predominant contributor of OT-elicited

reduction of weight gain and adiposity.

Adipocyte size

There was a near significant effect of denervation to reduce adipocyte size in IWAT

(P=0.093) but no effect of OT to significantly impact IWAT adipocyte size in either the

sham or denervation group (P=NS) (**Figure 7A**). There was also a significant effect of

denervation to reduce adipocyte size in EWAT (P=0.089; **Figure 7B**). In addition, there

was a near significant effect of 4V OT to reduce adipocyte size in sham (P=0.141) and

denervated rats (P=0.089) (**Figure 7B**).

Plasma hormone concentrations

To characterize the endocrine and metabolic effects between sham and denervated DIO

rats, we measured blood glucose levels and plasma concentrations of leptin, insulin, FGF-

21, irisin, adiponectin, TC, and FFAs at various times post-sham and denervation

procedure. At 1-week post-denervation/sham, there was a significant reduction of blood

glucose in the denervation group relative to the sham group (P<0.05). In contrast, there

were no differences in any of the other metabolic measures between respective sham

and denervated animals at 1, 6 or 7-weeks post-sham or denervation surgery (**Table 1**).

We also characterized the endocrine and metabolic effects of 4V OT (16 nmol/day) in

sham and denervated DIO rats. We found OT treatment was associated with a reduction

of plasma leptin in the sham group [(F(1,19) = 7.903, P=0.011)] which coincided with OT-

elicited reductions in fat mass. In addition, OT treatment was associated with a significant

reduction of plasma adiponectin in the denervation group [(F(1,15) = 5.284, P=0.036)]

(**Table 2**).

Study 6B: Determine the success of the surgical denervation procedure at 6-8

weeks post-sham or denervation in DIO rats by measuring thermogenic gene

expression.

The goal of this study was to 1) confirm the success of the SNS denervation of IBAT

procedure in DIO rats using an independent and complementary marker to IBAT NE

content (IBAT thermogenic gene expression), 2) verify that these changes would persist

out to more prolonged periods of time (6–8 weeks) and 3) determine if changes in

thermogenic gene expression in EWAT occur in response to IBAT denervation.

IBAT:

There was a significant reduction of IBAT Dio2 mRNA expression (P=0.016) in

denervated rats relative to IBAT from sham operated rats (P<0.05; **Table 3**). In addition,

there tended to be a near significant reduction of IBAT UCP-1 (P=0.057) and Prdm16

mRNA (P=0.052) expression from denervated rats relative to sham operated rats (**Table**
3).

EWAT:

There was an increase of EWAT UCP-1 (P=0.024) and β3-AR (P=0.032) mRNA
expression in denervated rats compared to sham rats. In addition, there was a near
significant reduction of IBAT β2-AR (P=0.073) and Prdm16 (P=0.078) mRNA expression
in denervated rats compared to sham operated rats (**Table 3**).

Collectively, these findings indicate that IBAT denervation results in significant (*Dio2*) or
near significant reductions (*Ucp1* and *Prdm16*) of thermogenic gene markers in IBAT
similar to what we recently reported following IBAT denervation in mice [29]. In addition,
these findings indicate that IBAT denervation results in significant increases in the
thermogenic markers, *Ucp1* and *Adrb3* in EWAT (indicative of browning/beiging of WAT).

Study 6C: Determine the extent to which IBAT denervation contributes to the ability
of OT to impact thermogenic gene expression in IBAT and EWAT in DIO rats.
The goal of this study was to determine 1) if 4V OT elicits changes in thermogenic gene
expression in IBAT and EWAT in sham operated rats, and 2) if 4V OT elicits changes in
thermogenic gene expression in IBAT and EWAT in IBAT denervated rats.

IBAT:

We found that chronic 4V OT treatment elicited a significant increase in the expression of

IBAT Cox8b (P=0.004) and β1-AR mRNA expression (P=0.004) but this effect was

blocked in denervated rats. 4V OT also elicited a near significant increase in IBAT Gpr120

(P=0.063), Cidea (P=0.075) and CN1 (P=0.069) mRNA expression and a reduction of

IBAT Prdm16 mRNA expression (P=0.068) in sham operated rats. There was also a

significant effect of 4V OT to increase IBAT Dio2 (P=0.041) and Bmp8b (P=0.019) mRNA

in denervated rats.

In addition, 4V OT produced significant (*Cox8b*, *Adrb1*) or near significant increases

(*Gpr120*, *Cidea* and *CN1*) in thermogenic markers in IBAT in sham rats, some of which

were blocked in response to denervation of IBAT (*Cox8b*, *Adrb1*). In contrast, 4V OT

increased mRNA expression of Dio2 and Bmp8b in denervated rats but not in sham rats

raising the possibility that activation of these thermogenic markers might contribute to the

effects of 4V OT on BAT thermogenesis in denervated animals.

EWAT:

4V OT elicited a significant increase in EWAT β3-AR (P=0.018) and Gpr120 (P=0.03)

mRNA expression in sham rats while it also reduced EWAT CN1 mRNA expression

(P=0.051) in sham rats. In addition, 4V OT also increased EWAT UCP-3 (P=0.042) mRNA

expression in denervated rats.

4V OT resulted in a significant increase (*Adrb3, Gpr120*) or decrease (*CN1*) of

thermogenic markers in EWAT of sham operated rats while it also increased the

thermogenic marker (*Ucp3*) in denervated rats. These findings raise the possibility that

different thermogenic markers in EWAT may contribute, in part, to the metabolic effects

of 4V OT in sham (*Adrb3, Gpr120, CN1*) and denervated rats (*Ucp3*).

Study 7A: Determine the extent to which 4V OT increases gross motor activity,

core temperature and T$_{IBAT}$ in DIO rats.

The goal of this study was to determine if hindbrain administration of OT increases gross

motor activity at doses that also increase BAT thermogenesis.

Gross motor activity: 4V OT (5 μg/μL) increased activity at 1.5-h post-injection (*P*<0.05)

and tended to stimulate activity at 0.5, 0.75-h post-injection (0.05<*P*<0.1) (**Figure 8A**).

The lower dose of OT (1 μg/μL) also tended to increase activity at 0.75-h post-injection

(0.05<*P*<0.1). 4V OT (5 μg/μL) also reduced activity at 4-h post-injection. However, 4V

OT (5 μg/μL) also appeared to elicit a modest increase in gross motor activity when the

data were averaged over the 1 (*P*=0.118) or 2-h (*P*=0.172) post-injection period but these

effects were not significant (data not shown).

Core temperature: 4V OT (5 μg/μL) increased core temperature at 0.75, 1, 1.5, 1.75, 2,

and 3-h post-injection (*P*<0.05) (**Figure 8B**) and tended to increase core temperature at

1.25-h post-injection (0.05<*P*<0.1). The lower dose of OT (1 μg/μL) also increased core

temperature at 2-h post-injection (*P*<0.05) and tended to increase core temperature at

1.75-h post-injection ($0.05 < P < 0.1$). While there was no effect of 4V to increase core

temperature when averaged over 1-h post-injection period ($CT_{1-hour\ AVE}$), there tended to

be an overall effect of 4V OT to increase core temperature when the data were averaged

over the 2-h post-injection period ($CT_{2-hour\ AVE}$) [($F(2,12) = 3.779$, $P=0.053$)] (data not

shown). Specifically, the high dose of OT (5 μg/μL) was able to stimulate

$CT_{2-hour\ AVE}$ ($P<0.05$) while the lower dose was ineffective (P=NS).

T_{IBAT}: 4V OT (5 μg/μL) increased T_{IBAT} at 0.5, 0.75, 1, 1.25, 1.5, 1.75, and 2-h post-

injection ($P<0.05$) (**Figure 8C**). The lower dose of OT (1 μg/μL) also increased T_{IBAT} at

1.75 and 2-h post-injection ($P<0.05$) and tended to increase T_{IBAT} at 0.75-h post-injection

($0.05 < P < 0.1$). There also tended to be an overall effect of 4V OT to increase T_{IBAT} when

the data were averaged over the 1 ($T_{IBAT\ 1-hour\ AVE}$) [($F(2,12) = 3.513$, $P=0.063$)] and 2-h

post-injection ($T_{IBAT\ 2-hour\ AVE}$) period [($F(2,12) = 8.815$, $P=0.006$)] (data not shown).

Specifically, the high dose of OT (5 μg/μL) was able to stimulate $T_{IBAT\ 1-hour\ AVE}$ and $T_{IBAT\ 2-hour\ AVE}$

($P<0.05$) while the lower dose had no significant effect (P=NS).

Energy intake:

Animals used in the telemetry studies were used in these studies. The goal was to confirm

that 4V OT reduced energy intake at doses that impacted BAT thermogenesis and activity

in DIO rats. There was an overall effect of OT dose to reduce energy intake at 0.5-h post-

injection [($F(2,12) = 7.867$, $P=0.007$)]. 4V administration of OT at the low dose (1 μg)

tended to suppress energy intake ($P=0.056$) while the high dose (5 μg) reduced energy

intake by 78.3±5.4% ($P<0.05$) (**Figure 8D**). 4V administration of OT at the high dose also

tended to suppress energy intake at 1-h post-injection ($P=0.053$).

Overall, these findings indicate that 4V OT increases gross motor activity at doses that

stimulate T_{IBAT}, core temperature and energy intake. Furthermore, 4V OT appeared to

increase T_{IBAT} prior to core temperature suggesting that OT-elicited changes in T_{IBAT} may

also contribute to OT-driven changes in core temperature.

1003

Study 7B: Determine the extent to which 4V OT activates an early marker of

neuronal activation, pERK1/2, within the hindbrain NTS in DIO rats.

The goal of this study was to determine if 4V OT treatment is associated with activation

of pERK1/2 within hindbrain NTS neurons. Consistent with previous work reporting

elevations of pERK1/2-immunoreactive cells at Levels 65–70 in the *Brain Maps 4.0* rat

brain atlas [56] in the rat NTS in response to glycemic challenge [73; 74], we identified

such elevations in the NTS that mapped to Levels 69 and 70 at the level of the *area*

postrema (>1840) (Level 70 data shown in **Figure 9A**). Fifteen minutes after receiving

OT into the 4th ventricle, the NTS and DMX display elevated numbers of cells

immunoreactive for pERK1/2 (**Figure 9C**). Specifically, phospho-ERK1/2-

immunoreactivity (-ir) was observed in fusiform-shaped cells in the NTS, where it also

marked slender neurites including what appeared to be axons (*arrowheads* in **Figure 9C**).

While further confirmation is needed to determine if these axonal fibers include vagal

afferents, pERK1/2 localization to NTS axons is consistent with previous work that

identified NTS vagal afferents displaying pERK1/2-ir in response to CCK treatment [59].

The perikaryal NTS labeling we observed appeared to fall within the boundaries of what

Swanson has defined as the *caudal subzone (>1840)* of the medial NTS (NTSmc; see

Methods for explanation of Swanson's nomenclature of these regions), although Nissl

material from an adjacent tissue series was not always available to help distinguish the

boundary zone between the *commissural part (>1840)* (NTSco) and NTSmc at this level.

In the DMX, immunoreactivity was observed mainly along the ventral margin of the

nucleus in larger, goblet-shaped cells and in smaller, faintly staining triangular-shaped

cells (**Figure 9C**). In contrast, control subjects receiving only vehicle solution did not

display pERK1/2 immunoreactivity in either brain region (**Figure 9B**). Thus, as with other

reported stimuli, 4V OT is also associated with rapid activation of NTS neurons.

1029

Discussion

The goal of the current studies was to determine if sympathetic innervation of IBAT is

required for OT to increase BAT thermogenesis and reduce BW and adiposity in male

DIO rats. To assess if OT-elicited increases in BAT thermogenesis require intact SNS

outflow to IBAT, we examined the effects of acute 4V OT (1, 5 μg) on T_{IBAT} in DIO rats

following sham or bilateral surgical SNS denervation to IBAT. We found that the high dose

of 4V OT (5 μg ≈ 4.96 nmol) stimulated T_{IBAT} similarly between sham rats and denervated

rats and that the effects of 4V OT to stimulate T_{IBAT} did not require β3-AR signaling. We

subsequently determined if OT-elicited reductions of BW and adiposity require intact SNS

outflow to IBAT. To accomplish this, we determined the effect of bilateral surgical or sham

denervation of IBAT on the ability of chronic 4V OT (16 nmol/day) or vehicle administration

to reduce BW, adiposity and food intake in DIO rats. We found that chronic 4V OT

produced comparable reductions of weight gain, adiposity, leptin levels and energy intake

in sham and denervated rats. Lastly, we found that increased activity is not likely to have

contributed to the effect of 4V OT to increase both T_{IBAT} and core temperature. 4V OT

appeared to increase T_{IBAT} prior to core temperature suggesting that OT-induced

elevations in T_{IBAT} may also contribute to OT-induced elevations in core temperature.

Collectively, our findings support the hypothesis that sympathetic innervation of IBAT is

not a predominant mediator of 4V OT-elicited BAT thermogenesis and reductions of BW

and adiposity in male DIO rats.

Our finding showing that the effect of chronic 4V OT to elicit weight loss does not

require SNS innervation of IBAT is similar to what we have demonstrated in the mouse

model [29]. These findings suggest that OT stimulates IBAT and elicits weight loss

through a mechanism that does not require SNS activation of IBAT across rodent species.

One outstanding question is how 4V OT activates IBAT if not through SNS outflow to

IBAT and signaling through the β3-AR. We found that systemic OT, at a dose that was

effective when given into the 4V was ineffective at recapitulating the effects of 4V OT on

weight loss and T_{IBAT} in DIO mice [29], suggesting that these effects are not likely a result

of 4V OT acting at OTRs in the periphery. One possibility is that 4V OT-elicited stimulation

of hindbrain and/or spinal cord OTRs results in the release of epinephrine from the

adrenal gland (adrenal medulla) and subsequent stimulation of IBAT. It is possible that

epinephrine could be acting through the β1-AR or β2-AR, as both receptor subtypes are

expressed in IBAT in both mice [75] and rats [76; 77] (for review see [78]). Others have

shown that the β1-AR is important in the control of thermogenesis in rats [79] and mice

[80] with the β2-AR receptor potentially playing a more minor role [79] in rodents

compared to humans [81; 82; 83], respectively. In addition, L-epinephrine has equal

affinity for both β1-AR and β2-AR in CHO cells [84] and it has been shown that

epinephrine application to brown adipocytes derived from rat IBAT stimulates respiration

and release of fatty acids [85]. While mice that are deficient in epinephrine are still capable

of maintaining body temperature in response to cold stress, they are unable to show an

elevation of PGC1-alpha or UCP-1 within IBAT, suggesting that epinephrine may be

important in mitochondrial uncoupling in IBAT [86]. However, while the hindbrain and

spinal cord are a component of a multi-synaptic projection to the adrenal gland [87; 88],

only 1% of PVN OT neurons within either the parvocellular or magnocellular PVN were

found to have multi-synaptic projections to the adrenal gland [87]. Future studies that

examine the effects of 4V OT on BAT thermogenesis and weight loss in adrenal

demedullated rats will be helpful in being able to address the extent to which OT-elicited

stimulation of epinephrine from adrenal gland is required for these effects.

We found that IBAT denervation resulted in a significant or near significant reduction

of the thermogenic genes, *Dio2*, *UCP-1* and *Prdm16*, from IBAT in DIO rats. These

findings are consistent with what others have reported with IBAT UCP-1 protein

expression from hamsters [72] and mice [71] following chemical (6-OHDA)-induced

denervation of IBAT relative to control animals. Similarly, we and others found a reduction

of UCP-1 mRNA expression in IBAT following unilateral or bilateral surgical denervation

in mice [29; 68; 69] and hamsters [89], respectively. Our finding showing a reduction of

IBAT *Dio2* following IBAT denervation in rats is also consistent with previous findings in

mice from our lab and with those of Fischer and colleagues [29; 68].

We also found that chronic 4V OT treatment produced a significant increase of mRNA

expression of the thermogenic genes, *Cox8b* and *Adrb1*, in IBAT of sham rats. In addition,

4V OT also tended to increase mRNA expression of other thermogenic markers (*Gpr120*,

Cidea and *CN1*) in IBAT in sham rats, some of which were blocked in response to

denervation of IBAT (*Cox8b*, *Adrb1*). In contrast, 4V OT increased expression of both

thermogenic markers, *Dio2* and *Bmp8b*, in denervated rats but not in sham rats. These

findings raise the possibility that activation of these thermogenic markers might contribute

to the effects of 4V OT on BAT thermogenesis in denervated animals.

In addition to changes in IBAT gene expression in response to IBAT denervation, we

also found that IBAT denervation was associated with browning of WAT. IBAT

denervation increased mRNA expression of the thermogenic genes, *Adrb3* and *UCP-1*,

in EWAT (indicative of browning/beiging of WAT). However, our finding showing an

elevation of EWAT UCP-1 in response to IBAT denervation in rats is in contrast to the

findings from Nguyen, who failed to find any change in UCP-1 protein in EWAT between

sham-operated and IBAT-denervated hamsters [72]. These different outcomes may

potentially be explained by species differences. While UCP-1 mRNA expression from

IWAT was not examined in our study, the findings pertaining to EWAT UCP-1 gene

expression in surgically denervated rats are consistent with findings from other white

adipose tissues, namely IWAT, which show an elevation of IWAT UCP-1 protein in both

hamsters [72] and mice [71] following chemical (6-OHDA)-induced denervation of IBAT

relative to control animals. Another study found what appeared to be a marked increase

in browning of IWAT (as indicated by UCP-1 immunohistochemistry) following bilateral

surgical denervation of IBAT in mice but the data were not quantified [70]. In addition,

Nguyen also found that IBAT denervation resulted in an elevation of IWAT temperature

(functional readout of WAT thermogenesis) in response to cold stress relative to sham

animals [72]. 4V OT resulted in increased expression of *Adrb3* and *Gpr120* as well as

decreased expression of CN1 in EWAT of sham-operated rats while it also increased

UCP-3 mRNA expression in denervated rats. It remains to be determined whether

different thermogenic markers in EWAT contribute, in part, to the metabolic effects of 4V

OT in sham (*Adrb3*, *Gpr120*, *CN1*) and denervated rats (*UCP-3*).

One of the remaining questions from this study is whether the changes in

thermogenic gene expression in other WAT depots in response to IBAT denervation may

impact the effectiveness of OT to stimulate BAT thermogenesis and elicit weight loss.

Previous findings have identified both distinct and overlapping CNS circuits that regulate

SNS outflow to IBAT and IWAT [72]. Namely, OT neurons within the parvocellular PVN

send multi-synaptic projections to IBAT [90; 91], EWAT [92; 93] and IWAT [90; 92]. A

small subset of OT neurons within the parvocellular PVN also overlap and project to both

IBAT and IWAT [90]. Thus, PVN OT neurons are anatomically positioned to impact SNS

outflow to IBAT and IWAT and increase both BAT and WAT thermogenesis, respectively.

It is not clear if these effects are mediated by the same OT neurons or through distinct

OT neurons that project to the hindbrain nucleus tractus solitarius (NTS) [94; 95] and/or

spinal cord [95], both of which are areas that can alter SNS outflow to IBAT and BAT

thermogenesis [96; 97]. Recent findings from Ong reported that viral-elicited knockdown

of OTR mRNA within the hindbrain dorsal vagal complex (DVC) failed to block the effects

of 4V OT to stimulate core temperature [98], suggesting that OTRs within other hindbrain

sites and/or spinal cord may be important in mediating these effects in rats. Future studies

should examine the extent to which OTRs in regions of the hindbrain NTS not targeted

by Ong and colleagues [98], other hindbrain sites such as the raphe pallidus [33; 97; 99;

100; 101] or spinal cord [30], contribute to the effect of 4V OT on BAT thermogenesis.

Our findings suggest that increased locomotor activity is not likely to be the main

driver of 4V OT on T_{IBAT}. 4V OT-elicited changes in both T_{IBAT} and core temperature

preceded significant changes in gross motor activity, as did the phospho-ERK1/2

activation patterns, which we observed in the NTS within 15 minutes of 4V OT

administration. It is possible that changes in activity in response to the higher dose of OT

(5 μg) could be potentially important in contributing or maintaining the increase in T_{IBAT}

and core temperature at 1.5-h post-injection, but this does not appear to be the case at

the earlier time points. Consistent with our findings linking CNS OT to activity, Sakamoto

and colleagues reported that intracerebroventricular (ICV) administration of a lower dose

of OT (0.5 μg) increased activity in mice [102]. In addition, Noble and colleagues found

that localized administration of OT (1 nmol ≈ 1.0072 μg) into the ventromedial

hypothalamus also increased physical activity at 1-h post-injection in rats [13]. Sutton and

colleagues also reported that chemogenetic activation of PVN OT neurons produced an

elevation in locomotor activity, energy expenditure and subcutaneous IBAT temperature

[30] in *Oxytocin-Ires* Cre mice. This finding raises the possibility that release of

endogenous OT following activation of PVN OT neurons may also elicit increases in

activity. While our limited sample size and restraint stress (see below) might have

hampered our ability to measure more robust effects of 4V OT on activity at earlier time

points, our findings suggest that locomotor activity does not contribute to the effects of 4V

OT on T_{IBAT} in DIO rats.

Similar to what we observed with our recent studies in mice, one limitation of our

behavioral studies is that handling or restraint stress may have limited our ability to

observe more pronounced differences between drug and vehicle on T_{IBAT} [103] and

activity during the time period when the effect of the drug on T_{IBAT} or activity is relatively

short-lived or small. As a way to minimize the effect of restraint stress on T_{IBAT}, we

adapted the rats to handling and mock injections one week prior to the experiment and

by administering the drugs during the early part of the light cycle when T_{IBAT} [52] and

catecholamine levels are lower [53]. While the effects of CL 316243 continued well

beyond the short-lived effects of restraint/vehicle stress on T_{IBAT} in our rat studies (~30-

45 minutes), the effects of OT and SR 59230A in **Study 5** were relatively short-lived and

may have been masked by restraint stress. Thus, stress-induced epinephrine, from the

adrenal medulla [104; 105; 106], may have activated β3-AR in IBAT to stimulate T_{IBAT},

even in IBAT-denervated rats.

We acknowledge that other BAT depots (axillary, cervical, mediastinal and perirenal

depots), all of which show cold-induced elevations of UCP-1 [107], might act to have

contributed, in part, to the effects of 4V OT to elicit weight loss in IBAT-denervated rats.

We maintained the focus on IBAT given that this depot is the best characterized of all

BAT depots [108]. However, IBAT is thought to contribute to ≥ 70% of total BAT mass

[109] or approximately 45% of total thermogenic capacity of BAT [68]. Moreover, we

acknowledge the potential contribution of beiging of WAT to OT-elicited weight loss and

BAT thermogenesis, particularly EWAT, given that chronic 4V OT was able to elevate

EWAT UCP-3 in IBAT of denervated rats. As indicated above, there is crosstalk between

SNS circuits that innervate WAT and IBAT [72]. In addition, there is increased NE turnover

and IWAT UCP-1 mRNA expression in IBAT-denervated hamsters [72]. It will be

important to 1) confirm our EWAT gene expression findings and to include an analysis of

IWAT in IBAT-denervated rats and 2) develop a model to better understand whether the

other BAT and specific WAT depots contribute to 4V OT-elicited weight loss.

In summary, our findings demonstrate that acute 4V OT (5 μg) produced comparable

increases in T_{IBAT} in both denervated and sham rats. We subsequently found that chronic

4V OT produced similar reductions of BW gain and adiposity in both sham and denervated

rats. Importantly, our findings suggest that there is no change or obvious functional

impairment in the response of the β3-AR agonist, CL 316243, to activate IBAT in rats with

impaired SNS innervation of IBAT in comparison to sham-operated rats. Furthermore, we

acknowledge that increased activity is not likely to have contributed to the effect of 4V OT

to increase both T_{IBAT} and core temperature. In addition, 4V OT appeared to increase

T_{IBAT} prior to core temperature suggesting that OT-induced elevations in T_{IBAT} may also

contribute to OT-induced elevations in core temperature. Together, these findings support

the hypothesis that sympathetic innervation of IBAT is not a predominant mediator of OT-

elicited increases in BAT thermogenesis and reductions of BW and adiposity in male DIO

rats.

1196

1197

1198

References

[1] G. Gimpl, and F. Fahrenholz, The oxytocin receptor system: structure, function, and regulation. Physiol Rev 81 (2001) 629-83.

[2] J.G. Veening, T.R. de Jong, M.D. Waldinger, S.M. Korte, and B. Olivier, The role of oxytocin in male and female reproductive behavior. European journal of pharmacology 753 (2015) 209-28.

[3] M. Kosfeld, M. Heinrichs, P.J. Zak, U. Fischbacher, and E. Fehr, Oxytocin increases trust in humans. Nature 435 (2005) 673-6.

[4] N. Striepens, K.M. Kendrick, W. Maier, and R. Hurlemann, Prosocial effects of oxytocin and clinical evidence for its therapeutic potential. Frontiers in neuroendocrinology 32 (2011) 426-50.

[5] J.E. Blevins, and D.G. Baskin, Translational and therapeutic potential of oxytocin as an anti-obesity strategy: Insights from rodents, nonhuman primates and humans. Physiol Behav (2015).

[6] E.A. Lawson, The effects of oxytocin on eating behaviour and metabolism in humans. Nature reviews. Endocrinology 13 (2017) 700-709.

[7] E.A. Lawson, P.K. Olszewski, A. Weller, and J.E. Blevins, The role of oxytocin in regulation of appetitive behaviour, BW and glucose homeostasis. J Neuroendocrinol 32 (2020) e12805.

[8] S.E. McCormack, J.E. Blevins, and E.A. Lawson, Metabolic Effects of Oxytocin. Endocrine reviews 41 (2020).

[9] J. Altirriba, A.L. Poher, and F. Rohner-Jeanrenaud, Chronic Oxytocin Administration as a Treatment Against Impaired Leptin Signaling or Leptin Resistance in Obesity. Frontiers in endocrinology 6 (2015) 119.

[10] N. Deblon, C. Veyrat-Durebex, L. Bourgoin, A. Caillon, A.L. Bussier, S. Petrosino, F. Piscitelli, J.J. Legros, V. Geenen, M. Foti, W. Wahli, V. Di Marzo, and F. Rohner-Jeanrenaud, Mechanisms of the anti-obesity effects of oxytocin in diet-induced obese rats. PloS one 6 (2011) e25565.

[11] G.J. Morton, B.S. Thatcher, R.D. Reidelberger, K. Ogimoto, T. Wolden-Hanson, D.G. Baskin, M.W. Schwartz, and J.E. Blevins, Peripheral oxytocin suppresses food intake and causes weight loss in diet-induced obese rats. Am J Physiol-Endoc M 302 (2012) E134-E144.

[12] J.E. Blevins, J.L. Graham, G.J. Morton, K.L. Bales, M.W. Schwartz, D.G. Baskin, and P.J. Havel, Chronic oxytocin administration inhibits food intake, increases energy expenditure, and produces weight loss in fructose-fed obese rhesus monkeys. Am J Physiol Regul Integr Comp Physiol 308 (2015) R431-8.

[13] E.E. Noble, C.J. Billington, C.M. Kotz, and C. Wang, Oxytocin in the ventromedial hypothalamic nucleus reduces feeding and acutely increases energy expenditure. Am J Physiol Regul Integr Comp Physiol 307 (2014) R737-45.

[14] G. Zhang, H. Bai, H. Zhang, C. Dean, Q. Wu, J. Li, S. Guariglia, Q. Meng, and D. Cai, Neuropeptide exocytosis involving synaptotagmin-4 and oxytocin in hypothalamic programming of BW and energy balance. Neuron 69 (2011) 523-35.

[15] G. Zhang, and D. Cai, Circadian intervention of obesity development via resting-stage feeding manipulation or oxytocin treatment. Am J Physiol Endocrinol Metab 301 (2011) E1004-12.

[16] B. Cannon, and J. Nedergaard, Brown adipose tissue: function and physiological significance. Physiol Rev 84 (2004) 277-359.

[17] S.F. Morrison, C.J. Madden, and D. Tupone, Central neural regulation of brown adipose tissue thermogenesis and energy expenditure. Cell Metab 19 (2014) 741-56.

[18] M. Periasamy, J.L. Herrera, and F.C.G. Reis, Skeletal Muscle Thermogenesis and Its Role in Whole Body Energy Metabolism. Diabetes Metab J 41 (2017) 327-336.

[19] C.M. Kotz, C.E. Perez-Leighton, J.A. Teske, and C.J. Billington, Spontaneous Physical Activity Defends Against Obesity. Current obesity reports 6 (2017) 362-370.

[20] W.G. Haynes, D.A. Morgan, S.A. Walsh, A.L. Mark, and W.I. Sivitz, Receptor-mediated regional sympathetic nerve activation by leptin. The Journal of clinical investigation 100 (1997) 270-8.

[21] Y. Li, K. Schnabl, S.M. Gabler, M. Willershauser, J. Reber, A. Karlas, S. Laurila, M. Lahesmaa, U.D. M, A. Bast-Habersbrunner, K.A. Virtanen, T. Fromme, F. Bolze, L.S. O'Farrell, J. Alsina-Fernandez, T. Coskun, V. Ntziachristos, P. Nuutila, and M. Klingenspor, Secretin-Activated Brown Fat Mediates Prandial Thermogenesis to Induce Satiation. Cell 175 (2018) 1561-1574 e12.

[22] S. Laurila, L. Sun, M. Lahesmaa, K. Schnabl, K. Laitinen, R. Klen, Y. Li, M. Balaz, C. Wolfrum, K. Steiger, T. Niemi, M. Taittonen, U.D. M, T. Valikangas, L.L. Elo, O. Eskola, A.K. Kirjavainen, L. Nummenmaa, K.A. Virtanen, M. Klingenspor, and P. Nuutila, Secretin activates brown fat and induces satiation. Nat Metab 3 (2021) 798-809.

[23] C. Clemmensen, S. Jall, M. Kleinert, C. Quarta, T. Gruber, J. Reber, S. Sachs, K. Fischer, A. Feuchtinger, A. Karlas, S.E. Simonds, G. Grandl, D. Loher, E. Sanchez-Quant, S. Keipert, M. Jastroch, S.M. Hofmann, E.B.M. Nascimento, P. Schrauwen, V. Ntziachristos, M.A. Cowley, B. Finan, T.D. Muller, and M.H. Tschop, Publisher Correction: Coordinated targeting of cold and nicotinic receptors synergistically improves obesity and type 2 diabetes. Nature communications 9 (2018) 4975.

[24] S. Stanic, K. Bardova, P. Janovska, M. Rossmeisl, J. Kopecky, and P. Zouhar, Prolonged FGF21 treatment increases energy expenditure and induces weight loss in obese mice independently of UCP1 and adrenergic signaling. Biochemical pharmacology 221 (2024) 116042.

[25] A. Hernandez, and M.J. Obregon, Triiodothyronine amplifies the adrenergic stimulation of uncoupling protein expression in rat brown adipocytes. Am J Physiol Endocrinol Metab 278 (2000) E769-77.

[26] R. Cereijo, J. Villarroya, and F. Villarroya, Non-sympathetic control of brown adipose tissue. International Journal of Obesity Supplments 5 (2015) S40-S44.

[27] M. Lopez, C.V. Alvarez, R. Nogueiras, and C. Dieguez, Energy balance regulation by thyroid hormones at central level. Trends Mol Med 19 (2013) 418-27.

[28] Z.S. Roberts, T.H. Wolden-Hanson, M.E. Matsen, V. Ryu, C.H. Vaughan, J.L. Graham, P.J. Havel, D.W. Chukri, M.W. Schwartz, G.J. Morton, and J.E. Blevins, Chronic Hindbrain Administration of Oxytocin is Sufficient to Elicit Weight Loss in Diet-Induced Obese Rats. Am J Physiol Regul Integr Comp Physiol (2017) ajpregu 00169 2017.

[29] M.M. Edwards, H.K. Nguyen, A.D. Dodson, A.J. Herbertson, T. Wolden-Hanson, T. Wietecha, M.K. Honeycutt, J.D. Slattery, K.D. O'Brien, J.L. Graham, P.J. Havel, T.O. Mundinger, C. Sikkema, E.R. Peskind, V. Ryu, G.J. Taborsky, and J.E. Blevins, Sympathetic innervation of interscapular brown adipose tissue is not a predominant

mediator of oxytocin-elicited reductions of BW and adiposity in male diet-induced obese mice. Frontiers in endocrinology 15 (2024).

[30] A.K. Sutton, H. Pei, K.H. Burnett, M.G. Myers, Jr., C.J. Rhodes, and D.P. Olson, Control of food intake and energy expenditure by Nos1 neurons of the paraventricular hypothalamus. J Neurosci 34 (2014) 15306-18.

[31] J. Yuan, R. Zhang, R. Wu, Y. Gu, and Y. Lu, The effects of oxytocin to rectify metabolic dysfunction in obese mice are associated with increased thermogenesis. Mol Cell Endocrinol 514 (2020) 110903.

[32] Y. Kasahara, K. Sato, Y. Takayanagi, H. Mizukami, K. Ozawa, S. Hidema, K.H. So, T. Kawada, N. Inoue, I. Ikeda, S.G. Roh, K. Itoi, and K. Nishimori, Oxytocin receptor in the hypothalamus is sufficient to rescue normal thermoregulatory function in male oxytocin receptor knockout mice. Endocrinology 154 (2013) 4305-15.

[33] Y. Kasahara, Y. Tateishi, Y. Hiraoka, A. Otsuka, H. Mizukami, K. Ozawa, K. Sato, S. Hidema, and K. Nishimori, Role of the Oxytocin Receptor Expressed in the Rostral Medullary Raphe in Thermoregulation During Cold Conditions. Frontiers in endocrinology 6 (2015) 180.

[34] C. Harshaw, J.K. Leffel, and J.R. Alberts, Oxytocin and the warm outer glow: Thermoregulatory deficits cause huddling abnormalities in oxytocin-deficient mouse pups. Hormones and behavior 98 (2018) 145-158.

[35] D. Xi, C. Long, M. Lai, A. Casella, L. O'Lear, B. Kublaoui, and J.D. Roizen, Ablation of Oxytocin Neurons Causes a Deficit in Cold Stress Response. Journal of the Endocrine Society 1 (2017) 1041-1055.

[36] Z. Wu, Y. Xu, Y. Zhu, A.K. Sutton, R. Zhao, B.B. Lowell, D.P. Olson, and Q. Tong, An obligate role of oxytocin neurons in diet induced energy expenditure. PloS one 7 (2012) e45167.

[37] C. Camerino, Low sympathetic tone and obese phenotype in oxytocin-deficient mice. Obesity 17 (2009) 980-4.

[38] Y. Takayanagi, Y. Kasahara, T. Onaka, N. Takahashi, T. Kawada, and K. Nishimori, Oxytocin receptor-deficient mice developed late-onset obesity. Neuroreport 19 (2008) 951-5.

[39] Guide for the Care and Use of Laboratory Animals: Eighth Edition, The National Academies Press, Washington, D.C., 2011.

[40] A. Thureson-Klein, B. Mill-Hyde, T. Barnard, and H. Lagercrantz, Ultrastructrual effects of chemical sympathectomy on brown adipose tissue. J Neurocytol 5 (1976) 677-90.

[41] F. Depocas, D.O. Foster, G. Zaror-Behrens, S. Lacelle, and B. Nadeau, Recovery of function in sympathetic nerves of interscapular brown adipose tissue of rats treated with 6-hydroxydopamine. Canadian journal of physiology and pharmacology 62 (1984) 1327-32.

[42] J.E. Blevins, M.W. Schwartz, and D.G. Baskin, Evidence that paraventricular nucleus oxytocin neurons link hypothalamic leptin action to caudal brain stem nuclei controlling meal size. Am J Physiol Regul Integr Comp Physiol 287 (2004) R87-96.

[43] J.E. Blevins, B.W. Thompson, V.T. Anekonda, J.M. Ho, J.L. Graham, Z.S. Roberts, B.H. Hwang, K. Ogimoto, T.H. Wolden-hanson, J.O. Nelson, K.J. Kaiyala, P.J. Havel, K.L. Bales, G.J. Morton, M.W. Schwartz, and D.G. Baskin, Chronic CNS oxytocin signaling preferentially induces fat loss in high fat diet-fed rats by enhancing satiety responses and increasing lipid utilization. Am J Physiol-Reg I (2016).

[44] G.J. Morton, M.E. Matsen, D.P. Bracy, T.H. Meek, H.T. Nguyen, D. Stefanovski, R.N.
Bergman, D.H. Wasserman, and M.W. Schwartz, FGF19 action in the brain induces
insulin-independent glucose lowering. The Journal of clinical investigation 123 (2013)
4799-808.
[45] G. Paxinos, and C. Watson, (Eds.), The rat brain in stereotaxic coordinates, Academic Press,
Burlington, 2007.
[46] M.D. Dorfman, J.E. Krull, J.D. Douglass, R. Fasnacht, F. Lara-Lince, T.H. Meek, X. Shi, V.
Damian, H.T. Nguyen, M.E. Matsen, G.J. Morton, and J.P. Thaler, Sex differences in
microglial CX3CR1 signalling determine obesity susceptibility in mice. Nature
communications 8 (2017) 14556.
[47] M.N. Brito, N.A. Brito, D.J. Baro, C.K. Song, and T.J. Bartness, Differential activation of the
sympathetic innervation of adipose tissues by melanocortin receptor stimulation.
Endocrinology 148 (2007) 5339-47.
[48] C.H. Vaughan, Y.B. Shrestha, and T.J. Bartness, Characterization of a novel melanocortin
receptor-containing node in the SNS outflow circuitry to brown adipose tissue involved in
thermogenesis. Brain Res 1411 (2011) 17-27.
[49] L.Y. Wang, R.R. Murphy, B. Hanscom, G. Li, S.P. Millard, E.C. Petrie, D.R. Galasko, C.
Sikkema, M.A. Raskind, C.W. Wilkinson, and E.R. Peskind, Cerebrospinal fluid
norepinephrine and cognition in subjects across the adult age span. Neurobiology of aging
34 (2013) 2287-92.
[50] D.L. Williams, R.R. Bowers, T.J. Bartness, J.M. Kaplan, and H.J. Grill, Brainstem
melanocortin 3/4 receptor stimulation increases uncoupling protein gene expression in
brown fat. Endocrinology 144 (2003) 4692-7.
[51] J. Suarez, P. Rivera, S. Arrabal, A. Crespillo, A. Serrano, E. Baixeras, F.J. Pavon, M.
Cifuentes, R. Nogueiras, J. Ballesteros, C. Dieguez, and F.R. de Fonseca,
Oleoylethanolamide enhances beta-adrenergic-mediated thermogenesis and white-to-
brown adipocyte phenotype in epididymal white adipose tissue in rat. Disease models &
mechanisms 7 (2014) 129-141.
[52] M.M. Edwards, H.K. Nguyen, A.J. Herbertson, A.D. Dodson, T. Wietecha, T. Wolden-
Hanson, J.L. Graham, K.D. O'Brien, P.J. Havel, and J.E. Blevins, Chronic Hindbrain
Administration of Oxytocin Elicits Weight Loss in Male Diet-Induced Obese Mice. Am J
Physiol Regul Integr Comp Physiol (2021).
[53] S.F. De Boer, and J. Van der Gugten, Daily Variations in Plasma Noradrenaline, Adrenaline
and Corticosterone Concentrations in Rats Physiol Behav 40 (1987) 323-328.
[54] V. Davidovic, and V.M. Petrovic, Diurnal variations in the catecholamine content in rat
tissues
Effects of exogenous noradrenaline. Archives Iniernationales de PhysioIogie et de Biochimie 89
(1981) 457-460.
[55] Y. Ootsuka, K. Kulasekara, R.C. de Menezes, and W.W. Blessing, SR59230A, a beta-3
adrenoceptor antagonist, inhibits ultradian brown adipose tissue thermogenesis and
interrupts associated episodic brain and body heating. Am J Physiol Regul Integr Comp
Physiol 301 (2011) R987-94.
[56] S. Bexis, and J.R. Docherty, Role of alpha 1- and beta 3-adrenoceptors in the modulation by
SR59230A of the effects of MDMA on body temperature in the mouse. British journal of
pharmacology 158 (2009) 259-66.

[57] P.J. Enriori, P. Sinnayah, S.E. Simonds, C. Garcia Rudaz, and M.A. Cowley, Leptin action in the dorsomedial hypothalamus increases sympathetic tone to brown adipose tissue in spite of systemic leptin resistance. J Neurosci 31 (2011) 12189-97.

[58] G.M. Sutton, L.M. Patterson, and H.R. Berthoud, Extracellular signal-regulated kinase 1/2 signaling pathway in solitary nucleus mediates cholecystokinin-induced suppression of food intake in rats. J Neurosci 24 (2004) 10240-7.

[59] C.A. Campos, J.S. Wright, K. Czaja, and R.C. Ritter, CCK-induced reduction of food intake and hindbrain MAPK signaling are mediated by NMDA receptor activation. Endocrinology 153 (2012) 2633-46.

[60] L.W. Swanson, Brain maps 4.0-Structure of the rat brain: An open access atlas with global nervous system nomenclature ontology and flatmaps. J Comp Neurol 526 (2018) 935-943.

[61] A.M. Khan, A.H. Grant, A. Martinez, G. Burns, B.S. Thatcher, V.T. Anekonda, B.W. Thompson, Z.S. Roberts, D.H. Moralejo, and J.E. Blevins, Mapping Molecular Datasets Back to the Brain Regions They are Extracted from: Remembering the Native Countries of Hypothalamic Expatriates and Refugees. Adv Neurobiol 21 (2018) 101-193.

[62] A.M. Khan, Controlling feeding behavior by chemical or gene-directed targeting in the brain: what's so spatial about our methods? Front Neurosci 7 (2013) 182.

[63] L.W. Swanson, Neuroanatomical Terminology: A Lexicon of Classical Origins and Historical Foundations, Oxford University Press, Oxford, 2015.

[64] A.A. Bremer, K.L. Stanhope, J.L. Graham, B.P. Cummings, W. Wang, B.R. Saville, and P.J. Havel, Fructose-fed rhesus monkeys: a nonhuman primate model of insulin resistance, metabolic syndrome, and type 2 diabetes. Clinical and translational science 4 (2011) 243-52.

[65] J.E. Blevins, D.H. Moralejo, T.H. Wolden-Hanson, B.S. Thatcher, J.M. Ho, K.J. Kaiyala, and K. Matsumoto, Alterations in activity and energy expenditure contribute to lean phenotype in Fischer 344 rats lacking the cholecystokinin-1 receptor gene. Am J Physiol Regul Integr Comp Physiol 303 (2012) R1231-40.

[66] B.P. Cummings, E.K. Digitale, K.L. Stanhope, J.L. Graham, D.G. Baskin, B.J. Reed, I.R. Sweet, S.C. Griffen, and P.J. Havel, Development and characterization of a novel rat model of type 2 diabetes mellitus: the UC Davis type 2 diabetes mellitus UCD-T2DM rat. Am J Physiol Regul Integr Comp Physiol 295 (2008) R1782-93.

[67] K.J. Livak, and T.D. Schmittgen, Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods 25 (2001) 402-8.

[68] A.W. Fischer, C. Schlein, B. Cannon, J. Heeren, and J. Nedergaard, Intact innervation is essential for diet-induced recruitment of brown adipose tissue. Am J Physiol Endocrinol Metab 316 (2019) E487-E503.

[69] M. Bajzer, M. Olivieri, M.K. Haas, P.T. Pfluger, I.J. Magrisso, M.T. Foster, M.H. Tschop, K.A. Krawczewski-Carhuatanta, D. Cota, and S. Obici, Cannabinoid receptor 1 (CB1) antagonism enhances glucose utilisation and activates brown adipose tissue in diet-induced obese mice. Diabetologia 54 (2011) 3121-31.

[70] S.M. Labbe, A. Caron, W.T. Festuccia, R. Lecomte, and D. Richard, Interscapular brown adipose tissue denervation does not promote the oxidative activity of inguinal white adipose tissue in male mice. Am J Physiol Endocrinol Metab 315 (2018) E815-E824.

[71] Q. Cao, J. Jing, X. Cui, H. Shi, and B. Xue, Sympathetic nerve innervation is required for beigeing in white fat. Physiological reports 7 (2019) e14031.

[72] N.L. Nguyen, C.L. Barr, V. Ryu, Q. Cao, B. Xue, and T.J. Bartness, Separate and shared
sympathetic outflow to white and brown fat coordinately regulate thermoregulation and
beige adipocyte recruitment. Am J Physiol Regul Integr Comp Physiol (2016) ajpregu
00344 2016.
[73] L.M. Gorton, A.M. Khan, M. Bohland, G. Sanchez-Watts, C.M. Donovan, and A.G. Watts,
A role for the forebrain in mediating time-of-day differences in glucocorticoid
counterregulatory responses to hypoglycemia in rats. Endocrinology 148 (2007) 6026-39.
[74] G.P. Tapia, L.J. Agostinelli, S.D. Chenausky, J.V.S. Padilla, V.I. Navarro, A. Alagh, G. Si,
R.H. Thompson, S. Balivada, and A.M. Khan, Glycemic Challenge Is Associated with the
Rapid Cellular Activation of the Locus Ceruleus and Nucleus of Solitary Tract:
Circumscribed Spatial Analysis of Phosphorylated MAP Kinase Immunoreactivity. J Clin
Med 12 (2023).
[75] V.S. Susulic, R.C. Frederich, J. Lawitts, E. Tozzo, B.B. Kahn, M.E. Harper, J. Himms-Hagen,
J.S. Flier, and B.B. Lowell, Targeted disruption of the beta 3-adrenergic receptor gene. The
Journal of biological chemistry 270 (1995) 29483-92.
[76] N.J. Rothwell, M.J. Stock, and D.K. Sudera, Beta-adrenoreceptors in rat brown adipose tissue:
proportions of beta 1- and beta 2-subtypes. Am J Physiol 248 (1985) E397-402.
[77] B.E. Levin, and A.C. Sullivan, Beta-1 receptor is the predominant beta-adrenoreceptor on rat
brown adipose tissue. The Journal of pharmacology and experimental therapeutics 236
(1986) 681-8.
[78] J. Galitzky, C. Carpene, A. Bousquet-Melou, M. Berlan, and M. Lafontan, Differential
activation of beta 1-, beta 2- and beta 3-adrenoceptors by catecholamines in white and
brown adipocytes. Fundamental & clinical pharmacology 9 (1995) 324-31.
[79] C. Atgie, F. D'Allaire, and L.J. Bukowiecki, Role of beta1- and beta3-adrenoceptors in the
regulation of lipolysis and thermogenesis in rat brown adipocytes. Am J Physiol 273 (1997)
C1136-42.
[80] C.B. Ueta, G.W. Fernandes, L.P. Capelo, T.L. Fonseca, F.D. Maculan, C.H. Gouveia, P.C.
Brum, M.A. Christoffolete, M.S. Aoki, C.L. Lancellotti, B. Kim, A.C. Bianco, and M.O.
Ribeiro, beta(1) Adrenergic receptor is key to cold- and diet-induced thermogenesis in
mice. The Journal of endocrinology 214 (2012) 359-65.
[81] M.E. Straat, C.A. Hoekx, F.H.P. van Velden, L.M. Pereira Arias-Bouda, L. Dumont, D.P.
Blondin, M.R. Boon, B. Martinez-Tellez, and P.C.N. Rensen, Stimulation of the beta-2-
adrenergic receptor with salbutamol activates human brown adipose tissue. Cell Rep Med
4 (2023) 100942.
[82] D.P. Blondin, S. Nielsen, E.N. Kuipers, M.C. Severinsen, V.H. Jensen, S. Miard, N.Z.
Jespersen, S. Kooijman, M.R. Boon, M. Fortin, S. Phoenix, F. Frisch, B. Guerin, E.E.
Turcotte, F. Haman, D. Richard, F. Picard, P.C.N. Rensen, C. Scheele, and A.C. Carpentier,
Human Brown Adipocyte Thermogenesis Is Driven by beta2-AR Stimulation. Cell Metab
32 (2020) 287-300 e7.
[83] Y. Ishida, M. Matsushita, T. Yoneshiro, M. Saito, S. Fuse, T. Hamaoka, M. Kuroiwa, R.
Tanaka, Y. Kurosawa, T. Nishimura, M. Motoi, T. Maeda, and K. Nakayama, Genetic
evidence for involvement of beta2-adrenergic receptor in brown adipose tissue
thermogenesis in humans. Int J Obes (Lond) 48 (2024) 1110-1117.
[84] K.M. Tate, M.M. Briend-Sutren, L.J. Emorine, C. Delavier-Klutchko, S. Marullo, and A.D.
Strosberg, Expression of three human beta-adrenergic-receptor subtypes in transfected
Chinese hamster ovary cells. European journal of biochemistry / FEBS 196 (1991) 357-61.

[85] N. Reed, and J.N. Fain, Stimulation of respiration in brown fat cells by epinephrine, dibutyryl-
3',5'-adenosine monophosphate, and m-chloro(carbonyl cyanide)phenylhydrazone. The
Journal of biological chemistry 243 (1968) 2843-8.
[86] R.I. Sharara-Chami, M. Joachim, M. Mulcahey, S. Ebert, and J.A. Majzoub, Effect of
epinephrine deficiency on cold tolerance and on brown adipose tissue. Mol Cell Endocrinol
328 (2010) 34-9.
[87] A.M. Strack, W.B. Sawyer, K.B. Platt, and A.D. Loewy, CNS cell groups regulating the
sympathetic outflow to adrenal gland as revealed by transneuronal cell body labeling with
pseudorabies virus. Brain Res 491 (1989) 274-96.
[88] R.P. Dum, D.J. Levinthal, and P.L. Strick, The mind-body problem: Circuits that link the
cerebral cortex to the adrenal medulla. Proc Natl Acad Sci U S A 116 (2019) 26321-26328.
[89] M. Klingenspor, A. Meywirth, S. Stohr, and G. Heldmaier, Effect of unilateral surgical
denervation of brown adipose tissue on uncoupling protein mRNA level and cytochrom-c-
oxidase activity in the Djungarian hamster. J Comp Physiol B 163 (1994) 664-70.
[90] B. Doslikova, D. Tchir, A. McKinty, X. Zhu, D.L. Marks, V.E. Baracos, and W.F. Colmers,
Convergent neuronal projections from paraventricular nucleus, parabrachial nucleus, and
brainstem onto gastrocnemius muscle, white and brown adipose tissue in male rats. J Comp
Neurol 527 (2019) 2826-2842.
[91] B.J. Oldfield, M.E. Giles, A. Watson, C. Anderson, L.M. Colvill, and M.J. McKinley, The
neurochemical characterisation of hypothalamic pathways projecting polysynaptically to
brown adipose tissue in the rat. Neuroscience 110 (2002) 515-26.
[92] H. Shi, and T.J. Bartness, Neurochemical phenotype of sympathetic nervous system outflow
from brain to white fat. Brain research bulletin 54 (2001) 375-85.
[93] S. Stanley, S. Pinto, J. Segal, C.A. Perez, A. Viale, J. DeFalco, X. Cai, L.K. Heisler, and J.M.
Friedman, Identification of neuronal subpopulations that project from hypothalamus to
both liver and adipose tissue polysynaptically. Proc Natl Acad Sci U S A 107 (2010) 7024-
9.
[94] L. Rinaman, Oxytocinergic inputs to the nucleus of the solitary tract and dorsal motor nucleus
of the vagus in neonatal rats. J Comp Neurol 399 (1998) 101-9.
[95] P.E. Sawchenko, and L.W. Swanson, Immunohistochemical identification of neurons in the
paraventricular nucleus of the hypothalamus that project to the medulla or to the spinal
cord in the rat. J Comp Neurol 205 (1982) 260-72.
[96] M. Bamshad, C.K. Song, and T.J. Bartness, CNS origins of the sympathetic nervous system
outflow to brown adipose tissue. Am J Physiol 276 (1999) R1569-78.
[97] G. Cano, A.M. Passerin, J.C. Schiltz, J.P. Card, S.F. Morrison, and A.F. Sved, Anatomical
substrates for the central control of sympathetic outflow to interscapular adipose tissue
during cold exposure. J Comp Neurol 460 (2003) 303-26.
[98] Z.Y. Ong, D.M. Bongiorno, M.A. Hernando, and H.J. Grill, Effects of Endogenous Oxytocin
Receptor Signaling in Nucleus Tractus Solitarius on Satiation-Mediated Feeding and
Thermogenic Control in Male Rats. Endocrinology 158 (2017) 2826-2836.
[99] D. Kong, Q. Tong, C. Ye, S. Koda, P.M. Fuller, M.J. Krashes, L. Vong, R.S. Ray, D.P. Olson,
and B.B. Lowell, GABAergic RIP-Cre neurons in the arcuate nucleus selectively regulate
energy expenditure. Cell 151 (2012) 645-57.
[100] S.F. Morrison, and K. Nakamura, Central neural pathways for thermoregulation. Frontiers
in bioscience 16 (2011) 74-104.
[101] S.F. Morrison, Central control of body temperature. F1000Research 5 (2016).

[102] T. Sakamoto, S. Sugimoto, and T. Uekita, Effects of intraperitoneal and
intracerebroventricular injections of oxytocin on social and emotional behaviors in pubertal
male mice. Physiol Behav 212 (2019) 112701.
[103] Y. Ootsuka, W.W. Blessing, and E. Nalivaiko, Selective blockade of 5-HT2A receptors
attenuates the increased temperature response in brown adipose tissue to restraint stress in
rats. Stress 11 (2008) 125-33.
[104] W.H. Harris, D.O. Foster, S.W. Ma, S. Yamashiro, and L.A. Langlais-Burgess, The
noradrenaline content and innervation of brown adipose tissue in the young rabbit.
Canadian journal of physiology and pharmacology 64 (1986) 561-7.
[105] G.J. Taborsky, Jr., Q. Mei, D.J. Hackney, D.P. Figlewicz, R. LeBoeuf, and T.O. Mundinger,
Loss of islet sympathetic nerves and impairment of glucagon secretion in the NOD mouse:
relationship to invasive insulitis. Diabetologia 52 (2009) 2602-11.
[106] L.K. Gilliam, J.P. Palmer, and G.J. Taborsky, Jr., Tyramine-mediated activation of
sympathetic nerves inhibits insulin secretion in humans. J Clin Endocrinol Metab 92 (2007)
4035-8.
[107] J.M. de Jong, O. Larsson, B. Cannon, and J. Nedergaard, A stringent validation of mouse
adipose tissue identity markers. Am J Physiol Endocrinol Metab 308 (2015) E1085-105.
[108] T.J. Bartness, C.H. Vaughan, and C.K. Song, Sympathetic and sensory innervation of brown
adipose tissue. Int J Obes (Lond) 34 Suppl 1 (2010) S36-42.
[109] N.C. Bal, S.K. Maurya, S. Singh, X.H. Wehrens, and M. Periasamy, Increased Reliance on
Muscle-based Thermogenesis upon Acute Minimization of Brown Adipose Tissue
Function. The Journal of biological chemistry 291 (2016) 17247-57.
[110] K.F.V.B.u.L.d.G. Burdach, Dyk'schen Buchhandlung, Leipzig, 1822.

Figure legends

Figure 1. Effect of IBAT surgical denervation procedure on IBAT NE content at 1-week post-sham or IBAT denervation in lean rats. Rats were maintained on chow (13% kcal from fat) for approximately 1 week prior to undergoing a sham (N=5/group) or bilateral surgical IBAT denervation (N=5/group) procedures and were euthanized at 1-week post-sham or denervation procedure. NE content was measured from IBAT, IWAT and liver. Data are expressed as mean ± SEM. *$P<0.05$, †$0.05<P<0.1$ denervation vs sham.

Figure 2: Effect of IBAT surgical denervation procedure on IBAT NE content at 1-, 6- and 7-weeks post-sham or IBAT denervation in DIO rats. Rats were maintained on a HFD (60% kcal from fat) for approximately 5 months prior to undergoing a sham (N=5/group) or bilateral surgical IBAT denervation (N=5/group) procedures and were euthanized at 1-, 6- and 7-weeks post-sham (N=15/group) or denervation procedure (N=15/group). NE content was measured from IBAT, IWAT, liver and pancreas. Data are expressed as mean ± SEM. *$P<0.05$, †$0.05<P<0.1$ denervation vs sham.

Figure 3A-F: Effect of systemic β3-AR agonist (CL 31643) administration (0.1 and 1 mg/kg) on IBAT temperature (T_{IBAT}), energy intake and BW post-sham or IBAT denervation in DIO rats. Rats were maintained on HFD (60% kcal from fat; N=6–10/group) for approximately 4.25 months prior to undergoing a sham or bilateral surgical IBAT denervation and implantation of temperature transponders underneath IBAT. Animals were subsequently adapted to a 4-h fast prior to receiving IP injections of CL 316243 (0.1 or 1 mg/kg, IP) or vehicle (sterile water) where each animal received each treatment at approximately 7-day intervals. *A/C*, Effect of CL 316243 on T_{IBAT} in *A)* sham operated or *C)* IBAT denervated DIO rats; *B/D*, Effect of CL 316243 on change in T_{IBAT} relative to baseline T_{IBAT} (delta T_{IBAT}) in *B)* sham operated or *D)* IBAT denervated DIO rats; *E*, Effect of CL 316243 on change in energy intake in sham or IBAT denervated DIO rats; *F*, Effect of CL 316243 on change in BW in sham or IBAT denervated DIO rats. Data are expressed as mean ± SEM. *$P<0.05$, †$0.05<P<0.1$ CL 316243 vs. vehicle.

Figure 4A–D: Effect of acute 4V OT administration (1 and 5 μg) on T$_{IBAT}$ post-sham or IBAT denervation in male DIO rats. Rats were maintained on HFD (60% kcal from fat; N=6–8/group) for approximately 4.25 months prior to undergoing a sham or bilateral surgical IBAT denervation and implantation of temperature transponders underneath IBAT. Rats were subsequently implanted with 4V cannulas and allowed to recover for 2 weeks prior to receiving acute 4V injections of OT or vehicle. Animals were subsequently adapted to a 4-h fast prior to receiving acute 4V injections of OT or vehicle *A/C*, Effect of acute 4V OT on T$_{IBAT}$ in *A)* sham operated or *C)* IBAT denervated DIO rats; *B/D*, Effect of acute 4V OT on change in T$_{IBAT}$ relative to baseline T$_{IBA}$T (delta T$_{IBAT}$) in *B)* sham operated or *D)* IBAT denervated DIO rats; Data are expressed as mean ± SEM. *$P<0.05$, †$0.05<P<0.1$ OT vs. vehicle.

Figure 5A–B: Effect of β3-AR antagonism on ability of acute 4V OT to increase T$_{IBAT}$ in DIO rats. Rats were maintained on HFD (60% kcal from fat; N=7/group) for approximately 4 months prior to being implanted with temperature transponders underneath the left IBAT depot. Rats were subsequently implanted with 4V cannulas and allowed to recover for 4 weeks. Animals were subsequently adapted to a 4-h fast and handling during the week prior to the experiment. On an experimental day, 4-h fasted rats received the β3-AR antagonist, SR 59230A (0.5 mg/kg, IP) or vehicle (DMSO) approximately 20 min prior to 4V administration of either OT (5 μg/μl) or vehicle. *A*, Effect of systemic pre-treatment with the β3-AR antagonist, SR 59230A, on the ability of acute 4V OT to stimulate T$_{IBAT}$ at 0.5-h post-injection in DIO rats; B), Effect of systemic pre-treatment with the β3-AR antagonist, SR 59230A, on the ability of acute 4V OT to stimulate T$_{IBAT}$ at 0.75-h post-injection in DIO rats. Data are expressed as mean ± SEM. *$P<0.05$, †$0.05<P<0.1$.

Figure 6A–D: Effect of chronic 4V OT infusions (16 nmol/day) on BW, adiposity and energy intake post-sham or IBAT denervation in male DIO rats. *A*, Rats were maintained on HFD (60% kcal from fat; N=8–11/group) for approximately 4.5 months prior to undergoing a sham or bilateral surgical IBAT denervation. Rats were subsequently

implanted with 4V cannulas and allowed to recover for 2 weeks prior to being implanted with subcutaneous minipumps that were subsequently attached to the 4V cannula. *A*, Effect of chronic 4V OT or vehicle on BW in sham operated or IBAT denervated DIO rats; *B*, Effect of chronic 4V OT or vehicle on BW change in sham operated or IBAT denervated DIO rats; *C*, Effect of chronic 4V OT or vehicle on adiposity in sham operated or IBAT denervated DIO rats; *D*, Effect of chronic 4V OT or vehicle on adiposity in sham operated or IBAT denervated DIO rats. Data are expressed as mean ± SEM. *$P<0.05$, †$0.05<P<0.1$ OT vs. vehicle.

Figure 7A–B: Effect of chronic 4V OT infusions (16 nmol/day) on adipocyte size post-sham or IBAT denervation in male DIO rats. *A*, Adipocyte size (pixel^2) was measured in IWAT from rats that received chronic 4V infusion of OT (16 nmol/day) or vehicle in sham or IBAT denervated DIO rats (N=4–5/group). *B*, Adipocyte size was measured in EWAT from rats that received chronic 4V infusion of OT (16 nmol/day) or vehicle in sham operated or IBAT denervated rats (N=4–5/group). Data are expressed as mean ± SEM. *$P<0.05$ OT vs. vehicle.

Figure 8A–D: Effect of acute 4V OT on gross motor activity, core temperature, T_{IBAT} and energy intake in DIO rats. Rats were maintained on HFD (60% kcal from fat; N=7/group) for approximately 6 months prior to being implanted with temperature transponders underneath IBAT, intra-abdominal telemetry devices and 4V cannulas. Animals were subsequently adapted to a 4-h fast prior to receiving acute 4V injections of OT or vehicle. Animals remained fasted for additional 4 h during which time gross motor activity, T_{IBAT} and core temperature was measured. Energy intake data was collected separately from the same animals following a 4-h fast *A*, Effect of acute 4V OT on gross motor activity in DIO rats; *B*, Effect of acute 4V OT on T_{IBAT} in DIO rats; C, Effects of acute 4V OT on core temperature in DIO rats, and D, Effect of acute 4V OT on energy intake in DIO rats. Data are expressed as mean ± SEM. *$P<0.05$, †$0.05<P<0.1$ OT vs. vehicle.

Figure 9A–C: Elevated recruitment of hindbrain nucleus of the solitary tract (NTS) neurons in male DIO rats 15 min after 4V OT administration. *A*, Sagittal view of the rat

brain adapted from *BM4.0* depicting the dorsal medullary atlas level area surveyed here. Midline shown in blue. *B/C,* Immunostained hindbrain sections of acute 4V VEH- *(B)* or *(C)* OT-treated subjects. The immunoreactive patterns were used to define the regional boundaries and are represented with dotted lines. Scale bars = 100 μm. Images shown were captured at ✕20 magnification. The header notes the inferred anteroposterior stereotaxic coordinate, expressed as millimeters from the bregma suture on the skull (β). *Note that the area postrema is damaged from histological processing. Abbreviations: AP, *Area postrema (>1840)*; Cmd, *Central canal of medulla (>1840)*; DMX, *Dorsal motor nucleus of vagus nerve (>1840)*; fus, *Funiculus separans (>1840)*; GRp, *gracile nucleus, principal part (>1840)*; NTSco, *Nucleus of solitary tract, commissural part (>1840)*; NTSmc, *Nucleus of solitary tract, medial central part (>1840)*; tcbf, *Transverse cerebellar fissure [110]*; ts, *Solitary tract (>1840)*. All *BM4.0* atlas maps were modified and/or reproduced with permission under the conditions outlined by a Creative Commons BY NC 4.0 license (https://creativecommons.org/licenses/by-nc/4.0/).

Table 1. Changes in plasma hormones at 1, 6, and 7-weeks post-sham or IBAT denervation in DIO rats. Blood was collected by tail vein (blood glucose) and cardiac stick (plasma hormones) following a 4-h fast. Different letters denote significant differences between respective sham and denervation groups at 1, 6 or 7-weeks post-sham or denervation. Shared letters denote no significant differences between respective sham and denervation groups. Data are expressed as mean ± SEM (N=5/group).

Table 2. Plasma measurements following 4V infusions of OT (16 nmol/day) or vehicle in sham and IBAT denervated DIO rats. Blood was collected by tail vein (blood glucose) and cardiac stick (plasma hormones) following a 4-h fast. Different letters denote significant differences between treatments. Shared letters denote no significant differences between treatments. Data are expressed as mean ± SEM (N=8-11/group).

Table 3. Changes in IBAT and EWAT gene expression following 4V infusions of OT or vehicle in male sham or IBAT denervated DIO rats. IBAT and EWAT were collected following a 4-h fast. *A,* Changes in IBAT mRNA expression following 4V infusions of OT

1683 or vehicle in male sham or IBAT denervated DIO rats; *B*, Changes in EWAT mRNA

1684 expression following 4V infusions of OT or vehicle in male sham or IBAT denervated DIO

1685 rats. Different letters denote significant differences between treatments. Shared letters

1686 denote no significant differences between treatments. Data are expressed as mean ±

1687 SEM. *P<0.05 (N=3-10/group).

1688

1689

1690

1691

1692

1693

1694

1695

1696

1697

1698

1699

1700

1701

1702

1703

1704

1705

1706

1707

1708

1709

1710

1711

1712

1713

Supplemental Study 1: Determine if surgical denervation of IBAT blocked the ability of 4V OT to increase T$_{IBAT}$ in lean rats. The goal of this study was to determine if OT- elicited increase in T$_{IBAT}$ requires intact SNS outflow to IBAT in lean rats. By design, rats were lean as determined by both BW (511.6 ± 2.4 g) and adiposity (3.3 ± 0.3 g fat mass; 18.1 ± 11.1% adiposity) after maintenance on the chow (13% kcal from fat; N=6/group) for approximately 1 week prior to denervation procedures and implantation of temperature transponders underneath IBAT. Rats were otherwise treated identically to those used in **Study 4**.

Supplemental Study 1:

Tissue samples from **Supplemental Study 1** were screened for NE content and compared with a previous cohort of sham-operated rats from **Study 1**. However, there was no age-matched cohort to use for comparison purposes to calculate reduction of IBAT NE content as the cohort used in **Supplemental Study 1** was heavier than the sham-operated cohort used in **Study 1** (354 ± 4.6 g) [(F(1,9) = 608.553, *P*<0.001]. IBAT NE content from sham-operated animals was only used to screen whether IBAT NE content was above the acceptable threshold for a successful IBAT denervation. NE content was reduced by 93.8 ±2.6% in IBAT from denervated rats relative to IBAT from sham-operated control rats [(F(1,9) = 16.390, *P*<0.01]. IWAT NE content was also reduced in the denervated group [(F(1,9) = 7.411, *P*<0.05)] but this might have been due, in part, to the differences in age of the animals between cohorts. In contrast, NE content was unchanged in pancreas but elevated in liver [(F(1,9) = 8.684, *P*<0.05] in denervated rats relative to sham rats.

In denervated rats, 4V OT (5 μg/μL) increased T$_{IBAT}$ at 0.5, 0.75, 1, 1.25, 1.5 and 1.75-h post-injection (*P*<0.05; **Supplemental Figure 1A**). The high dose (5 μg/μL) also stimulated T$_{IBAT}$ when measuring change in T$_{IBAT}$ relative to baseline T$_{IBAT}$ at 0.5, 0.75, 1, 1.25, 1.5, 1.75 and 2-h post-injection (*P*<0.05; **Supplemental Figure 1B**). The low dose (1 μg/μL) increased T$_{IBAT}$ at 1.5-h post-injection (*P*<0.05; **Supplemental Figure 1A**). The low dose (1 μg/μL) tended to stimulate T$_{IBAT}$ when measuring change in T$_{IBAT}$ relative to baseline T$_{IBAT}$ at 1- and 1.5-h post-injection (0.05<*P*<0.1; **Supplemental Figure 1B**).

Supplemental Study 2: Determine the extent to which β3-AR-induced activation of IBAT requires activation of β3-AR to increase T_{IBAT} in DIO rats.

Rats were fed *ad libitum* and maintained on HFD for approximately 4.5 months prior to receiving 4V cannulas and/or temperature transponders [28]. Rats were allowed to recover for at least 4 weeks during which time they were adapted to a daily 4-h fast, handling and mock injections.

The goal of this study was to determine a dose of the β3-AR antagonist that was sufficient to block the effects of the β3-AR agonist on T_{IBAT} in DIO rats. The dose of the β3-AR antagonist would then be used in **Study 6** to determine if 4V OT required β3-AR signaling to stimulate T_{IBAT} in DIO rats. By design, DIO rats were obese as determined by both BW (712 ± 23.1 g) and adiposity (217 ± 12.1 g fat mass; 31 ± 1.0% adiposity) after maintenance on the HFD for 4.5 months prior to sham/denervation procedures.

In the absence of the β3-AR antagonist, the β3-AR agonist increased T_{IBAT} at 0.25, 0.5, 0.75, 1, 1.5, 0.75, 2 and 4-h post-injection ($P<0.05$) and tended to stimulate T_{IBAT} at 180-min post-injection ($0.05<P<0.1$) (**Supplemental Figure 2**). These effects were blocked at 0.5-h post-injection of the β3-AR agonist. Together, these findings confirm the specificity of a β3-AR mediated effect of the β3-AR agonist on T_{IBAT} and identified a dose of the β3-AR antagonist that was sufficient to block the effects of the β3-AR agonist to stimulate T_{IBAT}.

Supplemental Study 3: Determine the extent to which OT-induced activation of sympathetic outflow to IBAT contributes to its ability to elicit weight loss in rats with more pronounced diet-induced obesity.

Rats were fed *ad libitum* and maintained on HFD for approximately 8.75 months prior to receiving 4V cannulas and minipumps to infuse vehicle or OT (16 nmol/day) over 29 days as previously described [28]. Daily food intake and BW were also tracked for 29 days. Animals were euthanized by rapid conscious decapitation at 9 weeks post-sham or denervation procedure. Trunk blood and tissues (IBAT, EWAT, IWAT, liver and pancreas) were collected from 4-h fasted rats and tissues were subsequently analyzed for IBAT NE

1776 content to confirm success of denervation procedure relative to sham operated animals

1777 and other tissues (EWAT, IWAT, liver and pancreas).

1778

The goal of this study was to determine if OT-elicited weight loss requires intact SNS

outflow to IBAT. By design, DIO rats were obese as determined by both BW (790.7 ± 23.8

g) and adiposity (281.9 ± 16.3 g fat mass; 35.3 ± 0.1.3% adiposity) after maintenance on

the HFD for 8.75 months prior to sham/denervation procedures. IBAT NE content was

reduced in denervated rats by 84.8±4.3% in denervated rats relative to sham-operated

control rats [(F(1,16) = 128.544, P=0.000)]. In contrast, NE content was unchanged in

IWAT, EWAT or pancreas in denervated rats relative to sham rats (P=NS). In contrast,

there was a reduction of NE content in liver in denervated rats relative to sham rats

[(F(1,16) = 4.654, P=0.047)]. There was no significant difference in BW between sham

and denervation groups at the end of the study (P=NS; data not shown).

1789

In sham-operated rats, as expected, 4V vehicle resulted in 4.3 ± 1.0% weight gain relative

to vehicle pre-treatment [(F(1,4) = 15.268, P=0.017]. In contrast, 4V OT reduced BW by

4.2 ± 1.2% relative to OT pre-treatment [(F(1,4) = 10.931, P=0.030] (**Supplemental**

Figure 3A) and it also reduced weight gain throughout the 29-day infusion period

(**Supplemental Figure 3B**). OT treatment reduced weight gain on days 4-29 (P<0.05)

and tended to reduce weight gain on day 3 (P=0.068). 4V OT reduced relative fat mass

(pre- vs post-intervention; **Supplemental Figure 3C**) (P<0.05) and also produced a

reduction in the relative lean body mass (pre- vs post-intervention; P<0.05) without

impacting total fat mass or lean body mass (P=NS). These effects that were mediated, at

least in part, by a modest reduction of energy intake that persisted throughout the first

two weeks of the treatment period (**Supplemental Figure 3D**; P<0.05). 4V OT also

tended to reduce energy intake during weeks 3 and 4 of the treatment period

(**Supplemental Figure 3D**; 0.05<P<0.1).

1803

In denervated rats, as expected, 4V vehicle resulted in 4.7 ± 0.8% weight gain relative to

vehicle pre-treatment [(F(1,3) = 44.734, P=0.007] (**Supplemental Figure 3A**). In contrast,

4V OT failed to reduce BW relative to OT pre-treatment but it reduced weight gain

throughout the 29-day infusion period (**Supplemental Figure 3B**). OT treatment reduced

weight gain on days 4-29 ($P<0.05$) and it tended to reduce weight gain on day 3 ($P=0.05$).

4V OT reduced relative fat mass (pre- vs post-intervention; **Supplemental Figure 3C**)

($P<0.05$) without effecting total fat mass or lean body mass ($P=$NS). These effects that

were mediated, at least in part, by a modest reduction of energy intake that was evident

during weeks 1 and 2 of the treatment period (**Supplemental Figure 3D**; $P<0.05$). 4V OT

also tended to reduce energy intake during week 3 of the treatment period

(**Supplemental Figure 3D**; $P=0.103$).

1815

1816 Overall, these findings demonstrate that sympathetic innervation of IBAT is not a

1817 predominant mediator of 4V OT-elicited reductions of BW and adiposity in male DIO rats.

1818

1819

1820

1821

1822

1823

1824

1825

1826

1827

1828

1829

1830

1831

1832

1833

1834

1835

1836

1837

 Figure legend

Supplemental Figure 1A–D: Effect of 4V OT on IBAT temperature (T$_{IBAT}$) in lean

rats. Rats were fed *ad libitum* and maintained on low fat chow diet (13% kcal from fat)

(N=6/group) for approximately 1 week prior to underdoing SNS denervation procedures

and implantation of temperature transponders underneath the left IBAT depot. Rats

subsequently received 4V cannulations. Rats were allowed to recover for at least 2 weeks

during which time they were adapted to a daily 4-h fast, handling and mock injections.

Rats subsequently received injections of 4V OT (1 or 5 µg/µL) or vehicle where each

animal received each treatment at least 48-h intervals. *A*, Effect of 4V OT on T$_{IBAT}$ in IBAT

denervated lean rats; *B*, Effect of 4V OT on change in T$_{IBAT}$ relative to baseline T$_{IBAT}$ (delta

T$_{IBAT}$) in IBAT denervated lean rats. Data are expressed as mean ± SEM. *$P<0.05$,

†$0.05<P<0.1$ 4V OT vs. vehicle.

1850

Supplemental Figure 2A–B: Effect of β3-AR antagonist, SR59230A, on the ability

of the β3-AR agonist, CL 316243, to increase T$_{IBAT}$ in DIO rats. Rats were maintained

on HFD (60% kcal from fat; N=17/group) for approximately 4.25 months prior to being

implanted with temperature transponders underneath IBAT. Animals were subsequently

adapted to a 4-h fast prior to receiving acute IP injections of the β3-AR agonist,

SR59230A or vehicle approximately 20 minutes prior to IP injections of the β3-AR agonist,

CL 316243. *A*, Effect of the β3-AR antagonist (SR-59230A) pre-treatment on the ability

of the β3-AR agonist, CL 316243, to stimulate T$_{IBAT}$ at 0.5-h post-injection; *B*, Effect of the

β3-AR antagonist (SR-59230A) pre-treatment on the ability of the β3-AR agonist, CL

316243, to stimulate T$_{IBAT}$ at 0.75-h post-injection. Data are expressed as mean ± SEM.

*$P<0.05$, †$0.05<P<0.1$ OT vs. vehicle.

1862

1863 **Supplemental Figure 3A–D: Effect of chronic 4V OT infusions (16 nmol/day) on**

1864 **BW, adiposity and energy intake post-sham or IBAT denervation in male DIO rats**

1865 **with more pronounced diet-induced obesity.** *A*, Rats were maintained on HFD (60%

1866 kcal from fat; N=4-5/group) for approximately 8.75 months prior to undergoing a sham or

1867 bilateral surgical IBAT denervation. Rats were subsequently implanted with 4V cannulas

1868 and allowed to recover for 2 weeks prior to being implanted with subcutaneous minipumps

1869 that were subsequently attached to the 4V cannula. *A*, Effect of chronic 4V OT or vehicle

1870 on BW in sham operated or IBAT denervated DIO rats; *B*, Effect of chronic 4V OT or

1871 vehicle on BW change in sham operated or IBAT denervated DIO rats; *C*, Effect of chronic

1872 4V OT or vehicle on adiposity in sham operated or IBAT denervated DIO rats; *D*, Effect

1873 of chronic 4V OT or vehicle on adiposity in sham operated or IBAT denervated DIO rats.

1874 Data are expressed as mean ± SEM. *$P<0.05$, †$0.05<P<0.1$ OT vs. vehicle.

1875

Figure 1

Figure 2A

Figure 2B

Figure 2C

Figure 3A

SHAM

Figure 3B

Figure 3C

Figure 3D

DENERVATION

Figure 3E

Figure 3F

SHAM

Figure 4A

Figure 4B

DENERVATED

Figure 4C

Figure 4D

Figure 5A

Figure 5B

Figure 6A

Figure 6B

Figure 6C

Figure 6D

Figure 7A

Figure 7B

Figure 8A

Figure 8B

Figure 8C

Figure 8D

A

BM4.0 Atlas Level 70 | β −14.16 mm

B

Vehicle

C

Oxytocin

pERK1/2-immunoreactive cells 15 min after intra-4V injection

Figure 9

Table 1.   Plasma Measurements at 1, 6 and 7 Weeks Post-SNS Denervation of IBAT In DIO Rats

	1 week		6 weeks		7 weeks	
	Sham	Denervation	Sham	Denervation	Sham	Denervation
Leptin (ng/mL)	25.7 ± 4.4[a]	31.9 ± 5.8[a]	59.7 ± 10.8[a]	53.4 ± 10.0[a]	50.2 ± 5.3[a]	55.3 ± 9.3[a]
Insulin (ng/mL)	3.4 ± 0.6[a]	5.0 ± 1.8[a]	5.6 ± 0.7[a]	9.1 ± 3.8[a]	6.0 ± 0.7[a]	9.5 ± 2.3[a]
FGF-21 (pg/mL)	181.5 ± 20.4[a]	211.7 ± 68.1[a]	219.3 ± 39.3[a]	180.9 ± 53.2[a]	212 ± 63.2[a]	240.5 ± 86.2[a]
Irisin (μg/mL)	11.4 ± 2.6[a]	10.3 ± 0.8[a]	10.6 ± 1.2[a]	11.3 ± 0.8[a]	10.6 ± 0.3[a]	7.7 ± 0.9[a]
Adiponectin (μg/mL)	7.4 ± 1.0[a]	7.1 ± 0.4[a]	8.1 ± 0.5[a]	7.5 ± 0.5[a]	7.2 ± 0.7[a]	7.3 ± 0.8[a]
Blood Glucose (mg/dL)	169.8 ± 11.9[a]	140.4 ± 6.2[b]	152 ± 8.2[a]	169.6 ± 7.2[a]	156 ± 11.4[a]	161.2 ± 5.1[a]
TG	78.5 ± 11.4[a]	97.3 ± 13.1[a]	119.9 ± 11.6[a]	150.9 ± 40.2[a]	101.9 ± 12.4[a]	130.7 ± 8.6[a]
FFA (mEq/L)	0.19 ± 0.01[a]	0.23 ± 0.03[a]	0.22 ± 0.02[a]	0.25 ± 0.03[a]	0.27 ± 0.06[a]	0.22 ± 0.04[a]
Total Cholesterol (mg/dL)	80.1 ± 3.1[a]	96.8 ± 9.3[a]	137.3 ± 8.7[a]	126.5 ± 7.3[a]	126.4 ± 15.5[a]	130.2 ± 9.3[a]

Table 2. Plasma Measurements Following 4V VEH or OT in Sham or Denervated DIO Rats

4V	VEH	OT	VEH	OT
	Sham	Sham	Denervation	Denervation
Leptin (ng/mL)	44.9 ± 4.8^a	29 ± 3.2^b	39.2 ± 6.5^{ab}	27.9 ± 6.6^b
Insulin (ng/mL)	6.0 ± 1.0^a	5.0 ± 0.6^a	5.0 ± 0.8^a	3.8 ± 0.7^a
FGF-21 (pg/mL)	153.2 ± 29.3^a	166.2 ± 40.7^a	146.5 ± 27.6^a	124.3 ± 20.7^a
Irisin (μg/mL)	8.2 ± 0.8^a	8.5 ± 0.6^a	6.4 ± 0.4^a	6.5 ± 0.9^a
Adiponectin (μg/mL)	7.4 ± 0.6^a	6.6 ± 0.4^{ab}	7.6 ± 0.3^a	6.0 ± 0.6^b
Blood Glucose (mg/dL)	183.5 ± 7.3^a	188.4 ± 4.4^a	185.8 ± 9.2^a	179.1 ± 7.9^a
FFA (mEq/L)	0.24 ± 0.03^a	0.24 ± 0.02^a	0.28 ± 0.05^a	0.23 ± 0.03^a
Total Cholesterol (mg/dL)	116.4 ± 7.1^a	113.1 ± 3.3^a	110.2 ± 9.9^a	102.8 ± 7.0^a

Table 3A. Changes in IBAT mRNA Expression Following Chronic 4V Infusions of OT or VEH in Male DIO Rats

4V Treatment	VEH	OT	VEH	OT
	Sham	Sham	Denervated	Denervated
IBAT				
Adra2a	1.0 ± 0.3^a	2.5 ± 1.4^a	1.0 ± 0.2^a	2.6 ± 1.1^a
adrb1	1.0 ± 0.2^a	4.5 ± 1.0^b	2.2 ± 0.9^a	2.1 ± 0.5^a
adrb2	1.0 ± 0.1^a	1.8 ± 0.2^a	2.1 ± 0.7^a	2.6 ± 1.1^a
adrb3	1.0 ± 0.6^a	0.3 ± 0.1^a	0.3 ± 0.1^a	0.3 ± 0.01^a
UCP1	1.0 ± 0.5^a	0.6 ± 0.1^a	0.3 ± 0.1^a	0.4 ± 0.1^a
UCP3	1.0 ± 0.3^a	1.2 ± 0.2^a	0.6 ± 0.2^a	0.8 ± 0.3^a
Gpr120	1.0 ± 0.3^a	4.0 ± 1.9^a	0.9 ± 0.4^a	0.8 ± 0.3^a
Dio2	1.0 ± 0.3^a	0.9 ± 0.2^a	0.2 ± 0.1^b	0.9 ± 0.2^{ac}
Cidea	1.0 ± 0.4^a	2.3 ± 0.6^a	0.8 ± 0.3^a	1.2 ± 0.5^a
Prdm16	1.0 ± 0.5^a	0.1 ± 0.04^a	0.1 ± 0.01^a	0.2 ± 0.1^a
Cox8b	1.0 ± 0.3^a	10.2 ± 3.6^b	0.8 ± 0.4^a	1.3 ± 0.6^a
Bmp8b	1.0 ± 0.1^{ab}	1.3 ± 0.4^{ab}	0.6 ± 0.3^a	2.1 ± 0.5^b
Cn1	1.0 ± 0.3^a	2.7 ± 0.8^a	1.3 ± 0.5^a	1.2 ± 0.6^a

Table 3B. Changes in EWAT mRNA Expression Following Chronic 4V Infusions of OT or VEH in Male DIO Rats

4V Treatment	VEH	OT	VEH	OT
	Sham	Sham	Denervated	Denervated
EWAT				
Adra2a	1.0 ± 0.2^a	0.9 ± 0.2^a	0.8 ± 0.1^a	1.0 ± 0.2^a
adrb1	1.0 ± 0.2^a	0.7 ± 0.2^a	0.7 ± 0.2^a	0.6 ± 0.2^{ab}
adrb2	1.0 ± 0.2^a	0.6 ± 0.1^b	0.6 ± 0.1^{ab}	0.7 ± 0.1^{ab}
adrb3	1.0 ± 0.2^a	1.9 ± 0.3^b	1.8 ± 0.3^b	2.3 ± 0.1^b
UCP1	1.0 ± 0.3^a	1.6 ± 0.4^{ab}	9.8 ± 4.2^b	4.9 ± 2.5^{ab}
UCP3	1.0 ± 0.3^{ab}	0.6 ± 0.1^{ab}	0.5 ± 0.1^a	1.2 ± 0.3^b
Gpr120	1.0 ± 0.2^a	3.6 ± 1.7^b	0.6 ± 0.1^a	1.1 ± 0.2^{ab}
Ppargc1	1.0 ± 0.3^a	1.2 ± 0.3^a	0.9 ± 0.2^a	0.4 ± 0.1^a
Dio2	1.0 ± 0.3^a	2.8 ± 1.1^a	2.0 ± 0.7^a	2.8 ± 1.3^a
Cidea	1.0 ± 0.2^a	1.1 ± 0.4^a	0.5 ± 0.1^{ab}	0.5 ± 0.1^a
Prdm16	1.0 ± 0.3^a	0.8 ± 0.2^a	0.4 ± 0.1^{ab}	0.4 ± 0.1^a
Cox8b	1.0 ± 0.4^a	1.7 ± 0.8^a	1.0 ± 0.4^a	0.9 ± 0.3^a
Cn1	1.0 ± 0.2^a	0.4 ± 0.1^b	0.8 ± 0.6^{ab}	0.8 ± 0.3^{ab}

Supplemental Figure 1A

Supplemental Figure 1B

Supplemental Figure 2A

Supplemental Figure 2B

Supplemental Figure 3A

Supplemental Figure 3B

Supplemental Figure 3C

Supplemental Figure 3D