1

4   The bipolar-to-multipolar transition of dendrite extension of cerebellar granule neurons

5   requires MTCL2 to link the Golgi apparatus to the microtubule cage.

7   Running title: MTCL2 promotes multipolar extension of dendrites.

10   Mari Minekawa and Atsushi Suzuki*

Molecular Cellular Biology Laboratory, Yokohama City University Graduate School of

Medical Life Science, 1-7-29 Suehiro-cho, Tsurumi-ku, Yokohama 230-0045, Japan.

* Corresponding author: Atsushi Suzuki; email: abell@yokohama-cu.ac.jp

Number of pages: 38

Number of Figures: 7,

Number of tables: 0

Number of words for abstract: 251

Number of words for introduction: 653

Number of words for discussion :   1085

Conflict of interest: The authors declare no competing financial interests.

Acknowledgments: We thank Dr. Tomoko Satake for her initial support in the

establishment of the primary cerebellar granule cell culture. This research was

supported by KAKENHI (to A.S.) (Grant Numbers 19H03228 and 22H02621) of the

Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan.

Author contributions: Designed research: A.S., M.M., Performed research: M.M.,

A.S., Analyzed data: M.M. A.S, Wrote the paper M.M., A.S., Supervision: A.S.;

Funding acquisition: A.S.

Keywords: MTCL2, Golgi apparatus, cerebellar granule neurons, dendrite, polarity

1   **Abstract**

The dynamic regulation of neuronal polarity is essential for the formation of neural

networks during brain development. Primary cultures of rodent neurons recapitulate

several aspects of this polarity regulation, providing valuable insights into the molecular

mechanisms underlying axon specification, dendrite formation, and neuronal migration.

However, the process by which the preexisting bipolarity of migrating neurons is

disrupted to form multipolar dendrites remains to be elucidated. In this study, we

demonstrate that MTCL2, a microtubule-crosslinking protein associated with the Golgi

apparatus, plays a crucial role in this type of polarity transformation observed during the

differentiation of cerebellar granule neurons (CGNs). MTCL2 is highly expressed in

CGNs and gradually accumulates in dendrites as the cells develop polarity. MTCL2

knockdown resulted in the generation of longer and fewer dendrites by suppressing the

bipolar-to-multipolar transition of dendrite extension observed in the normal

polarization process. During this process, the Golgi apparatus shifts from the base of the

preexisting bipolar neurites to the lateral or apical side of the nucleus in the cell body.

There, it forms a close association with the microtubule cage that wraps around the

nucleus. The resulting upward extension of the Golgi apparatus is tightly coupled with

the randomization of its position in the x-y plane. Knockdown and rescue experiments

demonstrated that MTCL2 promotes these Golgi position changes in a microtubule- and

Golgi-binding activity-dependent manner. These results suggest that MTCL2 promotes

the development of multipolar short dendrites by sequestering the Golgi apparatus into

the microtubule cage from the base of the preexisting neurite.

Significance Statement

Previous studies have shown that neurons polarize the localization of the Golgi apparatus, thereby facilitating the development of asymmetrically elongated dendrites. In contrast, the present study demonstrated that to symmetrically extend multiple dendrites, cerebellar granule neurons sequester the Golgi apparatus within a microtubule cage surrounding the nucleus. This represents a novel mechanism through which neurons suppress polarized vesicle transport. These findings also reveal an entirely novel function of microtubule cages, which have only been reported for their role in neuronal cell migration. The present study also revealed the involvement of the novel microtubule-regulating protein MTCL2, which links the Golgi apparatus to microtubules, and the LINC complex, which links microtubules to the nuclear membrane, in this process.

Introduction

Neurons are highly polarized cells in the human body that develop two types of functionally distinct processes: a long, thin axon and short dendrites. To establish elaborate neuronal circuits in the brain, developing neurons gradually establish neuronal polarity while migrating to appropriate regions.

Neuronal polarization is intrinsically programmed and can be recapitulated *in vitro* to some extent, even without specific cell-to-cell contacts or extracellular matrix molecules (Tahirovic and Bradke, 2009). Rodent hippocampal neurons are the most intensively studied model system for investigating neuronal polarity (Dotti et al., 1988; Fletcher and Banker, 1989). Cells isolated from late embryos first develop a multipolar morphology with several short neurites that exhibit rapid elongation and retraction. Remarkably, one of these initial neurites begins to grow rapidly and becomes a thin, long axon 1-2 days after plating. The remaining short neurites then develop into dendrites. Most studies have focused on the initial event through which neurons break the initial multipolarity, as similar axon specification processes have been observed *in vivo* in the developing neocortex (Miyata et al., 2004; Noctor et al., 2004; Takano et al., 2015).

Dissociation cultures of cerebellar granule neurons (CGNs) reflect the characteristic developmental process of this type of neurons *in vivo,* and provide an additional model for studying neuronal development (Tahirovic and Bradke, 2009; Powell et al., 1997; Zmuda and Rivas, 1998; Solecki et al., 2006). In contrast to hippocampal neurons, the initial development of CGNs isolated from early postnatal rodents is characterized by the formation of bipolar axons without the elongation of

multiple neurites. Subsequently, the cells transform the bipolar axon into a single T-shaped axon by either migrating perpendicular to the initial bipolar axis or by retracting one of the two axons. Finally, they extend short dendrites in multiple directions to complete their final morphology (Powell et al., 1997; Solecki et al., 2006; Tahirovic and Bradke, 2009; Zmuda and Rivas, 1998). Researchers initially employed this distinctive property of CGNs in culture to demonstrate the close correlation between the position of the centrosome and the Golgi apparatus with the site of axon elongation (Zmuda and Rivas, 1998). This culture system was then used to investigate the molecular mechanisms underlying axonogenesis (Arakawa et al., 2003; Oishi et al., 2021) and dendritogenesis (Huang et al., 2007; Litterman et al., 2011; Tan et al., 2013). However, no studies have focused on the unique polarity change, the transition from bipolar to multipolar, exhibited by CGNs, but not by hippocampal neurons. Consequently, the process by which CGNs disrupt preexisting bipolarity to extend multipolar dendrites remains unclear.

MTCL2, formerly known as SOGA1, is a recently identified microtubule-associated protein (Matsuoka et al., 2022). It dimerizes through its central coiled-coil region and interacts with microtubules (MTs) via its C-terminal region. The N-terminal portion of the coiled-coil region also exhibits additional activity to associate with the Golgi membrane through interactions with CLASPs and Giantin. These abilities enable MTCL2 to accumulate MTs around the Golgi apparatus, thereby promoting the lateral assembly of Golgi stacks and forming a compact Golgi ribbon. Using a wound-healing assay with a cultured cell line, Matsuoka et al. (2022) further suggested that these MTCL2 functions are essential for maintaining directional migration.

In this study, we demonstrated that MTCL2 is highly expressed in CGNs *in vivo*.

In cultured CGNs, MTCL2 depletion resulted in abnormally polarized cells with longer

and fewer dendrites. Several experiments, including live-cell imaging, revealed that

MTCL2 is necessary for the bipolar-to-multipolar transition of dendrite extensions, a

process observed during the normal development of CGNs. Immunofluorescence

analysis and knockdown/rescue experiments have revealed that MTCL2 sequesters the

Golgi apparatus from the base of the bipolar neurites to the MT cage that wraps around

the nucleus. The importance of the MT cage was also demonstrated by inhibiting the

LINC complex, which links the nuclear membrane to MTs. Taken together, these results

demonstrate MTCL2's novel role in promoting the bipolar-to-multipolar transition of

dendrite extensions by linking the Golgi apparatus and MTs.

<center>**Results**</center>

MTCL2 shows preferential enrichment in the dendrites of cerebellar granule neurons.

A previous study by Matsuoka et al. (2022) showed preferential expression of MTCL2 in the lung, testis, ovary, cerebrum, and cerebellum using western blotting of mouse tissue extracts. High expression of MTCL2 in the cerebellum is also evident in public mRNA expression databases for human and mouse organs (https://www.ebi.ac.uk/gxa/home and https://www.proteinatlas.org), as well as *in situ* hybridization data for the mouse brain (https://mouse.brain-map.org/) (Fig. S1A). We therefore performed an immunostaining analysis of MTCL2 in the mouse cerebellum and found that MTCL2 is preferentially expressed in granule cells and, to a lesser extent, in Bergmann glial cells, but not in Purkinje cells (Fig. 1A ; A. Suzuki, unpublished observation). The absence of MTCL2 in Purkinje cells is consistent with the public *in situ* hybridization data for the mouse brain (Fig. S1), whereas it is in sharp contrast to its paralogue, MTCL1, which is expressed in Purkinje cells and plays critical functions (Satake et al., 2017). Based on these results, in subsequent studies, we analyzed MTCL2 functions in cerebellar granule neurons (CGNs) using dissociated cultures of mouse CGNs (Krämer and Minichiello, 2010; Powell et al., 1997).

Immunofluorescence analysis of CGNs isolated and cultured for 5 days *in vitro* (DIV5) revealed strong anti-MTCL2 signals in the cell bodies and in some restricted neurites (Fig. 1B). The disappearance of these signals following MTCL2 knockdown (Fig. S2) indicates that they represent the true localization of endogenous MTCL2. Neurites with a strong MTCL2 signal showed a high degree of correlation with the

<center>8</center>

MAP2-positive dendrites (Fig. 1B). In fact, the intensities of MTCL2 and MAP2

staining on the neurites were highly correlated (Fig. 1C; Pearson's r = 0.83). However,

weak MTCL2 signals were also detected in MAP2-negative thin axons (Fig. 1B, C).

These results suggest that, while MTCL2 is not restricted to MAP2-positive dendrites, it

is preferentially enriched in these dendrites.

Next, we examined the timing of MTCL2 enrichment in dendrites during CGN

polarization. To this end, we compared the intensities and patterns of the

immunostaining for dephosphorylated tau (tau1), MAP2, and MTCL2 in CGNs cultured

for 1–5 days (Mandell and Banker, 1996). Consistent with previous reports, CGNs

extended tau1-positive short processes 1 day after seeding (Fig. 1D, arrowheads)

(Zmuda and Rivas, 1998). Most of these processes acutely elongated after DIV2 and

gradually became axon-like neurites with longer and thinner morphologies. Conversely,

MAP2 signals were very weak until DIV3, when they began to be detected in the short

neurites. In contrast, MTCL2 signals were strongly detected even at DIV1, at levels

comparable to those observed at DIV5 (~80%). This level of expression was maintained

thereafter, with slight decreases at DIV2 and 3 (Fig. 1D,E). High expression of MTCL2

in DIV1 CGNs was also confirmed by western blotting (Fig. 1F). MTCL2 signals in

CGNs at DIV1 and 2 were detected in cell bodies and in some tau1-positive initial

neurites (Fig. 1D, arrowheads). Around DIV3-4, however, when MAP2-positive

neurites appeared, MTCL2 gradually exhibited selective enrichment in these neurites

(Fig. 1D). These results suggest that MTCL2 enrichment in dendrites occurs during the

late stage of CGN polarization in parallel with MAP2 expression.

Depletion of MTCL2 resulted in the development of fewer and longer dendrites.

These results suggest a possibility that MTCL2 plays an important role in the

dendritic development of CGNs. To investigate this possibility, we examined the effects

of MTCL2 knockdown in primary cultured CGNs. Eight hours after seeding, we

transfected a plasmid expressing shRNA and EGFP into the cells (Fig. S2). As shown in

Fig. 2A, CGNs lacking MTCL2 appeared to differentiate normally into neurons with

long, thin axons at DIV5. Quantification of axon length and branching revealed no

significant differences between MTCL2-depleted cells and control cells (Fig. 2B; Fig.

S3A). Furthermore, MTCL2 knockdown did not prevent the appearance of MAP2-

positive dendrites in DIV6 CGNs (Fig. 2C). However, detailed analyses revealed that

MTCL2 knockdown resulted in fewer dendrites than control cells in terms of both the

number of dendrite tips (Fig. 2D) and bases (Fig. S3B). Additionally, the average

dendrite length of MTCL2-depleted CGNs (25.0 ± 11.5 µm) was greater than that of

control cells (15.4 ± 4.0 µm) (Fig. 2E). Analysis of the length distribution of the total

dendrites revealed that the increase in average length was primarily due to the existence

of long dendrites in the MTCL2-depleted cells, which were rarely observed in the

control cells (Fig. S3C). For instance, the maximum dendritic length of individual

CGNs was 35.7 ± 11.5 µm on average for control cells and 54.0 ± 21.6 µm for the

MTCL2-depleted cells (Fig. S3D). The proportion of dendrites longer than 40 µm in the

total number of dendrites was $3.14 \pm 0.13\%$ in control cells and $14.36 \pm 0.10\%$ in

MTCL2-depleted cells (Fig. 2F). These changes in dendrite numbers and length

distribution in MTCL2-depleted cells were reversed by expressing RNAi-resistant

mouse MTCL2 (Fig. 2D-F; Fig. S3B-D). These results suggest that MTCL2 depletion

leads to abnormal CGN development, resulting in fewer longer dendrites.

In MTCL2-depleted CGNs, dendrite extensions are biased in the direction opposite to axon elongation.

While characterizing MTCL2-depleted CGNs, we observed that their dendrite extensions were biased in the opposite direction of axon elongation (see Fig. 2A,C). To confirm this tendency, we quantitatively estimated the direction of dendrite extension in the CGNs at DIV6. Angles were measured based on the positions of the base of each dendrite (MAP2-positive neurite) and an axon (MAP2-negative, thin neurite) on the cell body (see Fig. 3A for an illustration). Consistent with previous reports (Powell et al., 1997; Zmuda and Rivas, 1998), control cells extended dendrites evenly in all directions (Fig. 3B). Conversely, dendrite extension in MTCL2-depleted CGNs was weakly or strongly biased toward the same or opposite directions of axon elongation, respectively, when focusing on dendrites longer than 15 μm (Fig. 3B). The directional bias (r), defined as the ratio of the number of dendrites extending forward or backward versus left or right, was 1.22 ± 0.05 and 1.66 ± 0.10 for control cells and MTCL2-depleted cells, respectively (Fig. 3C). However, expressing RNAi-resistant MTCL2 in MTCL2-depleted cells suppressed the directional bias of dendritic extension to 1.01 ± 0.03 (Fig. 3B,C). These results suggest that MTCL2-depleted CGNs exhibit bipolarity in relation to the direction of axon elongation, extending fewer and longer dendrites in that direction (Fig. S3E).

MTCL2 is necessary for the transition of dendrite extensions from bipolar to multipolar.

Many studies have examined dendrite growth in CGNs using *in vitro* culture systems (Huang et al., 2007; Litterman et al., 2011; Tan et al., 2013; Kawaji et al.,

2004). However, to our knowledge, no study has examined the polarity of dendrite extensions during CGN differentiation. Thus, we next examined whether the bipolar dendrite extensions observed in MTCL2-depleted cells corresponded to an intermediate state of normal CGN differentiation or an aberrant state of the cells that is not observed in normal CGN differentiation. To this end, we assessed the polarity of CGN dendrite extensions at the middle stage (DIV3) when MAP2-positive dendrites begin to appear (Fig. 1D; Fig. 4). Immunostaining with the axon marker Ankyrin G and the dendrite marker MAP2 revealed that all of the CGNs that we analyzed had a single, long axon and multiple dendrites (an average of 3.4 dendrites per cell) (Fig. 4A). This indicates that these cells have exited the initial bipolar stage with two axons and entered a stage in which they elongate multiple dendrites. In these DIV3 CGNs, the directions of the dendrite extensions were strongly biased toward the same axis as the axon elongation, as was observed in MTCL2-depleted cells (Fig. 4A,B; Fig. S4A). Interestingly, DIV3 CGN dendrites were fewer but longer than DIV5 CGN dendrites (Fig. 4C; Fig. S4B). This also corresponded to the situation observed in MTCL2-depleted cells. These results support the notion that MTCL2 knockdown arrests the bipolar-to-multipolar transition of dendrite extension, which normally occurs during CGN differentiation.

To directly examine this notion, we subjected GFP-expressing CGNs, with or without MTCL2 knockdown, to live imaging at DIV3. We recorded the morphological changes every 2 min for 48 h (Fig. 4E,F; Movie 1 and 2). Consistent with the aforementioned results, the cells exhibited extended bipolar neurites and a thin, long axon (arrowheads) at the beginning of the recording (Fig. 4E,F). The control cells exhibited a thick neurite extending in the direction opposite to the axon (arrow in Fig. 4E), which remained relatively stable for a few hours. In addition, several short neurites

grew and shrank dynamically in the same direction as axon elongation. A few hours

after recording began, the thick neurite started shrinking (6.2-9.2 h). Simultaneously,

short dynamic neurites began to elongate from various sites on the cell body (after 8.2h),

and the cell eventually acquired a multipolar morphology with short dendrites (Fig. 4E;

Fig. S4D; Movie 1). At this stage, the initial thick neurite became indistinguishable

from the other neurites in terms of length, thickness, and dynamic nature. These results

support the notion that CGN dendrites undergo a bipolar-to-multipolar transition during

their differentiation. Similar shrinkage of an initial bipolar neurite (arrow) (13.2-17.2 h)

and *de novo* elongation of unstable short dendrites (15.2-21.2 h) were observed in

MTCL2-depleted CGNs (Fig. 4F). However, the cells could not steadily extend short

dendrites longer than the width of their cell bodies (~ 15μm) in multiple directions.

Instead, these cells finally elongated new, long, stable dendrites along the initial bipolar

axis (25.2 h ~) (Fig.4F; Fig. S4D; Movie 2). These results suggest that MTCL2 is

essential for disrupting the bipolarity of dendrite extension that occurs during normal

CGN differentiation.

MTCL2 promotes the bipolar-to-multipolar transition of dendrite extension

through its MT- and Golgi-binding activity.

The localized positions of the Golgi apparatus are closely associated with the

development of various neurite processes, including axons, dendrites, and leading

processes involved in neuronal migration (Horton et al., 2005; Matsuki et al., 2010;

O'Dell et al., 2012; Tanabe et al., 2010; Zmuda and Rivas, 1998). Because MTCL2 is a

microtubule-regulating protein that binds to the Golgi membrane (Matsuoka et al.,

2020), we next assessed the possibility that MTCL2 facilitates the bipolar-to-multipolar

transition of dendrite extension polarity by affecting Golgi localization. To this end, we first examined the localization of the Golgi apparatus in MTCL2-depletecd cells (Fig. 3D-F). Consistent with a previous report (Zmuda and Rivas, 1998), the Golgi apparatus did not exhibit any specific localization in DIV6 CGNs. However, in MTCL2-depleted CGNs, the Golgi position was clearly biased toward the direction opposite to axon elongation, similarly to dendrite elongation. This abnormality was randomized again by expressing RNAi-resistant MTCL2. Interestingly, CGNs at DIV3 exhibited a similarly biased Golgi localization (Fig. 4D; Fig. S4C). These results support the possibility that MTCL2 disrupts the bipolarity of dendrite extensions by affecting the Golgi position.

To evaluate this possibility, we examined the ability of MTCL2 mutants to rescue the defects exhibited by MTCL2-depleted CGNs (Fig. 2; Fig. 3; Fig. S3). Unlike wild-type MTCL2, MTCL2 mutants lacking MT- (ΔKR) or Golgi- (4LA) binding activity (Matsuoka et al., 2022) failed to rescue the biased polarity of dendrite elongation and the Golgi localization (Fig. 3B,C and E,F). In addition, they did not restore the reduced number of dendrites observed in MTCL2-depleted CGNs (Fig. 2D; Fig.S3B). The only exception was that the 4LA mutant, but not the ΔKR mutant, exhibited substantial rescue activity for dendrite length (Fig. 2E,F; Fig. S3C,D). Consequently, MTCL2-depleted CGNs expressing RNAi-resistant MTCL2 4LA developed a few short dendrites with biased polarity (Fig. S3E). This indicates that the sustained destabilization of a bipolar long neurite, one of the two events observed in the bipolar-to-multipolar transition of dendrite extensions (Fig.4E; Movie 1), does not require MTCL2's Golgi-binding activity. In any case, these results suggest that the MT- and Golgi-binding activities of MTCL2 are crucial for disrupting the preexisting bipolarity and inducing multidirectional dendrite development in CGNs.

During the late stage of CGN polarization, the Golgi apparatus exhibited upward extension along the microtubule cage surrounding the nucleus.

To investigate how MTCL2 promotes the bipolar-to-multipolar transition of dendrite extension, we next examined the subcellular localization of MTCL2 in greater detail in relation to MTs and the Golgi apparatus.

Consistent with our previous observations in HeLa-K cells, basal confocal images of mature CGNs (DIV5) revealed intermittent granular MTCL2 stainings on MT bundles passing through MAP2-positive dendrites (Fig. S5A). In addition, numerous MTCL2 signals were observed in the cell bodies (Fig. 5A; Fig. S5A). These signals frequently colocalized with an MT network in the cell bodies that was efficiently stained by anti-tyrosinated but not anti-acetylated tubulin antibodies (Fig. S5B,C). Serial analysis of z-stack confocal images (Fig. 5A,B) and their reconstructed z-sectional views (Fig. 5C; Movie 3) revealed that the MT network completely encircles the nucleus. This indicates that it corresponds to the MT cage, the presence of which has been reported in CGNs (Rivas and Hatten, 1995; Solecki et al., 2004; Umeshima et al., 2007; Xie et al., 2003). Preferential staining with anti-tyrosinated, but not anti-acetylated, tubulin antibodies corroborated this conclusion (Umeshima et al., 2007).

Interestingly, the Golgi apparatus in CGNs at DIV5 is often associated with the MT cage on the lateral or apical surfaces of the nucleus (Fig. 5A,B). Consequently, the Golgi apparatus was drawn toward the nucleus, extending its apical end toward the upward direction along the nuclear surface (Fig. 5C). In the most extreme cases, the entire apparatus was placed on top of the nucleus (see arrows in Fig. 5C; Movie 3). This differs greatly from the situation observed in CGNs at DIV2. At this stage, the Golgi

apparatus was found in the basal position, biased toward the base of one of the bipolar

processes, as previously reported (Fig. 5D; Fig. S6A,B) (Zmuda and Rivas, 1998).

Some DIV2 cells contained cortical MTs underneath the apical membranes of their cell

bodies (Fig. 5D, see the arrowhead in Fig. S6B). However, these MTs were not

organized into a dense, isotropic network. Rather, they tended to align parallel to the

preexisting bipolar axis (Fig. 5D). Even at the CGNs at DIV3, where isotropic network-

like MT structures were detected (Fig. 5D, Fig. S6A), the upward shift of the Golgi

apparatus was not evident (Fig. 5D; Fig. S6B). Quantitative analysis demonstrated that

the cell heights of the CGNs at DIV3 and DIV5 did not differ significantly (Fig. S7A,B).

However, the height of the apical end of the Golgi apparatus relative to the

corresponding cell height increased as the polarization proceeded (Fig. 5E). These

results suggest that the final CGN polarization is tightly coupled with the sequestration

of the Golgi apparatus from the base of the bipolar neurites into an MT cage around the

nucleus. This is consistent with a previous report that the Golgi apparatus in mature

CGNs is not localized to the base of any neurites, including axons and dendrites (Zmuda

and Rivas, 1998)

MTCL2 promotes the sequestration of the Golgi apparatus into the MT cage.

Importantly, in the CGNs at DIV5, many MTCL2 signals substantially overlapped

with those of the Golgi and MT signals around the nucleus (Fig. 5A-C). Since both the

MT- and the Golgi-binding activities of MTCL2 were necessary to disrupt the Golgi's

biased localization in the x-y plane (Fig. 3E,F), these results suggest that MTCL2 plays

a crucial role in the Golgi's upward extension by facilitating its association with the MT

cage. To verify this possibility, we reexamined the positions of the Golgi apparatus in

MTCL2-depleted cells in the z direction (Fig. 6). As expected, the positions of the Golgi

apparatus were significantly lowered in MTCL2-depleted cells (Fig. 6A). Quantitative

analysis demonstrated that MTCL2 knockdown did not change the cell height (Fig.

S7C). However, the apical ends of the Golgi apparatus were lowered to a level similar

to that observed in bipolar CGNs at DIV3 (Fig. 6C). This defect was restored by

expressing RNAi-resistant MTCL2 wild type but not by expressing MTCL2 ΔKR or

4LA (Fig. 6C). These results support the idea that MTCL2 extends the Golgi apparatus

upward along the nucleus by connecting it to the MT cage (Fig. 7A).

The LINC complex activity is necessary for sequestering the Golgi apparatus into

the MT cage.

The initial observation of the MT cage in CGNs was 30 years ago (Rivas and Hatten,

1995), but its molecular basis remains unclear. The LINC complex is a potential

candidate associated with the MT cage because it links the nuclear membrane to MTs

(Chang et al., 2015; Kengaku, 2018). Recent studies have revealed the indispensable

role of the LINC complex in nuclear rotation, deformation, and translocation during

CGN migration (Wu et al., 2018; Zhou et al., 2024). Thus, we investigated the role of

the LINC complex in the bipolar-to-multipolar transition of dendrite elongation to

confirm the involvement of the MT cage in this process.

The Nesprin protein is a component of the LINC complex located in the outer nuclear

membrane. It interacts with MT motors via its N-terminal cytoplasmic stretch. Another

component is the inner nuclear membrane protein SUN, which interacts with the nuclear

lamina and chromatin within nucleus via its N-terminus (Fig. 7A). Nesprins interact

with SUN proteins through their C-terminal KASH domains in the perinuclear space

(Fig. 7A). Consequently, the overexpression of a Nesprin-1 mutant lacking the cytoplasmic stretch (KASH1) inhibits LINC complex activity by dominantly interfering with the interaction between endogenous Nesprin-1/2 and SUN proteins (Stewart-Hutchinson et al., 2008; Wu et al., 2018). When expressed in primary cultured CGNs, GFP-KASH1 localized to the nuclear membrane, as expected (Fig. 7B). Close inspection revealed a network-like distribution of GFP-KASH1 on the nuclear membrane, likely reflecting the underlying nuclear lamina pattern via SUN proteins (Fig. 7C; Fig. S8A). These cells exhibited the formation of the MT cage itself (Fig. 7C; Fig. S8B) and the colocalization of MTCL2 with the Golgi apparatus and the MT cage (Fig. 7B,C; Fig. S8A). However, the randomization of the Golgi position in the x-y plane (Fig. 7D; Fig. S8C) and the upward extension of the Golgi apparatus in the z direction (Fig. 7B,E) were specifically suppressed in these cells, as observed in MTCL2-depleted CGNs.

Importantly, when we examined the polarity of dendrite extensions, cells expressing GFP-KASH1 exhibited a reduced number of dendrites with a biased extension direction to the preexisting bipolar axis (Fig. 7F,G; Fig. S8D). As these cells did not exhibit a significant increase in dendritic length (Fig. 7H; Fig. S8E), they perfectly mimicked the phenotypes exhibited by MTCL2-depleted CGNs expressing the MTCL2 4LA mutant, which lacks Golgi-binding activity (Fig. S3E, Fig. S8F). These results support the hypothesis that the MTCL2-mediated sequestration of the Golgi apparatus within the MT cage plays a pivotal role in the multipolar development of dendrites observed in CGNs during the final stage of their polarization.

Discussion

Previously, we reported that MTCL2 is a unique MT-regulating protein with Golgi-binding activity (Matsuoka et al., 2022). In this study, we demonstrated that MTCL2 is highly expressed in CGNs and plays an indispensable role in the later stage of polarization, during which the dendrite extensions undergo a bipolar-to-multipolar transition. We provided data indicating that MTCL2 facilitates this transition by mediating the sequestration of the Golgi apparatus into the MT cage surrounding the nucleus.

The unique polarization process observed in isolated CGN cultures was first documented in 1997 (Powell et al., 1997;Kollins et al., 1999; Zmuda and Rivas, 1998). These pioneering studies also demonstrated the significance of the Golgi apparatus's positioning in the development of initial bipolar axons (Zmuda and Rivas, 1998). However, the mechanism by which the initial bipolarity is disrupted to result in multipolar dendrite extension remains to be elucidated. In this paper, to understand the molecular basis of the MTCL2 function in CGNs, we focused on this issue and demonstrated that the initial extension of multiple dendrites exhibits bipolarity (Fig. 4A, B). The live imaging analysis of CGNs from DIV3 to DIV5 not only corroborated this conclusion but also revealed that this bipolarity is disrupted through two distinct processes: destabilization of the stable dendrites elongating opposite to an axon, and the *de novo* generation of unstable dendrites from the multiple sites of the cell body (Fig. 4E,F). We further found that MTCL2 knockdown impeded both processes, resulting in a reduced number of longer dendrites that exhibit an extension bias directed in the opposite direction of an axon (Fig. 3A-C). The rescue experiments further indicated that

the MT-binding activity of MTCL2 was required for both processes, whereas the Golgi-

binding activity was dispensable for the sustained destabilization of a bipolar long

neurite and only required for the persistent generation of multipolar dendrites from the

cell body. Actual CGNs *in vivo* undergo a second bipolar phase between the initial

bipolar phase (characterized by two axons) and the final complex morphology

(characterized by multipolar dendrites). During the second bipolar phase when CGNs

migrate deep into the internal granular layer along the radially aligned Bergmann glial

cells, by extending a leading process in the migration direction and drawing the axon

that elongates to the opposite direction (trailing process) (Gao and Hatten, 1993; Solecki

et al., 2006). Importantly, Kawaji et al. (2004) demonstrated that the leading process

observed in this phase exhibits dendritic properties and ultimately contracts, resulting in

differentiation into one of the multipolar dendrites. Therefore, it is highly plausible that

the bipolarity observed in DIV3 CGNs with single axons mimics that exhibited by

migrating CGNs. Furthermore, the stable dendrites that elongate in the direction

opposite to the axon likely correspond to the radial leading process of CGNs *in vivo*.

Thus, the present results support the notion that MTCL2 is required to disrupt the

bipolarity of CGNs established during migration.

At present, the mechanism by which the MT-binding activity of MTCL2 is required

for the destabilization of long bipolar dendrites observed at DIV3 remains unclear.

Further research is necessary to elucidate this issue. However, the present study

provides data indicating how the MT- and Golgi-binding activities of MTCL2 promote

the multipolar development of *de novo* dendrites. Our novel findings, which include the

observation of upward extension of the Golgi apparatus of CGNs in the later stage of

polarization along the nucleus in an MTCL2-dependent manner (Fig. 5), support the

proposal that MTCL2 facilitates the bipolar-to-multipolar transition of dendrite extension by mediating the association between the Golgi apparatus and the MT cage. This, in turn, promotes the sequestering of the Golgi apparatus to the MT cage from the base of preexisting bipolar dendrites, thus inhibiting the polarized vesicle transport required for polarized dendrite extension (Fig. 7A). This hypothesis was further substantiated by the finding that the inhibition of the normal formation of the LINC complex also resulted in the suppression of the bipolar-to-multipolar transition of the dendritic extension. Notably, the inhibition of the LINC complex did not affect the formation of the MT-cage; rather, it suppressed the upward extension of the Golgi apparatus. The results indicate that the pulling forces generated by the MT motors may contribute to the attraction of the Golgi apparatus to the MT cage, thereby inducing its upward extension (Fig.7A).

The critical role of the Golgi apparatus in dendrite development has been extensively studied (Horton et al., 2005; Nakagawa and Iwasato, 2023; Tanabe et al., 2010; Wu et al., 2015). However, most of these studies have focused on the analysis of pyramidal neurons located in the cerebral cortex and hippocampus, as well as on Purkinje cells in the cerebellum. These neurons all develop thick, apically polarized primary dendrites, also known as apical dendrites. These studies demonstrate that polarized Golgi localization is required for asymmetric dendrite growth. In contrast, the present study distinguished itself by analyzing CGNs that exhibit near-symmetric short dendrites with uniform length, and demonstrated that sequestering the Golgi apparatus from the base of any dendrite is essential for developing multipolar dendrites devoid of any asymmetry. Furthermore, during this analysis, the MT cage was identified as the target structure to which the Golgi apparatus was sequestered. Since the MT cage had

previously been studied only in terms of its role in nucleokinesis observed in neural cell

migration (Rivas and Hatten, 1995; Solecki et al., 2004; Xie et al., 2003), the present

finding reveals an unexpected function of this higher-order MT structure. The molecular

mechanism suggested in this study may be specific to CGNs. Indeed, the presence of

the MT cage has primarily been described in CGNs, but not reported conspicuously in

other neurons (Rivas and Hatten, 1995; Solecki et al., 2004; Umeshima et al., 2007).

However, many nonpyramidal neurons such as GABAergic interneurons lack apical

dendrites and develop symmetrical, uniform dendrites (Horton et al., 2005; Müller et al.,

1984). In addition, some pyramidal neurons remodel their asymmetric dendrites by

withdrawing their apical dendrites and acquiring a multipolar morphology (Koester and

O'Leary, 1992). Therefore, neurons other than CGNs may utilize the MT-cage-

dependent regulation of the Golgi position, at least in part.

In summary, the results of this study shed light on how CGNs extend dendrites in

multiple directions. Our results highlight the significance of the MT cage surrounding

the nucleus for removing the Golgi apparatus from the base of preexisting neurites. The

critical importance of the MTCL2-mediated association between the Golgi apparatus

and the MT cage, as well as that of the LINC complex-mediated fixation of MT motors

to the nuclear membrane, was also suggested.

1   **Materials and Methods**

Animals

All protocols were approved by and performed in accordance with the guidelines

established by the Institutional Animal Care and Use Committee of Yokohama City

University, Medical Life Science (Approval No. T-A-19-006 and T-A-22-007).

Antibodies

The primary antibodies used in this study were as follows: anti-SOGA1 to detect

MTCL2 (HPA043992), anti-tau1 (MAB3420), anti-calbindin (C9848), anti-Ankyrin G

(MABN466) from Merk Sigma-Aldrich ; anti-MAP2 (sc-32791), anti-α-tubulin

(DM1A: sc-32293), anti-acetylated tubulin (sc-23950), anti-tyrosinated tubulin (sc-

53029), anti-GFP (sc-9996) from Santa Cruz Biotechnology; anti-GM130 (610822) and

anti-GS28 (611184) from BD Transduction Laboratories; anti-GAPDH (5G4) from

HyTest Ltd; and anti-NeuN (MAB377) from Merck Millipore.

Primary Neuron Culture and Transfection

Primary cerebellar granule neurons (CGNs) were isolated from P6 neonatal mice and

prepared as previously described (Krämer and Minichiello, 2010). Neurons were plated

on coverslips coated with poly-D-lysin (0.1 mg/mL; Merk Sigma-Aldrich or Thermo

Fisher Scientific) at a density of 180,000 cells/well, 300,000 cells/well, or 360,000

cells/well (24 well plates) and cultured for 8 h in Dulbecco's modified Eagle medium

(DMEM) (Life Technologies Corporation) containing 10% fetal bovine serum,

penicillin (100 U/ml), streptomycin (0.1 mg/ml), 3.5 mM glutamine, and 0.9 mM

sodium pyruvate (nacalai) under controlled temperature and CO_2 conditions (37℃, 5%).

Then, cells were subjected to plasmid transfection using a CalPhos™ Mammalian

Transfection Kit (Takara) according to the manufacturer's instruction for 8~16 h,

followed by incubation with the original medium supplemented with 0.01 mM Ara-C

(Tokyo Chemical Industry Co., Ltd.) and 20 mM KCl until fixation.

Plasmids

For knockdown experiments, pEB-super-GFP vectors simultaneously expressing EGFP

and appropriate shRNA were first constructed (Masuda-Hirata et al., 2009). Then, the

EBNA1 cDNA sequence and the associated CMV promoter were removed to reduce

vector size. For mouse MTCL2 knockdown, the RNAi target sequence 5′-

gacggtcagtgtaggtcttca-3′ was inserted under the H1 promoter as a hairpin

oligonucleotide sequence in a tail-to-tail fashion (Satake et al., 2017). The control

shRNA sequence used was 5′-cagtcgcgtttgcgactgg-3′. To generate RNAi-resistant

mouse MTCL2 expression vectors, silent mutations (a̲ ac̲t gt̲a t̲c̲a̲ gtc ggc̲ tt̲g cag a) in

the RNAi target sequence were introduced in the cDNAs of V5-tagged wild-type or

mutant mouse MTCL2 (4LA or ΔKR) subcloned in the pCAGGS expression vector

(Matsuoka et al., 2022). In rescue experiments, the shRNA expression vector for mouse

MTCL2 and appropriate expression vectors for RNA-resistant mouse MTCL2 wild type

or mutant MTCL2 were co-transfected into CGNs. To construct a GFP-KASH1

expression vector, a human Nesprin-1 cDNA fragment encoding the C-terminal region

(8726-8799 a.a.) was amplified by RT-PCR from total RNA from HeLa-K cells and

cloned into pEGFP-C1 at BglII/BamHI sites (Stewart-Hutchinson et al., 2008). The

resulting GFP-KASH1 cDNA fragment was subcloned into the pCAGGS vector to

change the promoter.

Immunocytochemistry

The CGNs were fixed in 4% paraformaldehyde (PFA) at room temperature for 15 min

at 37℃. After fixation and blocking, samples were incubated with appropriate primary

antibodies diluted in TBST (10 mM Tris–HCl pH 7.5, 150 mM NaCl, 0.01% [v/v]

Tween 20) containing 0.1% (w/v) BSA overnight at 4 °C. After washing with PBST,

samples were incubated with the appropriate secondary antibodies conjugated with

Alexa Fluor 488, 555, or 647 (Life Technologies Corporation) for 60 min at room

temperature. Dilutions of the used antibodies are as follows: anti-SOGA1 (1/2000), anti-

α-tubulin (1:1000), anti-acetylated tubulin (1:2000), anti-tyrosinated tubulin (1:2000),

anti-GM130 (1:1000), anti-GS28 (1:300), anti-GFP (1:2000), anti-MAP2 (1:1000), anti-

tau1 (1:1000), and anti-Ankyrin G (1:1000). All secondary antibodies were used at a

1:2000 dilution. The nuclei were counterstained with 4', 6-diamidino-2-phenylindole

(MBL, Japan) at a 1:2000 dilution in PBST during the final wash. For image acquisition,

samples on coverslips were mounted onto glass slides in Prolong Diamond Antifade

Mountant (Thermo Fisher Scientific).

Immunohistochemistry

Mice under deep isoflurane anesthesia were fixed by transcardial perfusion with 4%

paraformaldehyde (PFA) in 0.1 M PB (0.1 M NaH_2PO_4 and 0.1 M Na_2HPO_4, pH 7.4).

Brains were removed from the skull, post-fixed for 3–5 h at room temperature, and

sectioned at 50-μm thickness using a Vibratome (VT1200S; Leica Microsystems,

Wetzlar, Germany). Following permeabilization with 0.1% Triton-X-100 in 0.1 M PB

with gentle shaking for 30 min, cerebellar vermis slices were blocked with 10% calf

serum in TBST for 30 min at room temperature. Immunofluorescence staining was

performed as previously described.

Image acquisition and processing

Most images were acquired using a Leica SP8 laser scanning confocal microscope

equipped with an HC PL APO 63x/1.40 Oil 2 objective using the Hybrid Detector in

photon-counting mode. To obtain super-resolution images, HyVolution2 imaging was

performed on the same system using Huygens Essential software (Scientific Volume

Imaging). Single confocal images are shown unless otherwise indicated. The same

confocal system was used to obtain time-lapse images, except that an HC PL APO

20x/0.75 objective was used. Cells seeded onto 35-mm glass-bottom dishes and

subjected to appropriate transfection were placed in a closed heat-controlled chamber

(Leica) with a CO_2 supply system (Tokken) on the microscope stage for 2 days under

autofocus control. Wide-view or low-resolution images for quantification (Figs. 2, 3,

and 4) were obtained using an AxioImager ZI microscope (Carl Zeiss, Oberkochen,

Germany) equipped with a Plan-Apochromat 40×/0.95 objective and an Orca II CCD

camera (Hamamatsu Photonics, Shizuoka, Japan). To obtain images of brain slices and

to measure the heights of cells and the apical end of the Golgi apparatus, an AxioImager

Z1 microscope equipped with a CSU10 disc confocal system (Yokogawa Electric

Corporation, Tokyo, Japan) and a 20x/1.4 NA or 100x/1.46 NA Plan-apochromat

objective was used. Images were acquired using MetaMorph software (Molecular

Devices, Sunnyvale, CA) and processed using ImageJ software to obtain appropriate

brightness and contrast.

Quantification

To quantify the enrichment of MTCL2 and MAP2 in CGN neurites, immunofluorescence signal intensities of anit-MTCL2 and anti-MAP2 staining were measured in ROIs of 1 μm^2 circles placed on anti-acetylated tubulin- or tau1-positive neurites. To analyze the morphology of CGNs, the positions of bases, tips, or branching points of neurites (axons or dendrites) were marked manually, and their lengths and numbers of tips were automatically quantified using the NeuronJ plugin for ImageJ. Essential functions of ImageJ were used to manually analyze the polarity of the dendrite extension direction, Golgi position in the x–y plane, and number of bases of dendrites. The dendritic lengths of CGNs subjected to live imaging were also analyzed manually. The cell height and position of the Golgi apical end were determined from the focal plane at which the fluorescent signals of tyrosinated tubulin and GS28 disappeared in z-stack images taken at an interval of 0.3 μm. The 3D video shown in Movie 3 was created using software (LAS X) associated with the Leica SP8 laser scanning confocal system.

Cell extraction and western blotting

CGN extracts were prepared by adding RIPA buffer (25 mM Tris/HCl [pH 7.5], 150 mM NaCl, 1% NP40, 1% deoxycholic acid, 0.1% SDS) containing a protease inhibitor cocktail (Merk Sigma-Aldrich, P8340), followed by brief sonication and centrifugation (15,000 × g, 15 min). Samples were separated using SDS-PAGE and transferred to polyvinylidene fluoride membranes. Blots were incubated in blocking buffer containing 5% (w/v) dried skim milk in PBST (8.1 mM $Na_2HPO_4.12H_2O$, 1.47 mM KH_2PO_4, 137

mM NaCl, 2.7 mM KCl, and 0.05% Tween 20), followed by overnight incubation with

anti-SOGA1 or anti-GAPDH antibodies diluted in blocking buffer. The diluted anti-

SOGA1 and anti-GAPDH antibodies were 1:1000 and 1:5000, respectively. Blots were

then exposed to horseradish peroxidase (HRP)-conjugated secondary antibodies (GE

Healthcare) diluted in blocking buffer for 60 min at room temperature and washed again.

Blots were visualized using an Immobilon Western Chemiluminescent HRP Substrate

(Millipore) or ECL western blotting detection system (GE Healthcare).

Chemiluminescence was quantified using an ImageQuant LAS4000 Luminescent Image

Analyzer (GE Healthcare).

References

Arakawa, Y., Bito, H., Furuyashiki, T., Tsuji, T., Takemoto-Kimura, S., Kimura, K., Nozaki, K., Hashimoto, N. and Narumiya, S. (2003). Control of axon elongation via an SDF-1α/Rho/mDia pathway in cultured cerebellar granule neurons. *Journal of Cell Biology* **161**, 381–391.

Chang, W., Worman, H. J. and Gundersen, G. G. (2015). Accessorizing and anchoring the LINC complex for multifunctionality. *J Cell Biol* **208**, 11–22.

Dotti, C. G., Sullivan, C. A. and Banker, G. A. (1988). The establishment of polarity by hippocampal neurons in culture. *J Neurosci* **8**, 1454–1468.

Fletcher, T. L. and Banker, G. A. (1989). The establishment of polarity by hippocampal neurons: The relationship between the stage of a cell's development in situ and its subsequent development in culture. *Dev Biol* **136**, 446–454.

Gao, W. Q. and Hatten, M. E. (1993). Neuronal differentiation rescued by implantation of weaver granule cell precursors into wild-type cerebellar cortex. *Science (1979)* **260**, 367–369.

Horton, A. C., Rácz, B., Monson, E. E., Lin, A. L., Weinberg, R. J. and Ehlers, M. D. (2005). Polarized secretory trafficking directs cargo for asymmetric dendrite growth and morphogenesis. *Neuron* **48**, 757–771.

Huang, J., Furuya, A. and Furuichi, T. (2007). Very-KIND, a KIND domain containing RasGEF, controls dendrite growth by linking Ras small GTPases and MAP2. *J Cell Biol* **179**, 539–552.

Kengaku, M. (2018). Cytoskeletal control of nuclear migration in neurons and non-neuronal cells. *Proc Jpn Acad Ser B Phys Biol Sci* **94**, 337–349.

Koester, S. E. and O'Leary, D. D. M. (1992). Functional classes of cortical projection neurons develop dendritic distinctions by class-specific sculpting of an early common pattern. *Journal of Neuroscience* **12**, 1382–1393.

Kollins, K. M., Powell, S. K. and Rivas, R. J. (1999). GPI-anchored human placental alkaline phosphatase has a nonpolarized distribution on the cell surface of mouse cerebellar granule neurons in vitro. *J Neurobiol* **39**, 119–141.

Krämer, D. and Minichiello, L. (2010). Cell culture of primary cerebellar granule cells. *Methods Mol Biol* **633**, 233–239.

Litterman, N., Ikeuchi, Y., Gallardo, G., O'Connell, B. C., Sowa, M. E., Gygi, S. P., Harper, J. W. and Bonni, A. (2011). An OBSL1-CUl7Fbxw8 ubiquitin ligase signaling mechanism regulates golgi morphology and dendrite patterning. *PLoS Biol* **9**, 17–19.

Mandell, J. W. and Banker, G. A. (1996). A spatial gradient of tau protein phosphorylation in nascent axons. *Journal of Neuroscience* **16**, 5727–5740.

Masuda-Hirata, M., Suzuki, A., Amano, Y., Yamashita, K., Ide, M., Yamanaka, T., Sakai, M., Imamura, M. and Ohno, S. (2009). Intracellular polarity protein PAR-1 regulates extracellular laminin assembly by regulating the dystroglycan complex. *Genes to Cells* **14**, 835–850.

Matsuki, T., Matthews, R. T., Cooper, J. A., Van Der Brug, M. P., Cookson, M. R., Hardy, J. A., Olson, E. C. and Howell, B. W. (2010). Reelin and Stk25 have

opposing roles in neuronal polarization and dendritic Golgi deployment. *Cell* **143**, 826–836.

Matsuoka, R., Miki, M., Mizuno, S., Ito, Y., Yamada, C. and Suzuki, A. (2022). MTCL2 promotes asymmetric microtubule organization by crosslinking microtubules on the Golgi membrane. *J Cell Sci* **135**,.

Miyata, T., Kawaguchi, A., Saito, K., Kawano, M., Muto, T. and Ogawa, M. (2004). Asymmetric production of surface-dividing and non-surface-dividing cortical progenitor cells. *Development* **131**, 3133–3145.

Müller, L. J., Pattiselanno, A., Cardozo, B. N. and Vrensen, G. (1984). Development of synapses on pyramidal and multipolar non-pyramidal neurons in the visual cortex of rabbits. A combined Golgi-electron microscope study. *Neuroscience* **12**, 1045–1069.

Nakagawa, N. and Iwasato, T. (2023). Golgi polarity shift instructs dendritic refinement in the neonatal cortex by mediating NMDA receptor signaling. *Cell Rep* **42**, 112843.

Noctor, S. C., Martinez-Cerdeño, V., Ivic, L. and Kriegstein, A. R. (2004). Cortical neurons arise in symmetric and asymmetric division zones and migrate through specific phases. *Nature Neuroscience 2004 7:2* **7**, 136–144.

O'Dell, R. S., Ustine, C. J. M., Cameron, D. A., Lawless, S. M., Williams, R. M., Zipfel, W. R. and Olson, E. C. (2012). Layer 6 cortical neurons require Reelin-Dab1 signaling for cellular orientation, Golgi deployment, and directed neurite growth into the marginal zone. *Neural Dev* **7**, 25.

Oishi, Y., Hashimoto, K., Abe, A., Kuroda, M., Fujii, A. and Miyamoto, Y. (2021). Vitronectin regulates the axon specification of mouse cerebellar granule cell precursors via αvβ5 integrin in the differentiation stage. *Neurosci Lett* **746**, 135648.

Powell, S. K., Rivas, R. J., Rodriguez-Boulan, E. and Hatten, M. E. (1997). Development of polarity in cerebellar granule neurons. *J Neurobiol* **32**, 223–236.

Rivas, R. J. and Hatten, M. E. (1995). Motility and cytoskeletal organization of migrating cerebellar granule neurons. *Journal of Neuroscience* **15**, 981–989.

Satake, T., Yamashita, K., Hayashi, K., Miyatake, S., Tamura□Nakano, M., Doi, H., Furuta, Y., Shioi, G., Miura, E., Takeo, Y. H., et al. (2017). MTCL1 plays an essential role in maintaining Purkinje neuron axon initial segment. *EMBO J* **36**, 1227–1242.

Solecki, D. J., Model, L., Gaetz, J., Kapoor, T. M. and Hatten, M. E. (2004). Par6α signaling controls glial-guided neuronal migration. *Nat Neurosci* **7**, 1195–1203.

Solecki, D. J., Govek, E. E., Tomoda, T. and Hatten, M. E. (2006). Neuronal polarity in CNS development. *Genes Dev* **20**, 2639–2647.

Stewart-Hutchinson, P. J., Hale, C. M., Wirtz, D. and Hodzic, D. (2008). Structural requirements for the assembly of LINC complexes and their function in cellular mechanical stiffness. *Exp Cell Res* **314**, 1892–1905.

Tahirovic, S. and Bradke, F. (2009). Neuronal polarity. *Cold Spring Harb Perspect Biol* **1**, 1–18.

Takano, T., Xu, C., Funahashi, Y., Namba, T. and Kaibuchi, K. (2015). Neuronal polarization. *Development (Cambridge)* **142**, 2088–2093.

Tan, M., Ma, S., Huang, Q., Hu, K., Song, B. and Li, M. (2013). GSK-3α/β-mediated phosphorylation of CRMP-2 regulates activity-dependent dendritic growth. *J Neurochem* **125**, 685–697.

Tanabe, K., Kani, S., Shimizu, T., Bae, Y. K., Abe, T. and Hibi, M. (2010). Atypical protein kinase C regulates primary dendrite specification of cerebellar Purkinje cells by localizing golgi apparatus. *Journal of Neuroscience* **30**, 16983–16992.

Umeshima, H., Hirano, T. and Kengaku, M. (2007). Microtubule-based nuclear movement occurs independently of centrosome positioning in migrating neurons. *Proc Natl Acad Sci U S A* **104**, 16182–16187.

Wu, Y. K., Fujishima, K. and Kengaku, M. (2015). Differentiation of apical and basal dendrites in pyramidal cells and granule cells in dissociated hippocampal cultures. *PLoS One* **10**,.

Wu, Y. K., Umeshima, H., Kurisu, J. and Kengaku, M. (2018). Nesprins and opposing microtubule motors generate a point force that drives directional nuclear motion in migrating neurons. *Development* **145**,.

Xie, Z., Sanada, K., Samuels, B. A., Shih, H. and Tsai, L. H. (2003). Serine 732 phosphorylation of FAK by Cdk5 is important for microtubule organization, nuclear movement, and neuronal migration. *Cell* **114**, 469–482.

Zhou, C., Wu, Y. K., Ishidate, F., Fujiwara, T. K. and Kengaku, M. (2024). Nesprin-2 coordinates opposing microtubule motors during nuclear migration in neurons. *J Cell Biol* **223**,.

Zmuda, J. F. and Rivas, R. J. (1998). The Golgi Apparatus and the Centrosome Are Localized to the Sites of Newly Emerging Axons in Cerebellar Granule Neurons In Vitro. *Cell Motil Cytoskeleton* **41**, 18–38.

Figure Legends

Figure 1. MTCL2 is preferentially enriched in the dendrites of cerebellar granule

neurons.

(A) Cerebellar sections from 4-week-old mice are stained for calbindin and NeuN (left

panel) or MTCL2 and α-tubulin (middle and right panels). Scale bars: 250 μm. (B)

Representative images of isolated CGNs cultured for 5 days and stained for acetylated

tubulin, MTCL2, and MAP2. Scale bar: 20 μm. (C) Correlation between the

immunofluorescence intensities of MTCL2 and MAP2 in acetylated tubulin-positive

neurites. A total of 100 neurites from 3 independent experiments were analyzed. (D)

Representative images of CGNs at DIV1–DIV5 stained for tau1, MTCL2, and MAP2.

In the merged view, nucleus staining is included to clarify the positions of cell bodies.

The arrowheads indicate tau1-positive neurites exhibiting MTCL2 signals in immature

CGNs. Scale bar: 20 μm. (E) Time-dependent increase in the relative intensities of the

MTCL2 (solid line) and MAP2 (dotted line) signals in dendrites. Neurites exhibiting

MAP2 signal intensities three times greater than the background were analyzed as

dendrites. Data represent the means ± S.D. from 3 independent experiments (totally,

more than 67 neurites were analyzed for each DIV stage). The paired t-test was used to

examine statistical significance (**$p < 0.01$; ***$p < 0.005$; n.s., non-significant). (F)

Western blot analysis of MTCL2 and MAP2 expression in primary cultured CGNs from

various DIVs. A representative result of 2 independent experiments is presented.

Figure 2. Depletion of MTCL2 resulted in the development of fewer and longer

dendrites.

(A) Full views of single CGNs at DIV6 visualized by the GFP signal. Scale bar: 50 μm.

(B) Distribution of total axonal length of individual CGNs subjected to the indicated

RNAi and fixed at DIV6. (C) Dendritic development of DIV6 CGNs transfected with

control or MTCL2 knockdown vectors and visualized by GFP and MAP2 signals. The

arrowheads indicate the dendrites of GFP-positive cells. Scale bar: 10 μm. (D)

Distribution of total number of dendritic tips of individual CGNs subjected to the

indicated RNAi/rescue conditions and fixed at DIV6. (E) Distribution of averaged

dendritic lengths of the individual CGNs analyzed in (D). (F) Percentage of dendrites

longer than 40 μm in all dendrites of the same CGNs analyzed in (D) and (E). Each data

point in (B), (D), and (E) represents the result of one neuron in 3 independent

experiments. Mean and standard deviations are shown for each scatter plot. The bars in

(F) represent mean and standard errors of 3 independent experiments. For all data, the

paired t-test was used to examine statistical significance (*$p < 0.05$; **$p < 0.01$; ***$p <$

0.005; n.s., non-significant).

Figure 3. Dendrite extensions and Golgi apparatus positions of MTCL2-depleted

CGNs biased in the direction opposite to axonal elongation.

(A) The extension directions of dendrites longer than 15 μm were defined as the angle

(θ) formed between the roots of the dendrite and an axon, as depicted. (B) θ distribution

of individual dendrites of CGNs subjected to the indicated RNAi/rescue conditions and

fixed at DIV6. The θ values were categorized in 4 directions and presented in polar

histograms as indicated. More than 350 dendrites from the indicated number (n) of cells

from 3 independent experiments were analyzed for each condition. (C) Comparison of

dendritic direction bias (r) between the conditions. r =1 indicates no direction bias. (D)

The position of the Golgi apparatus for individual CGNs was estimated as the angle (θ)

formed between the center of the Golgi apparatus and the roots of an axon, as depicted.

(E) θ distribution of the Golgi position in CGNs subjected to the indicated RNAi/rescue

conditions and fixed at DIV6. The θ values were categorized in 4 directions and

presented in polar histograms as indicated. The numbers (n) of cells from 3 independent

experiments were analyzed. (F) Comparison of direction bias (r) of the Golgi position

under each condition. Mean values and standard errors are shown for all quantitative

data for each condition. In (C) and (F), the paired t-test was used to examine statistical

significance (*p < 0.05; ***p < 0.005; n.s., non-significant).

Figure 4. MTCL2 is required for the bipolar-to-multipolar dendrite extension

transition.

(A) Representative morphologies of CGNs at DIV3 visualized by GFP. Axons and

dendrites were identified by staining with Ankyrin G and MAP2, respectively. Arrows

and arrowheads indicate dendrites and an axon of the GFP-expressing cell, respectively.

Scale bar: 20 μm. (B) θ distribution of individual dendrites of CGNs at DIV3 and 5.

Data are presented as polar histograms for >350 dendrites of the indicated number (n) of

cells from 3 independent experiments for each condition, as in Fig. 3B. (C) Distribution

of the average dendritic length of individual CGNs analyzed in (B). Each data point

represents the result of one neuron in 3 independent experiments. Mean values and

standard deviations are shown for each scatter plot. (D) Comparison of the direction

bias (r) of the Golgi position in the GCNs at DIV3 and DIV5. The mean values and

standard errors of 3 independent experiments are shown for each condition. In (C) and

(D), the paired t-test was used to examine statistical significance (*p < 0.05; ***p <

0.005). (E, F) Representative live imaging data of GFP-expressing CGNs transfected

with control (E) or MTCL2 (F) knockdown vectors. In each case, bipolar CGNs were

selected at DIV3, and recordings were performed for 48 h. The frames showing the

characteristic changes in the dendrite extensions are only presented for each CGN (for

comprehensive images, refer to Movies 1 and 2). Arrowheads indicate axons. Arrows

indicate initial stable dendrites. Note that the MTCL2-depleted CGN exhibited

shrinkage of the preexisting bipolar dendrites but failed to persistently extend short

dendrites in multiple directions. Instead, the cell stably extended long bipolar dendrites,

again. Scale bars: 50 μm.

Figure 5. The Golgi apparatus in the differentiated CGNs was sequestered into the

MT cage surrounding the nucleus.

(A) High-resolution confocal images of mature CGNs (DIV5) stained for tyrosinated

tubulin, MTCL2, and GS28 (Golgi membrane protein). Three images of the same field

in different confocal planes are presented from the basal to apical planes, as indicated.

Scale bar: 5 μm. (B) Boxed regions in (A) were enlarged to show substantial

colocalization of tyrosinated tubulin, MTCL2, and the Golgi apparatus. Scale bar: 5 μm.

(C) Cross-sectional views of the z-stack images along the dotted line in (A). Two types

of merged images with different antibody combinations are presented. The arrows

indicate the Golgi apparatus at the top of the nucleus. Scale bar: 2 μm. (D) High-

resolution confocal images of immature CGNs at DIV2 and DIV3 after staining with

tyrosinated tubulin and GS28. Scale bar: 5 μm. (E) Height distribution of the apical

edge of the Golgi apparatus estimated as a percentage of the cell heights (see Fig. S7).

Each data point represents the result of one neuron observed in 3 independent

experiments. Mean values and standard deviations are shown for each scatter plot. The

paired t-test was used to examine statistical significance (**p < 0.01; ***p < 0.005).

Figure 6. MTCL2 promotes the upward extension of the Golgi apparatus in an

MT- and Golgi-binding activity-dependent manner.

(A) High-resolution confocal images of CGNs subjected to the indicated RNAi, fixed at

DIV6, and stained for tyrosinated tubulin and GS28. On the left, projected views of the

z-stack images indicate cells expressing the indicated shRNA vector together with GFP.

Images obtained at three different confocal planes are shown on the right. Scale bar: 5

μm. (B) Cross-sectional views of the z-stack images along the dotted lines in (A).

Asterisks indicate GFP-positive cells. Scale bar: 5 μm. (C) Height distribution of the

Golgi apical edge in CGNs subjected to the indicated RNAi/rescue conditions and fixed

at DIV6. Each data point represents the result of one neuron observed in 3 independent

experiments. Mean values and standard deviations are shown for each scatter plot. The

paired t-test was used to examine statistical significance (***p < 0.005; n.s., non-

significant).

Figure 7. Inhibition of LINC complex activity mimicked most of the phenotypes

shown by MTCL2 knockdown.

(A) Schematic illustration of our working hypothesis. At the late phase of CGN

polarization, the MTCL2-mediated association between the Golgi apparatus and the MT

cage induces an upward extension of the Golgi apparatus along the nucleus. This

recruitment to the MT cage results in sequestration of the Golgi apparatus from the

roots of the preexisting bipolar dendrites. The molecular organization of the LINC

complex on the nuclear membrane is illustrated on the right. (B) A typical cross-

sectional view of CGNs expressing GFP-KASH1, fixed at DIV6, and stained for

MTCL2 and GS28. The white dotted lines indicate the focus plane from which the

confocal image shown in Fig. S8A was obtained. Scale bar: 5 μm. (C) An apical

confocal section of a GFP-KASH1-expressing cell stained for MTCL2 and tyrosinated

tubulin. Scale bar: 5 μm. (D) θ distribution of the Golgi position in CGNs expressing

GFP or GFP-KASH1, examined at DIV6. Data are presented as polar histograms

as shown in Fig. 3E for the indicated number (n) of cells from 3 independent

experiments. (E) Hight distribution of the Golgi apical edge in the same CGNs analyzed

in (C). (F) θ distribution of individual dendrites of CGNs expressing GFP or GFP-

KASH1 and examined at DIV6. Data are presented as polar histograms as shown in

Fig. 3B for >400 dendrites of the indicated number (n) of cells from 3 independent

experiments. (G, H) Distribution of the number of dendritic tips (G) and averaged

dendritic length (H) of the same CGNs analyzed in (F). Each data point in (E), (G) and

(H) represents the result of a single neuron. Mean values and standard deviations are

shown for each scatter plot. The paired t-test was used to examine statistical

significance (***$p < 0.005$; n.s., non-significant).

Movie Legends

Movie 1. Morphological changes in the control CGN from DIV3 to DIV5.

GFP fluorescence images were taken every 2 min for 40 h. The video speed was 30 fps.

The representative frames of this movie are shown in Fig. 4E.

Movie 2. Morphological changes in the MTCL2-depleted CGN from DIV3 to DIV5.

GFP fluorescence images were taken every 2 min for 40 h. The video speed was 30 fps.

The representative frames of this movie are shown in Fig. 4F.

Movie 3. 3D reconstructed images of polarized CGNs at DIV5.

The z-stack images of the CGNs shown in Fig. 5A were reconstructed into 3D

stereoscopic images with different combinations of staining.

Figure 1

A

Figure 2

Figure 3

Figure 4

Figure 5

A

basal ↑ → apical

0 μm   1.3 μm   2.6 μm

MTCL2 / GS28   Tyrosinated tub   GS28 (Golgi)   Tyr-tub / MTCL2 / GS28

B

Tyrosinated tub   MTCL2   GS28   Tyr-tub / MTCL2 / GS28

C

Tyrosinated tub / GS28   Tyr-tub / MTCL2 / GS28 / Nuclei

D

basal ← → apical

0 μm   1.4 μm   2.8 μm

DIV2

DIV3

Tyrosinated tub / GS28

E

Golgi apical end position / cell height (%)

DIV2   DIV3   DIV5

**   ***

Figure 6

A

basal →→→ apical

Control RNAi | 0 μm | 2.1μm | 2.6 μm
MTCL2 RNAi

GFP / GS28

Tyr-tub / GS28

B

Control RNAi

MTCL2 RNAi

Tyr-tub / GS28

C

Golgi apical end position / cell height (%)

	n.s.		***	
***		***		
		***	n.s.	

none | none | wt | ΔKR | 4LA
Cont. RNAi | MTCL2 RNAi

Figure 7