MtvS Regulates the *Francisella* Type V-A CRISPR-Cas System

Maj Brodmann and Luciano A. Marraffini[*]

Laboratory of Bacteriology, The Rockefeller University, New York, NY 10065, USA

[*] Howard Hughes Medical Institute, The Rockefeller University, New York, NY 10065, USA

Correspondence: mbrodmann@rockefeller.edu, marraffini@rockefeller.edu

Abstract

CRISPR-Cas systems endow bacteria and archaea with adaptive immune systems against
mobile genetic elements, playing a fundamental role in shaping microbial communities. Many
organisms harbor more than one CRISPR-Cas system, however whether and how these are
differentially regulated is not known, in many instances due to the impossibility of studying
CRISPR immunity in native hosts. Here we studied the regulation of endogenous type II-B
and type V-A CRISPR-Cas systems in opportunistic human pathogen *Francisella novicida*
U112. Fluorescence microscopy and transcriptomics experiments revealed that while the type
II-B system is constitutively expressed, the type V-A CRISPR-Cas system is differentially
expressed at stationary phase and high cell density. Using mass spectrometry and genetics we
identified MtvS as a factor required for the differential expression of the type V-A CRISPR-
Cas locus. Surprisingly, MtvS-dependent expression of the type V-A CRISPR-Cas system at
high cell density is linked quorum sensing like behavior, even though *F. novicida* U112 lacks
canonical quorum sensing genes. Our findings provide an example of how bacteria harboring
multiple CRISPR systems regulate their expression.

Introduction

CRISPR (Clustered regularly short interspaced palindromic repeats)-Cas (CRISPR associated genes) systems protect bacteria and archaea against foreign nucleic acids such as phages and genetic mobile elements[1,2]. CRISPR-Cas systems are adaptive immune systems that can store memories of previous infections in the form of short invader-derived sequences that are inserted in between the repeats of the CRISPR array during infection[3–5]. CRISPR arrays are transcribed into cr(CRISPR)RNAs that guide Cas effectors to complementary sequences on the invading foreign elements[6,7] and destroy them [8–13]. Prokaryotic organisms often encode multiple CRISPR-Cas loci, which may be differently regulated[14]. Previous studies have reported Cas proteins, crRNAs, transcriptional regulators, quorum sensing and two component signaling systems involved in the regulation CRISPR-Cas systems [15–35]. Detailed investigation of regulatory networks governing CRISPR expression, however, has been difficult due to the lack of natural hosts that can be used as experimental systems for CRISPR-Cas immunity.

Here we studied the endogenous type II-B and type V-A CRISPR-Cas systems in the opportunistic human pathogen *Francisella tularensis* subspecies *novicida* Utah 112 (here *F. novicida* U112) (Fig. S1A-B)[13,36,37]. It has been shown previously that both *F. novicida* systems are functional and capable of restricting foreign DNA[38,39]. Interestingly, *F. novicida* Cas9 is also required for virulence and enhances membrane stability by repressing an operon containing a lipoprotein with unknown function[40–43]. Using florescence microscopy, we found that the two *F. novicida* CRISPR-Cas systems are differentially regulated and expressed in non-infection conditions. While the type II-B CRISPR-Cas system is constitutively expressed, the type V-A CRISPR-Cas system displays significantly higher expression during stationary phase. Using mass spectrometry and genetics we identified MtvS (Modulator of the type V-A CRISPR-Cas system expression in stationary phase, FTN_1238), a protein with previously unknown function in *Francisella* species, as a factor required for the differential expression of the type V-A CRISPR-Cas locus. Last, we showed that type V-A CRISPR-Cas system expression is induced upon introduction to spent overnight medium suggesting that *F. novicida* is capable of quorum sensing despite lacking canonical quorum sensing genes.

Results

*The F. novicida Type V-A CRISPR-*cas *locus is upregulated during stationary phase*

To investigate the expression pattern of the type II-B and type V-A CRISPR-Cas systems in *F. novicida* U112, their RNA-guided nucleases were C-terminally tagged to generate Cas9-msfGFP and Cas12-mCherry2, respectively. The fluorescent tags did not affect their nuclease function, since plasmid interference assays confirmed that the tagged nucleases prevented plasmid conjugation as well as the native ones (Fig. S1C). Fluorescence live-cell imaging of exponentially growing bacteria and cells in stationary phase revealed that fluorescence intensity was comparable in both growth phases for Cas9-msfGFP while fluorescent intensity for Cas12-mCherry2 was higher in stationary phase (Fig. 1A). We then performed RNA-seq to test if the distinct expression patterns of the two effector nucleases originate from different transcription levels. Comparing RNA levels of genes between exponentially growing and stationary phase bacteria confirmed that *cas12* and *cas1$_{typeV-A}$* RNA levels are significantly higher during stationary phase than in exponential growth phase (Fig. 1B). In comparison, all components from the type II-B CRISPR-Cas system, including *cas9*, had comparable RNA levels in the two tested conditions (Fig. 1B). Calculated expression changes between exponential and stationary growth for all genes are listed in the Supplementary Data File. Targeted measurement of RNA levels of *cas9* and *cas12* by RT-qPCR confirmed the invariability of RNA levels of *cas9* in the two tested growth conditions and the distinct temporal expression of *cas12* in stationary growth phase (Fig. 1C and D). Together these results show that the two effector nucleases from the *Francisella* type II-B and type V-A CRISPR-Cas systems have different expression patterns and that *cas12* expression levels significantly increase in stationary phase.

MtvS is required for upregulation of the Type V-A CRISPR-cas locus in stationary phase

Type V-A CRISPR-Cas systems have been shown to autoregulate in an RNA-guided and Cas effector dependent manner[30]. Moreover, *F. novicida* U112 *cas12* promoter sequence contains a predicted small repeat unit (SRU, with a similar but not identical sequence to the canonical CRISPR repeat, Fig. 2A), which was proposed to have a regulatory function[29]. To test if Cas12 is responsible for the upregulation of the type V-CRISPR-*cas* locus we replaced *cas12* with *mCherry2*. However, the fluorescence pattern did not change in absence of *cas12* (Fig. 2B). In addition, deletion of the predicted regulatory *SRU* sequence in the *cas12* promoter sequence did not result in any significant changes in *cas12* RNA levels or targeting efficiency (Fig. 2C-D). To identify other transcription factors, we performed a protein pull down using the *cas12* promoter DNA sequence and an intragenic *cas12* control sequence as bait. Lysates from exponentially growing and stationary bacteria were probed and pull-down proteins were identified proteins by LC-MS/MS. All identified proteins are listed in the Supplementary Data

File. An enrichment score was calculated by comparing protein abundance in pull downs from

the *cas12* promoter sequence to the control sequence. In exponentially growing cells, one of

the most significantly enriched proteins was the LysR-family transcriptional regulator

FTN_0392 (Fig. 3A, Supplementary Data File). However, neither Cas12-mCherry2

expression nor targeting capability was decreased in a *FTN_0392* deletion mutant (Fig. S2A-

B). On the other hand, in the stationary growth sample, FTN_1238 was significantly enriched

(Fig. 3B). Importantly, there was no enrichment for FTN_1238 when using a lysate from

exponentially growing bacteria (Fig. 3A). *FTN_1238* is the first gene of an operon encoding

for genes required for the respiratory chain (*ubiH*, *ubiF*), a putative phage protein

(*FTN_1235*) and a tRNA modification enzyme (*queA*) (Fig. S2C). FTN_1238 has no assigned

function in *Francisella* yet but is homologous to YgfB (Fig. S2D), which is conserved in

many γ-proteobacteria [44].

To test if FTN_1238 was involved in regulation of *cas12* expression, we made a non-sense

mutation by introducing three stop codons after aspartate at position 11 and analyzed the

mutant cultures via fluorescence microscopy. We found decreased fluorescence of Cas12-

mCherry2 in stationary phase for this mutant (Fig. 3C) and the phenotype could be rescued by

introducing a wild-type copy of *FTN_1238* with its native promoter at the Tn7 introduction

site downstream of gene *glmS* (*FTN_0485*) (Fig. 3C). Disruption of FTN_1238 significantly

affected the RNA levels of all type V-A CRISPR-Cas system genes in stationary phase as

measured by RT-qPCR (Fig. 3D-G). RNA levels were also restored with an additional wild-

type copy of *FTN_1238* (Fig. 3D-G) except for *cas2*, the last gene of the operon (Fig. 3G).

Importantly, *cas9* RNA levels were not affected (Fig. S3B), a result that demonstrates that

FTN_1238 acts specifically on the type V-A locus. Interestingly, while disruption of

FTN_1238 had a detrimental effect on *cas12* RNA levels in stationary phase, targeting

capability of Cas12 in plasmid interference assays was not affected (Fig. S2E), most likely

due to the presence of low levels of Cas12 that are sufficient to clear the invading plasmid.

Finally, while *FTN_1238* RNA levels were comparable in both growth phases (Fig. S2F,

Supplementary Data File), fluorescence microscopy suggested higher proteins levels of

FTN_1238-mCherry2 in stationary phase (Fig. S2G). In summary, our results indicate that

FTN_1238 is a <u>m</u>odulator of the <u>t</u>ype <u>V-A</u> CRISPR-Cas system expression in <u>s</u>tationary

phase, and therefore we renamed it MtvS.

Quorum sensing signals participate in the regulation of the Type V-A CRISPR-cas *locus*

During stationary phase the cell density is high (Figs. 1 and 3). Interestingly, microscopy of

exponentially growing cells in microfluidic plates showed that Cas12-mCherry2 is also

expressed in a MtvS dependent manner in this growing phase within densely populated areas

(Fig. 4A-B, Movies 1 and 3), but not in less dense regions belonging to the same flow

chamber (Fig. 4A-B, Movies 2-4). These results led us to hypothesize that quorum sensing

could play a role in regulating the type V-A CRISPR-Cas system. Therefore, we tested if the

addition of spent overnight medium could increase *cas12* expression in exponential cultures

of *F. novicida* U112, possible containing signals that would trigger expression. We replaced

the medium of exponentially growing cells with filtered spent overnight medium in

microfluidic plates and monitored individual cells by microscopy. We observed drastic

changes, with bacteria changing from rod to coccus shapes, and an increased Cas12-

mCherry2 fluorescence (Fig. 4C, Movie 5). Fresh medium reversed these changes and cells

resumed growth as rod-shaped and Cas12-mCherry2 expression decreased (Fig. 4D, Movie

5). Heat-treatment of the spent overnight medium eliminated its activity on exponentially

growing bacteria (Fig. S3A, Movie 6). When exposed to overnight spent medium, *mtvS*

nonsense mutant cultures also suffered rod-to-coccus shape changes, but displayed much

lower Cas12-mCherry fluorescent signal, indicating that Cas12-mCherry2 was less expressed

(Fig. 4E-F, Movie 7). This result was confirmed by RT-qPCR, sampling *mtvS* nonsense

mutant cells before, 0.5 h and 1 h after switching to spent overnight medium and probed for

cas12 and *cas9* RNA levels. While *cas12* RNA levels increased over 1 h after the medium

change, *cas9* RNA levels did not significantly change (Fig. 4H and Fig. S3B). Together these

results suggest that *F. novicida* U112 likely integrates quorum sensing signals into the MtvS

dependent regulation of Cas12 expression.

145 Discussion

Here we compared the expression of the two native type II-B and type V-A CRISPR-Cas

systems in *F. novicida* U112 in the absence of phage infection. We found that while the type

II-B CRISPR-Cas system is continuously expressed throughout growth in liquid culture, the

type V-A CRISPR-Cas system is significantly upregulated at high cell density and stationary

growth phase. We envision different possibilities behind this unique regulation. First, it is

known that bacteria enhance their anti-phage defenses in high cell density environments when

they are more vulnerable to phage predation[15,16,45–51]. Given that the type II-B system is

constitutively expressed, the upregulation of the type V-A system will equip *F. novicida* U112

with additional defenses in stationary phase. Moreover, our RNA-seq analysis also found upregulation of two of the six annotated restriction-modification systems in stationary growth phase (Supplementary Data File 1)[52]. It is also possible that Cas12 or other components of the type V-A CRISPR-Cas system have a role beyond immunity in stationary growth phase, analogous to the function of *F. novicida* Cas9 in the regulation of lipoprotein expression[40–43]. Our RNA-seq experiment showed drastic changes in transcription in stationary phase, indicating the need to adapt to this growth condition, an adaptation in which Cas12 could play a part that would justify its upregulation in this growth phase.

We identified MtvS (FTN_1238), a protein with previously unknown function in *F. novicida* U112, as a factor required for the upregulation of the type V-A CRISPR-Cas system in stationary phase and upon addition of spent overnight medium to the exponentially growing cultures. MtvS belongs to the YecA/YgfB-like protein family and is conserved in many γ-proteobacteria[44]. Besides the structure of *Haemophilus influenzae* YgfB, little is known about the function of this family of proteins[53]. In *Pseudomonas aeruginosa*, YgfB increases β-lactam resistance [54] through interactions with the antitermination factor AlpA that lead to the upregulation of AmpC expression[55]. We believe that it is unlikely that MtvS interacts with an antitermination factor to upregulate the type V-A CRISPR locus in *F. novicida* U112 since *cas12* transcription decreases in the *mtvS* mutant (Fig. 3). We were unable to detect specific binding to the *cas12* promoter region by purified MtvS (data not shown), therefore we hypothesize that MtvS either stimulates *cas12* transcription through interaction with other factors or with RNA polymerase or through the inhibition of a termination factor. Intriguingly, mass spectrometry analysis of pull-downs showed enrichment for NusA, which can act both as transcriptional termination and antitermination factor depending on the bacterial species, as well as transcriptional termination factor Rho (Supplementary Data File 1). Unfortunately, due to the essential nature of NusA and Rho in *F. novicida* U112[56], we were unable to create knock out strains to investigate a possible role for these factors in the modulation of the type V-A CRISPR locus. Finally, *mtvS* RNA levels were comparable in both growth conditions but MtvS-mCherry2 levels increased in stationary phase. Therefore, it is likely that MtvS is post-transcriptionally regulated.

F. novicida U112 cells reacted strongly to spent overnight medium, displaying a growth arrest, morphology changes and an increase in Cas12 expression, the latter also dependent on the presence of MtvS. Changes in gene expression as reaction to spent overnight medium is a hallmark of quorum sensing[57]. However, *F. novicida* U112 lacks canonical quorum sensing

187 genes[58]. Our results suggest that the type V-A CRISPR-Cas system and MtvS could be part of

188 a quorum sensing circuit that monitors and responds to changes in cell density. Future studies

189 will investigate this possibility, relevant not only for the understanding of the phage defense

190 strategies but also for the pathogenesis of *Francisella*.

Author Contributions. M.B. and L.A.M. conceived the study. M.B. performed all experiments. M.B. and L.A.M. wrote the manuscript. All authors read and approved the manuscript.

Acknowledgments. We thank D.M. Monack (Stanford University) for the conjugation plasmid pDMK3, A. Harms (ETH Zurich) and C. Dehio (Biozentrum, University of Basel) for the *E. coli* conjugation strain, M. Basler (Biozentrum, University of Basel) for plasmid pFNMB2 *msfGFP*, E. Campbell (The Rockefeller University) for helpful scientific discussions and H. Molina and C. Peralta (Proteomics Resource Center, The Rockefeller University) for performing and analyzing the LC-MS/MS experiment. Proteomics data was generated by the Proteomics Resource Center at The Rockefeller University (RRID:SCR_017797) using instrumentation funded by the Sohn Conferences Foundation and the Leona M. and Harry B. Helmsley Charitable Trust. Thanks also goes to C.F. Baca (The Rockefeller University) for scientific advice and support and to all members of the Marraffini lab (The Rockefeller University) for scientific discussions and feedback. M.B. was supported by the SNSF Early Postdoc.Mobility (P2BSP3_195673) and SNSF Postdoc.Mobility (P500PB_210998) fellowships. L.A.M. is an investigator of the Howard Hughes Medical Institute.

Competing interests. LAM is a cofounder and Scientific Advisory Board member of Intellia Therapeutics, a cofounder of Eligo Biosciences and a Scientific Advisory Board member of Ancilia Biosciences.

Material and Methods

Bacterial strains and growth conditions

Francisella novicida U112 (*F. novicida* U112) and associated mutants were grown aerobically in brain heart infection (BHI, BD Biosciences) broth or on BHI agar plates supplemented with 0.1 % L-cysteine (Sigma Alderich) and 50 µg/ml ampicillin (GoldBio). F. *novicida* strains carrying plasmids were grown with 15 µg/ml kanamycin (GoldBio).

To receive exponentially growing cultures, stationary phase cells were diluted 1:20 in fresh media and grown for 2-3 h until they reached an OD_{600} of 0.5-0.8 and had a rod-shaped morphology. Stationary phase cells were grown over night for at least 16 hours until they reached an OD_{600} of 3.5-4.5 and displayed a more coccus-like cell shape. Cell morphology was checked by microscopy.

Escherichia coli OneShotpir (*E. coli,* ThermoFisher) and derivative strains were aerobically grown in Luria broth (LB, BD Biosciences) or on LB agar plates supplemented with 50 µg/ml kanamycin at 37 °C. For conjugation *E. coli* JKE201 1.2 mM 2,6-Diaminopimelic acid (DAP, Sigma Alderich) was added to all growth media since this strain is auxotrophic for mDAP. Strains used for this study are listed in table 1.

Bacterial mutagenesis

To generate chromosomal in-frame *F. novicida* mutants, suicide vector pDMK3 was used [59]. For fluorescent tags, an Ala-Ala-Ala-Gly-Gly-Gly linker was added and after the stop codon the 30 bp sequence of the gene was repeated to avoid downstream polar affects. For the His-tags only a Gly linker was added. In-frame deletions yielded in peptide scars containing the first and last 10 amino acids. Mutagenesis and conjugation was carried out as reported previously [60,61]. For conjugation, donor *E. coli* JKE201 from A. Harms and C. Dehio [62] and recipient *F. novicida* strains were grown until a OD_{600} of 1 was reached. 1 ml of each culture was mixed in 20 µl of LB and spotted on a LB agar plate supplemented with 1.2 mM DAP and incubated over night at 25 °C. The mixture was plated on Mueller Hinton (BD Biosciences) agar plates supplemented with 0.1 % L-cysteine, 0.1 % D-glucose (Sigma Alderich), 0.1 % heat inactivated fetal calf serum (Gibco), 50 µg/ml ampicillin and 15 µg/ml kanamycin and incubated at 37 °C for 2 days. *F. novicida* colonies containing the suicide vector were restreaked on BHI agar plates supplemented with 15 µg/ml kanamycin. To select against pDMK3, individual colonies were restreaked on LB agar plates supplemented with 5 % sucrose (Sigma Alderich), 0.1 % L-cysteine and 50 µg/ml ampicillin and incubated at 25 °C until colonies lost the suicide vector. Each mutant was verified by colony PCR using primers binding outside of the recombined region and sequencing of these regions. In addition, they were restreaked on kanamycin again to check for loss of pDMK3.

All primers and plasmids are listed in table 2.

Fluorescence live cell imaging

A Nikon Ti-E inverted motorized microscope equipped with Perfect Focus System and a CFI Plan Apochromat DM Lambda 100X Oil objective lens (NA 1.4) was used for live-cell imaging. A SOLA Light Engine (Lumencor) set to 5 % power and C-FL GFP HC HISN Zero Shift filter (excitation: 470/40 nm (450–490 nm), emission: 525/50 nm (500–550 nm), dichroic mirror: 495 nm) and C-FL DS RED HC HISN Zero Shift filters (excitation: 545/30nm (530-560nm), emission: 620/60nm (590-630nm), dichroic mirror: 570nm), respectively, were used to excite and filter fluorescence from msfGFP and mCherry2. The

exposure time for each channel was set to 500 ms and images were collected with a Zyla 4.2 sCMOS (Andor, pixel size 65 nm) or an ORCA-FusionBT sCMOS camera (Hamamatsu, pixel size 65 nm). Phase contrast was imaged with the SOLA light Engine set to 2% power and 10 ms excitation. The microscope was controlled by the software NIS-Elements AR version 5.21.03.

Exponentially grown (rod-shaped) and stationary phase (roundish) *F. novicida* were spun down and resuspended in phosphate saline buffer (PBS, Gibco) and spotted on pads consisting of 1% agarose in PBS. Snapshots of phase contrast, RFP and GFP channels were taken immediately after.

Long-term imaging experiments which include switching of media were carried out in CellASIC ONIX microfluidic plates (Millipore). Exponentially growing bacteria were resuspended in BHI supplemented with 0.2 % L-cysteine and loaded into the microfluidic plate along with fresh BHI supplemented with 0.2 % L-cysteine for initial growth and the tested medium as specified in the Fig. legend. For generating spent overnight medium, cells in stationary phase were pelleted for 1.5 min at 16000 g and supernatant was filtered with 0.22 µm filters (Millipore). For heat-treated spent overnight medium, filtered spent overnight medium was incubated at 95 °C for 10-20 min. After connecting the plate to the CellASIC ONIX2 Microfluidic System, the plates were incubated at 37 °C for 1 h using a Tokai HIT thermal box (Zeiss). Images of phase contrast, RFP and GFP channel were taken every 5 minutes after loading cells into the imaging chamber. First, fresh BHI was flown over for 1 h before switching to the tested medium for 4h. Afterwards, fresh BHI was flown over again for 3 h. For all media, the flow rate was set to 13.8 kPa.

Image analysis

Image analysis was carried out with Fiji software [63]. For comparison of fluorescent signal intensities, the contrast was set to same values for sets of compared images in same sub panel. If compared across sub panels, it is stated in Fig. legend which contrasts are set to same values.

For long-term imaging, images were aligned to the first image using NIS-Elements AR version 5.21.03.

RNA-seq

RNA was extracted from exponentially growing and stationary phase cells in three biological replicates for each condition. Cells yielding in a total OD_{600} of 3 were pelleted and resuspended in 80 µl of PBS and 20 µl of 10 % Sarcosyl (Sigma Alderich). Then, RNA was extracted with TRI Reagent according to the manufacturer's protocol (Zymo Research). An in-column DNA digest during RNA extraction was performed. A second DNA digestion was carried out with 10 µg of RNA using Turbo DNase (Invitrogen) for 2x 30 min. The DNA-free RNA was cleaned up with the RNA Clean & Concentrator-5 kit (Zymo Research) and eluted in 15 µl of water. RNA quality was checked by running a bleach agarose gel[64]. RNA concentrations were measured with a Qubit 4 fluorometer (Invitrogen) using the Qubit RNA Broad Range Assay Kit.

Ribosomal RNA was depleted with the NEBNext rRNA depletion (bacteria) kit (NEB). An input of 1 µg of RNA was used. Then, the RNA was prepared for sequencing using the TruSeq Stranded mRNA kit (Illumina) according to manufacturer's instruction. For the first strand synthesis, Superscript IV reverse transcriptase (Invitrogen) was used with an annealing temperature of 50 °C. Final DNA concentrations and lengths were measured with a Qubit 4

fluorometer (Invitrogen) using the Qubit dsDNA High Sensitivity Assay Kit and the
Tapestation 4200 bioanalyzer (Agilent) with the Agilent D500 ScreenTape Assay (Agilent).

The DNA libraries were diluted to 12 pM and mixed with 2.5% 20 pM PhiX and subjected to
single read sequencing with a 150-cycle MiSeq Reagent Kit v3 (Illumina) on the Illumina
MiSeq platform.

The reads were mapped to the *F. novicida* U112 genome (RefSeq NC_008601[65]) and RPKM
values for each gene was calculated using coordinates of coding genes (ptt files) and of
rRNAs and tRNAs (rtt files) with EDGE-Pro v1.3.1[66]. EDGE-Pro statistics for each sample
can be found in table 5. Significant changes in gene expression between used conditions were
calculated with the R (version 4.3.0) package DESeq2 (version 1.40.2)[67]. Results can be
found in table 6. *P*-values of 0.01 and log2 foldchanges of 2 were defined significant.

RT-qPCR

RNA was extracted as reported for RNA-seq. The RNA was reverse transcribed to cDNA
with Superscript IV reverse transcriptase (Invitrogen) using 1 μg of RNA, random hexamer
primers, and a reaction temperature of 50 °C.

qPCR was performed in the QuantStudio 3 Real-Time PCR System (Applied Biosystems)
using the PowerUp SYBR Green Master Mix (Applied Biosystems). Reactions were set up in
a total volume of 20 μl containing 50 ng of cDNA template and 500 nM primers. Each
reaction was set up in three technical replicates. All primers used for RT-qPCR are found in
table 3. The annealing and amplification temperature was set to 60 °C.

The geometrical mean of the three technical replicates for each biological replicate was used
for further analysis. RNA levels of probed genes were normalized to the DNA 3'-5' helicase
gene *uvrD* (FTN_1594), which has been used previously for normalization and does not
change its expression in the used conditions as determined by RNA-seq[68]. The average value
of the three biological *uvrD* replicates was used to the ΔCt value for each gene. Relative RNA
abundance of probed genes was calculated as $2^{-\Delta Ct}$.

Statistical analysis for RT-qPCR data was performed with Prism10.2.3 (403) (GraphPad
Software). To test if relative RNA abundances determined by RT-qPCR are significantly
different, two-way ANOVA (α=0.05) with correction for multiple comparison (Tukey's
multiple comparisons test) was performed. Adjusted *P*-values are given in the Fig. legends.

Plasmid interference assay

Strains were generated with the native Type II-B or Type V-A CRISPR array replaced by a
targeting or non-targeting spacer matching expression plasmid pFNMB2 *msfGFP*. These
strains were grown exponentially together with the donor *E. coli* strain harboring pFNMB2
msfGFP. Cells yielding a total OD_{600} of 1 were pelleted and washed once with LB. Then,
recipient *F. novicida* strains were mixed with donor *E. coli* in a final volume of 20 μl LB and
spotted on nitrocellulose filter membranes with a 0.22 μm pore size (Millipore) located on LB
agar plates supplemented with 1.2 mM DAP. The plates were incubated for 2 h at 37 °C
before the bacteria were resuspended in a final volume of 100 μl MH broth (BD Biosciences).
Serial dilutions in MH broth were carried out and 5 μl of each dilution was spotted in
technical duplicates on MH agar plates supplemented with 50 μg/ml ampicillin for calculation
of total number of colony forming units (CFUs) and on MH agar plates supplemented with 50
μg/ml ampicillin and 15 μg/ml kanamycin for calculation of transformants. The average

number of CFUs of the two technical replicates was used for calculating CFU/ml. Three
biological replicates for each strain were analyzed.

Promoter sequence pull-down

Pull-down of proteins binding the *cas12* promoter and the intragenic *cas12* control sequence
was carried out as reported previously[69,70].

In short, primers with a 5' Biotin-TEG label were used to generate the bait DNA sequences
(table 4) from *F. novicida* gDNA. PCR products were checked by agarose gel and cleaned up
with QIAquick PCR Purification Kit (QIAGEN) and concentrated with Amicon Ultra-0.5 mL
Centrifugal Filters (Millipore Sigma) to 200 ng/µl in water. Washed Dynabeads MyOne
Streptavidin C1 (Thermo Fisher Scientific) were loaded with 40 µg of DNA for 20 minutes.
DNA concentrations in the removed supernatants were measured to ensure that most DNA
bound to the Dynabeads. Exponentially growing and stationary phase cells yielding a total
OD_{600} of 800 were resuspended in 5 ml 1x FastBreak cell lysis buffer (Promega) for 30
minutes on ice and then sonicated afterwards. 200 µl of precleared lysates was added to the
resuspended beads supplemented with 0.03 µg/µl salmon sperm DNA (Thermo Fisher
Scientific) and incubated for 1 h with agitation. The beads were washed 2x with a wash buffer
(100 mM NaCl (Sigma Alderich), 25 mM Tris-HCl (VWR, Thermo Scientific), pH 7.5) and
then switched to new tubes before a last wash. The dried beads were stored at 4 °C until they
were subjected to LC-MS/MS analysis.

LC-MS/MS analysis

Beads were resuspended in 8 M Urea and proteins were recovered from beads. After
alkylation, proteins were digested with trypsin and LysC. The reaction was stopped with
trifluoroacetic acid. Generated peptides were desalted and concentrated with solid-phase
extraction prior to analysis by LC-MS/MS (12cm packed-in-emitter column, 70min gradient,
QE-HF). The raw data was searched and quantified using ProteomeDiscoverer/Mascot and
MaxQuant (version 2.0.3.0) using Uniprot database for *F. tularensis* subspecies *novicida*
U112 (taxon ID: 401614) with additional common contaminants generating the LFQ and
iBAQ values. Statistical analysis was carried out with Perseus software. Contaminants and
proteins that have been identified in less than two biological replicates of a sample were
removed. iBAQ were median normalized. The Log2 foldchange was calculated between
proteins pulled-down with the *cas12* promoter sequence and proteins pulled-down with the
intragenic *cas12* control sequence for both exponential and stationary conditions. An FDR-
based (FDR = 0.05) two sample t-test was used to determine which changes were significant.
P-values of 0.01 and log2 foldchanges of 4 were defined significant.

Homology prediction with Foldseek

387 Foldseek[71] was used to find structural homologs for the predicted structure of FTN_1238
388 (Uniprot ID: A0A6I4RTR9).

Data availability

391 Source data are provided in this paper.

References

1.   Marraffini, L.A., and Sontheimer, E.J. (2008). CRISPR interference limits horizontal gene transfer in staphylococci by targeting DNA. Science *322*, 1843–1845. https://doi.org/10.1126/science.1165771.

2.   Barrangou, R., Fremaux, C., Deveau, H., Richards, M., Boyaval, P., Moineau, S., Romero, D.A., and Horvath, P. (2007). CRISPR provides acquired resistance against viruses in prokaryotes. Science *315*, 1709–1712. https://doi.org/10.1126/science.1138140.

3.   Bolotin, A., Quinquis, B., Sorokin, A., and Ehrlich, S.D. 2005 Clustered regularly interspaced short palindrome repeats (CRISPRs) have spacers of extrachromosomal origin. Microbiology *151*, 2551–2561. https://doi.org/10.1099/mic.0.28048-0.

4.   Mojica, F.J.M., Díez-Villaseñor, C., García-Martínez, J., and Soria, E. (2005). Intervening Sequences of Regularly Spaced Prokaryotic Repeats Derive from Foreign Genetic Elements. J. Mol. Evol. *60*, 174–182. https://doi.org/10.1007/s00239-004-0046-3.

5.   Pourcel, C., Salvignol, G., and Vergnaud, G.Y. 2005 CRISPR elements in Yersinia pestis acquire new repeats by preferential uptake of bacteriophage DNA, and provide additional tools for evolutionary studies. Microbiology *151*, 653–663. https://doi.org/10.1099/mic.0.27437-0.

6.   Brouns, S.J.J., Jore, M.M., Lundgren, M., Westra, E.R., Slijkhuis, R.J.H., Snijders, A.P.L., Dickman, M.J., Makarova, K.S., Koonin, E.V., and Oost, J. van der (2008). Small CRISPR RNAs Guide Antiviral Defense in Prokaryotes. Science. https://doi.org/10.1126/science.1159689.

7.   Hale, C., Kleppe, K., Terns, R.M., and Terns, M.P. (2008). Prokaryotic silencing (psi)RNAs in Pyrococcus furiosus. RNA N. Y. N *14*, 2572–2579. https://doi.org/10.1261/rna.1246808.

8.   Abudayyeh, O.O., Gootenberg, J.S., Konermann, S., Joung, J., Slaymaker, I.M., Cox, D.B.T., Shmakov, S., Makarova, K.S., Semenova, E., Minakhin, L., et al. (2016). C2c2 is a single-component programmable RNA-guided RNA-targeting CRISPR effector. Science *353*, aaf5573. https://doi.org/10.1126/science.aaf5573.

9.   Gasiunas, G., Barrangou, R., Horvath, P., and Siksnys, V. (2012). Cas9-crRNA ribonucleoprotein complex mediates specific DNA cleavage for adaptive immunity in bacteria. Proc. Natl. Acad. Sci. U. S. A. *109*, E2579-2586. https://doi.org/10.1073/pnas.1208507109.

10.   Hale, C.R., Zhao, P., Olson, S., Duff, M.O., Graveley, B.R., Wells, L., Terns, R.M., and Terns, M.P. (2009). RNA-Guided RNA Cleavage by a CRISPR RNA-Cas Protein Complex. Cell *139*, 945–956. https://doi.org/10.1016/j.cell.2009.07.040.

11.   Jinek, M., Chylinski, K., Fonfara, I., Hauer, M., Doudna, J.A., and Charpentier, E. (2012). A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. Science *337*, 816–821. https://doi.org/10.1126/science.1225829.

12.   Westra, E.R., van Erp, P.B.G., Künne, T., Wong, S.P., Staals, R.H.J., Seegers, C.L.C., Bollen, S., Jore, M.M., Semenova, E., Severinov, K., et al. (2012). CRISPR immunity relies on the consecutive binding and degradation of negatively supercoiled invader DNA by Cascade and Cas3. Mol. Cell *46*, 595–605. https://doi.org/10.1016/j.molcel.2012.03.018.

13.   Zetsche, B., Gootenberg, J.S., Abudayyeh, O.O., Slaymaker, I.M., Makarova, K.S.,
Essletzbichler, P., Volz, S.E., Joung, J., van der Oost, J., Regev, A., et al. (2015). Cpf1 is a
single RNA-guided endonuclease of a class 2 CRISPR-Cas system. Cell *163*, 759–771.
https://doi.org/10.1016/j.cell.2015.09.038.

14.   Makarova, K.S., Wolf, Y.I., Iranzo, J., Shmakov, S.A., Alkhnbashi, O.S., Brouns, S.J.J.,
Charpentier, E., Cheng, D., Haft, D.H., Horvath, P., et al. (2020). Evolutionary classification of
CRISPR-Cas systems: a burst of class 2 and derived variants. Nat. Rev. Microbiol. *18*, 67–83.
https://doi.org/10.1038/s41579-019-0299-x.

15.   Patterson, A.G., Jackson, S.A., Taylor, C., Evans, G.B., Salmond, G.P.C., Przybilski,
R., Staals, R.H.J., and Fineran, P.C. (2016). Quorum Sensing Controls Adaptive Immunity
through the Regulation of Multiple CRISPR-Cas Systems. Mol. Cell *64*, 1102–1108.
https://doi.org/10.1016/j.molcel.2016.11.012.

16.   Høyland-Kroghsbo, N.M., Paczkowski, J., Mukherjee, S., Broniewski, J., Westra, E.,
Bondy-Denomy, J., and Bassler, B.L. (2017). Quorum sensing controls the Pseudomonas
aeruginosa CRISPR-Cas adaptive immune system. Proc. Natl. Acad. Sci. U. S. A. *114*, 131–
135. https://doi.org/10.1073/pnas.1617415113.

17.   Lucas-Elío, P., Molina-Quintero, L.R., Xu, H., and Sánchez-Amat, A. (2021). A
histidine kinase and a response regulator provide phage resistance to Marinomonas
mediterranea via CRISPR-Cas regulation. Sci. Rep. *11*, 20564. https://doi.org/10.1038/s41598-
021-99740-9.

18.   Borges, A.L., Castro, B., Govindarajan, S., Solvik, T., Escalante, V., and Bondy-
Denomy, J. (2020). Bacterial alginate regulators and phage homologs repress CRISPR-Cas
immunity. Nat. Microbiol. *5*, 679–687. https://doi.org/10.1038/s41564-020-0691-3.

19.   Perez-Rodriguez, R., Haitjema, C., Huang, Q., Nam, K.H., Bernardis, S., Ke, A., and
DeLisa, M.P. (2011). Envelope stress is a trigger of CRISPR RNA-mediated DNA silencing in
Escherichia coli. Mol. Microbiol. *79*, 584–599. https://doi.org/10.1111/j.1365-
2958.2010.07482.x.

20.   Pul, U., Wurm, R., Arslan, Z., Geissen, R., Hofmann, N., and Wagner, R. (2010).
Identification and characterization of E. coli CRISPR-cas promoters and their silencing by H-
NS. Mol. Microbiol. *75*, 1495–1512. https://doi.org/10.1111/j.1365-2958.2010.07073.x.

21.   Westra, E.R., Pul, U., Heidrich, N., Jore, M.M., Lundgren, M., Stratmann, T., Wurm,
R., Raine, A., Mescher, M., Van Heereveld, L., et al. (2010). H-NS-mediated repression of
CRISPR-based immunity in Escherichia coli K12 can be relieved by the transcription activator
LeuO. Mol. Microbiol. *77*, 1380–1393. https://doi.org/10.1111/j.1365-2958.2010.07315.x.

22.   Kang, D.-Y., Kim, A., and Kim, J.N. (2023). CcpA and CodY Regulate CRISPR-Cas
System   of   Streptococcus   mutans.   Microbiol.   Spectr.,   e0182623.
https://doi.org/10.1128/spectrum.01826-23.

23.   Smith, L.M., Hampton, H.G., Yevstigneyeva, M.S., Mahler, M., Paquet, Z.S.M., and
Fineran, P.C. (2023). CRISPR-Cas immunity is repressed by the LysR-type transcriptional
regulator PigU. Nucleic Acids Res., gkad1165. https://doi.org/10.1093/nar/gkad1165.

24. Zhang, Y., Yang, J., and Bai, G. (2018). Regulation of the CRISPR-Associated Genes by Rv2837c (CnpB) via an Orn-Like Activity in Tuberculosis Complex Mycobacteria. J. Bacteriol. *200*, e00743-17. https://doi.org/10.1128/JB.00743-17.

25. Patterson, A.G., Chang, J.T., Taylor, C., and Fineran, P.C. (2015). Regulation of the Type I-F CRISPR-Cas system by CRP-cAMP and GalM controls spacer acquisition and interference. Nucleic Acids Res. *43*, 6038–6048. https://doi.org/10.1093/nar/gkv517.

26. Smith, L.M., Jackson, S.A., Malone, L.M., Ussher, J.E., Gardner, P.P., and Fineran, P.C. (2021). The Rcs stress response inversely controls surface and CRISPR-Cas adaptive immunity to discriminate plasmids and phages. Nat. Microbiol. *6*, 162–172. https://doi.org/10.1038/s41564-020-00822-7.

27. Medina-Aparicio, L., Rebollar-Flores, J.E., Gallego-Hernández, A.L., Vázquez, A., Olvera, L., Gutiérrez-Ríos, R.M., Calva, E., and Hernández-Lucas, I. (2011). The CRISPR/Cas immune system is an operon regulated by LeuO, H-NS, and leucine-responsive regulatory protein in Salmonella enterica serovar Typhi. J. Bacteriol. *193*, 2396–2407. https://doi.org/10.1128/JB.01480-10.

28. He, F., Vestergaard, G., Peng, W., She, Q., and Peng, X. (2017). CRISPR-Cas type I-A Cascade complex couples viral infection surveillance to host transcriptional regulation in the dependence of Csa3b. Nucleic Acids Res. *45*, 1902–1913. https://doi.org/10.1093/nar/gkw1265.

29. Shmakov, S.A., Barth, Z.K., Makarova, K.S., Wolf, Y.I., Brover, V., Peters, J.E., and Koonin, E.V. (2023). Widespread CRISPR repeat-like RNA regulatory elements in CRISPR-Cas systems. Preprint at bioRxiv, https://doi.org/10.1101/2023.03.03.530964 https://doi.org/10.1101/2023.03.03.530964.

30. Liu, C., Wang, R., Li, J., Cheng, F., Shu, X., Zhao, H., Xue, Q., Yu, H., Wu, A., Wang, L., et al. (2023). Widespread RNA-based cas regulation monitors crRNA abundance and anti-CRISPR proteins. Cell Host Microbe *31*, 1481-1493.e6. https://doi.org/10.1016/j.chom.2023.08.005.

31. Workman, R.E., Pammi, T., Nguyen, B.T.K., Graeff, L.W., Smith, E., Sebald, S.M., Stoltzfus, M.J., Euler, C.W., and Modell, J.W. (2021). A natural single-guide RNA repurposes Cas9 to autoregulate CRISPR-Cas expression. Cell *184*, 675-688.e19. https://doi.org/10.1016/j.cell.2020.12.017.

32. Liu, T., Li, Y., Wang, X., Ye, Q., Li, H., Liang, Y., She, Q., and Peng, N. (2015). Transcriptional regulator-mediated activation of adaptation genes triggers CRISPR de novo spacer acquisition. Nucleic Acids Res. *43*, 1044–1055. https://doi.org/10.1093/nar/gku1383.

33. Maharajan, A.D., Hjerde, E., Hansen, H., and Willassen, N.P. (2022). Quorum Sensing Controls the CRISPR and Type VI Secretion Systems in Aliivibrio wodanis 06/09/139. Front. Vet. Sci. *9*, 799414. https://doi.org/10.3389/fvets.2022.799414.

34. Bilger, R., Migur, A., Wulf, A., Steglich, C., Urlaub, H., and Hess, W.R. (2024). A type III-Dv CRISPR-Cas system is controlled by the transcription factor RpaB and interacts with the DEAD-box RNA helicase CrhR. Cell Rep. *43*, 114485. https://doi.org/10.1016/j.celrep.2024.114485.

35. Kim, H., and Marraffini, L.A. (2024). Cas9 interaction with the tracrRNA nexus modulates the repression of type II-A CRISPR-cas genes. Nucleic Acids Res., gkae597. https://doi.org/10.1093/nar/gkae597.

36. Schunder, E., Rydzewski, K., Grunow, R., and Heuner, K. (2013). First indication for a functional CRISPR/Cas system in Francisella tularensis. Int. J. Med. Microbiol. IJMM *303*, 51–60. https://doi.org/10.1016/j.ijmm.2012.11.004.

37. Fonfara, I., Richter, H., Bratovič, M., Le Rhun, A., and Charpentier, E. (2016). The CRISPR-associated DNA-cleaving enzyme Cpf1 also processes precursor CRISPR RNA. Nature *532*, 517–521. https://doi.org/10.1038/nature17945.

38. Ratner, H.K., and Weiss, D.S. (2020). F. novicida CRISPR-Cas systems can functionally complement each other in DNA defense while providing target flexibility. J. Bacteriol. https://doi.org/10.1128/JB.00670-19.

39. Ratner, H.K., and Weiss, D.S. (2021). crRNA complementarity shifts endogenous CRISPR-Cas systems between transcriptional repression and DNA defense. RNA Biol., 1–14. https://doi.org/10.1080/15476286.2021.1878335.

40. Sampson, T.R., Saroj, S.D., Llewellyn, A.C., Tzeng, Y.-L., and Weiss, D.S. (2013). A CRISPR/Cas system mediates bacterial innate immune evasion and virulence. Nature *497*, 254–257. https://doi.org/10.1038/nature12048.

41. Sampson, T.R., Saroj, S.D., Llewellyn, A.C., Tzeng, Y.-L., and Weiss, D.S. (2019). Author Correction: A CRISPR/Cas system mediates bacterial innate immune evasion and virulence. Nature *570*, E30–E31. https://doi.org/10.1038/s41586-019-1253-9.

42. Sampson, T.R., Napier, B.A., Schroeder, M.R., Louwen, R., Zhao, J., Chin, C.-Y., Ratner, H.K., Llewellyn, A.C., Jones, C.L., Laroui, H., et al. (2014). A CRISPR-Cas system enhances envelope integrity mediating antibiotic resistance and inflammasome evasion. Proc. Natl. Acad. Sci. U. S. A. *111*, 11163–11168. https://doi.org/10.1073/pnas.1323025111.

43. Ratner, H.K., Escalera-Maurer, A., Le Rhun, A., Jaggavarapu, S., Wozniak, J.E., Crispell, E.K., Charpentier, E., and Weiss, D.S. (2019). Catalytically Active Cas9 Mediates Transcriptional Interference to Facilitate Bacterial Virulence. Mol. Cell *75*, 498-510.e5. https://doi.org/10.1016/j.molcel.2019.05.029.

44. Bell, R.T., Wolf, Y.I., and Koonin, E.V. (2020). Modified base-binding EVE and DCD domains: striking diversity of genomic contexts in prokaryotes and predicted involvement in a variety of cellular processes. BMC Biol. *18*, 159. https://doi.org/10.1186/s12915-020-00885-2.

45. Tan, D., Svenningsen, S.L., and Middelboe, M. (2015). Quorum Sensing Determines the Choice of Antiphage Defense Strategy in Vibrio anguillarum. mBio *6*, e00627. https://doi.org/10.1128/mBio.00627-15.

46. Gao, R., Krysciak, D., Petersen, K., Utpatel, C., Knapp, A., Schmeisser, C., Daniel, R., Voget, S., Jaeger, K.-E., and Streit, W.R. (2015). Genome-wide RNA sequencing analysis of quorum sensing-controlled regulons in the plant-associated Burkholderia glumae PG1 strain. Appl. Environ. Microbiol. *81*, 7993–8007. https://doi.org/10.1128/AEM.01043-15.

47. Dela Ahator, S., Liu, Y., Wang, J., and Zhang, L.-H. (2022). The virulence factor regulator and quorum sensing regulate the type I-F CRISPR-Cas mediated horizontal gene

transfer in Pseudomonas aeruginosa. Front. Microbiol. *13*, 987656. https://doi.org/10.3389/fmicb.2022.987656.

48.   Nuñez, J.K., Bai, L., Harrington, L.B., Hinder, T.L., and Doudna, J.A. (2016). CRISPR Immunological Memory Requires a Host Factor for Specificity. Mol. Cell *62*, 824–833. https://doi.org/10.1016/j.molcel.2016.04.027.

49.   Ali Azam, T., Iwata, A., Nishimura, A., Ueda, S., and Ishihama, A. (1999). Growth phase-dependent variation in protein composition of the Escherichia coli nucleoid. J. Bacteriol. *181*, 6361–6370. https://doi.org/10.1128/JB.181.20.6361-6370.1999.

50.   Kretz, J., Börner, J., Friedrich, T., McIntosh, M., Procida-Kowalski, T., Gerken, F., Wilhelm, J., and Klug, G. (2024). Function of the RNA-targeting class 2 type VI CRISPR Cas system of Rhodobacter capsulatus. Front. Microbiol. *15*, 1384543. https://doi.org/10.3389/fmicb.2024.1384543.

51.   Høyland-Kroghsbo, N.M., Maerkedahl, R.B., and Svenningsen, S.L. (2013). A quorum-sensing-induced bacteriophage defense mechanism. mBio *4*, e00362-00312. https://doi.org/10.1128/mBio.00362-12.

52.   Gallagher, L.A., McKevitt, M., Ramage, E.R., and Manoil, C. (2008). Genetic dissection of the Francisella novicida restriction barrier. J. Bacteriol. *190*, 7830–7837. https://doi.org/10.1128/JB.01188-08.

53.   Galkin, A., Sarikaya, E., Lehmann, C., Howard, A., and Herzberg, O. (2004). X-ray structure of HI0817 from Haemophilus influenzae: protein of unknown function with a novel fold. Proteins *57*, 874–877. https://doi.org/10.1002/prot.20260.

54.   Sonnabend, M.S., Klein, K., Beier, S., Angelov, A., Kluj, R., Mayer, C., Groß, C., Hofmeister, K., Beuttner, A., Willmann, M., et al. (2020). Identification of Drug Resistance Determinants in a Clinical Isolate of Pseudomonas aeruginosa by High-Density Transposon Mutagenesis. Antimicrob. Agents Chemother. *64*, e01771-19. https://doi.org/10.1128/AAC.01771-19.

55.   Eggers, O., Renschler, F.A., Michalek, L.A., Wackler, N., Walter, E., Smollich, F., Klein, K., Sonnabend, M.S., Egle, V., Angelov, A., et al. (2023). YgfB increases β-lactam resistance in Pseudomonas aeruginosa by counteracting AlpA-mediated ampDh3 expression. Commun. Biol. *6*, 254. https://doi.org/10.1038/s42003-023-04609-4.

56.   Gallagher, L.A., Ramage, E., Jacobs, M.A., Kaul, R., Brittnacher, M., and Manoil, C. (2007). A comprehensive transposon mutant library of Francisella novicida, a bioweapon surrogate. Proc. Natl. Acad. Sci. U. S. A. *104*, 1009–1014. https://doi.org/10.1073/pnas.0606713104.

57.   Surette, M.G., and Bassler, B.L. (1998). Quorum sensing in Escherichia coli and Salmonella typhimurium. Proc. Natl. Acad. Sci. U. S. A. *95*, 7046–7050. https://doi.org/10.1073/pnas.95.12.7046.

58.   van Hoek, M.L., Hoang, K.V., and Gunn, J.S. (2019). Two-Component Systems in Francisella Species. Front. Cell. Infect. Microbiol. *9*, 198. https://doi.org/10.3389/fcimb.2019.00198.

59. Lindgren, H., Shen, H., Zingmark, C., Golovliov, I., Conlan, W., and Sjöstedt, A. (2007). Resistance of Francisella tularensis strains against reactive nitrogen and oxygen species with special reference to the role of KatG. Infect. Immun. *75*, 1303–1309. https://doi.org/10.1128/IAI.01717-06.

60. Brodmann, M., Dreier, R.F., Broz, P., and Basler, M. (2017). Francisella requires dynamic type VI secretion system and ClpB to deliver effectors for phagosomal escape. Nat. Commun. *8*, 15853. https://doi.org/10.1038/ncomms15853.

61. Brodmann, M., Heilig, R., Broz, P., and Basler, M. (2018). Mobilizable Plasmids for Tunable Gene Expression in Francisella novicida. Front. Cell. Infect. Microbiol. *8*. https://doi.org/10.3389/fcimb.2018.00284.

62. Harms, A., Segers, F.H.I.D., Quebatte, M., Mistl, C., Manfredi, P., Körner, J., Chomel, B.B., Kosoy, M., Maruyama, S., Engel, P., et al. (2017). Evolutionary Dynamics of Pathoadaptation Revealed by Three Independent Acquisitions of the VirB/D4 Type IV Secretion System in Bartonella. Genome Biol. Evol. *9*, 761–776. https://doi.org/10.1093/gbe/evx042.

63. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image analysis. Nat. Methods *9*, 676–682. https://doi.org/10.1038/nmeth.2019.

64. Aranda, P.S., LaJoie, D.M., and Jorcyk, C.L. (2012). Bleach gel: a simple agarose gel for analyzing RNA quality. Electrophoresis *33*, 366–369. https://doi.org/10.1002/elps.201100335.

65. Rohmer, L., Fong, C., Abmayr, S., Wasnick, M., Larson Freeman, T.J., Radey, M., Guina, T., Svensson, K., Hayden, H.S., Jacobs, M., et al. (2007). Comparison of Francisella tularensis genomes reveals evolutionary events associated with the emergence of human pathogenic strains. Genome Biol. *8*, R102. https://doi.org/10.1186/gb-2007-8-6-r102.

66. Magoc, T., Wood, D., and Salzberg, S.L. (2013). EDGE-pro: Estimated Degree of Gene Expression in Prokaryotic Genomes. Evol. Bioinforma. Online *9*, 127–136. https://doi.org/10.4137/EBO.S11250.

67. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550. https://doi.org/10.1186/s13059-014-0550-8.

68. Llewellyn, A.C., Jones, C.L., Napier, B.A., Bina, J.E., and Weiss, D.S. (2011). Macrophage replication screen identifies a novel Francisella hydroperoxide resistance protein involved in virulence. PloS One *6*, e24201. https://doi.org/10.1371/journal.pone.0024201.

69. Chaparian, R.R., and van Kessel, J.C. (2021). Promoter Pull-Down Assay: A Biochemical Screen for DNA-Binding Proteins. Methods Mol. Biol. Clifton NJ *2346*, 165–172. https://doi.org/10.1007/7651_2020_307.

70. Jutras, B.L., Verma, A., and Stevenson, B. (2012). Identification of novel DNA-binding proteins using DNA-affinity chromatography/pull down. Curr. Protoc. Microbiol. *Chapter 1*, Unit1F.1. https://doi.org/10.1002/9780471729259.mc01f01s24.

635 71. van Kempen, M., Kim, S.S., Tumescheit, C., Mirdita, M., Lee, J., Gilchrist, C.L.M.,
636 Söding, J., and Steinegger, M. (2023). Fast and accurate protein structure search with Foldseek.
637 Nat. Biotechnol., 1–4. https://doi.org/10.1038/s41587-023-01773-0.

Figure 1
CRISPR nucleases Cas9 and Cas 12 are differentially expressed during exponential and stationary growth.
(A) Fluorescence microscopy of *cas9-msfGFP* and *cas12-mCherry2*. Fields of view of 13 x 13 μm are shown. Upper panel is a merge of phase contrast and GFP channel, lower panel is a merge of phase contrast and RFP channel. Scale bars represent 1 μm.
(B) RNA-seq data of differentially expressed genes in stationary and exponential growth phase. Green: *type II-B CRISPR-Cas* genes. Magenta: *type V-A CRISPR-Cas* genes. Dotted horizontal line: *P*-value of 0.01. Dotted vertical line: 4-fold difference.
(C-D) Determination of mRNA levels for *cas9* **(C)** and *cas12* **(D)** by RT-qPCR. Grey bars: Parental strain. Mean with standard deviation is displayed. Green bars: *cas9-msfGFP*. Magenta bars: *cas12-mCherry2*. Significant differences were determined with Tukey's multiple comparison test. **** *AdjP*-value < 0.0001.

Supplemental figure 1

C-terminal fluorescent tags do not inhibit Cas9- and Cas12-mediated plasmid targeting.

(A) Type II-B CRISPR-Cas system in *F. novicida U112*. Green arrows: *cas* genes. Grey box: l*eader/tracr sequence*. Black diamonds: *repeats*. Orange box: *spacers*. White box: *scaRNA*. Drawn in scale.

(B) Type V-A CRISPR-Cas system in *F. novicida U112*. White diamond: *single repeat unit*. Magenta arrows: *cas* genes. Black diamonds: *repeats*. Orange box: *spacers*. Drawn in scale.

(C) Assessment of interference capability of wildtype and C-terminal tagged Cas9 and Cas12 in a conjugation-based plasmid inhibition assay. Strains have either both native type II-B and type V-A CRISPR arrays or have either CRISPR array replaced by a spacer matching a protospacer on the conjugated plasmid (targeting spacer) or one that does not match (non-targeting spacer). Mean with standard deviation is displayed. Filled circles: total number of cells. Empty circles: transformants. Grey circles: Parental strain. Green circles: *cas9-msfGFP*. Magenta circles: *cas12-mCherry2*.

Figure 2

Cas12 and SRU are not involved in regulating *cas12* in stationary phase.

(A) CRISPR-like single repeat unit (SRU) is partially homologous to the type V-A CRISPR repeats. Red box: CRISPR-repeat stem loop; green box: cleavage site of Cas12 during pre-crRNA processing.

(B) Fluorescence microscopy of *cas12::mCherry2*. A merge of phase contrast and RFP channel in 13 x 13 µm fields of view are shown. Scale bars represent 1 µm.

(C) Determination of mRNA levels for *cas12* by RT-qPCR. Grey bars: Parental strain. Olive bars: Δ*SRU* mutant. Mean with standard deviation. Significant differences were determined with Tukey's multiple comparison test. ** *AdjP*-value < 0.005, *** *AdjP*-value < 0.001.

(D) Assessment of interference capability of parental strain and a Δ*SRU* mutant in a conjugation-based plasmid inhibition assay. Strains have either a native type V-A CRISPR array or have the CRISPR array replaced by a spacer matching a protospacer on the conjugated plasmid (targeting spacer) or one that does not match (non-targeting spacer). Mean with standard deviation is displayed. Filled circles: total number of cells. Empty circles: transformants. Grey circles: Parental strain. Olive circles: Δ*SRU* mutant.

Figure 3

MtvS modulates expression of the Type V-A CRISPR-Cas system in *F. novicida* U112.

(A) Pull-down of proteins from exponential growth phase lysate binding specifically to the *cas12* promoter DNA sequence. Turquoise circle: MtvS. Dark blue circle: FTN_0392. Dotted horizontal line: *P*-value of 0.01. Dotted vertical line: 4-fold difference.

(B) Pull-down of proteins from stationary phase lysate binding specifically to the *cas12* promoter DNA sequence. Turquoise circle: MtvS. Dark blue circle: FTN_0392. Dotted horizontal line: *P*-value of 0.01. Dotted vertical line: 4-fold difference.

(C) Fluorescence microscopy of *cas12-mCherry2* in parental strain, *mtvS nonsense* mutant and *mtvS nonsense* mutant with wild-type *mtvS* at Tn7 insertion site. *mtvS nonsense* mutant results in decreased fluorescence intensity in stationary phase compared to parental strain and rescued mutant. Merge of phase contrast and RFP channel in 13 x 13 μm fields of view are shown. Scale bars represent 1 μm.

(D-G) Determination of mRNA levels for *cas12* **(C)** *cas4* **(D)**, *cas1* **(E)** and *cas2* **(F)** by RT-qPCR. Grey bars: Parental strain. Turquoise bars: *mtvS nonsense* mutant. Light blue bars: *mtvS nonsense* mutant with wild-type *mtvS* at Tn7 insertion site. Mean with standard deviation is displayed. Significant differences were determined with Tukey's multiple comparison test. * *AdjP*-value < 0.05, *** *AdjP*-value < 0.001, **** *AdjP*-value < 0.0001.

Supplemental figure 2
MtvS is a YgfB/YecA-like protein.

(A) Fluorescence microscopy of *cas12-mCherry2* in parental strain and *ΔFTN_0392* mutant. Comparable fluorescent intensities in stationary growth were detected. Merge of phase contrast and RFP channel in 13 x 13 μm fields of view are shown. Scale bars represent 1 μm.

(B) Assessment of interference capability of parental strain and *ΔFTN_0392* mutant in a conjugation-based plasmid inhibition assay. Strains have either a native type V-A CRISPR array or have the CRISPR array replaced by a spacer matching a protospacer on the conjugated plasmid (targeting spacer) or one that does not match (non-targeting spacer). Mean with standard deviation is displayed. Filled circles: total number of cells. Empty circles: transformants. Grey circles: Parental strain. Dark blue circles: *ΔFTN_0392* mutant.

(C) *mtvS* is the first gene of a conserved operon which also includes *ubiH* (2-octaprenyl-6-methoxyphenyl hydroxylase), *ubiF* (2-octaprenyl-3-methyl-6-methoxy-1,4-benzoquinol hydroxylase), *FTN_1235* (putative phage protein) and *queA* (S-adenosylmethionine:tRNA ribosyltransferase-isomerase).

(D) Comparison of Alphafold MtvS structure (blue) with *E. coli* YecA (yellow) by Foldseek.

(E) Assessment of interference capability of parental strain and *mtvS nonsense* mutant in a conjugation-based plasmid inhibition assay. Strains have either a native type V-A CRISPR array or have the CRISPR array replaced by a spacer matching a protospacer on the conjugated plasmid (targeting spacer) or one that does not match (non-targeting spacer). Mean with standard deviation is displayed. Filled circles: total number of cells. Empty circles: transformants. Grey circles: Parental strain. Turquoise circles: *mtvS nonsense* mutant.

(F) Fluorescence microscopy of *mtvS-mCherry2* in parental strain. Fluorescence intensity is higher in stationary phase than in exponential growth phase. Merge of phase contrast and RFP channel in 13 x 13 μm fields of view are shown. Scale bars represent 1 μm.

(G) Determination of mRNA levels for *mtvS* in the parental strain by RT-qPCR. Grey bars: Parental strain. Mean with standard deviation is displayed. Significant differences were determined with Tukey's multiple comparison test.

Figure 4

MtvS modulates *cas12* expression in a cell-density dependent manner

(A) Fluorescence microscopy of *cas12-mCherry2* in microfluidic plate. BHI was flown through for 8 h. Merge of phase contrast and RFP channel in 13 x 13 µm fields of view are shown. Same contrast settings as in **(B)**. Scale bars represent 1 µm.

(B) Fluorescence microscopy of *cas12-mCherry2* in *mtvS nonsense* mutant in microfluidic plate. BHI was flown through for 8 h. Merge of phase contrast and RFP channel in 13 x 13 µm fields of view are shown. Same contrast settings as in (**A**). Scale bars represent 1 µm.

(C-D) Fluorescence microscopy of *cas12-mCherry2* in microfluidic plate. BHI was flown through for 1 h before switching to spent overnight medium of parental strain for 4h. Afterwards, fresh BHI was flown through for 3 h. **(C)** Three examples of 240 min time point (180 min after switching to spent ON medium) and **(D)** three examples of 480 min (180 min after switching back to fresh BHI) are shown. Merge of phase contrast and RFP channel in 4 x 4 µm fields of view are shown. Same contrast settings as in (**E-F**) and **supplemental figure 3 (A)**. Scale bars represent 1 µm.

(E-F) Fluorescence microscopy of *cas12-mCherry2* in *mtvS nonsense* mutant in microfluidic plate. BHI was flown through for 1 h before switching to spent overnight medium of parental strain for 4h. Afterwards, fresh BHI was flown through for 3 h. **(E)** Three examples of 240 min time point (180 min after switching to spent ON medium) and **(F)** three examples of 480 min (180 min after switching back to fresh BHI) are shownare shown. Merge of phase contrast and RFP channel in 4 x 4 µm fields of view are shown. Same contrast settings as in (**C-D**) and **supplemental figure 3 (A)**. Scale bars represent 1 µm.

(G) Determination of mRNA levels for *cas12* after switching to spent overnight medium by RT-qPCR. Samples were taken 0 h, 0.5 h and 1 h after medium change. Grey bars: Parental strain. Turquoise bars: *mtvS nonsense* mutant. Light blue bars: *mtvS nonsense* mutant with wild-type *mtvS* at Tn7 insertion site. Mean with standard deviation is displayed. Significant differences were determined with Tukey's multiple comparison test. * *AdjP*-value < 0.05, *** *AdjP*-value < 0.001, **** *AdjP*-value < 0.0001.

Supplement figure 3

Spent overnight medium is heat sensitive and does not affect *cas9* expression.

(A) Fluorescence microscopy of *cas12-mCherry2* in microfluidic plate. BHI was flown through for 1 h before switching to heat-treated spent overnight medium of parental strain for 4h. Afterwards fresh BHI was flown through for 3 h. Three examples of 240 min time point (180 min after switching to spent ON medium) are shown. Merge of phase contrast and RFP channel in 4 x 4 µm fields of view are shown. Same contrast settings as in Figs. **3 (C-F)**. Scale bars represent 1 µm.

(B) Determination of mRNA levels for *cas9* after switching to spent overnight medium by RT-qPCR. Samples were taken 0 h, 0.5 h and 1 h after medium change. Grey bars: Parental strain. Turquoise bars: *mtvS nonsense* mutant. Light blue bars: *mtvS nonsense* mutant with wild-type *mtvS* at Tn7 insertion site. Mean with standard deviation is displayed. Significant differences were determined with Tukey's multiple comparison test.

Table 1: Strains used in this study, related to Material and methods

Organism	Genotype	Plasmid	Relevant features	Source
Francisella novicida U112			Parental strain	Bei resources (NR-13)
	typeII_tet_spacer_targ		Type II-B wild-type CRISPR array was replaced with targeting spacer matching *tetR* on plasmid pFNMB2	This study
	typeII_tet_spacer_nontarg		Type II-B wild-type CRISPR array was replaced with non-targeting spacer matching *tetR* on plasmid pFNMB2	This study
	typeV_tet_spacer_targ		Type V-A wild-type CRISPR array was replaced with targeting spacer matching *tetR* on plasmid pFNMB2	This study
	typeV_tet_spacer_nontarg		Type V-A wild-type CRISPR array was replaced with non-targeting spacer matching *tetR* on plasmid pFNMB2	This study
	mtvS nonsense		Three stop codons introduced at position 11 of *mtvs*	This study
	mtvS nonsense, Tn7_mtvS		Wild-type *mtvS* was introduced at the canonical Tn7 insertion site downstream of *glmS*	This study
	mtvS nonsense, typeV_tet_spacer_targ		Type V-A wild-type CRISPR array was replaced with targeting spacer matching *tetR* on plasmid pFNMB2	This study
	mtvs nonsense, typeV_tet_spacer_nontarg		Type V-A wild-type CRISPR array was replaced with non-targeting spacer matching *tetR* on plasmid pFNMB2	This study
	Δ*SRU*		Deletion of single repeat unit (SRU) in *cas12* promoter	This study
	Δ*SRU, typeV_tet_spacer_targ*		Type V-A wild-type CRISPR array was replaced with targeting spacer matching *tetR* on plasmid pFNMB2	This study
	Δ*SRU, typeV_tet_spacer_nontarg*		Type V-A wild-type CRISPR array was replaced with non-targeting spacer matching *tetR* on plasmid pFNMB2	This study
	ΔFTN_0392		Deletion of FTN_0392	This study
	ΔFTN_0392, *typeV_tet_spacer_targ*		Type V-A wild-type CRISPR array was replaced with targeting spacer matching *tetR* on plasmid pFNMB2	This study
	ΔFTN_0392, *typeV_tet_spacer_nontarg*		Type V-A wild-type CRISPR array was replaced with non-targeting spacer matching *tetR* on plasmid pFNMB2	This study
	cas9-msfGFP, cas12-mCherry2		C-terminal chromosomal fusion of *msfgfp* to *cas9* and c-terminal chromosomal fusion of *mCherry2* to *cas12*	This study
	cas9-msfGFP, cas12-mCherry2, typeV_tet_spacer_targ		Type V-A wild-type CRISPR array replaced with targeting spacer matching *tetR* on plasmid pFNMB2	This study
	cas9-msfGFP, cas12-mCherry2, typeV_tet_spacer_nontarg		Type V-A wild-type CRISPR array replaced with non-targeting spacer matching *tetR* on plasmid pFNMB2	This study
	cas9-msfGFP, cas12-mCherry2, typeII_tet_spacer_targ		Type II-B wild-type CRISPR array replaced with targeting spacer matching *tetR* on plasmid pFNMB2	This study
	cas9-msfGFP, cas12-mCherry2, typeII_tet_spacer_nontarg		Type II-B wild-type CRISPR array replaced with non-targeting spacer matching *tetR* on plasmid pFNMB2	This study
	cas9-msfGFP, cas12-mCherry2, mtvs nonsense		Three stop codons introduced at position 11 of *mtvS*	This study
	cas12::mCherry2		*cas12* was replaced with *mCherry2*	This study
Escherichia coli OneShotpir			Cloning strain	ThermoFisher
Escherichia coli JKE 201			Donor *E. coli* used for conjugation into *F. novicida*	[1]
		pFNMB2 *msfGFP*	Donor *E. coli* containing mobilizable *Francisella* plasmid pFNMB2 *msfGFP*	[2]

Table 2: Plasmids and primers used to generate mutants, related to Material and Methods.

Plasmid Name	Spacer/Peptide sequence	Primers	Sequence 5'-3' [base pairs]
pDMK3 *typeII_tet_spacer_targ*	TCGCGATGACTTAGTAAAGCACATCTAAAACTTT	TypeII_spacer_1.FOR	CTGAATTGTCGACCCTCGAGATAGTTGACTGGGTTATAG
		TypeII_spacer_tet_targ_1.REV	AAAGTTTTAGATGTGCTTTACTAAGTCATCGCGAGTTTCAGTTGCTGAATTATTTGGTAAACTACTGTTAGAGCCAAAACAAACCTAGTGT
		TypeII_spacer_tet_targ_2.FOR	TCGCGATGACTTAGTAAAGCACATCTAAAACTTTCTAACAGTAGTTTACCAAATAATTCAGCAACTGAAACTTAATTAATTTTGTAAATTA
		TypeII_spacer_2.REV	ATCTATCTAGAAGGGCCCCGTTTCACTAAATTCTTA
		TypeII_spacer_Det.FOR	ATGGGTTGCTCAAGGTGAAG
		TypeII_spacer_Det.REV	TCTCATGCTGGTTACGCTTG
pDMK3 *typeII_tet_spacer_nontarg*	AAAGTTTTAGATGTGCTTTACTAAGTCATCGCGA	TypeII_spacer_1.FOR	CTGAATTGTCGACCCTCGAGATAGTTGACTGGGTTATAG
		TypeII_spacer_tet_nontarg_1.REV	TCGCGATGACTTAGTAAAGCACATCTAAAACTTTGTTTCAGTTGCTGAATTATTTGGTAAACTACTGTTAGAGCCAAAACAAACCTAGTGT
		TypeII_spacer_tet_nontarg_2.FOR	AAAGTTTTAGATGTGCTTTACTAAGTCATCGCGACTAACAGTAGTTTACCAAATAATTCAGCAACTGAAACTTAATTAATTTTGTAAATTA
		TypeII_spacer_2.REV	ATCTATCTAGAAGGGCCCCGTTTCACTAAATTCTTA
		TypeII_spacer_Det.FOR	ATGGGTTGCTCAAGGTGAAG
		TypeII_spacer_Det.REV	TCTCATGCTGGTTACGCTTG
pDMK3 *typeV_tet_spacer_targ*	TCTAACATCTCAATGGCTAAGGCGTCGAG	TypeV_spacer_1.FOR	TCTATCGCCTTCTTGACGAGTTCTTCTGAATTGTCGACCCTCGAGATGCTGTTTCCAAGC
		TypeV_spacer_tet_targ_1.REV	CTCGACGCCTTAGCCATTGAGATGTTAGAGTCTAAGAACTTTAAATAATTTGTCTGTATATTATTGATTTCTAAATTAGAATTTT
		TypeV_spacer_tet_targ_2.FOR	TCTAACATCTCAATGGCTAAGGCGTCGAGATCTACAACAGTAGAAATTATTTAAAGTTCTTAGACCCGTTTTTGCCTAAATCAGC
		TypeV_spacer_2.REV	TTGTGAGCGGATAACAATTTGTGGAATTCCCGGGAGAGCTCCCTAGTTTGTGACTTTGTT
		TypeV_spacer_Det.FOR	GCCAGCGAATAAAAGTCCTG
		TypeV_spacer_Det.REV	CAAGCCTAGCGGAGCTAATG
pDMK3 *typeV_tet_spacer_nontarg*	CTCGACGCCTTAGCCATTGAGATGTTAGA	TypeV_spacer_1.FOR	TCTATCGCCTTCTTGACGAGTTCTTCTGAATTGTCGACCCTCGAGATGCTGTTTCCAAGC
		TypeV_spacer_tet_nontarg_1.REV	TCTAACATCTCAATGGCTAAGGCGTCGAGGTCTAAGAACTTTAAATAATTTGTCTGTATATTATTGATTTCTAAATTAGAATTTT
		TypeV_spacer_tet_nontarg_2.FOR	CTCGACGCCTTAGCCATTGAGATGTTAGAATCTACAACAGTAGAAATTATTTAAAGTTCTTAGACCCGTTTTTGCCTAAATCAGC
		TypeV_spacer_2.REV	TTGTGAGCGGATAACAATTTGTGGAATTCCCGGGAGAGCTCCCTAGTTTGTGACTTTGTT
		TypeV_spacer_Det.FOR	GCCAGCGAATAAAAGTCCTG
		TypeV_spacer_Det.REV	CAAGCCTAGCGGAGCTAATG
pDMK3 *mtvS nonsense*	MVKLA***DEFNP	FTN_1238_ns_1.FOR	CTTCTGAATTGTCGACCCTCGAGCCAAATAAGCCTACTCCT
		FTN_1238_ns_1.REV	AGATTAGTAGTAGGCATTAAAAGTAATGCAA
		FTN_1238_ns_2.FOR	TACTTTTAATGCCTACTACTAATCTTCAAAATTAGGTTTTT
		FTN_1238_ns_2.REV	ATAACAATTTGTGGAATTCCCGGGAGAGCTCAGGTTACGCGGCCGCCCACCCATGCCCATT
		FTN_1238_ns_Det.FOR	CCAACTATCCCACCCCCAAC

		FTN_1238_ns_Det.REV	TTTGCTGCTCATCAAAGGCT
		FTN_1238_ns_Det2.FOR	CCTACTACTAATCTTCAAAATTAG
pDMK3 *Tn7_mtvS*		FTN_0485_Tn7_1.FOR	CTTCTGAATTGTCGACCCTCGAGAATGTGCCAAATAGTTCG
		FTN_1238_Tn7_1.REV	TTTAGGTATTAACTTGGAAAATAAATGATAAGAAAGTAAAA
		FTN_1238_Tn7_2.FOR	TTTATTTTCCAAGTTAATACCTAAATTTTTTCACTAT
		FTN_1238_Tn7_2.REV	ATTTAGCTAAGTTTAAACTCTTGATTACATAGCC
		FTN_1238_Tn7_3.FOR	ATCAAGAGTTTAAACTTAGCTAAATCTGTAACC
		FTN_0485_Tn7_3.REV	AGAGCTCAGGTTACGCGGCCGCGTATCAAGCAAGAGCTTAG
		FTN_0485_Tn7_Det.FOR	TCGTTGCATGTGGAACTAGCT
		FTN_0485_Tn7_Det.REV	TGGTGAAACACTAGCTGAAGGT
pDMK3 Δ*SRU*	Deleted: TATTATAAGAATCTTTATTTAGTTTA TAATTTTGTAGAT	dSRU_1.FOR	CTTCTGAATTGTCGACCCTCGAGAATTTGTTCATAGTTTATAGCTT
		dSRU_1.REV	CAAAATTTTTTAGACTTTATATATAATCGAAATGTAGAG
		dSRU_2.FOR	TATATATAAAGTCTAAAAAATTTTGCCAAACTCA
		dSRU_2.REV	AGAGCTCAGGTTACGCGGCCGCATTTTGATAAAGGTAAAAGAGC
		dSRU_Det.FOR	ACCGCTGTACCAATAACACTAT
		dSRU_Det.REV	ACACTTGCTGATGTTGGTGA
pDMK3 Δ*FTN_0392*	MRITLKQLQVFIEKDINGCDIRLENFSI *	dFTN_0392_1.FOR	TCTATCGCCTTCTTGACGAGTTCTTCTGAATTGTCGACCCTCGAGAAGTGTTC TTACGTACA
		dFTN_0392_1.REV	CTCTAAAACAGCTTCAAGTGTTTATAGAGAAAGATATAAATGGCTG
		dFTN_0392_2.FOR	TTTATATCTTTCTCTATAAACACTTGAAGCTGTTT
		dFTN_0392_2.REV	ATAACAATTTGTGGAATTCCCGGGAGAGCTCAGGTTACGCGGCCGCGTGAG ACGAACCAAAAAC
		dFTN_0392_Det.FOR	TGCAAATTCACTATAGCGCCA
		dFTN_0392_Det.REV	TTGCAACATCCTCAACCCCA
pDMK3 *cas9-msfGFP*		FTN_0757-fusion_1.FOR	ATTGTCGACCCTCGAGCAAGCAGGTTTTGGATAAG
		FTN_0757-fusion_1.REV	GCGGCCGCATTATTAGATGTTTCATTATAAATACCT
		FTN_0757-fusion_2.FOR	TCTAATAATGCGGCCGCAGGA
		FTN_0757-msfGFP_2.REV	TACCAAGCATTTTGTAGAGCTCATCCATG
		FTN_0757-msfGFP_3.FOR	GCTCTACAAAATGCTTGGTATGAAATTAG
		FTN_0757-fusion_3.REV	ACAATTTGTGGAATTCCCGGGCGATACTCCTCCATTAGA
		FTN_0757-fusion_Det.FOR	TCTAAAAACCCAAAAATTACATGAG
		FTN_0757-fusion_Det.REV	TGTTCACCAGCAAATTCTCCAG
pDMK3 *cas12-mCherry2*		FTN_1397-fusion_1.FOR	CTGAATTGTCGACCCTCGAGCAAAAATATTAACAAAAACTATATTCTTT
		FTN_1397-mCherry2_1.REV	TGTACAAGTTATTTGAGTTCGTGCAGA
		FTN_1397-mCherry2_2.FOR	ACGAACTCAAATAACTTGTACAGCTCGT
		FTN_1397-fusion_2.REV	GAATAACGCGGCCGCAGGA
		FTN_1397-fusion_3.FOR	CGGCCGCGTTATTCCTATTCTGCACG
		FTN_1397-fusion_3.REV	CAATTTGTGGAATTCCCGGGGTGCTTAGAGCTTATCAG
		FTN_1397-fusion_Det.FOR	TCTCTTGCTGCTCTGGAGTT
		FTN_1397-fusion_Det.REV	AAAAGAGGGCGTTTCAAGGT
pDMK3 *cas12::mCherry2*		FTN_1397::mCherry2_1.FOR	CTTCTGAATTGTCGACCCTCGAGAGTCCGTTTTATCACATAAG
		FTN_1397::mCherry2_1.REV	TGTACAAGTAAGGTAGGATCAAAAATAATCA
		FTN_1397::mCherry2_2.FOR	TGATCCTACCTTACTTGTACAGCTCGTC
		FTN_1397::mCherry2_2.REV	AGTCTTTATCATGGTGAGCAAGGGC
		FTN_1397::mCherry2_3.FOR	GCTCACCATGATAAAGACTCCTTATAAAATT
		FTN_1397::mCherry2_3.REV	TCAGTAGCGGCCGCATTTTGATAAAGGTAAAAGAGC

FTN_1397::mCherry2_Det.FOR	TCTCTTGCTGCTCTGGAGTT
FTN_1397::mCherry2_Det.REV	ACACTTGCTGATGTTGGTGA

Table 3: Primers used for RT-qPCR, related to Material and methods.

Primers	Sequence 5'-3' [base pairs]
FTN_0757_qPCR. FOR	ATGCTAACACTTGTGCAGTTTG
FTN_0757_qPCR.REV	GTCTCTGAGCTTTGGCACTTA
FTN_1397_qPCR.FOR	AAGCTCTGCCTCACCATTTAG
FTN_1397_qPCR.REV	AAGGGCGACCAAATCTACATAC
FTN_1396_qPCR.FOR	GCCCATCTCTTGGAGACAAA
FTN_1396_qPCR.REV	GGGCTTACTTGTGGAGAGAAA
FTN_1395_qPCR.FOR	TATTCTAGGTTTGCGCCCTTG
FTN_1395_qPCR.REV	CAAGCCTAGCGGAGCTAATG
FTN_1394_qPCR.FOR	TCACATCTTTATCTGGAGCATT
FTN_1394_qPCR.REV	TGGTGTACGTTTACAATATTCGG
FTN_1238_qPCR.FOR	CACCCTCTTCAATCGGCTTAG
FTN_1238_qPCR.REV	GGACTGCTTTGTGCTTTGTTT

Table 4: Primers used for promoter sequence pull-down

Primers	Modification	Sequence 5'-3' [base pairs]
Promoter-FTN_1397.FOR		CTAGCTAGTTTAAAAGGACAA
Promoter-FTN_1397_biotin.REV	5' Biotin-TEG	TAACTCAAATCTTAGAGTTTTACTT
FTN_1397_pull-down.FOR		GCTGTACCAATAACACTATAAT
FTN_1397_biotin.REV	5' Biotin-TEG	AGAAAAATCTATTAAAGAAACACTATC

Table 5: Sequencing and EDGE-Pro statistics. RPKM reads were calculated as "Mapped reads"- "rRNA reads" and were the basis to calculate RPKM values. Related to Results.

	Parental strain Exponential growth Replicate 1	Parental strain Exponential growth Replicate 2	Parental strain Exponential growth Replicate 3	Parental strain Stationary phase Replicate 1	Parental strain Stationary phase Replicate 2	Parental strain Stationary phase Replicate 3
Total reads	3724938	3843094	3742820	2405842	2339574	3527853
Mapped reads	3596390	3768817	3666864	2234312	2095603	3262475
Unique reads	2586825	3630328	3559936	2170528	1391360	2017405
Multiple reads	1009565	138489	106928	63784	704243	1245070
rRNA reads	950125	27813	1105	62713	701355	1241682
RPKM reads	2646265	3741004	3665759	2171599	1394248	2020793

References

1. Harms, A., Segers, F.H.I.D., Quebatte, M., Mistl, C., Manfredi, P., Körner, J., Chomel, B.B., Kosoy, M., Maruyama, S., Engel, P., et al. (2017). Evolutionary Dynamics of Pathoadaptation Revealed by Three Independent Acquisitions of the VirB/D4 Type IV Secretion System in *Bartonella*. Genome Biol. Evol. *9*, 761–776. https://doi.org/10.1093/gbe/evx042.

2. Brodmann, M., Heilig, R., Broz, P., and Basler, M. (2018). Mobilizable Plasmids for Tunable Gene Expression in *Francisella novicida.* Front. Cell. Infect. Microbiol. *8*. https://doi.org/10.3389/fcimb.2018.00284.