Contributions of mechanical loading and hormonal changes to eccentric hypertrophy during volume overload: a Bayesian analysis using logic-based network models.

Johane H. Bracamonte[1], Lionel Watkins[2], Betty Pat [3,4], Louis J. Dell'Italia[3,4], Jeffrey J. Saucerman[2], Jeffrey W. Holmes[1,4,5,*]

(1) Department of Biomedical Engineering, University of Alabama at Birmingham, Birmingham, Alabama, United States of America.

(2) Department of Biomedical Engineering, University of Virginia, Charlottesville, Virginia, United States of America.

(3) Birmingham Veterans Affairs Health Care System, Birmingham, Alabama, United States of America.

(4) Division of Cardiovascular Disease, Heersink School of Medicine, University of Alabama at Birmingham, Birmingham, Alabama, United States of America.

(5) Division of Cardiothoracic Surgery, Heersink School of Medicine, University of Alabama at Birmingham, Birmingham, Alabama, United States of America.

(*) Corresponding author. Email: holmesjw@uab.edu

Abstract

Primary mitral regurgitation (MR) is a pathology that alters mechanical loading on the left ventricle, triggers an array of compensatory neurohormonal responses, and induces a distinctive ventricular remodeling response known as eccentric hypertrophy. Drug therapies may alleviate symptoms, but only mitral valve repair or replacement can provide significant recovery of cardiac function and dimensions. Questions remain about the optimal timing of surgery, with 20% of patients developing systolic dysfunction post-operatively despite being treated according to the current guidelines. Thus, better understanding of the hypertrophic process in the setting of ventricular volume overload (VO) is needed to improve and better personalize the management of MR. To address this knowledge gap, we employ a Bayesian approach to combine data from 70 studies on experimental volume overload in dogs and rats and use it to calibrate a logic-based network model of hypertrophic signaling in myocytes. The calibrated model predicts that growth in experimental VO is mostly driven by the neurohormonal response, with an initial increase in myocardial tissue stretch being compensated by subsequent remodeling fairly early in the time course of VO. This observation contrasts with a common perception that volume-overload hypertrophy is driven primarily by increased myocyte strain. The model reproduces many aspects of 43 studies not used in its calibration, including infusion of individual hypertrophic agonists alone or in combination with various drugs commonly employed to treat heart failure, as well as administration of some of those drugs in the setting of experimental volume overload. We believe this represents a promising approach to using the known structure of an intracellular signaling network to integrate information from multiple studies into quantitative

43   predictions of the range of expected responses to potential interventions in the complex

44   setting of cardiac hypertrophy driven by a combination of hormonal and mechanical

45   factors.

Author Summary

Mitral valve regurgitation is a common heart disease in which a malfunctioning valve allows part of the blood pumped by the heart to flow in the wrong direction. This condition overloads the heart by making it pump more blood volume than normal; the heart temporarily adapts by growing in mass and volume, but if untreated the condition can ultimately lead to heart failure and death. The most effective treatment is to surgically repair the valve; however, in many patients heart function deteriorates even after a successful surgery. Many researchers have studied this condition by experimentally overloading the hearts of dogs and rats, producing large amounts of data on the resulting geometric, mechanical, and biologic changes. Yet it has been difficult to translate those studies into effective selection and timing of treatment in all patients. In this work we integrate experimental data reported from 70 research articles on experimental volume overload through a simple model of heart mechanics and a more complex model of the molecular signaling pathways inside heart cells. We use a statistical approach to calibrate the computational model, so that it can predict not only average responses but also the degree of expected uncertainty for each prediction. We then use the model to explore how the heart responds to different combinations of drugs and hormones, and potential treatments during overload.

Statements and Declarations

Funding:

This work is supported by the American Heart Association postdoctoral fellowship grant 23POST1026645 (https://doi.org/10.58275/AHA.23POST1026645.pc.gr.161204) to J.H.B; the National Institutes of Health and National Heart, Lung, and Blood Institute (NIH/NHLBI) grants R01HL159945 to J.W.H, R01HL162925 to J.J.S, and P01 HL051952 and P50HL077100 to L.J.D; Department of Veteran Affairs for Merit Review grant 1CX000993-01 to L.J.D, and Department of Veteran Affairs John B. Barnwell Award to L.J.D.

Competing interest:

The authors have no competing interests to declare that are relevant to the content of this article.

Availability of data and source codes:

Links to GitHub repositories containing all data and codes are provided within the text.

Authors contributions:

Conceptualization: Johane H. Bracamonte, Jeffrey W. Holmes

Data Curation: Johane H. Bracamonte, Betty Pat, Lou J. Dell'Italia

Formal Analysis: Johane H. Bracamonte, Lionel Watkins

Funding Acquisition: Johane H. Bracamonte, Jeffrey W. Holmes, Jeffrey J. Saucerman, Lou J. Dell'Italia

Investigation: Johane H. Bracamonte, Jeffrey W. Holmes, Betty Pat, Lou J. Dell'Italia

Methodology: Johane H. Bracamonte, Jeffrey W. Holmes, Jeffrey J. Saucerman

Project Administration: Jeffrey W. Holmes

Resources: Johane H. Bracamonte, Jeffrey W. Holmes, Jeffrey J. Saucerman, Betty Pat, Lou J. Dell'Italia

Software: Johane H. Bracamonte, Lionel Watkins, Jeffrey J. Saucerman

Supervision: Jeffrey W. Holmes

Validation: Johane H. Bracamonte, Jeffrey W. Holmes

Visualization: Johane H. Bracamonte, Lionel Watkins

Writing-original draft preparation: Johane H. Bracamonte

Writing-review and editing: Johane H. Bracamonte, Jeffrey W. Holmes, Jeffrey J. Saucerman, Lou J. Dell'Italia

1. Introduction

Mitral valve regurgitation affects around 5 million people in America, and about 2% of the general population, with prevalence steeply increasing in individuals over 50 years of age [1]. In primary mitral regurgitation (MR) the dysfunction of one or more components of the valvular apparatus allows part of the blood volume pumped by the left ventricle to flow back to the low-pressure atrial compartment, making the heart pump a larger than usual volume of blood against a lower-than-normal resistance. The unique loading conditions imposed by MR and the resulting neurohormonal responses induce a distinctive ventricular remodeling response known as eccentric hypertrophy, consisting of the lengthening of individual cardiomyocytes by addition of sarcomeres in series, leading to an organ-scale dilation of the left ventricle volume with little change in its wall thickness [2,3]. The neurohormonal response to volume overload is characterized by the activation of the sympathetic and renin-angiotensin systems, similar to other forms of cardiac overloading [4–6]. Drug therapies for heart failure due to primary MR alleviate symptoms and slow its progression, but only mitral valve repair can provide significant recovery of cardiac function and dimensions [2,7]. If MR is severe enough or if it remains untreated for long enough, the condition can transition from a compensated asymptomatic stage into irreversible heart failure with systolic dysfunction, a condition where the heart is unable to supply sufficient cardiac output to the body [8,9]. This risk has led clinicians to operate earlier in the natural course of primary MR; yet 20% of patients still develop systolic dysfunction post-operatively despite being treated according to the current guidelines [10,11]. This fact highlights our incomplete understanding of eccentric hypertrophy due to primary MR and its transition

into systolic dysfunction and heart failure. A better understanding of this process is

needed to improve and better personalize the management of MR.

Most computational models of growth and remodeling during volume overload have

focused on the role of myocyte overstretch in driving eccentric hypertrophy [12,13]. By

contrast, molecular studies have shown reduced activity in stretch-sensitive myocyte

signaling pathways during experimental volume overload, the opposite of what would be

expected if remodeling is driven by stretch [14–18]. We hypothesized that this apparent

paradox might stem from complex interactions between the hypertrophic signaling

pathways triggered by stretching and those that respond to other hormones and growth

factors known to be upregulated during volume overload.

Here, we employ a Bayesian approach to combine the wealth of available data on

experimental volume overload in dogs and rats using a logic-based network model of

hypertrophic signaling in myocytes, with the goal of better understanding the relative

influence of multiple factors that influence eccentric hypertrophy. We employed data

from 70 studies of experimental volume overload to estimate the relative influence of

multiple input parameters for a network model of hypertrophic signaling in

cardiomyocytes during volume overload, accounting for evolving levels of mechanical

strain and circulating hormones such as norepinephrine (NE), angiotensin II (AngII),

endothelin 1 (ET1), and atrial ANP and brain (BNP) natriuretic peptides. We then

validated the ability of the calibrated model to reproduce features of volume overload

not included in the calibration, as well as experimental responses to relevant

independent experiments such as infusion of hormones that induce myocyte

hypertrophy, alone or in combination with receptor blockers used clinically to treat heart

failure, from 43 independent research articles.

The calibrated and validated model developed here represents a probabilistic, model-

driven meta-analysis of a large body of data on volume-overload hypertrophy. Our

analysis suggests that elevated levels of circulating hormones drive much of the

hypertrophic response during late stages of experimental volume overload, whereas

hormone-driven growth frequently reduces myocyte strain levels below baseline despite

elevated left ventricular volumes. These results contrast with the assumption of most

computational models that elevated myocyte stretch drives eccentric hypertrophy but

agree with much of the available molecular and signaling data. The performance of the

model in simulations of multiple independent experiments not used in its construction

suggest that the approach presented here is promising for better representing the

multifactorial complexity of a condition like volume overload, as well as the responses to

drugs administered in this complex setting. However, these simulations also highlight

that responses to some interventions such as β adrenergic blockade will require a

multiscale approach that considers both direct effects on hypertrophic signaling as well

as indirect effects through changes in mechanics and hemodynamics.

2. Methods

2.1. <u>Data collection</u>

We reviewed and collected data from 37 research articles on experimental mitral

regurgitation in dogs and 33 articles on experimental volume overload by aorto-caval

shunt in rats. All data employed for our quantitative analysis were reported as a mean

value and standard deviation, so we assumed a normal probability distribution function

(PDF) for all measurement-derived variables. Any single experimental data in the form

of a mean and standard deviation will be called a dataset in this work. For the

estimation of myocardial stretch at tissue scale, we focused on canine experiments to

avoid confounding effects of growth in body size common during experimental volume

overload in rodents. We collected data on changes in left ventricular mass (LVM), end-

diastolic volume (V_{ED}), and free wall thickness (h_{ED}), as well as previously reported

estimates of end-diastolic myofiber stretch in healthy dogs (λ_{ED}^0).

Both experimental volume overload (VO) and naturally occurring MR in dogs trigger

elevated circulating levels of multiple hormones relevant to hypertrophic signaling,

including AngII, NE, ET-1, ANP and BNP, all of which are included as inputs to the

cardiomyocte signaling network model employed here. We collected all data on plasma

or serum concentrations of these hormones reported in the reviewed articles. We also

collected data from the same studies on activity and phosphorylation levels of

intracellular signaling proteins represented as intermediate nodes in the cardiomyocyte

signaling model. Specifically, we collected Western blotting data on focal adhesion

kinase (FAK), Akt, ERK5, ERK12, ELK1, cGMP, p38, and JNK from myocardial tissue

extractions collected at several stages of VO. Additionally, we collected data on the

abundance of several proteins synthesized by myocytes in tissue samples extracted

following chronic VO including SERCA, Myosin heavy chain isoforms αMHC and βMHC,

ANP and BNP [19]; the network model predicts expression of the corresponding genes

as outputs. A detailed list of sources for all the collected quantitative data is summarized

in the supplementary material S1.1.

2.2.   Integration of canine and rat experimental data

Plots of the experimental fold change of normalized left ventricular mass to body mass

(LVM/BM) during volume overload showed very similar shapes for dogs and rats, but

hypertrophy occurred much faster in rats [9]. When we fitted data from each species

with an exponential function and normalized the time axis by the time constant of that fit,

we found that data from both species aligned (Fig 1a, the full list of data sources is

shown in supplementary material S1.1). We therefore normalized all time course data in

this study by the fitted time constant for each species. This allowed the use of combined

data from both animal models in our quantitative analysis.

Fig 1. Integration of experimental data from MR in dogs and VO in rats. Each plot shows fold changes in one
reported measure, plotted as a function of time normalized by the time constant of an exponential equation fitted to
the LVM/BW data. a) LVM/BM data from sixteen articles (41 datasets) on experimental MR in dogs [15,16,20–33]
and seventeen articles (45 datasets) on experimental VO in rats [34–50]. b) 6 measurements of serum angiotensin II
(ANGII) from three articles on experimental MR in dogs [29,30,51] and 7 measurements from six articles on
experimental VO in rats [38,41–43,52,53]. c) 3 measurements of serum atrial natriuretic peptide (ANP) levels from
three studies in dogs [51,54–56] and 9 measurements from five studies in rats [39,53,57–59]. d) Norepinephrine
(NE): 10 datasets from four MR studies in dogs [21,31,60,61] and 7 datasets from six studies in rats
[39,44,48,52,59,62]. e) Endothelin 1 (ET1): two datasets from two studies in dogs [63,64] and two datasets from two
studies in rats [36,42]. f) Brain natriuretic peptide (BNP): three datasets from three articles on experimental canine
MR [51,54–56]; two datasets from two articles on experimental VO in rats [38,57]. Probability distribution functions
(PDFs) are derived from fits to equations listed in table S1.3. PDFs are presented in shaded gray, median in solid line
and quantiles in dashed lines.

2.3.   <u>Time-varying hormonal input curves</u>

We found baselines or control values of relevant hormone concentrations in blood were

consistent across animal sizes and species, suggesting a common homeostatic range

of circulating concentrations for each hormone. In this work, we assume that fold-

changes in concentration levels of those circulating hormones represented the intensity

of the neurohormonal response and would trigger proportional changes in the hormone-

receptor reaction input in the cardiomyocyte signaling model. The time resolved data of

serum concentrations were normalized to their corresponding baseline or control

concentration and plotted as a function of characteristic growth time (t/τ). We confirmed

that data from both species followed similar trends and fitted the integrated

experimental data with the simplest function (linear or exponential) that captured the

temporal trends (Fig 1). Details of the fitting process and the specific functions

employed for fitting are provided in the supplementary material section S1.2. and table

S1.3 respectively. The fitted, time-varying probability distribution functions (PDFs) for

each hormone were used as inputs to the signaling network model in subsequent

simulations of volume overload.

2.4.   <u>Time-varying myoStrain input curve.</u>

In addition to the many hormonal changes discussed above, experimental volume

overload alters left ventricular (LV) mechanics acutely, followed by further evolution due

to the ongoing process of eccentric hypertrophy. The network model provides an input

labeled myoStrain as a generic representation of the stretch-activated signaling

pathways though to be involved in transducing changes in myocyte mechanics into

hypertrophy. Because a number of successful models of eccentric hypertrophy during

VO have employed end-diastolic strain relative to an unloaded reference state as a

growth stimulus [12,13], we assumed here that the myoStrain network input varies

proportionally with end-diastolic strain in the LV. We computed a time-varying PDF for

the fold change in end-diastolic strain using published data and a simple geometric

model of a thin-walled sphere, as outlined below.

2.4.1. Calculating end-diastolic strain relative to an unloaded state.

Let the circumference of a mechanically unloaded sphere (i.e., floating in saline without

any applied pressure) be $2\pi r_0$, where r is the radius of the sphere and the subscript

zero indicates the unloaded or reference state. Once inflated to end-diastole, the

sphere's circumference is $2\pi r_{ED}$. The stretch *(λ)* around its circumference is defined as

the ratio of loaded to unloaded length:

$$\lambda = \frac{2\pi r_{ED}}{2\pi r_0} = \frac{r_{ED}}{r_0} \quad (1)$$

In a spherical model of the LV, myocytes with any orientation parallel to the surface will

also stretch by this same fraction. While an actual heart is never fully unloaded unless it

is arrested and removed from the body, most mechanics models of growth use the

unloaded state as a mathematical reference point because it is easy to define and

doesn't change with beat-to-beat variations in hemodynamics.

Since many studies reviewed here tracked changes in volumes, we found it convenient

to express stretch and strain in terms of volumes rather than radius. Recognizing that

the volume of a sphere is proportional to r^3, we can rewrite the equation for stretch as:

$$\lambda = \frac{r_{ED}}{r_0} = \left(\frac{V_{ED}}{V_0}\right)^{1/3} \quad (2)$$

256 Finally, for large deformations measured relative to a fixed reference state, the most

257 appropriate measure of strain is the Lagrangian strain [65]:

$$\varepsilon_f = \frac{1}{2}(\lambda^2 - 1) = \frac{1}{2}\left(\left(\frac{V_{ED}}{V_0}\right)^{2/3} - 1\right) \quad (3)$$

2.4.2. Effect of growth on end-diastolic strain.

Returning to the unloaded sphere, if each myocyte grows longer by adding sarcomeres

in series – as occurs during eccentric hypertrophy – then the unloaded radius r_0 and

volume V_0 will be larger, and the stretch at a given end-diastolic volume will be smaller

(Equation 3). In other words, growth changes the amount of stretch each myocyte

experiences, which alters the stimulus for further growth. One of the key features of the

model presented here is that it accounts for this critical feedback between growth and

the hypertrophic signaling that produces it. Fig 2a illustrates this point graphically. In a

spherical model of a normal heart, stretch increases as a nonlinear function of volume

during simulated inflation. Imposing a sudden volume overload produces a big increase

in end-diastolic volume (blue circle) relative to baseline (black circle). A modest amount

of growth (10% volume increase) shifts the entire curve rightward, so that stretch is

lower at a given volume, and inflation to a larger volume is required to achieve the same

stretch (red triangle).

Fig 2. Effects of inflation and growth on end-diastolic stretch relative to unloaded state in a spherical model and in
hearts fixed at control volumes. a) Spherical model predicts that stretch increases with inflation (black solid line),
while eccentric hypertrophy shifts the stretch-volume curve downwards, to lower stretch at any given volume (red
dashed line). b) Experimental measurements of sarcomere stretch in dog hearts fixed at controlled volumes similarly
show increased stretch between physiological loading conditions (Baseline) and acute volume overload (Acute
dilation), and a shift in the stretch-volume relationship following 10% growth in LVM during long-term volume
overload. Figure adapted from Ross et al. (1971) [66].

A thin-walled sphere is a very simplistic representation of the geometry of the left

ventricle, and a more sophisticated geometric model would produce a different

quantitative relationship between volume and stretch. But that was not an important

limitation for this study, for three reasons. First, the spherical model does remarkably

well when compared to actual sarcomere lengths measured by Ross et al. in normal

and hypertrophied hearts fixed at controlled volumes (Fig 2b), suggesting its use here is

reasonable. Second, the trends we predict here – an acute increase in end-diastolic

strain followed by a gradual decline to baseline levels at 6 weeks – were experimentally

verified by Emery and Omens (1997) using markers attached to the LV surface [67].

Finally, it is important to note that the stretch and strain values we calculate from the

spherical model (or any alternative model) are ultimately mapped onto a normalized

input to the signaling network, and the relative influence of that input is calibrated

against data (see below). Therefore, a different geometric model would just produce a

different calibration; the overall behavior of the system would not be affected.

Consistent with the approach described above for integrating data on neurohormonal

alterations, we estimated the time-varying PDF of strain by integrating the spherical

model with data from sixteen articles reporting changes in LVM [15,16,20–33] and

twenty-one articles reporting changes in V_{ED} during experimental MR in dogs, as well as

five articles reporting end-diastolic myocardial or sarcomere stretch relative to baseline

in healthy dogs [66,68–71]. Details of this process are described in detail in

supplementary section S1.3.

2.4.3. Mapping end-diastolic strain onto the network input myoStrain.

The mechanical input to the cardiomyocyte network model (myoStrain) is a normalized

variable ranging from 0 to 1 (Section 2.5). However, in practice, increasing this input

from its default baseline of 0.06 to just 0.2 produces the maximum possible CellArea

output (CellArea=1) in the published version of the model employed here [72]. Thus, we

sought to design a function that maps the range of expected values for organ-scale end-

diastolic strain at baseline or during volume overload into myoStrain input values within

the 0-0.2 working range. We also wanted the mapping function to produce a zero

myoStrain input when organ-scale strain was zero (i.e., when the LV is in a completely

unloaded state). We decided to use an exponential mapping function of the form:

$$w_{myoStrain}^{i} = C_{myo} \left(e^{D_{myo}\, \varepsilon_f^i / \varepsilon_f^0} - 1 \right) \quad (4)$$

where superindex i refers to any given time, while superindex 0 is reserved for the

baseline (pre-VO) state, ε_f and $w_{myoStrain}$ are the maximum fiber strain and its

corresponding mapped myoStrain, and C_{myo} and D_{myo} are mapping parameters to be

fitted by a Markov Chain Monte Carlo (MCMC) algorithm (Section 2.6). To reduce the

order of the MCMC parameter space, we fixed the baseline myoStrain weight

$\left(w^0_{myoStrain}\right)$ at a single value for each MCMC run and repeated the process for

$w^0_{myoStrain}$ values of 0.02, 0.04, 0.05 0.055, 0.06, 0.065, 0.07, 0.08, and 0.09. On each

MCMC iteration we assign a random value to C_{myo} and calculate D_{myo} with equation (4),

the prescribed value of $w^0_{myoStrain}$, and reported baseline values of strain $\left(\varepsilon^0_f\right)$ [66,68–

71].

2.5.   <u>Model of cardiomyocyte hypertrophic signaling pathways</u>

We employed a published computational model of the hypertrophy signaling network

that integrates many established pathways implicated in cardiac myocyte growth. The

model consists of a logic-based network where the activity of each node follows a

normalized Hill equation with possible activity values ranging from 0 and 1 [19,73]. The

network consists of 106 nodes and 192 reactions. The nodes represent chemical and

mechanical inputs, intracellular signaling proteins, transcription factors, and genes

relevant to hypertrophy. The model has been used previously in the study of ventricular

hypertrophy and was recently optimized in the context of β adrenergic stimulation [72].

The set of network parameters is summarized in supplementary table S2. In

supplementary Fig S1.1 we show a representation of the network model highlighting the

nodes for which experimental data are available from experimental VO in rats or dogs.

The influence of a reaction on the downstream nodes is modulated by the weight

parameter *w*, which was left at the default value for all nodes except the inputs AngII,

ANP/BNP, ET1, NE, and stretch. The characteristic time constant governing the speed

of changes in node activity was chosen as 0.005τ for all intracellular reactions, 0.02τ for

output nodes reflecting gene expression, and τ for CellArea, where τ is the fitted time

constant for the exponential rise in LV mass (LVM), as discussed above. The network

model was solved with Netflux (https://github.com/saucermanlab/Netflux). More detail

about the network model formulation and solution method can be found in [74].

2.6.   <u>Bayesian inference parameter estimation</u>

All parameter estimations required in our data processing pipeline were performed

within a Bayesian inference framework. The Bayesian inference tool utilized for this

study was a standard Markov Chain Monte Carlo (MCMC) algorithm with Metropolis-

Hasting selection criteria and Gibbs sampling to navigate the multiparametric space.

Briefly, the algorithm iteratively solves a numerical model while randomly varying its

input parameters over a predetermined probability distribution, known as the prior

probability distribution function (prior PDF) of the parameter space. On each iteration,

the likelihood of the model's outputs is evaluated against experimental data. If the

likelihood of the outputs with the current parameter set is larger than the likelihood of

the previous iteration, the parameter set is saved. If the outputs for the current

parameter set are less likely, the decision on whether to save the current parameter set

is made randomly. After sufficient iterations, the collection of saved parameter sets

converges to a new probability distribution, or posterior PDF of the parameter space,

which are associated with probability distributions for the model predictions [75,76].

In this study, each MCMC algorithm was applied in two stages, first assuming a uniform

probability distribution of the parameters within their physiologically plausible limits for

10,000 iterations. The resulting posterior PDF was then used as the prior PDF for a

second run of the MCMC algorithm for an additional 20,000 iterations, with a check to

verify the convergence of the solutions every 5,000 iterations (Supplementary material

S1.4). The MCMC algorithm was programmed in MATLAB. The source code including

arrays containing the experimental data is available at

https://github.com/cardiacbiomechanicsgroup/MCMC_cardiomyocyte_VO_growth .

2.6.1. Probability distribution of hypertrophy network input weights

The normalized time-varying curves of hormone concentrations and mechanical strain

provide information on how these stimuli vary over time during volume overload, but not

on their relative influence in driving hypertrophy. One advance of the current work over

previous applications of this network model is that we allow the key hypertrophic stimuli

to have different weights, and calibrate those weights using experimental data. In the

cardiomyocyte signaling network model, the baseline weight of the hormone-receptor

reaction determines its relative influence on the network because, the magnitude of the

input at any given time is calculated as the product of its baseline weight $\left(w^0_{species}\right)$ and

its fold-change from baseline at the current time (sections 2.3 and 2.4). We employed

an MCMC to estimate the PDF of the baseline weights of hormone-receptor input

reactions as follows. We first assumed a uniform prior PDF for the input weights of

ANGII $\left(w^0_{AngII}\right)$, NE $\left(w^0_{NE}\right)$, and ET1 $\left(w^0_{ET1}\right)$ reactions. Sampling was constrained within

the range for which the CellArea output is sensitive to those inputs. Specifically,

$$0.01 < w^0_{ANGII} < 0.15$$

$$0.01 < w^0_{NE} < 0.24 \quad (5)$$

$$0.01 < w^0_{ET1} < 0.17$$

For the rest of the input reactions, we assign a single "background" reaction weight,

sampled within the $0.01 < w_{background} < 0.4$ range. A preliminary study revealed that,

within the range of interest, the input reaction weights of ANP and BNP to Guanylate

Cyclase A (GCA) receptors have only marginal effects on predicted changes in

CellArea; we therefore prescribed ANP and BNP the same background weight as the

other inputs for which limited experimental data were available. We assigned null weight

to the input reactions for the exogenous drugs phenylephrine and isoproterenol (ISO)

except when simulating drug infusions.

On each step of the MCMC, the algorithm randomly samples the w^0_{ANGII}, w^0_{NE}, w^0_{ET1},

$w_{background}$, and C_{myo} parameter space and randomly selects time-varying curves for

each stimulus from their respective PDFs. The likelihood of each model run was

evaluated against experimental data on FAK (7 datasets, 3 studies) [14,15,25], Akt (2

datasets, 2 studies) [15,77], ERK5 (1 dataset, 1 study) [78], ERK12 (7 datasets, 3

studies) [15,50,79], ELK1 (1 dataset, 1 study) [50], cGMP (4 datasets, 4 studies)

[53,78,80,81], p38 (6 datasets, 4 studies) [15,78,79], and JNK (5 datasets, 2 studies)

[15,79] activity and cardiomyocyte growth, CellArea in our model, (11 datasets, 9

studies) [16,21,23,28,29,34,82–84] from dog and rat experiments. We added a

condition assigning larger likelihoods to parameter sets that produce a baseline

CellArea activity near 0.5, in the most responsive region of the sigmoidal curve. Long-

term experiments agree that LVM plateaus at a new level in chronic stages of VO [26].

We therefore assumed that continued growth at late time points and negative growth

(reversal of hypertrophy) at any time point were very unlikely. After convergence of the

MCMC, we filtered out these very unlikely solutions and recorded the posterior PDFs of

the activity of the network nodes of interest.

2.7.   <u>Sensitivity analysis</u>

We evaluated the sensitivity of network outputs (expression of ANP, BNP, αMHC,

βMHC and SERCA) to the network inputs (ANGII, NE, ET1, ANP and BNP) by a

standard correlation matrix based on statistical linear regression. The Pearson

correlation coefficient (PCC) was calculated to quantify the parameter sensitivity. This

method exploits the wealth of samples produced during the MCMC runs to yield

sensitivity estimates that are meaningful within the expected range of network activity.

2.8.   <u>Validation</u>

Volume overload is a complex process driven by multiple stimuli that trigger intersecting

signaling cascades. We selected VO for the calibration of the hypertrophy network

model to take advantage of the wealth of published data on how these different

pathways respond over time. We calibrated the relative influence of the hormonal and

mechanical stimuli by adjusting the weight of the network input reactions to match data

on LVM/BM and the activity of intracellular signaling proteins from *in vivo* VO studies.

The output of the Bayesian calibration is a PDF for the baseline weight of each input

reaction.

We validated the calibration using independent studies of the infusion of individual

agonists that can stimulate hypertrophy as well as combinations of those agonists with

pharmacologic blockers of various signaling pathways. For each validation simulation,

we performed a Monte Carlo simulation (MC), with one thousand iterations (N=1000).

On each iteration, we selected a set of baseline reaction weights

$\left(w_{AngII}^0, w_{NE}^0, w_{ET1}^0, w_{backgroud}^0 \right)$ from their calibrated PDFs. We simulated the infusion of

agonists by increasing the weight of the corresponding reaction while keeping the rest

of reactions at their selected baselines. When simulating the infusion of agonists at

saturation doses, we increased the weight of the corresponding reaction to its maximum

value of one. Otherwise, we randomly selected a fold-change value from a normal

distribution matching the experimentally reported mean and deviation of the agonist

concentration in plasma and increased the input reaction weight by that same fold

changes. To simulate the effect of pharmacologic blockers, we set the maximum activity

of the target receptor to zero [85]. For the simulation of VO, we randomly picked time-

varying curves of AngII, NE, ET1, ANP, BNP, and ε_f from their respective PDFs on

each iteration. Although some of the agonist and blockers had significant effects on

heart rate (HR) and blood pressure (BP) in the corresponding studies, we did not model

those effects here.

In prior studies using large-scale signaling networks, our group compared model-

predicted changes (increase, decrease, or no change) for individual outputs or signaling

intermediates to published measurements (significant increase, significant decrease, or

no significant change) not used for model construction and calibration [85]. Here, we

adapted that approach slightly to account for the probabilistic nature of the Bayesian

calibration. Each model simulation now produces a distribution of possible outcomes; if

at least 75% of those simulations predicted an increase, we treated that as a model-

predicted increase. Predicted decreases were handled similarly. We considered the

model prediction to be validated if most available studies on the simulated intervention

reported a significant increase or decrease in the same output. As a more quantitative

metric, we also counted how many published studies reported a mean change that fell

within the 50% and 90% confidence intervals (CI50 and CI90) of the model prediction.

2.8.1. Simulation of infusion-induced hypertrophy.

We performed *in silico* simulations of infusion experiments of ISO and ANGII in rats,

and NE in dogs. We quantified the hypertrophic effect of infusion as the relative growth

with respect to baseline:

453
$$Normalized\ Growth = \begin{cases} \frac{LVM/BM^{final} - LVM/BM^{initial}}{LVM/BM^{initial}} \ (experiment) \\ \frac{CellArea^{final} - CellArea^{initial}}{CellArea^{initial}} \ (model) \end{cases} \quad (6)$$

We collected fifteen experimental datasets (mean±SD) from twelve independent studies

of ISO infusion in rats at rates >1mg/kg/day, a rapid-growth inducing saturation dose

[86–97]. We simulated these experiments by imposing the maximum input weight of 1

on the ISO node to simulate its infusion at saturating rates and comparing the steady-

state growth prediction to experimental data collected at termination times longer than

one week, at which maximum growth was reached in all reviewed experiments.

AngII infusion in rats at rates of 100-200 ng/kg/day is a standard rodent model of

hypertension. At this infusion rate, the plasma concentration of AngII increases two- to

five-fold [98–104]. We collected 20 experimental datasets from 14 independent studies

in rats, measured at 1 week, 2 weeks, or 4 weeks of infusion [98–111]. We simulated

these experimental conditions by increasing the AngII stimulation by 3.5 ± 1.5 fold.

We found four independent reports of experimental NE infusion in dogs, each with a

different dosage and termination time [112–115]. Each of the experiments in dogs

reported the plasma concentrations achieved, and we therefore simulated each

individual experiment by imposing the same fold increase on the NE network input as

the reported fold change in plasma NE, for the same duration as the experiment.

2.8.2. Replication of drug treatment of infusion-induced hypertrophy.

To validate the capacity of our model to reproduce cross-talk between different signaling

pathways, we also simulated experiments where agonist infusions were paired with β-

adrenergic receptor blockers (βB), angiotensin receptor blockers (ARB), or endothelin

receptor antagonists (ERA). In each case, we increased the input weight of the infused

hormone while keeping the rest of the inputs at their baseline and setting the activity of

the targeted receptor at zero. We quantified the drug effect on infusion-induced

hypertrophy as the chronic difference between the growth during infusion plus blocker

and growth during infusion alone measured at the same endpoint, normalized to the

initial dimension:

$$Normalized\ Drug\ Effect = \begin{cases} \dfrac{LVM/BM_{infusion+drug}^{final} - LVM/BM_{infusion}^{final}}{LVM/BM^{initial}} \ (experiment) \\ \dfrac{CellArea_{infusion+drug}^{final} - CellArea_{infusion}^{initial}}{CellArea^{initial}} \ (model) \end{cases} \quad (7)$$

We collected eight datasets from four independent articles studying the effect of βB on

ISO-induced hypertrophy [88,95,116,117], five experimental datasets from five

independent publications studying the effect of ARB on ISO-induced hypertrophy

[86,96,97,117,118], three experimental datasets from three experimental studies on the

effect of ERA on AngII-induced hypertrophy [111,119,120], and one experimental point

from a publication studying the effect of ERA on NE-induced hypertrophy [121]. All of

these experiments were performed in rat models.

2.8.3. Validation of additional model outputs.

Cardiac hypertrophy is typical associated with changes in expression of ANP, BNP,

αMHC, βMHC, and SERCA that are collectively termed the fetal gene program

[19,122,123]. The network model employed here predicts each of these outputs, but no

data on their changes were included in the Bayesian calibration process. As an

additional independent validation of the performance of the calibrated network model,

we therefore compared model-predicted changes with experimental reports of

significant changes in ANP, BNP, αMHC, βMHC, and SERCA protein abundance in

tissue extracted following infusion experiments. Because the model predicts gene

expression while the experiments measured protein abundance, we did attempt

quantitative comparisons for these outputs but rather simply assessed whether they

increased significantly, decreased significantly, or did not change significantly.

2.9.   <u>Administration of receptor blockers during VO</u>

Several drugs currently employed in patients with heart failure have been tested in

animal models of experimental VO. While we expect that some of those drugs have

important secondary effects not represented in the current model, we simulated

available experiments to further explore its capabilities and limitations. Model

predictions were compared to data from treatment of experimental VO with βB, ARB

and ERA. We collected six experimental data points on the early ($t/\tau \leq 1$) and two on

the chronic ($t/\tau > 1$) effects of βB in VO from three articles on experimental MR in dogs

[15,16,80] and two articles on experimental VO in rats [14,124]. We collected five

experimental datasets on the effect of AngII on VO-induced hypertrophy from three VO

experiments in rats [42,46,50], and one from MR experiments in dogs [29]. The effect of

ERA on VO was compared to four experimental datasets from four independent articles

on experimental VO in rats [125–128].

3. Results

3.1.   <u>Cross-species data integration and probability distribution of hormonal stimuli.</u>

The best-fit characteristic time constant for mass growth was (τ_M) 1095 hours for dogs

and 283 hours for rats. The ratio of the species time constant was $\tau_{M_{dog}}/\tau_{M_{rat}} = 3.86$.

When scaled by this ratio, data from rats and dogs were reasonably congruent;

variability within individual studies and/or between different studies in the same animal

model were generally larger than the differences between means measured from the

two different species at similar relative times (Fig 1). Interestingly, the ratio of growth

time constants computed here is close to the ratio of the reported heart rates (HR),

which is often used for allometric scaling [129]. With an average HR of $363 \pm 14\ bpm$ for

rats and $95 \pm 4\ bpm$ for dogs, the HR ratio was $HR_{rat}/HR_{dog} = 3.82 \pm 0.22$.

3.2.   <u>Probability distributions of fiber strain during VO.</u>

We combined available data on normal heart mechanics and volume-overload

hypertrophy in dogs with a simple geometric model to estimate the probability

distribution of end-diastolic strain (ε_f) relative to an unloaded state over the course of

experimental VO. The PDF displays the expected trends over time: an acute increase in

strain owing to the sudden increase of V_{ED} when mitral regurgitation is first introduced,

followed by a gradual decrease driven by the compensatory hypertrophic response (Fig

3). As VED and LVM curves reach a plateau in the chronic stages of VO, ε_f also

stabilizes. Models in which mechanical strain is the sole driver of cardiomyocyte growth

only produce a stable hypertrophied state if strain returns to its baseline level, or the

homeostatic strain level is allowed to adapt [13,130]. Interestingly, our data analysis

suggests that in 75% of the cases, strain falls below its original baseline level in chronic

stages of experimental VO (Fig 3c), with 50% of the cases passing below this threshold

relatively early (*t/τ ≤ 1* corresponding to the first 12 days in rats and 6 weeks in dogs).

The final chronic level of strain showed a strong inverse correlation to chronic LVM fold

change (PCC=-0.78), suggesting that myofiber stretch is more likely to fall below

baseline in cases with the greatest mass increase (Fig 3d).

Fig 3. Changes in volume, mass, and end-diastolic strain during experimental MR in dogs. a) Fitted data on fold
changes show a) a 36% increase in LVM [15,16,20–33] and b) 67% increase in LV end-diastolic volume (VED)
[16,17,20–26,29,32,33,61,78,80,82,83,131–136] on average at steady-state. c) In a spherical model, these changes
imply a 20% increase followed by a gradual decrease below baseline in end-diastolic strain on average, with a wide
range of trajectories possible for different combinations of mass and volume change. d) Calculated chronic strain
levels are lower when LVM increases more. Insets in panels (a-c) show acute changes plotted on a magnified time
axis.

3.3.   Calibration of relative weights for input stimuli to the hypertrophy network

model

In this work we allowed the key network inputs to have different weights, and calibrated

those weights using experimental data. Preliminary screenings of the parameter space

revealed that the baseline weight of the myoStrain input $\left(w_{myoStrain}^0\right)$ strongly influenced

the likelihood of producing unlikely predictions (i.e., those that contradict experimental

observations) such as spontaneous growth reversal despite continued overload and

runaway growth that never stabilized. We therefore ran a series of MCMCs at fixed

values of $w_{myoStrain}^0$ while randomly sampling the rest of the input reaction weights as

described in section 2.6.1. We found that constraining the baseline weight of the

myoStrain input within the range $0.05 \leq w_{myoStrain}^0 \leq 0.06$ minimized the number of

unlikely solutions. For $w_{myoStrain}^0 > 0.06$, the solutions tend to be dominated by the

myoStrain input, resulting in a reversal of growth at later time points despite continuing

simulated overload, while the solutions for $w_{myoStrain}^0 < 0.05$ were dominated by

adrenergic stimulation, increasing the chances of runaway growth (Supplemental

material S1.5). The results presented here were therefore obtained with a fixed

$w_{myoStrain}^0 = 0.06$, which was associated with the highest mean likelihood among the

myoStrain values we tested.

The posterior PDF of the remaining input reaction weights converged to $w_{ANGII} =$

0.010 ± 0.002, $w_{NE} = 0.033 \pm 0.014$, $w_{ET1} = 0.056 \pm 0.025$, and $w_{background} = 0.031 \pm$

0.022. We calculated correlations among these input weights across the 20,000

simulations from the final round of MCMC simulations. The strongest correlation was

between w_{ET1}^0 and $w_{background}^0$ with a Pearson PCC=-0.42, while all other PCC

magnitudes were below 0.3 (Supplementary material S1.5). The PDF of the strain-to-

Myostrain mapping parameter C_{myo} converged to $C_{myo}=5.86 \pm 1.38$. Among all nodes in

the network, the mean predicted chronic activation level at $t/\tau = 3$ (Fig 4) increased most

for Ca++/calmodulin-dependent kinase (CaMK) and calcineurin (CaN) and decreased

most for integrin and focal-adhesion kinase (FAK).

Fig 4. Predicted chronic activation of the cardiomyocyte signaling network during VO. Colors indicate the logarithm of the fold change in activity of each node in chronic stages of VO relative to baseline, with red indicating an increase and blue indicating a decrease.

3.4. <u>Validation</u>

First, we generated a forward MC simulation of untreated VO as described in section

2.7. Briefly, we performed N=1000 simulations of growth until steady-state ($t/\tau = 5$) in

untreated VO. For each iteration we selected a set of baseline reaction weights from

their calibrated (posterior) PDFs, and a set of time-varying curves for neurohormonal

stimulation and stretch from their respective PDFs [134]. The model predicted a wide

range of possible growth curves for untreated VO (Fig 5a). On average, normalized

growth in CellArea (Equation 6) was 47%, with 50% of simulations (CI50) predicting

growth between 34% and 60%, and 90% of simulations (CI90) predicting growth

between 17% to 83%. More than 75% of our simulations predicted chronic increases in

ANP, BNP, and βMHC and decreases of SERCA and αMHC; these predictions agreed

well with experimental measurements of protein abundance from cardiac tissue

extractions in experimental VO (Fig 5b).

Fig 5. Results of MC simulations of VO and agonist infusion. Top panels show normalized growth in CellArea for 100
representative simulations (gray lines), with terminal values indicated by black circles; adjacent boxplots reflect the
mean, CI50, and CI90 of terminal values for a) Untreated VO, c) ISO infusion, and e) AngII infusion. N indicates the
number of simulations employed for the graphical representation. Bottom panels contain heatmaps comparing the
predictions of changes in signaling network outputs (SERCA, αMHC, βMHC, ANP, BNP, and CellArea) to experimental
data following b) Untreated VO, d) ISO infusion, and f) AngII Infusion. Red indicates that more than 75% of simulations
predicted an increase in the output, or that the majority of studies reported a significant increase. Blue indicates a
decrease in >75% of simulations or the majority of experiments. Gray indicates the absence of consistent changes in
simulations or significant changes in experiments. White numbers indicate the number of studies reporting a significant
change and the total number of studies reporting data for each output.

3.4.1. Simulation of infusion-induced hypertrophy.

We validated the calibrated model against data from independent experiments not used

in its calibration that infused hypertrophic agonists over periods of weeks and measured

the resulting hypertrophy. Our model predicts that infusion of ISO at saturating doses

produces a mean LV growth of 40%, with 50% of simulations (CI50) falling between 33

and 50% growth, and 90% of simulations (CI90) falling in the range 19-68% (Fig 5c).

These predictions show good agreement with experimental data, with all available

studies reporting a significant hypertrophic effect of ISO at infusion rates >1 mg/kg/day

(Fig 5d). Four out of 15 studies reported mean increases in LVM/BM that fell within our

CI50, and 12 of 15 reported means within the predicted CI90 (Fig 6a) [86–97]. Our

model also predicted increases in ANP, BNP, and βMHC and decreases in SERCA and

αMHC in more than 75% of simulations of ISO infusion (Fig 5d), which is consistent with

the significant changes reported in all experimental measurements of ANP, βMHC,

SERCA, and αMHC protein abundance during ISO infusion (Fig 5d) [88,93,95]. Our

model predicted that increasing the serum concentration of AngII by *3.5 ± 1.5*-fold would

produce long-term growth of 13% on average, with 50% of simulations (CI50) in the

range [7%,19%], and CI90 of [1.8%,31%] (Fig 5e). In more than 75% of simulations,

AngII activated the fetal gene program, decreasing SERCA and αMHC while increasing

βMHC, ANP, and BNP (Fig 5f). This prediction agreed with all experimental data

collected for validation, except for a single study that found no significant difference in

βMHC protein abundance relative to control [105,106,109]. When comparing to data

collected at different time points during the course of AngII-induced hypertrophy, more

than 75% of simulations predicted growth at every time point, and the magnitude of

predicted growth increased gradually with time (Fig 6b).

Fig 6. Validation of model predictions of growth during infusion of hypertrophic agonists with and without receptor blockers. a) Boxplot shows the mean, 50% confidence interval, and 90% confidence interval for predicted normalized growth in 1000 MC simulations of ISO infusion with the calibrated network. Experimental means (plotted as individual points with error bars) from 13 of 14 published studies [86–97] fell within the CI90. b) Model-predicted CI90 for growth during AngII infusion encompassed most published values at the earliest time point and all reported experimental means at later time points [98–111]. c) Simulations of four NE infusion experiments in dogs with different doses, fold increases in plasma concentrations (Conc.), and termination times (t/τ) showed reasonable agreement with the limited available data [112–115]. d) Simulations of agonist infusion plus receptor blockers replicated reported significant effects of β blockers (βB) on ISO-induced hypertrophy infusion [88,95,116,117], and smaller effects that were significant in some but not all studies for endothelin receptor antagonist (ERA) administration during ISO [86,96,97,117,118] or AngII [111,119,120] infusion. By contrast, the model prediction of nearly equal likelihood that ERA increases or decreases NE-induced growth appears to contradict a significant reported decrease in the only available experimental study [121]. Red coloring of boxplots indicates increases in >75% of simulations, blue indicates decreases in more than 75%, and gray indicates mixed results. Points with error bars indicate experimental mean±SD for studies reporting statistically significant effects (filled markers) or non-significant effects (open markers), from studies in dogs (circles) or rats (triangles).

These predictions compared well to independent experimental data: at the earliest time

point, 7 of 9 experimental means fell within the predicted CI90; while at later time points

all experimental means fell within the CI90. The simulations appeared to under-predict

growth somewhat, with no experimental means falling inside the CI50 at 1 week ($t/\tau =$

0.6), 4 out of 8 datasets within the CI50 at 2 weeks ($t/\tau = 1.2$), and 1 out of 3

experimental points collected at one month of AngII infusion ($t/\tau = 2.4$) falling within the

predicted CI50 [98–111]. Variability in the experimental data was large compared to the

effect size of AngII treatment, and the CI90 of the model predictions spanned a similar

range to the experimental data, illustrating the potential advantages of the Bayesian

approach.

Simulations of NE infusion showed reasonable agreement with experimental data,

displaying growth in >75% of simulations at all plasma concentrations and increasing

growth with higher concentration or longer durations (Fig 6c). The experimental means

for all simulated infusion protocols fell within the CI90 of the model predictions, and 2

out of 4 fell within our CI50. We note that the lowest plasma concentration simulated

here fell within the range of NE levels observed during experimental VO, while the other

simulations fell well above the calibrated range.

3.4.2. Simulations of agonist infusion plus receptor blockade.

In addition to testing model predictions of the effects of infusing individual hypertrophic

agonists, we validated the model against experimental data on combinations of those

agonists with various receptor blockers. Our model predicted that βB administration

would reduce ISO-induced hypertrophy by 27%, with 50% of simulations falling in the

range [-36%, -22%] and 90% of simulations falling in the range [-47%, -18%] (Fig 6d).

These predictions showed good agreement with experiments, with 7 out of 8 reported

means falling within our CI90, 5 falling within the CI50, and all experiments reporting a

statistically significant effect of βB treatment [88,95,116,117].

The model predicted a much smaller effect of ARB administration on ISO-induced

growth, with a mean reduction in CellArea of 4.5%, CI50 of [-8%, -1.5%], and CI90 of [-

12%, +3%] (Fig 6d). Consistent with the fact that our CI90 included more predicted

decreases but some predicted increases, only 3 out of 5 studies reported a significant

effect of ARB administration on hypertrophy. All reported means fell within the model-

predicted CI90, and 3 out of 5 within CI50 (Fig 6d) [86,96,97,117,118]. Similarly, our

model predicted that ERA would reduce AngII-induced hypertrophy by an average of

9%, with a CI50 of [-13%, -4.7%] and a CI90 of [-20%, +2%] (Fig 6d), which was

consistent with significant reported reductions in 2 of 3 studies [111,119,120]. 2 out of 3

experimental means fell within the model-predicted CI50, and all reported means fell

within the model CI90

When simulating the combination of NE infusion and ERA administration, our model

predicted a negligible impact on hypertrophy, with a 3% reduction on average with respect

to untreated NE-infusion, a CI50 of [-10%, +3%] and a CI90 of [-23%, +12%]. This

contrasts with the sole experimental report of a 23% decrease, although the mean of that

study did still fall within our CI90 [121].

3.5. Administration of receptor blockers during VO.

We explored the capacity of the model to predict the effect of β blockers (βB),

angiotensin receptor blockers (ARB), and endothelin receptor antagonists (ERA) in the

context of experimental VO. In these simulations we ignored any possible secondary

effects of the blockers on circulating hormone levels or myocardial strain. Experiments

find no effect of β-blockers on ventricular mass in early stages of VO ($t/\tau < 0.7$) in dogs

or rats [14,15,80,124], but chronic use of this drug ($t/\tau = 2.7$) appears to exacerbate

hypertrophy in both animal models [14,16,124]. In our simulations, β-blockers produced

increases and decreases in mass with equal frequency, producing a negligible mean

effect on average at both early and late stages (Fig 7a). Although our simulations would

not have anticipated the significant reported increase in mass in the two studies of

chronic administration, experimental means from all studies and time points did fall

within the model-predicted CI90.

Fig 7. Effects of receptor blockers on VO-induced hypertrophy in published experiments and the calibrated model. a)
Simulations of β blockers (βB) at both early ($t/\tau \leq 0.6$) and late stages ($t/\tau = 3$) of VO, simulations (boxplots) were
equally likely to predict increases or decreases, with no net effect on average. MR experiments in dogs (circles)
[15,16,80] and VO in rats (triangles) [14,124] also found no significant effect early but a further increase in growth at
later timepoints relative to untreated VO . b) Simulated administration of angiotensin receptor blockers (ARB) also
produced mixed results, consistent with mixed results in published studies [29,42,46,50]. By contrast, simulated ERA
administration decreases VO-induced hypertrophy in the majority of simulations, and in all four available experiments
[125–128]. Blue coloring of boxplot indicates predicted decrease in >75% of simulations, gray boxplots indicate mixed
simulation results, filled markers indicate experimental means associated with statistically significant effects and open
markers indicate means associated with non-significant effects.

When simulating the effect of ARB administration on VO-induced hypertrophy, our

model produced mixed results, with only 62% of the simulations predicting a partial

reduction in hypertrophy and a mean normalized drug effect of -11%. These mixed

predictions of ARB effects are consistent with the fact that two VO experiments (one on

rats and one on dogs [29,50]) reported no significant effect of ARB on hypertrophy while

three other rat experiments reported a significant decrease [42,46,50]. Experimental

means from all of these studies fell within our predicted CI50 [-36%,+16%].

Our model predicted that ERA dampens VO-induced hypertrophy in 90% of the cases,

with a mean effect of -31%, CI50 of [-51%,-13%] and CI90 of [-77%,8%]. These

predictions agree with the significant hypertrophy-inhibiting effects of ERA reported in all

rat studies reviewed here [125–128]. Experimental means from all 4 studies fell within

our predicted CI90, and half within our CI50.

3.6. <u>Sensitivity analysis</u>

The sensitivity analysis revealed that at late simulated time points, all network outputs

were far more sensitive to ET-1 than to any of the other network inputs examined in this

study (Fig 8). This suggests that, in chronic stages of VO, variation in the circulating

level of ET-1 within its reported range has the largest influence on cardiomyocyte size

and gene expression of any individual hormonal or mechanical growth factor. According

to this analysis, higher levels of ET-1 in late stages of VO are associated with greater

cardiomyocyte growth, greater myocardial production of βMHC, ANP and BNP, and

reduced myocyte synthesis of SERCA and αMHC.

Fig 8. Sensitivity analysis of network outputs to calibrated network inputs in chronic stages of VO. Each marker
represents the final state of one VO experiment from 1,000 Monte Carlo simulations. A linear regression model
(dashed line) is fitted to each output-input pair, and the Pearson PCC is displayed on the upper left corner of each.
Among the inputs known to vary substantially over the course of VO, ET1 displayed the largest influence on all
outputs in the late stages of overload.

4. Discussion

This study used a Bayesian approach to integrate a large body of experimental VO data

in rats and dogs and calibrate a well-established model of hypertrophic signaling in the

heart. The resulting model captures both the temporal variations in neurohormonal and

mechanical stimuli during VO and the relative responsiveness of cardiomyocyte

signaling pathways to those stimuli. As a result, the model not only reproduced the data

on hypertrophy used to calibrate it, but also correctly predicted responses not used in

calibration, such as the re-expression of the fetal gene program during VO.

Furthermore, following calibration against data from 70 studies of VO, the model

performed remarkably well when validated against data from independent studies of

other interventions that trigger some of the same signaling pathways, such as infusion

of hypertrophic agonists with and without administration of receptor blockers. More

specifically, 50 of 56 (89%) reported mean changes in LVM/BW collected from those

studies at various doses and times fell within the 90% confidence interval predicted by

the model for its CellArea output.

Our primary motivation in developing a systems-level model of hypertrophy during VO is

that clinical conditions such as mitral regurgitation trigger changes in multiple

hypertrophic stimuli, including myocyte stretch/stress as well as circulating levels of

hormones known to modulate hypertrophy. Furthermore, most patients receive multiple

therapies – such as administration of β-blockers and mitral valve repair or replacement

– that themselves modify different hypertrophic stimuli. Thus, predicting the effects of

any potential treatment or combination of treatments requires a model that can

anticipate the crosstalk amongst relevant signaling pathways. In this respect, the

validations shown here against experiments where agonist infusion was combined with

a receptor blocker (Fig 6d) are particularly promising. In most cases, both the mean

response and the range of expected responses predicted by the model agreed with

available experimental data; the one exception was the combination of NE + ARBs,

where our model prediction disagreed with the lone study available for comparison.

We also explored model predictions of drug treatment during experimental VO, but did

not treat those simulations as formal validations because a multiscale model would be

needed to capture the expected secondary effects of the drugs on hemodynamics and

ventricular mechanics. While simulations of ARB and ERA treatment during VO

matched data fairly well, the limitations of the model were more apparent when

simulating β-blockade (Fig 7). Chronic β-blockade during experimental VO appears to

exacerbate hypertrophy after several months in dogs and rats [16,21,33,124]. By

contrast, in our calibrated model NE had limited influence on hypertrophic signaling

during VO, so simulated β-blockade had little effect on hypertrophy. The exaggerated

NE concentration levels required by Laks et al. (1973), King et al. (1987) and Stewart et

al. (1992) to produce LVM increments above 20% support the idea that serum levels

measured during VO should exert limited influence on hypertrophy [112–114], and β-

blockade does have minimal impact in isolated cardiomyocytes and short-term VO

experiments [14,15,80]. We therefore hypothesize that worsening of hypertrophy by

long-term β-blockade arises indirectly through their depression of LV contractility, an

effect that was not included in the model employed here. Evidence supporting that

hypothesis includes the fact that β-blockade partially restores FAK phosphorylation

levels during experimental VO, suggesting increased activation of stretch-modulated

hypertrophic pathways [14–16,60]. A further complication when modeling β-blockers is

that they appear to improve LV function, symptoms, and survival in patients with chronic

MR and heart failure [137,138] due to remodeling of the signaling pathways themselves,

another effect not modeled in the present study.

The Bayesian calibration approach proposed here results in a model that predicts a

distribution of expected responses for any simulated intervention. When using a model

to pre-screen potential therapies, this approach could have the important advantage of

providing information on both the expected effect size and the expected variability in the

response. This information could be used to prioritize for further development potential

treatments that are predicted to produce the desired effect in the vast majority of cases,

something not possible with traditional models that predict only a mean response.

Alternatively, if measurable pre-treatment factors distinguish a subpopulation of

simulations with large predicted responses for an otherwise unimpressive potential

therapy, this information might motivate further exploration of its potential in a specific

subgroup of patients. In the model presented here, the variability in any predicted

response reflects both the known topology of the signaling network being modeled and

the variability in the data used for calibration. In most cases where sufficient comparison

data were available to make an assessment, the spread of data from independent

studies used for validation was very similar to the range of responses predicted by the

model; this suggests that the calibration process worked as intended. Viewed through

this lens, even the one clear validation failure (Fig 7, NE+ARB) might not be so bad.

The model predicted a very small effect on average but high variability in that response;

although the one published study reported a significant reduction of NE-induced

hypertrophy by ARBs, its mean fell well within the model-predicted range. If the

experiment were repeated several more times, would this finding persist? Or would

other experiments produce a mix of significant and non-significant findings, as occurred

for some of the other combinations modeled.

The calibrated and validated model suggests some insights regarding hypertrophic

signaling in chronic VO that are not apparent from the structure of the signaling network

alone. For instance, although any of the agonists discussed here are capable of

stimulating cardiac hypertrophy, our sensitivity analysis showed ET1 to be the primary

driver of hypertrophy during chronic stages of VO (Fig 8). Consequently, the model

predicted that ERA treatment reduced VO hypertrophy in the majority of simulations,

while other blockers produced mixed results (Fig 7). These predictions not only agree

with the experimental data we collected for validation [29,42,46,50,125–128], but also

with the observations by Leskinen et al. 1997 and Fareh et al. 1996 on combined

treatment of VO in rats with ARB and ERA, that suggests ET1 stimulation is more

important in regulating the long-term myocyte adaptative response to VO than AngII or

stretch [139,140]. Despite the apparent benefits of ERA in experimental VO and its

proven vasodilator effect, their clinical use has been limited by potentially severe side

effects, such as alterations of liver function, anemia, and edema. The most common

clinical application of ERA is to treat pulmonary artery hypertension, and recent efforts

point to the development of selective ERAs for treating persistent hypertension [141].

One of the most interesting implications of our analysis is our prediction that in most

cases of VO, the combination of early overstretch and sustained neurohormonal

activation trigger sufficient hypertrophy to drive stretch below its baseline levels. This

prediction did not derive from the structure of the signaling network but rather from the

Bayesian approach to integrating published data on observed LV mass and end-

diastolic volume increases across a large number of experimental studies (Fig 2,3). This

prediction may explain the otherwise puzzling depression in the activity of mechano-

transduction pathways in the context of volume overload. While FAK phosphorylation is

elevated in pressure overload and aortic valve regurgitation relative to baseline, FAK

phosphorylation is reduced in VO despite elevated LV volumes that are commonly

assumed to indicate elevated levels of myocyte stretch [15,16]. In previous work, we

identified Ras as a relevant hub responsible for the crosstalk of multiple pathways

[19,73]. In the VO simulations presented here, Ras was a critical node integrating the

competing effects of mechanical stretch and neurohormonal inputs. During early VO,

elevated stretch and neurohormonal stimulation combined to drive strong activation of

Ras, while in chronic VO reduced stretch and FAK activity counteracted continuing

neurohormonal stimulation, producing a low level of Ras activation (Fig 4,

Supplementary material 3).

5. Limitations and future directions

The ability to predict not only mean responses but also the uncertainty around those

predictions is a major advantage of Bayesian approaches. However, the approach

described here may overestimate variability of the predicted responses, for several

reasons. First, we treated the levels of the various circulating hormones as independent

of each other; for example, a given VO simulation might randomly combine input curves

specifying very large changes in one input and very small changes in the others.

However, in reality most reports suggest that the expression of circulating

neurohormones is correlated to the severity of cardiac insult, and therefore to each

other. The current model also lacks output-to-input feedback loops. For example, the

strain-time curve is imposed through random sampling, with no previous knowledge of

the growth response, and the likelihood of a given CellArea prediction is evaluated

against all available data on mass increases, not just those occurring at a similar level

of strain. To address these sources of variability, we could introduce additional

experimental data and covariance relations to the likelihood evaluation. Another source

of variability that would be more difficult to address is the fact that the data used to

calibrate the model were gathered from studies in different animal models, performed by

different groups. These studies typically began with healthy animals and introduced a

severe or moderately severe overload at a known point in time. However, factors such

as variations in overload severity, species differences, and sex differences likely

contributed to the variability in predictions in the final, calibrated model.

Another limitation worth mentioning is that the simulations of both agonist infusions and

receptor blockers ignored secondary effects on hemodynamics, LV mechanics, and

strain. Known experimental responses that were neglected due to this simplification

include an increase in heart rate during ISO and NE infusion, an increase in systemic

blood pressure with ISO infusion in some studies [92,96,116] and in large infusion

doses of NE [114], vasoconstriction by AngII, vasodilation by ARB and pulmonary

vasodilation by ERA, and effects of adrenergic agonists and receptor blockers on LV

contractility and mechanics. One way to address these limitations would be to

incorporate the calibrated myocyte signaling model developed here into a multiscale

model of cardiovascular function. A multiscale model could use tissue and organ-scale

ventricular models to update the estimations of myocardial strain as growth proceeds,

and represent known feedback between systemic hemodynamics, neurohormonal

alterations, drug effects, and heart loading.

We assumed that changes in neurohormonal circulating concentrations are

proportionally transduced into receptor activity, which is likely true only when receptor

availability is high and binding is non-competitive, neglects changes in receptor

abundance, and ignores possible differences in hormonal concentrations between the

bloodstream and the immediate cellular environment. Hormone concentrations in

myocardial tissue extractions can be several orders of magnitude larger than circulating

concentrations. However, they show similar trends, suggesting that the circulating and

local values are at least correlated (Dell'Italia et al., 1995, 1997; Tallaj et al., 2003).

Ultimately, calibrating the model to widely accessible data such as serum

concentrations improves its translational value.

6. Conclusions

We employed a Bayesian approach to combine data from 70 studies on experimental

volume overload in dogs and rats and use it to calibrate a network model of

hypertrophic signaling in cardiomyocytes containing nearly 200 reactions. The

calibrated model reproduced many key results from 43 independent studies not used in

its calibration, including infusion of hypertrophic agonists alone or in combination with

receptor blockers and administration of multiple heart failure drugs in the setting of

experimental VO. In nearly all cases, if the majority of model runs for a given condition

predicted a consistent effect, the majority of experimental studies found the same effect;

where enough studies of a given intervention were available for comparison, the

variability among those studies was also similar to the range of responses predicted by

the model. These results suggest that the approach presented here could be useful in

future simulations of novel potential treatments for VO. The model suggests that

changes in cell size and activation of the fetal gene program in the chronic stage of VO

are particularly sensitive to Endothelin1 receptor activity. The calibrated model also

suggests that growth in experimental VO is mostly driven by the neurohormonal

response, with eccentric hypertrophy reducing initially elevated myocardial strain values

below baseline fairly quickly in most cases. This prediction provides a plausible

explanation for the depression of mechano-transduction signaling pathway in

experimental VO, despite the widespread conception of volume overload hypertrophy

as driven by myocyte overstretch.

7. Acknowledgements

The authors thank Dr. Pim Oomen, University of California Irvine, for his support and guidance on the selection and development of the Bayesian analysis method.

8. References

[1] Wu S, Chai A, Arimie S, Mehra A, Clavijo L, Matthews R V., et al. Incidence and treatment of severe primary mitral regurgitation in contemporary clinical practice. Cardiovascular Revascularization Medicine 2018;19:960–3. https://doi.org/10.1016/J.CARREV.2018.07.021.

[2] Carabello BA. The Current Therapy for Mitral Regurgitation. J Am Coll Cardiol 2008;52:319–26. https://doi.org/10.1016/J.JACC.2008.02.084.

[3] Lorell BH, Carabello BA. Left Ventricular Hypertrophy. Circulation 2000;102:470–9. https://doi.org/10.1161/01.CIR.102.4.470.

[4] Oyama MA. Neurohormonal activation in canine degenerative mitral valve disease: implications on pathophysiology and treatment. Journal of Small Animal Practice 2009;50:3–11. https://doi.org/10.1111/J.1748-5827.2009.00801.X.

[5] Komajda M, Pousset F, Isnard R, Lechat P. The role of the neurohormonal system in heart failure. Heart 1998;79:S17. https://doi.org/10.1136/HRT.79.2008.17S.

[6] Le Tourneau T, De Groote P, Millaire A, Foucher C, Savoye C, Pigny P, et al. Effect of mitral valve surgery on exercise capacity, ventricular ejection fraction and neurohormonal activation in patients with severe mitral regurgitation. J Am Coll Cardiol 2000;36:2263–9. https://doi.org/10.1016/S0735-1097(00)01015-9.

[7] El Sabbagh A, Reddy YNV, Nishimura RA. Mitral Valve Regurgitation in the Contemporary Era: Insights Into Diagnosis, Management, and Future Directions. JACC Cardiovasc Imaging 2018;11:628–43. https://doi.org/10.1016/J.JCMG.2018.01.009.

[8] Chen HH, Schrier RW. Pathophysiology of Volume Overload in Acute Heart Failure Syndromes. Am J Med 2006;119:S11–6. https://doi.org/10.1016/J.AMJMED.2006.09.012.

[9] Wilson K, Lucchesi PA. Myofilament dysfunction as an emerging mechanism of volume overload heart failure. Pflugers Arch 2014;466:1065–77. https://doi.org/10.1007/s00424-014-1455-9.

[10] Zheng J, Li Y, Billor N, Ahmed MI, Fang Y-HD, Pat B, et al. Understanding post-surgical decline in left ventricular function in primary mitral regurgitation using regression and machine learning models. Front Cardiovasc Med 2023;10. https://doi.org/10.3389/fcvm.2023.1112797.

[11] Otto CM, Nishimura RA, Bonow RO, Carabello BA, Erwin JP, Gentile F, et al. 2020 ACC/AHA Guideline for the Management of Patients With Valvular Heart Disease: A

Report of the American College of Cardiology/American Heart Association Joint
Committee on Clinical Practice Guidelines. Circulation 2021;143:E72–227.
https://doi.org/10.1161/CIR.0000000000000923.

[12] Kerckhoffs RCP, Omens JH, McCulloch AD. A single strain-based growth law predicts
concentric and eccentric cardiac growth during pressure and volume overload. Mech Res
Commun 2012;42:40–50. https://doi.org/10.1016/j.mechrescom.2011.11.004.

[13] Witzenburg CM, Holmes JW. A Comparison of Phenomenologic Growth Laws for
Myocardial Hypertrophy. J Elast 2017;129:257–81. https://doi.org/10.1007/s10659-017-
9631-8.

[14] Seqqat R, Guo X, Rafiq K, Kolpakov MA, Guo J, Koch WJ, et al. Beta1-adrenergic
receptors promote focal adhesion signaling downregulation and myocyte apoptosis in
acute volume overload. J Mol Cell Cardiol 2012;53:240–9.
https://doi.org/10.1016/J.YJMCC.2012.05.004.

[15] Sabri A, Rafiq K, Seqqat R, Kolpakov MA, Dillon R, Dell'italia LJ. Sympathetic activation
causes focal adhesion signaling alteration in early compensated volume overload
attributable to isolated mitral regurgitation in the dog. Circ Res 2008;102:1127–36.
https://doi.org/10.1161/CIRCRESAHA.107.163642.

[16] Pat B, Killingsworth C, Denney T, Zheng J, Powell P, Tillson M, et al. Dissociation
between cardiomyocyte function and remodeling with β-adrenergic receptor blockade in
isolated canine mitral regurgitation. American Journal of Physiology-Heart and Circulatory
Physiology 2008;295:H2321–7. https://doi.org/10.1152/ajpheart.00746.2008.

[17] Zheng J, Chen Y, Pat B, Dell'Italia DLA, Tillson M, Dillon AR, et al. Molecular cardiology
microarray identifies extensive downregulation of noncollagen extracellular matrix and
profibrotic growth factor genes in chronic isolated mitral regurgitation in the dog.
Circulation 2009;119:2086–95. https://doi.org/10.1161/CIRCULATIONAHA.108.826230.

[18] Mohamed BA, Schnelle M, Khadjeh S, Lbik D, Herwig M, Linke WA, et al. Molecular and
structural transition mechanisms in long-term volume overload. Eur J Heart Fail
2016;18:362–71. https://doi.org/10.1002/EJHF.465.

[19] Frank DU, Sutcliffe MD, Saucerman JJ. Network-based predictions of in vivo cardiac
hypertrophy. J Mol Cell Cardiol 2018;121:180–9.
https://doi.org/10.1016/j.yjmcc.2018.07.243.

[20] Nakano K, Swindle MM, Spinale F, Ishihara K, Kanazawa S, Smith A, et al. Depressed
contractile function due to canine mitral regurgitation improves after correction of the
volume overload. J Clin Invest 1991;87:2077–86. https://doi.org/10.1172/JCI115238.

[21] Tsutsui H, Spinale FG, Nagatsu M, Schmid PG, Ishihara K, DeFreyte G, et al. Effects of
chronic beta-adrenergic blockade on the left ventricular and cardiocyte abnormalities of
chronic canine mitral regurgitation. Journal of Clinical Investigation 1994;93:2639–48.
https://doi.org/10.1172/JCI117277.

[22] Dell'Italia LJ, Meng QC, Balcells E, Straeter-Knowlen IM, Hankes GH, Dillon R, et al.
Increased ACE and chymase-like activity in cardiac tissue of dogs with chronic mitral

regurgitation. Https://DoiOrg/101152/Ajpheart19952696H2065 1995;269.
https://doi.org/10.1152/AJPHEART.1995.269.6.H2065.

[23] Dell'Italia LJ, Balcells E, Meng QC, Su X, Schultz D, Bishop SP, et al. Volume-overload
cardiac hypertrophy is unaffected by ACE inhibitor treatment in dogs.
Https://DoiOrg/101152/Ajpheart19972732H961 1997;273:961–70.
https://doi.org/10.1152/AJPHEART.1997.273.2.H961.

[24] Katayama K, Tajimi T, Guth BD, Matsuzaki M, Lee JD, Seitelberger R, et al. Early
diastolic filling dynamics during experimental mitral regurgitation in the conscious dog.
Circulation 1988;78:390–400. https://doi.org/10.1161/01.CIR.78.2.390.

[25] Pat B, Chen Y, Killingsworth C, Gladden JD, Shi K, Zheng J, et al. Chymase inhibition
prevents fibronectin and myofibrillar loss and improves cardiomyocyte function and LV
torsion angle in dogs with isolated mitral regurgitation. Circulation 2010;122:1488–95.
https://doi.org/10.1161/CIRCULATIONAHA.109.921619.

[26] Matsuo T, Carabello BA, Nagatomo Y, Koide M, Hamawaki M, Zile MR, et al.
Mechanisms of cardiac hypertrophy in canine volume overload. American Journal of
Physiology-Heart and Circulatory Physiology 1998;275:H65–74.
https://doi.org/10.1152/ajpheart.1998.275.1.H65.

[27] Carabello BA, Zile MR, Tanaka R, Cooper IV G. Left ventricular hypertrophy due to
volume overload versus pressure overload.
Https://DoiOrg/101152/Ajpheart19922634H1137 1992;263:H1137–44.
https://doi.org/10.1152/AJPHEART.1992.263.4.H1137.

[28] Kihara Y, Sasayama S, Miyazaki S, Onodera T, Susawa T, Nakamura Y, et al. Role of
the left atrium in adaptation of the heart to chronic mitral regurgitation in conscious dogs.
Circ Res 1988;62:543–53. https://doi.org/10.1161/01.RES.62.3.543.

[29] Perry GJ, Wei CC, Hankes GH, Dillon SR, Rynders P, Mukherjee R, et al. Angiotensin II
receptor blockade does not improve left ventricular function andremodeling in subacute
mitral regurgitation in the dog. J Am Coll Cardiol 2002;39:1374–9.
https://doi.org/10.1016/S0735-1097(02)01763-1.

[30] Tallaj J, Wei C-C, Hankes GH, Holland M, Rynders P, Dillon AR, et al. β $_1$ -Adrenergic
Receptor Blockade Attenuates Angiotensin II–Mediated Catecholamine Release Into the
Cardiac Interstitium in Mitral Regurgitation. Circulation 2003;108:225–30.
https://doi.org/10.1161/01.CIR.0000079226.48637.5A.

[31] Hankes GH, Ardell JL, Tallaj J, Wei C-C, Aban I, Holland M, et al. β1-Adrenoceptor
blockade mitigates excessive norepinephrine release into cardiac interstitium in mitral
regurgitation in dog. American Journal of Physiology-Heart and Circulatory Physiology
2006;291:H147–51. https://doi.org/10.1152/ajpheart.00951.2005.

[32] Nemoto S, Razeghi P, Ishiyama M, De Freitas G, Taegtmeyer H, Carabello BA. PPAR-γ
agonist rosiglitazone ameliorates ventricular dysfunction in experimental chronic mitral
regurgitation. American Journal of Physiology-Heart and Circulatory Physiology
2005;288:H77–82. https://doi.org/10.1152/ajpheart.01246.2003.

[33] Nemoto S, Hamawaki M, De Freitas G, Carabello BA. differential effects of the angiotensin-converting enzyme inhibitor lisinopril versus the beta-adrenergic receptor blocker atenolol on hemodynamics and left ventricular contractile function in experimental mitral regurgitation. J Am Coll Cardiol 2002;40:149–54. https://doi.org/10.1016/S0735-1097(02)01926-5.

[34] Jobe LJ, Meléndez GC, Levick SP, Du Y, Brower GL, Janicki JS. TNF-α inhibition attenuates adverse myocardial remodeling in a rat model of volume overload. American Journal of Physiology-Heart and Circulatory Physiology 2009;297:H1462–8. https://doi.org/10.1152/ajpheart.00442.2009.

[35] Lu H, Meléndez GC, Levick SP, Janicki JS. Prevention of adverse cardiac remodeling to volume overload in female rats is the result of an estrogen-altered mast cell phenotype. American Journal of Physiology-Heart and Circulatory Physiology 2012;302:H811–7. https://doi.org/10.1152/ajpheart.00980.2011.

[36] Gardner JD, Murray DB, Voloshenyuk TG, Brower GL, Bradley JM, Janicki JS. Estrogen attenuates chronic volume overload induced structural and functional remodeling in male rat hearts. American Journal of Physiology-Heart and Circulatory Physiology 2010;298:H497–504. https://doi.org/10.1152/ajpheart.00336.2009.

[37] Mclarty JL, Meléndez GC, Levick SP, Bennett S, Sabo-Attwood T, Brower GL, et al. Estrogenic modulation of inflammation-related genes in male rats following volume overload. Physiol Genomics 2012;44:362–73. https://doi.org/10.1152/PHYSIOLGENOMICS.00146.2011/SUPPL_FILE/SUPPMAT.PDF.

[38] Wang XH, Zhuo XZ, Ni YJ, Gong M, Wang TZ, Lu Q, et al. Improvement of cardiac function and reversal of gap junction remodeling by Neuregulin-1β in volume-overloaded rats with heart failure. J Geriatr Cardiol 2012;9:172. https://doi.org/10.3724/SP.J.1263.2012.03271.

[39] Willenbrock R, Stauss H, Scheuermann M, Osterziel KJ, Unger T, Dietz R. Effect of chronic volume overload on baroreflex control of heart rate and sympathetic nerve activity. Am J Physiol 1997;273:H2580-5. https://doi.org/10.1152/ajpheart.1997.273.6.H2580.

[40] Lachance D, Dhahri W, Drolet M-C, Roussel É, Gascon S, Sarrhini O, et al. Endurance training or beta-blockade can partially block the energy metabolism remodeling taking place in experimental chronic left ventricle volume overload. BMC Cardiovasc Disord 2014;14:190. https://doi.org/10.1186/1471-2261-14-190.

[41] Shuji H, Takuroh I, Takeshi M, Yuichiro I, Johji K, Kazuo K, et al. Differential responses of circulating and tissue adrenomedullin and gene expression to volume overload. J Card Fail 2000;6:120–9. https://doi.org/10.1016/S1071-9164(00)90014-9.

[42] Ishiye M, Umemura K, Uematsu T, Nakashima M. Effects of Losartan, an Angiotensin II Antagonist, on the Development of Cardiac Hypertrophy Due to Volume Overload. Biol Pharm Bull 1995;18:700–4. https://doi.org/10.1248/BPB.18.700.

[43] Bauer P, Regitz-Zagrosek V, Kallisch H, Linz W, Schoelkens B, Hildebrandt AG, et al. Myocardial angiotensin receptor type 1 gene expression in a rat model of cardiac volume

overload. Basic Res Cardiol 1997;92:139–46.
https://doi.org/10.1007/BF00788631/METRICS.

[44] Oka T, Nishimura H, Ueyama M, Kubota J, Kawamura K. Lisinopril reduces cardiac
hypertrophy and mortality in rats with aortocaval fistula. Eur J Pharmacol 1993;234:55–
60. https://doi.org/10.1016/0014-2999(93)90705-M.

[45] Isgaard J, Wåhlander H, Adams MA, Friberg P. Increased expression of growth hormone
receptor mRNA and insulin-like growth factor-I mRNA in volume-overloaded hearts.
Hypertension 1994;23:884–8. https://doi.org/10.1161/01.HYP.23.6.884.

[46] Ruzicka M, Yuan B, Harmsen E, Leenen FHH. The renin-angiotensin system and volume
overload-induced cardiac hypertrophy in rats. Effects of angiotensin converting enzyme
inhibitor versus angiotensin II receptor blocker. Circulation 1993;87:921–30.
https://doi.org/10.1161/01.CIR.87.3.921.

[47] Pu M, Gao Z, Zhang X, Liao D, Pu DK, Brennan T, et al. Impact of mitral regurgitation on
left ventricular anatomic and molecular remodeling and systolic function: implication for
outcome. American Journal of Physiology-Heart and Circulatory Physiology
2009;296:H1727–32. https://doi.org/10.1152/ajpheart.00882.2008.

[48] Kristen A V., Kreusser MM, Lehmann L, Kinscherf R, Katus HA, Haass M, et al.
Preserved Norepinephrine Reuptake but Reduced Sympathetic Nerve Endings in
Hypertrophic Volume-Overloaded Rat Hearts. J Card Fail 2006;12:577–83.
https://doi.org/10.1016/J.CARDFAIL.2006.05.006.

[49] Stefano LM De, Matsubara LS, Matsubara BB. Myocardial dysfunction with increased
ventricular compliance in volume overload hypertrophy. Eur J Heart Fail 2006;8:784–9.
https://doi.org/10.1016/J.EJHEART.2006.02.005.

[50] Zhang W, Elimban V, Xu YJ, Zhang M, Nijjar MS, Dhalla NS. Alterations of Cardiac
ERK1/2 Expression and Activity Due to Volume Overload Were Attenuated by the
Blockade of RAS. Https://DoiOrg/101177/1074248409356430 2010;15:84–92.
https://doi.org/10.1177/1074248409356430.

[51] Ichiki T, Boerrigter G, Huntley BK, Sangaralingham SJ, McKie PM, Harty GJ, et al.
Differential expression of the pro-natriuretic peptide convertases corin and furin in
experimental heart failure and atrial fibrosis. American Journal of Physiology-Regulatory,
Integrative and Comparative Physiology 2013;304:R102–9.
https://doi.org/10.1152/ajpregu.00233.2012.

[52] Suzuki H, Maehara K, Yaoita H, Maruyama Y. Altered Effects of Angiotensin II Type 1
and Type 2 Receptor Blockers on Cardiac Norepinephrine Release and Inotropic
Responses During Cardiac Sympathetic Nerve Stimulation in Aorto-Caval Shunt Rats.
Circulation Journal 2004;68:683–90. https://doi.org/10.1253/CIRCJ.68.683.

[53] Pagel I, Langenickel T, Höhnel K, Philipp S, Nüssler AK, Blum WF, et al. Cardiac and
renal effects of growth hormone in volume overload-induced heart failure: role of NO.
Hypertension 2002;39:57–62. https://doi.org/10.1161/HY0102.098323.

[54]   Asano K, Masuda K, Okumura M, Kadosawa T, Fujinaga T. Plasma Atrial and Brain Natriuretic Peptide Levels in Dogs with Congestive Heart Failure. Journal of Veterinary Medical Science 1999;61:523–9. https://doi.org/10.1292/jvms.61.523.

[55]   Asano K, Masuda K, Okumura M, Kadosawa T, Fujinaga T. Association between Exogenous Atrial Natriuretic Peptide and Hemodynamics in Dogs with Congestive Heart Failure Produced by Experimental Mitral Regurgitation. Journal of Veterinary Medical Science 2001;63:243–50. https://doi.org/10.1292/JVMS.63.243.

[56]   Hori Y, Sano N, Kanai K, Hoshi F, Itoh N, Higuchi SI. Acute cardiac volume load-related changes in plasma atrial natriuretic peptide and N-terminal pro-B-type natriuretic peptide concentrations in healthy dogs. The Veterinary Journal 2010;185:317–21. https://doi.org/10.1016/J.TVJL.2009.06.008.

[57]   Langenickel T, Pagel I, Höhnel K, Dietz R, Willenbrock R. Differential regulation of cardiac ANP and BNP mRNA in different stages of experimental heart failure. American Journal of Physiology-Heart and Circulatory Physiology 2000;278:H1500–6. https://doi.org/10.1152/ajpheart.2000.278.5.H1500.

[58]   Abassi Z, Goltsman I, Karram T, Winaver J, Hoffman A. Aortocaval fistula in rat: A unique model of volume-overload congestive heart failure and cardiac hypertrophy. J Biomed Biotechnol 2011;2011. https://doi.org/10.1155/2011/729497.

[59]   Huang M, Hester RL, Guyton AC. Hemodynamic changes in rats after opening an arteriovenous fistula. Https://DoiOrg/101152/Ajpheart19922623H846 1992;262. https://doi.org/10.1152/AJPHEART.1992.262.3.H846.

[60]   Nagatsu M, Zile MR, Tsutsui H, Schmid PG, DeFreyte G, Cooper IV G, et al. Native beta-adrenergic support for left ventricular dysfunction in experimental mitral regurgitation normalizes indexes of pump and contractile function. Circulation 1994;89:818–26. https://doi.org/10.1161/01.CIR.89.2.818.

[61]   Lee JD, Sasayama S, Kihara Y, Ohyagi A, Fujisawa A, Yui Y, et al. Adaptations of the left ventricle to chronic volume overload induced by mitral regurgitation in conscious dogs. Heart Vessels 1985;1:9–15. https://doi.org/10.1007/BF02066481/METRICS.

[62]   Kristen A V., Just A, Haass M, Seller H. Central hypercapnic chemoreflex modulation of renal sympathetic nerve activity in experimental heart failure. Basic Res Cardiol 2002;97:177–86. https://doi.org/10.1007/S003950200009/METRICS.

[63]   Ray L, Mathieu M, Jespers P, Hadad I, Mahmoudabady M, Pensis A, et al. Early increase in pulmonary vascular reactivity with overexpression of endothelin-1 and vascular endothelial growth factor in canine experimental heart failure. Exp Physiol 2008;93:434–42. https://doi.org/10.1113/EXPPHYSIOL.2007.040469.

[64]   Cavero PG, Miller WL, Heublein DM, Margulies KB, Burnett JC. Endothelin in experimental congestive heart failure in the anesthetized dog. Https://DoiOrg/101152/Ajprenal19902592F312 1990;259. https://doi.org/10.1152/AJPRENAL.1990.259.2.F312.

[65]   Humphrey JD. Review Paper: Continuum biomechanics of soft biological tissues. Proceedings of the Royal Society of London Series A: Mathematical, Physical and Engineering Sciences 2003;459:3–46. https://doi.org/10.1098/rspa.2002.1060.

[66]   Ross J, Sonnenblick EH, Taylor RR, Spotnitz HM, Covell JW. Diastolic Geometry and Sarcomere Lengths in the Chronically Dilated Canine Left Ventricle. Circ Res 1971;28:49–61. https://doi.org/10.1161/01.RES.28.1.49.

[67]   Emery JL, Omens JH. Mechanical regulation of myocardial growth during volume-overload hypertrophy in the rat. American Journal of Physiology-Heart and Circulatory Physiology 1997;273:H1198–204. https://doi.org/10.1152/ajpheart.1997.273.3.H1198.

[68]   Ross J, McCullagh WH. Nature of Enhanced Performance of the Dilated Left Ventricle in the Dog during Chronic Volume Overloading. Circ Res 1972;30:549–56. https://doi.org/10.1161/01.RES.30.5.549.

[69]   Spotnitz HM, Sonnenblick EH, Spiro D. Relation of Ultrastructure to Function in the Intact Heart: Sarcomere Structure Relative to Pressure Volume Curves of Intact Left Ventricles of Dog and Cat. Circ Res 1966;18:49–66. https://doi.org/10.1161/01.RES.18.1.49.

[70]   Fomovsky GM, Clark SA, Parker KM, Ailawadi G, Holmes JW. Anisotropic reinforcement of acute anteroapical infarcts improves pump function. Circ Heart Fail 2012;5:515–22. https://doi.org/10.1161/CIRCHEARTFAILURE.111.965731.

[71]   Witzenburg CM, Holmes JW. Predicting the Time Course of Ventricular Dilation and Thickening Using a Rapid Compartmental Model. J Cardiovasc Transl Res 2018;11:109–22. https://doi.org/10.1007/S12265-018-9793-1.

[72]   Khalilimeybodi A, Paap AM, Christiansen SLM, Saucerman JJ. Context-specific network modeling identifies new crosstalk in β-adrenergic cardiac hypertrophy. PLoS Comput Biol 2020;16:e1008490. https://doi.org/10.1371/journal.pcbi.1008490.

[73]   Ryall KA, Holland DO, Delaney KA, Kraeutler MJ, Parker AJ, Saucerman JJ. Network reconstruction and systems analysis of cardiac myocyte hypertrophy signaling. Journal of Biological Chemistry 2012;287:42259–68. https://doi.org/10.1074/jbc.M112.382937.

[74]   Clark AP, Chowkwale M, Paap A, Dang S, Saucerman JJ. Logic-based modeling of biological networks with Netflux. BioRxiv 2024:2024.01.11.575227. https://doi.org/10.1101/2024.01.11.575227.

[75]   Efendiev Y, Hou T, Luo W. Preconditioning Markov Chain Monte Carlo Simulations Using Coarse-Scale Models. Https://DoiOrg/101137/050628568 2006;28:776–803. https://doi.org/10.1137/050628568.

[76]   Dhamala J, Arevalo HJ, Sapp J, Horácek BM, Wu KC, Trayanova NA, et al. Quantifying the uncertainty in model parameters using Gaussian process-based Markov chain Monte Carlo in cardiac electrophysiology. Med Image Anal 2018;48:43–57. https://doi.org/10.1016/J.MEDIA.2018.05.007.

[77]   Dent MR, Das S, Dhalla NS. Alterations in both death and survival signals for apoptosis in heart failure due to volume overload. J Mol Cell Cardiol 2007;43:726–32. https://doi.org/10.1016/J.YJMCC.2007.09.001.

[78] Liu Y, Dillon AR, Tillson M, Makarewich C, Nguyen V, Dell'Italia L, et al. Volume overload
induces differential spatiotemporal regulation of myocardial soluble guanylyl cyclase in
eccentric hypertrophy and heart failure. J Mol Cell Cardiol 2013;60:72–83.
https://doi.org/10.1016/J.YJMCC.2013.03.019.

[79] Kolpakov MA, Seqqat R, Rafiq K, Xi H, Margulies KB, Libonati JR, et al. Pleiotropic
effects of neutrophils on myocyte apoptosis and left ventricular remodeling during early
volume overload. J Mol Cell Cardiol 2009;47:634–45.
https://doi.org/10.1016/J.YJMCC.2009.08.016.

[80] Trappanese DM, Liu Y, McCormick RC, Cannavo A, Nanayakkara G, Baskharoun MM, et
al. Chronic β1-adrenergic blockade enhances myocardial β3-adrenergic coupling with
nitric oxide-cGMP signaling in a canine model of chronic volume overload: new insight
into mechanisms of cardiac benefit with selective β1-blocker therapy. Basic Res Cardiol
2015;110:456. https://doi.org/10.1007/s00395-014-0456-3.

[81] Arnal JF, Philippe M, Laboulandine I, Michel JB. Effect of perindopril in rat cardiac volume
overload. Am Heart J 1993;126:776–82. https://doi.org/10.1016/0002-8703(93)90929-4.

[82] Urabe Y, Mann DL, Kent RL, Nakano K, Tomanek RJ, Carabello BA, et al. Cellular and
ventricular contractile dysfunction in experimental canine mitral regurgitation. Circ Res
1992;70:131–47. https://doi.org/10.1161/01.RES.70.1.131.

[83] Spinale FG, Ishihra K, Zile M, DeFryte G, Crawford FA, Carabello BA. Structural basis for
changes in left ventricular function and geometry because of chronic mitral regurgitation
and after correction of volume overload. J Thorac Cardiovasc Surg 1993;106:1147–57.
https://doi.org/10.1016/S0022-5223(19)33992-3.

[84] Du Y, Plante E, Janicki JS, Brower GL. Temporal Evaluation of Cardiac Myocyte
Hypertrophy and Hyperplasia in Male Rats Secondary to Chronic Volume Overload. Am J
Pathol 2010;177:1155–63. https://doi.org/10.2353/ajpath.2010.090587.

[85] Zeigler AC, Chandrabhatla AS, Christiansen SL, Nelson AR, Holmes JW, Saucerman JJ.
Network model-based screen for FDA-approved drugs affecting cardiac fibrosis. CPT
Pharmacometrics Syst Pharmacol 2021;10:377–88. https://doi.org/10.1002/PSP4.12599.

[86] Leenen FHH, White R, Yuan B. Isoproterenol-induced cardiac hypertrophy: role of
circulatory versus cardiac renin-angiotensin system. American Journal of Physiology-
Heart and Circulatory Physiology 2001;281:H2410–6.
https://doi.org/10.1152/ajpheart.2001.281.6.H2410.

[87] Ennis IL, Escudero EM, Console GM, Camihort G, Dumm CG, Seidler RW, et al.
Regression of Isoproterenol-Induced Cardiac Hypertrophy by Na+/H+ Exchanger
Inhibition. Hypertension 2003;41:1324–9.
https://doi.org/10.1161/01.HYP.0000071180.12012.6E.

[88] Kitagawa Y, Yamashita D, Ito H, Takaki M. Reversible effects of isoproterenol-induced
hypertrophy on in situ left ventricular function in rat hearts. American Journal of
Physiology-Heart and Circulatory Physiology 2004;287:H277–85.
https://doi.org/10.1152/ajpheart.00073.2004.

[89] Murad N, Tucci PJ. Isoproterenol-Induced Hypertrophy May Result In Distinct Left
Ventricular Changes. Clin Exp Pharmacol Physiol 2000;27:352–7.
https://doi.org/10.1046/j.1440-1681.2000.03254.x.

[90] Nagano M, Higaki J, Nakamura F, Higashimori K, Nagano N, Mikami H, et al. Role of
cardiac angiotensin II in isoproterenol-induced left ventricular hypertrophy. Hypertension
1992;19:708–12. https://doi.org/10.1161/01.HYP.19.6.708.

[91] Brand T, Sharma HS, Schaper W. Expression of nuclear proto-oncogenes in
isoproterenol-induced cardiac hypertrophy. J Mol Cell Cardiol 1993;25:1325–37.
https://doi.org/10.1006/jmcc.1993.1145.

[92] Al-Rasheed NM, Al-Oteibi MM, Al-Manee RZ, Al-Shareef SA, Al-Rasheed NM, Hasan IH,
et al. Simvastatin prevents isoproterenol-induced cardiac hypertrophy through modulation
of the JAK/STAT pathway. Drug Des Devel Ther 2015;9:3217–29.
https://doi.org/10.2147/DDDT.S86431.

[93] Chowdhury D, Tangutur AD, Khatua TN, Saxena P, Banerjee SK, Bhadra MP. A
proteomic view of isoproterenol induced cardiac hypertrophy: Prohibitin identified as a
potential biomarker in rats. J Transl Med 2013;11:130. https://doi.org/10.1186/1479-5876-
11-130.

[94] Takeshita D, Shimizu J, Kitagawa Y, Yamashita D, Tohne K, Nakajima-Takenaka C, et al.
Isoproterenol-Induced Hypertrophied Rat Hearts: Does Short-Term Treatment
Correspond to Long-Term Treatment? The Journal of Physiological Sciences
2008;58:179–88. https://doi.org/10.2170/PHYSIOLSCI.RP004508.

[95] Boluyt MO, Long X, Eschenhagen T, Mende U, Schmitz W, Crow MT, et al. Isoproterenol
infusion induces alterations in expression of hypertrophy-associated genes in rat heart.
Https://DoiOrg/101152/Ajpheart19952692H638 1995;269.
https://doi.org/10.1152/AJPHEART.1995.269.2.H638.

[96] Golomb E, Abassi ZA, Cuda G, Stylianou M, Panchal VR, Trachewsky D, et al.
Angiotensin II maintains, but does not mediate, isoproterenol-induced cardiac
hypertrophy in rats. Https://DoiOrg/101152/Ajpheart19942674H1496 1994;267.
https://doi.org/10.1152/AJPHEART.1994.267.4.H1496.

[97] Miyoshi T, Nakamura K, Miura D, Yoshida M, Saito Y, Akagi S, et al. Effect of LCZ696, a
dual angiotensin receptor neprilysin inhibitor, on isoproterenol-induced cardiac
hypertrophy, fibrosis, and hemodynamic change in rats. Cardiol J 2019;26:575–83.
https://doi.org/10.5603/CJ.A2018.0048.

[98] Zou L-X, Imig JD, Von Thun AM, Hymel A, Ono H, Navar LG. Receptor-Mediated
Intrarenal Angiotensin II Augmentation in Angiotensin II–Infused Rats. Hypertension
1996;28:669–77. https://doi.org/10.1161/01.HYP.28.4.669.

[99] Van Eekelen JAM, Phillips MI. Plasma angiotensin II levels at moment of drinking during
angiotensin II intravenous infusion. Https://DoiOrg/101152/Ajpregu19882553R500
1988;255. https://doi.org/10.1152/AJPREGU.1988.255.3.R500.

[100]   Griffin SA, Brown WCB, MacPherson F, McGrath JC, Wilson VG, Korsgaard N, et al.
        Angiotensin II causes vascular hypertrophy in part by a non-pressor mechanism.
        Hypertension 1991;17:626–35. https://doi.org/10.1161/01.HYP.17.5.626.

[101]   Dostal DE, Baker KM. Angiotensin II Stimulation of Left Ventricular Hypertrophy in Adult
        Rat HeartMediation by the AT1 Receptor. Am J Hypertens 1992;5:276–80.
        https://doi.org/10.1093/AJH/5.5.276.

[102]   Dilley R. Heparin inhibits mesenteric vascular hypertrophy in angiotensin II-infusion
        hypertension in rats. Cardiovasc Res 1998;38:247–55. https://doi.org/10.1016/S0008-
        6363(98)00004-2.

[103]   Cassis LA, Marshall DE, Fettinger MJ, Rosenbluth B, Lodder RA. Mechanisms
        contributing to angiotensin II regulation of body weight. American Journal of Physiology-
        Endocrinology and Metabolism 1998;274:E867–76.
        https://doi.org/10.1152/ajpendo.1998.274.5.E867.

[104]   Fabris B, Candido R, Bortoletto M, Zentilin L, Sandri M, Fior F, et al. Dose and time-
        dependent apoptotic effects by angiotensin II infusion on left ventricular cardiomyocytes.
        J Hypertens 2007;25:1481–90. https://doi.org/10.1097/HJH.0B013E328121AAE7.

[105]   Baltatu O, Silva JA, Ganten D, Bader M. The Brain Renin-Angiotensin System Modulates
        Angiotensin II–Induced Hypertension and Cardiac Hypertrophy. Hypertension
        2000;35:409–12. https://doi.org/10.1161/01.HYP.35.1.409.

[106]   Kim S, Ohta K, Hamaguchi A, Yukimura T, Miura K, Iwao H. Angiotensin II Induces
        Cardiac Phenotypic Modulation and Remodeling In Vivo in Rats. Hypertension
        1995;25:1252–9. https://doi.org/10.1161/01.HYP.25.6.1252.

[107]   Mishra JS, More AS, Gopalakrishnan K, Kumar S. Testosterone plays a permissive role
        in angiotensin II-induced hypertension and cardiac hypertrophy in male rats. Biol Reprod
        2019;100:139–48. https://doi.org/10.1093/BIOLRE/IOY179.

[108]   Grobe JL, Mecca AP, Lingis M, Shenoy V, Bolton TA, Machado JM, et al. Prevention of
        angiotensin II-induced cardiac remodeling by angiotensin-(1–7). American Journal of
        Physiology-Heart and Circulatory Physiology 2007;292:H736–42.
        https://doi.org/10.1152/ajpheart.00937.2006.

[109]   Goldspink PH, McKinney RD, Kimball VA, Geenen DL, Buttrick PM. Angiotensin II
        induced cardiac hypertrophy in vivo is inhibited by cyclosporin A in adult rats. Mol Cell
        Biochem 2001;226:83–8. https://doi.org/10.1023/A:1012789819926/METRICS.

[110]   Wu R, Laplante M-A, De Champlain J. Prevention of angiotensin II-induced hypertension,
        cardiovascular hypertrophy and oxidative stress by acetylsalicylic acid in rats. J
        Hypertens 2004;22:793–801. https://doi.org/10.1097/01.hjh.0000098277.36684.6b.

[111]   Herizi A, Jover B, Bouriquet N, Mimran A. Prevention of the Cardiovascular and Renal
        Effects of Angiotensin II by Endothelin Blockade. Hypertension 1998;31:10–4.
        https://doi.org/10.1161/01.HYP.31.1.10.

[112] Laks MM, Morady F, Swan HJC. Myocardial Hypertrophy Produced by Chronic Infusion of Subhypertensive Doses of Norepinephrine in the Dog. Chest 1973;64:75–8. https://doi.org/10.1378/chest.64.1.75.

[113] King BD, Sack D, Kichuk MR, Hintze TH. Absence of hypertension despite chronic marked elevations in plasma norepinephrine in conscious dogs. Hypertension 1987;9:582–90. https://doi.org/10.1161/01.HYP.9.6.582.

[114] Stewart JM, Patel MB, Wang J, Ochoa M, Gewitz M, Loud A V., et al. Chronic elevation of norepinephrine in conscious dogs produces hypertrophy with no loss of LV reserve. Https://DoiOrg/101152/Ajpheart19922622H331 1992;262:331–9. https://doi.org/10.1152/AJPHEART.1992.262.2.H331.

[115] Raum WJ, Laks MM, Garner D, Swerdloff RS. β-Adrenergic receptor and cyclic AMP alterations in the canine ventricular septum during long-term norepinephrine infusion: Implications for hypertrophic cardiomyopathy. Circulation 1983;68:693–9. https://doi.org/10.1161/01.CIR.68.3.693.

[116] Hanada K, Asari K, Saito M, Kawana J ichi, Mita M, Ogata H. Comparison of pharmacodynamics between carvedilol and metoprolol in rats with isoproterenol-induced cardiac hypertrophy: Effects of carvedilol enantiomers. Eur J Pharmacol 2008;589:194–200. https://doi.org/10.1016/J.EJPHAR.2008.04.055.

[117] Grimm D, Holmer SR, Riegger GAJ, Kromer EP. Effects of Beta-Receptor Blockade and Angiotensin II Type I Receptor Antagonism in Isoproterenol - Induced Heart Failure in the Rat. Cardiovascular Pathology 1999;8:315–23. https://doi.org/10.1016/S1054-8807(99)00021-6.

[118] Goyal BR, Mehta AA. Benefi cial role of spironolactone, telmisartan and their combination on isoproterenol-induced cardiac hypertrophy. Acta Cardiol 2012;67:203–11. https://doi.org/10.1080/AC.67.2.2154211.

[119] Ficai S, Herizi A, Mimran A, Jover B. Endothelin Blockade In Angiotensin Ii Hypertension: Prevention And Treatment Studies In The Rat. Clin Exp Pharmacol Physiol 2001;28:1100–3. https://doi.org/10.1046/J.1440-1681.2001.03568.X.

[120] Park JB, Schiffrin EL. Cardiac and vascular fibrosis and hypertrophy in aldosterone-infused rats: role of endothelin-1. Am J Hypertens 2002;15:164–9. https://doi.org/10.1016/S0895-7061(01)02291-9.

[121] Kaddoura S, Firth JD, Boheler KR, Sugden PH, Poole-Wilson PA. Endothelin-1 Is Involved in Norepinephrine-Induced Ventricular Hypertrophy in Vivo Acute Effects of Bosentan, an Orally Active, Mixed Endothelin ETA and ETB Receptor Antagonist. Circulation 1996;93:2068–79. https://doi.org/10.1161/01.CIR.93.11.2068/FORMAT/EPUB.

[122] Saucerman JJ, Tan PM, Buchholz KS, McCulloch AD, Omens JH. Mechanical regulation of gene expression in cardiac myocytes and fibroblasts. Nat Rev Cardiol 2019;16:361. https://doi.org/10.1038/S41569-019-0155-8.

[123] Taegtmeyer H, Sen S, Vela D. Return to the fetal gene program: A suggested metabolic
link to gene expression in the heart. Ann N Y Acad Sci 2010;1188:191.
https://doi.org/10.1111/J.1749-6632.2009.05100.X.

[124] Kobayashi M, Machida N, Tanaka R, Yamane Y. Effects of β-Blocker on Left Ventricular
Remodeling in Rats with Volume Overload Cardiac Failure. Journal of Veterinary Medical
Science 2008;70:1231–7. https://doi.org/10.1292/JVMS.70.1231.

[125] Murray DB, McMillan R, Brower GL, Janicki JS. ET $_A$ selective receptor antagonism
prevents ventricular remodeling in volume-overloaded rats. American Journal of
Physiology-Heart and Circulatory Physiology 2009;297:H109–16.
https://doi.org/10.1152/ajpheart.00968.2008.

[126] Murray DB, Gardner JD, Brower GL, Janicki JS. Effects of nonselective endothelin-1
receptor antagonism on cardiac mast cell-mediated ventricular remodeling in rats.
American Journal of Physiology-Heart and Circulatory Physiology 2008;294:H1251–7.
https://doi.org/10.1152/ajpheart.00622.2007.

[127] Lee DS, Kim DK, Choi SM, Kim YK, Ko BH, Jung YW. Bosentan Attenuates
Compensatory Left Ventricular Hypertrophy Induced by Aortocaval Fistula in Rats.
Korean Circ J 2005;35:665. https://doi.org/10.4070/kcj.2005.35.9.665.

[128] Francis B, Winaver J, Karram T, Hoffman A, Abassi Z. Renal and systemic effects of
chronic blockade of ETA or ET B in normal rats and animals with experimental heart
failure. J Cardiovasc Pharmacol 2004;44.
https://doi.org/10.1097/01.FJC.0000166214.42791.F2.

[129] Caggiano LR, Holmes JW, Witzenburg CM. Individual variability in animal-specific
hemodynamic compensation following myocardial infarction. J Mol Cell Cardiol
2022;163:156–66. https://doi.org/10.1016/j.yjmcc.2021.10.008.

[130] Yoshida K, Holmes JW. Computational models of cardiac hypertrophy. Prog Biophys Mol
Biol 2021;159:75–85. https://doi.org/10.1016/J.PBIOMOLBIO.2020.07.001.

[131] Kleaveland JP, Kussmaul WG, Vinciguerra T, Diters R, Carabello BA. Volume overload
hypertrophy in a closed-chest model of mitral regurgitation. American Journal of
Physiology-Heart and Circulatory Physiology 1988;254:H1034–41.
https://doi.org/10.1152/ajpheart.1988.254.6.H1034.

[132] Nakano K, Sugawara M, Ishihara K, Kanazawa S, Corin WJ, Denslow S, et al. Myocardial
stiffness derived from end-systolic wall stress and logarithm of reciprocal of wall
thickness. Contractility index independent of ventricular size. Circulation 1990;82:1352–
61. https://doi.org/10.1161/01.CIR.82.4.1352.

[133] Carabello BA, Nakano K, Corin W, Biederman R, Spann JF. Left ventricular function in
experimental volume overload hypertrophy.
Https://DoiOrg/101152/Ajpheart19892564H974 1989;256.
https://doi.org/10.1152/AJPHEART.1989.256.4.H974.

[134] Berko B, Gaasch WH, Tanigawa N, Smith D, Craige E. Disparity between ejection and
end-systolic indexes of left ventricular contractility in mitral regurgitation. Circulation
1987;75:1310–9. https://doi.org/10.1161/01.CIR.75.6.1310.

[135]   Carabello BA, Nakano K, Ishihara K, Kanazawa S, W Biederman RW, Spann Jr JF.
1388   Coronary blood flow in dogs with contractile dysfunction due to experimental volume
1389   overload. Circulation 1991;83:1063–75. https://doi.org/10.1161/01.CIR.83.3.1063.

1390   [136]   Stewart JA, Wei CC, Brower GL, Rynders PE, Hankes GH, Dillon AR, et al. Cardiac mast
1391   cell- and chymase-mediated matrix metalloproteinase activity and left ventricular
1392   remodeling in mitral regurgitation in the dog. J Mol Cell Cardiol 2003;35:311–9.
1393   https://doi.org/10.1016/S0022-2828(03)00013-0.

1394   [137]   Ahmed A, Dell'Italia LJ. Use of Beta-Blockers in Older Adults with Chronic Heart Failure.
1395   Am J Med Sci 2004;328:100–11. https://doi.org/10.1097/00000441-200408000-00006.

1396   [138]   Ahmed MI, Aban I, Lloyd SG, Gupta H, Howard G, Inusah S, et al. A Randomized
1397   Controlled Phase IIb Trial of Beta1-Receptor Blockade for Chronic Degenerative Mitral
1398   Regurgitation. J Am Coll Cardiol 2012;60:833–8.
1399   https://doi.org/10.1016/J.JACC.2012.04.029.

1400   [139]   Fareh J, Touyz RM, Schiffrin EL, Thibault G. Endothelin-1 and Angiotensin II Receptors
1401   in Cells From Rat Hypertrophied Heart. Circ Res 1996;78:302–11.
1402   https://doi.org/10.1161/01.RES.78.2.302.

1403   [140]   Leskinen H, Vuolteenaho O, Ruskoaho H. Combined Inhibition of Endothelin and
1404   Angiotensin II Receptors Blocks Volume Load–Induced Cardiac Hormone Release. Circ
1405   Res 1997;80:114–23. https://doi.org/10.1161/01.RES.80.1.114.

1406   [141]   Enevoldsen FC, Sahana J, Wehland M, Grimm D, Infanger M, Krüger M. Endothelin
1407   Receptor Antagonists: Status Quo and Future Perspectives for Targeted Therapy.
1408   Journal of Clinical Medicine 2020, Vol 9, Page 824 2020;9:824.
1409   https://doi.org/10.3390/JCM9030824.

1410

$t/\tau = 3.0$

VO

a) N=50 (graph with Normalized Growth vs t/τ) N=1000 (Chronic VO box plot)

b)

Simulation Experiments	SERCA	αMHC	βMHC	ANP	BNP	CellArea
	3/3	3/4	3/3	5/5	6/6	13/16

ISO Infusion

c) N=50 (graph with Normalized Growth vs t/τ) N=1000 (Chronic ISO box plot)

d)

Simulation Experiments	SERCA	αMHC	βMHC	ANP	BNP	CellArea
	4/4	1/1	1/1	1/1		14/14

AngII Infusion

e) N=50 (graph with Normalized Growth vs t/τ) N=1000 (Chronic AngII box plot)

f)

Simulation Experiments	SERCA	αMHC	βMHC	ANP	BNP	CellArea
	1/1	0/1	2/2	2/2		3/3

Legend:

#/# Number of reports with significant change / number of reports

Increase in >75% of simulations or significant increase in experiments

Mixed simulation result or non-significant effects in experiments

Decrease in >75% of simulations or significant decrease in experiments

No available data