Identification of SPP1-positive macrophages by single-cell spatial analysis in human lung tissues with mycobacterial infection

Harutaka Katano[1], Akira Hebisawa[2], Yuko Sato[1], Yoshihiko Hoshino[3]

[1]Department of Pathology, National Institute of Infectious Diseases, Tokyo, Japan.
[2]Clinical Research Center, National Hospital Organization Tokyo National Hospital, Tokyo, Japan.
[3]Department of Mycobacteriology, Leprosy Research Center, National Institute of Infectious Diseases, Tokyo, Japan.

Address for correspondence: Harutaka Katano, Department of Pathology, National Institute of Infectious Diseases, 1-23-1 Toyama, Shinjuku, Tokyo 162-8640, Japan. e-mail: katano@niid.go.jp and Yoshihiko Hoshino, Department of Mycobacteriology, Leprosy Research Center, National Institute of Infectious Diseases, 4-2-1 Aoba, Higashi-Murayama, Tokyo, 189-0002, Japan e-mail: yhoshino@niid.go.jp

Abstract

Tuberculosis and non-tuberculous mycobacterial (NTM) diseases are infections caused

by *Mycobacterium tuberculosis* and non-tuberculous mycobacteria such as

Mycobacterium avium complex, leading to the formation of granulomatous lesions with

caseous necrosis in the lungs. Although granulomatous tissues are infiltrated by numerous

inflammatory cells, including macrophages, lymphocytes, and neutrophils, the

mechanisms underlying granuloma formation by mycobacteria remain unclear. In this

study, we performed single-cell spatial analysis on lung tissue samples from patients with

tuberculosis and NTM diseases to investigate the infiltrating cell populations. We

analyzed seven lung lesions and identified individual cell types infiltrating the

granulomatous tissue. Based on gene expression profiles, at least four macrophage

subtypes were identified. Notably, SPP1-positive macrophages predominantly infiltrated

the granulomatous tissue. Langhans giant cells expressed SPP1, and many SPP1-positive

macrophages without giant cell morphology were also present around the granuloma.

RNA-seq analysis revealed elevated SPP1 expression in mycobacterium-infected tissues.

The SPP1-CD44 signaling pathway was active in SPP1-positive macrophages and their

neighboring cells in mycobacterium-infected tissues. SPP1-positive macrophages were

also observed around granulomas in other granulomatous diseases, such as

granulomatosis with polyangiitis and sarcoidosis. These findings suggest that SPP1-

positive macrophages may play a key role in granuloma formation in granulomatous

diseases, including mycobacterial infections.

INTRODUCTION

Tuberculosis and non-tuberculous mycobacterial (NTM) diseases, caused by

Mycobacterium tuberculosis and various non-tuberculous mycobacteria—particularly the

Mycobacterium avium intracellulare complex (MAC)—represent significant public

health challenges[1]. These infections predominantly affect the lungs, leading to the

formation of granulomatous lesions, which are characterized by caseous necrosis[2].

Despite extensive research efforts, the mechanisms underlying granuloma formation in

mycobacterial infections remain largely enigmatic and poorly understood[3-5].

Granulomas are highly organized structures that form in response to persistent pathogens.

These cellular conglomerates, composed of immune cells such as macrophages,

lymphocytes, and neutrophils, aim to contain and neutralize the pathogen, serving as a

barrier to prevent its spread[2,4,6]. Immunohistochemical and special staining techniques

have detected mycobacterial antigens within the caseous necrotic tissue of granulomas,

supporting the hypothesis that granulomas play a crucial role in controlling the

dissemination of mycobacteria[3]. However, the precise formation and dynamic regulation

of these structures, particularly in the context of mycobacterial infections, remain

incompletely understood.

Various immune cells, including macrophages, B cells, T cells, NK cells, dendritic cells,

and neutrophils, participate in granuloma formation.[2,7]. Among these, macrophages are

pivotal, exhibiting remarkable plasticity that allows them to adapt to a wide range of

environmental stimuli. During mycobacterial infections, macrophages can assume

diverse functional states, contributing both to pathogen clearance and, paradoxically, to

pathogen survival[3,4,8,9]. Macrophages, along with dendritic cells and alveolar epithelial

cells, phagocytize mycobacteria during the initial stages of infection. The immune

response to mycobacteria triggers several signaling pathways, including IFN-γ and TLR2

signaling, which promote the production of pro-inflammatory cytokines such as IL-12,

IL-23, and TNF-α. These signals, particularly through TLR2, ultimately contribute to the

formation of granulomas.[8,10-13].

To gain insight into the cellular responses to mycobacterial infections in the lung, it is

critical to identify the specific cell types infiltrating granulomas and to characterize their

dynamic interactions. To this end, we employed the cutting-edge Nanostring CosMx™

platform to perform single-cell spatial analysis on lung tissue samples from patients with

MTB or NTM diseases.[14]. This advanced technology enabled us to map and identify

individual cell types surrounding the granulomatous tissue. In addition, single-cell spatial

analysis allowed for detailed gene expression profiling within individual cells, providing
insights into the localization of cells and the identification of key signaling pathways
involved in cell-cell communication within the tissue microenvironment. This innovative
approach holds promise for unraveling the complex interplay of immune cells during
MTB and NTM infections.
In our study, we applied single-cell spatial analysis to explore the cellular landscape of
lung tissue samples from patients with MTB or NTM infections, with a particular
emphasis on macrophages and their critical role in granuloma formation.
85
86

RESULTS
Cell typing in lung tissues with mycobacterial infection
Seven lung tissue samples from patients with MTB (n = 3) and NTM (n = 4) infections
were analyzed using the Nanostring CosMx™ platform (Supplementary Table 1). Each
slide contained granulomas with inflammatory cell infiltrates, including foreign body
macrophages (Langhans giant cells, Supplementary Figures 1a and 1b).
Immunohistochemistry and Ziehl-Neelsen staining demonstrated that mycobacteria
bacilli were predominantly detected in the center of the granulomas, but were absent in
the surrounding inflammatory cell infiltrates in both NTM and MTB samples
(Supplementary Figure 2). CosMx™ analysis identified a total of 300,841 cells across the
seven slides (Supplementary Table 2). Cluster analysis revealed 26 distinct clusters based
on gene expression profiles (Figure 1). Differentially expressed gene (DEG) analysis
identified 520 cluster-specific marker genes with adjusted p-values. These DEGs (gene
lists available in Supplementary Data: marker_heatmap_nb_clus_4.csv) were used to
annotate subpopulations. The cell types within each cluster were determined by SingleR.
Neutrophil and B-cell lineages were manually annotated based on gene expression data
from the Human Gene Atlas (Gene Set Enrichment Analysis, GSEA, https://www.gsea-
msigdb.org/gsea/index.jsp). The proportions of cell types varied across samples and fields
of view (FoVs) (Figure 1E and Supplementary Figures 3-5).
Through in silico analysis based on gene expression, at least four subtypes (a-d) of
macrophages were identified. Macrophage subtypes were categorized based on the
expression of specific markers such as CD14, CD163, CD68, and LYZ. Macrophage
subtypes *a*, *b*, and *c* were classified conventionally as M1, M2, and M2 macrophages,
respectively (Figures 2A and 3B). Macrophage subtype *d*, which expressed CCR7 and
did not express CD163 (suggesting an M1 phenotype), predominantly expressed SPP1
and MARCO (Figures 2A-D). Consequently, this subtype was designated as SPP1-

positive macrophages. SPP1-positive macrophages were present in all NTM and MTB
slides, although their abundance varied among the samples (Figure 1E, Supplementary
Table 3).

SPP1 macrophages infiltrate around MTB and NTM granuloma

CosMx™ spatial imaging revealed that SPP1-positive macrophages predominantly
infiltrated the periphery of NTM-positive granulomatous tissue (Figure 3, Supplementary
Figures 6a and 6b). In MTB-infected tissues, SPP1-positive macrophages were also
observed in the interstitial regions and alveolar spaces (Figure 4, Supplementary Figure
6c). Both CosMx™ spatial images and immunohistochemistry using an anti-SPP1
antibody demonstrated that SPP1 was localized in the cytoplasm of Langhans giant cells
(Figures 3 and 4), suggesting that these giant cells are a form of SPP1-positive
macrophages.

Additionally, CosMx™ spatial imaging and immunohistochemistry indicated that SPP1
was present in macrophages without giant cell morphology, located around
mycobacterium-positive granulomas in proximity to Langhans giant cells (Figure 3). This
observation implies that SPP1-positive macrophages initially infiltrate around the
granulomas, forming a barrier of macrophages, and that some of these cells may
subsequently fuse to form Langhans giant cells.

Bulk RNA-seq analysis of RNA extracted from mycobacterium-infected tissues also
revealed elevated expression of SPP1 in infected tissues compared to controls (Figures
5A-5C). Gene ontology analysis indicated that cytokine-mediated signaling pathways and
extracellular matrix organization were upregulated in mycobacterium-infected samples
(Figure 5D). These processes involved high expression of SPP1 and CD44, respectively.
Collectively, these findings suggest that SPP1-positive macrophages may play a critical
role in granuloma formation during mycobacterial infections.

Ligand-receptor interaction of SPP1

Using spatial transcriptome data, the R package InSituCor was employed for spatial
correlation analysis to detect significant ligand-receptor interactions in the samples. This
analysis identified SPP1, FN1, and MARCO as co-expression markers of SPP1-positive
macrophages (Figure 6A). Furthermore, InSituCor detected SPP1-CD44 and FN1-CD44
pairs as active ligand-receptor interactions in both NTM and MTB samples (Figures 6B-
C). Spatial imaging of these samples demonstrated that CD44-positive cells were
localized near SPP1-positive macrophages, though CD44-positive cells were distributed
more broadly compared to SPP1-positive macrophages. Spatial analysis also revealed co-
expression of CD44 and SPP1 in SPP1-positive macrophages (macrophage_d) of tissues
with mycobacterial infection (Figures 6D and 6E). Dot plot analysis showed that CD44

was expressed by a variety of cell types, including macrophages, mast cells, CD8+ T cells,
and dendritic cells around the granulomatous lesions, while SPP1 expression was
predominantly confined to the macrophage_d subtype (Figure 7A).
Immunohistochemistry for SPP1 and CD44 showed SPP1 and CD44 expression in
Langhans giant cells and various cells around SPP1-positive macrophages, respectively
(Figure 7B). Spatial images of CosMx™ confirmed that CD44 was expressed not only in
macrophages but also in various cell types, including B cells, T cells, and epithelial cells
surrounding SPP1-positive macrophages (Figure 7C). The SPP1-CD44 interaction was
also identified using the CellChat package (Figure 8). SPP1, expressed by macrophage_d,
was shown to interact with various cell types. Circle plots and heatmaps of the SPP1
signaling pathway network indicated that the receiver cells for the SPP1-CD44 signal
included macrophage_d, macrophage_c, and occasionally CD8+ cytotoxic T cells, while
the sender cell was macrophage_d (SPP1-positive macrophage) (Figures 8B, 8C, 8E, and
8G). The receiver cells of the SPP1-CD44 signaling pathway varied among samples
(Figure 8C). Although macrophages were the primary receiver cells, B cells, CD8+ T
cells, and alveolar epithelial cells also acted as receivers in certain cases. Beyond the
SPP1-CD44 signaling pathway, other significant cell communication pathways identified
by CellChat included collagen, MHC-I, MIF, FN1, integrin β2, PECAM1, and ICAM
pathways (Figure 8D, Supplementary Figure 9). The incoming and outgoing interactions
of the SPP1-CD44 pathway by macrophage_d were relatively strong among the cell types
and other pathways in mycobacterium-infected tissues (Figures 8A, 8F, and 8G,
Supplementary Figure 9). In the SPP1-CD44 signaling pathway, SPP1 expressed by
macrophage_d communicated with macrophage_d itself, mast cells, and CD8+ T cells
(Figures 8E-G and Supplementary Figure 9).

SPP1-positive macrophages in granuloma associated with non-mycobacterial
infection

To determine whether SPP1-positive macrophages are associated with granulomatous
diseases beyond mycobacterial infections, pathological samples from granulomatous
lesions in patients with rheumatoid nodule (RN), granulomatosis with polyangiitis (GPA),
sarcoidosis (SA), Aspergillus pneumonia (AP), and aspiration pneumonia (Aspi-P) were
examined. Immunohistochemistry revealed that SPP1 was present in some foreign body
giant macrophages surrounding the granulomatous lesions across all tested samples
(Figure 9 and Supplementary Figure10). RNA-seq analysis of RNA extracted from these
granulomatous samples demonstrated elevated expression of SPP1 in the granulomatous
lesions, similar to the expression levels observed in mycobacterial infections, compared
with control lung and tonsil samples (Figures 5A-5C). Expression of CD40L(CD154) and

CD40 pair, another Ligand-Receptor pair reported as activation in granulomatous
diseases[15], did not show significant difference between granulomatous samples and
controls (Figure 5B). Furthermore, immunohistochemistry revealed CD44 expression in
SPP1-positive macrophages and neighboring cells within the lesions of GPA, SA, and RN
(Figure 9A and 9B). Immunofluorescence imaging also demonstrated that a subset of
SPP1-positive macrophages expressed CD44 (Figure 9C). These findings suggest that
SPP1-positive macrophages are broadly associated with granulomatous diseases, and that
the SPP1-CD44 interaction is observed commonly in granulomatous pathology.
193
194

DISCUSSION
A pivotal discovery in our study was the identification of multiple macrophage subtypes
within the tissues of granulomatous diseases. Based on gene expression patterns, we
identified at least four distinct macrophage cell types. Among these, SPP1-positive
macrophages predominantly infiltrated the regions surrounding NTM- and MTB-positive
granulomatous tissue. Additionally, we found that Langhans giant cells were a specialized
form of these SPP1-positive macrophages. Notably, besides these giant cells, we observed
a significant number of SPP1-positive macrophages that did not exhibit giant cell
morphology, indicating a diverse range of morphologies for these cells.
Our spatial transcriptomics data demonstrated that at least 26 types of cells infiltrated the
lungs with mycobacterial infection (Fig.1). Although the percentages of infiltrating cell
types varied among samples, macrophages were one of the major cell types in all seven
samples. Along with macrophages, T cells, B cells, plasma cells, and dendritic cells were
also observed as major cell types. These cell populations are consistent with those
reported in MTB and NTM tissues in previous studies [2-9]. Macrophages have been shown
to be key player in granuloma formation [5,7,10]. Various macrophage subtypes infiltrate
granulomas alongside T and B cells in mycobacterial infections, with M1/M2
macrophage polarization inducing NF-kappa B activation, which is associated with
granuloma formation and outcome[10].
SPP1, also known as osteopontin, is a multifunctional secreted phosphorylated
glycoprotein produced by macrophages and T cells. It is associated with bone remodeling,
T cell and B cell activation, and chemotaxis[16-20]. SPP1 expression in foreign body
macrophages, such as Langhans giant cells, has been previously reported in pathological
samples from tuberculosis and sarcoidosis [16,21-24]. Recent studies using single cell
transcriptomics and/or spatial omics assay have highlighted the potential role of SPP1-
positive macrophages in infections, inflammatory diseases, and cancer [25,26]. For instance,

in tissue samples from a patient with granulomatous slack skin, CD11c+/LYZ+
macrophages exhibited an M1-like phenotype and strongly expressed MMP9, MMP12,
CHI3L1, CHIT1, COL1A1, TIMP1, and SPP1, all of which are involved in extracellular
matrix degradation and tissue remodeling [27]. SPP1-positive macrophages have also been
detected in bronchoalveolar lavage fluid from tuberculosis patients [28]. Interestingly,
plasma osteopontin levels are higher in patients with tuberculosis and sarcoidosis
compared to healthy donors [29,30]. A study on perivascular adipose tissue in heart transplant
recipients demonstrated the presence of SPP1-positive macrophages in the perivascular
tissues, with an identified interaction between secreted SPP1 and CD44 [31]. These data
suggest that SPP1-positive macrophages secrete SPP1 and induce SPP1-CD44 signaling
in inflammation. In a *Schistosoma mansoni* egg-induced granuloma animal model,
granulomas from osteopontin-knockout animals were smaller and contained significantly
fewer macrophages, as well as fewer macrophage-derived epithelioid cells and giant cells
[32]. Thus, the presence of SPP1-positive macrophages in granulomatous tissues suggests
an important role for SPP1 in granuloma formation through its interactions.
In our study, we demonstrated that the SPP1-CD44 pathway was one of the main ligand-
receptor pairs involved in cell-cell communication in granulomatous lesions (Fig. 8A-
8D). Recent studies have shown that SPP1 is associated with cancer cell growth and
resistance to chemoradiotherapy, inducing epithelial-mesenchymal transition, autophagy,
aberrant glucose metabolism, epigenetic alterations, and reduced drug uptake through the
SPP1-CD44 signaling [33]. SPP1-positive macrophages have been identified in colorectal
cancer and squamous cell carcinoma[34,35]. In hepatocellular cancer, SPP1-positive
macrophages infiltrated hypoxic tumor regions with cancer stem cells, where the SPP1
signaling pathway was highly active in cell-cell communication with SPP1-CD44 and
SPP1-ITGA/ITGB identified as the main ligand-receptor pairs [36]. SPP1 also binds to
integrins α4β1, α9β1, and α9β4, inducing cell adhesion and migration. CD44 has also
been identified as a receptor for SPP1, and the SPP1-CD44 signaling pathway has been
reported in various cancers[37-39]. SPP1-positive macrophages have been detected in
cancers as tumor-associated macrophages, and the SPP1-CD44 pathway has been
identified as a critical mediator of communication between tumor-associated macrophage
subpopulations [40-42]. The SPP1-CD44 pathway has been linked to cancer chemoresistance
in solid tumors and to cell-to-cell communication between cancer cells and tumor-
associated macrophages [36,43,44]. The presence of SPP1-positive macrophages in cancers
correlates with worse prognosis and poor immunotherapy response, leading to reduced
survival [36,40]. Some therapeutic approaches targeting SPP1-CD44 signaling have been
reported in vitro and in vivo. In a mouse model, SPP1 was shown to induce cancer cell

chemoresistance by activating the CD44 receptor, and that treatment with anti-SPP1 and
anti-CD44 antibodies improved cancer cell sensitivity to cisplatin [45]. Importantly, the
SPP1-CD44 signaling pathway has been identified as a potential immune checkpoint
target[46,47]. CD44 expression in tumor cells was positively correlated with PD-L1
expression and the numbers of infiltrated CD68- and CD168-positive macrophages[48]. It
was also demonstrated that SPP1-CD44 immune checkpoint regulates CD8+ T cell
activation and tumor immune evasion in interferon regulatory factor 8 deficient mice [46].
Inhibition of the SPP1-CD44 pathway, therefore, may be a potential therapeutic strategy
to reduce granuloma formation in granulomatous diseases, such as mycobacterial
infections. In tuberculosis and MAC infections, granulomas encapsulate mycobacteria in
necrotic tissue to prevent their spread. Since granuloma formation is a major characteristic
of mycobacterial infections, further studies will be required to explore the inhibition of
the SPP1-CD44 pathway in macrophages.
In conclusion, our study sheds light on the intricate cellular dynamics involved in
granuloma formation in mycobacterial diseases. The prominence of SPP1-positive
macrophages, including Langhans giant cells, suggests that these cells may play a key
role in this process, opening avenues for further research and potential therapeutic
interventions.
275
MATERIALS AND METHODS
Tissue samples
Formalin-fixed paraffin embedded (FFPE) samples of resected lung tissues from patients
with active mycobacterial diseases, granulomatous diseases, or other controls were used
(S Table 1). For CosMx™ SMI analysis[14], 7 lung resections were obtained from 3 MTB
and 4 NTM patients undergoing lung resection surgery. Total 28 lung tissues from 25
patients were applied to histological analysis, immunohistochemistry and RNAseq
analysis by next generation sequencing. In addition, five intact lung tissues from patients
with cancer and four tonsillar tissues were used as controls. All MTB and NTM were
diagnosed as active disease by sputum culture and/or PCR method. The study was
approved by the Ethics Committee of National Institute of Infectious Diseases (approval
no. 1781) and National Hospital Organization Tokyo National Hospital (approval no. 408).
288
CosMx™ spatial molecular imaging (SMI)
CosMx™ Human Universal Cell Characterization RNA Panel (1000-plex; Nanostring,
Seattle, WA) was applied to 7 FFPE samples: 3 tuberculosis, 3 MAC-positive pneumonia,
and 1 MAC-negative bronchiectasis samples with 15-45 fields of view (FoVs) per sample

(216 total FoVs) (Supplementary Table 2). The mean area explored was $4.60\,mm^2$ per
sample. FOVs were placed on the tissue to match regions of interest identified by H&E
staining of an adjacent serial section. FoVs were selected from areas around granuloma
or inflammatory cell infiltration to avoid granulomatous necrotic tissues. Sample
preparation and CosMx™ SMI instrument run was performed as described previously
[14,25]. In addition to RNA readout, tissue samples were incubated with a fluorophore-
conjugated antibody cocktail against CD298/B2M (488 nm), PanCK (532 nm), CD45
(594 nm), and either CD3 or CD68 (647 nm) proteins and DAPI stain. Cell segmentation,
assigning transcripts at the cell-level, and obtaining a transcript by cell count matrix were
performed as previously described [14,25]. Following total count normalization across all
detected cells, highly variable genes (HVGs) were identified. Then, principal component
analysis (PCA) was computed on the HVGs. Finally, uniform manifold approximation
and projection (UMAP) was performed on these data[14,25].
306
307 **Cell type annotation**
308 To identify subpopulation markers, we ran findMarkers (scran package, version 1.22.1)[49],
309 blocking by sample, considering HVGs only, and testing for positive log-fold changes.
310 The 100 top ranked genes were selected and passed to SingleR (version 1.8.1)[50].
311
Co-expression and cellular communication assay
Spatial correlation analysis such as co-expression and receptor-ligand assay was
performed in the CosMx™ dataset with a R package InSituCor [51]. InSituCor was run on
normalized count data, adjusted for the effects of cell type, tissue ID and single cell counts,
to uncover spatial correlations beyond those accounted for by these variables. Cellular
communication for cell types were also analyzed with CellChat (http://www.cellchat.org/).
In CellChat, to convert spatial transcriptomics data of CosMx™, conversion.factor = 0.18
was used for setting spatial factors. In function "computeCommunProb", contact.range =
50 and scale.distance = 20 were used.
321
Bulk RNA-seq analysis of lung tissues
Total RNA was extracted from FFPE samples (Supplementary Table 1, 5 MTB, 7 NTM,
12 granulomatous diseases including 2 RN, 3 AP, 3 GPA, 3 SA, and 1 Aspi-P, 4 control
lung, and 4 tonsils) with Quick-DNA/RNA FFPE Miniprep Kit (Zymo Research).
Barcoded RNA-seq library was established with NEBNext Ultra II RNA Library Prep Kit
for Illumina (New England BioLabs, Ipswich, MA) according to the manufacturer's
instructions. Libraries were subjected to pair-end sequencing (300 cycles) on a Nextseq

1000 platform (Illumina, San Diego, CA). Quality filtering, adapter trimming, and

RNAseq analyzing was performed using CLC genomics workbench (version 23, Qiagen,

Hilden, Germany). Homo sapience refseq GRCh38.p13 was used as a reference genome

of RNA sequences. PCA, volcano blot, and box plots were performed using the R (version

4.3.1) statistics package with RStudio (version 2023.06.1+524). Gene ontology assay was

performed in biojupies (https://maayanlab.cloud/biojupies/analyze/).

Immunohistochemistry (IHC) and immunofluorescence (IF)

Immunostaining was performed in the tissue sections of mycobacterial infection and other

granulomatous diseases. Anti-SPP1 rabbit polyclonal antibody (HPA027541, Sigma-

Aldrich, St. Louis, MO) and anti-CD44 mouse monoclonal antibody (UMAB134,

OriGene Technologies, Rockville, MD) were used as primary antibodies. For the second

and third phase immunostaining reagents, a biotinylated F(ab')2 fragment of rabbit anti-

mouse immunoglobulin (Dako, Copenhagen, Denmark) and peroxidase-conjugated

streptavidin (Dako) were used. 3-3'diaminobenzidine was used as a chromogen and the

slides were counterstained with hematoxylin. Nuclear counterstaining was performed

with DAPI on IF samples. Image acquisition was performed on virtual slide system

(VS200, Evident, Tokyo, Japan).

Data availability

Bulk RNA-seq and CosMx™ SMI raw data generated in this study have been deposited

in the GEO Omnibus database under accession code GSE276060 and GSE276083,

respectively. Seurat object, Giotto object, and DEG gene list

(marker_heatmap_nb_clus_4.csv) are registered in figshare (https://figshare.com/).

Code availability

Full code for analysis of CosMx™ SMI data can be found at

https://github.com/katanoh/Lung_CosMx_analysis.

Conflict of Interest

The authors declare that they have no conflict of interest.

Acknowledgment

The authors thank Mike McKenna, Claire Williams, Patrick Danaher, Joseph M. Beechem,

Nanostring USA, Fumiyo Kurokawa, Takayuki Taniya, and Takashi Uematsu, Nanostring

Japan, and Ikuyo Yamamoto, Department of Pathology, National Institute of Infectious

Diseases, for technical supports. The authors also thank Akiko Yamashita, Yukari Nogi, and Ginko Kaneda, Department of Mycobacteriology, National Institute of Infectious Diseases, for their assistance. This study was supported in part by grants from the Japan Agency for Medical Research and Development (AMED) to HK (JP24fk0108701) and to YH (JP23fk0108608, JP24fk0108673, JP24gm1610003, JP24gm1610007, JP24wm0125007, JP24wm0225022, JP24wm0325054, and JP24fk0108701). This study was also supported in part by grants from the Japan Society for the Promotion of Science (JSPS) for International Collaborative Research to YH (JP63KK0138), and for Scientific Research (A) to YH (JP24H00331), and for Scientific Research (C) to YH (JP23K07665 and 23K07958).

References

Furin, J., Cox, H. & Pai, M. Tuberculosis. *Lancet* **393**, 1642-1656, (2019).

Hunter, R. L. Pathology of post primary tuberculosis of the lung: an illustrated critical review. *Tuberculosis (Edinb)* **91**, 497-509, (2011).

Ulrichs, T. & Kaufmann, S. H. New insights into the function of granulomas in human tuberculosis. *J Pathol* **208**, 261-269, (2006).

Ehlers, S. & Schaible, U. E. The granuloma in tuberculosis: dynamics of a host-pathogen collusion. *Front Immunol* **3**, 411, (2012).

Russell, D. G., Cardona, P. J., Kim, M. J., Allain, S. & Altare, F. Foamy macrophages and the progression of the human tuberculosis granuloma. *Nat Immunol* **10**, 943-948, (2009).

Carow, B. *et al.* Immune mapping of human tuberculosis and sarcoidosis lung granulomas. *Front Immunol* **14**, 1332733, (2023).

Marakalala, M. J. *et al.* Inflammatory signaling in human tuberculosis granulomas is spatially organized. *Nat Med* **22**, 531-538, (2016).

Park, H. E. *et al.* Modulating macrophage function to reinforce host innate resistance against Mycobacterium avium complex infection. *Front Immunol* **13**, 931876, (2022).

Lin, P. L. *et al.* Sterilization of granulomas is common in active and latent tuberculosis despite within-host variability in bacterial killing. *Nat Med* **20**, 75-79, (2014).

Marino, S. *et al.* Macrophage polarization drives granuloma outcome during Mycobacterium tuberculosis infection. *Infect Immun* **83**, 324-338, (2015).

Smith, D., Hansch, H., Bancroft, G. & Ehlers, S. T-cell-independent granuloma formation in response to Mycobacterium avium: role of tumour necrosis factor-

alpha and interferon-gamma. *Immunology* **92**, 413-421, (1997).

12   Murray, P. J. Macrophage Polarization. *Annu Rev Physiol* **79**, 541-566, (2017).

13   Yoshida, M. *et al.* Transcriptome analysis of long non-coding RNAs in
Mycobacterium avium complex-infected macrophages. *Front Immunol* **15**,
1374437, (2024).

14   He, S. *et al.* High-plex imaging of RNA and proteins at subcellular resolution in
fixed tissue by spatial molecular imaging. *Nat Biotechnol* **40**, 1794-1806, (2022).

15   Villeneuve, J. *et al.* A Role for CD154, the CD40 Ligand, in Granulomatous
Inflammation. *Mediators Inflamm* **2017**, 2982879, (2017).

16   O'Regan, A. W. *et al.* Osteopontin is associated with T cells in sarcoid granulomas
and has T cell adhesive and cytokine-like properties in vitro. *J Immunol* **162**,
1024-1031, (1999).

17   Choi, S. T. *et al.* Osteopontin might be involved in bone remodelling rather than
in inflammation in ankylosing spondylitis. *Rheumatology (Oxford)* **47**, 1775-1779,
(2008).

18   Reinholt, F. P., Hultenby, K., Oldberg, A. & Heinegard, D. Osteopontin--a
possible anchor of osteoclasts to bone. *Proc Natl Acad Sci U S A* **87**, 4473-4475,
(1990).

19   Wang, K. X. & Denhardt, D. T. Osteopontin: role in immune regulation and stress
responses. *Cytokine Growth Factor Rev* **19**, 333-345, (2008).

20   Koh, A. *et al.* Role of osteopontin in neutrophil function. *Immunology* **122**, 466-
475, (2007).

21   Carlson, I., Tognazzi, K., Manseau, E. J., Dvorak, H. F. & Brown, L. F.
Osteopontin is strongly expressed by histiocytes in granulomas of diverse etiology.
Lab Invest **77**, 103-108, (1997).

22   Tsai, A. T. *et al.* The role of osteopontin in foreign body giant cell formation.
Biomaterials **26**, 5835-5843, (2005).

23   Nau, G. J. *et al.* A chemoattractant cytokine associated with granulomas in
tuberculosis and silicosis. *Proc Natl Acad Sci U S A* **94**, 6414-6419, (1997).

24   Wang, D. *et al.* The association between osteopontin and tuberculosis: A
systematic review and meta-analysis. *PLoS One* **15**, e0242702, (2020).

25   Garrido-Trigo, A. *et al.* Macrophage and neutrophil heterogeneity at single-cell
spatial resolution in human inflammatory bowel disease. *Nat Commun* **14**, 4506,
(2023).

26   MacDonald, L. *et al.* COVID-19 and RA share an SPP1 myeloid pathway that
drives PD-L1+ neutrophils and CD14+ monocytes. *JCI Insight* **6**, (2021).

27   Feng, Y. *et al.* Spatial transcriptomics reveals heterogeneity of macrophages in the
438   tumor microenvironment of granulomatous slack skin. *J Pathol* **261**, 105-119,
439   (2023).

440   28   Yang, Q. *et al.* The interaction of macrophages and CD8 T cells in bronchoalveolar
441   lavage fluid is associated with latent tuberculosis infection. *Emerg Microbes Infect*
442   **12**, 2239940, (2023).

443   29   Koguchi, Y. *et al.* High plasma osteopontin level and its relationship with
444   interleukin-12-mediated type 1 T helper cell response in tuberculosis. *Am J Respir*
445   *Crit Care Med* **167**, 1355-1359, (2003).

446   30   Kadota, J. *et al.* High plasma concentrations of osteopontin in patients with
447   interstitial pneumonia. *Respir Med* **99**, 111-117, (2005).

448   31   Fu, M. *et al.* Single-Cell RNA Sequencing of Coronary Perivascular Adipose
449   Tissue From End-Stage Heart Failure Patients Identifies SPP1(+) Macrophage
450   Subpopulation as a Target for Alleviating Fibrosis. *Arterioscler Thromb Vasc Biol*
451   **43**, 2143-2164, (2023).

452   32   O'Regan, A. W. *et al.* Abnormal pulmonary granuloma formation in osteopontin-
453   deficient mice. *Am J Respir Crit Care Med* **164**, 2243-2247, (2001).

454   33   Kretschmer, C. *et al.* Identification of early molecular markers for breast cancer.
455   *Mol Cancer* **10**, 15, (2011).

456   34   Bill, R. *et al.* CXCL9:SPP1 macrophage polarity identifies a network of cellular
457   programs that control human cancers. *Science* **381**, 515-524, (2023).

458   35   Qi, J. *et al.* Single-cell and spatial analysis reveal interaction of FAP(+) fibroblasts
459   and SPP1(+) macrophages in colorectal cancer. *Nat Commun* **13**, 1742, (2022).

460   36   Fan, G. *et al.* Single-cell and spatial analyses revealed the co-location of cancer
461   stem cells and SPP1+ macrophage in hypoxic region that determines the poor
462   prognosis in hepatocellular carcinoma. *NPJ Precis Oncol* **8**, 75, (2024).

463   37   Weber, G. F., Ashkar, S., Glimcher, M. J. & Cantor, H. Receptor-ligand interaction
464   between CD44 and osteopontin (Eta-1). *Science* **271**, 509-512, (1996).

465   38   Liu, L. *et al.* Construction of TME and Identification of crosstalk between
466   malignant cells and macrophages by SPP1 in hepatocellular carcinoma. *Cancer*
467   *Immunol Immunother* **71**, 121-136, (2022).

468   39   Wang, C. *et al.* The heterogeneous immune landscape between lung
469   adenocarcinoma and squamous carcinoma revealed by single-cell RNA
470   sequencing. *Signal Transduct Target Ther* **7**, 289, (2022).

471   40   Xu, L. *et al.* Tumor-associated macrophage subtypes on cancer immunity along
472   with prognostic analysis and SPP1-mediated interactions between tumor cells and

macrophages. *PLoS Genet* **20**, e1011235, (2024).

41   He, H. *et al.* Multi-dimensional single-cell characterization revealed suppressive
immune microenvironment in AFP-positive hepatocellular carcinoma. *Cell*
Discov **9**, 60, (2023).

42   He, C. *et al.* Single-Cell Transcriptomic Analysis Revealed a Critical Role of
SPP1/CD44-Mediated Crosstalk Between Macrophages and Cancer Cells in
Glioma. *Front Cell Dev Biol* **9**, 779319, (2021).

43   Xie, W. *et al.* Multi-Transcriptomic Analysis Reveals the Heterogeneity and
Tumor-Promoting Role of SPP1/CD44-Mediated Intratumoral Crosstalk in
Gastric Cancer. *Cancers (Basel)* **15**, (2022).

44   Nallasamy, P. *et al.* Pancreatic Tumor Microenvironment Factor Promotes Cancer
Stemness via SPP1-CD44 Axis. *Gastroenterology* **161**, 1998-2013 e1997, (2021).

45   Qian, J. *et al.* Cancer-associated mesothelial cells promote ovarian cancer
chemoresistance through paracrine osteopontin signaling. *J Clin Invest* **131**,
(2021).

46   Klement, J. D. *et al.* An osteopontin/CD44 immune checkpoint controls CD8+ T
cell activation and tumor immune evasion. *J Clin Invest* **128**, 5549-5560, (2018).

47   Lu, C. *et al.* WDR5-H3K4me3 epigenetic axis regulates OPN expression to
compensate PD-L1 function to promote pancreatic cancer immune escape. *J*
Immunother Cancer **9**, (2021).

48   Liu, S. *et al.* CD44 is a potential immunotherapeutic target and affects
macrophage infiltration leading to poor prognosis. *Sci Rep* **13**, 9657, (2023).

49   Haghverdi, L., Lun, A. T. L., Morgan, M. D. & Marioni, J. C. Batch effects in
single-cell RNA-sequencing data are corrected by matching mutual nearest
neighbors. *Nat Biotechnol* **36**, 421-427, (2018).

50   Aran, D. *et al.* Reference-based analysis of lung single-cell sequencing reveals a
transitional profibrotic macrophage. *Nat Immunol* **20**, 163-172, (2019).

51   Danaher, P. *et al.* Single cell spatial transcriptomic profiling of childhood-onset
lupus nephritis reveals complex interactions between kidney stroma and
infiltrating immune cells. *bioRxiv*, (2023).

 Figure legends

Figure 1. Single cell analysis of CosMx™ in mycobacterium-infected tissues. A.

Overview of the study design for CosMx™ SMI, RNA-seq, and histological analysis.

Lung samples including MTB, NTM, granulomatous diseases and controls were

processed using RNA-seq (n = 37 samples) and CosMx™ SMI (n = 7 samples). This

figure was created with BioRender.com. B. Heatmap of gene expression and cell types

based on single cell data from CosMx™. C. Dot plots showing expression of cell-typing

markers in the indicated cell clusters. The dot size represents the percentage of cells

expressing the genes in each cluster. The average expression intensity of the markers is

shown using blue/red color. D. UMAP plots of 300,841 cells from lung tissues with 4

NTM or 3 MTB patients. Each cluster is shown in a different color. E. Stacked bar graph

showing the distribution of cell types across 7 tissue samples.

Figure 2. Subtyping of macrophages. A. Violin plot showing macrophage-associated

protein expression in macrophages and dendritic cells. B. Dot plot illustrating

macrophage-associated protein expression specially in macrophages. C. UMAP plot

depicting four macrophage subtypes. D. UMAP plot displaying the expression of

macrophage-associated proteins.

Figure 3. Spatial images of SPP1-positive macrophages (macrophage_d) in MAC-

positive lung tissues (Case NTM2). A. HE staining (left panel), immunofluorescence

(second panel), and CosMx™ spatial images (right two panels) of macrophage_d. Colors

representing different cell types are shown in the right panel. In CosMx™ images, SPP1

and MARCO signals are indicated by green and red dots, respectively. B. Enlarged images

of the boxed regions in A. C. Immunohistochemistry for SPP1 in the same tissue near the

granuloma.

Figure 4. CosMx™ spatial images of SPP1 and MARCO expression in macrophage_d in

MTB tissue (Case MTB1). A. Macrophage_d in the interstitial region expressed SPP1

and MARCO. B. Enlarged images of the boxed regions in A. C. Immunohistochemistry

for SPP1 in the same tissue near the granuloma.

Figure 5. RNAseq Results for granulomatous diseases and mycobacterial infection. A.

Principal component analysis (PCA) of RNAseq data for 37 samples. B. Volcano plot

comparing granulomatous diseases (granulomatous diseases and mycobacterial

infections) versus control groups. The expression of SPP1, FN1, and MARCO is

significantly higher in the granulomatous disease group. C. Box plot showing SPP1, FN1,
and MARCO expression in control lung (CTL-Lung, n=5), granulomatous diseases
(Granul, n=12), and the mycobacterial infection group (MTB+NTM, Mycob, n=16).
Transcripts per million (TPM) values for each gene are plotted. P values are displayed on
comparison lines. An asterisk (*) indicates a significant difference. D. Gene Ontology
analysis comparing mycobacterial infection tissues to control lungs, with upregulated
pathways listed in the left panel.

Figure 6. Coexpression and ligand-receptor assay in InSituCor. A. Heatmap of
coexpressed transcripts across different cell types. Coexpression of FN1, SPP1, and
MARCO was specifically observed in macrophage_d (highlighted by the broken red line).
B. Conditional correlation of ligand-receptor pairs in the ligand-receptor assay. The SPP1-
CD44 and FN1-CD44 pairs show strong conditional correlations. C. Correlation network
of SPP1 and CD44-associated transcripts. D. Spatial images showing SPP1-CD44
correlations in MTB and NTM tissues. Yellow indicates high correlation in the images
(left: MTB1, right: NTM2). E. Spatial images from CosMx™ analysis of MTB and NTM
samples. High expression of SPP1, CD44, and both are indicated by yellow, blue, and red
dots, respectively.

Figure 7. Expression of SPP1 and CD44 in macrophages around granuloma tissue with
mycobacterial infection. A. Dot plot showing the expression of SPP1, MARCO, FN1, and
CD44 across all cell types. B. Immunohistochemistry for SPP1 and CD44 in tissues from
patients with MTB (upper panels) and NTM (lower panels) infections. C.
Immunofluorescence and CosMx™ spatial images for SPP1 and CD44. SPP1 was
expressed in the cytoplasm of multinuclear giant cells (immunofluorescence assay, left
panels). CD44 expression was observed on the membrane of multinucleated giant cells
and their neighboring cells. CosMx™ spatial images (right panels) showed that
macrophage_d expresses both SPP1 and CD44. Colors representing different cell types
are shown in the bottom right panel.

Figure 8. Analysis of cellular communication among cell types using CellChat. A and B.
Circular plots showing interactions among cell types in all pathways (A) and specifically
in the SPP1 signaling pathway (B) across all seven cases analyzed with CosMx™. C.
Circular plots depicting interactions among cell types in the SPP1 signaling pathway in
each individual case. D. Histogram showing the relative contribution of each ligand-
receptor (L-R) pair across all seven cases analyzed with CosMx™. Only significant L-R

pairs are shown, with SPP1 interacting exclusively with CD44. E. Heatmap illustrating
the signal intensity of the sender and receiver in the SPP1 signaling pathway across all
samples. F. Dot plots showing the outgoing communication patterns of secreting cells.
Significant outgoing signals in the identified signaling pathways are represented by dots.
G. Dot plots showing the incoming communication patterns of secreting cells. Significant
incoming signals are represented by dots.
583
584   **Figure 9**. SPP1 and CD44 expression in tissues from patients with granulomatous
585   diseases. A. HE staining. B. Immunohistochemistry for SPP1. C. Immunofluorescence of
586   SPP1 and CD44.   Tissues from patients with sarcoidosis (left), granulomatosis with
587   polyangiitis (GPA, middle), and rheumatoid nodule (RN, right) were investigated.
588
589

Fig.1

Fig.2

Fig.3

A

PanCk
CD45
CD3
DAPI

SPP1
MARCO

B

C
SPP1
SPP1
50 μm
200 μm

CD4+.T.cell
B.cell.linage
fibroblast_b
neutrophil
plasmacytoid.dendritic.cell
lymph.vessel.cell
alveolar.epithelial.cell.type.2
blood.vessel.cell
alveolar.epithelial.cell.type.1
natural.killer.cell
fibroblast_c
B.cell
mast.cell
dendritic.cell.type.2
plasma.cell
macrophage_b
regulatory.T.cell
fibroblast_a
macrophage_d
activated.dendritic.cell
CD8+.cytotoxic.T.cell
macrophage_c
macrophage_a
dendritic.cell.type.1
ciliated.cell
muscle.cell

Fig.4

Fig.5

Fig.6

A

B

C

D   MTB   NTM

SPP1-CD44
correlation

E

high SPP1
high CD44
high both

Fig.7

Fig.8

Fig.9

Supplementary Data

Identification of SPP1-positive macrophages by single-cell spatial analysis in human lung tissues with mycobacterial infection

Harutaka Katano, Akira Hebisawa, Yuko Sato, Yoshihiko Hoshino

Supplementary Table 1: patient and sample list. AP: Aspergillus pneumonia, Aspi P: aspiration pneumonia, BCG: Bacille Calmette-Guerin, CTL: control, FFPE: formalin-fixed paraffin-embedded sample, GPA: granulomatosis with polyangiitis, IHC: immunohistochemistry, MAC: mycobacterium avium intracellulae complex, MTB: mycobacterium tuberculosis, NTM: non tuberculosis mycobacterium infection, ND: no data, RN: rheumatoid nodule, SA: sarcoidosis.

ID	Age	Gender	Group	Clinical diagnosis	BCG-IHC in FFPE	Real-time PCR in FFPE	CosMx
MTB1	49	M	MTB	MTB	+	MTB	MTB1
MTB2	70	M	MTB	MTB	+	MTB	MTB2
MTB3	64	M	MTB	MTB	+	MTB	MTB3
MTB4	38	M	MTB	MTB	-	MTB	
MTB5	49	M	MTB	MTB	-	-	
MTB6	49	M	MTB	MTB	-	MTB	
NTM1	71	M	NTM	NTM	+	MAC	NTM1
NTM2	56	F	NTM	NTM	+	MAC	NTM2
NTM3	63	F	NTM	NTM	+	MAC	NTM3
NTM4	51	F	NTM	NTM	-	-	NTM4
NTM5	51	F	NTM	NTM	-	-	
NTM6	51	F	NTM	NTM	+	M.Kansasii	
NTM7	66	F	NTM	NTM	+	MAC	
NTM8	66	F	NTM	NTM	+	MAC	
NTM9	73	M	NTM	NTM	-	MAC	
NTM10	68	F	NTM	NTM	-	-	
Ctl-Lung1	62	F	CTL-Lung	Lung cancer (control)	-	-	
Ctl-Lung2	63	M	CTL-Lung	Lung cancer (control)	-	-	
Ctl-Lung3	70	M	CTL-Lung	Lung cancer (control)	-	-	
Ctl-Lung4	80	M	CTL-Lung	Lung cancer (control)	-	-	
Ctl-Lung5	54	M	CTL-Lung	Lung cancer (control)	-	-	
Ctl-Tonsil1	ND	ND	CTL-Tonsil	tonsillitis (control)	-	-	
Ctl-Tonsil2	ND	ND	CTL-Tonsil	tonsillitis (control)	-	-	
Ctl-Tonsil3	ND	ND	CTL-Tonsil	tonsillitis (control)	-	-	
Ctl-Tonsil4	ND	ND	CTL-Tonsil	tonsillitis (control)	-	-	
Granulo1	70	M	Granulo	RN	-	-	
Granulo2	77	M	Granulo	RN	-	-	
Granulo3	50	M	Granulo	AP	-	-	
Granulo4	52	M	Granulo	AP	-	-	
Granulo5	69	F	Granulo	AP	-	-	
Granulo6	58	F	Granulo	GPA	-	-	
Granulo7	71	M	Granulo	GPA	-	-	
Granulo8	77	F	Granulo	GPA	-	-	
Granulo9	56	F	Granulo	SA	-	-	
Granulo10	37	F	Granulo	SA	-	-	
Granulo11	37	M	Granulo	SA	-	-	
Granulo12	62	M	Granulo	Aspi P	-	-	

Supplementary Table 2: Summary of CosMx assay on 7 slides of MTB and NTM samples. FOV: focus of view.

tissue	MTB1	MTB2	MTB3	NTM1	NTM2	NTM3	NTM4	Total	Mean
Number_of_FOVs	21	47	45	21	15	22	45	216	30.9
Total_Tissue_Area_mm2	8.33	4.65	4.16	3.63	2.09	4.28	5.08	32.2	4.6
Number_of_Cells_in_Final_Analysis	70649	40962	34799	31874	23692	51176	47689	300841	42977
Total_Transcripts_Assigned_to_Cells	8768373	7013069	4749647	4267383	1882347	5634413	7876583	40191815	5741688
Mean_Transcripts_per_Cell	124.1	171.2	136.5	133.9	79.5	110.1	165.2	920.5	131.5
Maximum_Transcripts_per_Cell	2226	2079	1697	2339	728	1362	2944	13375.0	1910.7
Mean_Unique_Genes_Per_Cell	70.14	65.82	58.73	81.5	54.69	69.42	79.18	479.5	68.5
Mean_Transcripts_per_um2	0.0357	0.0222	0.0178	0.0458	0.0323	0.0466	0.0228	0.2	0.0

Supplementary Table 3: Cell population in CosMx assay on 7 slides of MTB and NTM samples.

Cell types	MTB1	MTB2	MTB3	NTM1	NTM2	NTM3	NTM4	overall_percentage
activated.dendritic.cell	0.71	0.15	0.25	0.36	1.75	0.34	1.46	0.72
alveolar.epithelial.cell.type.1	0.94	0.53	1.36	1.4	5.18	1.05	14.12	2.76
alveolar.epithelial.cell.type.2	6.48	8.67	4.26	9.7	5.88	7.49	3.95	6.72
B.cell	3.97	10.71	6.36	19.78	9.82	2.82	6.77	7.87
B.cell.lineage	5.63	17.05	16.73	5.54	11.32	14.11	10.89	11.19
blood.vessel.cell	3.38	2	1.62	6.5	6.13	4.81	5.28	4.16
CD4+.T.cell	3.98	5.24	5.39	6.38	1.4	5.09	0.97	4.07
CD8+.cytotoxic.T.cell	3.24	1.42	1.76	2.75	7.23	1.2	3.16	3.12
ciliated.cell	0.88	0.1	0.24	0.84	0.39	0.22	4.96	0.83
dendritic.cell.type.1	1.06	2.68	1.86	1.49	1.1	0.69	2.05	1.45
dendritic.cell.type.2	1.71	3.13	4.55	5.09	1.64	4.15	1.5	2.95
fibroblast_a	1.07	4.53	4.58	0.61	4.46	4.59	5.24	3.36
fibroblast_b	2.06	11.41	17.16	17.7	16.12	19.73	10.74	12.61
fibroblast_c	21.16	1.29	1.69	0.37	0.43	2.82	0.84	5.97
lymph.vessel.cell	1.48	1.81	1.21	0.95	0.89	0.78	0.81	1.18
macrophage_a	15.32	0.16	0.24	0.14	0.2	1.17	0.84	3.95
macrophage_b	12.67	0.11	0.3	0.53	0.63	0.89	0.35	3.36
macrophage_c	0.86	6.17	3.45	1.53	7.26	5.11	9.36	4.39
macrophage_d	0.25	2.36	6.79	7.16	5.03	1.16	2.78	3.18
mast.cell	0.95	0.42	0.28	1.33	0.97	0.95	0.92	0.84
muscle.cell	0.31	1.14	0.64	2.04	2.67	0.45	2.92	1.27
natural.killer.cell	0.99	3.42	6.39	1.16	1.91	7.19	3.5	3.3
neutrophil	1.17	1.39	2.53	2.44	0.38	6.36	1.11	2.18
plasma.cell	1.51	11.47	8.01	1.36	3.1	4.33	2.21	4.38
plasmacytoid.dendritic.cell	1.61	2.38	1.94	2.08	1.59	2.08	1.3	1.85
regulatory.T.cell	6.61	0.24	0.4	0.75	2.51	0.41	1.96	2.36

Supplementary Fig.1a. CosMx FOV placements.

Supplementary Fig.1b. CosMx FOV placements.

Supplementary Fig. 2: Immunohistochemistry for BCG. Blue and red boxes (right panels) show the enlarged images of left panels.

Supplementary Fig. 3: Each cell type in UMAP.

Supplementary Fig. 4: UMAP and population of each cell type. A. UMAP of each tissue. B. Populations of cell types in each tissue.

Supplementary Fig. 5: Distribution of cell type in FOVs of each slide

Supplementary Fig. 6a. CosMx spatial images of SPP1 and MARCO expression in macrophage_d in MAC+ tissue (NTM2 FOV9). A. Macrophage_d in the interstitial region expressed SPP1 and MARCO. B. Enlarged images of A. Colors indicating cell types are shown in the bottom panel.

Supplementary Fig. 6b. CosMx spatial images of SPP1 and MARCO expression in macrophage_d in MAC+ tissue (NTM2 FOV8). SPP1-positive macrophages were found in alveolar space. A. Macrophage_d in the alveolar space expressed SPP1 and MARCO. B. Enlarged images of A. Colors indicating cell types are shown in the bottom panel.

Supplementary Fig. 6c. CosMx spatial images of SPP1 and MARCO expression in macrophage_d in MTB+ tissue (MTB1, FOV24). A. Macrophage_d in the interstitial region expressed SPP1 and MARCO. B. Enlarged images of A. Colors indicating cell types are shown in the bottom panel.

Supplementary Fig. 7. CosMx spatial images of SPP1 and CD44 expression in MTB and NTM tissues. Colors indicating cell types are shown in the bottom panel.

Supplementary Fig. 8: InSituCor analysis in MTB and NTM samples. A. Example of a cell's nearestneighbors, used to define its "environment expression" profile. B. Subset of the environment expression matrix. C. Raw correlation matrix of the environment expression matrix showing near-ubiquitous correlations (subset of 300 random genes).D. Correlation matrix of the environment expression matrix (B) conditional on the confounding matrix, over the same subset of genes as (C). E. Raw vs. conditional correlation of environment gene expression. Selected pairs of marker genes are highlighted. F. Network representation of correlation between all genes in all modules. Genes with correlation > 0.2 are connected. The module from Fig.7A is highlighted. G. Correlation structure of 51 ligands assigned to modules. Edges show conditional correlations > 0.1,and color shows module membership.

Supplementary Fig. 9: Outgoing and incoming communication patterns of secreting cells by cellchat analysis in MTB and NTM samples. A. River plot of cell (left panel) and cell-communication (right panel) pattern for outgoing signal pathways. B. Heatmaps of cell (upper panel) and cell-communication (lower panel) pattern for outgoing signal pathways. C. River plot of cell (left panel) and cell-communication (right panel) pattern for incoming signal pathways. D. Heatmaps of cell (upper panel) and cell-communication (lower panel) pattern for incoming signal pathways. E. Dot plots of incoming and outgoing interaction strength of each cell-cell communications in all ligand-receptor interactions.

Supplementary Fig. 10: SPP1 expression in a tissue from a patient with Aspergillus pneumonia. A. HE. B. Immunohistochemistry for SPP1.