The SET1/COMPASS subunit RBBP5 orchestrates epigenetic control of global proteostasis and the 12h oscillator to safeguard metabolic and cellular homeostasis

Syeda Kubra[1, #], Michelle Sun[1, #], William Dion[1], Ahmet Catak[1], Hannah Luong[1], Haokun Wang[1], Yinghong Pan[2, ¥], Jia-Jun Liu[3,4], Aishwarya Ponna[1], Lijun Liu [3,5], Ian Sipula[6], Jian-Hua Luo [3,5], Michael J. Jurczak[6,7], Silvia Liu[3,4,8], and Bokai Zhu[1,3,6] *

[1] Aging Institute of UPMC, University of Pittsburgh School of Medicine, Pittsburgh, PA, U.S.A.

[2] UPMC Genome Center, UPMC, Pittsburgh, PA, U.S.A.

[3] Pittsburgh Liver Research Center, University of Pittsburgh, Pittsburgh, PA, U.S.A.

[4] Organ Pathobiology and Therapeutics Institute, University of Pittsburgh School of Medicine, Pittsburgh, PA, U.S.A.

[5] Department of Pathology, University of Pittsburgh School of Medicine, Pittsburgh, PA, U.S.A.

[6] Division of Endocrinology and Metabolism, Department of Medicine, University of Pittsburgh School of Medicine, Pittsburgh, PA, U.S.A.

[7] Center for Metabolism and Mitochondrial Medicine, University of Pittsburgh School of Medicine, Pittsburgh, PA, U.S.A.

[8] Department of Pharmacology and Chemical Biology, University of Pittsburgh School of Medicine, Pittsburgh, PA, U.S.A.

# These authors contribute equally.

¥ Present address: UPMC Magee-Womens Hospital, Pittsburgh, PA, U.S.A.

* Corresponding author: Bokai Zhu: bzhu@pitt.edu

One sentence summary:

RBBP5 regulates global mammalian proteostasis.

Abstract:

Proteostasis is vital for cellular health, with disruptions leading to aging, neurodegeneration and metabolic disorders. Traditionally, proteotoxic stress responses were studied as acute reactions to various noxious factors; however, recent evidence reveals that many proteostasis genes exhibit ~12h ultradian rhythms under physiological conditions in mammals, driven by an XBP1s-dependent 12h oscillator. By examining the chromatin landscape of this oscillator, we identified RBBP5 as an essential epigenetic regulator of global proteostasis dynamics. Mechanistically, as the core subunit of the SET1/COMPASS complex, RBBP5 co-activates XBP1s to facilitate dynamic proteostasis gene expression by marking promoter-proximal H3K4me3, which further recruits the Integrator Complex and SWI/SNF chromatin remodelers. Functionally, RBBP5 is indispensable for regulating both the 12h oscillator and acute transcriptional response to various proteotoxic stresses, including ER stress and nutrient deprivation. RBBP5 ablation causes increased susceptibility to proteotoxic stress, chronic inflammation, and hepatic steatosis in mice, along with impaired autophagy and reduced cell survival *in vitro*. In humans, lower *RBBP5* expression is associated with reduced adaptive stress-response gene expression and hepatic steatosis. Our findings not only highlight a previously unrecognized epigenetic timing mechanism distinct from circadian regulation but also establish RBBP5 as a central regulator of proteostasis, essential for cellular resilience and organismal health.

Introduction:

Proteostasis, the maintenance of proper protein folding, trafficking, and turnover, is a major challenge for the cell (1). Much of the cellular energy expenditure is dedicated to maintaining a healthy proteome, as millions of molecules are produced every single minute (2). Challenges to proteostasis, exemplified by the accumulation of unfolded or misfolded proteins, can lead to a range of cellular dysfunctions and pathological conditions, such as aging, neurodegenerative diseases, cancer, and metabolic disorders (1,3-6). To cope with proteotoxic stress, organisms have developed a wide range of stress response mechanisms, including the heat shock response, the unfolded protein response (UPR), and the more generalized integrated stress response (ISR) (7-9). Although the detailed signaling molecules vary, these pathways generally involve transmitting the information of cellular stress to the nucleus by activating various transcription factors (TF) containing the basic leucine zipper domain (bZIP), such as spliced form of XBP1 (XBP1s), ATF4 and ATF6 and heat shock factor 1 (HSF1). After binding the stress response element in the promoters/enhancers, these TFs in turn upregulate a plethora of genes aimed to restore proteostasis (10,11).

While proteotoxic stress responses were traditionally viewed and studied as distinct acute responses to different "noxious" factors, recent evidence by our group indicates that at the physiological level (and in the absence of evident extrinsic stress), the expression of hundreds of proteotoxic stress response regulatory and output genes exhibit cell-autonomous ultradian rhythms cycling with a ~12h period in multiple mammalian cell-types and tissues, including humans (12-21). These proteostasis genes encompass both output genes involved in protein processing in the ER/Golgi, redox regulation, protein folding, ER-associated protein degradation (ERAD), autophagy, and upstream regulatory molecules and TFs like *Xbp1*, *Atf4*, *Atf6*, *Ddit3*, *Perk* and *Ire1α* (12,22). Many of these ~12h ultradian rhythms are established by an XBP1s-dependent 12h ultradian oscillator separate from both the ~24h circadian clock and the cell cycle (21,23,24). It is hypothesized that the mammalian 12h oscillator may have evolved from the circatidal clock of marine animals, and later co-opted to adapt to the ~12h cycle of metabolic stress peaking at transition times at dawn and dusk in terrestrial animals (13,16,21,25). The 12h oscillator is well-studied in the liver, and mice with liver-specific 12h oscillator ablation exhibited accelerated liver aging and steatosis (19), manifested with impaired proteostasis, lipid metabolism, and

mitochondria dysfunction (19,20,26). Since the 12h oscillator integrates multiple proteotoxic
stress signaling, it provides us with a unique opportunity to study proteostasis using a holistic
approach: rather than studying individual stress response separately, by investigating the 12h
oscillator we could gain new mechanistic insights for universal proteostasis regulation.

While significant work has been done on upstream proteotoxic stress sensing and protein-
folding mechanisms in the ER and cytosol, there remains a lack of fundamental understanding of
the transcriptional regulation of proteostasis in general, particularly concerning the temporal
epigenome dynamics, chromatin landscapes, and co-regulatory networks underlying global
proteostasis control. In this study, we seek to address an important unanswered question: is there
a designated epigenetic regulator that orchestrates global proteostasis dynamics? By analyzing
the chromatin landscape of the murine 12h hepatic oscillator, we identified RBBP5—an essential
subunit of the Complex Proteins Associated with Set1 (COMPASS) complex responsible for
depositing H3K4me3 (27) —as a pivotal epigenetic regulator that governs global proteostasis
dynamics. By contrast, histone acetyltransferase (HAT) and associated H3K9Ac are dispensable
for dynamic proteostasis gene expression. RBBP5 plays a critical role in regulating both the 12h
oscillator and transcriptional response to a plethora of acute proteostasis stress, functioning as a
co-activator for XBP1s, to mark H3K4me3 at the promoter of proteostasis genes. Proximity
labeling of H3K4me3 further reveals the chromatin-associated proteomic architecture during
proteostatic stress, including subunits of the Integrator complex and SWI/SNF chromatin
remodelers. Loss of RBBP5 leads to heightened susceptibility to proteotoxic stress, driving
chronic inflammation and hepatic steatosis in mice, while also impairing autophagy and cell
survival *in vitro*. In humans, reduced RBBP5 expression is linked to a weakened adaptive stress
response and the onset of hepatic steatosis. Our findings not only highlight a previously
unrecognized epigenetic timing mechanism distinct from circadian regulation but also establish
RBBP5 as a central regulator of proteostasis, essential for cellular resilience and organismal
health.

Materials and Methods:

Plasmids

pCDH-EF1a-mCherry-EGFP-LC3B was a gift from Sang-Hun Lee (Addgene plasmid # 170446;
http://n2t.net/addgene:170446; RRID: Addgene_170446) (28). pLenti-X1-hygro-mCherry-RAMP4
was a gift from Jacob Corn (Addgene plasmid # 118391; http://n2t.net/addgene:118391;
RRID:Addgene_118391) (29).

The generation of *Rbbp5* (flox/flox) mice

The Rbbp5-flox allele was generated by the Mouse Embryo Services Core (University of
Pittsburgh, Department of Immunology). The targeting strategy is based on the Easi-CRISPR
method (30). In brief, fertilized embryos from C57BL/6J background, produced by natural mating,
were microinjected in the pronuclei with a mixture of 0.67 μM EnGen Cas9 protein (New England
Biolabs, Cat.No. M0646T), two Cas9 guides RNA: Rbbp5-695rev and Rbbp5-951forw (21.23
ng/μl each) and a long single stranded oligonucleotides Rbbp5-flox-ssODN (5 ng/μl). The long
single stranded oligonucleotide encoding the donor sequence was synthesized by Integrated DNA
Technologies (IDT).

To produce the sgRNA, a double strand linear DNA template is created by annealing of the
following target specific oligonucleotides with a common primer (gRNA-Scaffold-R: 5'-
AAAAAAGCACCGACTCGGTGCCACTTTTTCAAGTTGATAACGGACTAGCCTTATTTTAACTTG
CTATTTCTAGCTCTAAAAC-3') containing the full tracrRNA sequence and then PCR amplified as

previously described (31). The sgRNA was synthesized using HiScribe T7 Quick High Yield RNA
Synthesis Kit (New England Biolabs, Cat. No. E2050S).

T7-Rbbp5-695rev sequence:

TAATACGACTCACTATAGGAAGCATTCTAAGATTAAACCGTTTTAGAGCTAGAAATAGCA

T7-Rbbp5-951forw sequence:

TAATACGACTCACTATAGGAGTAACAGATAGTATTCCGAGTTTTAGAGCTAGAAATAGCA

Rbbp5-flox-ssODN sequence:

gcagttaataaacttgaagtgtttatatagaaaatacatcaggatggcaaccaaacccttagatacctgagatttattctcttatcccag
gtATAACTTCGTATAGCATACATTATACGAAGTTATgaattcttaatcttagaatgcttagtctgtttcatgctgaagata
agttgtcagaactcacattgttactgattttcgttcagtgactcgtctgttttgtcataggaaaattggagtgcatttgcaccagacttcaaag
agttggatgaaaatgtagaatatgaggaaagagaatcagaatttgatattgaggatgaagataagagtgagcctgagcaaacaggt
gatgcctctcagtaacagatagtattgaattcATAACTTCGTATAGCATACATTATACGAAGTTATccgaagggaag
gactgtcttactcaatgctctttgaattagtgaatgcatttatttactttgtgttgtaactgtgctcaaagcttcatctgccacgtttccctcacttgt
acatagttttcacagtgcttgaagagtcttactg

The injected zygotes were cultured overnight, the next day the embryos that developed to the
2-cell stage were transferred to the oviducts of pseudo pregnant CD1 female surrogates.
Following optimization of the cycling conditions, the potential founder mice were genotype with
the following primers pair, using Q5 High-Fidelity DNA Polymerase (NEW ENGLAND BIOLABS,
Cat. no. M0491L). The body weight of male adult mice was measured weekly.

Rbbp5 genotyping forward primer: AGTTCAGAGCTGGTTTCCAAC

Rbbp5 genotyping reverse primer: AGGTAATGCCCATGTGAGCC

Biological rhythm study in mice

All mice used for biological rhythm study are in C57BL/6J background, male and between 3 and
4 months of age. Wild-type C57BL/6J mice (n=12), *Rbbp5* (flox/flox); Alb-CRE (-/-) (n=48) and
Rbbp5 (flox/flox); Alb-CRE (+/-) mice (n=48) [*Rbbp5* (flox/flox); Alb-CRE (-/-) and *Rbbp5* (flox/flox);
Alb-CRE (+/-) mice were littermates] were first entrained under LD12:12 conditions for 2 weeks
before transferred to constant darkness for 24hrs. Mice were then sacrificed via cervical
dislocation at a 2h interval [for *Rbbp5* (flox/flox); Alb-CRE (-/-) and *Rbbp5* (flox/flox); Alb-CRE (+/-)
mice] or 4h interval (for wild-type C57BL/6J mice) for a total of 48 hours under constant darkness.
Mice were fed *ad libitum* during the entire experiment. The animal studies were carried out in
accordance with the National Institutes of Health guidelines and were granted formal approval by
the University of Pittsburgh's Institutional Animal Care and Use Committee (approval number
IS00013119 and IS00013119).

Food intake and locomotor activity monitoring

Promethion Multi-plexed Metabolic Cage System was used for real-time measuring of food intake
and locomotor activity. Male *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice between 3 and 4 months of age were
acclimated to the chambers and ad libitum food intake was monitored for 48 hours under LD12:12
followed by 48 hours of constant darkness. n=6 for *Rbbp5* ^{Flox} and n=8 for *Rbbp5* ^{LKO} mice for
LD12:12 study. n=5 for *Rbbp5* ^{Flox} and n=3 for *Rbbp5* ^{LKO} mice for DD study.

ER stress induction study in mice

For acute ER stress induction experiment, male littermates *Rbbp5* (flox/flox); Alb-CRE (-/-) and
Rbbp5 (flox/flox); Alb-CRE (+/-) mice between 5 and 7 months of age were randomly divided into
two groups, and intraperitoneally injected with 0.05mg/kg body weight of tunicamycin dissolved
in 500ul 3% DMSO in PBS or vehicle control (3% DMSO in PBS), respectively. Mice were injected
with Tu between 9~10am in the morning and 8 hours later dissected for transcriptomic and
western blot analysis for different tissues. The sample size is n=3 for *Rbbp5* ^{Flox} DMSO, n=3 for
Rbbp5 ^{Flox} Tu, n=4 for *Rbbp5* ^{LKO} DMSO, n=4 for *Rbbp5* ^{LKO} Tu.

For experiments involving RBBP5 liver-overexpressing mice, AAV8-TBG-EGFP and AAV8-
TGB-mRBBP5 viruses were customarily designed and produced from Vector Biolab (Malvern,
PA). Male wildtype C57BL/6J mice between 2 and 3 months of age were tail-vein injected with
0.5x10^11 genome copies of AAV8-TBG-EGFP or AAV8-TGB-mRBBP5 virus diluted in 100 µl of
PBS. 30 days later, mice injected with either viruses were randomly divided into two cohorts and
intraperitoneally injected with 0.05mg/kg body weight of tunicamycin dissolved in 500ul 3% DMSO
in PBS or vehicle control (3% DMSO in PBS). Mice were injected with Tu between 9~10am in the
morning and 8 hours later dissected for transcriptomic and western blot analysis for different
tissues. The sample size is n=3 for AAV8-TBG-EGFP DMSO, n=4 for AAV8-TBG-EGFP Tu, n=4
for AAV8-TGB-mRBBP5 DMSO and n=4 for AAV8-TGB-mRBBP5 Tu.

For chronic ER stress induction experiment, male littermates *Rbbp5* (flox/flox); Alb-CRE (-/-)
and *Rbbp5* (flox/flox); Alb-CRE (+/-) mice between 5 and 7 months of age were intraperitoneally
injected with 0.025mg/kg body weight of tunicamycin dissolved in 500ul 3% DMSO in PBS or
vehicle control (3% DMSO in PBS) daily for six consecutive days. Mice were injected with Tu
between 9~10am in the morning and 24 hours after the last injection dissected for transcriptomic,
western blot analysis, histology for liver tissues. The sample size is n=4 for *Rbbp5* ^{Flox} DMSO, n=4
for *Rbbp5* ^{Flox} Tu, n=2 for *Rbbp5* ^{LKO} DMSO, n=3 for *Rbbp5* ^{LKO} Tu.

Oil Red O Staining

Frozen sections were rinsed with PBS and then fixed with 10% neutral buffered formalin for 30
min at RT. After washing twice with deionized water, 60% isopropanol was applied to the fixed
cells for 5 min, followed by a freshly prepared working solution of 1.5 mg/ml Oil Red O in
isopropanol for 5 min at RT. The stained, fixed tissues were then rinsed with tap water until clear,
covered with tap water and viewed on a phase contrast microscope. Size and areas of lipid droplet
were quantified by customarily written pipelines in CellProfiler (v4.1.3).

Generation of stable autophagy reporter and ER-phagy reporter cell line

Primary MEFs were isolated from *Rbbp5* (flox/flox); Rosa26 CRE (+/+) mice and immortalized
with SV40 lentivirus as previously described (32). For mCherry-EGFP-LC3B-expressing
autophagy reporter MEFs, lentiviruses packaged in HEK293T cells with co-transfection of pCDH-
EF1a-mCherry-EGFP-LC3B, pMD2.G and psPAX2 plasmids were used to infect *Rbbp5* (flox/flox);
Rosa26 CRE (+/+) MEFs with a multiplicity of infection (MOI) of 3 for three times to achieve near
complete infection. For mCherry-RAMP4-expressing ER-phagy reporter MEFs, lentiviruses
packaged in HEK293T cells with co-transfection of pLenti-X1-hygro-mCherry-RAMP4, pMD2.G
and psPAX2 plasmids were used to infect *Rbbp5* (flox/flox); Rosa26 CRE (+/+) MEFs with a
multiplicity of infection (MOI) of 3. Stable MEFs were selected in the presence of 200µg/ml
hygromycin.

siRNA Transient Transfections

Immortalized MEFs isolated from wild-type C57BL/6J mice were transfected with 10µM of
different siRNAs for 24~48 hours with Lipofectamine RNAiMAX reagents (Life technologies) per
the manufacturer's instructions. Source of siRNA are as follows: siGENOME non-targeting siRNA

pool (Dharmacon, D-001206-1305), siGENOME SMARTpool mouse *Rbbp5* siRNA (Dharmacon,
L-054560-01-0005), siGENOME SMARTpool mouse *Ash2l* siRNA (Dharmacon, L-048754-01-
0005), siGENOME SMARTpool mouse *Setd1a* siRNA (Dharmacon, L-051358-01-0005),
siGENOME SMARTpool mouse *Kat2a/Gcn5* siRNA (Dharmacon, L-040665-01-0005),
siGENOME SMARTpool mouse *Kat2b/Pcaf* siRNA (Dharmacon, L-049885-01-0005) and
siGENOME SMARTpool mouse *Xbp1* siRNA (Dharmacon, L-040825-00-0005). MEFs were
transfected with 10µM of different siRNAs for 48 hours and treated with DMSO or 100ng/ml of
tunicamycin for 8h.

Real-time Luminescence Assay

Stable *Manf*-dluc MEFs were transfected with non-targeting, *Rbbp5,* or *Ash2l* siRNA for 48 hours
in DMEM (4.5g/L glucose) supplemented with 10% FBS and treated with 25ng/ml tunicamycin in
DMEM for 2h before subjected to real-time luminescence assay using a Lumicycle (Actimetrics)
as previously described (33). Briefly, after tunicamycin treatment, MEFs were washed with 1x
PBS and cultured with DMEM (4.5g/L glucose) supplemented with 0.1 mM luciferin and 10mM
HEPES buffer in 35 mm tissue culture dishes in the absence of serum and transferred immediately
to Lumicycle for real-time luminescence analysis. Periods of oscillation were identified by
embedded Periodogram function.

Immunofluorescence (IF)

Immunofluorescence was performed as previously described (17). Briefly, liver OCT sections
were fixed in cold acetone for 10 mins at -20 ºC. The sections were then air dried, rehydrated with
PBS and permeabilized with PBS+ 0.1% Triton X-100. The sections were then blocked with 10%
goat serum at room temperature for 1 hour. Primary antibody against RBBP5 (Bethyl, A300-109A)
was added to the OCT section at 1:1000 dilution overnight at 4 ºC. Next day, sections were
washed three times with PBS and stained with Alexa-555 anti-rabbit secondary antibody at room
temperature for 2 hours. After that, the sections were washed with PBS three times and
counterstained with DAPI before mounting (with ProlongGold Glass) and imaging using Leica
SP8 lightening confocal microscope (Leica Microsystems).

Time-lapse microscopy for autophagy reporter MEFs

Time-lapse imaging was performed using SP8 lightening confocal microscope (Leica
Microsystems) with Okolab stage top incubator to maintain constant CO_2 (5%), temperature (37
ºC) and humidity (90%). mCherry-EGFP-LC3B-expressing *Rbbp5* (flox/flox); Rosa26 CRE (+/+)
MEFs were plated into 8-well chamber slides in full DMEM media and treated with (to deplete
RBBP5) or without tamoxifen (3µM/ml) for seven days before replacing with full DMEM or leucine-
free media counterstained with DAPI and images were taken every 2 hours for a total of 16 hours.
The number of cytosolic autophagosomes (GFP and mCherry double positive foci) and
autolysosomes (GFP negative and mCherry positive foci) per cell were quantified manually.

Immunoblot

Nuclear extracts were made from liver according to previously published protocol (34). Whole cell
lysates were lysed in RIPA buffer as previously described (17). Protein concentrations were
determined by Bradford assays (Bio-Rad), and aliquots were snap-frozen in liquid nitrogen and
stored at -80°C until usage. Immunoblot analyses were performed as described previously (35).
Briefly, 25~50µg proteins separated by 4~20% gradient SDS-PAGE gels (Biorad) were transferred
to nitrocellulose membranes, blocked in TBST buffer supplemented with 5% bovine serum
albumin (BSA) or 5% fat-free milk and incubated overnight with primary antibody anti-PERK (Cell
signaling, #3192), anti-phospho-PERK (Thr908) (Thermo Fisher, MA5-15033), anti-ATF4 (Cell
signaling, #11815), anti-IRE1α (Cell signaling, #3294), anti-phospho-IRE1α (Ser724) (ABclonal,

AP0878), anti-XBP1s (Biolegend, 658802), anti-RBBP5 (Bethyl, A300-109A), anti-RBBP5 (D3I6P)
(Cell signaling, #13171), anti-WDR5 (Cell signaling, #13105), anti-ASH2L (Bethyl, A300-489A),
anti-ASH2L (Bethyl, A300-112A), anti-SETD1A (Diagenode, CS-117-100), anti-SETD1B
(Diagenode, CS-118-100), anti-MLL1-c (D6G8N) (Cell signaling, #14197), anti-mCherry (E5D8F)
(Cell signaling, #43590), anti-p62/SQSTM1 (Santa Cruz, sc-28359) and anti-β-ACTIN (Cell
signaling, #12620) at 4°C. Blots were incubated with an appropriate secondary antibody coupled
to horseradish peroxidase at room temperature for 1 hour, and reacted with ECL reagents per the
manufacturer's (Thermo) suggestion and detected by Biorad ChemiDoc MP Imaging System.

Co-Immunoprecipitation (Co-IP)

Nuclear extracts (NE) were made from liver according to our published protocol (34). Protein
concentrations were determined by Bradford assays (Bio-Rad), and aliquots were snap-frozen in
liquid nitrogen and stored at -80°C until usage. 200 µg of NE was used for per IP. 5 µg of different
antibodies or control IgG were coupled to 1.5 mg of magnetic Dynabeads (Life Technologies) for
every IP using Dynabeads Antibody Coupling Kit (Life Technologies) per manufactures' protocol.
Co-IP was essentially carried out as previously described (36) except that coupled antibody
Dynabeads was added to the NEs for incubation. The antibodies used for Co-IP are as follows:
anti-XBP1s (Biolegend, 658802).

qRT-PCR

Total mRNA was isolated from murine embryonic fibroblasts (MEFs) or mouse liver with PureLink
RNA mini kit (Life Technologies) with additional on-column DNase digestion step to remove
genomic DNA per the manufacturer's instructions. Reverse transcription was carried out using
5µg of RNA using Superscript III (Life Technologies) per the manufacturer's instructions. For gene
expression analyses, cDNA samples were diluted 1/30-fold (for all other genes except for 18sRNA)
and 1/900-fold (for 18sRNA). qPCR was performed using the SYBR green system with sequence-
specific primers. All data were analyzed with 18S or *β-actin* as the endogenous control. qPCR
primer sequences are as follows and all primers span introns, except for primers for quantifying
pre-mRNAs:
Mouse *Rbbp5* forward primer: CAGAGCCTATGCAGAAGCTGCA
Mouse *Rbbp5* reverse primer: CTTCACCAGGTTGCCAATGCTC
Mouse *Ddit3* forward primer: GGAGGTCCTGTCCTCAGATGAA
Mouse *Ddit3* reverse primer: GCTCCTCTGTCAGCCAAGCTAG
Mouse *Creld2* forward primer: CTACACCAAGGAGAGTGGACAG
Mouse *Creld2* reverse primer: TCTGTCTCCTCAAAGCCTTCCG
Mouse *Hspa5* forward primer: TGTCTTCTCAGCATCAAGCAAGG
Mouse *Hspa5* reverse primer: CCAACACTTCCTGGACAGGCTT
Mouse *Dnajb11* forward primer: TGTGACCGTCTCACTGGTTGAG
Mouse *Dnajb11* reverse primer: CCCTTTCTTCCACAGCTTGGCT
Mouse *Pdia4* forward primer: TGGGCTCTTTCAGGGAGATGGT
Mouse *Pdia4* reverse primer: GGGAGACTTTCAGGAACTTGGC
Mouse *Hsp90b1* forward primer: GTTTCCCGTGAGACTCTTCAGC
Mouse *Hsp90b1* reverse primer: ATTCGTGCCGAACTCCTTCCAG
Mouse *Syvn1* forward primer: CCAACATCTCCTGGCTCTTCCA
Mouse *Syvn1* reverse primer: CAGGATGCTGTGATAAGCGTGG
Mouse *Gdf15* forward primer: ACTCAGTCCAGAGGTGAGATTG
Mouse *Gdf15* reverse primer: GGGGCCTAGTGATGTCCCAG
Mouse *Il1a* forward primer: ACGGCTGAGTTTCAGTGAGACC
Mouse *Il1a* reverse primer: CACTCTGGTAGGTGTAAGGTGC
Mouse *Il1b* forward primer: TGGACCTTCCAGGATGAGGACA

Mouse *Il1b* reverse primer: GTTCATCTCGGAGCCTGTAGTG
Mouse *Fga* forward primer: GGATTCTAACTCACTGACCAGGA
Mouse *Fga* reverse primer: CCTCAGGATCTCAATTCTGCGC
Mouse *C3* forward primer: CGCAACGAACAGGTGGAGATCA
Mouse *C3* reverse primer: CTGGAAGTAGCGATTCTTGGCG
Mouse *Tifa* forward primer: AGCAAGAAAACCAGTTTGATGGTAG
Mouse *Tifa* reverse primer: GCAACAGGAACTGATACTCTCCG
Mouse *Fgl1* forward primer: GGAAACTGTGCTGAGGAAGAGC
Mouse *Fgl1* reverse primer: TCCGTTTCTGCCCTGTAGGAAC
Mouse *Fgb* forward primer: TGACACCTCCATCAAGCCGTAC
Mouse *Fgb* reverse primer: GGTCCCATTTCCTGCCAAAGTC
Mouse *Il6* forward primer: CTTCCATCCAGTTGCCTTCT
Mouse *Il6* reverse primer: CTCCGACTTGTGAAGTGGTATAG
Mouse *Herpud1* forward primer: acctgagccgagtctaccc
Mouse *Herpud1* reverse primer: aacagcagcttccagaataaa
Mouse *Nbr1* forward primer: GGATTTAAAGCACCTCCTGATTCC
Mouse *Nbr1* reverse primer: ATTGGGTCCCACTCAGGTCT
Mouse *Sesn2* forward primer: CTGGCCGAACTCATCCAAG
Mouse *Sesn2* reverse primer: CTGCCTCATGCGTTCCATC
Mouse *Sqstm1* forward primer: AACAGATGGAGTCGGGAAAC
Mouse *Sqstm1* reverse primer: AGACTGGAGTTCACCTGTAGA
Mouse *Stbd1* forward primer: CTGGGAAGTAGATGGGAAAGTG
Mouse *Stbd1* reverse primer: TCGGTGTTTGTGGTGTAGTG
Mouse *Sirt1* forward primer: TGACAGAACGTCACACGCCA
Mouse *Sirt1* reverse primer: ATTGTTCGAGGATCGGTGCCA
Mouse *Gcn5* forward primer: TTGATTGAGCGCAAACAGGC
Mouse *Gcn5* reverse primer: CAGCCTGTCTCTCGAATGCC
Mouse *Pcaf* forward primer: CGTGAAGAAGGCGCAGTTG
Mouse *Pcaf* reverse primer: CAGGACTCCTCTGCCTTGC
Mouse *Ash2l* forward primer: gctgtgtctgctagtgggaac
Mouse *Ash2l* reverse primer: catcttgctgcttacgcttg
Mouse *Setd1a* forward primer: gggtagcaccccctactctc
Mouse *Setd1a* reverse primer: gggtttgaaggaggttgaagt
Mouse *Eif2ak3* forward primer: ccttggtttcatctagcctca
Mouse *Eif3ak3* reverse primer: atccagggaggggatgat
Mouse total *Xbp1* forward primer: gggtctgctgagtcc
Mouse total *Xbp1* reverse primer: cagactcagaatctgaagagg
Mouse *Sec23b* forward primer: tgaccaaactggacttctgga
Mouse *Sec23b* reverse primer: aaagaatctcccatcaccatgt
Mouse *Manf* forward primer: gacagccagatctgtgaactaaaa
Mouse *Manf* reverse primer: tttcacccggagcttcttc
Mouse *Hyou1* forward primer: GAGGCGAAACCCATTTTAGA
Mouse *Hyou1* reverse primer: GCTCTTCCTGTTCAGGTCCA
Mouse *Atf4* forward primer: CCACTCCAGAGCATTCCTTTAG
Mouse *Atf4* reverse primer: CTCCTTTACACATGGAGGGATTAG
Mouse *Atf6* forward primer: CATGAAGTGGAAAGGACCAAATC
Mouse *Atf6* reverse primer: CAGCCATCAGCTGAGAATTAGA
Mouse 18s RNA forward primer: ctcaacacgggaaacctcac
Mouse 18s RNA reverse primer: cgctccaccaactaagaacg
Mouse *β-actin* forward primer: aaggccaaccgtgaaaagat
Mouse *β-actin* reverse primer: gtggtacgaccagaggcatac

RNA-Seq to identify RBBP5-dependent 12h hepatic transcriptome

Mouse liver tissues were collected from *Rbbp5* *Flox* (n=2) and *Rbbp5* *LKO* (n=2) mice at 2h intervals for a total of 48 hours under constant darkness condition. Total RNA was isolated from mouse liver with miRNeasy Mini Kit (Qiagen) per the manufacturer's instructions. RNA was assessed for quality using an Agilent TapeStation 4150/Fragment Analyzer 5300 and RNA concentration was quantified on a Qubit FLEX fluorometer. Libraries were generated with the Illumina Stranded mRNA Library Prep kit (Illumina: 20040534) according to the manufacturer's instructions. Briefly, 200 ng of input RNA was used for each sample. Following adapter ligation, 12 cycles of indexing PCR were completed, using IDT for Illumina RNA UD Indexes (Illumina: 20040553-6). Library quantification and assessment was done using a Qubit FLEX fluorometer and an Agilent TapeStation 4150/Fragment Analyzer 5300. The prepared libraries were pooled and sequenced using NoveSeq 6000 (Illumina), generating an average of 40 million 2×100 bp paired end reads per sample. RNA-Seq library preparation and sequencing were performed at UPMC Genome Center. Raw RNA-seq FASTQ files were analyzed by FastQC for quality control. Adaptors and low-quality reads were filtered by Trimmomatic (37). Then the processed reads were aligned by HISAT2 (38) against mouse reference mm10. Gene and isoform FKPM values were calculated by Cufflinks. Since the background gene expression level is FPKM=0.1 in mouse liver (21), only those genes with average gene expression across 48 hours in Rbbp5 *Flox* mice larger than 0.1 were included for rhythm-identification analysis. Averaged FPKM values at each time were used for cycling transcripts identification by the eigenvalue/pencil method. Raw temporal data was subject to polynomial detrend (n=2) first, and superimposed oscillations were identified using previously described eigenvalue/pencil method on the detrended dataset (23,39). Specifically, three oscillations were identified from each gene. Criterion for circadian gene is period between 21h to 27h, decay rate between 0.9 and 1.1; for ~12hr genes: period between 10h to 13h, decay rate between 0.9 and 1.1. The relative amplitude is calculated via dividing the amplitude by the mean expression value of each gene. All analysis were performed in MatlabR2017A. Heat maps were generated by Gene Cluster 3.0 and TreeView 3.0 alpha 3.0 using log2 mean-normalized values. RAIN analysis was performed as previously described in Bioconductor (3.4) (http://www.bioconductor.org/packages/release/bioc/html/rain.html) (40) using the full detrended dataset with duplicated at each time point.

For the eigenvalue method, every gene consists of multiple superimposed oscillations. Therefore, we define a circadian gene as any gene that exhibits a superimposed circadian rhythm, regardless of its relative amplitude compared to other superimposed oscillations. Similar definitions apply to 12hr genes. Under this definition, a gene can be both a circadian and 12hr-cycling gene. By comparison, we define a dominant circadian gene as any gene whose superimposed circadian rhythm has the largest amplitude among all oscillations. Similar definitions also apply to 12hr genes. Under this definition, dominant circadian and dominant 12hr genes are mutually exclusive.

RNA-seq for identifying RBBP5-dependent acute transcriptional response to ER stress

Mouse liver tissues were collected from *Rbbp5* *Flox* and *Rbbp5* *LKO* mice intraperitoneally treated with or without Tu for 8 hours. Total mRNA was isolated from mouse liver with PureLink RNA mini kit (Life Technologies) with additional on-column DNase digestion step to remove genomic DNA per the manufacturer's instructions. Detailed procedures for transcriptome sequencing were described previously(41) and were supported by the Genomics and Systems Biology Core of the Pittsburgh Liver Research Center. Briefly, the mRNA samples per mouse were processed into short-read libraries using the Bio-Rad SEQuoia Dual Indexed Primers. Next, the circularization was performed using the Element Biosciences Adept Library Compatibility Kit v1.1 based on the manufacturer's instructions, followed by the quantification using qPCR with the provided standard.

Finally, the libraries were measured by the Element Biosciences AVITI system to sequence paired end reads with 75 bp each. The raw FASTQ files from RNA-seq were trimmed by Trimmomatic (42) to filter out low-quality reads. The surviving reads were then aligned to the mouse mm10 reference genome by STAR aligner(43). Fragments Per Kilobase of transcript per Million mapped reads (FPKM) values per library were quantified by the tool Cufflinks(44). Log 2 normalized fold induction and adjusted p values between *Rbbp5* *Flox* DMSO and *Rbbp5* *LKO* DMSO, *Rbbp5* *Flox* Tu and *Rbbp5* *LKO* Tu, *Rbbp5* *Flox* DMSO and *Rbbp5* *Flox* Tu, and *Rbbp5* *LKO* DMSO and *Rbbp5* *LKO* Tu were calculated by using the iDEP (integrated Differential Expression and Pathway analysis) online application (45).

Chromatin Immunoprecipitation (ChIP)-Seq and ChIP-qPCR

ChIP for RBBP5 was performed using anti-RBBP5 antibody (Bethyl, A300-109A) as previously described (21). Briefly, mouse liver samples were submerged in PBS + 1% formaldehyde, cut into small (~1 mm3) pieces with a razor blade and incubated at room temperature for 15 minutes. Fixation was stopped by the addition of 0.125 M glycine (final concentration). The tissue pieces were then treated with a TissueTearer and finally spun down and washed twice in PBS. Chromatin was isolated by the addition of lysis buffer, followed by disruption with a Dounce homogenizer. The chromatin was enzymatically digested with MNase. Genomic DNA (Input) was prepared by treating aliquots of chromatin with RNase, Proteinase K and heated for reverse-crosslinking, followed by ethanol precipitation. Pellets were resuspended and the resulting DNA was quantified on a NanoDrop spectrophotometer. An aliquot of chromatin (30 µg) was precleared with protein A agarose beads (Invitrogen). Genomic DNA regions of interest were isolated using 4 µg of antibody. Complexes were washed, eluted from the beads with SDS buffer, and subjected to RNase and proteinase K treatment. Crosslinking was reversed by incubation overnight at 65 °C, and ChIP DNA was purified by phenol-chloroform extraction and ethanol precipitation. The DNA libraries were prepared at the University of Pittsburgh and sequenced at Gene by Gene, Ltd per standard protocols. DNA libraries were prepared with Ovation® Ultralow V2 DNA-Seq library preparation kit (NuGen) using 1ng input DNA. The size selection for libraries was performed using SPRIselect beads (Beckman Coulter) and purity of the libraries were analyzed using the High Sensitivity DNA chip on Bioanalyzer 2100 (Agilent). The prepared libraries pooled and sequenced using Nova-Seq 6000 (Illumina), generating ~30 million 75 bp single-end reads per sample. ChIP-qPCR for MEFs were essentially performed the same way as previously described with anti-RBBP5 (Bethyl, A300-109A) and anti-H3K4me3 (Active Motif, 61379) antibodies, except that the MEFs were directly fixed with 1% formaldehyde before subject to nuclei isolation and chromatin immunoprecipitation. The primers used for ChIP-qPCR are as follows:
Negative control region 1 forward primer: GCAACAACAACAGCAACAATAAC
Negative control region 1 reverse primer: CATGGCACCTAGAGTTGGATAA
Manf promoter forward primer: CCCTTAAATGGGTCAACGTCTC
Manf promoter reverse primer: GGCGCTAAACCCAAGGAAA
Xbp1 promoter forward primer: TCCGTACGGTGGGTTAGAT
Xbp1 promoter reverse primer: ACCTTCTTCTGTGCCTGTG
Hyou1 promoter forward primer: CCGTGTGGGTACGTCCT
Hyou1 promoter reverse primer: GACGGCTGCTCCATCCT
Sesn2 promoter forward primer: ACAGGAGGCCGGGACTA
Sesn2 promoter reverse primer: CTGGGCTGAAAGGAGTGTCTAT

ChIP-Seq analysis

The sequences identified were mapped to the mouse genome (UCSC mm10) using BOWTIE function in Galaxy Deeptool (https://usegalaxy.org/) (46). Only the sequences uniquely mapped with no more than 2 mismatches were kept and used as valid reads. PCR duplicates were also

removed. Peak calling was carried out by MACS2 (version 2.1.1.20160309) in Galaxy (options --
mfold 5, 50 --pvalue 1e-4), on each ChIP-seq file against input. To account for the different
sequencing depths between samples, the BAM files generated from MACS2 were RPKM
normalized to sequencing depth using the bamCoverage function in Galaxy Deeptool and the
bigwig files were generated accordingly. The relative intensity of each RBBP5 binding site is
further calculated via the computeMatrix function with the RPKM normalized bigwig files and bed
files from the peak calling as inputs by calculating the area under the curve.

Proximity labeling

Proximity labeling was performed using Protein A-TurboID as previously described (47). Briefly,
target cells were grown to 90-100% confluency in 15 cm dishes, harvested in PBS using a cell
scraper, and pelleted by centrifugation. Nuclei was isolated by resuspending the cell pellet in ice-
cold HLB buffer, incubating on ice, and washing through sequential centrifugation steps. Isolated
nuclei were incubated with digitonin buffer containing ~2 μg of IgG or H3K4me3 antibody (Active
Motif, 61379) for 20 minutes at room temperature with shaking, followed by incubation with 1.4
μg of Protein A-TurboID for 30 minutes at 4°C. After washing, biotinylation was induced by
incubating samples in biotinylation buffer (containing ATP, $MgCl_2$, and 500 μM biotin) for 10
minutes at 37°C. Nuclei were lysed in RIPA buffer at 4°C overnight, sonicated, and centrifuged to
collect the supernatant. Streptavidin affinity purification was performed by incubating lysates with
M-270 streptavidin beads for 2 hours at room temperature or overnight at 4°C. Beads were
washed extensively with RIPA buffer, and bound proteins were eluted by boiling at 100°C for 15
minutes in 70 μL of 1.5X LDS buffer. Samples were clarified by centrifugation and magnetic
separation.

For protein identification, half of each sample was separated ~1.5 cm on a 10% Bis-Tris Novex
mini-gel (Invitrogen) using the MES buffer system. The gel was stained with Coomassie, and each
lane was excised into ten equally sized segments. Gel pieces were processed using a DigestPro
robot (CEM) with the following protocol: washes with 25 mM ammonium bicarbonate and
acetonitrile, reduction with 10 mM dithiothreitol at 60°C, alkylation with 50 mM iodoacetamide at
room temperature, digestion with trypsin (Promega) at 37°C for 4 hours, and quenching with
formic acid. The resulting peptides were analyzed by nano LC-MS/MS using a Waters M-class
HPLC system interfaced with a ThermoFisher Fusion Lumos mass spectrometer. Peptides were
loaded onto a trapping column and eluted over a 75 μm analytical column at 350 nL/min with a
30-minute gradient. MS and MS/MS spectra were acquired in the Orbitrap at resolutions of 60,000
and 15,000 FWHM, respectively. Data was searched using Mascot against the SwissProt Mouse
database with a 10 ppm peptide mass tolerance and a 20 ppm fragment mass tolerance. Search
parameters included trypsin digestion, fixed carbamidomethylation of cysteines, and variable
modifications such as oxidation (M), acetylation (protein N-term), deamidation (NQ), and pyro-glu
(N-term Q). Data was processed using Scaffold software with a 1% false discovery rate (FDR) at
both the protein and peptide levels, requiring at least two unique peptides per protein.

Gene ontology analysis

DAVID (Version 2021) (48) (https://david.ncifcrf.gov) was used to perform Gene Ontology
analyses. Briefly, gene names were first converted to DAVID-recognizable IDs using Gene
Accession Conversion Tool. The updated gene list was then subject to GO analysis using all
Homo Sapiens as background and with Functional Annotation Chart function. GO_BP_DIRECT,
KEGG_PATHWAY or UP_KW_BIOLOGICAL_PROCESS was used as GO categories. Only GO
terms with a p value less than 0.05 were included for further analysis.

Motif analysis

501 Motif analysis was performed with the SeqPos motif tool (version 0.590) (49) embedded in Galaxy
502 Cistrome using all motifs within the mouse reference genome mm10 as background. LISA
503 analysis was performed using webtool (http://lisa.cistrome.org/).

Statistical Analysis

505 Data were analyzed and presented with GraphPad Prism software. Plots show individual data
506 points and bars at the mean and ± the standard error of the mean (SEM). One-tailed t-tests were
507 used to compare means between groups, with significance set at $p < 0.05$. In instances where
508 the p value is not shown, *, **, ***, and **** represent $p < 0.05$, 0.01, 0.001, and 0.0001,
509 respectively.

Results:

Global RBBP5 binding to chromatin exhibits a ~12h ultradian rhythm, concomitant with the promoter-proximal ~12h H3K4me3, but not H3K9Ac/H3K27Ac rhythms in mouse liver.

The 12h oscillator is transcriptionally regulated by the UPR TF XBP1s in mice (18,19,21), and
likely under control by additional bZIP TFs, such as ATF4 and ATF6 (50). However, decades of
research of eukaryotic gene regulation demonstrated that TFs themselves are not sufficient to
drive transcription initiation and elongation, and additional so-called "co-regulators" are required
for full gene activation by shaping the epigenetic landscape (51-53). Currently, the epigenetic
landscape of both the 12h oscillator and proteotoxic stress responses are poorly characterized.
To identify epigenetic regulators of global proteostasis control, we first examined a published
temporal hepatic epigenome dataset for histone modifications associated with the active
promoters of ~12h genes (54). We found ~12h rhythms of histone H3 trimethylated at lysine 4
(H3K4me3), but not two histone acetylation markers, histone H3 acetylated at lysine 9 (H3K9Ac)
and lysine 27 (H3K27Ac), at the promoters of ~12h genes (**Fig. 1A, B**). This is particularly
prominent for those ~12h genes with rhythmic expressions peaking at ZT0 and ZT12 that are
enriched for proteostasis pathways (21), with H3K4me3 level also peaking at these two time
points (**Fig. 1A, B**).

In eukaryotes, H3K4 is tri-methylated by the COMPASS complex, which includes regulatory
subunits RBBP5, ASH2L, WDR5, DPY30 and catalytic subunits SETD1A/B/MLL1-4 (**Fig. S1A**)
(55-57). Via writing H3K4me3, COMPASS is essential for the eviction of paused RNA polymerase
II (RNAPII) and promotes transcriptional elongation (58,59). Landscape In Silico deletion Analysis
(LISA) (60) further inferred COMPASS subunits WDR5, ASH2L, DYP30 and RBBP5 as putative
epigenetic regulators of hepatic ~12h, but not circadian genes (**Fig. S1B**). Being the commonly
shared regulatory subunits for all MLL family H3K4 methyl transferase complex (including
SETD1A/B and MLL1-4), RBBP5/ASH2L forms the minimal heterodimer sufficient to activate all
MLL histone methyl transferase **(61,62)** and are essential for the assembly of the whole
COMPASS complex **(57,63)**. Between the two essential subunits, we initially focused on RBBP5.
Eigenvalue/pencil method (64) revealed that the levels of both hepatic *Rbbp5* mRNA and nuclear
RBBP5 protein exhibit ~12h oscillations, with the nuclear protein level peaking at ~CT0 and
~CT12 (**Fig. S1C, D**), in line with the acrophases of H3K4me3 and gene expression oscillations
(**Fig. 1A, B**).

To further evaluate the potential role of RBBP5 in establishing the epigenetic landscape of
hepatic ~12h rhythms, we performed temporal RBBP5 ChIP-seq at a 4-h interval for a total of 48
hours in the liver of male C57BL/6 mice housed under constant darkness condition fed *ad libitum*.
Chromatin occupancies of RBBP5 align well with nucleosome positioning downstream of the
transcription start sites (TSS) and the H3K4me3 epigenome, with 'ridges' corresponding to 130bp,
310bp, 490bp, 670bp downstream of TSS at all time points, thus demonstrating the high quality

of our ChIP-seq dataset (**Fig. 1C-F**). In total, we identified 7,963 hepatic RBBP5 binding sites that
are within 2kb (up and downstream) of TSS of 6,451 genes (**Fig. 1C and Table S1**). The average
binding intensities of RBBP5 on all post-TSS nucleosomes displayed robust 12h rhythms, with
peaks occurring at CT0, CT12, CT24, and CT36 (**Fig. 1C-E**). These rhythms aligned with the 12h
oscillations of promoter-proximal H3K4me3 (**Fig. 1C-E**), but not with those of H3K9Ac or
H3K27Ac (**Fig. S1E**). This pattern is exemplified by *Xbp1*, one of the major UPR TFs and the
transcriptional regulator of the 12h oscillator, and *Hsph1*, a canonical 12h oscillator output gene
encoding a heat shock protein (**Fig. S1F**). Using the eigenvalue/pencil and RAIN (40) algorithms,
we identified 3,028 and 1,864 (p<0.05) RBBP5 binding sites that cycle with a ~12h period,
respectively (**Fig. S1G, H and Table S1**). Collectively, these results establish the epigenetic
landscape of hepatic 12h oscillator is marked by RBBP5-H3K4me3.

Hepatic RBBP5 cistrome coincides with that of XBP1s and hepatic ~12h transcriptome.

To investigate if RBBP5 regulates hepatic ~12h rhythms of gene expression, we first examined
the correlation between RBBP5 proximal promoter binding status and ~12h rhythms of gene
expression. As illustrated in **Fig. 2A**, genes with proximal promoter RBBP5 binding are enriched
for ~12h rhythms of gene expression (P=1.1e-39 by Chi-squared test). For those 2,525 ~12h
genes with RBBP5 binding, the phases of RBBP5 chromatin binding center around CT0 and CT12,
slightly preceding the phases of gene expression observed at ~CT1 and ~CT13 (**Fig. 2B**). This
temporal relationship suggests that RBBP5 acts as a driver, rather than a consequence of ~12h
rhythms of gene expression.

We next performed Gene Ontology (GO) analysis on those hepatic genes with proximal
promoter RBBP5 binding and revealed strong enrichment of proteostasis pathways including
protein processing in the ER and ubiquitin-mediated proteolysis (**Fig. 2C**). Additionally, pathways
involved in mRNA processing, such as spliceosome, RNA transport, and mRNA surveillance, were
also enriched (**Fig. 2C**). These findings align with previous observations that both mRNA and
protein metabolism pathways are enriched in genes exhibiting ~12h rhythms in both mice and
humans (16,17,65). The coupling of mRNA and protein metabolism is orchestrated by a SON-
XBP1s axis and involves the liquid-liquid phase separation (LLPS) dynamics of nuclear speckles
(17). Similar pathways are also enriched in genes exhibiting both ~12h rhythms of RBBP5
promoter binding and gene expression (**Fig. 2C**).

XBP1s is the major transcriptional factor regulating the hepatic 12h oscillator and also exhibits
a global 12h rhythm of chromatin binding to the promoter regions peaking at CT0, 12, 24 and 36
(19,21). Comparing the cistromes of RBBP5 and XBP1s revealed a significant overlap between
the two, with 920 genes sharing RBBP5 and XBP1s proximal promoter binding (**Fig. 2D**). Co-
immunoprecipitation in the hepatic nuclear extracts at CT0 confirmed the physical interaction
between XBP1s and subunits of SET1/COMPASS, including RBBP5, ASH2L, WDR5, and two
histone methyltransferase SETD1A and SETD1B (**Fig. 2E**). Motif analysis of the DNA sequences
around the 6,149 RBBP5 binding sites that do not overlap with XBP1s cistrome identified binding
motifs for GABPA, KLF, ATF6, NFYA and ATF4 (**Fig. 2F**). These findings suggest that, in addition
to acting as a co-activator for XBP1s, RBBP5 may also work with other transcription factors to
shape the hepatic 12h epigenome.

RBBP5 is an epigenetic regulator of the ~12h oscillator, but not the canonical ~24h core
circadian clock.

To establish the causality between RBBP5 and hepatic 12h rhythms of gene expression, we
generated RBBP5 liver hepatocyte-specific knockout (RBBP5 [LKO]) mice using the CRE-loxP
system. Exon 10 of mouse *Rbbp5,* which encodes the SET/ASH2L binding domain, was flanked
by loxP sites (**Fig. 3A**). CRE-mediated deletion of Exon 10 is expected to lead to a frameshift and

nonsense-mediated RNA decay of truncated *Rbbp5* transcript (**Fig. S2A**). Crossing homozygous
floxed *Rbbp5* mice with Albumin-CRE mice resulted in liver-specific deletion of RBBP5, which
was confirmed by DNA genotyping, IF against RBBP5 and western blot analysis (**Figs. 3B and
S2B-D**). The resulting RBBP5 ^{LKO} mice weighed slightly less than wild-type counterparts (**Fig.
S3A**) but maintained normal rhythmic locomotor activity and fasting-feeding cycles under both
12h:12h L/D and constant darkness conditions (**Fig. S3B-I**).

To identify the RBBP5-dependent oscillating hepatic transcriptome, we performed bulk RNA
sequencing (RNA-Seq) analysis in the liver of male *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice at 2-h intervals
for a total of 48 hour under constant darkness in duplicates (**Table S2**). To identify genes cycling
with a period ranging from 6 to 36 hours, we initially applied the RAIN algorithm (40) to each
genotype's temporal transcriptome (40) (**Table S3**). Compared to alternative methods like
JTK_CYCLE, RAIN detects rhythms with arbitrary waveforms and therefore more robustly
uncovers ultradian rhythms (21,23,39). Consistent with past studies (18,20,21,66,67), two
populations of hepatic transcripts cycling with periods of ~12h and ~24h were identified in *Rbbp5*
^{Flox} mice liver (**Fig. 3C, D**). By contrast, in *Rbbp5* ^{LKO} mice, the ~12h rhythms are all but abolished
and the circadian oscillations remain largely intact, which include all core circadian clock genes
(such as *Bmal1*, *Per1/2/3*, and *Nr1d1/2*) displaying robust circadian rhythms with the same
phases and amplitudes as those observed in *Rbbp5* ^{Flox} mice (**Figs. 3C-F and S4A**). The
maintenance of ~24h circadian rhythms and abolishment of ~12h ultradian rhythms in *Rbbp5* ^{LKO}
mice is further confirmed by the principal component analysis (PCA) (**Fig. S4B, C**). Specifically,
using RAIN with an FDR cutoff of 0.01 and 0.05, we identified a total of 3,549 and 6,025 hepatic
transcripts cycling at periods between 10-12h in *Rbbp5* ^{Flox} mice, respectively. By contrast, only
61 and 813 10-12h genes were observed in *Rbbp5* ^{LKO} mice with the same FDR thresholds,
indicating over 93% ~12h hepatic transcriptome was abolished with liver RBBP5 ablation (**Figs.
3G, H, S4D and Table S3**). GO analysis confirmed that these RBBP5-depenent ~12h hepatic
transcriptomes are enriched in various mRNA processing and proteostasis pathways, distinct from
those of RBBP5-independent circadian genes enriched in lipid metabolism and phosphorylation
(**Fig. 3I**).

To determine the robustness of our results to different analytic methods, we also performed
spectrum analysis with the eigenvalue/pencil method (**Table S4**) (19-21,23,26,39), which unlike
statistical methods such as JTK_CYCLE and RAIN does not require pre-assignment of period
range, enabling unbiased identification of multiple superimposed oscillations for any given gene
(19-21,23,26,39). Revealing up to three oscillations from each gene, eigenvalue/pencil analyses
also revealed prevalent ~24h and ~12h oscillations in *Rbbp5* ^{Flox} mice, along with a third
population cycling at a period around 8h (**Fig. S5A-E**). Hepatic ~8h oscillations were also
previously reported but their regulation and function remained elusive (22,66). *Rbbp5* ^{LKO} mice,
on the other hand, exhibited a drastically different spectrum, characterized by the near complete
loss of the ~12h rhythms, *de novo* gains of many faster ultradian rhythms with periods between 6
and 7h, and the maintenance of most circadian rhythms (**Fig. S5A-E**). Similar findings were also
observed for dominant oscillations (which is the one with the largest amplitude among the three
superimposed oscillations for each gene) for each gene (**Fig. S5A-E**). For the ~12h genes
originally identified in *Rbbp5* ^{Flox} mice but lost in *Rbbp5* ^{LKO} mice, the period spectrum in the *Rbbp5*
^{LKO} mice displayed a diverse distribution. This ranged from a complete lack of rhythmicity (~39%)
to shorter ultradian periods of 6-7 hours, and even to longer periods exceeding 24 hours (**Fig.
S5F**). RBBP5-dependent ~12h transcriptome uncovered by the eigenvalue/pencil showed
substantial overlap with the results obtained through the RAIN method. This convergence was
evident both in the specific genes identified (**Fig. S5C**) and in the enriched biological pathways
(**Fig. S5G, H**), thereby confirming the robustness of our findings. These results thus establish

RBBP5 as a central epigenetic regulator of the hepatic 12h oscillator, with more than 90% ~12h hepatic transcriptome abolished in its absence.

In addition to liver, cell-autonomous ~12h rhythms of proteostasis gene expression including *Xbp1* can also be observed in mouse embryonic fibroblasts (MEFs) synchronized by low concentration of ER stress inducer tunicamycin (Tu) **(21,23)**. We observed a ~12h rhythm of *Rbbp5* expression in Tu-synchronized MEFs, which is independent of BMAL1 but require XBP1s (**Fig. S6A**), indicating RBBP5 itself is also under 12h oscillator control. To test the functional role of COMPASS in regulating the cell-autonomous ~12h rhythms, we knocked down either *Rbbp5* or *Ash2l* via siRNA in a previously described 12h oscillator reporter MEFs that stably express *Manf* promoter-driven destabilized luciferase (dluc) (21), and found that both siRNAs significantly dampen the ~12h oscillation of luciferase activity as well as shorten the period (**Fig. S6B**). By contrast, neither *RBBP5* nor *ASH2L* knocking down affects the circadian oscillation of *Bmal1* promoter activity in human U2OS cells in a previous genome-wide siRNA screen study (68) (**Fig. S6C**). Collectively, our results demonstrate that RBBP5 is an integral component of the 12h oscillator, but dispensable for the canonical circadian clock both *in vivo* and *in vitro*.

RBBP5 modulates the hepatic transcriptional response to acute proteotoxic stress *in vivo*.

Before the discovery of the mammalian 12h oscillator, the dynamics of proteostasis were primarily studied as transient responses to various proteotoxic stresses, with a classic example being the UPR triggered by ER stress. In response to the accumulation of misfolded proteins in the ER, the UPR initiates a signaling cascade from the ER to the nucleus, ultimately activating three key transcription factors: XBP1s, ATF4, and ATF6. This activation leads to a beneficial adaptive response, upregulating proteostasis genes to restore ER homeostasis (**Fig. 4A**) (69-71). However, ER stress can also trigger maladaptive and deleterious responses, such as the activation of pro-inflammatory and immune gene expression through the TRAF2-mediated activation of AP1 (comprised of c-Fos and c-Jun), NF-κB or STAT3 TFs (**Fig. 4A**) (72). Our findings that RBBP5 can co-activate XBP1s to regulate the 12h oscillator suggest that RBBP5 is essential for the adaptive UPR TFs-mediated activation of proteostasis genes but is not required for the maladaptive activation of immune genes by immune-regulatory TFs, whose binding motifs are absent from RBBP5 binding sites (**Fig. 2H**).

To test this hypothesis, we intraperitoneally injected male *Rbbp5* [Flox] and *Rbbp5* [LKO] mice with vehicle control or 0.05mg/kg Tu to induce acute ER stress and performed RNA-seq on hepatic transcriptome isolated 8 hours post injection (**Table S5**). This lower dose of Tu was selected for its ability to induce the UPR, but that was significantly less toxic than the more common experimental dose of 1 mg/kg—which is known to elicit hepatocellular death and severe toxicity (73). Western blot analysis confirmed the depletion of RBBP5 and a reduction of H3K4me3 level in the liver of *Rbbp5* [LKO] mice (**Fig. 4B**). PCA demonstrated the separation of the four groups on the transcriptome space (**Fig. 4C**). Using an adjusted p value (FDR) of 0.10 as cut-off, we identified 139 and 45 genes significantly induced by Tu in the liver of *Rbbp5* [Flox] and *Rbbp5* [LKO] mice, respectively, with 21 commonly shared between the two (**Figs. 4D, S7A-G and Table S6**). In *Rbbp5* [Flox] mice, these 139 genes are enriched in pathways of proteostasis and immune functions as expected, with pathways of 'response to ER stress' and 'ERAD' among the top two GO terms (**Fig. 4D, E**). By contrast, in *Rbbp5* [LKO] mice, these 45 genes are only enriched in immune functions, with proteostasis genes not being significantly induced by Tu (**Fig. 4D, E**). In general, proteostasis genes were induced by Tu to a much greater fold in *Rbbp5* [Flox] mice (such as *Creld2*, *Xbp1*, *Hsp90b1*), while the opposite was found for blood coagulation and immune genes in *Rbbp5* [LKO] mice (such as *S100a9* and *Fgb*) (**Figs. 4D, E and S7E, F, H**). Further integrating RBBP5 ChIP-seq results, we found that, on average, the ablation of RBBP5 markedly diminished the fold induction by Tu for genes with RBBP5 promoter binding, while it had no effects

for those without RBBP5 chromatin recruitment (**Fig. 4D, F**). To rule out the possibility that RBBP5
regulates UPR via influencing the upstream ER stressing in the ER, we performed western blot
analysis of p-IRE1α and p-PERK in the liver of *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice and did not observe
any notable difference in the increase of their phosphorylated level in response to Tu between the
two genotypes (**Fig. 4G**).

To determine if RBBP5 overexpression is also sufficient to amplify UPR, we tail-vein injected
AAV8-TBG-GFP and AAV8-TBG-RBBP5 viruses into wild-type male mice followed by
intraperitoneal injection with vehicle control or 0.025mg/kg Tu 30 days later. Thyroxine binding
globulin (TBG) promoter is liver-specific and directs RBBP5 transgene expression in hepatocytes
(74). qPCR and western blot analysis confirmed an increase of total hepatic expression of RBBP5
of ~1.4 fold in AAV8-TBG-RBBP5 group under vehicle control condition (**Fig. 4H, I**). Consequently,
RBBP5 hepatic overexpression significantly amplified the UPR of proteostasis genes including
Manf, *Hspa5* (*BiP*), *Pdia4*, *Hyou1*, and *Syvn1* without affecting the upstream ER stress sensing
(**Fig. 4H, I**). In addition, we found that RBBP5 overexpression attenuated the expression of
several immune and blood coagulation genes such as *C3*, *Tifa*, *Fgl1* and *Fgb* under ER stress
conditions (**Fig. 4I**). Taken together, these results demonstrate that RBBP5 is both necessary and
sufficient for the transcriptional regulation of hepatic response to acute proteotoxic stress *in vivo*.
RBBP5 activates the expression of proteostasis genes while repressing those involved in immune
functions.

RBBP5 protects mice from chronic ER stress-induced hepatic inflammation and steatosis.

Contrary to acute ER stress, chronic ER stress, experimentally reconstituted by repeated injection
of Tu, paradoxically leads to feedback-mediated suppression of the UPR signaling, to even below
basal levels (73). Proposed mechanisms of this suppression include silencing of the ATF6a
branch of UPR, enhancement of mRNA degradation via IRE1-dependent decay (RIDD), and the
direct inhibition of UPR stress sensors by the ER chaperone HSPA5 (BiP) (73). To investigate
whether RBBP5 also modulates the feedback suppression of UPR under chronic ER stress
conditions, we repeatedly intraperitoneally injected male *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice with
vehicle control or 0.025mg/kg Tu for six consecutive days and harvested liver tissues 24 hours
after the last injection for analysis (**Fig. 5A**). Western blot analysis revealed attenuation of p-
IRE1α and ATF4 level in both *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice with repeated Tu injection (**Fig. 5B**).
In agreement with the Western blot result, qPCR analysis also confirmed the downregulation of
proteostasis genes *Atf4*, *Hyou1*, and *Manf* in both mice under chronic ER stress condition (**Fig.
5C**). By contrast, compared to *Rbbp5* ^{Flox} mice, the expression of pro-inflammatory cytokine genes
Il1α and *Il1b* were higher in Tu-injected *Rbbp5* ^{LKO} mice, consistent with higher levels of p-IRE1α
and p-c-Jun (Ser73) observed in *Rbbp5* ^{LKO} mice (**Fig. 5B, C**). The persistent elevated levels of
p-IRE1α/p-c-JUN and pro-inflammatory cytokine gene expression in *Rbbp5* ^{LKO} mice suggest that
these mice exhibit heightened susceptability to Tu-induced ER stress and inflammation. In
summary, our results indicate that while wild-type mice can sustain persistent cycles of UPR
activation and deactivation in response to ER stress, as previously described (73), the loss of
RBBP5 disrupts this balance. Although RBBP5-deficient mice retain the ability to attenuate UPR,
they are unable to properly activate it. Consequently, *Rbbp5* ^{LKO} mice exhibit consistently lower
expression of adaptive proteostasis genes under ER stress condition, leading to heightened
proteotoxic stress and chronic inflammation likely via activating the p-IRE1α-TRAF2-AP1
signaling cascade.

Both chronic ER stress and inflammation can result in hepatic steatosis (75-78). We therefore
performed Oil Red O staining, which stained all neutral lipids, on liver tissues from *Rbbp5* ^{Flox} and
Rbbp5 ^{LKO} mice after repeated vehicle or Tu injection. *Rbbp5* ^{LKO} mice liver displayed an increased
number of lipid droplets and total lipid content, compared to wild-type counterparts under both

basal and chronic ER stress conditions (**Fig. 5D**). Our data revealed that RBBP5 protects mice
from chronic ER stress-induced hepatic inflammation and steatosis.

Loss of RBBP5 compromises the cell-autonomous transcriptional response to proteotoxic
stress and nutrient-deprivation, resulting in impaired autophagy and reduced cell survival.

To determine whether RBBP5 regulates transcriptional responses to proteotoxic stress in a cell-
autonomous manner beyond the liver, we examined MEFs, which exhibit both a robust cell-
autonomous XBP1s-dependent 12h oscillator and strong transcriptional responses to proteotoxic
stress (17,18,21,79). *Rbbp5* mRNA expression was induced by Tu (**Fig. 6A**), and transient
knockdown of RBBP5 using siRNA (**Fig. S8A**) significantly dampened the UPR of proteostasis
gene expressions in response to Tu, mirroring the effects observed with XBP1s knockdown
(**Fig.6A**). Similarly, knocking down ASH2L also markedly dampened the UPR, whereas knocking
down SETD1A, one of the H3K4me3 methyltransferases in the SET1/COMPASS complex, had
only modest effects, which were likely due to redundancy among the many H3K4me3
methyltransferases that exist (**Fig. S8B**). As a negative control, we further knocked down both
GCN5 and PCAF, proteins that function to write H3K9Ac as subunits in two distinct histone
acetyltransferase (HAT) complexes, SAGA (Spt-Ada-Gcn5 acetyltransferase) and ATAC (Ada2a-
containing) (80-82) and found it had no effects on the induction of *Xbp1* and *Manf* expression in
response to ER stress (**Fig. 6B**). The lack of effects on UPR for HAT subunits is consistent with
the absence of ~12h rhythms of both H3K9Ac and H3K27Ac levels around the promoter of ~12h
proteostasis genes in mouse liver (**Fig. 1A, B**). ChIP-qPCR analysis confirmed a gradual,
temporal increase in the recruitment of RBBP5 to the proximal promoters of select proteostasis
genes in response to ER stress, concomitant with an increase in promoter-proximal H3K4me3
levels (**Fig. 6C**). Since XBP1s is the major TF driving UPR, we further performed IF on XBP1s
and RBBP5 in MEFs and found a strong recruitment of RBBP5 to XBP1s in response to ER stress
(**Fig. 6D, E**).

To broadly assess the function of RBBP5 in the transcription regulation of proteotoxic stress
response, we utilized orthogonal approaches to both deplete RBBP5 expression and trigger
cellular stress. We took advantage of our homozygous *Rbbp5 ^Flox^* mice by crossing them with
homozygous ROSA26 Cre-ERT2 mice, then isolated and immortalized primary MEFs from
homozygous F2 embryos (**Fig. S8C**). Treating these *Rbbp5* (flox/flox); ROSA26 Cre-ERT2 (+/+)
MEFs with tamoxifen for 7 days resulted in the complete deletion of both floxed *Rbbp5* alleles
(**Fig. S8D**) and a subsequent ~90% reduction in RBBP5 protein and a reduction of H3K4me3
level (**Fig. 6F**). The remaining RBBP5 protein is likely extremely long-lived and remains stable
even after the deletion of underlying coding genes. To trigger ER stress, we used a different
stimulus: Dithiothreitol (DTT), which disrupts protein disulfide formation in the ER (83,84). Similar
to Tu, RBBP5 depletion significantly dampened the induction of proteostasis gene expression in
response to DTT (**Fig. 6G**).

Next, we tested two additional proteotoxic stressors that are more physiologically relevant:
leucine deprivation to simulate amino acid restriction, and a more severe nutrient deprivation
stress by culturing cells in Earle's Balanced Salt Solution (EBSS) for 16 hours. Both treatments
not only trigger UPR but also activate the broader integrated stress response (ISR), resulting in
the induction of autophagy (85). RBBP5 depletion impaired the transcriptional response to both
leucine deprivation and EBSS, leading to a reduced induction of autophagy-promoting genes (**Fig.**
7A). Transcriptome profiling by RNA-seq confirmed the globally defective adaptive stress
response to nutrient deprivation (EBSS) in the absence of RBBP5, where only 327 out of 2,351
genes can still be induced (**Figs. 7B-D, S8E, and Table S7, 8**). To investigate the physiological
consequence of RBBP5 deletion, we quantified the autophagic flux by utilizing a tandem LC3
reporter mCherry-GFP-LC3 where an increase in the number of red-fluorescent cytosolic puncta

indicates increased autolysosome formation and autophagic flux (**Fig. 7E**) (85). Consistent with
the qPCR results, RBBP5 depleted MEFs failed to induce a robust general autophagy in response
to leucine deprivation and further exhibited lower basal level of autophagic flux (**Figs. 7F and**
S8F). Western blot analysis of p62/SQSTM1, a marker of autophagic flux due to its degradation
during autophagy (86), showed a significant reduction in wild-type, but not RBBP5 depleted cells
following leucine deprivation, thus confirming the defective autophagy phenotype associated with
RBBP5 depletion (**Fig. S8G, H**). Since prolonged nutrient deprivation, such as 16h of EBSS
treatment, can robustly induce both general autophagy and autophagy of the ER (ER-phagy) (87),
we further measured ER-phagy by utilizing a previously published ER-phagy reporter system
where a stably expressed ER-targeting mCherry-RAMP4 fusion is cleaved to free mCherry during
starvation-induced ER-phagy (29) (**Fig. 7G**). As shown in **Fig. 7H, I**, RBBP5 depletion prevented
the cleavage of mCherry in response to EBSS, indicating an impaired ER-phagy. Western blot
analysis of p62/SQSTM1 further showed undetectable levels in wild-type cells following EBSS
treatment, whereas a significant level remained in RBBP5 depleted cells under the same
conditions, thus confirming a reduction of the general autophagy in RBBP5 depleted cells in
response to EBSS (**Fig. 7H**). Consistent with weakened stress response and impaired autophagy,
RBBP5 depleted MEFs displayed reduced cell survival under nutritional stress (**Fig. S8I**). Taken
together, our results demonstrate that RBBP5 is indispensable for regulating the cell-autonomous
transcriptional responses to various proteotoxic stresses as a core subunit of the H3K4me3-
writing COMPASS complex.

806　Proximity labeling reveals H3K4me-associated proteomic architecture of the proteostatic
807　stress response.

H3K4me3 is widely recognized as an active promoter mark, but its precise role in transcriptional
regulation remains a subject of debate. Initially, it was believed that H3K4me3 facilitated
transcription initiation by recruiting the RNAPII preinitiation complex (88), However, recent
evidence suggests that H3K4me3 primarily regulates RNAPII pause-release and contributes to
transcriptional elongation, likely through the recruitment of Integrator Complex subunit 11 (INTS11)
(58,59). This discrepancy may stem from the context-specific functions of H3K4me3 in recruiting
different transcriptional regulators across different biological processes (89). To shed light on the
potential mechanisms by which H3K4me3 modulates proteostasis stress response, we performed
quantitative TurboID-based proximity labeling in MEFs (in triplicates per condition), utilizing anti-
H3K4me3 antibody (with anti-IgG as negative control) followed by recombinant protein A-TurboID
labeling in isolated nuclei of MEFs before (T=0h) and 2.5h and 8h after ER stress induction by Tu
(**Figs. 8A, S9A**) (47). While the total level of H3K4me3 remains largely unchanged during ER
stress (**Fig. 8B**), we identified a total of 239 proteins (p<0.1) whose recruitment to H3K4me3
peaked at 8h post Tu treatment (**Table S9**). Pathway analysis of these 239 proteins revealed
expected enrichment of COMPASS subunits (**Figs. 8C, S9B, C**), RNAPII subunits (**Fig. S9D**), as
well as previously identified Integrator Complex subunits (such as INTS11) (**Fig. S9E**). The
previously identified H3K4me3 demethylase KDM5A (59) revealed a much more constant
recruitment to H3K4me3 during ER stress, with only a slight increase at 8h after ER stress
induction (**Fig. S9C**).

Besides previously identified H3K4me3 writer/eraser/readers, we also revealed strong
enrichment of SWI/SNF chromatin remodelers on H3K4me3 peaking at 8h after ER stress
induction (**Fig. 8C, D**). These include BAF-specific (ARID1A/B), PBAF-specific (ARID2, PBRM1
and BRD7) and shared subunits (SMARCD1/2/3 and SMARCC2) (**Fig. 8C, D**) (90). The timing of
SWI/SNF recruitment to H3K4me3 during ER stress closely parallels that of COMPASS and
Integrator subunits but lags behind RNAPII subunits (**Fig. 8E**). This temporal pattern suggests
that SWI/SNF might potentially play a role in facilitating transcriptional elongation during stress-
induced gene activation.

To uncover further evidence supporting the recruitment of SWI/SNF to H3K4me3 during proteostatic stress, we turned back to the hepatic 12h oscillator and examined the temporal expression and cistrome of SWI/SNF subunits in mouse liver. Intriguingly, even before our discovery of the hepatic 12h oscillator in 2017, we had previously reported the 12h rhythmic expression of nuclear PBRM1 (BAF180), ARID2 (BAF200), and SMARCA4 (BRG1) proteins in wild-type mouse liver (36), though at the time, we were unaware of their potential implications in proteostasis control. To validate the hepatic 12h expression of nuclear PBRM1 (BAF180), we performed western blot analysis and confirmed its rhythmic peaks at CT10 and CT22 (**Fig. S10A**). Next, we conducted a post-hoc analysis of our previously published hepatic BAF180 ChIP-seq datasets at CT22, using two different antibodies (Bethyl and Active Motif) (35), and compared them with RBBP5 and H3K4me3 cistromes. Globally, genes with promoter-proximal BAF180 binding (±1 kb from the TSS) were significantly enriched in proteostasis-related pathways, including ER stress, autophagy, and proteolysis (**Fig. 8F, G**). Concordantly, we observed co-enrichment of BAF180, RBBP5, and H3K4me3 at the promoter regions of the 6,451 genes identified in **Fig. 1C,** with a strong positive correlation between their binding intensities on these genes (**Figs. 8H, S10B, C**). Although the causal relationship between SWI/SNF subunits and proteostasis gene expression remains to be determined, our collective findings in both MEFs and liver establish both COMPASS and SWI/SNF as critical components of the chromatin landscape governing dynamic proteostasis gene regulation.

Reduced RBBP5 expression is associated with aging in mice and hepatic steatosis in humans.

The roles of RBBP5 in diseases are incompletely understood (56,91). Since the decline of proteostasis and stress response is causally associated with both aging and metabolic diseases (92-95), we aim to uncover potential implications of RBBP5 in these two processes. Intriguingly, in a recent study, RBBP5 (and DPY30) is predicted to be a positive regulator of maximum lifespan across 26 different species, with its major effect predicted to be related to liver (96). Consistent with this prediction, an overall reduction of *Rbbp5* mRNA level is observed in the liver of aging mice (97), largely concordant with the expression dynamics of 97 stress response genes (those genes exhibiting 12h expression, directly bound by RBBP5 and can be induced by ER stress) (**Fig. 9A**).

Leveraging human gene-derived correlations across tissues (GD-CAT) dataset (98), we identified in human liver, RBBP5 expression is positively correlated with those involved in proteostasis (such as ribosome biogenesis, translation regulation) and mRNA metabolism (mRNA splicing and processing) and negatively associated with those implicated in immune response (acute phase response, complement activity, and immunoglobin activity) (**Fig. S11**). These results are in line with our RNA-seq and ChIP-seq data obtained from *Rbbp5 Flox* and *Rbbp5 LKO* mice (**Figs. 2C, 3I, 4D**). In humans with NAFLD/MAFLD (99), compared to healthy obese, RBBP5 mRNA level is significantly reduced in steatosis subjects (**Fig. 9B**), concordant with reduced stress gene expression in the same cohorts (**Fig. 9C-F**). Overall, a positive correlation between *Rbbp5* and the eigengene expression of stress response genes [eigengene (100) is the principle component of 97 stress response genes expression in each subject] is observed across all subjects (**Fig. 9D**). Collectively, these data suggest that the downregulation of *Rbbp5* and proteostatic stress response genes expression is associated with aging and MAFLD in mice and humans, respectively.

Discussion:

While the epigenetic landscape of the mammalian ~24h circadian clock was well-studied (54,101-104), the epigenome of the mammalian ~12h ultradian oscillator remains completely unexplored. In this study, we uncovered that unlike the circadian clock, which was predominantly regulated by

histone acetyltransferase (HAT) (35,105,106) and histone deacetylase (HDAC) (103,107,108), the epigenome of the 12h oscillator is marked by COMPASS-mediated histone methylation, specifically H3K4me3 at the proximal promoters. Loss of one of the essential subunits of COMPASS, RBBP5, abolished more than 90% of the hepatic ~12h transcriptome, while having no effects on the core circadian clock. Knocking down RBBP5 in MEFs further dampened the cell-autonomous 12h oscillations, but not the circadian rhythms. We further expanded our findings to the epigenetic regulation of response to acute and chronic proteotoxic stress and found both increased RBBP5 promoter recruitment and H3K4me3 level are associated with the transcription activation of adaptive stress response genes. Subsequently, depletion of RBBP5 significantly dampened the adaptive transcriptional responses to various proteotoxic stresses.

In our study, we uncovered distinct roles for H3K4me3 and H3K9Ac/H3K27Ac in regulating the 12h oscillator and transcriptional responses to acute proteotoxic stress. While all three histone modifications—H3K4me3, H3K9Ac, and H3K27Ac—are traditionally linked to transcriptional activation (81,82,109), our findings suggest that H3K9Ac and H3K27Ac are dispensable for the transcriptional regulation of proteostasis under both basal (the 12h oscillator) and stress conditions. Interestingly, recent research by (110) demonstrated that among various epigenetic marks associated with transcriptional activation, the installation of H3K4me3 at promoters resulted in the most potent transcriptional activation. This not only underscores the causal role of H3K4me3 in gene activation but also highlights the differential impacts of specific histone modifications on transcription. We speculate that among the many histone modifications, H3K4me3 has been evolutionarily selected as the primary epigenetic modification for stress response due to its ability to trigger the most robust gene activation, thereby providing a rapid adaptive advantage under adverse environmental conditions. Future investigations are needed to further test this hypothesis.

The precise role of H3K4me3 in transcriptional regulation remains a topic of debate. Recent evidence suggests that H3K4me3 positively contributes to transcriptional elongation by regulating the pause-release of RNAP II, likely through the recruitment of the Integrator subunit INST11 (89). Our H3K4me3 proximity labeling not only supports this finding but also unexpectedly reveals increased recruitment of the SWI/SNF complex to H3K4me3 during the UPR in MEFs. Additionally, we confirmed the co-occurrence of RBBP5, H3K4me3, and the PBAF subunit BAF180 in chromatin binding in vivo in mouse liver. Interestingly, the temporal dynamics of SWI/SNF recruitment to H3K4me3 coincide with the recruitment of the Integrator complex but occur after the initial increase in RNAP II subunits, which are likely involved in transcription initiation immediately following stress. This suggests that contrary to the widely accepted view that SWI/SNF primarily facilitates transcription initiation by increasing chromatin accessibility and transcription factor binding to promoter, it may also play a crucial yet underexplored role in transcription elongation. Although the literature on this aspect is rather limited, some earlier studies hint at this possibility (111,112). Future research is needed to firmly establish a causal relationship between SWI/SNF function, 12h oscillator regulation, and the proteostasis stress response, as well as to uncover the underlying molecular mechanisms.

The physiological and pathological roles of mammalian COMPASS subunits, particularly RBBP5, remain incompletely understood (91). While extensive research has primarily focused on their contributions to organogenesis, early development, and neurodevelopmental disorders, our study significantly broadens the scope of RBBP5's physiological roles by uncovering its involvement in proteostasis and hepatic metabolism. This expansion not only advances our understanding of RBBP5's function but also opens new avenues for exploring its broader impact on cellular and systemic processes. Of particular interest is a recent study that identified heterozygous loss-of-function mutations in *RBBP5* in humans, which were primarily linked to neurodevelopmental disorders (113). This finding raises intriguing questions about the full

spectrum of RBBP5's physiological roles. Given our discovery of RBBP5's involvement in metabolic regulation, it is tempting to hypothesize that individuals with RBBP5 loss-of-function mutations may also exhibit metabolic defects. Such a connection would have significant implications, suggesting that RBBP5 plays a multifaceted role in maintaining both neural and metabolic homeostasis. Future research should investigate whether these mutations influence metabolic pathways, potentially contributing to a broader spectrum of clinical symptoms in affected individuals. This line of inquiry could ultimately lead to a deeper understanding of how COMPASS subunits like RBBP5 integrate signals across different tissues to maintain overall physiological balance.

Finally, we propose that the 12h oscillator serves as an effective 'discovery tool' for uncovering previously unknown mechanisms of proteostasis regulation. By leveraging this approach, we recently identified a novel XBP1s-SON axis, which links nuclear speckle LLPS dynamics with the UPR (26). Herein, using a similar approach, we identified RBBP5 as an epigenetic regulator of proteostasis. We believe that continued exploration of the 12h oscillator will unveil further hidden principles governing proteostasis.

Acknowledgements:

We would like to thank Dr. Sebastien Gingras (University of Pittsburgh, Department of Immunology) for the design of the targeting strategy and identification of the founders carrying the properly targeted Rbbp5-flox allele. We would like to thank Chunming Bi and Zhaohui Kou of the Mouse Embryo Services Core (University of Pittsburgh, Department of Immunology) for microinjection of zygotes and production of Rbbp5-flox founder mice. We would also like to thank Dr. Heather Ballance, Dr. Naveen GV Kumar, Saad Irfan, C. Dupont, Dr. Bugra Zengin and Casey Edwards for assistance with mice colony maintenance, dissection, intraperitoneal injection and genotyping. We would also like to thank Dr. Michiel Vermeulen for kindly sharing the pK19HispATurbo plasmid. For time course transcriptome study, the library generation was performed in the Health Sciences Sequencing Core at UPMC Children's Hospital of Pittsburgh, Rangos Research Center. Services and instruments used in this project were graciously supported, in part, by the University of Pittsburgh, the Office of the Senior Vice Chancellor for Health Sciences, the Department of Pediatrics, the Institute for Precision Medicine, and the Richard K Mellon Foundation for Pediatric Research.

W.D. was supported by fellowship F31 AG080998 through NIA. B. Zhu was supported by grants 1DP2GM140924 and 1R21AG071893 through the NIH, and a grant from Richard King Mellon foundation. This research was supported in part by the University of Pittsburgh Center for Research Computing through the resources provided. Specifically, this work used the HTC cluster, which is supported by NIH award number S10OD028483. This research project was supported in part by the Pittsburgh Liver Research Centre supported by NIH/NIDDK Digestive Disease Research Core Center grant P30DK120531.

Author contributions:

Conceptualization, B.Z.; Methodology, M.J. and B.Z.; Investigation, S.K., M.S., W.D., A.C., H.L., H.W., A.P., I.S., S.L., Y.P., L.L., J.J.L., and B.Z.; Writing – Original Draft, B.Z.; Writing – Review & Editing, all authors; Funding Acquisition, S.L., and B.Z.; Resources, M.J., J.H.L and B.Z.; Supervision, M.J., J.H.L and B.Z.

Disclosure and competing interests:

The authors have no competing interests to declare.

Data availability:

All raw and processed sequencing data generated in this study have been submitted to the NCBI
Gene Expression Omnibus (GEO; http://www.ncbi.nlm.nih.gov/geo/) under accession numbers:
GSE276155, GSE276157 and GSE276159. All data needed to evaluate the conclusions in the
paper are present in the paper and/or the Supplementary Materials.

References:

1.   Labbadia, J. and Morimoto, R.I. (2015) The Biology of Proteostasis in Aging and
Disease. *Annual Review of Biochemistry*, **84**, 435-464.

2.   Buttgereit, F. and Brand, M.D. (1995) A hierarchy of ATP-consuming processes in
mammalian cells. *Biochemical Journal*, **312**, 163-167.

3.   Zhao, L. and Ackerman, S.L. (2006) Endoplasmic reticulum stress in health and disease.
Current Opinion in Cell Biology, **18**, 444-452.

4.   Klaips, C.L., Jayaraj, G.G. and Hartl, F.U. (2018) Pathways of cellular proteostasis in
aging and disease. *J Cell Biol*, **217**, 51-63.

5.   Martinez, G., Duran-Aniotz, C., Cabral-Miranda, F., Vivar, J.P. and Hetz, C. (2017)
Endoplasmic reticulum proteostasis impairment in aging. *Aging Cell*, **16**, 615-623.

6.   Taylor, R.C. and Hetz, C. (2020) Mastering organismal aging through the endoplasmic
reticulum proteostasis network. *Aging Cell*, **19**, e13265.

7.   Costa-Mattioli, M. and Walter, P. (2020) The integrated stress response: From
mechanism to disease. *Science*, **368**, eaat5314.

8.   Hetz, C., Zhang, K. and Kaufman, R.J. (2020) Mechanisms, regulation and functions of
the unfolded protein response. *Nature Reviews Molecular Cell Biology*, **21**, 421-438.

9.   Richter, K., Haslbeck, M. and Buchner, J. (2010) The heat shock response: life on the
verge of death. *Mol Cell*, **40**, 253-266.

10.   Ron, D. and Walter, P. (2007) Signal integration in the endoplasmic reticulum unfolded
protein response. *Nature Reviews Molecular Cell Biology*, **8**, 519-529.

11.   Balch, W.E., Morimoto, R.I., Dillin, A. and Kelly, J.W. (2008) Adapting proteostasis for
disease intervention. *Science*, **319**, 916-919.

12.   Ballance, H. and Zhu, B. (2021) Revealing the hidden reality of the mammalian 12-h
ultradian rhythms. *Cellular and Molecular Life Sciences*.

13.   Asher, G. and Zhu, B. (2022) Beyond circadian rhythms: Emerging roles of ultradian
rhythms in control of liver functions. *Hepatology*, **n/a**.

14.   Scott, M.R., Zong, W., Ketchesin, K.D., Seney, M.L., Tseng, G.C., Zhu, B. and McClung,
C.A. (2023) Twelve-hour rhythms in transcript expression within the human dorsolateral
prefrontal cortex are altered in schizophrenia. *PLOS Biology*, **21**, e3001688.

15. Zhu, B., Liu, S., David, N.L., Dion, W., Doshi, N.K., Siegel, L.B., Amorim, T., Andrews,
R.E., Kumar, G.N., Irfan, S. *et al.* (2023) Evidence for conservation of a primordial 12-
hour ultradian gene program in humans. *bioRxiv*, 2023.2005.2002.539021.

16. Zhu, B., Liu, S., David, N.L., Dion, W., Doshi, N.K., Siegel, L.B., Amorim, T., Andrews,
R.E., Kumar, G.V.N., Li, H. *et al.* (2024) Evidence for ~12-h ultradian gene programs in
humans. *npj Biological Timing and Sleep*, **1**, 4.

17. Dion, W., Ballance, H., Lee, J., Pan, Y., Irfan, S., Edwards, C., Sun, M., Zhang, J.,
Zhang, X. and Liu, S. (2022) Four-dimensional nuclear speckle phase separation
dynamics regulate proteostasis. *Science Advances*, **8**, eabl4150.

18. Zhu, B., Zhang, Q., Pan, Y., Mace, E.M., York, B., Antoulas, A.C., Dacso, C.C. and
O'Malley, B.W. (2017) A cell-autonomous mammalian 12 hr clock coordinates metabolic
and stress rhythms. *Cell metabolism*, **25**, 1305-1319. e1309.

19. Meng, H., Gonzales, N.M., Lonard, D.M., Putluri, N., Zhu, B., Dacso, C.C., York, B. and
O'Malley, B.W. (2020) XBP1 links the 12-hour clock to NAFLD and regulation of
membrane fluidity and lipid homeostasis. *Nature Communications*, **11**, 6215.

20. Meng, H., Gonzales, N.M., Jung, S.Y., Lu, Y., Putluri, N., Zhu, B., Dacso, C.C., Lonard,
D.M. and O'Malley, B.W. (2022) Defining the mammalian coactivation of hepatic 12-h
clock and lipid metabolism. *Cell Reports*, **38**, 110491.

21. Pan, Y., Ballance, H., Meng, H., Gonzalez, N., Kim, S.-M., Abdurehman, L., York, B.,
Chen, X., Schnytzer, Y., Levy, O. *et al.* (2020) 12-h clock regulation of genetic
information flow by XBP1s. *PLOS Biology*, **18**, e3000580.

22. Asher, G. and Zhu, B. (2022) Beyond circadian rhythms: Emerging roles of ultradian
rhythms in control of liver functions. *Hepatology*.

23. Zhu, B., Zhang, Q., Pan, Y., Mace, E.M., York, B., Antoulas, A.C., Dacso, C.C. and
O'Malley, B.W. (2017) A Cell-Autonomous Mammalian 12 hr Clock Coordinates
Metabolic and Stress Rhythms. *Cell Metab*, **25**, 1305-1319 e1309.

24. Zhu, B. and Liu, S. (2023) Preservation of ~12-hour ultradian rhythms of gene
expression of mRNA and protein metabolism in the absence of canonical circadian
clock. *bioRxiv*, 2023.2005.2001.538977.

25. Zhang, H., Yahagi, T., Miyamoto, N., Chen, C., Jiang, Q., Qian, P.-Y. and Sun, J. (2025)
Circatidal control of gene expression in the deep-sea hot vent shrimp <i>Rimicaris
leurokolos</i>. *Proceedings of the Royal Society B: Biological Sciences*, **292**, 20242970.

26. Dion, W., Ballance, H., Lee, J., Pan, Y., Irfan, S., Edwards, C., Sun, M., Zhang, J.,
Zhang, X., Liu, S. *et al.* (2022) Four-dimensional nuclear speckle phase separation
dynamics regulate proteostasis. *Science Advances*, **8**, eabl4150.

27. Shilatifard, A. (2012) The COMPASS family of histone H3K4 methylases: mechanisms of
regulation in development and disease pathogenesis. *Annu Rev Biochem*, **81**, 65-95.

28.   Wulansari, N., Darsono, W.H.W., Woo, H.J., Chang, M.Y., Kim, J., Bae, E.J., Sun, W.,
Lee, J.H., Cho, I.J., Shin, H. *et al.* (2021) Neurodevelopmental defects and
neurodegenerative phenotypes in human brain organoids carrying Parkinson's disease-
linked DNAJC6 mutations. *Sci Adv*, **7**.

29.   Liang, J.R., Lingeman, E., Ahmed, S. and Corn, J.E. (2018) Atlastins remodel the
endoplasmic reticulum for selective autophagy. *J Cell Biol*, **217**, 3354-3367.

30.   Quadros, R.M., Miura, H., Harms, D.W., Akatsuka, H., Sato, T., Aida, T., Redder, R.,
Richardson, G.P., Inagaki, Y., Sakai, D. *et al.* (2017) Easi-CRISPR: a robust method for
one-step generation of mice carrying conditional and insertion alleles using long ssDNA
donors and CRISPR ribonucleoproteins. *Genome Biol*, **18**, 92.

31.   Pelletier, S., Gingras, S. and Green, D.R. (2015) Mouse genome engineering via
CRISPR-Cas9 for study of immune function. *Immunity*, **42**, 18-27.

32.   Xu, J. (2005) Preparation, culture, and immortalization of mouse embryonic fibroblasts.
Curr Protoc Mol Biol, **Chapter 28**, Unit 28.21.

33.   Zhu, B., Gates, L.A., Stashi, E., Dasgupta, S., Gonzales, N., Dean, A., Dacso, C.C.,
York, B. and O'Malley, B.W. (2015) Coactivator-Dependent Oscillation of Chromatin
Accessibility Dictates Circadian Gene Amplitude via REV-ERB Loading. *Mol Cell*.

34.   Malovannaya, A., Lanz, R.B., Jung, S.Y., Bulynko, Y., Le, N.T., Chan, D.W., Ding, C., Shi,
Y., Yucer, N., Krenciute, G. *et al.* (2011) Analysis of the human endogenous coregulator
complexome. *Cell*, **145**, 787-799.

35.   Zhu, B., Gates, L.A., Stashi, E., Dasgupta, S., Gonzales, N., Dean, A., Dacso, C.C.,
York, B. and O'Malley, B.W. (2015) Coactivator-Dependent Oscillation of Chromatin
Accessibility Dictates Circadian Gene Amplitude via REV-ERB Loading. *Mol Cell*, **60**,
769-783.

36.   Zhu, B., Gates, L.A., Stashi, E., Dasgupta, S., Gonzales, N., Dean, A., Dacso, C.C.,
York, B. and O'Malley, B.W. (2015) Coactivator-dependent oscillation of chromatin
accessibility dictates circadian gene amplitude via REV-ERB loading. *Molecular cell*, **60**,
769-783.

37.   Bolger, A.M., Lohse, M. and Usadel, B. (2014) Trimmomatic: a flexible trimmer for
Illumina sequence data. *Bioinformatics*, **30**, 2114-2120.

38.   Kim, D., Langmead, B. and Salzberg, S.L. (2015) HISAT: a fast spliced aligner with low
memory requirements. *Nat Methods*, **12**, 357-360.

39.   Antoulas, A.C., Zhu, B., Zhang, Q., York, B., O'Malley, B.W. and Dacso, C.C. (2018) A
novel mathematical method for disclosing oscillations in gene transcription: A
comparative study. *PLoS One*, **13**, e0198503.

40.   Thaben, P.F. and Westermark, P.O. (2014) Detecting rhythms in time series with RAIN. *J
Biol Rhythms*, **29**, 391-400.

41. Liu, S., Obert, C., Yu, Y.P., Zhao, J., Ren, B.G., Liu, J.J., Wiseman, K., Krajacich, B.J.,
Wang, W., Metcalfe, K. *et al.* (2024) Utility analyses of AVITI sequencing chemistry. *BMC*
Genomics, **25**, 778.

42. Bolger, A.M., Lohse, M. and Usadel, B. (2014) Trimmomatic: a flexible trimmer for
Illumina sequence data. *Bioinformatics*, **30**, 2114-2120.

43. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M. and Gingeras, T.R. (2013) STAR: ultrafast universal RNA-seq aligner.
Bioinformatics, **29**, 15-21.

44. Trapnell, C., Williams, B.A., Pertea, G., Mortazavi, A., Kwan, G., van Baren, M.J.,
Salzberg, S.L., Wold, B.J. and Pachter, L. (2010) Transcript assembly and quantification
by RNA-Seq reveals unannotated transcripts and isoform switching during cell
differentiation. *Nature Biotechnology*, **28**, 511-U174.

45. Ge, S.X., Son, E.W. and Yao, R. (2018) iDEP: an integrated web application for
differential expression and pathway analysis of RNA-Seq data. *BMC bioinformatics*, **19**,
1-24.

46. Ramirez, F., Ryan, D.P., Gruning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne, S.,
Dundar, F. and Manke, T. (2016) deepTools2: a next generation web server for deep-
sequencing data analysis. *Nucleic Acids Res*, **44**, W160-165.

47. Santos-Barriopedro, I., van Mierlo, G. and Vermeulen, M. (2021) Off-the-shelf proximity
biotinylation for interaction proteomics. *Nature Communications*, **12**, 5015.

48. Huang da, W., Sherman, B.T. and Lempicki, R.A. (2009) Systematic and integrative
analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc*, **4**, 44-57.

49. He, H.H., Meyer, C.A., Shin, H., Bailey, S.T., Wei, G., Wang, Q., Zhang, Y., Xu, K., Ni,
M., Lupien, M. *et al.* (2010) Nucleosome dynamics define transcriptional enhancers.
Nature Genetics, **42**, 343-347.

50. Zhu, B. (2020) Decoding the function and regulation of the mammalian 12h-clock.
Journal of Molecular Cell Biology.

51. Dasgupta, S., Lonard, D.M. and O'Malley, B.W. (2014) Nuclear receptor coactivators:
master regulators of human health and disease. *Annu Rev Med*, **65**, 279-292.

52. Roeder, R.G. (2005) Transcriptional regulation and the role of diverse coactivators in
animal cells. *FEBS Lett*, **579**, 909-915.

53. Spiegelman, B.M. and Heinrich, R. (2004) Biological control through regulated
transcriptional coactivators. *Cell*, **119**, 157-167.

54. Koike, N., Yoo, S.H., Huang, H.C., Kumar, V., Lee, C., Kim, T.K. and Takahashi, J.S.
(2012) Transcriptional architecture and chromatin landscape of the core circadian clock
in mammals. *Science*, **338**, 349-354.

55. Takahashi, Y.H., Westfield, G.H., Oleskie, A.N., Trievel, R.C., Shilatifard, A. and Skiniotis,
G. (2011) Structural analysis of the core COMPASS family of histone H3K4 methylases
from yeast to human. *Proc Natl Acad Sci U S A*, **108**, 20526-20531.

56. Shilatifard, A. (2012) The COMPASS family of histone H3K4 methylases: mechanisms of
regulation in development and disease pathogenesis. *Annual review of biochemistry*, **81**,
65-95.

57. Qu, Q., Takahashi, Y.H., Yang, Y., Hu, H., Zhang, Y., Brunzelle, J.S., Couture, J.F.,
Shilatifard, A. and Skiniotis, G. (2018) Structure and Conformational Dynamics of a
COMPASS Histone H3K4 Methyltransferase Complex. *Cell*, **174**, 1117-1126.e1112.

58. Hu, S., Song, A., Peng, L., Tang, N., Qiao, Z., Wang, Z., Lan, F. and Chen, F.X. (2023)
H3K4me2/3 modulate the stability of RNA polymerase II pausing. *Cell Research*, **33**,
403-406.

59. Wang, H., Fan, Z., Shliaha, P.V., Miele, M., Hendrickson, R.C., Jiang, X. and Helin, K.
(2023) H3K4me3 regulates RNA polymerase II promoter-proximal pause-release.
Nature, **615**, 339-348.

60. Qin, Q., Fan, J., Zheng, R., Wan, C., Mei, S., Wu, Q., Sun, H., Brown, M., Zhang, J.,
Meyer, C.A. *et al.* (2020) Lisa: inferring transcriptional regulators through integrative
modeling of public chromatin accessibility and ChIP-seq data. *Genome Biology*, **21**, 32.

61. Li, Y., Han, J., Zhang, Y., Cao, F., Liu, Z., Li, S., Wu, J., Hu, C., Wang, Y., Shuai, J. *et al.*
(2016) Structural basis for activity regulation of MLL family methyltransferases. *Nature*,
530, 447-452.

62. Ruthenburg, A.J., Allis, C.D. and Wysocka, J. (2007) Methylation of lysine 4 on histone
H3: intricacy of writing and reading a single epigenetic mark. *Mol Cell*, **25**, 15-30.

63. Han, J., Li, T., Li, Y., Li, M., Wang, X., Peng, C., Su, C., Li, N., Li, Y., Xu, Y. *et al.* (2019)
The internal interaction in RBBP5 regulates assembly and activity of MLL1
methyltransferase complex. *Nucleic Acids Research*, **47**, 10426-10438.

64. Antoulas, A.C., Zhu, B., Zhang, Q., York, B., O'Malley, B.W. and Dacso, C.C. (2018) A
novel mathematical method for disclosing oscillations in gene transcription: a
comparative study. *PLoS One*, **13**, e0198503.

65. Zhu, B. and Liu, S. (2023) Preservation of ~12-h ultradian rhythms of gene expression of
mRNA and protein metabolism in the absence of canonical circadian clock. *Frontiers in
Physiology*, **14**.

66. Hughes, M.E., DiTacchio, L., Hayes, K.R., Vollmers, C., Pulivarthy, S., Baggs, J.E.,
Panda, S. and Hogenesch, J.B. (2009) Harmonics of circadian gene transcription in
mammals. *PLoS Genet*, **5**, e1000442.

67. van der Veen, D.R. and Gerkema, M.P. (2017) Unmasking ultradian rhythms in gene
expression. *The FASEB Journal*, **31**, 743-750.

68. Zhang, E.E., Liu, A.C., Hirota, T., Miraglia, L.J., Welch, G., Pongsawakul, P.Y., Liu, X.,
Atwood, A., Huss, J.W., 3rd, Janes, J. *et al.* (2009) A genome-wide RNAi screen for
modifiers of the circadian clock in human cells. *Cell*, **139**, 199-210.

69. Walter, P. and Ron, D. (2011) The unfolded protein response: from stress pathway to
homeostatic regulation. *Science*, **334**, 1081-1086.

70. Metcalf, M.G., Higuchi-Sanabria, R., Garcia, G., Tsui, C.K. and Dillin, A. (2020) Beyond
the cell factory: Homeostatic regulation of and by the UPR^{ER}. *Science*
Advances, **6**, eabb9614.

71. Wiseman, R.L., Mesgarzadeh, J.S. and Hendershot, L.M. (2022) Reshaping
endoplasmic reticulum quality control through the unfolded protein response. *Molecular*
Cell, **82**, 1477-1491.

72. So, J.S. (2018) Roles of Endoplasmic Reticulum Stress in Immune Responses. *Mol*
Cells, **41**, 705-716.

73. Rutkowski, D.T., Wu, J., Back, S.H., Callaghan, M.U., Ferris, S.P., Iqbal, J., Clark, R.,
Miao, H., Hassler, J.R., Fornek, J. *et al.* (2008) UPR pathways combine to prevent
hepatic steatosis caused by ER stress-mediated suppression of transcriptional master
regulators. *Dev Cell*, **15**, 829-840.

74. Yan, Z., Yan, H. and Ou, H. (2012) Human thyroxine binding globulin (TBG) promoter
directs efficient and sustaining transgene expression in liver-specific pattern. *Gene*, **506**,
289-294.

75. Guo, B. and Li, Z. (2014) Endoplasmic reticulum stress in hepatic steatosis and
inflammatory bowel diseases. *Frontiers in Genetics*, **5**.

76. Ajoolabady, A., Kaplowitz, N., Lebeaupin, C., Kroemer, G., Kaufman, R.J., Malhi, H. and
Ren, J. (2023) Endoplasmic reticulum stress in liver diseases. *Hepatology*, **77**, 619-639.

77. Gong, H., He, Q., Zhu, L., Feng, Z., Sun, M., Jiang, J., Yuan, X., Shen, Y. and Di, J.
(2024) Associations between systemic inflammation indicators and nonalcoholic fatty
liver disease: evidence from a prospective study. *Frontiers in Immunology*, **15**.

78. Gehrke, N. and Schattenberg, J.M. (2020) Metabolic Inflammation—A Role for Hepatic
Inflammatory Pathways as Drivers of Comorbidities in Nonalcoholic Fatty Liver Disease?
Gastroenterology, **158**, 1929-1947.e1926.

79. Dion, W., Tao, Y., Chambers, M., Zhao, S., Arbuckle, R.K., Sun, M., Kubra, S., Nie, Y.,
Ye, M., Larsen, M.B. *et al.* (2024) Nuclear speckle rejuvenation alleviates
proteinopathies at the expense of YAP1. *bioRxiv*, 2024.2004.2018.590103.

80. Jin, Q., Yu, L.R., Wang, L., Zhang, Z., Kasper, L.H., Lee, J.E., Wang, C., Brindle, P.K.,
Dent, S.Y. and Ge, K. (2011) Distinct roles of GCN5/PCAF-mediated H3K9ac and
CBP/p300-mediated H3K18/27ac in nuclear receptor transactivation. *Embo j*, **30**, 249-
262.

81.   Gates, L.A., Foulds, C.E. and O'Malley, B.W. (2017) Histone Marks in the 'Driver's Seat':
Functional Roles in Steering the Transcription Cycle. *Trends Biochem Sci*, **42**, 977-989.

82.   Gates, L.A., Shi, J., Rohira, A.D., Feng, Q., Zhu, B., Bedford, M.T., Sagum, C.A., Jung,
S.Y., Qin, J., Tsai, M.J. *et al.* (2017) Acetylation on histone H3 lysine 9 mediates a switch
from transcription initiation to elongation. *J Biol Chem*, **292**, 14456-14472.

83.   Higa, A., Taouji, S., Lhomond, S., Jensen, D., Fernandez-Zapico, M.E., Simpson, J.C.,
Pasquet, J.M., Schekman, R. and Chevet, E. (2014) Endoplasmic reticulum stress-
activated transcription factor ATF6α requires the disulfide isomerase PDIA5 to modulate
chemoresistance. *Mol Cell Biol*, **34**, 1839-1849.

84.   G, G. and Singh, J. (2022) Dithiothreitol causes toxicity in C. elegans by modulating the
methionine–homocysteine cycle. *eLife*, **11**, e76021.

85.   Pakos-Zebrucka, K., Koryga, I., Mnich, K., Ljujic, M., Samali, A. and Gorman, A.M.
(2016) The integrated stress response. *EMBO Rep*, **17**, 1374-1395.

86.   Bjørkøy, G., Lamark, T., Pankiv, S., Øvervatn, A., Brech, A. and Johansen, T. (2009)
Monitoring autophagic degradation of p62/SQSTM1. *Methods Enzymol*, **452**, 181-197.

87.   Liang, J.R., Lingeman, E., Luong, T., Ahmed, S., Muhar, M., Nguyen, T., Olzmann, J.A.
and Corn, J.E. (2020) A Genome-wide ER-phagy Screen Highlights Key Roles of
Mitochondrial Metabolism and ER-Resident UFMylation. *Cell*, **180**, 1160-1177.e1120.

88.   Gates, L.A., Shi, J., Rohira, A.D., Feng, Q., Zhu, B., Bedford, M.T., Sagum, C.A., Jung,
S.Y., Qin, J. and Tsai, M.-J. (2017) Acetylation on histone H3 lysine 9 mediates a switch
from transcription initiation to elongation. *Journal of Biological Chemistry*, **292**, 14456-
14472.

89.   Wang, H. and Helin, K. (2025) Roles of H3K4 methylation in biology and disease. *Trends
in Cell Biology*, **35**, 115-128.

90.   Hodges, C., Kirkland, J.G. and Crabtree, G.R. (2016) The Many Roles of BAF
(mSWI/SNF) and PBAF Complexes in Cancer. *Cold Spring Harb Perspect Med*, **6**.

91.   Cenik, B.K. and Shilatifard, A. (2021) COMPASS and SWI/SNF complexes in
development and disease. *Nature Reviews Genetics*, **22**, 38-58.

92.   Ghemrawi, R., Battaglia-Hsu, S.F. and Arnold, C. (2018) Endoplasmic Reticulum Stress
in Metabolic Disorders. *Cells*, **7**.

93.   Mohan, S., R, P.R.M., Brown, L., Ayyappan, P. and G, R.K. (2019) Endoplasmic
reticulum stress: A master regulator of metabolic syndrome. *European Journal of
Pharmacology*, **860**, 172553.

94.   Kourtis, N. and Tavernarakis, N. (2011) Cellular stress response pathways and ageing:
intricate molecular relationships. *The EMBO journal*, **30**, 2520-2531.

95. Chen, K., Shen, W., Zhang, Z., Xiong, F., Ouyang, Q. and Luo, C. (2020) Age-dependent
decline in stress response capacity revealed by proteins dynamics analysis. *Scientific*
Reports, **10**, 15211.

96. Lu, J.Y., Simon, M., Zhao, Y., Ablaeva, J., Corson, N., Choi, Y., Yamada, K.Y.H., Schork,
N.J., Hood, W.R., Hill, G.E. *et al.* (2022) Comparative transcriptomics reveals circadian
and pluripotency networks as two pillars of longevity regulation. *Cell Metabolism*, **34**,
836-856.e835.

97. Almanzar, N., Antony, J., Baghel, A.S., Bakerman, I., Bansal, I., Barres, B.A., Beachy,
P.A., Berdnik, D., Bilen, B., Brownfield, D. *et al.* (2020) A single-cell transcriptomic atlas
characterizes ageing tissues in the mouse. *Nature*, **583**, 590-595.

98. Zhou, M., Tamburini, I., Van, C., Molendijk, J., Nguyen, C.M., Chang, I.Y.-Y., Johnson,
C., Velez, L.M., Cheon, Y., Yeo, R. *et al.* (2024) Leveraging inter-individual transcriptional
correlation structure to infer discrete signaling mechanisms across metabolic tissues.
eLife, **12**, RP88863.

99. Ahrens, M., Ammerpohl, O., von Schönfels, W., Kolarova, J., Bens, S., Itzel, T., Teufel,
A., Herrmann, A., Brosch, M., Hinrichsen, H. *et al.* (2013) DNA Methylation Analysis in
Nonalcoholic Fatty Liver Disease Suggests Distinct Disease-Specific and Remodeling
Signatures after Bariatric Surgery. *Cell Metabolism*, **18**, 296-302.

100. Langfelder, P. and Horvath, S. (2007) Eigengene networks for studying the relationships
between co-expression modules. *BMC Systems Biology*, **1**, 54.

101. Masri, S., Kinouchi, K. and Sassone-Corsi, P. (2015) Circadian clocks, epigenetics, and
cancer. *Curr Opin Oncol*, **27**, 50-56.

102. Takahashi, J.S. (2017) Transcriptional architecture of the mammalian circadian clock.
Nat Rev Genet, **18**, 164-179.

103. Nakahata, Y., Kaluzova, M., Grimaldi, B., Sahar, S., Hirayama, J., Chen, D., Guarente,
L.P. and Sassone-Corsi, P. (2008) The NAD^+-Dependent Deacetylase
SIRT1 Modulates CLOCK-Mediated Chromatin Remodeling and Circadian Control. *Cell*,
134, 329-340.

104. Aguilar-Arnal, L. and Sassone-Corsi, P. (2015) Chromatin landscape and circadian
dynamics: Spatial and temporal organization of clock transcription. *Proceedings of the*
National Academy of Sciences, **112**, 6863-6870.

105. Doi, M., Hirayama, J. and Sassone-Corsi, P. (2006) Circadian Regulator CLOCK Is a
Histone Acetyltransferase. *Cell*, **125**, 497-508.

106. Stashi, E., Lanz, R.B., Mao, J., Michailidis, G., Zhu, B., Kettner, N.M., Putluri, N.,
Reineke, E.L., Reineke, L.C. and Dasgupta, S. (2014) SRC-2 is an essential coactivator
for orchestrating metabolism and circadian rhythm. *Cell reports*, **6**, 633-645.

107. Feng, D., Liu, T., Sun, Z., Bugge, A., Mullican, S.E., Alenghat, T., Liu, X.S. and Lazar,
M.A. (2011) A circadian rhythm orchestrated by histone deacetylase 3 controls hepatic
lipid metabolism. *Science*, **331**, 1315-1319.

108. Alenghat, T., Meyers, K., Mullican, S.E., Leitner, K., Adeniji-Adele, A., Avila, J., Bućan,
M., Ahima, R.S., Kaestner, K.H. and Lazar, M.A. (2008) Nuclear receptor corepressor
and histone deacetylase 3 govern circadian metabolic physiology. *Nature*, **456**, 997-
1000.

109. Chen, Y.C., Koutelou, E. and Dent, S.Y.R. (2022) Now open: Evolving insights to the
roles of lysine acetylation in chromatin organization and function. *Mol Cell*, **82**, 716-727.

110. Policarpi, C., Munafò, M., Tsagkris, S., Carlini, V. and Hackett, J.A. (2024) Systematic
epigenome editing captures the context-dependent instructive function of chromatin
modifications. *Nature Genetics*, **56**, 1168-1180.

111. Mazina, M.Y., Nikolenko, Y.V., Krasnov, A.N. and Vorobyeva, N.E. (2016) SWI/SNF
protein complexes participate in the initiation and elongation stages of Drosophila hsp70
gene transcription. *Russian Journal of Genetics*, **52**, 141-145.

112. Schwabish, M.A. and Struhl, K. (2007) The Swi/Snf complex is important for histone
eviction during transcriptional activation and RNA polymerase II elongation in vivo. *Mol
Cell Biol*, **27**, 6987-6995.

113. Huang, Y., Jay, K.L., Huang, A.Y.-W., Wan, J., Jangam, S.V., Chorin, O., Rothschild, A.,
Barel, O., Mariani, M., Iascone, M. *et al.* Loss-of-function in RBBP5 results
in a syndromic neurodevelopmental disorder associated with microcephaly. *Genetics in
Medicine*.

Fig. legends

1289

Fig. 1. Global ~12h RBBP5 cistrome is associated with promoter-proximal ~12h H3K4me3
epigenome in mouse liver. (A) Heatmap of ~12h hepatic transcriptome aligned with H3K4me3,
H3K9Ac and H3K27Ac cistromes. **(B)** Quantification of temporal H3K4me3, H3K9ac and
H3K27Ac. **(C)** Heatmap showing RBBP5 and H3K4me3 chromatin occupancy 1kb ± of TSS of
6,451 genes. **(D)** RBBP5 chromatin occupancy aligned with nucleosome positioning. **(E)**
Quantification of temporal average RBBP5 binding intensity on different nucleosomes. **(F)** RBBP5
chromatin occupancy aligned with H3K4me3 occupancy.

1297

1298

1299

Fig. 2. Hepatic RBBP5 cistrome coincides with that of XBP1s and hepatic ~12h transcriptome. (A) The table shows the number of genes with or without ~12h rhythms and with or without promoter RBBP5 binding. **(B)** Heatmap showing RBBP5 binding intensity along with -$\log_{10}$ transformed P values for having 1~2h rhythm by RAIN and ~12h rhythms of gene expression on 2,525 genes. **(C)** Heat map summary of GO analysis demonstrating the -$\log_{10}$ transformed P value of different enriched pathways for different genes. **(D)** Venn diagram demonstrating distinct and common cistromes for RBBP5 and XBP1s. **(E)** Western blot showing co-IP in liver nuclear extracts from CT0 using anti-XBP1s and normal IgG control antibody. **(F)** Motif analysis of RBBP5 binding sites of 6,149 genes.

1309

1310

1311

Fig. 3. RBBP5 is an epigenetic regulator of the hepatic ~12h oscillator, but not the canonical ~24h circadian clock. (A) Schematic of the design of *Rbbp5^Flox* mice. **(B)**

Immunofluorescence of RBBP5 in the liver of *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice. **(C)** Histograms
showing the period distributions of all rhythmic genes uncovered from *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO}
mice (with adjP values or FDR <0.05) (n=2 at each time point for either genotype). **(D, E)**
Cumulative distribution of 10~12h **(D)** and 22~28h **(E)** genes in *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice
with different FDR cut-offs. **(F)** Heat map of 15 core circadian clock (top) and canonical ~12h gene
expression (bottom) in the liver of *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice. Log$_2$ normalized fold change of
average gene expression across 48 hours between *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice was also
shown. **(G)** Heat map of 6,025 ~12h gene expression (FDR<0.05) in the liver of *Rbbp5* ^{Flox} and
Rbbp5 ^{LKO} mice, with 5,601 of them lost in *Rbbp5* ^{LKO} mice, along with -log$_{10}$ transformed adjP
values for having 10-12h rhythm by RAIN. Log$_2$ normalized fold change of average gene
expression across 48 hours between *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice was also shown. **(H)** RNA-
Seq data for representative proteostasis and mRNA processing genes in the liver of *Rbbp5* ^{Flox}
and *Rbbp5* ^{LKO} mice. **(I)** Heat map summary of GO analysis demonstrating the -log$_{10}$ transformed
P values of different enriched pathways for different genes. Data: Mean ± S.E.M.
1328

1329

1330

1331

1332

1333

1334

1335

1336

1337

1338

1339

1340

1341

1342

1343

1344

1345

1346

1347

1348

1349

Fig. 4. RBBP5 regulates the hepatic transcriptional response to proteotoxic stress. (A) The
diagram of UPR. **(B)** Western blot analysis in the liver of *Rbbp5 Flox* and *Rbbp5 LKO* mice with or
without Tu treatment. **(C)** PCA showing the hepatic transcriptional response to Tu in *Rbbp5 Flox*
and *Rbbp5 LKO* mice. **(D)** Representative RNA-seq data for proteostasis (top) and immune genes
(bottom) with RBBP5 promoter-binding status also shown below. n=3 for *Rbbp5 Flox* DMSO and

Rbbp5 ^{Flox} Tu, and n=4 for *Rbbp5* ^{LKO} DMSO and *Rbbp5* ^{LKO} Tu. **(E)** Heat map summary of GO

analysis demonstrating the -$\log_{10}$ transformed P values of different enriched pathways for different

genes. **(F)** Log $_2$ normalized fold induction by Tu in *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice for genes with

or without RBBP5 promoter binding. **(G)** Western blot analysis of different proteins in *Rbbp5* ^{Flox}

and *Rbbp5* ^{LKO} mice treated with or without Tu. **(H, I)** Mice tail-vein injected with AAV8-TGB-GFP

or AAV8-TGB-RBBP5 were treated with or without Tu. Western blot analysis **(H)** and qPCR **(I)**.

n=3 for AAV8-TBG-EGFP DMSO, n=4 for AAV8-TBG-EGFP Tu, n=4 for AAV8-TGB-mRBBP5

DMSO and n=4 for AAV8-TGB-mRBBP5 Tu. Data: Mean ± S.E.M.

1363

1364

Fig. 5. RBBP5 protects mice from chronic ER stress-induced hepatic inflammation and steatosis. (A) A schematic of experimental design. **(B)** Western blot and quantification of the expression of different proteins in *Rbbp5* [Flox] and *Rbbp5* [LKO] mice with or without repeated Tu injection. **(C)** qPCR analysis of different proteostasis and immune gene expressions in the same cohort of mice. **(D)** Representative images of Oil Red O staining and quantification of lipid droplet size, number and total lipid content (total lipid droplet area). n=4 for *Rbbp5* [Flox] DMSO, n=4 for *Rbbp5* [Flox] Tu, n=2 for *Rbbp5* [LKO] DMSO, n=3 for *Rbbp5* [LKO] Tu. Data: Mean ± S.E.M.

1372

Fig. 6. RBBP5 is required for cell-autonomous transcriptional response to ER stress. (A)
qPCR of different genes in MEFs transfected with scrambled, *Rbbp5* or *Xbp1* siRNAs and treated
with DMSO or Tu (100ng/ml) for 8 hours. **(B)** qPCR of different genes in MEFs transfected with
scrambled, or the combination *Gcn5* and *Pcaf* siRNAs and treated with DMSO or Tu (100ng/ml)
for 8 hours. (**C**) Selected genes aligned for RBBP5 and H3K4me3 ChIP-seq signal from CT0
mouse liver and ChIP-qPCR of RBBP5 and H3K4me3 in MEFs treated with Tu for different hours.
(**D, E**) Immunofluorescence of XBP1s and RBBP5 in MEFs treated with vehicle control (DMSO)
or Tu (100ng/ml) for 8 hours (**D**) and quantification of Manders' coefficient of colocalization of
XBP1s with RBBP5 (**E**). (**F**) *Rbbp5* (fl/fl) ROSA26-CreERT2 (+/+) MEFs were treated with
tamoxifen for 5 or 7 days and western blot analysis was performed. (**G**) qPCR of different genes
in *Rbbp5* (fl/fl) ROSA26-CreERT2 (+/+) MEFs treated with vehicle (WT) or tamoxifen (*Rbbp5* KO)
followed by treatment with vehicle control or 1mM DTT for 5 hours. Data: Mean ± S.E.M.

1385

Fig. 7. RBBP5 is required for cell-autonomous transcriptional response to nutrient deprivation. (A) qPCR analysis of *Rbbp5* (fl/fl) ROSA26-CreERT2 (+/+) MEFs pre-treated with (RBBP5 KO) or without (WT) tamoxifen and then treated with full DMEM media, leucine-free media or EBSS for 16 hours. **(B-D)**. Transcriptome profiling of *Rbbp5* (fl/fl) ROSA26-CreERT2 (+/+) MEFs pre-treated with (RBBP5 KO) or without (WT) tamoxifen and then treated with full DMEM media or EBSS for 16 hours by RNA-seq. Heatmap of all genes that are induced by EBSS in WT cells by EBSS (adjP<0.05) **(B)**. GO analysis of EBSS-induced genes in only WT or both genotypes **(C)**, and representative gene expression **(D)**. **(E)** Diagram of the mCherry-GFP-LC3 autophagy reporter. **(F)** Quantification of autophagic flux in mCherry-GFP-LC3-expressing *Rbbp5* (fl/fl) ROSA26-CreERT2 (+/+) MEFs treated with vehicle (WT) or tamoxifen (*Rbbp5* KO) followed by treatment with full DMEM media or leucine-free media for 16 hours. **(G)** Diagram of the ER-phagy reporter using the mCherry cleavage from ER assay. **(H, I)** Western blot of different proteins **(H)** and quantification **(I)** of mCherry cleavage in RAMP4-mCherry-expressing *Rbbp5* (fl/fl) ROSA26-CreERT2 (+/+) MEFs treated with vehicle (WT) or tamoxifen (*Rbbp5* KO) followed by treatment with full DMEM media or EBSS for 16 hours. Data: Mean ± S.E.M.

1401

Fig. 8. Proximity labeling reveals H3K4me-associated proteomic architecture of the proteostatic stress response. (**A**) A diagram illustrating the procedure of performing H3K4me3 proximity labeling via using the recombinant Protein A-TurboID. (**B**) Western blot analysis showing the nuclear level of total (input) or biotinylated (after streptavidin beads pull-down) level of H3K4me3 in MEF treated with 100ng/ml Tu for 0, 2.5 or 8h before subject to anti-IgG and anti-H3K4me3 antibody-based proximity labeling. (**C**) Enriched biological network of 239 proteins. (**D**) Heatmap of fold enrichment of different SWI/SNF subunits (over IgG control) at different times after ER stress. (**E**) Quantification of the average relative enrichment of proteins in different complexes at different times after ER stress. (**F**) GO analysis of BAF180 hepatic cistrome. (**G**) Selected UPR genes aligned for BAF180, RBBP5, H3K4me3 and XBP1s ChIP-seq signal in mouse liver. (**H**) Heatmap showing RBBP5, H3K4me3 and BAF180 chromatin occupancy 2kb ± of TSS of 6,451 genes. Data: Mean ± S.E.M.

1414

**Fig. 9. Reduced RBBP5 expression is associated with aging in mice and hepatic steatosis
in humans. (A)** Z score-normalized expression of mouse *Rbbp5* and average expression of
stress response genes across mouse life span according to Tabua Muris dataset (97). **(B-F)** Data
in humans with NAFLD/MAFLD was from (99). Expression of *RBBP5* (**B**) and the eigengene of
97 stress genes (**C**). (**D**) Scatter plot showing a positive correlation between *RBBP5* and
eigengene expression. **(E)** Heatmap of 97 stress genes expression in different cohorts of human
subjects. (**F**) GSEA analysis showing stress gene expression are downregulated in steatosis
subjects.