Epigenetic regulation of global proteostasis dynamics by RBBP5 ensures mammalian organismal health

Syeda Kubra[1, #], Michelle Sun[1, #], William Dion[1], Ahmet Catak[1], Hannah Luong[1], Haokun Wang[1], Yinghong Pan[2, ¥], Jia-Jun Liu[3,4], Aishwarya Ponna[1], Ian Sipula[5], Michael J. Jurczak[5,6], Silvia Liu[3,4,7], and Bokai Zhu[1,3,5] *

[1] Aging Institute of UPMC, University of Pittsburgh School of Medicine, Pittsburgh, PA, U.S.A.

[2] UPMC Genome Center, UPMC, Pittsburgh, PA, U.S.A.

[3] Pittsburgh Liver Research Center, University of Pittsburgh, Pittsburgh, PA, U.S.A.

[4] Organ Pathobiology and Therapeutics Institute, University of Pittsburgh School of Medicine, Pittsburgh, PA, U.S.A.

[5] Division of Endocrinology and Metabolism, Department of Medicine, University of Pittsburgh School of Medicine, Pittsburgh, PA, U.S.A.

[6] Center for Metabolism and Mitochondrial Medicine, University of Pittsburgh School of Medicine, Pittsburgh, PA, U.S.A.

[7] Department of Pharmacology and Chemical Biology, University of Pittsburgh School of Medicine, Pittsburgh, PA, U.S.A.

# These authors contribute equally.

¥ Present address: UPMC Magee-Womens Hospital, Pittsburgh, PA, U.S.A.

* Corresponding author: Bokai Zhu: bzhu@pitt.edu

Abstract:

Proteostasis is vital for cellular health, with disruptions leading to pathologies including aging, neurodegeneration and metabolic disorders. Traditionally, proteotoxic stress responses were studied as acute reactions to various noxious factors; however, recent evidence reveals that many proteostasis stress-response genes exhibit ~12-hour ultradian rhythms under physiological conditions in mammals. These rhythms, driven by an XBP1s-dependent 12h oscillator, are crucial for managing proteostasis. By exploring the chromatin landscape of the murine 12h hepatic oscillator, we identified RBBP5, a key subunit of the COMPASS complex writing H3K4me3, as an essential epigenetic regulator of proteostasis. RBBP5 is indispensable for regulating both the hepatic 12h oscillator and transcriptional response to acute proteotoxic stress, acting as a co-activator for proteostasis transcription factor XBP1s. RBBP5 ablation leads to increased sensitivity to proteotoxic stress, chronic inflammation, and hepatic steatosis in mice, along with impaired autophagy and reduced cell survival *in vitro*. In humans, lower *RBBP5* expression is associated with reduced adaptive stress-response gene expression and hepatic steatosis. Our findings establish RBBP5 as a central regulator of proteostasis, essential for maintaining mammalian organismal health.

Introduction:

Proteostasis, the maintenance of proper protein folding, trafficking, and turnover, is a major challenge for the cell (Labbadia & Morimoto, 2015). Much of the cellular energy expenditure is dedicated to maintaining a healthy proteome, as millions of molecules are produced every single minute (Buttgereit & Brand, 1995). Challenges to proteostasis, exemplified by the accumulation of unfolded or misfolded proteins, can lead to a range of cellular dysfunctions and pathological conditions, such as aging, neurodegenerative diseases, cancer, and metabolic disorders (Klaips *et al*, 2018; Labbadia & Morimoto, 2015; Martinez *et al*, 2017; Taylor & Hetz, 2020; Zhao & Ackerman, 2006). To cope with proteotoxic stress, organisms have developed a wide range of stress response mechanisms, including the heat shock response, the unfolded protein response (UPR), and the more generalized integrated stress response (ISR) (Costa-Mattioli & Walter, 2020; Hetz *et al*, 2020; Richter *et al*, 2010). Although the detailed signaling molecules vary, these pathways generally involve transmitting the information of cellular stress to the nucleus by activating various transcription factors (TF) containing the basic leucine zipper domain (bZIP), such as UPR TFs spliced form of XBP1 (XBP1s), ATF4 and ATF6 and heat shock factor 1 (HSF1). After binding to the stress response element in the promoters/enhancers, these TFs in turn upregulate a plethora of genes aimed to restore proteostasis (Balch *et al*, 2008; Ron & Walter, 2007).

While proteotoxic stress responses were traditionally viewed and studied as distinct acute responses to different "noxious" factors, recent evidence by our group indicates that at the physiological level (and in the absence of evident extrinsic stress), the expression of hundreds of proteotoxic stress response regulatory and output genes exhibit cell-autonomous ultradian rhythms cycling with a ~12h period in multiple mammalian cell-types and tissues, including humans (Asher & Zhu, 2022b; Ballance & Zhu, 2021; Dion *et al*, 2022a; Meng *et al*, 2022; Meng *et al*, 2020; Pan *et al*, 2020; Scott *et al*, 2023; Zhu *et al*, 2023; Zhu *et al*, 2024; Zhu *et al*, 2017b). These proteostasis genes encompass both output genes involved in protein processing in the ER/Golgi, redox regulation, protein folding, ER-associated protein degradation (ERAD), autophagy, and upstream regulatory molecules and TFs like *Xbp1*, *Atf4*, *Atf6*, *Ddit3*, *Perk* and *Ire1α* (Asher & Zhu, 2022a; Ballance & Zhu, 2021). Many of these ~12h ultradian rhythms are

established by an XBP1s-dependent 12h oscillator separate from both the ~24h circadian clock
and the cell cycle (Pan *et al.*, 2020; Zhu & Liu, 2023a; Zhu *et al*, 2017a). It is hypothesized that
the mammalian 12h oscillator may have evolved from the circatidal clock of marine animals, and
later co-opted to adapt to the ~12h cycle of metabolic stress peaking at transition times at dawn
and dusk in terrestrial animals (Asher & Zhu, 2022b; Pan *et al.*, 2020; Zhu *et al.*, 2024). The 12h
oscillator is well-studied in the liver, and mice with liver-specific 12h oscillator ablation exhibited
accelerated liver aging and steatosis (Meng *et al.*, 2020), manifested with impaired proteostasis,
lipid metabolism, and mitochondria dysfunction (Dion *et al*, 2022b; Meng *et al.*, 2022; Meng *et al.*,
2020). Since the 12h oscillator integrates multiple proteotoxic stress signaling, it provides us with
a unique opportunity to study proteostasis using a holistic approach: rather than studying
individual stress response separately, by investigating the 12h oscillator we could gain new
mechanistic insights for universal proteostasis regulation.

While significant work has been done on upstream proteotoxic stress sensing and protein-
folding mechanisms in the ER and cytosol, there remains a lack of knowledge regarding
transcriptional regulation in proteostasis in general, particularly concerning the temporal
epigenome dynamics, chromatin landscapes, and co-regulatory networks underlying global
proteostasis control. In this study, we seek to address an important unanswered question: is there
a designated epigenetic regulator that orchestrates global proteostasis? By examining the
chromatin landscape of the murine 12h hepatic oscillator, we identified RBBP5—an essential
subunit of the Complex Proteins Associated with Set1 (COMPASS) complex responsible for
depositing H3K4me3 (Shilatifard, 2012a) —as a pivotal epigenetic regulator that governs global
proteostasis dynamics. RBBP5 plays an indispensable role in regulating both the 12h oscillator
and transcriptional response to acute proteostasis stress, functioning as a co-activator for key
proteostasis master TFs, such as XBP1s. Ablation of RBBP5 leads to heightened sensitivity to
proteotoxic stress, resulting in chronic inflammation and hepatic steatosis in mice, as well as
defective autophagy and impaired cell survival *in vitro*. In humans, reduced RBBP5 expression is
linked to a diminished adaptive stress response and the onset of hepatic steatosis. Our findings
establish RBBP5 as a crucial epigenetic regulator of proteostasis dynamics, essential for
maintaining organismal health.

Results:

Global RBBP5 binding to chromatin exhibits a ~12h ultradian rhythm, correlated with the
promoter-proximal ~12h H3K4me3 rhythms in mouse liver.

The 12h oscillator is transcriptionally regulated by the UPR TF XBP1s in mice (Meng *et al.*, 2020;
Pan *et al.*, 2020; Zhu *et al.*, 2017b), and likely under control by additional bZIP TFs, such as ATF4
and ATF6 (Zhu, 2020). However, decades of research of eukaryotic gene regulation demonstrated
that TFs themselves are not sufficient to initiate transcription, and additional so-called "co-
regulators" are required for full gene activation by shaping the epigenetic landscape (Dasgupta *et*
al, 2014; Roeder, 2005; Spiegelman & Heinrich, 2004). Currently, the epigenetic landscape of
both the 12h oscillator and proteotoxic stress responses are poorly characterized. To identify
epigenetic regulators of global proteostasis control, we first examined a published temporal
hepatic epigenome dataset for histone modifications associated with the active promoters of ~12h
genes (Koike *et al*, 2012). We found ~12h rhythms of histone H3 trimethylated at lysine 4
(H3K4me3), but not two histone acetylation markers, histone H3 acetylated at lysine 9 (H3K9Ac)
and lysine 27 (H3K27Ac), at the promoters of ~12h genes (**Fig. 1A, B**). This is particularly
prominent for those ~12h genes with rhythmic expressions peaking at ZT0 and ZT12 that are

enriched for proteostasis pathways (Pan *et al.*, 2020), with H3K4me3 level also peaking at these
two time points (**Fig. 1A, B**).

In eukaryotes, H3K4 is tri-methylated by the COMPASS complex, which includes regulatory
subunits RBBP5, ASH2L, WDR5, DPY30 and catalytic subunits SETD1A/B/MLL1-4 (**Fig. S1A**)
(Qu *et al*, 2018; Shilatifard, 2012b; Takahashi *et al*, 2011). Via writing H3K4me3, COMPASS is
essential for the eviction of paused RNA polymerase II (RNAPII) and promotes transcriptional
elongation (Hu *et al*, 2023; Wang *et al*, 2023). Landscape In Silico deletion Analysis (LISA) (Qin
et al, 2020) further inferred COMPASS subunits WDR5, ASH2L, DYP30 and RBBP5 as putative
epigenetic regulators of hepatic ~12h, but not circadian genes (**Fig. S1B**). Being the commonly
shared regulatory subunits for all MLL family H3K4 methyl transferase complex (including
SETD1A/B and MLL1-4), RBBP5/ASH2L forms the minimal heterodimer sufficient to activate all
MLL histone methyl transferase (Li *et al*, 2016; Ruthenburg *et al*, 2007) and are essential for the
assembly of the whole COMPASS complex (Han *et al*, 2019; Qu *et al.*, 2018). Between the two
essential subunits, we initially focused on RBBP5. Eigenvalue/pencil method (Antoulas *et al*,
2018b) revealed that the levels of both hepatic *Rbbp5* mRNA and nuclear RBBP5 protein exhibit
~12h oscillations, with the nuclear protein level peaking at ~CT0 and ~CT12 (**Fig. S1C, D**), in line
with the acrophases of H3K4me3 and gene expression oscillations (**Fig. 1A, B**).

To further evaluate the potential role of RBBP5 in establishing the epigenetic landscape of
hepatic ~12h rhythms, we performed temporal RBBP5 ChIP-seq at a 4-h interval for a total of 48
hours in the liver of male C57BL/6 mice housed under constant darkness condition fed *ad libitum*.
Chromatin occupancies of RBBP5 align well with nucleosome positioning downstream of the
transcription start sites (TSS) and the H3K4me3 epigenome, with 'ridges' corresponding to 130bp,
310bp, 490bp, 670bp downstream of TSS at all time points, thus demonstrating the high quality
of our ChIP-seq dataset (**Fig. 1C-F**). In total, we identified 7,963 hepatic RBBP5 binding sites that
are within 2kb (up and downstream) of TSS of 6,451 genes (**Fig. 1C and Table S1**). The average
binding intensities of RBBP5 on all post-TSS nucleosomes displayed robust 12h rhythms, with
peaks occurring at CT0, CT12, CT24, and CT36 (**Fig. 1C-E**). These rhythms aligned with the 12h
oscillations of promoter-proximal H3K4me3 (**Fig. 1C-E**), but not with those of H3K9Ac or
H3K27Ac (**Fig. S1E**). This pattern is exemplified by *Xbp1*, one of the major UPR TFs and the
transcriptional regulator of the 12h oscillator, and *Hsph1*, a canonical 12h oscillator output gene
encoding a heat shock protein (**Fig. S1F**). Using the eigenvalue/pencil and RAIN (Thaben &
Westermark, 2014) algorithms, we identified 3,028 and 1,864 (p<0.05) RBBP5 binding sites that
cycle with a ~12h period, respectively (**Fig. S1G, H and Table S1**). Collectively, these results
establish the epigenetic landscape of hepatic 12h oscillator is marked by RBBP5-H3K4me3.

Hepatic RBBP5 cistrome coincides with that of XBP1s and hepatic ~12h transcriptome.

To investigate if RBBP5 regulates hepatic ~12h rhythms of gene expression, we first examined
the correlation between RBBP5 proximal promoter binding status and ~12h rhythms of gene
expression. As illustrated in **Fig. 2A**, genes with proximal promoter RBBP5 binding are enriched
for ~12h rhythms of gene expression (P=1.1e-39 by Chi-squared test). For those 2,525 ~12h
genes with RBBP5 binding, the phases of RBBP5 chromatin binding center around CT0 and CT12,
slightly preceding the phases of gene expression observed at ~CT1 and ~CT13 (**Fig. 2B**). This
temporal relationship suggests that RBBP5 acts as a driver, rather than a consequence of ~12h
rhythms of gene expression.

We next performed Gene Ontology (GO) analysis on those hepatic genes with proximal promoter RBBP5 binding and revealed strong enrichment of proteostasis pathways including protein processing in the ER and ubiquitin-mediated proteolysis (**Fig. 2C**). Additionally, pathways involved in mRNA processing, such as spliceosome, RNA transport, and mRNA surveillance, were also enriched (**Fig. 2C**). These findings align with previous observations that both mRNA and protein metabolism pathways are enriched in genes exhibiting ~12h rhythms in both mice and humans (Dion *et al.*, 2022a; Zhu & Liu, 2023b; Zhu *et al.*, 2024). The coupling of mRNA and protein metabolism is orchestrated by a SON-XBP1s axis and involves the liquid-liquid phase separation (LLPS) dynamics of nuclear speckles (Dion *et al.*, 2022a). Similar pathways are also enriched in genes exhibiting both ~12h rhythms of RBBP5 promoter binding and gene expression (**Fig. 2C**).

Recent studies indicated that by depositing H3K4me3, RBBP5 (as a part of COMPASS) is essential for the eviction of paused RNAPII and promotes transcriptional elongation (Hu *et al.*, 2023; Wang *et al.*, 2023). To investigate whether the ~12h rhythms of RBBP5 promoter binding may translate into ~12h rhythms of RNAPII pause release, we performed a post-hoc analysis of a published Global run-on sequencing (GRO-Seq) dataset in mouse liver (Fang *et al*, 2014) that measured nascent hepatic RNA transcription at a 3h interval for a total of 24 hours. While the sampling frequency and duration of this dataset are not optimal for a comprehensive and rigorous detection of ~12h rhythms, we believe it may still offer valuable mechanistic insights into the transcriptional regulation of the hepatic 12-hour oscillator. By calculating the temporal pausing index of each ~12h hepatic gene (**Fig. 2D**), we observed that, on average, genes with RBBP5 promoter binding exhibited a greater amplitude in pausing index oscillations, peaking at ZT10 and ZT22, compared to those without RBBP5 binding (**Fig. 2E**).

XBP1s is the major transcriptional factor regulating the hepatic 12h oscillator and also exhibits a global 12h rhythm of chromatin binding to the promoter regions peaking at CT0, 12, 24 and 36 (Meng *et al.*, 2020; Pan *et al.*, 2020). Comparing the cistromes of RBBP5 and XBP1s revealed a significant overlap between the two, with 920 genes sharing RBBP5 and XBP1s proximal promoter binding (**Fig. 2F**). Co-immunoprecipitation in the hepatic nuclear extracts at CT0 confirmed the physical interaction between XBP1s and subunits of COMPASS, including RBBP5, ASH2L, WDR5, and two histone methyltransferase SETD1A and SETD1B (**Fig. 2G**). Motif analysis of the DNA sequences around the 6,149 RBBP5 binding sites that do not overlap with XBP1s cistrome identified binding motifs for GABPA, KLF, ATF6, NFYA and ATF4 (**Fig. 2H**). These findings suggest that, in addition to acting as a co-activator for XBP1s, RBBP5 may also work with other transcription factors to shape the hepatic 12-hour epigenome.

RBBP5 is an epigenetic regulator of the ~12h oscillator, but not the canonical ~24h circadian clock.

To establish the causality between RBBP5 and hepatic 12h rhythms of gene expression, we generated RBBP5 liver hepatocyte-specific knockout (RBBP5 ^{LKO}) mice using the CRE-loxP system. Exon 10 of mouse *Rbbp5,* which encodes the SET/ASH2L binding domain, was flanked by loxP sites (**Fig. 3A**). CRE-mediated deletion of Exon 10 is expected to lead to a frameshift and nonsense-mediated RNA decay of truncated *Rbbp5* transcript (**Fig. S2A**). Crossing homozygous floxed *Rbbp5* mice with Albumin-CRE mice resulted in liver-specific deletion of RBBP5, which was confirmed by DNA genotyping, immunofluorescence against RBBP5 and western blot analysis (**Figs. 3B and S2B-D**). The resulting RBBP5 ^{LKO} mice weighed slightly less than wild-

type counterparts (**Fig. S3A**) but maintained normal rhythmic locomotor activity and fasting-feeding cycles under both 12h:12h L/D and constant darkness conditions (**Fig. S3B-I**).

To identify the RBBP5-dependent oscillating hepatic transcriptome, we performed bulk RNA sequencing (RNA-Seq) analysis in the liver of male *Rbbp5 Flox* and *Rbbp5 LKO* mice at 2-h intervals for a total of 48 hour under constant darkness in duplicates (**Table S2**). To identify genes cycling with a period ranging from 6 to 36 hours, we initially applied the RAIN algorithm (Thaben & Westermark, 2014) to each genotype's temporal transcriptome (Thaben & Westermark, 2014) (**Table S3**). Compared to alternative methods like JTK_CYCLE, RAIN detects rhythms with arbitrary waveforms and therefore more robustly uncovers ultradian rhythms (Antoulas *et al*, 2018a; Pan *et al.*, 2020; Zhu *et al.*, 2017a). Consistent with past studies (Hughes *et al*, 2009; Meng *et al.*, 2022; Pan *et al.*, 2020; van der Veen & Gerkema, 2017; Zhu *et al.*, 2017b), two populations of hepatic transcripts cycling with periods of ~12h and ~24h were identified in *Rbbp5 Flox* mice liver (**Fig. 3C, D**). By contrast, in *Rbbp5 LKO* mice, the ~12h rhythms are all but abolished and the circadian oscillations remain largely intact, which includes all core circadian clock genes (such as *Bmal1*, *Per1/2/3*, and *Nr1d1/2*) displaying robust circadian rhythms with the same phases and amplitudes as those observed in *Rbbp5 Flox* mice (**Figs. 3C-F and S4A**). The maintenance of ~24h circadian rhythms and abolishment of ~12h ultradian rhythms in *Rbbp5 LKO* mice is further confirmed by the principal component analysis (PCA) (**Fig. S4B, C**). Specifically, using RAIN with an FDR cutoff of 0.01 and 0.05, we identified a total of 3,549 and 6,025 hepatic transcripts cycling at periods between 10-12h in *Rbbp5 Flox* mice, respectively. By contrast, only 61 and 813 10-12h genes were observed in *Rbbp5 LKO* mice with the same FDR thresholds, indicating over 93% ~12h hepatic transcriptome was abolished with liver RBBP5 ablation (**Figs. 3G, H, S4D and Table S3**). GO analysis confirmed that these RBBP5-depenent ~12h hepatic transcriptome is enriched in various mRNA processing and proteostasis pathways, distinct from those of RBBP5-independent circadian genes enriched in lipid metabolism and phosphorylation (**Fig. 3I**).

To determine the robustness of our results to different analytic methods, we also performed spectrum analysis with the eigenvalue/pencil method (**Table S4**) (Antoulas *et al.*, 2018a; Dion *et al.*, 2022b; Meng *et al.*, 2022; Meng *et al.*, 2020; Pan *et al.*, 2020; Zhu *et al.*, 2017a), which unlike statistical methods such as JTK_CYCLE and RAIN does not require pre-assignment of period range, enabling unbiased identification of multiple superimposed oscillations for any given gene (Antoulas *et al.*, 2018a; Dion *et al.*, 2022b; Meng *et al.*, 2022; Meng *et al.*, 2020; Pan *et al.*, 2020; Zhu *et al.*, 2017a). Revealing up to three oscillations from each gene, eigenvalue/pencil analyses also revealed prevalent ~24h and ~12h oscillations in *Rbbp5 Flox* mice, along with a third population cycling at a period around 8h (**Fig. S5A-E**). Hepatic ~8h oscillations were also previously reported but their regulation and function remained elusive (Asher & Zhu, 2022a; Hughes *et al.*, 2009). *Rbbp5 LKO* mice, on the other hand, exhibited a drastically different spectrum, characterized by the near complete loss of the ~12h rhythms, *de novo* gain of many faster ultradian rhythms with periods between 6 and 7h, and the maintenance of most circadian rhythms (**Fig. S5A-E**). Similar findings were also observed for dominant oscillations (which is the one with the largest amplitude among the three superimposed oscillations for each gene) for each gene (**Fig. S5A-E**). For the ~12h genes originally identified in *Rbbp5 Flox* mice but lost in *Rbbp5 LKO* mice, the period spectrum in the *Rbbp5 LKO* mice displayed a diverse distribution. This ranged from a complete lack of rhythmicity (~39%) to shorter ultradian periods of 6-7 hours, and even to longer periods exceeding 24 hours (**Fig. S5F**). RBBP5-dependent ~12h transcriptome uncovered by the eigenvalue/pencil showed substantial overlap with the results obtained through the RAIN method.

This convergence was evident both in the specific genes identified (**Fig. S5C**) and in the enriched
biological pathways (**Fig. S5G, H**), thereby confirming the robustness of our findings. These
results thus establish RBBP5 as a central epigenetic regulator of the hepatic 12h oscillator, with
more than 90% ~12h hepatic transcriptome abolished in its absence.

In addition to liver, cell-autonomous ~12h rhythms of proteostasis gene expression including
Xbp1 can also be observed in mouse embryonic fibroblasts (MEFs) synchronized by low
concentration of ER stress inducer tunicamycin (Tu) (Pan *et al.*, 2020; Zhu *et al.*, 2017a). We
observed a ~12h rhythm of *Rbbp5* expression in Tu-synchronized MEFs, which is independent of
BMAL1 but require XBP1s (**Fig. S6A**), indicating RBBP5 itself is also under 12h oscillator control.
To test the functional role of COMPASS in regulating the cell-autonomous ~12h rhythms, we
knocked down either *Rbbp5* or *Ash2l* via siRNA in a previously described 12h oscillator reporter
MEFs that stably express *Manf* promoter-driven destabilized luciferase (dluc) (Pan *et al.*, 2020),
and found that both siRNAs significantly dampen the ~12h oscillation of luciferase activity as well
as shorten the period (**Fig. S6B**). By contrast, neither *RBBP5* nor *ASH2L* knocking down affects
the circadian oscillation of *Bmal1* promoter activity in human U2OS cells in a previous genome-
wide siRNA screen study (Zhang *et al*, 2009) (**Fig. S6C**). Collectively, our results demonstrate
that RBBP5 is an integral component of the 12h oscillator, but dispensable for the canonical
circadian clock both *in vivo* and *in vitro*.

RBBP5 modulates the hepatic transcriptional response to acute proteotoxic stress *in vivo*.

Before the discovery of the mammalian 12h oscillator, the dynamics of proteostasis were primarily
studied as transient responses to various proteotoxic stresses, with a classic example being the
UPR triggered by ER stress. In response to the accumulation of misfolded proteins in the ER, the
UPR initiates a signaling cascade from the ER to the nucleus, ultimately activating three key
transcription factors: XBP1s, ATF4, and ATF6. This activation leads to a beneficial adaptive
response, upregulating proteostasis genes to restore ER homeostasis (**Fig. 4A**) (Metcalf *et al*,
2020; Walter & Ron, 2011; Wiseman *et al*, 2022). However, ER stress can also trigger maladaptive
and deleterious responses, such as the activation of pro-inflammatory and immune gene
expression through the TRAF2-mediated activation of AP1 (comprised of c-Fos and c-Jun), NF-
κB or STAT3 TFs (**Fig. 4A**) (So, 2018). Our findings that RBBP5 can co-activate XBP1s to regulate
the 12h oscillator suggest that RBBP5 is essential for the adaptive UPR TFs-mediated activation
of proteostasis genes but is not required for the maladaptive activation of immune genes by
immune-regulatory TFs, whose binding motifs are absent from RBBP5 binding sites (**Fig. 2H**).

To test this hypothesis, we intraperitoneally injected male *Rbbp5* [Flox] and *Rbbp5* [LKO] mice with
vehicle control or 0.05mg/kg Tu to induce acute ER stress and performed RNA-seq on hepatic
transcriptome isolated 8 hours post injection (**Table S5**). This lower dose of Tu was selected for
its ability to induce the UPR, but that was significantly less toxic than the more common
experimental dose of 1 mg/kg—which is known to elicit hepatocellular death and severe toxicity
(Rutkowski *et al*, 2008). PCA demonstrated the separation of the four groups on the transcriptome
space (**Fig. 4B**). Using an adjusted p value (FDR) of 0.10 as cut-off, we identified 139 and 45
genes significantly induced by Tu in the liver of *Rbbp5* [Flox] and *Rbbp5* [LKO] mice, respectively, with
21 commonly shared between the two (**Figs. 4C, S7A-G and Table S6**). In *Rbbp5* [Flox] mice, these
139 genes are enriched in pathways of proteostasis and immune functions as expected, with
pathways of 'response to ER stress' and 'ERAD' among the top two GO terms (**Fig. 4C, D**). By
contrast, in *Rbbp5* [LKO] mice, these 45 genes are only enriched in immune functions, with
proteostasis genes not being significantly induced by Tu (**Fig. 4C, D**). In general, proteostasis

genes were induced by Tu to a much greater fold in *Rbbp5* ^{Flox} mice (such as *Creld2*, *Xbp1*,
Hsp90b1), while the opposite was found for blood coagulation and immune genes in *Rbbp5* ^{LKO}
mice (such as *S100a9* and *Fgb*) (**Figs. 4C, D and S7E, F, H**). Further integrating RBBP5 ChIP-
seq results, we found that, on average, the ablation of RBBP5 markedly diminished the fold
induction by Tu for genes with RBBP5 promoter binding, while it had no effects for those without
RBBP5 chromatin recruitment (**Fig. 4C, E**). To rule out the possibility that RBBP5 regulates UPR
via influencing the upstream ER stressing in the ER, we performed western blot analysis of p-
IRE1α and p-PERK in the liver of *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice and did not observe any notable
difference in the increase of their phosphorylated level in response to Tu between the two
genotypes (**Fig. 4F**).

To determine if RBBP5 overexpression is also sufficient to amplify UPR, we tail-vein injected
AAV8-TBG-GFP and AAV8-TBG-RBBP5 viruses into wild-type male mice followed by
intraperitoneal injection with vehicle control or 0.025mg/kg Tu 30 days later. Thyroxine binding
globulin (TBG) promoter is liver-specific and directs RBBP5 transgene expression in hepatocytes
(Yan *et al*, 2012). qPCR and western blot analysis confirmed an increase of total hepatic
expression of RBBP5 of ~1.4 fold in AAV8-TBG-RBBP5 group under vehicle control condition (**Fig.
4G, H**). Consequently, RBBP5 hepatic overexpression significantly amplified the UPR of
proteostasis genes including *Manf*, *Hspa5* (*BiP*), *Pdia4*, *Hyou1*, and *Syvn1* without affecting the
upstream ER stress sensing *(**Fig. 4G, H***). In addition, we found that RBBP5 overexpression
attenuated the expression of several immune and blood coagulation genes such as *C3*, *Tifa*, *Fgl1*
and *Fgb* under ER stress conditions (**Fig. 4H**). Taken together, these results demonstrate that
RBBP5 is both necessary and sufficient for the transcriptional regulation of hepatic response to
acute proteotoxic stress *in vivo*. RBBP5 activates the expression of proteostasis genes while
repressing those involved in immune functions.

RBBP5 protects mice from chronic ER stress-induced hepatic inflammation and steatosis.

Contrary to acute ER stress, chronic ER stress, experimentally reconstituted by repeated injection
of Tu, paradoxically leads to feedback-mediated suppression of the UPR signaling, to even below
basal levels (Rutkowski *et al.*, 2008). Proposed mechanisms of this suppression include silencing
of the ATF6a branch of UPR, enhancement of mRNA degradation via IRE1-dependent decay
(RIDD), and the direct inhibition of UPR stress sensors by the ER chaperone HSPA5 (BiP)
(Rutkowski *et al.*, 2008). To investigate whether RBBP5 also modulates the feedback suppression
of UPR under chronic ER stress conditions, we repeatedly intraperitoneally injected male *Rbbp5*
^{Flox} and *Rbbp5* ^{LKO} mice with vehicle control or 0.025mg/kg Tu for six consecutive days and
harvested liver tissues 24 hours after the last injection for analysis (**Fig. 5A**). Western blot analysis
revealed attenuation of p-IRE1α and ATF4 level in both *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice with
repeated Tu injection (**Fig. 5B**). In agreement with the Western blot result, qPCR analysis also
confirmed the downregulation of proteostasis genes *Atf4*, *Hyou1*, and *Manf* in both mice under
chronic ER stress condition (**Fig. 5C**). By contrast, compared to *Rbbp5* ^{Flox} mice, the expression
of pro-inflammatory cytokine genes *Il1α* and *Il1b* were higher in Tu-injected *Rbbp5* ^{LKO} mice,
consistent with higher levels of p-IRE1α and p-c-Jun (Ser73) observed in *Rbbp5* ^{LKO} mice (**Fig.
5B, C**). The persistent elevated levels of p-IRE1α/p-c-JUN and pro-inflammatory cytokine gene
expression in *Rbbp5* ^{LKO} mice suggest that these mice exhibit heightened sensitivity to Tu-induced
ER stress and inflammation. In summary, our results indicate that while wild-type mice can sustain
persistent cycles of UPR activation and deactivation in response to ER stress, as previously
described (Rutkowski *et al.*, 2008), the loss of RBBP5 disrupts this balance. Although RBBP5-

deficient mice retain the ability to attenuate UPR, they are unable to properly activate it. Consequently, *Rbbp5* *LKO* mice exhibit consistently lower expression of adaptive proteostasis genes under ER stress condition, leading to heightened proteotoxic stress and chronic inflammation likely via activating the p-IRE1α-TRAF2-AP1 signaling cascade.

Both chronic ER stress and inflammation can result in hepatic steatosis (Ajoolabady *et al*, 2023; Gehrke & Schattenberg, 2020; Gong *et al*, 2024; Guo & Li, 2014). We therefore performed Oil Red O staining, which stained all neutral lipids, on liver tissues from *Rbbp5* *Flox* and *Rbbp5* *LKO* mice after repeated vehicle or Tu injection. *Rbbp5* *LKO* mice liver displayed an increased number of lipid droplets and total lipid content, compared to wild-type counterparts under both basal and chronic ER stress conditions (**Fig. 5D**). Our data thus revealed that RBBP5 protects mice from chronic ER stress-induced hepatic inflammation and steatosis.

Ablation of RBBP5 in MEFs compromises the transcriptional response to proteotoxic stress and nutrient-deprivation, resulting in impaired autophagy and reduced cell survival.

To determine whether RBBP5 regulates transcriptional responses to proteotoxic stress in a cell-autonomous manner beyond the liver, we examined MEFs, which exhibit both a robust cell-autonomous XBP1s-dependent 12h oscillator and strong transcriptional responses to proteotoxic stress (Dion *et al.*, 2022a; Dion *et al*, 2024; Pan *et al.*, 2020; Zhu *et al.*, 2017b). *Rbbp5* mRNA expression was induced by Tu (**Fig. 6A**), and transient knockdown of RBBP5 using siRNA (**Fig. S8A**) significantly dampened the UPR of proteostasis gene expressions in response to Tu, mirroring the effects observed with XBP1s knockdown (**Fig.6A**). Similarly, knocking down ASH2L also markedly dampened the UPR, whereas knocking down SETD1A, one of the H3K4me3 methyltransferases in the COMPASS complex, had only modest effects—likely due to the redundancy among the many H3K4me3 methyltransferases that exist (**Fig. S8B**). As a negative control, we further knocked down both GCN5 and PCAF, proteins that function to catalyze H3K9Ac as subunits in two distinct histone acetyltransferase (HAT) complexes, SAGA (Spt-Ada-Gcn5 acetyltransferase) and ATAC (Ada2a-containing) (Gates *et al*, 2017a; Gates *et al*, 2017b; Jin *et al*, 2011) and found it had no effects on the induction of *Xbp1* and *Manf* expression in response to ER stress (**Fig. S8C**). The lack of effects on UPR for HAT subunits is consistent with the absence of ~12h rhythms of both H3K9Ac and H3K27Ac levels around the promoter of ~12h proteostasis genes in mouse liver (**Fig. 1A, B**). ChIP-qPCR analysis confirmed a gradual, temporal increase in the recruitment of RBBP5 to the proximal promoters of select proteostasis genes in response to ER stress, preceding the rise in H3K4me3 levels (**Fig. 6B**).

To broadly assess the function of RBBP5 in the transcription regulation of proteotoxic stress response, we utilized orthogonal approaches to both deplete RBBP5 expression and trigger cellular stress. We took advantage of our homozygous *Rbbp5* *Flox* mice by crossing them with homozygous ROSA26 Cre-ERT2 mice, then isolated and immortalized primary MEFs from homozygous F2 embryos (**Fig. S8D**). Treating these *Rbbp5* (flox/flox); ROSA26 Cre-ERT2 (+/+) MEFs with tamoxifen for 7 days resulted in the complete deletion of both floxed *Rbbp5* alleles (**Fig. S8E**) and a subsequent ~90% reduction in RBBP5 protein level (**Fig. S8F**). The remaining RBBP5 protein is likely extremely long-lived and remains stable even after the deletion of underlying coding genes. To trigger ER stress, we used a different stimulus: Dithiothreitol (DTT), which disrupts protein disulfide formation in the ER (G & Singh, 2022; Higa *et al*, 2014). Similar to Tu, RBBP5 depletion significantly dampened the induction of proteostasis gene expression in response to DTT (**Fig. 6C**).

Finally, we tested two additional proteotoxic stressors that are more physiologically relevant:
leucine deprivation to simulate amino acid restriction, and a more severe nutrient deprivation by
culturing cells in Earle's Balanced Salt Solution (EBSS) for 16 hours. Both treatments not only
trigger the UPR but also activate the broader integrated stress response (ISR), resulting in the
induction of autophagy (Pakos-Zebrucka *et al*, 2016). RBBP5 depletion impaired the
transcriptional response to both leucine deprivation and EBSS, leading to a reduced induction of
autophagy-promoting genes (**Fig. 6D**). To quantify the autophagic flux, we utilized a tandem LC3
reporter mCherry-GFP-LC3 where an increase in the number of red-fluorescent cytosolic puncta
indicates increased autolysosome formation and autophagic flux (**Figs. 6E and S8G**) (Pakos-
Zebrucka *et al.*, 2016). Consistent with the qPCR results, RBBP5 depleted MEFs failed to induce
a robust general autophagy in response to leucine deprivation, and further exhibited lower basal
level of autophagic flux (**Fig. 6F**). Western blot analysis of p62/SQSTM1, a marker of autophagic
flux due to its degradation during autophagy (Bjørkøy *et al*, 2009), showed a significant reduction
in wild-type, but not RBBP5 depleted cells following leucine deprivation, thus confirming the
defective autophagy phenotype associated with RBBP5 depletion (**Fig. S8H, I**). Since prolonged
nutrient deprivation, such as 16h of EBSS treatment, can robustly induce both general autophagy
and autophagy of the ER (ER-phagy) (Liang *et al*, 2020), we further measured ER-phagy by
utilizing a previously published ER-phagy reporter system where a stably expressed ER-targeting
mCherry-RAMP4 fusion is cleaved to free mCherry during starvation-induced ER-phagy (Liang *et
al*, 2018) (**Fig. 6G**). As shown in **Fig. 6H, I**, RBBP5 depletion prevented the cleavage of mCherry
in response to EBSS, indicating an impaired ER-phagy. Western blot analysis of p62/SQSTM1
further showed undetectable levels in wild-type cells following EBSS treatment, whereas a
significant level remained in RBBP5 depleted cells under the same conditions, thus confirming a
reduction of the general autophagy in RBBP5 depleted cells in response to EBSS (**Fig. 6H**).
Consistent with weakened stress response and impaired autophagy, RBBP5 depleted MEFs
displayed reduced cell survival under nutritional stress (**Fig. S8J**). In sum, our results demonstrate
RBBP5 is indispensable for regulating the transcriptional responses to a plethora of proteotoxic
stresses in a cell-autonomous manner.

Reduced RBBP5 expression is associated with aging in mice and hepatic steatosis in
humans.

The roles of RBBP5 in diseases are incompletely understood (Cenik & Shilatifard, 2021;
Shilatifard, 2012b). Since the decline of proteostasis and stress response is causally associated
with both aging and metabolic diseases (Chen *et al*, 2020; Ghemrawi *et al*, 2018; Kourtis &
Tavernarakis, 2011; Mohan *et al*, 2019), we aim to uncover potential implications of RBBP5 in
these two processes. Intriguingly, in a recent study, RBBP5 (and DPY30) is predicted to be a
positive regulator of maximum lifespan across 26 different species, with its major effect predicted
to be related to liver (Lu *et al*, 2022). Consistent with this prediction, an overall reduction of *Rbbp5*
mRNA level is observed in the liver of aging mice (Almanzar *et al*, 2020), largely concordant with
the expression dynamics of 97 stress response genes (those genes exhibiting 12h expression,
directly bound by RBBP5 and can be induced by ER stress) (**Fig. 7A**).

Leveraging human gene-derived correlations across tissues (GD-CAT) dataset (Zhou *et al*,
2024), we identified in human liver, RBBP5 expression is positively correlated with those involved
in proteostasis (such as ribosome biogenesis, translation regulation) and mRNA metabolism
(mRNA splicing and processing) and negatively associated with those implicated in immune
response (acute phase response, complement activity, and immunoglobin activity) (**Fig. S9**).

These results are in line with our RNA-seq and ChIP-seq data obtained from *Rbbp5* [Flox] and *Rbbp5*
[LKO] mice (**Figs. 2C, 3I, 4D**). In humans with NAFLD/MAFLD (Ahrens *et al*, 2013), compared to
healthy obese, RBBP5 mRNA level is significantly reduced in steatosis subjects (**Fig. 7B**),
concordant with reduced stress gene expression in the same cohorts (**Fig. 7C-F**). Overall, a
positive correlation between *Rbbp5* and the eigengene expression of stress response genes
[eigengene (Langfelder & Horvath, 2007) is the principle component of 97 stress response genes
expression in each subject] is observed across all subjects (**Fig. 7D**). Collectively, these data
suggests that downregulation of *Rbbp5* and proteostatic stress response genes expression is
associated with aging and MAFLD in mice and humans, respectively.

Discussion:

While the epigenetic landscape of the mammalian ~24h circadian clock was well-studied (Aguilar-
Arnal & Sassone-Corsi, 2015; Koike *et al.*, 2012; Masri *et al*, 2015; Nakahata *et al*, 2008;
Takahashi, 2017), the epigenome of the mammalian ~12h ultradian oscillator remains completely
unexplored. In this study, we uncovered that unlike the circadian clock, which was predominantly
regulated by histone acetyltransferase (HAT) (Doi *et al*, 2006; Stashi *et al*, 2014; Zhu *et al*, 2015a)
and histone deacetylase (HDAC) (Alenghat *et al*, 2008; Feng *et al*, 2011; Nakahata *et al.*, 2008),
the epigenome of the 12h oscillator is marked by COMPASS-mediated histone methylation,
specifically H3K4me3 at the proximal promoters. Loss of one of the essential subunits of
COMPASS, RBBP5, abolished more than 90% of the hepatic ~12h transcriptome, while having
no effects on the core circadian clock. Knocking down RBBP5 in MEFs further dampened the cell-
autonomous 12h oscillations, but not the circadian rhythms. We further expanded our findings to
the epigenetic regulation of response to acute and chronic proteotoxic stress and found both
increased RBBP5 promoter recruitment and H3K4me3 level are associated with the transcription
activation of adaptive stress response genes. Subsequently, depletion of RBBP5 significantly
dampened the adaptive transcriptional responses to various proteotoxic stresses.

In our study, we uncovered distinct roles for H3K4me3 and H3K9Ac/H3K27Ac in regulating
the 12h oscillator and transcriptional responses to acute proteotoxic stress. While all three histone
modifications—H3K4me3, H3K9Ac, and H3K27Ac—are traditionally linked to transcriptional
activation (Chen *et al*, 2022; Gates *et al.*, 2017a; Gates *et al.*, 2017b), our findings suggest that
H3K9Ac and H3K27Ac are dispensable for the transcriptional regulation of proteostasis under
both basal (the 12h oscillator) and stress conditions. Interestingly, recent research by (Policarpi
et al, 2024) demonstrated that among various epigenetic marks associated with transcriptional
activation, the installation of H3K4me3 at promoters resulted in the most potent transcriptional
activation. This not only underscores the causal role of H3K4me3 in gene activation but also
highlights the differential impacts of specific histone modifications on transcription. We speculate
that among the many histone modifications, H3K4me3 has been evolutionarily selected as the
primary epigenetic modification for stress response due to its ability to trigger the most robust
gene activation, thereby providing a rapid adaptive advantage under adverse environmental
conditions. Future investigations are needed to further test this hypothesis.

The physiological and pathological roles of mammalian COMPASS subunits, particularly
RBBP5, remain incompletely understood (Cenik & Shilatifard, 2021). While extensive research
has primarily focused on their contributions to organogenesis, early development, and
neurodevelopmental disorders, our study significantly broadens the scope of RBBP5's
physiological roles by uncovering its involvement in proteostasis and hepatic metabolism. This

expansion not only advances our understanding of RBBP5's function but also opens new avenues
for exploring its broader impact on cellular and systemic processes.

Of particular interest is a recent study that identified heterozygous loss-of-function mutations
in *RBBP5* in humans, which were primarily linked to neurodevelopmental disorders (Huang *et al*).
This finding raises intriguing questions about the full spectrum of RBBP5's physiological roles.
Given our discovery of RBBP5's involvement in metabolic regulation, it is tempting to hypothesize
that individuals with RBBP5 loss-of-function mutations may also exhibit metabolic defects. Such
a connection would have significant implications, suggesting that RBBP5 plays a multifaceted role
in maintaining both neural and metabolic homeostasis. Future research should investigate
whether these mutations influence metabolic pathways, potentially contributing to a broader
spectrum of clinical symptoms in affected individuals. This line of inquiry could ultimately lead to
a deeper understanding of how COMPASS subunits like RBBP5 integrate signals across different
tissues to maintain overall physiological balance.

Finally, we propose that the 12h oscillator serves as an effective 'discovery tool' for uncovering
previously unknown mechanisms of proteostasis regulation. By leveraging this approach, we
recently identified a novel XBP1s-SON axis, which links nuclear speckle LLPS dynamics with the
UPR (Dion et al., 2022b). Herein, using a similar approach, we identified RBBP5 as an epigenetic
regulator of proteostasis. We believe that continued exploration of the 12h oscillator will unveil
further hidden principles governing proteostasis.

Methods:

Plasmids

pCDH-EF1a-mCherry-EGFP-LC3B was a gift from Sang-Hun Lee (Addgene plasmid # 170446;
http://n2t.net/addgene:170446; RRID: Addgene_170446) (Wulansari *et al*, 2021). pLenti-X1-
hygro-mCherry-RAMP4 was a gift from Jacob Corn (Addgene plasmid # 118391;
http://n2t.net/addgene:118391; RRID:Addgene_118391) (Liang *et al.*, 2018).

The generation of *Rbbp5* (flox/flox) mice

The Rbbp5-flox allele was generated by the Mouse Embryo Services Core (University of
Pittsburgh, Department of Immunology). The targeting strategy is based on the Easi-CRISPR
method (Quadros *et al*, 2017). In brief, fertilized embryos from C57BL/6J background, produced
by natural mating, were microinjected in the pronuclei with a mixture of 0.67 μM EnGen Cas9
protein (New England Biolabs, Cat.No. M0646T), two Cas9 guides RNA: Rbbp5-695rev and
Rbbp5-951forw (21.23 ng/μl each) and a long single stranded oligonucleotides Rbbp5-flox-
ssODN (5 ng/μl). The long single stranded oligonucleotide encoding the donor sequence was
synthesized by Integrated DNA Technologies (IDT).

To produce the sgRNA, a double strand linear DNA template is created by annealing of the
following target specific oligonucleotides with a common primer (gRNA-Scaffold-R: 5'-
AAAAAAGCACCGACTCGGTGCCACTTTTTCAAGTTGATAACGGACTAGCCTTATTTTAACTTG
CTATTTCTAGCTCTAAAAC-3') containing the full tracrRNA sequence and then PCR amplified as
previously described (Pelletier *et al*, 2015). The sgRNA was synthesized using HiScribe T7 Quick
High Yield RNA Synthesis Kit (New England Biolabs, Cat. No. E2050S).

T7-Rbbp5-695rev sequence:

TAATACGACTCACTATAGGAAGCATTCTAAGATTAAACCGTTTTAGAGCTAGAAATAGCA

T7-Rbbp5-951forw sequence:

TAATACGACTCACTATAGGAGTAACAGATAGTATTCCGAGTTTTAGAGCTAGAAATAGCA

Rbbp5-flox-ssODN sequence:

gcagttaataaacttgaagtgtttatatagaaaatacatcaggatggcaaccaaacccttttagatacctgagatttttattctcttatcccag
gtATAACTTCGTATAGCATACATTATACGAAGTTATgaattcttaatcttagaatgcttagtctgtttcatgctgaagata
agttgtcagaactcacattgttactgatttttcgttcagtgactcgtctgttttgtcataggaaaattggagtgcatttgcaccagacttcaaag
agttggatgaaaatgtagaatatgaggaaagagaatcagaatttgatattgaggatgaagataagagtgagcctgagcaaacaggt
gatgcctctcagtaacagatagtattgaattcATAACTTCGTATAGCATACATTATACGAAGTTATccgaagggaag
gactgtcttactcaatgctctttgaattagtgaatgcatttatttactttgtgttgtaactgtgctcaaagcttcatctgccacgtttccctcacttgt
acatagttttcacagtgcttgaagagtcttactg

The injected zygotes were cultured overnight, the next day the embryos that developed to the
2-cell stage were transferred to the oviducts of pseudo pregnant CD1 female surrogates.
Following optimization of the cycling conditions, the potential founder mice were genotype with
the following primers pair, using Q5 High-Fidelity DNA Polymerase (NEW ENGLAND BIOLABS,
Cat. no. M0491L). The body weight of male adult mice was measured weekly.

Rbbp5 genotyping forward primer: AGTTCAGAGCTGGTTTCCAAC

Rbbp5 genotyping reverse primer: AGGTAATGCCCATGTGAGCC

Biological rhythm study in mice

All mice used for biological rhythm study are in C57BL/6J background, male and between 3 and
4 months of age. Wild-type C57BL/6J mice (n=12), *Rbbp5* (flox/flox); Alb-CRE (-/-) (n=48) and
Rbbp5 (flox/flox); Alb-CRE (+/-) mice (n=48) [*Rbbp5* (flox/flox); Alb-CRE (-/-) and *Rbbp5* (flox/flox);
Alb-CRE (+/-) mice were littermates] were first entrained under LD12:12 conditions for 2 weeks
before transferred to constant darkness for 24hrs. Mice were then sacrificed via cervical
dislocation at a 2h interval [for *Rbbp5* (flox/flox); Alb-CRE (-/-) and *Rbbp5* (flox/flox); Alb-CRE (+/-)
mice] or 4h interval (for wild-type C57BL/6J mice) for a total of 48 hours under constant darkness.
Mice were fed *ad libitum* during the entire experiment. The animal studies were carried out in
accordance with the National Institutes of Health guidelines and were granted formal approval by
the University of Pittsburgh's Institutional Animal Care and Use Committee (approval number
IS00013119 and IS00013119).

Food intake and locomotor activity monitoring

551   Promethion Multi-plexed Metabolic Cage System was used for real-time measuring of food intake
552   and locomotor activity. Male *Rbbp5 ^{Flox}* and *Rbbp5 ^{LKO}* mice between 3 and 4 months of age were
553   acclimated to the chambers and ad libitum food intake was monitored for 48 hours under LD12:12
554   followed by 48 hours of constant darkness. n=6 for *Rbbp5 ^{Flox}* and n=8 for *Rbbp5 ^{LKO}* mice for
555   LD12:12 study. n=5 for *Rbbp5 ^{Flox}* and n=3 for *Rbbp5 ^{LKO}* mice for DD study.

ER stress induction study in mice

For acute ER stress induction experiment, male littermates *Rbbp5* (flox/flox); Alb-CRE (-/-) and
Rbbp5 (flox/flox); Alb-CRE (+/-) mice between 5 and 7 months of age were randomly divided into

two groups, and intraperitoneally injected with 0.05mg/kg body weight of tunicamycin dissolved
in 500ul 3% DMSO in PBS or vehicle control (3% DMSO in PBS), respectively. Mice were injected
with Tu between 9~10am in the morning and 8 hours later dissected for transcriptomic and
western blot analysis for different tissues. The sample size is n=3 for *Rbbp5* ^{Flox} DMSO, n=3 for
Rbbp5 ^{Flox} Tu, n=4 for *Rbbp5* ^{LKO} DMSO, n=4 for *Rbbp5* ^{LKO} Tu.

For experiments involving RBBP5 liver-overexpressing mice, AAV8-TBG-EGFP and AAV8-
TGB-mRBBP5 viruses were customarily designed and produced from Vector Biolab (Malvern,
PA). Male wildtype C57BL/6J mice between 2 and 3 months of age were tail-vein injected with
0.5x10^11 genome copies of AAV8-TBG-EGFP or AAV8-TGB-mRBBP5 virus diluted in 100 µl of
PBS. 30 days later, mice injected with either viruses were randomly divided into two cohorts and
intraperitoneally injected with 0.05mg/kg body weight of tunicamycin dissolved in 500ul 3% DMSO
in PBS or vehicle control (3% DMSO in PBS). Mice were injected with Tu between 9~10am in the
morning and 8 hours later dissected for transcriptomic and western blot analysis for different
tissues. The sample size is n=3 for AAV8-TBG-EGFP DMSO, n=4 for AAV8-TBG-EGFP Tu, n=4
for AAV8-TGB-mRBBP5 DMSO and n=4 for AAV8-TGB-mRBBP5 Tu.

For chronic ER stress induction experiment, male littermates *Rbbp5* (flox/flox); Alb-CRE (-/-)
and *Rbbp5* (flox/flox); Alb-CRE (+/-) mice between 5 and 7 months of age were intraperitoneally
injected with 0.025mg/kg body weight of tunicamycin dissolved in 500ul 3% DMSO in PBS or
vehicle control (3% DMSO in PBS) daily for six consecutive days. Mice were injected with Tu
between 9~10am in the morning and 24 hours after the last injection dissected for transcriptomic,
western blot analysis, histology for liver tissues. The sample size is n=4 for *Rbbp5* ^{Flox} DMSO, n=4
for *Rbbp5* ^{Flox} Tu, n=2 for *Rbbp5* ^{LKO} DMSO, n=3 for *Rbbp5* ^{LKO} Tu.

Oil Red O Staining

Frozen sections were rinsed with PBS and then fixed with 10% neutral buffered formalin for 30
min at RT. After washing twice with deionized water, 60% isopropanol was applied to the fixed
cells for 5 min, followed by a freshly prepared working solution of 1.5 mg/ml Oil Red O in
isopropanol for 5 min at RT. The stained, fixed tissues were then rinsed with tap water until clear,
covered with tap water and viewed on a phase contrast microscope. Size and areas of lipid droplet
were quantified by customarily written pipelines in CellProfiler (v4.1.3).

Generation of stable autophagy reporter and ER-phagy reporter cell line

Primary MEFs were isolated from *Rbbp5* (flox/flox); Rosa26 CRE (+/+) mice and immortalized
with SV40 lentivirus as previously described (Xu, 2005). For mCherry-EGFP-LC3B-expressing
autophagy reporter MEFs, lentiviruses packaged in HEK293T cells with co-transfection of pCDH-
EF1a-mCherry-EGFP-LC3B, pMD2.G and psPAX2 plasmids were used to infect *Rbbp5* (flox/flox);
Rosa26 CRE (+/+) MEFs with a multiplicity of infection (MOI) of 3 for three times to achieve near
complete infection. For mCherry-RAMP4-expressing ER-phagy reporter MEFs, lentiviruses
packaged in HEK293T cells with co-transfection of pLenti-X1-hygro-mCherry-RAMP4, pMD2.G
and psPAX2 plasmids were used to infect *Rbbp5* (flox/flox); Rosa26 CRE (+/+) MEFs with a
multiplicity of infection (MOI) of 3. Stable MEFs were selected in the presence of 200µg/ml
hygromycin.

siRNA Transient Transfections

Immortalized MEFs isolated from wild-type C57BL/6J mice were transfected with 10µM of
different siRNAs for 24~48 hours with Lipofectamine RNAiMAX reagents (Life technologies) per
the manufacturer's instructions. Source of siRNA are as follows: siGENOME non-targeting siRNA
pool (Dharmacon, D-001206-1305), siGENOME SMARTpool mouse *Rbbp5* siRNA (Dharmacon,
L-054560-01-0005), siGENOME SMARTpool mouse *Ash2l* siRNA (Dharmacon, L-048754-01-
0005), siGENOME SMARTpool mouse *Setd1a* siRNA (Dharmacon, L-051358-01-0005),
siGENOME SMARTpool mouse *Kat2a/Gcn5* siRNA (Dharmacon, L-040665-01-0005),
siGENOME SMARTpool mouse *Kat2b/Pcaf* siRNA (Dharmacon, L-049885-01-0005) and
siGENOME SMARTpool mouse *Xbp1* siRNA (Dharmacon, L-040825-00-0005). MEFs were
transfected with 10µM of different siRNAs for 48 hours and treated with DMSO or 100ng/ml of
tunicamycin for 8h.

Real-time Luminescence Assay

Stable *Manf*-dluc MEFs were transfected with non-targeting, *Rbbp5,* or *Ash2l* siRNA for 48 hours
in DMEM (4.5g/L glucose) supplemented with 10% FBS and treated with 25ng/ml tunicamycin in
DMEM for 2h before subjected to real-time luminescence assay using a Lumicycle (Actimetrics)
as previously described (Zhu *et al*, 2015b). Briefly, after tunicamycin treatment, MEFs were
washed with 1x PBS and cultured with DMEM (4.5g/L glucose) supplemented with 0.1 mM
luciferin and 10mM HEPES buffer in 35 mm tissue culture dishes in the absence of serum and
transferred immediately to Lumicycle for real-time luminescence analysis. Periods of oscillation
were identified by embedded Periodogram function.

Immunofluorescence (IF)

Immunofluorescence was performed as previously described (Dion *et al.*, 2022a). Briefly, liver
OCT sections were fixed in cold acetone for 10 mins at -20 ºC. The sections were then air dried,
rehydrated with PBS and permeabilized with PBS+ 0.1% Triton X-100. The sections were then
blocked with 10% goat serum at room temperature for 1 hour. Primary antibody against RBBP5
(Bethyl, A300-109A) was added to the OCT section at 1:1000 dilution overnight at 4 ºC. Next day,
sections were washed three times with PBS and stained with Alexa-555 anti-rabbit secondary
antibody at room temperature for 2 hours. After that, the sections were washed with PBS three
times and counterstained with DAPI before mounting (with ProlongGold Glass) and imaging using
Leica SP8 lightening confocal microscope (Leica Microsystems).

Time-lapse microscopy for autophagy reporter MEFs

Time-lapse imaging was performed using SP8 lightening confocal microscope (Leica
Microsystems) with Okolab stage top incubator to maintain constant CO_2 (5%), temperature (37
ºC) and humidity (90%). mCherry-EGFP-LC3B-expressing *Rbbp5* (flox/flox); Rosa26 CRE (+/+)
MEFs were plated into 8-well chamber slides in full DMEM media and treated with (to deplete
RBBP5) or without tamoxifen (3µM/ml) for seven days before replacing with full DMEM or leucine-
free media counterstained with DAPI and images were taken every 2 hours for a total of 16 hours.
The number of cytosolic autophagosomes (GFP and mCherry double positive foci) and
autolysosomes (GFP negative and mCherry positive foci) per cell were quantified manually.

Immunoblot

Nuclear extracts were made from liver according to previously published protocol (Malovannaya
et al, 2011). Whole cell lysates were lysed in RIPA buffer as previously described (Dion *et al.*,
2022a). Protein concentrations were determined by Bradford assays (Bio-Rad), and aliquots were

snap-frozen in liquid nitrogen and stored at -80°C until usage. Immunoblot analyses were
performed as described previously (Zhu *et al.*, 2015a). Briefly, 25~50µg proteins separated by
4~20% gradient SDS-PAGE gels (Biorad) were transferred to nitrocellulose membranes, blocked
in TBST buffer supplemented with 5% bovine serum albumin (BSA) or 5% fat-free milk and
incubated overnight with primary antibody anti-PERK (Cell signaling, #3192), anti-phospho-PERK
(Thr908) (Thermo Fisher, MA5-15033), anti-ATF4 (Cell signaling, #11815), anti-IRE1α (Cell
signaling, #3294), anti-phospho-IRE1α (Ser724) (ABclonal, AP0878), anti-XBP1s (Biolegend,
658802), anti-RBBP5 (Bethyl, A300-109A), anti-RBBP5 (D3I6P) (Cell signaling, #13171), anti-
WDR5 (Cell signaling, #13105), anti-ASH2L (Bethyl, A300-489A), anti-ASH2L (Bethyl, A300-
112A), anti-SETD1A (Diagenode, CS-117-100), anti-SETD1B (Diagenode, CS-118-100), anti-
MLL1-c (D6G8N) (Cell signaling, #14197), anti-mCherry (E5D8F) (Cell signaling, #43590), anti-
p62/SQSTM1 (Santa Cruz, sc-28359) and anti-β-ACTIN (Cell signaling, #12620) at 4°C. Blots
were incubated with an appropriate secondary antibody coupled to horseradish peroxidase at
room temperature for 1 hour, and reacted with ECL reagents per the manufacturer's (Thermo)
suggestion and detected by Biorad ChemiDoc MP Imaging System.

Co-Immunoprecipitation (Co-IP)

Nuclear extracts (NE) were made from liver according to our published protocol (Malovannaya *et*
al., 2011). Protein concentrations were determined by Bradford assays (Bio-Rad), and aliquots
were snap-frozen in liquid nitrogen and stored at -80°C until usage. 200 µg of NE was used for
per IP. 5 µg of different antibodies or control IgG were coupled to 1.5 mg of magnetic Dynabeads
(Life Technologies) for every IP using Dynabeads Antibody Coupling Kit (Life Technologies) per
manufactures' protocol. Co-IP was essentially carried out as previously described (Zhu *et al*,
2015c) except that coupled antibody Dynabeads was added to the NEs for incubation. The
antibodies used for Co-IP are as follows: anti-XBP1s (Biolegend, 658802).

qRT-PCR

Total mRNA was isolated from murine embryonic fibroblasts (MEFs) or mouse liver with PureLink
RNA mini kit (Life Technologies) with additional on-column DNase digestion step to remove
genomic DNA per the manufacturer's instructions. Reverse transcription was carried out using
5µg of RNA using Superscript III (Life Technologies) per the manufacturer's instructions. For gene
expression analyses, cDNA samples were diluted 1/30-fold (for all other genes except for 18sRNA)
and 1/900-fold (for 18sRNA). qPCR was performed using the SYBR green system with sequence-
specific primers. All data were analyzed with 18S or *β-actin* as the endogenous control. qPCR
primer sequences are as follows and all primers span introns, except for primers for quantifying
pre-mRNAs:
Mouse *Rbbp5* forward primer: CAGAGCCTATGCAGAAGCTGCA
Mouse *Rbbp5* reverse primer: CTTCACCAGGTTGCCAATGCTC
Mouse *Ddit3* forward primer: GGAGGTCCTGTCCTCAGATGAA
Mouse *Ddit3* reverse primer: GCTCCTCTGTCAGCCAAGCTAG
Mouse *Creld2* forward primer: CTACACCAAGGAGAGTGGACAG
Mouse *Creld2* reverse primer: TCTGTCTCCTCAAAGCCTTCCG
Mouse *Hspa5* forward primer: TGTCTTCTCAGCATCAAGCAAGG
Mouse *Hspa5* reverse primer: CCAACACTTCCTGGACAGGCTT
Mouse *Dnajb11* forward primer: TGTGACCGTCTCACTGGTTGAG
Mouse *Dnajb11* reverse primer: CCCTTTCTTCCACAGCTTGGCT
Mouse *Pdia4* forward primer: TGGGCTCTTTCAGGGAGATGGT

Mouse *Pdia4* reverse primer: GGGAGACTTTCAGGAACTTGGC
Mouse *Hsp90b1* forward primer: GTTTCCCGTGAGACTCTTCAGC
Mouse *Hsp90b1* reverse primer: ATTCGTGCCGAACTCCTTCCAG
Mouse *Syvn1* forward primer: CCAACATCTCCTGGCTCTTCCA
Mouse *Syvn1* reverse primer: CAGGATGCTGTGATAAGCGTGG
Mouse *Gdf15* forward primer: ACTCAGTCCAGAGGTGAGATTG
Mouse *Gdf15* reverse primer: GGGGCCTAGTGATGTCCCAG
Mouse *Il1a* forward primer: ACGGCTGAGTTTCAGTGAGACC
Mouse *Il1a* reverse primer: CACTCTGGTAGGTGTAAGGTGC
Mouse *Il1b* forward primer: TGGACCTTCCAGGATGAGGACA
Mouse *Il1b* reverse primer: GTTCATCTCGGAGCCTGTAGTG
Mouse *Fga* forward primer: GGATTCTAACTCACTGACCAGGA
Mouse *Fga* reverse primer: CCTCAGGATCTCAATTCTGCGC
Mouse *C3* forward primer: CGCAACGAACAGGTGGAGATCA
Mouse *C3* reverse primer: CTGGAAGTAGCGATTCTTGGCG
Mouse *Tifa* forward primer: AGCAAGAAACCAGTTTGATGGTAG
Mouse *Tifa* reverse primer: GCAACAGGAACTGATACTCTCCG
Mouse *Fgl1* forward primer: GGAAACTGTGCTGAGGAAGAGC
Mouse *Fgl1* reverse primer: TCCGTTTCTGCCCTGTAGGAAC
Mouse *Fgb* forward primer: TGACACCTCCATCAAGCCGTAC
Mouse *Fgb* reverse primer: GGTCCCATTTCCTGCCAAAGTC
Mouse *Il6* forward primer: CTTCCATCCAGTTGCCTTCT
Mouse *Il6* reverse primer: CTCCGACTTGTGAAGTGGTATAG
Mouse *Herpud1* forward primer: acctgagccgagtctaccc
Mouse *Herpud1* reverse primer: aacagcagcttcccagaataaa
Mouse *Nbr1* forward primer: GGATTTAAAGCACCTCCTGATTCC
Mouse *Nbr1* reverse primer: ATTGGGTCCCACTCAGGTCT
Mouse *Sesn2* forward primer: CTGGCCGAACTCATCCAAG
Mouse *Sesn2* reverse primer: CTGCCTCATGCGTTCCATC
Mouse *Sqstm1* forward primer: AACAGATGGAGTCGGGAAAC
Mouse *Sqstm1* reverse primer: AGACTGGAGTTCACCTGTAGA
Mouse *Stbd1* forward primer: CTGGGAAGTAGATGGGAAAGTG
Mouse *Stbd1* reverse primer: TCGGTGTTTGTGGTGTAGTG
Mouse *Sirt1* forward primer: TGACAGAACGTCACACGCCA
Mouse *Sirt1* reverse primer: ATTGTTCGAGGATCGGTGCCA
Mouse *Gcn5* forward primer: TTGATTGAGCGCAAACAGGC
Mouse *Gcn5* reverse primer: CAGCCTGTCTCTCGAATGCC
Mouse *Pcaf* forward primer: CGTGAAGAAGGCGCAGTTG
Mouse *Pcaf* reverse primer: CAGGACTCCTCTGCCTTGC
Mouse *Ash2l* forward primer: gctgtgtctgctagtgggaac
Mouse *Ash2l* reverse primer: catcttgctgcttacgcttg
Mouse *Setd1a* forward primer: gggtagcaccccctactctc
Mouse *Setd1a* reverse primer: gggtttgaaggaggttgaagt
Mouse *Eif2ak3* forward primer: ccttggtttcatctagcctca
Mouse *Eif3ak3* reverse primer: atccagggaggggatgat
Mouse total *Xbp1* forward primer: gggtctgctgagtcc
Mouse total *Xbp1* reverse primer: cagactcagaatctgaagagg

Mouse *Sec23b* forward primer: tgaccaaactggacttctgga
Mouse *Sec23b* reverse primer: aaagaatctcccatcaccatgt
Mouse *Manf* forward primer: gacagccagatctgtgaactaaaa
Mouse *Manf* reverse primer: tttcacccggagcttcttc
Mouse *Hyou1* forward primer: GAGGCGAAACCCATTTTAGA
Mouse *Hyou1* reverse primer: GCTCTTCCTGTTCAGGTCCA
Mouse *Atf4* forward primer: CCACTCCAGAGCATTCCTTTAG
Mouse *Atf4* reverse primer: CTCCTTTACACATGGAGGGATTAG
Mouse *Atf6* forward primer: CATGAAGTGGAAAGGACCAAATC
Mouse *Atf6* reverse primer: CAGCCATCAGCTGAGAATTAGA
Mouse 18s RNA forward primer: ctcaacacgggaaacctcac
Mouse 18s RNA reverse primer: cgctccaccaactaagaacg
Mouse *β-actin* forward primer: aaggccaaccgtgaaaagat
Mouse *β-actin* reverse primer: gtggtacgaccagaggcatac

RNA-Seq to identify RBBP5-dependent 12h hepatic transcriptome

Mouse liver tissues were collected from *Rbbp5* ^{Flox} (n=2) and *Rbbp5* ^{LKO} (n=2) mice at 2h intervals
for a total of 48 hours under constant darkness condition. Total RNA was isolated from mouse
liver with miRNeasy Mini Kit (Qiagen) per the manufacturer's instructions. RNA was assessed for
quality using an Agilent TapeStation 4150/Fragment Analyzer 5300 and RNA concentration was
quantified on a Qubit FLEX fluorometer. Libraries were generated with the Illumina Stranded
mRNA Library Prep kit (Illumina: 20040534) according to the manufacturer's instructions. Briefly,
200 ng of input RNA was used for each sample. Following adapter ligation, 12 cycles of indexing
PCR were completed, using IDT for Illumina RNA UD Indexes (Illumina: 20040553-6). Library
quantification and assessment was done using a Qubit FLEX fluorometer and an Agilent
TapeStation 4150/Fragment Analyzer 5300. The prepared libraries were pooled and sequenced
using NoveSeq 6000 (Illumina), generating an average of 40 million 2×100 bp paired end reads
per sample. RNA-Seq library preparation and sequencing were performed at UPMC Genome
Center. Raw RNA-seq FASTQ files were analyzed by FastQC for quality control. Adaptors and
low-quality reads were filtered by Trimmomatic (Bolger *et al*, 2014b). Then the processed reads
were aligned by HISAT2 (Kim *et al*, 2015) against mouse reference mm10. Gene and isoform
FKPM values were calculated by Cufflinks. Since the background gene expression level is
FPKM=0.1 in mouse liver (Pan *et al.*, 2020), only those genes with average gene expession
across 48 hours in Rbbp5 ^{Flox} mice larger than 0.1 were included for rhythm-identification analysis.
Averaged FPKM values at each time were used for cycling transcripts identification by the
eigenvalue/pencil method. Raw temporal data was subject to polynomial detrend (n=2) first, and
superimposed oscillations were identified using previously described eigenvalue/pencil method
on the detrended dataset (Antoulas *et al.*, 2018a; Zhu *et al.*, 2017a). Specifically, three oscillations
were identified from each gene. Criterion for circadian gene is period between 21h to 27h, decay
rate between 0.9 and 1.1; for ~12hr genes: period between 10h to 13h, decay rate between 0.9
and 1.1. The relative amplitude is calculated via dividing the amplitude by the mean expression
value of each gene. All analysis were performed in MatlabR2017A. Heat maps were generated
by Gene Cluster 3.0 and TreeView 3.0 alpha 3.0 using log2 mean-normalized values. RAIN
analysis   was   performed   as   previously   described   in   Bioconductor   (3.4)
(http://www.bioconductor.org/packages/release/bioc/html/rain.html) (Thaben & Westermark,
2014) using the full detrended dataset with duplicated at each time point.

For the eigenvalue method, every gene consists of multiple superimposed oscillations.
Therefore, we define a circadian gene as any gene that exhibits a superimposed circadian rhythm,
regardless of its relative amplitude compared to other superimposed oscillations. Similar
definitions apply to 12hr genes. Under this definition, a gene can be both a circadian and 12hr-
cycling gene. By comparison, we define a dominant circadian gene as any gene whose
superimposed circadian rhythm has the largest amplitude among all oscillations. Similar
definitions also apply to 12hr genes. Under this definition, dominant circadian and dominant 12hr
genes are mutually exclusive.

RNA-seq for identifying RBBP5-dependent acute transcriptional response to ER stress

Mouse liver tissues were collected from *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice intraperitoneally treated
with or without Tu for 8 hours. Total mRNA was isolated from mouse liver with PureLink RNA mini
kit (Life Technologies) with additional on-column DNase digestion step to remove genomic DNA
per the manufacturer's instructions. Detailed procedures for transcriptome sequencing were
described previously(Liu *et al*, 2024) and were supported by the Genomics and Systems Biology
Core of the Pittsburgh Liver Research Center. Briefly, the mRNA samples per mouse were
processed into short-read libraries using the Bio-Rad SEQuoia Dual Indexed Primers. Next, the
circularization was performed using the Element Biosciences Adept Library Compatibility Kit v1.1
based on the manufacturer's instructions, followed by the quantification using qPCR with the
provided standard. Finally, the libraries were measured by the Element Biosciences AVITI system
to sequence paired end reads with 75 bp each. The raw FASTQ files from RNA-seq were trimmed
by Trimmomatic (Bolger *et al*, 2014a) to filter out low-quality reads. The surviving reads were then
aligned to the mouse mm10 reference genome by STAR aligner(Dobin *et al*, 2013). Fragments
Per Kilobase of transcript per Million mapped reads (FPKM) values per library were quantified by
the tool Cufflinks(Trapnell *et al*, 2010). Log 2 normalized fold induction and adjusted p values
between *Rbbp5* ^{Flox} DMSO and *Rbbp5* ^{LKO} DMSO, *Rbbp5* ^{Flox} Tu and *Rbbp5* ^{LKO} Tu, *Rbbp5* ^{Flox}
DMSO and *Rbbp5* ^{Flox} Tu, and *Rbbp5* ^{LKO} DMSO and *Rbbp5* ^{LKO} Tu were calculated by using the
iDEP (integrated Differential Expression and Pathway analysis) online application (Ge *et al*, 2018).

Chromatin Immunoprecipitation (ChIP)-Seq and ChIP-qPCR

ChIP for RBBP5 was performed using anti-RBBP5 antibody (Bethyl, A300-109A) as previously
described (Pan *et al.*, 2020). Briefly, mouse liver samples were submerged in PBS + 1%
formaldehyde, cut into small (~1 mm3) pieces with a razor blade and incubated at room
temperature for 15 minutes. Fixation was stopped by the addition of 0.125 M glycine (final
concentration). The tissue pieces were then treated with a TissueTearer and finally spun down
and washed twice in PBS. Chromatin was isolated by the addition of lysis buffer, followed by
disruption with a Dounce homogenizer. The chromatin was enzymatically digested with MNase.
Genomic DNA (Input) was prepared by treating aliquots of chromatin with RNase, Proteinase K
and heated for reverse-crosslinking, followed by ethanol precipitation. Pellets were resuspended
and the resulting DNA was quantified on a NanoDrop spectrophotometer. An aliquot of chromatin
(30 µg) was precleared with protein A agarose beads (Invitrogen). Genomic DNA regions of
interest were isolated using 4 µg of antibody. Complexes were washed, eluted from the beads
with SDS buffer, and subjected to RNase and proteinase K treatment. Crosslinking was reversed
by incubation overnight at 65 °C, and ChIP DNA was purified by phenol-chloroform extraction and
ethanol precipitation. The DNA libraries were prepared at the University of Pittsburgh and
sequenced at Gene by Gene, Ltd per standard protocols. DNA libraries were prepared with
Ovation® Ultralow V2 DNA-Seq library preparation kit (NuGen) using 1ng input DNA. The size

selection for libraries was performed using SPRIselect beads (Beckman Coulter) and purity of the
libraries were analyzed using the High Sensitivity DNA chip on Bioanalyzer 2100 (Agilent). The
prepared libraries pooled and sequenced using Nova-Seq 6000 (Illumina), generating ~30 million
75 bp single-end reads per sample. ChIP-qPCR for MEFs were essentially performed the same
way as previously described with anti-RBBP5 (Bethyl, A300-109A) and anti-H3K4me3 (Active
Motif, 61379) antibodies, except that the MEFs were directly fixed with 1% formaldehyde before
subject to nuclei isolation and chromatin immunoprecipitation. The primers used for ChIP-qPCR
are as follows:
Negative control region 1 forward primer: GCAACAACAACAGCAACAATAAC
Negative control region 1 reverse primer: CATGGCACCTAGAGTTGGATAA
Manf promoter forward primer: CCCTTAAATGGGTCAACGTCTC
Manf promoter reverse primer: GGCGCTAAACCCAAGGAAA
Xbp1 promoter forward primer: TCCGTACGGTGGGTTAGAT
Xbp1 promoter reverse primer: ACCTTCTTCTGTGCCTGTG
Hyou1 promoter forward primer: CCGTGTGGGTACGTCCT
Hyou1 promoter reverse primer: GACGGCTGCTCCATCCT
Sesn2 promoter forward primer: ACAGGAGGCCGGGACTA
Sesn2 promoter reverse primer: CTGGGCTGAAAGGAGTGTCTAT

ChIP-Seq analysis

The sequences identified were mapped to the mouse genome (UCSC mm10) using BOWTIE
function in Galaxy Deeptool (https://usegalaxy.org/) (Ramirez *et al*, 2016). Only the sequences
uniquely mapped with no more than 2 mismatches were kept and used as valid reads. PCR
duplicates were also removed. Peak calling was carried out by MACS2 (version 2.1.1.20160309)
in Galaxy (options --mfold 5, 50 --pvalue 1e-4), on each ChIP-seq file against input. To account
for the different sequencing depths between samples, the BAM files generated from MACS2 were
RPKM normalized to sequencing depth using the bamCoverage function in Galaxy Deeptool and
the bigwig files were generated accordingly. The relative intensity of each RBBP5 binding site is
further calculated via the computeMatrix function with the RPKM normalized bigwig files and bed
files from the peak calling as inputs by calculating the area under the curve.

Gene ontology analysis

DAVID (Version 2021) (Huang da *et al*, 2009) (https://david.ncifcrf.gov) was used to perform Gene
Ontology analyses. Briefly, gene names were first converted to DAVID-recognizable IDs using
Gene Accession Conversion Tool. The updated gene list was then subject to GO analysis using
all Homo Sapiens as background and with Functional Annotation Chart function. GO_BP_DIRECT,
KEGG_PATHWAY or UP_KW_BIOLOGICAL_PROCESS was used as GO categories. Only GO
terms with a p value less than 0.05 were included for further analysis.

Motif analysis

Motif analysis was performed with the SeqPos motif tool (version 0.590) (He *et al*, 2010)
embedded in Galaxy Cistrome using all motifs within the mouse reference genome mm10 as
background. LISA analysis was performed using webtool (http://lisa.cistrome.org/).

Statistical Analysis

Data were analyzed and presented with GraphPad Prism software. Plots show individual data points and bars at the mean and ± the standard error of the mean (SEM). One-tailed t-tests were used to compare means between groups, with significance set at $p < 0.05$. In instances where the p value is not shown, *, **, ***, and **** represent $p < 0.05$, 0.01, 0.001, and 0.0001, respectively.

Acknowledgements:

We would like to thank Dr. Sebastien Gingras (University of Pittsburgh, Department of Immunology) for the design of the targeting strategy and identification of the founders carrying the properly targeted Rbbp5-flox allele. We would like to thank Chunming Bi and Zhaohui Kou of the Mouse Embryo Services Core (University of Pittsburgh, Department of Immunology) for microinjection of zygotes and production of Rbbp5-flox founder mice. We would also like to thank Dr. Heather Ballance, Dr. Naveen GV Kumar, Saad Irfan, C. Dupont, Dr. Bugra Zengin and Casey Edwards for assistance with mice colony maintenance, dissection, intraperitoneal injection and genotyping. For time course transcriptome study, the library generation was performed in the Health Sciences Sequencing Core at UPMC Children's Hospital of Pittsburgh, Rangos Research Center. Services and instruments used in this project were graciously supported, in part, by the University of Pittsburgh, the Office of the Senior Vice Chancellor for Health Sciences, the Department of Pediatrics, the Institute for Precision Medicine, and the Richard K Mellon Foundation for Pediatric Research.

W.D. was supported by fellowship F31 AG080998 through NIA. B. Zhu was supported by grants 1DP2GM140924 and 1R21AG071893 through the NIH, and a grant from Richard King Mellon foundation. This research was supported in part by the University of Pittsburgh Center for Research Computing through the resources provided. Specifically, this work used the HTC cluster, which is supported by NIH award number S10OD028483. This research project was supported in part by the Pittsburgh Liver Research Centre supported by NIH/NIDDK Digestive Disease Research Core Center grant P30DK120531.

Author contributions:

Conceptualization, B.Z.; Methodology, M.J. and B.Z.; Investigation, S.K., M.S., W.D., A.C., H.L., H.W., A.P., I.S., S.L., Y.P., J.J.L., and B.Z.; Writing – Original Draft, B.Z.; Writing – Review & Editing, all authors; Funding Acquisition, S.L., and B.Z.; Resources, M.J. and B.Z.; Supervision, M.J. and B.Z.

Disclosure and competing interests:

The authors have no competing interests to declare.

Data availability:

All raw and processed sequencing data generated in this study have been submitted to the NCBI Gene Expression Omnibus (GEO; http://www.ncbi.nlm.nih.gov/geo/) under accession numbers: GSE276155, GSE276157 and GSE276159. All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials.

References:

Aguilar-Arnal L, Sassone-Corsi P (2015) Chromatin landscape and circadian dynamics: Spatial
and temporal organization of clock transcription. *Proceedings of the National Academy of*
Sciences 112: 6863-6870

Ahrens M, Ammerpohl O, von Schönfels W, Kolarova J, Bens S, Itzel T, Teufel A, Herrmann A,
Brosch M, Hinrichsen H *et al* (2013) DNA Methylation Analysis in Nonalcoholic Fatty Liver
Disease Suggests Distinct Disease-Specific and Remodeling Signatures after Bariatric Surgery.
Cell Metabolism 18: 296-302

Ajoolabady A, Kaplowitz N, Lebeaupin C, Kroemer G, Kaufman RJ, Malhi H, Ren J (2023)
Endoplasmic reticulum stress in liver diseases. *Hepatology* 77: 619-639

Alenghat T, Meyers K, Mullican SE, Leitner K, Adeniji-Adele A, Avila J, Bućan M, Ahima RS,
Kaestner KH, Lazar MA (2008) Nuclear receptor corepressor and histone deacetylase 3 govern
circadian metabolic physiology. *Nature* 456: 997-1000

Almanzar N, Antony J, Baghel AS, Bakerman I, Bansal I, Barres BA, Beachy PA, Berdnik D,
Bilen B, Brownfield D *et al* (2020) A single-cell transcriptomic atlas characterizes ageing tissues
in the mouse. *Nature* 583: 590-595

Antoulas AC, Zhu B, Zhang Q, York B, O'Malley BW, Dacso CC (2018a) A novel mathematical
method for disclosing oscillations in gene transcription: A comparative study. *PLoS One* 13:
e0198503

Antoulas AC, Zhu B, Zhang Q, York B, O'Malley BW, Dacso CC (2018b) A novel mathematical
method for disclosing oscillations in gene transcription: a comparative study. *PLoS One* 13:
e0198503

Asher G, Zhu B (2022a) Beyond circadian rhythms: Emerging roles of ultradian rhythms in
control of liver functions. *Hepatology*

Asher G, Zhu B (2022b) Beyond circadian rhythms: Emerging roles of ultradian rhythms in
control of liver functions. *Hepatology* n/a

Balch WE, Morimoto RI, Dillin A, Kelly JW (2008) Adapting proteostasis for disease intervention.
Science 319: 916-919

Ballance H, Zhu B (2021) Revealing the hidden reality of the mammalian 12-h ultradian
rhythms. *Cellular and Molecular Life Sciences*

Bjørkøy G, Lamark T, Pankiv S, Øvervatn A, Brech A, Johansen T (2009) Monitoring autophagic
degradation of p62/SQSTM1. *Methods Enzymol* 452: 181-197

Bolger AM, Lohse M, Usadel B (2014a) Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics* 30: 2114-2120

Bolger AM, Lohse M, Usadel B (2014b) Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics* 30: 2114-2120

Buttgereit F, Brand MD (1995) A hierarchy of ATP-consuming processes in mammalian cells.
Biochemical Journal 312: 163-167

Cenik BK, Shilatifard A (2021) COMPASS and SWI/SNF complexes in development and
disease. *Nature Reviews Genetics* 22: 38-58

Chen K, Shen W, Zhang Z, Xiong F, Ouyang Q, Luo C (2020) Age-dependent decline in stress
response capacity revealed by proteins dynamics analysis. *Scientific Reports* 10: 15211

Chen YC, Koutelou E, Dent SYR (2022) Now open: Evolving insights to the roles of lysine
acetylation in chromatin organization and function. *Mol Cell* 82: 716-727

Costa-Mattioli M, Walter P (2020) The integrated stress response: From mechanism to disease.
Science 368: eaat5314

Dasgupta S, Lonard DM, O'Malley BW (2014) Nuclear receptor coactivators: master regulators
of human health and disease. *Annu Rev Med* 65: 279-292

Dion W, Ballance H, Lee J, Pan Y, Irfan S, Edwards C, Sun M, Zhang J, Zhang X, Liu S (2022a)
Four-dimensional nuclear speckle phase separation dynamics regulate proteostasis. *Science*
Advances 8: eabl4150

Dion W, Ballance H, Lee J, Pan Y, Irfan S, Edwards C, Sun M, Zhang J, Zhang X, Liu S *et al*
(2022b) Four-dimensional nuclear speckle phase separation dynamics regulate proteostasis.
Science Advances 8: eabl4150

Dion W, Tao Y, Chambers M, Zhao S, Arbuckle RK, Sun M, Kubra S, Nie Y, Ye M, Larsen MB *et*
al (2024) Nuclear speckle rejuvenation alleviates proteinopathies at the expense of YAP1.
bioRxiv: 2024.2004.2018.590103

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras
TR (2013) STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29: 15-21

Doi M, Hirayama J, Sassone-Corsi P (2006) Circadian Regulator CLOCK Is a Histone
Acetyltransferase. *Cell* 125: 497-508

Fang B, Everett LJ, Jager J, Briggs E, Armour SM, Feng D, Roy A, Gerhart-Hines Z, Sun Z,
Lazar MA (2014) Circadian enhancers coordinate multiple phases of rhythmic gene transcription
in vivo. *Cell* 159: 1140-1152

Feng D, Liu T, Sun Z, Bugge A, Mullican SE, Alenghat T, Liu XS, Lazar MA (2011) A circadian
rhythm orchestrated by histone deacetylase 3 controls hepatic lipid metabolism. *Science* 331:
1315-1319

G G, Singh J (2022) Dithiothreitol causes toxicity in C. elegans by modulating the methionine–
homocysteine cycle. *eLife* 11: e76021

Gates LA, Foulds CE, O'Malley BW (2017a) Histone Marks in the 'Driver's Seat': Functional
Roles in Steering the Transcription Cycle. *Trends Biochem Sci* 42: 977-989

Gates LA, Shi J, Rohira AD, Feng Q, Zhu B, Bedford MT, Sagum CA, Jung SY, Qin J, Tsai MJ *et*
al (2017b) Acetylation on histone H3 lysine 9 mediates a switch from transcription initiation to
elongation. *J Biol Chem* 292: 14456-14472

Ge SX, Son EW, Yao R (2018) iDEP: an integrated web application for differential expression
and pathway analysis of RNA-Seq data. *BMC bioinformatics* 19: 1-24

Gehrke N, Schattenberg JM (2020) Metabolic Inflammation—A Role for Hepatic Inflammatory
Pathways as Drivers of Comorbidities in Nonalcoholic Fatty Liver Disease? *Gastroenterology*
158: 1929-1947.e1926

Ghemrawi R, Battaglia-Hsu SF, Arnold C (2018) Endoplasmic Reticulum Stress in Metabolic
Disorders. *Cells* 7

Gong H, He Q, Zhu L, Feng Z, Sun M, Jiang J, Yuan X, Shen Y, Di J (2024) Associations
between systemic inflammation indicators and nonalcoholic fatty liver disease: evidence from a
prospective study. *Frontiers in Immunology* 15

Guo B, Li Z (2014) Endoplasmic reticulum stress in hepatic steatosis and inflammatory bowel
diseases. *Frontiers in Genetics* 5

Han J, Li T, Li Y, Li M, Wang X, Peng C, Su C, Li N, Li Y, Xu Y *et al* (2019) The internal
interaction in RBBP5 regulates assembly and activity of MLL1 methyltransferase complex.
Nucleic Acids Research 47: 10426-10438

He HH, Meyer CA, Shin H, Bailey ST, Wei G, Wang Q, Zhang Y, Xu K, Ni M, Lupien M *et al*
(2010) Nucleosome dynamics define transcriptional enhancers. *Nature Genetics* 42: 343-347

Hetz C, Zhang K, Kaufman RJ (2020) Mechanisms, regulation and functions of the unfolded
protein response. *Nature Reviews Molecular Cell Biology* 21: 421-438

Higa A, Taouji S, Lhomond S, Jensen D, Fernandez-Zapico ME, Simpson JC, Pasquet JM,
Schekman R, Chevet E (2014) Endoplasmic reticulum stress-activated transcription factor
ATF6α requires the disulfide isomerase PDIA5 to modulate chemoresistance. *Mol Cell Biol* 34:
1839-1849

Hu S, Song A, Peng L, Tang N, Qiao Z, Wang Z, Lan F, Chen FX (2023) H3K4me2/3 modulate
the stability of RNA polymerase II pausing. *Cell Research* 33: 403-406

Huang da W, Sherman BT, Lempicki RA (2009) Systematic and integrative analysis of large
gene lists using DAVID bioinformatics resources. *Nat Protoc* 4: 44-57

Huang Y, Jay KL, Huang AY-W, Wan J, Jangam SV, Chorin O, Rothschild A, Barel O, Mariani M,
Iascone M *et al* Loss-of-function in RBBP5 results in a syndromic
neurodevelopmental disorder associated with microcephaly. *Genetics in Medicine*

Hughes ME, DiTacchio L, Hayes KR, Vollmers C, Pulivarthy S, Baggs JE, Panda S, Hogenesch
JB (2009) Harmonics of circadian gene transcription in mammals. *PLoS Genet* 5: e1000442

Jin Q, Yu LR, Wang L, Zhang Z, Kasper LH, Lee JE, Wang C, Brindle PK, Dent SY, Ge K (2011)
Distinct roles of GCN5/PCAF-mediated H3K9ac and CBP/p300-mediated H3K18/27ac in
nuclear receptor transactivation. *Embo j* 30: 249-262

Kim D, Langmead B, Salzberg SL (2015) HISAT: a fast spliced aligner with low memory
requirements. *Nat Methods* 12: 357-360

Klaips CL, Jayaraj GG, Hartl FU (2018) Pathways of cellular proteostasis in aging and disease. *J Cell Biol* 217: 51-63

Koike N, Yoo SH, Huang HC, Kumar V, Lee C, Kim TK, Takahashi JS (2012) Transcriptional architecture and chromatin landscape of the core circadian clock in mammals. *Science* 338: 349-354

Kourtis N, Tavernarakis N (2011) Cellular stress response pathways and ageing: intricate molecular relationships. *The EMBO journal* 30: 2520-2531

Labbadia J, Morimoto RI (2015) The Biology of Proteostasis in Aging and Disease. *Annual Review of Biochemistry* 84: 435-464

Langfelder P, Horvath S (2007) Eigengene networks for studying the relationships between co-expression modules. *BMC Systems Biology* 1: 54

Li Y, Han J, Zhang Y, Cao F, Liu Z, Li S, Wu J, Hu C, Wang Y, Shuai J *et al* (2016) Structural basis for activity regulation of MLL family methyltransferases. *Nature* 530: 447-452

Liang JR, Lingeman E, Ahmed S, Corn JE (2018) Atlastins remodel the endoplasmic reticulum for selective autophagy. *J Cell Biol* 217: 3354-3367

Liang JR, Lingeman E, Luong T, Ahmed S, Muhar M, Nguyen T, Olzmann JA, Corn JE (2020) A Genome-wide ER-phagy Screen Highlights Key Roles of Mitochondrial Metabolism and ER-Resident UFMylation. *Cell* 180: 1160-1177.e1120

Liu S, Obert C, Yu YP, Zhao J, Ren BG, Liu JJ, Wiseman K, Krajacich BJ, Wang W, Metcalfe K *et al* (2024) Utility analyses of AVITI sequencing chemistry. *BMC Genomics* 25: 778

Lu JY, Simon M, Zhao Y, Ablaeva J, Corson N, Choi Y, Yamada KYH, Schork NJ, Hood WR, Hill GE *et al* (2022) Comparative transcriptomics reveals circadian and pluripotency networks as two pillars of longevity regulation. *Cell Metabolism* 34: 836-856.e835

Malovannaya A, Lanz RB, Jung SY, Bulynko Y, Le NT, Chan DW, Ding C, Shi Y, Yucer N, Krenciute G *et al* (2011) Analysis of the human endogenous coregulator complexome. *Cell* 145: 787-799

Martinez G, Duran-Aniotz C, Cabral-Miranda F, Vivar JP, Hetz C (2017) Endoplasmic reticulum proteostasis impairment in aging. *Aging Cell* 16: 615-623

Masri S, Kinouchi K, Sassone-Corsi P (2015) Circadian clocks, epigenetics, and cancer. *Curr Opin Oncol* 27: 50-56

Meng H, Gonzales NM, Jung SY, Lu Y, Putluri N, Zhu B, Dacso CC, Lonard DM, O'Malley BW (2022) Defining the mammalian coactivation of hepatic 12-h clock and lipid metabolism. *Cell Reports* 38: 110491

Meng H, Gonzales NM, Lonard DM, Putluri N, Zhu B, Dacso CC, York B, O'Malley BW (2020) XBP1 links the 12-hour clock to NAFLD and regulation of membrane fluidity and lipid homeostasis. *Nature Communications* 11: 6215

Metcalf MG, Higuchi-Sanabria R, Garcia G, Tsui CK, Dillin A (2020) Beyond the cell factory: Homeostatic regulation of and by the UPR^{ER}. *Science Advances* 6: eabb9614

Mohan S, R PRM, Brown L, Ayyappan P, G RK (2019) Endoplasmic reticulum stress: A master regulator of metabolic syndrome. *European Journal of Pharmacology* 860: 172553

Nakahata Y, Kaluzova M, Grimaldi B, Sahar S, Hirayama J, Chen D, Guarente LP, Sassone-Corsi P (2008) The NAD$^+$-Dependent Deacetylase SIRT1 Modulates CLOCK-Mediated Chromatin Remodeling and Circadian Control. *Cell* 134: 329-340

Pakos-Zebrucka K, Koryga I, Mnich K, Ljujic M, Samali A, Gorman AM (2016) The integrated stress response. *EMBO Rep* 17: 1374-1395

Pan Y, Ballance H, Meng H, Gonzalez N, Kim S-M, Abdurehman L, York B, Chen X, Schnytzer Y, Levy O *et al* (2020) 12-h clock regulation of genetic information flow by XBP1s. *PLOS Biology* 18: e3000580

Pelletier S, Gingras S, Green DR (2015) Mouse genome engineering via CRISPR-Cas9 for study of immune function. *Immunity* 42: 18-27

Policarpi C, Munafò M, Tsagkris S, Carlini V, Hackett JA (2024) Systematic epigenome editing captures the context-dependent instructive function of chromatin modifications. *Nature Genetics* 56: 1168-1180

Qin Q, Fan J, Zheng R, Wan C, Mei S, Wu Q, Sun H, Brown M, Zhang J, Meyer CA *et al* (2020) Lisa: inferring transcriptional regulators through integrative modeling of public chromatin accessibility and ChIP-seq data. *Genome Biology* 21: 32

Qu Q, Takahashi YH, Yang Y, Hu H, Zhang Y, Brunzelle JS, Couture JF, Shilatifard A, Skiniotis G (2018) Structure and Conformational Dynamics of a COMPASS Histone H3K4 Methyltransferase Complex. *Cell* 174: 1117-1126.e1112

Quadros RM, Miura H, Harms DW, Akatsuka H, Sato T, Aida T, Redder R, Richardson GP, Inagaki Y, Sakai D *et al* (2017) Easi-CRISPR: a robust method for one-step generation of mice carrying conditional and insertion alleles using long ssDNA donors and CRISPR ribonucleoproteins. *Genome Biol* 18: 92

Ramirez F, Ryan DP, Gruning B, Bhardwaj V, Kilpert F, Richter AS, Heyne S, Dundar F, Manke T (2016) deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic Acids Res* 44: W160-165

Richter K, Haslbeck M, Buchner J (2010) The heat shock response: life on the verge of death. *Mol Cell* 40: 253-266

Roeder RG (2005) Transcriptional regulation and the role of diverse coactivators in animal cells. *FEBS Lett* 579: 909-915

Ron D, Walter P (2007) Signal integration in the endoplasmic reticulum unfolded protein response. *Nature Reviews Molecular Cell Biology* 8: 519-529

Ruthenburg AJ, Allis CD, Wysocka J (2007) Methylation of lysine 4 on histone H3: intricacy of
writing and reading a single epigenetic mark. *Mol Cell* 25: 15-30

Rutkowski DT, Wu J, Back SH, Callaghan MU, Ferris SP, Iqbal J, Clark R, Miao H, Hassler JR,
Fornek J *et al* (2008) UPR pathways combine to prevent hepatic steatosis caused by ER stress-
mediated suppression of transcriptional master regulators. *Dev Cell* 15: 829-840

Scott MR, Zong W, Ketchesin KD, Seney ML, Tseng GC, Zhu B, McClung CA (2023) Twelve-
hour rhythms in transcript expression within the human dorsolateral prefrontal cortex are altered
in schizophrenia. *PLOS Biology* 21: e3001688

Shilatifard A (2012a) The COMPASS family of histone H3K4 methylases: mechanisms of
regulation in development and disease pathogenesis. *Annu Rev Biochem* 81: 65-95

Shilatifard A (2012b) The COMPASS family of histone H3K4 methylases: mechanisms of
regulation in development and disease pathogenesis. *Annual review of biochemistry* 81: 65-95

So JS (2018) Roles of Endoplasmic Reticulum Stress in Immune Responses. *Mol Cells* 41: 705-
716

Spiegelman BM, Heinrich R (2004) Biological control through regulated transcriptional
coactivators. *Cell* 119: 157-167

Stashi E, Lanz RB, Mao J, Michailidis G, Zhu B, Kettner NM, Putluri N, Reineke EL, Reineke
LC, Dasgupta S (2014) SRC-2 is an essential coactivator for orchestrating metabolism and
circadian rhythm. *Cell reports* 6: 633-645

Takahashi JS (2017) Transcriptional architecture of the mammalian circadian clock. *Nat Rev*
Genet 18: 164-179

Takahashi YH, Westfield GH, Oleskie AN, Trievel RC, Shilatifard A, Skiniotis G (2011) Structural
analysis of the core COMPASS family of histone H3K4 methylases from yeast to human. *Proc*
Natl Acad Sci U S A 108: 20526-20531

Taylor RC, Hetz C (2020) Mastering organismal aging through the endoplasmic reticulum
proteostasis network. *Aging Cell* 19: e13265

Thaben PF, Westermark PO (2014) Detecting rhythms in time series with RAIN. *J Biol Rhythms*
29: 391-400

Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, Salzberg SL, Wold BJ,
Pachter L (2010) Transcript assembly and quantification by RNA-Seq reveals unannotated
transcripts and isoform switching during cell differentiation. *Nature Biotechnology* 28: 511-U174

van der Veen DR, Gerkema MP (2017) Unmasking ultradian rhythms in gene expression. *The*
FASEB Journal 31: 743-750

Walter P, Ron D (2011) The unfolded protein response: from stress pathway to homeostatic
regulation. *Science* 334: 1081-1086

Wang H, Fan Z, Shliaha PV, Miele M, Hendrickson RC, Jiang X, Helin K (2023) H3K4me3 regulates RNA polymerase II promoter-proximal pause-release. *Nature* 615: 339-348

Wiseman RL, Mesgarzadeh JS, Hendershot LM (2022) Reshaping endoplasmic reticulum quality control through the unfolded protein response. *Molecular Cell* 82: 1477-1491

Wulansari N, Darsono WHW, Woo HJ, Chang MY, Kim J, Bae EJ, Sun W, Lee JH, Cho IJ, Shin H *et al* (2021) Neurodevelopmental defects and neurodegenerative phenotypes in human brain organoids carrying Parkinson's disease-linked DNAJC6 mutations. *Sci Adv* 7

Xu J (2005) Preparation, culture, and immortalization of mouse embryonic fibroblasts. *Curr Protoc Mol Biol* Chapter 28: Unit 28.21

Yan Z, Yan H, Ou H (2012) Human thyroxine binding globulin (TBG) promoter directs efficient and sustaining transgene expression in liver-specific pattern. *Gene* 506: 289-294

Zhang EE, Liu AC, Hirota T, Miraglia LJ, Welch G, Pongsawakul PY, Liu X, Atwood A, Huss JW, 3rd, Janes J *et al* (2009) A genome-wide RNAi screen for modifiers of the circadian clock in human cells. *Cell* 139: 199-210

Zhao L, Ackerman SL (2006) Endoplasmic reticulum stress in health and disease. *Current Opinion in Cell Biology* 18: 444-452

Zhou M, Tamburini I, Van C, Molendijk J, Nguyen CM, Chang IY-Y, Johnson C, Velez LM, Cheon Y, Yeo R *et al* (2024) Leveraging inter-individual transcriptional correlation structure to infer discrete signaling mechanisms across metabolic tissues. *eLife* 12: RP88863

Zhu B (2020) Decoding the function and regulation of the mammalian 12h-clock. *Journal of Molecular Cell Biology*

Zhu B, Gates LA, Stashi E, Dasgupta S, Gonzales N, Dean A, Dacso CC, York B, O'Malley BW (2015a) Coactivator-Dependent Oscillation of Chromatin Accessibility Dictates Circadian Gene Amplitude via REV-ERB Loading. *Mol Cell* 60: 769-783

Zhu B, Gates LA, Stashi E, Dasgupta S, Gonzales N, Dean A, Dacso CC, York B, O'Malley BW (2015b) Coactivator-Dependent Oscillation of Chromatin Accessibility Dictates Circadian Gene Amplitude via REV-ERB Loading. *Mol Cell*

Zhu B, Gates LA, Stashi E, Dasgupta S, Gonzales N, Dean A, Dacso CC, York B, O'Malley BW (2015c) Coactivator-dependent oscillation of chromatin accessibility dictates circadian gene amplitude via REV-ERB loading. *Molecular cell* 60: 769-783

Zhu B, Liu S (2023a) Preservation of ~12-hour ultradian rhythms of gene expression of mRNA and protein metabolism in the absence of canonical circadian clock. *bioRxiv*: 2023.2005.2001.538977

Zhu B, Liu S (2023b) Preservation of ~12-h ultradian rhythms of gene expression of mRNA and protein metabolism in the absence of canonical circadian clock. *Frontiers in Physiology* 14

Zhu B, Liu S, David NL, Dion W, Doshi NK, Siegel LB, Amorim T, Andrews RE, Kumar GN, Irfan
S *et al* (2023) Evidence for conservation of a primordial 12-hour ultradian gene program in
humans. *bioRxiv*: 2023.2005.2002.539021

Zhu B, Liu S, David NL, Dion W, Doshi NK, Siegel LB, Amorim T, Andrews RE, Kumar GVN, Li
H *et al* (2024) Evidence for ~12-h ultradian gene programs in humans. *npj Biological Timing and*
Sleep 1: 4

Zhu B, Zhang Q, Pan Y, Mace EM, York B, Antoulas AC, Dacso CC, O'Malley BW (2017a) A
Cell-Autonomous Mammalian 12 hr Clock Coordinates Metabolic and Stress Rhythms. *Cell*
Metab 25: 1305-1319 e1309

Zhu B, Zhang Q, Pan Y, Mace EM, York B, Antoulas AC, Dacso CC, O'Malley BW (2017b) A
cell-autonomous mammalian 12 hr clock coordinates metabolic and stress rhythms. *Cell*
metabolism 25: 1305-1319. e1309

1169

1170

1171

1172

1173

1174 **Fig. legends**

1175

Fig. 1. Global ~12h RBBP5 cistrome is associated with promoter-proximal ~12h H3K4me3 epigenome in mouse liver. (A) Heatmap of ~12h hepatic transcriptome aligned with H3K4me3, H3K9Ac and H3K27Ac cistromes. **(B)** Quantification of temporal H3K4me3, H3K9ac and H3K27Ac. **(C)** Heatmap showing RBBP5 and H3K4me3 chromatin occupancy 1kb ± of TSS of 6,451 genes. **(D)** RBBP5 chromatin occupancy aligned with nucleosome positioning. **(E)** Quantification of temporal average RBBP5 binding intensity on different nucleosomes. **(F)** RBBP5 chromatin occupancy aligned with H3K4me3 occupancy.

1183

1184

Fig. 2. Hepatic RBBP5 cistrome coincides with that of XBP1s and hepatic ~12h transcriptome. (A) The table showing the number of genes with or without ~12h rhythms and with or without promoter RBBP5 binding. **(B)** Heatmap showing RBBP5 binding intensity along with -log$_{10}$ transformed P values for having 1~2h rhythm by RAIN and ~12h rhythms of gene expression on 2,525 genes. **(C)** Heat map summary of GO analysis demonstrating the -log$_{10}$ transformed P value of different enriched pathways for different genes. **(D)** A representative diagram depicting a typical Gro-Seq signal from TSS to transcription stop site (TTS) of a gene and using AUC to calculate the pausing index. **(E)** Log$_2$ mean-normalized temporal pausing index calculated from the GRO-Seq data for ~12h genes with or without promoter RBBP5 binding. **(F)** Venn diagram demonstrating distinct and common cistromes for RBBP5 and XBP1s. **(G)** Western blot showing co-IP in liver nuclear extracts from CT0 using anti-XBP1s and normal IgG control antibody. **(H)** Motif analysis of RBBP5 binding sites of 6,149 genes.

1200

Fig. 3. RBBP5 is an epigenetic regulator of the hepatic ~12h oscillator, but not the canonical ~24h circadian clock. (A) Schematic of the design of *Rbbp5^Flox* mice. **(B)**

Immunofluorescence of RBBP5 in the liver of *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice. **(C)** Histograms
showing the period distributions of all rhythmic genes uncovered from *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO}
mice (with adjP values or FDR <0.05) (n=2 at each time point for either genotype). **(D, E)**
Cumulative distribution of 10~12h **(D)** and 22~28h **(E)** genes in *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice
with different FDR cut-offs. **(F)** Heat map of 15 core circadian clock (top) and canonical ~12h gene
expression (bottom) in the liver of *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice. Log$_2$ normalized fold change of
average gene expression across 48 hours between *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice was also
shown. **(G)** Heat map of 6,025 ~12h gene expression (FDR<0.05) in the liver of *Rbbp5* ^{Flox} and
Rbbp5 ^{LKO} mice, with 5,601 of them lost in *Rbbp5* ^{LKO} mice, along with -log$_{10}$ transformed adjP
values for having 10-12h rhythm by RAIN. Log$_2$ normalized fold change of average gene
expression across 48 hours between *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice was also shown. **(H)** RNA-
Seq data for representative proteostasis and mRNA processing genes in the liver of *Rbbp5* ^{Flox}
and *Rbbp5* ^{LKO} mice. **(I)** Heat map summary of GO analysis demonstrating the -log$_{10}$ transformed
P values of different enriched pathways for different genes. Data: Mean ± S.E.M.
1217

1218

1219

1220

1221

1222

1223

1224

1225

1226

1227

1228

1229

1230

1231

1232

1233

1234

1235

1236

1237

1238

Fig. 4. RBBP5 regulates the hepatic transcriptional response to proteotoxic stress. (A) The
diagram of UPR. **(B)** PCA showing the hepatic transcriptional response to Tu in *Rbbp5 Flox* and
Rbbp5 LKO mice. **(C)** Representative RNA-seq data for proteostasis (top) and immune genes
(bottom) with RBBP5 promoter-binding status also shown below. n=3 for *Rbbp5 Flox* DMSO and
Rbbp5 Flox Tu, and n=4 for *Rbbp5 LKO* DMSO and *Rbbp5 LKO* Tu. **(D)** Heat map summary of GO

analysis demonstrating the -$\log_{10}$ transformed P values of different enriched pathways for different

genes. **(E)** Log $_2$ normalized fold induction by Tu in *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice for genes with

or without RBBP5 promoter binding. **(F)** Western blot analysis of different proteins in *Rbbp5* ^{Flox}

and *Rbbp5* ^{LKO} mice treated with or without Tu. **(G, H)** Mice tail-vein injected with AAV8-TGB-GFP

or AAV8-TGB-RBBP5 were treated with or without Tu. Western blot analysis **(G)** and qPCR **(H)**.

n=3 for AAV8-TBG-EGFP DMSO, n=4 for AAV8-TBG-EGFP Tu, n=4 for AAV8-TGB-mRBBP5

DMSO and n=4 for AAV8-TGB-mRBBP5 Tu. Data: Mean ± S.E.M.

1251

1252

Fig. 5. RBBP5 protects mice from chronic ER stress-induced hepatic inflammation and steatosis. (A) A schematic of experimental design. **(B)** Western blot and quantification of the expression of different proteins in *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice with or without repeated Tu injection. **(C)** qPCR analysis of different proteostasis and immune gene expressions in the same cohort of mice. **(D)** Representative images of Oil Red O staining and quantification of lipid droplet size, number and total lipid content (total lipid droplet area). n=4 for *Rbbp5* ^{Flox} DMSO, n=4 for *Rbbp5* ^{Flox} Tu, n=2 for *Rbbp5* ^{LKO} DMSO, n=3 for *Rbbp5* ^{LKO} Tu. Data: Mean ± S.E.M.

Fig. 6. RBBP5 is required for transcriptional responses to diverse proteotoxic stresses. (A) qPCR of different genes in MEFs transfected with scrambled, *Rbbp5* or *Xbp1* siRNAs and treated with DMSO or Tu (100ng/ml) for 8 hours. **(B)** Selected genes aligned for RBBP5 and H3K4me3 ChIP-seq signal from CT0 mouse liver and ChIP-qPCR of RBBP5 and H3K4me3 in MEFs treated with Tu for different hours. (**C, D**) qPCR of different genes in *Rbbp5* (fl/fl) ROSA26-CreERT2 (+/+) MEFs treated with vehicle (WT) or tamoxifen (*Rbbp5* KO) followed by treatment with vehicle control or 1mM DTT for 5 hours **(C)** or treatment with full DMEM media, leucine-free media or EBSS for 16 hours **(D)**. **(E)** Diagram of the mCherry-GFP-LC3 autophagy reporter. **(F)** Quantification of autophagic flux in mCherry-GFP-LC3-expressing *Rbbp5* (fl/fl) ROSA26-CreERT2 (+/+) MEFs treated with vehicle (WT) or tamoxifen (*Rbbp5* KO) followed by treatment with full DMEM media or leucine-free media for 16 hours. **(G)** Diagram of the ER-phagy reporter using the mCherry cleavage from ER assay. **(H, I)** Western blot of different proteins **(H)** and quantification **(I)** of mCherry cleavage in RAMP4-mCherry-expressing *Rbbp5* (fl/fl) ROSA26-CreERT2 (+/+) MEFs treated with vehicle (WT) or tamoxifen (*Rbbp5* KO) followed by treatment with full DMEM media or EBSS for 16 hours. Data: Mean ± S.E.M.

1276

Fig. 7. Reduced RBBP5 expression is associated with aging in mice and hepatic steatosis
in humans. (A) Z score-normalized expression of mouse *Rbbp5* and average expression of
stress response genes across mouse life span according to Tabua Muris dataset (Almanzar *et al.*,
2020). **(B-F)** Data in humans with NAFLD/MAFLD was from (Ahrens *et al.*, 2013). Expression of
RBBP5 (**B**) and the eigengene of 97 stress genes (**C**). (**D**) Scatter plot showing a positive
correlation between *RBBP5* and eigengene expression. **(E)** Heatmap of 97 stress genes
expression in different cohorts of human subjects. (**F**) GSEA analysis showing stress gene
expression are downregulated in steatosis subjects.

Supplemental Figures

1286

Fig. S1. Global ~12h RBBP5 cistrome is associated with promoter-proximal ~12h H3K4me3 epigenome in mouse liver. (A) COMPASS writes H3K4me3 using S-Adenosyl methionine (SAM) as the substrate. **(B)** LISA revealing putative TFs and co-regulators for hepatic top 500 most robust (the smallest p values by RAIN analysis) circadian (y-axis) and 12h (x-axis) genes. **(C)** Temporal hepatic *Rbbp5* expression assayed by RNA-seq. Period is calculated by the eigenvalue/pencil method. **(D)** Western blot and quantification (n=2) of temporal nuclear RBBP5 level in mouse liver at different CTs. Period is calculated by eigenvalue/pencil method. **(E)**

Heatmap showing RBBP5, H3K9Ac and H3K27Ac chromatin occupancy 1kb ± of TSS of 6,451
genes. **(F)** Snapshot of target genes selected for alignment of hepatic RBBP5, H3K4me3,
H3K9Ac and H3K27Ac at different CTs. **(G)** Heatmap of temporal RBBP5 binding intensity for
7,963 binding sites, along with $-\log_{10}$ transformed P value for having 12h rhythm by RAIN. **(H)**
Cumulative distribution of the number of ~12h and circadian RBBP5 binding sites ranked by p
value or false discovery rate (FDR). Data: Mean ± S.E.M.

1300

1301

1302

1303

1304

1305

1306

1307

1308

1309

1310

1311

1312

1313

1314

1315

1316

1317

1318

1319

1320

1321

1322

1323

1324

1325

1326

1. Wild-type C57BL/6J mouse liver
2. *Rbbp5* (fl/fl); Alb-CRE (-/-) mouse liver
3. *Rbbp5* (fl/fl); Alb-CRE (+/-) mouse heart
4. *Rbbp5* (fl/fl); Alb-CRE (+/-) mouse kidney
5. *Rbbp5* (fl/fl); Alb-CRE (+/-) mouse lung
6. *Rbbp5* (fl/fl); Alb-CRE (+/-) mouse liver

1327

1328 **Fig. S2. The generation of RBBP5 ^{LKO} mice. (A, B)** Expected **(A)** and actual genotyping result
1329 **(B)** for different mice. **(C, D)** Western blot **(C)** and quantification **(D)** of RBBP5 in different tissues
1330 in *Rbbp5* (fl/fl); Alb-CRE (+/-) and *Rbbp5* (fl/fl); Alb-CRE (-/-) mice. n=2 per genotype. Data: Mean
1331 ± S.E.M.

1332

1333

1334

1335

1336

1337

1338

1339

1340

1341

1342

1343

1344

1345

1346

1347

Fig. S3. Liver-specific deletion of RBBP5 does not alter rhythmic locomotor activity nor
fasting-feeding cycles in mice. (A) Average body weight of male *Rbbp5* *Flox* and *Rbbp5* *LKO* mice
at different ages. Mean ± 95% confidence interval. N=3~71 for *Rbbp5* *Flox* and n=6~54 for *Rbbp5*
LKO mice at each week. P<0.0001 by One-way ANOVA. **(B)** Real-time locomotor activity monitoring
in *Rbbp5* *Flox* and *Rbbp5* *LKO* mice under 12hr light/12hr dark conditions. **(C)** Averaged
measurements within the light and dark phase as described in **B. (D)** Real-time measurement of
food intake in *Rbbp5* *Flox* and *Rbbp5* *LKO* mice under 12hr light/12hr dark conditions. **(E)** Averaged
measurements within the light and dark phase as described in **D.** n=6 for *Rbbp5* *Flox* and n=8 for
Rbbp5 *LKO* mice. **(F)** Real-time locomotor activity monitoring in *Rbbp5* *Flox* and *Rbbp5* *LKO* mice
under constant dark conditions. **(G)** Averaged measurements within the rest and active phase as
described in **F**. **(H)** Real-time measurement of food intake in *Rbbp5* *Flox* and *Rbbp5* *LKO* mice under
constant dark conditions. **(I)** Averaged measurements within the rest and active phase as
described in **H.** n=5 for *Rbbp5* *Flox* and n=3 for *Rbbp5* *LKO* mice. Data: Mean ± S.E.M.

Fig. S4. RBBP5 is an epigenetic regulator of the hepatic ~12h oscillator, but not the canonical ~24h circadian clock. (A) Heat map of 8,340 circadian gene expression (FDR<0.05) in the liver of *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice, with 5,043 of them shared between the two, along with -log$_{10}$ transformed adjP values for having 22-28h rhythm by RAIN. Log$_2$ normalized fold change of average gene expression across 48 hours between *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice was also shown. **(B, C)** PCA of hepatic temporal transcriptome in *Rbbp5* ^{Flox} **(B)** and *Rbbp5* ^{LKO} **(C)** mice. **(D)** A table listing the number of ~12h and ~24h circadian genes in *Rbbp5* ^{Flox} and *Rbbp5* ^{LKO} mice with different statistical thresholds.

1370

Fig. S5. Eigenvalue/pencil method analysis of RBBP5-dependent hepatic ~12h transcriptome. (A, B) Distributions of all **(A)** and dominant **(B)** oscillations uncovered by the eigenvalue/pencil method in *Rbbp5 ^{Flox}* and *Rbbp5 ^{LKO}* mice. **(C)** Venn diagram comparing unique and common ~12h transcriptomes uncovered by the eigenvalue/pencil and RAIN methods with two different FDR cut-offs. **(D)** Heat map of all ~12h gene expression (or lack thereof) in *Rbbp5 ^{Flox}* and *Rbbp5 ^{LKO}* mice with both raw data and superimposed ~12h rhythms shown. 4,522 ~12h genes were abolished, 238 ~12h genes dampened, and 392 ~12h genes were enhanced in *Rbbp5 ^{LKO}* mice, respectively. Log$_2$ normalized fold change of average gene expression across

48 hours between *Rbbp5 Flox* and *Rbbp5 LKO* mice was also shown. **(E)** Heat map of dominant
~12h gene expression (or lack thereof) in *Rbbp5 Flox* and *Rbbp5 LKO* mice with both raw data and
superimposed ~12h rhythms shown. 2,450 ~12h genes were abolished, 70 ~12h genes
dampened, and 90 ~12h genes were enhanced in *Rbbp5 LKO* mice, respectively. Log_2 normalized
fold change of average gene expression across 48 hours between *Rbbp5 Flox* and *Rbbp5 LKO* mice
was also shown. **(F)** The periods distribution in *Rbbp5 LKO* mice of those ~12h genes originally
identified in *Rbbp5 Flox* mice but lost in *Rbbp5 LKO* mice. **(G, H)** Heat map summary of GO analysis
demonstrating the $-log_{10}$ transformed P values of different enriched pathways for all **(G)** and
dominant **(H)** ~12h genes.

1388

1389

1390

1391

1392

1393

1394

1395

1396

1397

1398

1399

1400

1401

1402

1403

1404

1405

1406

1407

1408

1409

1410

1411

1412

Fig. S6. RBBP5 is a cell-autonomous epigenetic regulator of the ~12h oscillator, but not the canonical ~24h circadian clock. (A) qPCR analysis of *Rbbp5* expression in Tu (25ng/ml)-synchronized MEFs with scrambled, *Bmal1* and *Xbp1* siRNAs. P values for having 12h rhythms were calculated by RAIN. **(B)** Real-time luminescence of MEFs expressing *Manf* promoter-driven dluc (Zhu *et al.*, 2017a) transfected with different siRNAs and quantified periods. **(C)** Real-time luminescence traces of Bmal1-dluc U2OS cells transfected with different siRNAs as reported in (Zhang *et al.*, 2009). Data: Mean ± S.E.M.

1420

Fig. S7. RBBP5 regulates the hepatic transcriptional response to proteotoxic stress. (A-D)
Volcano plot illustrating the log 2 normalized fold change vs -log2 transformed adjusted p values
for different comparisons. **(E)** Scatter plot comparing the log 2 transformed fold change of gene
expression by Tu in *Rbbp5* *Flox* (x- axis) and *Rbbp5* *LKO* (y-axis) mice. **(F)** Scatter plot comparing
the -log 10 transformed adjusted p values for gene expression induced by Tu in *Rbbp5* *Flox* (x- axis)
and *Rbbp5* *LKO* (y-axis) mice. **(G)** Venn diagram comparing distinct and common genes induced
by Tu in *Rbbp5* *Flox* and *Rbbp5* *LKO* mice with FDR<0.1. **(H)** Heat map summary of GO analysis
demonstrating the -log$_{10}$ transformed P values of different enriched pathways for different groups
of genes.

Fig. S8. RBBP5 is required for transcriptional responses to diverse proteotoxic stresses.
(A) Western blot of RBBP5 in MEFs transfected with *scrambled* or *Rbbp5* siRNA for 48 hours. **(B)** qPCR of different genes in MEFs transfected with scrambled, *Ash2l* or *Setd1a* siRNAs and treated with DMSO or Tu (100ng/ml) for 8 hours. **(C)** qPCR of different genes in MEFs transfected with scrambled, or the combination *Gcn5* and *Pcaf* siRNAs and treated with DMSO or Tu (100ng/ml) for 8 hours. **(D-F)** Isolation and immortalization of *Rbbp5* (fl/fl) ROSA26-CreERT2 (+/+) MEFs **(D)**, which were treated with vehicle (WT) or tamoxifen (*Rbbp5* KO) for 5 or 7 days and genotyping **(E)** and western blot analysis **(F)** to confirm the depletion of RBBP5. **(G)** Representative images of autophagic flux in mCherry-GFP-LC3-expressing *Rbbp5* (fl/fl) ROSA26-CreERT2 (+/+) MEFs treated with vehicle (WT) or tamoxifen (*Rbbp5* KO) followed by treatment with full DMEM media or leucine-free media for 16 hours. **(H, I)** Western blot of different proteins **(H)** and quantification **(I)** of p62 in *Rbbp5* (fl/fl) ROSA26-CreERT2 (+/+) MEFs treated with vehicle (WT) or tamoxifen (*Rbbp5* KO) followed by treatment with full DMEM media or leucine free media for 16 hours**. (J)** Percentage of cells normalized to before treatment for mCherry-GFP-LC3-expressing *Rbbp5* (fl/fl) ROSA26-CreERT2 (+/+) MEFs treated with vehicle (WT) or tamoxifen (*Rbbp5* KO) followed by treatment with full DMEM media, leucine-free media or EBSS for up to 16 hours. Data: Mean ± S.E.M.

1460

1461 **Fig. S9. RBBP5 expression is positively correlated with those involved in proteostasis and**
1462 **mRNA metabolism and negatively associated with those implicated in immune response**
1463 **in human liver.** Data are from the human GD-CAT dataset (Zhou *et al.*, 2024).

1464

1465

1466

1467

1468

1469

1470

1471

1472

1473

 Supplemental Tables

Table S1. FPKM quantification of temporal hepatic RBBP5 ChIP-seq in wild-type mouse
liver.

Table S2. FPKM quantification of temporal hepatic RNA-seq in *Rbbp5* ^{*Flox*} and *Rbbp5* ^{*LKO*}
mice.

Table S3. RAIN analysis of 10~12h hepatic transcriptome identified in *Rbbp5* ^{*Flox*} and *Rbbp5*
^{*LKO*} **mice.**

Tab 1: 10~12h oscillations identified in *Rbbp5* ^{*Flox*} mice.

Tab 2: 10~12h oscillations identified in *Rbbp5* ^{*LKO*} mice.

Table S4. Eigenvalue/pencil analysis of oscillating hepatic transcriptome identified in
***Rbbp5* ^{*Flox*} and *Rbbp5* ^{*LKO*} mice.**

Tab 1: All oscillations identified in *Rbbp5* ^{*Flox*} mice.

Tab 2: All dominant oscillations identified in *Rbbp5* ^{*Flox*} mice.

Tab 3: All 10~13h oscillations identified in *Rbbp5* ^{*Flox*} mice.

Tab 4: All dominant 10~13h oscillations identified in *Rbbp5* ^{*Flox*} mice.

Tab 5: All oscillations identified in *Rbbp5* ^{*LKO*} mice.

Tab 6: All dominant oscillations identified in *Rbbp5* ^{*LKO*} mice.

Tab 7: All 10~13h oscillations identified in *Rbbp5* ^{*LKO*} mice.

Tab 8: All dominant 10~13h oscillations identified in *Rbbp5* ^{*LKO*} mice.

Table S5. RNA-seq quantification of hepatic transcriptome in *Rbbp5* ^{*Flox*} and *Rbbp5* ^{*LKO*} mice
with or without Tu injection.

Tab 1: Raw counts

Tab 2: CPM quantification

Tab 3: FKPM quantification

Table S6. Differentially expressed gene (DEG) analysis of hepatic transcriptome in *Rbbp5*
^{*Flox*} **and *Rbbp5* ^{*LKO*} mice with or without Tu injection.**

1500