MANUSCRIPT TITLE

Diversification and conservation of DNA binding specificities of SPL family of

transcription factors

AUTHORS

Miaomiao Li[1,2,3]*, Tao Yao[2,3], Mary Galli[4], Luigi F. Di Costanzo[5], Wanru Lin[1], Yilin Zhou[1], Jin-Gui

Chen[2,3], Andrea Gallavotti[4] and Shao-shan Carol Huang[1],*

[1] Center for Genomics and Systems Biology, Department of Biology, New York University, New

York, NY 10003, USA

[2] Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA

[3] Center for Genomics and Systems Biology, Department of Biology, New York University, New
York, NY 10003, USA

[4] Waksman Institute of Microbiology, Rutgers University, Piscataway, NJ 08854-8020, USA.

[5] Department of Agricultural Sciences, University of Naples Federico II, Via Università 100,

80055 Portici, NA, Italy

*Correspondence: s.c.huang@nyu.edu, lim7@ornl.gov

GRAPHICAL ABSTRACT

ABSTRACT

SQUAMOSA Promoter-Binding Protein-Like (SPL) transcription factors are key regulators of

plant development and stress responses. Here, we present a comprehensive DNA affinity

purification sequencing (DAP-seq) analysis of 14 of the 16 SPL members in *Arabidopsis*

thaliana, revealing two major classes based on distinct DNA-binding motifs. Gene Ontology

enrichment of target genes indicated both shared and specialized functions among AtSPLs,

including roles in hormone signaling, metal ion homeostasis, and developmental transitions.

Comparative analysis of closely related paralogs, AtSPL9 and AtSPL15, uncovered divergence

in their genomic binding locations and target gene regulation, particularly in auxin and abscisic

acid pathways. These differences were supported by motif analysis and protoplast reporter

assays. Extending our study across species, we analyzed DAP-seq data from maize (*Zea*

mays) and wheat (*Triticum aestivum)* SPL homologs, identifying conserved target genes but

species-specific DNA motifs. Notably, wheat SPLs preferentially bind to longer, palindromic

motifs distinct from the GTAC core motif in *Arabidopsis*. Using AlphaFold3 structure modeling,

we show that these motif differences may arise from enhanced protein-protein interactions that

stabilize dimeric binding. Our results highlight how SPL family members diversify through DNA-

binding evolution and suggest that transcription factor dimerization contributes to motif

complexity and regulatory specialization across plant genomes.

INTRODUCTION

Transcription factors (TFs) are crucial regulators of gene expression, playing a central role in

controlling various biological processes by binding to specific DNA sequences and modulating

the transcription of target genes. The expansion of TF families through gene duplication events

and subsequent functional diversification are a key evolutionary mechanism that enhances the

complexity of gene regulation in plants (1). This expansion allows for the development of novel

regulatory networks and the fine-tuning of gene expression in response to environmental and

developmental cues. This process, known as functional diversification, can result in the

emergence of TFs with altered DNA-binding specificities, thereby contributing to the evolution of

complex regulatory circuits (1–3). A comprehensive analysis of DNA-binding specificities across

an entire TF family is a powerful way to understand the full range of their regulatory potential.

This can reveal how different TFs within a family contribute to the regulation of shared and

distinct biological pathways, offering insights into the mechanisms underlying the evolution of

gene regulatory networks.

As a plant-specific TF family, *SQUAMOSA PROMOTER BINDING PROTEIN-LIKEs* (SPLs)

have highly conserved SQUAMOSA Promoter-Binding Protein (SBP) domains, which are

approximately 76 amino acids long and contain two zinc finger-like structures and a nuclear

localization signal motif (4). The first two SBP proteins discovered, *AmSBP1* and *AmSBP2*,

were identified in *Antirrhinum majus* floral meristem and found to act during early flower

development (5). In *Arabidopsis*, the SPL genes are central to the age-dependent pathway of

flowering (6–8). Many SPL members in *Arabidopsis* are targeted by miR156/157, and this

miR156/SPL module plays an important role in diverse development stages including shoot

meristem growth, flower development and the phase change from vegetative growth to

reproductive growth, as well as the response to abiotic environmental stress (9–12). SPLs in

Arabidopsis directly activate the expression of critical flowering genes, such as *SUPPRESSOR*

OF OVEREXPRESSION OF CONSTANS 1 (SOC1), *FLOWERING LOCUS T (FT)*, *FRUITFULL*

(FUL), and *APETALA1 (AP1)* (6, 13, 14), as well as auxin related genes. They also repress the

expression of *AUXIN RESPONSE FACTOR (ARF)* genes by binding to their promoters such as

those of *ARF6* and *ARF8*, thereby balancing auxin signaling (15).

In *Arabidopsis*, *AtSPL9* is one of the most extensively studied members of the SPL family (6, 7,

16). *rSPL9* is a miR156-insensitive mutant, showing reduced juvenile period and less response

to aging (6, 17, 18). Within the *Arabidopsis* SPL family, *AtSPL9* and *AtSPL15* are the closest

paralogs resulting from genome duplication and have redundant biological functions (19, 20).

The *spl9/spl15* double mutants exhibit alternations in branching and flowering patterns,

highlighting the critical role of AtSPL9 and AtSPL15 in regulating shoot branching (21).

However, AtSPL15 and AtSPL9 coordinate distinct flowering phenotypes under short-day

conditions (22). SPL homologs in crops play critical roles in controlling plant architecture,

flowering time, and reproductive development. In rice (*Oryza sativa*), one of the most studied

SPL genes, *OsSPL14*, also known as *Ideal Plant Architecture1 (IPA1)* and closely related to

AtSPL9 and *AtSPL15*, is a key regulator of tiller number and panicle size (8, 23). In maize (*Zea*

mays), *ZmSBP8* and *ZmSBP30* are homologous to *AtSPL9/15*. The *zmsbp8/30* mutants exhibit

notable changes in kernel row number, ear size, and cob architecture, ultimately affecting grain

yield (24, 25). In hexaploid bread wheat (*Triticum aestivum*), the *AtSPL9/15* homologs across

the three subgenomes are *TaSPL7A/B/D* and *TaSPL13A/B/D.* CRISPR/Cas9 knockout of

TaSPL7A/B/D and *TaSPL15A/B/D* resulted in increased tiller number, decreased spike length,

and reduced spikelet number (23, 26). These proteins are thus essential regulators of

agronomic traits that directly impact crop yield, so identifying and editing the targets of the TFs

could provide powerful tools for crop improvement.

Despite extensive research on the SPL family, key questions remain about the differences in

DNA binding specificities among paralogs and the functional similarities and differences of these

genes across plant species, especially in crops. In this study, we used DNA affinity purification-

sequencing (DAP-seq) (27, 28), to map the genome-wide binding sites of the entire SPL family

in *Arabidopsis* and systematically analyze their DNA binding specificities. Additionally, we

utilized additional DAP-seq datasets for homologous SPLs in maize and wheat to examine the

conservation of potential target genes SPLs across the three species, uncovering changes of

their DNA binding specificities that could be mediated by a dimerization mechanism. By

focusing on the expansion of the SBP/SPL TF family, our study provides insights into how TF

duplication and subsequent functional diversification, mediated by evolution of DNA binding

specificities, have contributed to the complexity of gene regulation in plants.

MATERIALS AND METHODS

Plant materials

Arabidopsis reference accession Col-0 (CS70000) was grown in soil at 21 °C under long-day

(16 h light and 8 h dark) conditions for four weeks. Rosette leaves were collected and flash

frozen in liquid nitrogen prior to genomic DNA (gDNA) isolation for DAP-seq DNA library

preparation.

DAP-seq experiments

The *Arabidopsis* SPL DAP and ampDAP-seq experiments were performed according to

published protocols of (27, 29). First, frozen leaf tissues were ground into a fine powder using a

mortar and pestle, dissolved in extraction buffer (0.1 M TRIS base, 0.5 M sodium chloride, 0.05

M EDTA, pH 8.0), and subject to phenol-chloroform extraction to remove proteins and other

contaminants, resulting in purified gDNA. DAP-seq gDNA library was then prepared as a

standard DNA sequencing library for the Illumina platform. gDNA was fragmented to an average

of 200 bp using Covaris S220 Sonicator. Fragmented gDNA was end-repaired using the End-It

DNA End-Repair Kit (Lucigen, ER81050, ordered in 2021) and incubated at room temperature

for 45 min, followed by an A-tailing reaction using Klenow (3′→5′ exo-) (NEB, M0212) incubated

at 37 °C for 30 min. A-tailed DNA fragments and annealed adapters were ligated in a ligation

reaction using T4 DNA Ligase (Promega, M1804) incubated at room temperature for 3 h. The

DNA was precipitated and suspended in elution buffer (10 mM Tris-Cl, pH 8.5), resulting in the

gDNA libraries used in DAP-seq. A SPL protein expression reaction was prepared using TNT

SP6 Coupled Reticulocyte Lysate System (Promega, L4600) incubated with 1000 ng pIX-HALO-

SPL plasmids for 3 h at 30 °C. 10 µl of Magne HaloTag Beads (Promega, G7282) and 50 µl of

wash buffer (PBS with 0.05% NP40) were added to the reaction, bringing the final volume to

100 µl, and the mixture was incubated on a rotator for 1 hour at room temperature. The proteins

immobilized on beads were then washed five times on a magnet with 100 µl wash buffer to

purify HaloTag-fused SPL protein. After washing, the protein-bound beads were incubated with

100 ng adapter-ligated gDNA library in 100 µl wash buffer for 2 h on a rotator. The mixture was

then washed five times with wash buffer to remove unbound ligated DNA fragments. The beads

were suspended in 30 µl elution buffer, heated at 98 °C for 10 min, and put on ice immediately

for more than 5 min to denature the protein and release the bound DNA fragments. 25 µl of the

supernatant was used for the PCR enrichment step. The ampDAP library is generated by

amplifying DNA from the gDNA library by PCR, followed by incubation with the purified TF

protein as for the DAP-seq protocol. The PCR reactions for making ampDAP-seq libraries and

sequencing libraries following protein binding were prepared as 1 µl of Phusion DNA

Polymerase (New England Biolabs, M0530), 10 µl of 5x Phusion HF Buffer, 2.5 µl of 10 mM

dNTPs, 1 µl each 25 µM index primers, 25 µl of eluted DNA, add water to 50 µl. The PCR

condition is as follows: 98 °C for 2 min, 15-19 cycles of 98 °C for 15 s, 60 °C for 30 s, and 72 °C

for 1 min, final extension at 72 °C for 10 min. PCR-amplified samples were run in 1% agarose

gel and cut to purify fragments from 200 bp to 600 bp with Zymoclean Gel DNA Recovery Kit

(Zymo Research, D4007). The purified DNA fragments were measured by Qubit HS dsDNA

Assay Kit (ThermoFisher, Q32854) and sequenced on the Illumina platform.

Reporter assay in *Arabidopsis* protoplasts

To generate the effector plasmids used in the reporter assay, *AtSPL9* and *AtSPL15* CDS

fragments were cloned in *pUC19-35S-DC* using LR reactions. A 758 bp sequence from the

AtDI21 promoter (Supplementary Table S1) was synthesized by Twist BioScience and

subcloned into *pUC19-DC-GUS* as reporter plasmid. The combinations of effector plasmid,

reporter plasmid, and reference plasmid (*pUC19-35S-LUC*) were transformed into *Arabidopsis*

leaf protoplasts. Middle sections of four-week-old fully expanded leaves were cut out, sliced into

strips and immersed into enzyme solution containing 0.4 M mannitol, 20 mM KCl, 20 mM MES,

10 mM $CaCl_2$, 5 mM β-mercaptoethanol, 0.1% BSA, 0.4% macerozyme R10, and 1.5% cellulase

R10. The mixture was incubated at room temperature for 2 h before filtering the protoplasts

through a 75 μm nylon mesh and washing them with W5 solution (154 mM NaCl, 125 mM $CaCl_2$,

5 mM KCl, and 2 mM MES). After centrifuging at 1000 rpm for 3 min at 4 °C, the protoplasts

were resuspended with MMg solution (0.8 M mannitol, 1 M $MgCl_2$ and 0.2 M MES) to obtain a

concentration of 200,000 cells per ml. Next, 6 μg effector, 3 μg reporter and 100 ng reference

plasmids were co-transfected into 100 μl of protoplasts using the PEG-calcium mediated

transfection method, followed by incubation in darkness for 18 to 20 h at room temperature. The

GUS activity assay was conducted (30) and measured using a Fluoroskan microplate reader

(30). The MUG (4-Methylumbelliferyl β-d-glucuronide) (Sigma-Aldrich, M9130) and luciferase

assay system (Promega, E1500) were used to perform GUS and LUC activity assays,

respectively. Relative GUS activity was calculated via normalization to LUC activity, and the

data are presented as three independent biological replicates.

DAP-seq data processing and analysis

Read processing, peak calling and motif discovery

The *Arabidopsis* DAP-seq libraries were sequenced on an Illumina NextSeq 500 instrument.

The reads were basecalled using Picard IlluminaBasecallsToFastq (31) version 2.23.8, with

APPLY_EAMSS_FILTER set to false. Following basecalling, the reads were demultiplexed

using Pheniqs (32) version 2.1.0. The entire process was executed using a custom nextflow

pipeline, GENEFLOW (33). Adapter sequences were trimmed from the single read FASTQ files

by Trim Galore (version 0.6.6) and Cutadapt version 3.1 (34) with quality cutoff of 20. The

trimmed reads were mapped to the *Arabidopsis* reference genome sequence TAIR10 using

Bowtie2 (35)version 2.2.9 with default parameters. Aligned reads were filtered by mapping

quality (MAPQ) score of at least 30 using SAMtools (36) version 1.11. Peak calling was done by

the GEM peak caller (37) version 3.3 on the filtered mapped reads with the default read

distribution, TAIR10 nuclear chromosome sequences, q-value threshold of 0.01 (--q 2), and

parameters "--f SAM --t 1 --min 200 --k_min 5 --k_max 14 --k_seqs 2000 --k_neg_dinu_shuffle --

outNP --outBED --outMEME --outJASPAR --outHOMER --print_bound_seqs --

print_aligned_seqs". Peaks were called for each replicate individually or by merging the two

replicates using GEM's multi-replicate mode, with samples from experiments of empty vector

pIXHALO as control. To create a blacklist of regions that contained highly enriched but artifact

signals, peak calling was done for a set of six pIXHALO empty vector controls using MACS3

(38), which could find broader regions of read enrichment compared to the point-source binding

events reported by GEM. The peaks called by MACS3 (version 3.0.0a5) for these control

samples with parameters "--keep-dup auto --nomodel --extsize 200 -q 0.1" were merged and the

peak regions shared by at least two of these control samples were designated to be blacklist

regions. DAP-seq peaks overlapping these regions were removed prior to all downstream

analysis. BigWig files of normalized read signals were created using the MAPQ 30 filtered

alignment BAM files by the bamCoverage program in the deepTools package (39) version 3.5.0

with the following parameters: "--binSize 1 --normalizeUsing RPKM --ignoreForNormalization Mt

Pt". The read normalized bigwig files were visualized in JBrowse 2 (40). For motif discovery, the

top 1000 binding events for each TF were obtained by sorting the GEM output narrowPeak files

first by increasing q-value then by decreasing signal value. Sequences of 200 bp centered at

the peak summits were extracted from the TAIR10 genome sequence and used as input for

MEME motif discovery (41) with a Markov background model of order 2 computed from the

peak sequences and parameters "-mod zoops -nmotifs 5 -minw 5 -maxw 15 -dna -searchsize 0

-revcomp -csites 1000". The resulting PWM motifs were imported into R by the universalmotif

package (42), aligned by the DiffLogo package (43), and plotted by the ggseqlogo package (44).

The maize DAP-seq peak dataset, with blacklist regions already removed, was obtained from

Zenodo under DOI 10.5281/zenodo.14991916 (45). The wheat DAP-seq data were obtained

from NCBI GEO accession GSE188699 (26). Adapter sequences were trimmed from the pair-

end FASTQ files by Trim Galore (version 0.6.6) and Cutadapt version 3.1 (34). The trimmed

pair-end reads were mapped to the *T. aestivum cv. Chinese Spring v2.1* genome sequence

assembly (46) using Bowtie2 (35) version 2.2.9 with default parameters. Aligned reads were

filtered by MAPQ score of at least 20 using SAMtools (36) version 1.11. Peak calling was done

by the MACS3 peak caller (38) version 3.0.0a5 with input library as the control and parameters

"-f BAMPE -g 11453938932.0 --keep-dup auto --nomodel --call-summits -q 0.05". Blacklist

regions of artifactual read enrichment were created by calling peaks on two replicates of the

input libraries individually, merging the peaks and computing the regions called in both

replicates of the input. DAP-seq peaks overlapping with blacklist regions were removed prior to
all downstream analysis.

Arabidopsis DAP-seq sample clustering

MANorm2_utils (version 1.0.0) (47) were first used to find the number of DAP-seq reads of all
Arabidopsis AtSPL DAP-seq samples contained in the GEM peak regions that occurred in at
least one replicate. MAnorm2 R package (47) version 1.2.2 was used on this count matrix to
perform normalization by a hierarchical normalization approach: the replicates for each TF were
first normalized, then all the TFs were normalized to a pseudo-reference, a mean-variance
curve was then fitted using parametric regression and occupied regions only. Peak regions
ranked among the top 2,000 intervals by normalized signal in more than one sample were used
to compute pairwise distances between TFs using the distBioCond function. The pairwise
distance matrix was then used in hierarchical clustering of TFs with complete linkage method.

Peak annotation and Gene Ontology Enrichment Analysis

To associate *Arabidopsis* DAP-seq peaks to genes, the top 3000 merged replicate GEM peaks
for each TF were annotated with a gene that had the closest transcription start site (TSS) using
the annotatePeak function in the ChIPseeker package (48) with the protein coding genes in
Araport11 (49). Genes that had a DAP-seq peak within -1000 bp upstream and 500 bp
downstream from the TSS were designated potential target genes. The R package
clusterProfiler (50) was used to identify the top 10 most enriched GO terms for the target genes
of each TF in the Biological Process ontology annotated in the org.At.tair.db database, and the
enrichment P-values of these GO terms for all the TFs were computed. The enrichment P-
values were corrected by the Benjamini & Hochberg method and a matrix of −log10 adjusted P-
values was computed with the enriched GO terms on the rows and DAP-seq TFs on the
columns and plotted as a heatmap by the ComplexHeatmap package (51). Clustering
dendrogram was obtained by hierarchical clustering of the −log10 adjusted P-value matrix with
Euclidean distance between rows and columns as distances and the complete linkage method.

***spl7* RNA-seq data processing and Gene Set Enrichment Analysis**

RNA-seq gene counts matrix for *spl7* mutant and Col-0 wild-type roots in low copper treatment
and control conditions were downloaded from NCBI GEO accession GSE104916 (52). Using
the R package DESeq2 (53), the count matrix was imported into R, pre-filtered by keeping only
genes that had read counts more than 10 in at least 3 replicates, normalized and analyzed for
differential expression with the design formula genotype+treatment+genotype:treatment.
Differentially expressed genes between *spl7* and Col-0 wild-type (WT) in control condition were

extracted from the contrast for the genotype factor between *spl7* and WT, with adjusted P-value

threshold of 0.1. The genes were then ranked by the log2 fold change between WT and *spl7*

and compared to the DAP-seq predicted target genes by the Gene Set Enrichment Analysis test

(GSEA) function in clusterProfiler (50).

Analysis of differential binding between *Arabidopsis* AtSPL9 and AtSPL15

Differential binding analysis between *Arabidopsis* AtSPL9 and AtSPL15 DAP-seq samples was

performed by the R package DiffBind (54) version 3.6.5. The samples were normalized by the

native DESeq2 normalization method RLE using reads in peaks, and differentially bound peaks

were calculated by DESeq2 with the AtSPL9 as the reference and q-value threshold of 0.05.

The significantly differentially bound peaks that have had a positive fold change were

designated as AtSPL15-preferred and those that had a negative fold change were designated

as AtSPL9-preferred. Sequences were extracted from the TAIR10 reference genome for the top

1000 AtSPL15-preferred or AtSPL9-preferred peaks sorted by adjusted P-values for motif

discovery using MEME-ChIP (55) version 5.3.0 with the parameters "-meme-mod anr -meme-

searchsize 0 -minw 5". Genes were associated with the top 3000 AtSPL15-preferred or AtSPL9-

preferred peaks and GO enrichment was performed as described above.

Cross-species comparison of AtSPL9/15 and homologs in maize and wheat

DAP-seq peaks for maize and wheat were annotated using the R package ChIPseeker (48),

with the Zm00001eb.1 genome annotation for maize (56) and the IWGSC RefSeq v2.1

annotation for wheat (46). Genes with DAP-seq peaks located within 10,000 bp upstream and

500 bp downstream of the TSS were designated as potential target genes. Homologs between

maize and *Arabidopsis* and between wheat and *Arabidopsis* were taken from the

Best.hit.arabi.name column in the PhytozomeV13 annotation_info file for maize and wheat,

respectively (57). Conserved target genes between maize and *Arabidopsis* were defined as

genes associated with peaks shared between ZmSBP8 and ZmSBP30 that are homologous to

genes associated with peaks shared between AtSPL9 and AtSPL15. For wheat and

Arabidopsis, conserved targets were defined as genes associated with peaks shared by at least

two of TaSPL7A/B/D and at least two of TaSPL13A/B/D that were homologous to those genes

associated with peaks shared between AtSPL9 and AtSPL15. To identify DNA-binding motifs,

sequences within the DAP-seq peaks associated with conserved target genes were extracted

from their respective genomes. Motif discovery was then performed using MEME-ChIP with the

parameters "-ccut 0 -meme-mod anr -minw 4 -meme-nmotifs 5".

Structure modeling of SPL-DNA interaction

To investigate the structural basis of differential DNA motif binding among SPL homologs from

Arabidopsis, maize and wheat, we used AlphaFold3 (AF3) (58) to predict structures of the SPL

TFs interacting with DNA sequences containing the different motif types. Our goal was to

determine whether PPIs could explain why SPL homologs in different species prefer distinct

DNA motifs, i.e., GTAC for *Arabidopsis* SPLs and GTASTAC for wheat SPLs, by evaluating

differences in TF-TF and TF-DNA interaction potential. To this end, we modeled both one-copy

and two-copy TF-DNA complexes for each protein in combination with either its species-

preferred binding motif or the motif preferred by an orthologous SPL in a contrasting species.

For protein inputs, we used full-length amino acid sequences corresponding to the isoforms

analyzed in the DAP-seq experiments (26, 45). For DNA inputs, we focused on DAP-seq peaks

associated with conserved SPL target genes across species. Within these peaks, we scanned

for motif matches corresponding to the preferred motif of the SPL in question, for example,

GTAC in *Arabidopsis* SPL peaks and GTASTAC in wheat SPL peaks. We selected the ten

highest-scoring motif matches for each protein as representative binding sites. For each

selected site, we extracted a 20 bp DNA sequence centered on the motif match, yielding a set

of DNA fragments we refer to as "species-preferred motif sequences". To create a comparison

set, we then searched genomic regions nearby (within 10 kb of the preferred motif) for high-

scoring matches to the cross-species motif, i.e., the motif preferred by a homologous SPL in

another species. Importantly, these cross-species motif matches were selected to not overlap

any DAP-seq peaks, to reduce the likelihood that they are bound by the SPL under study. We

extracted 20 bp DNA fragments centered on these motifs to create the "cross-species motif

sequences". Thus, for each SPL protein, we generated two matched DNA input sets: one

representing motifs preferred and bound by this SPL and one representing motifs not preferred

or bound by this SPL but preferred by a homologous SPL in another species.

Each protein-DNA combination was then modeled using AF3 with at least six replicate

prediction runs using different random seeds. Each run used either a one-copy or two-copy

version of the TF sequence paired with one of the twenty 20 bp-long DNA sequences including

the forward and reverse complement (RC) strands, as well as two zinc ions for each copy of the

protein. The resulting structures were visualized using ChimeraX v1.10.1 (59). Confidence in

structural predictions was assessed using AF3's standard metrics: predicted template modelling

(pTM) score, the interface predicted template modeling (ipTM) score, and predicted aligned

error (PAE). Local Interaction Score (LIS) calculations were performed by modifying the Jupyter

notebook provided using the recommended parameters (60). For each combination of protein

(one- or two-copy) and motif (species-preferred or cross-species motif) condition, we computed

the median LIS score across all replicate runs for each of the ten DNA sequence instances.

Wilcoxon rank-sum tests were used to compare the LIS score distributions of the ten motif

matches in the species-preferred motif sequences versus the cross-species motif sequences.

RESULTS

Genome-wide binding site analysis of *Arabidopsis* SPL family by DAP-seq

We applied DAP-seq to all 16 members of the SPL family in *Arabidopsis,* successfully

identifying between 854 to 34,301 peaks (read enrichment q-value ≤ 0.01) for 14 members

(Figure 1A). By using ampDAP-seq, which assays protein binding on a PCR-amplified DNA

library, we obtained 2,189 peaks for AtSPL16 (Figure 1A). With the exception of AtSPL11 and

AtSPL13, all other AtSPLs had thousands of peaks, indicating strong DNA binding activities. In

contrast, AtSPL11 (854 peaks) or AtSPL13 (no peaks) may have potential DNA binding

activities that are less detectable under the tested experimental conditions in DAP-seq, or they

might require cofactors to enhance their DNA binding activity. The genome-wide binding site

maps of the AtSPL members showed both conserved and variable binding signals at the

promoter regions of the well-known downstream targets, for example, the flowering

development gene *SOC1* and an auxin response gene *ARF8* (Figure 1B and 1C). The

distribution of binding events relative to genomic features was similar for all the AtSPL

members, with around 30-40% of the binding events located in proximal promoter regions of -

1000 bp to 500 bp from the transcription start sites (TSS) (Figure 1D).

To gain insight into the potential biological functions of the AtSPL family members based on

their DNA binding profiles, we performed Gene Ontology (GO) enrichment analysis of genes

associated with the top 3000 binding events ranked by DAP-seq binding site scores. The top ten

enriched biological processes across family members revealed their shared and distinct

functions (Figure 1E). Notably, hormonal responses, such as response to auxin and abscisic

acid (ABA), were highly enriched for AtSPL15 and AtSPL9. Hypoxia and oxygen-related

responses were more enriched for AtSPL15, AtSPL1, AtSPL2, AtSPL4 and AtSPL12. In

contrast, the GO term "photosynthesis, light harvesting in photosystem I" was uniquely enriched

for AtSPL8, suggesting a potential role in capturing light energy. Interestingly, metal ion

homeostasis processes, such as cellular copper ion homeostasis and transition metal ion

transport, were predominantly enriched for AtSPL7 DAP-seq targets, consistent with its

previously characterized function in regulating copper uptake and distribution, which is essential

for plants' adaptation to low copper conditions (13, 52, 61, 62).

We next evaluated the potential for DAP-seq reported binding events to regulate gene

expression by testing the association between the DAP-seq predicted target genes and

differentially expressed genes (DEG) in *spl* mutants, in particular, in the root of *spl7* mutant from

a published dataset (52). Using Gene Set Enrichment Analysis (GSEA) tests (63, 64), we found

that among the AtSPL members that had strong GO term enrichment, DAP-seq targets of

AtSPL7 were the most enriched for down-regulated DEGs between *spl7* mutant and wild-type

Col-0 (up-regulated comparing wild-type and mutant; Figure 1F), suggesting that DAP-seq can

predict transcriptionally regulated genes by the relevant TFs for further functional studies.

Genome-wide binding and DNA sequence motifs support two major classes in the AtSPL

family

To investigate the relationships between the *Arabidopsis* SPL family members in terms of their

DNA binding specificities, we first clustered the AtSPLs based on their genome-wide binding

patterns reported by DAP-seq. To do this, we first determined the regions of the highest DAP-

seq read signals among all the AtSPL binding profiles and used the read enrichment signal in

these regions to calculate the pairwise distances between the individual TFs (47). Hierarchical

clustering of the distance matrix revealed two major subgroups (Figure 2A): Class A containing

four SPLs (AtSPL2, 10, 7 and 11) and Class B containing the rest of the SPLs (AtSPL9, 15, 1,

12, 4, 14, 5, 6, 3, 8). We then performed *de novo* motif discovery for each AtSPL using

sequences in the 1000 peaks with the highest significance of DAP-seq read enrichment in each

experiment. We found that the most significant motifs recognized by the Class A members

shared a core GTAC motif, flanked by a weaker T base on either side, while Class B members

preferred a core GTACGG motif (Figure 2B). These results suggest Class A and Class B

members could bind to distinct sets of genes. For example, the well-known gene *HSL2* (also

known as *VAL2*), which promotes vegetative phase change (65), showed strong binding signals

at its promoter region from all Class B members that preferred the GTACGG motif. In contrast,

relatively low binding signals were observed for Class A members that preferred the GTAC motif

(Figure 2C). Additionally, in Class B, motifs for AtSPL9 and AtSPL15 (close relatives derived

from a duplication event) contained weaker GG consensus sequence adjacent to the GTAC

core, suggesting these two paralogs bind DNA with more flexibility outside of the core motif than

other Class B members.

DAP-seq revealed differential DNA binding specificities between paralogs AtSPL9 and AtSPL15

To investigate the functional coherence or divergence of the evolutionarily conserved AtSPL9 and AtSPL15, we conducted differential binding analysis (54) between the AtSPL9 and AtSPL15 DAP-seq data (54). We identified 5,066 AtSPL15-prefered and 3,122 AtSPL9-prefered binding sites at FDR threshold of 0.05 (Figure 3A; Supplementary Table S2), about 18% and 15% of the binding sites for these two SPLs, indicating potential functional distinctions between them (Figure 3A). Strikingly, distinct biological functions were enriched for genes associated with the AtSPL15- and AtSPL9-preferred binding events (Figure 3B). AtSPL15-preferred gene targets were strongly enriched in processes related to ion homeostasis, positive regulation of transcription, response to oxygen and hypoxia, and auxin response. In contrast, AtSPL9-preferred gene targets were mainly enriched in secondary cell wall biogenesis, responses to photooxidative stress and regulation of the abscisic acid (ABA)-activated signaling pathway. These results suggest that beyond the conserved functions, AtSPL15 and AtSPL9 could have distinct roles in regulating different biological processes, and DNA binding preferences in the genome could mediate their specialized functions in plant development and stress responses.

To understand the mechanism underlying the differential DNA binding preferences between AtSPL9 and AtSPL15, we performed *de novo* motif discovery using the sequences in the 1000 differentially bound peaks with the highest increase in binding by AtSPL15 and AtSPL9, respectively (Figure 3C). We found that the most significant motifs in the AtSPL15-preferred binding sites contained the consensus sequences GTACgg and TGTACT, while AtSPL9 preferred motifs contained the consensus sequences GNCGG and GTAC. As an example, the differentially bound peak at the promoter region of the gene *DROUGHT-INDUCED 21* (*AtDI21*; AT4G15910) showed strong binding by AtSPL9 but not by AtSPL15, and it contained a binding site sequence GTAC (Figure 3D). To validate that the differential binding events could influence gene expression, we performed reporter assays (66) by cloning a 758 bp sequence from the promoter region of *DI21*, including the 200 bp AtSPL9-preferred DAP-seq peak, into the reporter plasmid *pUC-GUS* (*DI21pro::GUS*) to drive the GUS reporter gene expression. When we transiently expressed AtSPL9 and AtSPL15 by transfecting SPL9 and SPL15 separately, we found that AtSPL9 induced significantly higher reporter activation compared to AtSPL15 (Figure 3E). These results support our hypothesis that AtSPL15 and AtSPL9 have distinct binding preferences to DNA, resulting in differences in target gene regulation.

Comparative analysis of SPL target genes in *Arabidopsis*, maize, and wheat suggests
conserved functions

To characterize the DNA binding specificities among SPL homologs across plant species, we
compared the conserved genes associated with DAP-seq peaks - referred to as "target genes" -
of AtSPL9/15 in *Arabidopsis*, ZmSBP8/30 in maize (67) and TaSPL7/13 in wheat (including
three paralogs from the A/B/D subgenome) (26) from DAP-seq experiments conducted on their
respective genomes (26, 67). By comparing the target genes of maize ZmSBP8/30 to those of
Arabidopsis AtSPL9/15, we identified 4,557 unique target genes for AtSPL9/15, 1,125 unique
target genes for ZmSBP8/30, and 492 conserved target genes shared between the two species
(Figure 4A; Supplementary Table S3). Similarly, when comparing wheat TaSPL7/13 target
genes with the target genes of AtSPL9/15 in *Arabidopsis*, we found 4,017 unique target genes
for *Arabidopsis* AtSPL9/15, 2,709 unique target genes for wheat TaSPL7/13, and 939
conserved target genes (Figure 4B; Supplementary Table S4).

To gain insight into the function of these conserved target genes, we performed GO enrichment
analysis and found four major clusters of enriched biological processes (Figure 4C). Cluster 2
and Cluster 4 highlighted functions conserved across the three species, primarily related to cell
wall biogenesis, transcriptional regulation, and pattern specification. Cluster 1 also revealed
functions that were conserved among all three species, such as xylem and cell wall
development, while also suggesting *Arabidopsis*-wheat-specific conserved processes such as
cell wall modification, secondary metabolite biosynthesis, and lignin metabolic processes. These
GO terms suggest the potential role of TaSPL7/13 in specific aspects of wheat architecture,
possibly related to structural integrity and stress responses. Cluster 3 GO terms were
significantly enriched for SPL targets conserved between *Arabidopsis* and maize, including
processes associated with different developmental stages, such as floral organ development,
cotyledon development, and post-embryonic development. Additionally, we observed processes
such as plant organ specification and gibberellin response, which are closely related to
flowering time and overall plant development. Overall, these results suggest that different
biological functions and developmental processes are targeted by the SPL shared target genes
in different species, offering insights into the evolutionary trajectories of the SPL homologs in
the three species spanning monocot to dicot.

SPL homologs bind to variations of the GTAC motifs

Since conservation of binding motifs is often used as the basis for cross-species prediction of
cis-regulatory elements, we analyzed the enriched DNA sequence motifs in the DAP-seq peaks
associated with the conserved target genes in the three species. Unexpectedly, we identified
three different types of DNA binding motifs, although they all shared or partially shared the core
GTAC sequence (Figure 4D-E, red highlights). In the *Arabidopsis* genome, AtSPL9/15
recognizes motifs with a GTAC core adjacent to a GG dinucleotide. In the maize genome,
ZmSBP8/30 recognizes a back-to-back repeat of the GTAC sequence, GTACGTRC, although
one of the GTAC elements is weak. In the wheat genome, TaSPL7/13 binds to a motif
containing two strong GTAC elements overlapping by one base pair, resulting in the consensus
GTASTAC (GTACTAC or GTAGTAC). These different yet related motif architectures suggest
differences in genome-wide DNA recognition mechanisms among the SPL homologs.
Importantly, the more complex back-to-back or overlapping motifs in maize and wheat may
contribute to enhancing binding specificities in the much larger and complex genomes of maize
(2.3 Gbp) and wheat (17 Gbp) relative to *Arabidopsis* (135 Mbp).

Structure modeling links protein-protein interaction preferences to DNA binding motif
variation across SPL homologs

To explore the potential structural mechanisms underlying the motif variation among SPL
homologs, we performed structural prediction of the SPL-DNA binding complexes using
AlphaFold3 (AF3) (58). We began by modeling the interaction between full-length SPL proteins
and DNA sequences containing their species-preferred binding motifs. These sequences were
drawn from DAP-seq peaks associated with conserved target genes, representing binding
events that are conserved across species and therefore likely to be functionally important. As
shown in Figure 5A for the representative homologs, TaSPL13A interacting with the TaSPL-
preferred GTASTAC motif and AtSPL9 interacting with the AtSPL-preferred GTAC motif, the
predicted structures had low overall confidence scores (ipTM: 0.51-0.59; pTM: 0.29-0.31), likely
due to high fractions of disordered regions outside the conserved SBP DNA-binding domain
(fraction disordered: 0.80-0.81). Nevertheless, when we superimposed the AF3-predicted
structures with the experimentally solved X-ray crystal structure of AtSPL5 (PDB 8J49 Chain A
(68)), we observed strong structural similarity in the SBP domains, including two well-conserved
zinc-binding pockets (RMSD 0.685-0.672 Å, 63 Cα atoms; Figure 5B). Furthermore, the
predicted structures consistently positioned the zinc finger regions in the C-terminal portion of
the SBP domain in proximity to the DNA (Figure 5B), consistent with prior solution studies
showing that this zinc-binding site is required for DNA binding (69). While these results

confirmed structural conservation of the SBP domain across species, they did not readily

explain the observed motif divergence across species.

We next hypothesized that the wheat SPLs' strong preference for the overlapping GTAC motifs

(GTASTAC) may arise from enhanced protein-protein interactions (PPIs) between SPL

monomers. This hypothesis was motivated by three observations. First, DAP-seq has been

shown to capture dimeric protein binding to DNA for several TF families in multiple species (45,

70, 71). Secondly, DNA-guided TF interaction is known to produce overlapping or composite

motifs that partially reflect specificities of individual TFs (70, 72). Third, dimerization has been

observed in several SPLs in different species. In *Arabidopsis*, structural and biochemical studies

show that the SBP domains of *Arabidopsis* SPLs bind DNA as monomers (69, 73, 74), while

full-length proteins like AtSPL7 and AtSPL9 are capable of dimerization, though the functional

role of this dimerization in DNA binding remains unclear (75, 76). In particular, AtSPL7

dimerization may modulate its regulatory activity *in vivo*, possibly by preventing its translocation

from the cytoplasm to the nucleus (75). In maize, the SBP-box TF ZmTGA1 forms both

monomeric and dimeric complexes to bind DNA. Notably, the dimeric form of ZmTGA1 exhibits

higher stability than that of its teosinte ortholog, and this increased dimer stability was implicated

as a key molecular mechanism underlying the naked kernel phenotype that distinguishes

domesticated maize from its wild progenitor, teosinte (77). Together, these observations

suggest that PPI-driven DNA binding could underlie the emergence of complex binding motifs in

SPLs.

To test this hypothesis, we used AF3 to predict SPL-DNA complex structures with two copies of

each SPL interacting with their preferred binding motif (species-preferred motif) or with a motif

preferred by SPL homologs from a different species (cross-species motif) (see MATERIALS

AND METHODS). These DNA sequences were drawn from genomic regions associated with

conserved SPL target genes and centered on high-scoring motif matches either within a SPL's

own DAP-seq peaks (species-preferred motif sequences) or from nearby unbound regions

containing the motif of the SPL homolog in a different species (cross-species motif sequences).

This design allowed us to compare predicted structural interactions in sequence contexts that

are biologically relevant while minimizing confounding effects from local genomic features.

Figure 5C shows representative results for TaSPL13A, comparing its interaction with the

species-preferred GTASTAC motif (structure i) versus the cross-species GTAC motif preferred

by AtSPLs (structure ii). In the predicted aligned error (PAE) matrices, a strong low-error block

was present between the two TaSPL13A monomers in the GTASTAC-bound complex but

absent in the GTAC-bound complex (cyan arrows), suggesting a more stable dimer interface

between the two TaSPLA monomers in the presence of the species-preferred motif.

Since AF3 is better at predicting the overall geometry of TF-TF-DNA complexes than localizing

the precise TF-TF or TF-DNA contacts (70, 78), we did not analyze the direct contacts in the

predicted structures. Instead, we calculated the local interaction score (LIS) between the

subunits in the predicted structures (60). LIS has improved sensitivity in detecting PPIs,

particularly for those that involve flexible and small interfaces where traditional pTM scores are

less informative (60). The LIS heatmaps shown in Figure 5C confirmed that both monomer-

monomer (cyan boxes) and monomer-DNA (blue boxes) interactions were stronger in the

TaSPL13A complex bound to the GTASTAC motif (structure i) compared to that bound to the

GTAC motif (structure ii). In contrast, control structures predicted using a single copy of

TaSPL13A (structures iii and iv; Figure 5C) did not show differences in LIS values for the

monomer-DNA interaction (purple boxes) that are consistent with the motif preferences,

supporting the idea that PPIs between TaSPL13A monomers stabilize the binding to the

GTASTAC motif.

To generalize these results, we similarly modeled one- and two-copy SPL-DNA complexes for

AtSPL9, AtSPL15, six wheat TaSPL7/13 paralogs, and two maize ZmSBPs ZmSBP8 and

ZmSBP30. For each protein, we predicted structures with both the species-preferred motif

sequences and cross-species motif sequences (see MATERIALS AND METHODS), and

computed the LIS values for the monomer-monomer and monomer-DNA interactions. The

resulting LIS distributions across ten motif instances for species-preferred and cross-species

motifs are shown as dotplots and boxplots in Figure 5D. Statistical comparisons between

predictions for species-preferred versus cross-species motifs (Wilcoxon rank-sum test, $p < 0.05$;

orange background panels in Figure 5D) revealed a consistent and striking pattern. *Arabidopsis*

AtSPL9 and AtSPL15 showed significantly higher LIS values for the GTAC motif only in the one-

copy models. In contrast, most wheat SPL paralogs exhibited significantly stronger monomer-

monomer and monomer-DNA interaction scores when modeled with the GTASTAC motif, but

only in the two-copy structures. These results support a model in which PPIs between TaSPL

monomers enhance binding to the overlapping GTASTAC motif, whereas AtSPLs bind DNA as

monomers with a preference for the short GTAC motif.

Interestingly, applying the same analysis to maize ZmSBP8 and ZmSBP30 did not show

consistent preferences between one-copy and two-copy structures for either the GTAC or

GTACGTRC motifs (AtSPL-preferred or ZmSBP-preferred) (Figure S1). This may reflect the
flexibility of maize SBPs in forming monomeric or dimeric complexes, such as those found for
ZmTGA1 (77).

DISCUSSION

TFs regulate gene expression by binding to specific DNA sequences, and variations in their
binding sites can lead to differential regulation of genes within a TF family in a species and
across different species. Our analysis of the DAP-seq binding profiles of the *Arabidopsis* SPL
family revealed two major classes of binding site motifs (Figure 2), yet the enriched GO terms of
the target genes near the most enriched binding sites were much more diverse among the
family members (Figure 1E). This suggests that even closely related TFs, such as AtSPL9 and
AtSPL15, can develop distinct regulatory roles through subtle differences in their DNA-binding
preferences. By analyzing this comprehensive cistrome dataset for SPL TF family in
Arabidopsis, our study offers insights into how specific DNA-binding characteristics contribute to
the functional diversity within the SPL family of TFs.

Gene duplication is a key feature of genomic architecture, especially within TF families, leading
to opportunities for subfunctionalization and neofunctionalization (79–82). Some duplicated TFs
retain ancestral functions by maintaining their original DNA binding preferences, while others
acquire mutations that allow them to bind different DNA sequences and thus regulate different
sets of genes (2, 83, 84). Such changes can enhance a species' ability to respond to broader
environmental challenges including biotic or abiotic stresses, or activate new developmental
pathways for adaptation. DAP-seq offers a powerful approach to investigate this functional
divergence mediated by DNA binding, allowing for detailed scrutiny of DNA binding motifs and
target genes among paralogous TFs, even when genetic redundancy makes functional studies
challenging. For example, our analysis revealed that even slight changes in DNA binding motifs
between the duplicated homologs AtSPL9 and AtSPL15 led to distinct target gene sets, with
AtSPL15 showing preferences for genes involved in auxin responses, while AtSPL9 favoring
targets related to ABA signaling. These variations help differentiate the roles of closely related
TFs in regulating downstream targets and ensuring functional specificity. We expect that the
same analysis can be applied to other TF homologs within the SBP/SPL family or extend to
other TF families to gain a broader understanding of TF functional diversity.

The evolution of TF-DNA interactions is a complex and dynamic process that involves multiple
factors, including changes in the TF proteins, their DNA-binding motifs, and the genomic context

of the binding sites such as DNA shape and chemical modifications (3). In this study, we

observed that the conserved functions of the SPL/SBP homologs based on their DNA binding

targets were distributed differently in *Arabidopsis*, maize, and wheat: while some functions

maintained across all three species, others were shared only between specific pairs, such as

Arabidopsis and maize or *Arabidopsis* and wheat. For example, common regulatory functions

related to developmental processes were found between *Arabidopsis* and maize, while

Arabidopsis and wheat shared functions in stress response and structural regulation. Analysis of

DAP-seq binding sequences revealed species-specific preferences for related but distinct motif

variants, which may contribute to the divergence of regulatory networks across species. These

findings provide insights into how the SPL/SBP TFs have evolved to balance the core,

conserved functions with species-specific adaptations, highlighting the power of DAP-seq to

shed light on the evolutionary trajectories of TF networks across plant species.

A particularly interesting finding emerged when we compared the DNA binding motifs among

the SPL/SBP homologs across species, revealing that the SPL/SBP homologs in maize and

wheat bind to longer motif types that are variations of the core GTAC sequences recognized by

the *Arabidopsis* protein. Specifically, the GTACGTRC motif in maize is a back-to-back

palindromic sequence of two GTAC units, while the GTASTAC motif in wheat is a palindromic

sequence with one base pair overlap between two GTAC units (Figure 4E). Palindromic motifs

have also been observed for dimers of the bZIP family of TFs, suggesting such motifs can be a

general mechanism for refining TF-DNA recognition (64). Many TF families, including

AP2/ERF, bZIP, ARF, homeodomain and MADS-box, are known to form dimers or higher-order

complexes to bind DNA and regulate gene expression (28, 85, 86). Using AF3 structural

modeling, we found that TaSPL7/13 in wheat may recognize the palindromic GTASTAC motif

through DNA-guided dimerization, enabling binding to a more complex motif configuration than

the GTAC motif preferred by AtSPLs. The ability to recognize longer, more complex motifs

through PPI could be a mechanism that allows SBP/SPL TFs to increase DNA binding

specificities to achieve precise gene regulation, particularly in the context of complex plant

genomes like those of maize and wheat. Our findings underscore the potential role of PPI in the

evolution of DNA recognition mechanisms by TFs, offering insights into how TFs adapt to

regulate diverse regulatory programs in plants with highly divergent genome complexities.

AUTHOR CONTRIBUTIONS

S.C.H. and M.L.: Conceptualization, Formal analysis, Visualization, Methodology, Validation,
Writing-original draft. W.L. and Y.Z.: Validation. M.G. and A.G.: Formal analysis. T.Y. and J.C.:
Methodology, Validation. L.F.D.C.: Investigation, Methodology. S.C.H. and A.G.: Funding
acquisition. All authors: review & editing.

ACKNOWLEDGEMENTS

We thank Joseph Ecker for the gift of TF expression plasmids. We also thank Aurelia Li for
technical assistance. We acknowledge the Zegar Family Foundation for their generous support.
This work was supported in part through the NYU IT High Performance Computing resources,
services, and staff expertise.

FUNDING

This work was supported by the NIH award R35GM138143 to S.C.H. and NSF Plant Genome
Research Project grant IOS-1916804 to S.C.H. and A.G. This material was also in part based
on work supported by the Center for Bioenergy Innovation (CBI), U.S. Department of Energy,
Office of Science, Biological and Environmental Research Program under Award Number
ERKP886; Biological and Environmental Research Program under Award Number ERKPA88;
and the Oak Ridge National Laboratory Director's R&D (DRD) Program 12093. Oak Ridge
National Laboratory is managed by UT-Battelle, LLC for the Office of Science of the U.S.
Department of Energy under Contract Number DE-AC05-00OR22725.

CONFLICT OF INTEREST

The authors declare they have no competing interests.

Figure 1: DAP-seq identifies genome-wide binding events for SPL TFs that are biologically relevant. (**A**) Number of DAP-seq peaks for all SPL members in *Arabidopsis*. ** indicates ampDAP-seq. (**B**) *Arabidopsis* AtSPL DAP-seq binding signal at the promoter region of a known target gene *SOC1*. (**C**) *Arabidopsis* SPL DAP-seq binding signal at the promoter of *ARF8*. (**D**) Distribution of AtSPL DAP-seq peaks at genome annotation features. Promoter is defined as -1000 bp upstream to 500 bp downstream of TSS. (**E**) Enriched Gene Ontology biological process terms for DAP-seq predicted targets of *Arabidopsis* SPLs. (**F**) GSEA of AtSPL DAP-seq targets and the differentially expressed genes in *spl7* mutant vs. Col-0 wild type (WT).

Figure 2: Genome-wide binding profile comparison and DNA motifs of the *Arabidopsis* SPL

family. (**A**) Hierarchical clustering of AtSPL DAP-seq binding profiles splits the family members

into Class A and Class B. (**B**) PWM models of the enriched motifs from the top 1000 DAP-seq

peaks for each AtSPL. (**C**) Class A and Class B AtSPLs show different DAP-seq binding signals

at the promoter of phase change related gene *AtHSL2* (*AtVAL2*).

Figure 3: Differential binding events, target genes and motifs comparing *Arabidopsis* AtSPL9 versus AtSPL15. (**A**) DAP-seq read signals at differentially bound peaks of AtSPL15 and AtSPL9. (**B**) Top 10 enriched GO Biological Process terms for genes associated with AtSPL15-preferred and AtSPL9-preferred peaks. (**C**) PWM motif models enriched in AtSPL15- and AtSPL9-preferred binding sites. (**D**) AtSPL9-preferred binding site upstream of gene *AtDI21*. (E) Transient expression of AtSPL9 and AtSPL15 induced significantly different expressions of the reporter AtDI21::GUS. Error bars are standard error of the mean (n=3) and P-values were computed by two sample t-tests followed by Bonferroni correction.

Figure 4: Comparative analysis of DAP-seq target genes and binding site sequences revealed

variation of DNA binding mechanism among SPL homologs. (**A**) Upset plot comparing the target

gene overlap between *Arabidopsis* AtSPL9/15 and maize ZmSBP8/30. (**B**) UpSet plot

comparing the target gene overlap between *Arabidopsis* AtSPL9/15 and wheat TaSPL7/13. (**C**)

GO enrichment analysis showing the biological processes enriched for targets conserved

between SPL homologs between *Arabidopsis* and maize and between *Arabidopsis* and wheat.

(**D**) DNA binding motifs discovered by MEME for DAP-seq peaks associated with conserved

SPL targets in *Arabidopsis*, maize and wheat.

Figure 5: AlphaFold3 (AF3) structure modeling of DNA binding by *Arabidopsis* and wheat SPL

proteins. (**A**) Representative AF3-predicted structures of TaSPL13A binding to its preferred

DNA motif GTASTAC (left) and AtSPL9 binding to its preferred motif GTAC (right). (**B**)

Superposition of the SBP domains from the predicted structures in (A) with the X-ray crystal

structure of AtSPL5 (PDB: 8J49), highlighting the conserved zinc-binding regions. (**C**) AF3-

predicted complex structures, Predicted aligned error (PAE) matrices, and Local Interaction

Score (LIS) heatmaps for one or two copies of TaSPL13A interacting with either the species-

preferred GTASTAC motif or the cross-species GTAC motif (see MATERIALS AND

METHODS). Colored boxes highlight predicted monomer-monomer (cyan), monomer-DNA

(blue) interactions in two-copy protein structures, and monomer-DNA interactions (purple) in

one-copy protein structures. RC: Reverse Complement. (**D**) Summary of LIS values across

predicted SPL-DNA structures. Each point represents the median LIS from n≥6 replicate

predictions for a given motif instance. For *Arabidopsis* SPLs (AtSPL9 and AtSPL15), structures

were modeled with one or two protein copies interacting with either GTAC motifs (species-

preferred motifs bound in DAP-seq) or GTASTAC motifs (cross-species motifs, unbound). For

wheat SPLs (TaSPL7A/B/D and TaSPL13A/B/D), structures were modeled analogously with

GTASTAC (species-preferred motifs bound in DAP-seq) and GTAC (cross-species motifs,

unbound) motifs. Boxplots summarize monomer-monomer and monomer-DNA interaction

scores for each configuration. P-values were computed by two sample Wilcoxon rank-sum tests

between species-preferred and cross-species motif conditions and orange-shaded panels

indicate statistically significant differences (p < 0.05).

REFERENCES

1. Panchy,N., Lehti-Shiu,M. and Shiu,S.-H. (2016) Evolution of Gene Duplication in Plants. *Plant*
681   *Physiol.*, **171**, 2294–2316.

682   2. McKeown,A.N., Bridgham,J.T., Anderson,D.W., Murphy,M.N., Ortlund,E.A. and
683   Thornton,J.W. (2014) Evolution of DNA specificity in a transcription factor family
684   produced a new gene regulatory module. *Cell*, **159**, 58–68.

685   3. Rogers,J.M. and Bulyk,M.L. (2018) Diversification of transcription factor-DNA interactions and
686   the evolution of gene regulatory networks. *Wiley Interdiscip. Rev. Syst. Biol. Med.*, **10**,
687   e1423.

688   4. Birkenbihl,R.P., Jach,G., Saedler,H. and Huijser,P. (2005) Functional dissection of the plant-
689   specific SBP-domain: overlap of the DNA-binding and nuclear localization domains. *J.*
690   *Mol. Biol.*, **352**, 585–596.

691   5. Klein,J., Saedler,H. and Huijser,P. (1996) A new family of DNA binding proteins includes
692   putative transcriptional regulators of the Antirrhinum majus floral meristem identity gene
693   SQUAMOSA. *Mol. Gen. Genet.*, **250**, 7–16.

694   6. Wang,J.-W., Czech,B. and Weigel,D. (2009) miR156-regulated SPL transcription factors
695   define an endogenous flowering pathway in Arabidopsis thaliana. *Cell*, **138**, 738–749.

7. Xu,M., Hu,T., Smith,M.R. and Poethig,R.S. (2016) Epigenetic Regulation of Vegetative Phase
Change in Arabidopsis. *Plant Cell*, **28**, 28–41.

8. Wang,J., Zhou,L., Shi,H., Chern,M., Yu,H., Yi,H., He,M., Yin,J., Zhu,X., Li,Y., *et al.* (2018) A
single transcription factor promotes both yield and immunity in rice. *Science*, **361**, 1026–
1028.

9. Zheng,C., Ye,M., Sang,M. and Wu,R. (2019) A Regulatory Network for miR156-SPL Module
in. *Int. J. Mol. Sci.*, **20**.

10. Yu,N., Niu,Q.-W., Ng,K.-H. and Chua,N.-H. (2015) The role of miR156/SPLs modules in
Arabidopsis lateral root development. *Plant J.*, **83**, 673–685.

11. The miR156/SPL Module, a Regulatory Hub and Versatile Toolbox, Gears up Crops for
Enhanced Agronomic Traits (2015) *Mol. Plant*, **8**, 677–688.

12. Jerome Jeyakumar,J.M., Ali,A., Wang,W.-M. and Thiruvengadam,M. (2020) Characterizing
the Role of the miR156-SPL Network in Plant Development and Stress Response.
Plants, **9**.

13. Yamasaki,H., Hayashi,M., Fukazawa,M., Kobayashi,Y. and Shikanai,T. (2009) SQUAMOSA
Promoter Binding Protein-Like7 Is a Central Regulator for Copper Homeostasis in
Arabidopsis. *Plant Cell*, **21**, 347–361.

14. Kim,J.J., Lee,J.H., Kim,W., Jung,H.S., Huijser,P. and Ahn,J.H. (2012) The microRNA156-
SQUAMOSA PROMOTER BINDING PROTEIN-LIKE3 module regulates ambient
temperature-responsive flowering via FLOWERING LOCUS T in Arabidopsis. *Plant
Physiol.*, **159**, 461–478.

15. Nagpal,P., Ellis,C.M., Weber,H., Ploense,S.E., Barkawi,L.S., Guilfoyle,T.J., Hagen,G.,
Alonso,J.M., Cohen,J.D., Farmer,E.E., *et al.* (2005) Auxin response factors ARF6 and
ARF8 promote jasmonic acid production and flower maturation. *Development*, **132**,
4107–4118.

16. Wu,G., Park,M.Y., Conway,S.R., Wang,J.-W., Weigel,D. and Poethig,R.S. (2009) The
sequential action of miR156 and miR172 regulates developmental timing in Arabidopsis.
Cell, **138**, 750–759.

17. Hu,L., Qi,P., Peper,A., Kong,F., Yao,Y. and Yang,L. (2023) Distinct function of SPL genes in
age-related resistance in Arabidopsis. *PLoS Pathog.*, **19**, e1011218.

18. Wang,J.-W., Schwab,R., Czech,B., Mica,E. and Weigel,D. (2008) Dual Effects of miR156-
Targeted SPL Genes and CYP78A5/KLUH on Plastochron Length and Organ Size in
Arabidopsis thaliana. *Plant Cell*, **20**, 1231–1243.

19. Yang,Z., Wang,X., Gu,S., Hu,Z., Xu,H. and Xu,C. (2008) Comparative study of SBP-box
gene family in Arabidopsis and rice. *Gene*, **407**, 1–11.

20. Preston,J.C. and Hileman,L.C. (2013) Functional Evolution in the Plant SQUAMOSA-
PROMOTER BINDING PROTEIN-LIKE (SPL) Gene Family. *Front. Plant Sci.*, **4**, 80.

21. Schwarz,S., Grande,A.V., Bujdoso,N., Saedler,H. and Huijser,P. (2008) The microRNA
regulated SBP-box genes SPL9 and SPL15 control shoot maturation in Arabidopsis.

Plant Mol. Biol., **67**, 183–195.

22. Hyun,Y., Richter,R., Vincent,C., Martinez-Gallegos,R., Porri,A. and Coupland,G. (2016)
    Multi-layered Regulation of SPL15 and Cooperation with SOC1 Integrate Endogenous
    Flowering Pathways at the Arabidopsis Shoot Meristem. *Dev. Cell*, **37**, 254–266.

23. Song,X., Lu,Z., Yu,H., Shao,G., Xiong,J., Meng,X., Jing,Y., Liu,G., Xiong,G., Duan,J., *et al.*
    (2017) IPA1 functions as a downstream transcription factor repressed by D53 in
    strigolactone signaling in rice. *Cell Res.*, **27**, 1128–1141.

24. Chuck,G.S., Brown,P.J., Meeley,R. and Hake,S. (2014) Maize SBP-box transcription factors
    unbranched2 and unbranched3 affect yield traits by regulating the rate of lateral
    primordia initiation. *Proc. Natl. Acad. Sci. U. S. A.*, **111**, 18775–18780.

25. Wei,H., Zhao,Y., Xie,Y. and Wang,H. (2018) Exploiting SPL genes to improve maize plant
    architecture tailored for high-density planting. *J. Exp. Bot.*, **69**, 4675–4688.

26. Pei,H., Teng,W., Gao,L., Gao,H., Ren,X., Liu,Y., Jia,J., Tong,Y., Wang,Y. and Lu,Z. (2023)
    Low-affinity SPL binding sites contribute to subgenome expression divergence in
    allohexaploid wheat. *Sci. China Life Sci.*, **66**, 819–834.

27. Bartlett,A., O'Malley,R.C., Huang,S.-S.C., Galli,M., Nery,J.R., Gallavotti,A. and Ecker,J.R.
    (2017) Mapping genome-wide transcription-factor binding sites using DAP-seq. *Nat.
    Protoc.*, **12**, 1659–1672.

28. O'Malley,R.C., Huang,S.-S.C., Song,L., Lewsey,M.G., Bartlett,A., Nery,J.R., Galli,M.,
    Gallavotti,A. and Ecker,J.R. (2016) Cistrome and Epicistrome Features Shape the
    Regulatory DNA Landscape. *Cell*, **165**, 1280–1292.

29. Li,M. and Huang,S.-S.C. (2022) DNA Affinity Purification Sequencing (DAP-Seq) for
    Mapping Genome-Wide Transcription Factor Binding Sites in Plants. *Accelerated
    Breeding of Cereal Crops*, 10.1007/978-1-0716-1526-3_15.

30. Tiwari,S.B., Hagen,G. and Guilfoyle,T. (2003) The roles of auxin response factor domains in
    auxin-responsive transcription. *Plant Cell*, **15**, 533–543.

31. Broad Institute Picard Tools. *Broad Institute, GitHub Repository*.
    http://broadinstitute.github.io/picard/.

32. Galanti,L., Shasha,D. and Gunsalus,K.C. (2021) Pheniqs 2.0: accurate, high-performance
    Bayesian decoding and confidence estimation for combinatorial barcode indexing. *BMC
    Bioinformatics*, **22**, 359.

33. New York University GENEFLOW. *New York University Center for Genomics and System
    Biology Genomics Core, GitHub Repository*.
    https://github.com/gencorefacility/GENEFLOW.

34. Martin,M. (2011) Cutadapt removes adapter sequences from high-throughput sequencing
    reads. *EMBnet.journal*, **17**, 10–12.

35. Langmead,B. and Salzberg,S.L. (2012) Fast gapped-read alignment with Bowtie 2. *Nat.
    Methods*, **9**, 357–359.

36. Danecek,P., Bonfield,J.K., Liddle,J., Marshall,J., Ohan,V., Pollard,M.O., Whitwham,A.,
Keane,T., McCarthy,S.A., Davies,R.M., *et al.* (2021) Twelve years of SAMtools and
BCFtools. *Gigascience*, **10**.

37. Guo,Y., Mahony,S. and Gifford,D.K. (2012) High resolution genome wide binding event
finding and motif discovery reveals transcription factor spatial binding constraints. *PLoS*
Comput. Biol., **8**, e1002638.

38. Zhang,Y., Liu,T., Meyer,C.A., Eeckhoute,J., Johnson,D.S., Bernstein,B.E., Nusbaum,C.,
Myers,R.M., Brown,M., Li,W., *et al.* (2008) Model-based Analysis of ChIP-Seq (MACS).
Genome Biol., **9**, 1–9.

39. Ramírez,F., Ryan,D.P., Grüning,B., Bhardwaj,V., Kilpert,F., Richter,A.S., Heyne,S.,
Dündar,F. and Manke,T. (2016) deepTools2: a next generation web server for deep-
sequencing data analysis. *Nucleic Acids Res.*, **44**, W160-5.

40. Diesh,C., Stevens,G.J., Xie,P., De Jesus Martinez,T., Hershberg,E.A., Leung,A., Guo,E.,
Dider,S., Zhang,J., Bridge,C., *et al.* (2023) JBrowse 2: a modular genome browser with
views of synteny and structural variation. *Genome Biol.*, **24**, 74.

41. Bailey,T.L. and Elkan,C. (1994) Fitting a mixture model by expectation maximization to
discover motifs in biopolymers. *Proc. Int. Conf. Intell. Syst. Mol. Biol.*, **2**, 28–36.

42. Tremblay,B.J.-M. (2024) universalmotif: An R package for biological motif analysis. *Journal*
of Open Source Software, **9**, 7012.

43. Nettling,M., Treutler,H., Grau,J., Keilwagen,J., Posch,S. and Grosse,I. (2015) DiffLogo: a
comparative visualization of sequence motifs. *BMC Bioinformatics*, **16**, 1–9.

44. Wagih,O. (2017) ggseqlogo: a versatile R package for drawing sequence logos.
Bioinformatics, **33**, 3645–3647.

45. Galli,M., Chen,Z., Ghandour,T., Chaudhry,A., Gregory,J., Feng,F., Li,M., Schleif,N.,
Zhang,X., Dong,Y., *et al.* (2025) Transcription factor binding divergence drives
transcriptional and phenotypic variation in maize. *Nat Plants*, **11**, 1205–1219.

46. Zhu,T., Wang,L., Rimbert,H., Rodriguez,J.C., Deal,K.R., De Oliveira,R., Choulet,F., Keeble-
Gagnère,G., Tibbits,J., Rogers,J., *et al.* (2021) Optical maps refine the bread wheat
Triticum aestivum cv. Chinese Spring genome assembly. *Plant J.*, **107**, 303–314.

47. Tu,S., Li,M., Chen,H., Tan,F., Xu,J., Waxman,D.J., Zhang,Y. and Shao,Z. (2021) MAnorm2
for quantitatively comparing groups of ChIP-seq samples. *Genome Res.*, **31**, 131–145.

48. Wang,Q., Li,M., Wu,T., Zhan,L., Li,L., Chen,M., Xie,W., Xie,Z., Hu,E., Xu,S., *et al.* (2022)
Exploring Epigenomic Datasets by ChIPseeker. *Curr Protoc*, **2**, e585.

49. Cheng,C.-Y., Krishnakumar,V., Chan,A.P., Thibaud-Nissen,F., Schobel,S. and Town,C.D.
(2017) Araport11: a complete reannotation of the Arabidopsis thaliana reference
genome. *Plant J.*, **89**, 789–804.

50. Xu,S., Hu,E., Cai,Y., Xie,Z., Luo,X., Zhan,L., Tang,W., Wang,Q., Liu,B., Wang,R., *et al.*
(2024) Using clusterProfiler to characterize multiomics data. *Nat. Protoc.*,
10.1038/s41596-024-01020-z.

51. Gu,Z., Eils,R. and Schlesner,M. (2016) Complex heatmaps reveal patterns and correlations
in multidimensional genomic data. *Bioinformatics*, **32**, 2847–2849.

52. Kastoori Ramamurthy,R., Xiang,Q., Hsieh,E.-J., Liu,K., Zhang,C. and Waters,B.M. (2018)
New aspects of iron-copper crosstalk uncovered by transcriptomic characterization of
Col-0 and the copper uptake mutant spl7 in Arabidopsis thaliana. *Metallomics*, **10**, 1824–
1840.

53. Love,M.I., Huber,W. and Anders,S. (2014) Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol.*, **15**, 550.

54. Ross-Innes,C.S., Stark,R., Teschendorff,A.E., Holmes,K.A., Ali,H.R., Dunning,M.J.,
Brown,G.D., Gojis,O., Ellis,I.O., Green,A.R., *et al.* (2012) Differential oestrogen receptor
binding is associated with clinical outcome in breast cancer. *Nature*, **481**, 389–393.

55. Machanick,P. and Bailey,T.L. (2011) MEME-ChIP: motif analysis of large DNA datasets.
Bioinformatics, **27**, 1696–1697.

56. Hufford,M.B., Seetharam,A.S., Woodhouse,M.R., Chougule,K.M., Ou,S., Liu,J., Ricci,W.A.,
Guo,T., Olson,A., Qiu,Y., *et al.* (2021) De novo assembly, annotation, and comparative
analysis of 26 diverse maize genomes. *Science*, **373**, 655–662.

57. Goodstein,D.M., Shu,S., Howson,R., Neupane,R., Hayes,R.D., Fazo,J., Mitros,T., Dirks,W.,
Hellsten,U., Putnam,N., *et al.* (2012) Phytozome: a comparative platform for green plant
genomics. *Nucleic Acids Res.*, **40**, D1178-86.

58. Abramson,J., Adler,J., Dunger,J., Evans,R., Green,T., Pritzel,A., Ronneberger,O.,
Willmore,L., Ballard,A.J., Bambrick,J., *et al.* (2024) Accurate structure prediction of
biomolecular interactions with AlphaFold 3. *Nature*, **630**, 493–500.

59. Meng,E.C., Goddard,T.D., Pettersen,E.F., Couch,G.S., Pearson,Z.J., Morris,J.H. and
Ferrin,T.E. (2023) UCSF ChimeraX: Tools for structure building and analysis. *Protein*
Sci, **32**, e4792.

60. Kim,A.-R., Hu,Y., Comjean,A., Rodiger,J., Mohr,S.E. and Perrimon,N. (2024) Enhanced
Protein-Protein Interaction Discovery via AlphaFold-Multimer. *bioRxiv*,
10.1101/2024.02.19.580970.

61. Araki,R., Mermod,M., Yamasaki,H., Kamiya,T., Fujiwara,T. and Shikanai,T. (2018) SPL7
locally regulates copper-homeostasis-related genes in Arabidopsis. *J. Plant Physiol.*,
224–225, 137–143.

62. Bernal,M., Casero,D., Singh,V., Wilson,G.T., Grande,A., Yang,H., Dodani,S.C.,
Pellegrini,M., Huijser,P., Connolly,E.L., *et al.* (2012) Transcriptome sequencing identifies
SPL7-regulated copper acquisition genes FRO4/FRO5 and the copper dependence of
iron homeostasis in Arabidopsis. *Plant Cell*, **24**, 738–761.

63. Korotkevich,G., Sukhov,V., Budin,N., Shpak,B., Artyomov,M.N. and Sergushichev,A. (2021)
Fast gene set enrichment analysis. *bioRxiv*, 10.1101/060012.

64. Subramanian,A., Tamayo,P., Mootha,V.K., Mukherjee,S., Ebert,B.L., Gillette,M.A.,
Paulovich,A., Pomeroy,S.L., Golub,T.R., Lander,E.S., *et al.* (2005) Gene set enrichment

851  analysis: a knowledge-based approach for interpreting genome-wide expression profiles.
852  *Proc. Natl. Acad. Sci. U. S. A.*, **102**, 15545–15550.

853 65. Fouracre,J.P., He,J., Chen,V.J., Sidoli,S. and Poethig,R.S. (2021) VAL genes regulate
854  vegetative phase change via miR156-dependent and independent mechanisms. *PLoS*
855  *Genet.*, **17**, e1009626.

856 66. Yoo,S.-D., Cho,Y.-H. and Sheen,J. (2007) Arabidopsis mesophyll protoplasts: a versatile
857  cell system for transient gene expression analysis. *Nat. Protoc.*, **2**, 1565–1572.

858 67. Ricci,W.A., Lu,Z., Ji,L., Marand,A.P., Ethridge,C.L., Murphy,N.G., Noshay,J.M., Galli,M.,
859  Mejía-Guerra,M.K., Colomé-Tatché,M., *et al.* (2019) Widespread long-range cis-
860  regulatory elements in the maize genome. *Nat Plants*, **5**, 1237–1249.

861 68. Yan,X., Yuan,X., Lv,J., Zhang,B., Huang,Y., Li,Q., Ma,J., Li,Y., Wang,X., Li,Y., *et al.* (2024)
862  Molecular basis of SAP05-mediated ubiquitin-independent proteasomal degradation of
863  transcription factors. *Nat Commun*, **15**, 1170.

864 69. Yamasaki,K., Kigawa,T., Inoue,M., Yamasaki,T., Yabuki,T., Aoki,M., Seki,E., Matsuda,T.,
865  Tomo,Y., Terada,T., *et al.* (2006) An Arabidopsis SBP-domain fragment with a disrupted
866  C-terminal zinc-binding site retains its tertiary structure. *FEBS Lett*, **580**, 2109–2116.

867 70. Xie,Z., Sokolov,I., Osmala,M., Yue,X., Bower,G., Pett,J.P., Chen,Y., Wang,K., Cavga,A.D.,
868  Popov,A., *et al.* (2025) DNA-guided transcription factor interactions extend human gene
869  regulatory code. *Nature*, **641**, 1329–1338.

870 71. Li,M., Yao,T., Lin,W., Hinckley,W.E., Galli,M., Muchero,W., Gallavotti,A., Chen,J.-G. and
871  Huang,S.-S.C. (2023) Double DAP-seq uncovered synergistic DNA binding of interacting
872  bZIP transcription factors. *Nat Commun*, **14**, 2600.

873 72. Rodríguez-Martínez,J.A., Reinke,A.W., Bhimsaria,D., Keating,A.E. and Ansari,A.Z. (2017)
874  Combinatorial bZIP dimers display complex DNA-binding specificity landscapes. *Elife*, **6**.

875 73. Yamasaki,K., Kigawa,T., Inoue,M., Tateno,M., Yamasaki,T., Yabuki,T., Aoki,M., Seki,E.,
876  Matsuda,T., Nunokawa,E., *et al.* (2004) A novel zinc-binding motif revealed by solution
877  structures of DNA-binding domains of Arabidopsis SBP-family transcription factors. *J*
878  *Mol Biol*, **337**, 49–63.

879 74. Liang,X., Nazarenus,T.J. and Stone,J.M. (2008) Identification of a consensus DNA-binding
880  site for the Arabidopsis thaliana SBP domain transcription factor, AtSPL14, and binding
881  kinetics by surface plasmon resonance. *Biochemistry*, **47**, 3645–3653.

882 75. Garcia-Molina,A., Xing,S. and Huijser,P. (2014) Functional characterisation of Arabidopsis
883  SPL7 conserved protein domains suggests novel regulatory mechanisms in the Cu
884  deficiency response. *BMC Plant Biol.*, **14**, 231.

885 76. Li,R.-J., Li,L.-M., Liu,X.-L., Kim,J.-C., Jenks,M.A. and Lü,S. (2019) Diurnal Regulation of
886  Plant Epidermal Wax Synthesis through Antagonistic Roles of the Transcription Factors
887  SPL9 and DEWAX. *Plant Cell*, **31**, 2711–2733.

888 77. Wang,H., Studer,A.J., Zhao,Q., Meeley,R. and Doebley,J.F. (2015) Evidence That the
889  Origin of Naked Kernels During Maize Domestication Was Caused by a Single Amino

Acid Substitution in tga1. *Genetics*, **200**, 965–974.

78. Lin,P.-Y., Huang,S.-C., Chen,K.-L., Huang,Y.-C., Liao,C.-Y., Lin,G.-J., Lee,H. and Chen,P.-
Y. (2025) Analysing protein complexes in plant science: insights and limitation with
AlphaFold 3. *Botanical Studies*, **66**, 1–12.

79. Flagel,L.E. and Wendel,J.F. (2009) Gene duplication and evolutionary novelty in plants.
New Phytol., **183**, 557–564.

80. Magadum,S., Banerjee,U., Murugan,P., Gangapur,D. and Ravikesavan,R. (2013) Gene
duplication as a major force in evolution. *J. Genet.*, **92**, 155–161.

81. Rastogi,S. and Liberles,D.A. (2005) Subfunctionalization of duplicated genes as a transition
state to neofunctionalization. *BMC Evol. Biol.*, **5**, 28.

82. Ohno,S. (2013) Evolution by Gene Duplication Springer Science & Business Media.

83. Shiu,S.-H., Shih,M.-C. and Li,W.-H. (2005) Transcription factor families have much higher
expansion rates in plants than in animals. *Plant Physiol.*, **139**, 18–26.

84. Gera,T., Jonas,F., More,R. and Barkai,N. (2022) Evolution of binding preferences among
whole-genome duplicated transcription factors. *Elife*, **11**.

85. Strader,L., Weijers,D. and Wagner,D. (2022) Plant transcription factors - being in the right
place with the right company. *Curr. Opin. Plant Biol.*, **65**, 102136.

86. Galli,M., Khakhar,A., Lu,Z., Chen,Z., Sen,S., Joshi,T., Nemhauser,J.L., Schmitz,R.J. and
Gallavotti,A. (2018) The DNA binding landscape of the maize AUXIN RESPONSE
FACTOR family. *Nat. Commun.*, **9**, 4526.