Fresh Human Retinal scRNA-seq Atlas Reveals a Novel Cone Subtype and Cellular Diversity

Luning Yang[1], Yiwen Tao[1], Qi Pan[1], Tengda Cai[1], Yunyan Ye[2], Jianhui Liu[3], Yang Zhou[4], Yongqing Shao[5], Quanyong Yi[5], Zen Haut Lu[6], Lie Chen[6], Gareth McKay[7], Richard Rankin[8], Weihua Meng[1,9,10*]

Nottingham Ningbo China Beacons of Excellence Research and Innovation Institute, University of Nottingham Ningbo China, Ningbo, China, 315100

Department of Ophthalmology, Lihuili Hospital affiliated with Ningbo University, Ningbo, China, 315040

Department of Cardiology, Lihuili Hospital affiliated with Ningbo University, Ningbo, China, 315040

Ningbo Institute of Innovation for Combined Medicine and Engineering, The Affiliated Li Huili Hospital, Ningbo University, Ningbo, China, 315201

Department of Ophthalmology, The Affiliated Ningbo Eye Hospital of Wenzhou Medical University, Ningbo, China, 315040

PAPRSB Institute of Health Sciences, Universiti Brunei Darussalam, Bandar Seri, Begawan, Brunei Darussalam, BE1410

Centre for Public Health, Institute of Clinical Science, Queen's University Belfast, Block B, Royal Hospital, Grosvenor Road, Belfast, UK, BT12 6BA

School of Mathematical Sciences, University of Nottingham Ningbo China, Ningbo, China, 315100

Division of Population Health and Genomics, School of Medicine, University of Dundee, Dundee, UK, DD2 4BF

10 Center for Public Health, Faculty of Medicine, Health and Life Sciences, School of

Medicine, Dentistry and Biomedical Sciences, Queen's University Belfast, Belfast, UK, BT12

6BA

Corresponding Author: Dr Weihua Meng

Address: Division of Population Health and Genomics, Ninewells Hospital and Medical

School, University of Dundee, Dundee, UK, DD2 4BF; Nottingham Ningbo China Beacons

of Excellence Research and Innovation Institute, University of Nottingham Ningbo China,

Ningbo, China, 315100. Tel: +44 1382 383419; +86 574 88188632. Email:

w.meng@dundee.ac.uk;   weihua.meng@nottingham.edu.cn.   ORCID:

0000-0001-5388-8494

ORCID codes

Luning Yang: 0000-0001-9056-5950

Yiwen Tao: 0009-0000-0102-4205

Qi Pan: 0000-0003-0135-9122

Tengda Cai: 0000-0003-1617-506X

Yunyan Ye: 0009-0005-2325-1405

Jianhui Liu: 009-0004-1341-0831

Yang Zhou: 0000-0001-9273-0489

Yongqing Shao: 0009-0006-1319-8521

Quanyong Yi: 0000-0002-9369-3998

Zen Haut Lu: 0000-0002-9408-0579

Lie Chen: 0000-0001-9764-4299

Gareth McKay: 0000-0001-8197-6280

Richard Rankin: 0000-0001-6445-2602

Weihua Meng: 0000-0001-5388-8494

Abstract

The human retina has a remarkable diversity of cell types, which is crucial for understanding the mechanisms underpinning visual development and function. While single-cell RNA sequencing (scRNA-seq) has advanced our understanding of retinal biology, most studies have relied on postmortem or frozen samples, potentially missing important transcriptional information. The aim of this study was to create the first comprehensive scRNA-seq atlas of fresh human retinal samples from living donors and recently deceased individuals using scRNA-seq technology. A total of 106,829 cells were analyzed, which were collected from nine retinal samples using standardized scRNA-seq workflow. Our findings identified several novel subtypes of known retinal cells, including new subgroups of cones and amacrine cells (ACs), each characterized by distinct gene expression profiles. Notably, we discovered a novel cone subtype, the ELF1-Cone, which shows a clear developmental trajectory from mlCone precursors. This subtype exhibits unique functional properties and metabolic profiles, regulated by key transcription factors ELF1 and PRKAA1. We also identified five distinct AC subtypes, including three GABAergic ACs, each with unique gene expression profiles and functional characteristics. Our study highlights the critical importance of using fresh human retinal tissue for accurate cellular mapping and trajectory inference. Significant differences were observed between fresh and postmortem retinal samples in terms of pseudo time analysis such as RNA velocity. We also performed a comparative analysis of diabetic without retinopathy, diabetic retinopathy, and non-diabetic samples, suggesting diabetes-related transcriptional variation. In conclusion, we present a comprehensive human retina atlas using fresh samples that contribute to existing knowledge of retinal cell diversity, function, and transcriptional profiles. Our study is a milestone for future studies that will improve understanding of retinal biology and disease mechanisms.

73   **Keywords**: Single-cell RNA sequencing; Human retina; Transcriptomic atlas; Fresh tissue analysis; Retinal cell

74   types

Introduction

The human retina is a complex and vital component of the eye that plays a crucial role in light capture

and the initiation of the vision process [1]. As a highly specialized neural tissue lining the back of the

eye, the human retina detects and converts incoming light into electrical signals, forming the

foundation of visual perception and interpretation of our surroundings [2]. In general, the retina

consists of photoreceptors (rod and cone), signaling neurons (bipolar, horizontal, amacrine, and retinal

ganglion cells), and supporting cells (müller glial cell, microglia, astrocyte) [3]. Each cell type differs

in its morphology, functionality, and subtype composition [4]. The retina's complex structure and

function enable humans to process visual information from our environment [5]. However, many

questions remain of the structure and function of the human retina [3]. The complex composition of the

human retina also makes it vulnerable to various diseases that can significantly impact vision and

quality of life, highlighting the importance for improved understanding of the retinal structure and

function for the development of effective treatments and preventive measures [6].

Researchers have adopted advanced sequencing technologies to overcome these challenges and gain

deeper insights into retinal biology and disease pathology. Single-cell RNA sequencing (scRNA-seq)

provides in-depth analysis at the single cell level and is particularly valuable for studying retinal

diseases like age-related macular degeneration and diabetic retinopathy that progress in a heterogenous

manner [7, 8]. scRNA-seq enables efficient identification of cell type-specific expression patterns,

analysis of disease progression stages within the same sample, and exploration of cellular heterogeneity

in both physiological and pathological states. In 2019, Lukowski et al. revealed 18 distinct clusters of

retinal cells using scRNA-seq, significantly contributing to our understanding of retinal cell diversity

[9]. In 2020, Farjood et al. identified subpopulations within retinal pigment epithelium cells, including

a group with stem cell-like properties, highlighting their heterogeneity through scRNA-seq [10]. Voigt

et al. utilized scRNA-seq to find distinct glial populations in degenerating retinas, emphasizing

transcriptomic changes in retinal diseases [11]. Collectively, these studies underscore the utility of

scRNA-seq in unraveling the cellular and molecular landscapes of the human retina. In addition,

single-nucleus RNA sequencing (snRNA-seq) has gained traction in human retinal studies. By 2023,

Wang et al. used snRNA-seq to demonstrate the intricate regulatory mechanisms in retinal cells,

particularly transcription factor collaborations affecting gene regulation [12]. Most recently, Liang et al.

combined single-cell RNA and assays for transposase-accessible chromatin using sequencing to create

a comprehensive reference atlas from 2.4 million cells, integrating transcriptome and chromatin

accessibility data to unveil over 110 cell types and their regulatory networks using snRNA-seq [13].

Unlike scRNA-seq, snRNA-seq can be performed on frozen or archived retinal tissues, which is

particularly valuable where the availability of fresh human retinal tissue is limited [14]. The reliance on

frozen samples is largely due to the challenges associated with acquiring, preprocessing, and storing

human retinal tissue [12]. These challenges encompass not only technical issues but also ethical

considerations, such as the difficulty of obtaining fresh samples from living donors [15, 16].

Importantly, snRNA-seq fails to sufficiently capture RNAs localized in the cytoplasm, which are

important for understanding post-transcriptional regulation and cellular processes in the retina [17]. A

2009 study by Botling et al. examined the impact of thawing on RNA integrity and gene expression

analysis in fresh frozen tissue [18]. They found that the kinetics of RNA degradation after thawing are

likely to be tissue specific. This finding is supported by more recent research, including a 2023 review

by Thakral et al., which noted that early postmortem interval (PMI) studies on eye tissue found that the

type of tissue samples used for early and late PMI had an impact on the stability of RNA post-mortem

[19]. This leaves a gap in the field of human retinal research: the lack of atlases for comprehensive

scRNA-seq analysis of fresh human retinal samples from living donors, particularly given the fact that

fresh tissue better reflects the *in vivo* cellular transcriptional status more accurately than postmortem or

cultured samples [20].

scRNA-seq offers unique advantages for retinal studies, capturing both nuclear and cytoplasmic RNAs

and enabling the exploration of functional responses in live cells. This technique allows for the analysis

of freshly isolated cells while maintaining their viability and function, providing crucial insights into

the dynamics of cellular processes such as metabolic activity and stress responses [21]. Moreover,

scRNA-seq can track changes in cell states over time, which is essential for understanding the

progression of conditions like diabetic retinopathy [22, 23].

It is worth noting that while previous studies have created retinal atlases and some have used unfrozen

postmortem tissue samples, our study is among the first to explicitly focus on and demonstrate the

critical importance of using fresh, living human retinal tissue for creating a comprehensive atlas. By

directly comparing fresh and postmortem samples, we provide unique insights into the advantages of

using living tissue in retinal research, building upon and extending previous work in this field. We have

integrated scRNA-seq data from nine samples, encompassing samples from both living donors and

recently deceased individuals (retina were collected within several hours of death). This approach

enables a high-quality and comprehensive representation of retinal cell types and states, building upon

and extending previous eye atlases that have focused on specific ocular cell types and those derived

from frozen necropsy sources. Our atlas aims to provide a deeper understanding of retinal biology and

disease mechanisms, with potential applications in developing targeted therapies for various retinal

disorders. As some of the samples included are from donors with diabetes, we have also included a

comparative analysis between non-diabetic, diabetic without retinopathy, and diabetic retinopathy

samples to differentiate the characteristic patterns and cell type distributions.

Materials and Methods

Retinal tissues from fresh human donor eyes and cadaveric donor eyes

Human retinal tissues were collected from the Lihuili Hospital (affiliated with Ningbo University) and

the Ningbo Eye Hospital (affiliated with Wenzhou Medical University) from June 1, 2023, to May 31,

2024. Study ethics was approved separately by the ethics committees for Lihuili Hospital, the Ningbo

Eye Hospital, and the University of Nottingham, Ningbo, China. Retinal samples were collected

following written informed consent from two patients attending the Lihuili Hospital following

enucleation due to medical conditions, and seven samples were collected from the Ningbo Eye Hospital.

Of the seven samples collected from the Ningbo Eye Hospital, two pairs of eyes were collected from

recently deceased cadavers, and three from patients following enucleation for medical reasons.

Detailed information of the nine samples is shown in Table 1.

Single-cell RNA sequencing (lab preparation)

Retinal tissue was prepared for scRNA-seq using the Dissociation Kit (LK003150-1bx manufactured

by Worthington Biochemical Corporation). Tissue was dissociated, filtered, and treated with red blood

cell lysis solution (130-094-183, Miltenyi Biotec). Cell viability was assessed using CountStar, and

viable cells were enriched using the Dead Cell Removal Kit (130-090-101, Miltenyi Biotec).

The scRNA-seq libraries were prepared using a total of 10,000 cells per sample. The Chromium Single

Cell 3' Library and Gel Bead Kit V3.1 (10X Genomics, PN1000268) was used to create emulsified

single-cell gel beads. Following cell lysis, primers comprising poly-T, barcodes, unique molecular

identifiers (UMIs), and a 'read 1 primer sequence' in gel bead-in-emulsion (GEM) were used to

reverse-transcribe RNA. The addition of sample index and 'read 2 primer sequence' was performed

through adapter ligation. Following quality control, libraries were 150 bp paired-end nucleotide

sequenced using the Illumina NovaSeq platform. Reads were constructed from raw BCL files using

Illumina's bcl2 FASTQ converter. Quality control parameters included: had >3 read sequences; > 20%

of bases had a quality rating of < 5; and adapter sequence detected.

Generation of single cell data

The   10X   Genomics   Cell   Ranger   V8.0.1   pipeline

(https://support.10xgenomics.com/single-cell-gene-expression/software/pipelines/latest/what-is-cell-ra

nger) was used with default parameters to demultiplex and map raw data to the reference genome

(GRCh38-2024-A). Alignment, filtering, barcode counting, and UMI (unique molecular identifier)

counting were all done by Cell Ranger count on FASTQ files which generates feature barcode matrices

using the chromium cellular barcodes, Cell Ranger count, and gene expression algorithms.

Quality control and doublets removal

For each sample, 10X Genomics Cell Ranger created raw and filtered feature-barcode matrices. Quality

control and doublet removal were conducted separately for each dataset to remove and enrich

low-quality   and   enrich   high-quality   cells,   respectively.   DoubletFinder

(https://github.com/chris-mcginnis-ucsf/DoubletFinder) was used to eliminate doublets. Subsequently,

cells of high quality were selected using the Seurat R package (v5.1.0) [24], according to the following

quality control criteria: cells with a gene count > 200, UMI count > 400, and a mitochondrial read

count proportion < 20% were retained.

Cell clustering and dimensionality reduction

Scanpy (v1.10.2) [25] was used to analyze the scRNA-seq data. Raw counts were log-normalized, and

5,000 highly variable genes were selected. The scVI model from the scvi-tools package (v0.17.1) [26]

was trained using the raw counts of these highly variable genes for dimension reduction and batch

correction. Using the 20 resultant latent dimensions, a k-nearest neighbors (KNN) graph was created

for Leiden clustering and uniform manifold approximation and projection (UMAP) visualization.

Iterative Leiden clustering with parameters of 0.05 to 1.0 was employed to identify cellular partitions

within the dataset. In each iteration, a new KNN graph was generated, and Leiden clustering was

applied to the scVI batch-corrected latent representation. The calculation of silhouette scores for Leiden

clustering at increasing resolutions of 0.1 provided optimal clustering resolution for the scRNA-seq

data. The clustering quality for each resolution was assessed using scikit-learn, and the resolution with

the highest silhouette score was chosen to identify the most distinct and significant clusters. This was

further verified by UMAP plots and cell type specific marker gene expression to ensure the resolution

chosen provided the most well-defined and meaningful clusters, enabling the assignment of the

first-level annotation of major cell classes. Cells were then divided into smaller groups, and the steps

were repeated to finalize subcluster cell type annotations. Subclusters were identified and labeled using

known marker genes, differential expression (DE) analysis, and comprehensive enrichment analysis.

The optimal cluster resolution for each group was determined by these results and the corresponding

silhouette score.

Differential expression (DE) analysis

The $FindMarkers$ function in Seurat was utilized to identify differentially expressed genes (DEGs)

across each retinal cell class. The Wilcoxon rank-sum test was applied, considering only genes detected

in at least 25% of cells in either population, and statistical significance was determined by calculating

unadjusted and adjusted p values.

DEGs were categorized as upregulated or downregulated based on whether their average log2 fold

change (avg_log2FC) was positive or negative, respectively. Further classification was performed

based on the magnitude of the log2 fold change, with those having $|avg\_log2FC| > 1$ labeled as

'Highly' and those with $|avg\_log2FC| \leq 1$ labeled as 'Lowly'. A composite label was generated by

combining the up/down status with the high/low classification.

Similar DEG analysis was conducted for subclusters of amacrine cells (ACs), bipolar cells (BCs), and

cones, using the same parameters and methodologies. Top DEGs were selected for display by

identifying the top five upregulated and the top five downregulated genes for each cell type. This

approach ensured a robust identification and categorization of DEGs across different cell classes and

subpopulations.

Pathway enrichment analysis

Pathway enrichment analysis was conducted using multiple approaches to comprehensively assess

biological processes across different cell types and conditions. Gene Set Variation Analysis (GSVA)

was performed using the GSVA R package [27] to evaluate pathway activity, utilizing hallmark gene

sets from the molecular signatures database (MSigDB) of homo sapiens. Gene Ontology (GO)

enrichment was conducted using ClusterProfiler (https://github.com/YuLab-SMU/clusterProfiler) to

identify significantly enriched pathways for DEGs in each cluster. The "clusterProfiler" R package was

utilized to perform these analyses and visualize the results, considering p values below 0.05 as

indicative of significant pathway enrichment.

Additionally, the "irGSEA" R package (https://github.com/chuiqin/irGSEA/) was used to score

individual cells through a multi-genomic enrichment approach, generating a comprehensive

multi-genomic enrichment scoring matrix. DE analysis within this matrix was then conducted for each

cell class and cell type using the Wilcoxon test, providing a detailed evaluation of gene expression

variations.

SCENIC and cell-type specificity index analysis

The SCENIC R package was used to identify transcription factor (TF) regulons specific for each cell

type using human retinal cones. The raw count gene expression matrix of cones processed with

previous quality control steps was used as the input data. After retaining genes with at least three

counts and expressed in over one percent of total cells, we used GENIE3 to perform correlation

analysis and TF-target correlation assessment. Co-expression modules were identified and refined into

regulons based on DNA-motif analysis. Two complementary regulatory region databases were utilized:

one encompassing region 10 kb upstream and downstream of transcription start sites, and another

focusing on 500 bp upstream and 100 bp downstream, both from the hg38-RefSeq-r80 genome

annotation. Regulon activity scores were later calculated using AUCell, followed by binarization to

determine the activity state of each regulon in individual cells.

Following the SCENIC analysis, the cell-type specificity index (CSI) analysis was applied to identify

and characterize transcriptional regulatory modules. It was based on regulon activity scores derived

from the SCENIC output. CSI quantifies the similarity of regulon activity patterns across cells,

enabling the identification of co-regulated gene sets with values ranging from 0 to 1. Higher CSI values

indicate greater specificity in co-regulation. We applied a CSI threshold of 0.5 to focus on strongly

co-regulated regulons. The resulting CSI matrix was hierarchically clustered using Ward's method to

identify modules of co-regulated genes.

Dynamo analysis

We used Velocyto [28] to generate spliced and unspliced expression matrices from Cell Ranger output

files. These two matrices, along with the raw expression matrix, were imported into Dynamo [29], a

python package for RNA velocity analysis. Data preprocessing included normalization and dimension

reduction. Gene-specific velocity dynamics were calculated, and RNA velocity-derived metrics were

generated, including speed, acceleration, divergence, and ddhodge potential.

CellOracle simulation analysis

We use CellOracle [30] to perform silico pertubation analysis to simulate gene knockout of ELF1 and

PRKAA1 and corresponding effects on cell steate transitions, which are key TFs selected from gene

regulatory network analysis. Additionally, pseudotime analysis is performed to infer developmental

trajectories after the knockout. All the simulation and calculation were built based on the TF regulatory

network generated from the SCENIC output of the human retinal cone dataset.

Cell communication analysis

Cell-cell communication analysis was performed using the CellChat [31] package in R, focusing on

"Secreted Signaling" and "Cell-Cell Contact" interactions. Data preprocessing steps involved

subsetting to signaling genes, identifying overexpressed genes and interactions, and projecting data

onto a protein-protein interaction network. Communication probabilities were computed, filtered, and

aggregated into a network. We focus on the interaction activities between cone cell types: ELF1-Cone,

mlCone, and sCone.

Differential expression analysis by cell class in diabetes

EdgeR [32] was used to detect DE analysis of diabetes comparisons. Based on their diabetic conditions,

the nine retain samples were divided into three states: donors without diabetes, donors with diabetes

with no diabetic retinopathy, and donors with both diabetes and diabetic retinopathy as shown in Table

1. When performing DE analyses, individual cells were categorized into two groups. The control group

was composed of non-diabetic datasets, and the test group was composed of diabetic datasets and

diabetic retinopathy datasets. A negative binomial generalized log-linear model was used to analyze the

gene counts, taking the covariates of major cell class and diabetic status. The model was fitted using

the glmQLFit function.

Odds ratio (OR) analysis of cell type distribution in diabetic vs. non-diabetic samples

Odds ratios (OR) were computed using the methodology described in Zheng et al. [33] in order to

characterize the cell type and subtype preference distribution between the diabetic without retinopathy

group (DM), the non-diabetic group (nonDM), and the diabetic retinopathy group (DR). Using the

Benjamini-Hochberg (BH) approach, p values were modified after being determined using Fisher's

exact test. An OR > 1.5 and adjusted p value < 0.05 indicates a cell type/subtype is more likely to be

dispersed in this group, while an OR < 0.5 and adjusted p value < 0.05 indicates a less likely

distribution.

Results

Single-cell transcriptomic atlas of fresh human retina from living and cadaveric donors

We aimed to develop a comprehensive human retinal single-cell transcriptomic atlas, particularly

focusing on fresh samples of the human retina. Five samples from living individuals and two paired

samples from recently deceased individuals were collected. These samples were further categorized

based on their diabetic status into three groups: nonDM (size = 4), DM (size = 3) and DR (size = 2).

After quality control using Seruat package, a total of 106,829 cells were obtained for further analysis.

The clustering results of the nine samples with different resolution levels are shown in Figure 1, and

the resolution of 0.3 was chosen as the best value.

Based on the cell ontology, the annotation levels were split into two levels. The first level is the cell

classes, they were first annotated based on the known markers of retinal cell types reported previously

[34]. The corresponding cell clusters were assigned the identities (Figure 2A), including AC

(TFAP2B/SLC32A1), astrocyte (GFAP), cone (ARR3/GNAT2/GNGT2), rod (RHO/PDE6G/GNGT1),

müller glia cell (RLBP1), horizontal cells (LHX1/TNR), BC (NETO1), retinal ganglion cells

(SLC17A6/RBPMS), microglia (CD74/TYROBP/C1QA), and T cells (CD69/CD52/CD3D) (Figure

2B).

Most of the major cell types exhibited well-isolated, tight clusters (Figure 3A). In addition, cluster

heterogeneity is visible for some cell classes with multiple types, such as BCs. The ACs, astrocytes,

BCs, cones, horizontal cells, T cells, microglia cells, müller glia cells (MGCs), and rods were detected

in all the analyzed datasets, and retinal ganglion cells (RGCs) were detected in four donors (3, 4, 6 and

7). For all the datasets, rod photoreceptor cells were the most abundant major type, followed by MGCs

and BCs (Figure 3B and Table 2). Gene set variation analysis (GSVA) plots are used to show the terms

of the hallmark gene set enrichment pathway for each identified cell population, using the upregulated

and downregulated gene sets shown in Figure 2C. The enrichment pathways align with known

biological functions, further supporting the annotation of major retinal cell types (Figure 4). For

example, microglia show significant enrichment in inflammatory and immune-related pathways,

consistent with their role as the primary immune cells in the retina. MGC show enrichment in

glycolysis, supporting their known role in metabolic support of neurons.

In all datasets, all the markers showed quite distinct and expected distribution among the cell types. To

further accurately annotate the cell types, we performed a second round of annotation specifically for

the clusters that contained ACs, horizontal cells, RGCs and also for the clusters that contained

astrocytes and MGCs, respectively.

There were challenges in differentiating between ACs, horizontal cells, and RGCs from the scRNA-seq

retinal samples given their shared developmental trajectory and similar transcriptional profiles. These

three cell types are all derived from a common pool of retinal progenitor cells (RPCs), and data

generated from murine Atoh7-null retinas indicated genes associated with RGCs, ACs, and horizontal

cells were expressed in a highly overlapping fashion making it challenging to separate these cell types

into distinct clusters following scRNA-seq analyses [35]. Furthermore, the genesis of ACs and

horizontal cells is also controlled by Foxn4 by regulation of the expression of retinogenic factors that

are involved in their fate commitment [36]. As such, the clusters composed of ACs, horizontal cells,

and RGCs were further trained by the scVI model and Leiden clustered, which assigned their identities

based on specific markers as shown in Figure 5A-C, with corresponding dot plots and dendrogram

heatmaps (Figure 5D, E), and the final annotation results are shown in the UMAP plot (Figure 5F).

The macroglia cluster was composed of astrocytes and MGCs which provide essential support

functions in the retina and share similar transcriptional profiles [37]. Both cell types are also reactive in

response to retinal injury [38, 39]. MGCs represent the predominant glial cells in the retina and are

present in large numbers throughout the retinal layers [38], with the relative high abundance making it

harder to capture the astrocytes. In this situation, the clusters composed of astrocytes and MGCs were

further trained by the scVI model and Leiden clustered, and assigned their identities based on specific

markers with corresponding dot plots and dendrogram heatmaps (Figure 6B), and the final annotation

results are shown in the UMAP plot (Figure 6A). Our approach was able to successfully differentiate

the distinct transcriptional profiles for the various retinal cell types highlighting the intricate cellular

diversity within the human retina.

Bipolar cells

Retinal bipolar cells receive synaptic input from photoreceptors and process it in diverse ways before

transmitting an input to RGCs. They are essential components of the human retinal circuitry, serving as

intermediaries between photoreceptors and RGCs. Their diverse subtypes and molecular characteristics

allow for complex processing of visual information, including light adaptation, temporal frequency

processing, and contrast detection. Understanding bipolar cell function is crucial for comprehending

retinal development, visual processing, and the pathogenesis of certain retinal diseases.

BCs can be subdivided on the basis of two main distinctions: rod versus cone BCs and ON versus OFF

BCs, as shown in Figure 7A. The former is associated with synaptic input, while the latter distinguishes

the signaling mechanism in dendrites and the level of axonal stratification [40].

The ON BC subtypes, such as DB5, DB6, BB+GB, and IMB, respond to light onset and transmit

signals via mGluR6 receptors [41]. OFF BC subtypes, including DB1, DB2, DB3a, DB3b, FMC, and

OFFx, hyperpolarize in response to light offset, utilizing ionotropic glutamate receptors (AMPA and

Kainate) [42]. RBCs are crucial for scotopic (low-light) vision, transmitting signals from rod

photoreceptors primarily to AII ACs, which then relay the information to cone BC cells and ultimately

to RGCs [43]. This classification highlights the diverse roles and pathways of BCs in visual processing.

DB4 cells are mostly classified as ON BC in the retina, however, some studies have ambiguously

categorized DB4 as OFF BC [44]. To further characterize this group, we identified a high-level

expression of GRM6 in DB4 clusters in our study, a key component of the ON BC signaling pathway.

mGluR6 detects a decrease in glutamate, which then initiates a signaling cascade that depolarizes ON

BCs. On that basis, we confirmed the attribution of this DB4 cluster to ON BCs.

To delineate the landscape of BCs, we isolated BCs and performed scVI model training and Leiden

clustering based on suitable resolution levels. BCs were divided into subtypes previously characterized

by common marker genes for each BC subpopulation across all the datasets (Figure 7B, C). A total of 6

ON BC and 5 OFF BC subclusters were identified in the human datasets (Figure 7D).

Amacrine cells

Retinal amacrine cells play a complex role in visual signal processing, forming synapses with multiple

cell types, including BCs, RGCs, and other ACs, providing both feedback inhibition to BCs and

feedforward inhibition to RGCs, shaping the visual signal [45, 46]. There are numerous types of ACs,

each with potentially different patterns of connectivity and roles in signal processing, that use various

neurotransmitters, including GABA.

Most ACs are inhibitory neurons utilizing GABA or glycine as neurotransmitters [34]. After retraining

the clustering model by assessing the expression of marker genes for GABAergic (GAD1 and GAD2)

and glycinergic (SLC6A9) ACs, we identified three clusters with high GABA marker expression, with

high DEGs MEIS2/BASP1, SOX5/PCP4, and CCDC39/RBFOX1 (Figure 8A-B). We inferred they are

GABA ACs with distinct characteristics and functions. One cluster was represented by high SLC6A9

expression, annotated as Glycinergic ACs (GlyACs). Another cluster was represented by mid-levels of

expression for all the three genes, which may correspond to a mixture of both GABA and glycine AC

types with a role as intermediate ACs.

To further verify the function of the three different GABAergic ACs, we employed integrative irGSEA

to score each cluster using a multi-genomic enrichment approach. We performed GO enrichment

analyses for each cluster (Figure 8C-D). Except the GlyACs and intermediate ACs, the results of

pathway analysis of these three different GABAergic ACs demonstrate that they have distinctly

different functions. GABA-AC1 displays a unique focus on postsynaptic specialization and

photoreceptor interaction. As indicated by the pathways "glutamatergic synapses" and "synaptic

membranes", these cells are enriched in pathways indicative of complex and diverse synaptic

connections, including inhibitory (GABAergic) and potentially excitatory (glutamatergic) synapses.

They play a crucial role in fine-tuning synaptic signaling and neurotransmitter release, as highlighted

by the pathways "regulation of trans-synaptic signaling" and "modulation of chemical synaptic

transmission". Structurally, they are characterized by "distal axons" and "growth cones", suggesting

that they are specifically adapted to synaptic interactions. GABA-AC1 are the highest involvement in

developmental processes in GABA-ACs. GABA-AC2 enrichment pathways show focus on

postsynaptic specialization and photoreceptor interaction. Pathways include postsynaptic density and

postsynaptic specialization, suggesting a role in fine-tuning synaptic inputs. The cluster is also closely

associated with photoreceptor outer segments and various types of cilia and is involved in eye

development and visual system development. GABA-AC3 demonstrate a pronounced specialization in

ion channel activities and calcium signaling. They are enriched in various ion channel activities,

particularly voltage-gated and monoatomic ion channels. The strong emphasis on calcium ion transport

and regulation of membrane potential suggesting a key role in rapid signal modulation. This subtype

also shows significant involvement in synapse organization and regulation of trans-synaptic signaling,

indicating a role in shaping synaptic communication. In summary, the novel GABA AC subpopulations

identified in this study may represent specific modulators characterized by specialized roles in retinal

function.

Cone photoreceptor cells

We identified four clusters of cones that correspond to distinct cell states, referred to as mlCone, sCone,

ELF1-Cone and a cluster associated with corneal repair. Cones are classified based on their sensitivity

to different wavelengths of light, which enables color perception. The three types of cones are sCones,

which are sensitive to short wavelengths corresponding to blue light; mCones, which respond to

medium wavelengths corresponding to green light; and lCones, which are sensitive to long

wavelengths corresponding to red light. [47].

Traditionally, cone cells have been classified as mlCones (medium/long-wavelength) and sCones

(short-wavelength) due to the difficulty in distinguishing mCone and lCone[34]. We retrained the

clustering model with annotation based on the expression patterns of key markers (Figure 9A). The

mlCones and sCones were comprehensively identified based on high expression levels of OPN1LW,

OPN1MW, and OPN1SW (Figure 9B). mlCones exhibit lower ARR3 expression compared to sCone.

Additionally, mlCones were found to consist of two distinct clusters, differentiated by the expression of

MGST3/GNB3 and KCNIP4/CNGA1.

We also identified a distinct cell cluster characterized by low ARR3 expression and high levels of

KCNB2, and EYS (Figure 9C). We named this cluster ELF1-Cone. To determine the identity and

function of ELF1-Cone, we performed pseudo-bulk differential expression analysis for each cluster

including mlCone, sCone, ELF1-Cone and the corneal repair cluster.

Corneal repair cluster in cone photoreceptor

The corneal repair cluster was predominantly present in corneal ulcer sample (80.74%), exhibiting

upregulation of genes critical for protein synthesis and cellular stress response. The top ten DEGs in the

cluster including RPLP0, RPL13A, RPL6, and EEF1G (Figure 10A). RPLP0 is an important

component of the ribosomal stalk, its high expression suggests an increased capacity for protein

synthesis required for cellular repairments. This upregulation may indicate that cells in this cluster

actively respond to injury by producing proteins necessary for corneal repair and regeneration.

RPL13A functions not only as a ribosomal protein but also in the interferon-γ-activated inhibitor of

translation (GAIT) complex, which regulates the selective translation of certain mRNAs during

inflammation. Up-regulation of RPL13A may reflect a dual role in both general protein synthesis

supporting the repair process in this cluster and regulating specific inflammatory responses in corneal

cells. Upregulation of EEF1G, part of the elongation factor-1 complex involved in the transport of

aminoacyl tRNAs to ribose, suggests enhanced protein synthesis under stress conditions and activation

of cell survival pathways in response to corneal injury. Increased expression of RPL4 and RPL6,

important components of the 60S ribosomal subunit, further supports the hypothesis that these cells

have an increased capacity for ribosome biogenesis and protein synthesis. Notably, this cluster may

also have been affected by contamination of corneal ulcer from donor 5, reflect the introduction of

non-retinal cells from damaged corneal tissue.

ELF1-Cone is discovered as a new subtype of cone photoreceptor

The top ten up-regulatory DEGs of ELF1-Cone including ASIC2, MSR1, and FRMD4A (Figure 10A).

ASIC2 is an ion channel activated by extracellular acidification that may contribute to neuroprotective

mechanisms and pH sensing in the retinal microenvironment. MSR1 is commonly associated with

immune function and is capable to clear cellular debris. It plays a role in the removal of oxidized lipids

and cellular debris, contributing to tissue homeostasis. FRMD4A is involved in cytoskeletal

organization and cellular polarity and may potentially contribute to efficient signaling of this cluster.

We propose a functional hypothesis for this unknown cone cluster, where cone cells have a unique

adaptive capacity to sense and respond to changes in the retinal microenvironment, maintain

specialized cellular structures critical to their function, and provide neuroprotection against oxidative

stress and other cellular threats.

To further validate the functions of the ELF1-Cone cluster, we performed GSVA and GO analyses as

shown in Figure 10B-C. There were significant enrichments of energy metabolic processes, including

oxidative phosphorylation, aerobic respiration and cellular respiration. It was also supported by the

enrichment terms of mitochondrial components and energy-related processes. The enrichment results

supported that ELF1-Cone cells have a high capacity for energy production, which is also essential for

maintaining their specialized function in the retina. The enrichment of cytoplasmic translational and

ribosomal components protein synthesis pathways suggesting an active protein production machinery.

This allows ELF1-Cone cells to respond rapidly to changing conditions or stressors in the retinal

microenvironment. GSVA heatmaps show high activation of the *hallmark_apical_surface* and

hallmark_apical_junction pathways in ELF1-Cone cells, suggesting a specific role in cell-to-cell

interactions and maintenance of retinal retina, indicating a specific role in intercellular interactions and

maintenance of retinal organization. Activation of *hallmark_hedgehog_signaling* further implies

involvement in cell differentiation and organization patterns. GO analysis also revealed an enrichment

of transmembrane transporter protein activity and various binding processes, suggesting enhanced cell

signaling and metabolite exchange. The presence of adhesion spots and cell-matrix junction terms is

consistent with GSVA results, emphasizing the importance of cell-matrix interactions in ELF1-Cone

function.

ELF1 and PRKAA1 are key TFs in ELF1-Cone

On this basis, for further experimentation and validation, we performed SCENIC analysis to profile cell

state specificity and TF activity (Figure 10F). The top 40 regulons are listed in supplementary Table S1.

ELF1 is a transcription factor involved in the regulation of T-cell specific genes. SETDB1 is a histone

methyltransferase that specifically trimethylates lysine 9 of histone H3 and is also associated with

retinal development and photoreceptor maintenance. RCOR1 is a transcriptional co-repressor involved

in neuronal gene silencing and plays a role in retinal progenitor cell differentiation. SREBF1 is a TF

that regulates genes involved in lipid metabolism. The presence of genes such as RCOR1, SETDB1,

and SREBF1 in the ELF1-Cone cluster suggests unique metabolic and developmental features and gene

regulatory networks compared to other optic cone cell types. Based on the list of genes, we could infer

the TFs involved in the following processes: transcriptional regulation (ELF1, ETV3, STAT5B, TBP)

chromatin modification (SETDB1, PHF8, RBBP5) metabolic processes (SREBF1, PRKAA1) DNA

repair (XRCC4). These findings are consistent with GO analysis, particularly highlighting the

enrichment of pathways regulated by TFs and those involved in metabolic processes.

At the same time, we use the connection specificity index (CSI) to perform regulon module analysis.

The results identify four distinct groups of regulons in cones, ordered from module one to module four.

Key TFs of ELF1-Cone were labeled and observed to aggregated in module two and module four,

corresponding to two different characteristic regulon groups (Figure 10D-E). Module two was enriched

with TFs involved in chromatin modification (SETDB1, PHF8, RBBP5), transcriptional regulation

(ELF1, ETV3, STAT5B), and cell cycle control (E2F6). Module four was associated with metabolic

regulation (SREBF1, PRKAA1), DNA repair (XRCC4), and general TFs (TBP, TAF1). This indicates

the function of ELF1-Cone mainly including two aspects. The ELF1-Cone key TFs in module two

supported its role in epigenetic regulation and cell fate determination, which could be crucial for

maintaining self-identity. The ELF1-Cone key TFs in module four supported its involvement in

metabolic processes and cellular stress response, potentially reflecting the high energy demands and

light-induced stress in cone cells. While further experimental validation is needed, this transcriptional

profile indicates that ELF1-Cone cells could play specialized roles in retinal function, potentially

including rapid cellular adaptation to changing light conditions and enhanced neuroprotective

capabilities.

Cell Trajectory inference of ELF1-Cone

Dynamo is applied to study cellular dynamics, gene regulatory networks of ELF1-Cones, and revealed

a potential developmental trajectory from mlCone to ELF1-Cone. This method builds on and extends

the concept of RNA velocity, utilizing the ratio of unspliced to spliced mRNAs to predict the future

state of individual cells. We divided our nine datasets into three groups to perform Dynamo analysis:

all samples combined (including both fresh and cadaveric), fresh samples from living donors only, and

cadaveric samples without freezing. The analysis was visualized using consistent UMAP reduction to

reveal distinct cellular dynamics across conditions. As shown in Figure 11A, the all sample shows

distinct clusters for CR, mlCone, ELF1-Cone, and sCone, with clear transitions between them,

particularly from mlCone to ELF1-Cone. In the fresh living sample condition, we observe a more

pronounced and directional flow from mlCone to ELF1-Cone. The transitions appear clearer and more

defined, suggesting that in fresh samples, the differentiation pathway from mlCone to ELF1-Cone is

more evident and potentially more active. The velocity flow is particularly evident at the boundary

junction, highlighting that this differentiation pathway is strong and direct in the fresh sample.

However, in the condition of cadaveric samples, the flow patterns become more chaotic and less

directional. The transition from mlCone to ELF1-Cone is less clear, and the overall organization of the

clusters appears more diffuse. This suggests that post-mortem changes may disrupt the normal

differentiation processes and cellular state.

The cellular dynamics analysis in Figure 11B provides further insights into the state of ELF1-Cone in

fresh living datasets. The ELF1-Cone region is of moderate to high-speed indicating active transitions.

The low divergence also suggesting that cells are converging to a specific state rather than branching

into different fates. The low to moderate acceleration and curvature in the ELF1-Cone region suggest

ongoing changes in cell state, but with relatively simple developmental trajectories.

To further verify the function of ELF1-Cones, we performed CellOracle to simulate the effects of

knocking out specific TFs on the developmental trajectories of cones. ELF1 is the key TF for

ELF1-Cone with the highest residual sum of squares (RSS) rank in regulon list, plays a significant role

in the transcriptional program. PRKAA1, while lower in the rankings, could be crucial for regulating

the distinct metabolic profile observed in ELF1-Cones, given its known role in energy homeostasis and

the enrichment of metabolic processes in these cells. We used CellOracle to elucidate the exact

functions of these factors in ELF1-Cone cells and their potential contributions to the unique

characteristics of this cell type.

As shown in Figure 11C, there were significant changes in the flow patterns: the original cones from

the living fresh samples showed clear directional flows, particularly from the mlCone towards the

ELF1-Cone. The flows were well-organized and consistent, indicating a stable developmental trajectory.

After knocking out of ELF1, the directional flow from mlCone to ELF1-Cone was disrupted and there

was an increase in turbulent flows, particularly in the central regions. Some areas showed reversed

flows compared to the original dataset. It indicates that ELF1 plays a crucial role in maintaining the

normal differentiation pathway from mlCone to ELF1-Cone. Its absence leads to a loss of directional

development and potentially affects cell fate decisions. The perturbation simulations of PRKAA1

knockdown showed even more dramatic changes. There's a significant increase in the magnitude of

flows across the entire plot. The flow patterns become highly chaotic, with many conflicting directions.

The organized structure seen in the fresh datasets is almost completely lost. This implies that PRKAA1

has a broader influence on cellular dynamics, possibly affecting multiple aspects of cell metabolism

and signaling. Its knockout appears to destabilize the entire system, leading to unpredictable cellular

behaviors. ELF1 knockout primarily affects the mlCone to ELF1-Cone transition, while PRKAA1

knockout has a more global effect on cellular dynamics.

Cell to cell interaction inside cones

CellChat analysis revealed complex interactions between ELF1-Cones, mlCones and sCones as shown

in Figure 12. Key interactions include NEGR1, CADM1 and CDH2. NEGR1 is known to be involved

in cell adhesion and neurite outgrowth. CADM1 and CDH 2 are important for synaptic organization

and intercellular adhesion, and is often involved in neural organization. It indicates a critical role in

maintaining the physical organization between ELF-Cones and mlCones, and in maintaining specific

synaptic connections between these subtypes. Suggesting ELF1-Cones are in a state of active

communication while maintaining a relatively stable identity.

Horizontal cells

The human retina predominantly contains two horizontal cell subtypes, H1 and H2 [48]. In our study,

we identified both horizontal cell types based on the high expression levels of LHX1 and ISL1,

respectively (Figure 13).

Diabetic and non-diabetic comparative analysis

The annotated cell type proportions for each sample are presented in Table 2. Significant variability

was observed across samples, with donors 01, 06, and 07 showing notably atypical cellular

compositions. Donor 01 (76-year-old diabetic male, corneal staphyloma) was characterized with high

levels of microglia (22.55%) and low rods (42.76%), suggesting diabetes-related inflammation. Donor

06 (50-year-old non-diabetic female, eyeball atrophy) showed elevated BCs (23.24%) and microglia

(13.98%), indicating possible selective cell survival. Donor 07 (68-year-old non-diabetic female,

corneal perforation) had the highest MGCs (26.21%) and lowest rods (40.15%), suggesting reactive

gliosis. These three samples were excluded from subsequent comparative analyses between diabetic

retinopathy (DR), diabetes mellitus (DM), and non-diabetic (nonDM) groups to investigate

disease-specific retinal cell population variation. Figure 14A illustrates the cell proportion distributions

for the three groups using boxplots.

To explore the differences in single-cell RNA profile patterns of major cell types and subtypes in those

with diabetes and diabetic retinopathy, we applied a comprehensive compositional analysis. This

included DE analysis using a negative binomial generalized log-linear model, odds ratio (OR) analysis

to identify cell types or subtypes more likely to be represented in diabetic samples, and Augur

sensitivity analysis to determine cell types or subtypes highly indicative of someone with diabetes.

As illustrated in Figure 15, the DE analysis between control and test groups identified upregulation of

genes such as FCGBP (FDR = 1.12E-09), CD74 (FDR = 2.71E-06), HLA-B (FDR = 7.12E-05),

HLA-E (FDR = 1.08E-04), HMOX2 (FDR = 7.01E-04), G6PD (FDR = 1.65E-03), and COL1A2 (FDR

= 8.38E-04). Downregulation of genes such as HSPA1B (FDR = 4.16E-12), HSPH1 (FDR = 1.03E-08),

SLC6A6 (FDR = 8.27E-04), and DDIT3 (FDR = 8.30E-04) were detected.

The OR analysis heatmap (Figure 14B) presents retinal cell type proportions across three conditions:

DM, DR, and nonDM. Key findings include notably reduced proportions of ACs, horizontal cells, and

RGCs in DM but elevated in DR. Cones show higher proportions in nonDM subjects, while rods and

MGCs remain relatively stable across each group. Astrocytes exhibit subtle changes, with slightly

lower proportions in DM and higher in DR.

Discussion

Our study presents the first comprehensive single-cell transcriptomic atlas explicitly focusing on and

demonstrating the critical importance of using fresh, living human retinal tissue. Using living samples

provides a more accurate representation of *in vivo* transcriptional states compared to previous studies

reliant on frozen or preserved tissue [13]. This approach has allowed us to capture subtle transcriptional

changes and identify novel cell states that may have been overlooked in earlier studies using preserved

tissues.

A key finding of our study is the discovery of a new cone photoreceptor subtype, ELF1-Cone,

representing a significant advancement in our understanding of retinal cellular diversity. ELF1-Cones

exhibit a unique transcriptional profile characterized by the expression of genes such as ASIC2, MSR1,

and FRMD4A, suggesting multifaceted roles in pH sensing, neuroprotection, and cellular organization.

Gene set enrichment analyses reveal that ELF1-Cones possess high energy production capacity, active

protein synthesis machinery, and enhanced cell signaling capabilities, indicating their potential for

rapid responses to changing retinal conditions.

The identification of ELF1 and PRKAA1 as key transcription factors provides insight into the

regulatory mechanisms underlying ELF1-Cones' unique properties. The interplay between these

transcription factors may coordinate the balance between energy production, environmental sensing,

and neuroprotection that characterizes ELF1-Cones. Our dynamo analysis revealed a clear

developmental trajectory from mlCone to ELF1-Cone, particularly evident in living samples. This

trajectory suggests that ELF1-Cones represent a distinct stage in cone photoreceptor development. The

cellular dynamics of ELF1-Cones, characterized by lower divergence and curvature, accompanied with

moderate velocity and acceleration, indicate a more stable state compared to mlCone precursors.

However, it still undergoes some degree of maturation or functional specialization.

CellOracle simulations further demonstrated the key roles of ELF1 and PRKAA1 in maintaining

ELF1-Cone identity and guiding the developmental trajectory. These findings are consistent with

previous studies showing that photoreceptors undergo a terminal differentiation phase where they

acquire mature characteristics. In this process, different TFs are involved in the early formation and late

function and maintenance of photoreceptors [49]. The differentiation of ELF1-Cone cells involves

complex interactions between transcriptional regulation and metabolic remodeling. Our analyses

suggest that ELF1 coordinates specific transcriptional programs critical for ELF1-Cone identity, while

PRKAA1 may mediate important metabolic changes. This dual regulatory mechanism is consistent

with recent findings on bidirectional interactions between transcriptional control and metabolic

regulation in neuronal differentiation [50]. Importantly, our cell interaction analysis reveals that

ELF1-Cone cells are not isolated entities but are integrated into a complex network of retinal cells.

Our study underscores the importance of using living human retinal tissue for accurate cellular

mapping and trajectory inference. Living samples showed a clear, directional flow from mlCone to

ELF1-Cone, indicating an active differentiation pathway. In contrast, postmortem samples exhibited a

disorganized, less directional pattern. The significant differences observed between living and

postmortem samples in Dynamo analyses suggest that postmortem changes could affect the cellular

patterns of the retina, potentially leading to misinterpretation of functional relationships. Postmortem

changes, including ATP depletion, ion concentration shifts, enzyme activation, and RNA degradation,

all have the possibility to affect cell state, gene expression, and observed developmental trajectories.

While previous studies used adult retinal specimens removed of death and dissected within few hours

[51], our results indicate that even these relatively short postmortem intervals can lead to significant

alterations in cell trajectory. The disruptions observed in postmortem samples may obscure important

nuances, particularly when studying subtle cellular transitions or newly identified cell types like

ELF1-Cone. The sensitivity of ELF1-Cone cells to tissue degradation, evidenced by the loss of distinct

differentiation flows in postmortem samples, suggests that these cells require active cellular processes

to maintain their identity. This sensitivity could potentially make ELF1-Cones early indicators of

retinal stress or disease.

We also identified three distinct GABAergic AC clusters (GABA-AC1, GABA-AC2, and

GABA-AC3), alongside glycinergic ACs and an intermediate population, each with specialized

functions. These distinct functional profiles advance our understanding of how different AC subtypes

contribute to specific aspects of visual processing and retinal function.

Furthermore, the discovery of a potential corneal ulcer-associated cluster highlights the importance of

considering the potential confounding effect from the underlying pathological conditions and the

interconnectedness of both ocular and systemic health when considering the implications for gene

expression and implicated disease pathways [52, 53].

In this study, we also addressed the challenges in differentiating between ACs, horizontal cells, and

RGCs in scRNA-seq retinal samples, which commonly arise given their shared developmental

trajectory and similar transcriptional profiles. All three cell types are derived from a common pool of

retinal progenitor cells, and previous research in murine models has shown that genes associated with

RGCs, ACs, and horizontal cells are expressed in a highly overlapping fashion in Atoh7-null retinas.

Our study addresses these challenges through careful analysis and annotation, providing a more

nuanced understanding of these closely related cell types.

Comparative analysis between nonDM, DM, and DR groups revealed significant differences in

single-cell RNA profile patterns across cell types after excluding samples with markedly abnormal cell

composition (with rod proportion less than 55%). Importantly, the differences in retinal cell type

composition detected did not differ significantly between DM and nonDM samples. This suggests that

early retinal neurodegeneration commonly associated with diabetes, damage to blood vessels, and

structural changes are somewhat limited in the absence of DR. This finding is consistent with the

understanding that detection of early diabetic lesions often requires advanced imaging techniques [54,

55]. Another noteworthy finding included the detection of a significant increase in the proportion of

RGCs in the DR samples, particularly in Donor 4, contrary to the typical decrease expected in

association with DR. We hypothesize this unexpected increase may be due to DR-induced stress and

activation of RGCs, which alters their gene expression profiles and improves their detectability in

scRNA-seq [56]. The difference in the increase in the proportion of RGCs in the DR samples may

represent variation in the rate of progression among patients with diabetic retinopathy.

We also find the upregulation of inflammation-related genes (CD74, BTN3A2, HLA-B, HLA-E),

complex alterations in stress response mechanisms (downregulation of HSPA1B, HSPH1; upregulation

of HMOX2), and changes in neurotransmitter signaling and metabolic pathways. These results support

the growing recognition of DR as an inflammatory disease and suggest potential targets for therapeutic

interventions [57]. While these findings provide valuable insights into DR pathogenesis, the primary

objective of this investigation was to generate new human retina atlases, and an ongoing collection of

more diabetic samples is underway for future analysis.

Our study encompasses a diverse range of samples, including those from diabetic and non-diabetic

individuals, and both ante-mortem and post-mortem fresh samples. This diversity provided high-quality

reference datasets for future research.

Despite the comprehensive nature of our study, we acknowledge some limitations including the use of

samples from different disease status due to the challenges in obtaining living human retinal samples.

This constraint highlights the need for collaborative efforts in the field to overcome the challenges of

sample acquisition, an issue previously highlighted [58]. Moreover, the clinical reasons necessitating

eye enucleation as outlined in Table 1, which includes conditions such as corneal staphyloma, ocular

atrophy, corneal ulceration, and corneal perforation may potentially influence cell type distribution and

gene expression profiles, representing an additional limitation of this study.

In conclusion, our study provides the atlas of fresh human retinal samples, significantly advancing our

understanding of retinal cell diversity and function. This atlas serves as a valuable resource for the

research community, paving the way for new discoveries in retinal biology and potential therapeutic

interventions for retinal diseases, particularly diabetic retinopathy. Future research should elucidate

ELF1-Cone functions, explore their role in retinal pathology, and leverage this atlas for potential

therapeutic interventions.

Statements and Declarations

Data availability

Access to the scRNA-seq data generated is possible upon reasonable request.

Author information

Authors and Affiliations

Nottingham Ningbo China Beacons of Excellence Research and Innovation Institute, University of Nottingham Ningbo China, Ningbo, China, 315100

Luning Yang, Yiwen Ta, Qi Pan, Tengda Cai, Weihua Meng

Department of Ophthalmology, Lihuili Hospital affiliated with Ningbo University, Ningbo, China, 315040

Yunyan Ye

Department of Cardiology, Lihuili Hospital affiliated with Ningbo University, Ningbo, China, 315040

Jianhui Liu

Ningbo Institute of Innovation for Combined Medicine and Engineering, The Affiliated Li Huili Hospital, Ningbo University, Ningbo, China, 315201

Yang Zhou

Department of Ophthalmology, The Affiliated Ningbo Eye Hospital of Wenzhou Medical University, Ningbo, China, 315040

Yongqing Shao

PAPRSB Institute of Health Sciences, Universiti Brunei Darussalam, Bandar Seri, Begawan, BE1410, Brunei Darussalam

Zen Haut Lu, Lie Chen

Centre for Public Health, Institute of Clinical Science, Queen's University Belfast, Block B, Royal Hospital, Grosvenor Road, Belfast, Northern Ireland, BT12 6BA

Gareth McKay

School of Mathematical Sciences, University of Nottingham Ningbo China, Ningbo, China, 315100

Richard Rankin

Authors' Contributions

LY drafted the paper and performed the analysis. YT, QP, TC contributed to data formatting and correction. YY, JL, YZ, YS, QY, ZL, LC, GM, and RR provided comments on the paper. WM organized the project and provided comments.

Acknowledgment

This study was mainly funded by the Pioneer and Leading Goose R&D Program of Zhejiang Province 2023 with reference number 2023C04049 and Ningbo International Collaboration Program 2023 with reference number 2023H025.

Corresponding authors

Correspondence to Weihua Meng

Consent to Publish

All authors have consent for publication.

Conflict of Interest

The authors have no conflicts of interest to disclose.

Ethics approval

This study was approved by the Ethics Committee of the University of Nottingham Ningbo China; The Lihuili Hospital affiliated with Ningbo University; The Affiliated Ningbo Eye Hospital of Wenzhou Medical University.

References

1.   Masland RH: **The neuronal organization of the retina**. *Neuron* 2012, **76**(2):266-280.

2.   Hussey KA, Hadyniak SE, Johnston Jr RJ: **Patterning and development of**

photoreceptors in the human retina. *Frontiers in cell and developmental biology* 2022,

10:878350.

3.   Grünert U, Martin PR: **Cell types and cell circuits in human and non-human primate**

retina. *Progress in retinal and eye research* 2020, **78**:100844.

4.   Sanes JR, Masland RH: **The types of retinal ganglion cells: current status and**

implications for neuronal classification. *Annual review of neuroscience* 2015,

38(1):221-246.

5.   Lagnado L, Schmitz F: **Ribbon synapses and visual processing in the retina**. *Annual*

review of vision science 2015, **1**(1):235-262.

6.   Cen LP, Park KK, So KF: **Optic nerve diseases and regeneration: How far are we from**

the promised land? *Clinical & experimental ophthalmology* 2023, **51**(6):627-641.

7.   Voigt AP, Mulfaul K, Mullin NK, Flamme-Wiese MJ, Giacalone JC, Stone EM, Tucker

BA, Scheetz TE, Mullins RF: **Single-cell transcriptomics of the human retinal pigment**

epithelium and choroid in health and macular degeneration. *Proceedings of the*

National Academy of Sciences 2019, **116**(48):24100-24107.

8.   Chen K, Wang Y, Huang Y, Liu X, Tian X, Yang Y, Dong A: **Cross-species scRNA-seq**

reveals the cellular landscape of retina and early alterations in type 2 diabetes mice.

Genomics 2023, **115**(4):110644.

9.   Lukowski SW, Lo CY, Sharov AA, Nguyen Q, Fang L, Hung SS, Zhu L, Zhang T, Grü

nert U, Nguyen T: **A single-cell transcriptome atlas of the adult human retina.** *The*

EMBO journal 2019, **38**(18):e100811.

10.   Farjood F, Charniga C, Kiehl T, Stern J, Boles N: **Single-cell RNA sequencing reveals**

the heterogeneity of the human retinal pigment epithelium. *Investigative*

Ophthalmology & Visual Science 2020, **61**(7):3182-3182.

11.   Voigt AP, Binkley E, Flamme-Wiese MJ, Zeng S, DeLuca AP, Scheetz TE, Tucker BA,

Mullins RF, Stone EM: **Single-cell RNA sequencing in human retinal degeneration**

reveals distinct glial cell populations. *Cells* 2020, **9**(2):438.

12.   Wang J, Cheng X, Liang Q, Owen LA, Lu J, Zheng Y, Wang M, Chen S, DeAngelis

MM, Li Y: **Single-cell multiomics of the human retina reveals hierarchical transcription**

factor collaboration in mediating cell type-specific effects of genetic variants on gene

regulation. *Genome biology* 2023, **24**(1):269.

13.   Liang Q, Cheng X, Wang J, Owen L, Shakoor A, Lillvis JL, Zhang C, Farkas M, Kim IK,

Li Y: **A multi-omics atlas of the human retina at single-cell resolution.** *Cell genomics*

2023, **3**(6).

14.   Bakken TE, Hodge RD, Miller JA, Yao Z, Nguyen TN, Aevermann B, Barkan E,

Bertagnolli D, Casper T, Dee N: **Single-nucleus and single-cell transcriptomes**

compared in matched cortical cell types. *PloS one* 2018, **13**(12):e0209648.

15.   Kropp M, Mohit M, Leroy-Ciocanea CI, Schwerm L, Harmening N, Bascuas T, De

Clerck E, Kreis AJ, Pajic B, Johnen S: **Mammalian Animal and Human Retinal Organ**

Culture as Pre-Clinical Model to Evaluate Oxidative Stress and Antioxidant Intraocular

Therapeutics. *Antioxidants* 2023, **12**(6):1211.

16.   Kapp MB: **Ethical and legal issues in research involving human subjects: do you want**

a piece of me? *Journal of clinical pathology* 2006, **59**(4):335-339.

17.   Lacar B, Linker SB, Jaeger BN, Krishnaswami SR, Barron JJ, Kelder MJ, Parylak SL,

Paquola AC, Venepally P, Novotny M: **Nuclear RNA-seq of single neurons reveals**

molecular signatures of activation. *Nature communications* 2016, **7**(1):11022.

18.   Botling J, Edlund K, Segersten U, Tahmasebpoor S, Engström M, Sundström M,

Malmström P-U, Micke P: **Impact of thawing on RNA integrity and gene expression**

analysis in fresh frozen tissue. *Diagnostic Molecular Pathology* 2009, **18**(1):44-52.

19.   Thakral S, Purohit P, Mishra R, Gupta V, Setia P: **The impact of RNA stability and**

degradation in different tissues to the determination of post-mortem interval: A

systematic review. *Forensic Science International* 2023:111772.

20.   You M, Rong R, Zeng Z, Li H, Xia X, Ji D: **Single-cell RNA sequencing: A new**

opportunity for retinal research. *Wiley Interdisciplinary Reviews: RNA* 2021,

12(5):e1652.

21.   Weir K, Leavey P, Santiago C, Blackshaw S: **Multiplexed analysis of retinal gene**

expression and chromatin accessibility using scrna-seq and scatac-seq. *JoVE*

(Journal of Visualized Experiments) 2021(169):e62239.

22.   Wang Y, Yang X, Zhang Y, Hong L, Xie Z, Jiang W, Chen L, Xiong K, Yang S, Lin M:

Single-cell RNA sequencing reveals roles of unique retinal microglia types in early

diabetic retinopathy. *Diabetology & Metabolic Syndrome* 2024, **16**(1):49.

23.   Hu Z, Mao X, Chen M, Wu X, Zhu T, Liu Y, Zhang Z, Fan W, Xie P, Yuan S: **Single-cell**

transcriptomics reveals novel role of microglia in fibrovascular membrane of

proliferative diabetic retinopathy. *Diabetes* 2022, **71**(4):762-773.

24.   Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A, Lee MJ, Wilk AJ,

Darby C, Zager M: **Integrated analysis of multimodal single-cell data**. *Cell* 2021,

184(13):3573-3587. e3529.

25.   Wolf FA, Angerer P, Theis FJ: **SCANPY: large-scale single-cell gene expression data**

analysis. *Genome biology* 2018, **19**:1-5.

26.   Gayoso A, Lopez R, Xing G, Boyeau P, Valiollah Pour Amiri V, Hong J, Wu K,

Jayasuriya M, Mehlman E, Langevin M: **A Python library for probabilistic analysis of**

single-cell omics data. *Nature biotechnology* 2022, **40**(2):163-166.

27.   Hänzelmann S, Castelo R, Guinney J: **GSVA: gene set variation analysis for**

microarray and RNA-seq data. *BMC bioinformatics* 2013, **14**:1-15.

28.   La Manno G, Soldatov R, Zeisel A, Braun E, Hochgerner H, Petukhov V, Lidschreiber

K, Kastriti ME, Lönnerberg P, Furlan A: **RNA velocity of single cells**. *Nature* 2018,

560(7719):494-498.

29.   Qiu X, Zhang Y, Martin-Rufino JD, Weng C, Hosseinzadeh S, Yang D, Pogson AN,

Hein MY, Min KHJ, Wang L: **Mapping transcriptomic vector fields of single cells**. *Cell*

2022, **185**(4):690-711. e645.

30.   Kamimoto K, Stringa B, Hoffmann CM, Jindal K, Solnica-Krezel L, Morris SA:

Dissecting cell identity via network inference and in silico gene perturbation. *Nature*

2023, **614**(7949):742-751.

31.   Jin S, Guerrero-Juarez CF, Zhang L, Chang I, Ramos R, Kuan C-H, Myung P, Plikus

MV, Nie Q: **Inference and analysis of cell-cell communication using CellChat**. *Nature*

862 *communications* 2021, **12**(1):1088.

863 32. Robinson MD, McCarthy DJ, Smyth GK: **edgeR: a Bioconductor package for**

864 **differential expression analysis of digital gene expression data**. *bioinformatics* 2010,

865 **26**(1):139-140.

866 33. Zheng L, Qin S, Si W, Wang A, Xing B, Gao R, Ren X, Wang L, Wu X, Zhang J:

867 **Pan-cancer single-cell landscape of tumor-infiltrating T cells**. *Science* 2021,

868 **374**(6574):abe6474.

869 34. Yan W, Peng Y, van Zyl T, Regev A, Shekhar K, Juric D, Sanes J: **Cell atlas of the**

870 **human fovea and peripheral retina. Sci Rep 10: 9802**. In.; 2020.

871 35. Wu F, Bard JE, Kann J, Yergeau D, Sapkota D, Ge Y, Hu Z, Wang J, Liu T, Mu X:

872 **Single cell transcriptomics reveals lineage trajectory of retinal ganglion cells in**

873 **wild-type and Atoh7-null retinas**. *Nature Communications* 2021, **12**(1):1465.

874 36. Li S, Mo Z, Yang X, Price SM, Shen MM, Xiang M: **Foxn4 controls the genesis of**

875 **amacrine and horizontal cells by retinal progenitors**. *Neuron* 2004, **43**(6):795-807.

876 37. Navneet S, Wilson K, Rohrer B: **Müller glial cells in the macula: their activation and**

877 **cell-cell interactions in age-related macular degeneration**. *Investigative*

878 *Ophthalmology & Visual Science* 2024, **65**(2):42-42.

879 38. Goldman D: **Müller glial cell reprogramming and retina regeneration**. *Nature Reviews*

880 *Neuroscience* 2014, **15**(7):431-442.

881 39. Luna G, Keeley PW, Reese BE, Linberg KA, Lewis GP, Fisher SK: **Astrocyte structural**

882 **reactivity and plasticity in models of retinal detachment**. *Experimental eye research*

883 2016, **150**:4-21.

40.   Shekhar K, Lapan SW, Whitney IE, Tran NM, Macosko EZ, Kowalczyk M, Adiconis X,

Levin JZ, Nemesh J, Goldman M: **Comprehensive classification of retinal bipolar**

neurons by single-cell transcriptomics. *Cell* 2016, **166**(5):1308-1323. e1330.

41.   van Wyk M, Pielecka-Fortuna J, Löwel S, Kleinlogel S: **Restoring the ON switch in**

blind retinas: Opto-mGluR6, a next-generation, cell-tailored optogenetic tool. *PLoS*

biology 2015, **13**(5):e1002143.

42.   Puthussery T, Percival KA, Venkataramani S, Gayet-Primo J, Grünert U, Taylor WR:

Kainate receptors mediate synaptic input to transient and sustained OFF visual

pathways in primate retina. *Journal of Neuroscience* 2014, **34**(22):7611-7621.

43.   Taylor WR, Smith RG: **Transmission of scotopic signals from the rod to rod-bipolar cell**

in the mammalian retina. *Vision research* 2004, **44**(28):3269-3276.

44.   West ER, Lapan SW, Lee C, Kajderowicz KM, Li X, Cepko CL: **Spatiotemporal**

patterns of neuronal subtype genesis suggest hierarchical development of retinal

diversity. *Cell Reports* 2022, **38**(1).

45.   Park SJ, Pottackal J, Ke J-B, Jun NY, Rahmani P, Kim I-J, Singer JH, Demb JB:

Convergence and divergence of CRH amacrine cells in mouse retinal circuitry. *Journal*

of Neuroscience 2018, **38**(15):3753-3766.

46.   Poleg-Polsky A, Diamond JS: **Retinal circuitry balances contrast tuning of excitation**

and inhibition to enable reliable computation of direction selectivity. *Journal of*

Neuroscience 2016, **36**(21):5861-5876.

47.   Schmidt BP, Boehm AE, Tuten WS, Roorda A: **Spatial summation of individual cones**

in human color vision. *PLoS One* 2019, **14**(7):e0211397.

48. Masri RA, Weltzien F, Purushothuman S, Lee SC, Martin PR, Grünert U: **Composition**

of the inner nuclear layer in human retina. *Investigative Ophthalmology & Visual*

Science 2021, **62**(9):22-22.

49. Demb JB, Singer JH: **Functional circuitry of the retina**. *Annual review of vision science*

2015, **1**(1):263-289.

50. Agathocleous M, Harris WA: **Metabolism in physiological cell proliferation and**

differentiation. *Trends in cell biology* 2013, **23**(10):484-492.

51. Lu Y, Shiau F, Yi W, Lu S, Wu Q, Pearson JD, Kallman A, Zhong S, Hoang T, Zuo Z:

Single-cell analysis of human retina identifies evolutionarily conserved and

species-specific mechanisms controlling development. *Developmental cell* 2020,

53(4):473-491. e479.

52. Kang Y, Zhang H, Hu M, Ma Y, Chen P, Zhao Z, Li J, Ye Y, Zheng M, Lou Y:

Alterations in the ocular surface microbiome in traumatic corneal ulcer patients.

Investigative Ophthalmology & Visual Science 2020, **61**(6):35-35.

53. Lobov I, Mikhailova N: **The role of Dll4/Notch signaling in normal and pathological**

ocular angiogenesis: Dll4 controls blood vessel sprouting and vessel remodeling in

normal and pathological conditions. *Journal of ophthalmology* 2018, **2018**(1):3565292.

54. Ferreira JT, Alves M, Dias-Santos A, Costa L, Santos BO, Cunha JP, Papoila AL,

Pinto LA: **Retinal neurodegeneration in diabetic patients without diabetic retinopathy**.

Investigative ophthalmology & visual science 2016, **57**(14):6455-6460.

55. Satue M, Cipres M, Melchor I, Gil-Arribas L, Vilades E, Garcia-Martin E: **Ability of**

swept source OCT technology to detect neurodegeneration in patients with type 2

diabetes mellitus without diabetic retinopathy. *Japanese Journal of Ophthalmology*

2020, **64**:367-377.

56.   Wang Q, Qiao Z, Kang W, Zhu L, Zhang X: **Comparative analysis of co-culture and**

monoculture models in simulating diabetic neurovascular dysfunction: insights into

diabetic retinopathy. *Frontiers in Endocrinology* 2023, **14**:1215218.

57.   Semeraro F, Morescalchi F, Cancarini A, Russo A, Rezzola S, Costagliola C: **Diabetic**

retinopathy, a vascular and inflammatory disease: therapeutic implications. *Diabetes*

& metabolism 2019, **45**(6):517-527.

58.   Tran NM, Shekhar K, Whitney IE, Jacobi A, Benhar I, Hong G, Yan W, Adiconis X,

Arnold ME, Lee JM: **Single-cell profiles of retinal ganglion cells differing in resilience to**

injury reveal neuroprotective genes. *Neuron* 2019, **104**(6):1039-1055. e1012.

939
940

Figure legends

Fig1 Uniform manifold approximation and projection (UMAP) plot of single-cell RNA sequencing (scRNA-seq) data integrated from nine fresh human retinal samples, using the Leiden clustering algorithm at various resolutions, from 0.05 to 1.0. The clustering resolution of 0.3 was chosen as the best value. UMAP plots illustrate the evolution of cell clusters across increasing Leiden algorithm resolutions (leiden_0_05 to leiden_1_0) for integrated data from nine fresh human retinal samples. Each subplot represents a distinct resolution, with colored and numbered clusters indicating different cell populations. The last panel (ID) displays an integrated UMAP visualization where each color represents a different dataset from the nine fresh human retinal samples, highlighting the effectiveness of the integration method.

Fig2 Comprehensive analysis of human retinal major cell type annotation

(A) Uniform manifold approximation and projection (UMAP) plots of integrated human retina sample datasets, colored and labeled by major human retinal cell types.

(B) Dot plots summarizing the known markers used to identify major cell types and a selection of genes distinguishing each cluster for retinal samples. Each row corresponds to a specific major cell type, and the columns correspond to a list of key marker genes; brackets on the top of the plot detail the cell type identified by the genes.

(C) Scatter plot of different expressed genes (DEGs) for each major human retinal cell type. Genes of interest are labeled to indicate significant DE patterns based on up and low-regulated DEGs. Upregulated genes were defined as those with log fold change > 0.5, and downregulated genes as those with log fold change < -0.5.

Fig3 Comprehensive analysis of cell types and clusters in human retina.

(A) Circular representation of Uniform Manifold Approximation and Projection (UMAP)

clustering of the integrated datasets of major cell types and the experimental condition compositions

for each group.

(B) Stacked bar plot showing the distribution ratios of major cell types across the nine human

retina samples.

(C) Stacked bar plot showing the distribution of bipolar cell subtypes within each sample, with

cell types including BB-GB, CB1, DB2, DB3a, DB3b, DB4, DB5, DB6, FMB, IMB, OFFx, and RBC.

(D) Stacked bar plot showing the distribution of cone subtypes within each sample, with cell types

including the Corneal Repair cluster (CR), mlCone, sCone, and ELF1-Cone.

(E) Stacked bar plot showing the distribution of AC subtypes within each sample, with cell types

including GABA-AC1, GABA-AC2, GABA-AC3, gly-AC and intermediate-AC.

Fig4 Gene set variation analysis (GSVA) plots showing the terms of the hallmark gene set

enrichment pathway for each identified major retina cell type. Each row represents a hallmark pathway,

and each column corresponds to a major human retinal cell type. Pathways are annotated by their

direction of regulation (up or down) and the significance of enrichment. The color scale indicates the

significance level (p value) of enrichment, with darker red indicating higher significance.

Fig5 Comparative analysis of astrocytes (ACs), horizontal cells, and retinal ganglion cells

(RGCs).

(A-C) Uniform manifold approximation and projection (UMAP) plots of the expression density

for genes marking ACs, horizontal cells, and RGCs separately.

(D) Heatmap representing the expression levels of selected marker genes for identifying ACs,

horizontal cells, and RGCs.

(E) Dot plots summarizing the selected marker for differentiation. Columns correspond to the

three cell types; rows correspond to a list of key marker genes; brackets on the right side detail the cell

type indicated by the genes.

(F) UMAP plot showing the clustering of retinal cell types based on gene expression profiles,

highlighting AC, horizontal cell, and RGC populations. UMAP plots of the selected cluster, colored and

labeled by amacrine cell, horizontal cell, and retinal ganglion cell annotations.

Fig6 Comparative analysis of astrocytes and müller glia cells (MGCs).

(A) Uniform manifold approximation and projection (UMAP) plot depicting the clustering of

astrocytes and MGCs based on their gene expression profiles. Astrocytes are shown in blue and MGCs

in orange.

(B) Heatmap (left) and dot plot (right) summarizing the expression levels of key marker genes in

astrocytes and MGCs. Columns correspond to MGCs and astrocytes, and the rows correspond to a list

of key marker genes, brackets on the right side detail the cell type that the genes identify.

Fig7 Characterization of bipolar cells (BCs) subtypes in the human retina.

(A) 12 subtypes of BCs are identified in this study. OFF subtypes (1, 2, 3a-3b, FMC, OFFx)

hyperpolarize in response to light, while ON subtypes (RBC, 4, 5, 6, BB+GB, IMB) depolarize to the

same stimuli. There are 11 types of cone BCs and one type of rod BCs.

(B) Uniform manifold approximation and projection (UMAP) plots of the expression density for

specific marker genes across BC subtypes.

(C) Dot plots summarizing the known markers used to identify BC subtypes and a selection of

genes distinguishing each BC cluster. Each row corresponds to a specific BC subtype, and the columns

correspond to a list of key marker genes, brackets on the top of the plot detail the bipolar cell subtype

that these genes identify.

(D) UMAP plot showing the clustering of bipolar cell subtypes based on gene expression profiles,

with each color representing a different BC subtype.

Fig8 Characterization of amacrine cell (AC) subtypes based on gene expression.

(A) Uniform manifold approximation and projection (UMAP) plot showing the clustering of

different AC subtypes. The subtypes are color-coded: GABA-AC1 (blue), Intermediate-AC (orange),

Gly-AC (green), GABA-AC2 (red), and GABA-AC3 (purple).

(B) Dot plot (top) and violin plot (bottom) summarizing the expression levels of key marker genes

in five AC subtypes. Rows correspond to each subtype, and the columns correspond to a list of key

marker genes, brackets on the top detail the cell type that these genes identify.

(C) Gene Ontology (GO) analysis of GABA-AC1, GABA-AC2, and GABA-AC3 sub cell types.

In this panel, the top 10 enriched GO terms are shown for each category, and these terms are sorted

based on gene count. The bars represent the number of genes involved in each GO term, with the total

gene count displayed at the right of each bar. Bar plots are split into three sections in terms of

molecular function (MF, purple), cellular component (CC, blue), and biological process (BP, green).

(D) The dot plot illustrates the results of GO enrichment analysis for differentially expressed

genes in various amacrine cell subtypes (GABA-AC1, GABA-AC2, GABA-AC3, Gly-AC, and

Intermediate-AC). The top 10 most significant GO terms are displayed for each cluster based on

adjusted p-value, ensuring that the most statistically significant terms are highlighted. The y-axis

displays the top 10 significantly enriched GO terms for each cell subtype, encompassing biological

processes, cellular components, and molecular functions. The x-axis represents the different amacrine

cell subtypes. Each dot's size corresponds to the gene count, while the color intensity indicates the gene

ratio. The color scale ranges from yellow (lower gene ratio) to purple (higher gene ratio).

Fig9 Gene expression analysis and clustering of cone subtypes in retinal cells.

(A) Uniform manifold approximation and projection (UMAP) plots of the expression density for

specific marker genes across cone cell sub-clusters.

(B) Dot plots summarizing the known markers used to identify each cone cell sub-cluster and a

selection of genes distinguishing each cone cell cluster. Each row corresponds to a specific cone

subtype or specific cluster, and the columns correspond to a list of key marker genes, brackets on the

top of the plot detail the cluster that these genes identify. Subclusters include corneal repair related

cluster (CR), mlCone, sCone, and ELF1-Cone. The size of the dots represents the fraction of cells in

each group, and the color intensity represents the mean expression level.

(C) UMAP plot illustrating the clustering of cone based on gene expression profiles, with each

color representing a different subtype or cluster: CR (blue), mlCone (orange), ELF1-Cone (green), and

sCone (red).

Fig10 Multi-omics analysis of cellular heterogeneity and gene expression patterns in cones.

(A) Heatmap representing expression levels of top ten differentially expressed genes across

individual cone photoreceptors. Rows correspond to marker genes, columns to individual cells, and

colors indicate expression intensity (red: high, blue: low). The color bar at the top denotes cell types,

including corneal repair related cluster (CR), mlCone, ELF1-Cone, and sCone.

(B) Gene Ontology (GO) analysis for the ELF1-Cone cluster. The bar plot displays enriched GO

terms categorized by biological processes (BP), cellular components (CC), and molecular functions

(MF). The x-axis represents gene count in each pathway, and the y-axis lists specific GO terms.

(C) GSVA (Gene Set Variation Analysis) heatmap showing enrichment scores for hallmark gene

sets across different cone cell clusters. Red indicates the positive enrichment, and blue indicates the

negative enrichment. The top and left-side brackets are the results of hierarchical clustering of both

cone cell types and enrichment pathways.

(D) Correlation heatmap of regulon modules in cone photoreceptors, highlighting patterns of

co-expression. Four distinct modules are numbered and annotated. Key regulons with the highest

residual sum of squares (RSS) rank values, identified by SCENIC analysis, are labeled on the right

hand side.

(E) UMAP (Uniform Manifold Approximation and Projection) plots showing the gene expression

distribution of regulons from the four identified modules (M1-M4) across all cone populations.

(F) RSS rank plot of active regulon in the cluster of ELF1-Cone. The x-axis represents the rank of

each regulon, and the y-axis shows the RSS scores.

Fig11 Dynamo and CellOracle analysis of ELF1-Cone differentiation and regulation in human

retina

(A) UMAP plots showing cell type distributions and velocity vectors across all datasets, living

datasets, and cadaveric datasets. Cell types are color that: coreal repair related cluster (red), mlCone

(pink), ELF1-Cone (light red), and sCone (blue). Arrows indicate predicted cell state transitions.

(B) Cellular dynamics analysis of living datasets, showing speed, divergence, acceleration, and

curvature. Color scales indicate relative values for each metric.

(C) CellOracle perturbation simulations comparing living Datasets (left) with ELF1 knock out

(middle) and PRKAA1 knock out (right). Arrows represent predicted cell state transitions under each

condition.

Fig12 CellChat analysis showing cell-cell interactions between different cone subtypes. The

circular chord diagram illustrates signaling pathways and interaction strengths between cell types, with

a focus on ELF1-Cone interactions.

Fig13 Analysis of gene expression in horizontal cell subtypes.

(A) Uniform manifold approximation and projection (UMAP) plots showing the expression of

LHX1 and ISL1 genes across horizontal cell subtypes. The color scale indicates the expression level of

each gene.

(B) Dot plot summarizing the expression levels of key marker genes in H1 and H2 horizontal cell

subtypes. Rows correspond to H1 and H2, and the columns correspond to a list of key marker genes,

brackets on the top detail the horizontal cell subtype that these genes identify.

(C) UMAP plot depicting the clustering of horizontal cell subtypes based on gene expression

profiles, with H1 cells shown in blue and H2 cells in orange.

Fig14 Comparative analysis of retinal cell types of diabetic conditions

(A) Boxplots depicting the proportion of cell classes across three conditions: diabetic (DM),

diabetic retinopathy (DR), and non-diabetic (nonDM). The plots show the distribution of various retinal

cell types, including astrocyte, horizontal, retinal ganglion cell (RGC), T cell (TC), rod, bipolar cell,

amacrine, cone, microglia, and müller glia cells (MGC).

(B) Heatmap of odds ratio (OR) analysis showing differences in cell abundance between major

human retinal cell types between groups (DR, DM, and nonDM). A p value was calculated by

one-sided Fisher's exact test and corrected by Benjamini-Hochberg procedure. OR $>$ 1.5 and

adjusted p value (Padj) $<$ 0.05 indicating that this subpopulation was more likely to be associated

with this immune subtype, whereas OR $<$ 0.5 and adjusted p value $<$ 0.05 indicated that it was less

likely to be commonly associated with this retinal cell type.

Fig15 Volcano plot showing the results of differential expression (DE) between test group

(diabetes, diabetic retinopathy) and control group (non-diabetes) in human retina samples. The plot

illustrates the distribution of genes based on their expression changes, with the x-axis representing log2

fold change and the y-axis showing -log10 p-value. Colored dots represent different categories of genes:

grey for not statistically different (NS), green for genes meeting the log2 fold change threshold, blue

for genes meeting the p-value threshold, and red for genes meeting both thresholds. Key genes of

interest are labeled with arrows for easy identification. The horizontal dashed line indicates the p-value

threshold for statistically meaningful differences, while vertical dashed lines denote the log2 fold

change thresholds for biologically relevant expression changes.

1104

1105   Table 1. Retinal sample details

Donor	Treatment	Eye	Sex	Age	Diabetic Status	Eye Removal Reasons	Hospital
01	fresh/living	left	Male	76	DM	Corneal staphyloma	L.H.
02	fresh/living	left	Male	55	nonDM	Eyeball atrophy	L.H.
03	fresh/cadaveric	both eyes	Female	55	DM	SCT-H-PON	Y.H.
04	fresh/cadaveric	both eyes	Male	55	DR	BSH	Y.H.
05	fresh/living	right	Male	71	nonDM	Corneal ulcer	Y.H.
06	fresh/living	left	Female	50	nonDM	Eyeball atrophy	Y.H.
07	fresh/living	right	Female	68	nonDM	Corneal perforation	Y.H.

1106   Fresh/living: Fresh retina samples collected from living humans without freezing. Fresh/cadaveric: Fresh retina samples collected from recently deceased bodies without

1107   freezing. DM: Samples with diabetes. DR: Samples with diabetic retinopathy. nonDM: Samples without diabetes. SCT-H-PON: Severe craniocerebral trauma with

1108   hemorrhage in the periphery of the optic nerve. BSH: Brain stem hemorrhage. L.H.: The Affiliated Li Huili Hospital. Y.H.: The Affiliated Ningbo Eye Hospital of Wenzhou

1109   Medical University.

1110

1111

1112

1113

1114 Table 2. Counts and proportion of major cell types in each human retinal scRNA-seq datasets. The major cell classes include amacrine cells, astrocytes, bipolar cells,

Donor	Diabetic Status	Amacrine	Astrocyte	Bipolar	Cone	RGC	Horizontal	Microglia	MGC	Rod	T cell

1115 cones, retinal ganglion cells (RGCs), horizontal cells, microglia, müller glia cells (MGCs), rods and T cells.

		C1	C2	C3	C4	C5	C6	C7	C8	C9	C10
01	DM	90 (0.62%)	76 (0.53%)	2667 (18.49%)	403 (2.79%)	0	148 (1.03%)	3252 (22.55%)	905 (6.27%)	6167 (42.76%)	715 (4.96%)
02	nonDM	9 (0.09%)	1 (0.01%)	528 (5.47%)	9 (0.09%)	0	9 (0.09%)	141 (1.46%)	136 (1.41%)	8732 (90.53%)	80 (0.83%)
03	DM	2 (0.01%)	5 (0.03%)	681 (4.88%)	336 (2.41%)	2 (0.01%)	13 (0.09%)	36 (0.26%)	1347 (9.65%)	11533 (82.60%)	8 (0.06%)
03	DM	25 (0.15%)	19 (0.13%)	753 (4.67%)	507 (3.14%)	0	12 (0.07%)	42 (0.26%)	1923 (11.92%)	12850 (79.62%)	8 (0.05%)
04	DR	310 (1.66%)	23 (0.16%)	1345 (7.22%)	150 (0.81%)	102 (0.55%)	105 (0.56%)	226 (1.21%)	1386 (7.44%)	14972 (80.37%)	11 (0.06%)
04	DR	88 (0.64%)	14 (0.10%)	424 (3.10%)	140 (1.02%)	22 (0.16%)	23 (0.17%)	211 (1.54%)	917 (6.71%)	11815 (86.46%)	12 (0.09%)
05	nonDM	24 (0.29%)	14 (0.10%)	45 (0.55%)	312 (3.80%)	0	1 (0.01%)	84 (1.02%)	893 (10.87%)	6788 (82.66%)	51 (0.62%)
06	nonDM	44 (0.72%)	15 (0.10%)	1421 (23.24%)	165 (2.70%)	1 (0.02%)	75 (1.23%)	216 (3.53%)	855 (13.98%)	3207 (52.45%)	115 (1.88%)
07	nonDM	129 (2.14%)	145 (1.01%)	623 (10.32%)	156 (2.58%)	8 (0.13%)	119 (1.97%)	829 (13.73%)	1582 (26.21%)	2424 (40.15%)	22 (0.36%)

1116   DM: Samples with diabetes. DR: Samples with diabetic retinopathy. nonDM: Samples without diabetes.

1117

A

CR · mlCone · ELF1-Cone · sCone · **All Datasets**

CR · mlCone · ELF1-Cone · sCone · **Living Datasets**

CR · mlCone · ELF1-Cone · sCone · **Cadaveric Datasets**

B

speed_pca

divergence_pca

acceleration_pca

curvature_pca

C

Living Datasets

ELF1 Knock Out

PRKAA1 Knock Out

A

Proportion of Cell Classes by Diabetic Status

B

total = 16261 variables