Translation profiling of stress-induced small proteins reveals a novel link among signaling systems

Sangeevan Vellappan[1,2,3], Junhong Sun[1], John Favate [2,3], Pranavi Jagadeesan[1], Debbie Cerda[1,2], Premal Shah [2,3], Srujana S. Yadavalli[1,2]*

[1]Waksman Institute of Microbiology, Rutgers University, Piscataway, NJ USA

[2]Department of Genetics, School of Arts and Sciences, Rutgers University, Piscataway, NJ USA

[3]Human Genetics Institute of New Jersey, Rutgers University, Piscataway, New Jersey, USA

*Corresponding author

Waksman Institute of Microbiology

Rutgers University

190 Frelinghuysen Rd

Piscataway NJ 08854

Phone: 848-445-0251

Email: sam.yadavalli@rutgers.edu

Abstract

Signaling networks allow adaptation to stressful environments by activating genes that counteract stressors. Small proteins (≤ 50 amino acids long) are a rising class of stress response regulators. *Escherichia coli* encodes over 150 small proteins, most of which lack phenotypes and their biological roles remain elusive. Using magnesium limitation as a stressor, we identify stress-induced small proteins using ribosome profiling, RNA sequencing, and transcriptional reporter assays. We uncover 17 small proteins with increased translation initiation, several of them transcriptionally upregulated by the PhoQ-PhoP two-component signaling system, crucial for magnesium homeostasis. Next, we describe small protein-specific deletion and overexpression phenotypes, underscoring their physiological significance in low magnesium stress. Most remarkably, we elucidate an unusual connection via a small membrane protein YoaI, between major signaling networks – PhoR-PhoB and EnvZ-OmpR in *E. coli*, advancing our understanding of small protein regulators in cellular signaling.

Keywords

Stress response / translation-profiling / small proteins / magnesium starvation / bacterial signaling

46
47
48 • Ribo-RET identifies 17 small proteins induced under low magnesium (Mg^{2+}) stress in *E. coli*
49 • Many of these proteins are transcriptionally activated by PhoQP signaling system
50 • Half of the stress-induced small proteins localize to the membrane
51 • Deletion or overexpression of specific small proteins affects growth under stress
52 • Small protein YoaI connects PhoR-PhoB and EnvZ-OmpR signaling networks
53
54

Introduction

Living organisms sense and respond to environmental stressors through a wide variety of gene
regulatory mechanisms (Carlberg & Molnar, 2016). In bacteria, signal transduction is primarily
carried out by two-component signaling systems (Goulian, 2010; Stock *et al*, 2000; Laub, 2014).
Gene expression regulation can occur at various stages including transcription, post-
transcription, translation, and post-translation (Baumberg, 1999; Banerjee, 2022; Tollerson &
Ibba, 2020; Cain *et al*, 2014). While some of these mechanisms involving regulators such as
transcription factors and small RNAs have been extensively studied (Gottesman, 2019; Storz *et*
al, 2011; Seshasayee *et al*, 2011), other regulatory pathways essential for stress adaptation
remain less well understood. Small proteins (≤50 amino acids in prokaryotes and ≤100 amino
acids in eukaryotes), encoded by authentic small open reading frames, are emerging as key
65

regulators of stress response (Burton *et al*, 2024; Hemm Matthew R. *et al*, 2020; Yadavalli &
Yuan, 2022), highlighting a need for further investigation into their roles and mechanisms of
action.

Advances in bioinformatics, gene expression studies, and ribosome profiling have identified
hundreds of previously unannotated small proteins in all kingdoms of life including humans
(Hemm Matthew R. *et al*, 2020; Gray *et al*, 2022; Pueyo *et al*, 2016; Patraquim *et al*, 2020;
Sberro *et al*, 2019; Kastenmayer *et al*, 2006; Steinberg & Koch, 2021). In *Escherichia coli,* more
than 150 small proteins have been documented (Hemm Matthew R. *et al*, 2020; Gray *et al*,
2022). While there has been significant progress in identifying small proteins, a major gap in our
knowledge is the lack of understanding of the functions of most of these proteins (Gray *et al*,
2022; Burton *et al*, 2024). Small proteins are typically non-essential, at least under standard
growth conditions used in the labs, and their requirement is likely conditional when the activity of
a larger target needs fine-tuning. In support of this idea, it was previously observed that some
small proteins accumulate at higher levels in a specific growth phase, medium, or stress
condition (Hemm *et al*, 2008, 2010). Magnesium (Mg^{2+}) is an important cofactor affecting the
catalysis and stability of numerous proteins and RNAs involved in vital cellular processes,
including translation (Groisman & Chan, 2021). Given the central role of Mg^{2+}, its deprivation
poses a major stress to the cells. The signal transduction and physiological response to low
Mg^{2+} is fairly well understood, which involves the activation of the master regulator, PhoQP two-
component signaling system in gammaproteobacteria (Groisman *et al*, 2021; Groisman & Chan,
2021; Groisman, 2001).

So far, three small proteins are known to mediate stress response to magnesium limitation in
bacteria. Firstly, MgrB, a 48-aa transmembrane protein negatively regulates the sensor kinase
PhoQ of the PhoQP signaling system (Yadavalli Srujana S. *et al*, 2020; Lippa & Goulian, 2009).
During magnesium starvation, the absence of MgrB leads to hyperactivation of the PhoQP
pathway, resulting in cell division inhibition and filamentation in *E. coli* (Yadavalli *et al*, 2016). In
addition to its role in Mg^{2+} stress, loss of MgrB in *E. coli* increases tolerance to trimethoprim, an
antibiotic commonly used to treat bacterial infections (Patel & Matange, 2021). In clinical
isolates of *Klebsiella pneumoniae*, disruption of MgrB-mediated inhibition of PhoQ also leads to
acquired colistin resistance (Cannatelli *et al*, 2014). A second small membrane protein, MgtS is
also induced under low Mg^{2+} stress and is important for magnesium homeostasis (Wang *et al*,
2017; Yin *et al*, 2019). MgtS protects the magnesium transporter, MgtA, from degradation when
Mg^{2+} is limited (Wang *et al*, 2017). PmrR is the third small membrane protein shown to be
expressed under low Mg^{2+} stress in *Salmonella enterica* (Kato *et al*, 2012; Hong *et al*, 2018).
PmrR is activated by the PmrAB two-component system (Chen *et al*, 2021; Kato *et al*, 2012;
Tian *et al*, 2021) in a PhoP-dependent manner and inhibits the lipopolysaccharide modification
enzyme, LpxT.

Here we ask, what subset of the ~150 small proteins in *E. coli* are induced under magnesium
limitation? We hypothesized that, like MgrB, MgtS, and PmrR, many other small proteins with
potential regulatory roles are yet to be identified as part of this stress response. We leveraged
the translation-initiation profiling method, Ribo-RET (Meydan *et al*, 2019; Weaver *et al*, 2019) to

identify small proteins induced by low magnesium stress, revealing a set of 17 proteins
representing a substantial proportion (~11%) of the documented small proteins in *E. coli*. To
investigate the transcriptional regulation of the stress-induced small proteins, we utilized RNA-
Seq and transcriptional reporter assays, we investigated the transcriptional regulation of these
stress-induced small proteins, shedding light on the genomic regions responsible for their
expression and discerning any regulatory influence from the PhoQP system. Additionally, using
epitope-tagging, microscopy, and biochemical analysis, we meticulously examined the
localization, overexpression, and loss-of-function phenotypes of these small proteins. Through
these experiments, we unexpectedly uncovered how the small membrane protein YoaI,
transcriptionally controlled by the PhoRB signaling pathway (Gardner & McCleary, 2019),
displays increased protein levels under magnesium stress and activates another well-studied
two-component system EnvZ-OmpR (Kenney & Anand, 2020) in *E. coli*.
122

Results

Identification of low magnesium stress-induced small proteins in *E. coli* using
translation-initiation profiling
To identify small proteins that accumulate under low magnesium stress, we adapted the
translation initiation profiling method called Ribo-RET (Fig. 1A) – a modified ribosome profiling
approach utilizing translation initiation inhibitors for the identification of small proteins in bacteria
(Cope *et al*, 2022). Retapamulin (RET) is an antimicrobial compound that can stall ribosomes at
the translation start sites to help identify previously unannotated open reading frames (Meydan
et al, 2019; Weaver *et al*, 2019). In this study, we prepared Ribo-RET libraries of wild-type *E.*
coli cells grown in media with or without magnesium to examine small proteins expressed in
response to magnesium starvation.
134

In our Ribo-RET data, we observe an increase in reads mapping to a specific distance of the
ribosomal P-site codon from the 3' end of the reads in our libraries, ~6-10 nucleotides (nt)
downstream of the expected start codon (Fig. S1), similar to the pattern observed in previous
translation initiation profiling studies (Weaver *et al*, 2019). We calculated ribosome density at
annotated translation initiation sites using reads mapped from 4 to 20 nt downstream of the first
nucleotide in the start codon, using a broad window of 16 nt to capture all relevant footprint
sizes. In our analysis, the ribosome footprint sizes range from 16-24 nt, as illustrated in Fig. S1,
as anticipated for the bacterial systems (Glaub *et al*, 2020).
143

The PhoQP signaling system is upregulated under Mg^{2+} limitation (Kato *et al*, 1999) (Fig. S2A-
B). As expected, the read counts at the translation start site for PhoP and PhoQ were more than
4-fold and 10-fold higher, respectively, under stress conditions compared to no stress. Three
small proteins known to be induced under low Mg^{2+} stress – MgrB, MgtS, and PmrR, show a 9-
fold, 5-fold, and 5-fold increase in the reads mapping to the translation start sites, respectively,
under stress vs. no stress (Fig. 1B-C, S2C). Overall, in our Ribo-RET data, we see a moderate
correlation between the reads mapping to the start sites of all proteins expressed under these
two conditions (Fig. 1C), which is indicative of the anticipated changes in the translation
initiation of proteins during stress response.

153
We detected 379 proteins with ≥3-fold increase in read counts under stress, several among
these are small proteins <50 aa in length (Table S4). We narrowed down a set of 17 small
proteins namely MgtS, MgrB, PmrR, MgtT, YmiC, YmiA, YoaI, YobF, YddY, YriB, YadX, YkgS,
YriA, YdgU, YqhI, YadW and DinQ for further analysis (Table 1). Ribosome footprints at the
start site of an open reading frame are a good proxy for the expression of the corresponding
protein. However, it is critical to establish whether a given small open reading frame (sORF)
results in a protein product of that specific size. All but one protein YoaI were previously
validated for expression via genomic epitope-tagging (Table S5). In the case of YoaI, the protein
could not be detected in rich media or specific growth conditions, including envelope stress, acid
stress, and heat shock (Hemm *et al*, 2008, 2010). Given the strong signal for expression of YoaI
in our Ribo-RET data (~12-fold increase in RPM for stress vs. no stress) (Fig. S4A), we
wondered if YoaI is conditionally expressed during magnesium limitation. To examine the
expression of YoaI, we utilized a strain carrying chromosomally-encoded fusion protein YoaI-
SPA (Hemm *et al*, 2008, 2010). We grew cells in media containing different levels of Mg^{2+},
prepared membrane and cytoplasmic fractions, and performed western blot analysis. Consistent
with our hypothesis, we detected expression of YoaI in a magnesium-dependent manner, where
the protein level is highest at the lowest concentration of magnesium (Fig. S4B). In addition, this
result also confirms that YoaI is predominantly associated with the membranes, consistent with
the bioinformatic prediction that it is a membrane protein (Table S7).
173
14 of the 17 shortlisted small proteins were not previously associated with expression under low
magnesium stress. Small proteins YobF (Hemm *et al*, 2010; Hobbs *et al*, 2010) and DinQ
(Weel-Sneve *et al*, 2013) have been linked to other stress responses in *E. coli*, here we find the
expression of these two proteins in low magnesium stress as well. Many of these small proteins
have not been characterized biochemically and have no documented function yet. In the
following sections, we systematically study these low-Mg^{2+}-induced small proteins by
investigating their transcriptional regulation and cellular localization and characterizing their
overexpression and deletion phenotypes.
182
Transcriptional regulation of low magnesium stress-induced small proteins
To determine changes in the levels of transcripts encoding stress-induced small proteins, we
performed RNA-Seq using cells grown to early exponential phase in high and low Mg^{2+}
conditions. It is well-established that in cells depleted for Mg^{2+}, the PhoQP signaling system is
stimulated, leading to PhoP-mediated transcriptional activation of hundreds of genes necessary
for the stress response (Groisman *et al*, 2021). Not surprisingly, we see an increase in the
mRNA abundances for *phoQ* and *phoP* (Fig. S3). Differential expression analysis of our RNA-
seq data revealed significant upregulation (fold change threshold of >2 and a p-value threshold
of <0.05) in transcription under low Mg^{2+} stress for at least 10 of the 17 small proteins
shortlisted, including MgtS, MgtT, and MgrB (Fig. 2A). It is worth noting that YadX also showed
a small increase (1.7-fold) although it falls below the >2-fold threshold we applied in our data
analysis. Intriguingly, YriA, whose open reading frame overlaps with YriB by ~60% did not show
an increase in the transcript levels. In addition, we did not see a significant change in the mRNA
levels corresponding to YoaI, YobF, YkgS, and DinQ. In these cases, it is possible that the

change in transcript abundances is generally low and therefore fell below the significance
threshold. Alternatively, gene expression might be regulated at a post-transcriptional or
translational level under low magnesium stress. Indeed, *yobF* expression is induced post-
transcriptionally during heat shock response (Hemm *et al,* 2010). The expression of *yobF* under
magnesium limitation may be controlled via a similar post-transcriptional regulatory mechanism.

To delve deeper into the transcriptional control mechanism of the small proteins we sought to:
1) identify the regulatory regions upstream of each sORF, and 2) determine if the expression is
PhoQ-dependent. To identify the regions containing the putative promoters of low Mg^{2+}-induced
small proteins, we carefully selected ~200 to 500 base pairs upstream of each sORF (Table
S6). In general, we did not include any annotated full-length ORFs or non-coding RNAs
occurring within the regulatory region being tested, to avoid interference from the ectopic
expression of the corresponding product. In cases where a sORF appeared to be in the middle
of an operon, we selected two regions to test: one upstream of the entire operon and another
immediately upstream of the sORF of interest. We designed transcriptional reporter constructs
to measure the activity of each putative regulatory region, as shown in the schematic (Fig. 2B).
To analyze which of these sORFs are regulated by the PhoQP two-component system, we
measured the transcriptional reporter activities in a *phoQ* deletion strain and compared them to
that of the wild type (Fig. 2C).
215
Consistent with our RNA-seq data, we saw an increase in the transcriptional activity for
regulatory regions corresponding to 11 sORFs with this upregulation under low Mg^{2+} condition
being PhoQ-dependent (Fig. 2C). Two sets of overlapping (out-of-frame) sORFs including
mgtS-mgtT and *yriA-yriB*, strongly suggest their association within the same operon. Three
other genes, *yadW, ymiC,* and *ydgU* are likely regulated as part of the operons, *yadX-clcA-*
yadW, *ymiA-yciX-ymiC*, and *asr-ydgU*, respectively. For all the sORFs occurring within an
established or putative operon, the region upstream of the operon showed a higher
transcriptional reporter activity in response to stress, while a second region immediately
upstream of the given sORF had negligible activity (Table S6, Fig. 2C). Based on the reporter
assay, the transcriptional activity for the 11 sORFs (*mgrB*, *mgtS*, *mgtT*, *ymiA*, *ymiC*, *yddY*, *yriA*,
yriB, yadX, *yadW*, and *ydgU*) peaks at ~6-10 hours. The RNA-Seq data represent
transcriptomic abundances at a given time point of growth (early exponential phase). This
transcriptional reporter assay captures temporal changes in the reporter activity and serves as a
complementary approach to RNA-Seq, where changes in low-abundance transcripts would be
missed. For instance, we did not observe a significant upregulation of *yadX* and *yadW* in the
RNA-Seq data, but we see a PhoQ-dependent increase in the reporter activity for the putative
regulatory region $P_{yadX-clcA-yadW}$ region under low magnesium stress. Conversely, for *pmrR* and
yqhI, while the RNA-Seq data suggested upregulation, we did not detect a significant increase
in the transcriptional reporter activity under stress suggesting that there may be distal or
additional regulatory factors, outside of our reporter construct, involved in controlling their
transcription. Indeed, *pmrR* is shown to be transcriptionally activated by the PmrA-PmrB
signaling system during magnesium limitation through the PhoQP-regulated connector protein
PmrD in *Salmonella enterica* (Kato *et al*, 2012) . This connection between the PhoQ-PhoP and
PmrA-PmrB signaling pathways is also active in *E. coli* under specific conditions including high

stimulation of the PhoQ-PhoP system (Chen *et al*, 2021; Rubin *et al*, 2015). We did not observe
a reporter activity for *dinQ, ykgS, yobF*, and *yoaI* in either the wild-type or Δ*phoQ* cells,
consistent with the results from the RNA-Seq.
243
244 In summary, our RNA-Seq and transcriptional reporter assays reveal transcriptional
245 upregulation of 13 of the 17 small proteins, of which PhoQ regulates at least 11 under
246 magnesium starvation.
247

Regulation of small membrane protein YoaI under Mg^{2+} stress
YoaI is known to be regulated by the PhoRB two-component system (Fitzgerald *et al*, 2023;
Yoshida *et al*, 2012; Chekabab *et al*, 2014), which responds to phosphate starvation (Gardner &
McCleary, 2019). PhoRB signaling is also activated under low magnesium stress due to the
slowdown of protein synthesis and reduction of cytoplasmic phosphate levels (Pontes &
Groisman, 2018). Consistently, we noticed a 10-fold increase in reads mapping to the PhoB
translation start site under magnesium stress (Fig. S2D). We then tested our transcriptional
reporter for *yoaI* (P$_{yoaI}$-YFP) in a low-phosphate medium and saw increased transcriptional
activity for P$_{yoaI}$ when compared to no-stress condition, in a PhoB-dependent manner (Fig. 3A).
This transcriptional activation is PhoQ-independent as we see an amplification of P$_{yoaI}$ reporter
in Δ*phoQ* cells as well (Fig. 3B). Taken together, these results show that *yoaI* transcription is
controlled by PhoB, and there is no significant effect of low magnesium stress on *yoaI*
transcription based on RNA-Seq and transcriptional reporter data (Fig. 2A, C). Yet YoaI protein
levels increase under low magnesium conditions, and we wondered if PhoQ controls YoaI
abundance. To test this possibility, we examined YoaI protein expression in wild-type and
Δ*phoQ* cells in high and low magnesium conditions by western blotting. Our results demonstrate
that YoaI's expression remains magnesium-dependent in a Δ*phoQ* strain, mirroring the pattern
seen in the wild type (Fig. 3C), suggesting that PhoQ does not affect YoaI levels. Therefore,
YoaI expression is induced in a PhoQ-independent manner under magnesium limitation
potentially via a post-transcriptional regulatory mechanism that enhances translation initiation.
268

About half of the low Mg^{2+} stress-induced small proteins localize to the membrane
As a first step to classifying the 17 stress-induced small proteins, we checked if a given small
protein is likely associated with the membrane or cytoplasm. Most of these proteins have limited
or no documented information, so their cellular localization can provide insights into their
potential mechanisms of action and interactions within the cell. Using bioinformatic tools
TMHMM (Möller *et al*, 2001), TMPred (Ikeda *et al*, 2003), and Phobius (Käll *et al*, 2004) to
predict membrane helices, we find that at least 9 of our candidates are putative membrane
proteins. Three of them, MgrB, MgtS, and DinQ, were previously shown to localize to the
membrane in *E. coli* (Hemm *et al*, 2008; Weel-Sneve *et al*, 2013). Additionally, PmrR from *S.*
enterica was shown to be membrane-bound (Kato *et al*, 2012). Our bioinformatic analyses
predict that small proteins YdgU, YmiA, YmiC, YoaI, and YobF also localize to the membrane
(Table S7).
281
Next, to confirm the membrane association and visualize the small protein localization we used
epitope-tagging with GFP. Taking into account the predicted orientation of the putative

transmembrane protein, we created either an N- or C-terminal fusion of GFP by tagging the end
that is expected to face the cytoplasm (Fig. S5A, Table S7) to ensure proper folding of GFP
(Feilmeier *et al*, 2000). Accordingly, YdgU, YmiA, and YmiC were tagged at the N-terminus, and
YoaI was tagged at the C-terminus. Upon observation by fluorescence microscopy, cells
expressing GFP-tagged YdgU, YmiA, YmiC, and YoaI revealed bright fluorescence at the cell
periphery indicative of membrane localization, similar to the known membrane protein, MgrB
(Fig. S5B). As a control, cells expressing GFP alone showed uniform fluorescence indicative of
cytoplasmic localization.
292
For PmrR, we constructed an N-terminal fusion with GFP, where the tag was expected to
localize to the cytoplasm. However, we did not observe membrane localization (Fig. S5B).
Considering the predicted membrane helices, this suggests that the tag interferes with PmrR
localization. Alternately, we tagged the C-terminus of *E. coli* PmrR with a hexahistidine (6XHis)
tag (Fig. S5C) and performed western blotting using membrane and cytoplasmic fractions.
PmrR-6XHis strongly associated with the membrane fraction (Fig. S5D), consistent with the
bioinformatic prediction and data from *S. enterica* PmrR. As in the case of PmrR, YobF-GFP
localized to the cytoplasm, which was inconsistent with its predicted membrane localization
(Table S7). Therefore, we used a 6XHis-tagged version of YobF, which showed a greater
association with the membranes than the cytoplasmic fraction (Fig. S5D). Overall, among the 17
small proteins induced under low Mg^{2+}, we find that 9 of them are membrane proteins.
304

Three small proteins exhibit stress-specific growth defects upon gene deletion
To identify potential defects in growth and cellular morphology caused by the absence of each
small protein in the cell, we utilized single gene deletions of *mgtS*, *mgtT*, *pmrR*, *yadW*, *yadX*,
yddY, *ykgS*, *ymiC*, *yqhI*, *yobF, dinQ*, *ydgU*, *ymiA*, and *yoaI* in the *E. coli* genome. *mgtT* lies in an
operon with *mgtS* (Weaver *et al*, 2019), with an out-of-frame overlap between the stop codon of
MgtS and the start codon of MgtT. Therefore, we carefully generated an *mgtS* deletion without
affecting the start codon of MgtT, and similarly, an *mgtT* deletion while keeping the stop codon
for MgtS intact (for details see materials and methods, Table SI). In the case of *yriAB*, the two
small proteins YriA and YriB have overlapping open reading frames, where ~80% of the *yriB* gene
overlaps with that of *yriA* (Fig. 4D, schematic), so, we first deleted the combined *yriAB* ORF to
see if there is a phenotype. 12 out of the 17 small proteins displayed no discernible growth defects
upon deletion relative to the wild-type strain when grown under magnesium limitation. However,
four mutants – namely *ΔpmrR, ΔyobF, ΔyqhI,* and *ΔyriAB* – exhibit reduced growth yields when
grown over 24 hours (Fig. 4A). The cells carrying these deletions entered the stationary phase
earlier than the wild type. Notably, PmrR and YobF are membrane-bound proteins, while YqhI,
YriA, and YriB are putative soluble proteins (Fig. S5D, Table S7).
321

To determine whether these growth defects are specifically related to the loss of these small
proteins, we complemented the deletion strains with plasmids encoding the corresponding small
proteins and induced with IPTG. Successful rescue of growth is observed for *ΔpmrR*, and *ΔyobF*
(Fig. 4B-C; top panel). To confirm that this complementation is specific to the small protein and
not due to a cryptic ORF or regulatory element present within the small protein coding region, we
prepared plasmids carrying *pmrR* and *yobF* where their start codons are mutated to stop codons.

When tested for complementation in Δ*pmrR* and Δ*yobF*, respectively, the start codon mutants
retained the growth defect similar to that observed for the empty vector (Fig. 4B-C; top panel),
reinforcing that the growth phenotype is specific to the small proteins themselves. In addition, we
investigated whether the growth defect observed with *pmrR* and *yobF* deletions is specific to low
magnesium stress. Our results indicate that deletions of *pmrR* and *yobF* do not cause growth
defects in the absence of stress (Fig. 4B-C; bottom panel), and the phenotype is specific to
magnesium-limited conditions. Intriguingly, expression of plasmid-encoded PmrR in the absence
of stress resulted in a growth defect (Fig. 4B, bottom panel), suggesting that increased expression
of PmrR under no-stress conditions may be detrimental.
337
Given the growth phenotype observed for Δ*yriAB*, we performed complementation with either
plasmid-encoded YriA and/or YriB to establish if the loss of either protein is responsible for the
growth defect. Intriguingly, complementation with either YriA or YriB alone failed to restore growth
but a plasmid expressing the region from *yriB* to *yriA,* mimicking the native genomic arrangement
(*yriAB*), partially rescued the growth defect (Fig. 4D, top panel). In our plasmid construct encoding
YriB and YriAB, there is a possibility of a spurious small protein produced due to the intact, out-
of-frame YriA start codon within the *yriB* ORF (Fig. 4D, schematic) that potentially interferes with
the complementation. To address this issue, we created a modified *yriB* construct where YriA
start codon ATG is mutated to TAA, this substitution changes codons 4 (Asp) and 5 (Val) in YriB
to 4 (Ile) and 5 (Ile). This variant of YriB confers full complementation (Fig. 4D, top panel),
indicating that loss of YriB contributes to the growth phenotype, at least in part. In addition, we
see that the growth defect associated with *yriAB* deletion is specific to low magnesium stress
condition (Fig. 4D, bottom panel).
351
Finally, the growth defect observed in Δ*yqhI* is not complemented with a plasmid encoding YqhI
(Fig. 4E), which suggests that this phenotype is not directly related to the YqhI protein itself.
Intriguingly, we did not see a growth defect in Δ*mgtS* strain when grown in a low magnesium
medium (Fig. 4A), contrasting with the previous report (Wang *et al*, 2017). This discrepancy might
be due to differences in strain construction and/or growth media. Together, our results underscore
the complexities in studying overlapping small ORFs and their phenotypes.
358

Seven small proteins affect growth and cell morphology upon overexpression
As a parallel approach to analyzing loss-of-function phenotypes, we explored the phenotypes
associated with the overexpression of small proteins. We cloned the genes corresponding to the
untagged small proteins into either an IPTG-inducible (all except DinQ as we were unable to
generate a DinQ clone in the IPTG-inducible pEB52 background) or arabinose-inducible (DinQ)
vector. Upon induction, slow growth or a prolonged lag phase is observed for 7 out of the 17 small
proteins tested: MgtS, MgrB, PmrR, YmiA, YoaI, YkgS, and DinQ (Fig. 5). Notably, six of these
proteins are membrane-bound (MgtS, MgrB, PmrR, YmiA, YoaI, and DinQ). In the case of MgtS,
MgrB, PmrR, YmiA, and YoaI, induction of expression with IPTG leads to an extended lag phase
lasting for ~8-10 hours, after which the cells grow exponentially at growth rates similar to that of
control cells carrying empty vector, eventually reaching saturation at comparable levels.
Interestingly, cells expressing small proteins YkgS and DinQ did not display a long lag phase but
grew at a slower rate, eventually catching up to the same growth yield as the control cells. The

growth defect due to DinQ overexpression aligns with previous studies indicating that its
overexpression disrupts membrane potential and depletes intracellular ATP levels (Weel-Sneve
et al, 2013). It is worth noting that overexpression of the other 10 small proteins, including 3
membrane proteins YdgU, YmiC, and YobF had a negligible impact on cell growth under the same
condition, suggesting that cells may tolerate high amounts of these proteins (Fig. 5). More
importantly, the lack of a phenotype for most of these small proteins indicates that the growth
defects observed here are not due to a generic burden on the cell from plasmid-driven protein
overexpression.
380
To be cautious in interpreting the overexpression data, we considered the possibility that the
observed growth defects might be attributed to a cryptic small open reading frame or regulatory
element within the small ORF being overexpressed. To check if the growth defects are indeed
small protein-dependent, we prepared plasmid constructs containing variants of the small protein
where the start codon (ATG) is substituted with a stop codon (TAA). For all 7 small proteins
displaying overexpression phenotypes, we repeated the growth assay with the start codon
mutants of each small protein. Expression of the mutant small proteins rescued the growth defect
with the cells exhibiting similar growth patterns to those carrying the empty vector (Fig. 6A),
reinforcing that the observed growth defects are specific to the overexpression of these small
proteins.
391
In addition to the impact on growth, we explored whether overexpression of these 7 small proteins
(MgtS, MgrB, PmrR, YmiA, YoaI, YkgS, and DinQ) influences cell size and morphology, especially
given the delayed growth (Fig. 5, 6A). We hypothesized that cells overexpressing these small
proteins may undergo morphological changes as part of adaptation during the extended lag and
log phases of growth. Using phase contrast microscopy, we imaged cells induced for expression
of the small proteins during the lag and exponential growth phases. As controls, we used wild-
type cells carrying either an empty vector or a plasmid containing a variant of the small ORF
where its start codon (ATG) is mutated to a stop (TAA). The control cells exhibited a roughly
uniform cell size with volumes around 1.1-1.3 fL and 1.2-1.4 fL in lag and log phases, respectively.
Intriguingly, overexpression of small proteins MgtS, MgrB, PmrR, YkgS, YoaI, and YmiA causes
distinct changes in cell size (Fig. 6B, S6). Specifically, overexpression of MgtS results in smaller
cell size (Fig. 6B), with a reduction in cellular volume to ~30-36% of the control cells carrying
either empty vector or start codon mutants in both lag and log phases (Fig. S6). Similarly,
overexpression of PmrR and YoaI leads to a ~29% decrease in cell size and volume during the
lag phase, however, the cell size reverts to that of control cells during the exponential phase (Fig.
6B, S6). In contrast, overexpression of MgrB leads to an increase in cell size, with an average
volume of 2.2 fL, ~1.6-fold larger than the controls in the lag phase (Fig. 6B, S6), akin to the
Δ*phoQ* cells grown in low magnesium conditions (Fig. S7). This finding is remarkable because
MgrB is known to inhibit PhoQ (Lippa & Goulian, 2009; Yadavalli Srujana S. *et al*, 2020),
consistently, plasmid overexpression of MgrB likely leads to strong inhibition of PhoQ, resulting
in a cellular morphology similar to that of Δ*phoQ*. These cells eventually return to normal size (cell
volume of 1.2 fL) in the exponential phase. Overexpression of YkgS and YmiA causes varying
extents of cell elongation affecting cell size and volume (Fig. 6B, S6). Cells overexpressing DinQ
did not show a significant change in cell size.

Overall, the slow or delayed growth observed here along with the diverse morphological
phenotypes at distinct growth stages reflects how cells adapt in response to the overexpression
of small proteins mentioned above.

Small protein YoaI connects distinct signaling systems – PhoQ-PhoP, PhoR-PhoB, and
EnvZ-OmpR

We wondered whether the growth delay observed upon overexpression of the 7 small proteins
described above is PhoQ-dependent, given the pivotal role of PhoQ in low magnesium stress. To
explore this, we evaluated the effects of overexpressing these proteins in a Δ*phoQ* strain under
magnesium limitation. Notably, the growth phenotype associated with protein overexpression is
retained in the Δ*phoQ* cells, similar to those observed in the wild-type strain for 6 out of 7
candidates – MgrB, MgtS, PmrR, YmiA, YkgS, and DinQ (Fig. 6C), indicating that these effects
are PhoQ-independent. Interestingly, in the case of YoaI overexpression, the extended lag phase
observed in wild-type cells is completely abolished in the Δ*phoQ* cells, suggesting a direct or
indirect link between PhoQ and YoaI where PhoQ functions upstream of YoaI in this regulatory
pathway.

To check if YoaI and PhoQ physically interact with each other, we employed a bacterial two-hybrid
(BACTH) assay based on reconstituting split adenylyl cyclase (CyaA) (Battesti & Bouveret, 2012;
Karimova *et al*, 1998). In this system, CyaA activity is restored when the two fragments T18 and
T25 come close leading to an increase in cyclic AMP (cAMP) levels and subsequent expression
of β-galactosidase (β-gal) from the *lac* promoter. We fused the T18 fragment to YoaI and tested
it against T25-PhoQ. We included the T18-MgrB and T25-PhoQ pair as a positive control and two
other histidine kinases, PhoR and EnvZ, each fused to the T25 fragment as additional controls.
In this experiment, YoaI did not show an interaction with either PhoQ or PhoR, however, to our
surprise there was a strong signal between YoaI and EnvZ that is comparable to the positive
control (Fig. 7A). EnvZ is primarily known to be an osmosensor, integral to the EnvZ-OmpR two-
component system, which regulates porins in response to changes in osmolarity, pH,
temperature, and growth phase (Kenney & Anand, 2020). Several small membrane proteins are
known to modulate the activities of sensor kinases (Yadavalli & Yuan, 2022). A 127-amino acid
protein, MzrA regulates EnvZ resulting in higher levels of phosphorylated OmpR (Gerken *et al*,
2009). Based on the BACTH interaction between YoaI and EnvZ, we suspected YoaI might
influence the activity of the EnvZ-OmpR signaling system. To address this possibility, we utilized
an EnvZ-OmpR-regulated transcriptional reporter strain with chromosomal fusion of mScarlet to
P*omrB* (Bastet *et al*, 2024) and measured *omrB* promoter activity in cells grown in low magnesium
vs. no-stress media. Interestingly, we observe an increase of 2.5-fold in EnvZ-regulated
transcription under stress in wild-type cells (Fig. 7B). The significantly high activity of *omrB*
reporter in low Mg^{2+} conditions suggests that upregulation of YoaI protein levels may mediate this
effect (Fig. 3C, S4). Upon deletion of *yoaI*, the *omrB* transcription reduced by ~1.4-1.5-fold in both
low and high Mg^{2+} media, indicating that YoaI affects EnvZ-OmpR activity regardless of stress.
Together with our observation that YoaI-SPA levels increase under low Mg^{2+} stress (Fig. 3C,
S4B), the higher transcriptional activity of the EnvZ-regulated promoter can be attributed to the
positive interaction of YoaI with EnvZ. To further support the idea that an increase in YoaI levels

is responsible for modulating the activity of the EnvZ-OmpR system, we examined *omrB*
transcription upon plasmid overexpression of YoaI in wild-type *E. coli* cells. We observed a greater
than 6-fold increase in P$_{omrB}$-*lacZ* reporter activity in cells expressing YoaI on a plasmid (Fig. S8A)
relative to the control cells carrying an empty vector. We also tested two other reporters to genes
ompF, P$_{ompF}$-YFP and *ompC*, P$_{ompC}$-CFP (Batchelor *et al*, 2004), and found a ~2-fold change in
their transcriptional activities (Fig. S8B-C), consistent with the activation of EnvZ (Alphen &
Lugtenberg, 1977; Batchelor *et al*, 2005; Kenney & Anand, 2020). In summary, these findings
suggest that YoaI interacts with and stimulates EnvZ activity, potentially enhancing OmpR-P
levels.
469
470

Discussion
In this study, we identified 17 small proteins (≤50 amino acids) induced by low magnesium
stress from approximately 150 documented small proteins in *E. coli*, using Ribo-RET (Meydan
et al, 2019; Weaver *et al*, 2019). Only 3 of these proteins, MgrB, MgtS, and PmrR were
previously studied in the context of magnesium starvation (Lippa & Goulian, 2009; Wang *et al*,
2017; Kato *et al*, 2012). For all the stress-induced small proteins except one (YoaI), expression
from the genomic locus was verified *in vivo* (Table S5). In a previous report when cells were
grown in a rich medium, expression of YoaI tagged with SPA epitope from a genomic translation
fusion was undetected (Hemm *et al*, 2008). Interestingly, our observation of cells grown in low
Mg^{2+} detected a robust expression of YoaI expression using the same construct, highlighting the
condition-specific expression of this protein (Fig. 3C, S4B). Using RNA-Seq, we found that
many of these stress-induced proteins are transcriptionally activated in low magnesium media.
Further, we identified the regions upstream of these small ORFs responsible for transcriptional
regulation using operon fusions and reporter assays. Few hits, including YoaI and YobF, do not
show a change in transcription levels under low Mg^{2+}, suggesting that they are regulated post-
transcriptionally.
487
Lacking any functional or phenotypic information for most of these stress-induced proteins, we
began their characterization by predicting their subcellular localization using bioinformatic tools
(Tables 1, S7). By epitope tagging and imaging or western blotting, we empirically determined
that 9 of the 17 shortlisted stress-induced small proteins localize to the membrane (Fig. S5).
These membrane-bound small proteins may interact with, stabilize, or fine-tune the activities of
large transmembrane proteins, such as sensor kinases, channel proteins, and drug efflux pumps,
thereby mediating cellular adaptation and stress responses (Yadavalli & Yuan, 2022). Among the
9 candidates, three proteins – MgrB, MgtS, and PmrR are shown to have specific membrane
protein targets, as described earlier. To further investigate their biological roles, we performed
targeted deletion and protein overexpression analysis.
498
Loss-of-function analysis of small proteins can be a useful approach to understanding their
physiological roles. For example, deleting *mgrB* causes hyperactivation of the PhoQP two-
component system, leading to cell division inhibition and filamentation of cells grown under low
magnesium stress (Yadavalli *et al*, 2016). In clinical isolates of *Klebsiella pneumoniae,* the
absence of MgrB-mediated PhoQ inhibition has been linked to acquired colistin resistance

(Cannatelli *et al*, 2014; Olaitan *et al*, 2014). In the current work, we found deletion phenotypes
associated with the following small proteins—PmrR, YobF, YriA/B, and YqhI—leading to an early
entry into the stationary phase and reduced yield at saturation under magnesium limitation. Based
on plasmid complementation, growth defects caused by PmrR and YobF are specific to the
respective small proteins. The overlapping ORFs of YriA and YriB present a unique challenge in
interpreting the Δ*yriAB* growth phenotype observed under low magnesium stress. While
expressing the native *yriAB* region partially rescues the defect, individual expression of YriA or
YriB fails to do so; however, modifying YriB by mutating the out-of-frame YriA start codon fully
restores growth, emphasizing that the loss of YriB is a critical factor in the observed phenotype.
In the case of YqhI, the deletion phenotype could not be fully complemented by plasmid-encoded
small protein, pointing to putative cryptic ORF or regulatory factors within the small ORF that
warrant further investigation.
516
We also analyzed overexpression phenotypes associated with each of the stress-induced small
proteins to gain insights into cellular pathways or components functionally linked to these proteins.
For instance, overexpression of MgtS is shown to activate the PhoRB two-component system by
regulating the phosphate transporter, PitA (Yin *et al*, 2019). Similarly, DinQ overexpression
disrupts membrane potential and depletes intracellular ATP levels (Weel-Sneve *et al*, 2013). We
observed overexpression phenotypes for 7 of the 17 small proteins induced under magnesium
stress—MgtS, MgrB, PmrR, YmiA, YoaI, YkgS, and DinQ—resulting in either a long lag phase or
slow growth (Fig. 5). These overexpression phenotypes are specific to each small protein as we
see control plasmids carrying a mutated start codon do not cause growth defects (Fig. 6A).
Analysis of the cell morphology during the lag and exponential phases of growth revealed
interesting variations in cell sizes and volumes that reflect how cells adapt to the overexpression
of these small proteins. Remarkably for one of the small proteins, PmrR, both the deletion and
overexpression caused growth defects under low magnesium stress. It is unclear how cells
balance small protein levels when needed during specific conditions vs. when not needed, say in
the absence of stress. In this context, complementation of the Δ*pmrR* strain with plasmid-encoded
PmrR restored normal growth under magnesium limitation but caused a growth defect under no-
stress condition (Fig. 4B). This suggests that PmrR levels may be controlled precisely in cells
undergoing stress, as both its absence and its overexpression disrupt the balance necessary for
cellular homeostasis.
536
In *Salmonella enterica,* PmrR has been shown to inhibit the inner membrane protein LpxT, which
plays a role in lipopolysaccharide modification (Kato *et al*, 2012; Tian *et al*, 2021; Hong *et al*,
2018). We explored whether the growth defects associated with PmrR deletion and
overexpression in *E. coli* are mediated through its interaction with LpxT. Interestingly, expressing
PmrR in a Δ*lpxT* background did not alleviate the growth defects regardless of stress (Fig. S9A).
It is possible that higher LpxT levels, resulting from the absence of PmrR, could lead to growth
defects. However, both single deletions (Δ*lpxT* and Δ*pmrR*) and the double deletion (Δ*pmrR*
Δ*lpxT*) exhibited growth defects under magnesium stress (Fig. S9B). This result suggests that the
growth defect upon *pmrR* deletion is independent of PmrR's interaction with LpxT. PmrR may be
involved in other pathways or regulatory mechanisms where it has additional interacting partners.
It is not uncommon for small proteins to have multiple binding partners, as seen with MgtS, which

interacts with both MgtA and PitA (Yin *et al*, 2019; Wang *et al*, 2017). Whether PmrR associates
with other targets contributing to phenotypes observed here warrants further investigation.
550
Finally, we would like to discuss the interesting case of the small membrane protein YoaI,
whose transcription is controlled by the PhoR-PhoB signaling system (Fitzgerald *et al*, 2023;
Yoshida *et al*, 2012; Chekabab *et al*, 2014). YoaI levels are quite low or undetectable when cells
are grown in a rich medium or specific stress conditions (Hemm *et al*, 2008, 2010). However, we
find that YoaI protein levels strongly increase under magnesium limitation and it becomes
detectable (Fig. 3C). Overexpression of YoaI leads to a long delay in growth (Fig. 5, 6A, S4B).
This phenotype is abolished in a *phoQ* mutant (Fig. 6C), hinting at an unknown PhoQ-
dependent factor that may contribute to the growth defect observed upon YoaI overexpression.
Through a bacterial two-hybrid assay, we found a direct interaction of YoaI with the EnvZ sensor
kinase (Fig. 7A), which controls osmoregulation (Kenney & Anand, 2020) suggesting that this
small protein acts as a connector between the PhoR-PhoB and EnvZ-OmpR signaling networks.
Activation of EnvZ by YoaI is supported by the data from our transcriptional reporter analysis of
promoters regulated by the EnvZ-OmpR system (Fig. 7B, S8). Modulation of sensor kinases via
small protein connectors appears to be a general theme that many small membrane proteins of
unknown function may fall under (Yadavalli & Yuan, 2022). For instance, under acidic
conditions, a 65-amino acid protein SafA is activated by the EvgS-EvgA two-component system,
which subsequently interacts with and activates the PhoQ-PhoP system (Eguchi *et al*, 2007). In
a second example, an 88-amino acid protein PmrD connects the PmrB-PmrA and PhoQ-PhoP
signaling systems in *Salmonella* (Kox *et al*, 2000; Kato & Groisman, 2004). Another 127-amino
acid protein MzrA is known to bind and activate EnvZ, which links the CpxA-CpxR and EnvZ-
OmpR pathways in *E. coli* (Gerken *et al*, 2009). Taken together, our findings outlining the
transcriptional regulation of YoaI by PhoB (Fig. 3A), PhoQ-dependent growth phenotype upon
YoaI overexpression (Fig. 6C), and YoaI's physical interaction with EnvZ and modulation of
EnvZ-regulated promoter activity (Fig. 7B, S8) reveal how this 34-amino acid small protein acts
as a connector between distinct two-component signaling systems – PhoRB, PhoQP, and EnvZ-
OmpR to integrate diverse stress responses (Fig. 7C).
577
Most small proteins (≤50 amino acids in prokaryotes) are of unknown function and little is known
about when/if they are expressed in the cell and what their physiological relevance is. Small
proteins are thought to fine-tune the activity and/or stability of larger target molecules in the cell,
adding a layer of gene regulation. This type of conditional regulation by small proteins may be
especially important when cells are grown in non-ideal or stressful environments. Overall, our
work describing the identification of condition-specific small proteins and targeted phenotypic
characterization provides a deeper understanding of small protein regulation and the
physiological consequences of altering their expression levels, laying a foundation for
deciphering their functions and regulatory mechanisms under relevant stress conditions.
587
588
589

Methods

Media and growth conditions

Bacterial cultures were grown in either Luria-Bertani (LB) Miller medium (IBI Scientific) or minimal A medium (Miller & Others, 1992) supplemented with 0.2% glucose, 0.1% casamino acids, and the indicated concentration of $MgSO_4$, with aeration at 37°C, unless otherwise specified. In general, overnight cultures are grown in supplemented minimal A medium containing 1 mM $MgSO_4$ and antibiotic(s) as applicable. Saturated cultures are diluted 1:500 into fresh medium. For low Mg^{2+} stress, no Mg^{2+} was added when overnight cultures were diluted into fresh medium. Media containing 10 mM Mg^{2+} was used as the no-stress condition. For phosphate stress experiments, MOPS minimal medium (Teknova) supplemented with 0.2% glucose, 0.1% casamino acids, and either 1 mM K_2HPO_4 (no-stress) or no K_2HPO_4 (phosphate stress). For routine growth on solid medium, LB containing 1.5% bacteriological-grade agar (VWR Lifesciences) was used. For antibiotic selection, carbenicillin, kanamycin, and chloramphenicol were added at a final concentration of 50-100 μg ml^{-1}, 25- 50 μg ml^{-1}, and 6-25 μg ml^{-1}. b-isopropyl-D-thiogalactoside (IPTG) was used at a final concentration of 500 μM, unless otherwise specified. Arabinose was used at final concentrations of 0.5% and 30 mM for pBAD24-based vectors and pMR120-based vectors, respectively, to induce protein expression when needed.

Strains, plasmids, and cloning

See supplemental information for details of the strains (Table S1), plasmids (Table S2), and primers (Table S3) used in this study. All the strains used were derived from *Escherichia coli K-12* MG1655. Strain JW2162 was a gift from Dr. Bryce Nickels; plasmids pEB52, pAL38, pAL39, pSMART, pMR120 and strains AML67, MDG147, TIM92, TIM100, and TIM202 were gifts from Dr. Mark Goulian. Strains GSO195, GSO219, GSO225, GSO232, and GSO317 were gifts from Dr. Gisela Storz. Strains JM2110 and JM2113 were kind gifts from Dr. Maude Guillier.

A modified version of the method described by (Datsenko & Wanner, 2000) was used to generate in-frame gene deletions corresponding to each small protein. Briefly, the entire small open reading frame (sORF) (including the stop codon) was deleted from the MG1655 genome and substituted with a kanamycin resistance cassette. While constructing the gene deletions, we avoided deleting any overlapping annotated ORFs, non-coding RNAs, or putative regulatory regions in the vicinity of the target gene. In the case of *yriA-yriB (yriAB)*, the two small protein ORFs significantly overlapped; therefore, we deleted the combined *yriAB* ORF. Detailed information on the genomic coordinates of deleted regions can be found in Table S1. Genomic deletions and reporter constructs carrying the kanamycin resistance cassettes were transferred between strains using P1$_{vir}$ transduction (Miller & Others, 1992) and validated by PCR. When necessary, the kanamycin cassette was excised from the chromosome by FLP-mediated recombination using pCP20, as previously described (Cherepanov & Wackernagel, 1995). PCR and loss of antibiotic resistance confirmed the excision of the kanamycin cassette.

Plasmids expressing translational fusions of GFP-A206K to the small open reading frames (sORFs), specifically pPJ1-13 and pSV25, were generated through inverse PCR (Ochman *et al*,

1990) using pAL39 as a template. Plasmids encoding C-terminally 6XHis-tagged-YobF and -
PmrR (sORF-GGSG linker-6XHis) were created using inverse PCR with pEB52 as a template,
resulting in pSV14 and pSV26, respectively. Plasmids encoding untagged small proteins
(pSV30-pSV37, pSV58, pSV60, and pJS1-pJS7) were also generated by inverse PCR with
pEB52 as a template. pSV54 was generated by inverse PCR using pBAD24 as a template.
Plasmids pSV38-pSV45, pSV55, pSV59, and pSV61, encoding variants of sORFs with start
codon substitutions to stop codons (TAA), were generated by site-directed mutagenesis with
inverse PCR using templates pSV35, pSV34, pJS7, pJS4, pSV36, pJS5, pSV54, pSV37, and
pSV60 as applicable. For pSV39-pSV42, another in-frame start codon within five amino acids of
the annotated start was also substituted to TAA. All constructs were verified by Sanger
sequencing. To generate pSV47, *yoaI* was PCR amplified from the MG1655 genome, digested
with *XbaI/EcoRI* restriction enzymes, and then cloned into pUT18C at *XbaI/EcoRI* sites.
646
To examine the transcriptional regulation of the sORFs and identify the putative regulatory
regions, ~200-500 bp upstream region of the start codon of each sORF was selected for cloning
(see details in Table S6). If a given sORF was suspected to be part of an operon, two putative
regions were chosen for analysis, the first directly upstream of the gene and a second region
upstream of the entire operon. Each DNA segment was amplified from MG1655 genomic DNA
by PCR and inserted upstream of the *yfp* reporter gene in pMR120 into the restriction sites
(*XhoI/BamHI* to construct pSV16, while *EcoRI-BamHI* sites were used for the remaining
constructs). The pMR120 vector also includes a *cfp* reporter regulated by a constitutive *tetA*
promoter, serving as an internal plasmid control. The resulting plasmids (pPJ14, pPJ16-pPJ20,
pPJ23, pSV16-21, pSV24, pSV28, pSV29) were confirmed by Sanger sequencing and
transformed into MG1655 and TIM202 (*ΔphoQ*). Note that pSMART (Lucigen), pMR120, and
the derivatives are single-copy plasmids when transformed into *E. coli* (Godiska R, Patterson M,
Schoenfeld T, and Mead DA, 2005). *E. cloni* replicator cells (Lucigen) are transformed with
these plasmids and cultured with arabinose induction to increase the copy number for plasmid
isolation.
662
Translation initiation profiling (Ribo-RET) and RNA-Seq
Overnight cultures of wild-type *E. coli* grown in supplemented minimal A medium and 1 mM
$MgSO_4$ were diluted 1:500 in 100 ml of fresh medium and the specified concentrations of
$MgSO_4$. Once the cultures reached an OD600 of ~0.4, they were split into two 50 ml conical
tubes. Cells from one tube were pelleted and submitted to Genewiz for RNA-Seq, while the cells
from the second tube were used for Ribo-RET. For Ribo-RET, cells were treated with
retapamulin at a final concentration of 100 µg ml^{-1} for 5 minutes at 37°C with continuous
shaking. After retapamulin treatment, cells were filtered through a nitrocellulose membrane with
a pore size of 0.45 µm, quickly scraped off the surface, and flash-frozen in liquid nitrogen. The
resulting samples were processed, and libraries were prepared as described previously (Cope
et al, 2022) with the following modifications. Specifically, RNA fragments of 15-30 nucleotides
(nt) in length were isolated by performing a tight gel size selection, and rRNA depletion was
performed using the RiboCop for Gram-negative bacteria kit (Lexogen). The resulting RNA
fragments were then pooled, and libraries were prepared, with a final library structure consisting
of a 5' adapter – 4 random bases – insert – 5 random bases – sample barcode – 3' adapter. The

randomized bases serve as Unique Molecular Identifiers (UMIs) for deduplication. The
multiplexed library was sequenced on an Illumina HiSeq 4000 with a paired end reading of 150
bases (PE150 runs) at a sequencing depth of 40-60 million raw reads per sample.
681

Sequencing data analysis
The raw sequencing data for this project has been deposited in the GEO database with the
accession number GSE276379. An R package with codes for data processing and analysis has
been made available on GitHub (https://github.com/yadavalli-lab/Small-proteins-induced-under-
low-magnesium-stress-in-E.coli). The raw sequencing data was first demultiplexed, and the
adaptors were removed using cutadapt (Martin, 2011). The reads were then depleted of rRNA
and tRNA and deduplicated using umi_tools dedup (Smith *et al*, 2017). The Ribo-RET reads
were aligned to the *E. coli* MG1655 genome NC_000913.3 (RefSeq assembly accession:
GCF_000005845.2) using hisat2 (Kim *et al*, 2019), and the ribosome density was assigned to
the 3' end of the reads. Initiation peak density was calculated for the annotated translation start
sites by summing up the normalized reads (reads per million, RPM) within 4 to 20 nt
downstream of the first nucleotide in the start codon. For transcript quantification of RNA-Seq
reads, Kallisto (Bray *et al*, 2016) was used. DESeq2 (Love *et al*, 2014) was used for the
differential expression analysis of gene expression in the RNA-Seq data. Normalization of the
data for differential expression was carried out using the "apeglm" method (Zhu *et al*, 2019).
Fold-changes were estimated by comparing two replicates of the low Mg^{2+} dataset to two
replicates of the no-stress dataset, with filtering criteria set at more than two-fold changes and a
p-value below 0.05.
700

Measurement of growth (optical density) and fluorescence using a microplate reader
The optical density (OD_{600}) measurements for small protein overexpression and deletion
constructs were performed in a clear flat bottom 96-well microplate (Corning) containing 150 µL
culture per well using a microplate reader (Agilent BioTek Synergy Neo2S). Fluorescence
measurements for transcriptional reporters were carried out similarly, using black-walled, clear-
flat bottom 96-well microplates (Corning). Saturated cultures grown in supplemented minimal A
or MOPS minimal medium were normalized to the same initial OD_{600} values across all samples
and diluted 1:50 in fresh medium containing specified concentrations of $MgSO_4$ or K_2HPO_4,
along with appropriate antibiotics and inducers, as indicated. The microplates were incubated at
37 °C with double orbital shaking before the real-time measurements of optical density (at 600
nm), YFP (excitation: 500 nm; emission: 540 nm), or CFP (excitation: 420 nm; emission: 485
nm) were acquired every hour for 20-24 hours.
713

Microscopy, and image analysis
For localization of GFP-tagged small proteins, cells were grown to an OD_{600} of ~0.2 (~3 hours)
and then rapidly cooled in an ice slurry. Streptomycin was added to a final concentration of 250
µg ml^{-1}. GFP fluorescence was captured with a 475 nm excitation wavelength and a 540 nm
emission filter, with an exposure time of 30 ms and 20% intensity. While the exposure times
were typically set to these specified values, slight variations were occasionally made to optimize
imaging based on the levels of fluorescence expression observed.
721

For overexpression analysis, overnight cultures were diluted 1:500 into supplemented minimal
medium containing no added MgSO$_4$, 50 µg ml^{-1} carbenicillin, and either 500 µM IPTG or 0.5%
arabinose, as indicated. Cells were grown to the lag phase (OD$_{600}$ ~0.005), then rapidly cooled
in an ice slurry before phase contrast microscopy. For empty vector control and start codon
variants, we captured lag phase micrograph images approximately 30-40 minutes post-
inoculation and -induction. For MgtS, MgrB, PmrR, YoaI, and YmiA overexpression, lag phase
measurements were performed at 2-3 hours, and for YkgS and DinQ, at 1-hour post-inoculation
and -induction. To concentrate the cells, 1 mL of the culture was pelleted and resuspended in
30-50 µL minimal A salt solution. For exponential phase analysis (OD$_{600}$ between 0.2-0.4) in
cultures carrying empty vector control and start codon variants images were taken at ~3-4
hours. In the case of cells expressing MgtS, MgrB, PmrR, YoaI, and YmiA, images were taken
at 16-18 hours, and YkgS and DinQ at 6-8 hours. Microscope slides were prepared using 1%
agarose pads as described previously (Yadavalli *et al*, 2016), with approximately 5-7 µL of the
resuspended cells. A 40 ms exposure time and 20% intensity were applied to capture the
phase-contrast images.
737
For single-cell fluorescence measurements of wild type and Δ*yoaI* cells carrying P$_{omrB}$-*mScarlet*
reporter, overnight cultures were diluted 1:500 into fresh supplemented minimal A medium
containing either 10 mM or no added MgSO$_4$. Cells were grown to exponential (OD$_{600}$ ~0.3 for
both stress and no stress conditions) and stationary phase (OD$_{600}$ ~1.0 for stress condition and
~3.5 for no stress condition), and then rapidly cooled in an ice slurry. Streptomycin was added
to a final concentration of 250 µg ml^{-1}. Fluorescence was measured using the mCherry channel
with an excitation wavelength of 570 nm and an emission filter of 645 nm. The exposure time
was set to 100 ms, with an intensity of 20%. Background fluorescence was determined by
imaging MG1655 cells grown under the same conditions.
747
For single-cell fluorescence measurements upon YoaI overexpression (to measure reporter
activities of P$_{ompF}$-*yfp* and P$_{ompC}$-*cfp*), the overnight cultures were diluted 1:500 into fresh
supplemented minimal A medium containing 10 mM MgSO$_4$. Cells were grown to exponential
phase (OD$_{600}$ between 0.2-0.4), then rapidly cooled in an ice slurry, and streptomycin was added
at 250 µg ml^{-1}. YFP fluorescence was measured using a 500 nm excitation wavelength and a
535 nm emission filter. CFP fluorescence was measured using a 435 nm excitation wavelength
and a 480 nm emission filter. The exposure time for YFP and CFP measurements was set to 50
ms, respectively, with an intensity of 20%. The background fluorescence was determined by
imaging MG1655/pEB52 cells grown under the same conditions.
757
Image visualization and acquisition were performed using a Nikon Ti-E epifluorescence
microscope equipped with a TI2-S-HU attachable mechanical stage. The images were captured
with a Prime 95B sCMOS camera from Teledyne Photometrics with 1x1 binning. All image
acquisition was managed using Metamorph software (Molecular Devices), version 7.10.3.279. A
minimum of 50 cells per replicate were analyzed for cell volume and single-cell fluorescence
quantification. The distributions of cellular volume were assumed to follow the sphero-cylindrical
shape of *E. coli.* The volume was calculated using the formula $V = \frac{pi}{4} * W^2 * \left(L - \frac{W}{3}\right)$. Images

were analyzed, and the cellular length, width, and fluorescence intensity were quantified using
ImageJ (Schneider *et al*, 2012) and MicrobeJ plugin (Jiang *et al*, 2014).
767

Preparation of membrane and cytoplasmic fractions
Membrane and cytoplasmic protein fractions were prepared as described previously (Yadavalli
Srujana S. *et al*, 2020). Briefly, saturated cultures of MG1655/pSV14 (PmrR-GGSG-6XHis),
MG1655/pSV26 (YobF-GGSG-6XHis), and MG1655/pEB52 (empty vector) were diluted 1:500
in 4 ml of LB media containing 100 µg ml^{-1} carbenicillin and grown at 37 °C. After 4 hours of
growth, 0.5 mM IPTG was added to the cultures, and the cells were harvested after 2 hours of
induction. Cell pellets were resuspended in 50 µl of cold resuspension buffer containing 20%
sucrose, 30 mM Tris pH 8.0, and 1X protease inhibitor cocktail (Sigma, cOmplete, EDTA-free).
Then, 50 µl of 10 mg ml^{-1} lysozyme, freshly prepared in 0.1 M EDTA pH 7.0, was added to the
cell suspension, and the mixture was incubated on ice for 30 minutes. Next, 1 ml of 3 mM EDTA
pH 7.5 was added to each sample, followed by sonication (10s pulse, 10s gap, 6 times) using
VCX-130 Vibra-Cell Ultrasonic sonicator (Fisher Scientific). Samples were briefly centrifuged at
6,000 rpm for 10 minutes at 4 °C to remove cellular debris, and the supernatant was collected
and spun at 21,000 ×g for 30 minutes at 4 °C. The supernatant from this step is separated and
stored as the cytoplasmic fraction. The pellet, representing the membrane fraction, was
resuspended in a storage buffer containing 20 mM Tris pH 8.0/20% glycerol. For strains
carrying genomic YoaI-SPA translational fusion, cultures were grown to an OD$_{600}$ of ~0.2-0.4
before harvesting, and membrane and cytoplasmic fractions were prepared as mentioned
above.
787

Western blot analysis
The membrane and cytoplasmic fractions were resuspended in 4X Laemmli sample buffer
containing 5% 2-mercaptoethanol and no dyes. The samples were heated at 70 °C for 10
minutes and run on a 12% Bis-Tris gel (NuPage, Invitrogen) using MES running buffer
(Invitrogen) at 160V for 80 minutes. After electrophoresis, proteins were transferred to a PVDF
membrane (Amersham™ Hybond™) with a 0.2 µm pore size using a semi-dry transfer cell (Bio-
Rad Trans-Blot SD). The membranes were blocked with 5% w/v milk in Tris-buffered saline, pH
7.4, with 1% Tween 20 (TBS-T). Primary antibodies – mouse M2 anti-FLAG (Sigma-Aldrich) and
rabbit anti-6XHis tag antibody (Rockland) were used at 1:1000 dilution to detect SPA- and
6XHis-tagged constructs, respectively. IRDye 800CW goat anti-mouse and IRDye 680RD
donkey anti-rabbit antibodies (LiCOR) were used for secondary detection. SeaBlue™ Plus2 Pre-
stained Protein Standard (Invitrogen) was used as a ladder to visualize lower molecular weight
protein bands. The proteins were visualized using a LI-COR Odyssey imager.
801

Bacterial two-hybrid and β-galactosidase reporter gene assays
For the bacterial two-hybrid assay, several clones from the transformation plate were picked to
reduce heterogeneity (Battesti & Bouveret, 2012) and inoculated in LB containing 100 µg ml^{-1}
ampicillin and 50 µg ml^{-1} kanamycin. Cultures were grown overnight at 30°C with shaking. The
cultures were then diluted 1:1000 into LB supplemented with 500 µM IPTG and grown at 30°C
until OD$_{600}$ reached between 0.2-0.4, after which β-galactosidase activities (Miller units) were
measured as previously described (Thibodeau *et al*, 2004).

For the β-galactosidase reporter gene assay, strains carrying P$_{omrB}$-*lacZ* with either pSV34
(pYoaI) or pEB52 (empty vector) were grown overnight in LB medium with 100 µg ml^{-1}
ampicillin. The next day, the cultures were diluted 1:100 in the same medium and grown at 37
°C. After 3 hours of growth, 0.5 mM IPTG was added to the cultures, and the cells were
harvested after 2 hours of induction. For high-throughput measurements in a 96-well format, the
protocol was modified as previously described (Thibodeau *et al*, 2004).
816

Resource availability

Lead contact
Further information and requests for resources and reagents should be directed to and will be
fulfilled by the lead contact, Srujana S. Yadavalli (sam.yadavalli@rutgers.edu).

Materials availability
Requests for strains and plasmids generated in this study should be directed to and will be
fulfilled by the lead contact, Srujana S. Yadavalli (sam.yadavalli@rutgers.edu).

Data and code availability
Next-generation sequencing data generated in this study are deposited at Gene Expression
Omnibus (GEO) with accession number GSE276379. All data generated or analyzed during this
study are included in the manuscript and its supporting files. Source data files for all figures
have been provided. The code used for data processing and analysis is available in a series of
R Markdown documents hosted on GitHub (https://github.com/yadavalli-lab/Small-proteins-
induced-under-low-magnesium-stress-in-E.coli).

834
835

Acknowledgments
The authors would like to thank Drs. Mark Goulian, Bryce Nickels, Gisela Storz, and Maude
Guillier for generously sharing strains and plasmids. We thank Dr. Goulian, and the past and
present members of the Yadavalli, Storz, and Shah Labs for their input and helpful discussions.
Funding: S.S.Y. is supported by the National Institutes of Health - National Institute of General
Medical Sciences (NIH-NIGMS) ESI-MIRA R35 GM147566 and institutional start-up funds from
Rutgers. P.S. was supported by NIH/NIGMS grant R35 GM124976 and start-up funds from the
Human Genetics Institute of New Jersey at Rutgers University. S.V. is supported by the
Waksman Institute Busch Predoctoral fellowship (2022-24). The funders did not play any role in
the study design, data collection and analysis, decision to publish, or preparation of the
manuscript.

847
848

849 Author contributions
850 S.V., P.S., and S.S.Y. conceived the experiments. All authors planned the experiments,
851 designed the methodology, and performed data analysis. S.V., P.S., and S.S.Y. wrote the initial
852 draft of the manuscript, P.S. contributed to revisions and editing, and all authors reviewed the
853 final manuscript.

854
855

Disclosure and competing interests statement

The authors declare no competing interests. P.S. is a director at an RNA-therapeutics startup.

References

Alphen WV & Lugtenberg B (1977) Influence of osmolarity of the growth medium on the outer membrane protein pattern of Escherichia coli. *J Bacteriol* 131: 623–630

Banerjee T (2022) Regulation of Gene Expression in Prokaryotes. In *Genetics Fundamentals Notes* pp 569–596. Singapore: Springer Nature Singapore

Bastet L, Korepanov AP, Jagodnik J, Grondin JP, Lamontagne A-M, Guillier M & Lafontaine DA (2024) Riboswitch and small RNAs modulate btuB translation initiation in Escherichia coli and trigger distinct mRNA regulatory mechanisms. *Nucleic Acids Res* 52: 5852–5865

Batchelor E, Silhavy TJ & Goulian M (2004) Continuous control in bacterial regulatory circuits. *J Bacteriol* 186: 7618–7625

Batchelor E, Walthers D, Kenney LJ & Goulian M (2005) The Escherichia coli CpxA-CpxR envelope stress response system regulates expression of the porins ompF and ompC. *J Bacteriol* 187: 5723–5731

Battesti A & Bouveret E (2012) The bacterial two-hybrid system based on adenylate cyclase reconstitution in Escherichia coli. *Methods* 58: 325–334

Baumberg S ed. (1999) Prokaryotic Gene Expression. doi:10.1093/oso/9780199636044.001.0001 [PREPRINT]

Bray NL, Pimentel H, Melsted P & Pachter L (2016) Near-optimal probabilistic RNA-seq quantification. *Nat Biotechnol* 34: 525–527

Burton AT, Zeinert R & Storz G (2024) Large Roles of Small Proteins. *Annu Rev Microbiol*

Cain JA, Solis N & Cordwell SJ (2014) Beyond gene expression: the impact of protein post-translational modifications in bacteria. *J Proteomics* 97: 265–286

Cannatelli A, Giani T, D'Andrea MM, Di Pilato V, Arena F, Conte V, Tryfinopoulou K, Vatopoulos A, Rossolini GM & COLGRIT Study Group (2014) MgrB inactivation is a common mechanism of colistin resistance in KPC-producing Klebsiella pneumoniae of clinical origin. *Antimicrob Agents Chemother* 58: 5696–5703

Carlberg C & Molnar F (2016) Mechanisms of Gene Regulation 2nd ed. Dordrecht, Netherlands: Springer

Chekabab SM, Jubelin G, Dozois CM & Harel J (2014) PhoB activates Escherichia coli O157:H7 virulence factors in response to inorganic phosphate limitation. *PLoS One* 9: e94285

Chen AI, Albicoro FJ, Zhu J & Goulian M (2021) Effects of regulatory network organization and
environment on PmrD connector activity and polymyxin resistance in Klebsiella
pneumoniae and Escherichia coli. *Antimicrob Agents Chemother* 65

Cherepanov PP & Wackernagel W (1995) Gene disruption in Escherichia coli: TcR and KmR
cassettes with the option of Flp-catalyzed excision of the antibiotic-resistance
determinant. *Gene* 158: 9–14

Cope AL, Vellappan S, Favate JS, Skalenko KS, Yadavalli SS & Shah P (2022) Exploring
Ribosome-Positioning on Translating Transcripts with Ribosome Profiling. *Methods Mol
Biol* 2404: 83–110

Datsenko KA & Wanner BL (2000) One-step inactivation of chromosomal genes in Escherichia
coli K-12 using PCR products. *Proc Natl Acad Sci U S A* 97: 6640–6645

Eguchi Y, Itou J, Yamane M, Demizu R, Yamato F, Okada A, Mori H, Kato A & Utsumi R (2007)
B1500, a small membrane protein, connects the two-component systems EvgS/EvgA
and PhoQ/PhoP in *Escherichia coli*. *Proc Natl Acad Sci U S A* 104: 18712–18717

Feilmeier BJ, Iseminger G, Schroeder D, Webber H & Phillips GJ (2000) Green fluorescent
protein functions as a reporter for protein localization in Escherichia coli. *J Bacteriol* 182:
4068–4076

Fitzgerald DM, Stringer AM, Smith C, Lapierre P & Wade JT (2023) Genome-Wide Mapping of
the Escherichia coli PhoB Regulon Reveals Many Transcriptionally Inert, Intragenic
Binding Sites. *MBio* 14: e0253522

Gardner SG & McCleary WR (2019) Control of the phoBR Regulon in Escherichia coli. *EcoSal
Plus* 8

Gerken H, Charlson ES, Cicirelli EM, Kenney LJ & Misra R (2009) MzrA: a novel modulator of
the EnvZ/OmpR two-component regulon. *Mol Microbiol* 72: 1408–1422

Glaub A, Huptas C, Neuhaus K & Ardern Z (2020) Recommendations for bacterial ribosome
profiling experiments based on bioinformatic evaluation of published data. *J Biol Chem*
295: 8999–9011

Godiska R, Patterson M, Schoenfeld T, and Mead DA (2005) Beyond pUC: Vectors for Cloning
Unstable DNA. In *DNA Sequencing: Optimizing the Process and Analysis*, Kieleczawa J
(ed) pp 55–76. Sudbury, Massachusetts: Jones & Bartlett Learning

Gottesman S (2019) Trouble is coming: Signaling pathways that regulate general stress
responses in bacteria. *J Biol Chem* 294: 11685–11700

Goulian M (2010) Two-component signaling circuit structure and properties. *Curr Opin Microbiol*
13: 184–189

Gray T, Storz G & Papenfort K (2022) Small Proteins; Big Questions. *J Bacteriol* 204: e0034121

Groisman EA (2001) The pleiotropic two-component regulatory system PhoP-PhoQ. *J Bacteriol*
183: 1835–1842

Groisman EA & Chan C (2021) Cellular Adaptations to Cytoplasmic Mg2+ Limitation. *Annu Rev*
Microbiol 75: 649–672

Groisman EA, Duprey A & Choi J (2021) How the PhoP/PhoQ System Controls Virulence and
Mg2+ Homeostasis: Lessons in Signal Transduction, Pathogenesis, Physiology, and
Evolution. *Microbiol Mol Biol Rev* 85: e0017620

Hemm Matthew R., Weaver Jeremy, Storz Gisela, Lovett Susan T. & Hinton Deborah (2020)
Escherichia coli Small Proteome. *EcoSal Plus* 9

Hemm MR, Paul BJ, Miranda-Ríos J, Zhang A, Soltanzad N & Storz G (2010) Small stress
response proteins in Escherichia coli: proteins missed by classical proteomic studies. *J*
Bacteriol 192: 46–58

Hemm MR, Paul BJ, Schneider TD, Storz G & Rudd KE (2008) Small membrane proteins found
by comparative genomics and ribosome binding site models. *Mol Microbiol* 70: 1487–
1501

Hobbs EC, Astarita JL & Storz G (2010) Small RNAs and small proteins involved in resistance
to cell envelope stress and acid shock in Escherichia coli: analysis of a bar-coded
mutant collection. *J Bacteriol* 192: 59–67

Hong X, Chen HD & Groisman EA (2018) Gene expression kinetics governs stimulus-specific
decoration of the *Salmonella* outer membrane. *Sci Signal* 11

Ikeda M, Arai M, Okuno T & Shimizu T (2003) TMPDB: a database of experimentally-
characterized transmembrane topologies. *Nucleic Acids Res* 31: 406–409

Jiang C, Brown PJB, Ducret A & Brun YV (2014) Sequential evolution of bacterial morphology
by co-option of a developmental regulator. *Nature* 506: 489–493

Käll L, Krogh A & Sonnhammer ELL (2004) A combined transmembrane topology and signal
peptide prediction method. *J Mol Biol* 338: 1027–1036

Karimova G, Pidoux J, Ullmann A & Ladant D (1998) A bacterial two-hybrid system based on a
reconstituted signal transduction pathway. *Proc Natl Acad Sci U S A* 95: 5752–5756

Kastenmayer JP, Ni L, Chu A, Kitchen LE, Au W-C, Yang H, Carter CD, Wheeler D, Davis RW,
Boeke JD, *et al* (2006) Functional genomics of genes with small open reading frames
(sORFs) in S. cerevisiae. *Genome Res* 16: 365–373

Kato A, Chen HD, Latifi T & Groisman EA (2012) Reciprocal control between a bacterium's
regulatory system and the modification status of its lipopolysaccharide. *Mol Cell* 47:
897–908

Kato A & Groisman EA (2004) Connecting two-component regulatory systems by a protein that
protects a response regulator from dephosphorylation by its cognate sensor. *Genes Dev*
18: 2302–2313

Kato A, Tanabe H & Utsumi R (1999) Molecular characterization of the PhoP-PhoQ two-
component system in Escherichia coli K-12: identification of extracellular Mg2+-
responsive promoters. *J Bacteriol* 181: 5516–5520

Kenney LJ & Anand GS (2020) EnvZ/OmpR Two-Component Signaling: An Archetype System
That Can Function Noncanonically. *EcoSal Plus* 9

Kim D, Paggi JM, Park C, Bennett C & Salzberg SL (2019) Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* 37: 907–915

Kox LF, Wösten MM & Groisman EA (2000) A small protein that mediates the activation of a
two-component system by another two-component system. *EMBO J* 19: 1861–1872

Laub MT (2014) The role of two-component signal transduction systems in bacterial stress
responses. In *Bacterial Stress Responses* pp 45–58. Washington, DC, USA: ASM Press

Lippa AM & Goulian M (2009) Feedback inhibition in the PhoQ/PhoP signaling system by a
membrane peptide. *PLoS Genet* 5: e1000788

Love MI, Huber W & Anders S (2014) Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* 15: 550

Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal 17: 10–12

Meydan S, Marks J, Klepacki D, Sharma V, Baranov PV, Firth AE, Margus T, Kefi A, Vázquez-
Laslop N & Mankin AS (2019) Retapamulin-Assisted Ribosome Profiling Reveals the
Alternative Bacterial Proteome. *Mol Cell* 74: 481-493.e6

Miller JH & Others (1992) A Short Course in Bacterial Genetics: A Laboratory Manual and
Handbook for Escherichia Coli and Related Bacteria CSHL Press

Möller S, Croning MD & Apweiler R (2001) Evaluation of methods for the prediction of
membrane spanning regions. *Bioinformatics* 17: 646–653

Ochman H, Medhora MM, Garza D & Hartl DL (1990) Amplification of flanking sequences by
inverse PCR. *PCR protocols: A guide to methods and applications*: 219–227

Olaitan AO, Morand S & Rolain J-M (2014) Mechanisms of polymyxin resistance: acquired and
intrinsic resistance in bacteria. *Front Microbiol* 5: 643

Patel V & Matange N (2021) Adaptation and compensation in a bacterial gene regulatory
network evolving under antibiotic selection. *Elife* 10

Patraquim P, Mumtaz MAS, Pueyo JI, Aspden JL & Couso J-P (2020) Developmental regulation
of canonical and small ORF translation from mRNAs. *Genome Biol* 21: 128

Pontes MH & Groisman EA (2018) Protein synthesis controls phosphate homeostasis. *Genes
Dev* 32: 79–92

Pueyo JI, Magny EG & Couso JP (2016) New peptides under the s(ORF)ace of the genome.
Trends Biochem Sci 41: 665–678

Rubin EJ, Herrera CM, Crofts AA & Trent MS (2015) PmrD is required for modifications to
escherichia coli endotoxin that promote antimicrobial resistance. *Antimicrob Agents
Chemother* 59: 2051–2061

Sberro H, Fremin BJ, Zlitni S, Edfors F, Greenfield N, Snyder MP, Pavlopoulos GA, Kyrpides
NC & Bhatt AS (2019) Large-Scale Analyses of Human Microbiomes Reveal Thousands
of Small, Novel Genes. *Cell* 178: 1245-1259.e14

Schneider CA, Rasband WS & Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image
analysis. *Nat Methods* 9: 671–675

Seshasayee ASN, Sivaraman K & Luscombe NM (2011) An overview of prokaryotic
transcription factors : a summary of function and occurrence in bacterial genomes.
Subcell Biochem 52: 7–23

Smith T, Heger A & Sudbery I (2017) UMI-tools: modeling sequencing errors in Unique
Molecular Identifiers to improve quantification accuracy. *Genome Res* 27: 491–499

Steinberg R & Koch H-G (2021) The largely unexplored biology of small proteins in pro- and
eukaryotes. *FEBS J* 288: 7002–7024

Stock AM, Robinson VL & Goudreau PN (2000) Two-component signal transduction. *Annu Rev
Biochem* 69: 183–215

Storz G, Vogel J & Wassarman KM (2011) Regulation by small RNAs in bacteria: expanding
frontiers. *Mol Cell* 43: 880–891

Thibodeau SA, Fang R & Joung JK (2004) High-throughput beta-galactosidase assay for
bacterial cell-based reporter systems. *Biotechniques* 36: 410–415

Tian X, Manat G, Gasiorowski E, Auger R, Hicham S, Mengin-Lecreulx D, Boneca IG & Touzé T
(2021) LpxT-Dependent Phosphorylation of Lipid A in Escherichia coli Increases
Resistance to Deoxycholate and Enhances Gut Colonization. *Front Microbiol* 12: 676596

Tollerson R 2nd & Ibba M (2020) Translational regulation of environmental adaptation in
bacteria. *J Biol Chem* 295: 10434–10445

Wang H, Yin X, Wu Orr M, Dambach M, Curtis R & Storz G (2017) Increasing intracellular
magnesium levels with the 31-amino acid MgtS protein. *Proc Natl Acad Sci U S A* 114:
5689–5694

Weaver J, Mohammad F, Buskirk AR & Storz G (2019) Identifying Small Proteins by Ribosome
Profiling with Stalled Initiation Complexes. *MBio* 10

Weel-Sneve R, Kristiansen KI, Odsbu I, Dalhus B, Booth J, Rognes T, Skarstad K & Bjørås M
(2013) Single transmembrane peptide DinQ modulates membrane-dependent activities.
PLoS Genet 9: e1003260

Yadavalli Srujana S., Goh Ted, Carey Jeffrey N., Malengo Gabriele, Vellappan Sangeevan,
Nickels Bryce E., Sourjik Victor, Goulian Mark, Yuan Jing & Silhavy Thomas J. (2020)
Functional Determinants of a Small Protein Controlling a Broadly Conserved Bacterial
Sensor Kinase. *J Bacteriol* 202: e00305-20

Yadavalli SS, Carey JN, Leibman RS, Chen AI, Stern AM, Roggiani M, Lippa AM & Goulian M
(2016) Antimicrobial peptides trigger a division block in Escherichia coli through
stimulation of a signalling system. *Nat Commun* 7: 1–10

Yadavalli SS & Yuan J (2022) Bacterial Small Membrane Proteins: the Swiss Army Knife of
Regulators at the Lipid Bilayer. *J Bacteriol* 204: e0034421

Yin X, Wu Orr M, Wang H, Hobbs EC, Shabalina SA & Storz G (2019) The small protein MgtS
and small RNA MgrR modulate the PitA phosphate symporter to boost intracellular
magnesium levels. *Mol Microbiol* 111: 131–144

Yoshida Y, Sugiyama S, Oyamada T, Yokoyama K & Makino K (2012) Novel members of the
phosphate regulon in Escherichia coli O157:H7 identified using a whole-genome
shotgun approach. *Gene* 502: 27–35

Zhu A, Ibrahim JG & Love MI (2019) Heavy-tailed prior distributions for sequence count data:
removing the noise and preserving large differences. *Bioinformatics* 35: 2084–2092

1087 Figures and tables
1088
1089

1090
1091
Figure 1. Identification of small proteins induced under low magnesium stress in *E. coli*
by translation-initiation profiling (Ribo-RET). (A) Schematic diagram showing the Ribo-RET
and RNA-Seq experimental setup used to identify small proteins induced by low magnesium
stress. Wild-type *E. coli* K-12 MG1655 cells were grown in supplemented minimal A medium
containing $MgSO_4$ at either 10 mM (no stress) or ~1 µM (no added magnesium, low Mg^{2+}
stress). (B) Scatterplot showing the correlation between the Ribo-RET reads mapping to the
annotated start sites of proteins expressed under low Mg^{2+} stress and no stress. The red dots
represent annotated small proteins of ≤ 50 amino acids in length, and the gray dots represent
proteins > 50 amino acids long. Small proteins, MgrB, MgtS, and PmrR, known to be induced
under magnesium starvation, are highlighted. Pearson's coefficient, r= 0.85. (C) Ribo-RET data
for a representative small protein MgrB induced under low Mg^{2+} stress. A purple dashed line
indicates the translation start site, and the 8 nucleotide (nt) indicates the distance between the
highest ribosome density peak and the first nucleotide of the start codon. (Also see Tables 1,
S4, Fig. S1 and S2)
1106
1107

1109

1110

Figure 2. Transcriptional regulation of small proteins induced under low magnesium

stress. (A) RNA-Seq analysis of transcripts representing small proteins induced under low

magnesium stress. Volcano plot illustrating differential expression of small proteins induced

under low magnesium stress from RNA-Seq data. Transcripts either upregulated or

downregulated are highlighted in red or blue, respectively, and those with no significant change

are represented in gray. The criteria for significance include a fold change threshold of >2 and a

p-value of <0.05. The -$\log_{10}$ P-value of *mgrB* and *mgtS* were 75 and 140 respectively. To

prevent distortion from their exceptionally high values dominating the plot, the transformed p-

values were capped at 10 to enhance visualization. Data represent differential expression

analysis from two independent replicates. (B) Schematic representation of the transcriptional

reporter fusion construct. Regions (200-500 bp) upstream of the small open reading frames of

interest were amplified from the MG1655 genome and cloned into a YFP-encoding reporter

plasmid (pMR120) resulting in plasmids pPJ14, pPJ16-pPJ20, pPJ23, pSV16-pSV21, pSV24,

pSV28-pSV29 (See Table S6 for genomic coordinates of the regulatory regions tested here and

Table S2 for plasmid details). (C) Measurement of transcriptional reporter activity (YFP:CFP) for

the indicated small proteins under low Mg^{2+} (no added magnesium). The solid and dashed lines

represent the wild-type *E. coli* MG1655 and $\Delta phoQ$ cells, respectively, harboring plasmids with

either transcriptional fusion for a putative regulatory region of a small protein with *yfp* (red line)

or *yfp* only (gray line). The schematic above each plot depicts the arrangement of the small

protein-encoding genes (shown in gray), including the putative regulatory regions and operons

where applicable. Transcriptional reporters showing an increased activity compared to the

empty vector control (pMR120) are highlighted in red. The cultures were grown in a

supplemented minimal medium with 6 µg ml^{-1} chloramphenicol. The data represent averages

and standard errors of the means for four independent replicates. (Also see Tables S4, S6, and

Fig. S3)

1136

1137

1138

1139

1140

Figure 3. Regulation of YoaI expression during magnesium limitation. (A-B) Transcriptional activity of *yoaI* was measured as a function of YFP reporter fluorescence in *E. coli* MG1655, $\Delta phoB$ (SV48), and $\Delta phoQ$ (TIM100) strains under phosphate stress as indicated. The red and gray lines represent cells carrying plasmid-encoded P_{yoaI}-*yfp* transcriptional reporter (pPJ18) and *yfp*-only control vector (pMR120), respectively. The cultures were grown in a supplemented MOPS minimal medium with 6 µg ml^{-1} chloramphenicol with either 1 mM K_2HPO_4 for no stress condition, or no added K_2HPO_4 for phosphate stress. The data illustrate average YFP:CFP fluorescence with standard error of the means, derived from four independent replicates. (C) Detection of YoaI protein in *E. coli* wild-type (GSO317) and $\Delta phoQ$ (SV60) strains containing *yoaI-SPA* genomic translational fusion. Cells were grown in supplemented minimal A medium, with $MgSO_4$ added at the specified concentration (10mM or no added $MgSO_4$). Membrane fractions were analyzed by western blotting using M2 anti-FLAG antibodies (top) and by Coomassie Brilliant Blue staining (bottom). The data represent results from two independent replicates.

1158

Figure 4. Effect of deleting genes encoding low Mg^{2+} stress-induced small proteins on
bacterial growth. (A) Growth curves of wild-type *E. coli* MG1655 (gray) and mutants
corresponding to the deletions of small protein-encoding genes (red) (strains SV64, AML67,
SV35, SV31, SV41, SV57, SV59, SV33, SV29, SV25, SV39, SV43, SV58, SV37, SV27, and
SV56, please see Table S1 and methods for strain details). Deletion mutants showing reduced
growth yield compared to wild-type cells are highlighted in red. The cultures were grown in
supplemented minimal A medium with no added magnesium and 50 µg ml^{-1} carbenicillin. Data
represent averages and standard errors of means for four independent cultures. (B-C)
Complementation of growth phenotypes associated with *pmrR,* and *yobF* deletions. Growth
curves of Δ*pmrR* (SV35), and Δ*yobF* (SV34) deletion strains carrying an empty vector (pEB52,
solid gray line) or complemented with plasmids encoding small proteins pPmrR (pSV35) and
pYobF (pJS5) (red line) or their variants pPmrR$_{ATG→TAA}$ (pSV38), and pYobF$_{ATG→TAA}$ (pSV45)
(black line), as indicated. Wild-type MG1655 cells containing the empty vector are included as a
control (dashed gray line). (D) Schematic of the overlapping genes *yriAB* followed by the growth
curves of gene deletion Δ*yriAB* (SV25) complemented with pYriAB (pSV58), pYriA (pJS1), pYriB
(pSV37), pYriB (YriA$_{ATG→TAA}$) (pSV59), or an empty vector (pEB52, as indicated. (E) Growth
curves of gene deletion Δ*yqhI* (SV37) carrying an empty vector (pEB52), or complemented with
a plasmid encoding YqhI. Wild-type cells containing an empty vector (pEB52, dashed gray line)
are included as a control. For panels A-D, cultures were grown in supplemented minimal A
medium containing 50 µg ml^{-1} carbenicillin and either no added magnesium or 10 mM MgSO$_4$ as
indicated. 500 µM IPTG was used for induction, except for PmrR derivatives (pSV35 and
pSV38), where 12.5 µM IPTG was used. Data represent averages and standard errors of
means for four independent cultures.
1182
1183

1184
1185
Figure 5. Effect of overexpression of low Mg^{2+} stress-induced small proteins on growth.
Growth curves of *E. coli* MG1655 cells expressing each of the 17 stress-specific small proteins
(encoded by plasmids pSV30-37, pSV54, pSV60, or pJS1-7; please see Table S2 for details)
are represented in red, while cells carrying the empty vector (pEB52 and pBAD24) are shown in
gray. The cells were grown under low Mg^{2+} stress (no added magnesium). Small proteins whose
expression causes a growth defect or delay are highlighted in red. The cultures were grown in
supplemented minimal A medium with 50 µg ml^{-1} carbenicillin. 500 µM IPTG was used as an
inducer for all P$_{trc}$-based vectors, in the case of pSV54and pBAD24, 0.5% arabinose was used.
Data represent averages and standard error of means for four independent cultures.

1195
1196
Figure 6. Analysis of growth defect and cell size changes associated with overexpression of small proteins, MgtS, MgrB, PmrR, YkgS, YoaI, YmiA and DinQ. (A) Growth curves of *E. coli* MG1655 cells expressing the indicated small proteins (red line) and empty vector control (pEB52 or pBAD24, gray line). The start codon (ATG) of each small protein is mutated to a stop codon (TAA), and the growth curves of cells expressing these constructs are depicted in dotted black. Plasmids pSV34-36, pSV38-42, pSV54-55, pSV60-61, or pJS4, pJS7 were used as applicable, please see Table S2 for details. Data represent averages and standard error of means for four independent cultures. (B) Representative phase contrast micrographs of the cells expressing small proteins during lag (top panel) and exponential phase (bottom panel) after induction, scale bar = 5 µm. Data is representative of three independent cultures. (C) Growth curves of Δ*phoQ* (TIM202) cells expressing the indicated small proteins (red line) and empty vector control (pEB52 or pBAD24, gray line). Plasmids pSV34-36, pSV54, pSV60, pJS4, and pJS7 were used as applicable, please see Table S2 for details. Data represent averages and standard error of means for four independent cultures. For panels A-C, the cultures were grown in supplemented minimal A medium containing no added magnesium, 50 µg ml^{-1} carbenicillin, and 500 µM IPTG for all P$_{trc}$-based vectors, except for pSV54 and pBAD24, where 0.5% arabinose was utilized instead. (Also see Fig. S6 and S7)

1214
1215
Figure 7. Effect of Yoal expression on the EnvZ-OmpR system. (A) BACTH assays were
conducted using cells harboring T18 and T25 fusion proteins of *B. pertussis* adenylyl cyclase, in
a *cyaA* mutant BACTH host strain, SAM85, which contains *lacI*^{q}. β-gal activity was measured to
assess interactions between T18-Yoal (pSV47) and T25-linked PhoQ (pAL27), PhoR (pSY68),
or EnvZ (pKK14). T25 alone was used as a negative control, and T18-MgrB (pAL33) combined
with T25-PhoQ (pAL27) was used as a positive control. The cultures were grown in LB medium
containing 50 μg ml^{-1} kanamycin, 100 μg ml^{-1} carbenicillin, and 500 μM IPTG. Each circle
represents a biological replicate, with the mean values from two independent experiments
indicated, expressed in Miller units (MU). (B) Transcriptional activity of *omrB* was measured as
a function of mScarlet reporter fluorescence in *E. coli* MG1655 (SV97), and Δ*yoaI* (SV95)
strains at the stationary phase (top panel) and exponential phase (bottom panel). Each circle
corresponds to a single cell. Δ*yoaI* cells are depicted in red, while wild-type cells are in gray.
The cultures were grown in supplemented minimal A medium containing either 10mM (no
stress) or no added magnesium (Mg^{2+} stress). Data are derived from three independent
replicates and the number of cells analyzed is indicated by (n). Mean fluorescence is shown
with blue bars and is noted in blue text. P-values indicate the results of a t-test when Δ*yoaI* cells
were compared to the wild-type cells, ****P ≤ 0.0001, ***P ≤ 0.001, and "ns" = P > 0.05. (C)
Schematic representation of small membrane protein Yoal as a connector of the two-component
systems, PhoR-PhoB and EnvZ-OmpR. Yoal is transcriptionally controlled by the PhoR-PhoB
system, under low Mg^{2+} Yoal protein levels increase and it binds sensor kinase EnvZ and
modulates signaling through the EnvZ-OmpR pathway. (Also see Fig. S9)

Table 1. Summary of the 17 small proteins induced under low magnesium stress in _E. coli_. Small proteins with at least a 3-fold increase in
reads mapping to the translation initiation site under low Mg^{2+} relative to no stress are indicated below. The analysis of transcriptional regulation,
localization, and characterization of deletion and overexpression phenotypes is summarized.

Small protein		Transcriptional Regulation	Membrane localization	Phenotypic characterization	
				Deletion	Overexpression
MgtS	+	PhoQP-dependent, in an operon _mgtS-mgtT_	Y	No effect	Growth defect (prolonged lag phase) & decrease in cell size during lag and exponential phase
MgrB	+	PhoQP-dependent	Y	Increase in cell length, and change in colony morphology (larger, rough, translucent colonies) (Yadavalli _et al_, 2016)	Growth defect (prolonged lag phase) & increase in cell size during lag phase
PmrR	+	BasSR-dependent (Chen _et al_, 2021; Kato _et al_, 2012)	Y	Growth defect (lower yield at saturation)	Growth defect (prolonged lag phase) & decrease in cell size during lag phase
MgtT	+	PhoQP-dependent, in an operon _mgtS-mgtT_	N	No effect	No effect
YmiC	+	PhoQP-dependent, in an operon _ymiA-yciX-ymiC_	Y	No effect	No effect
YmiA	+		Y	No effect	Growth defect (prolonged lag phase) & increase in cell length during lag phase
YoaI	-	PhoRB-dependent (Fitzgerald _et al_, 2023; Yoshida _et al_, 2012; Chekabab _et al_, 2014)	Y	No effect	Growth defect (prolonged lag phase) & decrease in cell size during lag phase
YobF	-		Y	Growth defect (lower yield at saturation)	No effect
YddY	+	PhoQP-dependent	N	No effect	No effect
YriB	+	PhoQP-dependent, in an operon _yriA-yriB_	N	Growth defect (lower yield at saturation)	No effect
YadX	+	PhoQP-dependent, in an operon _yadX-clcA-yadW_	N	No effect	No effect
YkgS	-		N	No effect	Growth defect (slow growth) & increase in cell length during exponential phase
YriA	+	PhoQP-dependent, in an operon _yriA-yriB_	N	No effect	No effect
YdgU	+	PhoQP-dependent, in an operon _asr-ydgU_	Y	No effect	No effect
YqhI	+		N	No effect	No effect
YadW	+	PhoQP-dependent, in an operon _yadX-clcA-yadW_	N	No effect	No effect
DinQ	-		Y	No effect	Growth defect (slow growth)

1240 '+' indicates transcriptional upregulation and '-' indicates no change in transcriptional activity under magnesium stress vs. no stress; 'Y' indicates proteins predicted to
1241 contain membrane helices and shown to localize to the membrane, 'N' indicates a lack of membrane helices based on bioinformatic prediction (see Methods for details)