NuRD chromatin remodeling is required to repair exogenous DSBs in the *Caenorhabditis elegans* germline

Deepshikha Ananthaswamy[1], Kelin Funes[1†], Thiago Borges[1†], Scott Roques[1], Nina Fassnacht[2], Sereen El Jamal[2], Paula M. Checchi[2], Teresa Wei-sy Lee[1]*

[1] Department of Biological Sciences, University of Massachusetts Lowell, 198 Riverside Dr. Lowell MA, 01854

[2] Department of Biology, Marist College, 3399 North Road, Poughkeepsie, NY 12601

† contributed equally

* for correspondence, teresa_lee@uml.edu

Running title: NuRD remodeling repairs exogenous DSBs

Keywords: DNA repair, NuRD (nucleosome remodeling and deacetylase complex), LET-418/CHD4, FCD-2/FANCD2, germline

ABSTRACT

1
Organisms rely on coordinated networks of DNA repair pathways to protect genomes against toxic
double-strand breaks (DSBs), particularly in germ cells. All repair mechanisms must successfully
negotiate the local chromatin environment in order to access DNA. For example, nucleosomes can be
repositioned by the highly conserved Nucleosome Remodeling and Deacetylase (NuRD) complex. In
Caenorhabditis elegans, NuRD functions in the germline to repair DSBs – the loss of NuRD's ATPase
subunit, LET-418/CHD4, prevents DSB resolution and therefore reduces fertility. In this study, we
challenge germlines with exogenous DNA damage to better understand NuRD's role in repairing DSBs.
We find that *let-418* mutants are sensitive to cisplatin and hydroxyurea: exposure to either mutagen
impedes DSB repair, generates aneuploid oocytes, and reduces fertility and embryonic survival. These
defects resemble those seen when the Fanconi anemia (FA) DNA repair pathway is compromised, and
we find that LET-418's activity is epistatic to that of the FA component FCD-2/FANCD2. We propose a
model in which NuRD is recruited to the site of DNA lesions to remodel chromatin and allow access for
FA pathway components. Together, these results implicate NuRD in the repair of both endogenous DSBs
and exogenous DNA lesions to preserve genome integrity in developing germ cells.

16
17

ARTICLE SUMMARY

Preserving genome integrity in germ cells is critical for the survival of individuals and species. Our
previous work shows that nucleosome remodeling plays an important role in repairing meiotic DNA
damage. Here, we further challenge genomes with toxic DNA damaging agents to test the requirement
for remodeling in the germline. We find that DNA damage accumulates in the absence of remodeling,
which drastically reduces oocyte quality, and also show that the requirement for remodeling is epistatic
to the Fanconi anemia DNA repair pathway. These findings demonstrate that local chromatin
environments must be remodeled in response to DNA damage to maintain oocyte quality.

INTRODUCTION

DNA damage poses a danger to all cells, but if left unrepaired in germ cells, it is deleterious for organismal survival. During meiosis, any DNA lesion that persists until chromosome segregation will generate an aneuploid gamete, which in turn causes infertility or embryonic lethality (Ceccaldi et al., 2016; Kim et al., 2016). The evolutionary pressure to protect genomes for future generations has generated multiple distinct, but overlapping, DNA repair pathways. These pathways handle both endogenous programmed double-strand breaks (DSBs) required for crossover recombination, along with any exogenous DNA damage incurred in the germline (Chapman et al., 2012; Gartner and Engebrecht, 2022; Keeney, 2007). Studies in organisms ranging from budding yeast to mammals have identified strong conservation among these repair pathways, which include error-free mechanisms like homologous recombination and error-prone mechanisms like non-homologous end joining (De Massy, 2013; Stinson and Loparo, 2021, 2021; Wilson et al., 1997). For example, the Fanconi anemia (FA) pathway is activated when replication forks collide with DNA interstrand crosslinks, which are one of the most toxic DNA lesions (Gartner and Engebrecht, 2022). Components of the FA pathway coordinate the excision of the crosslink and conversion to DSBs, which can then be resolved by the meiotic homologous recombination pathway (Adamo et al., 2010; Lachaud et al., 2016; Raghunandan et al., 2015; Schlacher et al., 2012). Defects in the FA pathway are linked to an elevated risk of cancer and cause chromosomal instability or meiotic catastrophe, hallmarks of the rare congenital disorder of Fanconi anemia (Auerbach, 2009, 1993; Ceccaldi et al., 2016; De Winter and Joenje, 2009; Nalepa and Clapp, 2018; Tischkowitz et al., 2003, 2004).

In eukaryotes, DNA repair must always happen in the context of chromatin, where DNA is wrapped around histone octamers to form nucleosomes (Allis and Jenuwein, 2016). Nucleosomes act as obstacles that control access to DNA, and therefore their spacing is regulated by nucleosome remodeling complexes (Becker and Workman, 2013). The highly conserved remodeling complex, Nucleosome remodeling and deacetylase (NuRD), includes ATPase remodelers from the chromodomain helicase DNA binding (CHD) family (Basta and Rauchman, 2015; Kehle, 1998; Tong et al., 1998; Wade et al., 1998; Woodage et al., 1997; Xue et al., 1998). In addition to CHD remodeling subunits, NuRD also consists of the histone deacetylases HDAC1/2, histone demethylase LSD1/KDM1A, and other components that allow NuRD to bind modified histones, methyl CpG, and other proteins (Basta and Rauchman, 2015; Wang et al., 2009; Whyte et al., 2012). Initially, NuRD's primary role was thought to be transcriptional repression; but studies have since identified a more nuanced involvement in gene expression, along with other biological functions like DNA repair (Basta and Rauchman, 2015). The

nematode *Caenorhabditis elegans* has two catalytic CHD paralogs associated with the core NuRD

complex: CHD-3 (also known as Mi-2α or CHD3 in mammals) and LET-418 (Mi-2β or CHD4) (Passannante

et al., 2010; von Zelewsky et al., 2000). Mutations in either the *chd-3* or the *let-418* genes severely

reduce fertility due to high levels of persistent meiotic DSBs (McMurchy et al., 2017; Turcotte et al.,

2018).

These studies highlight the importance of NuRD's role in the germline for proper DNA repair –

the loss of NuRD activity appears to shunt DSBs from error-free repair towards error-prone pathways

like NHEJ. The genomic instability of *chd-3* or *let-418* mutants resembles those seen in mutants

defective for the FA pathway (Rageul and Kim, 2020; Tsui and Crismani, 2019; Turcotte et al., 2018). One

hallmark of mutations in FA components is an increased sensitivity to mutagens like cisplatin, which

induces DNA interstrand crosslinks, or hydroxyurea, which stalls replication forks (Bailly and Gartner,

2011; Datta and Brosh Jr., 2019; Youds et al., 2009). Both of these lesions can generate DSBs if left

unrepaired, which activate the S-phase checkpoint and subsequently delays cell cycle progression

(Kessler and Yanowitz, 2014; Kim and Colaiácovo, 2015). Resolution of interstrand crosslinks by the FA

pathway involves recognition and binding by the core FA complex, which then recruits FANCD2 to

chromatin by mono-ubiquitination (Nakanishi et al., 2005; Niraj et al., 2019). Once at the DNA lesion,

FANCD2 orchestrates the repair of the lesion by converting it to DSBs, which are then resolved by

homologous recombination (Ceccaldi et al., 2016; Niraj et al., 2019).

NuRD is required for repairing DNA lesions in germ cells (McMurchy et al., 2017; Turcotte et al.,

2018), but it is not clear how or whether nucleosome remodeling interacts with repair processes like the

FA pathway. Here, we challenge the germline by inducing exogenous DNA damage to investigate the

relationship between the FA pathway and NuRD's catalytic component, LET-418/CHD4. *C. elegans* has

homologs for many core FA components, including FCD-2/FANCD2 (Kim et al., 2018; Lee et al., 2010).

One advantage of using *C. elegans* for this study is the germline's spatiotemporal organization, which

allows us to easily follow the effects of mutagen exposure at multiple downstream stages during

oogenesis (Jaramillo-Lambert et al., 2007; Kessler and Yanowitz, 2014). We show that LET-418 activity is

epistatic to FCD-2 and is necessary to repair germline DSBs and preserve fertility. These results support a

model where NuRD generates a permissive local chromatin environment at the site of DNA damage to

allow access to DSB repair machinery and therefore preserve gamete quality in the face of genotoxic

stress.

METHODS

Strains and husbandry: All strains were cultured using standard methods (Brenner, 1974) at 20°C on 6-

cm NGM agar plates spotted with OP50 *E. coli* grown overnight at 37°C in Luria Broth (LB). Gene and

sequence information was downloaded from WormBase (Davis et al., 2022). The *syb6818* allele is a

deletion of LET-418's ATPase domain and removes amino acids 590-1114. The *syb6708* allele is a

deletion of LET-418's PHD finger domain and removes amino acids 256-365. Both domains were

identified using an NCBI structure prediction and based on homology with human CHD4 (Farnung et al.,

2020; Sayers et al., 2022). TER6 is a homozygous sterile strain, and was maintained over an *nT1* balancer

chromosome by picking GFP-positive heterozygotes. The following strains were obtained from the

Caenorhabditis Genetics Center or were generated by SunyBiotech (*syb* alleles, which were backcrossed

five times to N2):

- N2: wild-type (Bristol isolate)

- MT14390: *let-418 (n3536) V*

- NB105: *fcd-2 (tm1298) IV*

- TG1660: *xpf-1 (tm2842) II*

- TER14: *fcd-2 (tm1298) IV; let-418 (n3536) V*

- TER6: *let-418 (syb6818)/nT1 [qIs51] IV:V*

- TER10: *let-418 (syb6708) V*

Mutagen exposure: All exposure assays were performed on hermaphrodites cultured at 20°C on 3-cm

NGM agar plates. Plates were first spotted with OP50 *E. coli*, then treated by spreading 250 μL of either

400 μM cisplatin (Sigma Aldrich) or 10 mM hydroxyurea (Thermo Scientific) onto agar at least 12 hours

prior to experiment. Synchronized young adult hermaphrodites (12 hours post-L4 larval stage) were

placed on mutagen plates for 24 hours, then moved to regular NGM plates to recover before assessing

nuclei in the appropriate meiotic stage; the timing for each assay is described below (Jaramillo-Lambert

et al., 2007; Kessler and Yanowitz, 2014).

Immunofluorescence and image processing: To assess RAD-51 foci in the mitotic region after mitotic

exposure, exposed and control hermaphrodites were allowed to recover on NGM plates for four hours

before dissection to produce better gonad extrusion. To assess RAD-51 foci in pachytene after mitotic

exposure, hermaphrodites were allowed to recover for 30-40 hours, allowing nuclei to progress to mid-

to late-pachytene. Gonad dissection, fixation, and immunostaining were performed as previously

described (Ananthaswamy et al., 2022). In brief, gonads were dissected using 25-gauge syringe needles,

fixed in 0.8% paraformaldehyde (Electron Microscopy Services) for five minutes at room temperature,

subjected to freeze crack, and dehydrated in 100% methanol for one minute at -20°C. Slides were

washed three times with PBS + 0.1% Tween-20 (Thermo Scientific) prior to overnight incubation at room

temperature with primary antibody. Slides were washed three times before incubation with secondary

antibody for four hours at room temperature. Samples were mounted in VectaShield (Vector

Laboratories) with 2 µg/ml DAPI (Thermo Life) before sealing with nail polish. Antibodies used were

rabbit anti-RAD-51 at 1:5000 (a gift from Diana Libuda (Kurhanewicz et al., 2020)) and donkey anti-rabbit

AlexaFluor594 at 1:400 (Invitrogen).

Immunofluorescence images were collected at 63x as Z-stacks with 0.25 µm intervals using a

Zeiss Axio Imager M2 microscope with Zen 3.8 software (Zeiss). All representative images shown are

max projections after processing in FIJI software (Schindelin et al., 2012). Unprocessed Z-stacks were

used to quantify RAD-51 foci. Three biological replicates were performed, with three to five gonads

imaged per condition in each replicate. Only one gonad arm was analyzed from each hermaphrodite. In

the mitotic region, all nuclei were scored for RAD-51 foci. In pachytene, only nuclei in the top row were

scored for RAD-51 foci; pachytene regions were determined by counting the number of rows after the

transition zone and before diplotene, and dividing into equal thirds. Some nuclei had a distinctive

apoptotic appearance with collapsed, DAPI-bright chromosomes and an excess of RAD-51 staining; these

were included in the total nuclei number, but not scored for RAD-51 foci. All statistical analyses were

performed in Prism 8 (GraphPad), with *P*-values determined using a Kruskal-Wallis non-parametric test

after a Dunn's post-hoc test used to account for pairwise comparisons.

Analysis of diakinesis DAPI bodies: Following exposure to cisplatin or hydroxyurea, hermaphrodites were

allowed to recover for 60 hours, allowing mitotic oocyte nuclei to progress through meiosis to diakinesis.

Gonads were dissected as previously described (Ananthaswamy et al., 2022) with the following

modifications: the primary and secondary antibody incubation steps were skipped, and slides were

washed at least three times in PBS + 0.1% Tween-20 (Thermo Scientific) before mounting in SlowFade

Diamond (Molecular Probes) with 2 µg/ml DAPI (Thermo Life). Images were collected at 63x as Z-stacks

with 0.25 µm intervals captured on Zeiss Axio Imager M2 using Zen 3.8 software (Zeiss). All

representative images shown are max projections after processing in FIJI (Schindelin et al., 2012).

Unprocessed z-stacks were used to quantify DAPI-stained bodies in the -1 to the -4 oocyte from the

spermatheca. Three biological replicates were performed, with each condition consisting of a minimum

of 100 nuclei from at least 27 hermaphrodites. All statistical analyses were performed in Prism 10

(GraphPad), with p-values determined using a Fisher's exact test on a 2x5 contingency table.

<u>Embryonic survival – 4-hr laying period:</u> After exposure to cisplatin, hydroxyurea, or nitrogen mustard,

hermaphrodites were allowed to recover on NGM plates for 60 hours, allowing mitotic oocyte nuclei to

progress through the gonad and be laid as fertilized embryos. Exposed mothers were then moved onto

freshly spotted 3-cm NGM plates (five to ten mothers per plate) and allowed to lay for four hours before

being removed from plates. The number of embryos per plate was immediately recorded after removal

of mothers, and were scored as fertilized, unfertilized, or dead. Hatched progeny were counted as adults

three days later. Percent survival was calculated by dividing the number of hatched progeny by the

number of embryos, multiplied by 100. Each condition consists of progeny from 125-150 mothers,

collected over three biological replicates. All statistical analyses were performed in Prism 8 (GraphPad),

with p-values determined from a one-way ANOVA after a Šídák correction used to account for multiple

comparisons.

<u>Embryonic survival – entire brood:</u> Individual hermaphrodites were isolated on separate plates as L4

larvae and transferred every 12 hours until the end of their fertile period (usually 5-6 days). Embryos

and unfertilized eggs were counted every 12 hours immediately after transfer of the mother. Total

brood was calculated for each mother as all embryos laid during the fertile period. Hatched progeny

were then counted two to three days later (when they were L4s or young adults). Percent survival was

calculated by dividing the number of hatched progeny by the number of embryos laid and multiplied by

100. Each biological replicate includes the broods of five mothers scored for each genotype (i.e. five

technical replicates). The following biological replicates were performed: seven of N2; six of *let-418*

(n3536); four of *let-418 (syb6818)*; and four of *let-418 (syb6708)*. Broods were censored from analysis if

the mother died of unnatural causes before the end of their fertile period (usually from vulva rupture or

crawling off the agar). All statistical analyses were performed in Prism 8 (GraphPad), with p-values

determined from a one-way ANOVA after a Šídák correction used to account for multiple comparisons.

RESULTS

<u>LET-418 activity is required to repair mitotic DNA lesions in mitosis and meiosis.</u>

Mutants lacking both NuRD paralogs (CHD3 and CHD4) have severely compromised fertility,

which makes downstream analyses difficult (Turcotte et al., 2018). Therefore, we focused only on the

impact of LET-418/CHD4, reasoning that it plays a larger role in repairing germline DSBs because its loss

causes more meiotic defects than that of CHD-3 (Turcotte et al., 2018). A previous study noted that at

pachytene entry, *let-418* mutants have more DSBs than wild-type (Turcotte et al., 2018), leading us to

wonder whether these DSBs represent unrepaired DNA damage acquired during replication. Because

DSBs are rare in mitotic nuclei, we used cisplatin and hydroxyurea to generate exogenous DNA damage

and further stress the DNA repair process. We examined *let-418 (n3536)* missense hypomorphic

mutants and *fcd-2 (tm1298)* deletion mutants, which lack the FA pathway component FCD-2/FANCD2.

We capitalized on the fact that the *C. elegans* germline is organized in a spatiotemporal manner, with

nuclei undergoing mitosis at the distal tip and moving proximally as they proceed through meiosis. The

timing of how nuclei progress through mitotic replication, meiosis, oocyte maturation, and fertilization

has been extensively characterized (Jaramillo-Lambert et al., 2007; Kessler and Yanowitz, 2014). We

therefore estimated the timing after exposure to capture the impact of DNA damage incurred during

mitosis.

Mitotic zone DSBs were assessed using an antibody for the RecA recombinase RAD-51, a

common cytological marker for DSBs (Alpi et al., 2003; Colaiácovo et al., 2003; Kurhanewicz et al., 2020).

Hermaphrodites were cultured on either cisplatin or hydroxyurea for twenty hours, allowed to recover

for four hours, and dissected for immunofluorescence (Jaramillo-Lambert et al., 2007; Kessler and

Yanowitz, 2014). Consistent with prior reports, unexposed wild-type animals accumulated few RAD-51

foci in the mitotic zone (Fig. 1A, C, Fig. 2A, C, and Table S1) (Alpi et al., 2003; Turcotte et al., 2018), which

was also true for all mutants examined, with foci in less than 8% of nuclei. Unexposed *let-418* single

mutants had a modest but significant increase in foci compared to wild-type animals ($P < 0.0001$,

Kruskal-Wallis test), but neither *fcd-2* single mutants nor *fcd-2; let-418* double mutants had more foci

than wild-type (Fig. 1A, $P > 0.08$ for both comparisons, Kruskal-Wallis test). Unexposed *fcd-2; let-418*

double mutants were not more affected than either single mutant ($P > 0.8$ for both comparisons,

Kruskal-Wallis test). We note that, although the average number of foci did not vary much between

genotypes, we occasionally observed nuclei with two or three foci in *let-418* single mutants (0.8% of

nuclei) and *fcd-2; let-418* double mutants (0.7%), which were rarely seen in wild-type germlines (0.2%)

(Fig. 1A and Table S1).

As expected, exposure to 400 µM cisplatin increased the number of RAD-51 foci across all

genotypes compared to unexposed controls (Fig. 1B, D and Table S1). This dose was chosen based on a

dose curve showing that 400 µM drastically reduced embryonic survival in both wild-type and mutant

populations (Fig. S1 and Table S6). Wild-type populations exposed to cisplatin accumulated over five

times more foci than their unexposed controls *P* < 0.0001, Kruskal-Wallis test) (Fig. 1A, B). Similarly, both exposed *fcd-2* and *let-418* mutants accumulated five times more foci than unexposed controls (*P* < 0.0001 for both comparisons, Kruskal-Wallis test) (Fig. 1A, B). In *let-418* mutants, the increase was driven by nuclei with multiple foci, sometimes as many as five per nucleus (10.8% of nuclei compared to 0.7%). Interestingly, exposed *fcd-2; let-418* double mutants strongly resembled exposed *let-418* single mutants, with over seven times more foci than unexposed controls (*P* < 0.0001, Kruskal-Wallis test) (Fig. 1B). Again, this increase was primarily driven by nuclei with multiple foci, especially those with four or five foci (9.6% of nuclei compared to 0.7%). Notably, nuclei with four or five foci were never observed in exposed wild-type or *fcd-2* single mutants. All mutants examined were also more sensitive to cisplatin than wild-type controls. Exposed *fcd-2* single mutants experienced 1.7 times more foci than exposed wild-type populations (*P* < 0.0001), whereas exposed *let-418* single mutants were even more affected, with 2.4 times more foci than exposed wild-type populations (*P* < 0.0001 compared to WT, *P* < 0.05 compared to *fcd-2*). The increase in exposed *fcd-2; let-418* double mutants resembled that seen in *let-418* single mutants (*P* > 0.9), with the double mutants having 2.7 times more foci than exposed wild-type populations (*P* < 0.0001).

We next challenged mutants using hydroxyurea, which stalls replication forks and generates DSBs if left unrepaired (Bailly and Gartner, 2011). A dose of 10 mM was chosen after examining embryonic survival in wild-type and *let-418* mutants across a dose curve (Fig. S3 and Table S6). Hydroxyurea affected all genotypes in a manner similar to cisplatin (Fig. 2 and Table S1). Once again, few DSBs were observed in each unexposed population, with less than 7% of mitotic nuclei containing a RAD-51 focus (Fig. 2A). As expected, exposure to hydroxyurea raised the average number of foci in all genotypes (*P* < 0.02 for all comparisons, Kruskal-Wallis test) (Fig. 2B). Exposed wild-type populations and *fcd-2* mutants had nearly two times more foci than unexposed controls (*P* < 0.02 for both comparisons, Kruskal-Wallis test), both primarily driven by an increase in the number of nuclei with one focus. Resembling what we observed after cisplatin exposure, both single mutants were more sensitive to hydroxyurea than wild-type populations: exposed *fcd-2* single mutants had 1.6 times more foci than exposed wild-type controls (*P* < 0.05) and *let-418* single mutants had twice as many foci as exposed wild-type populations (*P* < 0.0001). As with cisplatin exposure, the response in *fcd-2; let-418* double mutants resembled the response in *let-418* single mutants (*P* > 0.9), with 2.6 times more foci than exposed wild-type populations (*P* < 0.0001). We note that exposed *let-418* and *fcd-2; let-418* mutants each contained nuclei with as many as five RAD-51 foci, which were almost never observed in exposed wild-type or *fcd-2* mutant populations (Fig. 2B).

To examine whether these unrepaired exogenous DSBs persisted from mitosis into meiosis, we

assessed RAD-51 foci throughout pachytene (Fig. 3 and Table S2). Hermaphrodites were exposed to

cisplatin or hydroxyurea for twenty hours, allowed to recover for 30-40 hours (to allow mitotic nuclei to

progress into mid- or late-pachytene), and dissected for immunofluorescence (Jaramillo-Lambert et al.,

2007; Kessler and Yanowitz, 2014). At mid-pachytene, wild-type animals had an average of 3.3 foci per

nucleus, which is consistent with prior observations (Fig. 3A, B) (Achache et al., 2021; Cahoon et al.,

2023; Colaiácovo et al., 2003; Germoglio et al., 2020; Rosu et al., 2013; Stamper et al., 2013). All

unexposed mutants had significantly more RAD-51 foci than unexposed wild-type animals in mid-

pachytene ($P < 0.001$ for all mutants, Kruskal-Wallis test; see Table S2 for additional comparisons

between early-pachytene and late-pachytene), and the number of foci in *fcd-2; let-418* double mutants

resembled that in both single mutants ($P > 0.9$) (Fig. 3A, B).

Exposing mitotic nuclei to cisplatin significantly increased the number of foci during mid- or late-

pachytene across all genotypes compared to unexposed controls ($P < 0.0001$ for all comparisons) (Fig.

3C, D). Notably, all mutants had persistent damage even in late-pachytene (Fig. 3C and Table S2) ($P <$

0.05 for all genotypes compared to wild-type) – both *let-418* single mutants and *fcd-2; let-418* double

mutants were more affected than *fcd-2* single mutants, with each experiencing an average of over five

foci per nucleus in late-pachytene ($P < 0.05$ for each compared to *fcd-2*, $P > 0.9$ for double mutant

compared to *let-418*). Similarly, exposure to hydroxyurea significantly increased foci during mid-

pachytene in both *let-418* single mutants and *fcd-2; let-418* double mutants compared to unexposed

controls ($P < 0.0001$ for both comparisons) (Fig. 3E, F), and this increase persisted into late-pachytene

when compared to exposed wild-type animals or *fcd-2* single mutants ($P < 0.0001$ for all comparisons)

(Fig. 3E and Table S2).

Taken together, these data indicate that cisplatin or hydroxyurea generate mitotic DNA lesions

that require both LET-418 and FCD-2 for proper repair. In their absence, exogenous damage incurred

during mitosis persists until the end of pachytene. Additionally, when responding to mitotic DNA

damage in the germline, *fcd-2; let-418* double mutants phenocopy *let-418* single mutant for both

mutagens.

<u>Unrepaired mitotic DSBs affect oocyte quality in *let-418* mutants.</u>

Although *let-418* single mutants accumulate DSBs in the mitotic region and during pachytene,

they do not display aneuploidy or chromosome fragmentation, indicating that any remaining DSBs are

fully resolved by diakinesis (Turcotte et al., 2018). However, mutants lacking both CHD paralogs, CHD-3

and LET-418, suffer from meiotic catastrophe, genome fragmentation, and chromosome fusions

(Turcotte et al., 2018), suggesting that NuRD needs at least one CHD component to repair endogenous

DSBs. To see whether LET-418 is required for the resolution of exogenous DNA lesions, we exposed

hermaphrodites to cisplatin or hydroxyurea for twenty hours and allowed them to recover for sixty

hours (Jaramillo-Lambert et al., 2007; Kessler and Yanowitz, 2014). This timing allows us to induce DNA

damage in mitotic cells and assess its impact during diakinesis, where we expect to see six DAPI-staining

bodies, one for each homologous pair of chromosomes.

As expected, unexposed wild-type animals rarely produced aneuploid gametes: when scoring

aged mothers in their third day of adulthood, we found that only 9% of oocytes contained five DAPI-

stained bivalents (Fig. 4A, C, E, Fig. S2, and Table S3). This result is consistent with previous studies

showing that oocyte quality decreases with maternal age (Achache et al., 2021; Andux and Ellis, 2008;

Luo et al., 2010; Scharf et al., 2021). Each unexposed mutant produced more aneuploid oocytes than

wild-type controls, with at least 16% of nuclei containing fewer than six DAPI bodies ($P < 0.01$ for all

comparisons, Fisher's exact test), including oocytes with four DAPI bodies that were never observed in

wild-type animals (Fig. 4A, C, E and Fig. S2). We also noted that *let-418* single mutants generated a rare

oocyte with seven DAPI bodies (Fig. 4E). The presence of univalents in this nucleus reflected the lack of a

chiasmata and failure of crossover formation, which almost never occurs in wild-type animals (Dernburg

et al., 1998; Yu et al., 2016). When exposed to 400 μM cisplatin, both *fcd-2* single mutants and *fcd-2; let-*

418 double mutants suffered a significant deterioration in oocyte quality ($P < 0.04$ for both comparisons,

Fisher's exact test) (Fig. 4B). Each mutant produced oocytes with more than six DAPI bodies, and *fcd-2;*

let-418 double mutants also produced an oocyte with only three DAPI bodies (Fig. 4F). Although

exposure to cisplatin did not induce a statistically significant change in wild-type and *let-418* single

mutants ($P > 0.5$ for both comparisons, Fisher's exact test), both genotypes produced oocyte classes that

were never observed in unexposed controls (Fig. 4E, F and Table S3).

When compared to either single mutant, *fcd-2; let-418* double mutants were more affected by

cisplatin exposure ($P < 0.003$ for both comparisons, Fisher's exact test). This difference was driven by

oocytes with both more and fewer DAPI bodies: *fcd-2; let-418* double mutants produced more nuclei

with four or fewer DAPI bodies, but also nuclei with seven to ten DAPI bodies, indicating a severe

disruption in crossover formation (Fig. 4F). Given our findings for mitotic DSBs, where the double

mutant phenocopied the *let-418* single mutant (Fig. 1 and Fig. 2), we were surprised to see that

diakinesis DAPI body number was significantly more affected in *fcd-2; let-418* double mutants than in

let-418 single mutants. However, both the double mutant and *let-418* single mutant produced the rare

category of nuclei with three DAPI bodies (Fig. 4F). The severe reduction in DAPI bodies indicated that at

least half of the *C. elegans* genome has experienced chromosomal fusions, a phenotype that was also

previously observed in mutants missing both CHD paralogs (Turcotte et al., 2018).

Similar to cisplatin, exposure to 10 mM hydroxyurea also induced aneuploidy during diakinesis

(Fig. 4C, D and Table S3). Although exposure to hydroxyurea did not cause a statistically significant

change in wild-type population or *fcd-2* single mutants ($P > 0.3$ for both comparisons, Fisher's exact

test), both genotypes produced oocyte classes that were never observed in unexposed controls (Fig. S2

and Table S3). However, unlike what we observed after cisplatin exposure, we did not find that *fcd-2*

single mutants produced nuclei with more than six DAPI bodies after hydroxyurea exposure (Fig. 4D),

indicating that *fcd-2* mutants were more sensitive to cisplatin. Exposed *let-418* single mutants were

significantly affected compared to wild-type controls, producing oocytes with both more than seven and

fewer than four DAPI bodies, categories that were never observed in unexposed controls ($P < 0.006$,

Fisher's exact test).

Challenging *fcd-2; let-418* double mutants with hydroxyurea caused a large deterioration in

oocyte quality ($P < 0.009$, Fisher's exact test). Exposed *fcd-2; let-418* double mutants were significantly

more affected than either of the exposed single mutant controls ($P < 0.004$ for both comparisons,

Fisher's exact test). This decline in oocyte quality was partly driven by nuclei with more than six DAPI

bodies (sometimes as many as ten), a category that was only ever observed in exposed *fcd-2; let-418*

double mutants and *let-418* single mutants (Fig. S2B and Table S3). Once again, the existence of nuclei

with univalents indicates that hydroxyurea also causes a severe failure of crossover formation or

chromosome fragmentation (Fig. S2B). Taken together, this assessment of diakinesis nuclei indicates

that *fcd-2; let-418* double mutants are more sensitive to interstrand crosslinks and stalled replication

forks than either single mutant.

<u>Mutagen exposure reduces embryonic survival of *let-418* mutants.</u>

After having shown that LET-418 is necessary to repair exogenous DSBs during mitosis and

meiosis (Fig. 1 through 3) and prevent aneuploidy during diakinesis (Fig. 4), we next examined oocyte

viability by assessing embryonic survival (Fig. 5 and Table S4). To capture the impact of mutagen

exposure during mitosis, we assessed embryos 60 hours after exposure, which were produced by

mothers in their third day of adulthood (Jaramillo-Lambert et al., 2007; Kessler and Yanowitz, 2014). As

expected for aged mothers, unexposed wild-type populations had an average embryonic survival of 96%

(Fig. 5A). As a positive control for cisplatin sensitivity, we examined *xpf-1 (tm2842)* mutants, which lack

an endonuclease involved in the DNA damage response (Ward et al., 2007). Unexposed *xpf-1* mutants

have a lower average survival than unexposed wild-type populations ($P < 0.007$, ANOVA) (Fig. 5A), which

is reduced even further by cisplatin in a dose-dependent manner (Fig. S1B and Table S6) (Meier et al.,

2014; Ward et al., 2007). Survival in unexposed *fcd-2*, *let-418*, or *fcd-2; let-418* mutants did not

significantly differ from wild-type populations ($P > 0.1$ for all comparisons), although each mutant

displayed more variability and lower averages across replicates compared to wild-type: across both

cisplatin and hydroxyurea experiments, *fcd-2* single mutants averaged 86% survival, *let-418* single

mutants averaged 77% survival, and *fcd-2; let-418* double mutants averaged 75% survival (Table S4).

When challenged with 400 µM cisplatin, embryonic survival in wild-type populations was significantly

reduced to 73% ($P < 0.04$). As expected for a positive control, cisplatin exposure reduced *xpf-1* mutant

survival more severely than in wild-type populations, down to 39% survival ($P < 0.01$) (Fig. 5A).

Embryonic survival in exposed *fcd-2* single mutants was not significantly affected compared to

unexposed controls ($P > 0.8$, ANOVA) (Fig. 5A). Conversely, exposed *let-418* single mutants experienced

a significant decline in survival, down to 39% ($P < 0.0001$), a decrease 2.4 times more than wild-type and

similar to *xpf-1* mutants (Fig. 5A). We noted that cisplatin had a variable impact on *let-418* mutant

survival, across both technical and biological replicates. This observation was consistent with previous

variability observed in *let-418* mutants (Turcotte et al., 2018): some replicates were completely

unaffected, resembling unexposed controls, whereas others suffered complete embryonic lethality that

was never observed in wild-type or *xpf-1* positive controls. Again, the effects of cisplatin on embryonic

survival in *fcd-2; let-418* double mutants strongly resembled *let-418* single mutants (Fig. 5A). Exposed

fcd-2; let-418 double mutants experienced a significant drop in survival down to 33% ($P < 0.0001$). As

seen in *let-418* single mutants, this reduction was 2.7 times more than wild-type and similar to *xpf-1*

positive controls. However, in one departure from *let-418* single mutants, we found that cisplatin caused

less variability in *fcd-2; let-418* double mutant survival, which had a distribution that overlapped only the

lower half of that observed in *let-418* single mutants (Fig. 5A).

We next challenged hermaphrodites with 10 mM hydroxyurea to assess the impact on

embryonic survival (Fig. 5B). As expected, this low dose of exposure did not significantly reduce survival

in wild-type populations or *fcd-2* single mutants ($P > 0.09$ for both comparisons, ANOVA) (Kim et al.,

2018). However, *let-418* single mutants were sensitive to hydroxyurea – embryonic survival dropped

from 76% in unexposed controls to 46% after exposure ($P < 0.0003$), a reduction 2.1 times more than

experienced by wild-type populations (Fig. 5B). Similarly, survival in exposed *fcd-2; let-418* double

mutants dropped from 73% down to 36% ($P < 0.0001$), a reduction 2.8 times more than wild-type ($P <$

0.0001) (Fig. 5B).

Finally, we challenged hermaphrodites with an additional mutagen, nitrogen mustard, which

creates both interstrand crosslinks and DNA-protein crosslinks (Fig. S4 and Table S6) (Povirk and Shuker,

1994). A low dose of 100 μM did not affect wild-type embryonic survival ($P > 0.9$, ANOVA), whereas a

higher dose of 150 μM significantly decreased wild-type survival to 52% ($P < 0.0001$) (Fig. S4). The low

100 μM dose of nitrogen mustard decreased survival in both *let-418* and *fcd-2* single mutants ($P <$

0.0001 for both comparisons) and the high 150 μM dose further reduced survival to 25% ($P < 0.0001$ for

both comparisons, ANOVA). Nitrogen mustard affected *fcd-2; let-418* double mutants similarly to each

single mutant ($P > 0.6$ for all comparisons), with survival severely reduced to 10% by the high 150 μM

dose ($P < 0.0001$ compared to unexposed control) (Fig. S4). Taken together, the impact of mutagen

exposure on embryonic survival indicate that LET-418 is required to resolve exogenous DNA lesions to

maintain oocyte viability.

<u>LET-418's ATPase domain is necessary for its role in germline DSB repair.</u>

The protein sequences of *C. elegans* LET-418 and human CHD4 share extensive homology,

including three domains: an ATPase domain, a PHD finger domain, and a double chromodomain (Fig. 6A)

(Farnung et al., 2020; Käser-Pébernard et al., 2016; Passannante et al., 2010). Computational models of

CHD4 structure suggest that each domain has distinct functions: the ATPase domain slides nucleosomes

along DNA (Farnung et al., 2020), the PHD finger domains binds specific residues on histone H3 (Farnung

et al., 2020; Mansfield et al., 2011) and the chromodomains primarily bind DNA (Bouazoune et al.,

2002). Mutations discovered in human patients, along with *in vitro* experiments of CHD4 activity, have

implicated each of these domains in CHD4's nucleosome remodeling activity (Farnung et al., 2020;

Watson et al., 2012; Weiss et al., 2016), but the impact of each domain has yet to be assessed *in vivo*.

Because the kinetics of repair occur rapidly in response to DNA lesions (Kochan et al., 2017; Nair et al.,

2017), we hypothesized that LET-418's function in the germline requires its ATPase domain to slide

nucleosomes and expose the site of DNA damage.

Figures 1 through 5 of this paper relied on the canonical allele *let-418 (n3536)*, a point mutation

in the ATPase domain that is hypomorphic, but viable, at the normal husbandry temperature of 20°C

(Andersen et al., 2006; Turcotte et al., 2018). We engineered deletion alleles in the *let-418* gene using

CRISPR-Cas9 to remove either the PHD finger domain or the ATPase enzymatic domain (see Methods for

details). We then assessed the number of embryos produced throughout the entire laying period in each

let-418 mutant (a full brood) under normal husbandry conditions, using the canonical *n3536* allele as a

positive control (Fig. 6A and Table S5). Consistent with previous studies, wild-type mothers produced

broods that averaged 311 embryos (Fig 6C). Strikingly, mutants that lack the ATPase domain*, syb6818*,

were completely sterile and produced no embryos (Fig. 6C). In these ATPase deletion mutants, we also

noted a high occurrence of protruding vulvae and a clear body due to absence of germline, which

accounts for the complete sterility (Fig. 6B). Conversely, mutants that lack the PHD finger domain,

syb6708, produced an average brood of 304 embryos, which was not significantly lower than wild-type

broods ($P > 0.7$, ANOVA). Consistent with previous reports, average broods of canonical *n3636* mutants

were 188 embryos, which represents a significant reduction compared to wild-type broods ($P < 0.001$).

We also noted the variability in canonical *n3636* mutant broods: some mothers produced at wild-type

levels (~300 embryos), whereas others were completely sterile (McMurchy et al., 2017; Turcotte et al.,

2018).

Because LET-418 is also involved in development (Erdelyi et al., 2017; Passannante et al., 2010;

von Zelewsky et al., 2000), we assessed embryonic survival throughout the full laying period. Average

survival of canonical *n3636* mutants was 92%, which represented a slight but significant reduction

compared to average wild-type survival of 98% ($P < 0.01$, ANOVA). In *syb6708* PHD deletion mutants,

embryonic survival resembled wild-type survival at 99%, similar to what we observed with brood size (P

> 0.9) (Fig. 6D). Finally, we were unable to assess survival in *syb6818* ATPase deletion mutants, because

all mothers examined were sterile (N = 20). Altogether, our findings support a model in which LET-418's

ATPase domain, but not its PHD domain, is necessary for NuRD's role during oogenesis in otherwise

wild-type conditions. Future studies will examine whether the PHD domain is necessary for repairing

exogenous DNA damage.

DISCUSSION

The germline must preserve genome integrity for all future generations, a task that requires

extensive coordination between DNA repair machinery and local chromatin environments (Chen and

Tyler, 2022; Clouaire and Legube, 2019). At the site of DNA damage, chromatin landscapes help

orchestrate repair using either error-prone or error-free DSB repair pathways (Chen and Tyler, 2022).

Although the mechanisms of DNA repair have been well-described across species, the role of chromatin

remodelers remains poorly understood. In this study, we show that LET-418, the ATPase remodeling

subunit of NuRD, is required to maintain oocyte quality when genomes are challenged by exogenous

DNA damage. After exposure to cisplatin or hydroxyurea, mutants with reduced LET-418 activity

accumulate more DSBs in the germline (Figs. 1-3), generate more aneuploid oocytes (Fig. 4), and suffer

from reduced embryonic viability (Fig. 5 and Fig. 6) – phenotypes that strongly resemble those seen in

mutants defective for the FA repair pathway (Adamo et al., 2010; Gartner and Engebrecht, 2022; Lee et

al., 2010). When testing the genetic relationship between NuRD and the FA pathway, we showed that

let-418 gene activity is epistatic to the FA gene *fcd-2*, which suggests that NuRD nucleosome remodeling

is necessary to allow proper functioning of the FA repair pathway.

LET-418/CHD4 has been implicated in biological functions that include maintaining repressive

chromatin states, repairing DNA damage, and defining cell fate during embryonic development (De Vaux

et al., 2013; Guerry et al., 2007; Käser-Pébernard et al., 2016; Kunert et al., 2009; Saudenova and Wicky,

2018; Turcotte et al., 2018; Unhavaithaya et al., 2002; von Zelewsky et al., 2000). In mammals, CHD4's

role in resisting DNA damage has led to its upregulation in multiple types of cancer: for example, in

glioblastoma, CHD4 overexpression is associated with poor prognosis, likely due to its role in promoting

DNA damage repair to assure cancer cell survival (Chudnovsky et al., 2014; McKenzie et al., 2019). Other

work in ovarian cancer cells, acute myeloid leukemia cells, and mammary cells has shown that depleting

CHD4 impairs homologous recombination and sensitizes tumorigenic cells to DNA damaging agents (Pan

et al., 2012; Polo et al., 2010; Sperlazza et al., 2015). Similarly, in *C. elegans*, mutations in LET-418 or its

paralog CHD3 prevent DSB repair throughout meiotic pachytene, suggesting that LET-418/CHD4's role is

highly conserved across taxa (Turcotte et al., 2018).

Our work identifies a new role for LET-418/CHD4 in response to genotoxic threats – in the

absence of LET-418's ATPase activity, mutants cannot fully repair exogenous DNA damage incurred

during mitosis, which ultimately reduces gamete viability and impairs organismal survival. In germlines,

LET-418 is associated with other heterochromatin factors that work with small RNA pathways to silence

repetitive elements and protect the genome (McMurchy et al., 2017). The DSB repair defects observed

in *let-418* mutants may be caused by disruptions in heterochromatin formation or maintenance, leading

to higher sensitivity to DNA damaging agents. A similar effect has been observed with the loss of

another core component of NuRD, RBBP4 (LIN-53 in *C. elegans*), which sensitizes cells to DNA damaging

agents by establishing a permissive chromatin environment and downregulating RAD51 expression

(Kitange et al., 2016).

We propose that LET-418 and NuRD are required during at least two points in gametogenesis: to

repair exogenous damage accumulated in mitosis and to resolve endogenous DSBs formed during

meiosis (this study and Turcotte et al., 2018). In the absence of LET-418 or NuRD activity, mitotic

damage persists as nuclei enter meiosis, contributing to the increase of DSBs observed in pachytene

(this study and Turcotte et al., 2018). The accumulated damage persists until the end of pachytene, and

is reflected in aberrant diakinesis oocytes. The increase in meiotic DSBs may be caused by an inability of

the FA pathway to access the damage. During meiosis, the FA pathway continues to repair DNA damage

caused by replication fork stalling or interstrand crosslinks – after the lesion is recognized, FA

component FANCD2 is mono-ubiquitinated and recruited to the lesion, where it mediates a conversion

to DSBs and repair by homologous recombination (Ceccaldi et al., 2016; Niraj et al., 2019). Therefore,

the loss of LET-418 or NuRD activity may interfere with recognition of the lesion or with recruiting FCD-2

to the site of damage.

Cytological experiments show that FA components, including FCD-2/FANCD2, are recruited to

DNA lesions to mediate repair in the mitotic region of the germline (Collis et al., 2006; Kim et al., 2018;

Lee et al., 2010). Our results are consistent with FCD-2 playing a role in resolving mitotic DSBs – when

exposed to mutagens, *fcd-2* single mutants generated aneuploid oocytes that were never observed in

wild-type populations: some had chromosome fusions, whereas others experienced crossover failure

(Fig. 4 and Fig. S2). These defects represent genomic instability that is a consistent hallmark of defects

with the FA pathway, and is even used to diagnose human patients for Fanconi anemia (Adamo et al.,

2010, 2010; Auerbach, 2009; De Winter and Joenje, 2009; Lee et al., 2007; Rageul and Kim, 2020).

However, we were surprised to find that embryonic viability in *fcd-2* mutants was not affected by

exposure to either cisplatin or hydroxyurea (Fig. 5), since others have previously reported a cisplatin

sensitivity in this mutant background (Adamo et al., 2010; Collis et al., 2006; Germoglio et al., 2020).

After examining the timing of exposure and recovery used in those studies, we realized that they

primarily assessed how damage induced during late pachytene affected embryonic survival (Jaramillo-

Lambert et al., 2007; Kessler and Yanowitz, 2014). In contrast, we used a time course that allowed us to

examine the effects of damage induced earlier, during mitosis. Altogether, our results suggest that FCD-

2 and the FA pathway are involved in repairing exogenous DSBs, but may not be the sole, or even the

main, repair pathway at this point in gametogenesis. However, in the later stages of meiotic prophase,

all DSBs must be resolved before chromosome segregation. Therefore, as nuclei approach diplotene,

they rely on alternative error-prone DSB repair pathways like NHEJ to resolve remaining DSBs (Macaisne

et al., 2018; Smolikov et al., 2007). These pathways usually result in non-crossovers and disruption of

chiasmata formation, or can generate chromosome translocations or fusions (Gartner and Engebrecht,

2022); we observed both outcomes in diakinesis nuclei of *let-418* mutants.

We have also demonstrated that LET-418's activity is epistatic to that of FCD-2 and the FA

pathway – the *fcd-2; let-418* double mutant phenocopies the *let-418* single mutant when challenged by

exogenous DNA damage in its repair of DSBs and embryonic survival (Fig. 1-3, and Fig. 5). When

examining how mitotic DNA lesions affect diakinesis DAPI body formation, we observed that *fcd-2; let-*

418 double mutants were more severely affected by cisplatin and hydroxyurea than either single mutant

(Fig. 4 and Fig. S2). Strikingly, double mutant germlines contained classes of diakinesis nuclei that were

only seen in one or the other single mutant, suggesting an additive effect of losing both LET-418 and

FCD-2 activity. This result indicates that the FA pathway and NuRD may function in separate DSB repair

pathways. The use of distinct pathways may be influenced by the region of the germline in which

damage was incurred – in addition to the exogenous damage in mitosis caused by mutagen exposure,

let-418 mutants also accumulate damage endogenous DSBs during pachytene, which are not further

exacerbated by the loss of FCD-2 in unexposed double mutants (Fig. 4A, C) (Turcotte et al., 2018).

514   *C. elegans* LET-418 shares extensive homology with human CHD4, including an ATPase domain, a

PHD finger domain, and a double chromodomain (Käser-Pébernard et al., 2016; Passannante et al.,

2010). Computational models indicate that the ATPase domain repositions nucleosomes by sliding them

along DNA (Farnung et al., 2020). When examining human patient mutations identified in cancer or the

neurodevelopmental disorder Sifrim-Hitz-Weiss syndrome, we noticed that many missense mutations

occurred in the ATPase domain, indicating its importance for LET-418 and NuRD function. (Farnung et

al., 2020). To test the requirement for LET-418's ATPase activity *in vivo*, we engineered an endogenous

deletion of the ATPase domain (allele *syb6818*) and demonstrated that it is essential for LET-418's

function in the germline. ATPase deletion mutants were completely sterile, and all homozygote mutants

had defective vulvas (Fig. 6). Conversely, deletion of the PHD domain did not affect fertility or embryonic

survival, indicating that this domain is dispensable for LET-418's role in the germline in the absence of

exogenous DNA damage. Our findings are consistent with deletion of the PHD domain in other contexts.

In biochemical assays using recombinant human CHD4, deletion of the PHD domain causes a minor

decrease in remodeling, but we note that this study used unmodified histones (Watson et al., 2012). In

Drosophila, a hypomorphic point mutation to the PHD domain also slightly reduced remodeling ability,

but did not otherwise affect binding of Mi2/CHD4 to chromatin (Kovač et al., 2018).

530   The ATPase deletion phenotypes resemble those of *let-418* genetic null alleles, which were

previously used to characterize LET-418's role in vulval development (Guerry et al., 2007; von Zelewsky

et al., 2000). In somatic tissues, LET-418 belongs to two distinct complexes that have a conserved role in

maintaining cell identity – in addition to NuRD, it is also found in MEC, which consists of LET-418, the

histone deacetylase HDA-1, and the Krüppel-like protein MEP-1 (Hou et al., 2020; Käser-Pébernard et

al., 2016; Pfefferli et al., 2014; Unhavaithaya et al., 2002). As part of these complexes, LET-418 has roles

in specifying vulval fate, embryonic and larval development, and defining lifespan (De Vaux et al., 2013;

Erdelyi et al., 2017; Guerry et al., 2007; Käser-Pébernard et al., 2014; Passannante et al., 2010;

Saudenova and Wicky, 2018; von Zelewsky et al., 2000). Further characterization of these deletion

strains will establish their impact after exposure to DNA damaging agents in in roles outside of the

germline.

Based on our results, we propose the following model. When replication forks are blocked or

stalled, NuRD's remodeling activity creates a permissive chromatin environment that allows for the

recruitment of FA pathway components, like FCD-2, to repair the damage via homologous

recombination. In the absence of NuRD-mediated remodeling, the FA pathway is unable to repair DNA

lesions, causing the accumulation of mitotic and meiotic DSBs and their eventual resolution by error-

prone pathways later in meiosis. However, it is not clear from our findings whether NuRD functions

entirely upstream of the full FA pathway, or whether it mediates an intermediate step during interstrand

crosslink repair. For example, it is possible that DNA lesions are first recognized by the FA component

FNCM-1 (which acts upstream of FCD-2), which then recruits NuRD to the site of damage. In summary,

we've shown that LET-418's nucleosome remodeling activity is necessary to resolve germline DSBs by

error-free pathways like the FA pathway. This work highlights the important role of local chromatin

landscapes in coordinating DNA damage repair pathways during gametogenesis.

DATA AVAILIBILITY

Strains are available from the *Caenorhabditis* Genetics Center or upon request. The authors affirm that

all data necessary for confirming the conclusions of the article are present within the article, figures, and

tables.

ACKNOWLEDGEMENTS

We thank Diana Libuda and Monica Colaiacovo for generously sharing reagents, Cori Cahoon for

immunofluorescence advice, and WormBase for gene and sequence information (Davis et al., 2022). We

also thank two anonymous reviewers for suggestions that strengthened the manuscript. Strains were

provided by the *Caenorhabditis* Genetics Center, which is funded by the National Institutes of Health

(NIH) through the Office of Research Infrastructure Programs (P40 OD010440).

FUNDING

D. A. was supported by a UMass Lowell KCS Science Scholarship. T. B. and K. F. were supported by

fellowships from the UMass Lowell River Hawk Scholars Academy. K. F. was supported by the Urban

Massachusetts LSAMP program, which is funded by the National Science Foundation (EES 2308724). T.B.

was supported by the Society for Developmental Biology through a Choose Development! fellowship,

which is funded by the National Institutes of Health (R25HD105600). This work was supported by the

National Institutes of Health, with grants R15GM117479 and R15HD104115 to P.M.C. and grant

R15GM144861 to T.W.L.

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

REFERENCES

Achache, H., Falk, R., Lerner, N., Beatus, T., Tzur, Y.B., 2021. Oocyte aging is controlled by
mitogen-activated protein kinase signaling. Aging Cell 20, e13386.
https://doi.org/10.1111/acel.13386

Adamo, A., Collis, S.J., Adelman, C.A., Silva, N., Horejsi, Z., Ward, J.D., Martinez-Perez, E.,
Boulton, S.J., La Volpe, A., 2010. Preventing Nonhomologous End Joining Suppresses
DNA Repair Defects of Fanconi Anemia. Molecular Cell 39, 25–35.
https://doi.org/10.1016/j.molcel.2010.06.026

Allis, C.D., Jenuwein, T., 2016. The molecular hallmarks of epigenetic control. Nat Rev Genet 17,
487–500. https://doi.org/10.1038/nrg.2016.59

Alpi, A., Pasierbek, P., Gartner, A., Loidl, J., 2003. Genetic and cytological characterization of the
recombination protein RAD-51 in *Caenorhabditis elegans*. Chromosoma 112, 6–16.
https://doi.org/10.1007/s00412-003-0237-5

Ananthaswamy, D., Croft, J.C., Woozencroft, N., Lee, T.W., 2022. *C. elegans* Gonad Dissection
and Freeze Crack for Immunofluorescence and DAPI Staining. JoVE 64204.
https://doi.org/10.3791/64204

Andersen, E.C., Lu, X., Horvitz, H.R., 2006. *C. elegans* ISWI and NURF301 antagonize an Rb-like
pathway in the determination of multiple cell fates. Development 133, 2695–2704.
https://doi.org/10.1242/dev.02444

Andux, S., Ellis, R.E., 2008. Apoptosis Maintains Oocyte Quality in Aging *Caenorhabditis elegans*
Females. PLoS Genet 4, e1000295. https://doi.org/10.1371/journal.pgen.1000295

Auerbach, A.D., 2009. Fanconi anemia and its diagnosis. Mutation Research/Fundamental and
Molecular Mechanisms of Mutagenesis 668, 4–10.
https://doi.org/10.1016/j.mrfmmm.2009.01.013

Auerbach, A.D., 1993. Fanconi anemia diagnosis and the diepoxybutane (DEB) test. Exp
Hematol 21, 731–733.

Bailly, A., Gartner, A., 2011. *Caenorhabditis elegans* Radiation Responses, in: DeWeese, T.L.,
Laiho, M. (Eds.), Molecular Determinants of Radiation Response. Springer New York,
New York, NY, pp. 101–123. https://doi.org/10.1007/978-1-4419-8044-1_5

Basta, J., Rauchman, M., 2015. The nucleosome remodeling and deacetylase complex in
development and disease. Translational Research 165, 36–47.
https://doi.org/10.1016/j.trsl.2014.05.003
Becker, P.B., Workman, J.L., 2013. Nucleosome Remodeling and Epigenetics. Cold Spring Harbor
Perspectives in Biology 5, a017905–a017905.
https://doi.org/10.1101/cshperspect.a017905
Bouazoune, K., Mitterweger, A., Längst, G., Imhof, A., Akhtar, A., Becker, P.B., 2002. The dMi-2
chromodomains are DNA binding modules important for ATP-dependent nucleosome
mobilization. The EMBO Journal 21, 2430–2440.
https://doi.org/10.1093/emboj/21.10.2430
Brenner, S., 1974. THE GENETICS OF *CAENORHABDITIS ELEGANS*. Genetics 77, 71–94.
https://doi.org/10.1093/genetics/77.1.71
Cahoon, C.K., Richter, C.M., Dayton, A.E., Libuda, D.E., 2023. Sexual dimorphic regulation of
recombination by the synaptonemal complex in *C. elegans*. eLife 12, e84538.
https://doi.org/10.7554/eLife.84538
Ceccaldi, R., Rondinelli, B., D'Andrea, A.D., 2016. Repair Pathway Choices and Consequences at
the Double-Strand Break. Trends in Cell Biology 26, 52–64.
https://doi.org/10.1016/j.tcb.2015.07.009
Chapman, J.R., Taylor, M.R.G., Boulton, S.J., 2012. Playing the End Game: DNA Double-Strand
Break Repair Pathway Choice. Molecular Cell 47, 497–510.
https://doi.org/10.1016/j.molcel.2012.07.029
Chen, Z., Tyler, J.K., 2022. The Chromatin Landscape Channels DNA Double-Strand Breaks to
Distinct Repair Pathways. Front. Cell Dev. Biol. 10, 909696.
https://doi.org/10.3389/fcell.2022.909696
Chudnovsky, Y., Kim, D., Zheng, S., Whyte, W.A., Bansal, M., Bray, M.-A., Gopal, S., Theisen,
M.A., Bilodeau, S., Thiru, P., Muffat, J., Yilmaz, O.H., Mitalipova, M., Woolard, K., Lee, J.,
Nishimura, R., Sakata, N., Fine, H.A., Carpenter, A.E., Silver, S.J., Verhaak, R.G.W.,
Califano, A., Young, R.A., Ligon, K.L., Mellinghoff, I.K., Root, D.E., Sabatini, D.M., Hahn,
W.C., Chheda, M.G., 2014. ZFHX4 Interacts with the NuRD Core Member CHD4 and
Regulates the Glioblastoma Tumor-Initiating Cell State. Cell Reports 6, 313–324.
https://doi.org/10.1016/j.celrep.2013.12.032
Clouaire, T., Legube, G., 2019. A Snapshot on the Cis Chromatin Response to DNA Double-
Strand Breaks. Trends in Genetics 35, 330–345.
https://doi.org/10.1016/j.tig.2019.02.003
Colaiácovo, M.P., MacQueen, A.J., Martinez-Perez, E., McDonald, K., Adamo, A., La Volpe, A.,
Villeneuve, A.M., 2003. Synaptonemal Complex Assembly in *C. elegans* Is Dispensable
for Loading Strand-Exchange Proteins but Critical for Proper Completion of
Recombination. Developmental Cell 5, 463–474. https://doi.org/10.1016/S1534-
5807(03)00232-6
Collis, S.J., Barber, L.J., Ward, J.D., Martin, J.S., Boulton, S.J., 2006. *C. elegans* FANCD2 responds
to replication stress and functions in interstrand cross-link repair. DNA Repair 5, 1398–
1406. https://doi.org/10.1016/j.dnarep.2006.06.010

Datta, A., Brosh Jr., R.M., 2019. Holding All the Cards—How Fanconi Anemia Proteins Deal with
Replication Stress and Preserve Genomic Stability. Genes 10, 170.
https://doi.org/10.3390/genes10020170
Davis, P., Zarowiecki, M., Arnaboldi, V., Becerra, A., Cain, S., Chan, J., Chen, W.J., Cho, J., Da
Veiga Beltrame, E., Diamantakis, S., Gao, S., Grigoriadis, D., Grove, C.A., Harris, T.W.,
Kishore, R., Le, T., Lee, R.Y.N., Luypaert, M., Müller, H.-M., Nakamura, C., Nuin, P.,
Paulini, M., Quinton-Tulloch, M., Raciti, D., Rodgers, F.H., Russell, M., Schindelman, G.,
Singh, A., Stickland, T., Van Auken, K., Wang, Q., Williams, G., Wright, A.J., Yook, K.,
Berriman, M., Howe, K.L., Schedl, T., Stein, L., Sternberg, P.W., 2022. WormBase in
2022—data, processes, and tools for analyzing *Caenorhabditis elegans*. Genetics 220,
iyac003. https://doi.org/10.1093/genetics/iyac003
De Massy, B., 2013. Initiation of Meiotic Recombination: How and Where? Conservation and
Specificities Among Eukaryotes. Annu. Rev. Genet. 47, 563–599.
https://doi.org/10.1146/annurev-genet-110711-155423
De Vaux, V., Pfefferli, C., Passannante, M., Belhaj, K., Von Essen, A., Sprecher, S.G., Müller, F.,
Wicky, C., 2013. The *Caenorhabditis elegans* LET-418/ Mi2 plays a conserved role in
lifespan regulation. Aging Cell 12, 1012–1020. https://doi.org/10.1111/acel.12129
De Winter, J.P., Joenje, H., 2009. The genetic and molecular basis of Fanconi anemia. Mutation
Research/Fundamental and Molecular Mechanisms of Mutagenesis 668, 11–19.
https://doi.org/10.1016/j.mrfmmm.2008.11.004
Erdelyi, P., Wang, X., Suleski, M., Wicky, C., 2017. A Network of Chromatin Factors Is Regulating
the Transition to Postembryonic Development in *Caenorhabditis elegans*. G3
Genes|Genomes|Genetics 7, 343–353. https://doi.org/10.1534/g3.116.037747
Farnung, L., Ochmann, M., Cramer, P., 2020. Nucleosome-CHD4 chromatin remodeler structure
maps human disease mutations. eLife 9, e56178. https://doi.org/10.7554/eLife.56178
Gartner, A., Engebrecht, J., 2022. DNA repair, recombination, and damage signaling. Genetics
220, iyab178. https://doi.org/10.1093/genetics/iyab178
Germoglio, M., Valenti, A., Gallo, I., Forenza, C., Santonicola, P., Silva, N., Adamo, A., 2020. In
vivo analysis of FANCD2 recruitment at meiotic DNA breaks in *Caenorhabditis elegans*.
Sci Rep 10, 103. https://doi.org/10.1038/s41598-019-57096-1
Guerry, F., Marti, C.-O., Zhang, Y., Moroni, P.S., Jaquiéry, E., Müller, F., 2007. The Mi-2
nucleosome-remodeling protein LET-418 is targeted via LIN-1/ETS to the promoter of
lin-39/Hox during vulval development in *C. elegans*. Developmental Biology 306, 469–
479. https://doi.org/10.1016/j.ydbio.2007.03.026
Hou, T., Cao, Z., Zhang, J., Tang, M., Tian, Y., Li, Y., Lu, X., Chen, Y., Wang, Hui, Wei, F.-Z., Wang,
L., Yang, Y., Zhao, Y., Wang, Z., Wang, Haiying, Zhu, W.-G., 2020. SIRT6 coordinates with
CHD4 to promote chromatin relaxation and DNA repair. Nucleic Acids Res 48, 2982–
3000. https://doi.org/10.1093/nar/gkaa006
Jaramillo-Lambert, A., Ellefson, M., Villeneuve, A.M., Engebrecht, J., 2007. Differential timing of
S phases, X chromosome replication, and meiotic prophase in the *C. elegans* germ line.
Developmental Biology 308, 206–221. https://doi.org/10.1016/j.ydbio.2007.05.019
Käser-Pébernard, S., Müller, F., Wicky, C., 2014. LET-418/Mi2 and SPR-5/LSD1 Cooperatively
Prevent Somatic Reprogramming of *C. elegans* Germline Stem Cells. Stem Cell Reports 2,
547–559. https://doi.org/10.1016/j.stemcr.2014.02.007

Käser-Pébernard, S., Pfefferli, C., Aschinger, C., Wicky, C., 2016. Fine-tuning of chromatin
composition and Polycomb recruitment by two Mi2 homologues during *C. elegans* early
embryonic development. Epigenetics & Chromatin 9, 39.
https://doi.org/10.1186/s13072-016-0091-3
Keeney, S., 2007. Spo11 and the Formation of DNA Double-Strand Breaks in Meiosis, in: Egel, R.,
Lankenau, D.-H. (Eds.), Recombination and Meiosis, Genome Dynamics and Stability.
Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 81–123.
https://doi.org/10.1007/7050_2007_026
Kehle, J., 1998. dMi-2, a Hunchback-Interacting Protein That Functions in Polycomb Repression.
Science 282, 1897–1900. https://doi.org/10.1126/science.282.5395.1897
Kessler, Z., Yanowitz, J., 2014. Methodological considerations for mutagen exposure in *C.*
elegans. Methods 68, 441–449. https://doi.org/10.1016/j.ymeth.2014.04.011
Kim, H.-M., Beese-Sims, S.E., Colaiácovo, M.P., 2018. Fanconi Anemia FANCM/FNCM-1 and
FANCD2/FCD-2 Are Required for Maintaining Histone Methylation Levels and Interact
with the Histone Demethylase LSD1/SPR-5 in *Caenorhabditis elegans*. Genetics 209,
409–423. https://doi.org/10.1534/genetics.118.300823
Kim, H.-M., Colaiácovo, M.P., 2015. DNA Damage Sensitivity Assays in *Caenorhabditis elegans*.
Bio Protoc 5, e1487.
Kim, S., Peterson, S.E., Jasin, M., Keeney, S., 2016. Mechanisms of germ line genome instability.
Seminars in Cell & Developmental Biology 54, 177–187.
https://doi.org/10.1016/j.semcdb.2016.02.019
Kitange, G.J., Mladek, A.C., Schroeder, M.A., Pokorny, J.C., Carlson, B.L., Zhang, Y., Nair, A.A.,
Lee, J.-H., Yan, H., Decker, P.A., Zhang, Z., Sarkaria, J.N., 2016. Retinoblastoma Binding
Protein 4 Modulates Temozolomide Sensitivity in Glioblastoma by Regulating DNA
Repair Proteins. Cell Reports 14, 2587–2598.
https://doi.org/10.1016/j.celrep.2016.02.045
Kochan, J.A., Desclos, E.C.B., Bosch, R., Meister, L., Vriend, L.E.M., van Attikum, H., Krawczyk,
P.M., 2017. Meta-analysis of DNA double-strand break response kinetics. Nucleic Acids
Res 45, 12625–12637. https://doi.org/10.1093/nar/gkx1128
Kovač, K., Sauer, A., Mačinković, I., Awe, S., Finkernagel, F., Hoffmeister, H., Fuchs, A., Müller,
R., Rathke, C., Längst, G., Brehm, A., 2018. Tumour-associated missense mutations in the
dMi-2 ATPase alters nucleosome remodelling properties in a mutation-specific manner.
Nat Commun 9, 2112. https://doi.org/10.1038/s41467-018-04503-2
Kunert, N., Wagner, E., Murawska, M., Klinker, H., Kremmer, E., Brehm, A., 2009. dMec: a novel
Mi-2 chromatin remodelling complex involved in transcriptional repression. EMBO J 28,
533–544. https://doi.org/10.1038/emboj.2009.3
Kurhanewicz, N.A., Dinwiddie, D., Bush, Z.D., Libuda, D.E., 2020. Elevated Temperatures Cause
Transposon-Associated DNA Damage in *C. elegans* Spermatocytes. Current Biology 30,
5007-5017.e4. https://doi.org/10.1016/j.cub.2020.09.050
Lachaud, C., Moreno, A., Marchesi, F., Toth, R., Blow, J.J., Rouse, J., 2016. Ubiquitinated Fancd2
recruits Fan1 to stalled replication forks to prevent genome instability. Science 351,
846–849. https://doi.org/10.1126/science.aad5634

Lee, K.Y., Chung, K.Y., Koo, H.-S., 2010. The involvement of FANCM, FANCI, and checkpoint proteins in the interstrand DNA crosslink repair pathway is conserved in *C. elegans*. DNA Repair 9, 374–382. https://doi.org/10.1016/j.dnarep.2009.12.018

Lee, K.Y., Yang, I., Park, J.-E., Baek, O.-R., Chung, K.Y., Koo, H.-S., 2007. Developmental stage- and DNA damage-specific functions of *C. elegans* FANCD2. Biochemical and Biophysical Research Communications 352, 479–485. https://doi.org/10.1016/j.bbrc.2006.11.039

Luo, S., Kleemann, G.A., Ashraf, J.M., Shaw, W.M., Murphy, C.T., 2010. TGF-β and Insulin Signaling Regulate Reproductive Aging via Oocyte and Germline Quality Maintenance. Cell 143, 299–312. https://doi.org/10.1016/j.cell.2010.09.013

Macaisne, N., Kessler, Z., Yanowitz, J.L., 2018. Meiotic Double-Strand Break Proteins Influence Repair Pathway Utilization. Genetics 210, 843–856. https://doi.org/10.1534/genetics.118.301402

Mansfield, R.E., Musselman, C.A., Kwan, A.H., Oliver, S.S., Garske, A.L., Davrazou, F., Denu, J.M., Kutateladze, T.G., Mackay, J.P., 2011. Plant Homeodomain (PHD) Fingers of CHD4 Are Histone H3-binding Modules with Preference for Unmodified H3K4 and Methylated H3K9. Journal of Biological Chemistry 286, 11779–11791. https://doi.org/10.1074/jbc.M110.208207

McKenzie, L.D., LeClair, J.W., Miller, K.N., Strong, A.D., Chan, H.L., Oates, E.L., Ligon, K.L., Brennan, C.W., Chheda, M.G., 2019. CHD4 regulates the DNA damage response and RAD51 expression in glioblastoma. Sci Rep 9, 4444. https://doi.org/10.1038/s41598-019-40327-w

McMurchy, A.N., Stempor, P., Gaarenstroom, T., Wysolmerski, B., Dong, Y., Aussianikava, D., Appert, A., Huang, N., Kolasinska-Zwierz, P., Sapetschnig, A., Miska, E.A., Ahringer, J., 2017. A team of heterochromatin factors collaborates with small RNA pathways to combat repetitive elements and germline stress. eLife 6, e21666. https://doi.org/10.7554/eLife.21666

Meier, B., Cooke, S.L., Weiss, J., Bailly, A.P., Alexandrov, L.B., Marshall, J., Raine, K., Maddison, M., Anderson, E., Stratton, M.R., Gartner, A., Campbell, P.J., 2014. *C. elegans* whole-genome sequencing reveals mutational signatures related to carcinogens and DNA repair deficiency. Genome Res. 24, 1624–1636. https://doi.org/10.1101/gr.175547.114

Nair, N., Shoaib, M., Sørensen, C.S., 2017. Chromatin Dynamics in Genome Stability: Roles in Suppressing Endogenous DNA Damage and Facilitating DNA Repair. IJMS 18, 1486. https://doi.org/10.3390/ijms18071486

Nakanishi, K., Yang, Y.-G., Pierce, A.J., Taniguchi, T., Digweed, M., D'Andrea, A.D., Wang, Z.-Q., Jasin, M., 2005. Human Fanconi anemia monoubiquitination pathway promotes homologous DNA repair. Proc. Natl. Acad. Sci. U.S.A. 102, 1110–1115. https://doi.org/10.1073/pnas.0407796102

Nalepa, G., Clapp, D.W., 2018. Fanconi anaemia and cancer: an intricate relationship. Nat Rev Cancer 18, 168–185. https://doi.org/10.1038/nrc.2017.116

Niraj, J., Färkkilä, A., D'Andrea, A.D., 2019. The Fanconi Anemia Pathway in Cancer. Annu. Rev. Cancer Biol. 3, 457–478. https://doi.org/10.1146/annurev-cancerbio-030617-050422

Pan, M.-R., Hsieh, H.-J., Dai, H., Hung, W.-C., Li, K., Peng, G., Lin, S.-Y., 2012. Chromodomain Helicase DNA-binding Protein 4 (CHD4) Regulates Homologous Recombination DNA Repair, and Its Deficiency Sensitizes Cells to Poly(ADP-ribose) Polymerase (PARP)

779   Inhibitor Treatment. Journal of Biological Chemistry 287, 6764–6772.
780   https://doi.org/10.1074/jbc.M111.287037

781 Passannante, M., Marti, C.-O., Pfefferli, C., Moroni, P.S., Kaeser-Pebernard, S., Puoti, A.,
782   Hunziker, P., Wicky, C., Müller, F., 2010. Different Mi-2 Complexes for Various
783   Developmental Functions in *Caenorhabditis elegans*. PLoS ONE 5, e13681.
784   https://doi.org/10.1371/journal.pone.0013681

785 Pfefferli, C., Müller, F., Jaźwińska, A., Wicky, C., 2014. Specific NuRD components are required
786   for fin regeneration in zebrafish. BMC Biol 12, 30. https://doi.org/10.1186/1741-7007-
787   12-30

788 Polo, S.E., Kaidi, A., Baskcomb, L., Galanty, Y., Jackson, S.P., 2010. Regulation of DNA-damage
789   responses and cell-cycle progression by the chromatin remodelling factor CHD4. EMBO J
790   29, 3130–3139. https://doi.org/10.1038/emboj.2010.188

791 Povirk, L.F., Shuker, D.E., 1994. DNA damage and mutagenesis induced by nitrogen mustards.
792   Mutat Res 318, 205–226. https://doi.org/10.1016/0165-1110(94)90015-9

793 Rageul, J., Kim, H., 2020. Fanconi anemia and the underlying causes of genomic instability.
794   Environ and Mol Mutagen 61, 693–708. https://doi.org/10.1002/em.22358

795 Raghunandan, M., Chaudhury, I., Kelich, S.L., Hanenberg, H., Sobeck, A., 2015. FANCD2, FANCJ
796   and BRCA2 cooperate to promote replication fork recovery independently of the
797   Fanconi Anemia core complex. Cell Cycle 14, 342–353.
798   https://doi.org/10.4161/15384101.2014.987614

799 Rosu, S., Zawadzki, K.A., Stamper, E.L., Libuda, D.E., Reese, A.L., Dernburg, A.F., Villeneuve,
800   A.M., 2013. The *C. elegans* DSB-2 Protein Reveals a Regulatory Network that Controls
801   Competence for Meiotic DSB Formation and Promotes Crossover Assurance. PLoS Genet
802   9, e1003674. https://doi.org/10.1371/journal.pgen.1003674

803 Saudenova, M., Wicky, C., 2018. The Chromatin Remodeler LET-418/Mi2 is Required Cell Non-
804   Autonomously for the Post-Embryonic Development of *Caenorhabditis elegans*. JDB 7, 1.
805   https://doi.org/10.3390/jdb7010001

806 Sayers, E.W., Bolton, E.E., Brister, J.R., Canese, K., Chan, J., Comeau, D.C., Connor, R., Funk, K.,
807   Kelly, C., Kim, S., Madej, T., Marchler-Bauer, A., Lanczycki, C., Lathrop, S., Lu, Z., Thibaud-
808   Nissen, F., Murphy, T., Phan, L., Skripchenko, Y., Tse, T., Wang, J., Williams, R., Trawick,
809   B.W., Pruitt, K.D., Sherry, S.T., 2022. Database resources of the national center for
810   biotechnology information. Nucleic Acids Research 50, D20–D26.
811   https://doi.org/10.1093/nar/gkab1112

812 Scharf, A., Pohl, F., Egan, B.M., Kocsisova, Z., Kornfeld, K., 2021. Reproductive Aging in
813   *Caenorhabditis elegans*: From Molecules to Ecology. Front. Cell Dev. Biol. 9, 718522.
814   https://doi.org/10.3389/fcell.2021.718522

815 Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
816   Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J., Hartenstein, V., Eliceiri, K.,
817   Tomancak, P., Cardona, A., 2012. Fiji: an open-source platform for biological-image
818   analysis. Nat Methods 9, 676–682. https://doi.org/10.1038/nmeth.2019

819 Schlacher, K., Wu, H., Jasin, M., 2012. A Distinct Replication Fork Protection Pathway Connects
820   Fanconi Anemia Tumor Suppressors to RAD51-BRCA1/2. Cancer Cell 22, 106–116.
821   https://doi.org/10.1016/j.ccr.2012.05.015

Smolikov, S., Eizinger, A., Hurlburt, A., Rogers, E., Villeneuve, A.M., Colaiácovo, M.P., 2007.
Synapsis-defective mutants reveal a correlation between chromosome conformation
and the mode of double-strand break repair during *Caenorhabditis elegans* meiosis.
Genetics 176, 2027–2033. https://doi.org/10.1534/genetics.107.076968
Sperlazza, J., Rahmani, M., Beckta, J., Aust, M., Hawkins, E., Wang, S.Z., Zu Zhu, S., Podder, S.,
Dumur, C., Archer, K., Grant, S., Ginder, G.D., 2015. Depletion of the chromatin
remodeler CHD4 sensitizes AML blasts to genotoxic agents and reduces tumor
formation. Blood 126, 1462–1472. https://doi.org/10.1182/blood-2015-03-631606
Stamper, E.L., Rodenbusch, S.E., Rosu, S., Ahringer, J., Villeneuve, A.M., Dernburg, A.F., 2013.
Identification of DSB-1, a Protein Required for Initiation of Meiotic Recombination in
Caenorhabditis elegans, Illuminates a Crossover Assurance Checkpoint. PLoS Genet 9,
e1003679. https://doi.org/10.1371/journal.pgen.1003679
Stinson, B.M., Loparo, J.J., 2021. Repair of DNA Double-Strand Breaks by the Nonhomologous
End Joining Pathway. Annu. Rev. Biochem. 90, 137–164.
https://doi.org/10.1146/annurev-biochem-080320-110356
Tischkowitz, M., Ameziane, N., Waisfisz, Q., De Winter, J.P., Harris, R., Taniguchi, T., D'Andrea,
A., Hodgson, S.V., Mathew, C.G., Joenje, H., 2003. Bi-allelic silencing of the Fanconi
anaemia gene *FANCF* in acute myeloid leukaemia: *Short Report*. British Journal of
Haematology 123, 469–471. https://doi.org/10.1046/j.1365-2141.2003.04640.x
Tischkowitz, M.D., Morgan, N.V., Grimwade, D., Eddy, C., Ball, S., Vorechovsky, I., Langabeer, S.,
Stöger, R., Hodgson, S.V., Mathew, C.G., 2004. Deletion and reduced expression of the
Fanconi anemia FANCA gene in sporadic acute myeloid leukemia. Leukemia 18, 420–
425. https://doi.org/10.1038/sj.leu.2403280
Tong, J.K., Hassig, C.A., Schnitzler, G.R., Kingston, R.E., Schreiber, S.L., 1998. Chromatin
deacetylation by an ATP-dependent nucleosome remodelling complex. Nature 395,
917–921. https://doi.org/10.1038/27699
Tsui, V., Crismani, W., 2019. The Fanconi Anemia Pathway and Fertility. Trends in Genetics 35,
199–214. https://doi.org/10.1016/j.tig.2018.12.007
Turcotte, C.A., Sloat, S.A., Rigothi, J.A., Rosenkranse, E., Northrup, A.L., Andrews, N.P., Checchi,
P.M., 2018. Maintenance of Genome Integrity by Mi2 Homologs CHD-3 and LET-418 in
Caenorhabditis elegans. Genetics 208, 991–1007.
https://doi.org/10.1534/genetics.118.300686
Unhavaithaya, Y., Shin, T.H., Miliaras, N., Lee, J., Oyama, T., Mello, C.C., 2002. MEP-1 and a
Homolog of the NURD Complex Component Mi-2 Act Together to Maintain Germline-
Soma Distinctions in *C. elegans*. Cell 111, 991–1002. https://doi.org/10.1016/S0092-
8674(02)01202-3
von Zelewsky, T., Palladino, F., Brunschwig, K., Tobler, H., Hajnal, A., Müller, F., Brehm, A., 2000.
The *C. elegans* Mi-2 chromatin-remodelling proteins function in vulval cell fate
determination. Development 127, 5277–5284. https://doi.org/10.1242/dev.127.24.5277
Wade, P.A., Jones, P.L., Vermaak, D., Wolffe, A.P., 1998. A multiple subunit Mi-2 histone
deacetylase from Xenopus laevis cofractionates with an associated Snf2 superfamily
ATPase. Current Biology 8, 843–848. https://doi.org/10.1016/S0960-9822(98)70328-8
Wang, Y., Zhang, H., Chen, Y., Sun, Y., Yang, F., Yu, W., Liang, J., Sun, L., Yang, X., Shi, L., Li, R., Li,
Y., Zhang, Y., Li, Q., Yi, X., Shang, Y., 2009. LSD1 Is a Subunit of the NuRD Complex and

Targets the Metastasis Programs in Breast Cancer. Cell 138, 660–672.
https://doi.org/10.1016/j.cell.2009.05.050
Ward, J.D., Barber, L.J., Petalcorin, M.I., Yanowitz, J., Boulton, S.J., 2007. Replication blocking
lesions present a unique substrate for homologous recombination. EMBO J 26, 3384–
3396. https://doi.org/10.1038/sj.emboj.7601766
Watson, A.A., Mahajan, P., Mertens, H.D.T., Deery, M.J., Zhang, Wenchao, Pham, P., Du, X.,
Bartke, T., Zhang, Wei, Edlich, C., Berridge, G., Chen, Y., Burgess-Brown, N.A.,
Kouzarides, T., Wiechens, N., Owen-Hughes, T., Svergun, D.I., Gileadi, O., Laue, E.D.,
2012. The PHD and chromo domains regulate the ATPase activity of the human
chromatin remodeler CHD4. J Mol Biol 422, 3–17.
https://doi.org/10.1016/j.jmb.2012.04.031
Weiss, K., Terhal, P.A., Cohen, L., Bruccoleri, M., Irving, M., Martinez, A.F., Rosenfeld, J.A.,
Machol, K., Yang, Y., Liu, P., Walkiewicz, M., Beuten, J., Gomez-Ospina, N., Haude, K.,
Fong, C.-T., Enns, G.M., Bernstein, J.A., Fan, J., Gotway, G., Ghorbani, M., DDD Study, van
Gassen, K., Monroe, G.R., van Haaften, G., Basel-Vanagaite, L., Yang, X.-J., Campeau,
P.M., Muenke, M., 2016. De Novo Mutations in CHD4, an ATP-Dependent Chromatin
Remodeler Gene, Cause an Intellectual Disability Syndrome with Distinctive
Dysmorphisms. Am J Hum Genet 99, 934–941.
https://doi.org/10.1016/j.ajhg.2016.08.001
Whyte, W.A., Bilodeau, S., Orlando, D.A., Hoke, H.A., Frampton, G.M., Foster, C.T., Cowley,
S.M., Young, R.A., 2012. Enhancer decommissioning by LSD1 during embryonic stem cell
differentiation. Nature 482, 221–225. https://doi.org/10.1038/nature10805
Wilson, T.E., Grawunder, U., Lieber, M.R., 1997. Yeast DNA ligase IV mediates non-homologous
DNA end joining. Nature 388, 495–498. https://doi.org/10.1038/41365
Woodage, T., Basrai, M.A., Baxevanis, A.D., Hieter, P., Collins, F.S., 1997. Characterization of the
CHD family of proteins. Proc. Natl. Acad. Sci. U.S.A. 94, 11472–11477.
https://doi.org/10.1073/pnas.94.21.11472
Xue, Y., Wong, J., Moreno, G.T., Young, M.K., Côté, J., Wang, W., 1998. NURD, a Novel Complex
with Both ATP-Dependent Chromatin-Remodeling and Histone Deacetylase Activities.
Molecular Cell 2, 851–861. https://doi.org/10.1016/S1097-2765(00)80299-3
Youds, J.L., Barber, L.J., Boulton, S.J., 2009. *C. elegans*: A model of Fanconi anemia and ICL
repair. Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis
668, 103–116. https://doi.org/10.1016/j.mrfmmm.2008.11.007

899
900
901 **FIGURE LEGENDS**
902
903 **Figure 1: *let-418* mutants cannot repair mitotic DSBs when challenged with cisplatin.**
904 **(A, B)** Stacked histograms showing the percentage of nuclei with the indicated number of RAD-51 foci
905 for animals exposed to 0 μM (A) and 400 μM of cisplatin (B). Average number of foci per nucleus is
906 indicated under each genotype. Comparisons between wild-type and mutant populations at the same
907 dose are shown directly under averages for each genotype; other comparisons are indicated by

908 horizontal lines. For all genotypes, exposed populations differed significantly from unexposed controls

909 (not indicated on histogram, $P < 0.0001$ for all comparisons). Each condition includes at least 2000 nuclei

910 analyzed from 15 germlines across three biological replicates. (**C, D**) Representative images of mitotic

911 region nuclei in unexposed (C) and exposed (D) animals, with the distal tip oriented to the left (as

912 indicated by white asterisk). Germlines are stained for RAD-51 (in yellow) and DAPI (in blue). Scale bar

913 represents 2 μm. All statistical analyses were performed using a Kruskal-Wallis test with Dunn's post test

914 (n.s. = not significant, * $P < 0.05$, **** $P < 0.0001$). Summary statistics and data are included in Table S1.

916 **Figure 2: *let-418* mutants cannot repair mitotic DSBs when challenged with hydroxyurea.**

917 **(A, B)** Stacked histograms showing the percentage of nuclei with the indicated number of RAD-51 foci

918 for animals exposed to 0 mM (A) and 10 mM of hydroxyrea (B). Average number of foci per nucleus is

919 indicated under each genotype. Comparisons between wild-type and mutant populations at the same

920 dose are shown directly under averages for each genotype; other comparisons are indicated by

921 horizontal lines. For all genotypes, exposed populations differed significantly from unexposed

922 populations (not shown on graph, $P < 0.0001$ for all comparisons). Each condition includes at least 2000

923 nuclei analyzed from 15 germlines across three biological replicates. (**C, D**) Representative images of

924 mitotic region nuclei in unexposed (C) and exposed (D) animals, with the distal tip oriented to the left (as

925 indicated by white asterisk). Germlines are stained for RAD-51 (in yellow) and DAPI (in blue). Scale bar

926 represents 2 μm. All statistical analyses were performed using a Kruskal-Wallis test with Dunn's post test

927 (n.s. = not significant, * $P < 0.05$, ** $P < 0.01$, **** $P < 0.0001$). Summary statistics and data are included

928 in Table S1.

930 **Figure 3: Mutagen-induced DNA damage persists throughout pachytene in *let-418* mutants.**

931 **(A, C, E)** Quantification of number of nuclei with the indicated number of RAD-51 foci in unexposed (A),

932 cisplatin-exposed (C), and hydroxyurea-exposed (E) hermaphrodites. Dark line indicates average number

933 of foci for each stage of pachytene: early pachytene (EP), mid-pachytene (MP), and late pachytene (LP).

934 Each stage represents at least 100 nuclei scored from three gonads. Averages for mid-pachytene are

935 shown under each genotype. All comparisons shown are only for mid-pachytene regions (comparisons

936 for early- and late-pachytene regions are included in Table S2): comparison between exposed and

937 unexposed populations are shown under averages; other comparisons are indicated by horizontal lines.

938 **(B, D, F)** Representative immunofluorescence images of mid-pachytene nuclei stained for RAD-51

939 (yellow) and DAPI (blue) in unexposed (B), cisplatin-exposed (D), and hydroxyurea-exposed (F)

hermaphrodites. Scale bar represents 2 µm. Summary statistics and data are included in Table S2. All

statistical analyses were performed using a Kruskal-Wallis test with Dunn's post test (n.s. = not

significant, **** $P < 0.0001$).

Figure 4: *let-418* mutants have poor oocyte quality due to mutagen induced mitotic DSBs.

(A, B, C, D) Stacked histograms showing the percentage of nuclei with the indicated number of DAPI

bodies for animals exposed to 0 µM of cisplatin (A), 400 µM of cisplatin (B), 0 mM of hydroxyurea (C)

and 10 mM of hydroxyurea (D). Comparisons between wild-type and mutant populations at the same

dose are shown directly under averages for each genotype; other comparisons are indicated by

horizontal lines. Each condition includes at least 100 nuclei analyzed from 27 germlines across three

biological replicates. **(E, F)** Representative images of -1 oocytes in animals exposed to 0 uM (E) and 400

uM of cisplatin (F). See Fig. S2 for representative images of -1 oocytes exposed to hydroxyurea.

Germlines are stained with DAPI (in white). Scale bar represents 2 µm. All statistical analyses were

performed as a Fisher's exact test on number of DAPI bodies (n.s. = not significant, * $P < 0.05$, ** $P <$

0.01, **** $P < 0.0001$). Summary statistics and data are included in Table S3.

Figure 5: DNA damaging agents reduce embryonic viability in *let-418* mutants.

(**A, B**) Percent survival of indicated genotypes with and without exposure to cisplatin (A) or hydroxyurea

(B); dose is indicated with triangle above genotype names, with cisplatin in salmon (0 µM and 400 µM)

and hydroxyurea in blue (0 mM and 10 mM). For all conditions in (A) N=125 and (B) N=150. Dark line

represents the mean and whiskers represent S.E.M. Comparisons between wild-type and mutant

populations at the same dose are above each condition; other comparisons are indicated by horizontal

lines. All statistical analyses were performed using a one-way ANOVA with a Šídák correction (n.s. = not

significant, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$). Summary statistics and data are

found in Table S4.

Figure 6: The ATPase motor domain is necessary for LET-418 protein function in DSB repair.

(A) Protein alignment of human CHD4 and *C. elegans* LET-418 homologs showing conserved domains.

Primary structural alignments were performed according to NCBI-predicted domains using UniProt.

Locations of each allele are indicated above the alignment: *syb6708* PHD finger deletion (magenta),

canonical *n3536* missense mutation (pink), and *syb6818* ATPase deletion (teal). (**B**) Images of young

adult hermaphrodites. The ATPase deletion *syb6818*, has a protruding vulva phenotype (arrow to pVulv)

and a lack of germline as indicated by the white space in the body. Scale bar, 100 µm. (**C**) Total brood

was assessed in wild-type (gray), *n3536* canonical allele mutants (light pink), *syb6708* PHD finger

deletion mutants (magenta), and *syb6818* ATPase deletion mutants (teal). (**D**) Percent embryonic

survival, excluding *syb6818* due to complete sterility. For both graphs, dark line represents mean and

whiskers represent S.E.M. All statistical analyses were performed using ANOVA with a Šídák correction

(n.s. = not significant, *$P < 0.05$, **$P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$). Summary statistics and data

are included in Table S5.

Figure S1: Cisplatin affects embryonic survival in a dose-dependent manner.

(**A, B**) Mean percent survival of wild-type populations (gray) exposed to doses of cisplatin ranging from 0

µM to 700 µM (A) or 0 µM to 400 µM (B). Relative dose levels are indicated by triangles under the X-

axis. (B) Examining the effects of cisplatin on *let-418* mutants (pink), with *xpf-1* mutants (green) included

as a positive control for cisplatin sensitivity. Whiskers represent S.E.M. For each condition, at least 25

broods were evaluated. Summary statistics and data are included in Table S6.

Figure S2: *let-418* mutants have poor oocyte quality due to hydroxyurea induced mitotic DSBs.

(**A, B**) Representative images of -1 oocytes in unexposed (A) and exposed (B) animals. Germlines are

stained with DAPI (in white). Scale bar represents 2 µm. Summary statistics and data are included in

Table S3.

Figure S3: Hydroxyurea affects embryonic survival in a dose-dependent manner*.*

Mean percent survival of wild-type (gray) and *let-418* mutants (pink) exposed to increasing doses of

hydroxyurea (from 0 mM to 10 mM, with relative levels indicated by the blue triangle under the X-axis).

Whiskers represent S.E.M. For each condition, at least 25 broods were evaluated. Summary statistics

and data are included in Table S6.

Figure S4: Nitrogen mustard reduces embryonic survival in *let-418* and *fcd-2* mutants*.*

Percent survival of indicated genotypes with and without exposure nitrogen mustard; brown triangles

indicate dose (0 mM, 100 mM, and 150 mM). Dark line represents mean percent survival and whiskers

represent S.E.M. . Comparisons between wild-type and mutant populations at the same dose are above

each mutant; other comparisons are indicated by horizontal lines. All statistical analyses were

performed using ANOVA followed by Šídák's multiple comparisons (**$P < 0.01$, *** $P < 0.001$, **** $P <$

0.0001). Summary statistics and data are included in Table S6.

Figure 1: *let-418* mutants cannot repair mitotic DSBs when challenged with cisplatin.
(A, B) Stacked histograms showing the percentage of nuclei with the indicated number of RAD-51 foci for animals exposed to 0 µM (A) and 400 µM of cisplatin (B). Average number of foci per nucleus is indicated under each genotype. Comparisons between wild-type and mutant populations at the same dose are shown directly under averages for each genotype; other comparisons are indicated by horizontal lines. For all genotypes, exposed populations differed significantly from unexposed controls (not indicated on histogram, $P < 0.0001$ for all comparisons). Each condition includes at least 2000 nuclei analyzed from 15 germlines across three biological replicates. **(C, D)** Representative images of mitotic region nuclei in unexposed (C) and exposed (D) animals, with the distal tip oriented to the left (as indicated by white asterisk). Germlines are stained for RAD-51 (in yellow) and DAPI (in blue). Scale bar represents 2 µm. All statistical analyses were performed using a Kruskal-Wallis test with Dunn's post test (n.s. = not significant, * $P < 0.05$, **** $P < 0.0001$). Summary statistics and data are included in Table S1.

Figure 2: *let-418* mutants cannot repair mitotic DSBs when challenged with hydroxyurea.
(A, B) Stacked histograms showing the percentage of nuclei with the indicated number of RAD-51 foci for animals exposed to 0 mM (A) and 10 mM of hydroxyrea (B). Average number of foci per nucleus is indicated under each genotype. Comparisons between wild-type and mutant populations at the same dose are shown directly under averages for each genotype; other comparisons are indicated by horizontal lines. For all genotypes, exposed populations differed significantly from unexposed populations (not shown on graph, *P* < 0.0001 for all comparisons). Each condition includes at least 2000 nuclei analyzed from 15 germlines across three biological replicates. **(C, D)** Representative images of mitotic region nuclei in unexposed (C) and exposed (D) animals, with the distal tip oriented to the left (as indicated by white asterisk). Germlines are stained for RAD-51 (in yellow) and DAPI (in blue). Scale bar represents 2 μm. All statistical analyses were performed using a Kruskal-Wallis test with Dunn's post test (n.s. = not significant, * *P* < 0.05, ** *P* < 0.01, **** *P* < 0.0001). Summary statistics and data are included in Table S1.

Figure 3: Mutagen-induced DNA damage persists throughout pachytene in *let-418* mutants.
(A, C, E) Quantification of number of nuclei with the indicated number of RAD-51 foci in unexposed (A), cisplatin-exposed (C), and hydroxyurea-exposed (E) hermaphrodites. Dark line indicates average number of foci for each stage of pachytene: early pachytene (EP), mid-pachytene (MP), and late pachytene (LP). Each stage represents at least 100 nuclei scored from three gonads. Averages for mid-pachytene are shown under each genotype. All comparisons shown are only for mid-pachytene regions (comparisons for early- and late-pachytene regions are included in Table S2): comparison between exposed and unexposed populations are shown under averages; other comparisons are indicated by horizontal lines.
(B, D, F) Representative immunofluorescence images of mid-pachytene nuclei stained for RAD-51 (yellow) and DAPI (blue) in unexposed (B), cisplatin-exposed (D), and hydroxyurea-exposed (F) hermaphrodites. Scale bar represents 2 μm. Summary statistics and data are included in Table S2. All statistical analyses were performed using a Kruskal-Wallis test with Dunn's post test (n.s. = not significant, **** *P* < 0.0001).

Figure 4: *let-418* mutants have poor oocyte quality due to mutagen induced mitotic DSBs.
(A, B, C, D) Stacked histograms showing the percentage of nuclei with the indicated number of DAPI bodies for animals exposed to 0 µM of cisplatin (A), 400 µM of cisplatin (B), 0 mM of hydroxyurea (C) and 10 mM of hydroxyurea (D). Comparisons between wild-type and mutant populations at the same dose are shown directly under averages for each genotype; other comparisons are indicated by horizontal lines. Each condition includes at least 100 nuclei analyzed from 27 germlines across three biological replicates. **(E, F)** Representative images of -1 oocytes in animals exposed to 0 uM (E) and 400 uM of cisplatin (F). See Fig. S1 for representative images of -1 oocytes exposed to hydroxyurea. Germlines are stained with DAPI (in white). Scale bar represents 2 µm. All statistical analyses were performed as a Fisher's exact test on number of DAPI bodies (n.s. = not significant, * $P < 0.05$, ** $P < 0.01$, **** $P < 0.0001$). Summary statistics and data are included in Table S3.

Figure 5: DNA damaging agents reduce embryonic viability in *let-418* mutants.

(**A, B**) Percent survival of indicated genotypes with and without exposure to cisplatin (A) or hydroxyurea (B); dose is indicated with triangle above genotype names, with cisplatin in salmon (0 μM and 400 μM) and hydroxyurea in blue (0 mM and 10 mM). For all conditions in (A) N=125 and (B) N=150. Dark line represents the mean and whiskers represent S.E.M. Comparisons between wild-type and mutant populations at the same dose are above each condition; other comparisons are indicated by horizontal lines. All statistical analyses were performed using a one-way ANOVA with a Šídák correction (n.s. = not significant, *P < 0.05, ** P < 0.01, *** P < 0.001, **** P < 0.0001). Summary statistics and data are found in Table S4.

Figure 6: The ATPase motor domain is necessary for LET-418 protein function in DSB repair.
(A) Protein alignment of human CHD4 and *C. elegans* LET-418 homologs showing conserved domains.
Primary structural alignments were performed according to NCBI-predicted domains using UniProt.
Locations of each allele are indicated above the alignment: *syb6708* PHD finger deletion (magenta),
canonical *n3536* missense mutation (pink), and *syb6818* ATPase deletion (teal). **(B)** Images of young
adult hermaphrodites. The ATPase deletion *syb6818*, has a protruding vulva phenotype (arrow to pVulv)
and a lack of germline as indicated by the white space in the body. Scale bar, 100 µm. **(C)** Total brood
was assessed in wild-type (gray), *n3536* canonical allele mutants (light pink), *syb6708* PHD finger
deletion mutants (magenta), and *syb6818* ATPase deletion mutants (teal). **(D)** Percent embryonic
survival, excluding *syb6818* due to complete sterility. For both graphs, dark line represents mean and
whiskers represent S.E.M. All statistical analyses were performed using ANOVA with a Šídák correction
(n.s. = not significant, $*P < 0.05$, $**P < 0.01$, $*** P < 0.001$, $**** P < 0.0001$). Summary statistics and data
are included in Table S5.

Figure S1: Cisplatin affects embryonic survival in a dose-dependent manner.

(**A, B**) Mean percent survival of wild-type populations (gray) exposed to doses of cisplatin ranging from 0 µM to 700 µM (A) or 0 µM to 400 µM (B). Relative dose levels are indicated by triangles under the X-axis. (B) Examining the effects of cisplatin on *let-418* mutants (pink), with *xpf-1* mutants (green) included as a positive control for cisplatin sensitivity. Whiskers represent S.E.M. For each condition, at least 25 broods were evaluated. Summary statistics and data are included in Table S6.

Figure S2: *let-418* mutants have poor oocyte quality due to hydroxyurea induced mitotic DSBs.

(**A, B**) Representative images of -1 oocytes in unexposed (A) and exposed (B) animals. Germlines are stained with DAPI (in white). Scale bar represents 2 µm. Summary statistics and data are included in Table S3.

Figure S3: Hydroxyurea affects embryonic survival in a dose-dependent manner.
Mean percent survival of wild-type (gray) and *let-418* mutants (pink) exposed to increasing doses of hydroxyurea (from 0 mM to 10 mM, with relative levels indicated by the blue triangle under the X-axis). Whiskers represent S.E.M. For each condition, at least 25 broods were evaluated. Summary statistics and data are included in Table S6.

Figure S4: Nitrogen mustard reduces embryonic survival in *let-418* and *fcd-2* mutants.
Percent survival of indicated genotypes with and without exposure nitrogen mustard; brown triangles indicate dose (0 mM, 100 mM, and 150 mM). Dark line represents mean percent survival and whiskers represent S.E.M. . Comparisons between wild-type and mutant populations at the same dose are above each mutant; other comparisons are indicated by horizontal lines. All statistical analyses were performed using ANOVA followed by Šídák's multiple comparisons (**$P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$). Summary statistics and data are included in Table S6.