NuRD chromatin remodeling is required to repair exogenous DSBs in the *Caenorhabditis elegans* germline

Deepshikha Ananthaswamy[1], Kelin Funes[1†], Thiago Borges[1†], Scott Roques[1], Nina Fassnacht[2], Sereen El Jamal[2], Paula M. Checchi[2], Teresa Wei-sy Lee[1]*

[1] Department of Biological Sciences, University of Massachusetts Lowell, 198 Riverside Dr. Lowell MA, 01854

[2] Department of Biology, Marist College, 3399 North Road, Poughkeepsie, NY 12601

† contributed equally

* for correspondence, teresa_lee@uml.edu

Running title: NuRD remodeling repairs germline exogenous DSBs

Keywords: DNA repair, NuRD (nucleosome remodeling and deacetylase complex), LET-418/CHD4, FCD-2/FANCD2, germline

ABSTRACT

Organisms rely on coordinated networks of DNA repair pathways to protect genomes against toxic

double-strand breaks (DSBs), particularly in germ cells. All repair mechanisms must successfully

negotiate the local chromatin environment in order to access DNA. For example, nucleosomes can be

repositioned by the highly conserved Nucleosome Remodeling and Deacetylase (NuRD) complex. In

Caenorhabditis elegans, NuRD functions in the germline to repair DSBs – the loss of NuRD's ATPase

subunit, LET-418/CHD4, prevents DSB resolution and therefore reduces fertility. In this study, we

challenge germlines with exogenous DNA damage to better understand NuRD's role in repairing DSBs.

We find that *let-418* mutants are hypersensitive to cisplatin and hydroxyurea: exposure to either

mutagen impedes DSB repair, generates aneuploid oocytes, and severely reduces fertility and embryonic

survival. These defects resemble those seen when the Fanconi anemia (FA) DNA repair pathway is

compromised, and we find that LET-418's activity is epistatic to that of the FA component FCD-

2/FANCD2. We propose a model in which NuRD is recruited to the site of DNA lesions to remodel

chromatin and allow access for FA pathway components. Together, these results implicate NuRD in the

repair of both endogenous DSBs and exogenous DNA lesions to preserve genome integrity in developing

germ cells.

ARTICLE SUMMARY

Preserving genome integrity in germ cells is critical for the survival of individuals and species. Our

previous work shows that nucleosome remodeling plays an important role in repairing meiotic DNA

damage. Here, we further challenge genomes with toxic DNA damaging agents to test the requirement

for remodeling in the germline. We find that DNA damage accumulates in the absence of remodeling,

which drastically reduces oocyte quality, and also show that the requirement for remodeling is epistatic

to the Fanconi anemia DNA repair pathway. These findings demonstrate that local chromatin

environments must be remodeled in response to DNA damage to maintain oocyte quality.

INTRODUCTION

DNA damage poses a danger to all cells, but if left unrepaired in germ cells, it can prove

deleterious for organismal fitness. During meiosis, any DNA lesion that persists until chromosome

segregation will generate an aneuploid gamete, which in turn causes infertility or embryonic lethality

(Ceccaldi et al., 2016; Kim et al., 2016). The evolutionary pressure to protect genomes for future

generations has generated multiple distinct, but overlapping, DNA repair pathways. These pathways

handle both endogenous programmed double-strand breaks (DSBs) required for crossover

recombination, along with any exogenous DNA damage incurred in the germline (Chapman et al., 2012;

Gartner and Engebrecht, 2022; Keeney, 2007). Studies in organisms ranging from budding yeast to

mammals have identified strong conservation among these repair pathways, which include error-free

mechanisms like homologous recombination and error-prone mechanisms like non-homologous end

joining (De Massy, 2013; Stinson and Loparo, 2021, 2021; Wilson et al., 1997). For example, the Fanconi

anemia (FA) pathway is activated when replication forks collide with DNA interstrand crosslinks, which

are one of the most toxic DNA lesions (Gartner and Engebrecht, 2022). Components of the FA pathway

coordinate the excision of the crosslink and conversion to DSBs, which can then be resolved by the

meiotic homologous recombination pathway (Adamo et al., 2010; Lachaud et al., 2016; Raghunandan et

al., 2015; Schlacher et al., 2012). Defects in the FA pathway are linked to an elevated risk of cancer and

cause chromosomal instability or meiotic catastrophe, hallmarks of the rare congenital disorder of

Fanconi anemia (Auerbach, 2009, 1993; Ceccaldi et al., 2016; De Winter and Joenje, 2009; Nalepa and

Clapp, 2018; Tischkowitz et al., 2003, 2004).

In eukaryotes, DNA repair must always happen in the context of chromatin, where DNA is

wrapped around histone octamers to form nucleosomes (Allis and Jenuwein, 2016). Nucleosomes act as

obstacles that control access to DNA, and therefore their spacing is regulated by nucleosome

remodeling complexes (Becker and Workman, 2013). The highly conserved remodeling complex,

Nucleosome remodeling and deacetylase (NuRD), includes ATPase remodelers from the chromodomain

helicase DNA binding (CHD) family (Basta and Rauchman, 2015; Kehle, 1998; Tong et al., 1998; Wade et

al., 1998; Woodage et al., 1997; Xue et al., 1998). In addition to CHD remodeling subunits, NuRD also

consists of the histone deacetylases HDAC1/2, histone demethylase LSD1/KDM1A, and other

components that allow NuRD to bind modified histones, methyl CpG, and other proteins (Basta and

Rauchman, 2015; Wang et al., 2009; Whyte et al., 2012). Initially, NuRD's primary role was thought to be

transcriptional repression; but studies have since identified a more nuanced involvement in gene

expression, along with other biological functions like DNA repair (Basta and Rauchman, 2015). The

nematode *Caenorhabditis elegans* has two catalytic CHD paralogs associated with the core NuRD

complex: CHD-3 (also known as Mi-2α or CHD3 in mammals) and LET-418 (Mi-2β or CHD4) (Passannante

et al., 2010; von Zelewsky et al., 2000). Mutations in either the *chd-3* or the *let-418* genes severely

reduce fertility due to high levels of persistent meiotic DSBs, (McMurchy et al., 2017; Turcotte et al.,

2018).

These studies highlight the importance of NuRD's role in the germline for proper DNA repair –

the loss of NuRD activity appears to shunt DSBs from error-free repair towards error-prone pathways

like NHEJ. The genomic instability of *chd-3* or *let-418* mutants resembles those seen in mutants

defective for the FA pathway (Rageul and Kim, 2020; Tsui and Crismani, 2019). One hallmark of

mutations in FA components is an increased sensitivity to mutagens like cisplatin, which induces DNA

interstrand crosslinks, or hydroxyurea, which stalls replication forks (Bailly and Gartner, 2011; Datta and

Brosh Jr., 2019; Youds et al., 2009). Both of these lesions can generate DSBs if left unrepaired, which

activate the S-phase checkpoint and subsequently delays cell cycle progression (Kessler and Yanowitz,

2014; Kim and Colaiácovo, 2015). Resolution of interstrand crosslinks by the FA pathway involves

recognition and binding by the core FA complex, which then recruits FANCD2 to chromatin by mono-

ubiquitination (Nakanishi et al., 2005; Niraj et al., 2019). Once at the DNA lesion, FANCD2 orchestrates

the repair of the lesion by converting it to DSBs, which are then resolved by homologous recombination

(Ceccaldi et al., 2016; Niraj et al., 2019).

NuRD is required for repairing DNA lesions in germ cells (McMurchy et al., 2017; Turcotte et al.,

2018), but it is not clear how or whether nucleosome remodeling interacts with repair processes like the

FA pathway. Here, we challenge the germline by inducing exogenous DNA damage to investigate the

relationship between the FA pathway and NuRD's catalytic component, LET-418/CHD4. *C. elegans* has

homologs for many core FA components, including FCD-2/FANCD2 (Kim et al., 2018; Lee et al., 2010).

One advantage of using *C. elegans* for this study is the germline's spatiotemporal organization, which

allows us to easily follow the effects of mutagen exposure at multiple downstream stages during

oogenesis (Jaramillo-Lambert et al., 2007; Kessler and Yanowitz, 2014). We show that LET-418 activity is

epistatic to FCD-2 and is necessary to repair germline DSBs and preserve fertility. These results support a

model where NuRD generates a permissive local chromatin environment at the site of DNA damage to

allow access to DSB repair machinery and therefore preserve gamete quality in the face of genotoxic

stress.

METHODS

Strains and husbandry: All strains were cultured using standard methods (Brenner, 1974) at 20°C on 6-

cm NGM agar plates spotted with OP50 *E. coli* grown overnight at 37°C in Luria Broth (LB). The *syb6818*

allele is a deletion of LET-418's ATPase domain and removes amino acids 590-1114. The *syb6708* allele is

a deletion of LET-418's PHD finger domain and removes amino acids 256-365. Both domains were

identified using an NCBI structure prediction and based on homology with human CHD4 (Farnung et al.,

2020; Sayers et al., 2022). Sterile strains were maintained over an *nT1* balancer chromosome by picking

GFP-positive heterozygotes. The following strains were obtained from the *Caenorhabditis* Genetics

Center or were generated by SunyBiotech (*syb* alleles, which were backcrossed five times to N2):

- N2: wild-type (Bristol isolate)

- MT14390: *let-418 (n3536) V*

- NB105: *fcd-2 (tm1298) IV*

- TG1660: *xpf-1 (tm2842) II*

- TER14: *fcd-2 (tm1298) IV; let-418 (n3536) V*

- TER6: *let-418 (syb6818)/nT1 [qls51] IV:V*

- TER10: *let-418 (syb6708) V*

Mutagen exposure: All exposure assays were performed on hermaphrodites cultured at 20°C on 3-cm

NGM agar plates. Plates were first spotted with OP50 *E. coli*, then treated by spreading 250 µL of either

400 µM cisplatin (Sigma Aldrich) or 10 mM hydroxyurea (Thermo Scientific) onto agar at least 12 hours

prior to experiment. Synchronized young adult hermaphrodites (12 hours post-L4 larval stage) were

placed on mutagen plates for 24 hours, then moved to regular NGM plates to recover before assessing

nuclei in the appropriate meiotic stage (timing for each assay described below) (Jaramillo-Lambert et al.,

2007; Kessler and Yanowitz, 2014).

Immunofluorescence and image processing: Following exposure to cisplatin or hydroxyurea,

hermaphrodites were allowed to recover on NGM plates for four hours to produce better gonad

extrusion. Gonad dissection, fixation, and immunostaining were performed as previously described

(Ananthaswamy et al., 2022). In brief, gonads were dissected using 25-gauge syringe needles, fixed in

0.8% paraformaldehyde (Electron Microscopy Services) for five minutes at room temperature, subjected

to freeze crack, and dehydrated in 100% methanol for one minute at -20°C. Slides were washed 3X with

PBS + 0.1% Tween-20 (Thermo Scientific) prior to overnight incubation at room temperature with

primary antibody. Slides were washed three times before incubation with secondary antibody for four

hours at room temperature. Samples were mounted in VectaShield (Vector Laboratories) with 2 µg/ml

DAPI (Thermo Life) before sealing with nail polish. Antibodies used were rabbit anti-RAD-51 at 1:5000

(gifted by Diana Libuda, (Kurhanewicz et al., 2020)) and donkey anti-rabbit AlexaFluor594 at 1:400

(Invitrogen).

Immunofluorescence images were collected at 63x as Z-stacks with 0.25 µm intervals using a

Zeiss Axio Imager M2 microscope with Zen 3.8 software (Zeiss). All representative images shown are

max projections after processing in FIJI software (Schindelin et al., 2012). Raw z-stacks were used to

quantify RAD-51 foci. Three biological replicates were performed, with five gonads imaged per condition

in each replicate (each condition had a total of 15 gonads analyzed). Only one gonad arm was analyzed

from each hermaphrodite. All statistical analyses were performed in Prism 8 (GraphPad), with p-values

determined using a Kruskal-Wallis test with Dunn's post-hoc test.

Analysis of diakinesis DAPI bodies: Following exposure to cisplatin or hydroxyurea, hermaphrodites were

allowed to recover for 60 hours, which is the time required for a mitotic oocyte nucleus to progress

through meiosis to diakinesis. Gonads were dissected as previously described (Ananthaswamy et al.,

2022) with the following modifications: the primary and secondary antibody incubation steps were

skipped, and slides were washed at least three times in PBS + 0.1% Tween-20 (Thermo Scientific) before

mounting in SlowFade Diamond (Molecular Probes) with 2 µg/ml DAPI (Thermo Life). Images were

collected at 63x as Z-stacks with 0.25 µm intervals captured on Zeiss Axio Imager M2 using Zen 3.8

software (Zeiss). All representative images shown are max projections after processing in FIJI (Schindelin

et al., 2012). Raw z-stacks were used to quantify DAPI-stained bodies in the -1 to the -4 oocyte from the

spermatheca. Three biological replicates were performed, with each condition consisting of a minimum

of 100 nuclei from >27 hermaphrodites. P-values were determined using a Fisher's exact test.

Embryonic survival – 4-hr laying period: After exposure to cisplatin, hydroxyurea, or nitrogen mustard,

hermaphrodites were allowed to recover on NGM plates for 60 hours, which is the length of time for

mitotic oocyte nuclei to progress through the gonad and be laid as fertilized embryos. Exposed mothers

were then moved onto freshly spotted 3-cm NGM plates (five to ten per plate) and allowed to lay for

four hours before being removed from plates. The number of embryos per plate was immediately

recorded after removal of mothers, and were scored as fertilized, unfertilized, or dead. Hatched progeny

were counted as adults three days later. Percent survival was calculated by dividing the number of

hatched progeny by the number of embryos, multiplied by 100. Each condition consists of progeny from

125-150 mothers, collected over three biological replicates. All statistical analyses were performed in

Prism 8 (GraphPad), with p-values determined from a one-way ANOVA after a Šídák correction used to

account for multiple comparisons.

Embryonic survival – entire brood: Individual hermaphrodites were isolated on separate plates as L4

larvae and transferred every 12 hours until the end of their fertile period (usually 5-6 days). Embryos

and unfertilized eggs were counted every 12 hours immediately after transfer of the mother. Total

brood was calculated for each mother as all embryos laid during the fertile period. Hatched progeny

were then counted two to three days later (when they were L4s or young adults). Percent survival was

calculated by dividing the number of hatched progeny by the number of embryos laid and multiplied by

100. Each biological replicate includes the broods of five mothers scored for each genotype (i.e. five

technical replicates). The following biological replicates were performed: seven of N2; six of *let-418*

(n3536); four of *let-418 (syb6818)*; and one of *let-418 (syb6708)*. Broods were censored from analysis if

the mother died of unnatural causes before the end of their fertile period (usually from vulva rupture or

crawling off the agar). All statistical analyses were performed in Prism 8 (GraphPad), with p-values

determined from a one-way ANOVA after a Šídák correction used to account for multiple comparisons.

RESULTS

LET-418 activity is required to repair mitotic DNA lesions.

Mutants lacking both NuRD paralogs (CHD3 and CHD4) have severely compromised fertility,

making downstream analysis difficult (Turcotte et al., 2018). Therefore, we focused only on the impact

of LET-418/CHD4, reasoning that it plays a larger role in repairing germline DSBs because its loss causes

more meiotic defects than that of CHD-3 (Turcotte et al., 2018). At pachytene entry, *let-418* mutants

have significantly more DSBs than wild-type (Turcotte et al., 2018), leading us to wonder whether these

DSBs represent unrepaired DNA damage acquired during replication. We used cisplatin and hydroxyurea

to generate exogenous DNA damage in *let-418 (n3536)* missense mutants and *fcd-2 (tm1298)* deletion

mutants, which lack the FA pathway component FCD-2/FANCD2. We capitalized on the fact that the *C.*

elegans germline is organized in a spatiotemporal manner, with nuclei undergoing mitosis at the distal

tip and moving proximally as they proceed through meiosis. The timing of how nuclei progress through

mitotic replication, meiosis, oocyte maturation, and fertilization has been extensively characterized

(Jaramillo-Lambert et al., 2007; Kessler and Yanowitz, 2014). We therefore estimated the timing post

exposure to capture the impact of DNA damage incurred during mitosis.

Mitotic zone DSBs were assessed using an antibody for the RecA recombinase RAD-51, a

common cytological marker (Alpi et al., 2003; Colaiácovo et al., 2003; Kurhanewicz et al., 2020).

Hermaphrodites were cultured on either cisplatin or hydroxyurea for twenty hours, allowed to recover

for four hours, and dissected for immunofluorescence (Jaramillo-Lambert et al., 2007; Kessler and

Yanowitz, 2014). Consistent with prior reports, unexposed wild-type animals accumulated few RAD-51

foci in the mitotic zone (Fig. 1A, C, Fig. 2A, C, and Table S1) (Alpi et al., 2003; Turcotte et al., 2018), which

was also true for all mutants examined, with foci in less than 8% of nuclei. Unexposed *let-418* single

mutants had a modest but significant increase in foci compared to wild-type animals ($P < 0.0001$,

Kruskal-Wallis test), but neither *fcd-2* single mutants nor *fcd-2; let-418* double mutants had more foci

than wild-type (Fig. 1A, $P > 0.08$ for both comparisons, Kruskal-Wallis test). Unexposed *fcd-2; let-418*

double mutants were not more affected than either single mutant ($P > 0.8$ for both comparisons,

Kruskal-Wallis test). We note that, although average foci number did not vary much between genotypes,

we occasionally observed nuclei with two or three foci in *let-418* single mutants (0.8% of nuclei) and *fcd-*

2; let-418 double mutants (0.7%), which were rarely seen in wild-type germlines (0.2%) (Fig. 1A and

Table S1).

As expected, exposure to 400 µM cisplatin raised RAD-51 focus number across all genotypes

compared to unexposed controls (Fig. 1B, D and Table S1). This dose was chosen based on a dose curve

showing that 400 µM significantly impacted embryonic survival in both wild-type and mutant

populations (Fig. S2). Wild-type populations exposed to cisplatin accumulated over five times more foci

than their unexposed controls $P < 0.0001$, Kruskal-Wallis test) (Fig. 1A, B). Similarly, both exposed *fcd-2*

and *let-418* mutants accumulated five times more foci than unexposed controls ($P < 0.0001$ for both

comparisons, Kruskal-Wallis test) (Fig. 1A, B). In *let-418* mutants, the increase was driven by nuclei with

multiple foci, sometimes as many as five (10.8% of nuclei compared to 0.7%). Interestingly, exposed *fcd-*

2; let-418 double mutants strongly resembled exposed *let-418* single mutants, with over seven times

more foci than unexposed controls ($P < 0.0001$, Kruskal-Wallis test) (Fig. 1B). Again, this large increase

was primarily driven by nuclei with multiple foci, especially those with four or five (9.6% of nuclei

compared to 0.7%). Notably, nuclei with four or five foci were never observed in exposed wild-type or

fcd-2 single mutants.

We next challenged mutants using hydroxyurea, which stalls replication forks and generates

DSBs if left unrepaired (Bailly and Gartner, 2011). A dose of 10 mM was chosen after examining

embryonic survival in wild-type and *let-418* mutants across a dose curve (Fig. S3 and Table S5).

Hydroxyurea affected all genotypes in a manner similar to cisplatin (Fig. 2 and Table S1). Once again, few

DSBs were observed in each unexposed population, with less than 7% of mitotic nuclei containing a

RAD-51 focus (Fig. 2A). As expected, exposure to hydroxyurea raised the average number of foci in all

genotypes ($P < 0.02$ for all comparisons, Kruskal-Wallis test) (Fig. 2B). Exposed wild-type populations and

fcd-2 mutants had nearly two times more foci than unexposed controls ($P < 0.02$ for both comparisons,

Kruskal-Wallis test), both primarily driven by an increase in the number of nuclei with one focus.

Resembling what we observed after cisplatin exposure, *let-418* single mutants and *fcd-2; let-418* double

mutants responded to hydroxyurea in similar ways (Fig. 2B, D). Both mutants had two times more foci

than unexposed controls ($P < 0.0001$ for both comparisons, Kruskal-Wallis test). We note that exposed

let-418 and *fcd-2; let-418* mutants each contained nuclei with as many as five RAD-51 foci, which were

almost never observed in wild-type or *fcd-2* mutant populations (Fig. 2B). Taken together, these data

show how exposure to cisplatin or hydroxyurea generates DNA lesions that require the activity of the

let-418 gene and the *fcd-2* gene for proper repair. Additionally, *fcd-2; let-418* double mutants

phenocopy *let-418* single mutants when responding to mitotic DNA damage in the germline.

<u>Unrepaired mitotic DSBs affect oocyte quality in *let-418* mutants.</u>

Although *let-418* single mutants accumulate DSBs in the mitotic region and during pachytene,

they do not display aneuploidy or chromosome fragmentation, indicating that any remaining DSBs are

fully resolved by diakinesis (Turcotte et al., 2018). However, mutants lacking both CHD paralogs, CHD-3

and LET-418, suffer from meiotic catastrophe, genome fragmentation, and chromosome fusions

(Turcotte et al., 2018), suggesting that NuRD needs at least one CHD component to repair endogenous

DSBs. To see whether LET-418 is required for the resolution of exogenous DNA lesions, we exposed

hermaphrodites to cisplatin or hydroxyurea for twenty hours and allowed them to recover for sixty

hours (Jaramillo-Lambert et al., 2007; Kessler and Yanowitz, 2014). This timing allows us to induce DNA

damage in mitotic cells and assess its impact during diakinesis, where we expect to see six DAPI-staining

bodies, one for each homologous pair of chromosomes.

As expected, unexposed wild-type animals rarely produced aneuploid gametes: when scoring

aged mothers in their third day of adulthood, we found that only 9% of oocytes contained five DAPI-

stained bivalents (Fig. 3A, C, E, Fig. S1, and Table S2). This result is consistent with previous studies

showing that oocyte quality decreases with maternal age (Achache et al., 2021; Andux and Ellis, 2008;

Luo et al., 2010; Scharf et al., 2021). Each unexposed mutant produced more aneuploid oocytes than

wild-type controls, with at least 16% of nuclei containing fewer than six DAPI bodies ($P < 0.01$ for all

comparisons, Fisher's exact test), including oocytes with four DAPI bodies that were never observed in

wild-type animals (Fig. 3A, C, E and Fig. S1). We also noted that *let-418* single mutants generated a rare

oocyte with seven DAPI bodies (Fig. 3E). The presence of univalents in this nucleus reflected the lack of a

chiasmata and failure of crossover formation, which almost never occurs in wild-type animals (Dernburg

et al., 1998; Yu et al., 2016). When exposed to 400 µM cisplatin, both *fcd-2* single mutants and *fcd-2; let-*

418 double mutants suffered a significant deterioration in oocyte quality ($P < 0.04$ for both comparisons,

Fisher's exact test) (Fig. 3B). Each mutant produced oocytes with more than six DAPI bodies, and *fcd-2;*

let-418 double mutants also produced an oocyte with only three DAPI bodies (Fig. 3F). Although

exposure to cisplatin did not induce a statistically significant change in wild-type and *let-418* single

mutants ($P > 0.5$ for both comparisons, Fisher's exact test), both genotypes produced oocyte classes that

were never observed in unexposed controls (Fig. 3E, F and Table S2).

When compared to either single mutant, *fcd-2; let-418* double mutants were more affected by

cisplatin exposure ($P < 0.003$ for both comparisons, Fisher's exact test). This difference was driven by

oocytes with both more and fewer DAPI bodies: *fcd-2; let-418* double mutants produced more nuclei

with four or fewer DAPI bodies, but also nuclei with seven to ten DAPI bodies, indicating a severe

disruption in crossover formation (Fig. 3F). Given our findings for mitotic DSBs, where the double

mutant phenocopied the *let-418* single mutant (Fig. 1 and Fig. 2), we were surprised to see that

diakinesis DAPI body number was significantly more affected in *fcd-2; let-418* double mutants than in

let-418 single mutants. However, both the double mutant and *let-418* single mutant produced the rare

category of nuclei with three DAPI bodies (Fig. 3F). The severe reduction in DAPI bodies indicated that at

least half of the *C. elegans* genome has experienced chromosomal fusions, a phenotype that was also

previously observed in mutants missing both CHD paralogs (Turcotte et al., 2018).

Like cisplatin, 10 mM hydroxyurea also induced aneuploidy during diakinesis (Fig. 3C, D and

Table S2). Although exposure to hydroxyurea did not cause a statistically significant change in wild-type

population or *fcd-2* single mutants ($P > 0.3$ for both comparisons, Fisher's exact test), both genotypes

produced oocyte classes that were never observed in unexposed controls (Fig. S1 and Table S2).

However, unlike what we observed after cisplatin exposure, we did not find that *fcd-2* single mutants

produced nuclei with more than six DAPI bodies after hydroxyurea exposure (Fig. 3D), indicating that

fcd-2 mutants were more sensitive to cisplatin. Exposed *let-418* single mutants were significantly

affected compared to wild-type controls, producing oocytes with both more than seven and fewer than

four DAPI bodies, categories that were never observed in unexposed controls ($P < 0.006$, Fisher's exact

test).

Challenging *fcd-2; let-418* double mutants with hydroxyurea caused a large deterioration in

oocyte quality (*P* < 0.009, Fisher's exact test). Exposed *fcd-2; let-418* double mutants were significantly

more affected than either of the exposed single mutant controls (*P* < 0.004 for both comparisons,

Fisher's exact test). This decline in oocyte quality was partly driven by nuclei with more than six DAPI

bodies (sometimes as many as ten), a category only observed in exposed *fcd-2; let-418* double mutants

and *let-418* single mutants (Fig. S1B). Once again, the existence of nuclei with univalents indicates that

hydroxyurea also causes a severe failure of crossover formation or chromosome fragmentation (Fig.

S1B). Taken together, this assessment of diakinesis nuclei indicates that *fcd-2; let-418* double mutants

are more sensitive to interstrand crosslinks and stalled replication forks than either single mutant.

<u>Mutagen exposure reduces embryonic survival of *let-418* mutants.</u>

After having shown that LET-418 is necessary to repair exogenous DSBs during mitosis (Fig. 1

and 2) and prevent aneuploidy during diakinesis (Fig. 3), we next examined oocyte viability by assessing

embryonic survival (Fig. 4 and Table S3). To capture the impact of mutagen exposure during mitosis, we

assessed embryos 60 hours later, which were produced by mothers in their third day of adulthood

(Jaramillo-Lambert et al., 2007; Kessler and Yanowitz, 2014). As expected for aged mothers, unexposed

wild-type populations had an average embryonic survival of 96% (Fig. 4A). As a positive control for

cisplatin sensitivity, we examined *xpf-1 (tm2842)* mutants, which lack an endonuclease involved in the

DNA damage response (Ward et al., 2007). Unexposed *xpf-1* mutants have a lower average survival than

unexposed wild-type populations (*P* < 0.007, ANOVA), which is reduced even further by cisplatin in a

dose-dependent manner (Fig. S2B and Table S5) (Meier et al., 2014; Ward et al., 2007). Survival in the

remaining unexposed mutants did not significantly differ from wild-type populations (*P* > 0.1 for all

comparisons, ANOVA), although each mutant displayed more variability and lower averages across

replicates compared to wild-type: across both cisplatin and hydroxyurea experiments, *fcd-2* single

mutants averaged 86% survival, *let-418* single mutants averaged 77% survival, and *fcd-2; let-418* double

mutants averaged 75% survival (Table S3). When challenged with 400 µM cisplatin, embryonic survival

in wild-type populations was significantly reduced to 73% (*P* < 0.04, ANOVA). As expected for a positive

control, cisplatin exposure reduced *xpf-1* mutant survival more severely than in wild-type populations,

dropping survival to 39% (*P* < 0.01, ANOVA) (Fig. 4A).

Embryonic survival in exposed *fcd-2* single mutants was not significantly affected compared to

unexposed controls (*P* > 0.8, ANOVA) (Fig. 4A). Conversely, exposed *let-418* single mutants experienced

a significant decline in survival, down to 39% (*P* < 0.0001, ANOVA), a decrease 2.4 times more than wild-

type and similar to *xpf-1* mutants (Fig. 4A). We noted that cisplatin had a variable impact on *let-418*

mutant survival, across both technical and biological replicates. This observation was consistent with

previous variability observed in *let-418* mutants (Turcotte et al., 2018): some replicates were completely

unaffected, resembling unexposed controls, whereas others suffered complete embryonic lethality that

was never observed in wild-type or *xpf-1* positive controls. Again, the effects of cisplatin on embryonic

survival in *fcd-2; let-418* double mutants strongly resembled *let-418* single mutants (Fig. 4A). Exposed

fcd-2; let-418 double mutants experienced a significant drop in survival down to 33% ($P < 0.0001$,

ANOVA). As seen in *let-418* single mutants, this reduction was 2.7 times more than wild-type and similar

to *xpf-1* positive controls. However, in one departure from *let-418* single mutants, we found that

cisplatin caused less variability in *fcd-2; let-418* double mutant survival, which had a distribution that

overlapped only the lower half of that observed in *let-418* single mutants (Fig. 4A).

We next challenged hermaphrodites with 10 mM hydroxyurea to assess the impact on

embryonic survival (Fig. 4B). As expected, this low dose of exposure did not significantly reduce survival

in wild-type populations or *fcd-2* single mutants ($P > 0.09$ for both comparisons, ANOVA) (Kim et al.,

2018). However, *let-418* single mutants were sensitive to hydroxyurea – embryonic survival dropped

from 76% in unexposed controls to 46% after exposure ($P < 0.0003$, ANOVA), a reduction 2.1 times more

than experienced by wild-type populations (Fig. 4B). Exposed *fcd-2; let-418* double mutants suffered

from an even greater decrease in survival, from 73% down to 36% ($P < 0.0001$, ANOVA), a reduction 2.8

times more than wild-type (Fig. 4B).

Finally, we challenged hermaphrodites with an additional mutagen, nitrogen mustard, which

creates both interstrand crosslinks and DNA-protein crosslinks (Fig. S4 and Table S5) (Povirk and Shuker,

1994). A low dose of 100 µM did not affect wild-type embryonic survival ($P > 0.9$, ANOVA), whereas a

higher dose of 150 µM significantly decreased wild-type survival to 52% ($P < 0.0001$, ANOVA) (Fig. S4).

The low 100 µM dose of nitrogen mustard decreased survival in both *let-418* and *fcd-2* single mutants (P

< 0.0001 for both comparisons, ANOVA) and the high 150 µM dose further reduced survival to 25% ($P <$

0.0001 for both comparisons, ANOVA). Nitrogen mustard affected *fcd-2; let-418* double mutants

similarly to each single mutant ($P > 0.6$ for all comparisons), with survival severely reduced to 10% by

the high 150 µM dose ($P < 0.0001$ compared to unexposed control, ANOVA) (Fig. S4). Taken together,

the impact of mutagen exposure on embryonic survival indicate that LET-418 is required to resolve

exogenous DNA lesions to maintain oocyte viability.

<u>LET-418's ATPase domain is necessary for its role in germline DSB repair.</u>

The protein sequences of *C. elegans* LET-418 and human CHD4 share extensive homology,

including three domains: an ATPase domain, a PHD finger domain, and a double chromodomain (Fig. 5A)

(Farnung et al., 2020; Käser-Pébernard et al., 2016; Passannante et al., 2010). Computational models of

CHD4 structure suggest that each domain has distinct functions: the ATPase domain slides nucleosomes

along DNA (Farnung et al., 2020), the PHD finger domains bind modified histones, and the

chromodomains primarily bind DNA (Bouazoune et al., 2002). Mutations discovered in human patients,

along with *in vitro* experiments of CHD4 activity, have implicated each of these domains in CHD4's

nucleosome remodeling activity (Farnung et al., 2020; Watson et al., 2012; Weiss et al., 2016), but the

impact of each domain has yet to be assessed *in vivo*. Because the kinetics of repair occur rapidly in

response to DNA lesions (Kochan et al., 2017; Nair et al., 2017), we hypothesized that LET-418's function

in the germline requires its ATPase domain to slide nucleosomes and expose the site of DNA damage.

Figures 1 through 4 of this paper relied on the canonical allele *let-418 (n3536)*, a point mutation

in the ATPase domain that is hypomorphic, but viable, at the normal maintenance temperature of 20°C

(Andersen et al., 2006; Turcotte et al., 2018). We engineered separation-of function deletion alleles in

the *let-418* gene using CRISPR-Cas9 (see Methods for details). We then assessed the number of embryos

produced throughout the entire laying period in each *let-418* mutant (a full brood), using the canonical

n3536 allele as a positive control (Fig. 5A and Table S4). Consistent with previous studies, wild-type

mothers produced broods that averaged 303 embryos (Fig 5C). Strikingly, mutants that lack the ATPase

domain*, syb6818*, were completely sterile and produced no embryos (Fig. 5C). In these ATPase deletion

mutants, we also noted a high occurrence of protruding vulvae and a clear body due to the lack of a fully

proliferated germline (Fig. 5B). Conversely, mutants that lack the PHD finger domain, *syb6708*, produced

an average brood of 268 embryos, which was not significantly lower than wild-type broods ($P > 0.7$,

ANOVA). Consistent with previous reports, average broods of canonical *n3636* mutants were 219

embryos, which represents a significant reduction compared to wild-type broods ($P < 0.001$, ANOVA).

We also noted the variability in canonical *n3636* mutant broods: some mothers produced at wild-type

levels (~300 embryos), whereas others were completely sterile (McMurchy et al., 2017; Turcotte et al.,

2018).

Because LET-418 is also involved in development (Erdelyi et al., 2017; Passannante et al., 2010;

von Zelewsky et al., 2000), we assessed embryonic survival throughout the full laying period. Average

survival of canonical *n3636* mutants was 95%, which represented a slight but significant reduction

compared average wild-type survival of 99% ($P < 0.05$, ANOVA). Similar to what we observed with brood

size, embryonic survival was not affected by the *syb6708* PHD deletion (Fig. 5D). Finally, we were unable

to assess survival in *syb6818* ATPase deletion mutants, because all mothers examined were sterile (N =

20). Altogether, our findings support a model in which NuRD's role in DNA repair during oogenesis

specifically requires LET-418's ATPase activity.

DISCUSSION

The germline must preserve genome integrity for all future generations, a task that requires

extensive coordination between DNA repair machinery and local chromatin environments (Chen and

Tyler, 2022; Clouaire and Legube, 2019). At the site of DNA damage, chromatin landscapes help

orchestrate repair using either error-prone or error-free DSB repair pathways (Chen and Tyler, 2022).

Although the mechanisms of DNA repair have been well-described across species, the role of chromatin

remodelers remains poorly understood. In this study, we show that LET-418, the ATPase remodeling

subunit of NuRD, is required to maintain oocyte quality when genomes are challenged by exogenous

DNA damage. After exposure to cisplatin or hydroxyurea, mutants with reduced LET-418 activity

accumulate more mitotic DSBs (Fig. 1 and Fig. 2), generate more aneuploid oocytes (Fig. 3), and suffer

from reduced embryonic viability (Fig. 4 and Fig. 5) – phenotypes that strongly resemble those seen in

mutants defective for the FA repair pathway (Adamo et al., 2010; Gartner and Engebrecht, 2022; Lee et

al., 2010). When testing the genetic relationship between NuRD and the FA pathway, we showed that

let-418 gene activity is epistatic to the FA gene *fcd-2*, which suggests that NuRD nucleosome remodeling

is necessary to allow proper functioning of the FA repair pathway.

LET-418/CHD4 has been implicated in biological functions that include maintaining repressive

chromatin states, repairing DNA damage, and defining cell fate during embryonic development (De Vaux

et al., 2013; Guerry et al., 2007; Käser-Pébernard et al., 2016; Kunert et al., 2009; Saudenova and Wicky,

2018; Turcotte et al., 2018; Unhavaithaya et al., 2002; von Zelewsky et al., 2000). In mammals, CHD4's

role in resisting DNA damage has led to its upregulation in multiple types of cancer: for example, in

glioblastoma, CHD4 overexpression is associated with poor prognosis, likely due to its role in promoting

DNA damage repair to assure cancer cell survival (Chudnovsky et al., 2014; McKenzie et al., 2019). Other

work in ovarian cancer cells, acute myeloid leukemia cells, and mammary cells has shown that depleting

CHD4 impairs homologous recombination and sensitizes tumorigenic cells to DNA damaging agents (Pan

et al., 2012; Polo et al., 2010; Sperlazza et al., 2015). Similarly, in *C. elegans*, mutations in LET-418 or its

paralog CHD3 prevent DSB repair throughout meiotic pachytene, suggesting that LET-418/CHD4's role is

highly conserved across taxa (Turcotte et al., 2018).

Our work identifies a new role for LET-418/CHD4 in response to genotoxic threats – in the

absence of LET-418's ATPase activity, mutants cannot repair exogenous DNA damage incurred during

mitosis, which ultimately reduces gamete viability and impairs organismal survival. In germlines, LET-418

is associated with other heterochromatin factors that work with small RNA pathways to silence

repetitive elements and protect the genome (McMurchy et al., 2017). The DSB repair defects observed

in *let-418* mutants may be caused by disruptions in heterochromatin formation or maintenance, leading

to higher sensitivity to DNA damaging agents. A similar effect has been observed with the loss of

another core component of NuRD, RBBP4 (LIN-53 in *C. elegans*), which sensitizes cells to DNA damaging

agents by establishing a permissive chromatin environment and downregulating RAD51 expression

(Kitange et al., 2016).

We propose that LET-418 and NuRD are required during at least two points in gametogenesis: to

repair exogenous damage accumulated in mitosis and to resolve endogenous DSBs formed during

meiosis (this study and Turcotte et al., 2018). In the absence of LET-418 or NuRD activity, mitotic

damage persists as nuclei enter meiosis, contributing to the increase of DSBs observed in pachytene

(Turcotte et al., 2018). In meiosis, the FA pathway repairs DNA damage caused by replication fork

stalling or interstrand crosslinks – after the lesion is recognized, FA component FANCD2 is mono-

ubiquitinated and recruited to the lesion, where it mediates a conversion to DSBs and repair by

homologous recombination (Ceccaldi et al., 2016; Niraj et al., 2019).

Cytological experiments show that FA components, including FCD-2/FANCD2, are recruited to

DNA lesions to mediate repair in the mitotic region of the germline (Collis et al., 2006; Kim et al., 2018;

Lee et al., 2010). Our results are consistent with FCD-2 playing a role in resolving mitotic DSBs – exposed

fcd-2 single mutants generated aneuploid oocytes that were never observed in wild-type populations:

some had chromosome fusions, whereas others experienced crossover failure (Fig. 3 and Fig. S1). These

defects represent genomic instability that is a consistent hallmark of defects with the FA pathway, and is

even used to diagnose human patients for Fanconi anemia (Adamo et al., 2010, 2010; Auerbach, 2009;

De Winter and Joenje, 2009; Lee et al., 2007; Rageul and Kim, 2020). However, we were surprised to find

that embryonic viability in *fcd-2* mutants was not affected by exposure to either cisplatin or hydroxyurea

(Fig. 4), since others have previously reported a cisplatin sensitivity in this mutant background (Adamo

et al., 2010; Collis et al., 2006; Germoglio et al., 2020). After examining the timing of exposure and

recovery used in those studies, we realized that they primarily assessed how damage induced during

late pachytene affected embryonic survival (Jaramillo-Lambert et al., 2007; Kessler and Yanowitz, 2014).

In contrast, we used a time course that allowed us to examine the effects of damage induced earlier

during mitosis. Altogether, our results suggest that FCD-2 and the FA pathway are involved in repairing

exogenous DSBs, but may not be the sole, or even the main, repair pathway at this point in

gametogenesis. However, in the later stages of meiotic prophase, all DSBs must be resolved before

chromosome segregation. Therefore, as nuclei approach diplotene, they rely on alternative error-prone

DSB repair pathways like NHEJ to resolve remaining DSBs (Macaisne et al., 2018; Smolikov et al., 2007).

These pathways usually result in non-crossovers and disruption of chiasmata formation, or can generate

chromosome translocations or fusions (Gartner and Engebrecht, 2022); we observed both outcomes in

diakinesis nuclei of *let-418* mutants.

We have also demonstrated that LET-418's activity is epistatic to that of FCD-2 and the FA

pathway – the *fcd-2; let-418* double mutant phenocopies the *let-418* single mutant when challenged by

exogenous DNA damage (Fig. 1, Fig. 2, and Fig. 4). When examining how mitotic DNA lesions affect

diakinesis DAPI body formation, we observed that *fcd-2; let-418* double mutants were more severely

affected by cisplatin and hydroxyurea than either single mutant (Fig. 3 and Fig. S1). Strikingly, double

mutant germlines contained classes of diakinesis nuclei that were only seen in one or the other single

mutant, suggesting an additive effect of losing both LET-418 and FCD-2 activity.

C. elegans LET-418 shares extensive homology with human CHD4, including an ATPase domain, a

PHD finger domain, and a double chromodomain (Käser-Pébernard et al., 2016; Passannante et al.,

2010). Computational models indicate that the ATPase domain repositions nucleosomes by sliding them

along DNA (Farnung et al., 2020). When examining human patient mutations identified in cancer or the

neurodevelopmental disorder Sifrim-Hitz-Weiss syndrome, we noticed that many missense mutations

occurred in the ATPase domain, indicating its importance for LET-418 and NuRD function. (Farnung et

al., 2020). To test the requirement for LET-418's ATPase activity *in vivo*, we engineered an endogenous

deletion of this domain (allele *syb6818*) and demonstrated that it is essential for LET-418's function in

the germline. ATPase deletion mutants were completely sterile (Fig. 5), and all homozygote mutants had

defective vulvas (Fig. 5B). Conversely, deletion of the PHD domain did not affect fertility or embryonic

survival, indicating that this domain is dispensable for LET-418's role in the germline. The ATPase

deletion phenotypes resemble those of *let-418* genetic null alleles, which were previously used to

characterize LET-418's role in vulval development (Guerry et al., 2007; von Zelewsky et al., 2000). In

somatic tissues, LET-418 belongs to two distinct complexes that have a conserved role in maintaining

cell identity – in addition to NuRD, it is also found in MEC, which consists of LET-418, the histone

deacetylase HDA-1, and the Krüppel-like protein MEP-1 (Hou et al., 2020; Käser-Pébernard et al., 2016;

Pfefferli et al., 2014; Unhavaithaya et al., 2002). As part of these complexes, LET-418 has roles in

specifying vulval fate, embryonic and larval development, and defining lifespan (De Vaux et al., 2013;

Erdelyi et al., 2017; Guerry et al., 2007; Käser-Pébernard et al., 2014; Passannante et al., 2010;

Saudenova and Wicky, 2018; von Zelewsky et al., 2000). Further characterization of these deletion

strains will establish their impact outside of the germline.

Based on our results, we propose the following model. When replication forks are blocked or

stalled, NuRD's remodeling activity creates a permissive chromatin environment that allows for the

recruitment of FA pathway components, like FCD-2, to repair the damage via homologous

recombination. In the absence of NuRD-mediated remodeling, the FA pathway is unable to repair DNA

lesions, causing the accumulation of mitotic and meiotic DSBs and their eventual resolution by error-

prone pathways later in meiosis. However, it is not clear from our findings whether NuRD functions

entirely upstream of the full FA pathway, or whether it mediates an intermediate step during interstrand

crosslink repair. For example, it is possible that DNA lesions are first recognized by the FA component

FNCM-1 (which acts upstream of FCD-2), which then recruits NuRD to the site of damage. In summary,

we've shown that LET-418's nucleosome remodeling activity is necessary to resolve germline DSBs by

error-free pathways like the FA pathway. This work highlights the important role of local chromatin

landscapes in coordinating DNA damage repair pathways during gametogenesis.

DATA AVAILIBILITY

Strains are available upon request. The authors affirm that all data necessary for confirming the

conclusions of the article are present within the article, figures, and tables.

ACKNOWLEDGEMENTS

We thank Diana Libuda and Monica Colaiacovo for generously sharing reagents, Cori Cahoon for

immunofluorescence advice, and WormBase for gene and sequence information (Davis et al., 2022).

Strains were provided by the *Caenorhabditis* Genetics Center, which is funded by the National Institutes

of Health (NIH) through the Office of Research Infrastructure Programs (P40 OD010440).

FUNDING

D. A. was supported by a UMass Lowell KCS Science Scholarship. T. B. and K. F. were supported by

fellowships from the UMass Lowell River Hawk Scholars Academy. K. F. was supported by the Urban

Massachusetts LSAMP program, which is funded by the National Science Foundation (EES 2308724). T.B.

was supported by the Society for Developmental Biology through a Choose Development! fellowship,

which is funded by the NIH (R25HD105600). This work was supported by the NIH, with grants

R15GM117479 and R15HD104115 to P.M.C. and grant R15GM144861 to T.W.L.

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

REFERENCES

Achache, H., Falk, R., Lerner, N., Beatus, T., Tzur, Y.B., 2021. Oocyte aging is controlled by
mitogen-activated protein kinase signaling. Aging Cell 20, e13386.
https://doi.org/10.1111/acel.13386

Adamo, A., Collis, S.J., Adelman, C.A., Silva, N., Horejsi, Z., Ward, J.D., Martinez-Perez, E.,
Boulton, S.J., La Volpe, A., 2010. Preventing Nonhomologous End Joining Suppresses
DNA Repair Defects of Fanconi Anemia. Molecular Cell 39, 25–35.
https://doi.org/10.1016/j.molcel.2010.06.026

Allis, C.D., Jenuwein, T., 2016. The molecular hallmarks of epigenetic control. Nat Rev Genet 17,
487–500. https://doi.org/10.1038/nrg.2016.59

Alpi, A., Pasierbek, P., Gartner, A., Loidl, J., 2003. Genetic and cytological characterization of the
recombination protein RAD-51 in *Caenorhabditis elegans*. Chromosoma 112, 6–16.
https://doi.org/10.1007/s00412-003-0237-5

Ananthaswamy, D., Croft, J.C., Woozencroft, N., Lee, T.W., 2022. *C. elegans* Gonad Dissection
and Freeze Crack for Immunofluorescence and DAPI Staining. JoVE 64204.
https://doi.org/10.3791/64204

Andersen, E.C., Lu, X., Horvitz, H.R., 2006. *C. elegans* ISWI and NURF301 antagonize an Rb-like
pathway in the determination of multiple cell fates. Development 133, 2695–2704.
https://doi.org/10.1242/dev.02444

Andux, S., Ellis, R.E., 2008. Apoptosis Maintains Oocyte Quality in Aging *Caenorhabditis elegans*
Females. PLoS Genet 4, e1000295. https://doi.org/10.1371/journal.pgen.1000295

Auerbach, A.D., 2009. Fanconi anemia and its diagnosis. Mutation Research/Fundamental and
Molecular Mechanisms of Mutagenesis 668, 4–10.
https://doi.org/10.1016/j.mrfmmm.2009.01.013

Auerbach, A.D., 1993. Fanconi anemia diagnosis and the diepoxybutane (DEB) test. Exp
Hematol 21, 731–733.

Bailly, A., Gartner, A., 2011. *Caenorhabditis elegans* Radiation Responses, in: DeWeese, T.L.,
Laiho, M. (Eds.), Molecular Determinants of Radiation Response. Springer New York,
New York, NY, pp. 101–123. https://doi.org/10.1007/978-1-4419-8044-1_5

Basta, J., Rauchman, M., 2015. The nucleosome remodeling and deacetylase complex in
development and disease. Translational Research 165, 36–47.
https://doi.org/10.1016/j.trsl.2014.05.003

Becker, P.B., Workman, J.L., 2013. Nucleosome Remodeling and Epigenetics. Cold Spring Harbor
Perspectives in Biology 5, a017905–a017905.
https://doi.org/10.1101/cshperspect.a017905

Bouazoune, K., Mitterweger, A., Längst, G., Imhof, A., Akhtar, A., Becker, P.B., 2002. The dMi-2
chromodomains are DNA binding modules important for ATP-dependent nucleosome
mobilization. The EMBO Journal 21, 2430–2440.
https://doi.org/10.1093/emboj/21.10.2430
Brenner, S., 1974. THE GENETICS OF *CAENORHABDITIS ELEGANS*. Genetics 77, 71–94.
https://doi.org/10.1093/genetics/77.1.71
Ceccaldi, R., Rondinelli, B., D'Andrea, A.D., 2016. Repair Pathway Choices and Consequences at
the Double-Strand Break. Trends in Cell Biology 26, 52–64.
https://doi.org/10.1016/j.tcb.2015.07.009
Chapman, J.R., Taylor, M.R.G., Boulton, S.J., 2012. Playing the End Game: DNA Double-Strand
Break Repair Pathway Choice. Molecular Cell 47, 497–510.
https://doi.org/10.1016/j.molcel.2012.07.029
Chen, Z., Tyler, J.K., 2022. The Chromatin Landscape Channels DNA Double-Strand Breaks to
Distinct Repair Pathways. Front. Cell Dev. Biol. 10, 909696.
https://doi.org/10.3389/fcell.2022.909696
Chudnovsky, Y., Kim, D., Zheng, S., Whyte, W.A., Bansal, M., Bray, M.-A., Gopal, S., Theisen,
M.A., Bilodeau, S., Thiru, P., Muffat, J., Yilmaz, O.H., Mitalipova, M., Woolard, K., Lee, J.,
Nishimura, R., Sakata, N., Fine, H.A., Carpenter, A.E., Silver, S.J., Verhaak, R.G.W.,
Califano, A., Young, R.A., Ligon, K.L., Mellinghoff, I.K., Root, D.E., Sabatini, D.M., Hahn,
W.C., Chheda, M.G., 2014. ZFHX4 Interacts with the NuRD Core Member CHD4 and
Regulates the Glioblastoma Tumor-Initiating Cell State. Cell Reports 6, 313–324.
https://doi.org/10.1016/j.celrep.2013.12.032
Clouaire, T., Legube, G., 2019. A Snapshot on the Cis Chromatin Response to DNA Double-
Strand Breaks. Trends in Genetics 35, 330–345.
https://doi.org/10.1016/j.tig.2019.02.003
Colaiácovo, M.P., MacQueen, A.J., Martinez-Perez, E., McDonald, K., Adamo, A., La Volpe, A.,
Villeneuve, A.M., 2003. Synaptonemal Complex Assembly in *C. elegans* Is Dispensable
for Loading Strand-Exchange Proteins but Critical for Proper Completion of
Recombination. Developmental Cell 5, 463–474. https://doi.org/10.1016/S1534-
5807(03)00232-6
Collis, S.J., Barber, L.J., Ward, J.D., Martin, J.S., Boulton, S.J., 2006. *C. elegans* FANCD2 responds
to replication stress and functions in interstrand cross-link repair. DNA Repair 5, 1398–
1406. https://doi.org/10.1016/j.dnarep.2006.06.010
Datta, A., Brosh Jr., R.M., 2019. Holding All the Cards—How Fanconi Anemia Proteins Deal with
Replication Stress and Preserve Genomic Stability. Genes 10, 170.
https://doi.org/10.3390/genes10020170
Davis, P., Zarowiecki, M., Arnaboldi, V., Becerra, A., Cain, S., Chan, J., Chen, W.J., Cho, J., Da
Veiga Beltrame, E., Diamantakis, S., Gao, S., Grigoriadis, D., Grove, C.A., Harris, T.W.,
Kishore, R., Le, T., Lee, R.Y.N., Luypaert, M., Müller, H.-M., Nakamura, C., Nuin, P.,
Paulini, M., Quinton-Tulloch, M., Raciti, D., Rodgers, F.H., Russell, M., Schindelman, G.,
Singh, A., Stickland, T., Van Auken, K., Wang, Q., Williams, G., Wright, A.J., Yook, K.,
Berriman, M., Howe, K.L., Schedl, T., Stein, L., Sternberg, P.W., 2022. WormBase in
2022—data, processes, and tools for analyzing *Caenorhabditis elegans*. Genetics 220,
iyac003. https://doi.org/10.1093/genetics/iyac003

De Massy, B., 2013. Initiation of Meiotic Recombination: How and Where? Conservation and
Specificities Among Eukaryotes. Annu. Rev. Genet. 47, 563–599.
https://doi.org/10.1146/annurev-genet-110711-155423
De Vaux, V., Pfefferli, C., Passannante, M., Belhaj, K., Von Essen, A., Sprecher, S.G., Müller, F.,
Wicky, C., 2013. The *Caenorhabditis elegans* LET-418/ Mi2 plays a conserved role in
lifespan regulation. Aging Cell 12, 1012–1020. https://doi.org/10.1111/acel.12129
De Winter, J.P., Joenje, H., 2009. The genetic and molecular basis of Fanconi anemia. Mutation
Research/Fundamental and Molecular Mechanisms of Mutagenesis 668, 11–19.
https://doi.org/10.1016/j.mrfmmm.2008.11.004
Erdelyi, P., Wang, X., Suleski, M., Wicky, C., 2017. A Network of Chromatin Factors Is Regulating
the Transition to Postembryonic Development in *Caenorhabditis elegans*. G3
Genes|Genomes|Genetics 7, 343–353. https://doi.org/10.1534/g3.116.037747
Farnung, L., Ochmann, M., Cramer, P., 2020. Nucleosome-CHD4 chromatin remodeler structure
maps human disease mutations. eLife 9, e56178. https://doi.org/10.7554/eLife.56178
Gartner, A., Engebrecht, J., 2022. DNA repair, recombination, and damage signaling. Genetics
220, iyab178. https://doi.org/10.1093/genetics/iyab178
Germoglio, M., Valenti, A., Gallo, I., Forenza, C., Santonicola, P., Silva, N., Adamo, A., 2020. In
vivo analysis of FANCD2 recruitment at meiotic DNA breaks in *Caenorhabditis elegans*.
Sci Rep 10, 103. https://doi.org/10.1038/s41598-019-57096-1
Guerry, F., Marti, C.-O., Zhang, Y., Moroni, P.S., Jaquiéry, E., Müller, F., 2007. The Mi-2
nucleosome-remodeling protein LET-418 is targeted via LIN-1/ETS to the promoter of
lin-39/Hox during vulval development in *C. elegans*. Developmental Biology 306, 469–
479. https://doi.org/10.1016/j.ydbio.2007.03.026
Hou, T., Cao, Z., Zhang, J., Tang, M., Tian, Y., Li, Y., Lu, X., Chen, Y., Wang, Hui, Wei, F.-Z., Wang,
L., Yang, Y., Zhao, Y., Wang, Z., Wang, Haiying, Zhu, W.-G., 2020. SIRT6 coordinates with
CHD4 to promote chromatin relaxation and DNA repair. Nucleic Acids Res 48, 2982–
3000. https://doi.org/10.1093/nar/gkaa006
Jaramillo-Lambert, A., Ellefson, M., Villeneuve, A.M., Engebrecht, J., 2007. Differential timing of
S phases, X chromosome replication, and meiotic prophase in the *C. elegans* germ line.
Developmental Biology 308, 206–221. https://doi.org/10.1016/j.ydbio.2007.05.019
Käser-Pébernard, S., Müller, F., Wicky, C., 2014. LET-418/Mi2 and SPR-5/LSD1 Cooperatively
Prevent Somatic Reprogramming of *C. elegans* Germline Stem Cells. Stem Cell Reports 2,
547–559. https://doi.org/10.1016/j.stemcr.2014.02.007
Käser-Pébernard, S., Pfefferli, C., Aschinger, C., Wicky, C., 2016. Fine-tuning of chromatin
composition and Polycomb recruitment by two Mi2 homologues during *C. elegans* early
embryonic development. Epigenetics & Chromatin 9, 39.
https://doi.org/10.1186/s13072-016-0091-3
Keeney, S., 2007. Spo11 and the Formation of DNA Double-Strand Breaks in Meiosis, in: Egel, R.,
Lankenau, D.-H. (Eds.), Recombination and Meiosis, Genome Dynamics and Stability.
Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 81–123.
https://doi.org/10.1007/7050_2007_026
Kehle, J., 1998. dMi-2, a Hunchback-Interacting Protein That Functions in Polycomb Repression.
Science 282, 1897–1900. https://doi.org/10.1126/science.282.5395.1897

Kessler, Z., Yanowitz, J., 2014. Methodological considerations for mutagen exposure in *C.*
elegans. Methods 68, 441–449. https://doi.org/10.1016/j.ymeth.2014.04.011

Kim, H.-M., Beese-Sims, S.E., Colaiácovo, M.P., 2018. Fanconi Anemia FANCM/FNCM-1 and
FANCD2/FCD-2 Are Required for Maintaining Histone Methylation Levels and Interact
with the Histone Demethylase LSD1/SPR-5 in *Caenorhabditis elegans*. Genetics 209,
409–423. https://doi.org/10.1534/genetics.118.300823

Kim, H.-M., Colaiácovo, M.P., 2015. DNA Damage Sensitivity Assays in *Caenorhabditis elegans*.
Bio Protoc 5, e1487.

Kim, S., Peterson, S.E., Jasin, M., Keeney, S., 2016. Mechanisms of germ line genome instability.
Seminars in Cell & Developmental Biology 54, 177–187.
https://doi.org/10.1016/j.semcdb.2016.02.019

Kitange, G.J., Mladek, A.C., Schroeder, M.A., Pokorny, J.C., Carlson, B.L., Zhang, Y., Nair, A.A.,
Lee, J.-H., Yan, H., Decker, P.A., Zhang, Z., Sarkaria, J.N., 2016. Retinoblastoma Binding
Protein 4 Modulates Temozolomide Sensitivity in Glioblastoma by Regulating DNA
Repair Proteins. Cell Reports 14, 2587–2598.
https://doi.org/10.1016/j.celrep.2016.02.045

Kochan, J.A., Desclos, E.C.B., Bosch, R., Meister, L., Vriend, L.E.M., van Attikum, H., Krawczyk,
P.M., 2017. Meta-analysis of DNA double-strand break response kinetics. Nucleic Acids
Res 45, 12625–12637. https://doi.org/10.1093/nar/gkx1128

Kunert, N., Wagner, E., Murawska, M., Klinker, H., Kremmer, E., Brehm, A., 2009. dMec: a novel
Mi-2 chromatin remodelling complex involved in transcriptional repression. EMBO J 28,
533–544. https://doi.org/10.1038/emboj.2009.3

Kurhanewicz, N.A., Dinwiddie, D., Bush, Z.D., Libuda, D.E., 2020. Elevated Temperatures Cause
Transposon-Associated DNA Damage in C. elegans Spermatocytes. Current Biology 30,
5007-5017.e4. https://doi.org/10.1016/j.cub.2020.09.050

Lachaud, C., Moreno, A., Marchesi, F., Toth, R., Blow, J.J., Rouse, J., 2016. Ubiquitinated Fancd2
recruits Fan1 to stalled replication forks to prevent genome instability. Science 351,
846–849. https://doi.org/10.1126/science.aad5634

Lee, K.Y., Chung, K.Y., Koo, H.-S., 2010. The involvement of FANCM, FANCI, and checkpoint
proteins in the interstrand DNA crosslink repair pathway is conserved in *C. elegans*. DNA
Repair 9, 374–382. https://doi.org/10.1016/j.dnarep.2009.12.018

Lee, K.Y., Yang, I., Park, J.-E., Baek, O.-R., Chung, K.Y., Koo, H.-S., 2007. Developmental stage-
and DNA damage-specific functions of *C. elegans* FANCD2. Biochemical and Biophysical
Research Communications 352, 479–485. https://doi.org/10.1016/j.bbrc.2006.11.039

Luo, S., Kleemann, G.A., Ashraf, J.M., Shaw, W.M., Murphy, C.T., 2010. TGF-β and Insulin
Signaling Regulate Reproductive Aging via Oocyte and Germline Quality Maintenance.
Cell 143, 299–312. https://doi.org/10.1016/j.cell.2010.09.013

Macaisne, N., Kessler, Z., Yanowitz, J.L., 2018. Meiotic Double-Strand Break Proteins Influence
Repair Pathway Utilization. Genetics 210, 843–856.
https://doi.org/10.1534/genetics.118.301402

McKenzie, L.D., LeClair, J.W., Miller, K.N., Strong, A.D., Chan, H.L., Oates, E.L., Ligon, K.L.,
Brennan, C.W., Chheda, M.G., 2019. CHD4 regulates the DNA damage response and
RAD51 expression in glioblastoma. Sci Rep 9, 4444. https://doi.org/10.1038/s41598-
019-40327-w

McMurchy, A.N., Stempor, P., Gaarenstroom, T., Wysolmerski, B., Dong, Y., Aussianikava, D.,
Appert, A., Huang, N., Kolasinska-Zwierz, P., Sapetschnig, A., Miska, E.A., Ahringer, J.,
2017. A team of heterochromatin factors collaborates with small RNA pathways to
combat repetitive elements and germline stress. eLife 6, e21666.
https://doi.org/10.7554/eLife.21666
Meier, B., Cooke, S.L., Weiss, J., Bailly, A.P., Alexandrov, L.B., Marshall, J., Raine, K., Maddison,
M., Anderson, E., Stratton, M.R., Gartner, A., Campbell, P.J., 2014. *C. elegans* whole-
genome sequencing reveals mutational signatures related to carcinogens and DNA
repair deficiency. Genome Res. 24, 1624–1636. https://doi.org/10.1101/gr.175547.114
Nair, N., Shoaib, M., Sørensen, C.S., 2017. Chromatin Dynamics in Genome Stability: Roles in
Suppressing Endogenous DNA Damage and Facilitating DNA Repair. IJMS 18, 1486.
https://doi.org/10.3390/ijms18071486
Nakanishi, K., Yang, Y.-G., Pierce, A.J., Taniguchi, T., Digweed, M., D'Andrea, A.D., Wang, Z.-Q.,
Jasin, M., 2005. Human Fanconi anemia monoubiquitination pathway promotes
homologous DNA repair. Proc. Natl. Acad. Sci. U.S.A. 102, 1110–1115.
https://doi.org/10.1073/pnas.0407796102
Nalepa, G., Clapp, D.W., 2018. Fanconi anaemia and cancer: an intricate relationship. Nat Rev
Cancer 18, 168–185. https://doi.org/10.1038/nrc.2017.116
Niraj, J., Färkkilä, A., D'Andrea, A.D., 2019. The Fanconi Anemia Pathway in Cancer. Annu. Rev.
Cancer Biol. 3, 457–478. https://doi.org/10.1146/annurev-cancerbio-030617-050422
Pan, M.-R., Hsieh, H.-J., Dai, H., Hung, W.-C., Li, K., Peng, G., Lin, S.-Y., 2012. Chromodomain
Helicase DNA-binding Protein 4 (CHD4) Regulates Homologous Recombination DNA
Repair, and Its Deficiency Sensitizes Cells to Poly(ADP-ribose) Polymerase (PARP)
Inhibitor Treatment. Journal of Biological Chemistry 287, 6764–6772.
https://doi.org/10.1074/jbc.M111.287037
Passannante, M., Marti, C.-O., Pfefferli, C., Moroni, P.S., Kaeser-Pebernard, S., Puoti, A.,
Hunziker, P., Wicky, C., Müller, F., 2010. Different Mi-2 Complexes for Various
Developmental Functions in *Caenorhabditis elegans*. PLoS ONE 5, e13681.
https://doi.org/10.1371/journal.pone.0013681
Pfefferli, C., Müller, F., Jaźwińska, A., Wicky, C., 2014. Specific NuRD components are required
for fin regeneration in zebrafish. BMC Biol 12, 30. https://doi.org/10.1186/1741-7007-
12-30
Polo, S.E., Kaidi, A., Baskcomb, L., Galanty, Y., Jackson, S.P., 2010. Regulation of DNA-damage
responses and cell-cycle progression by the chromatin remodelling factor CHD4. EMBO J
29, 3130–3139. https://doi.org/10.1038/emboj.2010.188
Povirk, L.F., Shuker, D.E., 1994. DNA damage and mutagenesis induced by nitrogen mustards.
Mutat Res 318, 205–226. https://doi.org/10.1016/0165-1110(94)90015-9
Rageul, J., Kim, H., 2020. Fanconi anemia and the underlying causes of genomic instability.
Environ and Mol Mutagen 61, 693–708. https://doi.org/10.1002/em.22358
Raghunandan, M., Chaudhury, I., Kelich, S.L., Hanenberg, H., Sobeck, A., 2015. FANCD2, FANCJ
and BRCA2 cooperate to promote replication fork recovery independently of the
Fanconi Anemia core complex. Cell Cycle 14, 342–353.
https://doi.org/10.4161/15384101.2014.987614

Saudenova, M., Wicky, C., 2018. The Chromatin Remodeler LET-418/Mi2 is Required Cell Non-
Autonomously for the Post-Embryonic Development of *Caenorhabditis elegans*. JDB 7, 1.
https://doi.org/10.3390/jdb7010001
Sayers, E.W., Bolton, E.E., Brister, J.R., Canese, K., Chan, J., Comeau, D.C., Connor, R., Funk, K.,
Kelly, C., Kim, S., Madej, T., Marchler-Bauer, A., Lanczycki, C., Lathrop, S., Lu, Z., Thibaud-
Nissen, F., Murphy, T., Phan, L., Skripchenko, Y., Tse, T., Wang, J., Williams, R., Trawick,
B.W., Pruitt, K.D., Sherry, S.T., 2022. Database resources of the national center for
biotechnology information. Nucleic Acids Research 50, D20–D26.
https://doi.org/10.1093/nar/gkab1112
Scharf, A., Pohl, F., Egan, B.M., Kocsisova, Z., Kornfeld, K., 2021. Reproductive Aging in
Caenorhabditis elegans: From Molecules to Ecology. Front. Cell Dev. Biol. 9, 718522.
https://doi.org/10.3389/fcell.2021.718522
Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J., Hartenstein, V., Eliceiri, K.,
Tomancak, P., Cardona, A., 2012. Fiji: an open-source platform for biological-image
analysis. Nat Methods 9, 676–682. https://doi.org/10.1038/nmeth.2019
Schlacher, K., Wu, H., Jasin, M., 2012. A Distinct Replication Fork Protection Pathway Connects
Fanconi Anemia Tumor Suppressors to RAD51-BRCA1/2. Cancer Cell 22, 106–116.
https://doi.org/10.1016/j.ccr.2012.05.015
Smolikov, S., Eizinger, A., Hurlburt, A., Rogers, E., Villeneuve, A.M., Colaiácovo, M.P., 2007.
Synapsis-defective mutants reveal a correlation between chromosome conformation
and the mode of double-strand break repair during *Caenorhabditis elegans* meiosis.
Genetics 176, 2027–2033. https://doi.org/10.1534/genetics.107.076968
Sperlazza, J., Rahmani, M., Beckta, J., Aust, M., Hawkins, E., Wang, S.Z., Zu Zhu, S., Podder, S.,
Dumur, C., Archer, K., Grant, S., Ginder, G.D., 2015. Depletion of the chromatin
remodeler CHD4 sensitizes AML blasts to genotoxic agents and reduces tumor
formation. Blood 126, 1462–1472. https://doi.org/10.1182/blood-2015-03-631606
Stinson, B.M., Loparo, J.J., 2021. Repair of DNA Double-Strand Breaks by the Nonhomologous
End Joining Pathway. Annu. Rev. Biochem. 90, 137–164.
https://doi.org/10.1146/annurev-biochem-080320-110356
Tischkowitz, M., Ameziane, N., Waisfisz, Q., De Winter, J.P., Harris, R., Taniguchi, T., D'Andrea,
A., Hodgson, S.V., Mathew, C.G., Joenje, H., 2003. Bi-allelic silencing of the Fanconi
anaemia gene *FANCF* in acute myeloid leukaemia: *Short Report*. British Journal of
Haematology 123, 469–471. https://doi.org/10.1046/j.1365-2141.2003.04640.x
Tischkowitz, M.D., Morgan, N.V., Grimwade, D., Eddy, C., Ball, S., Vorechovsky, I., Langabeer, S.,
Stöger, R., Hodgson, S.V., Mathew, C.G., 2004. Deletion and reduced expression of the
Fanconi anemia FANCA gene in sporadic acute myeloid leukemia. Leukemia 18, 420–
425. https://doi.org/10.1038/sj.leu.2403280
Tong, J.K., Hassig, C.A., Schnitzler, G.R., Kingston, R.E., Schreiber, S.L., 1998. Chromatin
deacetylation by an ATP-dependent nucleosome remodelling complex. Nature 395,
917–921. https://doi.org/10.1038/27699
Tsui, V., Crismani, W., 2019. The Fanconi Anemia Pathway and Fertility. Trends in Genetics 35,
199–214. https://doi.org/10.1016/j.tig.2018.12.007

Turcotte, C.A., Sloat, S.A., Rigothi, J.A., Rosenkranse, E., Northrup, A.L., Andrews, N.P., Checchi,
P.M., 2018. Maintenance of Genome Integrity by Mi2 Homologs CHD-3 and LET-418 in
Caenorhabditis elegans. Genetics 208, 991–1007.
https://doi.org/10.1534/genetics.118.300686
Unhavaithaya, Y., Shin, T.H., Miliaras, N., Lee, J., Oyama, T., Mello, C.C., 2002. MEP-1 and a
Homolog of the NURD Complex Component Mi-2 Act Together to Maintain Germline-
Soma Distinctions in *C. elegans*. Cell 111, 991–1002. https://doi.org/10.1016/S0092-
8674(02)01202-3
von Zelewsky, T., Palladino, F., Brunschwig, K., Tobler, H., Hajnal, A., Müller, F., Brehm, A., 2000.
The *C. elegans* Mi-2 chromatin-remodelling proteins function in vulval cell fate
determination. Development 127, 5277–5284. https://doi.org/10.1242/dev.127.24.5277
Wade, P.A., Jones, P.L., Vermaak, D., Wolffe, A.P., 1998. A multiple subunit Mi-2 histone
deacetylase from Xenopus laevis cofractionates with an associated Snf2 superfamily
ATPase. Current Biology 8, 843–848. https://doi.org/10.1016/S0960-9822(98)70328-8
Wang, Y., Zhang, H., Chen, Y., Sun, Y., Yang, F., Yu, W., Liang, J., Sun, L., Yang, X., Shi, L., Li, R., Li,
Y., Zhang, Y., Li, Q., Yi, X., Shang, Y., 2009. LSD1 Is a Subunit of the NuRD Complex and
Targets the Metastasis Programs in Breast Cancer. Cell 138, 660–672.
https://doi.org/10.1016/j.cell.2009.05.050
Ward, J.D., Barber, L.J., Petalcorin, M.I., Yanowitz, J., Boulton, S.J., 2007. Replication blocking
lesions present a unique substrate for homologous recombination. EMBO J 26, 3384–
3396. https://doi.org/10.1038/sj.emboj.7601766
Watson, A.A., Mahajan, P., Mertens, H.D.T., Deery, M.J., Zhang, Wenchao, Pham, P., Du, X.,
Bartke, T., Zhang, Wei, Edlich, C., Berridge, G., Chen, Y., Burgess-Brown, N.A.,
Kouzarides, T., Wiechens, N., Owen-Hughes, T., Svergun, D.I., Gileadi, O., Laue, E.D.,
2012. The PHD and chromo domains regulate the ATPase activity of the human
chromatin remodeler CHD4. J Mol Biol 422, 3–17.
https://doi.org/10.1016/j.jmb.2012.04.031
Weiss, K., Terhal, P.A., Cohen, L., Bruccoleri, M., Irving, M., Martinez, A.F., Rosenfeld, J.A.,
Machol, K., Yang, Y., Liu, P., Walkiewicz, M., Beuten, J., Gomez-Ospina, N., Haude, K.,
Fong, C.-T., Enns, G.M., Bernstein, J.A., Fan, J., Gotway, G., Ghorbani, M., DDD Study, van
Gassen, K., Monroe, G.R., van Haaften, G., Basel-Vanagaite, L., Yang, X.-J., Campeau,
P.M., Muenke, M., 2016. De Novo Mutations in CHD4, an ATP-Dependent Chromatin
Remodeler Gene, Cause an Intellectual Disability Syndrome with Distinctive
Dysmorphisms. Am J Hum Genet 99, 934–941.
https://doi.org/10.1016/j.ajhg.2016.08.001
Whyte, W.A., Bilodeau, S., Orlando, D.A., Hoke, H.A., Frampton, G.M., Foster, C.T., Cowley,
S.M., Young, R.A., 2012. Enhancer decommissioning by LSD1 during embryonic stem cell
differentiation. Nature 482, 221–225. https://doi.org/10.1038/nature10805
Wilson, T.E., Grawunder, U., Lieber, M.R., 1997. Yeast DNA ligase IV mediates non-homologous
DNA end joining. Nature 388, 495–498. https://doi.org/10.1038/41365
Woodage, T., Basrai, M.A., Baxevanis, A.D., Hieter, P., Collins, F.S., 1997. Characterization of the
CHD family of proteins. Proc. Natl. Acad. Sci. U.S.A. 94, 11472–11477.
https://doi.org/10.1073/pnas.94.21.11472

Xue, Y., Wong, J., Moreno, G.T., Young, M.K., Côté, J., Wang, W., 1998. NURD, a Novel Complex
with Both ATP-Dependent Chromatin-Remodeling and Histone Deacetylase Activities.
Molecular Cell 2, 851–861. https://doi.org/10.1016/S1097-2765(00)80299-3
Youds, J.L., Barber, L.J., Boulton, S.J., 2009. *C. elegans*: A model of Fanconi anemia and ICL
repair. Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis
668, 103–116. https://doi.org/10.1016/j.mrfmmm.2008.11.007

812
813
814 **FIGURE LEGENDS**
815
Figure 1: *let-418* mutants cannot repair mitotic DSBs when challenged with cisplatin.

(A, B) Stacked histograms showing the percentage of nuclei with the indicated number of foci for

animals exposed to 0 µM of cisplatin (A) and 400 µM of cisplatin (B). Average number of foci per nucleus

is indicated under each genotype. For all genotypes, exposed populations were significantly different

from unexposed controls (not indicated on histogram, $P < 0.0001$ for all comparisons). Each condition

includes at least 2000 nuclei analyzed from 15 germlines across three biological replicates. (**C, D**)

Representative images of mitotic region nuclei in unexposed (C) and exposed (D) animals, with the distal

tip oriented to the left (as indicated by white asterisk). Germlines are stained for RAD-51 (in yellow) and

DAPI (in blue). Scale bar represents 2 µm. All statistical analyses were performed using a Kruskal-Wallis

test with Dunn's post test (ns = not significant, * $P < 0.05$, **** $P < 0.0001$). Summary statistics are

included in Table S1.

Figure 2: *let-418* mutants cannot repair mitotic DSBs when challenged with hydroxyurea.

(A, B) Stacked histograms showing the percentage of nuclei with the indicated number of foci for

animals exposed to 0 mM of hydroxyurea (A) and 10 mM of cisplatin (B). Average number of foci per

nucleus is indicated under each genotype. For all genotypes, exposed populations were significantly

different from unexposed populations (not shown on graph, $P < 0.0001$ for all comparisons). Each

condition includes at least 2000 nuclei analyzed from 15 germlines across three biological replicates. (**C,**

D) Representative images of mitotic region nuclei in unexposed (C) and exposed (D) animals, with the

distal tip oriented to the left (as indicated by white asterisk). Germlines are stained for RAD-51 (in

yellow) and DAPI (in blue). Scale bar represents 2 µm. All statistical analyses were performed using a

Kruskal-Wallis test with Dunn's post test (ns = not significant, * $P < 0.05$, **** $P < 0.0001$). Summary

statistics are included in Table S1.

Figure 3: *let-418* mutants have poor oocyte quality due to mutagen induced mitotic DSBs.

(A, B, C, D) Stacked histograms showing the percentage of nuclei with the indicated number of DAPI

bodies for animals exposed to 0 µM of cisplatin (A), 400 µM of cisplatin, 0 mM of hydroxyurea (C) and 10

mM of hydroxyurea (D). Each condition includes at least 100 nuclei analyzed from 27 germlines across

three biological replicates. **(E, F)** Representative images of -1 oocytes in animals exposed to 0 uM of

cisplatin (E) and 400 uM of cisplatin (F). See Fig. S1 for representative images of -1 oocytes exposed to

cisplatin. Germlines are stained with DAPI (in white). Scale bar represents 2 µm. All statistical analyses

were performed as a Fisher's exact test on number of DAPI bodies (* $P < 0.05$, ** $P < 0.01$, **** $P <$

0.0001). Summary statistics are included in Table S2.

Figure 4: DNA damaging agents reduce embryonic viability in *let-418* mutants.

(**A, B**) Percent survival of indicated genotypes with and without exposure to cisplatin (A) or hydroxyurea

(B); dose is indicated with triangle above genotype names, with cisplatin in salmon (0 µM and 400 µM)

and hydroxyurea in blue (0 mM and 10 mM). For all conditions in (A) N=125 and (B) N=150. Dark line

represents the mean and whiskers represent S.E.M. All statistical analyses were performed using a one-

way ANOVA with a Šídák correction (*$P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$). Summary

statistics and data are found in Table S3.

Figure 5: The ATPase motor domain is necessary for LET-418 protein function in DSB repair. (A) Protein

alignment of human CHD4 and *C. elegans* LET-418 homologs showing conserved domains. Primary

structural alignments were performed according to NCBI-predicted domains using UniProt. The location

of the canonical *n3536* allele is indicated with a line. (**B**) Images of young adult hermaphrodites. The

ATPase deletion *syb6818*, has a protruding vulva (pVulv) phenotype (arrow) and a lack of germline as

indicated by the white space in the body. Scale bar, 100 µm. (**C**) Total brood was assessed in wild-type

(gray), the canonical allele *n3536* (light pink), the PHD finger deletion mutant *syb6708* (magenta), and

the ATPase deletion mutant *syb6818* (blue). (**D**) Percent embryonic survival, with the exception of *let-*

418(syb6818) due to its complete sterility. For both graphs, dark line represents mean and whiskers

represent S.E.M. All statistical analyses were performed using ANOVA with a Šídák correction (*$P < 0.05$,

*** $P < 0.001$, **** $P < 0.0001$). Summary statistics are included in Table S4.

Figure S1: let-418 mutants have poor oocyte quality due to hydroxyurea induced mitotic DSBs. (**A, B**)

Representative images of -1 to -4 oocytes in unexposed (C) and exposed (D) animals. Germlines are

stained with DAPI (in white). Scale bar represents 2 µm. Summary statistics are included in Table S2.

Figure S2: Cisplatin affects embryonic survival in a dose-dependent manner.

(**A, B**) Mean percent survival of wild-type populations (gray) exposed to doses of cisplatin ranging from 0 µM to 700 µM (A) or 0 µM to 400 µM (B). Relative dose levels are indicated by the triangles under the X-axis. (B) Examining the effects of cisplatin on *let-418* mutants (pink). *xpf-1* mutants (green) are included as a positive control for cisplatin sensitivity. Whiskers represent S.E.M. For each condition, at least 25 broods were evaluated. Summary statistics are included in Table S5.

Figure S3: Hydroxyurea affects embryonic survival in a dose-dependent manner. Mean percent survival of wild-type (gray) and *let-418* mutants (pink) exposed to increasing doses of hydroxyurea (from 0 mM to 10 mM, with relative levels indicated by the blue triangle under the X-axis). Whiskers represent S.E.M. For each condition, at least 25 broods were evaluated. Summary statistics are included in Table S5.

Figure S4: Nitrogen mustard reduces embryonic survival in *let-418* and *fcd-2* mutants. Percent survival of indicated genotypes with and without exposure nitrogen mustard; brown triangles indicate dose (0 mM, 100 mM, and 150 mM). Dark line represents mean percent survival and whiskers represent S.E.M. Statistical comparisons to wild-type at the same dose are indicated with asterisks above mutant data. All statistical analyses were performed using ANOVA followed by Šídák's multiple comparisons (**$P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$). Summary statistics are included in Table S5.

FIGURES WITH LEGENDS

Figure 1: *let-418* mutants cannot repair mitotic DSBs when challenged with cisplatin.
(A, B) Stacked histograms showing the percentage of nuclei with the indicated number of foci for animals exposed to 0 µM of cisplatin (A) and 400 µM of cisplatin (B). Average number of foci per nucleus is indicated under each genotype. For all genotypes, exposed populations were significantly different from unexposed controls (not indicated on histogram, $P < 0.0001$ for all comparisons). Each condition includes at least 2000 nuclei analyzed from 15 germlines across three biological replicates. **(C, D)** Representative images of mitotic region nuclei in unexposed (C) and exposed (D) animals, with the distal tip oriented to the left (as indicated by white asterisk). Germlines are stained for RAD-51 (in yellow) and DAPI (in blue). Scale bar represents 2 µm. All statistical analyses were performed using a Kruskal-Wallis test with Dunn's post test (ns = not significant, * $P < 0.05$, **** $P < 0.0001$). Summary statistics are included in Table S1.

Figure 2: *let-418* mutants cannot repair mitotic DSBs when challenged with hydroxyurea.
(A, B) Stacked histograms showing the percentage of nuclei with the indicated number of foci for animals exposed to 0 mM of hydroxyurea (A) and 10 mM of cisplatin (B). Average number of foci per nucleus is indicated under each genotype. For all genotypes, exposed populations were significantly different from unexposed populations (not shown on graph, $P < 0.0001$ for all comparisons). Each condition includes at least 2000 nuclei analyzed from 15 germlines across three biological replicates. **(C, D)** Representative images of mitotic region nuclei in unexposed (C) and exposed (D) animals, with the distal tip oriented to the left (as indicated by white asterisk). Germlines are stained for RAD-51 (in yellow) and DAPI (in blue). Scale bar represents 2 μm. All statistical analyses were performed using a Kruskal-Wallis test with Dunn's post test (ns = not significant, * $P < 0.05$, **** $P < 0.0001$). Summary statistics are included in Table S1.

Figure 3: *let-418* mutants have poor oocyte quality due to mutagen induced mitotic DSBs.
(A, B, C, D) Stacked histograms showing the percentage of nuclei with the indicated number of DAPI bodies for animals exposed to 0 µM of cisplatin (A), 400 µM of cisplatin, 0 mM of hydroxyurea (C) and 10 mM of hydroxyurea (D). Each condition includes at least 100 nuclei analyzed from 27 germlines across three biological replicates. **(E, F)** Representative images of -1 oocytes in animals exposed to 0 uM of cisplatin (E) and 400 uM of cisplatin (F). See Fig. S1 for representative images of -1 oocytes exposed to cisplatin. Germlines are stained with DAPI (in white). Scale bar represents 2 µm. All statistical analyses were performed as a Fisher's exact test on number of DAPI bodies (* $P < 0.05$, ** $P < 0.01$, **** $P < 0.0001$). Summary statistics are included in Table S2.

Figure 4: DNA damaging agents reduce embryonic viability in *let-418* mutants.

(**A, B**) Percent survival of indicated genotypes with and without exposure to cisplatin (A) or hydroxyurea (B); dose is indicated with triangle above genotype names, with cisplatin in salmon (0 µM and 400 µM) and hydroxyurea in blue (0 mM and 10 mM). For all conditions in (A) N=125 and (B) N=150. Dark line represents the mean and whiskers represent S.E.M. All statistical analyses were performed using a one-way ANOVA with a Šídák correction (*$P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$). Summary statistics and data are found in Table S3.

Figure 5: The ATPase motor domain is necessary for LET-418 protein function in DSB repair. (A) Protein alignment of human CHD4 and *C. elegans* LET-418 homologs showing conserved domains. Primary structural alignments were performed according to NCBI-predicted domains using UniProt. The location of the canonical *n3536* allele is indicated with a line. (**B**) Images of young adult hermaphrodites. The ATPase deletion *syb6818*, has a protruding vulva (pVulv) phenotype (arrow) and a lack of germline as indicated by the white space in the body. Scale bar, 100 µm. (**C**) Total brood was assessed in wild-type (gray), the canonical allele *n3536* (light pink), the PHD finger deletion mutant *syb6708* (magenta), and the ATPase deletion mutant *syb6818* (blue). (**D**) Percent embryonic survival, with the exception of *let-418(syb6818)* due to its complete sterility. For both graphs, dark line represents mean and whiskers represent S.E.M. All statistical analyses were performed using ANOVA with a Šídák correction (*$P < 0.05$, *** $P < 0.001$, **** $P < 0.0001$). Summary statistics are included in Table S4.

Figure S1: let-418 mutants have poor oocyte quality due to hydroxyurea induced mitotic DSBs. (A, B) Representative images of -1 to -4 oocytes in unexposed (C) and exposed (D) animals. Germlines are stained with DAPI (in white). Scale bar represents 2 μm. Summary statistics are included in Table S2.

Figure S2: Cisplatin affects embryonic survival in a dose-dependent manner.
(A, B) Mean percent survival of wild-type populations (gray) exposed to doses of cisplatin of ranging from 0 μM to 700 μM (A) or 0 μM to 400 μM (B). Relative dose levels are indicated by the triangles under the X-axis. (B) Examining the effects of cisplatin on *let-418* mutants (pink). *xpf-1* mutants (green) are included as a positive control for cisplatin sensitivity. Whiskers represent S.E.M. For each condition, at least 25 broods were evaluated. Summary statistics are included in Table S5.

Figure S3: Hydroxyurea affects embryonic survival in a dose-dependent manner. Mean percent survival of wild-type (gray) and *let-418* mutants (pink) exposed to increasing doses of hydroxyurea (from 0 mM to 10 mM, with relative levels indicated by the blue triangle under the X-axis). Whiskers represent S.E.M. For each condition, at least 25 broods were evaluated. Summary statistics are included in Table S5.

Figure S4: Nitrogen mustard reduces embryonic survival in *let-418* and *fcd-2* mutants. Percent survival of indicated genotypes with and without exposure nitrogen mustard; brown triangles indicate dose (0 mM, 100 mM, and 150 mM). Dark line represents mean percent survival and whiskers represent S.E.M. Statistical comparisons to wild-type at the same dose are indicated with asterisks above mutant data. All statistical analyses were performed using ANOVA followed by Šídák's multiple comparisons (**$P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$). Summary statistics are included in Table S5.