PIP5K-Ras bistability initiates plasma membrane symmetry breaking to regulate cell polarity and migration

Yu Deng[1,2], Tatsat Banerjee[1,2,6], Dhiman Sankar Pal[1,6], Parijat Banerjee[3], Huiwang Zhan[1], Jane Borleis[1], Pablo A. Igleias[1,4], Peter N. Devreotes[1,5]*

[1]Department of Cell Biology and Center for Cell Dynamics, School of Medicine, Johns Hopkins University, Baltimore, MD, USA.

[2]Department of Chemical and Biomolecular Engineering, Whiting School of Engineering, Johns Hopkins University, Baltimore, MD, USA.

[3]Department of Physics & Astronomy, Johns Hopkins University, Baltimore, MD, USA.

[4]Department of Electrical and Computer Engineering, Whiting School of Engineering, Johns Hopkins University, Baltimore, MD, USA.

[5]Department of Biological Chemistry, School of Medicine, Johns Hopkins University, Baltimore, MD, USA.

[6]These authors contributed equally to this work

*Email for correspondence: pnd@jhmi.edu

Abstract

Symmetry breaking, polarity establishment, and spontaneous cell protrusion formation are fundamental but poorly explained cell behaviors. Here, we demonstrate that a biochemical network, where the mutually inhibitory localization of PIP5K and Ras activities plays a central role, governs these processes. First, in resting cells devoid of cytoskeletal activity, PIP5K is uniformly elevated on the plasma membrane, while Ras activity remains minimal. Symmetry is broken by spontaneous local displacements of PIP5K, coupled with simultaneous activations of Ras and downstream signaling events, including PI3K activation. Second, knockout of PIP5K dramatically increases both the incidence and size of Ras-PI3K activation patches, accompanied by branched F-actin assembly. This leads to enhanced cortical wave formation, increased protrusive activity, and a shift in migration mode. Third, high inducible overexpression of PIP5K virtually eliminates Ras-PI3K signaling, cytoskeletal activity, and cell migration, while acute recruitment of cytosolic PIP5K to the membrane induces contraction and blebs in cancer cells. These arrested phenotypes are reversed by reducing myosin II activity, indicating myosin's involvement in the PIP5K-Ras-centered regulatory network. Remarkably, low inducible overexpression of PIP5K unexpectedly facilitates polarity establishment, highlighting PIP5K as a highly sensitive master regulator of these processes. Simulations of a computational model combining an excitable system, cytoskeletal loops, and dynamic partitioning of PIP5K recreates the experimental observations. Taken together, our results reveal that a bistable, mutually exclusive localization of PIP5K and active Ras on the plasma membrane triggers the initial symmetry breaking. Coupled actomyosin reduction and increased actin polymerization lead to intermittently extended protrusions and, with

40 feedback from the cytoskeleton, self-organizing, complementary gradients of PIP5K
41 versus Ras steepen, raising the threshold of the networks at the rear and lowering it at
42 the front to generate polarity for cell migration.

Introduction

The diverse morphological behaviors displayed by cells all depend on the ability to
achieve an asymmetric form spontaneously or in response to an external or internal cue
[1-6]. During migration, for example, a cell must generate and maintain a distinct front
and back [7-10]. Signaling and cytoskeletal components and activities self-organize
precisely into specific front or back regions of the cortex/plasma membrane [2, 6, 11-20].
This complementary organization is conserved across various processes, including
macropinocytosis, phagocytosis, cytokinesis, and apical-basal polarity in epithelial cells
[21-25]. Our studies focus on the role of PIP5K in regulating this elegant spatiotemporal
organization during migration. Still, the insights gained should apply broadly to symmetry-
breaking processes.

Symmetry-breaking is a critical initial step, but further events must follow to achieve
effective cell migration [6, 21, 26-33]. In human leukocytes and epithelial cells, as well as
Dictyostelium amoebae, protrusions underlying cell movement are driven by
spontaneously triggered waves of Ras, PI3K, which couple to cytoskeletal events at the
front, while PTEN, RhoA, and myosin II assembly retreat from these active areas [22, 26,
34-42]. Additional important aspects of migration are directional sensing and polarity,
where cells consistently display the same front-back complementary relationship of all
these molecular events [36]. Uncovering the relationship that links the downstream
processes to the initial symmetry-breaking events would provide a deep understanding
of the basis of dynamic cellular morphology.

The initial symmetry-breaking event in migration, which can occur even in the
absence of cytoskeletal activities, is the spontaneous local activation of Ras [35, 43-45];
however, the mechanism that controls Ras activation at specific membrane regions
remains obscure. Whatever the mechanism, it must amplify the signal locally and
simultaneously restrict it to prevent the activation elsewhere. Recent studies show that
PIP2 levels are depleted on the spontaneous protrusions [19]. Globally lowering PI(4,5)P2
activates Ras and increases the size of protrusions in *Dictyostelium* [46], and causes
spreading in human cells [34]. Reasoning that PIP5K could be the central regulator of
PIP2 production [47-50], we focused on the roles of this enzyme in fine-tuning Ras
activation.

Here, we report remarkably conserved localizations and outsized roles of PIP5Ks
in cell migration in amoebae, neutrophils, macrophages, and cancer cells and delineate
the function of these enzymes in cellular behavior. A central finding is a bistable, mutually
exclusive localization of PIP5Ks and active Ras on the plasma membrane, leading to
symmetry-breaking and locally regulated signal transduction network activities. These
activities are coupled to and amplified by interactions and feedback from the cytoskeleton.
The dramatic effects of deletion or overexpression of the enzymes on protrusions, cell
migration, and polarity demonstrate the significance of the spatial-temporal regulation of

the PIP5Ks. These findings are incorporated into a new model that seamlessly links
symmetry-breaking, cell polarity, and migration.

Results

Lowering PIP2 increases signal transduction, cytoskeletal, and protrusive
activities

The inconsistency of a report that PIP5K had little effect on random cell migration
with our previous reports of strong phenotypes caused by lowering PI(4,5)P2, prompted
us to reexamine this issue [51]. More detailed observation of *pi5k- (piki-)* cells revealed a
heterogeneous spectrum of migratory patterns (Figure 1a-h). While a minority of cells
retained wild-type amoeboid behavior, most exhibited fan-shaped, keratocyte-like
movements or oscillatory spreading/contracting movements, consistent with the
previously reported phenotype induced by abrupt reduction of PI(4,5)P2 in wild-type cells
[46]. These observations were supported by color-coded temporal overlay profiles (Figure
1a-c). Tracking experiments showed that fan-shaped cells migrate rapidly, with speeds
reaching $16.9 \pm 2.2\,\mu m\,min^{-1}$ (Figure 1e, 1g), in contrast to the $6.8 \pm 2.8\,\mu m\,min^{-1}$
observed in WT cells (Figure 1d, 1g). Since the spreading phases of the oscillatory cells
were asymmetric, these cells also displayed increased mobility, moving at a rate of around
$11.2 \pm 3\,\mu m\,min^{-1}$ (Figure 1f-g). Quantification of cell area of the entire population (Figure
1h) revealed an overall 4-fold increase, primarily due to flattening, although some of the
increase was due to a multinuclear phenotype of *pi5k-* cells (Figure S1c-d). These defects
in migration probably explain the inability of *pi5k-* cells to aggregate and form multicellular
structures in a timely fashion (Figure S1e). Most of these studies were carried out on the
original *pi5k-* line generated by homologous recombination [51]. To validate that the
phenotypes were attributable to loss of PI5K activity, we used CRISPR-mediated
disruption of PIP5K to create an independent loss-of-function cell line. These cells
displayed a consistent heterogeneous phenotype, primarily consisting of fan-shaped and
oscillatory cell populations, with cells more spread than wild-type cells (Figure S1a-b).

To further understand the basis of these phenotypes, we examined a series of
signal transduction and cytoskeletal activities using biosensors. As expected, PH-PLCδ
was mainly found on the membrane of WT cells, suggesting a significant level of PI(4,5)P2
[19], but there was no apparent membrane association of the biosensor in most of the
pi5k- cells (Figure 1i, 1k, Video S1). Cell boundary-to-cytosol quantification showed an
apparent 78% decrease of PH-PLCδ on the membrane in *pi5k-* (Figure 1k). Curiously, a
few outlier cells showed a bright signal comparable to WT cells (Figure 1k); these cells
are under investigation. Next, we examined the spatial distribution of Ras activity using a
Ras binding domain (RBD) biosensor in WT and *pi5k-* cells. Compared with WT cells,
which have characteristic patches at the cell protrusions, *pi5k-* cells have much broader
RBD patches, nearly a 2.9-fold increase (Figure 1j, 1l, Video S2). These observations
were supported by membrane kymograph analyses (Figure 1m-n). Similar results were
observed using the PHcrac biosensor to examine PIP3 accumulation between *pi5k-* cells
and WT cells, with a 2.1-fold increase in *pi5k-* cells (Figure S2a, 2c, Video S2). LimE, a
biosensor reflecting newly formed F-actin, also displayed much broader patches in *pi5k-*
cells, with a 3-fold increase in *pi5k-* cells (Figure S2b, 2d, Video S2). The dynamic

behavior of cytoskeletal activities was captured in membrane kymographs (Figure S2h-
i), which also revealed that the signaling activities oscillate in oscillatory cells (Figure S1f-
g). When cells spread, the signaling activities increase, while they disappear upon cell
shrinking. The three biosensors examined are typically associated with the active, "front-
state" of the cell, and the elevated levels suggest that *pi5k-* cells are vastly more activated
than wild-type cells.

Next, we examined signal transduction and cytoskeletal activities typically
associated with the "back-state" of cells. CynA is a biosensor that reports the level of
PI(3,4)P2 [52]. Strikingly, there generally was no detectable signal from CynA biosensor
in *pi5k-* cells. This reduction is indirect since PI(3,4)P2 is not a product of PI5K. (Figure
S2f, Video S1). However, in oscillatory cells, CynA did display a transient patch when cells
contracted, and this activity disappeared upon cell spreading (Figure S1h), indicating that
signaling activities at the cell back can also oscillate. Similar results were observed when
we examined cytoskeletal activities at the cell back with Myosin II (Figure S2e, Video S1).
These results suggested that cell activities are low at the cell back.

We also observed the same biosensors within ventral waves in the *pi5k-* cells. As
explained above, cellular protrusions consist of spontaneously initiated waves of signal
transduction and cytoskeletal activities. The waves are more conveniently observed along
the basal surface than in confocal slices. On the ventral surface, biosensors for Ras and
PI3K activity and F-actin appear as broad propagating regions, while "back-state"
sensors, CynA and myosin II, leave these zones, creating traveling "shadow waves." As
shown in Figure S2n-o, ventral waves are not apparent in single WT cells and are typically
visualized in electro-fused giant cells. However, they are readily apparent without
electrofusion in the flat, multinucleated *pi5k-* cells [41, 46, 53-55] (Figure S2j-k, Video S3).

Since the phenotypes of *pi5k-* cells were similar to those induced by reduction of
PI(4,5)P2 in *Dictyostelium* cells [46], we used recruitment of Inp54p to extend our studies
to differentiated HL-60 neutrophils and macrophages. We developed a recruitable Inp54p
by fusing it with CRY2PHR-mCherry, which enabled light-induced association with
membrane anchor CIBN-CAAX (Figure 1r). Globally recruiting Inp54p to the plasma
membrane induced fan-shaped cells and increased F-actin activities at the protrusions
reported by the LifeAct biosensor in both differentiated HL-60 neutrophils (Figure 1o,
Video S4) and macrophages (Figure S3a, Video S4). Recruiting an empty vector did not
cause a cell shape change (Figure S3e, Video S4) [44]. These observations were
supported by membrane kymograph analyses (Figure S3j, S3l) and color-coded temporal
overlay profiles (Figure S3k, S3m). Across the population, upon Inp54p recruitment,
differentiated HL-60 neutrophils and macrophages induced a 1.43-fold and 1.5-fold
increase in the cell area and aspect ratio, respectively (Figure 1p, Figure S3b). Cell
migration speed also increased from $4.0 \pm 1.4 \, \mu m \, min^{-1}$ (488 nm OFF) to $6.3 \pm$
$2.2 \, \mu m \, min^{-1}$ (488 nm ON) for differentiated HL-60 neutrophils (Figure 1q, S3f-g) and 4.0
$\pm 1.4 \, \mu m \, min^{-1}$ (488 nm OFF) to $6.3 \pm 2.2 \, \mu m \, min^{-1}$ (488 nm ON) for differentiated HL-60
macrophages (Figure S3c, S3h-i). Although cells became fan-shaped with one broad
lamellipodia structure at the cell front, cell polarity quantified by aspect ratio did not
change significantly (Figure S2g, S3d).

In summary, lowering PI(4,5)P2 levels by depleting PIP5K or recruiting Inp54p has
profound effects on the overall state of cells, generally causing *Dictyostelium* cells,
neutrophils, and macrophages to spread with broader protrusions and increased speed
(Figure 1s). Increased "front-state" and reduced "back-state" signal transduction and
cytoskeletal activities all suggest that cells are highly activated. These results imply that
PI5K, by regulating its product, PI(4,5)P2, is a key negative regulator of cell behavior.

PIP5Ks localize to back-state regions of the membrane

To investigate the localization of PIP5K and whether it could shed light the
decreases in PI(4,5)P2 at protrusions [19], we co-expressed PIP5K with a typical front-
state biosensor PHcrac in *Dictyostelium* cells. The distribution of PIP5K was
complementary to PHcrac and was depleted from protrusions in migrating single cells
(Figure 2a, Video S5). The back-state localization of PIP5Ks was conserved in HL60-
differentiated neutrophils. Both PIP5K1B and PIP5K1C isomers displayed a higher signal
on the back-state of the membrane in neutrophils (Figure 2b-c, Video S5) during random
migration, which was consistent with previously reported PIP5K1B and PIP5K1C back
localization during chemotaxis to N-formyl-met-leu-phe (fMLP) [49, 56].

We then examined ventral wave activities of the same cell lines in Figure 2a and
found, consistently, that PI5K was depleted in the front-state regions marked by high PIP3
and high Ras (Figure 2d, S4a, Video S6). The complementary pattern of PIP5K and PIP3
or PIP5K and Ras was maintained dynamically as the waves traveled across the basal
surface of the cells. Line scans showed about 80% less PIP5K in the front-state regions
compared with the back-state regions in *Dictyostelium* cells. Next, we co-expressed
PIP5K and PI(4,5)P2 biosensor PH-PLCδ and observed ventral wave activities. PIP5K
and PH-PLCδ travel together as "shadow" waves; they co-localize at the back-state
regions of the membrane (Figure 2e, Video S6). Line scans showed that PIP5K and PH-
PLCδ are largely co-localized, suggesting that the redistribution of PIP5K is the basis
PI(4,5)P2 is localized at the back-state of the membrane. Reciprocal patterns of PIP5K1B
and LifeAct were observed during frustrated phagocytosis in HL-60-differentiated
macrophages (Figure 2g, Video S6) and in RAW 264.7 macrophages, as previously
reported [54]. These results suggest that the localization of PIP5K to the rear and back-
state of the cells is a highly conserved phenomenon.

Previous studies have shown that signal transduction events can be triggered, and
the membrane can be spontaneously segregated into front- and back-states in the
absence of F-actin [40, 46, 57-59]. To test whether the spatiotemporal separation of
PIP5K depends on the existence of actin barrier between front and back states, we
treated *Dictyostelium* cells with Latrunculin A. We observed that on the ventral surface of
these cytoskeleton-impaired cells, the asymmetric waves of PI5K can propagate,
maintaining consistent complementarity to Ras-rich domains (Figure 2f, Videos S6).
Furthermore, the data in *Dictyostelium* indicates that the localization of PIP5K is
independent of the cytoskeletal network.

Typically, when cells are activated with global chemoattractant stimulation, the
front-state biosensors, such as PHcrac, transiently translocate to the membrane while
back-state proteins, such as PTEN, dissociate [60-64]. We therefore anticipated that,

based on its back-state localization, PIP5K would transiently move to the cytosol during
stimulation. However, upon adding cAMP to differentiated cells, PIP5K remained on the
membrane even as the front-state indicator PHcrac displayed its typical transient
translocation to the membrane, demonstrating robust receptor activation. (Figure S4b-c,
Video S7). PH-PLCδ biosensor also did not dramatically translocate to the cytosol,
although there was a 5% shift upon global stimulation (Figure S4d-e, Video S7).

When PIP5K is depleted in the front region as protrusions form, it remains on the
membrane, suggesting that it must redistribute to higher levels elsewhere. To more
definitively test this, we designed a KikGR-tagged PIP5K to photo convert a specific
region on the plasma membrane. As Figure 2h and Video S8 shows, PIP5K remains on
the membrane where the region was converted and gradually diffused along the
membrane. The red fluorescence intensity on the membrane increased in three steps
corresponding to three conversions in different regions (Figure 2i). Eventually the red
fluorescence on the membrane was uniformly distributed. At no time, the red fluorescence
was redistributed to the cytosol. This suggests that the amount of PIP5K on the
membrane is conserved, and depletion in one region must produce an increase in another
region. Since PIP5K controls signaling activities, its redistribution provides a powerful
mechanism for symmetry breaking.

To seek regions of PIP5K that control membrane binding and trailing edge
accumulation in *Dictyostelium* cells, we generated a series of truncation constructs to
examine their localization (Figure S4f). Despite its robust membrane association, PIP5K
does not contain a typical membrane association or transmembrane domain. $PIP5K_{301-718aa}$
was still localized as WT cells, although a small fraction of cells showed a relatively
smaller localization at the back-state. (Figure S4g, Video S9). $PIP5K_{316-718aa}$ appeared
significantly in the cytosol, and the protein remaining on the membrane was not localized
at the back. Upon recruitment to a uniformly localized protein cAR1-iLiD, this fragment
also did not relocalize cAR1-iLiD to the back (Figure S4h, Video S8). N-terminal 315aa
was not on the membrane and did not relocate cAR1-FKBP-FKBP to the back (Figure
S4i, Video S8). These results indicate that a region between 301-316 aa is important for
localization to the membrane, but the back localization region was not identified. Back
localization may require a coincidental association of two independent sequences within
the protein. Future experiments will address these possibilities.

Expressing PIP5Ks suppresses cell protrusions and alters migration

In previous experiments (Figure 2), where PIP5K was expressed from a
constitutive vector and the expression level was relatively low, we noticed a slight
impairment of migration. Reasoning that the highly expressing cells were selected against
and lost before observation, we developed a conditional doxycycline-inducible version of
PIP5K. Without doxycycline (DOX) induction, *Dictyostelium* cells show their normal
amoeboid migration behavior (Figure 3a, Video S10). Upon overnight DOX induction, two
striking phenotypes appeared. About 60% of the cells became much rounder, lacked
protrusions, and failed to migrate (Figure 3b, Video S10). About 40% of the cells displayed
many tiny, broken protrusions, resembling short filopodia, along the cell perimeter and
migration was also impaired (Figure S5a, Video S10). These observations were
supported by color-coded temporal overlay profiles (Figure 3i-l, S5l). Combining data from

the "rounded" and "spiky" cells, tracking experiments showed that the migration speed of
was reduced to 1.52 ± 0.7 µm min^{-1} (Figure 3g, S5b), in contrast to the 8.8 ± 2.3 µm min^{-1}
observed in cells without DOX incubation (Figure 3g, S5c). The cells were flatter and
occupied about 1.7-fold more surface area (Figure S5d). Z-stack imaging of cells showed
that upon PIP5K expression, the z-axis values drop from 12 µm to 4.87 µm (Figure S5g).
In addition, FITC-dextran uptake measurements showed that macropinocytosis was
significantly reduced, consistent with the nearly absent or tiny protrusions (Figure S5h-j)
[37, 65, 66].

We could titrate PIP5K activity in several ways. Protein expression levels varied
with doxycycline incubation times (Figure S5e). Surprisingly, cells with low PIP5K
expression after 2h DOX incubation became very polarized (Figure S5k), with a 1.32-fold
increase in aspect ratio (Figure S5f). Tracking experiments showed that these cells
migrated much faster than uninduced cells, with speeds reaching 19.2 ± 3 µm min^{-1}
(Figure 3g, S5o). These observations were supported by color-coded temporal overlay
profiles (Figure S5m). A similar increased speed was observed in cells expressing a
PIP5K mutant (K681N, K682N) predicted to have impaired kinase activity, with speeds
reaching 12.5 ± 3.3 µm min^{-1} (Figure 3g, S5n, S5r, Video S10) [67].

Next, we extended our studies to differentiated HL-60 neutrophils and
macrophages to examine whether overexpressing PIP5Ks inhibits cell migration in
leukocytes. We first overexpressed either PIP5K1B or PIP5K1C in differentiated HL-60
neutrophils, as these two isomers, according to previous literature, are important to
neutrophil chemotaxis and directed migration [49, 56]. Like *Dictyostelium* cells,
differentiated HL60 WT neutrophils showed a typical flat and polar phenotype (Figure 3c,
Video S11). Upon overexpressing either PIP5K1B or PIP5K1C, two phenotypes existed
in the basal state populations: round cells with PIP5Ks broadly distributed on the
membrane (Figure 3d, S6a, Video S11) or polarized cells with PIP5Ks enriched on the tail
(Figure S6b-c, Video S11). These observations were supported by color-coded temporal
overlay profiles (Figure S6d-e, S6i-j). Tracking experiments showed that the migration
speed of cells overexpressing PIP5Ks is impaired, with speeds reaching 2.92 ±
1.3 µm min^{-1} for PIP5K1B and 1.23 ± 0.6 µm min^{-1} for PIP5K1C (Figure 3h, S5q, S6h), in
contrast to the 9.0 ± 2.4 µm min^{-1} observed in WT cells (Figure 3h, S6g). The overlay
suggested that even though the polarized cells were active at the leading edge, they did
not move well because the tail appeared to adhere more strongly to the surface (Figure
S6d-e). We also measured the mean membrane intensity between the polarized and
round phenotype cells. Cells with rounder shapes have higher PIP5K expression levels,
consistent with the *Dictyostelium* results (Figure S6n-o), although the round-shaped cells
are not as flat as *Dictyostelium* cells. Cell areas were about 33% and 40% reduced for
PIP5K1B and PIP5K1C cells, respectively (Figure S6f). We also overexpressed PIP5K1B
and PIP5K1C in differentiated HL-60 macrophages. This led to a very flat and round-
shaped phenotype, abolishing protrusions compared with WT macrophages (Figure 3e-f,
Figure S6p, Video S12). These observations were supported by color-coded temporal
overlay profiles (Figure S6k-m).

Next, we extended our inquiry to epithelial-derived cancer cells by studying MDA-
MB-231 cells. Since these cells are very heterogeneous, we developed a system to
immediately observe the effects caused by activating PIP5K in an individual cell. We

subcloned a previously developed recruitable PIP5K1C [68] into an optogenetic system
by fusing it with crimson-SspB, which enabled light-induced association with membrane
anchor iLiD-CAAX [69] (Figure S6q). Globally recruiting cytosolic PIP5K1C to the plasma
membrane in MDA-MB-231 cells induced cell retraction with retraction fibers at the cell
rear, possibly due to the increased contractility (Figure 3k, Video S13). About 5 min after
recruitment, cells started to display retraction fibers. Contraction was formed after about
20 minutes (Figure 3k, S7a). Recruiting an empty vector did not cause a cell shape
change (Figure S7e, Video S13). Across the population, upon PIP5K1C recruitment,
MDA-MB-231 cells induced a 25% and 59% reduction in the cell area and aspect ratio,
respectively (Figure 3l-m), while recruiting an empty vector did not induce any cell area
or polarity change (Figure S7f-g). These observations were supported by color-coded
temporal overlay profiles (Figure S7b-d). In some recruited cells, blebbing was induced
in addition to retraction (Figure 3k), indicating these cells potentially had even higher
contractility upon PIP5K1C recruitment.

In summary, the phenotypes of overexpressing PIP5Ks are generally conserved
across *Dictyostelium,* leukocytes, and epithelial cells. Cells show different migratory
behaviors upon different PIP5K expression levels. We suggest that moderate expression
of PIP5Ks tends to polarize cells by quenching extraneous protrusions, while strongly
expressing PIP5Ks eliminates protrusive activities and severely impairs cell migration.

Expressing PIP5Ks suppresses signal transduction and cytoskeletal activities

To further understand the basis of the overexpression phenotype, we examined a
series of signal transduction and cytoskeletal activities using biosensors. We first
examined the spatial distribution of Ras activity using a Ras binding domain (RBD)
biosensor in *Dictyostelium* cells with different DOX incubation times. As expected, cells
showed characteristic patches at the cell protrusions without DOX incubation (Figure 4a,
Video S14), whereas cells incubated with DOX overnight had much fewer RBD patches.
Notably, there were no RBD patches in rounded, non-moving cells (Figure 4b, Video S14).
In the "spiky" cells, there were small, narrow RBD patches (Figure S8e, Video S14),
although these were increased in number (Figure 4e). These observations were
supported by membrane kymograph analyses (Figure S8a-b). There was an inverse
correlation between patch size and number (Figure 4g). Combining data from "rounded"
and "spiky" cells, there was an overall 44% reduction in total Ras patches (Figure 4b,
S8e, 4e). Similar results were observed using the PHcrac biosensor to examine PIP3
activities in cells with different DOX incubation times, with an overall 58% reduction in
cells with overnight DOX incubation (Figure 4c-d, 4f, S8f, Video S14). These observations
were supported by membrane kymograph analyses (Figure S8c-d). We also observed
the ventral wave differences for cells with or without DOX incubation. We found that
uninduced cells displayed broad, sustained waves, and cells with overnight DOX
incubation showed no waves or smaller, slower-moving waves, indicating low Ras
signaling activities (Figure 4h, Video S15).

We further examined cytoskeletal activities in the PIP5K overexpressing cells.
LimE displayed much smaller or no patches in cells following overnight DOX induction,
with a 60% reduction in these cells (Figure S8g-i, S9a, Video S16). The dynamic behavior
of these activities was captured in t-stack kymographs (Figure S9b-c), indicating that the

tiny LimE patches in cells expressing PIP5K are more transient than patches in untreated cells. Since branched actin is activated by a Rac1/Arp2/3 complex/branched actin pathway, we used a Pak1-GBD and ArpC biosensors to examine these activities. The Rac1 activity paralleled localization of LimE (Figure S9d-f, Video S16). Induced cells showed a 70% reduction in Rac1 patch size (Figure S9g). The ArpC patches were also much smaller in PI5K-expressing cells (Figure S9h-i, Video S16).

We have previously reported that cells lacking actin-sequestering actobindins A,B,C (*abnABC-*) have increased Ras and PI3K activity, suggesting that branched actin is in a positive feedback loop with the these STEN activities [70]. We examined the extent to which the inhibitory effect of PIP5K expression can interrupt this feedback loop by overexpressing PIP5K in *abnABC-* cells [70]. Without DOX incubation, cells showed broad RBD patches as expected (Figure S10j, Video S17). The RBD patches decreased in length and increased in numbers dramatically upon overnight DOX incubation (Figure S10k, Video S17), indicating competition between the positive feedback from branched actin and inhibition by PIP5K. However, unlike WT, in the *abnABC-* background there were no completely round cells containing no RBD patches, again suggesting a balance between activation by positive feedback from F-actin and inhibition by PIP5K.

In previous experiments (Figure 3), we showed that cells showed a much more polarized phenotype and migrated faster when incubated only 2h with DOX than without DOX incubation. We examined both Ras and PIP3 signaling activities in these cells and found the Ras and PIP3 signaling activities were confined to tiny patches at the leading edge or undetectable (Figure 4f, S10a-b, Video S14). Similar Ras signaling activities were low or undetectable in fast-moving cells expressing mutant PIP5K (K681N, K682N) (Figure S10c, Video S18). The LimE patches were smaller and confined at the front of these cells (Figure S10d, Video S18). Our lab previously reported that a RasGAP C2GAPB in *Dictyostelium* cells can also inhibit Ras activities but increase polarity and migration speed [43], which showed a similar phenotype compared to the cells expressing a low level of PI5K activity. To assess whether C2GAPB and PI5K are acting in the same pathway, we co-expressed PIP5K and C2GAPB and found that both cell shape and migration speed mimic expressing PIP5K alone (Figure S11 a-b, Video S19). To investigate whether attenuating Ras activity in a highly activated cell could restore polarity and migration, we overexpressed C2GAPB in *pi5k-* cells. Indeed, upon expressing C2GAPB, the *pi5k-* cells changed from fan-shaped or oscillatory cells to more polarized cells, with migration speed increase from $5.1 \pm 2.2\,\mu m\,min^{-1}$ (-C2GAPB) to $22 \pm 7\,\mu m\,min^{-1}$ (+ C2GAPB) (Figure S11 c-e, Video S19).

PI(4,5)P2 has been considered to promote cell migration by interacting with many actin-binding proteins [71-75]. To assess whether the inhibitory role of PIP5K in our studies is dependent of cytoskeletal dynamics, we treated cells with Latrunculin A to remove all F-actin activities and caffeine to induce the crescent patches of RBD or PHcrac. Without DOX incubation, cells displayed the characteristic crescent RBD or PHcrac patches (Figure 4i, 4K, Video S20). However, with overnight DOX incubation, 60% cells did not have any patches, while the rest, showed a 31% or 45% reduction in the RBD or PHcrac crescent length, respectively (Figure 4j-l, 4m-n). These observations were supported by membrane kymograph analyses (Figure S10e-h). These responses of the STEN patches to PIP5K-mediated inhibition parallel the PI5K-induced conversion of

389 amoeboid cells to "rounded" or "spiky" cells seen in the absence of latrunculin A. These
390 observations suggest that the PIP5K-induced cell shape, migration speed, and
391 cytoskeletal phenotypes reported above can be traced back to the inhibitory effects of
392 PIP5K expression on STEN activities (Figure 4o), although there may also be additional
393 effects on feedback from the cytoskeleton.

Expressing PIP5Ks increases the threshold for STEN activation

In previous experiments (Figure 4), we showed that expressing PIP5Ks inhibits
Ras and PI3K activities and that cells expressing PIP5Ks showed smaller, shorter-lived
waves. Reduced spontaneous signaling and curtailed wave activity in an excitable
system indicate an increased threshold [20, 31, 34, 40, 41, 46]. To assess the threshold
of the network more directly, we monitored the accumulation of PIP3 with biosensor PH-
Akt in response to chemoattractant C5a in RAW 264.7 cells with or without PIP5K
expression (Figure 5a-c, Video 21). In WT macrophages stimulated with 1 μM C5a, PIP3
transiently increased and cells extended protrusions. In contrast, at this dose, there was
no PIP3 increase in cells expressing PIP5K1C; cells stayed rounded and inactivated
(Figure 5a-b, Video 21). Compared with WT cells, the dose-response spanning 0.1 μM to
100 μM C5a for PIP5KIC-expressing cells was shifted to higher concentrations (Figure
5c, Video 21). The PH-Akt translocation responses at each C5a concentration are shown
in Figure S13. The PH-Akt translocation in cells expressing PIP5K1C was evident only at
100 μM C5a (Figure S12h, Video 21).

Similar results were observed in *Dictyostelium* cells stimulated with folic acid (FA).
At 1 nM FA, untreated cells showed about 25% RBD translocation response (Figure 5d).
Following induction of PIP5K by overnight incubation, there was no RBD translocation
response at 1 nM (Figure 5e). The line scan results support these observations (Figure
S13a-b). Compared to untreated cells, the dose-response curve from 0.1 nM to 100 nM
FA for PI5K-expressing cells was shifted to higher concentrations (Figure 5f). Figure
S12c-d shows typical images of the 5% and 20% PH-Akt or RBD responses. Dividing the
cells into < 5% or > 20% response groups showed clearly that there many fewer PIP5K-
expressing cells in the > 20% response set even with the highest concentrations of C5aR
and FA (Figure S13e-f).

Since both the C5a receptor (C5aR) in macrophages and the folic acid receptor
(fAR1) in *Dictyostelium* are G-protein-coupled receptors (GPCRs) [76, 77], we wondered
whether the inhibitory effects of PIP5Ks are still evident when GPCR is uncoupled from
the signal transduction network. In *Dictyostelium* cells lacking Gβ, conditional expression
of PIP5K, spontaneous LimE patches still decreased in size and increased in numbers as
previously observed in WT cells (Figure 5g-h, Video S22, and see Figure S8g-i).
Considering that LimE is a downstream indicator of network activity, these data indicate
the inhibitory role of PI5K can act downstream of GPCR signaling.

Since expressing PI5K caused severe impairment of random migration and shifted
the dose-response curve to higher concentrations, we expected that directional migration
would also be inhibited. Indeed, untreated *Dictyostelium* cells mounted a chemotactic
response towards folic acid, while the insensitive, rounded PIP5K-expressing cells largely
failed to migrate towards higher concentrations of (Figure S13g, S14a, Video 23). There

was a weak response in the "spiky" or less affected cells. Overall, cells expressing PIP5K
have less directional migration speed and polarity (Figure S14a-c).

In summary, expressing PIP5Ks inhibits signal transduction activities in response
to chemoattractant stimulation, but this inhibitory role does not require G-protein,
indicating that it acts at a point downstream. We suggest that the PIP5K inhibitory effect
raises the threshold of the network and reduces its response to internal spontaneous
activation as well as receptor inputs. We might reasonably expect responses to other
external cues, such as electric fields or shear forces, to also be attenuated.

Inhibiting myosin II activity counteracts PIP5K-induced phenotypes

In previous experiments (Figures 3 and 4), we showed that expressing PIP5Ks can
eliminate signal transduction and cytoskeletal activities and cell migration. We
hypothesized that the inhibitory effects of PIP5Ks on cell morphology and migration might
be related to increased Myosin II activities [78] and asked the extent to which lowering
Myosin II would counteract PIP5K expression and restore migration. To test this, we co-
expressed PIP5K and a recruitable Myosin heavy chain kinase C (MHCKC) previously
shown to abruptly decrease Myosin II bipolar thick filament assembly [70] (Figure 6a). In
PIP5K-expressing "rounded" cells, global recruitment of MHCKC to the plasma
membrane caused cells to polarize and migrate faster (Figure 6b, Video S24). These
observations were illustrated by color-coded temporal overlay profiles and line scans
(Figure 6c, S15a-b). Across the population, upon MHCKC recruitment, migration speed
increased from 1.6 ± 0.6 µm min^{-1} to 5.4 ± 0.9 µm min^{-1} (Figure 6h, S15c-d). The aspect
ratio also increased by 1.23-fold (Figure 6i). A heat map of individual cell migration speed
at each time illustrates the increase in migration speed after MHCKC recruitment, which
took about 10 mins to establish (Figure 6d).

Corresponding results were observed in propagating waves of Ras activity on the
ventral surface of fused giant cells. As shown in Figure 6e and Video S25, in this example,
expressing PIP5K completely suppressed waves of Ras activity. Strikingly, upon adding
Myosin II inhibitor, Blebbistatin, the RBD waves started to burst and gradually increased
until the entire surface displayed activity. Activity then decreased and reappeared. These
results suggest that the mutual antagonism between Ras and PIP5K activities can be
modulated by Myosin II. These results are consistent with an independent study from our
laboratory demonstrating negative feedback from Myosin II to Ras [70].

We then extended these studies to differentiated HL-60 neutrophils and RAW
264.7 macrophage-like cells and assessed the effects of Myosin II reduction caused by
recruitment of protein phosphatase 1 regulatory subunit 12A, Mypt1 in cells expressing
PIP5Ks. We took advantage of a recruitable Mypt169, which enabled light-induced
association with the membrane (Figure 6f) [79]. Globally recruiting Mypt169 to the plasma
membrane in WT cells as the control induced cell spreading and polarity in differentiated
HL-60 macrophages (Figure S15e-f, Video S26). Differentiated HL-60 neutrophils co-
expressing PIP5K1B and the recruitable Mypt169 in initially displayed the same rounded
phenotype as those expressing PIP5K1B alone (Figure 6g, Video S26). Globally
recruiting Mypt169 to the plasma membrane, combined with Rock inhibitor Y27632,
induced cell spreading and a polarized cell shape (Figure 6g, Video S26). These

observations were supported by color-coded temporal overlay profiles (Figure 6l). Cell
migration did not show a significant change (Figure 6l). Across the population, upon
Mypt169 recruitment, differentiated HL-60 neutrophils induced a 1.53-fold and 1.23-fold
increase in the cell area and polarity, respectively (Figure 6j-k). Similarly, globally
recruiting Mypt169 to the plasma membrane in RAW 264.7 cells co-expressing PIP5K1C
induced cell spreading and protrusion formation (Figure S15g, Video S26). These
observations were supported by color-coded temporal overlay profiles (Figure S15h).

Since there is a shift from front towards "back" activities in PIP5K overexpressing
cells, we reasoned that there might be a selective inhibition of branched actin relative to
linear, cortical actin. ABD120 is a protein that labels total F-actin in *Dictyostelium* cells
[80], while LimE selectively labels branched actin. We expressed both constructs in a
PIP5K-overexpressing cell line and used the ABD120-LimE patch ratio to indicate the
relative level of inhibition. Before inducing PIP5K, ABD120 is primarily labeled uniformly
on the plasma membrane, with some enrichment at the protrusions, whereas LimE is
localized almost exclusively to protrusions (Figure S16a, Video S27). Upon inducing
PIP5K, cells changed to a rounder shape, and LimE patches disappeared (Figure S16b,
Video S27). However, ABD120 was still uniformly distributed on the plasma membrane,
resulting in a higher ABD120-LimE ratio (Figure S16e), suggesting a selective inhibition
of branched actin. Consistent results were observed when staining cells with phalloidin.
In uninduced cells, phalloidin mainly stained at the protrusions, while in cells expressing
PIP5K, phalloidin staining colocalized with PIP5K around the perimeter (Figure S16c-d).
Note that earlier (Figure 4), we found that PIP5K-expressing cells that showed the "spiky"
phenotype changed to the "rounded" phenotype with CK666, further indicating that round-
shaped cells have less branched actin (Figure S16f, Video S28).

The stochastic, reaction-diffusion model recreates experimental observations

We next sought a simple mathematical model that could account for the extensive
observations of the effects loss- and gain-of-function of PIP5Ks on cell migration and
polarity and explain the observed symmetry breaking in the absence of cytoskeletal
activity (Methods). The previously reported core of the model included interactions
between Ras, PIP2, and PKBA, which gave rise to excitable behavior [36, 46]. Based on
our new findings, we adjusted the inhibitory interaction from Ras to PIP2 to act through
PIP5K. This, together with the negative regulation of Ras by PI(4,5)P2, closed a double-
negative feedback loop, functionally equivalent to a positive feedback loop as in the
original core model (Fig. 7A). However, based on the new findings, we made a critical
change by implementing this loop with a dynamic partitioning scheme, where PIP5K
remained on the membrane and was pushed away from regions of high Ras ([18];
Methods). We included two further feedback loops representing the interactions of the
cytoskeleton with Ras activity, which were supported by an independently published study
as well as the current findings. The first involved the positive contribution of actin to Ras
activity, mediated through PKBA [40]. The second was the contribution of myosin,
regulated by PI(4,5)P2, also in a double-negative feedback motif [70]. Simulations of this
system on a one-dimensional periodic domain, representative of the cell boundary, gave
rise to long-lived periods of localized, elevated Ras activity (Fig. 7B). Kymographs of the
other model components showed that most signaling events either colocalized with Ras
(e.g., Actin and PKBA) or formed a corresponding "shadow wave" where the signaling

molecule was absent (e.g., PI(4,5)P2, PIP5K, and myosin). Notably, because of the
dynamic partitioning, PIP5K was highest at the boundary between the wave and shadow
wave (red arrow in Fig. 7B). In this case, the enzyme could not stay in the region of high
Ras, but could not leave the membrane and it was elevated elsewhere. As shown the
total PIP5K across the perimeter remained constant over time. As PIP5K regulated
PI(4,5)P2 and subsequently myosin, we observed these bands of high activity in both
markers.

We then used the model to recreate several perturbation experiments. Following
the elimination of PIP5K, we observed Ras and actin activity increase greatly,
transitioning from localized streaks to high levels that extended almost throughout the cell
perimeter (Fig. 7C). In contrast, modeling PIP5K overexpression by increasing the PIP5K
levels eliminated the Ras streaks completely (Fig. 7D). Simulations of latrunculin
treatment showed that although the duration and persistence of the streak diminished,
the model could still generate persistent streaks (Fig. 7E). Overall, the simulations
strongly agreed with the experiments.

Discussion

Our studies reveal that spatiotemporal regulation of PIP5K on the membrane is a
primary determinant of polarity, protrusion formation, and resulting mode of migration
(Figure 8). The phenotypes we observed are consistent with the kymographs generated
in the computational simulations. In the absence of cytoskeletal activities (Figure 8a), cells
are largely in a "back" state where PIP5K is elevated and Ras activity is low. Local
spontaneous depletions of PIP5K and coordinated activations of Ras build up, breaking
symmetry (Figure 8b). Coupled myosin reduction and increased actin polymerization lead
to intermittently extended protrusions of relatively defined size and cell polarity (Figure 8c
and S17). When PIP5K is absent in *pi5k-* cells, Ras is highly activated and signal
transduction events spread across most of the cortex. Myosin is depleted, actin is
polymerized, protrusions are wide, and cells switch their migratory behaviors from
amoeboid to fan-shaped and oscillatory phenotypes (Figure 8d and S17). When PIP5K
is slightly overexpressed in WT, signal transduction activities and protrusions are curtailed
and cells, can polarize and migrate more rapidly (Figure 8e and S17). When PIP5K is
increased further, Ras, PI3K, and F-actin activities are confined to small patches, or
extinguished, and the entire cortex is shifted toward the back state. Protrusions become
smaller and spikier, or disappear entirely and cells stop migration. (Figure 8f and S17). In
summary, different phenotypes can be explained as lying along a spectrum from low-to-
high PIP5K expression levels, which are inversely related to high-to-low signal
transduction network activities (Figure S17).

As indicated in Figure 8, our results suggested that PIP5K is a *negative* regulator
of signal transduction and cytoskeletal activities, which was not previously recognized.
We propose that the chemotaxis defects previously reported in *pi5k-* cells can be traced
to removal of the normal PIP5K-mediated constraint on Ras, leading to elevated,
unregulated signal transduction and cytoskeletal activities which interfere with the cell's
ability to selectively direct protrusions towards chemoattractants. We found that although
the overall speed of the *pi5k-* cells on average does not decrease, the actual behaviors
of the cells as fans and oscillators are very defective, which was overlooked in the

previous report [51]. In neutrophils and macrophages, we found that lowering PIP2 by recruiting Inp54p induced more protrusions, fan-shaped movement, and increased actin polymerization, consistent with the phenotype of the *pi5k-* cells and previous results in *Dictyostelium* [46]. Furthermore, previous reports that a dominant negative PIP5K1C interferes with the uropod in neutrophils and disrupts chemotaxis may also have been related to the loss of PIP5K function, leading to excessive Ras and associated protrusive activity [56]. The effects of PIP5K overexpression or recruitment in *Dictyostelium*, neutrophils, macrophages, and cancer-derived MDA-MB-231 cells provided further evidence that the enzyme is a negative regulator: 1) Cell protrusions and migration speed are severely suppressed; 2) Ras, PI3K, and F-actin activities are strongly inhibited. 3) The dose-response curves for Ras and PI3K activation by chemoattractant are shifted to higher concentrations. The effects of overexpression of PIP5K could be due to elevating PIP2 levels or preventing it from going down, as it and other back localized components normally do as protrusions form.

PIP5K plays a crucial role in symmetry breaking and polarity establishment. As migrating *Dictyostelium* cells and human neutrophils migrate, they occasionally shift between more circular and polarized profiles whereupon PIP5K shifts between a uniform localization and a back-to-front gradient. Since the enzyme is tightly associated with the membrane (confirmed in centrifugation assay), it must diffuse along the membrane to the back [18]. Protrusions are formed at the nascent front as PIP5K moves away from a local region. As it does not shuttle to the cytosol, PIP5K leaving the front accumulates at the rear, suppressing Ras, PI3K, and branched actin polymerization activities at the back and promoting a shift toward actomyosin. Thus, the spatiotemporal relocalization of PIP5K provides a way to activate the cell front and suppress the rear simultaneously, an elegant mechanism for symmetry breaking and self-organizing polarity. The regulation of signal transduction by PIP5K can occur without cytoskeletal activities in the presence of latrunculin, but there are significant interactions with actomyosin. In untreated cells, feedback from actomyosin is an essential part of the suppressive mechanism at the back since inhibiting myosin II can overcome the PIP5K-mediated suppression and reactivate the signal transduction activities.

Various models have been put forth suggesting that cell polarity arises from the competition of local positive and global negative feedback loops. Whereas positive feedback helps to amplify initial cellular asymmetries, negative feedback prevents other sites from developing, ensuring that the cell has a singular region of high activity, thus enabling persistent directional motion. Most of these models suggest that the positive feedback loop incorporates actin-mediated activation of various signaling molecules [81, 82]. The origin of the negative feedback is less clear, though mechanisms by which membrane or cortical tension inhibit protrusions have been suggested [17, 29, 83]. Computational models of these complementary motifs generate simulations of cell motility that closely resemble the shape and tracks of real cells [84, 85]. The dynamic partitioning of PIP5K reported here suggests an alternative actin-independent mechanism by which the same molecule can provide both localized positive feedback and global negative feedback. At the regions of high Ras activity, the removal of PIP5K lowers the levels of the Ras-inhibiting PIP2. This double-negative feedback loop provides localized positive feedback. Additionally, the dispersal of this membrane-bound PIP5K to regions away

from high Ras activity, increases PIP2 levels and lowers the probability that Ras
excitability is triggered elsewhere – thus completing a global negative feedback.

There are numerous reported interactions of PIP2 with various actin-binding
proteins, but as some are reported to be activators and others are inhibitors, the overall
effects of altering PIP2 in cells are challenging to understand. Some of the studies
suggest that PIP2 sequesters critical actin-binding proteins. This mechanism would align
with our proposal that PIP5K is a negative regulator: *Van Rheenen et al.* reported that
EGF-induces PIP2 reduction in metastatic tumor cells and promotes actin polymerization
[86]; *Sengelaub et al.* reported that in metastatic breast cancer cells, lowering PI(4,5)P2
levels releases cofilin to enhance actin turnover and cell migration [87]. Alternative studies
suggested that PIP2 is an activator: It can directly interact with WASP to promote Arp2/3
complex nucleation [74] and it can bind to proteins like talin, vinculin, and focal adhesion
kinases (FAK), to promote focal adhesion formation [88-93]. Our study indicates that
inhibition to Ras by PIP5K, transmitted downstream, shifts the balance from branched F-
actin to actomyosin and that the inhibitory role of PIP5Ks occurs in cells that do not form
stable focal adhesions. Thus, the overall direction of the behavior in living cells favors
PIP5K as a negative regulator regardless of potential direct interactions of PI(4,5)P2 with
individual actin-binding proteins.

The counterintuitive effects of PIP5K inhibition on cell migration can be explained
by the Ras-PIP5K mutual inhibitory feedback loop [46], with additional feedback loops
from the cytoskeletal network, shown in Figure 7 and 8, but further studies are needed to
sort out the detailed mechanism. Even without the cytoskeletal network, there is a strong
reciprocal connection between Ras and PIP5K. It has been reported that Ras can directly
interact with PIP5K [94]; if that is the case, the interaction must push Ras towards the
inactive form to be consistent with our observations. It is possible that PIP5K or PIP2
interacts with Ras GAPs or competes with Ras GEFs to inhibit Ras activities. We are
currently testing these possibilities. In the presence of the cytoskeletal network, PIP2 may
regulate the interaction between the membrane and the actomyosin cortex [38]. While
raising PIP2 would increase the interaction, lowering PIP2 may loosen the linkage, and
could initiate a shift from actomyosin to branched actin. Such a mechanism would be
consistent with the model of front and back-promoting feedback loops we recently
reported [46, 70].

Our results suggest that phenotypes resulting from loss-of-function or gain-of-
function alterations of PIP5K in tissue will likely depend on the initial state of the affected
cells and may be difficult to predict. Without this study, PIP5K might be considered an
oncogene since it is the substrate for producing second messengers, such as PIP3, DAG,
and IP3, which are classic tumor promoters. The depletion of PIP5K and its product PIP2
from the front of the cells is puzzling since these messengers are localized at the front.
However, even when PIP2 is lowered by as much as 90%, PIP3 is more elevated, and
when PIP5K is overexpressed, Ras and PI3K activity are severely inhibited. Based on
this, one might expect PIP5K to act like a tumor suppressor and consider activating or
increasing PIP5K as a therapeutic mechanism to suppress activated oncogenic Ras. Our
results show that a slight increase in PIP5K and a decrease in PIP3 activity can enhance
polarity, while a large rise in PIP5K and elimination in PIP3 completely shut down the

movement. Therefore, if PIP5K were to be considered a deterrent to Ras-mediated
growth, caution would be needed to monitor the effects on cell migration.

Acknowledgments

We thank all members of the Devreotes, Iglesias, and Robinson laboratories
(Schools of Medicine and Engineering, JHU) for helpful discussions and for providing
resources. We thank Jonathan Kuhn and Yiyan Lin (Devreotes Lab) for providing
constructs. We acknowledge Orion Weiner (UCSF) for providing HL-60 cell line. We thank
Sean Collins (UC Davis) for providing transposon plasmids. We thank Stephen Gould
(School of Medicine, JHU) for help with instrumentation. We appreciate services of
DictyBase and Addgene for providing plasmids. This work was supported by NIH grant
R35 GM118177 (to P.N.D.), DARPA HR0011-16-C-0139 (to P.A.I. and P.N.D.), AFOSR
MURI FA95501610052 (to P.N.D.), as well as NIH grant S10OD016374 (to S. Kuo of the
JHU Microscope Facility).

Author Contributions

Y.D. and P.N.D. conceived and designed the project with inputs from T.B., D.S.P.,
and P.A.I.; Y.D. engineered constructs/stable cell lines, designed and executed
experiments, and performed the majority of data analyses with input from other authors;
P.B. and P.A.I. devised and conducted computational simulations, and deposited codes.
T.B. executed kymograph analyses; D.S.P. performed neutrophils and macrophages
Inp54p recruitment experiments and provided some stable cell lines; H.Z. generated
heatmap analysis; J.B. made *pi5k-* cell line and some constructs; P.N.D. and Y.D.
prepared initial drafts, and wrote and revised the final version of manuscript with help
from T.B., H.Z., and P.A.I.; PND supervised the study.

Competing Interests
677 The authors declare no competing interests.

Statistics and reproducibility

Statistical analyses were executed using unpaired or paired two-tailed
nonparametric tests on GraphPad Prism 10. Results are expressed as mean ± s.d. from
at least three independent experiments. NS, $P > 0.05$, *$P \leq 0.05$, **$P \leq 0.01$, ***$P \leq 0.001$,
***$P \leq 0.0001$. Tukey's convention was used to plot box-and-whisker plots. Statistical test
details are indicated in the figure legends. No statistical methods were used to
predetermine sample sizes, but our sample sizes are similar to those reported in previous
publications[4,35,79]. Each representative image or image series is from n > 3 independent
experiments. Any variations observed between treatment groups are not attributed to
sampling bias. No data were excluded from the analyses.

Data availability

689 All data are provided in the main or supplementary text. Requests for additional
690 information on this work are to be made to the corresponding author.

Methods

Preparation of reagents and inhibitors

Rapamycin (Millipore Sigma, 553210) was dissolved in DMSO to prepare the 5 mM stock solution. Then, 1 µl aliquots were diluted 1:100 in the development buffer (DB) to a 10x concentration (50 µM). CK666 (Millipore Sigma, 182515) was dissolved in DMSO to prepare the 100 mM stock solution. Then, 0.5 µl aliquots were diluted 1:100 in the warm development buffer (DB) to a 10x concentration (1 mM). 237 µM latrunculin A in ethanol (Cayman, 10010630) was diluted in the development buffer (DB) to prepare the 50 µM (10x) stock solution. Folic acid (Sigma-Aldrich, 329823065) was dissolved in sterile water, with the addition of 2 M NaOH, to prepare the 1.25 mM stock solution. Caffeine (Millipore-Sigma,1085003) was dissolved to 1 M in sterile water and then diluted to a 10x concentration (40 mM) in the development buffer (DB). Fibronectin (Sigma-Aldrich, F4759-2MG) was dissolved in 2ml sterile water, followed by dilution in 8ml PBS to prepare the 200 µg ml^{-1} stock solution. N-Formyl-Met-Leu-Phe (fMLP; Sigma-Aldrich, 47729) was dissolved in dimethylsulfoxide (DMSO) (Sigma-Aldrich, D2650) to prepare the 50 mM stock solution. Then, 1 µl aliquots were diluted in RPMI to a 10x concentration (2 µM). Phorbol 12-myristate 13-acetate (PMA; Sigma-Aldrich, P8139) was dissolved in DMSO to prepare the 1 mM stock solution. Then, 1 µl aliquots were diluted in RPMI to a 10x concentration (320 nM). FKP-(D-Cha)-Cha-r (ChaCha peptide, Anaspec; 65121) was dissolved in PBS to prepare the 2.5 mM stock solution. Then, 1 µl aliquots were diluted in DMEM to a 10x concentration (320 nM). A stock solution of EGF (Sigma-Aldrich, E9644) was prepared by dissolving it in 10 mM acetic acid to a final concentration of 1 mg ml^{-1}. . The anti-BSA mouse monoclonal antibody was acquired from Sigma-Aldrich (SAB4200688, clone BSA-33). Blebbistatin (Peprotech, 8567182) was dissolved in DMSO to prepare the 50mM stock solution. Then, 1 µl aliquots were diluted 1:100 in the development buffer (DB) to a 10x concentration (500 µM). 5 mM Y-27632 (Sigma-Aldrich, 688001) was dissolved in DMSO to prepare the 5 mM stock solution. Then, 1 µl aliquots were diluted in RPMI to a 10x concentration (100 µM). Hygromycin B (Thermo Fisher Scientific, 10687010) or G418 sulfate (Thermo Fisher Scientific, 10131035) was purchased as 50 mg ml^{-1} stock solution. Then, 10 µl Hygromycin B or G418 sulfate was added in a 10ml cell culture. Blasticidin S (Sigma-Aldrich, 15205) or puromycin (Sigma-Aldrich, P8833) was dissolved in sterile water to prepare the stock solutions of 10 mg ml^{-1} or 2.5 mg ml^{-1}, respectively. Then, 10 µl Blasticidin S or 4 µl puromycin was added in a 10ml cell culture. Doxycycline hyclate (Sigma, D9891-1G) was dissolved in sterile water to prepare the stock solution of 5 mg ml^{-1}. Then, 100 µl Doxycycline hyclate was added in a 10ml cell culture before the experiment. TRITC-dextran (Sigma-Aldrich, T1162) was dissolved in sterile water to prepare the 50 mg ml^{-1} stock solution. Then, the stock solution was further diluted to 2 mg ml^{-1} in the HL5 culture medium before the experiment. All stock solutions were aliquoted and stored at −20 °C.

Cells and plasmids

WT *Dictyostelium discoideum* cells of the AX2 strain (dictyBase, strain ID DBS0235521) were obtained from the Kay laboratory (MRC Laboratory of Molecular Biology) and cultured in HL5 medium (laboratory stock) at 22 °C [95] for a maximum of 2

months after thawing from frozen stock. *Pi5k-* (piki-) cells were purchased from Dictybase (strain ID DBS0350270). pikI− cells were grown on a Klebsiella aerogenes lawn on an SM plate and transferred to HL5 medium supplemented with heat-killed Klebsiella aerogenes before the experiment. Gβ- cells were generated in the Devreotes Lab previously [96]. abnABC- cells were generated previously in the Devreotes Lab [97]. Growth-stage cells were used for all the experiments. GFP-arpC cells were purchased from Dictybase (strain ID DBS2036065).

The human HL-60 cell line (ATCC CCL-240; RRID:CVCL_0002) was obtained from the Weiner laboratory (UCSF) and cultured in RPMI 1640 medium (Gibco, 22400-089) supplemented with 15% heat-inactivated fetal bovine serum (Thermo Fisher, 16140071)4,75. To obtain migration-competent neutrophils, WT or stable lines were differentiated in the presence of 1.3% DMSO with a total of 1.5 million cells over 5-7 days [98]. Differentiated cells are an effective model to study human neutrophils [99]. To differentiate HL-60 cells into macrophages, cells were incubated with 32 nM PMA for 48-72h [100]. Cells were grown in humidified conditions at 5% CO2 and 37 °C.

RAW 264.7 macrophage-like cells were obtained from the N. Gautam laboratory (Washington University School of Medicine in St. Louis). RAW 264.7 cells were cultured in Dulbecco's modified Eagle's medium (DMEM) containing 4,500 mg l^{-1} glucose, l-glutamine, sodium pyruvate and sodium bicarbonate (Sigma-Aldrich, D6429), supplemented with 10% heat-inactivated fetal bovine serum (ThermoFisher Scientific, 16140071) and 1% penicillin–streptomycin (ThermoFisher Scientific, 15140122). Cells were grown in humidified conditions at 5% CO2 and 37 °C.

All DNA oligonucleotides were purchased from Sigma-Aldrich. *Dictyostelium* PIP5K (below named PI5K) was PCR amplified from genome DNA and cloned in pCV5-GFP expression plasmid to generate PI5K-GFP (pCV5). Two-point mutations K681N and K582N were introduced to PI5K-GFP (pCV5) to generate PI5K (K581N, K682N)-GFP (pCV5). PI5K-GFP was then PCR amplified and cloned into doxycycline-inducible pDM335 plasmid (dictyBase, ID no. 523) using BglII/SpeI restriction digestion to generate mRFPmars-PI5K (pDM335). mRFPmars-PI5K was than PCR amplified and cloned into doxycycline-inducible pDM359 plasmid (dictyBase, ID no. 518) using BglII/SpeI restriction digestion to generate mRFPmars-PI5K (pDM359). mRFPmars was then replaced with EGFP to generate EGFP-PI5K (pDM359). EGFP was then replaced with KikGR to generate KikGR-PI5K (pDM359). DNA sequences encoding the 1-315aa, or 316-718aa, or 369-718aa or 1-718aa amino acids of PI5K, were PCR-amplified from PI5K-GFP (pCV5) and cloned into mCherry-FRB-MCS (pCV5) expression plasmid previously created in the Devreotes Lab using NheI/XhoI restriction digestion to generate mCherry–FRB-PI5K (1-315aa) (pCV5), mCherry–FRB- PI5K (316-718aa) (pCV5), mCherry–FRB-PI5K (369-718aa) (pCV5) and mCherry–FRB-PI5K (1-718aa) (pCV5). DNA sequences encoding the 316-718aa amino acids of PI5K, were PCR amplified and cloned into mRFPmars-SspBR73Q-MCS (pCV5) using NheI/NotI restriction digestion to generate mRFPmara-SspBR73Q-PI5K (316-718aa). DNA sequences encoding the 301-718aa amino acids of PI5K, were PCR amplified and cloned into pCV5-GFP expression plasmid to generate PI5K (301-718aa)-GFP (pCV5). PH-PLCδ-YFP (pCV5), CynA-KikGR (KF2), RBD-EGFP (pDM358), PHcrac-RFP (pDRH), LimEΔcoil-mCherry (pDM181), PHcrac-YFP (pDM358), LimEΔcoil-RFP (pDRH), cAR1-FKBP-FKBP (pDM358 or pCV5), LimE-

GFP-FKBP-FKBP (pDM358), mCherry-FRB-MHCKC (pCV5), and PKBR1$_{N150}$-iLiD
(pDM358) were previously created in the Devreotes Lab. Myosin II-GFP (pDM181) was
obtained from the Robinson laboratory (School of Medicine, JHU). PAK1(GBD)-YFP
(pDEXH) was a gift from the C. Huang lab (JHU). RBD-RFP (pDM358) GFP-ABD120
(pDXA) was purchased from Dictybase (ID no. 472).

For transit transfection in RAW 264.7 macrophage-like cells and MDA-MB-231
cells, PHAkt-mCherry and CIBN-CAAX (Addgene #79574) were from Devreotes Lab.
EGFP-PIP5K1B (Addgene #202722), GFP-PIP5K1gama90 (Addgene #22299) , CRY2-
mCherry-MYPT169 (Addgene #178526), TagBFP-FKBP-PIP5K1C (Addgene #220090)
(recruitable PIP5K1C), and mCherry-NES-SSPB-MCS-iLiD-CAAX (Addgene #173869)
were purchased from Addgene. mCherry was replaced with Crismon fluorescent protein,
and a point mutation R73Q was introduced to SapB to generate Crismon-NES-
SapBR73Q-MCS-iLiD-CAAX. Recruitable PIP5K1C was PCR amplified and cloned into
the XhoI/EcoRI sites of the Crismon-NES-SSPBR73Q-MCS-iLiD-CAAX expression
plasmid to generate Crismon-NES-SspBR73Q-PIP5K1C-iLiD-CAAX.

For stable lines in HL-60 cells, stable lines co-expressing LifeAct–miRFP703
(pLJM1) (Addgene #201750) and CIBN-CAAX (pLJM1) (Addgene #201749) or
expressing RFP-PHAkt (pFUW2) were previously created in the Devreotes Lab [44].
EGFP-PIP5K1B and EGFP-PIP5K1C were PCR amplified and cloned into the BspEI/SalI
sites of the PiggyBac transposon plasmid to generate EGFP-PIP5K1B (pPB) and EGFP-
PIP5K1C (pPB). Two-point mutations K359N, K260K or K407N, K408N were introduced
to EGFP-PIP5K1B (pPB) or EGFP-PIP5K1C (pPB), to generate EGFP-PIP5K1B (K359N,
K360N) and EGFP-PIP5K1C (K407N, K408N), respectively. Recruitable PIP5K1C were
PCR amplified and cloned into the BspEI/NotI sites of the PiggyBac transposon plasmid
to generate CRY2PHR-mCherry-PIP5K1C (pPB). MYPT169 was PCR amplified and
cloned into the BsiWI/NotI sites of the mCherry-CRY2PHR (pHR) expression plasmid ,
which was previously modified from construct mCherry-CRY2PHR (pHR) (Addgene
#201750) by introducing two restriction enzyme sites BstZ17I and BsiWI, to generate
mCherryCRY2PHR (pHR). Constructs were verified by diagnostic restriction digestion
and sequenced at the JHMI Synthesis and Sequencing Facility.

Transfection

Dictyostelium AX2 cells were transfected using a standard electroporation
protocol. Briefly, for each transfection, 5×10^6 cells were pelleted, washed twice with ice-
cold H-50 buffer (20 mM HEPES, 50 mM KCl, 10 mM NaCl, 1 mM MgSO4, 5 mM
NaHCO3, 1 mM NaH2PO4, pH adjusted to 7.0) and subsequently resuspended in 100 µl
ice-cold H-50 buffer. 2 µg of total DNA was mixed with the cell suspension, which was
then transferred to an ice-cold 0.1-cm-gap cuvette (Bio-Rad, 1652089) for two rounds of
electroporation at 850 V and 25 µF with an interval of 5 s (Bio-Rad Gene Pulser Xcell
Electroporation Systems). After a 10 min incubation on ice, the electroporated cells were
transferred to a 10-cm Petri dish containing HL-5 medium. The cells were selected by the
addition of hygromycin B (50 µg ml−1) and/or G418 (20–30 µg ml−1) after 24h as per the
antibiotic resistance of the vectors.

RAW 264.7 cells and MDA-MB-231 cells were transitly transfected by
nucleofection in an Amaxa Nucleofector II device using Amaxa Cell line kit V (Lonza,
VACA-1003) following a pre-existing protocol [101]. For each transfection, 3×10^6 cells
were pelleted, resuspended in 100 µl supplemented Nucleofector Solution V. A total of 2-
3 µg of DNA mixture was added and immediately transferred to a Lonza cuvette for
electroporation using the program setting D-032 and X-013 for RAW 264.7 cells and
MDA-MB-231 cells, respectively. Pre-warmed pH-adjusted culture medium (500 µl) was
added to the electroporated cells in the cuvette. The cell suspension was then transferred
to a 1.5 ml vial and incubated at 37 °C and 5% CO2 for 10 min and 30min for RAW 264.7
cella and MDA-MB-231 cells, respectively. Next, 50–100 µl solution containing cells was
transferred to a coverslip chamber and allowed to adhere for 1 h. Finally, approximately
400 µl of pre-warmed pH-adjusted culture medium was added to each chamber and the
cells were further incubated for 4–6 h before imaging for RAW 264.7 cells. MDA-MB-231
cells were imaged the next day.

Stable expression lines in HL-60 cells were generated by a combination of
3rdgeneration lentiviral- and PiggyBac[TM] transposon-integration based approaches [44,
102]. For transposon integration in HL-60 cell line, 5 µg transposon plasmid was co-
electroporated with an equal amount of transposase expression plasmid into two million
cells using Neon[TM] transfection kit (Invitrogen; MPK10025B). Cells and DNA mix were
resuspended in buffer 'R' before electroporation in 100 µl pipettes at 1350V for 35 ms in
Neon[TM] electroporation system (Invitrogen; MPK5000). Cells were resuspended in mixed
culture medium in a 6-well plate and allowed to recover for 24 hours. Post-recovery,
transfected cells were selected in presence of 10 mg/mL blasticidine S for 5-6 days. Once
blasticidine S was removed, resistant cells were transferred to 48-well cell culture plate
(Sarstedt; 83.3923), and further grown over 3-4 weeks into stable cell lines. Stable cells
were maintained throughout in Blasticidine S.

For virus transfection, 2[nd] generation virus was prepared in HEK293T cells grown
to about 80% confluency in 10 cm cell culture dish (Greiner Bio-One; 664160). For each
reaction, a mixture of 2 µg pMD2.G (Addgene #12259), 8 µg psPAX2 (Addgene #12260),
and 10 µg mCherry-CRY2PHR-MYPT19 (pHR) construct were transfected using
Lipofectamine 3000 as per manufacturer's instructions (Invitrogen; L3000-008). After 96
hours, virus containing culture medium was harvested at 3000 rpm for 20 mins at 4°C. In
a 6-well plate (Greiner Bio-One; 657160), entire viral medium was added to 4×10^6 HL-
60 cells (seeded at a density of 0.25×10^6 cells/mL) in presence of 10 µg /mL polybrene
(Sigma; TR1003). After 24 hours, viral medium was removed, and cells were introduced
to a mix of fresh and conditioned (mixed) culture medium. For selecting expressors,
infected cells were sorted after 5 days, and subsequently, grown to confluency. For high-
speeding sorting, we used 561 nm excitation laser to sort for mCherry expression using
SH800S cell sorter (Sony). Briefly, cells were pelleted, resuspended in sorting buffer (1x
PBS, Ca2+/Mg2+ free; 0.9% heat-inactivated FBS; 2% penicillin-streptomycin) at a
density of 15×10^6 cells/ml, and sorted using 100 mm microfluidic sorting chip. High
expressors (top 1%-5%) were collected in fresh culture medium (containing 2% penicillin-
streptomycin) and grown to confluency.

Confocal microscopy and Live-cell imaging

All *Dictyostelium* experiments were performed on a 22 °C stage. All RAW 264.7,
MDA-MB-231 and neutrophil experiments were performed inside a 37 °C chamber with
5% CO2 supply. All time-lapse live-cell imaging experiments were performed using one
of the following microscopes: (1) Zeiss LSM 780-FCS Single-point, laser scanning
confocal microscope (Zeiss Axio Observer with 780-Quasar; 34-channel spectral, high-
sensitivity gallium arsenide phosphide detectors), (2) Zeiss LSM800 GaAsP Single-point,
laser scanning confocal microscope with wide-field camera, (3) Nikon Eclipse Ti-E
dSTROM total internal reflection fluorescence (TIRF) microscope (Photometrics Evolve
EMCCD camera). The Zeiss 780 confocal microscope was controlled using ZEN Black
software; Zeiss 800 confocal microscope was controlled using ZEN Blue software; the
Nikon TIRF was controlled using Nikon NIS-Element software. The 40X/1.30 or 63X/1.4
Plan-Neofluar oil objective (with proper digital zoom) was used in all microscopes. The
488 nm (Ar laser) excitation was used for GFP and YFP, 561 nm (solid-state) excitation
was used for RFP and mCherry, and 633nm (diode laser) excitation was used for
miRFP703 in Zeiss 780 and 800 confocal microscopes. The 488nm (Ar laser) excitation
was used for GFP and 561 nm (0.5W fiber laser) excitation was used for mCherry and
RFP in Nikon TIRF.

To prepare imaging, vegetative *Dictyostelium* or differentiated macrophages were
adhered on an eight-well coverslip chamber for 40 min. Differentiated neutrophils, pre-
treated with heat-killed Klebsiella aerogenes, were adhered to fibronectin-coated
chambers for 40 min. Next, fresh DB (5 mM Na2HPO4 and 5 mM KH2PO4 supplemented
with 2 mM MgSO4 and 0.2 mM CaCl2) or RPMI 1640 medium was added to attached
cells and used for imaging. To induce PI5K expression in *Dictyostelium*, doxycycline
(50 µg ml^{-1}) was added 8 h before imaging. For macropinocytosis assay, cells were
incubated with 2 mg/ml TRITC-dextran for 10 min, washed twice with DB and imaged
subsequently. All live- or fix- cell imaging was acquired with 0.3–0.8% (*Dictyostelium*) or
1–8% (HL-60) laser intensity. In single cells, confocal imaging was performed at a middle
plane of the cells. To visualize cortical/ventral waves in *Dictyostelium* and RAW 264.7
cells, confocal microscopes were focused on the very bottom of the cell to capture the
substrate-attached ventral surface of the cell. For the inhibitor experiments, neutrophils
were treated with 10 µM Y-27632 for at least 10 min before imaging. In *Dictyostelium,*
50 µM blebbistatin or 100 µM CK666 was added during imaging, while 5 µM latrunculin A
and 4 mM caffeine were added at least 20 min before imaging.

Chemically induced dimerization

The plasmids and experimental details of the chemically induced dimerization
system have mostly been previously described [40, 46]. Here, to generate Figure S4i,
S5a-c, and 6a-d data, the two following combinations were used to express the systems:
(1) cAR1-FKBP-FKBP (pDM358) as the membrane anchor, mCherry–FRB-PI5K (1-
315aa) (pCV5) as the recruitee; (2) LimE-GFP-FKBP-FKBP (pDM358) as the membrane
anchor, mCherry–FRB- PI5K (1-718aa) (pCV5), or mCherry–FRB- PI5K (316-718aa)
(pCV5), or mCherry–FRB- PI5K (369-718aa) (pCV5), as the recruitee; (3) cAR1-FKBP-
FKBP (pCV5) as the membrane anchor, mCherry–FRB-MHCKC (pCV5) as the recruitee.
Dictyostelium cells in the growth phase were transferred to an eight-well coverslip
chamber and incubated for 10–15 min to allow them to adhere well. The HL-5 medium
was then replaced with 450 µl DB buffer. Image acquisition was started 15–20 min after

the medium change. After imaging a certain number of frames, rapamycin was added gently to the chamber (to a final concentration of 5 µM) during image acquisition to facilitate the recruitment of Inp54p to the plasma membrane.

Optogenetics

Optogenetics experiments were performed using slightly modified protocols [103] for *Dictyostelium*, HL-60 cells or MDA-MB-231 cells. Throughout image acquisition, a solid-state laser (561 nm excitation and 579–632 nm emission) was used for visualizing proteins or recruitable effectors fused to an mCherry, mRFPmars, or crismon tag, whereas a diode laser (633 nm excitation and 659–709 nm emission) was used to capture miRFP703 expression. Images were acquired for 5–10 min, before which a 450/488 nm excitation laser was switched on globally to activate recruitment and 10-20 min after. Image acquisition and photoactivation were performed at 7s intervals. Using the T-PMT associated with the red channel, we acquired DIC images. For recruiting MYPT169 or PIP5K1C in RAW264.7, MDA-MB-231, and HL-60 cells, 10 µM C5aR agonist, 10 ng/ml EGF, and 200 nM fMLP were added at least 15mins before imaging, respectively.

Frustrated phagocytosis and osmotic shock

We slightly modified a pre-existing protocol to visualize ventral waves in PMA-differentiated macrophages [54]. Briefly, Nunc Lab-Tek eight-well coverslip chambers were pre-washed with 30% nitric acid, coated with 1 mg ml^{-1} BSA for 3 h, washed with PBS and finally incubated with 5 µg ml^{-1} anti-BSA (1:200 dilution) for 2 h. The chambers were finally washed twice with PBS to remove excess antibodies. Before imaging, transfected PMA-differentiated macrophage cells were starved in suspension in 1×Ringer's buffer (150 mM NaCl, 5 mM KCl, 1 mM CaCl2, 1 mM MgCl2, 20 mM HEPES and 2 g l^{-1} glucose, pH 7.4) for 30 min. Next, these cells were introduced to the opsonized chambers and allowed to spread on the antibody-coated surface for 5–10 min, and then hypotonic shock was applied using 0.5×Ringer's solution before imaging.

For development on non-nutrient agar, cells were washed with developmental buffer (DB) and plated on solidified DB agar at a density of 5.2×10^5 cells/cm^2. For most experiments, development in suspension was carried out by pulsing cells resuspended in DB at 2×10^7 cells/ml with 50 to 100 nM cAMP every 6 min starting from 1 hour. For the experiments presented in Fig. 6 (A to D) and fig. S10 (A to C), *crac*$^-$ or GFP-GtaC/*crac*$^-$ cells were pulsed with different concentrations of cAMP or at different intervals starting from 2 hours. For development in the presence of sp-cAMPS, 1 µM was included in DB or DB agar.

Cell differentiation

For *Dictyostelium* cell development, 1×10^8 cells in the exponential growth phase were collected from suspension culture and pelleted as previously described [104, 105]. After washing with DB twice, the cells were resuspended in 4 ml DB and shaken at 105 r.p.m. for 1 h. The cells were then pulsed with 100 nM cAMP (5s pulse every 6 min) using a time-controlled peristaltic pump for 5-6 h with continual shaking. This allowed the cells to develop and polarize. After development, Around 10–15 µL of differentiated cells (in DB media) was collected were transferred from the shaker to an eight-well coverslip

chamber, resuspended thoroughly in 450 µl DB and incubated for 20–30 min before
imaging. For development on non-nutrient agar, cells were washed with developmental
buffer (DB) and plated on solidified DB agar at a density of 5.2×10^5 cells/cm^2.

Global receptor activation assay

Cells were first differentiated as described in cell differentiation protocol. Cells were
subsequently incubated with 5 µM Latrunculin A for around 20 min before imaging. Using
a confocal laser scanning microscope, 10-20 frames were first acquired to record the
basal activity of the proteins, then cAMP was added to the chamber (to a final
concentration of 1 µM) to activate all the cAR1 receptors, and the image acquisition was
continued. An imaging frequency of 3 s/frame was maintained throughout the experiment.

For vegetative *Dictyostelium* cells, cells were seeded on the eight-well coverslip
chamber and incubated for 20–30 min. Then, the HL5 medium was replaced with DB and
incubated cells with an additional 20 minutes before the imaging. Cells were subsequently
incubated with 5 µM Latrunculin A for around 20 min before imaging. Using a confocal
laser scanning microscope, 10-20 frames were first acquired to record the basal activity
of the proteins, then folic acid was added to the chamber (to a final concentration of 0.1,
1, 10, or 100 nM) to activate all the fAR1 receptors, and the image acquisition was
continued. An imaging frequency of 3 s/frame was maintained throughout the experiment.

For RAW 264.7 cells, transfected cells were previously incubated on the eight-well
coverslip chamber for 4-6h. The culture medium was replaced with HBSS
buffer supplemented with 1 g/L glucose and incubate with another 30 min before the
imaging. Using a confocal laser scanning microscope, 10-20 frames were first acquired
to record the basal activity of the proteins, then C5aR agonist was added to the chamber
(to a final concentration of 0.1, 1, 10, or 100 µM) to activate all the C5a receptors, and
the image acquisition was continued. An imaging frequency of 15 s/frame was maintained
throughout the experiment.

2D Chemotaxis Assay

We slightly modified a pre-existing protocol to perform chemotaxis assay for
vegetative *Dictyostelium* cells [106]. Briefly, cells were seeded on the 1-well Nunc Lab-
Tek chamber as a drop at the center and incubated for 20–30 min. Then 2 mL DB buffer
was added to the chamber and cells were dispersed by pipetting multiple times. A
Femtotip microinjection needle (Eppendorf) was loaded with 10 µM of filtered folic acid
solution and then connected to a FemtoJet microinjector (Eppendorf). The microinjector
was operated in continuous injection mode with a compensation pressure of 15 hPa. To
initiate the chemotaxis, the micropipette was brought to the (x,y,z) coordinate of cells
using a programmed micromanipulator. The imaging was continued with an acquisition
frequency of 30 s/frame.

CRISPR/Cas9 to generate *pi5k-* cell line

We slightly modified a pre-existing protocol for using CRISPR/Cas9 system to
target specific genes and generate KO mutants in Dictyostelium cells [107]. gRNA guide
sequences targeting pi5k gene were designed using Cas-Designer. Customized gRNA

oligos were purchased from Sigma-Aldrich. This selected gRNA sequence AAACAACTCATTGGATTCAGATGC (forward) was then cloned into pTM1285 plasmid obtained from the Robinson laboratory (School of Medicine, JHU). AX2 cells were transformed with pTM1285 plasmid containing pi5k gRNA using electroporation, using the transfection protocol described above. After 8–16 h of transformation, we replaced the media with HL-5 media containing G418 (20–30 μg ml−1) and cultured the transformed cells for 3 more days. After that, floating dead cells were washed off with HL-5 medium. Remaining live cells were plated on SM plates with cultured Klebsiella aerogenes and incubated at room temperature for 3–4 d until visible, isolated plaques formed. These individual plaques were separately transferred to 96-well plates containing HL-5 medium in the absence of G418. After approximately 2–3 d when colonies grew, genomic DNA was extracted for PCR verification and then the rest cells were transferred to 6-well and then 10-cm Petri dishes and grown to confluency.

Multinucleation assay

Cells that have been in suspension at 200 rpm for 2–3 d were seeded at ~70% confluency on glass coverslips in HL-5 medium and allowed to adhere for 20 min. Next, HL-5 medium was replaced with ice-cold fixative solution (2% PFA + 0.01% Triton X-100 in 1.5× HL-5 medium) for 10 min. Cells were then washed quickly with 1× PBS, and the nuclei were stained with 10 μg/ml Hoechst for 10 min. Afterwards, cells were washed three times, 10 min each time, with 1× PBT (1× PBS + 0.01% Triton X-100). Before visualization, cells were mounted on coverslips using Invitrogen SlowFade™ Gold antifade reagent. Cells were imaged using Zeiss LSM 780-FCS Single-point, laser scanning confocal microscope with 405 nm laser.

Immunofluorescence

Cells were seeded on glass coverslips in HL-5 medium and allowed to adhere for 20 min. Next, HL-5 medium was replaced with fixative solution (2% buffered paraformaldehyde with 0.25% Glutaraldehyde and 0.1% Triton X-100 in HL-5 medium) for 10min at room temperature (RT), then quenched in 1mg/mL Sodium Borohydride for 10min, then washed three times with 1x PBT (1x PBD supplemented with 0.5% BSA and 0.05% Triton X-100), each time for 3min. Cells were then blocked in blocking buffer (3% BSA in 1× PBT) for 1 h at room temperature. Blocking buffer was removed and cells were washed with 1x PBT quicky then incubated with Invitrogen Alexa Fluor ™ 647 phalloidin (1: 200 dilution) for 1h at RT. Cells were then washed three times with 1x PBT, each time for 3min. Before visualization, cells were mounted on coverslips using Invitrogen SlowFade™ Gold antifade reagent. Cells were imaged using Zeiss LSM 780-FCS Single-point, laser scanning confocal microscope with 405 nm laser.

Image Analysis

All images were analysed with Fiji/ImageJ 1.52i (NIH), Python v.3.10, and MATLAB 2019b (MathWorks) software. We utilized GraphPad Prism 10 (GraphPad), OriginPro v.9.0 (Originlab Corporation) and Microsoft Excel (Microsoft) for plotting our results.

Kymographs. For the membrane kymographs, the cells were segmented against the background following standard image-processing steps with custom code written in

MATLAB 2019b. The kymographs were created from the segmented cells as previously
described, where consecutive lines over time were aligned by minimizing the sum of the
Euclidean distances between the coordinates in two consecutive frames using a custom-
written MATLAB function [19]. A linear colourmap was used for the normalized intensities
in the kymographs. In coloured kymographs, the lowest intensity is indicated by blue and
the highest by yellow. Line kymographs that accompanied ventral waves were generated
in Fiji/ImageJ by drawing a thick line with a line width of 8–12 pixels and processing the
entire stack in the 'KymographBuilder' plugin.

t-stacks. The ImageJ (NIH) software was used for image processing and analysis. For t-
stacks, TIRF time-lapse videos were opened and converted to greyscale in ImageJ. The
ImageJ 3D Viewer plugin was then used to stack the frames from the video. The
resampling factor is set to 1 to avoid blurring of activities between frames.

Line-scan intensity profile. Line scans were performed in Fiji/ImageJ (NIH) by drawing
a straight-line segment (using line tool) inside the cells with a line width of 8–12 pixels to
obtain an average intensity value. The intensity values along that particular line were
obtained in the green and red channels using the 'Plot Profile' option. The values were
saved and normalized in OriginPro 9.0 (OriginLab). The intensity profiles were graphed
and finally smoothened using the adjacent-averaging method, using proper boundary
conditions. The plots were then normalized by dividing by the maximum value. For a
particular line scan, the green and red profiles were smoothed and processed using the
exact same parameters to maintain consistency.

Membrane to cytosol ratio quantification. The cytosol of cells was first outlined with
freehand selection tool in Fiji/ImageJ (NIH). The mean intensity of the cytosol was
measured using analyze>>measure function. Then make band function was applied to
the outline and proper band size was applied according to the cells. Then mean
membrane intensity was measured using the same method. Using mean membrane to
cytosol intensity to divide mean cytosol intensity, we can get the membrane to cytosol
ratio for each biosensor. Each data was then plotted as box-and-whisker graphs in
GraphPad Prism 10.

Cell migration analysis Analysis was performed by segmenting *Dictyostelium* or
neutrophil cells in Fiji/ImageJ 1.52i software. For this, image stack was thresholded using
the 'Threshold' option. 'Calculate threshold for each image' box was unchecked, and
range was not reset. Next, cell masks were created by size-based thresholding using the
'Analyzed particles' option. To optimize binarized masks, 'Fill holes', 'Dilate', and 'Erode'
were done several times. For creating temporal color-coded cell outlines, 'Outline' was
applied on binarized masks, followed by 'Temporal-Color Code' option. Next, 'Centroid'
and 'Shape descriptors' boxes were checked in 'Set Measurements' option under
'Analyze' tab. This provided us with values for centroid coordinates and aspect ratio.
Mean and SEM from replicates of aspect ratio values were determined and plotted in
GraphPad Prism 10. The starting point for centroid values was set to zero for each track,
and these new coordinates were plotted in Microsoft Excel to generate migration tracks.
Velocity was calculated by computing displacement between two consecutive frames.
Displacement was then divided by time interval to obtain speed for each cell. These speed

values were then time-averaged over all frames to produce data points for cell speed
which were plotted as box-and-whisker graphs in GraphPad Prism 10.

Reaction-diffusion model

1082 ***Core excitable network.*** The simulations are based on a model, whose core consists of
1083 three interacting species, Ras, PI(4,5)P2, and PKB, that together form an excitable
1084 network [46]. The concentrations of each of these molecules is described by stochastic,
1085 reaction-diffusion partial differential equations (RD-PDEs):

$$\frac{\partial[\text{Ras}]}{\partial t} = -(a_1 + a_2[\text{PKB}])[\text{Ras}] + \frac{a_3}{1 + a_4^2[\text{PIP2}]^2} + a_5 + w_{\text{Ras}} + D_{\text{Ras}}\nabla^2[\text{Ras}]$$

$$\frac{\partial[\text{PIP2}]}{\partial t} = -(b_1 + b_2[\text{Ras}])[\text{PIP2}] + b_3 + w_{\text{PIP2}} + D_{\text{PIP2}}\nabla^2[\text{PIP2}]$$

$$\frac{\partial[\text{PKB}]}{\partial t} = -c_1[\text{PKB}] + c_2[\text{Ras}] + w_{\text{PKB}} + D_{\text{PKB}}\nabla^2[\text{PKB}]$$

1089 In each of these equations, the final term represents the diffusion of the species, where
1090 D_* is the respective diffusion coefficient and ∇^2 is the spatial Laplacian (in one dimension).
1091 The second-to-last terms represent the molecular noise. Our model assumes a Langevin
1092 approximation in which the size of the noise is based on the reaction terms [108]. For
1093 example, in the case of PKB, the noise is given by

$$w_{\text{PKB}}(t) = \alpha\sqrt{c_1[\text{PKB}] + c_2[\text{Ras}]} \times w(t)$$

where $w(t)$ is a zero mean, unit variance, Gaussian, white noise process. In the
simulations, the size of this noise was adjusted with the empirical parameter α.

PIP5K incorporation through dynamic partitioning. To this model we include the
contribution of PIP5K, and base it on a dynamic partition principle [18]. In this case, PI5PK
exists on the membrane in two states with corresponding low and high diffusivities (PI5K_loc
and PI5K_glo). The conversion from low to high diffusivity is mediated by Ras. Together,
they obey the following two RD-PDEs:

$$\frac{\partial[\text{PI5K}_{\text{loc}}]}{\partial t} = -\left(\frac{[\text{Ras}]^2}{d_1 + [\text{Ras}]^2}\right)[\text{PI5K}_{\text{loc}}] + d_2[\text{PI5K}_{\text{glo}}] + D_{\text{PI5K}_{\text{loc}}}\nabla^2[\text{PI5K}_{\text{loc}}]$$

$$\frac{\partial[\text{PI5K}_{\text{glo}}]}{\partial t} = -d_2[\text{PI5K}_{\text{glo}}] + \left(\frac{[\text{Ras}]^2}{d_1 + [\text{Ras}]^2}\right)[\text{PI5K}_{\text{loc}}] + D_{\text{PI5K}_{\text{glo}}}\nabla^2[\text{PI5K}_{\text{glo}}]$$

Note that we do not include noise in these equations. That the total concentration is
conserved can be seen by noting that the sum of the reaction terms is equal to zero. Thus,
the only changes in concentration appear through the diffusion terms. Integrating over
the complete domain and using the fact that boundary conditions are periodic guarantee
that the total concentration of PI5PK is constant. The main effect of PI5PK is to increase
the production of PIP2:

$$b_1 \mapsto b_1 \times (1 + d_3[\text{PI5K}_{\text{loc}}])$$

1111 ***Cytoskeleton feedback loops.*** Finally, we incorporated two other terms. The first
1112 represents feedback from branched actin onto Ras. Since the actin cytoskeleton is not
1113 directly modeled, we model the origin of this feedback from PKB, which is upstream to
1114 the cytoskeleton [40] and incorporated one actin term:

1117
$$\frac{\partial [P_{Actin}]}{\partial t} = -p_1[P_{Actin}] + p_2[PKB] + D_{P_{Actin}}\nabla^2[P_{Actin}]$$

1115 The second feedback loop accounts for the inhibitory regulation by actomyosin on Ras
1116 that originates from PIP2:

1119
$$\frac{\partial [P_{Myosin}]}{\partial t} = -p_3[P_{Myosin}] + p_4[PIP2] + D_{P_{Actin}}\nabla^2[P_{Actin}]$$

1118
1120 Together they modify the equations for Ras:

1121
$$(a_1 + a_2[PKB])[Ras] \mapsto (a_1 + a_2[PKB])[Ras] \times \left(1 + p_m[P_{Myosin}]\right)$$

1122
$$\frac{a_3}{1 + a_4^2[PIP2]^2} + a_5 \mapsto \left(\frac{a_3}{1 + a_4^2[PIP2]^2} + a_5\right) \times (1 + p_a[P_{Actin}])$$

1123
1124 ***Modeling perturbations.*** The perturbations in Fig. 7c-d were done by modifying the
1125 decay and production rates of the relevant species during the simulation. Specifically:
1126

a. Eliminating PIP5K: $\left(\frac{\partial[PI5K_*]}{\partial t}\right)_{new} = \left(\frac{\partial[PI5K_*]}{\partial t}\right)_{old} - 0.2[PI5K_*]$, for both types of
PIP5K
b. Increasing PIP5K: $\left(\frac{\partial[PI5K_*]}{\partial t}\right)_{new} = \left(\frac{\partial[PI5K_*]}{\partial t}\right)_{old} + 0.001 - 0.2[PI5K_*]$, for both
types of PIP5K.
c. Eliminating myosin: $p_4 = 0$ and $p_m = 0$.
d. Eliminating actin: $p_2 = 0$ and $p_a = 0$.
1133
Implementing the simulations. Simulations were run on MATLAB 2023b (MathWorks,
Natick, MA) on custom-code based on the Itô solution in the Stochastic Differential
Equation toolbox (http://sdetoolbox.sourceforge.net). The simulations aim to recreate the
membrane fluorescence observed in single-cell confocal images. The dimension is
therefore smaller, assuming a cell radius of 5 μm and a spacing of 0.25 μm, resulting in
$2\pi \times 5/0.25 \approx 126$ points along the perimeter, and periodic boundary conditions. Initial
conditions were set by solving the steady-state value of the core model. The PIP5K values
are normalized. All plots shown are after a 200 second interval so that the effect of the
initial values is minimized. For the kymographs, contrast for each species was done
through MATLAB's imadjust command.

Parameter	Value	Units	Parameter	Value	Units
a_1	0.122	s^{-1}	a_2	2.85	μM^{-1} s^{-1}
a_3	1.14	s^{-1}	a_4	440	μM^{-1}

a_5	1.02×10^{-3}	s^{-1}	b_1	0.0144	s^{-1}
b_2	2134	µM^{-1} s^{-1}	b_3	1.75	µM s^{-1}
c_1	0.0744	µM^{-1} s^{-1}	c_2	0.864	µM^{-1} s^{-1}
d_1	0.025	µM^{-1}	d_2	0.05	µM^{-1} s^{-1}
d_3	160	µM^{-1}			
p_1	0.05	s^{-1}	p_2	0.05	s^{-1}
p_3	0.05	s^{-1}	p_4	0.05	s^{-1}
D_{Ras}	0.03	µm^2/s	D_{PIP2}	0.03	µm^2/s
D_{PKB}	0.09	µm^2/s			
$D_{P_{\mathrm{Myosin}}}$	0.005	µm^2/s	$D_{P_{\mathrm{Actin}}}$	0.0025	µm^2/s
$D_{PI5K_{local}}$	0.0025	$µm2/s$	$D_{PI5K_{global}}$	0.5	µm^2/s
α	0.0833				

1144

Statistics and reproducibility

Statistical analyses were executed using unpaired or paired two-tailed
nonparametric tests on GraphPad Prism 10. Results are expressed as mean ± s.d. from
at least three independent experiments. NS, $P > 0.05$, *$P \leq 0.05$, **$P \leq 0.01$, ***$P \leq 0.001$,
***$P \leq 0.0001$. Tukey's convention was used to plot box-and-whisker plots. Statistical test
details are indicated in the figure legends. No statistical methods were used to
predetermine sample sizes, but our sample sizes are similar to those reported in previous
publications[4,35,79]. Each representative image or image series is from n > 3 independent
experiments. Any variations observed between treatment groups are not attributed to
sampling bias. No data were excluded from the analyses.

Data availability

1156 All data are provided in the main or supplementary text. Requests for additional
1157 information on this work are to be made to the corresponding author.

1158

1159

1160

1161

1162

1163

1164

1165

Reference

1. Sasaki, A.T., et al., *G protein-independent Ras/PI3K/F-actin circuit regulates basic cell motility.* J Cell Biol, 2007. **178**(2): p. 185-91.
2. Weiner, O.D., et al., *Spatial control of actin polymerization during neutrophil chemotaxis.* Nat Cell Biol, 1999. **1**(2): p. 75-81.
3. Cai, H. and P.N. Devreotes, *Moving in the right direction: how eukaryotic cells migrate along chemical gradients.* Semin Cell Dev Biol, 2011. **22**(8): p. 834-41.
4. Wang, Y., et al., *Identifying network motifs that buffer front-to-back signaling in polarized neutrophils.* Cell Rep, 2013. **3**(5): p. 1607-16.
5. Swaney, K.F., C.H. Huang, and P.N. Devreotes, *Eukaryotic chemotaxis: a network of signaling pathways controls motility, directional sensing, and polarity.* Annu Rev Biophys, 2010. **39**: p. 265-89.
6. Parent, C.A., *Making all the right moves: chemotaxis in neutrophils and Dictyostelium.* Curr Opin Cell Biol, 2004. **16**(1): p. 4-13.
7. SenGupta, S., C.A. Parent, and J.E. Bear, *The principles of directed cell migration.* Nat Rev Mol Cell Biol, 2021. **22**(8): p. 529-547.
8. Saha, S., T.L. Nagy, and O.D. Weiner, *Joining forces: crosstalk between biochemical signalling and physical forces orchestrates cellular polarity and dynamics.* Philos Trans R Soc Lond B Biol Sci, 2018. **373**(1747).
9. Shellard, A. and R. Mayor, *All Roads Lead to Directional Cell Migration.* Trends Cell Biol, 2020. **30**(11): p. 852-868.
10. Ridley, A.J., et al., *Cell migration: integrating signals from front to back.* Science, 2003. **302**(5651): p. 1704-9.
11. Hadjitheodorou, A., et al., *Directional reorientation of migrating neutrophils is limited by suppression of receptor input signaling at the cell rear through myosin II activity.* Nat Commun, 2021. **12**(1): p. 6619.
12. Ghabache, E., et al., *Coupling traction force patterns and actomyosin wave dynamics reveals mechanics of cell motion.* Mol Syst Biol, 2021. **17**(12): p. e10505.
13. Li, X., et al., *Excitable networks controlling cell migration during development and disease.* Semin Cell Dev Biol, 2020. **100**: p. 133-142.
14. Ladoux, B., R.M. Mege, and X. Trepat, *Front-Rear Polarization by Mechanical Cues: From Single Cells to Tissues.* Trends Cell Biol, 2016. **26**(6): p. 420-433.
15. Bagorda, A. and C.A. Parent, *Eukaryotic chemotaxis at a glance.* J Cell Sci, 2008. **121**(Pt 16): p. 2621-4.
16. Rickert, P., et al., *Leukocytes navigate by compass: roles of PI3Kgamma and its lipid products.* Trends Cell Biol, 2000. **10**(11): p. 466-73.
17. Houk, A.R., et al., *Membrane tension maintains cell polarity by confining signals to the leading edge during neutrophil migration.* Cell, 2012. **148**(1-2): p. 175-88.
18. Banerjee, T., et al., *A dynamic partitioning mechanism polarizes membrane protein distribution.* Nat Commun, 2023. **14**(1): p. 7909.
19. Banerjee, T., et al., *Spatiotemporal dynamics of membrane surface charge regulates cell polarity and migration.* Nature Cell Biology, 2022. **24**(10): p. 1499-1515.

20.   Huang, C.H., et al., *An excitable signal integrator couples to an idling cytoskeletal*
oscillator to drive cell migration. Nat Cell Biol, 2013. **15**(11): p. 1307-16.

21.   Ghose, D., T. Elston, and D. Lew, *Orientation of Cell Polarity by Chemical Gradients*.
Annu Rev Biophys, 2022. **51**: p. 431-451.

22.   Pal, D.S., et al., *The excitable signal transduction networks: movers and shapers of*
eukaryotic cell migration. The International Journal of Developmental Biology, 2019.
63(8-9-10): p. 407-416.

23.   Schink, K.O., K.W. Tan, and H. Stenmark, *Phosphoinositides in Control of Membrane*
Dynamics. Annu Rev Cell Dev Biol, 2016. **32**: p. 143-171.

24.   Shewan, A., D.J. Eastburn, and K. Mostov, *Phosphoinositides in cell architecture*.
Cold Spring Harb Perspect Biol, 2011. **3**(8): p. a004796.

25.   Carrasco-Rando, M., et al., *An acytokinetic cell division creates PIP2-enriched*
membrane asymmetries leading to slit diaphragm assembly in Drosophila
nephrocytes. Development, 2023. **150**(18).

26.   Wu, M. and J. Liu, *Mechanobiology in cortical waves and oscillations*. Curr Opin Cell
Biol, 2021. **68**: p. 45-54.

27.   Cramer, L.P., *Forming the cell rear first: breaking cell symmetry to trigger directed cell*
migration. Nat Cell Biol, 2010. **12**(7): p. 628-32.

28.   Teruel, M.N. and T. Meyer, *Translocation and reversible localization of signaling*
proteins: a dynamic future for signal transduction. Cell, 2000. **103**(2): p. 181-4.

29.   De Belly, H., et al., *Cell protrusions and contractions generate long-range membrane*
tension propagation. Cell, 2023. **186**(14): p. 3049-3061 e15.

30.   Carvalho, K., et al., *Actin polymerization or myosin contraction: two ways to build up*
cortical tension for symmetry breaking. Philos Trans R Soc Lond B Biol Sci, 2013.
368(1629): p. 20130005.

31.   Xiong, Y., et al., *Cells navigate with a local-excitation, global-inhibition-biased*
excitable network. Proc Natl Acad Sci U S A, 2010. **107**(40): p. 17079-86.

32.   Wang, X. and A. Harrison, *A general principle for spontaneous genetic symmetry*
breaking and pattern formation within cell populations. J Theor Biol, 2021. **526**: p.
110809.

33.   van Haastert, P.J.M., *Symmetry Breaking during Cell Movement in the Context of*
Excitability, Kinetic Fine-Tuning and Memory of Pseudopod Formation. Cells, 2020.
9(8).

34.   Zhan, H., et al., *An Excitable Ras/PI3K/ERK Signaling Network Controls Migration and*
Oncogenic Transformation in Epithelial Cells. Developmental Cell, 2020. **54**(5): p.
608-623.e5.

35.   Li, X., et al., *Mutually inhibitory Ras-PI(3,4)P(2) feedback loops mediate cell*
migration. Proc Natl Acad Sci U S A, 2018. **115**(39): p. E9125-E9134.

36.   Devreotes, P.N., et al., *Excitable Signal Transduction Networks in Directed Cell*
Migration. Annual Review of Cell and Developmental Biology, 2017. **33**(1): p. 103-125.

37.   Buckley, C.M., et al., *Coordinated Ras and Rac Activity Shapes Macropinocytic Cups*
and Enables Phagocytosis of Geometrically Diverse Bacteria. Curr Biol, 2020. **30**(15):
p. 2912-2926 e5.

38.   Bisaria, A., et al., *Membrane-proximal F-actin restricts local membrane protrusions
1252   and directs cell migration*. Science, 2020. **368**(6496): p. 1205-1210.
1253   39.   Tsai, T.Y., et al., *Efficient Front-Rear Coupling in Neutrophil Chemotaxis by Dynamic
1254   Myosin II Localization*. Dev Cell, 2019. **49**(2): p. 189-205 e6.
1255   40.   Miao, Y., et al., *Wave patterns organize cellular protrusions and control cortical
1256   dynamics*. Mol Syst Biol, 2019. **15**(3): p. e8585.
1257   41.   Gerhardt, M., et al., *Actin and PIP3 waves in giant cells reveal the inherent length
1258   scale of an excited state*. J Cell Sci, 2014. **127**(Pt 20): p. 4507-17.
1259   42.   Matsuoka, S. and M. Ueda, *Mutual inhibition between PTEN and PIP3 generates
1260   bistability for polarity in motile cells*. Nat Commun, 2018. **9**(1): p. 4481.
1261   43.   Lin, Y., et al., *Ras suppression potentiates rear actomyosin contractility-driven cell
1262   polarization and migration*. Nat Cell Biol, 2024.
1263   44.   Pal, D.S., et al., *Actuation of single downstream nodes in growth factor network steers
1264   immune cell migration*. Dev Cell, 2023. **58**(13): p. 1170-1188 e7.
1265   45.   Kolch, W., D. Berta, and E. Rosta, *Dynamic regulation of RAS and RAS signaling*.
1266   Biochem J, 2023. **480**(1): p. 1-23.
1267   46.   Miao, Y., et al., *Altering the threshold of an excitable signal transduction network
1268   changes cell migratory modes*. Nature Cell Biology, 2017. **19**(4): p. 329-340.
1269   47.   van den Bout, I. and N. Divecha, *PIP5K-driven PtdIns(4,5)P2 synthesis: regulation and
1270   cellular functions*. J Cell Sci, 2009. **122**(Pt 21): p. 3837-50.
1271   48.   Ling, K., et al., *Type I gamma phosphatidylinositol phosphate kinase modulates
1272   adherens junction and E-cadherin trafficking via a direct interaction with mu 1B
1273   adaptin*. J Cell Biol, 2007. **176**(3): p. 343-53.
1274   49.   Lacalle, R.A., et al., *Type I phosphatidylinositol 4-phosphate 5-kinase controls
1275   neutrophil polarity and directional movement*. J Cell Biol, 2007. **179**(7): p. 1539-53.
1276   50.   El Sayegh, T.Y., et al., *Phosphatidylinositol-4,5 bisphosphate produced by
1277   PIP5KIgamma regulates gelsolin, actin assembly, and adhesion strength of N-
1278   cadherin junctions*. Mol Biol Cell, 2007. **18**(8): p. 3026-38.
1279   51.   Fets, L., J.M. Nichols, and R.R. Kay, *A PIP5 kinase essential for efficient chemotactic
1280   signaling*. Curr Biol, 2014. **24**(4): p. 415-21.
1281   52.   Swaney, K.F., et al., *Novel protein Callipygian defines the back of migrating cells*. Proc
1282   Natl Acad Sci U S A, 2015. **112**(29): p. E3845-54.
1283   53.   Flemming, S., et al., *How cortical waves drive fission of motile cells*. Proc Natl Acad
1284   Sci U S A, 2020. **117**(12): p. 6330-6338.
1285   54.   Masters, T.A., M.P. Sheetz, and N.C. Gauthier, *F-actin waves, actin cortex
1286   disassembly and focal exocytosis driven by actin-phosphoinositide positive
1287   feedback*. Cytoskeleton (Hoboken), 2016. **73**(4): p. 180-96.
1288   55.   Zhan, H., et al., *Self-organizing glycolytic waves fuel cell migration and cancer
1289   progression*. bioRxiv, 2024.
1290   56.   Lokuta, M.A., et al., *Type Igamma PIP kinase is a novel uropod component that
1291   regulates rear retraction during neutrophil chemotaxis*. Mol Biol Cell, 2007. **18**(12): p.
1292   5069-80.
1293   57.   Shin, D.Y., et al., *Sphingomyelin metabolism underlies Ras excitability for efficient cell
1294   migration and chemotaxis*. Cell Struct Funct, 2023. **48**(2): p. 145-160.

58.   Fukushima, S., S. Matsuoka, and M. Ueda, *Excitable dynamics of Ras triggers spontaneous symmetry breaking of PIP3 signaling in motile cells*. J Cell Sci, 2019. **132**(5).

59.   Arai, Y., et al., *Self-organization of the phosphatidylinositol lipids signaling system for random cell migration*. Proc Natl Acad Sci U S A, 2010. **107**(27): p. 12399-404.

60.   Sobczyk, G.J., J. Wang, and C.J. Weijer, *SILAC-based proteomic quantification of chemoattractant-induced cytoskeleton dynamics on a second to minute timescale*. Nat Commun, 2014. **5**: p. 3319.

61.   Sasaki, A.T., et al., *Localized Ras signaling at the leading edge regulates PI3K, cell polarity, and directional cell movement*. J Cell Biol, 2004. **167**(3): p. 505-18.

62.   Iijima, M. and P. Devreotes, *Tumor suppressor PTEN mediates sensing of chemoattractant gradients*. Cell, 2002. **109**(5): p. 599-610.

63.   Parent, C.A., et al., *G protein signaling events are activated at the leading edge of chemotactic cells*. Cell, 1998. **95**(1): p. 81-91.

64.   Janetopoulos, C., et al., *Chemoattractant-induced phosphatidylinositol 3,4,5-trisphosphate accumulation is spatially amplified and adapts, independent of the actin cytoskeleton*. Proc Natl Acad Sci U S A, 2004. **101**(24): p. 8951-6.

65.   Li, X., et al., *Reverse fountain flow of phosphatidylinositol-3,4-bisphosphate polarizes migrating cells*. EMBO J, 2021. **40**(4): p. e105094.

66.   Veltman, D.M., et al., *A plasma membrane template for macropinocytic cups*. Elife, 2016. **5**.

67.   Kunz, J., et al., *The activation loop of phosphatidylinositol phosphate kinases determines signaling specificity*. Mol Cell, 2000. **5**(1): p. 1-11.

68.   Weckerly, C.C., et al., *Nir1-LNS2 is a novel phosphatidic acid biosensor that reveals mechanisms of lipid production*. bioRxiv, 2024.

69.   Inaba, H., Q. Miao, and T. Nakata, *Optogenetic control of small GTPases reveals RhoA mediates intracellular calcium signaling*. J Biol Chem, 2021. **296**: p. 100290.

70.   Kuhn, J., et al., *Complementary Cytoskeletal Feedback Loops Control Signal Transduction Excitability and Cell Polarity*. bioRxiv, 2024.

71.   Zhao, H., M. Hakala, and P. Lappalainen, *ADF/cofilin binds phosphoinositides in a multivalent manner to act as a PIP(2)-density sensor*. Biophys J, 2010. **98**(10): p. 2327-36.

72.   Bae, Y.H., et al., *Profilin1 regulates PI(3,4)P2 and lamellipodin accumulation at the leading edge thus influencing motility of MDA-MB-231 cells*. Proc Natl Acad Sci U S A, 2010. **107**(50): p. 21547-52.

73.   Papayannopoulos, V., et al., *A polybasic motif allows N-WASP to act as a sensor of PIP(2) density*. Mol Cell, 2005. **17**(2): p. 181-91.

74.   Rozelle, A.L., et al., *Phosphatidylinositol 4,5-bisphosphate induces actin-based movement of raft-enriched vesicles through WASP-Arp2/3*. Curr Biol, 2000. **10**(6): p. 311-20.

75.   Janmey, P.A. and T.P. Stossel, *Modulation of gelsolin function by phosphatidylinositol 4,5-bisphosphate*. Nature, 1987. **325**(6102): p. 362-4.

76.   Pan, M., et al., *A G-protein-coupled chemoattractant receptor recognizes lipopolysaccharide for bacterial phagocytosis*. PLoS Biol, 2018. **16**(5): p. e2005754.

77. Li, X.X., R.J. Clark, and T.M. Woodruff, *C5aR2 Activation Broadly Modulates the*
Signaling and Function of Primary Human Macrophages. J Immunol, 2020. **205**(4): p.
1102-1112.
78. Ren, Y., et al., *Mechanosensing through cooperative interactions between myosin II*
and the actin crosslinker cortexillin I. Curr Biol, 2009. **19**(17): p. 1421-8.
79. Yamamoto, K., et al., *Optogenetic relaxation of actomyosin contractility uncovers*
mechanistic roles of cortical tension during cytokinesis. Nat Commun, 2021. **12**(1):
p. 7145.
80. Shibata, K., et al., *Actin binding domain of filamin distinguishes posterior from*
anterior actin filaments in migrating Dictyostelium cells. Biophys Physicobiol, 2016.
13: p. 321-331.
81. Inoue, T. and T. Meyer, *Synthetic activation of endogenous PI3K and Rac identifies an*
AND-gate switch for cell polarization and migration. PLoS One, 2008. **3**(8): p. e3068.
82. Srinivasan, S., et al., *Rac and Cdc42 play distinct roles in regulating PI(3,4,5)P3 and*
polarity during neutrophil chemotaxis. J Cell Biol, 2003. **160**(3): p. 375-85.
83. Kabacoff, C., et al., *Dynacortin facilitates polarization of chemotaxing cells.* BMC
Biol, 2007. **5**: p. 53.
84. Shi, C., et al., *Interaction of motility, directional sensing, and polarity modules*
recreates the behaviors of chemotaxing cells. PLoS Comput Biol, 2013. **9**(7): p.
e1003122.
85. Neilson, M.P., et al., *Chemotaxis: a feedback-based computational model robustly*
predicts multiple aspects of real cell behaviour. PLoS Biol, 2011. **9**(5): p. e1000618.
86. van Rheenen, J., et al., *EGF-induced PIP2 hydrolysis releases and activates cofilin*
locally in carcinoma cells. J Cell Biol, 2007. **179**(6): p. 1247-59.
87. Sengelaub, C.A., et al., *PTPRN2 and PLCbeta1 promote metastatic breast cancer cell*
migration through PI(4,5)P2-dependent actin remodeling. EMBO J, 2016. **35**(1): p. 62-
76.
88. Chinthalapudi, K., E.S. Rangarajan, and T. Izard, *The interaction of talin with the cell*
membrane is essential for integrin activation and focal adhesion formation. Proc Natl
Acad Sci U S A, 2018. **115**(41): p. 10339-10344.
89. Thapa, N., et al., *PIPKIgamma and talin couple phosphoinositide and adhesion*
signaling to control the epithelial to mesenchymal transition. Oncogene, 2017. **36**(7):
p. 899-911.
90. Goni, G.M., et al., *Phosphatidylinositol 4,5-bisphosphate triggers activation of focal*
adhesion kinase by inducing clustering and conformational changes. Proc Natl Acad
Sci U S A, 2014. **111**(31): p. E3177-86.
91. Cai, X., et al., *Spatial and temporal regulation of focal adhesion kinase activity in living*
cells. Mol Cell Biol, 2008. **28**(1): p. 201-14.
92. Gilmore, A.P. and K. Burridge, *Regulation of vinculin binding to talin and actin by*
phosphatidyl-inositol-4-5-bisphosphate. Nature, 1996. **381**(6582): p. 531-5.
93. Fukami, K., et al., *alpha-Actinin and vinculin are PIP2-binding proteins involved in*
signaling by tyrosine kinase. J Biol Chem, 1994. **269**(2): p. 1518-22.
94. Adhikari, H. and C.M. Counter, *Interrogating the protein interactomes of RAS isoforms*
identifies PIP5K1A as a KRAS-specific vulnerability. Nat Commun, 2018. **9**(1): p. 3646.

95. Watts, D.J. and J.M. Ashworth, *Growth of myxameobae of the cellular slime mould Dictyostelium discoideum in axenic culture*. Biochem J, 1970. **119**(2): p. 171-4.

96. Lilly, P., et al., *A G-protein beta-subunit is essential for Dictyostelium development*. Genes Dev, 1993. **7**(6): p. 986-95.

97. Lampert, T.J., et al., *Shear force-based genetic screen reveals negative regulators of cell adhesion and protrusive activity*. Proc Natl Acad Sci U S A, 2017. **114**(37): p. E7727-E7736.

98. Millius, A. and O.D. Weiner, *Manipulation of neutrophil-like HL-60 cells for the study of directed cell migration*. Methods Mol Biol, 2010. **591**: p. 147-58.

99. Rincon, E., B.L. Rocha-Gregg, and S.R. Collins, *A map of gene expression in neutrophil-like cell lines*. BMC Genomics, 2018. **19**(1): p. 573.

100. Rovera, G., D. Santoli, and C. Damsky, *Human promyelocytic leukemia cells in culture differentiate into macrophage-like cells when treated with a phorbol diester*. Proc Natl Acad Sci U S A, 1979. **76**(6): p. 2779-83.

101. Meshik, X., P.R. O'Neill, and N. Gautam, *Optogenetic Control of Cell Migration*. Methods Mol Biol, 2018. **1749**: p. 313-324.

102. Bell, G.R.R., et al., *Optogenetic control of receptors reveals distinct roles for actin- and Cdc42-dependent negative signals in chemotactic signal processing*. Nat Commun, 2021. **12**(1): p. 6148.

103. Pal, D.S., et al., *Optogenetic modulation of guanine nucleotide exchange factors of Ras superfamily proteins directly controls cell shape and movement*. Frontiers in Cell and Developmental Biology, 2023. **11**.

104. Kamimura, Y., M. Tang, and P. Devreotes, *Assays for chemotaxis and chemoattractant-stimulated TorC2 activation and PKB substrate phosphorylation in Dictyostelium*. Methods Mol Biol, 2009. **571**: p. 255-70.

105. Cai, H., et al., *Nucleocytoplasmic shuttling of a GATA transcription factor functions as a development timer*. Science, 2014. **343**(6177): p. 1249531.

106. Aufderheide, K.J. and C. Janetopoulos, *Migration of Dictyostelium discoideum to the Chemoattractant Folic Acid*. Methods Mol Biol, 2016. **1407**: p. 25-39.

107. Sekine, R., T. Kawata, and T. Muramoto, *CRISPR/Cas9 mediated targeting of multiple genes in Dictyostelium*. Sci Rep, 2018. **8**(1): p. 8471.

108. Gillespie, D.T., *The chemical Langevin equation*. Journal of Chemical Physics, 2000. **113**(1): p. 297-306.

Main Figures

Figure 1. Lowering PIP2 increases signal transduction, cytoskeletal, and protrusive activities (a-c) Representative live-cell time-lapse confocal images (left, DIC) and color-coded temporal overlay profiles (right) of *Dictyostelium* AX2 (WT) cells **(a)**, *pi5k-* fan-shaped **(b)**, and *pi5k-* oscillatory cells **(c)**. A linear color map shows that green corresponds to 0 min and red corresponds to 15 min. The yellow outline corresponds to the frame of time-lapse images and cell boundaries. Scale bars represent 5 mm. **(d-f)** Centroid tracks of cells (n_c=22) showing random motility in AX2 (WT) cells **(d)**, *pi5k-* fan-shaped **(e)**, and *pi5k-* oscillatory cells **(f)**. Each track lasted at least 15 minutes and was reset to the same origin. **(g-h)** Box-and-whisker plots of **(g)** cell migration speed, **(h)** cell area. n_c=22 from at least 3 independent experiments; asterisks indicate significant difference, ****$P \leq 0.0001$ (Mann-Whitney test. Compare ranks). The median is at the center, and whiskers and outliers are graphed according to Tukey's convention (GraphPad Prism 10). **(i-j)** Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 (WT) cells (left), and *pi5k-* cells (right) expressing PHPLCδ-YFP (biosensor for PI(4,5)P2) **(i)** or RBD-GFP (biosensor for activated Ras) **(j)**. Scale bars represent 5 mm. **(k-l)** Box-and-whisker plots of **(k)** PHPLCd1 membrane-to-cytosol ratio, **(l)** RBD patch length/Cell Perimeter. n_c=22 from at least 3 independent experiments; asterisks indicate significant difference, ****$P \leq 0.0001$ (Mann-Whitney test. Compare ranks). The median is at the center, and whiskers and outliers are graphed according to Tukey's convention (GraphPad Prism 10). **(m-n)** Representative membrane kymograph of RBD intensity in AX2 **(m)** or *pi5k-* **(n)** cells, respectively. A linear color map shows that blue has the lowest RBD intensity, whereas yellow has the highest. **(o)** Time-lapse confocal images of differentiated HL-60 neutrophil expressing CIBN-CAAX, CRY2PHR-mCherry-Inp54p (magenta; upper panel) and LifeAct-miRFP703 (cyan; lower panel), before or after 488 nm laser was switched on globally. Time in min:sec format. Scale bars represent 5 μm. **(p-q)** Box-and-whisker plots of **(p)** cell area, **(q)** cell migration speed correspond to (o). n_c=10 from at least 3 independent experiments; asterisks indicate significant difference, ****$P \leq 0.0001$ (Mann-Whitney test. Compare ranks). The median is at the center, and whiskers and outliers are graphed according to Tukey's convention (GraphPad Prism 10). **(r)** Cartoon illustrating mechanism of opto-Inp54p global recruitment on differentiated HL-60 neutrophil membrane with the help of CRY2-CIBN optogenetic system. **(s)** Cartoon illustrating signal transduction activities in amoeboid, fan-shaped and oscillatory cells.

Figure 2 PIP5Ks localize to the back-state regions of the membrane. (a) Representative live-cell time-lapse images of Dictyostelium cells coexpressing PI5K-GFP and PHcrac–RFP (biosensor for PIP3) during migration showing PIP5K dynamically moves away from protrusions in migrating cells. Time in sec format. Scale bars represent 5 mm. **(b-c)** Representative live-cell time-lapse images of differentiated HL-60 neutrophils coexpressing PIP5K1C-GFP and LifeAct-iRFP703 (biosensor for F-actin) **(b)** or expressing PIP5K1B-GFP **(c)** during migration showing PIP5Ks dynamically moves away from protrusions in migrating cells. Time in min:sec format. Scale bars represent 5 mm. **(d-e)** Representative live-cell time-lapse images of Dictyostelium cells coexpressing PIP5K-GFP and PHcrac-RFP (biosensor for PIP3) **(d)** or PIP5K-mCherry and PHPLCδ–GFP (biosensor for PI(4,5)P2) **(e)** during ventral wave propagation, showing PI5K dynamically localizes to the back-state regions in ventral waves. Line-scan intensity profiles are shown in the bottommost panels. The red line represents PHcrac and PI5K in (d) and (e). The green line represents PIP5K and PHPLCδ. Time in min:sec format. Scale bars represent 5 mm. **(f)** Representative live-cell images of Dictyostelium cell co-expressing PIP5K-mCherry and RBD-GFP (biosensor for activated Ras), where the substrate-attached ventral surface was imaged, showing consistent complementarity between PIP5K and RBD during 2D wave propagation, even in the absence of actin cytoskeleton. Cells were pre-treated with actin polymerization inhibitor Latrunculin A (final concentration 5μM) for 20min. Line-scan intensity profiles are shown in the bottommost panels. Red line and green line represent PIP5K and RBD, respectively. Time in sec format. Scale bars represent 5 mm. **(g)** Live-cell images of a differentiated HL-60 macrophage coexpressing PIP5K1B and LifeAct demonstrating dynamic complementary distribution in its ventral waves. Line-scan intensity profiles are shown in the bottommost panels. Green line and Cyan line represent PIP5K1B and LifeAct, respectively. Time in min:sec format. Scale bars represent 5 mm. (h) Representative live-cell time-lapse images of Dictyostelium cells expressing KikGR-PIP5K. Yellow box indicates where the photo conversion happened. Top panel is GFP channel (488nm), while bottom panel is RFP channel (561nm). Time in min:sec format. Scale bars represent 5 mm. **(i)** Scatter plot of RFP channel fluorescence intensity vs. time. The red line indicated when the conversions happened. **(j)** Cartoon illustrating the front-back complementarity in migrating cell protrusions and ventral wave propagation.

Figure 3 Expressing PIP5Ks suppresses cell protrusions and alters migration. (a-b) Representative live-cell images of *Dictyostelium* cells expressing doxycycline-inducible PIP5K without DOX induction **(a)** or with overnight DOX induction (rounded) **(b)**. Time in sec format. Scale bars represent 5 μm. **(c-d)** Representative live-cell images of differentiated HL-60 neutrophils (WT) **(c)** and expressing PIP5K1B (rounded) **(d)**. Time in min:sec format. Scale bars represent 5 μm. **(e-f)** Representative live-cell images of differentiated HL-60 macrophages (WT) **(e)** and expressing PIP5K1B **(f)**. Time in min:sec format. Scale bars represent 5 μm. **(g-h)** Box-and-whisker plots of cell migration speed of Dictyostelium cells expressing PIP5K at different DOX incubation time **(g)**, or differentiated HL-60 neutrophils expressing PIP5Ks **(h)**. n_c=20 from at least 3 independent experiments; asterisks indicate significant difference, ****$P \leq 0.0001$ (Mann-Whitney test. Compare ranks). The median is at the center, and whiskers and outliers are graphed according to Tukey's convention (GraphPad Pris 10). **(i-j)** Color-coded temporal overlay profiles of Dictyostelium cells expressing doxycycline-inducible PIP5K without DOX incubation **(i)** or with overnight DOX incubation **(j)**. **(k)** Time-lapse confocal images of MDA-MB-231 cells expressing crimson-SspB-PIP5K1C-P2A-iLiD-CAAX, before or after 488 nm laser was switched on globally. Time in min:sec format. Scale bars represent 10 μm. Blue arrows indicate where retraction fibers or blebs are formed. Cells are pretreated with 10 ng/ml EGF for 10 mins. **(l-m)** Box-and-whisker plots of **(l)** cell area, **(m)** aspect ratio correspond to (m-n). n_c=10 from at least 3 independent experiments; asterisks indicate significant difference, ****$P \leq 0.0001$ (Mann-Whitney test. Compare ranks). The median is at the center, and whiskers and outliers are graphed according to Tukey's convention (GraphPad Prism 10).

Figure 4 Expressing PIP5Ks suppresses signal transduction and cytoskeletal activities. (a-b) Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 co-expressing RBD-GFP (biosensor for activated Ras) and doxycycline-inducible PI5K without DOX induction **(a)** or with overnight DOX induction (rounded) **(b)**. Time in sec format. Scale bars represent 5 mm. **(c-d)** Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 co-expressing PHcrac-YFP (biosensor for PIP3) and doxycycline-inducible PI5K without DOX induction **(c)** or with overnight DOX induction (rounded) **(d)**. Time in sec format. Scale bars represent 5 mm. **(e-f)** Box-and-whisker plots of RBD patch size (left axis) and RBD patch number (right axis) **(e),** or PHcrac patch size (left axis) and PHcrac patch number (right axis) **(f).** n_c=20 from at least 3 independent experiments; asterisks indicate significant difference, ****P ≤ 0.0001 (Mann-Whitney test. Compare ranks). The median is at the center, and whiskers and outliers are graphed according to Tukey's convention (GraphPad Prism 10). **(g)** Box-and-whisker plots individual RBD patch size at different patch numbers. Data is from Figure e. n_c=10 for column 1, n_c=26 for column 2, n_c=42 for column 3, n_c=45 for column 4, n_c=5 for column 5, and n_c=6 for column 6. All results are from at least 3 independent experiments; asterisks indicate significant difference, ****P ≤ 0.0001 (Mann-Whitney test. Compare ranks). The median is at the center, and whiskers and outliers are graphed according to Tukey's convention (GraphPad Prism 10). **(h)** Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 co-expressing RBD-GFP (biosensor for activated Ras) and doxycycline-inducible PI5K without DOX induction (upper panel) or with overnight DOX induction (lower panel) during ventral wave propagation. Time in min:sec format. Scale bars represent 10 mm. **(i-j)** Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 co-expressing RBD-GFP (biosensor for activated Ras) and doxycycline-inducible PI5K without DOX induction **(i)** or with overnight DOX induction **(j),** even in the absence of actin cytoskeleton. Cells were pre-treated with actin polymerization inhibitor Latrunculin A (final concentration 5μM) and caffeine (final concentration 4mM) for 20min. Time in sec format. Scale bars represent 5 mm. **(k-l)** Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 co-expressing PHcrac-YFP (biosensor for PIP3) and doxycycline-inducible PI5K without DOX induction **(k)** or with overnight DOX induction **(l),** even in the absence of actin cytoskeleton. Cells were pre-treated with actin polymerization inhibitor Latrunculin A (final concentration 5μM) and caffeine (final concentration 4mM) for 20min. Time in sec format. Scale bars represent 5 mm. **(m-n)** Box-and-whisker plots of RBD crescent size **(m),** or PHcrac crescent size **(n).** n_c=35 from at least 3 independent experiments; asterisks indicate significant difference, ****P ≤ 0.0001 (Mann-Whitney test. Compare ranks). The median is at the center, and whiskers and outliers are graphed according to Tukey's convention (GraphPad Prism 10). **(o)** Cartoon illustrating the localization of front signaling components in cell protrusions and cell morphology at different doxycycline-inducible time points.

a + 1 μM C5a agonist
PH-Akt: 00:00 | 08:12 | 26:12

b + 1 μM C5a agonist
PIP5K1C: 00:00 | 07:45 | 24:45
PH-Akt

c
Normalized PH-Akt Response vs C5a Concentration (μM, Log-axis)
WT
PIP5K1C
ns **** **** ***

d + 1 nM folic acid
RBD: 0 | 65s | 85s

e + 1 nM folic acid
PI5K: 0 | 68s | 134s
RBD

f
Normalized RBD Response vs Folic Acid Concentration (nM, Log-axis)
- DOX
+ DOX
ns *** ** ****

g (Gβ-) - DOX
LimE: 00:00 | 02:20 | 04:54

h (Gβ-) + DOX overnight
PI5K: 00:23 | 03:10 | 09:07
LimE

Figure 5 Expressing PIP5Ks increases the threshold for STEN activation (a-b)
Representative live-cell time-lapse confocal images of responses of PH-Akt-mCherry to global
simulation C5a agonist at 1 µM in RAW 264.7 WT cells **(a)** or cells overexpressing PIP5K1C **(b).**
Time in min:sec format. Scale bars represent 5 mm. **(c)** Normalized PH-Akt responses (a drop of
cytosolic intensity) to different doses of C5a agonist for WT (blue) and PIP5K1C cells (magenta).
From the lowest to highest concentration, n_c=15 for each column from at least 3 independent
experiments; asterisks indicate significant difference, ****$P \leq 0.0001$, ***$P \leq 0.001$, $^{ns}P = 0.0408$,
(Mann-Whitney test. Compare ranks). The median is at the center, and whiskers and outliers are
graphed according to Tukey's convention (GraphPad Prism 10). **(d-e)** Representative live-cell
time-lapse confocal images of responses of RBD-GFP to global simulation folic acid at 1 nM in
Dictyostelium AX2 expressing doxycycline-inducible PI5K without DOX induction **(d)** or with
overnight DOX induction **(e)**. Time in sec format. Scale bars represent 5 mm. **(f)** Normalized RBD
responses (a drop of cytosolic intensity) to different doses of folic acid for WT (blue) and PIP5K1C
cells (magenta). From the lowest to highest concentration, n_c=15 for each column from at least 3
independent experiments; asterisks indicate significant difference, ****$P \leq 0.0001$, ***$P \leq 0.001$,
**$P \leq 0.01$, $^{ns}P = 0.2854$, (Mann-Whitney test. Compare ranks). The median is at the center, and
whiskers and outliers are graphed according to Tukey's convention (GraphPad Prism 10). **(g-h)**
Representative live-cell time-lapse confocal images of *Dictyostelium* Gb- cells co-expressing
RBD-GFP and doxycycline-inducible PI5K without DOX induction **(g)** or with overnight DOX
induction **(h).** Time in min:sec format. Scale bars represent 5 mm.

1602

1603

1604

1605

1606

1607

1608

1609

1610

1611

1612

1613

1614

1615

1616

a

CAR1-FKBP-FKBP   mCherry-FRB-MHCKC

+ Rapamycin

b

+DOX overnight

PI5K   06:23   16:36

+ Rapa

MHCKC   30:40   35:47

c

37min

+ Rapa

d

+ Rapa

Cell Number

-7   0   Time (min)   9   18

e

+ 50 μM Blebbistatin

RBD   00:00   06:33   13:20   17:24   21:42   26:45

PI5K

f

CIBN-CAAX

CRY2PHR-mCherry-Mypt169 (488 nm OFF)

CRY2PHR-mCherry-Mypt169 (488 nm ON)

488 nm ON

g

488 nm ON and + 10 μM Y27632

-17:10   00:00   25:30   54:30

-11:51   CRY2-Mypt169

PIP5K1B

h

Cell migration speed (μm/min)

****

- Rapa   + Rapa

i

Aspect Ratio

***

- Rapa   + Rapa

j

Aspect Ratio

**

488 nm OFF   488 nm ON

k

Cell Area (μm²)

****

488 nm OFF   488 nm ON

l

70 min

488 nm ON

Figure 6 Inhibiting myosin II activity counteracts PIP5K-induced phenotypes (a) Cartoon
illustrating mechanism of MHCKC global recruitment on *Dictyostelium* membrane with the help of
FKBP-FRB CID system in cells expressing doxycycline-inducible PI5K with overnight DOX
incubation. **(b)** Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 co-
expressing CAR1-FKBP-FKBP, mCherry-FRB-MHCKC, and doxycycline-inducible PI5K with
overnight DOX induction before and after 5 μM rapamycin treatment. Time in min:sec format.
Scale bars represent 5 mm. **(c)** Color-coded temporal overlay profile corresponds to (b).
(d) Heatmap quantification of cell migration speed at individual time points (time frame
1min/frame) of *Dictyostelium* AX2 co-expressing CAR1-FKBP-FKBP, mCherry-FRB-MHCKC, and
doxycycline-inducible PI5K with overnight DOX induction before and after 5 μM rapamycin
treatment. n_c=13 from at least 3 independent experiments. **(e)** Representative live-cell time-lapse
images of Dictyostelium cells coexpressing RBD-GFP and doxycycline-inducible PI5K with
overnight DOX induction during ventral wave propagation, before and after 50 μM blebbistatin
treatment. Time in min:sec format. Scale bars represent 5 mm. **(f)** Cartoon illustrating mechanism
of opto-Mypt169 global recruitment on differentiated HL-60 neutrophil membrane with the help of
CRY2-CIBN optogenetic system in cells expressing PIP5K1B. **(g)** Representative live-cell time-
lapse confocal images of differentiated HL-60 neutrophils expressing CIBN-CAAX, CRY2PHR-
mCherry-Mypt169 (magenta) and PIP5K1B-GFP (green), before or after 488 nm laser was
switched on globally. Time in min:sec format. Scale bars represent 5 μm. Cells are pretreated with
10 μM Y27632 for 10 mins. **(h-i)** Box-and-whisker plots of cell migration speed **(h)** or aspect ratio
(i) correspond to Figure 6b. n_c=15 (e) or n_c=12 (f) from at least 3 independent experiments;
asterisks indicate significant difference, ****$P \leq 0.0001$, ***$P = 0.0005$ (Mann-Whitney test.
Compare ranks). The median is at the center, and whiskers and outliers are graphed according
to Tukey's convention (GraphPad Prism 10). **(j-k)** Box-and-whisker plots of aspect ratio **(j)** or cell
area **(k)** correspond to Figure 6g. n_c=16 (g-h) from at least 3 independent experiments; asterisks
indicate significant difference, ****$P \leq 0.0001$, **$P = 0.0049$ (Mann-Whitney test. Compare
ranks). The median is at the center, and whiskers and outliers are graphed according to Tukey's
convention (GraphPad Prism 10). **(l)** Color-coded temporal overlay profile corresponds to Figure
6g.

1646

1647

1648

1649

1650

1651

1652

1653

1654

1655

1656

1657

Figure 7 Stochastic, reaction-diffusion model recreates experimental observations (a)
Schematic demonstrating the interactions modeled. **(b)** Kymographs of each of the six model
elements for a wild-type 300 second simulation. The red arrows in the three "rear" signals (PIP2,
PI5K and myosin) show the accumulation of these elements at the boundary between front and
back. The bottom right panel shows the sum over the complete perimeter of PIP5K as a function
of time. **(c-e)** Kymographs of simulations involving perturbations. In all cases, the simulations
started with WT parameters, and the perturbation was made after 150 s (marked by the white,
dotted lines). Specifically, PIP5K levels were lowered **(c)** or increased **(d)**; actin was eliminated
(e). As in b, all kymographs show 300 s of simulated time.

1667

1668

1669

1670

1671

1672

1673

1674

1675

1676

1677

1678

1679

1680

1681

1682

1683

1684

1685

1686

1687

1688

Figure 8 Schematic illustration showing the effects of PIP5Ks on symmetry-breaking and
cell morphology. Schematic demonstrating the interactions modeled. Each cell morphology's
front components and front localizations are highlighted in magenta, while back components and
back localizations are highlighted in green. At rest state, back components, such as PIP5K will be
uniformly distributed on plasma membrane in Lat A treated cell (a). Symmetry breaking will
happen at one spot on the membrane and PIP5K moves away from the spot (b). Upon removing
Lat A, the cytoskeletal activities will be restored and cell polarizes, becoming WT (c). Deleting
PIP5K in WT cells will induce fan-shaped cells (d). Expressing low level PIP5K will promote cell
polarity (e). Expressing high-level PIP5K will either induce fiopodias or completely shut down cells
(f). This phenotype can be reversed to (e) through lowering myosin II activities.

1699

1700

1701

1702

1703

1704

1705

1706

1707

1708

1709

1710

1711

1712

1713

1714

1715

1716

1717

a

```
Query  436  CCAAA--CATTGGATTCAGATGCACCAGAGGAAACATCAAATTTTAAATAAAAAATCTGAA  493
            ||||||  ||||||||||||||||||||||||||||||||||||||||||  |||||||||
Sbjct  521  CCAAACTCATTGGATTCAGATGCACCAGAGGAAACATCAAATTTAAATAAAAAAATCTGAA  580
```

b

c

AX2

PI5K-null

10 µm

d

Normalize Cell Number

Nuclei number

■ piki- 65h   ■ (piki-)PI5K 65h

e

6h   18h   24h

AX2

PI5K-null

f

00:00   01:10   04:05   05:43

PHcrac

g

02:30   03:20   09:10   10:25

LimE

h

00:00   00:55   02:24   04:04

CynA

1718

1719

1720

1721

1722

Supplementary Information

Supplementary Figures

Figure S1 *pi5k-* cells display cytokinesis and development defects. (a-b) Verification of CRISPR-mediated disruption of PI5K. **(a)** Genomic PCR sequence analysis confirmed mutations on the *PI5K* coding genes of *PI5K* mutant clone. Sequences of WT (Sbjct) and mutation clone (Query) are presented. 2 bp deletions are highlighted in the green box, and induced stop Condon TAA is highlighted with a red box. **(b)** Representative live-cell time-lapse confocal images (DIC) of CRISPR-mediated *pi5k-* fan-shaped (left), and *pi5k-* oscillatory cells (right). **(c)** Images of DIC channel and nuclear staining by Hoescht (merged) reveal a big increase in number of nuclei in each cell in *pi5k-* cells. **(d)** Histogram quantification of normalized cell number at different nuclei numbers for (*pi5k-*) PI5K and *pi5k-* after 65h in suspension. **(e)** WT (top) and *pi5k⁻* cells (bottom) were plated on development buffer (DB) medium agar for starvation at 6h, 18h, and 24h. WT cells aggregate normally, while *pi5k-* cells fail to aggregate. **(f-h)** Representative live-cell time-lapse confocal images of *Dictyostelium pi5k-* oscillatory cells expressing PHcrac-RFP (biosensor for PIP3) **(f)** or LimE-mCherry (biosensor for F-actin polymerization) **(g)** or CynA-GFP (biosensor for PI(3,4)P2) **(h).** Scale bars represent 5 mm.

1739

1740

1741

1742

1743

1744

1745

1746

1747

1748

1749

1750

1751

1752

1753

Figure S2 *pi5k-* cells display less Myosin and PI(3,4)P2 activities (a-b) Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 (WT) cells (left), and *pi5k-* cells (right) expressing PHcrac-RFP (biosensor for PIP3) **(a)** or LimE-mCherry (biosensor for F-actin polymerization) **(b)**. Scale bars represent 5 μm. **(c-d)** Box-and-whisker plots of **(c)** PHcrac patch length/Cell Perimeter, **(d)** LimE patch length/Cell Perimeter. n_c=22 from at least 3 independent experiments; asterisks indicate significant difference, ****P ≤ 0.0001 (Mann-Whitney test. Compare ranks). The median is at the center, and whiskers and outliers are graphed according to Tukey's convention (GraphPad Prism 10). **(e-f)** Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 (WT) cells (left), and *pi5k-* cells (right) expressing Myosin II-GFP **(e)** or CynA-GFP (biosensor for PI(3,4)P2) **(f)**. Scale bars represent 5 μm. **(g)** Box-and-whisker plot of aspect ratio corresponds to Figure 1o, 488 nm OFF or 488 nm ON. n_c=10 from at least 3 independent experiments; 'ns' indicates non-significant difference, ns denotes P>0.05 (Mann-Whitney test. Compare ranks). The median is at the center, and whiskers and outliers are graphed according to Tukey's convention (GraphPad Prism 10). **(h-i)** Representative membrane kymograph of LimE intensity in AX2(WT) cells **(h)** and *pi5k-* cells **(i)** respectively. A linear color map shows that blue has the lowest LimE or Myosin II intensity, whereas yellow has the highest. **(j-k)** Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 (WT) cells **(j),** or *pi5k-* cells **(k)** expressing Myosin II-GFP and LimE-mCherry (biosensor for F-actin polymerization). Cells show ventral wave propagation in the substrate-attached surface of the cell in **(k).** Scale bars represent 5 μm.

1774

1775

1776

1777

1778

1779

1780

1781

1782

1783

a

488 nm ON

00:21   03:37   06:11   08:38   13:46

CRY2-Inp54p

LifeAct

b

Cell Area (μm²)

****

488 nm OFF   488 nm ON

c

Cell migration speed (μm/min)

***

488 nm OFF   488 nm ON

d

Aspect Ratio

ns

488 nm OFF   488 nm ON

e

488 nm ON

00:00   03:23   04:05   04:54   07:07

CRY2-CTRL

LifeAct

f

488 nm OFF

Y-Distance (μm)

X-Distance (μm)

g

488 nm ON

Y-Distance (μm)

X-Distance (μm)

h

488 nm OFF

Y-Distance (μm)

X-Distance (μm)

i

488 nm ON

Y-Distance (μm)

X-Distance (μm)

j

488 nm ON

20 μm

60s

High

Low

k

0   6 min

488 nm ON

l

488 nm ON

20 μm

60s

High

Low

m

0   17 min

488 nm ON

1784

1785

1786

1787

1788

Figure S3 Globally recruiting Inp54p in neutrophil and macrophage induced fan-shaped phenotype (a) Time-lapse confocal images of differentiated HL-60 macrophage expressing CRY2PHR-mCherry-Inp54p (magenta; upper panel) and LifeAct-miRFP703 (cyan; lower panel), before or after 488 nm laser was switched on globally. Time in min:sec format. Scale bars represent 5 μm. **(b-d)** Box-and-whisker plots of **(b)** cell area, **(c)** cell migration speed, and **(d)** aspect ratio correspond to (a), 488 nm OFF or 488 nm ON. n_c=10 from at least 3 independent experiments; ****P ≤ 0.0001, 'ns' indicates non-significant difference, ns denotes P>0.05 (Mann-Whitney test. Compare ranks). The median is at the center, and whiskers and outliers are graphed according to Tukey's convention (GraphPad Prism 10). **(e)** Time-lapse confocal images of differentiated HL-60 macrophage expressing CIBN-CAAX, empty vector CRY2PHR-mCherry-CTRL (magenta; upper panel) and LifeAct-miRFP703 (cyan; lower panel), before or after 488 nm laser was switched on globally. Time in min:sec format. Scale bars represent 5 μm. **(f-i)** Centroid tracks of differentiated HL-60 neutrophils **(f-g)** or macrophage **(h-i)** (n_c=10) showing random motility at 488 nm OFF **(f, h)**, or 488 nm ON **(g, i)**. Each track lasted at least 5 minutes and was reset to the same origin. **(j-k)** Representative kymograph of cortical LifeAct intensity in Inp54p-expressing neutrophil **(j)** or macrophage **(k)** before or after 488 nm laser was turned on. A linear color map shows that blue has the lowest LifeAct intensity, whereas yellow has the highest. Duration of the kymograph is 5 mins. **(l-m)** color-coded temporal overlay profiles of differentiated HL-60 neutrophil **(l)** and macrophage **(m)** expressing CRY2PHR-mCherry-Inp54p. Square brackets indicate the range of recruitment.

1809

1810

1811

1812

1813

1814

1815

1816

1817

1818

1819

1820

1821

1822

1823

Figure S4 PI5K displays dynamic partitioning upon cAMP global stimulation
(a) Representative live-cell time-lapse images of Dictyostelium cells coexpressing PI5K-mCherry
and RBD-GFP during ventral wave propagation, showing PI5K dynamically localizes to the back-
state regions in ventral waves. Line-scan intensity profiles are shown in the bottommost panels.
Red line and green line represent PI5K and RBD, respectively. Time in min:sec format. Scale bars
represent 5 μm. **(b-e)** Representative live-cell images of *Dictyostelium* cells co-expressing PI5K-
GFP and PHcrac-RFP **(b)** or PHPLCδ–GFP and PHcrac-RFP **(d)** upon global cAMP stimulation,
demonstrating that upon transient global activation of cAR1 receptors, PHcrac gets uniformly
recruited to membrane whereas PI5K and PHPLCδ remained steadily membrane-bound
throughout the entire time course of the experiment. PHPLCδ had about 5% response. At time
t=53s or 78s, 1 µM (final concentration) cAMP was added. Time series plot of normalized cytosolic
intensities of PI5K and PHCrac **(c)** or PHPLCδ and PHcrac **(e),** showing the kinetics of the
response upon global stimulation with cAMP In all these figures, vertical dashed lines are used to
indicate the time of stimulation. Mean ± SEM are shown for n_c=18 cells. **(f)** Schematic
representation of PI5K and the derived truncations. **(g)** Representative live-cell time-lapse images
of Dictyostelium cells expressing PI5K-GFP (301-718aa) during migration showing PI5K (301-
718aa) dynamically localizes at the trailing edge in migrating cells. Time in min:sec format. Scale
bars represent 5 μm. **(h)** Time-lapse confocal images of *Dictyostelium* cells expressing
mRFPmars-SspBR73Q-PI5K(316-718aa), before or after 488 nm laser was switched on globally.
Time in sec format. Scale bars represent 5 µm. **(i)** Time-lapse confocal images of *Dictyostelium*
cells expressing mCherry-FRB-PI5K(1-315aa), before or after 5 µM Rapamycin was added. Time
in min:sec format. Scale bars represent 5 µm.

1846

1847

1848

1849

1850

1851

1852

1853

1854

1855

1856

1857

1858

1859

Figure S5 Expressing PI5K induces macropinocytosis defects (a) Representative live-cell images of *Dictyostelium* cells expressing doxycycline-inducible PI5K with overnight DOX induction (spiky). Time in sec format. Scale bars represent 5 µm. **(b-c)** Centroid tracks of cells (n_c=20) showing random motility in cells expressing doxycycline-inducible PI5K without DOX induction **(b)** or with overnight DOX induction **(c)**. Each track lasted at least 10 minutes and was reset to the same origin. **(d-f)** Box-and-whisker plots cell area **(d),** mean intensity **(e),** and aspect ratio **(f).** n_c=20 from at least 3 independent experiments; asterisks indicate significant difference, ****$P \le 0.0001$ (Mann-Whitney test. Compare ranks). The median is at the center, and whiskers and outliers are graphed according to Tukey's convention (GraphPad Prism 10). **(g)** Representative z-stack imaging showing the height of *Dictyostelium* cells expressing doxycycline-inducible PI5K without DOX induction (left) or with overnight DOX induction (right). Scale bars represent 5 µm. **(h-i)** Representative confocal images of Dictyostelium cells without DOX induction **(h)** or with overnight DOX induction **(i).** Cells were treated with FITC-dextran (green) for 10mins before imaging. The yellow outline corresponds cell area. Scale bars represent 5 µm. **(j)** Quantification of macropinocytosis uptake. n_c=58 from at least 3 independent experiments; asterisks indicate significant difference, ****$P \le 0.0001$ (Mann-Whitney test. Compare ranks). The median is at the center, and whiskers and outliers are graphed according to Tukey's convention (GraphPad Prism 10). **(k, r)** Representative live-cell images of *Dictyostelium* cells expressing doxycycline-inducible PI5K with 2h DOX induction **(k)** or PI5K (K681N, K682N) **(r).** Time in sec format. Scale bars represent 5 µm. **(l-n)** Color-coded temporal overlay profiles of *Dictyostelium* cells expressing doxycycline-inducible PI5K with overnight DOX induction (spiky) **(l),** or doxycycline-inducible PI5K with 2h DOX induction **(m),** or PI5K (K681N, K682N) **(n). (o)** Centroid tracks of cells (n_c=20) showing random motility in cells expressing doxycycline-inducible PI5K with 2h DOX induction. Each track lasted at least 10 minutes and was reset to the same origin. **(p)** Color-coded temporal overlay profiles of differentiated HL-60 neutrophils expressing PIP5K1C (rounded). **(q)** Centroid tracks of cells (n_c=15) showing random motility in differentiated HL-60 neutrophils expressing PIP5K1B. Each track lasted at least 10 minutes and was reset to the same origin.

1888

1889

1890

1891

1892

1893

1894

a PIP5K1C 00:00 07:17

b PIP5K1B 00:00 04:20 13:00

c PIP5K1C 00:00 08:45 16:46
16:00

d PIP5K1B 13 min 0

e PIP5K1C 19 min 0

f
Cell Area (µm²)
**** ****
WT PIP5K1B PIP5K1C

g HL60 WT
Y-Distance (µm)
X-Distance (µm)

h PIP5K1C
Y-Distance (µm)
X-Distance (µm)

i HL60 WT 17 min 0

j PIP5K1B 13 min 0

k WT +PMA 48h 20 min 0

l PIP5K1B +PMA 48h 21 min 0

m PIP5K1C +PMA 48h 9 min 0

n PIP5K1B
Mean Membrane Intensity
****
Polarized Round

o PIP5K1C
Mean Membrane Intensity
****
Polarized Round

p PIP5K1C +PMA 48h 00:00 03:58

q
488 nm ON

iLiD-CAAX (488 nm OFF)
iLiD-CAAX (488 nm ON)
Crismon-SspB R73Q-PIP5K1C

1895

1896

1897

1898

Figure S6 Expressing PIP5Ks impairs cell migration and induces cell contraction (a-c) Representative live-cell images of differentiated HL-60 neutrophils expressing PIP5K1C (rounded) **(a),** PIP5K1C (polarized) **(b)** , or PIP5K1C (polarized) **(c).** Time in min:sec format. Scale bars represent 5 µm. **(d-e)** Color-coded temporal overlay profiles of differentiated HL-60 neutrophils expressing PIP5K1B (polarized) **(d),** or PIP5K1C (polarized) **(e). (f)** Box-and-whisker plots of cell migration speed. n_c=20 from at least 3 independent experiments; asterisks indicate significant difference, ****P ≤ 0.0001 (Mann-Whitney test. Compare ranks). The median is at the center, and whiskers and outliers are graphed according to Tukey's convention (GraphPad Prism 10). **(g-h)** Centroid tracks of cells (n_c=15) showing random motility in differentiated HL-60 neutrophils WT **(g),** or expressing PIP5K1C **(h).** Each track lasted at least 10 minutes and was reset to the same origin. **(i-j)** Color-coded temporal overlay profiles of differentiated HL-60 neutrophils (WT) **(i),** or expressing PIP5K1B **(j). (k-m)** Color-coded temporal overlay profiles of differentiated HL-60 macrophages (WT) **(k),** or expressing PIP5K1B **(l),** or expressing PIP5K1C **(m). (n-o)** Box-and-whisker plots of mean membrane intensity for differentiated HL-60 neutrophils expressing PIP5K1B **(n)** or PIP5K1C **(o).** n_c=20 from at least 3 independent experiments; asterisks indicate significant difference, ****P ≤ 0.0001 (Mann-Whitney test. Compare ranks). The median is at the center, and whiskers and outliers are graphed according to Tukey's convention (GraphPad Prism 10). **(p)** Representative live-cell images of differentiated HL-60 macrophages expressing PIP5K1C. Time in min:sec format. Scale bars represent 5 µm. **(q)** Cartoon illustrating mechanism of PIP5K1C global recruitment on MDA-MB-231 cell membrane with the help of iLiD-SspB optogenetic system.

1920

1921

1922

1923

1924

1925

1926

1927

1928

1929

1930

1931

a 488 nm ON

-12: 21 | -06: 26 | 01: 00 | 12: 06 | 28: 53 | 42: 30

SspB-PIP5K1C

b 55 min / 488 nm ON / 0

c 0 — 40 min / 488 nm ON

d 0 / 488 nm ON / 32 min

e 488 nm ON

-05: 05 | 00: 10 | 13: 18 | 27: 45

SspB-CTRL

f ns

Cell Area (μm²)

488 nm OFF | 488 nm ON

g ns

Aspect Ratio

488 nm OFF | 488 nm ON

1932

1933

1934

1935

Figure S7 Globally recruiting empty vector in MDA-MB-231 cells do not induce cell contraction (a) Time-lapse confocal images of MDA-MB-231 cells expressing crimson-SspB-PIP5K1C-P2A-iLiD-CAAX, before or after 488 nm laser was switched on globally. Time in min:sec format. Scale bars represent 10 μm. Blue arrows indicate where retraction fibers or blebs are formed. Cells are pretreated with 10 ng/ml EGF for 10 mins**. (b-d)** Color-coded temporal overlay profile corresponds to Figure S7a, 3k, and S7e. (e) Time-lapse confocal images of MDA-MB-231 cells expressing empty vector crimson-SspB-MCS-P2A-iLiD-CAAX, before or after 488 nm laser was switched on globally. Time in min:sec format. Scale bars represent 10 μm. **(f-g)** Box-and-whisker plots of **(f)** cell area, **(g)** aspect ratio correspond to (m-n). n_c=10 from at least 3 independent experiments; asterisks indicate significant difference, 'ns' indicates non-significant difference, ns denotes $P>0.05$ (Mann-Whitney test. Compare ranks). The median is at the center, and whiskers and outliers are graphed according to Tukey's convention (GraphPad Prism 10).

1948

1949

1950

1951

1952

1953

1954

1955

1956

1957

1958

1959

1960

1961

1962

1963

1964

1965

1966

1967

1968

Figure S8 Expressing PI5K can induce spiky filopods and suppresses signal transduction and cytoskeletal activities (a-b) Representative membrane kymograph of RBD intensity in *Dictyostelium* AX2 expressing doxycycline-inducible PI5K without DOX induction **(a)** or with overnight DOX induction (spiky) **(b).** A linear color map shows that blue has the lowest RBD intensity, whereas yellow has the highest. **(c-d)** Representative membrane kymograph of PHcrac intensity in *Dictyostelium* AX2 expressing doxycycline-inducible PI5K without DOX induction **(c)** or with overnight DOX induction (spiky) **(d).** A linear color map shows that blue has the lowest RBD intensity, whereas yellow has the highest. **(e-f)** Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 co-expressing doxycycline-inducible PI5K with overnight DOX induction (spiky) and RBD-GFP (biosensor for activated Ras) **(e)** or PHcrac-YFP (biosensor for PIP3) **(f)**. Time in sec format. Scale bars represent 5 μm. **(g-i)** Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 co-expressing LimE-mCherry (biosensor for F-actin) and doxycycline-inducible PI5K without DOX induction **(g)** or with overnight DOX induction (rounded) **(h),** or with overnight DOX induction (spiky) **(i).** Time in sec format. Scale bars represent 5 μm.

1983

1984

1985

1986

1987

1988

1989

1990

1991

1992

1993

1994

1995

1996

1997

1998

1999

2000

2001

2002

Figure S9 Expressing PI5K inhibits Rac1/Arp2/3 complex/F-actin axis (a) Box-and-whisker plots of LimE patch size (left axis) and LimE patch number (right axis). n_c=20 from at least 3 independent experiments; asterisks indicate significant difference, ****P ≤ 0.0001 (Mann-Whitney test. Compare ranks). The median is at the center, and whiskers and outliers are graphed according to Tukey's convention (GraphPad Prism 10). **(b-c)** *t*-stacks from a cell co-expressing LimE-mCherry and doxycycline-inducible PI5K without DOX induction **(b)** or with overnight DOX induction (rounded) **(c).** The white arrow corresponds to the time duration of the *t*-stack kymograph. **(d-f)** Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 co-expressing Pak1-GFP (biosensor for Rac1) and doxycycline-inducible PI5K without DOX induction **(d)** or with overnight DOX induction (rounded) **(e),** or with overnight DOX induction (spiky) **(f).** Time in min:sec format. Scale bars represent 5 μm. **(g)** Box-and-whisker plots of Pak1 patch size (left axis) and Pak1 patch number (right axis). n_c=20 from at least 3 independent experiments; asterisks indicate significant difference, ****P ≤ 0.0001 (Mann-Whitney test. Compare ranks). The median is at the center, and whiskers and outliers are graphed according to Tukey's convention (GraphPad Prism 10). **(h-i)** Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 co-expressing ArpC-GFP and doxycycline-inducible PI5K without DOX induction **(h)** or with overnight DOX induction **(i).** Time in min:sec format. Scale bars represent 5 μm. **(j-k)** Representative live-cell time-lapse confocal images of *Dictyostelium* abnABC- cells co-expressing RBD-GFP and doxycycline-inducible PI5K without DOX induction **(j)** or with overnight DOX induction **(k).** Time in min:sec format. Scale bars represent 5 μm.

2023

2024

2025

2026

2027

2028

2029

2030

2031

2032

2033

2034

2035

2036

2037

2038

Figure S10 Expressing less PI5K increases cell polarity but inhibits signal transduction
activities (a-b) Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 co-
expressing RBD-GFP **(a)** or PHcrac-YFP **(b)** and doxycycline-inducible PI5K with 2h DOX
induction. Time in min:sec format. Scale bars represent 5 μm. **(c-d)** Representative live-cell time-
lapse confocal images of *Dictyostelium* AX2 co-expressing RBD-RFP **(c)** or LimE-RFP **(d)** and
PI5K (K681N, K682N). Time in min:sec format. Scale bars represent 5 μm. **(e-f)** Representative
membrane kymograph of RBD intensity in *Dictyostelium* AX2 expressing doxycycline-inducible
PI5K without DOX induction **(e)** or with overnight DOX induction **(f),** even in the absence of actin
cytoskeleton. Cells were pre-treated with actin polymerization inhibitor Latrunculin A (final
concentration 5μM) and caffeine (final concentration 4mM) for 20min. A linear color map shows
that blue has the lowest RBD intensity, whereas yellow has the highest. **(g-h)** Representative
membrane kymograph of PHcrac intensity in *Dictyostelium* AX2 expressing doxycycline-inducible
PI5K without DOX induction **(g)** or with overnight DOX induction **(h),** even in the absence of actin
cytoskeleton. Cells were pre-treated with actin polymerization inhibitor Latrunculin A (final
concentration 5μM) and caffeine (final concentration 4mM) for 20min. A linear color map shows
that blue has the lowest PHcrac intensity, whereas yellow has the highest.

2055

2056

2057

2058

2059

2060

2061

2062

2063

2064

2065

2066

2067

2068

2069

2070

a

C2GAPB

PI5K

00: 00 08: 12 16: 50

b

+ C2GAPB

17 min

c

- C2GAPB   + C2GAPB

14 min   14 min

0   0

d

- C2GAPB

Y-Distance (µm)

X-Distance (µm)

+ C2GAPB

Y-Distance (µm)

X-Distance (µm)

e

Cell migration speed (µm/min)

****

- DOX + DOX

2071

2072

2073

2074 **Figure S11 Expressing C2GAPB in *pi5k-* cell increases cell polarity and migration (a)**
2075 Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 co-expressing
2076 doxycycline-inducible PI5K C2GAPB and doxycycline-inducible PI5K with overnight DOX
2077 induction. Time in min:sec format. Scale bars represent 5 μm. **(b)** Color-coded temporal overlay
2078 profile of the cell corresponds to **(a). (c)** Color-coded temporal overlay profiles of *Dictyostelium*
2079 *pi5k-* cells expressing doxycycline-inducible C2GAPB without DOX induction (left) or with
2080 overnight DOX induction (right). **(d)** Centroid tracks of cells (n_c=16) showing random motility in
2081 *pi5k-* cells expressing doxycycline-inducible C2GAPB without DOX induction (left) or with
2082 overnight DOX induction (right). Each track lasted at least 10 minutes and was reset to the same
2083 origin. **(e)** Box-and-whisker plot of cell migration speed corresponds to (c). n_c=16 from at least 3
2084 independent experiments; asterisks indicate significant difference, ****$P \leq 0.0001$ (Mann-Whitney
2085 test. Compare ranks). The median is at the center, and whiskers and outliers are graphed
2086 according to Tukey's convention (GraphPad Prism 10).

2087

2088

2089

2090

2091

2092

2093

2094

2095

2096

2097

2098

2099

2100

2101

2102

2103

2104

2105

2106

2107

Figure S12 Expressing PIP5Ks in RAW 264.7 cells increases the threshold for PI3K activation (a-h) Representative live-cell time-lapse confocal images of responses of PH-Akt-mCherry to global simulation C5aR agonist at 0.1 µM in RAW 264.7 WT cells **(a)** or cells overexpressing PIP5K1C **(b)**; or at 1 µM in RAW 264.7 WT cells **(c)** or cells overexpressing PIP5K1C **(d);** or at 10 µM in RAW 264.7 WT cells **(e)** or cells overexpressing PIP5K1C **(f);** or at 100 µM in RAW 264.7 WT cells **(g)** or cells overexpressing PIP5K1C **(h).** Time in min:sec format. Scale bars represent 5 µm.

2115

2116

2117

2118

2119

2120

2121

2122

2123

2124

2125

2126

2127

2128

2129

2130

2131

2132

2133

2134

2135

a

b

c +100 μM C5a   +100 μM C5a

PH-Akt

PIP5K1C

d +100 nM FA   +100 nM FA

RBD

PI5K + DOX

e

Normalized Cell Number

C5a concentration (μM), Log-axis

■ < 0.05 (PIP5K1C) ■ > 0.2 (PIP5K1C) ■ < 0.05 (control) ■ > 0.2 (control)

f

Normalized Cell Number

FA concentration (nM), Log-axis

■ < 0.05 (+ DOX) ■ > 0.2 (+ DOX) ■ < 0.05 (- DOX) ■ > 0.2 (- DOX)

g

PI5K -DOX

PI5K +DOX overnight
polarized shape

PI5K +DOX overnight
rounder shape

2136

2137

2138

Figure S13 Expressing PI5K induces chemotaxis defects (a-b) Line-scan intensity profiles correspond to Figure 5d-e. **(c)** Representative live-cell time-lapse confocal images of 5% (left) or 20% (right) responses of PH-Akt-mCherry to global simulation C5a agonist at 100 μM in RAW 264.7 cells overexpressing PIP5K1C. **(d)** Representative live-cell time-lapse confocal images of 5% (left) or 20% (right) responses of RBD-GFP to global simulation folic acid (FA) at 100 nM in *Dictyostelium* AX2 expressing doxycycline-inducible PI5K with overnight DOX induction. **(e)** Histogram quantification of normalized cell number at different doses of C5a agonist in RAW 264.7 WT cells or cells overexpressing PIP5K1C**.** Gray columns represent PIP5K1C cells that have < 5% responses. Yellow columns represent PIP5K1C cells that have > 20% responses. Blue columns represent WT cells that have < 5% responses. Orange columns represent WT cells that have > 20% responses. **(f)** Histogram quantification of normalized cell number at different doses of folic acid (FA) in in *Dictyostelium* AX2 expressing doxycycline-inducible PI5K without DOX induction (- DOX) or with overnight DOX induction (+ DOX). Green columns represent + DOX cells that have < 5% responses. Cyan columns represent + DOX cells that have > 20% responses. Dark blue columns represent - DOX cells that have < 5% responses. Orange columns represent - DOX cells that have > 20% responses. **(g)** Color-coded temporal overlay profiles of vegetative *Dictyostelium* AX2 expressing doxycycline-inducible PI5K without DOX (left), or polarized cells with overnight DOX induction (middle), or rounded cells with overnight DOX induction (right), chemotaxing to 10 μM folic acid. The green box is where the center of the chemoattractant source.

2158

2159

2160

2161

2162

2163

2164

2165

2166

2167

2168

2169

2170

2171

a

PI5K -DOX

PI5K +DOX overnight

b

c

2172

2173

2174

Figure S14 Quantifications of directed migration speed with or without PI5K expression (a)
Centroid tracks of cells (n_c=13) showing chemotaxis motility in cells expressing doxycycline-inducible PI5K without DOX induction (left) or with overnight DOX induction (right). Each track lasted 30 minutes, and the center of the chemoattractant source was reset to origin. **(b-c)** Scatter dot plots of directed cell migration speed (b) and aspect ratio (c) corresponds to Figure S12e. n_c=18 from at least 2 independent experiments; asterisks indicate significant difference, ****P ≤ 0.0001, ***P ≤ 0.001 (Mann-Whitney test. Compare ranks). The median is at the center, and whiskers and outliers are graphed according to Tukey's convention (GraphPad Prism 10).

2183

2184

2185

2186

2187

2188

2189

2190

2191

2192

2193

2194

2195

2196

2197

2198

2199

2200

2201

2202

2203

2204

Figure S15 Globally recruiting Mypt169 in neutrophil and macrophage induces cell polarity
and protrusive activities (a-b) Line-scan intensity profiles correspond to Figure 6b time point
30: 40 **(a)** or 35: 47 **(b). (c-d)** Centroid tracks of cells correspond to Figure 6b (n_c=15), showing
random motility before **(c)** and after **(d)** 5 µM rapamycin treatment. Each track lasted at least 15
minutes and was reset to the same origin. **(e)** Representative live-cell time-lapse confocal images
of differentiated HL-60 macrophage expressing CIBN-CAAX, CRY2PHR-mCherry-Mypt169
(magenta) and LifeAct-miRFP703 (Cyan), before or after 488 nm laser was switched on globally.
Blue arrows indicate where protrusions are formed. Time in min:sec format. Scale bars represent
5 µm. **(f)** Color-coded temporal overlay profile corresponds to (j). **(g)** Representative live-cell time-
lapse confocal images of RAW 264.7 cells expressing CIBN-CAAX, CRY2PHR-mCherry-Mypt169
(magenta) and PIP5K1C-GFP (green), before or after 488 nm laser was switched on globally.
Blue arrows indicate where protrusions are formed. Time in min:sec format. Scale bars represent
5 µm. Cells are pretreated with 10 µM C5a agonist for 10 mins. **(h)** Color-coded temporal overlay
profile corresponds to (l).

2219

2220

2221

2222

2223

2224

2225

2226

2227

2228

2229

2230

2231

2232

2233

2234

2235

2236

2237

2238

Figure S16 Expressing PI5K induces a shift to cortical actin at the cell rear (a-b) Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 co-expressing ABD120-GFP (green), LimE-Halo (magenta), and doxycycline-inducible PI5K without DOX induction **(l)** or with overnight DOX induction **(m).** Time in sec format. Scale bars represent 5 μm. **(c-d)** Phalloidin stain of *Dictyostelium* AX2 doxycycline-inducible PI5K without DOX induction **(c)** or with overnight DOX induction **(d). (e)** Box-and-whisker plot of ABD120-LimE ratio corresponds to (a-b). n_c=17 from at least 3 independent experiments; asterisks indicate significant difference, ****$P ≤ 0.0001$ (Mann-Whitney test. Compare ranks). The median is at the center, and whiskers and outliers are graphed according to Tukey's convention (GraphPad Prism 10). **(f)** Representative live-cell time-lapse images of Dictyostelium cells coexpressing LimE-mCherry and doxycycline-inducible PI5K with overnight DOX induction, before and after 100 μM CK666 treatment. Time in min:sec format. Scale bars represent 5 μm.

2251

2252

2253

2254

2255

2256

2257

2258

2259

2260

2261

2262

2263

2264

2265

2266

2267

2268

2269

2270

2271

Figure S17 Schematic illustration showing the effect of PIP5Ks on cell morphology, signal transduction, and cytoskeletal dynamics. Cells display different morphology at different PIP5Ks expression levels as shown on the left of this figure. The front region of the cell at each cell morphology is shown in yellow, while the back region of the cell is shown in green. The blue box on each cell represents the zoom-in region on the right side of the figure. The yellow or green-shaded lipid head groups at each zoomed-in box represent the inner leaflet membrane. The headgroups of the inner leaflet lipid molecules that are enriched in front-state are shown in yellow, while the headgroups that are enriched in back-state are shown in green. Actin or actomyosin structures are shown at the front-state or bac-state of the cell, respectively. The black box on each lipid bilayer represents the zoom-in region of the lipid bilayer and the signaling pathways within this region. In each condition, the lighter color icons represent the depleted signaling components. Green arrow represents the increased change of this signaling molecular, while orange arrow represents the decreased change.

2285

2286

2287

2288

2289

2290

2291

2292

2293

2294

2295

2296

2297

2298

2299

2300

2301

2302

2303

2304

Supplementary Video Legends

2305

Video S1

Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 (WT) cells and *pi5k-*
cells expressing PHPLCδ-YFP (biosensor for PI(4,5)P2), CynA-GFP (biosensor for PI(3,4)P2),
and mhcA-GFP. Top left corner shows time in min:sec format. Scale bar represents 10 μm.

Video S2

Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 (WT) cells and *pi5k-*
cells expressing RBD-GFP (biosensor for activated Ras), PHcrac-RFP (biosensor for PIP3), and
LimE-mCherry (biosensor for actin polymerization). Top left corner shows time in min:sec format.
Scale bar represents 10 μm.

Video S3

Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 (WT) cells and *pi5k-*
cells expressing mhcA-GFP (green) and LimE-mCherry (magenta). In *pi5k-* cells, ventral wave
activities of mhcA and LimE can be observed. Top left corner shows time in min:sec format. Scale
bar represents 10 μm in *pi5k-* cells and 5 μm in AX2 cells, respectively.

Video S4

Time-lapse confocal images of differentiated HL-60 neutrophil and macrophage expressing CIBN-
CAAX, CRY2PHR-mCherry-Inp54p (magenta) and LifeAct-miRFP703 (cyan), or differentiated
HL60 macrophage expressing untagged CIBN-CAAX, CRY2PHR-mCherry-empty vector
(magenta) and LifeAct-miRFP703 (cyan), before or after 488 nm laser was switched on globally.
Top left corner shows time in min:sec format. To start recruitment (magenta), the laser was
switched on at '03:09', or '03:51', or '03:30', once '488 nm ON' appears at the top of the video.
Cell was not exposed to chemoattractant during the experiment. Scale bar represents 5 μm.

Video S5

Representative live-cell time-lapse images of Dictyostelium cells coexpressing PI5K-GFP and
PHcrac–RFP (biosensor for PIP3), differentiated HL-60 neutrophil expressing PIP5K1B, or
PIP5K1C and LifeAct (Cyna) during migration showing PI5K dynamically moves away from
protrusions in migrating cells. Top left corner shows time in min:sec format. Scale bars represent
5 mm.

Video S6

Representative live-cell time-lapse images of Dictyostelium cells coexpressing PI5K-GFP and
PHcrac-RFP (biosensor for PIP3), PI5K-mCherry and RBD-GFP (biosensor for activated Ras),
PI5K-mCherry and PHPLCδ–GFP (biosensor for PI(4,5)P2), differentiated HL-60 macrophage
expressing PIP5K1B and LifeAct, and Dictyostelium cells coexpressing PI5K-mRFP and RBD-
GFP (biosensor for activated Ras) upon Latrunculin A treatment, during ventral wave propagation,
showing PI5K dynamically localizes to the back-state regions in ventral waves. Top left corner
shows time in min:sec format. Scale bars represent 5 mm.

Video S7

Representative live-cell images of *Dictyostelium* cells co-expressing PI5K-GFP and PHcrac-RFP,
or PHPLCδ–GFP and PHcrac-RFP upon global cAMP stimulation. Top left corner shows time in
min:sec format. To start global stimulation, cAMP was added at '00:53', or '01:17', once '+ cAMP'
appears at the top of the video. Scale bar represents 5 μm.

Video S8

Representative live-cell images of *Dictyostelium* cells expressing doxycycline inducible KikGR-
PI5K with overnight DOX incubation. Top left corner shows time in min:sec format. Photo
conversion happened at '01:50', '05:30', or '11:34'. Scale bar represents 5 μm.

Video S9

Representative live-cell time-lapse images of Dictyostelium cells expressing PI5K-GFP (301-
718aa) during migration showing PI5K (301-718aa) dynamically localizes at the trailing edge in
migrating cells, or expressing mRFPmars-SspBR73Q-PI5K(316-718aa), before or after 488 nm
laser was switched on globally, or expressing mCherry-FRB-PI5K(1-315aa), before or after 5 μM
Rapamycin was added. Top left corner shows time in min:sec format. To start recruitment, the
laser was switched on at '01:20', once '488 nm ON' appears at the top of the video, or Rapamycin
was added at '12:51', once '+ Rapamycin' appears at the top of the video. Scale bars represent
5 μm.

Video S10

Representative live-cell images of *Dictyostelium* cells expressing doxycycline-inducible PI5K
without DOX induction, or with 2h DOX incubation, or with overnight DOX induction (rounded), or
with overnight DOX induction (spiky), or expressing PI5K(K681N, K682N). Top left corner shows
time in min:sec format. Scale bars represent 5 mm for first 3 videos and 10 mm for last video.

Video S11

Representative live-cell images of differentiated HL-60 neutrophils (WT) expressing LifeAct as the
biosensor, or expressing PIP5K1B (rounded), or expressing PIP5K1B (polarized), or expressing
PIP5K1C (rounded), or expressing PIP5K1C (polarized). Top left corner shows time in min:sec
format. Scale bars represent 5 μm.

Video S12

Representative live-cell images of differentiated HL-60 macrophage (WT) expressing PH-Akt as
the biosensor, or expressing PIP5K1B (rounded), or expressing PIP5K1C (rounded). Top left
corner shows time in min:sec format. Scale bars represent 5 μm.

Video S13

Time-lapse confocal images of MDA-MB-231 cells expressing crimson-SspB-PIP5K1C-P2A-iLiD-
CAAX or expressing crimson-SspB-empty vector-P2A-iLiD-CAAX, before or after 488 nm laser
was switched on globally. Top left corner shows time in hour:min:sec or min:sec format. To start
recruitment, the laser was switched on at '00:12:21', or '09:44', or '04:55', once '488 nm ON'
appears at the top of the video. Scale bar represents 10 μm.

Video S14

Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 co-expressing RBD-
GFP (biosensor for activated Ras) or PHcrac-YFP (biosensor for PIP3) and doxycycline-inducible
PI5K without DOX induction, or with 2h DOX incubation, or with overnight DOX induction

(rounded), or with overnight DOX induction (spiky). Top left corner shows time in min:sec format.
Scale bars represent 5 µm.

Video S15

Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 co-expressing RBD-
GFP (biosensor for activated Ras) and doxycycline-inducible PI5K without DOX induction, or with
overnight DOX induction during ventral wave propagation. Top left corner shows time in min:sec
format. Scale bars represent 10 µm.

Video S16

Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 co-expressing Pak1-
GFP (biosensor for Rac1), or ArpC-GFP, or LimE-mCherry and doxycycline-inducible PI5K
without DOX induction, or with overnight DOX induction (rounded) or with overnight DOX
induction (spiky). Top left corner shows time in min:sec format. Scale bars represent 5 µm.

Video S17

Representative live-cell time-lapse confocal images of *Dictyostelium abnABC-* cells co-
expressing RBD-GFP and doxycycline-inducible PI5K without DOX induction, or with overnight
DOX induction. Top left corner shows time in min:sec format. Scale bars represent 5 µm.

Video S18

Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 co-expressing RBD-
RFP, or LimE-RFP and PI5K (K681N, K682N). Top left corner shows time in min:sec format. Scale
bars represent 5 µm.

Video S19

Representative live-cell time-lapse confocal images of Dictyostelium *pi5k-* cells expressing
doxycycline-inducible PI5K C2GAPB without DOX induction, or with overnight DOX induction, or
Dictyostelium AX2 co-expressing doxycycline-inducible PI5K C2GAPB and doxycycline-inducible
PI5K with overnight DOX induction. Top left corner shows time in min:sec format. Scale bars
represent 10 µm for first 2 videos, and 5 µm for last video.

Video S20

Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 co-expressing RBD-
GFP (biosensor for activated Ras), or PHcrac-YFP (biosensor for PIP3) and doxycycline-inducible
PI5K without DOX induction, or with overnight DOX induction upon Latrunculin A treatment. Top
left corner shows time in min:sec format. Scale bars represent 5 µm.

Video S21

Representative live-cell time-lapse confocal images of responses of PH-Akt-mCherry to global
simulation C5aR agonist in RAW 264.7 WT cells, or cells overexpressing PIP5K1C at at 0.1 µM-
100 µM. To start global stimulation, C5a agonist was added at '00:53', or '06:09', or '05:45', or
'06:40', or '16:42', or '03:42', or '06:21', or '05:55', or '12:52', once '+ C5a' appears at the top of
the video. Scale bar represents 5 µm.

Video S22

Representative live-cell time-lapse confocal images of *Dictyostelium Gb-* cells co-expressing
RBD-GFP and doxycycline-inducible PI5K without DOX induction, or with overnight DOX
induction. Top left corner shows time in min:sec format. Scale bars represent 5 µm.

Video S23

Representative live-cell time-lapse confocal images of vegetative *Dictyostelium* AX2 expressing
doxycycline-inducible PI5K without DOX, or with overnight DOX induction, chemotaxing to 10 mM
folic acid-filled micropipette. The white box is where the center of the chemoattractant source. Top
left corner shows time in min:sec format. Scale bars represent 20 µm.

Video S24

Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 co-expressing CAR1-
FKBP-FKBP, mCherry-FRB-MHCKC, and doxycycline-inducible PI5K with overnight DOX
induction before and after 5 µM rapamycin treatment. Top left corner shows time in min:sec format.
To start recruitment, Rapamycin was added at '14:03', once '+ Rapamycin' appears at the top of
the video. Scale bars represent 10 µm.

Videos S25

(e) Representative live-cell time-lapse images of Dictyostelium cells coexpressing RBD-GFP and
doxycycline-inducible PI5K with overnight DOX induction during ventral wave propagation, before
and after 50 µM blebbistatin treatment. Top left corner shows time in min:sec format. Blebbistatin
was added at '06:33', once '+ Blebbistatin' appears at the top of the video. Scale bars represent
10 µm.

Videos S26

Representative live-cell time-lapse confocal images of differentiated HL-60 macrophage
expressing untagged CIBN-CAAX, CRY2PHR-mCherry-Mypt169 (magenta), or differentiated HL-
60 neutrophil and macrophage expressing untagged CIBN-CAAX, CRY2PHR-mCherry-Mypt169
(magenta), and PIP5K1B-GFP (green), before or after 488 nm laser was switched on globally.
Top left corner shows time in min:sec or hour:min:sec format. To start recruitment (magenta), the
laser was switched on at '03:41', or '00:05:55', or '02:51', once '488 nm ON' appears at the top of
the video. Cell was not exposed to chemoattractant during the experiment. Scale bar represents
10 µm for the frist video, and 5 µm for the rest two videos.

Video S27

Representative live-cell time-lapse confocal images of *Dictyostelium* AX2 co-expressing ABD120-
GFP (green), LimE-Halo (magenta), and doxycycline-inducible PI5K without DOX induction, or
with overnight DOX induction. Top left corner shows time in min:sec format. Scale bars represent
5 µm.

Video S28

Representative live-cell time-lapse images of Dictyostelium cells coexpressing LimE-mCherry
and doxycycline-inducible PI5K with overnight DOX induction, before and after 100 µM CK666
treatment. Top left corner shows time in min:sec format. CK666 was added at '12:23', once '+
CK666' appears at the top of the video. Scale bars represent 5 µm.

2461

2462

2463

2464