Title

Genome divergence between European anchovy ecotypes fuelled by structural

variants originating from cross-hemispheric admixture

Authors

Laura Meyer[1]*, Pierre Barry[2,3], Alan Le Moan[4], Christine Arbiol[1], Rita Castilho[5], Carl Van der
Lingen[6,7], Malika Chlaïda[8], Niall J. McKeown[9], Bruno Ernande[10], François Bonhomme[1]**, Pierre-
Alexandre Gagnaire[1]**, Bruno Guinand[1]**
*Corresponding author: laura.meyer@umontpellier.fr
**Co-last authors

Affiliations:

[1] ISEM, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France
[2] CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO Laboratório
Associado, Campus de Vairão, Universidade do Porto, Vairão, Portugal
[3] BIOPOLIS Program in Genomics, Biodiversity and Land Planning, CIBIO, Campus de Vairão,
Vairão, Portugal
[4] CNRS-Sorbonne Université, Station Biologique de Roscoff, Roscoff, France
[5] CCMAR, Centro de Ciências do Mar, Universidade do Algarve, Campus de Gambelas, Faro,
Portugal
[6] Marine and Coastal Management, Department of Forestry, Fisheries and the Environment, Cape
Town, South Africa
[7] University of the Western Cape, Cape Town, South Africa
[8] Research and Development Unit on Marine Biology, National Institute of Fisheries Research,
Casablanca, Morocco
[9] Aberystwyth University, Department of Life Sciences, Ceredigion, Wales, UK
[10] MARBEC, Univ. Montpellier, Ifremer, CNRS, IRD, Montpellier, France

Abstract

The formation of ecotypes is shaped by mechanisms that reduce gene flow through complex

interactions between ecological, historical, and genomic factors. In the European anchovy (*Engraulis*

encrasicolus), marine and coastal ecotypes have been identified in the North-East Atlantic and

Mediterranean Sea, yet the genomic basis of their divergence remains unclear. Here, we present the

first genome-scale analysis of this species complex, integrating whole-genome sequencing (WGS)

and RAD-seq data from populations across its distribution range. In addition to the marine and

coastal ecotypes, we identify a previously undetected lineage which is present in southern Morocco,

the Canary Islands and even in South Africa. This southern Atlantic lineage exhibits a gradient of

admixture with northern populations near the Atlantic-Mediterranean transition zone. Genomic

differentiation landscapes reveal large regions of high linkage disequilibrium, likely corresponding to

thirteen structural variants (SVs) segregating within or between the lineages. Notably, three of the six

SVs contributing to the gene flow barrier between northern ecotypes originated in the southern

lineage, supporting a partially shared evolutionary history between the coastal ecotype and the

southern lineage. Our findings suggest that anchovy ecotype divergence has been shaped by a

combination of ancient structural variation, admixture, and local adaptation. This study highlights how

SVs that arose between geographically isolated lineages can act as key genetic elements in ecotype

formation, reinforcing reproductive isolation through distinct evolutionary pathways.

44 Introduction

Speciation is a complex, multi-level process shaped by temporal, spatial, ecological and genome

architectural factors (Abbott et al., 2013). These dimensions collectively influence a variety of

reproductive isolation mechanisms, which establish and accumulate barriers to gene flow,

strengthening reproductive isolation during the course of speciation (Kulmuni et al., 2020; Schluter &

Rieseberg, 2022). Although the multifaceted nature of speciation is widely recognized, many

comprehensive studies have focused on systems where divergence appears to be primarily driven by

one or a few dominant factors within a well-defined eco-evolutionary context. Ecotype formation is

one such example, where genomic variation associated with distinct ecological habitats has

traditionally been used to investigate ecologically driven speciation. In these cases, speciation is

often assumed to be rapidly initiated by local adaptation, provided divergent selection arises from

habitat variation and sufficient genetic variation is available for selection to act upon (Rundle & Nosil,

2005; Schluter, 2000; Schluter & Conte, 2009). However, recent advances in speciation genomics

present a more nuanced perspective, highlighting growing evidence that ecotype formation results

from the interplay of multiple factors (Van Belleghem et al., 2018; Marques et al., 2019).

A key finding from these genomic investigations relates to the temporal dimension of ecotype

formation, which reveals that ecotype divergence is often associated with ancient allelic variation.

Even when ecotype formation was initially attributed to the recent establishment of new habitats,

divergence times are frequently found to exceed those derived from paleoenvironmental

reconstructions or from coalescent time distributions for sequences from a panmictic population (Han

et al., 2017; Le Moan et al., 2021; Louis et al., 2020; Nelson & Cresko, 2018; Small et al., 2023). This

discrepancy between ecological and genetic divergence time frames (Van Belleghem et al., 2018)

can be explained by several mechanisms, including long-term balancing selection on the variants

underlying ecotype divergence, ancestral population structure or admixture between divergent

lineages.

Another recurrent finding is that genetic differences contributing to ecotype formation are

concentrated in genomic regions with low recombination rates, rather than being spread across the

genome. These regions often comprise structural variants (SVs) such as chromosomal inversions,

which appear to play a disproportionate role in ecotype divergence (Faria et al., 2019; Hager et al.,

2022; Lundberg et al., 2023; Todesco et al., 2020). Recent evidence suggests that SVs may form a

substantial barrier to gene flow, ensuring at least moderate reproductive isolation (RI) between

lineages, particularly when multiple SVs separate the lineages and when these variants are coupled

(Le Moan et al., 2024). Since recombination is largely suppressed in SVs such as inversions,

divergence accumulates more readily between haplotypes, which are protected from the

homogenizing effects of gene flow (Berdan et al., 2024; Rieseberg, 2001). While SVs are often

implicated in RI between ecotypes through their role in local adaptation by capturing multiple loci

under environmental selection (Kirkpatrick & Barton, 2006; Wellenreuther & Bernatchez, 2018), they

may also harbor coadapted gene complexes or Dobzhansky-Muller incompatibilities that contribute to

the isolation of nascent lineages carrying different haplotypes (Navarro & Barton, 2003).

Understanding the origin of the genetic variation that underlies ecotype divergence can be a first step

towards reconciling the different components of ecotype formation. This can aid in identifying the

most plausible evolutionary scenarios to explain how ecotypes have formed. One scenario that has

been proposed to account for both the temporal and ecological components of ecotype formation, is

admixture between geographical lineages. Admixture is not merely a homogenizing process; it has

also been linked to evolutionary diversification that may lead to ecotype formation through several

mechanisms. These include the transfer of locally adaptive variants via introgression (Abbott et al.,

2013), combinatorial speciation through the emergence of new allelic combinations selected in

different habitats (Marques et al., 2019), and the resolution of genomic conflicts caused by

incompatibilities, resulting in RI from both parental lineages (Schumer et al., 2015). The latter may be

followed by ecological niche segregation to reduce competitive interactions and avoid detrimental

hybridisation. In such scenarios, SVs may provide "pre-packaged" divergent haplotypes that can

readily contribute to RI following introgression (Edelman et al., 2019; Feder et al., 2003) or may

better resist re-homogenization, compared to the collinear genome, after the onset of gene flow

(Machado et al., 2007; Noor et al., 2001; Rafajlović et al., 2021). However, further empirical evidence

is required to better understand the role of admixture in driving ecotype divergence through the

transfer or assortment of SVs.

Here, we investigate the role of SVs in ecotype formation in the European anchovy, *Engraulis*

encrasicolus sensu lato. This species complex has formerly been shown to be subdivided into a

marine ecotype (offshore and pelagic, *E. engraulis s. stricto*) and a coastal ecotype (nearshore,

lagoonal and estuarine, *E. maeoticus*) that are able to co-exist in quasi-sympatry despite frequent

hybridisation (Le Moan et al., 2016). Partial RI between these ecotypes is evidenced by differences

in both their genetic makeup and phenotypic characteristics (see Bonhomme et al., 2022 for a

review). The divergence between ecotypes involves a condensed genomic architecture, suggesting a

potential role for SVs in ecotype formation (Le Moan et al. 2016). Moreover, the most likely

demographic model for their divergence involves secondary contact after a prolonged period of

geographic isolation. The current distribution of *E. encrasicolus* spans the North-East Atlantic,

Mediterranean, and Black seas. Classically, the southern range limit of *E. encrasicolus* along the

western African coast is considered as the south of the Gulf of Guinea, while *E. capensis* has been

described further south in the Benguela system off South Africa. However, some samples collected

near the Atlantic-Mediterranean transition zone, off the Moroccan coast or near the Canary Islands,

demonstrated genetic proximity with *E. capensis* (Silva et al., 2017; Zarraonaindia et al., 2012),

suggesting that these might present a third unknown ancestry. These findings warrant further

investigation into potential admixture scenarios between *E. capensis* and European anchovies, and

considering possible evolutionary impacts on the divergence between the marine and coastal

ecotypes.

We present the first study using whole-genome sequencing (WGS) in the *E. encrasicolus* species

complex to provide a detailed characterization of its genetic structure. By employing reference-based

mapping and anchoring to a chromosome-scale reference genome, we reveal the genomic

architecture underlying divergence between the different anchovy lineages. To complement the WGS

approach, we incorporated RAD-sequencing data to characterise the eco-geographic structure of

anchovies across a broad geographical range. This included individuals from both the marine and

coastal ecotypes in the northern part of the distribution, as well as anchovies from the Canary

Islands, the Moroccan coast, and South Africa. We aimed to determine whether and how these

lineages have genetically interacted over their evolutionary history and to evaluate the potential role

of SVs in driving their divergence.

Materials and methods

Sampling and DNA extraction

Samples were collected from multiple sites covering a large part of the species distribution area (Locations table, **Supplementary Table S2**) and were issued from various sampling expeditions and local fisheries (**Supplementary Table S1**, Type=="Tissue"). These samples were collected in different types of habitats, which were classified either as coastal or marine habitats. Also included in our sampling scheme were eight individuals collected off the South African coast (Gqeberha). Whole genomic DNA was extracted from muscle tissue or fin clips using commercial tissue kits (Qiagen and Macherey-Nagel). Extraction quality was checked on agarose gel for the presence of high molecular weight DNA, and double-stranded nucleic acid concentration was measured using Qubit 2.0 and standardised in concentration before library construction.

Reference genome assembly

We performed high-coverage linked-read (10X genomics) sequencing of a marine Atlantic *E. encrasicolus* individual from the Faro location (Algarve) to generate a new reference genome assembly (hereafter called Eencr_V1), following the same methodology as in Meyer et al. (2024). The *de novo* assembly obtained by analysing preprocessed linked-reads (raw coverage ~40X) with supernova v2.1.1 (Weisenfeld et al., 2017) reached a total length of ~926 Mb (925,873,119 bp; contig N50=13.08 kb; scaffold N50=20.36 kb). All downstream analyses required for variant calling from RAD-seq and WGS data were performed on the subset of scaffolds longer than 10 kb to account for assembly fragmentation. Genomic landscapes of differentiation and local PCA (see

149 below) were reconstructed after anchoring scaffolds to the recently released chromosome-level

150 assembly of an *E. encrasicolus* individual from the Black Sea (GenBank assembly accession:

151 GCA_034702125.1) (**Supplementary Fig. S1**). Whole-genome alignment between our Eencr_V1

152 reference genome and the new assembly was performed with Minimap2 (Li, 2018) and visualised

153 using D-GENIES (Cabanettes & Klopp, 2018).

154 Whole-genome resequencing data

Thirty-nine samples (**Supplementary Table S1**, WGS=="yes") were selected for whole-genome

sequencing (WGS), including samples from coastal and marine habitats in the Atlantic (*GAS*) and the

Mediterranean Sea (*GDL*, *SPN*). We also included samples from the Atlantic-Mediterranean

transition zone (*PRS*) and from South Africa to investigate genetic makeup in these localities.

Individual WGS libraries were prepared following the Illumina TruSeq DNA PCR-Free Protocol and

sequenced to an average depth of ~10-30X on an Illumina NovaSeq 6000. Raw demultiplexed reads

were processed using fastp (v0.19.05) (Chen et al., 2018) and aligned to our Eencr_V1 reference

genome using BWA-MEM (BWA v0.7.17; Li, 2013). Picard (v2.26.8) ("Picard toolkit", 2019) was used

for sorting read alignments, marking duplicates and adding read groups.

Variants were called using the GATK best practices workflow (McKenna et al., 2010; Van der Auwera

et al., 2013), without performing variant and base quality score recalibration steps. Firstly, individual

GVCF files were created from bam files with HaplotypeCaller (GATK v.4.3.0.0). This information was

then stored in a GVCF database using GenomicsDBImport, and VCF files (one file per scaffold) were

generated with GenotypeGVCFs. These files were concatenated into a single VCF file which was

filtered using vcftools v0.1.16 (Danecek et al., 2011) and bcftools v1.19 (Danecek et al., 2021) to

retain only high quality SNPs. This included recoding genotypes as missing for low-quality calls ("--

minGQ 20") and hard-filtering sites based on: normalised variant quality ("QD<2.0"), mapping quality

("MQ<40.0" and "MQRankSum"), the rank sum test for site position within reads

("ReadPosRankSum<-8.0"), strand bias ("FS>60.0" and "SOR>3.0"), average genotype quality

("AVG(FMT/GQ)<20"), and average depth (greater than 90X, corresponding to the 97.5th quantile of

the individual read depth distribution, i.e. "AVG(FMT/DP)>90.0"). The VCF was finally filtered for

indels (including nearby SNPs, "--SnpGap 5"), multiallelic SNPs ("--max-alleles 2") and sites

containing more than 15% of missing genotypes ("--max-missing 0.85"). The final VCF file (hereafter

referred to as the **WGS dataset**) contained ~5.9 M sites located on 9093 different scaffolds longer

than 10 kb.

RAD sequencing data

RAD-seq libraries were prepared for 243 samples (**Supplementary Table S1**, RAD=="yes" and

Type=="Tissue") by batches of 64 multiplexed individuals, following a similar protocol to Baird et al.

(2008) using the *Sbf*I restriction enzyme. Twenty-five of these samples were also used to produce

WGS data, providing a link to understand the genetic structure in both datasets. RAD library

sequencing was performed on an Illumina HiSeq2500 sequencer in 100 bp single-read mode. To

complement our sampling, we also included raw sequencing data for 128 anchovy samples from Le

Moan et al. (2016), which were generated using the same restriction enzyme. The data were

demultiplexed using process_radtags (Stacks v2.60) and reads were aligned to our Eencr_V1

reference genome using BWA-MEM (BWA v0.7.17; Li, 2013). The reference-based Stacks pipeline

was run in an integrated workflow developed by the MBB bioinformatic platform

(https://web.mbb.cnrs.fr/subwaw/workflowmanager.php) (Penaud et al., 2020). Gstacks was run

using default parameters ("--model marukilow --var-alpha 0.05 --gt-alpha 0.05 --max-clipped 0.2")

with the minimum PHRED-scaled mapping quality set to 20 ("--min-mapq 20"). Thereafter, genotypes

were exported in VCF format using the populations module ("--min-populations 2 --min-samples-per-

pop 0.7 --min-maf 0.05") and filtered to remove SNPs with more than 15% missing data across

individuals. The VCF was also filtered to only retain sites that were present in the WGS dataset,

since our objective was to describe the same genetic variation but at a larger geographic scale.

Lastly, all samples from the WGS dataset were integrated into the RAD VCF. The final VCF file

(hereafter referred to as the **RAD dataset**) contained genotype data for 385 samples at 3880

variable sites.

Population structure

To describe the genetic structure in both WGS and RAD datasets, we conducted genome-wide and

chromosome-wide Principal Component Analysis (PCA) using the R package SNPRelate (v1.28.0)

(Zheng et al., 2012). We used ADMIXTURE v1.3.0 (Alexander et al., 2009) to estimate individual

ancestry proportions in all samples using the RAD dataset, assuming K=3 parental ancestries and

using default parameters. Individual ancestry proportions were visualised in a triangle plot based on

their ternary coordinates, and samples were then classified into different non-admixed and admixed

ancestry categories based on their positions. The genetic composition of each sampling location was

summarised by representing the fraction of individuals assigned to each ancestry category, in order

to describe the eco-geographical distribution of the three ancestries.

Genetic differentiation (F_{ST}), nucleotide diversity (π), absolute genetic divergence (d_{XY}), and individual

heterozygosity were calculated for the WGS data in non-overlapping 5 kb windows (with "--minSites

15") using the popgenWindows.py script (Martin, 2018;

https://github.com/simonhmartin/genomics_general).

Identification and genotyping of structural variants

Our analyses of anchovy population structure revealed the presence of large SVs that were several

megabases in size and that segregate across the distribution range. We therefore used

chromosome-wide PCA to identify clusters of individuals representing alternate genotypes at each

SV, and assigned individuals' genotypes based on their cluster membership using both the WGS and

RAD datasets. The PCA axis representing structural variation was evidenced by the presence of

three separated clusters, and was in most cases supported by PC1. Samples which did not show

clear cluster membership based on their PCA coordinates were not genotyped. For samples that had

WGS data, we corroborated genotype assignment with the relative positions of each sample in local

PCA, which was conducted in non-overlapping windows of 5 kb using lostruct (v0.0.0.9; Li & Ralph,

2019). In total, we genotyped individuals at 13 large SVs that occur on different chromosomes.

Divergence history of structural variants

To study the evolutionary relationships between individuals carrying different haplotypes at the 13

SVs, we constructed neighbour-joining phylogenetic trees for each chromosome carrying a large SV.

Trees were obtained using the "phylo" command from VCF-Kit (Cook & Andersen, 2017) which uses

variable sites from a VCF to create a multiple sequence alignment to then calculate a difference

matrix using MUSCLE (Edgar, 2004). Because our chromosome-wide PCAs indicated low

recombination between alternate SV haplotypes on the 13 chromosomes, these SV trees can be

used to resolve the evolutionary relationships among haplotypes without the confounding effect of

recombination. For these analyses, we used a subset of high coverage individuals that are

homozygous for the SV to avoid phasing issues and to facilitate visualisation of haplotype

relationships. We rooted trees on the branch separating alternate homozygote groups. For

comparison, trees were also constructed for chromosomes not carrying SVs and rooted using a

South African sample ("ATL_MAR_ZDA_61_1162").

239 Results

240 Genetic structure in the *E. encrasicolus* species complex reveals

241 three-way admixture

We used a combination of both WGS and RAD-seq data as complementary approaches to study the

genetic structure of anchovies in the eastern Atlantic, Mediterranean and Black Seas. Our RAD

dataset, with mean per-sample coverage of 52.7X, revealed a clear picture of the overall genetic

structure in the *E. encrasicolus* species complex across its entire range distribution. These results,

obtained with a reduced representation SNP dataset, were very similar to those obtained using our

WGS dataset (per-sample coverage 10-30X) which contains a smaller subset of samples (detailed

results provided in the supplied HTML report, see **Supplementary Appendix**). Firstly, we observed

genetic differentiation between samples collected in marine and coastal habitats in the northern part

of the range, corresponding to the previously described marine and coastal ecotypes (Le Moan et al.

2016; Bonhomme et al. 2022). This can be observed along the second axis of variation (PCA 2 in

Fig. 1A), where most coastal samples are positioned at the top of the plot, while marine samples fall

in the bottom right corner. As for PCA 1, this axis shows a different signal that reflects geographic

structure rather than ecological structure. On the horizontal axis, South African samples and other

individuals collected off the African Atlantic coast (Morocco and the Canary Islands) are spread out

towards the left-hand side of the plot, while the majority of other samples group to the right. Hence,

PCA at a genome-wide scale shows the existence of three distinct genetic ancestries, which were

further confirmed using admixture analysis. Inferred individual ancestry was represented in a ternary

plot (**Fig. 1B**) showing the relative proportions of coastal (top), marine (right) and southern (left)

ancestry for each individual. Ongoing gene flow between the three ancestries was revealed by

considerable levels of admixture, in particular between the marine and coastal ancestries. A large

number of samples also fell in the central area of the plot due to having balanced proportions of the

three ancestry components, reflecting the existence of three-way admixture.

Based on their ternary coordinates, samples were classified as belonging to one of seven ancestry

categories, each corresponding to a different sub-area of the triangle plot (black demarcations **Fig.**

1B, triangle in **Fig. 1C**). A sample was thus considered to belong to a non-admixed genetic cluster

(coastal/*C*: green; southern/*S*: red; marine/*M*: blue) if that ancestry reached more than 80% of its

total genetic ancestry. Secondly, we distinguished three different admixed classes where individual

ancestry proportions were mainly made up of two ancestries (the third not amounting to more than

10%). These classes were *CS* (admixed between *C* and *S*; brown), *SM* (admixed between *S* and *M*;

purple) and *MC* (admixed between *M* and *C*; seagreen). A last admixed class, called *MCS*, consisted

of individuals with balanced proportions of all three ancestries (admixed between *M*, *C* and *S*; grey).

The eco-geographical distribution of the three ancestries revealed that individuals belonging to the *C*

cluster (green) were only found in coastal habitats (diamond symbols) in the northern part of the

range, while *M* individuals (blue) mainly occurred in marine environments (square symbols) of the

same region (**Fig. 1C**). This corresponds to the previously described ecotypic structure between

coastal and marine anchovies. This structure was particularly evident in the Mediterranean Sea,

where almost all coastal samples were part of the *C* cluster (e.g. coastal habitats in *SIC*, *TNO* and

GDL). However, this signal of ecotypic differentiation becomes diluted near the Atlantic-

Mediterranean boundary, where a gradient of increasing southern ancestry is observed. This

admixture gradient can be seen through the increasing proportion of *MCS* individuals (grey) in the

Alboran Sea (*ALB*), off the southern coast of Portugal (*PRS*) and in northern Morocco (MA4 and

MA3). Finally, we observed that samples from locations to the south of the Canary islands (*CNR*),

including the sampling site in South Africa (*ZDA*, inset map), all belonged to the *S* cluster (red).

After describing the three ancestries as well as their ecogeographic distribution patterns, we aimed to

study their genomic architecture of differentiation. Genomic differentiation landscapes reconstructed

between the coastal, marine and southern clusters using WGS data, yielded highly heterogeneous

patterns that strongly varied from chromosome to chromosome (**Fig. 2**). The background level of

differentiation between the marine and coastal clusters was lower (**Fig. 2A,** mean F_{ST} 0.013)

compared to the background F_{ST} between the southern and coastal clusters (**Fig. 2B**, mean F_{ST}

0.075) and between the southern and marine clusters (**Fig. 2C**, mean F_{ST} 0.066) (see

Supplementary Fig. S9 for F_{ST} distributions). Genomic landscapes of differentiation between marine

and coastal individuals in the Atlantic (**Fig. 2A,** *ATL*: first row of each comparison) and in the

Mediterranean Sea (**Fig. 2A,** *MED*: second row) were similar, even though some differences were

observed (e.g. on chromosomes CM068262 and CM068273).

Heterogeneous F_{ST} values across genomic differentiation landscapes were largely characterised by

the presence of high-differentiation regions (F_{ST} values above the 95[th] quantile) which clustered into

continuous F_{ST} plateaus. These patterns could point to the presence of large SVs that are associated

with ecotype or lineage divergence. To assess whether these putative SVs harbour divergent, non-

recombining haplotypes, we performed PCA on individual chromosomes using both the WGS and

RAD datasets (**Supplementary Fig. S2 and S4**). Chromosome-wide PCAs revealed consistent

patterns of clustering between the WGS and RAD datasets, but these varied greatly across

chromosomes, as did the proportion of genetic variation explained by the first two PC axes.

Representative examples of chromosome-wide PCAs (**Fig. 3A-H**) illustrate the main diversity of

observed patterns. While some chromosomes showed largely continuous ancestry gradients (**Fig.**

3A & E), others exhibited several discrete clusters where samples are tightly grouped (**Fig. 3B-D &**

F-H). These tight PCA clusters indicate that a large number of SNPs are in strong linkage

disequilibrium (LD), resulting in the segregation of a limited number of non-recombining haplotypes.

Combined with elevated levels of heterozygosity in the intermediate clusters (**Fig. 3K,**

Supplementary Fig. S11) and the continuous clustering of genotypes observed across numerous

consecutive local PCA windows (**Fig. 3L**), these patterns strongly support the presence of multiple

large SVs.

In total, we identified 13 chromosomes (indicated by asterisks in **Fig. 2**) that showed evidence for

SVs spanning at least 2.5% of the windows on the chromosome. Using individual coordinates from

chromosome-wide PCAs, these 13 SVs were successfully genotyped (**Supplementary Fig. S3 and**

S5) and individuals were classified as either *00* homokaryotes (pink), *01* heterokaryotes (salmon) or

11 homokaryotes (gold) (**Fig. 3I-L**). Individuals that could not be confidently assigned to any given

group were not genotyped (grey). For consistency, we always assigned the *00* genotype to the group

containing the most southern samples, in order to polarise the *0* haplotype with respect to southern

lineage ancestry. For the SV located on chromosome CM068256, the PCAs based on the WGS and

RAD datasets showed different results, which complicated our assignment of genotypes for this SV

(**Supplementary Fig. S4 and S6**). We opted to genotype this SV according to the variation captured

by PC1 in the WGS analysis, since this dataset included more markers and the structure was

consistent with the phylogenetic reconstruction (**Supplementary Fig. S7)**.

Anchovy lineages are differentiated at multiple SVs

Based on the assigned SV genotypes, we analysed the frequency patterns of the two haplotypes at

each SV (*0* and *1*) (**Supplementary Fig. S8**) as well as the genotype frequencies (*00*, *01* or *11*)

within each genetic cluster (**Fig. 4**). We observed that the coastal, marine and southern clusters

(background colours in **Fig. 4**) carried different sets of SV genotypes. The southern cluster (bottom

row of pie charts) largely harboured homokaryotic *00* genotypes at SVs (mean *00* frequency of 77%

across SVs, 6 out of 13 fixed), while marine individuals (top two rows) were mostly homokaryotic for

the *11* genotype (mean *11* frequency of 79% across SVs). In coastal individuals, on the other hand,

some SVs were nearly fixed for the *00* genotype, while others were fixed for the *11* genotype (see

details below). Overall, we found that the three clusters show substantial frequency differences at

multiple SVs (**Supplementary Table S3**), which is coherent with the F_{ST} plateaus that were observed

in the differentiation landscapes (**Fig. 2**). Admixed individuals that carry more than one type of

ancestry (e.g. *MCS*) were often heterokaryotes at SVs (**Supplementary Fig. S6**), consistent with

their admixed status.

Even though SV frequency differences between the coastal, marine and southern clusters are

observed, these SVs are not always fixed for a given haplotype but display a degree of haplotype

sharing between the three clusters. This is clearly visible on CM068262 and CM068273, for example,

where the SVs are polymorphic in almost all populations (**Fig 4**). Patterns of haplotype sharing can

also be observed in neighbour-joining trees of these SV regions (**Fig. 5B & F**), where samples group

according to their SV genotype (pink and gold branches) and not according to their ancestry category

(tip colour). By looking at the general patterns of haplotype distributions across different SVs, it can

be observed that southern ancestry haplotypes (*0*) are common in marine and/or coastal populations

in the north. Furthermore, these southern haplotypes are slightly more common in the Atlantic (first

and third rows in **Fig. 4**) than in the Mediterranean (second and fourth rows).

One of our main findings is that there is a substantial excess of haplotype sharing at SVs between

the southern and coastal clusters (mean F_{ST} 0.218 in SV regions), which is not observed between the

southern and marine clusters (mean F_{ST} 0.273 in SV regions). This is particularly evident on

CM068255, CM068258 and CM068265, where *0* haplotypes are highly predominant (sometimes

fixed) in the coastal samples. These three chromosomes are among the six showing high F_{ST}

between the marine and coastal clusters (asterisks in **Fig. 2A**), which can thus be explained by the

presence of southern haplotypes in coastal samples. This was confirmed in phylogenies of the SV

regions (**Fig. 5C & G**), since coastal and southern samples are grouped in the same branch (pink),

while marine samples (with the alternative haplotype) are in a separate branch (gold).

On the other hand, high F_{ST} values between coastal and marine samples on CM068256, CM068270

and CM068271 are not due to the presence of southern haplotypes. On these chromosomes,

northern populations almost exclusively carried *1* haplotypes (rows 1-4 in **Fig. 4**) that were, in

addition to their differentiation from the *0* haplotype, also divergent between *C* and *M*. The presence

of haplotype structure within the *1* haplogroup is indeed suggested by the separation of coastal and

marine samples on PCA 2 (**Fig. 3D & H**), and also supported by the subdivision between different *1*

haplotypes in the phylogenies (brown and gold branches in **Fig. 5D & H**). Certain SVs on other

chromosomes also showed evidence for three distinct haplotypes (e.g. CM068273), but only

CM068256, CM068270 and CM068271 were investigated (haplotype frequencies shown in

Supplementary Fig. S8) since these chromosomes were involved in ecotype differentiation (**Fig.**

2A). Overall, these patterns suggest that there are three distinct haplotypes segregating on

CM068256, CM068270 and CM068271, potentially resulting from the presence of multiple SVs on

each chromosome.

By studying the distribution of d_{XY} values (**Supplementary Fig. S10**) and the branch lengths

separating different haplotypes in phylogenies (**Supplementary Fig. S7)**, we gathered information

about divergence in different genomic regions. We aimed to evaluate their concordance with

alternative divergence scenarios and observed different types of patterns that are illustrated by the

four columns in **Fig. 5**. We found that chromosomes that do not carry large SVs are characterised by

short branch lengths and low levels of divergence (**Fig. 5A & E**). Such collinear regions did not show

pronounced genetic structure between the three clusters and supported highest genetic similarity

between the coastal and marine clusters. These patterns contrasted with phylogenies reconstructed

from SV regions, where long branches were found to separate samples carrying different genotypes

(**Fig. 5B-D & F-H**). Estimated d_{XY} values between alternate homokaryotes showed relatively similar

distributions across SVs, with mean d_{XY} between 0.15 and 0.20%. Divergence between coastal *1*

haplotypes and marine *1* haplotypes on CM068256, CM068270 and CM068271 was lower, as is

reflected by shorter branch lengths in phylogenies (**Fig. 5D & H**). We note, however, that SV block

delimitation (based on F_{ST} and local PCA patterns) was probably not precise enough to exclude all

recombinant regions and to provide precise divergence estimates.

Discussion

We present the first genome-scale study investigating genetic structure in the *E. encrasicolus* species complex, comprising anchovies from the eastern Atlantic, Mediterranean and Black Seas. Our findings reveal that two previously described ecotype lineages (marine and coastal) genetically interact with a third, southern lineage. This lineage has not previously been described as such, even though some anchovy samples from the northeast Atlantic and from southern Morocco were shown to display genetic proximity with *E. capensis* (Silva et al., 2017; Zarraonaindia et al., 2012). Although taxonomic considerations are beyond the scope of this study, our results show that the coastal, marine and southern lineages are primarily differentiated by genotype combinations at multiple large SVs, with the collinear parts of their genomes being weakly differentiated. We further found that these SVs most likely originated from admixture with the southern lineage, and that the coastal ecotype carries southern haplotypes at several SVs. Our study supports that SVs that arose in geographically isolated lineages can play an important role in driving ecotype formation following lineage admixture.

Previous genetic studies have shown that the European anchovy is subdivided into marine and coastal ecotypes that are present from the Bay of Biscay, through the Mediterranean to the Black Sea (reviewed in Bonhomme et al., 2022). Here, we show that there is a third component of genetic structure in this species complex, corresponding to an Atlantic lineage occurring off the African coastline. This southern lineage shows genetic homogeneity at a very large spatial scale, with genetic similarity between individuals sampled in Morocco, the Canary Islands and even as far as South Africa. However, from northern Morocco and southern Portugal into the Alboran Sea, we observe genetic admixture resulting in a gradient of decreasing southern ancestry. Previous studies

reporting various signals of spatial structure in this species may have unknowingly captured different

aspects of these complex admixed ancestries, leading to many different and conflicting

interpretations in the literature. For instance, Zarraonaindia et al. (2012) reported the presence of an

admixture gradient extending from the Canary islands to the northern Iberian peninsula. However,

this gradient was attributed to the interaction of two components corresponding to populations

inhabiting narrow versus wide shelf waters. We instead propose that this region corresponds to a

three-way contact zone between the southern lineage and the two northern marine and coastal

ecotype lineages, thus involving both an ecological component as well as admixture between

geographic lineages. We observe post-F1 introgressive hybridisation resulting in widespread

admixture and gene flow between the southern, coastal and marine genetic clusters, as is reflected

by gradual ancestry gradients in the PCA plot (**Fig. 1**). Almost all individuals that were identified as

belonging to the three-way admixed class (MCS) were sampled in the contact zone between the

three lineages (**Fig. 1C**). This pattern of three-way admixture also extends further north in the

Atlantic, where a few individuals carrying a MCS background were detected. The existence of gene

flow between coastal and marine ecotypes has already been illustrated in previous work (Le Moan et

al., 2016), but our results reveal that admixture with the southern lineage also contributes to global

diversity patterns.

We found evidence for multiple megabase-scale SVs that segregate in the marine, coastal and

southern anchovy lineages. Although the presence of SVs was only supported through indirect

evidence based on LD, divergence and heterozygosity patterns, these regions showed many of the

signals typically associated with chromosomal inversions (Mérot et al., 2020). In the literature, it has

often been reported that SVs play a role in differentiating evolutionary lineages within species

complexes, suggesting that they can play an important role in the formation of cryptic lineages or

ecotypes (Berdan et al., 2024; Zhang et al., 2021). Our results in anchovies seem to point in this

direction, since markers differentiating lineages and ecotypes were largely concentrated in SVs,

whereas collinear regions of the genome showed low differentiation levels which likely reflects the

homogenising effect of gene flow and recombination. By reconstructing the genomic landscape of

ecotype divergence, we determined that marine and coastal ecotypes were differentiated at six large

SVs. These SVs cover roughly 25% of the genome, which is in line with a previous study that

estimated that the barriers to gene flow between ecotypes span 20 to 25% of the genome (Le Moan

et al., 2016). The fine-scale structure of the SVs will need to be addressed by long-read sequencing

in order to directly identify breakpoints and to resolve the possibly complex variation at each SV. In

addition, there is also the possibility that small genomic islands of differentiation, located in the

collinear genome and not detected here, could also play a role in ecotype divergence.

Our results also revealed that the origin of the six SVs that differentiate ecotypes is closely linked to

the impacts of admixture between divergent geographic lineages. In particular, the coastal ecotype

carries the same haplotype as the southern lineage at a minimum of three SVs. If these SVs were

already segregating in the ancestral population to the marine, coastal and southern lineages,

different conflicting genealogies would have been expected due to incomplete lineage sorting (ILS).

However, we do not observe any SVs where the marine and southern lineages carry the same

haplotype and are divergent with the coastal ecotype. If these patterns are not due to ILS, it suggests

that the coastal and southern lineages could have a partly shared evolutionary history. Overall, the

patterns of population structure in the collinear genome as well as haplotype distributions at SVs

(whether shared between lineages or private) suggest that there are three alternative scenarios

which could underlie the observed genealogical patterns (**Fig. 5M, N & O**):

1) In the first scenario, SVs could have been introgressed from the southern lineage into the

coastal lineage. The deepest split in the tree represents the initial divergence between the

southern lineage and the northern ancestral branch giving rise to the marine and coastal

ecotypes (**Fig. 5M**). Following this split, genomic rearrangements likely accumulated in each

branch, and subsequent gene flow could have led to the introgression of SVs. In this

scenario, haplotypes from the southern lineage may have been introgressed into a

preexisting coastal lineage or may have contributed to the formation of the coastal ecotype.

This introgression likely occurred during an earlier period of contact, since current admixture

in the contact zone seems insufficient to explain why coastal samples from across the

distribution range almost exclusively carry southern haplotypes at these SVs. Evidence for

the introgression of SVs is documented in other systems (Jay et al., 2018; Todesco et al.,

2020), where they may serve as a source of divergent haplotypes involved in the evolution of

novel traits or ecotype formation.

2) The second scenario proposes that the coastal ecotype directly originated from the southern

lineage and does not share its most recent common ancestor with the marine lineage (**Fig.**

5N). In this case, recent common ancestry between the coastal and southern lineages would

only be observed at the SVs, as widespread introgression with the marine lineage has

eroded divergence in the collinear genome. This scenario does not require multiple SVs to

successfully pass through various selective filters to establish. Instead, it involves neutral

introgression in the background genome, as well as polymorphism at certain SVs that were

shared via gene flow.

3) Another alternative scenario is whole-genome lineage admixture, where the coastal ecotype

would have originated from an admixed population composed of both southern and northern

ancestries. This admixture could have been followed by selective reassortment of SVs, driven

either by adaptation to the coastal environment (Marques et al., 2019) or by the resolution of

genomic conflicts resulting from incompatibilities (Schumer et al., 2015).

The evolutionary history of the marine and coastal ecotypes has previously been studied, and

demographic inference has indicated that ecotype divergence probably occurred in a context of

allopatric isolation followed by secondary contact (Le Moan et al., 2016). Although the

aforementioned study did not include the third anchovy lineage in their analyses, the period of

allopatric divergence that was inferred between ecotypes may in fact correspond to the split between

northern and southern lineages (**Fig. 5M & N & O**). This divergence signal may have been preserved

within the SVs, as is suggested by their d_{XY} distributions and long internal branch lengths in

genealogies. By contrast, the signal of past divergence between the northern and southern lineages

may have been eroded by subsequent gene flow and recombination in the background genome.

In addition to the six SVs that separate coastal and marine anchovies, we also identified other SVs

(e.g. on CM068260) that are shared between ecotypes and where one haplotype has a southern

lineage origin. Here again, the lack of recombination between rearrangements has preserved the

historical divergence signal, despite the fact that these SVs have been subsequently exchanged

between ecotypes. This result closely mirrors what has been described by studies using

mitochondrial data, where two deeply divergent clades (called A and B) were found to segregate in

populations now known to correspond to the marine and coastal ecotypes (Magoulas et al., 1996;

Oueslati et al., 2014) and the southern lineage (Chahdi Ouazzani, 2016; W. S. Grant et al., 2005;

Silva et al., 2014, 2017). Therefore, the mitochondrial genome and the shared SVs likely represent

genomic regions that have retained the signal of north-south divergence despite gene flow following

secondary contact.

The last type of genealogical pattern that we identified for certain SVs, separates the southern

lineage from coastal and marine ecotypes that share more recent common ancestry (**Fig. 5D & H**),

thus displaying the same topology as for the collinear genome but with significantly longer branches.

This might imply that, in addition to north-south divergence, there is also more recent divergence

between the two northern anchovy ecotypes, possibly involving complex rearrangements such as

nested inversions (Joron et al., 2011; Kollar et al., 2024). Alternatively, these SVs could have a

similar evolutionary history as proposed above in scenarios 1, 2 & 3. Namely, coastal anchovy would

have carried a southern haplotype in the past, before it recombined with the northern haplotype due

to gene flow between coastal and marine ecotypes, consequently decreasing the level of divergence

between haplotypes in these populations. Gene flow between inversion arrangements has previously

been described in several other systems, in particular for large chromosomal inversions that are

impacted by gene conversion and double crossovers (Korunes & Noor, 2019; Matschiner et al.,

2022; Meyer et al., 2024; Navarro et al., 1997).

Overall, we found that secondary contact between the northern and southern anchovy lineages –

whether involving introgression of specific SVs (scenario 1), background genome introgression of the

southern lineage (scenario 2), or genome-wide admixture (3) – is sufficient to explain the diversity of

genealogical patterns observed at the genome-wide scale. However, our results do not allow us to

determine which of the three alternative evolutionary scenarios is the most likely. This would require

more in-depth demographic divergence analyses and complementary information regarding the past

geographic distributions of the different lineages, especially during the last glacial period. However,

we may hypothesise that the divergence between the northern and southern branches probably took

place across hemispheres, as it broadly reflects the current distribution of the extant lineages.

Alternatively, divergence may have occurred between the northeast and -west Atlantic, since

anchovy populations from the Gulf of Mexico may also form part of the *E. encrasicolus* species

complex (Silva et al. 2014; 2017). It is likely that several periods of secondary contact could have

taken place between ancient anchovy lineages, most likely during cooler periods when long-range

distributional range shifts were possible and when the lineages were confined to lower latitudes due

to ice sheets near the poles (Bouchenak-Khelladi et al., 2008; W. Grant & Bowen, 1998). In

particular, the population of southern anchovies off the Moroccan coast could have originated during

the last glacial episode, when anchovies from the southern hemisphere could migrate north and

come into contact with northern populations.

It remains unresolved, however, whether the ancestral anchovy lineages were already associated

with particular habitats or how/when this association took place to result in the current coastal and

marine ecotypes. Today, the southern lineage seems to be associated with marine habitats, at least

along the Moroccan, Mauritanian, Namibian and South African coasts (Chahdi Ouazzani et al., 2017;

Whitfield, 1994). In this light, it is not obvious at first sight how the southern SVs haplotypes, which

are carried by the coastal ecotype, could underlie adaptation to coastal environments. However, it is

possible that epistatic interactions with loci in the genomic background have affected ecological traits

that were not initially associated with these haplotypes. A classical example is the introgression of

the Denisovan *Epas1* variant involved in adaptation to high altitude in modern humans, without being

obviously involved in adaptation to altitude in the Denisova lineage (Huerta-Sánchez et al., 2014).

Alternatively, mutations could have accumulated in the SVs over time, conferring local adaptation to

the coastal environment, which was not initially the case in the southern lineage. Empirical studies

(Lamichhaney et al., 2016) and recent simulation work (Schaal et al., 2022) have indeed found

support for the gradual accumulation of local adaptation alleles in inversions under high gene flow

conditions.

Our study highlights the crucial role of SVs and admixture in the formation of ecotypes, supporting

that hybridisation between geographically isolated lineages can provide the genetic substrate

necessary for ecological specialisation and partial RI. SVs can act as "pre-packaged" divergent

genomic elements containing multiple genes that contribute to reproductive barriers through local

adaptation, coadapted gene complexes, or genetic incompatibilities. This supports a broader view

that ecotype formation is rarely driven by strict *in situ* adaptation alone, but often involves phases of

allopatric divergence and other historical contingencies (Van Belleghem et al., 2018; Bernatchez et

al., 1996; Bierne et al., 2011; Hendry, 2009; Rundell & Price, 2009). Following lineage admixture,

SVs can fuel ecotypic differentiation and resist gene flow over the long term, facilitating coupling

among different components of RI. These associations are necessary for the maintenance of

ecotypes and are instrumental if speciation between incipient ecotypes is ever to complete (Barton &

de Cara, 2009; Bierne et al., 2011; Butlin & Smadja, 2018). Furthermore, the role of coupled SVs

may be even more significant for the persistence of differentiated ecotypes in high gene flow marine

species (Han et al., 2020; Jones et al., 2012; Kirubakaran et al., 2016; Wilder et al., 2020). However,

whether these SVs alone are sufficient for marine ecotypes to speciate remains uncertain

(Johannesson et al., 2024), as the moving balance between gene flow and divergent selection

continuously reshapes their evolutionary trajectory. Future research should explore the broader role

of lineage admixture as a source of divergent SVs and investigate the evolutionary history of these

variants, to reveal how they contribute to the formation and maintenance of ecotypes across different

taxa.

565 Acknowledgements

We wish to thank the numerous colleagues that have provided us with samples and insightful

contributions over the many years of this long-term study, more particularly amongst them Jean-

Pierre Quignard, Karima Fadhlaoui-Zid, Petr Strelkov, Laurent Soulier, Lilia Bahri-Sfar and Nicolas

Bierne. This work was supported by the ANR grant CoGeDiv ANR-17-CE02-0 006-01 to Pierre-

Alexandre Gagnaire. We would like to thank the MGX-Montpellier GenomiX platform as well as the

Montpellier Bioinformatics Biodiversity platform (MBB) supported by the LabEx CeMEB.

Fig. 1. A) Principal Component Analysis (PCA) performed on the entire dataset of 385 anchovy samples. Sites used

in the analysis were high-quality variants present both in the WGS data as well as in the RAD data, corresponding to

a total of 3881 SNPs. Shapes indicate habitat type and colours reflect ancestry proportions as determined by

admixture analysis (see B). B) Ternary plot showing the admixture level between three genetic ancestries: coastal

(green), southern (red), and marine (blue) ancestry. Coordinates, as well as RGB colours, reflect the relative ancestry

proportions of samples along each of the three axes. Samples were classified as belonging to a genetic cluster (black

lines demarcating seven areas) based on their position in the plot. Clusters *C*, *S* and *M* represent "non-admixed"

parental lineage ancestries, while *CS*, *SM*, *MC* and *MCS* represent various levels of admixture. B) Map with sampling

locations where symbols represent habitat type and pie charts show the proportions of different genetic clusters

present. Numbers beneath pie charts indicate sample sizes. Locations are described in **Supplementary Table S2**.

Fig. 2. Genomic landscapes of differentiation (F_{ST}) calculated in 5 kb sliding windows between groups of samples (3 individuals per group) from different genetic clusters (see Fig. 1). Differentiation landscapes are shown for three different comparisons (A: Coastal vs. Marine; B: Coastal vs. southern; C: southern vs. Marine). Each panel consists of two rows, representing cases where Coastal/Marine samples either originated from the Atlantic (ATL) or from the Mediterranean Sea (MED). Orange points are windows where F_{ST} was higher than the 95th quantile, while red points are above the 99th quantile. Stars indicate chromosomes where more than 2.5% of windows showed F_{ST} higher than the 95th quantile. Grey and white rectangles delimit the 24 chromosomes of *E. encrasicolus*.

Fig. 3. Examples of relatedness patterns on different chromosomes. PCA conducted on whole-genome data (n=39) at

a chromosome-wide level (A-H) reveals that certain chromosomes show signs of reduced recombination through the

presence of tight PCA clusters (B-D, F-H). These chromosomes show evidence for the presence of structural variants

(SVs), which we aimed to genotype in all individuals (I-L). SV genotypes were assigned for whole-genome and RAD

data (n=385) based on PCA coordinates (I & J). For whole-genome data, these assignments were further

corroborated by individual heterozygosity levels (K) and local PCA patterns (L). In (L), PCA 1 coordinates plotted for

non-overlapping 5 kb windows along the chromosome show sustained clustering patterns in the SV region (ending

around ~40 Mb). Gaps represent regions on CM068255 for which we did not have data (scaffolds < 10 kb or not

present in our reference genome). Colours either reflect genome-wide ancestry proportions (A-H) or the SV genotype

that was assigned to each individual (I-L).

Fig 4. Genotype frequencies for 13 SVs on different chromosomes (columns). Pie charts show frequencies for the *00* (pink), *01* (salmon) and *11* (gold) genotypes in the Marine cluster (blue background), Coastal cluster (green background) and southern cluster (red background), with upper and lower rows corresponding to Atlantic and Mediterranean samples respectively (for *M* and *C*). Darker background colour in *M* and *C* indicates chromosomes that show elevated F_{ST} when comparing Marine and Coastal individuals (Fig. 2A). For three of these chromosomes (CM068255, CM068258 and CM068265), ecotype differentiation involves southern haplotypes (*00*) that are present at high frequency in the Coastal individuals, while this is not the case for CM068270 and CM068271.

Fig. 5. A-H) Neighbour-joining trees showing interindividual relationships on the same subset of chromosomes as in

Fig. 3. Short branch lengths for normally recombining chromosomes (A & E) contrast with long branches between

divergent haplotypes at SVs (B-D, F-H). These SVs show varying patterns of being shared or being private to the

coastal, marine or southern clusters. This is what we illustrate with schematic trees in (I-L), where pink or gold

colouration reflects divergence between haplotypes. In (L), brown colouration indicates the divergence of a third

haplotype differentiating coastal and marine samples (e.g. CM068270 and CM068271). We propose three

evolutionary scenarios (M, N & O) involving gene flow amongst lineages (horizontal arrows) which could explain the

observed patterns of haplotype distributions across SVs. For (A & E), trees were constructed using SNPs on the

entire chromosome, whereas in (B-D, F-H), the region was limited to that covered by the SV. For trees of SV regions,

intermediate samples that present heterokaryotes are not displayed. Leaf tips are coloured according to individual

ancestry categories. Trees were plotted on the same vertical scale.

References

Abbott, R., Albach, D., Ansell, S., Arntzen, J. W., Baird, S. J. E., Bierne, N., Boughman, J.,
   Brelsford, A., Buerkle, C. A., Buggs, R., Butlin, R. K., Dieckmann, U., Eroukhmanoff, F.,
   Grill, A., Cahan, S. H., Hermansen, J. S., Hewitt, G., Hudson, A. G., Jiggins, C., …
   Zinner, D. (2013). Hybridization and speciation. *Journal of Evolutionary Biology*, *26*(2),
   229-246. https://doi.org/10.1111/j.1420-9101.2012.02599.x

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in
   unrelated individuals. *Genome Research*, *19*(9), 1655-1664.
   https://doi.org/10.1101/gr.094052.109

Baird, N. A., Etter, P. D., Atwood, T. S., Currey, M. C., Shiver, A. L., Lewis, Z. A., Selker, E. U.,
   Cresko, W. A., & Johnson, E. A. (2008). Rapid SNP Discovery and Genetic Mapping
   Using Sequenced RAD Markers. *PLOS ONE*, *3*(10), e3376.
   https://doi.org/10.1371/journal.pone.0003376

Barton, N. H., & de Cara, M. A. R. (2009). The Evolution of Strong Reproductive Isolation.
   Evolution, *63*(5), 1171-1190.

Belleghem, S. M. V., Vangestel, C., Wolf, K. D., Corte, Z. D., Möst, M., Rastas, P., Meester, L.
   D., & Hendrickx, F. (2018). Evolution at two time frames : Polymorphisms from an ancient
   singular divergence event fuel contemporary parallel evolution. *PLOS Genetics*, *14*(11),
   e1007796. https://doi.org/10.1371/journal.pgen.1007796

Benjamin Penaud, Rémy Dernat, Iago Bonnici, & Khalid Belkhir. (2020). *MBB-Framework : A set
   of tools to build and run reproducible and portable workflows* [Logiciel].

https://web.mbb.cnrs.fr/subwaw/workflowmanager.php

Berdan, E. L., Aubier, T. G., Cozzolino, S., Faria, R., Feder, J. L., Giménez, M. D., Joron, M.,

Searle, J. B., & Mérot, C. (2024). Structural Variants and Speciation : Multiple Processes

at Play. *Cold Spring Harbor Perspectives in Biology*, *16*(3), a041446.

https://doi.org/10.1101/cshperspect.a041446

Bernatchez, L., Vuorinen, J. A., Bodaly, R. A., & Dodson, J. J. (1996). Genetic Evidence for

Reproductive Isolation and Multiple Origins of Sympatric Trophic Ecotypes of Whitefish

(Coregonus). *Evolution*, *50*(2), 624-635. https://doi.org/10.2307/2410836

Bierne, N., Welch, J., Loire, E., Bonhomme, F., & David, P. (2011). The coupling hypothesis :

Why genome scans may fail to map local adaptation genes: THE COUPLING

HYPOTHESIS. *Molecular Ecology*, *20*(10), 2044-2072. https://doi.org/10.1111/j.1365-

294X.2011.05080.x

Bonhomme, F., Meyer, L., Arbiol, C., Bănaru, D., Bahri-Sfar, L., Fadhlaoui-Zid, K., Strelkov, P.,

Arculeo, M., Soulier, L., Quignard, J.-P., & Gagnaire, P.-A. (2022). Systematics of

European coastal anchovies (genus Engraulis Cuvier). *Journal of Fish Biology*, *100*(2),

594-600. https://doi.org/10.1111/jfb.14964

Bouchenak-Khelladi, Y., Durand, J.-D., Magoulas, A., & Borsa, P. (2008). Geographic structure

of European anchovy : A nuclear-DNA study. *Journal of Sea Research*, *59*.

https://doi.org/10.1016/j.seares.2008.03.001

Butlin, R. K., & Smadja, C. M. (2018). Coupling, Reinforcement, and Speciation. *The American*

Naturalist, *191*(2), 155-172. https://doi.org/10.1086/695136

Cabanettes, F., & Klopp, C. (2018). D-GENIES : Dot plot large genomes in an interactive,

efficient and simple way. *PeerJ*, *6*, e4958. https://doi.org/10.7717/peerj.4958

Chahdi Ouazzani, K. (2016). Genetic population structure of the European anchovy (Engraulis

encrasicolus) based on mitochondrial DNA sequences along the Moroccan coast. *AACL*

Bioflux, *9*.

Chahdi Ouazzani, K., Benazzou ,Touria, Charouki ,Najib, Bonhomme ,François, & and Chlaida,

M. (2017). Genetic differentiation of European anchovy (Engraulis encrasicolus) along

the Moroccan coast reveals a phylogeographic break around the 25th parallel North.

Marine Biology Research, *13*(3), 342-350.

https://doi.org/10.1080/17451000.2016.1257811

Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). fastp : An ultra-fast all-in-one FASTQ

preprocessor. *Bioinformatics*, *34*(17), i884-i890.

https://doi.org/10.1093/bioinformatics/bty560

Cook, D. E., & Andersen, E. C. (2017). VCF-kit : Assorted utilities for the variant call format.

Bioinformatics, *33*(10), 1581-1582. https://doi.org/10.1093/bioinformatics/btx011

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker, R.

E., Lunter, G., Marth, G. T., Sherry, S. T., McVean, G., Durbin, R., & 1000 Genomes

Project Analysis Group. (2011). The variant call format and VCFtools. *Bioinformatics*

(Oxford, England), *27*(15), 2156-2158. https://doi.org/10.1093/bioinformatics/btr330

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham, A.,

Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve years of SAMtools

and BCFtools. *GigaScience*, *10*(2), giab008. https://doi.org/10.1093/gigascience/giab008

Edelman, N. B., Frandsen, P. B., Miyagi, M., Clavijo, B., Davey, J., Dikow, R. B., García-

Accinelli, G., Van Belleghem, S. M., Patterson, N., Neafsey, D. E., Challis, R., Kumar, S.,

Moreira, G. R. P., Salazar, C., Chouteau, M., Counterman, B. A., Papa, R., Blaxter, M.,

685  Reed, R. D., … Mallet, J. (2019). Genomic architecture and introgression shape a

686  butterfly radiation. *Science*, *366*(6465), 594-599.

687  https://doi.org/10.1126/science.aaw2090

688 Edgar, R. C. (2004). MUSCLE : Multiple sequence alignment with high accuracy and high

689  throughput. *Nucleic Acids Research*, *32*(5), 1792-1797.

690  https://doi.org/10.1093/nar/gkh340

691 Faria, R., Chaube, P., Morales, H. E., Larsson, T., Lemmon, A. R., Lemmon, E. M., Rafajlović,

692  M., Panova, M., Ravinet, M., Johannesson, K., Westram, A. M., & Butlin, R. K. (2019).

693  Multiple chromosomal rearrangements in a hybrid zone between Littorina saxatilis

694  ecotypes. *Molecular Ecology*, *28*(6), 1375-1393. https://doi.org/10.1111/mec.14972

695 Feder, J. L., Berlocher, S. H., Roethele, J. B., Dambroski, H., Smith, J. J., Perry, W. L.,

696  Gavrilovic, V., Filchak, K. E., Rull, J., & Aluja, M. (2003). Allopatric genetic origins for

697  sympatric host-plant shifts and race formation in Rhagoletis. *Proceedings of the National*

698  *Academy of Sciences*, *100*(18), 10314-10319. https://doi.org/10.1073/pnas.1730757100

699 Grant, W., & Bowen, B. (1998). Shallow population histories in deep evolutionary lineages of

700  marine fishes : Insights from sardines and anchovies and lessons for conservation.

701  *Journal of Heredity*, *89*(5), 415-426. https://doi.org/10.1093/jhered/89.5.415

702 Grant, W. S., Leslie, R. W., & Bowen, B. W. (2005). Molecular genetic assessment of bipolarity

703  in the anchovy genus Engraulis. *Journal of Fish Biology*, *67*(5), 1242-1265.

704  https://doi.org/10.1111/j.1095-8649.2005.00820.x

705 Hager, E. R., Harringmeyer, O. S., Wooldridge, T. B., Theingi, S., Gable, J. T., McFadden, S.,

706  Neugeboren, B., Turner, K. M., Jensen, J. D., & Hoekstra, H. E. (2022). A chromosomal

707  inversion contributes to divergence in multiple traits between deer mouse ecotypes.

Science, *377*(6604), 399-405. https://doi.org/10.1126/science.abg0718

Han, F., Jamsandekar, M., Pettersson, M. E., Su, L., Fuentes-Pardo, A. P., Davis, B. W.,

Bekkevold, D., Berg, F., Casini, M., Dahle, G., Farrell, E. D., Folkvord, A., & Andersson,

L. (2020). Ecological adaptation in Atlantic herring is associated with large shifts in allele

frequencies at hundreds of loci. *eLife*, *9*, e61076. https://doi.org/10.7554/eLife.61076

Han, F., Lamichhaney, S., Grant, B. R., Grant, P. R., Andersson, L., & Webster, M. T. (2017).

Gene flow, ancient polymorphism, and ecological adaptation shape the genomic

landscape of divergence among Darwin's finches. *Genome Research*, *27*(6), 1004-1015.

https://doi.org/10.1101/gr.212522.116

Hendry, A. P. (2009). Ecological speciation ! Or the lack thereof?This Perspective is based on

the author's J.C. Stevenson Memorial Lecture delivered at the Canadian Conference for

Fisheries Research in Halifax, Nova Scotia, January 2008. *Canadian Journal of Fisheries*

and Aquatic Sciences, *66*(8), 1383-1398. https://doi.org/10.1139/F09-074

Huerta-Sánchez, E., Jin, X., Asan, Bianba, Z., Peter, B. M., Vinckenbosch, N., Liang, Y., Yi, X.,

He, M., Somel, M., Ni, P., Wang, B., Ou, X., Huasang, Luosang, J., Cuo, Z. X. P., Li, K.,

Gao, G., Yin, Y., … Nielsen, R. (2014). Altitude adaptation in Tibetans caused by

introgression of Denisovan-like DNA. *Nature*, *512*(7513), 194-197.

https://doi.org/10.1038/nature13408

Jay, P., Whibley, A., Frézal, L., Rodríguez de Cara, M. Á., Nowell, R. W., Mallet, J.,

Dasmahapatra, K. K., & Joron, M. (2018). Supergene Evolution Triggered by the

Introgression of a Chromosomal Inversion. *Current Biology*, *28*(11), 1839-1845.e3.

https://doi.org/10.1016/j.cub.2018.04.072

Johannesson, K., Faria, R., Moan, A. L., Rafajlović, M., Westram, A. M., Butlin, R. K., &

Stankowski, S. (2024). Diverse pathways to speciation revealed by marine snails. *Trends*

in Genetics, *40*(4), 337-351. https://doi.org/10.1016/j.tig.2024.01.002

Jones, F. C., Grabherr, M. G., Chan, Y. F., Russell, P., Mauceli, E., Johnson, J., Swofford, R.,

Pirun, M., Zody, M. C., White, S., Birney, E., Searle, S., Schmutz, J., Grimwood, J.,

Dickson, M. C., Myers, R. M., Miller, C. T., Summers, B. R., Knecht, A. K., … Kingsley,

D. M. (2012). The genomic basis of adaptive evolution in threespine sticklebacks. *Nature*,

484(7392), Article 7392. https://doi.org/10.1038/nature10944

Joron, M., Frezal, L., Jones, R. T., Chamberlain, N. L., Lee, S. F., Haag, C. R., Whibley, A.,

Becuwe, M., Baxter, S. W., Ferguson, L., Wilkinson, P. A., Salazar, C., Davidson, C.,

Clark, R., Quail, M. A., Beasley, H., Glithero, R., Lloyd, C., Sims, S., … ffrench-Constant,

R. H. (2011). Chromosomal rearrangements maintain a polymorphic supergene

controlling butterfly mimicry. *Nature*, *477*(7363), 203-206.

https://doi.org/10.1038/nature10341

Kirkpatrick, M., & Barton, N. (2006). Chromosome Inversions, Local Adaptation and Speciation.

Genetics, *173*(1), 419-434. https://doi.org/10.1534/genetics.105.047985

Kirubakaran, T. G., Grove, H., Kent, M. P., Sandve, S. R., Baranski, M., Nome, T., De Rosa, M.

C., Righino, B., Johansen, T., Otterå, H., Sonesson, A., Lien, S., & Andersen, Ø. (2016).

Two adjacent inversions maintain genomic differentiation between migratory and

stationary ecotypes of Atlantic cod. *Molecular Ecology*, *25*(10), 2130-2143.

https://doi.org/10.1111/mec.13592

Kollar, L. M., Stanley, L. E., Raju, S. K. K., Lowry, D. B., & Niederhuth, C. E. (2024). *The*

evolutionary dynamics of locally adaptive chromosome inversions in Mimulus guttatus (p.

2023.12.06.570460). bioRxiv. https://doi.org/10.1101/2023.12.06.570460

Korunes, K. L., & Noor, M. A. F. (2019). Pervasive gene conversion in chromosomal inversion

heterozygotes. *Molecular Ecology*, *28*(6), 1302-1315. https://doi.org/10.1111/mec.14921

Kulmuni, J., Butlin, R. K., Lucek, K., Savolainen, V., & Westram, A. M. (2020). Towards the

completion of speciation : The evolution of reproductive isolation beyond the first barriers.

Philosophical Transactions of the Royal Society B: Biological Sciences, *375*(1806),

20190528. https://doi.org/10.1098/rstb.2019.0528

Lamichhaney, S., Fan, G., Widemo, F., Gunnarsson, U., Thalmann, D. S., Hoeppner, M. P.,

Kerje, S., Gustafson, U., Shi, C., Zhang, H., Chen, W., Liang, X., Huang, L., Wang, J.,

Liang, E., Wu, Q., Lee, S. M.-Y., Xu, X., Höglund, J., … Andersson, L. (2016). Structural

genomic changes underlie alternative reproductive strategies in the ruff (Philomachus

pugnax). *Nature Genetics*, *48*(1), Article 1. https://doi.org/10.1038/ng.3430

Le Moan, A., Bekkevold, D., & Hemmer-Hansen, J. (2021). Evolution at two time frames :

Ancient structural variants involved in post-glacial divergence of the European plaice

(Pleuronectes platessa). *Heredity*, *126*(4), Article 4. https://doi.org/10.1038/s41437-020-

00389-3

Le Moan, A., Stankowski, S., Rafajlović, M., Ortega-Martinez, O., Faria, R., Butlin, R. K., &

Johannesson, K. (2024). Coupling of twelve putative chromosomal inversions maintains

a strong barrier to gene flow between snail ecotypes. *Evolution Letters*, qrae014.

https://doi.org/10.1093/evlett/qrae014

Li, H. (2013). *Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM*

(No. arXiv:1303.3997). arXiv. http://arxiv.org/abs/1303.3997

Li, H. (2018). Minimap2 : Pairwise alignment for nucleotide sequences. *Bioinformatics*, *34*(18),

3094-3100. https://doi.org/10.1093/bioinformatics/bty191

Li, H., & Ralph, P. (2019). Local PCA Shows How the Effect of Population Structure Differs

Along the Genome. *Genetics*, *211*(1), 289-304.

https://doi.org/10.1534/genetics.118.301747

Louis, M., Galimberti, M., Archer, F., Berrow, S., Brownlow, A., Fallon, R., Nykänen, M., O'Brien,

J., Roberston, K. M., Rosel, P. E., Simon-Bouhet, B., Wegmann, D., Fontaine, M. C.,

Foote, A. D., & Gaggiotti, O. E. (2020). *Selection on ancestral genetic variation fuels*

parallel ecotype formation in bottlenose dolphins [Preprint]. Evolutionary Biology.

https://doi.org/10.1101/2020.10.05.325159

Lundberg, M., Mackintosh, A., Petri, A., & Bensch, S. (2023). Inversions maintain differences

between migratory phenotypes of a songbird. *Nature Communications*, *14*(1), Article 1.

https://doi.org/10.1038/s41467-023-36167-y

Machado, C. A., Haselkorn, T. S., & Noor, M. A. F. (2007). Evaluation of the Genomic Extent of

Effects of Fixed Inversion Differences on Intraspecific Variation and Interspecific Gene

Flow in Drosophila pseudoobscura and D. persimilis. *Genetics*, *175*(3), 1289-1306.

https://doi.org/10.1534/genetics.106.064758

Magoulas, A., Tsimenides, N., & Zouros, E. (1996). Mitochondrial DNA phylogeny and the

reconstruction of the population history of a species : The case of the European anchovy

(Engraulis encrasicolus). *Molecular Biology and Evolution*, *13*(1), 178-190.

https://doi.org/10.1093/oxfordjournals.molbev.a025554

Marques, D. A., Meier, J. I., & Seehausen, O. (2019). A Combinatorial View on Speciation and

Adaptive Radiation. *Trends in Ecology & Evolution*, *34*(6), 531-544.

https://doi.org/10.1016/j.tree.2019.02.008

Matschiner, M., Barth, J. M. I., Tørresen, O. K., Star, B., Baalsrud, H. T., Brieuc, M. S. O.,

Pampoulie, C., Bradbury, I., Jakobsen, K. S., & Jentoft, S. (2022). Supergene origin and

maintenance in Atlantic cod. *Nature Ecology & Evolution*, *6*(4), Article 4.

https://doi.org/10.1038/s41559-022-01661-x

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K.,

Altshuler, D., Gabriel, S., Daly, M., & DePristo, M. A. (2010). The Genome Analysis

Toolkit : A MapReduce framework for analyzing next-generation DNA sequencing data.

Genome Research, *20*(9), 1297-1303. https://doi.org/10.1101/gr.107524.110

Mérot, C., Oomen, R. A., Tigano, A., & Wellenreuther, M. (2020). A Roadmap for Understanding

the Evolutionary Significance of Structural Genomic Variation. *Trends in Ecology &*

Evolution, *35*(7), 561-572. https://doi.org/10.1016/j.tree.2020.03.002

Meyer, L., Barry, P., Riquet, F., Foote, A., Der Sarkissian, C., Cunha, R. L., Arbiol, C.,

Cerqueira, F., Desmarais, E., Bordes, A., Bierne, N., Guinand, B., & Gagnaire, P.-A.

(2024). Divergence and gene flow history at two large chromosomal inversions

underlying ecotype differentiation in the long-snouted seahorse. *Molecular Ecology*,

e17277. https://doi.org/10.1111/mec.17277

Moan, A. L., Gagnaire, P.-A., & Bonhomme, F. (2016). Parallel genetic divergence among

coastal–marine ecotype pairs of European anchovy explained by differential

introgression after secondary contact. *Molecular Ecology*, *25*(13), 3187-3202.

https://doi.org/10.1111/mec.13627

Navarro, A., & Barton, N. H. (2003). Accumulating postzygotic isolation genes in parapatry : A

new twist on chromosomal speciation. *Evolution; International Journal of Organic*

Evolution, *57*(3), 447-459. https://doi.org/10.1111/j.0014-3820.2003.tb01537.x

Navarro, A., Betrán, E., Barbadilla, A., & Ruiz, A. (1997). Recombination and Gene Flux Caused

823   by Gene Conversion and Crossing Over in Inversion Heterokaryotypes. *Genetics*, *146*(2),

824   695-709. https://doi.org/10.1093/genetics/146.2.695

825 Nelson, T. C., & Cresko, W. A. (2018). Ancient genomic variation underlies repeated ecological

826   adaptation in young stickleback populations. *Evolution Letters*, *2*(1), 9-21.

827   https://doi.org/10.1002/evl3.37

828 Noor, M. A. F., Grams, K. L., Bertucci, L. A., & Reiland, J. (2001). Chromosomal inversions and

829   the reproductive isolation of species. *Proceedings of the National Academy of Sciences*,

830   *98*(21), 12084-12088. https://doi.org/10.1073/pnas.221274498

831 Oueslati, S., Fadhlaoui-Zid, K., Kada, O., Augé, M. T., Quignard, J. P., & Bonhomme, F. (2014).

832   Existence of two widespread semi-isolated genetic entities within Mediterranean

833   anchovies. *Marine Biology*, *161*(5), 1063-1071. https://doi.org/10.1007/s00227-014-2399-

834   5

835 Picard toolkit. (2019). In *Broad Institute, GitHub repository*. Broad Institute.

836   https://broadinstitute.github.io/picard/

837 Rafajlović, M., Rambla, J., Feder, J. L., Navarro, A., & Faria, R. (2021). Inversions and genomic

838   differentiation after secondary contact : When drift contributes to maintenance, not loss,

839   of differentiation. *Evolution*, *75*(6), 1288-1303. https://doi.org/10.1111/evo.14223

840 Rieseberg, L. H. (2001). Chromosomal rearrangements and speciation. *Trends in Ecology &*

841   *Evolution*, *16*(7), 351-358. https://doi.org/10.1016/s0169-5347(01)02187-5

842 Rundell, R. J., & Price, T. D. (2009). Adaptive radiation, nonadaptive radiation, ecological

843   speciation and nonecological speciation. *Trends in Ecology & Evolution*, *24*(7), 394-399.

844   https://doi.org/10.1016/j.tree.2009.02.007

845 Rundle, H. D., & Nosil, P. (2005). Ecological speciation. *Ecology Letters*, *8*(3), 336-352.

846   https://doi.org/10.1111/j.1461-0248.2004.00715.x

847 Schaal, S. M., Haller, B. C., & Lotterhos, K. E. (2022). Inversion invasions : When the genetic

848   basis of local adaptation is concentrated within inversions in the face of gene flow.

849   *Philosophical Transactions of the Royal Society B: Biological Sciences*, *377*(1856),

850   20210200. https://doi.org/10.1098/rstb.2021.0200

851 Schluter, D. (2000). *The Ecology of Adaptive Radiation*. OUP Oxford.

852 Schluter, D., & Conte, G. L. (2009). Genetics and ecological speciation. *Proceedings of the*

853   *National Academy of Sciences*, *106*(supplement_1), 9955-9962.

854   https://doi.org/10.1073/pnas.0901264106

855 Schluter, D., & Rieseberg, L. H. (2022). Three problems in the genetics of speciation by

856   selection. *Proceedings of the National Academy of Sciences*, *119*(30), e2122153119.

857   https://doi.org/10.1073/pnas.2122153119

858 Schumer, M., Cui, R., Rosenthal, G. G., & Andolfatto, P. (2015). Reproductive Isolation of Hybrid

859   Populations Driven by Genetic Incompatibilities. *PLOS Genetics*, *11*(3), e1005041.

860   https://doi.org/10.1371/journal.pgen.1005041

861 Silva, G., Cunha, R. L., Ramos, A., & Castilho, R. (2017). Wandering behaviour prevents inter

862   and intra oceanic speciation in a coastal pelagic fish. *Scientific Reports*, *7*(1), Article 1.

863   https://doi.org/10.1038/s41598-017-02945-0

864 Silva, G., Horne, J. B., & Castilho, R. (2014). Anchovies go north and west without losing

865   diversity : Post-glacial range expansions in a small pelagic fish. *Journal of Biogeography*,

866   *41*(6), 1171-1182. https://doi.org/10.1111/jbi.12275

867 Simon H. Martin. (2018). *Genomics_general*. https://github.com/simonhmartin/genomics_general

868 Small, S. T., Costantini, C., Sagnon, N., Guelbeogo, M. W., Emrich, S. J., Kern, A. D., Fontaine,

M. C., & Besansky, N. J. (2023). Standing genetic variation and chromosome differences

drove rapid ecotype formation in a major malaria mosquito. *Proceedings of the National*

Academy of Sciences of the United States of America, *120*(11), e2219835120.

https://doi.org/10.1073/pnas.2219835120

Todesco, M., Owens, G. L., Bercovich, N., Légaré, J.-S., Soudi, S., Burge, D. O., Huang, K.,

Ostevik, K. L., Drummond, E. B. M., Imerovski, I., Lande, K., Pascual-Robles, M. A.,

Nanavati, M., Jahani, M., Cheung, W., Staton, S. E., Muños, S., Nielsen, R., Donovan, L.

A., … Rieseberg, L. H. (2020). Massive haplotypes underlie ecotypic differentiation in

sunflowers. *Nature*, *584*(7822), Article 7822. https://doi.org/10.1038/s41586-020-2467-6

Van der Auwera, G. A., Carneiro, M. O., Hartl, C., Poplin, R., del Angel, G., Levy-Moonshine, A.,

Jordan, T., Shakir, K., Roazen, D., Thibault, J., Banks, E., Garimella, K. V., Altshuler, D.,

Gabriel, S., & DePristo, M. A. (2013). From FastQ Data to High-Confidence Variant

Calls : The Genome Analysis Toolkit Best Practices Pipeline. *Current Protocols in*

Bioinformatics, *43*(1), 11.10.1-11.10.33. https://doi.org/10.1002/0471250953.bi1110s43

Weisenfeld, N. I., Kumar, V., Shah, P., Church, D. M., & Jaffe, D. B. (2017). Direct determination

of diploid genome sequences. *Genome Research*, *27*(5), 757-767.

https://doi.org/10.1101/gr.214874.116

Wellenreuther, M., & Bernatchez, L. (2018). Eco-Evolutionary Genomics of Chromosomal

Inversions. *Trends in Ecology & Evolution*, *33*(6), 427-440.

https://doi.org/10.1016/j.tree.2018.04.002

Whitfield, A. (1994). An estuary-association classification for the fishes of southern Africa. *South*

African Journal of Science, *90*, 411-417.

Wilder, A. P., Palumbi, S. R., Conover, D. O., & Therkildsen, N. O. (2020). Footprints of local

adaptation span hundreds of linked genes in the Atlantic silverside genome. *Evolution*

Letters, *n/a*(n/a). https://doi.org/10.1002/evl3.189

Zarraonaindia, I., Iriondo, M., Albaina, A., Pardo, M. A., Manzano, C., Grant, W. S., Irigoien, X.,

& Estonba, A. (2012). Multiple SNP Markers Reveal Fine-Scale Population and Deep

Phylogeographic Structure in European Anchovy (Engraulis encrasicolus L.). *PLOS*

ONE, *7*(7), e42201. https://doi.org/10.1371/journal.pone.0042201

Zhang, L., Reifová, R., Halenková, Z., & Gompert, Z. (2021). How Important Are Structural

Variants for Speciation? *Genes*, *12*(7), 1084. https://doi.org/10.3390/genes12071084

Zheng, X., Levine, D., Shen, J., Gogarten, S. M., Laurie, C., & Weir, B. S. (2012). A high-

performance computing toolset for relatedness and principal component analysis of SNP

data. *Bioinformatics (Oxford, England)*, *28*(24), 3326-3328.

https://doi.org/10.1093/bioinformatics/bts606