Sex-, development-, and nutrition-dependent expression of a major age-at-maturity gene in Atlantic salmon

Eirik Ryvoll Åsheim[1,2,3], Paul Vincent Debes[1,4], Andrew House[1,2,3], Petra Liljeström[1,3], Annukka Ruokolainen[1], Morgane Frapin[1], Iikki Donner[1], Ehsan Pashay Ahi[1], Jaakko Erkinaro[5], Jukka-Pekka Verta[1,6]*, Craig R Primmer[1,2]*

Organismal and Evolutionary Biology Research Programme, Faculty of Biological and Environmental Sciences, University of Helsinki, 00014 Helsinki, Finland.
Institute of Biotechnology, Helsinki Institute of Life Science (HiLIFE), University of Helsinki, Helsinki 00014, Finland.
Lammi Biological Station, Faculty of Biological and Environmental Sciences, University of Helsinki, 16900 Lammi, Finland.
Department of Aquaculture and Fish Biology, Hólar University, Sauðárkrókur 550, Iceland.
Natural Resources Institute Finland (LUKE), 90014 Oulu, Finland.
Current address: Faculty of Biosciences and Aquaculture, Nord University, Norway

*Authors contributed equally

ABSTRACT

Sexual maturation is a key process in the life history of an organism and is influenced by both genetic and environmental factors. In Atlantic salmon (Salmo salar), alternative alleles at the gene vgll3 have been found to associate with a large proportion of the variation in maturation age, yet the molecular and physiological mechanisms of this association are still largely unknown. Here, we investigated the mechanisms by which genetic variation in this major age-at-maturity gene translated into phenotypic variation. We established a time series of gonadal vgll3 mRNA expression in male and female Atlantic salmon in a common garden setting, starting at the prepubertal stage and spanning multiple spawning seasons. We found that vgll3 mRNA expression is reduced in early-maturing males and females. However, for the second spawning season, vgll3 expression increased with gonadal development in males, but eventually decreased in females. Expression of vgll3 had context-dependent correlations with expression of amh and lhr. We also found signs of reduced testicular vgll3 expression in a low-fat feed treatment. By significantly widening the scope of our knowledge on temporal vgll3 expression dynamics, our results exemplify that major-effect genes may use different pathways in sexes but achieve similar outcomes.

INTRODUCTION

Sexual maturation is a process that spans a considerable period of an organism's growth and development. This process is also tightly linked with the allocation of resources to reproduction, thus having a central role in determining organismal life history and ecology (Hutchings, 2021). Variations in the maturation process manifest themselves in different forms, one of which is the age at maturity – the amount of time an organism spends from its conception until it is reproductively mature. Atlantic salmon display remarkable variation in age at maturity, ranging from one to ten years (Hutchings & Jones, 1998). In European populations, a large fraction of this variation in age at maturity has been linked to variation at a single genetic locus – *vgll3* – for both males and females (Barson et al., 2015). Although such large-effect loci have been discovered for a range of life-history traits in various organisms (Johnston et al., 2013; Küpper et al., 2016; Lamichhaney et al., 2016; Pearse et al., 2019; Troth et al., 2018), knowledge on the functional connections (molecular and physiological) between their genotypes and respective phenotypes is largely lacking. Consequently, much remains to be learned about the broader implications of these genes, including other phenotypic effects, environmental interactions, and maintenance of genetic variation. The link between *vgll3* and age at maturity is a notable exception to this pattern, as several facets of the processes connecting *vgll3* genotype and age at maturity phenotype have been uncovered since its initial discovery.

In Atlantic salmon, two alleles of *vgll3* associate with either early (*vgll3**E) or late (*vgll3**L) maturation. This association was initially observed in wild European Atlantic salmon males and females (Barson et al., 2015), and has subsequently been confirmed in multiple laboratory common-garden studies (Åsheim et al., 2023; Debes et al., 2021; Sinclair-Waters et al., 2022; Verta et al., 2020). Furthermore, *vgll3* has been shown to have pleiotropic phenotypic effects, with alternative genotypes being associated with body condition (Debes et al., 2021; House et al., 2023), aerobic scope (Prokkola et al., 2021), as well as aggressive behaviour (Bangura et al., 2022). At the molecular level, *vgll3* genotype has been associated with differing expression of genes relating to the HPG axis in the pituitary and in the testis (Ahi et al., 2022, 2023), and *vgll3* alleles have been associated with alternative splicing (Verta et al., 2020). Correlated RNA expression of other genes has indicated that *vgll3* might be active in gene networks regulating cell differentiation and proliferation, such as the Hippo pathway (Kjærner-Semb et al., 2018; Kurko et al., 2020). The timing and location of *vgll3* expression itself has indicated an important and direct role in testis development (Kjærner-Semb et al., 2018; Verta et al., 2020). Interestingly, *vgll3* has been found to have similar associations in a range of other vertebrates. In humans, genetic variation at- or close to the *vgll3* locus has been associated with age at maturity (Cousminer et al., 2013; Elks et al., 2010; Perry et al., 2014), adiposity-related traits (Nakayama et al., 2017), cancer (Hélias-Rodzewicz et al., 2010), and autoimmune disease

(Liang et al., 2017). In mice, *Vgll3* has been identified as an inhibitor of adipocyte differentiation (Halperin et al., 2013), and is involved in muscle development (Figeac et al., 2019; Mielcarek et al., 2009) and testis development (McDowell et al., 2012). Overall, this strongly suggests a conserved function of *vgll3* across vertebrates, but also shows how *vgll3* seems to have a role that is active in multiple cellular processes, making it hard to pinpoint a single "starting point" for the influence of *vgll3* on age at maturity.

Gonadal tissues and in particular Sertoli and granulosa cells have been identified as key sites of *vgll3* expression in Atlantic salmon testis and ovaries (Kjærner-Semb et al. (2018)). Expression of *vgll3* was also found to be higher in immature- than in mature testes (Kjærner-Semb et al. (2018), Verta et al. (2020)). This has indicated that *vgll3* may function as an inhibitor of testis development. Specifically, based on the down-regulation of *vgll3* and Hippo pathway members in pubertal testis, Kjærner-Semb et al. (2018) hypothesized that *vgll3* controls the early Sertoli cell proliferation which is necessary for the further development of spermatogonial cells (França et al., 2015; Schulz et al., 2010). Significant knowledge gaps remain regarding how *vgll3* is expressed in gonads during the maturation process. For example, earlier studies only cover early-spawning males and no mature females. Thus, *vgll3* expression in relation to gonadal maturation for later-maturing males and females is not known. Further, the data available for *vgll3* RNA expression in wild-derived Atlantic salmon covers only early parr maturation (1st-year maturation, Kurko et al., 2020; Verta et al., 2020). Some data are also available for the post smolt stage (Kjærner-Semb et al., 2018), though only from aquaculture strain Atlantic salmon. Finally, some studies have suggested that the influence of *vgll3* on age at maturity might be mediated via effects on adiposity and resource acquisition (Debes et al., 2021; Halperin et al., 2013; House et al., 2023), but there have been no studies so far looking at how nutrient availability (or other environmental factors) influence *vgll3* RNA expression.

Here, we sought to expand the knowledge of the role of *vgll3* during the maturation process by quantifying its gonadal mRNA expression over multiple years in a single experimental cohort. The time series presented here covers gonadal *vgll3* expression levels in wild-derived Atlantic salmon, starting at the immature stage (second spring, two years old) and spanning multiple subsequent spawning seasons for both female and male individuals (up to four years old). This study also included a feed nutrition treatment designed to test if the influence of *vgll3* varies with environmental variation in nutrient availability. We found patterns of expression supporting previous findings of gonadal *vgll3* expression being associated with delayed maturation in males and found the same pattern for early-maturing females. Unexpectedly, we found striking differences in patterns of expression over gonadal development for third- and fourth-year spawners. These results suggest a different mode of action in testes and ovaries and provides a basis for development of new hypotheses and research on the mechanistic pathways of *vgll3* associated with maturation.

METHODS

Experimental design

This study used Atlantic salmon (*Salmo salar*, L. 1758) from the same rearing experiment as described in Åsheim et al. (2023). In brief, parents with known homozygous *vgll3* genotypes were crossed in late October 2017 to create a series of 2 x 2 factorials (one of each sex of each homozygous *vgll3* genotype), resulting in one family with all offspring having the *vgll3*EE* genotype, one family all *vgll3*LL* and two families all *vgll3*EL*. In total, 17 factorials were created using 67 parental individuals, resulting in 68 unique crosses (Figure 1). At first feeding, the fish were divided evenly (for families) among 12 circular flow-through tanks (1.00 m high, 2.77 m wide, up to 5.4 m^3) at Lammi Biological Research Station (University of Helsinki, Lammi, Finland). Water was pumped directly from the local lake Pääjärvi, and thus the water temperatures in the tanks closely followed the daily and seasonal lake water variation. The 12 tanks were divided among two temperature treatments, using a heat exchange system to divide the incoming water into a cold and a warm temperature treatment (2°C difference). The current study only used fish from the warm treatment, thus using fish from six of the 12 tanks, which experienced temperatures following those of the lake plus approximately 1°C. Note that in March 2021, after completion of the experiment including both the warm and cold temperature treatments, the cold-temperature tanks were cleared out and individuals from each of the six warm-treatment tanks were divided among 12 tanks for the final year of the experiment (Figure 1 B).

To study potential gene-environment interactions between *vgll3* genotype and nutrition, tanks were subjected to one of two feeding treatments, "control" and "low-fat". At the beginning of the study (3/2018), all fish were fed ad-libitum using body-size-matched pellets of commercial fish feed ("Hercules", Raisioaqua, Raisio, Finland). In July 2019 (summer of the second year post-fertilization), the feed for half of the tanks was changed to a custom-made fat-reduced feed of the same brand and pellet type. Thus, from July 2019, fish were either fed the control feed (17-26% fat, 18.10-20.40 kj g^{-1} depending on pellet size), or the low-fat feed (12-13% fat, 17.25 kj g^{-1}).

To keep track of fish identities, all fish were marked with a passive integrated transponder tag (PIT tag) at the first measurement in April 2019. Tags were inserted into the abdominal cavity, slightly behind (posterior to) their right-side pectoral fin. Additionally, a small fin clip was taken from their caudal fin for genotyping, sex determination, parental assignment and *vgll3* genotype confirmation (as in Debes et al., 2021).

Figure 1. Experimental design. Shows the crossing design (A) the allocation of fish used in this study to different tanks and treatments (B), and a photograph (C) of the experimental system (taken in March 2020).

Fish were reared until March 2022, after which almost all male and female fish had matured. At this point, all fish were euthanized by anaesthetic overdose (MS-222, Tricaine methanesulfonate, 0.250 g L^{-1}, sodium bicarbonate buffered).

The fish used in this study were offspring of first-generation hatchery fish. The parental broodstock is used for supplementary stocking of Finnish rivers and has its name "Neva" from the river where it was mainly sourced in the 1980s. Since that time, it is renewed every few years using fish that have completed a seaward migration after their release. Thus, the grandparents of our source experimental population had completed a full marine migration, while all the parents had experienced the same environment in the hatchery (thus minimizing potential maternal/epigenetic variation, such as the environmental variation experienced by migrating individuals).

Tissue sampling

Fish were measured and sampled four times per year, starting in April 2019 and continuing until the end of the study in March 2022, resulting in 13 time points (see Table 1 for an

overview). At these time points, phenotypic traits were recorded for each individual (details below). For each tank, eight individuals were euthanized for tissue sampling one day after measurements, except for time point 3, where they were sampled seven to 12 days after, and time point 4, where they were sampled on the same day. Measuring and sampling at each time point was conducted over a period of 14-16 days.

Table 1. Sample overview. Overview of tissue-sampled individuals that were included in this study. Sample sizes are shown per sex and *vgll3* genotype at each time point. Note that this is a subset of the total amount of sampled fish, as both heterozygous *vgll3* genotypes were sampled (both EL and LE) from each tank (n=6) and sex (n=2), but only half were included in the analysis (randomly selected). No fish were sampled for tissues at time point 1 (only measurements).

Time point	Year	Month	n	Male	EE	EL	LL	Female	EE	EL	LL
2	2019	July	2	11	3	4	4	14	6	5	3
3	2019	October	3	18	5	6	7	17	5	6	6
4	2019	November	3	18	6	6	6	18	6	6	6
5	2020	February	3	17	5	6	6	19	6	6	7
6	2020	May	3	16	5	6	5	17	5	6	6
7	2020	August	3	19	7	6	6	19	7	6	6
8	2020	November	3	15	5	5	5	17	6	6	5
9	2021	February	3	16	5	6	5	17	4	6	7
10	2021	May	3	20	6	8	6	19	5	9	5
11	2021	August	3	17	6	6	5	19	6	7	6
12	2021	November	3	18	6	6	6	17	6	6	5
13	2022	February	3	19	6	7	6	17	6	6	5

Phenotypes recorded were body mass, body length, maturation status, and whether fish showed external signs of smoltification (migration phenotype). Male maturation status was checked by stroking the abdomen towards the vent, recording any fish releasing milt as mature. Females were recorded as mature if releasing eggs. Migration phenotype, being either migrant (smolt) or resident (parr), was scored based on silvering of color and loss of parr marks. Individuals were recorded as having smolted from the time point following the last recording of resident (parr) characteristics. For measurements, fish were netted out of their tank and placed in an aerated anaesthetic bath (MS-222, 0.125 g L^{-1}, sodium bicarbonate buffered) within 1°C of the tank temperature. After fish stopped responding to touching or holding, their weights were measured down to the nearest 0.01 g (April and July 2019) and subsequently 0.1 g using a digital scale (Scout STX222 or STX6201, Ohaus, Parsippany, USA). Fork length (snout to fork of tail) was measured using a digital fish-measurement board (DCS5, Big Fin Scientific, Austin,

TX, USA). Finally, migration and maturation phenotypes were recorded before the fish were
returned to their tank.

A subset of fish was selected for tissue sampling at each measurement time point. From each
tank, one individual of each sex-*vgll3* genotype combination was selected for sampling, avoiding
selecting multiple fish from the same family within tanks, totalling 2 x 4 x 6 = 48 sampled
individuals per measurement period. At the first sampling time point (time point 2, Table 1),
the sex and genotypes of the fish were unknown, so 10 fish were randomly sampled from each
tank. For all time points following time point 4, fish selected for sampling were placed directly
into a separate holding tank after measurements and were taken from this holding tank for
sampling on the following day. On the preceding time points (2, 3 and 4), fish to be sampled
were netted from their original holding tank.

At sampling, fish were first euthanized using an MS-222 overdose (0.250 g L^{-1}, sodium
bicarbonate buffered), and were then immediately dissected for collection of tissue samples.
Sampled gonads were immediately placed in 1.5 ml Eppendorf tubes and flash-frozen in liquid
nitrogen before being stored at -80°C. Gonads were sampled whole, and in separate tubes for
as long as their size allowed this. For gonads that were too large to fit the 1.5 ml Eppendorf
tubes, only the tip of the gonad (anterior end) was sampled. Mature gonads were sampled
without stripping eggs or sperm, but sperm was removed from samples before RNA extraction.

RNA extraction and quantification

Before RNA extraction, testis samples were stripped for sperm by thawing them in room-
temperature phosphate-buffered saline (PBS) for ~1 minute and then squeezing and rolling the
tissues on a piece of paper tissue (coffee filter paper) until only solid tissue remained. This
procedure was done on all testis samples regardless of maturation status and lasted 2-3
minutes from start of thawing.

To extract total RNA, we used the NucleoSpin 96 RNA kit (MACHEREY-NAGEL, Dueren,
Germany). Samples were homogenized in RA1 buffer with added DTT (final concentration 10
mM) using a Bead Ruptor Elite (Omni Inc, Kennesaw, GA), 2 ml or 7 ml tubes, and 2.4 mm steel
beads. Residual DNA was removed with TURBO DNase (Invitrogen, Waltham, MA). To avoid any
batch effects confounding other variables, samples were randomly divided among the five
plates used for RNA extraction and subsequent RT-ddPCR (reverse transcription digital droplet
PCR).

We measured RNA concentrations using ThermoFisher Quant-it reagents and diluted the
samples to a 10 ng/µl working concentration. Dilution concentrations were measured with
ThermoFisher Qubit High Sensitivity reagents and were re-adjusted if needed. Final
concentrations were verified with Qubit HS. To quantify the absolute level of expression of *vgll3*

(chromosome 25), *amh (*anti- müllerian hormone*)*, and *lhr* (luteinizing hormone receptor), we
used one-step RT-ddPCR (BioRad) and TaqMan probes (Supplementary table 1). For allele-
specific quantification of *vgll3*, we used TaqMan probes specific to the *Early* and *Late vgll3*
alleles as in Verta et al. (2020). We used ~40 ng of total RNA as template for *vgll3*, and ~20 ng
for *amh* and *lhr*.

To extract counts of detected mRNA molecules, RT-ddPCR results were analysed in *BioRad*
Quantasoft v.1.7.4.0917. For most samples, we used an automatic selection of signal thresholds,
and used manually set thresholds for samples where the software was not able to select them
automatically. All 1-D amplitude plots were visually checked, and threshold settings verified.
We excluded samples with less than 5000 accepted droplets (unreliable estimates) as well as
samples with zero negative droplets (extreme values). Finally, we normalized the counts of
detected *mRNA* molecules to the amount of total template RNA so that mRNA expression for
vgll3, amh, and lhr was expressed as number of copies per ng of total RNA.

mRNA counts for ovaries were corrected for 5S rRNA amounts since the amount of this
ribosomal RNA changes during development and can have a significant influence on the total
RNA count (Shen et al., 2017). This was corrected by determining the proportion of 5S rRNA
from electropherograms acquired using a 2100 Agilent Bioanalyzer instrument, and then
subtracting this amount from the total RNA before calculating the number of copies (*vgll3*, *lhr*,
amh) per ng RNA. Thus, expression for ovaries is expressed as total RNA excluding 5S rRNA.

Overview of animals and samples.
The experiment was conducted under an animal experiment permit granted by the Finnish
Project Authorization Board (Permit ESAVI/2778/2018)

Over the entire study period, 562 fish were tissue-sampled. The 417 individuals analysed for
this study are a subsample of this due to only half of the *vgll3* heterozygotes being included.
Heterozygotes were randomly selected for each combination of tank, time-point, and sex (Table
1). For some statistical tests, the sample size was smaller than the original number of
individuals included due to missing data. Overall, the number of individuals with missing mRNA
expression data was 17 for *vgll3*, 27 for *lhr*, and 49 for *amh*.

In 2019, the starting number of Neva individuals in the warm-temperature treatment (the
subset of the source experimental population used here) was 1162 individuals, and 411
remained at the conclusion of the experiment in 2022. Of the 751 fish which died throughout
the experiment, 255 (22%) were unplanned mortalities which were not part of any sampling
or other procedure.

Figure 2 shows the number of individuals in different stages of maturation for each time point.
Most sampled individuals had undergone smoltification. Exceptions were for two, two and

three females in time points 2, 3 and 4, respectively, and two, three and one males in time points
3, 4 and 10, respectively having parr characteristics.

Statistical methods

Our statistical analysis approach aimed to test for *vgll3* mRNA expression differences between
fish at different developmental stages and to explore seasonal or developmental patterns in
expression of *vgll3* mRNA (and associations with *vgll3* genotype). Additionally, having access
to mRNA expression levels of *amh* and *lhr*, we explored potential correlations between *vgll3*
and *lhr / amh* expression. Finally, we tested if the association between *vgll3* genotype and age
at maturity was detectable in our sample of individuals. For all comparisons via models (except
correlations), we included feeding treatment as an explanatory factor to assess if this affected
any of the expression patterns.

Males and females started spawning in different years. To simplify comparisons between
sexes, we analysed the data relative to the first spawning season. Thus, we refer to the first and
second spawning seasons, which were in 2019 and 2020 for males, and 2020 and 2021 for
females (Figure 2, lines S1 and S2).

Description of analysis

Gonads were classified into one of three stages of development: 1) "undeveloped" (immature)
testes were identified as those resembling a thin transparent string running dorsally along the
entire length of the body cavity. In females, we identified undeveloped ovaries as those having
a length less than $1/4^{th}$ of the body cavity. 2). "Developing" testes were classified as those being
thicker than the undeveloped ones and having a white or grey hue. Developing ovaries were
identified as those being longer than $1/4^{th}$ of the body cavity. 3) "Mature" testes and ovaries
were identified as those releasing eggs or sperm directly from the fish. Figure 2 shows the
number of fish in each category for each sampling time point.

For all models described below, the IDs of each individual's dam, sire, and family were
included as random effects to account for parental- and family effects. Tank ID was also
included as a random effect to account for potential tank effects. Males and females were
analysed in separate models.

Model 1: Here we tested for differences in *vgll3* expression between maturation stage and
feed treatment in the first spawning season. We simplified the grouping of the gonads into two
categories, being either undeveloped or developing/mature. We then specified models testing
if the log of standardized *vgll3* mRNA expression (counts ng^{-1} RNA) changed with maturation
status, feeding treatment, or month of sampling (November or February). Interactions between
feed and both maturation status and month of sampling were also included.

Model 2: Here we tested the observed increase in *vgll3* expression in the time point leading
up to the second spawning season in developing gonads (Figure 2, lines "Pre S2"). To do this,
we specified models testing if the log of standardized *vgll3* expression (counts ng^{-1} RNA)
changed with month of sampling (2, 5, 8), feeding treatment, and whether the sampled
individual had been recorded as mature on a previous occasion. Interactions between month of
sampling and both feed and previous maturation status and were also included.

Model 3: To test for differences in maturation probability in the first spawning season
between *vgll3* genotypes and feed treatments, we used generalized linear mixed effect models
with Bernoulli error distribution and logit link function. Maturation was categorized in the
same two categories as in model 1, and the model tested if maturation probability differed
between *vgll3* genotypes and feed treatments. The body weight of each individual in the
proceeding spring, the time point where growth is expected to have its most significant effect
on future maturation (Rowe et al 1991), was included in the model to account for body-size
differences. Interactions between feed and both maturation status and month of sampling were
also included.

Correlations: Expression correlations between *vgll3* and *amh* or *lhr* were estimated using
Pearson correlations on log-adjusted standardized expression levels (counts ng^{-1} RNA).
Correlations were estimated separately on data grouped by sex, maturation status, and season
of the year (spring, summer, autumn, winter). In the final month before the second spawning
season, which was the time point where we observed the largest variance in gonadosomatic
index (GSI) in developing testes and ovaries, we checked whether this variation was associated
with expression of *vgll3* using Pearson correlations between log gene expression and GSI.

To verify that the feeding treatment had an impact on fish growth, we performed *t*-tests on
body condition between fish in the control- and low-fat feed treatment for each measurement
time point. Body condition was calculated as body mass (g) divided by the cube of the body
length (cm) multiplied by 10 000 (Fulton's K)(Ricker, 1975). This analysis confirmed that body
conditions between the two treatments were significantly different at every time point after
May 2020 (the low-fat feed treatment started in July 2019) (Supplementary figure 3).

For all gene expression level analyses, we log-transformed expression level of *vgll3*, *amh*, and
lhr (copies ng^{-1} RNA) to account for heteroscedasticity (higher variance at higher expression
levels). GSI was calculated as percent gonad mass of total body mass.

Technical model descriptions
The following formulae describe the models introduced above, showing the overall form of
the models, specifically which variables were included as explanatory and response variables,

and which interactions were included. Each model was fitted separately for each sex (testes,
ovaries). Not shown: parameters b_i for each term (i) of the models (except R and e).

Model 1: *vgll3* expression differences between maturation stages in the first spawning season.

$$316 \qquad log(count_{vgll3}) = M + T_{S1} + F + FM + FT_{S1} + R + e$$

317 Model 2: Gradual change in *vgll3* expression leading up to the second spawning season.

$$318 \qquad log(count_{vgll3}) = T_{PreS2} + F + Pr + FT_{PreS2} + PrT_{PreS2} + R + e$$

319 Model 3: Effects on maturation probability

$$320 \qquad log(odds_{maturation}) = G_{vgll3} + W_S + F + FG_{vgll3} + FW_s + R + e$$

Variable	Explanation
$count_{vgll3}$	*vgll3* expression (mean-centred, SD-scaled log of expression, copies ng-1 RNA).
$odds_M$	Odds of gonads having started maturing or not.
M	Gonad development having started or not (0=no, 1=yes).
T_{S1}, T_{PreS2}	Month of sampling, during the first spawning season (S1: 0=Dec, 1=Feb), or leading up to the second spawning season (PreS2: 0=Feb, 3=May, 6=Aug).
F	Feed treatment (0=low fat, 1=control).
Pr	Whether the sampled individual had been recorded as mature earlier or not (0=not previously mature, 1= previously mature).
G_{vgll3}	*vgll3* genotype (LL=-1, EL=0, EE =1).
W_s	Body weight in the preceding spring (mean-centred, SD-scaled log of weight).
$R + e$	The random effects on the intercept (dam ID, sire ID, their interaction, and Tank ID) and residual error.

Bayesian model parameters (b_i, R and e) were estimated using *stan*, which was interfaced via
the *R* package *brms*. All models were run for 5 000 iterations in four chains, discarding the first
500 iterations as warmup, resulting in 18 000 posterior samples of each parameter. All model
fits were verified using a visual posterior predictive check (function *pp_check* in R package
brms) and verified for influential points by inspecting pareto k diagnostic values (function *loo*
in R package *loo*) (Vehtari et al., 2017). Three models had high numbers of influential data
points (points for which k > 0.7), which were male model 2 (3 of 42 points with k > 0.7), female
model 2 (6 of 26 points with k > 0.7), and male model 1 (8 of 35 points with k > 0.7). For these
models, we tested alternative versions without random effects for parental- and family ID,
which reduced the number of influential points down to 0, 2, and 1, respectively, indicating that
most of these influential points were affecting the estimates of parental- and family effects. We
compared effect-size estimates of the models including- and excluding parental- and family

effects, found that the differences were small, and chose to keep the models including parental- and family effects.

We chose relatively non-informative priors for all parameters. We used normally distributed priors with a mean of 0 and SD of 1 for the intercept and for all effect-size parameters. For error terms, we used normally distributed priors with a mean of 0.2 and SD of 0.5.

A summary of the models' fixed-effect parameter estimates (b_i) is presented in Supplementary Figure 1. Where interactions were non-significant (95% credible intervals including zero), we simplified parameter estimates by calculating unconditional (marginal) estimates of the main effects (Supplementary figure 1).

All statistical analysis was done in the *Rstudio* v2022.7.2.576 (RStudio Team, 2020) software environment running *R* v4.2.2 (R Core Team, 2021) and *Rstan* v2.21.8 (Stan Development Team, 2022) R packages *brms* v2.18.0 (Bürkner, 2017) was used for creating *Rstan* models, *loo* v2.5.1 (Vehtari et al., 2022) for inspecting pareto k diagnostic values, *bayesplot* v1.10.0 (Gabry & Mahr, 2022) for posterior predictive checks, *ggplot2* v3.4.0 (Wickham, 2016) for producing plots, and *tidyverse* v1.3.2 (Wickham et al., 2019) for data workflow.

RESULTS

Expression of *vgll3* across 3 years of development

We measured the gonadal expression of *vgll3* mRNA (copies ng^{-1} RNA) in different seasons over multiple years, in both male and female Atlantic salmon. We found that *vgll3* expression was elevated in both undeveloped testes and ovaries in the first spawning season, but also that expression patterns were more complex for the following spawning seasons.

In the first spawning season, in both testes and ovaries (Figure 2, lines S1), gonadal *vgll3* expression was higher in undeveloped gonads than in those showing signs of maturation (i.e., gonads enlarged or fully mature). For ovaries, *vgll3* expression was 1.59 times higher [1.05, 2.34] (brackets indicate 95% credible intervals) in undeveloped than in developing or mature ovaries, with a predicted mean of 44.12 copies ng^{-1} RNA in developing or mature ovaries, and 69.88 copies ng^{-1} RNA in undeveloped ovaries (posterior predictions). For testes, *vgll3* expression was 2.05 times higher [1.44, 2.89] in undeveloped testes (same comparison), with a predicted mean of 40.05 copies ng^{-1} RNA for developing or mature testes, and 81.98 copies ng^{-1} RNA in undeveloped testes.

Figure 2. Gonadal maturation and *vgll3* expression over time. Gonadal expression of *vgll3* mRNA over the entire study period, measured as copies ng^{-1} of RNA in ovaries (A) and testes (B). Colours and shapes correspond to maturation status of the gonads: purple circles for undeveloped gonads (no sign of enlargement), orange squares for developing gonads (some sign of enlargement), and red diamonds for completely mature gonads (releasing sperm or eggs). Lines connect means between time points with at least two samples, and error bars indicate SE. Numbers below points indicate sample sizes for undeveloped, developing, and mature gonads. The y-axes are $\log_e$ scaled.

In the second spawning season (Figure 2, lines S2), we were unable to directly compare maturing and undeveloped individuals as most individuals were maturing. However, looking at the period leading up to the second spawning season (Figure 2, lines Pre S2), there was a trend of increasing *vgll3* expression up towards spawning. In testis, *vgll3* expression increased 1.20-fold [1.21, 1.40] for each month going from February (39.22 copies ng^{-1} RNA) to August 2020 (86.46 copies ng^{-1} RNA). For females, ovary *vgll3* expression increased 1.12-fold [1.02, 1.24] at each time point from February (33.1 copies ng^{-1} RNA) to May to August 2021 (69.8 copies ng^{-1} RNA). After this period, as maturation occurred, testis *vgll3* expression remained high, while ovary *vgll3* expression first dropped in November, but then increased again in the following February (Figure 2, lines S2). Thus, *vgll3* expression increased as testes and ovaries were developing, which contrasts with the first season where the undeveloped gonads had the

highest *vgll3* expression. Notably, in testes, this pattern was dependent on feed treatment and
was only present in the control feed treatment (detailed below). Whether fish had matured in
the first spawning season had no detectable effect on *vgll3* expression in the period leading up
to the second spawning season. In this analysis, 17 of 42 male individuals- and 9 of 26 female
individuals had matured previously.

GSI and *vgll3* correlated significantly at the last time-point before the second spawning season,
which was when GSI had been increasing the most (August, Figure 3 A and B). However, this
correlation was sex dependent. In testes, a higher GSI corresponded to a higher expression of
vgll3 (Pearson's R = 0.79, p < 0.001), while expression of *vgll3* decreased with GSI in ovaries (R
=-0.92, p = 0.001).

Figure 3. GSI and expression of *vgll3*. Relationship between gonadosomatic index (GSI, % gonad
mass of whole body-mass) and gonadal *vgll3* expression (copies ng^{-1} RNA) in ovaries (A) and testes (B)
at the three sampling time-points leading up to the second spawning season. Data are only shown for
developing gonads (i.e., not fully mature, but showing some physical sign of development). Lines, shown
only at the last time points, are for illustrative purposes and represent simple linear regression models
of relationships between *vgll3* and GSI. N indicates sample size, R represents the degree of correlation
(Spearman) and p the statistical significance. Asterisks (*) indicate p values below 0.05. The y-axes are
log$_e$ scaled.

The low-fat feed treatment, which started in July 2019, had a negative effect on body condition
from August 2020 (Supplementary figure 3). Feed effects on *vgll3* expression were detected,
but were dependent on time point and sex. In the first spawning season, there were no
detectable effects of feed treatment on *vgll3* expression. However, in the period leading up to

the second spawning season (as mentioned above), there was a significant effect of feed treatment on testicular *vgll3* expression. While there was an increase in *vgll3* expression over these three time points for the control-feed individuals, there was no statistically significant change over time for the low-fat treatment individuals. Specifically, for the low-fat treatment individuals, the estimated change in *vgll3* expression per month was 0.95-fold [0.80, 1.14]. Over the same time period (leading up to second spawning season), we detected 1.58 times higher [0.85, 2.80] expression of *vgll3* in the testes of control-feed treatment individuals, compared to the low-fat feed treatment. Given the 95% CI of this effect includes unity, there is some uncertainty about this effect, but 93.3% of the probability mass of this effect is above 1. Considering the following year (post second spawning season, leading up to the third spawning season), there were signs of a similar expression difference in testes between the feed treatments (Supplementary figure 3). In ovaries, no effects of the feed treatments on *vgll3* expression were detected.

Expression of *amh* and *lhr*

Anti-müllerian hormone (encoded by the gene *amh*) is known to act as an inhibitor of spermatogenesis (Morais et al., 2017; Pfennig et al., 2015). Expression of *amh* is therefore a useful marker for maturation in testes, and we included *amh* expression measurements to gain additional context on the maturation process. As expected, the expression of *amh* was reduced in developing testes in late summer (August) in both years (Figure 4 B), but higher in winters. Ovarian *amh* expression showed clear signs of decreased expression only in fully mature ovaries (Figure 4 A). Interestingly, we detected positive correlations between expression of *vgll3* and *amh* in both testes and ovaries. However, these correlations depended strongly on season and developmental stage. In ovaries (Figure 5 A), these correlations were significant for undeveloped ovaries only in summer and autumn, for developing ovaries only in autumn, and never for mature ones. In testes (Figure 5 B), the correlations between *vgll3* and *amh* were significant for undeveloped testis in summer, autumn, and winter; for developing testes only in summer and autumn; and for mature testes in all seasons where they were present.

The luteinizing hormone receptor (encoded by the gene *lhr*) can be used as a marker of maturation in ovaries as its expression gradually increases towards maturation (Andersson et al., 2013). As expected, *lhr* expression increased towards maturation in ovaries and was highly expressed also in mature testes (Figure 4 C and D). Some significant correlations between expression of *vgll3* and *lhr* were detected. In ovaries (Figure 5 C), correlations between the expression of *vgll3* and *lhr* were more sporadic, and only present in summer for undeveloped and developing ovaries, and in autumn for mature ovaries. In testes (Figure 5 D), expression of *vgll3* and *lhr* correlated positively for undeveloped testes in summer, autumn, and winter; for developing testes in summer and autumn, and for mature testes in autumn and winter.

Figure 4. Gonadal maturation and *amh* and *lhr* expression over time. Gonadal expression of *amh* (A, B) and *lhr* (C, D) mRNA over the entire study period in ovaries (A, C) and testes (B, D), measured as copies ng^{-1} of RNA. Colours and shapes correspond to the maturation status of the gonads: purple circles for undeveloped gonads (no sign of enlargement), orange squares for developing gonads (some sign of enlargement), and red diamonds for fully mature gonads (releasing sperm or eggs). Lines connect means between time points with at least two samples and error bars indicate SE. The numbers on top indicate sample sizes for undeveloped, developing, and mature gonads. The y-axes are log$_e$ scaled.

Figure 5. Correlated expression of *vgll3*, *amh* and *lhr*. Associations between gonadal RNA expression (copies ng⁻¹ RNA) of *vgll3* and either *amh* (A, B) or *lhr* (C, D) in for different seasons (over all years) in ovaries (A, C) and testes (C, D) of different maturation status. Colours and shapes correspond to the maturation status of the gonads: purple circles for undeveloped gonads (no sign of enlargement), orange squares for developing gonads (some sign of enlargement), and red diamonds for fully mature gonads (releasing sperm or eggs). Lines, for illustrative purposes, represent simple linear regression models of relationships between *vgll3* expression and expression of either *amh* or *lhr*. N indicates sample size, R represents the degree of correlation (Pearson) and p the statistical significance for the groups of the same colour. Asterisks (*) indicate $p < 0.05$. The x- and y-axes are $\log_e$ scaled.

Maturation, *vgll3* genotype and feed

The association between *vgll3* genotype and age at maturity is already well established, also

for the males of the source experimental population used in this experiment (Åsheim et al.,

2023). Despite the low sample sizes, we tested for a similar association in the individuals used

in the current study. An association between maturation probability and *vgll3* genotype was

indeed detected for males in the first spawning season, but not for females. In the second

spawning season, almost all individuals matured regardless of genotype. In the first spawning

season, for an average spring body weight, the mean predicted probability of having either

developing or mature testes (vs. undeveloped) was 24.9% and 80.0% for a *vgll3**LL and *vgll3**EE

individuals, respectively [95% CI of difference: 14.4, 79.7]. For females, there was no detectable

effect of *vgll3* genotype on maturation probability in the first spawning season. Using the same

comparison as above for females gave predicted probabilities of 49.1 and 48.9 % for *vgll3**LL

and *vgll3**EE individuals [95% CI of difference: -47.3, 47.4]. We detected no significant effects of

the feed treatments on maturation probability in the first spawning season for neither males

nor females.

Genotype and expression of *vgll3*

Studying expression variation between different *vgll3* genotypes within the maturation

categories is challenging as there are already *vgll3*-maturation and maturation-expression

associations present. To examine possible differences, we plotted *vgll3* expression over time

for different *vgll3* genotypes separately for each maturation category (undeveloped and

developing/mature) and sex (Supplementary figure 2). However, in both testes and ovaries, we

observed that associations between *vgll3* genotype and expression were overall weak and

differed between time points; no genotype (EE/EL/LL) was consistently expressed higher or

lower than the others. Thus, over the full time series, we could not identify consistent patterns

of associations between *vgll3* genotype and mRNA transcript abundance.

DISCUSSION

We have presented for the first time a continuous time series of gonadal development and

vgll3 expression in both male and female Atlantic salmon over a 3-year time period, covering

multiple spawning seasons. From these data, we identified several novel aspects of *vgll3*

expression patterns during development and maturation.

Higher *vgll3* expression in the ovaries and testes of individuals with delayed maturation

The observation that *vgll3* expression was higher in immature, undeveloped ovaries in the

first spawning season suggests that the same pattern of *vgll3*-associated inhibition of

maturation is present in both males and females. However, this pattern contrasts to what was

found by Kjærner-Semb et al. (2018), who observed an increase in ovarian *vgll3* expression

going from the oildrop prepubertal stage to the early pubertal stage. Compared to the
expression pattern we found, the pattern observed in Kjærner-Semb et al. (2018) may
correspond to the slight increase in *vgll3* expression we observed going from February to
August after the first spawning season. In males, the association between testicular *vgll3*
expression and maturation status has been suggested to be mediated via a *vgll3*-effect on the
Hippo pathway. In this scenario, *vgll3* competes with Yap1/Taz for access to TEAD elements,
affecting processes related to cellular differentiation (Kjærner-Semb et al., 2018; Kurko et al.,
2020; Verta et al., 2020). The Hippo pathway has been found to associate with both ovarian and
testicular development in other vertebrates (Clark et al., 2022; Kawamura et al., 2013; Lyu et
al., 2016; Sen Sharma et al., 2019; Xiang et al., 2015), and has also been found to be involved in
a range of other developmental processes (Meng et al., 2016). In any case, these results suggest
that *vgll3* expression is similarly involved in the early maturation of males and females, at least
in the sense that it involves a higher level of *vgll3* expression for immature individuals.

Expression during development reflects inhibition of early testis development
Transitioning from the first to the second spawning season, we observed a change in the
pattern of *vgll3* expression. For testes which were immature in the first spawning season, *vgll3*
expression was high but then dropped in the following winter, slowly increasing up towards
the next spawning season, while correlating positively with GSI as testis were gaining mass.
Testicular development happens in sequential stages with regards to the proliferation and
differentiation of spermatogonia and the Sertoli cells encysting them (Clermont, 1972; França
et al., 2015; Schulz et al., 2010). Following this, we interpret from the expression patterns
mentioned above that *vgll3* might be mainly involved in the inhibition of the earlier stages of
testicular development. This mechanism was similarly hypothesized by Kjærner-Semb et al.
(2018) on the grounds of histological observations as well as an observed downregulation of
vgll3 in maturing testes and a further increase in *vgll3* expression in regressing testis (post-
spawning). Here, our data provide further support to this hypothesis by showing that *vgll3*
expression is positively correlated with testis growth in later years, thus increasing not only
after maturation (as in Kjærner-Semb et al., 2018) but also in the process leading up to
maturation. Notably, for mature testes, the mean *vgll3* expression was low during the first
spawning season, but high during the second one. This could indicate that the age at spawning
affects how changes in expression of *vgll3* are timed, or that the influence of *vgll3* during
spawning changes with age or testicular development. Taken together, we hypothesize from
these results that that the earlier stages of testis development involve a downregulation of *vgll3*
expression, while later stages involve a gradual upregulation of *vgll3*, and that the extent or
timing of this upregulation may vary with age.

Sex-specific expression dynamics

While we observed an increase in *vgll3* expression with GSI in testes, the pattern was more complex for ovaries. After the initial drop in *vgll3* expression as the ovaries started to develop, there was a steady increase in its expression, but then a negative association between GSI and *vgll3* expression in the autumn as the ovaries were starting to gain mass. These opposing patterns of *vgll3* expression and GSI between testes and ovaries could indicate that *vgll3* is involved in processes that are differently regulated in males and females. An example of such a process is estrogen (17β-oestradiol) signalling, which in males is high in the early stages of testis development, but in females coincides with the later stages of ovary development (Billard & Breton, 1978; Clelland & Peng, 2009; T. Miura et al., 2003). In testis, estrogen promotes spermatogonial germ cell- and Sertoli cell proliferation in the early phases of testis development (Amer et al., 2001; T. Miura et al., 1999, 2003; Song & Gutzeit, 2003). In ovaries, current knowledge of the role of estrogen is that it plays an important part during the vitellogenic phase of oogenesis (i.e., the phase of secondary oocyte growth)(Clelland & Peng, 2009). Estrogen may also be important at the earliest stages of oogenesis (C. Miura et al., 2007; See also Wootton & Smith, 2014, Ch 5).

Changes in estrogen signalling could involve both changes in estrogen production and changes in the expression of the estrogen receptor (esr1). In humans, expression of *VGLL3* has been negatively linked to expression of *ESR1*, the gene encoding the estrogen receptor ERα (Ma et al., 2022). Notably, ERα has been found to be central for the role of Sertoli cells during the first stage of testis development (spermatogonial stem cell renewal) (T. Miura et al., 2003). Production of estrogen itself, on the other hand, can be inhibited by *amh* via an inhibiting of cyp19a1 (aromatase, or estrogen synthase), an enzyme that converts testosterone to estrogen (Rodríguez-Marí et al., 2005). In this study, we found *vgll3* to have correlated expression with *amh*, so if one assumes a negative association between *vgll3* and estrogen signalling, then it can also be hypothesised that *amh* is involved in the process by inhibiting the testosterone-estrogen conversion. However, this hypothesis does not fit well with the observation that testicular *amh* expression was at its lowest at the same time as estrogen signalling should have been low (Billard & Breton, 1978). Changes in the levels of estrogen and testosterone could potentially also alter behaviours that are modulated by these hormones (See: Borg, 1994; Dey et al., 2010; Fernald, 1976). In this context, it is worth noting that behavioural trials have found a higher frequency of aggressive behaviour in juvenile individuals homozygous for the late *vgll3* genotype (Bangura et al., 2022).

While the hypothetical link between estrogen signalling and *vgll3* expression is intriguing, we underscore that it is at this time highly speculative, and that we include it as an example of how the results of this study could generate further hypotheses regarding the pathways *vgll3* is involved in.

Expression and genotype of *vgll3*

To assess whether some of the effect of *vgll3* may be linked to genotype-specific expression levels, we examined whether any of the genotypes were more highly expressed than the others over the study period. This analysis was done after splitting the samples into groups of either undeveloped or developed/mature, accounting for a potential scenario where maturation status affects *vgll3* expression (reverse causality). Here, we found no significant and consistent association between genotype and expression. Previously, Verta et al. (2020), found a 14% higher expression of *vgll3* exon 2 (same exon as in this study) in *vgll3**LL individuals, but the variation was high with a highly overlapping range of expression between individuals of the two genotypes. Our sample sizes per genotype per time-point were overall low, so it is possible that we did not have sufficient statistical power to detect an effect, if it existed.

Feed effects

The difference in *vgll3* expression between feed treatments, which was only observed in testes in the time periods leading up to the second and third spawning season, provides the first indication that nutritional intake can affect the temporal dynamics of *vgll3* expression. Two potential interpretations of this interaction are 1) the feed treatment could be affecting the speed of development of the testis, thus resulting in different *vgll3* expression levels when measured at the same time point, or 2) pathways regulating *vgll3* expression are integrating the nutritional or energetic state (e.g., lipid storage) of the fish. This mirrors previous findings where we observed an interaction between *vgll3* genotype and body condition in affecting maturation (Åsheim et al., 2023) as well as an analysis indicating that some of the effect of *vgll3* on maturation might be mediated via body condition (Debes et al., 2021). Interestingly, here, the effect of feed treatment on *vgll3* expression was only observed in males, indicating that the relationship between nutrient availability and *vgll3* expression might be sex specific.

Conclusions

This study provides new insights into the temporal patterns of gonadal *vgll3* expression in Atlantic salmon throughout the maturation process, especially filling in knowledge-gaps related to maturation in females. We also demonstrate that differences in feed nutrient content can be reflected in the expression of *vgll3*, providing further insight on the connection between *vgll3*, resource acquisition and age at maturity. Our findings provide starting points for further research on the role of *vgll3* in the maturation process. For example, studies looking at the differences observed between males and females could target mechanisms with known opposite effects between sexes on maturation-related processes. Additionally, while we have uncovered novel patterns in whole-tissue expression of *vgll3*, further studies are necessary to identify underlying mechanisms for these patterns, such as changes in regulation of expression, or changes in proportions of various cell types during development. This could help in identifying new mechanisms related to the function of *vgll3*.

Author contributions:

Conceptualization: CRP, JPV, ERÅ, PVD

Data curation: ERÅ, AR, ID, MF, PVD

Formal analysis: ERÅ, PVD, JP

Funding acquisition: CRP

Investigation: ERÅ, AH, PL, AR, MF, ID, EPA, CRP

Methodology: JPV, PVD, CRP, ERÅ,

Project administration: CRP, JPV,

Resources: CRP, JE

Software: ERÅ, PVD,

Supervision: CRP, JPV,

Visualization: ERÅ,

Writing - original draft: ERÅ, JPV, CRP, PVD, EPA

Writing - review & editing: ERÅ, JPV, CRP, PVD, ID, MF, AR

Funding

Funding for this study was provided by the Research Council of Finland (grant numbers
314254, 314255, 327255 and 342851), the University of Helsinki, and the European Research
Council under the European Articles Union's Horizon 2020 and Horizon Europe research and
innovation programs (grant no. 742312 and 101054307). Views and opinions expressed are
those of the author(s) only and do not necessarily reflect those of the European Union or the
European Research Council Executive Agency. Neither the European Union nor the granting
authority can be held responsible for them.

Acknowledgements:

We acknowledge Valeria Valanne, Katja Maamela, Markus Lauha, and Suvi Ikonen for helping
with tissue dissections; Suvi Ikonen, Katja Maamela, Markus Lauha, Nikolai Piavchenko, Paul
Bangura, Tiemen Jansen, Shadi Jansouz, Teemu Mäkinen, Jack O'Callagan, Anna Toikkanen, Ilke
van Gesten, Pirta Palola, Fin Morrison, Jeferson Florez, Juho Kökkö, Matias Boehm, Mikko
Immonen and Simon Lecarte for their assistance with animal husbandry and data collection;
Jacqueline Moustakas-Verho, Kzenia Zueva, Marion Sinclair-Waters, Nico Lorenzen, Victoria
Pritchard, Yann Czorlich, Suvi Ikonen, Spiros Papakostas and the staff at the Laukaa and
Taivalkoski hatcheries for assistance with egg and sperm collection and/or fertilizations; Tutku
Aykanat for calling SNPs; Seija Tillanen, and Shadi Jansouz for help with *vgll3* genotyping and
sexing; Xindi Huang for discussions about ovary 5S rRNA variation; Susanna Airaksinen,
Thomas Ginström Raisioaqua Ltd for producing the control- and low-fat feeds; John Loehr for
assistance with logistics at Lammi Biological Station; and Natural Resources Institute Finland
(LUKE) for access to their broodstocks.

SUPPLEMENTARY MATERIALS

Supplementary Table 1. TaqMan assays.

vgll3 Exon 2 Forward Primer	*vgll3* Exon 2 Reverse Primer	*vgll3* Exon 2 *Early* Probe	*vgll3* Exon 2 *Late* probe
GAAGCTGGGTGTTTACAGTAGGAT	TGTCTCCGCCCTGGAAAC	TGCTGCTCCATGCTGT	TGCTGCTCCGTGCTGT

amh Forward Primer	*amh* Reverse Primer	*amh* probe
ATGAAAGAGTGACAGAAAAGCTGAGT	TGCAGCTCACACAGAAACGT	CCCTCCCCTGCTCCC

lhr Forward Primer	*lhr* Reverse Primer	*lhr* probe
GCGGCGGGCTTCCT	CAGAGTGATAGTCGACAACGTGTAC	CTCCCCACCGAACACA

Supplementary figure 3. Posterior distributions of fixed-effect parameter estimates (b_i) for all fixed-effects (i). See statistical methods for a descriptiption of model structures and parameters. Thick and thin bars indicate 50% and 97.5% credible intervals, respectively. Bars are colored gray if their intervals include zero. Blue distributions indicate that the parameter estimates are unconditional (marginal) of their non-significant interactions. Numbers indicate the mean parameter estiamte.

Supplementary figure 1. Gonadal *vgll3* genotype and expression over time. Expression of *vgll3* mRNA in gonads over the entire study period, measured as copies ng^{-1} of RNA, for all sampled gonads. Upper plots show values for undeveloped ovaries (A) and testes (B), and lower plots show values for developing or mature ovaries (A) and testes(B). Colours and shapes correspond to the genotype of the sampled individuals: purple (+) for *vgll3*EE, blue (✳) for *vgll3*EL, and green (x) for *vgll3*LL. Lines connect means between time-points with at least 2 samples and error bars indicate SE. The numbers on bottom indicate sample sizes for *vgll3* genotypes LL, EL, and EE. The y-axes are log$_e$ scaled.

Supplementary figure 2. Body condition and *vgll3* expression over time, for two feed treatments. Body condition (A) and gonadal expression of *vgll3* mRNA (B: ovaries, C: testes) for the entire study period for all sampled individuals. Dark points represent individuals from the control feed treatment, and light grey points the low-fat feed treatment. Larger points and error bars indicate means and SE. Red stars in (A) indicate the significance ($p > 0.05$) of *t*-tests between the two treatments at the corresponding time point. The y-axes for B and C are $\log_e$ scaled.

REFERENCES

Ahi, E. P., Sinclair-Waters, M., Donner, I., & Primmer, C. R. (2023). A pituitary gene network

linking vgll3 to regulators of sexual maturation in male Atlantic salmon. *Comparative*

Biochemistry and Physiology Part A: Molecular & Integrative Physiology, *275*, 111337.

https://doi.org/10.1016/j.cbpa.2022.111337

Ahi, E. P., Sinclair-Waters, M., Moustakas-Verho, J., Jansouz, S., & Primmer, C. R. (2022). Strong

regulatory effects of vgll3 genotype on reproductive axis gene expression in juvenile male

Atlantic salmon. *General and Comparative Endocrinology*, *325*, 114055.

https://doi.org/10.1016/j.ygcen.2022.114055

Amer, M. A., Miura, T., Miura, C., & Yamauchi, K. (2001). Involvement of Sex Steroid Hormones in

the Early Stages of Spermatogenesis in Japanese Huchen (*Hucho perryi*). *Biology of*

Reproduction, *65*(4), 1057–1066. https://doi.org/10.1095/biolreprod65.4.1057

Andersson, E., Schulz, R. W., Male, R., Bogerd, J., Patiña, D., Benedet, S., Norberg, B., & Taranger, G.

L. (2013). Pituitary gonadotropin and ovarian gonadotropin receptor transcript levels:

Seasonal and photoperiod-induced changes in the reproductive physiology of female

Atlantic salmon (*Salmo salar*). *General and Comparative Endocrinology*, *191*, 247–258.

https://doi.org/10.1016/j.ygcen.2013.07.001

Åsheim, E. R., Debes, P. V., House, A., Liljeström, P., Niemelä, P. T., Siren, J. P., Erkinaro, J., &

Primmer, C. R. (2023). Atlantic salmon (Salmo salar) age at maturity is strongly affected

by temperature, population and age-at-maturity genotype. *Conservation Physiology*, *11*(1),

coac086. https://doi.org/10.1093/conphys/coac086

Bangura, P. B., Tiira, K., Niemelä, P. T., Erkinaro, J., Liljeström, P., Toikkanen, A., & Primmer, C. R.

(2022). Linking vgll3 genotype and aggressive behaviour in juvenile Atlantic salmon

(Salmo salar). *Journal of Fish Biology*, *100*(5), 1264–1271.

https://doi.org/10.1111/jfb.15040

Barson, N. J., Aykanat, T., Hindar, K., Baranski, M., Bolstad, G. H., Fiske, P., Jacq, C., Jensen, A. J.,

Johnston, S. E., Karlsson, S., Kent, M., Moen, T., Niemelä, E., Nome, T., Næsje, T. F., Orell, P.,

Romakkaniemi, A., Sægrov, H., Urdal, K., ... Primmer, C. R. (2015). Sex-dependent

dominance at a single locus maintains variation in age at maturity in salmon. *Nature*,

528(7582), 405–408. https://doi.org/10.1038/nature16062

Billard, R., & Breton, B. (1978). Rhythms of reproduction in teleost fish. In *Rhythmic activity of*

fishes (p. np). Academic Press, INC. https://hal.science/hal-01600956

Borg, B. (1994). Androgens in teleost fishes. *Comparative Biochemistry and Physiology Part C:*

Pharmacology, Toxicology and Endocrinology, *109*(3), 219–245.

https://doi.org/10.1016/0742-8413(94)00063-G

Bürkner, P.-C. (2017). brms: An R Package for Bayesian Multilevel Models Using Stan. *Journal of*

Statistical Software, *80*, 1–28. https://doi.org/10.18637/jss.v080.i01

Clark, K. L., George, J. W., Przygrodzka, E., Plewes, M. R., Hua, G., Wang, C., & Davis, J. S. (2022).

Hippo Signaling in the Ovary: Emerging Roles in Development, Fertility, and Disease.

Endocrine Reviews, *43*(6), 1074–1096. https://doi.org/10.1210/endrev/bnac013

Clelland, E., & Peng, C. (2009). Endocrine/paracrine control of zebrafish ovarian development.

Molecular and Cellular Endocrinology, *312*(1), 42–52.

https://doi.org/10.1016/j.mce.2009.04.009

Clermont, Y. (1972). Kinetics of spermatogenesis in mammals: Seminiferous epithelium cycle and

spermatogonial renewal. *Physiological Reviews*, *52*(1), 198–236.

https://doi.org/10.1152/physrev.1972.52.1.198

Cousminer, D. L., Berry, D. J., Timpson, N. J., Ang, W., Thiering, E., Byrne, E. M., Taal, H. R., Huikari,

V., Bradfield, J. P., Kerkhof, M., Groen-Blokhuis, M. M., Kreiner-Møller, E., Marinelli, M.,

Holst, C., Leinonen, J. T., Perry, J. R. B., Surakka, I., Pietiläinen, O., Kettunen, J., … Early

Growth Genetics (EGG) Consortium. (2013). Genome-wide association and longitudinal

analyses reveal genetic loci linking pubertal height growth, pubertal timing and childhood

adiposity. *Human Molecular Genetics*, *22*(13), 2735–2747.

https://doi.org/10.1093/hmg/ddt104

Debes, P. V., Piavchenko, N., Ruokolainen, A., Ovaskainen, O., Moustakas-Verho, J. E., Parre, N.,

Aykanat, T., Erkinaro, J., & Primmer, C. R. (2021). Polygenic and major-locus contributions

to sexual maturation timing in Atlantic salmon. *Molecular Ecology*, *30*(18), 4505–4519.

https://doi.org/10.1111/mec.16062

Dey, C. J., O'Connor, C. M., Gilmour, K. M., Van Der Kraak, G., & Cooke, S. J. (2010). Behavioral and

physiological responses of a wild teleost fish to cortisol and androgen manipulation

during parental care. *Hormones and Behavior*, *58*(4), 599–605.

https://doi.org/10.1016/j.yhbeh.2010.06.016

Elks, C. E., Perry, J. R. B., Sulem, P., Chasman, D. I., Franceschini, N., He, C., Lunetta, K. L., Visser, J.

A., Byrne, E. M., Cousminer, D. L., Gudbjartsson, D. F., Esko, T., Feenstra, B., Hottenga, J.-J.,

Koller, D. L., Kutalik, Z., Lin, P., Mangino, M., Marongiu, M., … Murray, A. (2010). Thirty new

loci for age at menarche identified by a meta-analysis of genome-wide association studies.

Nature Genetics, *42*(12), Article 12. https://doi.org/10.1038/ng.714

Fernald, R. D. (1976). The Effect of Testosterone on the Behavior and Coloration of Adult Male

Cichlid Fish (Haplochromis burtoni, Günther). *Hormones*, *7*(3), 172–178.

https://doi.org/10.1159/000178726

Figeac, N., Mohamed, A. D., Sun, C., Schönfelder, M., Matallanas, D., Garcia-Munoz, A., Missiaglia, E.,

Collie-Duguid, E., De Mello, V., Pobbati, A. V., Pruller, J., Jaka, O., Harridge, S. D. R., Hong, W.,

Shipley, J., Vargesson, N., Zammit, P. S., & Wackerhage, H. (2019). VGLL3 operates via

TEAD1, TEAD3 and TEAD4 to influence myogenesis in skeletal muscle. *Journal of Cell*

Science, *132*(jcs225946). https://doi.org/10.1242/jcs.225946

França, L. R., Nóbrega, R. H., Morais, R. D. V. S., De Castro Assis, L. H., & Schulz, R. W. (2015).

Sertoli cell structure and function in anamniote vertebrates. In *Sertoli Cell Biology* (pp.

385–407). Elsevier. https://doi.org/10.1016/B978-0-12-417047-6.00013-2

Gabry, J., & Mahr, T. (2022). *bayesplot: Plotting for bayesian models*. https://mc-

stan.org/bayesplot/

Halperin, D. S., Pan, C., Lusis, A. J., & Tontonoz, P. (2013). Vestigial-like 3 is an inhibitor of

adipocyte differentiation[S]. *Journal of Lipid Research*, *54*(2), 473–481.

https://doi.org/10.1194/jlr.M032755

Hélias-Rodzewicz, Z., Pérot, G., Chibon, F., Ferreira, C., Lagarde, P., Terrier, P., Coindre, J.-M., &

Aurias, A. (2010). YAP1 and VGLL3, encoding two cofactors of TEAD transcription factors,

are amplified and overexpressed in a subset of soft tissue sarcomas. *Genes, Chromosomes*

and Cancer, *49*(12), 1161–1171. https://doi.org/10.1002/gcc.20825

House, A. H., Debes, P. V., Kurko, J., Erkinaro, J., & Primmer, C. R. (2023). *Genotype-specific*

variation in seasonal body condition at a large-effect maturation locus (p.

2023.01.15.524064). bioRxiv. https://doi.org/10.1101/2023.01.15.524064

Hutchings, J. A. (2021). *A Primer of Life Histories: Ecology, Evolution, and Application* (1st ed.).

Oxford University Press. https://doi.org/10.1093/oso/9780198839873.001.0001

Hutchings, J. A., & Jones, M. (1998). Life history variation and growth rate thresholds for maturity

in Atlantic salmon, Salmo salar. *Canadian Journal of Fisheries and Aquatic Sciences - CAN J*

FISHERIES AQUAT SCI, *55*, 22–47. https://doi.org/10.1139/cjfas-55-S1-22

Johnston, S. E., Gratten, J., Berenos, C., Pilkington, J. G., Clutton-Brock, T. H., Pemberton, J. M., &

Slate, J. (2013). Life history trade-offs at a single locus maintain sexually selected genetic

variation. *Nature*, *502*(7469), Article 7469. https://doi.org/10.1038/nature12489

Kawamura, K., Cheng, Y., Suzuki, N., Deguchi, M., Sato, Y., Takae, S., Ho, C., Kawamura, N., Tamura,

M., Hashimoto, S., Sugishita, Y., Morimoto, Y., Hosoi, Y., Yoshioka, N., Ishizuka, B., & Hsueh,

A. J. (2013). Hippo signaling disruption and Akt stimulation of ovarian follicles for

infertility treatment. *Proceedings of the National Academy of Sciences*, *110*(43), 17474–

17479. https://doi.org/10.1073/pnas.1312830110

Kjærner-Semb, E., Ayllon, F., Kleppe, L., Sørhus, E., Skaftnesmo, K., Furmanek, T., Segafredo, F. T.,

Thorsen, A., Fjelldal, P. G., Hansen, T., Taranger, G. L., Andersson, E., Schulz, R. W.,

Wargelius, A., & Edvardsen, R. B. (2018). Vgll3 and the Hippo pathway are regulated in

Sertoli cells upon entry and during puberty in Atlantic salmon testis. *Scientific Reports*,

8(1), Article 1. https://doi.org/10.1038/s41598-018-20308-1

Küpper, C., Stocks, M., Risse, J. E., dos Remedios, N., Farrell, L. L., McRae, S. B., Morgan, T. C.,

Karlionova, N., Pinchuk, P., Verkuil, Y. I., Kitaysky, A. S., Wingfield, J. C., Piersma, T., Zeng, K.,

Slate, J., Blaxter, M., Lank, D. B., & Burke, T. (2016). A supergene determines highly

divergent male reproductive morphs in the ruff. *Nature Genetics*, *48*(1), Article 1.

https://doi.org/10.1038/ng.3443

Kurko, J., Debes, P. V., House, A. H., Aykanat, T., Erkinaro, J., & Primmer, C. R. (2020). Transcription

Profiles of Age-at-Maturity-Associated Genes Suggest Cell Fate Commitment Regulation as

a Key Factor in the Atlantic Salmon Maturation Process. *G3 Genes|Genomes|Genetics*, *10*(1),

235–246. https://doi.org/10.1534/g3.119.400882

Lamichhaney, S., Fan, G., Widemo, F., Gunnarsson, U., Thalmann, D. S., Hoeppner, M. P., Kerje, S.,

Gustafson, U., Shi, C., Zhang, H., Chen, W., Liang, X., Huang, L., Wang, J., Liang, E., Wu, Q., Lee,

S. M.-Y., Xu, X., Höglund, J., ... Andersson, L. (2016). Structural genomic changes underlie

alternative reproductive strategies in the ruff (*Philomachus pugnax*). *Nature Genetics*,

48(1), Article 1. https://doi.org/10.1038/ng.3430

Liang, Y., Tsoi, L. C., Xing, X., Beamer, M. A., Swindell, W. R., Sarkar, M. K., Berthier, C. C., Stuart, P.

E., Harms, P. W., Nair, R. P., Elder, J. T., Voorhees, J. J., Kahlenberg, J. M., & Gudjonsson, J. E.

(2017). A gene network regulated by the transcription factor VGLL3 as a promoter of sex-

biased autoimmune diseases. *Nature Immunology*, *18*(2), Article 2.

https://doi.org/10.1038/ni.3643

Lyu, Z., Qin, N., Tyasi, T. L., Zhu, H., Liu, D., Yuan, S., & Xu, R. (2016). The Hippo/MST Pathway

Member SAV1 Plays a Suppressive Role in Development of the Prehierarchical Follicles in

Hen Ovary. *PLOS ONE*, *11*(8), e0160896. https://doi.org/10.1371/journal.pone.0160896

Ma, S., Tang, T., Probst, G., Konradi, A., Jin, C., Li, F., Gutkind, J. S., Fu, X.-D., & Guan, K.-L. (2022).

Transcriptional repression of estrogen receptor alpha by YAP reveals the Hippo pathway

as therapeutic target for ER+ breast cancer. *Nature Communications*, *13*(1), Article 1.

https://doi.org/10.1038/s41467-022-28691-0

McDowell, E. N., Kisielewski, A. E., Pike, J. W., Franco, H. L., Yao, H. H.-C., & Johnson, K. J. (2012). A

Transcriptome-Wide Screen for mRNAs Enriched in Fetal Leydig Cells: CRHR1 Agonism

Stimulates Rat and Mouse Fetal Testis Steroidogenesis. *PLOS ONE*, *7*(10), e47359.

https://doi.org/10.1371/journal.pone.0047359

Meng, Z., Moroishi, T., & Guan, K.-L. (2016). Mechanisms of Hippo pathway regulation. *Genes &*

Development, *30*(1), 1–17. https://doi.org/10.1101/gad.274027.115

Mielcarek, M., Piotrowska, I., Schneider, A., Günther, S., & Braun, T. (2009). VITO-2, a new SID

domain protein, is expressed in the myogenic lineage during early mouse embryonic

development. *Gene Expression Patterns*, *9*(3), 129–137.

https://doi.org/10.1016/j.gep.2008.12.002

Miura, C., Higashino, T., & Miura, T. (2007). A Progestin and an Estrogen Regulate Early Stages of

Oogenesis in Fish1. *Biology of Reproduction*, *77*(5), 822–828.

https://doi.org/10.1095/biolreprod.107.061408

Miura, T., Miura, C., Ohta, T., Nader, M. R., Todo, T., & Yamauchi, K. (1999). Estradiol-17β

Stimulates the Renewal of Spermatogonial Stem Cells in Males. *Biochemical and*

Biophysical Research Communications, *264*(1), 230–234.

https://doi.org/10.1006/bbrc.1999.1494

Miura, T., Ohta, T., Miura, C. I., & Yamauchi, K. (2003). Complementary Deoxyribonucleic Acid

Cloning of Spermatogonial Stem Cell Renewal Factor. *Endocrinology*, *144*(12), 5504–5510.

Scopus. https://doi.org/10.1210/en.2003-0800

Morais, R. D. V. S., Crespo, D., Nóbrega, R. H., Lemos, M. S., van de Kant, H. J. G., de França, L. R.,

Male, R., Bogerd, J., & Schulz, R. W. (2017). Antagonistic regulation of spermatogonial

differentiation in zebrafish (*Danio rerio*) by Igf3 and Amh. *Molecular and Cellular*

Endocrinology, *454*, 112–124. https://doi.org/10.1016/j.mce.2017.06.017

Nakayama, K., Ohashi, J., Watanabe, K., Munkhtulga, L., & Iwamoto, S. (2017). Evidence for Very

Recent Positive Selection in Mongolians. *Molecular Biology and Evolution*, *34*(8), 1936–

1946. https://doi.org/10.1093/molbev/msx138

Pearse, D. E., Barson, N. J., Nome, T., Gao, G., Campbell, M. A., Abadía-Cardoso, A., Anderson, E. C.,

Rundio, D. E., Williams, T. H., Naish, K. A., Moen, T., Liu, S., Kent, M., Moser, M., Minkley, D.

R., Rondeau, E. B., Brieuc, M. S. O., Sandve, S. R., Miller, M. R., … Lien, S. (2019). Sex-

dependent dominance maintains migration supergene in rainbow trout. *Nature Ecology &*

Evolution, *3*(12), 1731–1742. https://doi.org/10.1038/s41559-019-1044-6

Perry, J. R., Day, F., Elks, C. E., Sulem, P., Thompson, D. J., Ferreira, T., He, C., Chasman, D. I., Esko,

T., Thorleifsson, G., Albrecht, E., Ang, W. Q., Corre, T., Cousminer, D. L., Feenstra, B.,

Franceschini, N., Ganna, A., Johnson, A. D., Kjellqvist, S., … Ong, K. K. (2014). Parent-of-

origin-specific allelic associations among 106 genomic loci for age at menarche. *Nature*,

514(7520), 92–97. https://doi.org/10.1038/nature13545

Pfennig, F., Standke, A., & Gutzeit, H. O. (2015). The role of Amh signaling in teleost fish – Multiple

functions not restricted to the gonads. *General and Comparative Endocrinology*, *223*, 87–

107. https://doi.org/10.1016/j.ygcen.2015.09.025

Prokkola, J. M., Åsheim, E. R., Morozov, S., Bangura, P., Erkinaro, J., Ruokolainen, A., Primmer, C. R.,

& Aykanat, T. (2021). Genetic coupling of life-history and aerobic performance in juvenile

Atlantic salmon. *bioRxiv*, 2021.08.23.457324. bioRxiv.

https://doi.org/10.1101/2021.08.23.457324

R Core Team. (2021). *R: A Language and Environment for Statistical Computing*. R Foundation for

Statistical Computing. https://www.R-project.org/

Ricker, W. E. (1975). Computation and interpretation of biological statistics of fish populations.

Bull. Fish. Res. Bd. Can., *191*, 1–382.

Rodríguez-Marí, A., Yan, Y.-L., BreMiller, R. A., Wilson, C., Cañestro, C., & Postlethwait, J. H. (2005).

Characterization and expression pattern of zebrafish anti-Müllerian hormone (amh)

relative to sox9a, sox9b, and cyp19a1a, during gonad development. *Gene Expression*

Patterns, *5*(5), 655–667. https://doi.org/10.1016/j.modgep.2005.02.008

RStudio Team. (2020). *RStudio: Integrated Development Environment for R* (1.4.1717) [Computer

software]. RStudio, PBC. http://www.rstudio.com/

Schulz, R. W., de França, L. R., Lareyre, J.-J., LeGac, F., Chiarini-Garcia, H., Nobrega, R. H., & Miura,

T. (2010). Spermatogenesis in fish. *General and Comparative Endocrinology*, *165*(3), 390–

411. https://doi.org/10.1016/j.ygcen.2009.02.013

Sen Sharma, S., Vats, A., & Majumdar, S. (2019). Regulation of Hippo pathway components by FSH

in testis. *Reproductive Biology*, *19*(1), 61–66.

https://doi.org/10.1016/j.repbio.2019.01.003

Shen, Z.-G., Yao, H., Guo, L., Li, X.-X., & Wang, H.-P. (2017). Ribosome RNA Profiling to Quantify

Ovarian Development and Identify Sex in Fish. *Scientific Reports*, *7*(1), Article 1.

https://doi.org/10.1038/s41598-017-04327-y

Sinclair-Waters, M., Piavchenko, N., Ruokolainen, A., Aykanat, T., Erkinaro, J., & Primmer, C. R.

(2022). Refining the genomic location of single nucleotide polymorphism variation

affecting Atlantic salmon maturation timing at a key large-effect locus. *Molecular Ecology*,

31(2), 562–570. https://doi.org/10.1111/mec.16256

Song, M., & Gutzeit, H. O. (2003). Effect of 17-alpha-ethynylestradiol on germ cell proliferation in

organ and primary culture of medaka (*Oryzias latipes*) testis. *Development, Growth &*

Differentiation, *45*(4), 327–337. https://doi.org/10.1046/j.1440-169x.2003.00701.x

Stan Development Team. (2022). *RStan: The R interface to Stan* [Computer software]. https://mc-

stan.org/

Troth, A., Puzey, J. R., Kim, R. S., Willis, J. H., & Kelly, J. K. (2018). Selective trade-offs maintain

alleles underpinning complex trait variation in plants. *Science (New York, N.Y.)*, *361*(6401),

475–478. https://doi.org/10.1126/science.aat5760

Vehtari, A., Gabry, J., Magnusson, M., Yao, Y., Bürkner, P.-C., Paananen, T., & Gelman, A. (2022). *loo:*

Efficient leave-one-out cross-validation and WAIC for Bayesian models [Computer

software]. https://mc-stan.org/loo/

Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model evaluation using leave-one-

out cross-validation and WAIC. *Statistics and Computing*, *27*(5), 1413–1432.

https://doi.org/10.1007/s11222-016-9696-4

Verta, J.-P., Debes, P. V., Piavchenko, N., Ruokolainen, A., Ovaskainen, O., Moustakas-Verho, J. E.,

Tillanen, S., Parre, N., Aykanat, T., Erkinaro, J., & Primmer, C. R. (2020). Cis-regulatory

differences in isoform expression associate with life history strategy variation in Atlantic

salmon. *PLOS Genetics*, *16*(9), e1009055. https://doi.org/10.1371/journal.pgen.1009055

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

https://ggplot2.tidyverse.org

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Grolemund, G., Hayes,

A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache, S. M., Müller, K., Ooms, J.,

Robinson, D., Seidel, D. P., Spinu, V., … Yutani, H. (2019). Welcome to the tidyverse. *Journal*

of Open Source Software, *4*(43), 1686. https://doi.org/10.21105/joss.01686

Wootton, R. J., & Smith, C. (2014). *Reproductive Biology of Teleost Fishes: Wootton/Reproductive*

Biology of Teleost Fishes. John Wiley & Sons, Ltd.

https://doi.org/10.1002/9781118891360

Xiang, C., Li, J., Hu, L., Huang, J., Luo, T., Zhong, Z., Zheng, Y., & Zheng, L. (2015). Hippo Signaling

Pathway Reveals a Spatio-Temporal Correlation with the Size of Primordial Follicle Pool in

Mice. *Cellular Physiology and Biochemistry*, *35*(3), 957–968.

https://doi.org/10.1159/000369752