Phytic Acid (InsP$_6$) Activates HDAC3 Epigenetic Axis to Maintain Intestinal Barrier Function

Sujan Chetterjee[1], Zachary Sin[1], Nguyen Tran[1], Loretta Vierra[1,2],Anuj Shukla[3], Tam Tran[1], George Koshkaryan[1], Kevin Ritter[3], Xue Bessie Su[4], Saharat Jolak Ragsac[1], Seungman Park[5], Qian Liu[1,2], Richard Van[2], Mira Han[2], Adolfo Saiardi[4], Katherine Huang[6], Kayci Huff-Hardy[6], Richard Rood[6], Anas Gremida[6], Martin Gregory[6], Chien-Huan Chen[6], Parakkal Deepak[6],Henning J. Jessen[3], and Prasun Guha[1,2,@]

[1]Nevada Institute of Personalized Medicine, University of Nevada, Las Vegas, NV, USA;

[2]School of Life Sciences, University of Nevada, Las Vegas, NV, USA.

[3]Institute of Organic Chemistry, Faculty of Chemistry and Pharmacy, University of Freiburg, Albertstr. 21, 79104 Freiburg, Germany.

[4]Medical Research Council Laboratory for Molecular Cell Biology, University College London, London, WC1E 6BT, United Kingdom.

[5]Department of Mechanical Engineering, University of Nevada, Las Vegas, Las Vegas, NV, 89154, USA

[6]Division of Gastroenterology, Washington University School of Medicine in St. Louis, St. Louis, MO, USA

[@] **Corresponding Authors:**

Prasun Guha, University of Nevada-Las Vegas, 4505 S Maryland Pkwy, Harry Reid Center, Rm 183C, Las Vegas, NV 89154, prasun.guha@unlv.edu, 702-895-2033

Abstract.

While HDAC inhibition shows promise in cancer treatment, pan-HDAC inhibitors cause

gastrointestinal issues in 48% of patients. Understanding HDAC activation mechanisms

is crucial to treat diverse diseases beyond cancer. Our study reveals the essential role

of inositol polyphosphate multikinase (IPMK) and inositol hexakisphosphate ($InsP_6$ or

phytic acid), enriched in vegan diets, in activating the HDAC3 epigenetic axis and

maintaining intestinal barrier integrity. IPMK binds to HDAC3, driving $InsP_6$ synthesis,

which selectively activates HDAC3 at 10nM concentration by recruiting the DAD domain

of its corepressor protein. *IPMK* deletion diminishes HDAC3 activation, leading to

histone hyperacetylation and MMP gene transcription, compromising intestinal barrier

integrity. $InsP_6$ treatment is sufficient to rescue these effects. In inflammatory bowel

disease, diminished IPMK levels exacerbated intestinal permeability, while oral $InsP_6$

treatment mitigated gut permeability by restoring the HDAC3 epigenetic axis, indicating

the clinical implications of the IPMK-HDAC3 epigenetic axis and therapeutic potential of

phytic acid.

Introduction

HDAC3, a key regulator of histone acetylation[1], plays a crucial role in epigenetic

modification[1]. While nonspecific HDAC inhibitors show promise in cancer treatment [2],

their side effects[3] highlight the importance of understanding individual HDAC functions.

HDAC3, a prominent class I HDAC family member[1], has emerged as a critical regulator

of various physiological functions. Preclinical mouse model studies have demonstrated

that HDAC3 is pivotal in maintaining metabolic homeostasis and circadian rhythms,

particularly in the liver[4]. Furthermore, HDAC3 loss has been observed in inflammatory

bowel disease (IBD) patients[5], and its deletion in mouse intestinal epithelial cells

induces Crohn's disease-like symptoms, including increased intestinal permeability[5].

Despite its physiological importance, our understanding of the underlying mechanisms

regulating HDAC3's enzymatic activity remains limited.

What is known about the HDAC3 activation mechanism is mostly related to protein-

protein interaction, suggesting HDAC3 binds to the corepressor complex silencing

mediator for retinoid and thyroid hormone receptors (SMRT/NCoR2)[1] and nuclear

receptor co-repressor 1 (NCoR1)[1], essential for its enzymatic activity[1]. Recent studies

have illuminated the interaction between HDAC3 and inositol phosphates, particularly

$InsP_4$, through crystal structure analysis[1]. However, significant gaps in our

understanding persist. The physiological basis of InsP's role in activating HDAC3

remains unclear. The precise mechanism and cellular location (cytoplasm, nucleoplasm,

or chromatin) of InsP-mediated HDAC3 activation are unknown. Most crucially, the

downstream effects of InsP-activated HDAC3 on the epigenetic axis, gene regulation,

and overall cell function remain largely unclear.

While negatively charged $InsP_4$ binding to HDAC3 was charge-dependent[1,6], in a cellular

environment, it is produced in very small amounts[7]. In contrast, $InsP_6$, which is more

negatively charged than $InsP_4$, is produced in cells at higher micromolar concentrations

and is more abundant[8]. Although the crystal structure did not observe $InsP_6$ binding to

HDAC3[1], biochemical *in vitro* studies identified high-affinity binding of $InsP_6$ to HDAC3[6].

While *in vitro* studies show $InsP_4$, $InsP_5$, and $InsP_6$ can activate purified HDAC3

complexes from wild-type cells, these experiments involved dissociating InsPs in high-

salt conditions[6,9]. Although this suggests InsPs regulate HDAC3 outside the cellular

environment, it doesn't confirm that physiological InsPs influences HDAC3 function.

Thus, the biological significance of *in vivo* InsP-mediated HDAC3 regulation remains

unclear.

At the physiological level, intracellular InsPs production is regulated by several

enzymes, with IPMK (inositol polyphosphate multikinase) playing a critical role[8], as its

deletion diminishes $InsP_4$, $InsP_5$, and $InsP_6$ levels[7]. Here, we investigated the effect of

IPMK loss of function on HDAC3 activation to determine the importance of InsPs in

regulating HDAC3 activation and the precise molecular mechanism. We discovered

that IPMK physically interacts with HDAC3, and the interaction occurs specifically in

chromatin but not in other cellular compartments. This finding is particularly intriguing

given the known dynamic behavior of HDAC3, which shuttles between the cytoplasm

and nucleus and requires recruitment to chromatin for its deacetylation function[10]. This

chromatin-specific binding of IPMK to HDAC3 suggests a targeted regulatory

mechanism, potentially influencing HDAC3's activity precisely where it exerts its

epigenetic effects.

Our research further uncovered a critical role for $InsP_6$ in the activation of HDAC3.

Notably, treatment with $InsP_6$ to *IPMK* knockout cells and mouse models completely

restored HDAC3 activity to levels comparable to those observed in wild-type cells, firmly

establishing $InsP_6$ as a physiological small molecule activator of HDAC3.

Notably, $InsP_6$, also known as phytic acid, is abundant in plant-based diets, particularly

in rice and legumes[11]. This dietary connection becomes significant in light of our

findings. We discovered that in a mouse model of inflammatory bowel disease (IBD),

there was a notable loss of IPMK protein expression. This loss disrupted the IPMK-

HDAC3 epigenetic axis, leading to the acetylated histone lysine 16 (H4K16) promoter

enrichment and transcriptional upregulation of matrix metalloproteinase (MMP) genes

elevating gut permeability. MMPs are proteolytic enzymes that play a crucial role in

degrading cell-cell junctions, consequently exacerbating intestinal permeability, a

hallmark feature of IBD[12]. This aligns with genome-wide association studies (GWAS)

and our preclinical study that have identified IPMK as a risk gene for IBD[13]. While

transcriptional upregulation of MMPs in IBD is well-established[12,14], the mechanism of

epigenetic regulation of their expression was previously obscure. Our findings suggest

that disruption of the IPMK- HDAC3 epigenetic axis influence the transcriptional

upregulation of MMP family genes in IBD. Notably, we demonstrate that oral $InsP_6$

(Phytic acid) treatment is effective in reducing intestinal permeability. .

In conclusion, our study reveals that IPMK physically interacts with chromatin-bound

HDAC3 and identifies $InsP_6$ as a crucial physiological small molecule activator of

HDAC3. $InsP_6$ functions by recruiting the DAD domain of the corepressor to HDAC3,

thereby activating it and leading to the deacetylation of specific histone marks. This

mechanism plays a crucial role in modulating the epigenetic landscape, specifically by

repressing MMPs transcription. Consequently, it helps maintain cell-cell junctions and

preserve intestinal barrier integrity, highlighting its importance in gut homeostasis and

potentially in preventing or intercepting inflammatory bowel diseases.

Results

IPMK binds to the nuclear HDAC3 and is essential for HDAC3 activation regulating histone acetylation.

To investigate the physiological impact of HOIPs on HDAC3 activation, *IPMK* was deleted in HCT116 (colon), A549 (lung), and MEF (mouse embryonic fibroblast cell lines **(Extended Fig1 A)**. As the rate-limiting enzyme in the HOIP pathway **(Fig 1A)**, *IPMK* deletion is expected to cause a profound loss in the levels of all downstream HOIP pathway products[8]. Consistent with this expectation, we observed a significant reduction in the levels of $InsP_5$ and $InsP_6$ in *IPMK* knockout (KO) cells, with only a modest decline in $InsP_4$ **(Fig 1B)**. Additionally, HPLC data from the wild-type (WT) cells indicated that the overall concentrations of $InsP_5$ and $InsP_6$ are 3 to 4 times higher than that of $InsP_4$ **(Fig 1B)**. To investigate the impact of *IPMK* deletion on HDAC3's enzymatic activity, we immunopurified endogenous HDAC3 from respective WT and *IPMK* KO cells, followed by HDAC3 activity assay **(Fig 1C)**. We optimized the HDAC3 activity assay by immunoprecipitating HDAC3 from various protein concentrations ranging from 5μg, 20μg, 100μg and 500μg. Our results showed that HDAC3 purified from 20μg of protein exhibited strong deacetylase activity, with only modest increments in activity observed up to 500μg **(Extended Fig 1B)**. Based on these findings, we selected 20μg as the minimum protein concentration for all subsequent HDAC3 pulldown followed by enzymatic assays. This choice allowed us to achieve better resolution in detecting activity differences. Notably, *IPMK* deletion dramatically reduced HDAC3's enzymatic activity by over 90% in all tested cell lines compared to their WT counterparts **(Fig1 D),**

however, without affecting HDAC3 or corepressor protein levels **(Extended Fig 1C)**.

Strikingly, this reduction in HDAC3 activity was comparable to that observed in *HDAC3-*

KO cells **(Extended Fig 1D)** or WT cells treated with the HDAC3-specific inhibitor

RGFP966 **(Fig1 E).** The selectivity of RGFP966 for HDAC3 is precise, as it shows 200-

fold selectivity for HDAC3 compared to other HDACs and binds to HDAC3's enzymatic

site, inhibiting its deacetylase activity[15]. These findings demonstrate that at the

physiological level, *IPMK* deletion impairs HDAC3 activation to an extent similar to

complete *HDAC3* deletion or pharmacological inhibition, highlighting IPMK's crucial role

in maintaining HDAC3 function.

Given HDAC3's role as a deacetylase, the loss of its activity in *IPMK* KO cells was

expected to increase histone acetylation. Our findings confirmed that *IPMK* deletion

enhanced acetylation levels of specific histone residues, particularly H3K9/27 and

H4K16 **(Extended Fig 1E, F)**. Notably, the extent of H3K9/27 and H4K16 acetylation in

IPMK KO cells was comparable to that observed in *HDAC3* KO cells and wild-type cells

treated with the HDAC3 inhibitor RGFP966 **(Fig 1 F, G)**. To determine whether this

effect in *IPMK* deleted cells was due to the loss of HDAC3 activity, we treated *IPMK* KO

cells with the HDAC3-specific inhibitor RGFP966. Interestingly, the inhibitor failed to

induce any further increase in histone acetylation either in *IPMK* or *HDAC3* KO cells

(Fig 1 F, G) while profoundly increasing H3K9/27 and H4K16 acetylation when treated

to WT cells **(Fig 1 F, G)**. This differential response strongly suggests that HDAC3

activity was already substantially compromised in *IPMK*-deficient cells; consequently,

RGFP966 was unable to enhance histone acetylation in *IPMK* KO cells further.

Next, we explored the potential direct interaction between IPMK and HDAC3 under

physiological conditions. Considering HDAC3's known shuttling between the nucleus

and cytoplasm[10] and its requirement to bind chromatin for histone deacetylation in the

nucleus, we performed cell fractionation and chromatin immunopurification, revealing

that IPMK specifically associates with chromatin-bound HDAC3 **(Fig 1 H, I)**. We did not

find the interaction between IPMK and HDAC3 in the cytoplasm; however, it was

detected when chromatin-bound HDAC3 was immunopurified **(Fig 1H, I)**. To further

validate this interaction, we conducted an *in vitro* binding assay using recombinant

HDAC3 and IPMK proteins, which confirmed a direct physical interaction between these

two proteins **(Extended Fig 1G)**.

Overall, the study demonstrated that IPMK binds to nuclear HDAC3 and is crucial for its

enzymatic activity.

IPMK's inositol kinase activity is essential for HDAC3 activation.

To elucidate the role of IPMK's inositol kinase activity in regulating HDAC3's enzymatic

activity, we conducted experiments using *IPMK* knockout (KO) cells. We transfected

these cells with either wild-type (WT) or kinase-dead (KD) IPMK[7]. Our results showed

that rescuing *IPMK* KO cells with WT-IPMK restored HDAC3 deacetylase activity, while

KD-IPMK failed to do so **(Fig 2A)**. Interestingly, immunoprecipitation studies revealed

that both WT-IPMK and KD-IPMK could interact with HDAC3 complex proteins **(Fig 2B)**.

These findings suggest a crucial distinction: while IPMK's physical binding to HDAC3 is

insufficient for rescuing HDAC3's enzymatic activity, the kinase activity of IPMK is

required for HDAC3 activation.

Next, we tested the possibility of rescuing HDAC3's enzymatic activity with $InsP_4$, $InsP_5$,

and $InsP_6$ *in vitro* test tubes after immunopurification of HDAC3 from *IPMK* KO cells.

Biochemical analysis revealed that $InsP_6$, in a dose-dependent manner, rescued the

enzymatic activity of HDAC3 **(Fig 2 C)**. While 10 nM (0.01µM) $InsP_6$ restored >50% of

HDAC3 activity, 1 µM $InsP_6$ [InsP (1,2,3,4,5,6) P_6] fully restored activity comparable to

the WT **(Fig 2 C)**. In WT cells, 5 or 10 µM $InsP_6$ did not further increase HDAC3 activity,

suggesting $InsP_6$ saturation and a threshold effect at 1µM **(Fig 2B)**. Although $InsP_4$

[InsP (1,4,5,6) P_4] and $InsP_5$ [InsP (1,3,4,5,6) P_5] also restored HDAC3's activity at

1 µM **(Fig 2 D)**, $InsP_6$ showed the highest effect compared to all other HOIPs.

Furthermore, while 10 nM of $InsP_5$ and $InsP_6$ both restored around 50% of HDAC3

activity compared to WT cells **(Extended Fig 2 A)**, $InsP_4$ (10 nM) did not have any

effect on HDAC3 activation and was similar to the KO cells. Interestingly, a second

$InsP_4$ isomer [InsP (1,3,4,5) P_4] and $InsP_3$ [InsP (1,4,5) P_3] completely failed to rescue

HDAC3 activity, even at 10 µM dose **(Fig 2 D)**. While Wu et al.'s[16] previous study

indicated $InsP_3$ activating HDAC3, our experiments showed that $InsP_3$ could not

influence HDAC3 activity at either 1 µM or 10 µM concentrations **(Fig 2 D)**. This finding

is supported by previous crystal structure analysis[6], indicating the necessity of a

phosphate group at the 6^{th} position of higher order inositolphosphate (HOIP) activating

HDAC3. This crucial phosphate is missing in both $InsP_3$ and the $InsP_4$ isomer

[InsP(1,3,4,5)P$_4$], explaining their inability to activate HDAC3 in our study. As IPMK is a

physiological PI3 kinase [13]that also generates PIP3, we further tested the role of PIP3 in

rescuing HDAC3 activity when purified from *IPMK* KO cells. We found that even at

10 μM**,** PIP3 failed to restore HDAC3 activity and was similar to the *IPMK* KO **(Fig 2**

D). Based on the dose-dependent study **(Fig 2 C, D)** and previous research

demonstrating InsP$_6$'s health benefits[11], we opted to utilize InsP$_6$ for our subsequent

experiments.

Next, we explored the possibilities of rescuing HDAC3 activity *in vivo* by treating cells

with cell-permeable InsP$_6$ (cell-permeable InsP$_4$ and InsP$_5$ have not been successfully

synthesized). InsP$_6$ exhibits poor cell permeability in cultured cells as cations present in

cell culture media (such as calcium, magnesium, sodium, and potassium) may chelate

InsP$_6$, diminishing its availability for cellular uptake. Consequently, experiments often

require the use of high, non-physiological concentrations of InsP$_6$ to achieve desired

effects. To overcome this, we used a cell-permeable InsP$_6$ precursor molecule, where

the negative charges of InsP$_6$ are temporarily masked with biolabile groups, allowing it

to pass through the cell membrane and then releasing active InsP$_6$ after removal of the

biolabile groups by enzymatic digestion[17]. We treated *IPMK* KO cells with this cell-

permeable InsP$_6$ and measured HDAC3 activity. Remarkably, 24 hours of cell-

permeable InsP$_6$ treatment rescued HDAC3 activity in *IPMK* KO cells to levels

comparable to those of WT cells **(Fig 2 E)**, followed by a substantial decrease in histone

acetylation, which was also comparable to the WT **(Fig 2 F)**. However, CP-InsP$_6$ did not

exert any effect when treated to wild type cells **(Extended Fig 2B)**. Notably, the

treatment of HDAC3 inhibitor RGFP966 inhibited the rescue effect of cell-permeable

InsP$_6$ **(Fig 2 F)**, indicating that InsP$_6$ deacetylates acetylated H3K9/27 and H4K16 by

activating HDAC3.

Lastly, we evaluated the ability of CP-InsP6 to enhance intracellular InsP6 levels.

Following 24 hours of treatment with 50 µM CP-InsP6, mass spectrometric analysis

revealed a significant increase in intracellular InsP6 compared to wild-type cells

(Extended Fig 2C). We also observed a parallel rise in InsP5 and InsP4 levels,

suggesting dephosphorylation of InsP6 by phosphatases into lower-order inositol

phosphates. Nevertheless, InsP6 remained the most abundant among the detected

species.

***IPMK* deletion disrupts HDAC3's interaction with the DAD domain of its**

corepressor proteins, diminishing the activity of chromatin-bound HDAC3.

Next, we wondered about the mechanism underlying the loss of HDAC3's enzymatic

activity in *IPMK*-deleted cells. As mentioned earlier, the HDAC3 protein level remains

unaltered in *IPMK* KO cells **(Extended Fig 1B)**; hence, we wondered whether *IPMK*

deletion disrupts HDAC3 binding to its corepressor protein, which is essential for

HDAC3 function[1]. Immunoaffinity purification **(Extended Fig 3A)** and proximity ligation

assay (PLA) **(Fig 3A, B)** demonstrated that *IPMK* deletion did not disrupt the

association between HDAC3 and its corepressor protein, maintaining the overall

integrity of the complex. While the SANT domain of the corepressor protein maintains

the integrity of the HDAC3:corepressor complex[1,9], the corepressor's DAD domain

binding to HDAC3 induces structural changes[1,9], that are crucial for HDAC3's enzymatic

activity, although it is not necessary for preserving the complex's integrity **(Fig 3C)**.

Given this, it's plausible that *IPMK* deletion specifically compromised the interaction

between the DAD domain and HDAC3, thereby hindering HDAC3's enzymatic activity

without disrupting the overall complex integrity. To investigate whether *IPMK* deletion

interferes with DAD domain interaction with HDAC3, we overexpressed DAD-Flag in

IPMK knockout cells and subsequently performed a pull-down assay to isolate

endogenous HDAC3. DAD-Flag pulled down endogenous HDAC3 from WT cells but,

interestingly, failed when pulled down from *IPMK* KO cells **(Fig 3 D)**. Notably, the

impaired DAD binding to HDAC3 observed in *IPMK* knockout cells was successfully

rescued by treating the cells with cell-permeable $InsP_6$ **(Fig 3 E)**. This observation aligns

with previous crystal structure analyses, which revealed HOIP's role as a molecular

"glue" facilitating the interaction between the DAD domain of the corepressor protein

and HDAC3[1,6,9].

Given that HDAC3's chromatin recruitment is crucial for HDAC3-mediated histone

deacetylation next, we wondered whether *IPMK* deletion impaired HDAC3's chromatin

recruitment. Using HDAC3-to-histone H3 PLAs, or immunopurification of H3 or H4

chromatin fraction followed by HDAC3 western blot, we found a comparable amount of

HDAC3 chromatin recruitment in *IPMK* KO and WT cells **(Fig 3 F, G, Extended Fig**

3B), indicating that *IPMK* deletion did not impair HDAC3's chromatin recruitment.

Our data collectively support a functional model where HDAC3 recruitment to chromatin

and its binding to the corepressor complex occur independently of IPMK. However,

$InsP_6$ is crucial for the DAD domain's interaction with HDAC3, which triggers its

enzymatic activity and subsequent histone deacetylation **(Fig 3H)**.

To investigate whether $InsP_6$ serves as the final rate-limiting step in HDAC3-mediated

histone deacetylation, we employed a simplified cell-free system. We treated cell-free,

chromatin fraction in reaction buffer from *IPMK* knockout cells with $InsP_6$ for one hour, a

context where any observed effects would likely result from $InsP_6$'s direct action on

chromatin-bound HDAC3, independent of complex multistep processes of HDAC3

chromatin recruitment involving other proteins. Strikingly, $InsP_6$ treatment fully

suppressed histone acetylation in *IPMK* KO nuclear lysate **(Fig 3I)**, restoring it to wild-

type levels. The specificity of this effect was confirmed by co-treatment with the HDAC3-

selective inhibitor RGFP966, which negated $InsP_6$'s impact **(Fig 3I)**. However, the

overall integrity of the HDAC3 chromatin protein complex (including NCoR1/2, TBL1,

GPS2)[18], remains intact in *IPMK* KO cells and is comparable to the WT **(Extended Fig**

3C). These results provide strong evidence that $InsP_6$ directly activates chromatin-

bound HDAC3 by recruiting the DAD domain of the corepressor to HDAC3, functioning

as the critical final step in the histone deacetylation process.

IPMK generates InsP5 , while IPPK produces $InsP_6$. Given that IPMK interacts with

HDAC3, we examined whether IPPK also associates with HDAC3, potentially

contributing to a localized HOIP signaling event within the HDAC3 complex. Our

biochemical and microscopy analyses revealed that IPPK indeed interacts with nuclear

HDAC3 **(Extended Fig. 3II, A, B)**.

318
Disruption of the IPMK-HDAC3 epigenetic axis hypertranscribes MMP gene

family.

We aimed to investigate the role of the IPMK-HDAC3 epigenetic axis in transcriptional

regulation. Our bulk RNA sequencing analysis of *IPMK*-deleted HCT116 cells revealed

significant transcriptional changes compared to wild-type cells, with 4,014 genes

upregulated and 3,271 downregulated **(Fig 4A)**. Pathway analysis indicated that the

genes related to the extracellular matrix degradation, which includes the collagenase

gene family, were notably upregulated in *IPMK* KO cells **(Fig 4B)**. Specifically,

archetypal matrix metalloproteinases (MMPs), including MMP1, MMP10, MMP12, and

MMP13, were significantly upregulated in the RNA-seq data **(Fig 4A)**. We validated

these findings at the protein level using western blot analysis for MMP1, MMP10, and

MMP13, confirming the RNA-seq results **(Fig 4C)**. Due to a lack of high-quality

antibodies, we were unable to test MMP12.

Given that IPMK is recognized as a risk gene for Inflammatory bowel disease (IBD)[13],

we also analyzed transcriptomic data from human patients using the Inflammatory

Bowel Disease Transcriptome and Metatranscriptome Meta-Analysis (IBDTaMMA[19],

https://ibd-meta-analysis.herokuapp.com/)[19] **(Extended Fig 4 A)**. This analysis revealed

significant transcriptional upregulation of MMP1, MMP3, MMP10, MMP12, and MMP13,

in IBD patient cells similar to the findings in *IPMK* KO HCT116- intestinal cells.

Furthermore, separate analyses of Crohn's disease (CD) and ulcerative colitis (UC)

patients revealed a striking increase in major MMPs—such as MMP3, MMP13, and

MMP10—that were consistently elevated in patient tissues and similarly upregulated in

IPMK-depleted cells **(Extended Fig 4 II A,B)**.

Interestingly, *IPMK* KO cells rescued with *wild-type IPMK* showed a reduction in MMP

expression, whereas cells with kinase-dead IPMK did not exhibit this effect **(Extended**

Fig. 4B). Importantly, cell-permeable InsP$_6$ treatment to *IPMK* KO cells suppressed

MMP expression as well, making it comparable to that of the WT cells **(Fig 4D)**,

indicating that the loss of IPMK's inositol kinase activity upregulates MMP gene

expression. To determine whether InsP$_6$ treatment repressed MMP gene expression by

activating HDAC3, we co-treated HDAC3 inhibitor RGFP966, which abrogated the

rescue effect of InsP$_6$ **(Fig 4D)**.

Moreover, in the *HDAC3* KO cells, MMP genes **(Extended Fig 4C)** and protein

expression were markedly increased **(Fig 4E)** and comparable to the *IPMK* KO cells

(Fig 4E). In our second approach to determine whether identified MMP genes are

regulated via the IPMK-HDAC3 epigenetic axis, we examined the effects of the HDAC3

inhibitor RGFP966 on WT, *IPMK* KO, and *HDAC3* KO cells. We hypothesized that

RGFP966 would increase MMP expression in WT cells but not in *IPMK or HDAC3* KO

cells, as these cells lack HDAC3 activity. However, if MMP expression in *IPMK* KO cells

was influenced by pathways independent of HDAC3, RGFP966 treatment might further

increase MMP expression. As anticipated, RGFP966 significantly increased MMP

expression in WT cells but failed to induce further expression in *IPMK and HDAC3* KO

cells **(Fig 4F, G)**. Taken together, the results confirm that the disruption of the IPMK-

HDAC3 epigenetic axis is responsible for the transcriptional upregulation of MMPs,

ruling out the involvement of HDAC3-independent pathways in this process.

Next, we examined whether the promoters of matrix metalloproteinases (MMPs) exhibit

enrichment of histone acetylation. We selected MMP13 as a representative example for

this analysis. Our biochemical studies confirmed that *IPMK* deletion selectively

upregulates H3K9/27 and H4K16 acetylation at the protein level, an effect also

observed in *HDAC3* KO cells and those treated with HDAC3 inhibitors **(Fig 1F)**.

Intriguingly, ChIP-qPCR targeting the MMP13 promoter region **(Fig 4H)** revealed a

selective enrichment of acetylated H4K16 in the promoters of MMP13 in *IPMK* KO cells,

while H3K9/27 acetylation showed no significant change **(Fig 4 H)**. The ChIP data also

indicated an enrichment of HDAC3 as tested using MMP13 promoter regions

(Extended Fig 4 D).

A particularly fascinating finding emerged when we treated *IPMK KO* cells with cell-

permeable $InsP_6$. This treatment substantially reduced the enrichment of acetylated

H4K16 at MMP promoters **(Fig 4I)**. However, this effect was nullified when cells were

co-treated with the HDAC3 inhibitor RGFP966 **(Fig 4I)**. Notably, $InsP_6$ treatment did not

affect HDAC3 chromatin recruitment **(Extended Fig 4 D)**.

Given the known upregulation of MMP3 in IBD[20], we revisited our *IPMK* KO HCT116

RNA-seq data to understand the MMP3 log fold change. Although we found a 2.5-Log

2-fold increase in MMP3 expression in HCT116 cells, the p-value of 0.3 exceeded our

significance threshold of $p<0.05$. Raw data analysis revealed zero reads in all three WT

samples and a maximum of two reads in KO samples for MMP3, likely contributing to

the lack of statistical significance. While deeper sequencing might yield statistically

significant results, our qPCR **(Extended Fig 4E)** and protein-level data **(Extended Fig 4**

F - H) clearly demonstrated a substantial increase in MMP3 levels in both *IPMK* and

HDAC3 KO cells, which was reversed by InsP$_6$ treatment **(Extended Fig 4I)**.

Furthermore, studies using HDAC3 inhibitors and knockout models confirmed that

disrupting the IPMK-HDAC3 epigenetic axis substantially elevated MMP3 transcriptional

expression, providing additional validation of our findings.

Finally, to assess the overlap between the global transcripts regulated by HDAC3 and

IPMK, we first identified differentially expressed genes in *HDAC3* KO cells and

compared this dataset with the expression profile of *IPMK* KO cells. The results showed

that approximately 44% of HDAC3-regulated genes were also affected in *IPMK* KO cells

(Fig 4J). Importantly, HDAC3 exhibits a dual regulatory role in transcription: it functions

not only through its deacetylase activity but also as a cotranscriptional activator

independent of this enzymatic function[16]. This dual functionality likely explains why

certain genes are affected by *HDAC3* deletion but remain unaltered in *IPMK* KO cells.

Specifically, the genes impacted in *HDAC3* KO cells but not in *IPMK* KO cells may be

influenced by HDAC3's non-enzymatic roles, such as its function as a cotranscriptional

activator.

410
IPMK deletion is associated with increased cell and intestinal permeability in

mice.

The upregulation of MMP is associated with compromised intestinal barrier function by

disrupting tight junctions, leading to a "leaky gut", which is commonly observed in IBD

patients[12]. Given these associations, we investigated the consequences of *IPMK*

deletion stimulating increased MMP production in cell permeability. To assess that, we

employed a transwell assay using Caco2 cells, which is classically used for *in vitro* cell

permeability assay[21]. We cultured these cells in a confluent layer on a trans well insert

and treated them with 4kDa FITC-Dextran. Cell permeability would increase the leakage

of 4kDa FITC-Dextran from the upper cell chamber into the bottom chamber **(Fig 5 A)**.

Notably, conditioned media from *IPMK* KO or *HDAC3* KO cells elevated Caco2 cell

permeability **(Fig 5 B)**, while media transferred from WT cells failed to induce cell

permeability. The observed effect may be attributed to the secretory nature of MMPs. In

IPMK KO HCT116 cells, the conditioned media contained elevated levels of secreted

MMPs (Extended Fig 5 II A). When this MMP-enriched media was transferred to wild-

type Caco2 cells, it increased cellular permeability by disrupting the cell-cell

junction. This hypothesis was corroborated by the western blot study that identified the

loss of tight junction proteins ZO-1 and Occludin in CaCo2 cells treated with either

IPMK KO cell-conditioned media **(Extended Fig 5A)**. Remarkably, *IPMK or HDAC3* KO

cells pretreated with Pan-MMP inhibitor (Ilomastat) followed by media transfer

completely blocked the Caco2 cell permeability **(Fig 5 B)**, confirming that the effect was

due to MMP in the conditioned media.

Furthermore, *IPMK* KO cells treated with just cell permeable $InsP_6$ followed by media

transfer to Caco2 cells prevented the Caco2 cell permeability, but the RGFP966 co-

treatment abrogated $InsP_6$'s rescue effect **(Fig 5 B)**. This confirms that disruption of the

IPMK-HDAC3 epigenetic axis leads to MMP upregulation and consequent increased

cell permeability, effects successfully rescued by $InsP_6$ treatment.

In vivo, mice studies further supported the cell-level findings. We created intestine-

specific *IPMK* KO mice (IPMK$^{\Delta IEC}$) **(Extended Fig 5 B-E)** and tested gut permeability by

orally gavaged mice with FITC-dextran (4kD) and Rhodamine- dextran (70 kD). The

disruption of the tight junction only allows 4kD dextran but restricts 70 kD[22-24], activating

the leaky pathway, while significant destruction of the epithelial cell layer allows both

4kD and 70 kD dextran **(Fig 5 C)**. Notably, the IPMK$^{\Delta IEC}$ mouse showed a marked

increase in the leaky pathway, allowing 4kD dextran but without allowing 70 kD to pass

through **(Fig 5 D)**, indicating intestinal loss of *IPMK* selectively activates the leaky

pathway expected due to loss of tight junction disruption.

To determine the hyper-transcription of MMPs in IPMK$^{\Delta IEC}$ mice, we flow purified

(Extended Fig 5 B-E) pure intestinal epithelial cells (IEC) from IPMK$^{F/F}$ and IPMK$^{\Delta IEC}$

mice followed by RNA-seq. Data indicated a profound increase in MMP transcription

(Fig 5 E), also validated by qPCR, tested in two representative MMP genes (MMP3 and

MMP13) **(Fig 5 F).** Pathway analysis using Metascape **(Extended Fig. 5II G-I)** and **GSEA**

(Extended Fig. 5II J) confirmed enrichment of gene sets associated with MMP activity

and extracellular matrix organization. Protein level analysis by performing western blots

of crypts **(Fig 5G, Extended Fig 5 II B,C)** and immunohistochemistry **(Fig 5 H)**

reconfirmed transcriptomic data showing elevated MMPs in IPMK$^{\Delta IEC}$ mice.

Fascinatingly, oral administration of InsP$_6$ solution normalized to pH 7.0 , treated to

IPMK$^{\Delta IEC}$ mice completely reduced MMP levels, making them comparable to those

observed in IPMK$^{F/F}$ mice **(Fig 5 G,H, Extended Fig 5 II B,C).** Next, we studied the

levels of major tight junction proteins like ZO-1, Occludin, and claudin 2. Notably, ZO-1

and Occludin levels were significantly reduced in IPMK$^{\Delta IEC}$ mice as tested in Western

blot and IHC studies **(Fig 5 G,I,J, Extended Fig 5 II D,E)**. Interestingly, claudin 2 levels

were rather increased in IPMK$^{\Delta IEC}$ **(Extended Fig 5 II F)**, which agrees with previous

studies indicating increased claudin 2 is linked with elevated intestinal permeability[23].

Importantly, oral InsP6 treatment rescued all tight junction protein levels. Moreover, the

HDAC3 activity assay indicated a marked decline in HDAC3 activation in IPMK$^{\Delta IEC}$ **(Fig**

5 K). Notably, oral InsP$_6$ treatment restored HDAC3 activity to the level of IPMK$^{F/F}$ mice

(Fig 5 K. Protein level analysis showed that in IPMK$^{\Delta IEC}$ mice intestine H4K16

acetylation was markedly increased which oral InsP$_6$ restored back to the IPMK$^{F/F}$ level;

however, the IPMK protein was unaffected by $InsP_6$ treatment **(Fig 5 L, Extended Fig 5**

F), indicating oral $InsP_6$ treatment alone could restore the epigenetic change of H4K16

acetylation in IPMK$^{\Delta IEC}$ mice in absence of IPMK **(Fig 5L).** The protein level epigenetic

change was reconfirmed at the chromatin level by ChIP-qPCR analysis of MMP3, and

13, indicating in IPMK$^{\Delta IEC}$ mice marked increase of H4K16ac in MMP3, 13 promoters,

which were reduced and reversed back to the level of IPMK$^{F/F}$ mice by oral $InsP_6$

treatment **(Fig 5 M,N)**. As expected, MMP promoter recruitment of HDAC3 was evident

in IPMK$^{\Delta IEC}$ mice and comparable to the IPMK$^{F/F}$ mice, and it was unaffected by oral

$InsP_6$ treatment **(Extended Fig 5G).** While $InsP_6$ cellular uptake is poor in cell culture

systems, in our animal studies, a 2% body weight, $InsP_6$ treatment was sufficient to

demonstrate its effects. This observation aligns with previous research showing

intestinal absorption[25-27] and therapeutic efficacy of oral $InsP_6$[28] in mice and rats.

To determine whether the above finding has any clinical significance in relation to IBD,

we found that both CD and UC patient samples showed diminished HDAC3 activity **(Fig**

5 O, P). Remarkably, IPMK protein levels were markedly reduced in both CD and UC

patient samples **(Fig 5Q,R)**, but this did not affect IPPK protein levels, indicating

specificity. Moreover, we found a loss of ZO-1 and Occludin levels in CD and UC patient

samples, along with increased H4K16 acetylation **(Fig 5Q,R)**.

The collective findings suggest that the deletion of intestinal *IPMK* leads to a reduction

in HDAC3 activity, which in turn promotes the expression of (MMPs). This phenomenon

is evident in IBD patient samples, showing a pronounced loss of IPMK and its

downstream effects, as observed in IPMK$^{\Delta IEC}$ mice. This cascade of events disrupts cell

junctions, ultimately resulting in a leaky gut condition. Notably, oral administration of

InsP$_6$ effectively reversed leaky gut effects.

500
In a mouse model of IBD, *IPMK* expression and HDAC3 activity are reduced, and

oral InsP$_6$ treatment restores intestinal permeability.

We used a DSS (dextran sodium sulfate)-induced IBD mice model [29] to test the leaky

gut, where , marked elevation in intestinal permeability for 4kD dextran was evident as

early as 24h, not allowing 70kD **(Fig 6A)**. Remarkably, pre-treatment with oral InsP$_6$

dosed at 2% of the mouse body weight 72h prior to the DSS treatment, prevented leaky

gut comparable to the untreated mice **(Fig 6 A)**. Since InsP$_6$'s bioavailability is modest[25],

we pretreated mice for 72 hours. Notably, the 24h of DSS treatment selectively reduced

IPMK protein expression in the HOIP pathway **(Fig. 1 A)** while keeping IPPK protein

expression unaltered (**Fig 6B, Extende Fig. 6A)**, with a substantial loss of HDAC3

activity, which was restored in DSS-treated mice by oral InsP$_6$ pre-treatment **(Fig 6C)**.

Moreover, the western blot of colonic crypts indicated a substantial increase in MMP3

and 13 expressions after DSS treatment, which oral InsP$_6$ pretreatment attenuated and

restored to the level of untreated mice **(Fig 6D)**. We reconfirmed the western blot data

using immunohistochemistry of MMP3 in the colon and even in the ileum, showing that

both small and large intestines exhibited increased MMP3 expression **(Fig 6E-H)**.

Though acute DSS treatment mostly shows ulcerative colitis-like phenotype in mice

colon, the previous study also indicated its pathological effects in the ileum[30]; hence, we

also included the ileum to show the molecular effect is unrestricted throughout the

intestine after DSS treatment.

At the epigenetic level, DSS treatment markedly increased H4K16 acetylation in the

protein level **(Fig 6 G)** with increased H4K16 acetylation enriched MMP3 and MMP13

promoter **(Fig 6 H)** and transcriptional upregulation of MMP 3 and 13 **(Fig 6 I)**. These

were restored to the level of untreated mice by oral $InsP_6$ pretreatment **(Fig 6 G-I)**.

While HDAC3 promoter enrichment on the MMP3 and 13 promoter was evident in DSS-

treated mice, it was unaltered by oral $InsP_6$ pretreatment **(Extended Fig 6 A)**. Its

important to mention that oral $InsP_6$ also has no effects on IPMK protein expression so

works indepdnet of IPMK **(Extended Fig 6B)** .

This finding highlights that disruption of the IPMK-HDAC3 epigenetic axis in a preclinical

mouse model of IBD exacerbates intestinal permeability; however, oral $InsP_6$ (phytic

acid) treatment mitigates the effects of the leaky gut.

Oral $InsP_6$ treatment increased intestinal epithelial cell-specific $InsP_6$ level and

prevents DSS-induced intestinal injury.

To assess the preventative effect of $InsP_6$ on intestinal inflammation, wild-type mice

were exposed to 2% dextran sodium sulfate (DSS) in drinking water for 7 days, with or

without 2% oral InsP$_6$ co-treatment **(Fig 7A)**. For the InsP$_6$ group, mice were pretreated

with oral InsP$_6$ for 2 days prior to DSS exposure to enhance the protective effect **(Fig.**

7A). To induce colitis, mice were given 2% DSS in drinking water continuously for 7

days, followed by sacrifice on the morning of the 8th day **(Fig 7A)**. Each day, mice were

also gavaged with 200 microliters of DSS to ensure uniform exposure.

As expected, mice treated with DSS alone developed severe colitis, characterized by

body weight loss starting around 6th day **(Extended Fig. 7A)**, reduced colon length

(Fig. 7B), and mucosal erosion and tissue damage observed in H&E-stained colonic

sections **(Extended Fig. 7B)**. This inflammatory phenotype was further confirmed by

markedly elevated colonic IL-1β (Fig. 7**C)** and myeloperoxidase (MPO) activity (Fig.

7**D)**, indicating robust inflammatory cytokine production and neutrophil infiltration,

respectively.

Notably, oral co-treatment with InsP$_6$ completely prevented these DSS-induced

pathologies. Mice receiving InsP$_6$ maintained their body weight **(Extended Fig. 7A)** and

exhibited no reduction in colon length **(Fig. 7B)** compared to untreated controls.

Histological analysis revealed preserved tissue integrity with minimal damage

(Extended Fig. 7B), and both IL-1β **(Fig. 7C)** and MPO levels **(Fig. 7D)** remained at

baseline , comparable to those of untreated mice. Collectively, these data demonstrate

that oral InsP$_6$ provides robust protection against the development of DSS-induced

colitis.

Lastly, we assessed intestinal permeability. Both 4 kDa and 70 kDa dextran assays

showed increased permeability following DSS treatment **(Fig 7E)**, with the greatest

increase observed for 4 kDa dextran **(Fig 7E)**. Remarkably, oral $InsP_6$ treatment

robustly reduced intestinal permeability **(Fig 7E)** and effectively preserved barrier

integrity.

We next sought to determine if mice with an intestinal epithelial cell-specific deletion of

IPMK (IPMK$^{\Delta IEC}$) exhibit heightened susceptibility to DSS-induced inflammation and

whether oral $InsP_6$ could rescue this phenotype. Three groups of mice were studied:

IPMK $^{F/F}$ (controls), IPMK$^{\Delta IEC}$, and IPMK$^{\Delta IEC}$ + $InsP_6$. All groups were treated with 2%

DSS for five days and euthanized on day six **(Fig 7F)**. The IPMK$^{\Delta IEC}$ + $InsP_6$ group was

pre-treated with oral $InsP_6$ for three days before DSS co-treatment (Fig 7F), a strategy

intended to replenish the depleted endogenous $InsP_6$ pools resulting from IPMK loss.

Consistent with a critical role for IPMK in maintaining barrier function, IPMK$^{\Delta IEC}$ mice

displayed a markedly exacerbated inflammatory response to DSS compared to IPMK $^{F/F}$

controls, as evidenced by more severe tissue damage in H&E-stained sections

(Extended Fig 7E) and significantly elevated levels of IL-1β **(Fig 7G)** and MPO **(Fig**

7H). Furthermore, intestinal permeability to both 4 kDa and 70 kDa dextran was

profoundly increased in IPMK$^{\Delta IEC}$ mice after DSS treatment **(Fig 7I)**. Remarkably, oral

InsP$_6$ treatment almost completely abrogated these effects **(Fig 7I)**. The InsP$_6$-treated

IPMK$^{\Delta IEC}$ mice exhibited robust protection, with inflammation markers and intestinal

permeability comparable to those of the uninjured IPMK$^{F/F}$ controls. In this experiment

the duration of 6 days DSS treatment failed to induce any weight loss or colon size

(Extended Fig 7C,D).

While our data established that oral InsP$_6$ protects against intestinal inflammation, we

sought to determine if this effect stems from direct elevation of intestinal epithelial cell-

specific InsP$_6$ levels . Mass spectrometry of HOIP metabolites from isolated ileal crypts

(Fig 7J) revealed that InsP$_4$, InsP$_5$, and InsP$_6$ were markedly reduced in *IPMK$^{\Delta IEC}$* mice

relative to IPMK$^{F/F}$ controls, with InsP$_6$ levels diminished by approximately sevenfold

(Fig 7K). Remarkably, a single 24-hour oral dose of pH-calibrated (7.2) InsP$_6$ treated to

IPMK$^{\Delta IEC}$ mice restored InsP$_6$ to levels comparable to controls (Fig 7**K)**, while also

increasing InsP$_4$ and InsP$_5$, mirroring observations with cell-permeable InsP$_6$ treated to

cell lines **(Extended Fig 2C)**. These findings support a model of direct epithelial uptake

of orally administered InsP$_6$, which is then partially dephosphorylated. Although gut

microbial phytases can generate lower-order inositol phosphates and myoinositol for

uptake via transporters like SMIT2/SGLT6[34], the lack of IPMK in IPMK$^{\Delta IEC}$ mice would

prevent the subsequent re-phosphorylation to $InsP_5$ and $InsP_6$, thereby implying the

existence of a specific epithelial transport mechanism for InsP6 itself.

As we found that incorporation of $InsP_6$ at both cellular and tissue levels not only

increased $InsP_6$ concentration but also elevated $InsP_5$ and $InsP_4$, we sought to

determine which HOIP is physiologically critical for HDAC3 activation and MMP

expression. To specifically deplete $InsP_6$, we examined HDAC3 activity and MMP

expression in *IPPK* knockout cells. IPPK catalyzes the conversion of $InsP_5$ to $InsP_6$;

therefore, the loss of IPPK selectively reduces $InsP_6$ without affecting upstream

polyphosphates such as $InsP_5$ or $InsP_4$.

Intriguingly, HDAC3 activity was markedly reduced in *IPPK* knockout cells **(Fig 7L)**,

accompanied by elevated MMP levels **(Fig 7M),** however with out influencing HDAC3

expression itself. These findings indicate that $InsP_6$ plays a physiologically significant

role in HDAC3 activation compared to other HOIPs. In the clinical context, the *IPMK*

depletion model better reflects disease conditions such as inflammatory bowel disease

(IBD), where IPMK levels are reduced while IPPK remains unchanged **(Fig 5Q,R)**,

leading to lowered $InsP_6$ concentrations.

625
626

Discussion

Our study provides compelling evidence for the physiological role of $InsP_6$ as a crucial

activator of HDAC3, with significant implications for epigenetic regulation and intestinal

barrier function in IBD. While previous structural studies identified $InsP_4$ as a molecular

link between HDAC3 and its co-repressors[1], our findings demonstrate that $InsP_6$ is the

primary physiological activator of HDAC3, influencing epigenetic changes that regulate

MMP gene transcription.

We investigated HDAC3 activation and downstream epigenetic changes in *IPMK*-

deleted cell lines and mouse models to determine the physiological role of HOIPs in

regulating HDAC3 activation and its impact on the epigenetic axis. We targeted IPMK

because it is the rate-limiting enzyme in HOIP biosynthesis[8], and its deletion resulted in

reduced levels of $InsP_4$, $InsP_5$, and $InsP_6$ in cells **(Fig 1B)**. By focusing on IPMK, we

aimed to thoroughly evaluate the effects of HOIP depletion on HDAC3-mediated

epigenetic regulation.

While *IPMK* deletion markedly diminishes HDAC3's enzymatic activity **(Fig 1D)**, we

discovered that IPMK directly binds to HDAC3 in the chromatin complex **(Fig 1H)**.

However, through chemical genetic studies using kinase-dead IPMK and cell-permeable

$InsP_6$, we revealed that IPMK's binding to HDAC3 is dispensable for HDAC3's

enzymatic activity **(Fig 2A,B)**. These findings suggest that IPMK's kinase function,

rather than its physical interaction with HDAC3, is critical for maintaining HDAC3's

deacetylase activity. In the physiological context, we hypothesize that IPMK's

localization and association with chromatin-bound HDAC3 may facilitate the generation

of HOIPs in specific spatiotemporal regions. This localized production of HOIPs could

serve to maintain the efficiency of the HDAC3 regulatory mechanism, ensuring precise

control of histone deacetylation in chromatin.

While evaluating the concentration of HOIPs in cells, we made several intriguing

discoveries. We found that $InsP_4$ concentration was minimal compared to $InsP_5$ and

$InsP_6$ **(Fig 1B).** Notably, *IPMK* deletion primarily diminished $InsP_5$ and $InsP_6$ levels

rather than $InsP_4$ **(Fig 1B)**. Consistent with these observations, our *in vitro* rescue

experiments demonstrated that both $InsP_5$ and $InsP_6$ could restore a significant

proportion of HDAC3 activity in *IPMK* KO cells, with $InsP_6$ exhibiting the highest efficacy

(Fig 2D). Furthermore, we observed that cell-permeable $InsP_6$ alone was sufficient to

restore HDAC3 activity *in vivo* when administered to *IPMK* KO cells **(Fig 2 E)**. These

findings collectively underscore the crucial role of $InsP_6$ in HDAC3 activation and

suggest a direct mechanism of action that is independent of IPMK's physical presence.

At the mechanistic level, our study revealed that while *IPMK* loss did not disrupt the

formation of the HDAC3-corepressor complex **(Fig 3A,B and Extended Fig 3A),** it did

impair the DAD (deacetylase activation domain) domain of its corepressor interaction

with HDAC3 **(Fig 3D,E),** which is crucial for activating HDAC3's deacetylase activity

however, dispensable for structural integrity.

While HDAC3 recruitment to chromatin is a crucial step in the regulation of gene

expression, our study revealed that neither IPMK nor InsP$_6$ affects this recruitment

process **(Fig 3F, E and Extended Fig 3B)**. Remarkably, we discovered a direct

mechanism of HDAC3 activation in a cell-free system. By treating the chromatin fraction

from *IPMK* KO cells with InsP$_6$, we observed that InsP$_6$ facilitates the final step of

HDAC3 activation. This activation occurs through the recruitment of the (DAD) domain

to chromatin-bound HDAC3, subsequently triggering histone deacetylation **(Fig 3I)**.

These findings reveal that InsP$_6$ is crucial in activating chromatin-bound HDAC3,

serving as a key rate-limiting step. However, InsP$_6$ does not contribute to the initial

recruitment of HDAC3 to chromatin, highlighting its specific function in enzyme

activation rather than chromatin localization.

Building upon our cell model findings that demonstrated a significant role of IPMK and

InsP$_6$ in HDAC3 activation, we investigated the physiological relevance of the IPMK-

HDAC3 epigenetic axis, particularly in the context of IBD, for which IPMK has been

identified as a risk gene through GWAS and preclinical studies[13]. Our study revealed

that intestinal loss of *IPMK* impaired HDAC3 activity **(Fig 5 I)** in mice, leading to a leaky

gut condition **(Fig 5 D)**. At the epigenetic level, disruption of the IPMK-HDAC3 axis

increased H4K16 acetylation in MMP promoters **(Fig 4H and Fig 5 J, K)**, activating their

transcription **(Fig 4A and Fig 5E)** and resulting in degradation of tight junctions

(Extended Fig 5A), thereby exacerbating intestinal permeability **(Fig 5 D)**. Notably,

transcriptomic analysis of patients with IBD showed a significant increase in MMPs

(Extended Figure 4 A), mirroring observations in intestine-specific *IPMK* knockout mice

and suggesting a probable link. Our study further identified that in the mouse model of

IBD, the IPMK-HDAC3 epigenetic axis was disrupted due to the loss of IPMK protein

expression **(Fig 6 B)**, reflecting the epigenetic effects observed in IPMK$^{\Delta IEC}$ mice.

Remarkably, oral administration of InsP$_6$ (phytic acid) mitigated the leaky gut effect **(Fig**

6A) by restoring HDAC3 activation **(Fig 6 C)**, reducing H4K16 acetylation **(Fig 6 G)**, and

suppressing MMP gene and protein expression **(Fig 6 D - F and I)**, highlighting its

potential as an approach in intercepting pre-clinical disease given prior data showing

that increased intestinal permeability is associated with later development of Crohn's

Disease. This may also work as a therapeutic approach for IBD, with impaired intestinal

permeability shown to be associated with ongoing bowel symptoms and severity of

diarrhea in patients with IBD despite having achieved mucosal healing.

Previous studies have indicated that InsP$_6$ is absorbed in the gastrointestinal tract,

albeit with low bioavailability[25-27,31]. However, studies indicated the therapeutic efficacy of

InsP$_6$ to treat cancer in mice models[28]. One hypothesis suggests that gut microbiota-

derived phytase enzymes dephosphorylate InsP$_6$ to lower-order inositol phosphates,

which are more readily absorbed[29]. However, our biochemical analysis, corroborated by

Watson et al.[6], demonstrates that HDAC3 activity is exclusively influenced by InsP$_4$,

InsP$_5$, or InsP$_6$, with InsP$_3$ having no effect **(Fig 2 D)**. Notably, oral InsP$_6$ administration

in IPMK$^{\Delta IEC}$ mice, which are incapable of generating InsP$_4$, InsP$_5$, or InsP$_6$, exhibited a

rescue effect **(Fig 5 I, J)**. This observation suggests the possibility of direct InsP$_6$

absorption in the gastrointestinal tract, although the precise mechanism is yet to be

explored.

In conclusion, our study highlights the critical role of the IPMK-HDAC3 epigenetic axis in

regulating intestinal barrier integrity and its potential disruption in IBD **(Fig 8)**. We

identify InsP$_6$ as a key physiological activator of HDAC3 and demonstrate that

InsP$_6$supplementation can restore HDAC3 activity, reduce histone acetylation, and

improve intestinal barrier function in mouse models of IBD **(Fig 8)**.

Materials and Methods

Cell culture.

Human Colon Tumor 116 (HCT116), Mouse Embryonic Fibroblast (MEF), A549, HEK and CaCo2 cells were cultured in McCoy's 5A medium, Dulbecco's Modified Eagle's Medium (DMEM) or F12/K respectively. DMEM, F12/K media was supplemented with 10% FBS, 2 mM L-glutamine, 100 U/mL penicillin, and 100 mg/mL streptomycin was used for MEF or A549 cells. While McCoy's 5A medium was supplemented with 10% FBS, 100 U/mL penicillin, and 100 mg/mL streptomycin. *IPMK* wild-type (WT) and knockout (KO) HCT116 and A549 cells were generated using CRISPR/Cas9 technology, which partially deleted exon 5 and the coding region of exon 6. Afterward, clonal populations were isolated, purified, and screened for IPMK expression.

For the generation of *IPMK* KO MEF cells, floxed MEFs were immortalized through transfection with pSG5-Large T (a plasmid provided by William Hahn to Addgene) using the Polyfect transfection reagent (Qiagen) according to the manufacturer's instructions. After serial passaging five times to select against non-transformed cells, *IPMK* KO cells were produced by transducing floxed cells with an adenovirus carrying the Cre recombinase gene (obtained from the University of Iowa Gene Transfer Vector Core). The virus was combined with Gene Jammer (Stratagene) transfection reagent to improve transduction efficiency. Clonal populations were then purified and screened for IPMK expression.

Animals.

All protocols were approved by the Institutional Animal Care and Use Committee (IACUC, UNLV, IACUC-01204, IACUC-01206), University of Nevada, Las Vegas. Mice were housed according to institutional guidelines, in a controlled environment at a temperature of 22_C ± 1_C, under a 12-h dark-light period and provided with standard chow diet and water *ad libitum*. Male and female *IPMK* $^{F/F}$, *IPMK*$^{\Delta IEC}$ (between eight to 16 weeks) were used. Specifically, intestinal epithelial cells from male mice were used for biochemical and transcriptomic analysis. All mice were maintained in C57BL/6 mixed background.

Generation of intestine-specific *IPMK* knockout mice.

Ipmk$^{F/F}$ mice were generated as previously described[13]*. Ipmk*$^{F/F}$ mice were crossed with C57BL/6J mice carrying Cre expressed under the control of the murine villin promoter as described previously to create intestinal-specific *Ipmk* knockout mice (*Ipmk*$^{\Delta IEC}$). Homozygous *Ipmk*$^{F/F}$ mice were crossed with the *Ipmk*$^{\Delta IEC}$ mice, which mediate excision of floxed alleles in the intestine. All mice were maintained on a C57BL/6J background and were 10th-generation backcrossed. Validated by MiniMUGA. Background analysis was performed by Transnetyx, Cordova, USA Mice were housed in a 12-hour light/12-hour dark cycle at 22°C and fed standard rodent chow. All research involving mice was approved by the University of Nevada, Las Vegas's Institutional Animal Care and Use Committee (IACUC).

Proximity ligation assay and quantitation:

PLA was utilized to detect *in situ* protein–protein interactions as previously described.

After fixation and permeabilization, cells were blocked with blocking solution supplied by

PLA kit before incubation with primary antibodies as in routine immunofluorescence

staining. The cells were then processed using a PLA kit (NaveniFlex Cell MR Red Cat

no. NAV-NC.MR.100.RED) according to the manufacturer's instructions. The slides

after PLA were further processed for immunofluorescence staining against the DAPI

(Cat.8617, Cell Signaling). The slides were then mounted with Prolong Gold antifade

Reagent. A Zeiss LSM 750 confocal microscope detected PLA signals as discrete

punctate foci and provided the intracellular localization of complexes. ImageJ was used

to quantify and represent the number of PLA puncta per nucleus.

DSS-induced colitis model and oral $InsP_6$ treatment.

DSS (MP Biomedicals; relative molecular mass 36,000–50,000) was added to drinking

water at 3% weight/volume for 24 hr or 2% weight/volume for five or seven days. Phytic

acid ($InsP_6$) was added to 2% weight/volume (2 gm/100 ml) containing drinking water

after pH neutralization with $NaHCO\square$ for 48 hr after that, the mouse received 2%

weight/volume $InsP_6$ and 3% DSS for 24 hr or 2% weight/volume for five or seven days..

Intestinal permeability assay.

$IPMK^{F/F}$ and either $IPMK^{\triangle IEC}$ mice were fasted for 4 hours with access to water before

being gavaged with 4 kDa-FITC-Dextran and 70 kDa RhodamineB (Sigma-Aldrich, Cat

no: #FD4-1G) at a single dose of 0.6mg/gm body weight. Untreated control group

received an equal amount of PBS. Fluorescence intensity of plasma samples was

measured (excitation 485 nm/emission 535 nm) after 4 hours of gavage. Value was

represented as a fold change in respect without FITC Dextran and Rhodamine treated

control groups after blank subtraction. All measurements were performed using

GraphPad Prism (version 6.0, GraphPad Software, Inc)[22,].

Generation of IPMK antibody:

Custom anti-IPMK antibody was generated in collaboration with Pro-Sci. The anti-rabbit

IPMK antibody had been raised against human IPMK amino acids 295–311

(SKAYSRHRKLYAKKHQS). This sequences are also closely conserved in mice

SKMYA**RHRK**IYT**KKH**H**S** hence reacts to both mouse and human IPMK.

803
Western Blot analysis.

Cell lysates were prepared through syringe flush by using a salt-free (NaCl-free) Lysis

buffer (50 mM Tris pH 7.5, 50 mM potassium acetate, 5 % v/v glycerol, 0.3 % v/v Triton

X-100, one tablet of Roche complete protease inhibitor). We used salt-free (NaCl-free)

lysis buffer for all the assays because high salt concentration dissociates higher-order

inositol phosphate (HOIPs) from HDAC3, which is essential for their deacetylase

activity[25,27]. Samples were centrifuged at 15,000g for 10 min, and the protein

concentration of the supernatant was measured. Proteins were resolved by SDS-

polyacrylamide gel electrophoresis (NuPAGE Bis-Tris Midi GEL, Life Technologies, Cat

#: WG1402BX10) and transferred to Immobilion-P PVDF (Millipore-Sigma, Cat #:

IPVH00010) transfer membranes. The membranes were incubated with primary

antibody diluted in 3% BSA in Tris- Buffered Saline with Tween 20 (20 mM Tris-HCl, pH

7.4, 150 mM NaCl, and 0.02% Tween 20) overnight incubation at 4°C. Respective

antibodies for western blot anti-HDAC3 (Santa Cruz Biotechnology, Cat #: sc-376957),

IPMK (Generated in our lab) anti-beta actin (ProteinTech, Cat #: 81115-1-RR), anti-

lamin (ProteinTech, Cat #: 12987-1-AP), anti-NCoR1 (Bethyl Laboratories, Cat #: A301-

148A), anti-Ncor1 (Cell Signaling Technology, Cat #: 5948), anti-Rcor1 (ProteinTech,

Cat #: 26686-1-AP), Sin3A (ProteinTech, Cat #: 14638-1-AP), anti-DNTTIP1 (Bethyl

Laboratories, Cat #: A304-048A), anti-H3 (Cell Signaling Technology, Cat #: 12648),

anti-H3k8Ac (Cell Signaling Technology, Cat #: 2598), anti-H3k9Ac (Cell Signaling

Technology, Cat #: 9649), anti-H3k18Ac (Cell Signaling Technology, Cat #: 13998),

anti-H3k27Ac (Abcam, Cat #: ab4729) ,anti-H3k56Ac (Cell Signaling Technology, Cat #:

4243), anti-H4k8Ac (Cell Signaling Technology, Cat #: 2594), anti-H4k12Ac (Cell

Signaling Technology, Cat #: 13944), anti-H4k16Ac (Cell signaling Technology, Cat #:

13534S) and anti-Myc (Invitrogen, Cat #:46-0709). Following primary antibody

incubation, the PVDF membrane was washed three times with TRIS-buffered

saline/Tween-20, and incubated with HRP-conjugated secondary antibody (ECL, Cat

#s: NA934V and NA931V), and the bands visualized by chemiluminescence (Super

Signal West Pico, Pierce, Cat #: 34579). The depicted blots are representative

replicates selected from at least three experiments. Densitometric analysis was

performed using Image J software. Blots for acetylated histones were stripped using

Restore Western Blot stripping buffer (Thermo Scientific, Cat # 21059), then reprobed

with H3 antibody (Cell Signaling Technology, Cat #: 12648S) at room temperature for 2

h followed by secondary antibody incubation and development.

HDAC3 (histone deacetylase) activity.

To analyze deacetylase activity of specific Class I HDAC members, the HDAC 3 were

individually immuno-purified, followed by an activity assay using the following kit:

HDAC3 Assay Kit (BPS, Cat#: 10186-628). Since high salt concentration dissociates

higher-order inositol phosphates (HOIPs) from HDAC3, which are crucial for their

deacetylase activity, WT and IPMK-KO HCT 116 cells were lysed in salt-free (NaCl-

free) lysis buffer (50 mM Tris pH 7.5, 50 mM potassium acetate, 5 % v/v glycerol, 0.3 %

v/v Triton X-100, one tablet of Roche complete protease inhibitor). Then, the lysate was

centrifuged at 15,000 g for 10 min, and the protein concentration of the supernatant was

measured. Equal amount, around 20 µg total protein from the lysate was incubated

against anti-HDAC3 (Santa Cruz Biotechnology, Cat #: sc-81600) antibodies,

respectively, at 4°C for 1 hour followed by capturing antibody with EZview A/G beads

(Millipore Sigma, Cat #: E3403). Beads were washed 3x with an ice-cold reaction buffer

(supplied with the kit). After a specific substrate reaction, fluorescence was measured

using a Spectra max iD5 plate reader (Molecular Devices). IgG was immunoprecipitated

from Wild Type HCT 166 cells (WT) and used as a negative control. In each

experiment, separate IgG immunoprecipitation was performed, and the data was

expressed as a fold change relative to the respective IgG control. Although slight

variations in IgG values were observed across experiments, the overall data range

differed slightly in each case, but consistently showed a similar trend. All measurements

were performed in triplicate, and data was analyzed using GraphPad Prism (version 6.0,

GraphPad Software, Inc).

Intestinal Epithelial cell purification.

863 The ileum (similar length as colon moving proximal from the ileocecal junction) were

864 dissected from mice. After five times washing with ice cold PBS, the tissues were

865 opened longitudinally and cut into small fragments (2–3 cm in length). To prepare single

866 cell suspension after washing three time with PBS, small intestinal fragments were

867 incubated in 30 mM EDTA-PBS on ice for 30 min, during which the tissues were shaken

868 vigorously every 5 min. Isolated crypts were washed once with cold PBS and

869 dissociated using TrypLE Express (Invitrogen Cat# 12604021) for 10 min at room

870 temperature. Cell suspensions were passed through a 70-mm cell strainer and then

871 labeled with a cocktail of fluorescent antibodies specific for APC-CD45 (BioLegend,

872 Cat# 103113), FITC-CD31 (BioLegend, Cat# 102406), PE/Cy7-TER-119 (BioLegend,

873 Cat# 116222), and PE-EpCAM (Thermo Fisher Scientific, Cat# 12-5791-83). Dead cells

874 were excluded using Tail Cell Cycle Solution (Invitrogen, Cat# A10798).Following

875 doublet exclusion and single-cell population isolation through forward and side scatter

876 analysis, the Propidium Iodide- (Ex 535nm: Em 617nm) / APC-CD45- (Ex 651nm: Em

877 660nm) cell population was gated. Next, FITC-CD31- (Ex 498nm: Em 517nm) / PE-Cy7-

878 TER119- (Ex 565nm: Em 774nm) cells were gated and subtracted from the PI-/APC-

879 CD45- population. Remaining cells were gated as PE-EpCAM+ (Ex 565nm: Em 576nm)

880 and isolated as CD45-/CD31-/TER119-/EpCAM+ intestinal epithelial cells. CD45-/CD31-

881 /TER119-/EpCAM+ intestinal epithelial cells were sorted by using a SONY SH800S Cell

882 Sorter and collected in a post-sort buffer (RPMI-1640 with 20% FBS) .

884 To isolate crypt from colon, the Colon (from the cecum to the rectum) were dissected

885 out from mice as previously described[33]. After five times washing with ice cold PBS, the

tissues were opened longitudinally and cut into small fragments (2–3 cm in length). To

isolate crypts after washing three times with cold PBS, small ileum fragments were

incubated in 30 mM EDTA-PBS on ice for 30 min, during which the tissues were shaken

vigorously every 5 min. Isolated crypts were washed once with cold PBS and

dissociated using TrypLE Express (Invitrogen Cat# 12604021) for 10 min at room

temperature. Cell suspensions were passed through a 70-µm cell strainer.

InsP$_6$ Rescuing HDAC3 activity.

Wild Type and *IPMK*-Knock out (KO) HCT116 cells were lysed in salt-free (NaCl-free)

lysis buffer (50 mM Tris pH 7.5, 50 mM potassium acetate, 5 % v/v glycerol, 0.3 % v/v

Triton X-100, and one Roche complete protease inhibitor tablet). Samples were

centrifuged at 15,000 g for 10 min, and the protein concentration of the supernatant was

measured. Equal amount, 20 µg total protein from the lysate was incubated against anti-

HDAC3 antibodies respectively at 4°C for 1 hr, followed by capturing antibodies with

EZview A/G beads (Millipore Sigma, Cat #: E3403). Beads were washed 3x with an ice-

cold reaction buffer (supplied with the kit). Following washing, beads were resuspended

in reaction buffer then beads were incubated with 1nM, 10nM, 100nM, 500nM, 1µM,

5µM and 10µM concentrations of InsP$_6$ (Sigma Aldrich Cat #: P8810-100G) for 1 hr at

room temperature. After incubation, beads were washed 3x with a reaction buffer.

HDAC3 activity was measured by the addition of a fluorogenic substrate provided by the

HDAC Assay Kit (BPS, Cat #: 10186-628) and read using a Spectramax iD5 plate

reader (Molecular Devices). IgG was immunoprecipitated from IPMK Wild Type HCT

166 cells (WT) and used as a negative control. In each experiment, separate IgG

immunoprecipitation was performed, and the data was expressed as a fold change

relative to the respective IgG control. Although slight variations in IgG values were

observed across experiments, the overall data range differed slightly in each case, but

consistently showed a similar trend. All measurements were performed in triplicate data

and analyzed using GraphPad Prism (version 6.0, GraphPad Software, Inc)[15,13,16].

Analysis of rescuing HDAC3 activity using PIP3, InsP$_3$, Ins (1,3,4,5) P$_4$,

Ins(1,4,5,6)P$_4$, InsP$_5$ and InsP$_6$**:**

Wild type and *IPMK*-Knock out (KO) HCT116 cells were lysed in salt-free (NaCl-free)

lysis buffer (50 mM Tris pH 7.5, 50 mM potassium acetate, 5 % v/v glycerol, 0.3 % v/v

Triton X-100, and one Roche complete protease inhibitor tablet). Samples were

centrifuged at 15,000 g for 10 min, and the protein concentration of the supernatant was

measured. Equal amount, 20 μg total protein from the lysate was incubated against anti-

HDAC3 antibodies respectively at 4°C for 1 hr, followed by capturing antibodies with

EZview A/G beads (Millipore Sigma, Cat #: E3403). Beads were washed 3x with an ice-

cold reaction buffer (supplied with the kit). Following washing, beads were resuspended

in reaction buffer then beads were incubated with either 10μM concentration of PIP3,

InsP$_3$, Ins(1,3,4,5)P$_4$ or 1μM Ins(1,4,5,6)P$_4$, InsP$_5$ and InsP$_6$ (Sigma Aldrich Cat #:

P8810-100G) for 1 hr at room temperature. After incubation, beads were washed 3x

with a reaction buffer. HDAC3 activity was measured by the addition of a fluorogenic

substrate provided by the HDAC3 Assay Kit (BPS, Cat #: 10186-628) and read using a

Spectramax iD5 plate reader (Molecular Devices). IgG was immunoprecipitated from

Wild Type HCT 166 cells (WT) and used as a negative control. In each experiment,

separate IgG immunoprecipitation was performed, and the data was expressed as a

fold change relative to the respective IgG control. Although slight variations in IgG

values were observed across experiments, the overall data range differed slightly in

each case, but consistently showed a similar trend. All measurements were performed

in triplicate data and analyzed using GraphPad Prism (version 6.0, GraphPad Software,

Inc)[15,13,16].

Plasmids and Recombinant Proteins.

pMX-myc, pMX-*IPMK-WT*-myc and pMX-*IPMK-KSA*-myc plasmids transiently

transfected into *IPMK*-KO HCT 116 cells using Lipofectamine 3000 (Cat no. L3000001,

Thermo Scientific) according to the manufacturer protocol. Recombinant myc-IPMK

was purchased from Origene, (Cat #: TP309343) while recombinant GST-HDAC3 (Cat

#: H85-30G) and GST (Cat #: G52-30U) were purchased from Signal Chem.

Cell fractionation and isolation of chromatin fraction

Approximately 7x106 cells were isolated and washed three times with chilled PBS. After

that cells were lysed with ice cold cell lysis buffer (50 mM HEPES-KOH, 140 mM NaCl,1

mM EDTA, 10% glycerol, 0.5% NP-40, 0.25% Triton X-100, 1 mM DTT Supplemented

with 1x protease inhibitor cocktail). After centrifugation at 1100 g for 5 min supernatant

was separated as a cytoplasmic fraction. Pellets were the crude nuclei. After washing

three time with cell lysis buffer, nuclei was incubated 10 min in ice with nuclear lysis

buffer (10 mM Tris-HCl, 200 mM NaCl, 1 mM EDTA, 0.5 mM EGTA, Supplemented with

1x protease inhibitor cocktail). Gently resuspended 10 times. Then centrifuge at 1100 g

for 10 min. Supernatant was separated as a nuclear fraction and pellet was the crude

chromatin. After washing three times with a nuclear lysis buffer, crude unsheared

chromatin was used for further experiment.

Immunoprecipitation.

To analyze endogenous binding, HCT116 cells were lysed in salt-free lysis buffer (50

mM Tris pH 7.5, 50 mM potassium acetate, 5 % v/v glycerol, 0.3 % v/v Triton X-100,

and one Roche complete protease inhibitor tablet). For the endogenous

immunoprecipitation (IP), respective antibodies against HDAC3 were used, followed by

western blot of the binding partners. In brief, the IP was performed from 500 µug of

protein lysate. Protein lysates were incubated for 2 hr at 4^{o}C with respective antibodies,

then the antibody-protein complex was captured with EZview A/G beads (Millipore

Sigma, Cat #: E3403). Beads were pelleted and washed with lysis buffer 3x, followed by

elution in sample buffer. Immunoprecipitated samples were resolved on a NuPAGE Bis-

Tris gel, followed by western blotting [18]

ChIP-qPCR:

Approximately 7x106 cells were fixed with 1% final volume from fresh 16% stock

formaldehyde (Sigma-Aldrich, Cat#: F8775) at room temperature for 10 min followed by

ChIP using ChIP assay Kit from iDEAL Diagenode (Cat no.C01010051). Cells were

then harvested and lysed in 500 mL of ChIP lysis Buffer (50 mM Tris-HCl pH 8.0, 5 mM

EDTA, 150 mM NaCl, 0.5% Triton X-100, 0.5% SDS, 0.5% NP-40, 1 mM sodium

butyrate) containing protease inhibitor cocktail. The lysates were subjected to sonication

to shear DNA to a length of approximately between 150 and 900 bp. The lysate was

then diluted in 0.75 mL of ChIP dilution buffer and incubated with control IgG (Cell

Signaling Technology, Cat#: 2729S) or primary antibody H3K9 ac (Cells signalling

Technology, Cat#9649T) , H3K27 ac (Abcam cat# 4729) , H4k16ac (Cells signalling

Technology, Cat#13534) , HDAC3 (Santa Cruz Biotechnology, Cat# 39717) and

HDAC1 (Santa Cruz Biotechnology, Cat# 81598X) 4C overnight. Then the lysate was

incubated with Protein G magnetic bead (provided in kit) for 1 h at 4C. The beads were

washed sequentially with a wash buffer provided in the kit. The immune complexes

were eluted with 75 mL of elution buffer (1% SDS, 0.1 M NaHCO3) twice at 65C for 30

min. After elution, the cross-link was reversed by adding NaCl and incubated together

with Proteinase K provided with the kit overnight at 65C. ChIP DNA was purified using

the ChIP DNA purification kit Provided with the kit. The purified DNA was analyzed on a

StepOnePlus using the power SYBR Green Master Mix. The results are presented as a

fold change in respect with IgG after calculating the percentage of input. qPCR analyses

were done in triplicate. We used MMP3 5′-AAAATAGAGTAGCAGAGGCAGGTA -3′ and

5′-AGAGTGGTGGCAGTGATGTGAA -3′, and MMP 13 Fwd 5′

CAACCATGGGGCTCAATCCT3′ Rev 5′ CTTACGTGGCGACTTTTTCTTTTC3′ Primer

for human and MMP3 5′-TGTGGTCACTGATAGTGTGGACTGT -3′ and 5′-

CCAGAAGCCCAAGCTGAATACTGT -3′, and MMP 13 Fwd 5′

CTGCCACAAACCACACTTAGG3′ Rev 5′ AGTCACCACTTTGGGGTGTG3′ Primer for

mouse [18]

mRNA expression by qPCR:

After extracting the total RNA using RNeasy Mini Kit (Qiagen), and checking its integrity

by nanodrop, the cDNA was synthesized from 1 mg of purified total RNA using High

capacity cDNA Reverse Transcription Kit (Applied Biosystem). Expression of mouse

and human MMP1, MMP3 and MMP13, 18S for human and mouse GAPDH was

detected using suitably designed Taqman primers (Invitrogen). The experiments were

performed (real-time PCR Systems StepOne plus, Applied Biosystems) in triplicate.

Data were quantified for the above genes using the comparative Ct method, as

described in the Assays-on-Demand Users Manual (Applied Biosystems)[18]

1009

***In vitro* Binding assay.**

Equal amount of recombinant myc-IPMK (Origene, Cat #: TP309343) was co-incubated

with either GST-HDAC3 (Signal Chem, Cat #: H85-30G) or GST (Signal Chem, Cat #:

G52-30U) in lysis buffer and the complex was maintained for 30 min at 4°C. After the

addition of GST beads, incubation continued for an additional 45 min and washed 3x

with an ice cold lysis buffer. SDS sample buffer was added, and binding was confirmed

by western blotting of anti-Myc antibodies[18].

1017
Transwell assay.

Transwell filter inserts (pore size 1.0 µm, 12 mm diameter, polyester membrane,

Corning, New York, USA) were coated with fibronectin (10 µg/ml; Sigma) used as insert

for 24 well plate. For time dependent permeability assay CaCo2 cells were seeded

(0.25×10^6 cells/well) and grown on transwell filters for 48 hours until reaching

confluency. FITC-dextran 4 kDa-FITC-Dextran (Sigma-Aldrich, Cat no: #FD4-1G) was

added to the top well at 25 mg/ml. at 37°C, and fluorescence was measured after 10

min by using a Spectramax iD5 plate reader (Molecular Devices, (excitation 485

nm/emission 535 nm)[47].

To analyze crucially of IPMK or InsP$_6$ in cell permeability, CaCo2 cells were seeded

$(0.25 \times 10^6$ cells/well) and grown on transwell filters for 48 hours until reaching

confluency. Soup from WT and *IPMK*-KO HCT116 cells and confluent monolayers of

WT and *IPMK*-KO HCT 116 cells were treated with similar amounts of PBS (as a

negative control), Ilomastat (10µM) or cell permeable InsP$_6$ (50µM) or MMP Activator p-

Aminophenylmercuric acetate (5µM) overnight. Soup from those HCT 116 cell cultures

was transferred to mono layer of CaCo2 cells in the transwell plate. Incubate for four

hours. FITC-dextran 4 kDa-FITC-Dextran (Sigma-Aldrich, Cat no: #FD4-1G) was added

to the top well at 25 mg/ml for 10 min at 37°C, and fluorescence was measured using a

Spectramax iD5 plate reader (Molecular Devices, (excitation 485 nm/emission 535 nm).

1037

1038   **RGFP966 treatment**

1039   One million cells WT and *HDAC3* or *IPMK* knockout HCT 116 cells were plated and let

1040   them grow overnight. Next day cells were treated with HDAC3 specific inhibitor

1041   RGFP966 at a dose of 10µM for 24 hr.

1042

CP-InsP$_6$ treatment

Cell permeable InsP$_6$ was created and characterized in Prof. Dr. Henning J. Jessen Lab,

University of Freiburg, Germany[17]. After synthesis cell permeable InsP$_6$ was primarily

dissolved in DMSO to prepare a 10 mM primary stock. One million cells *IPMK* KO HCT

116 cells were plated and let them grow overnight. Next day cells were treated with CP-

InsP$_6$ at a dose of 50µM for 24 hr in media.

1049

Intracellular inositol content analysis using HPLC:

IPMK-WT and *KO* HCT116 cells were plated at a density of 250,000 cells per 60 mm plate, then labeled with 60 mCi (1 Ci = 37 GBq) [3H] myo-inositol (PerkinElmer) in conventional cell culture media for three days. To extract soluble inositol phosphates, cell pellets were suspended in 300 mL of ice-cold 0.6 M perchloric acid buffer (0.1 mg/mL IP6, 2 mM EDTA) and incubated on ice for 1 min. Ninety mL of 1 M potassium carbonate with 5 mM EDTA were added and incubated on ice for 1 h. Extracts were centrifuged at 12,000 rpm for 15 min. The supernatant was collected and analyzed by HPLC using a Partisphere SAX column (Whatman Inc.). The column was eluted with a gradient generated by mixing Buffer A (1 mM EDTA) and Buffer B (Buffer A plus 1.3 M (NH4)2HPO4, pH 3.8 with H3PO4). The 1 mL fractions were collected and counted using 5 mL of Ultima-Flo AP mixture (PerkinElmer).

Unlabelled Inositol phosphates purification from cells and tissue

Inositol phosphates (HOIP) purification was performed as previously described[35]. Briefly, for cell culture HCT116 cells total of around 30-40 million) was used and the total inositol content was normalized by the total cell count. The data is represented as picomole (pM) per 1 million cells. For the mouse intestinal crypts from the Ileum they were washed in cold PBS, and a fraction was set aside for normalization using total protein. The samples were resuspended with cold 1 M perchloric acid and incubated on ice. Titanium dioxide beads were prepared as described (4 mg/sample). Samples were centrifuged and the supernatant was transferred to Eppendorfs containing 4 mg of

1073 titanium dioxide beads and rotated at 4°C for 20 minutes. Samples were centrifuged,

1074 and the beads were washed with cold 1 M perchloric acid three times. Inositol

1075 phosphates were eluted twice by resuspending in ammonium hydroxide and rotating at

1076 4 C for 10 minutes. Purified inositol phosphates were dried on low heat using a

1077 centrifugal evaporator.

1078

Inositol phosphates quantification using mass spectrometry

The analysis was performed on a CE-QQQ system (Agilent 7100 CE-with Agilent

6495C Triple Quadrupole and Agilent Jet Stream electrospray ionization source,

adopting an Agilent CE-ESI-MS interface). An isocratic Agilent 1200 LC pump was used

to deliver the sheath liquid (50% isopropanol in water) with a final splitting flow speed of

10 µl/min via a splitter. All separation was performed with a bare fused silica capillary

with a length of 100 cm (50 µm internal diameter and 365 µm outer diameter). 40 mM

ammonium acetate titrated with ammonium hydroxide to pH 9.0 was used as

background electrolyte (BGE). Between runs of each sample, the capillary was flushed

with BGE for 400s. Samples were injected by applying 100 mbar pressure for 15s (30

nL). The MS source parameters were as follows: gas temperature was 150°C, gas flow

was 11 L/min, nebulizer pressure was 8 psi, sheath gas temperature was 175°C and

with a flow at 8 L/min, the capillary voltage was -2000V, the nozzle voltage was 2000V.

Negative high-pressure RF and negative low-pressure RF were 70 V and 40 V,

respectively. Set multiple reaction monitoring (MRM) as shown in the table below.

1094

Molecular name	Precursor Ion	Product Ion	dwell	Fragmentor (V)	Collision Energy (V)	Cell Accelerator Voltage	Polarity
$[^{13}C_6]$ InsP$_5$	292	504.9	80	166	9	3	Negative
InsP$_5$	289	498.9	80	166	9	3	Negative
$[^{13}C_6]$ InsP$_6$	331.9	79.1	80	166	53	4	Negative
InsP$_6$	328.9	79.1	80	166	53	4	Negative
$[^{13}C_6]$ InsP$_7$	371.9	322.9	80	166	9	3	Negative
$[^{18}O_2]$ 4-InsP$_7$	370.9	319.9	80	166	9	3	Negative
InsP$_7$	368.9	319.9	80	166	9	3	Negativ

e

InsP$_4$	249	418.9	80	166	5	1	Negative
InsP$_3$	418.9	320.8	80	166	17	4	Negative

1095

Internal standard (IS) stock solution of were spiked to samples for the assignment and

quantification of InsPs and PP-InsPs. Quantification of 1,5-InsP$_8$, 5-InsP$_7$, 1-InsP$_7$, and

InsP$_6$ was performed with known amounts of corresponding heavy isotopic references

spiked into the samples. Quantification of 4/6-InsP$_7$ was performed with 13C 5-InsP$_7$

and Quantification of InsP$_{3-4}$ of which no isotopic standards are available was performed

with spiked 13C InsP$_6$. After spiking, 20 µM [^{13}C$_6$] 2-OH InsP$_5$, 20 µM [^{13}C$_6$] InsP$_6$, 4 µM

[^{13}C$_6$] 5-InsP$_7$, 2 µM [^{13}C$_6$] 1-InsP$_7$ and 1 µM [^{13}C$_6$] 1,5-InsP$_8$ were the final concentration

inside samples. All 13C inositol references were kindly provided by Dorothea Fiedler[36].

1104

1105 **Histology and Immunohistochemistry:**

1106

10% Phosphate-buffered-paraformaldehyde fixed intestine and colon tissues was

subsequently opened lengthwise and washed in cold PBS, then cut into approximately

3-4 mm fragments for sectioning and paraffinization. After that tissue was dehydrated

through 70%, 90% 100% and xylene alcohol gradation and subjected to block

preparation in low melting point (65°C) paraffin wax. six micron tissue sections were cut

and spread over Mayer's albumin coated slides. Tissue slides were deparaffinized

and stained with hematoxylin and eosin following standard protocol for histological

evaluation under light microscope.

1115

Mice were euthanized by CO_2. Immediately following euthanasia, the intestine and

colon was removed and flushed with cold PBS. The intestine was subsequently opened

lengthwise and washed in cold PBS, then cut into approximately 3-4 mm fragments for

sectioning and paraffinization. Samples were deparaffinized with Histo-Clear (Thermo

Fisher) and rehydrated in successive washes of 100%, 90%, and 75% ethanol followed

by deionized (DI) water. Sections were unmasked via heat-induced epitope retrieval

(HIER) using citrate buffer, followed by blocking with 3% goat serum for 1 hour. Primary

antibodies were diluted per manufacturer's recommendation in 1% Triton-X containing

blocking solution, incubating overnight at 4°C in a humidified chamber. The following

day, samples were washed with TBST, followed by incubation with fluorescent-tagged

secondary antibodies for 60 minutes at 37°C. After five time washing with TBST,

samples were stained with 4′, 6-diamidino-2-phenylindole (DAPI) diluted in PBS (1

µg/mL). Slides were imaged using an Axiovert Zeiss epifluorescence microscope and

LSM800 confocal microscope. Subsequent histogram analysis in ImageJ produced

signal intensities from 0-255, with 0 = darkest intensity (positive staining) and 255

representing a white, unstained signal. A threshold of 0-230 was applied to eliminate

background staining.

1133

RNA isolation and library preparation.

1135 Total RNA form was isolated by using RNAeasy Mini Kit (Qiagen Cat no.74104) by

1136 following manufacturer's kit protocol. For RNA-seq, libraries were prepared from total

1137 RNA by using True Seq RNA Library Prep Kit V2 (Illumina).

1138

RNA seq analysis.

FASTQ files were retrieved from sequencing. All pre-alignment and post-alignment QA/QC and

data analysis were performed in Partek Flow (v.12.0.0). Reads were aligned to the whole

genome using hg19 assembly with STAR (v.2.7.8a). Aligned reads were quantified to the hg19

annotation model using HTSeq (v.0.11.0). Differential analysis was performed for *IPMK*

knockout versus wild-type, *HDAC3* knockout versus wild-type, and IPMK$^{\Delta IEC}$ versus

IPMK$^{F/F}$ using DESeq2 (v.3.5)31.

1146

Pathway analysis using Metascape.

Pathway analysis was performed on differentially expressed genes (FDR <= 0.05 and

log2FoldChange >= 2) for HCT116 IPMK knockout versus wild-type and Vilcre versus FF using

Metascape (v.3.5) with default settings. The genes were queried against GO Biological

Pathways, KEGG Pathway, Reactome Gene Sets, Canonical Pathways, WikiPathways, and

Panther pathways. Terms with a p-value < 0.01, a minimum count of 3, and an enrichment

factor > 1.5 were considered. All genes were used as a background gene list.

GSEA pathway analysis.

Gene set enrichment analysis (GSEA) was performed using the R package cluster Profiler[37] . We used all the genes after DESeq2 normalization to generate a ranked gene list sorted by Wald's test statistic, then queried relevant pathways containing MMP genes from the C2:CP:Reactome subcollection of the Molecular Signature Database (MSigDB)[38].

ELISA analysis of ILβ and MPO:

ELM-IL1b-1, ELM-MPO-1). Briefly, tissue lysates were incubated in wells precoated with the relevant antibody, and detection was carried out by measuring absorbance at 450nm. Final optical density values were blank-subtracted and normalized to total protein content.

Statistical analysis.

All plots and statistical analyses were performed with Prism 9 (GraphPad) software. Statistical significance was determined by either Student's t-test (two-tailed) for two groups or 1-way ANOVA for multiple groups with similar samples. Error bars represent the standard deviation of the mean and indicate replicates or the number of animals employed. Results were representative of at least three independent experiments (n). Graphs were generated by using either Prism 9 (GraphPad) software or R software.

Reporting summary

Further information on research design is available in the Nature Portfolio Reporting Summary linked to this article.

1176

1177 **Data availability.**

1178 Once the paper published the raw data will be included as supplemental figures and

1179 genomic data will be uploaded in GEO platform.

Acknowledgements:

This work was supported by NIH R16GM154726 and 5P20GM121325 COBRE grant and University of Nevada, Las Vegas start-up funds to Prasun Guha. This work was supported by UK Research and Innovation (UKRI)'s Medical Research Council (MRC) grant MR/T028904/1. We would like to thank NIPM's Genomic Core for assisting with instruments and experiments. The specimen collection and phenotyping was made possible by the Washington University (WU) DDRCC (NIDDK P30 DK052574). Parakkal Deepak was supported by the Helmsley Charitable Trust, the IBD Plexus of the Crohn's and Colitis Foundation, and the Leo & Carean Goss Crohn's Disease Research Fund. We would like to thank Marc C Jhonson, from University of Missouri-School of Medicine for generously providing *IPPK* Knock out cells. We would like to thank Mark Donowitz from Johns Hopkins Medicine, Subrata H Mishra from NIST and Brian Hedlund from UNLV for reading the paper and giving suggestions.

Competing interests:

Parakkal Deepak, MBBS MS has received research support under a sponsored research agreement unrelated to the data in the study and/or consulting from Johnson and Johnson, Pfizer, AbbVie, Arena Pharmaceuticals, Bristol Myers Squibb, CorEvitas LLC, Sandoz, Takeda Pharmaceuticals, Direct Biologics, Prometheus Biosciences, Lilly, Teva Pharmaceuticals, Merck, ExeGI Pharmaceuticals, AGMB, Landos Pharmaceuticals, Tr1X, and Boehringer Ingelheim. The remaining authors declare no conflict of interest. All authors reviewed the results and approved the final version of the manuscript.

Author Contributions:

PG conceived the study. PG and SC designed the experiments. SC performed most

experiments. LVP validated major biochemical experiments. SC performed wet-lab

experiments related to NGS studies. ZS and LVP performed most of the NGS data

analysis. KR and HJJ generated the cell-permeable InsP$_6$. XBS and AS analyzed

intercellular inositol content. NT and SC performed animal and microscopy related

experiments. QL performed NGS data validation. SJR maintained animal colonies and

isolated animal samples for genotyping. SP performed mathematical analysis of

transwell assay. SC and ZS designed the Figures. PG and SC wrote the manuscript.

SC takes responsibility of all wet lab data.

1214

1215

References

1. Watson PJ, Fairall L, Santos GM, Schwabe JWR. Structure of HDAC3 bound to co-repressor and inositol tetraphosphate. *Nature*. 2012;481(7381):335–340. doi: 10.1038/nature10728.

2. Li Y, Seto E. HDACs and HDAC inhibitors in cancer development and therapy. *Cold Spring Harb Perspect Med*. 2016;6(10):a026831. doi: 10.1101/cshperspect.a026831. doi: 10.1101/cshperspect.a026831.

3. Subramanian S, Bates SE, Wright JJ, Espinoza-Delgado I, Piekarz RL. Clinical toxicities of histone deacetylase inhibitors. *Pharmaceuticals (Basel)*. 2010;3(9):2751–2767. doi: 10.3390/ph3092751.

4. Feng D, Liu T, Sun Z, et al. A circadian rhythm orchestrated by histone deacetylase 3 controls hepatic lipid metabolism. *Science*. 2011;331(6022):1315–1319. doi: 10.1126/science.1198125.

5. Alenghat T, Osborne LC, Saenz SA, et al. Histone deacetylase 3 coordinates commensal-bacteria-dependent intestinal homeostasis. *Nature*. 2013;504(7478):153–157. doi: 10.1038/nature12687.

6. Watson PJ, Millard CJ, Riley AM, et al. Insights into the activation mechanism of class I HDAC complexes by inositol phosphates. *Nat Commun*. 2016;7:11262. doi: 10.1038/ncomms11262.

7. Guha P, Tyagi R, Chowdhury S, et al. IPMK mediates activation of ULK signaling and

transcriptional regulation of autophagy linked to liver inflammation and regeneration.

Cell Rep. 2019;26(10):2692–2703.e7. doi: 10.1016/j.celrep.2019.02.013.

8. Kim S, Bhandari R, Brearley CA, Saiardi A. The inositol phosphate signalling network

in physiology and disease. *Trends Biochem Sci*. 2024;49(11):969–985. doi:

10.1016/j.tibs.2024.08.005.

9. Millard CJ, Watson PJ, Celardo I, et al. Class I HDACs share a common mechanism

of regulation by inositol phosphates. *Mol Cell*. 2013;51(1):57–67. doi:

10.1016/j.molcel.2013.05.020.

10. He R, Liu B, Geng B, Li N, Geng Q. The role of HDAC3 and its inhibitors in

regulation of oxidative stress and chronic diseases. *Cell Death Discov*. 2023;9(1):131–

w. doi: 10.1038/s41420-023-01399-w.

11. Kumar A, Singh B, Raigond P, et al. Phytic acid: Blessing in disguise, a prime

compound required for both plant and human nutrition. *Food Res Int*. 2021;142:110193.

doi: 10.1016/j.foodres.2021.110193.

12. O'Sullivan S, Gilmer JF, Medina C. Matrix metalloproteinases in inflammatory bowel

disease: An update. *Mediators Inflamm*. 2015;2015:964131. doi: 10.1155/2015/964131.

13. Reilly L, Semenza ER, Koshkaryan G, et al. Loss of PI3k activity of inositol

polyphosphate multikinase impairs PDK1-mediated AKT activation, cell migration, and

intestinal homeostasis. *iScience*. 2023;26(5):106623. doi: 10.1016/j.isci.2023.106623.

14. Naito Y, Yoshikawa T. Role of matrix metalloproteinases in inflammatory bowel

disease. *Mol Aspects Med.* 2005;26(4-5):379–390. doi: 10.1016/j.mam.2005.07.009.

15. Rumbaugh G, Daws SE, Ozkan ED, et al. Pharmacological selectivity within class I

histone deacetylases predicts effects on synaptic function and memory rescue.

Neuropsychopharmacology. 2015;40(10):2307–2316. doi: 10.1038/npp.2015.93.

16. Nguyen HCB, Adlanmerini M, Hauck AK, Lazar MA. Dichotomous engagement of

HDAC3 activity governs inflammatory responses. *Nature.* 2020;584(7820):286–290. doi:

10.1038/s41586-020-2576-2.

17. Pavlovic I, Thakor DT, Bigler L, et al. Prometabolites of 5-diphospho-myo-inositol

pentakisphosphate. *Angew Chem Int Ed Engl.* 2015;54(33):9622–9626. doi:

10.1002/anie.201503094.

18. Ning L, Rui X, Bo W, Qing G. The critical roles of histone deacetylase 3 in the

pathogenesis of solid organ injury. *Cell Death Dis.* 2021;12(8):734–6. doi:

10.1038/s41419-021-04019-6.

19. Massimino L, Lamparelli LA, Houshyar Y, et al. The inflammatory bowel disease

transcriptome and metatranscriptome meta-analysis (IBD TaMMA) framework. *Nat

Comput Sci.* 2021;1(8):511–515. doi: 10.1038/s43588-021-00114-y.

20. Barberio B, D'Inca R, Facchin S, et al. Matrix metalloproteinase 3 predicts

therapeutic response in inflammatory bowel disease patients treated with infliximab.

Inflamm Bowel Dis. 2020;26(5):756–763. doi: 10.1093/ibd/izz195.

21. Iaia N, Canzoneri F, Biasi F, et al. The CaCo-2 cell junction derangement exerted by

the single addition of oxysterols commonly detected in foods is markedly quenched

when they are in mixture. *J Steroid Biochem Mol Biol*. 2025;246:106648. doi:

10.1016/j.jsbmb.2024.106648.

22. Horowitz A, Chanez-Paredes SD, Haest X, Turner JR. Paracellular permeability and

tight junction regulation in gut health and disease. *Nat Rev Gastroenterol Hepatol*.

2023;20(7):417–432. doi: 10.1038/s41575-023-00766-3.

23. Tsai P, Zhang B, He W, et al. IL-22 upregulates epithelial claudin-2 to drive diarrhea

and enteric pathogen clearance. *Cell Host Microbe*. 2017;21(6):671–681.e4. doi:

10.1016/j.chom.2017.05.009.

24. Odenwald MA, Turner JR. The intestinal epithelial barrier: A therapeutic target? *Nat*

Rev Gastroenterol Hepatol. 2017;14(1):9–21. doi: 10.1038/nrgastro.2016.169.

25. Grases F, Simonet BM, Vucenik I, et al. Absorption and excretion of orally

administered inositol hexaphosphate (IP(6) or phytate) in humans. *Biofactors*.

2001;15(1):53–61. doi: 10.1002/biof.5520150105.

26. Vucenik I, Shamsuddin AM. 3H]inositol hexaphosphate (phytic acid) is rapidly

absorbed and metabolized by murine and human malignant cells in vitro. *J Nutr*.

1994;124(6):861–868. doi: 10.1093/jn/124.6.861.

27. Grases F, Simonet BM, Vucenik I, Perello J, Prieto RM, Shamsuddin AM. Effects of

exogenous inositol hexakisphosphate (InsP(6)) on the levels of InsP(6) and of inositol

trisphosphate (InsP(3)) in malignant cells, tissues and biological fluids. *Life Sci.*

2002;71(13):1535–1546. doi: 10.1016/s0024-3205(02)01927-6.

28. Badodi S, Pomella N, Zhang X, et al. Inositol treatment inhibits medulloblastoma

through suppression of epigenetic-driven metabolic adaptation. *Nat Commun.*

2021;12(1):2148–7. doi: 10.1038/s41467-021-22379-7.

29. Wu S, Hashimoto-Hill S, Woo V, et al. Microbiota-derived metabolite promotes

HDAC3 activity in the gut. *Nature.* 2020;586(7827):108–112. doi: 10.1038/s41586-020-

2604-2.

30. Williams KL, Fuller CR, Dieleman LA, et al. Enhanced survival and mucosal repair

after dextran sodium sulfate-induced colitis in transgenic mice that overexpress growth

hormone. *Gastroenterology.* 2001;120(4):925–937. doi: 10.1053/gast.2001.22470.

31. Vucenik I, Shamsuddin AM. Cancer inhibition by inositol hexaphosphate (IP6) and

inositol: From laboratory to clinic. *J Nutr.* 2003;133(11 Suppl 1):3778S–3784S. doi:

10.1093/jn/133.11.3778S.

32. Tsai P, Zhang B, He W, et al. IL-22 upregulates epithelial claudin-2 to drive diarrhea

and enteric pathogen clearance. *Cell Host Microbe.* 2017;21(6):671–681.e4. doi:

10.1016/j.chom.2017.05.009.

33. Ge Y, Zadeh M, Mohamadzadeh M. Dissociation and flow cytometric isolation of

murine intestinal epithelial cells for multi-omic profiling. *STAR Protoc.* 2023;4(1):101936.

doi: 10.1016/j.xpro.2022.101936.

34. Aouameur R, Da Cal S, Bissonnette P, Coady MJ, Lapointe JY. SMIT2 mediates all

myo-inositol uptake in apical membranes of rat small intestine. Am J Physiol

Gastrointest Liver Physiol. 2007 Dec;293(6):G1300-7. doi: 10.1152/ajpgi.00422.2007.

Epub 2007 Oct 11. PMID: 17932225.

35. Wilson MS, Saiardi A. Inositol Phosphates Purification Using Titanium Dioxide

Beads. Bio Protoc. 2018 Aug 5;8(15):e2959. doi: 10.21769/BioProtoc.2959. PMID:

30148188; PMCID: PMC6108412.

36. Harmel RK, Puschmann R, Nguyen Trung M, Saiardi A, Schmieder P, Fiedler D.

Harnessing 13C-labeled myo-inositol to interrogate inositol phosphate messengers by

NMR. Chem Sci. 2019 Apr 23;10(20):5267-5274. doi: 10.1039/c9sc00151d. PMID:

31191882; PMCID: PMC6540952.

37. Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R package for comparing biological

themes among gene clusters. OMICS. 2012 May;16(5):284-7. doi:

10.1089/omi.2011.0118. Epub 2012 Mar 28. PMID: 22455463; PMCID: PMC3339379.

38. Liberzon A, Subramanian A, Pinchback R, Thorvaldsdóttir H, Tamayo P, Mesirov JP.

Molecular signatures database (MSigDB) 3.0. Bioinformatics. 2011 Jun 15;27(12):1739-

40. doi: 10.1093/bioinformatics/btr260. Epub 2011 May 5. PMID: 21546393; PMCID:

PMC3106198.

1333

1334

 Figure Legends

1336

Figure 1: IPMK binds chromatin-bound HDAC3, regulating its activation and

associated epigenetic changes. (A) Pictorial representation demonstrates the function

of IPMK in the higher-order inositol phosphate (HOIP) pathway. IPMK converts $InsP_3$ to

$InsP_4$ and $InsP_4$ to $InsP_5$, while IPPK converts $InsP_5$ to $InsP_6$. **(B)** HPLC analysis of

inositol phosphate level depicted *IPMK* deletion reduced $InsP_5$ and $InsP_6$ levels in HCT

116 cells. The blue line indicated Wild type HCT116 cells (WT), and the red line

indicated *IPMK*- Knock out HCT 116 cells (KO). Result was representative of two

individual experiments (n=2). **(C)** Schematic diagram of HDAC3 activity assay.

Immunoprecipitation of HDAC3 from cell lysate followed incubation with substrate and

measuring HDAC3 activity fluorometrically. **(D)** HDAC3 were immunoprecipitated from

Wild Type (WT) and *IPMK*- Knock out (KO) HCT116, MEF and A549 cells followed by

HDAC3 activity assay. HDAC3 activity was markedly decreased in *IPMK*-Knock out

(KO) HCT116, MEF and A549 cell lines compared to respective Type (WT) cells (HCT

116 n=12, MEF n=4, A549 n=4). Data was represented as a fold change compared to

IgG. Results are representative of minimum four individual experiments. Percentage

decrease of HDAC3 activity in IPMK KO cells compared to the WT was shown

respective to each cell line. **(E)** Deletion of *IPMK* diminished HDAC3 activity in HCT 116

cells which is comparable to CRISPR-knock out *HDAC3* cells or WT cells treated with

the HDAC3-specific inhibitor RGFP966 (10 µM for 24 hrs). Data was represented as a

fold change compared to IgG. Results are representative of minimum three individual

experiments (n=3). **(F)** Western blot analysis revealed that the deletion of *IPMK* or

HDAC3 led to profound increased acetylation of H3K9, H3K27, and H4K16 compared to

wild-type cells. The levels of acetylation observed in both mutant cell types were

comparable. An 8-hour treatment with the HDAC3-specific inhibitor RGFP966 at a dose

of 10 µM in HCT116 WT, *IPMK* KO, and *HDAC3* KO cells showed no further increase in

H3K9, H3K27, and H4K16 acetylation compared to the untreated *IPMK* KO and *HDAC3*

KO cells. Lysates were immunoblotted against acetylated anti-H3K9, anti-H3k27 , anti-

H4k16 and HDAC3 antibodies and followed by stripping and reprobing against anti-H3

or anti-H4 antibody and actin, respectively, used as loading controls. Data is

representative of three experimental replicates (n=3). **(G)** Representative densitometric

analysis of H3k9ac / H3, H3k27ac / H3 and H4k16ac / H3. (n=3). **(H)** Schematic

diagram of HDAC3 chromatin immuno precipitation. **(I)** HDAC3 was immunopurified

using an anti-HDAC3 antibody from the chromatin fraction, and cytoplasmic fraction,

followed by western blot analysis with an anti-IPMK antibody confirmed that IPMK

exclusively binds to HDAC3 in chromatin fraction but not with cytoplasmic HDAC3. IgG

was used as a negative control. Immunoblotting against anti-HDAC3, anti-IPMK, anti-H3

antibodies from the chromatin fraction isolated from *IPMK-* Wild type HCT116 (WT) and

IPMK- Knock out HCT 116 (KO) cells were used as input control while anti-RNA pol II

and anti- E-Cadherin used as cell fractionation control. Immunoblot with anti-HDAC3

from the bead used as loading control. The result was representative of three individual

experiments (n=3).

1378

Extended Figure 1: (A) Western blot analysis of the lysates showed complete loss of

IPMK protein in three CRISPR Knock out cell lines (HCT116, MEF and A5490 as

compared to lysate isolated from respective Wild type (WT) HCT116 cells. Actin was

used as a loading control. Results were representative of three individual experiments

(n=3). **(B)** HDAC3 were immunoprecipitated from 5µM, 20µM, 50µM, 100µM and

500µM Wild Type (WT) HCT 116 cell lysate followed by HDAC3 activity assay. Data

was represented as a fold change compared to IgG. Results are representative of

minimum three individual experiments. (n=3) **(C)** Western blot analysis of the lysates

isolated from WT and *IPMK* Knock out (KO) HCT 116 cells showed HDAC3 and its

corepressor protein NCoR1/2 levels were comparable in all genotypes. Actin was used

as a loading control. Results were representative of three individual experiments (n=3).

(D) Western blot analysis of the lysates showed complete loss of HDAC3 protein in

CRISPR Knock out cell lines HCT116, as compared to lysate isolated from respective

Wild type (WT) HCT116 cells. Actin was used as a loading control. Results were

representative of three individual experiments (n=3). **(E,F)** Western blot analysis

demonstrated that histone acetylation increased in *IPMK*- Knock out -HCT 116 cells

(KO) compared to the Wild Type HCT 116 cells (WT). *IPMK* deletion specifically

increased acetylated H3K9, and 27 and H4k16. Marked by * , denoted high alteration

level. Data represents three experimental replicates (n=3). Followed by densitometric

analysis. **(G)** After incubation with recombinant HDAC3-GST, and IPMK-myc the

sample was immunoprecipited with myc bead and western blotted against anti-GST

antibody. Result showed recombinant IPMK binds with HDAC3. GST protein was used

as control. Results was representative of three experimental replicates (n=3).

Figure 2: Kinase activity of IPMK is essential for activating HDAC3

(A , B) *IPMK*-Knockout (KO) HCT116 cells transiently overexpressed with *myc-control*, *WT-IPMK*-myc*,* and *KD-IPMK*-myc plasmid. Results demonstrated that *myc-control* and *IPMK-KD*-myc *were* unable to rescue HDAC3 activity while wt-IPMK-myc rescued. Data was represented as a fold change compared to IgG. Results were representative of three independent experiments (n=3).**(B)** Immunoprecipitation of IPMK-myc followed by western blot of NCoR1,NCoR2,HDAC3, GPS2, TBL1 indicates that the interaction of HDAC3,TBL1, GPS2 with NCoR1/2 in *IPMK* KO HCT 116 cells was comparable in *KD-IPMK*-myc as compared to *WT-IPMK*-myc. Data represents three experimental replicates (n=3). **(C)** To analyze the role of $InsP_6$ in activating HDAC3 activity *in vitro*, endogenous HDAC3 were immunopurified from Wild type HCT116 (WT) and *IPMK*-Knock out HCT 116 (KO) cells followed by dose-dependent (0.001 μM, 0.01 μM, 0.1 μM, 0.5 μM, 1 μM, 5 μM and 10 μM) *in vitro* $InsP_6$ treatment. Data was represented as a fold change compared to respective IgG. Results were representative of three independent experiments (n=3). 54.45% increase of HDAC3 activity after 0.01 μM of $InsP_6$ treatment in IPMK KO cells. **(D)** To analyze the impact of other HOIPs and PIP3 in HDAC3 activity, endogenous HDAC3 were immunopurified from Wild Type HCT116 (WT) and *IPMK*- Knock out HCT 116 (KO) cells followed by *in vitro* 10μM treatment with PIP3, $InsP_3$, Ins $(1,3,4,5)P_4$ and 1μM of $InsP_3$, Ins $(1,4,5,6)P_4$, InsP5 and $InsP_6$. Data was represented as a fold change compared to IgG, following blank subtraction and normalization. Results were representative of three independent experiments (n=3). **(E)** 24-hour treatment with 50 μM cell-permeable $InsP_6$ (CP-$InsP_6$) in IPMK-knockout (KO) HCT116 cells significantly restored HDAC3 activity comparable to untreated wild-type

(WT) HCT116 cells. The results showed that 95.64% of HDAC3 activity was rescued in

IPMK-KO cells after 24 hr of 50 µM CP-InsP$_6$ treatment, relative to IPMK KO cells. Data

was represented as a fold change compared to IgG. Results were representative of

three independent experiments (n=3). **(F)** Cell-permeable InsP$_6$ (CP-InsP$_6$) treatment at

a dose of 50 µM for 24 hr successfully rescued H3K9, H3K27 and H4K16 acetylation as

comparable with untreated WT cells which HDAC3 inhibitor RGFP966 treatment

abrogated. Data represents three experimental replicates (n=3).

Extended Figure 2: (A) Next, efficacy of 0.01µM InsP$_4$ and InsP$_5$ analyzed compared

to InsP$_6$ in activating HDAC3 activity *in vitro*. Data was represented as a fold change

compared to respective IgG. Results were representative of three independent

experiments (n=3). **(B)** 50 µM cell permeable InsP$_6$ for 24 hr failed to increase HDAC3

activity when treated to WT cells. Data was represented as a fold change compared to

IgG. Results are representative of three individual experiments (n=3). **(C)**Mass

spectrometric analysis of HOIP level after IPMK KO HCT116 cells treated with CP-InsP$_6$

for 24 hours.(n=3)

Figure 3: Mechanism of IPMK mediated HDAC3 activation *in vivo* and cell free chromatin fraction: (A,B) Proximity ligation assay using anti-HDAC3 and anti-NCoR2 antibody showed comparable amount of puncta (RED) in the WT and *IPMK* KO HCT 116 cells. Images were representative of three individual experiments (n=50). **(C)** Pictorial representation of NCoR2 protein domains, which includes the SANT1, SANT2 and DAD domains. The model represents that interaction of DAD of NCoR2 is crucial for the enzymatic activity of HDAC3, but does not influence HDAC3: corepressor complex integrity. **(D)** WT and *IPMK*-Knockout (KO) HCT116 cells transiently overexpressed with FLAG-control, and DAD-FLAG plasmid. FLAG was immunoprecipitated by WT and IPMK KO HCT 116 cells, followed by immunoblotting against anti-HDAC3 antibodies. Results demonstrated that DAD-FLAG failed to bind with endogenous HDAC3 when over expressed in *IPMK* KO cells while DAD-FLAG overexpressed in WT HCT 116 cells successfully to binds with endogenous HDAC3. Lysate control and bead inputs were represented. The result was representative of three individual experiments (n=3). **(E)** Treatment of *IPMK*-Knockout (KO) HCT116 cells with cell permeable $InsP_6$ (CP-$InsP_6$) at a dose of 50 µM for 24 hr,restored Flag-DAD binding to endogenous HDAC3. The result was representative of three individual experiments (n=3). **(F, G)** Proximity ligation assay with anti-HDAC3 and anti-H3 antibody showed comparable amount of puncta (RED) in the WT and *IPMK* KO HCT 116 cells. Results were representative of three individual experiments. Scale bar 5 µM (n=35). **(H)** Picture represents that HDAC3:NCoR2 complex binds to chromatin independent of $InsP_6$; however, remains enzymatically inactive because DAD domain does not interact with HDAC3; presence of $InsP_6$ drives DAD domain HDAC3 interaction

activating HDAC3's deacetylase activity. **(I)** Schematic diagram of chromatin

purification and *in vitro* InsP$_6$ treatment. InsP$_6$ treatment to cell free chromatin fraction

isolated from IPMK KO cells repressed H3k9/27 and H4k16 acetylation which HDAC3

inhibitor RGFP966 treatment abrogated. Immunoblotting against anti-RNA POL 2

(chromatin) and E Cadherin (cytoplasmic and cell membrane) antibody used as cell

fractionation control. Data represents three experimental replicates (n=3).

Extended Figure 3 I: (A) Immunoprecipitation of HDAC3 followed by western blot of

various corepressor proteins (NCoR1/2) indicates that the interaction of HDAC3 and

NCoR1/2 in *IPMK* KO HCT 116 cells was comparable to WT HCT 116 cells. The result

was representative of three individual experiments. (n=3). **(B)** The interaction of HDAC3

and H3/H4 in IPMK KO HCT 116 cells was comparable to WT HCT 116 cells. The result

was representative of three individual experiments. (n=3). **(C)** Immunoprecipitation of

NCoR2/1 from cell free chromatin followed by western blot of HDAC3, GPS2, TBL1

indicates that the interaction of HDAC3, TBL1, GPS2 with NCoR1/2 in *IPMK* KO HCT

116 cells was comparable to WT HCT 116 cells. Immunoblotting against anti RNA POL

2 (chromatin) and E Cadherin (cytoplasmic and cell membrane) antibody used as cell

fractionation control. Data represents three experimental replicates (n=3).

1480

Extended Figure 3 II: (A) Immunoprecipitation of HDAC3 followed by western blot of

IPPK indicates that the interaction of HDAC3 and IPPK in *IPMK* KO HCT 116 cells was

comparable to WT HCT 116 cells. The result was representative of three individual

experiments. (n=3). **(B)** Proximity ligation assay using anti-HDAC3 and anti-IPPK

antibodies showed RED PLA puncta in the nucleus of WT HCT 116 cells. Scale bar

5μM (n=35).

1487

1488

1489

1490

1491

1492

1493

1494

1495

1496

1497

1498

1499

Figure 4: Deletion of *IPMK* stimulates hypertranscription of MMP gene family

(A) Volcano plot (Log 10[*FDR*] vs. Log 2[Fold change]) displays differentially expressed genes after *IPMK* deletion in HCT116 cells (*IPMK-KO*) as compared to WT-HCT116 cells (*WT*). Red dots represent gene expression that was upregulated, while blue dots represent gene expression that was downregulated in *IPMK* KO cells. Y-axis denotes – Log10 FDR Value, while the X-axis denotes the Log2 fold change value. The result was representative of three individual experiments (n=3). **(B)** Pathway analysis of RNA-seq data identified common genes that are upregulated and down regulated following *IPMK* deletion in HCT 116 (KO) cells. **(C)** Western blot analysis showed MMP1, 10, and 13 markedly increased in *IPMK* KO HCT116 cells. (n=3) **(D)** Western blot of MMP1, 10 and 13 from IPMK WT and KO cells. 24 hr of 50 µM CP-InsP$_6$ treatment to KO cells successfully rescued MMP1, MMP10 and MMP 13 expression while co treatment with 10 µM RGFP966 abrogated InsP$_6$'s rescue effect. Actin used as loading control. Data represents three experimental replicates (n=3). **(E)** Lysate isolated from WT, *IPMK-KO*, and *HDAC3* KO HCT 116 cells were western blotted against anti-MMP1, anti-MMP10 and anti-MMP13 antibody. Result demonstrated as compared to WT HCT 116 cells (WT) MMP1, MMP10 and MMP 13 expression was increased in *HDAC3* KO HCT 116 cells (*HDAC3* KO) which is comparable with *IPMK* KO HCT116 (*IPMK* KO) cells. Actin used as loading control. Data represents three experimental replicates (n=3). **(F)** Western blot analysis revealed that the deletion of *IPMK or HDAC3* led to profound increased MMP1, MMP10 and MMP13 expression as compared to WT HCT116 cells. An 8-hour treatment with the HDAC3-specific inhibitor RGFP966 at a dose of 10 µM in HCT116 WT, *IPMK* KO, and *HDAC3* KO cells showed no further increase in MMP1,

MMP10, MMP13 expression compared to the untreated *IPMK* KO and *HDAC3* KO

HCT116 cells. Actin used as loading control. Data represents three experimental

replicates (n=3). **(G)** Representative densitometric analysis. Data represents three

experimental replicates (n=3). **(H)** Pictorial representation showing MMP13 primer

locations on the human genome. Chromatin immunoprecipitation against acetylated

H3K9, H3K27 and H4K16 followed by qPCR analysis using primers for MMP13

promoter. Results depicted that *IPMK* deletion selectively induced promoter enrichment

of acetylated H4k16, but not for acetylated H3k9/27. **(I)** MMP13 promoter enrichment

was significantly rescued in *IPMK*-KO cells after 24 hr 50 µM cell permeable InsP$_6$ (CP-

InsP$_6$) treatment. Co-treatment of 10 µM RGFP966 with 50 µM cell permeable InsP$_6$

(CP-InsP$_6$) abrogated InsP$_6$'s rescue effect (n=3). **(J)** Overlap of differentially expressed

RNA seq data compared with *IPMK* KO and *HDAC3* KO cells.

Extended Figure 4 I: (A) Volcano plot (Log 10[*FDR*] vs. Log 2[Fold change]) displays

differentially expressed genes in Crohn's disease patients from the Inflammatory Bowel

Disease Transcriptome and Metatranscriptome Meta-Analysis (IBD TaMMA) database

as compared with healthy individuals. Red dots represent gene expression upregulated

in Crohn's patient ileum, while blue dots represent down regulated gene expression.

The Y-axis denotes the –Log10 FDR Value, while the X-axis denotes the Log2 fold

change value. The result represented an expression of various MMP genes high in

Crohn's disease patient's ileum. **(B)** *IPMK*-Knockout (KO) HCT116 cells transiently

overexpressed with *myc-control*, *wt-IPMK*myc*,* and *KD-IPMK*-myc plasmid. Western blot

analysis demonstrated that *myc-control* and *KD-IPMK*-myc *were* unable to rescue

MMP1, MMP10 or MMP13 expression while wt-IPMK-myc was rescued*.* Actin, was

used as a loading control respectively. Results were representative of three

independent experiments (n=3). **(C)** qPCR analysis of mRNA isolated from WT and

HDAC3 KO HCT 116 (HDAC3 KO) cells indicated increased MMP1, MMP3 and MMP13

gene transcription. 18s used as loading control. Results represented three experimental

replicates (n=3). **(D)** ChIP-qPCR analysis using anti-HDAC3 and anti-HDAC1 antibodies

showed MMP13 promoter enrichment was exclusively on HDAC3 and comparable in

WT and *IPMK* KO cells. Data represents three experimental replicates (n=3). **(E)** qPCR

analysis of mRNA isolated from WT and *IPMK* KO HCT 116 (KO) cells indicated

increased MMP3 transcription. 18s used as loading control. Results represented three

experimental replicates (n=3). **(F)** Western Blot analysis with anti-MMP3 antibody

showed profound increase in MMP3, expression at protein level after *IPMK* deletion in

HCT 116 cells (KO) as compared to WT and **(G)** was comparable to *HDAC3* KO HCT

116 cells (HDAC3 KO). Actin used as loading control. Data represents three

experimental replicates (n=3). **(H)** Western blot analysis revealed that the deletion of

IPMK or HDAC3 led to profound increased MMP3 expression as compared to WT

HCT116 cells. An 8-hour treatment with the HDAC3-specific inhibitor RGFP966 at a

dose of 10 µM in HCT116 WT, IPMK KO, and HDAC3 KO cells showed no further

increase in MMP3 expression compared to the untreated *IPMK* KO and *HDAC3* KO

HCT116 cells. Actin used as loading control. Data represents three experimental

replicates (n=3). **(I)** Western blot of MMP3 from IPMK WT and KO cells. 24 hr of 50 µM

CP-InsP$_6$ treatment to KO cells successfully rescued MMP3 expression while co

treatment with 10 µM RGFP966 abrogated InsP$_6$'s rescue effect. Actin used as loading

control. Data represents three experimental replicates (n=3).

1569

1570

Extended Figure 4 II: (A,B) Volcano plot (Log 10[*FDR*] vs. Log 2[Fold change])

displays differentially expressed genes in Crohn's disease and ulcerative colitis patients

from the Inflammatory Bowel Disease Transcriptome and Metatranscriptome Meta-

Analysis (IBD TaMMA) database as compared with healthy individuals. Red dots

represent gene expression upregulated in patient tissue, while blue dots represent

downregulated gene expression. The Y-axis denotes the –Log10 FDR Value, while the

X-axis denotes the Log2 fold change value. The result represented   increased

expression of various MMP genes.

1579

1580

Figure 5: *IPMK* deletion markedly increased cell and intestinal permeability in

mice. (A) Schematic diagram of transwell cell permeability assay. **(B)** Conditioned

media transferred from *IPMK and HDAC3* KO cells to the Caco2 (WT) monolayer

elevated cell permeability. Treatment of MMP activator (p-Aminophenylmercuric

acetate) to WT HCT16 cell followed by conditioned media transfer also increased

Caco2 cell permeability. Treatment of HDAC3 inhibitor RGFP966 to WT HCT16 cell

followed by conditioned media transfer also increased Caco2 cell permeability.

Treatment of CP-InsP$_6$ to *IPMK* KO and *HDAC3* KO cells followed by conditioned media

transfer to CaCo2 failed to induce cell permeability. RGFP966 co-treatment with InsP$_6$

to *IPMK* KO cells followed by conditioned media transfer increased CaCo2 cell

permeability Result was representative of three independent experiments. (n=3). **(C)**

Pictorial representation of Pore, Leak and Unrestricted intestinal permeability pathway.

Pore pathway does not allow 4kDa Dextran while Leak pathway allows 4kDa dextran

permeability but not 70 kDa. Unrestricted pathway is activated due to cell death or cell

damage allows both 4kDa and 70 kDa Dextran. **(D)** Intestinal permeability assay

indicated a significant increase in 4 kDa FITC Dextran fluorescence (not 70 kDa

Rhodamine) in IPMK$^{\Delta IEC}$ mice serum compared to the IPMK $^{F/F}$ Interestingly, 3 days

pretreatment of 2% body weight of oral InsP$_6$ to IPMK$^{\Delta IEC}$ reduced intestinal 4 kDa FITC

Dextran leak. After 4 hr of fasting a single dose of 4 kDa FITC-Dextran was gavaged at

0.6mg/gm body weight. The untreated control group received an equal amount of PBS.

Fluorescence intensity of plasma samples was measured (excitation 485 nm/emission

535 nm) after 4 hours of gavage. Data were expressed as fold change relative to control

mice serum, which received only PBS gavaged mice without FITC or Rhodamine

Dextran. Result was representative of three independent experiments (n=5). **(E)**

Volcano plot (Log 10[*FDR*] vs. Log 2[Fold change]) displays differentially expressed

genes in IPMK$^{\Delta IEC}$ as compared with IPMK$^{F/F}$ mice. Red dots represent gene expression

that was upregulated in IPMK$^{\Delta IEC}$, while blue dots represent gene expression that was

downregulated. The Y-axis denotes –the Log10 FDR Value, while the X-axis denotes

the Log2 fold change value. The result was representative of three individual

experiments (n=3). **(F)** qPCR analysis of mRNA isolated from intestinal epithelial cells of

IPMK$^{F/F}$ and IPMK$^{\Delta IEC}$ cells indicated increased MMP3 and MMP13 gene transcription.

18s used as loading control. Results represented three experimental replicates (n=3).

(G) Western blot from intestinal crypts (Ileum) showed increased MMP3 and 13

expression in IPMK$^{\Delta IEC}$ mice while 72 hr 2% b.w InsP$_6$ treatment reduced it. ZO-1,

occludin level decreased with Caludin 2 level increased in IPMK$^{\Delta IEC}$ mice; however, the

level was restored by oral InsP6. The result was representative of three independent

experiments (n=3). **(H)** Immunofluorescence microscopy showed increased MMP3

expression (Green) in ileum IPMK$^{\Delta IEC}$ as compared to IPMK$^{F/F}$ .DNA (blue). 2% body

weight (B.W) of InsP$_6$ treatment for 72hr to IPMK$^{\Delta IEC}$ completely reduced MMP3

expression. Scale bar represents 50 µm.Data was representative of three individual

experiments. (n=3) (Total number of images=30). **(I,J)** Showing confocal images of

colon ZO-1 (red pseudo color) and Occludin (white pseudo color) showing loss of its

expression in IPMK$^{\Delta IEC}$ and restored by InsP$_6$. Scale bar 50 μm.**(K)** HDAC3 activity was

diminished in intestinal epithelial cells isolated from IPMK$^{\Delta IEC}$ compared with IPMK$^{F/F}$.

2% body weight (B.W) of InsP$_6$ treatment for 72hr to IPMK$^{\Delta IEC}$ completely reduced

HDAC3 activity. Data was represented as a fold change compared to IgG. Results are

representative of a minimum of three individual experiments. (n=14 for IPMK$^{F/F}$, n=14

for IPMK$^{\Delta IEC}$, n=5 for 2% body weight (B.W) of InsP$_6$ treatment for 72hr to IPMK$^{\Delta IEC}$).

(L) Western blot from intestinal crypts (Ileum) indicated elevated H4K16 acetylation in

IPMK$^{\Delta IEC}$ mice which was reduced by pretreatment of 2% oral InsP$_6$. Western blot

analysis further showed HDAC3 protein levels were comparable in all untreated and

untreated genotypes. Actin was used as a loading control. Results were representative

of three individual experiments (n=3). **(M,)** Pictorial representation showing MMP3 and

13 primer locations on the mouse genome. **(N)** ChIP-qPCR analysis indicated MMP3/13

promoter enrichment of acetylated H4K16 when purified from IPMK$^{\Delta IEC}$ crypts (Ileum),

which was diminished by pretreatment oral InsP$_6$ (2%). Data represents three

experimental replicates (n=3). **(O)** Reduced HDAC3 activity in Crohns patient tissue

(Control, n=4, CD,n=5) and **(P)** Ulcerative colitis patient tissue (Control, n=3, UC,n=5)**.**

(Q) and **(R)** showing loss of IPMK in CD and UC patient samples. IPPK level

unchanged. MMP levels increased in diseased tissue. ZO-1 and Occludin reduced in

CD and UC. H4K16acetylation induced in CD and UC.

Extended Figure 5 I: **(A)** Monolayer of CaCo2 cells were treated with conditioned

media from WT and *IPMK* KO HCT 116 cells for 8 hr. After treatment CaCo2 cells were

lysed and western blotted against anti-ZO-1 and anti-Occludin antibodies, actin was

used as loding control, followed by densitometric analysis. Data represents three

experimental replicates (n=3). **(B)** Pictorial representation of FACS based intestinal

epithelial cell purification procedure. **(C)** After isolation, intestinal cells were flow

purified, followed by staining with APC-CD45 (Ex 651nm: Em 660nm), FITC-CD31(Ex

498nm: Em 517nm), PE/Cy7-TER-119 (Ex 565nm: Em 774nm), and PE-EpCAM (Ex

565nm: Em 576nm). Dead cells were excluded using the Tail Cell Cycle Solution,

propidium iodide (PI) (Ex 535nm: Em 617nm). CD45/CD31/TER-119 /EpCAM. intestinal

epithelial cells were sorted using a SONY SH800S Cell Sorter. **(D)** CD45./CD31./TER-

119./EpCam+ intestinal epithelial cells were isolated from IPMK$^{\triangle IEC}$ and IPMK $^{F/F}$ mice

through FACS purification showing loss of IPMK protein in IPMK$^{\triangle IEC}$ *mice.* Actin was

used as a loading control. **(E)** Western blot analysis showed the purity of EpCAM $^+$

intestinal epithelial cell population. Lysate was isolated from unsorted and sorted

intestinal cells, followed by western blot against anti-CD31, anti-CD45, anti-TER-119,

and anti-EpCam. Actin was used as a loading control. Analysis showed that the sorted

cell population was devoid of CD31, CD45, and TER-119 cell population but positive for

EpCam. Results were representative of three individual experiments (n=3). **(F)**

Western blot from intestinal crypts (ileum) indicated elevated H4k16 acetylation in

IPMK$^{\triangle IEC}$ mice, which was reduced by pretreatment of 2% oral InsP$_6$. Western blot

analysis further showed IPPK protein levels were comparable in all untreated and

untreated genotypes. Lamin was used as a loading control. Results were

representative of three individual experiments (n=3). (G) Western blot analysis of

colonic crypts revealed elevated levels of MMP3, MMP13, and acetylated H4K16 in

IPMK$^{\Delta IEC}$ mice, which were rescued following pretreatment with 2% oral InsP6. IPMK

expression was diminished, while IPPK levels remained unaltered. Actin served as a

loading control. Data are of three independent experiments (n = 3). **(H)** ChIP-qPCR

analysis depicted HDAC3 promoter enrichment for the MMP3/13 gene in IPMK F/F,

IPMK$^{\Delta IEC}$ and IPMK$^{\Delta IEC}$ mice pretreated with oral InsP$_6$. Data represents three

experimental replicates (n=3).

Extended Figure 5II: (A) IPMK KO cell lysate and also conditioned media show MMP

13 and 3 enrichment. Densitometric analysis of western blot from Fig 5 G for (B) MMP3;

(C) MMP13; (D) ZO-1, Occludin (E), and (F) Claudin 2. Pathway analysis using the

Metascape tool for RNA-seq data shows (G) upregulated genes in $IPMK^{\Delta IEC}$ compared

to $IPMK^{F/F}$ and (H) downregulated pathways. (I) Relevant enriched gene set from GSEA

of RNA-seq datasets for *$IPMK^{F/F}$ vs $IPMK^{\Delta IEC}$* of the mouse model, using the MSigDB

C2:CP:Reactome subcollection.

igure 6: In the DSS-induced IBD mouse model, diminished IPMK expression and

HDAC3 activity led to increased intestinal permeability, which was rescued by

oral InsP$_6$ treatment. (A) Intestinal permeability assay indicated a significant increase

in 4 kDa FITC Dextran fluorescence in 24 hr 3% DSS treated mice serum without

altering 70 kDa Dextran (Rhodamine B) as compared with untreated mice. Data were

expressed as fold change relative to control mice serum, which received only PBS

gavage without FITC or Rhodamine Dextran. (n=15). **(B)** Western blot from colonic crypt

showed that 3% DSS treatment reduced IPMK expression while IPPK expression was

unaltered. Actin was used as loading control. (n=3) **(C)** 3% DSS treatment (24 hrs)

markedly reduced HDAC3 activity in colonic crypts which oral InsP$_6$ treatment rescued.

Result was representative of three independent experiments (n=6). **(D)** Western blot

from colonic crypts indicated increased MMP3 and 13 expression after 3% of DSS

treatment (24hrs) which pretreatment of oral InsP$_6$ reduced. Actin was used as loading

control. Data was representative of three individual experiments. (n=6) **(E)**

Immunofluorescence microscopy shows increased MMP3 (Green) in 24 hr 3% DSS

treated colon and **(F)** ileum as compared to untreated control. That was reduced by

$InsP_6$ pretreatment (72hr). Scale bar represents 50 μm. Data represents three

experimental replicates (n=3). (Total number of images=30) **(G)** Western blot from

colonic crypt showed that 3% DSS treatment induced H4k16 acetylation which $InsP_6$

pretreatment (72hrs) reduced. Actin was used as loading control. Result was

representative of three independent experiments (n=6). **(H)** ChIP-qPCR analysis

indicated MMP3/13 promoter enrichment of acetylated H4K16 when purified from DSS

(3%) treated colonic crypts, which was diminished by pretreatment oral $InsP_6$ (2%). Data

represents three experimental replicates (n=3). **(I)** qPCR analysis of MMP3 and 13

isolated from colonic crypt. 24 hrs of 3%DSS treated induced gene expression while 2%

of $InsP_6$ treatment reduced it. 18s used as loading control. Results represented three

experimental replicates (n=3).

Extended Figure 6: (A) Western blot analysis of colonic crypts diminished IPMK

expression after 24 hours of 3% DSS treatment, which was not restored by oral InsP6

pretreatment. While IPPK expression remained unchanged in both treated and

untreated groups. Lamin served as the loading control. Data are representative of three

independent experiments (n = 4). **(B)** ChIP-qPCR analysis depicted HDAC3 promoter

enrichment for MMP3/13 gene in IPMK $^{F/F}$, IPMK$^{\triangle IEC}$ and IPMK$^{\triangle IEC}$ mice pretreated with

oral InsP$_6$. Data represents three experimental replicates (n=3).

1714

1715 **Figure 7: InsP6's absorption in intestinal tissue prevents DSS-induced**

1716 **inflammation in both wild-type and IPMK$^{\Delta IEC}$ mice.**

1717

(A) Schematics of the treatment plan in wild-type mice. **(B)** In wild-type C57BL/6 mice,

oral DSS treatment reduces colon size, which InsP6 prevents; n=5. **(C)** ELISA from

tissue lysate for IL1β and **(D)** MPO; n=3 for each group. **(E)** Evaluation of intestinal

permeability using 4 kDa and 7 kDa dextran. DSS induces intestinal permeability, which

oral InsP6 prevents; n=5 in DSS and DSS+InsP6 groups, n=3 untreated. **(F)**

Schematics of the treatment plan in IPMK$^{F/F}$ and IPMK$^{\Delta IEC}$ mice. **(G)** ELISA from tissue

lysate for IL1β and **(H)** MPO; n=3 for each group, normalized by total protein. **(I)**

Evaluation of intestinal permeability using 4 kDa and 7 kDa dextran. DSS induces

permeability, which oral InsP6 prevents; n=3 in DSS and DSS+InsP6 groups. **(J)**

Schematics of HOIP measurement from intestinal crypts using mass spectrometry. **(K)**

Data from mass spectrometry indicates an increase in InsP6 and other HOIPs after oral

InsP6 treatment in IPMKΔIEC mice; n=3. **(L)** HDAC3 activity assay in *IPPK* KO cells.

(M) Western blot of MMPs from *IPPK KO* HEK293T cells.

1731

1732

1733

1734

1735

1736

1737

1738

Extended Figure 7. (A) Average body weight of wild-type mice. (B) H&E staining of the colon from untreated, DSS, and DSS+InsP6-treated mice. n=3.Scale bar represents 60 μm (C) Body weight plot of IPMK$^{F/F}$ and IPMK$^{\Delta IEC}$ mice. (D) Colon length of IPMKF/F and *IPMK$^{\Delta IEC}$* mice after DSS treatment. n=3. (E) H&E staining of the colon from IPMK$^{F/F}$ and IPMK$^{\Delta IEC}$ mice after DSS and DSS+InsP6 treatment. n=3. Scale bar represents 60 μm (F) MiniMUGA mouse backcross genotyping analysis by Transnetyx was performed using two *IPMK$^{F/F}$* mice, confirming that the mice were backcrossed for 10 generations to the C57BL/6 strain, achieving a congenic background of 97% or higher.(n=2)

FIGURE 2 MAIN

Figure 3 Main

Figure 4 Main

Figure 5 Main

Figure 6 Main

Figure 7 Main