Evaluating Best Practices for Isolating Pyrophilous Bacteria and Fungi from Burned Soil

Dylan J. Enright[1*], Aishwarya Veerabahu[1], Ryan J. Quaal[1], Marely Vega[1], Anna Nguyen[1], Jenna Grindeland[1], James W.J. Randolph[1], Maria E. Ordonez[1], and Sydney I. Glassman[1*]

Affiliations:

[1]University of California Riverside, Department of Microbiology and Plant Pathology, 900 University Ave. Riverside CA, 92521, USA

Corresponding authors: sydneyg@ucr.edu and denri009@ucr.edu

ORCIDs:

Dylan J. Enright: 0000-0002-5153-0728

Aishwarya Veerabahu: 0009-0000-5411-7093

Ryan J. Quaal: 0009-0002-1549-6670

Marely Vega: 0009-0003-3493-7790

Anna Nguyen: 0009-0005-2364-898X

Jenna Grindeland: 0000-0002-5693-455

James W.J. Randolph: 0009-0007-7653-2037

Maria E. Ordonez: 0009-0000-2055-4324

Sydney I. Glassman: 0000-0001-9115-3026

Key Words: fire, pyrophilous, culture, bacteria, fungi, soil.

Abstract

A live microbial culture is invaluable to assess traits and functions, yet culturing

immediately from fresh soil is logistically challenging and media selection is not trivial. Deeper

ecological understanding of pyrophilous microbes and their traits is hampered by a lack of

culture-based work causing researchers to rely heavily on community sequence analysis. To

improve our understanding of isolating pyrophilous bacteria and fungi after wildfires from

burned soil, we tested which: 1) soil storage method and 2) media retained highest genera

richness and highest culturable viability as measured by CFUs retained. We tested four soil

storage methods (dried, refrigerated at 4ºC, stored at -80ºC with or without glycerol) using a

well-homogenized soil sample, plated microbes on rich media, and compared to fresh soil

obtained 6 months after a severe California shrubland wildfire. From the fresh soils, we tested 3

media types: rich (Lysogeny Broth (LB) for bacteria; Malt Yeast Agar (MYA) for fungi),

oligotrophic (Reasoner's 2 Agar (R2A)) and media made from pyrogenic organic matter

(PyOM). For bacteria, storing soil frozen at -80ºC without glycerol yielded the highest genera

richness and with glycerol preserved the most species. For fungi, storing soil at -80ºC without

glycerol preserved the most species but preserved equivalent richness of genera as fresh and 4°C

soil. R2A yielded the highest bacterial and fungal genus richness but some species of interest

were only captured with PyOM. Using a combination of these storage and media methods along

with a few additional techniques (mushroom cultivation, smoke cultivation, etc.) from 2018-

2022, we cultured >500 isolates (286 bacteria; 258 fungi) after 7 California wildfires for testing

pyrophilous microbial traits.

Importance

Fires are increasing in frequency and severity across the globe, making the understanding of the traits of the pyrophilous microbes that survive and thrive post-fire and influence post-fire ecosystem regeneration critical and pressing. Yet, our ability to characterize the traits of pyrophilous microbes through genomics, transcriptomics, and phenotypic assays is hampered by lack of organisms in culture. Here, we debut a new medium created from pyrogenic organic matter that helps to cultivate fastidious pyrophilous microbes, and also describe a large-scale culture collection of pyrophilous bacteria and fungi (>500 isolates) that can be used for future collaborative research. We comprehensively test how soil storage and media type affect the cultivation of both bacteria and fungi, and for the first time, we test the best method to store soil to retain fungal diversity. Our work can be applied to improve culture collections of fastidious microbes from a variety of environments.

Introduction

Although advances in the fields of molecular biology and DNA sequencing have freed microbiologists from the need to work with culturable species exclusively (1), few laboratory resources are as valuable as live cultures. Pure cultures are essential components for high quality whole-genome assembly (2, 3), genetic engineering experiments (4), and the phenotypic profiling of microorganisms (5–10). Additionally, microbial isolates greatly improve the interpretation of metabolomics data (11), and possession of a diverse culture collection can unlock a plethora of applied microbiological research directions (4, 12, 13). Once isolated and properly stocked, culture collections become a permanent resource for future microbiological

research and generate new avenues for collaborative research (14–16). However, there are many

hurdles to creating microbial culture collections and each collection has unique challenges (14).

The various factors that can be modified when attempting to culture microorganisms are

as varied as there are microbes to isolate. For example, media selection may require grappling

with an endless number of variables including consideration of the proper carbon sources,

nitrogen and phosphorus content, salt concentrations, pH, and a host of other micronutrient

considerations (13, 17, 18). One must also consider incubation temperature and duration, water

and oxygen availability, and whether to alternate light and dark cycles or to make the media

nutrient rich or poor (oligotrophic) (17–19). This complexity contributes to many

microorganisms remaining uncultured (20). Though recent advances in our understanding of

microbiology have begun to make the adage of "99% of microbes are unculturable" outdated

(13, 18, 19, 21), culturing the majority of microorganisms still remains beyond our reach.

One group of understudied microorganisms that is of growing interest are the pyrophilous

(or "fire-loving") microbes. Pyrophilous microbes were rare or absent pre-fire but massively

increase in abundance post-fire, sometimes persisting for many years post-fire (22–24), and

influencing post-fire ecosystem recovery. While there are many studies identifying which

microbes become more abundant post-fire, there are very few studies characterizing how

pyrophilous microbes interact with their environment (25). Wildfires dramatically change

landscapes and create legacy effects that can impact soil microorganisms and their overall

ecosystems for many years (26–28). With rising global frequency and severity of wildfires across

many ecosystems (29, 30), understanding how pyrophilous microbes modify and interact with

their environments is critical. An important step to characterizing the ecology of pyrophilous

microbes is laboratory testing of cultured isolates to provide data on how individual pyrophilous

taxa compete for resources, cycle post-fire nutrients, and interact with the post-fire environment.

However, there are numerous technological and logistical challenges to creating a culture

collection of pyrophilous microbes.

Beyond the well-known hurdles of creating a microbial culture collection, attempting to

culture microbes associated with wildfires presents a unique set of logistical challenges including

proper soil storage. Wildfires are unpredictable natural phenomena. As such, it is often not

possible to have the infrastructure for large-scale culturing efforts (personnel available, media

selected and created, broth tubes or petri dishes ordered, culturing conditions decided and

established) in place when a wildfire occurs, so post-fire soil samples may have to be stored for

varying lengths of time prior to culturing. This is a logistical problem shared among many

environmental microbial culturing efforts ranging from marine systems (31) to Antarctic soils

(32). While many insights have been gained regarding these logistical hurdles such as culturing

from samples as quickly as possible (14, 33), and methods for freezing samples that improve

cultural viability, their efficacy is not always consistent and the viability of culturable

microorganisms may be reduced depending on the soil storage method (33, 34).

Wildfires also radically alter environmental nutrient and resource landscapes (27, 35),

making culture media selection challenging. Wildfire pyrolysis turns labile, easily accessible

carbon sources into difficult to degrade pyrogenic organic matter (PyOM) (28, 36). PyOM is

often enriched in polycyclic aromatic hydrocarbons (PAHs) that may require specialized

metabolic pathways to process (12, 37–40). This can lead to accumulated char persisting in the

environment for up to and exceeding 100 years (41). Additionally, wildfires deposit inorganic

nitrogen that was previously locked in plant biomass through combustion and ash deposition (26,

35, 42). This high-nitrogen environment may alter the nutrient strategies that microorganisms

employ post-fire and limit the microbes capable of surviving. For example, many species of

mycorrhizal fungi are known to be nitrophobic (43), preventing them from establishing essential

plant associations in high nitrogen environments. These nutrient shifts may result in commonly

used laboratory media not providing appropriate resources to cultivate pyrophilous microbes.

Here, our goal was to create a large culture collection of pyrophilous microbes to use to

test the traits that govern pyrophilous microbial adaptation to fire and the post-fire environment

via biophysical assays and genomics. Keeping in mind the above described logistic and resource

and nutrient-related challenges, we used soil from a local high severity California chaparral

wildfire to test the best 1) soil storage methods and 2) laboratory media to retain the greatest

richness and culturing viability of pyrophilous microbes. For soil storage, we tested culturing

immediately from fresh soils versus soils stored for 3 months in four ways (air-dried, refrigerated

at 4°C, and frozen at -80°C with and without glycerol). To reduce confounding variables, we

plated all soil storage experiments on rich media. For media type, we tested culturing

immediately from fresh soil onto rich (lysogeny broth (LB) agar for bacteria, malt-yeast agar

(MYA) for fungi), oligotrophic (Reasoner's 2 Agar (R2A)), or a media created in-house from

PyOM. We tested the impact of our treatments on abundance (CFUs: Colony Forming Units) and

microbial richness and composition (assessed via Sanger sequencing of 16S for bacteria or ITS

for fungi; Figure 1). We hypothesized (H1) that other than fresh soil, air-dried soil would retain

the greatest abundance and genera richness of pyrophilous bacteria and fungi since fires reduce

soil moisture (35, 44), so we expected air drying soils would be the least dramatic disturbance for

soil borne microorganisms. We further hypothesized (H2) that a combination of the rich and

PyOM media would yield the greatest richness of pyrophilous microbes as the rich media could

grow a wide range of bacteria and fungi and the PyOM media would capture those pyrophilous

species with unique metabolic requirements specific to post-fire burned environments. Although

we specifically sought to create a pyrophilous culture collection, the insights gleaned here may

also aid environmental microbiologists in decision making about storage and media selection for

many types of environmental culturing for fastidious microbes. Finally, the insights this study

provided helped to guide our related work of creating a large and diverse collection of

pyrophilous microorganisms of >540 isolates (286 bacteria; 258 fungi) from burned soils,

mushrooms, and smoke derived from 7 California wildfires.

Results

Microbes Obtained: Throughout our soil storage and media selection experiments we

isolated a visually diverse array of bacterial and fungal morphotypes (Figure 2). We cultured a

total of 170 bacterial isolates representing 4 phyla, 10 orders, 31 genera and 83 taxa from burned

soils collected 6 months after the 2020 El Dorado Fire. We also cultured 160 fungal isolates

representing 3 phyla, 9 orders, 15 genera and 35 taxa (Table S1). Of these isolates, 138/170

bacterial isolates composed of 17 genera and 71 taxa and 152/160 fungal isolates composed of 8

genera and 30 taxa had literature support for inclusion as putative or confirmed pyrophilous taxa

(22, 25, 27, 37, 38, 45–51) (Table S2). Putative pyrophilous taxa are those (or a close relative

taxa) which appear repeatedly post-fire but have not been designated as pyrophilous in one or

more publications yet. The putative pyrophilous taxa list includes more bacteria than fungi

because the research on pyrophilous bacteria is more nascent than on pyrophilous fungi (24, 25,

45, 49), which dates back to 1909. We used the total (Table S1) and pyrophilous isolates (Table

S2) to test the effect of media type and soil storage method on CFUs and microbial richness. In

addition to testing storage methods and media selection, here we present our culture collection of

544 pyrophilous microbes (286 bacteria; 258 fungi) that can be used for testing of pyrophilous

microbial traits, for example genomics, transcriptomics and phenotypic assays (52, 53), that is

the result of our related work (from 2018-2022) in which we isolated bacteria and fungi from

burned soils, mushrooms, and smoke of 7 California wildfires (Table S3). For bacteria, we had

286 isolates from 8 phyla, 16 orders, 35 genera, 76 taxa. For fungi, we had 258 isolates from 3

phyla, 11 orders, 13 genera, and 34 taxa. The full table includes the fires of origin, collection

times, substrate cultures derived from, isolate names, FASTA sequences, and BLAST summary

statistics (Table S3).

Effect of Storage Method Selection on CFUs: Soil storage method significantly impacted

the number of CFUs obtained per gram soil for both bacteria (ANOVA: $F_{1,4} = 19.97$, $p < 0.001$;

Figure 3A) and fungi (ANOVA: $F_{1,4} = 19.78$, $p < 0.001$; Figure 3B). For both bacteria and fungi,

soil stored air-dried (Bacteria: 2.31×10^5 CFUs/gram, Fungi: 1.33×10^4 CFUs/gram) and stored at -

80ºC with glycerol (Bacteria: 1.31×10^5 CFUs/gram, Fungi: 6.69×10^3 CFUs/gram) retained the

fewest average CFUs/gram soil. For bacteria, soils stored at 4ºC (6.98×10^5 CFUs/gram) or at -

80ºC without glycerol (6.39×10^5 CFUs/gram) were not significantly different from fresh soils

(5.73×10^5 CFUs). For fungi, fresh soils had the highest average CFUs/gram soil (3.07×10^4

CFUs/gram), followed by soil frozen at -80ºC without glycerol (2.66×10^4 CFUs/gram), and then

stored at 4ºC (2.16×10^4 CFUs/gram).

Effect of Media Selection on CFUs: Media type significantly impacted the number of

CFUs obtained per gram of fresh soil for both bacteria (ANOVA: $F_{1,2} = 39.43$, $p < 0.001$; Figure

3C) and fungi ($F_{1,2} = 7.77$, $p < 0.01$; Figure 3D). For both bacteria and fungi, PyOM media

yielded the fewest average CFUs per gram (Bacteria: 1.66×10^5; Fungi: 1.03×10^4), and rich media

(LB for bacteria and MYA for fungi) had the highest average CFUs per gram (Bacteria:

5.73×10^5; Fungi: 3.07×10^4). R2A media had the second highest per gram average CFU counts for

bacteria (3.20×10^5) and for fungi, CFU counts (2.20×10^4) were statistically equivalent to MYA.

Effect of Storage Method Selection on Bacterial Diversity: Soil storage method strongly

affected richness of bacterial genera with fresh soil retaining the highest total (Figure 4A) and

frozen soil retaining the highest pyrophilous richness (Figure 4B). Dried soil retained the lowest

richness of bacterial genera among both total (Figure 4A) and pyrophilous isolates (Figure 4B).

Across all soil storage treatments, we cultured a total of 21 bacterial genera (Figure S2) and 64

taxa (Figure S3), and we found literature support to classify 14 genera (Figure 6A) and 51 taxa

(Figure S4) as pyrophilous or putatively pyrophilous. Considering total bacterial taxa, fresh and

frozen soil were equally rich (62% of genera), followed by soil frozen with glycerol (57% of

genera), 4°C soil (48% of genera) with dried soil as least rich (38% of genera). Only 3 genera

(_Arthrobacter, Curtobacterium, Streptomyces_) were isolated across all 5 storage treatments

whereas 8 genera (_Brevibacterium, Fontibacillus, Glutamicibacter, Massilia,_

Ornithinimicrobium, Paeniglutamicibacter, an unidentified Firmicute, and _Pseudoarthobacter_)

were isolated from a single treatment (Figure S2). Finally, 64% of taxa were isolated one time

with only one taxon best identified as _Curtobacterium oceanosedimentum_ observed in all storage

treatments (Figure S3). Among the pyrophilous subset (Figure 4B, S4), frozen soil was the most

rich (85% of genera), followed by fresh soil (62% of genera), soil frozen with glycerol (54% of

genera), 4°C soil (54% of genera) with dried soil as least rich (38% of genera).

Effect of Storage Method Selection on Fungal Diversity: Soil storage method strongly

affected fungal richness with fresh soil retaining the highest total genera (Figure 4C) and fresh,

frozen and refrigerated retaining equal numbers of pyrophilous genera (Figure 4D). Soil frozen

with glycerol retained the lowest richness of fungal genera among total (Figure 4C) and

pyrophilous isolates (Figure 4D). Across all storage treatments, we cultured a total of 6 fungal

genera (Figure S5) and 23 fungal taxa (Figure S6), of which we found literature support as

pyrophilous for 3 genera (Figure 6B) and 20 taxa (Figure S7). For total fungi, fresh soil was the

most rich, capturing 100% of all 6 genera (*Aspergillus, Holtermaniella, Ochrocladiosporium*,

Penicillium, and 2 uncultured Helotiales), followed by frozen and refrigerated soil capturing 50%

of the genera, and 30% in dried soil (Figure S5). Soil frozen with glycerol performed the worst,

capturing only *Penicillium* (17% of the total; Figure S5). For pyrophilous fungi, fresh, frozen,

and refrigerated soil all captured 100% of the 3 genera (*Aspergillus, Holtermanniella,*

Penicillium), whereas dried captured 2 (*Aspergillus, Penicillium*) and frozen with glycerol only

captured 1 (*Pencillium*; Figure 6B).

Effect of Media Selection on Bacterial Diversity*: Media type strongly affected bacterial

richness with R2A and LB both capturing the highest numbers of total bacterial genera (Figure

5A), R2A yielding the highest richness of pyrophilous genera (Figure 5B), and PyOM capturing

the fewest total and pyrophilous bacterial genera. Across all media types from fresh soil, we

cultured 24 total bacterial genera (Figure S8), 61 total bacterial taxa (Figure S9), and 16

pyrophilous bacterial genera (Figure 7A) and 49 pyrophilous bacterial taxa (Figure S10). Among

total bacteria, LB and R2A obtained equally rich communities (58% of genera), with the least in

PyOM (33% of genera). Among pyrophilous bacteria, R2A captured the greatest richness (69%

of genera), followed by LB (56% of genera) with PyOM yielding the lowest richness (44% of

genera). Although PyOM media yielded the lowest diversity, it was the only media to isolate

some important pyrophilous bacterial genera like *Noviherbaspirillum* and *Peribacillus* (46, 47,

49, 52) and interestingly, R2A was the only media to capture *Massilia* (Figure 7A) suggesting a

more oligotrophic lifestyle than previously thought (45, 49).

236   <u>Effect of Media Selection on Fungal Diversity:</u> Media type affected fungal richness with

237 R2A capturing the most total (Figure 5C) and pyrophilous (Figure 5D) fungal genera whereas

238 MYA captured the fewest. Across all media types from fresh soil, for total fungi we cultured 14

239 genera (Figure S11), and 29 taxa (Figure S12), and 8 genera (Figure 7B) and 23 taxa for

240 pyrophilous fungi (Figure S13). Among total fungi, R2A media captured the highest richness

241 (57% of genera) while both MYA and PyOM were equally diverse (43% of genera). Among

242 pyrophilous fungi, R2A also captured the highest richness (89% of genera), followed by PyOM

243 (50% of genera), and MYA was the least rich (38% of genera). While PyOM did not capture the

244 most diversity, it was the only media to capture *Neurospora* (Figure 7B), a well-documented

245 post-fire fungus with relatively little knowledge about its ecology (54).

Discussion

248   Here, we present results of an experiment to test soil storage and media selection to

249 obtain the best diversity and abundance of bacterial and fungal isolates, which have informed our

250 continual culturing efforts resulting in a collection of 544 pyrophilous microbes (286 bacteria;

251 258 fungi) (Table S3). To our knowledge, this is the first study examining the impact of soil

252 storage on fungal diversity, and this is also the first large-scale culturing effort of pyrophilous

253 bacteria, and among the largest soil and mushroom based culturing efforts of pyrophilous fungi

254 since 1910 (22). For both bacteria and fungi, freezing soil without glycerol yielded the highest

255 number of pyrophilous genera on rich media, and from fresh soil, R2A yielded the most genera.

256 Although it did not capture the highest number of CFUs or highest number of taxa, novel solid

257 agar media we created from field-harvested PyOM was necessary to capture some key

258 pyrophilous genera of interest like *Neurospora* for fungi and *Noviherbaspirillium* for bacteria.

For bacteria, we conclude that freezing soil at -80°C is the best storage method relative to

fresh soil, yielding the highest number of CFUs and total and pyrophilous bacterial genera when

plated onto rich media. Interestingly, adding glycerol increased the number of taxa but reduced

the number of total and pyrophilous genera isolated. While CFU based analyses are useful in

giving an approximation of the total number of culturable cells present in the original sample and

a broad overview of the effectiveness of certain culturing practices at cultivating cells, it does not

capture the important distinction of isolate diversity. However, in this case, freezing soils without

glycerol captured both the highest number of CFUs and yielded the vast majority of the

pyrophilous bacterial genera of interest based on the post-fire bacterial succession literature (45–

47, 49, 55, 56). While culturing from fresh soil did capture the greatest number of total genera,

freezing soil captured the greatest richness of pyrophilous genera (Figure 4). Interestingly,

freezing soil resulted in more total and pyrophilous bacterial genera being isolated, but adding

glycerol increased the total number of taxa and was necessary to isolate *Massilia,* which is a

bacterial genus repeatedly observed post-fire (46, 47, 49, 57), with very little understanding

about its ecology, making it a prime candidate for further study. It is possible that taxa only

isolated when glycerol was added could be either taxa that are more sensitive to the thermal shift

that occurs with freezing or they could be taxa which do not sequester themselves into

aggregates and other microenvironments and therefore need the glycerol to buffer them against

lysing under freezing conditions. Therefore, it may be of value for researchers to freeze some soil

with glycerol as well as without, to capture some of these more potentially vulnerable taxa.

Finally, dried soils performed the worst in capturing total and pyrophilous bacterial diversity,

which surprised us given how dry post-fire soils usually are. While there is little similar existing

literature to compare to, our findings differ from one previous study that found that refrigeration

at 4ºC was better than freezing at -20ºC at retaining microbial biomass and respiration (33). In

contrast to our burned chapparal soils, this study examined unburned grassland and forest soils.

Different soil structure (prevalence of soil aggregates, water holding capacity, porosity, etc.) can

influence microbial activity (58) and therefore could influence how soil microenvironments

behave under storage. Although we know of no other cultivation studies to compare to, a study

examining how storage time (0-14 days) and temperature (room temp, 4°C, and -20°C) affected

16S sequencing results similarly found that freezing preserved the most bacterial diversity,

though they did not test freezing to -80°C as we did (34). Both of these studies only froze at -

20ºC rather than -80ºC, which may have resulted in fewer taxa retained (59).

Soil storage method seemed to matter less for pyrophilous fungi than bacterial genera.

While there was one method that stood out as the worst (freezing at -80°C with glycerol), the

remaining methods captured most of the richness observed (Figures 4D and 6B). This could be

due to the resilience of fungal spores to a variety of disturbance effects (60), but it is also

possible that MYA was not the ideal culture medium for pyrophilous fungi, given the success of

R2A and PyOM media at cultivating pyrophilous fungi (Figures 5D and 7B). Future work

repeating the storage experiments with R2A and PyOM might improve pyrophilous fungal

isolate retention. Despite this caveat, we were able to draw some very surprising conclusions

from our experiments. For example, our hypothesis that air-drying soil would be the most

effective storage method was fully unsupported by our results. Indeed, air-drying yielded the

lowest CFU counts and total and pyrophilous diversity. This is an important finding since drying

soils prior to laboratory use is not uncommon (34, 61, 62). Notably, we were unable to find other

studies exploring the impact of soil storage on fungal communities, indicating a strong need for

these methods to be tested across kingdoms.

Surprisingly, R2A was the most effective culturing media for capturing pyrophilous

bacteria and fungi. R2A produced the second highest number of CFUs after the rich medias for

bacteria (LB) and fungi (MYA) (Figure 3) and the most diverse pyrophilous bacterial (Figure 5B)

and fungal (Figure 5D) isolates. Since R2A was developed to culture bacteria from oligotrophic

potable drinking water (63), this suggests that many pyrophilous microbes prefer a more

oligotrophic environment. While shifting life-strategies of microbial communities under stress

can be extremely varied and nuanced (64), oligotrophic bacteria and fungi may be favored under

an increasing stress gradient (such as carbon stress, soil moisture stress, or heat stress) (65),

which might be ideal for capturing a wide diversity of microbes thriving in the stressful post-fire

environment. However, this is also a surprising result as post-fire environments tend to be high in

nitrogen and phosphorus, even if the carbon landscape has radically changed (26, 28). It is

therefore possible that these pyrophilous taxa are less affected by the high nitrogen and

phosphorus and instead conform more to the previously observed pattern of oligotrophy under a

stressful environment. As such, it is possible that availability of certain carbon forms plays a

more integral role in post-fire nutrient acquisition and should be studied further. While previous

studies exist examining the role of media conditions on microbial culturability (18, 32, 66, 67),

these studies often take either a fine-scale approach to tuning a specific culture medium or test

only CFU-based metrics rather than diversity or use only rich media. Continued exploration of

media conditions is essential to increasing the number of cultured microbial species, and insights

gained from our study may help open new paths in culture media development.

Finally, although our home-made PyOM media captured the lowest CFUs of bacteria and

fungi, and the lowest bacterial genus richness, it captured the second highest fungal richness and

was necessary to isolate key bacterial (i.e. *Noviherbaspirillum soli*) and fungal (i.e. *Neurospora*)

pyrophilous taxa of interest. In this case, isolating finicky pyrophilous taxa of interest was

improved by the low-density growth we observed in PyOM, which likely resulted from the lower

nutrient content similar to R2A media. Rich media like LB, MYA, and potato dextrose agar are

often plagued by fast growing ruderal species that can outpace taxa that grow at lower density

(68). These taxa are particularly interesting because *Noviherbaspirillum* has been repeatedly

observed increasing in abundance post-fire (45–47) and recent genomic work has shown

Noviherbaspirillum to have a diverse array of PyOM degradation pathways (52). Likewise,

Neurospora is famously found post-fire and is widely used as a genetics model organism despite

very little understanding about its ecological role in the environment (54) though recent genomic

work has also characterized potential PyOM degradative abilities (53). Therefore, we conclude

that R2A is best media to use to maximize pyrophilous bacterial and fungal diversity, but it

would be ideal to also include a PyOM media if specifically interested in isolating the most

finicky and interesting pyrophilous microbes.

While we made significant progress regarding testing effective culturing practices for

environmental microbes and specifically pyrophilous bacteria and fungi, much additional work

remains and caveats exist. Except for the home-made PyOM media, we limited ourselves to

commonly available laboratory culturing media. We acknowledge there is a near endless range of

possible medias to use (66), possible range of growth temperatures (50, 66), additional isolation

strategies (69), and media pH (17, 67), which might be more effective for the culturing of

pyrophilous microorganisms. However, we assert that the broad trends (rich vs. oligotrophic vs.

PyOM media) yield valuable insights into the culturing of these microorganisms. Additionally, as

this work is intended as a foundation for understanding the culturing of pyrophilous microbes,

we endeavored to stick to laboratory materials and methodologies that are commonly available to

a wide range of research institutions, and accessible for undergraduates, which included most of

our co-authors. The PyOM media uses a water-extraction based method of incorporating

nutrients, which might have precluded key nutrients required for post-fire metabolism that are

less water-soluble in PyOM than others (70), which might have missed some pyrophilous

microbial taxa. While we have some limited data on our PyOM media composition (GC mass

spectrometer analysis showed 10x more nitrogen present in PyOM compared to LB), additional

metabolic analysis is needed to fully characterize this environmentally-derived media.

Additionally, the culturing effort in this study all took place aerobically at room temperature in a

temperature controlled lab. We also acknowledge that our CFU calculations are not normalized

by dry soil weight, since we instead prioritized the preservation of potentially sensitive taxa over

drying soil prior to calculation, which may limit direct comparisons from our CFU data to that of

other studies. Finally, while we acknowledge that separating microbial cultures based on colony

morphology is difficult at best (71–73), we decided that choosing colonies based on morphotype

and then identifying with Sanger sequencing was an acceptable concession to the technical and

logistic demands of isolating, stocking, and identifying every single colony we grew. This does

mean that we could have missed some species-level diversity since some species can share

extremely similar colony morphologies (72, 73) and techniques like dilution to extinction

culturing might capture more isolates identified to species level (74–76). However, with the

difficulties of identifying taxa to species level with rRNA fragments alone coupled with the

potential morphological limitation just discussed, we believe our genus-level observations to be

more robust hence our exclusion of species-level analyses in this study. Despite these caveats, we

still obtained a wide diversity of total (Table S1) and pyrophilous (Table S2) bacteria and fungi

and found meaningful differences in media and storage types.

Conclusions

Here, we present the first study to systematically test how soil storage methods impact

fungal cultivability, and the largest scale test of how soil storage and media type affect bacterial

and fungal cultivation. We found that storing soils at -80°C without glycerol and utilizing a

combination of rich media and oligotrophic R2A media preserved the greatest diversity of

culturable pyrophilous bacteria and fungi compared to fresh soils from burned chaparral.

However, certain pyrophilous taxa of interest were only captured from soil frozen with glycerol,

like *Massilia*, and on our novel PyOM media, like *Neurospora* and *Noviherbaspirillium*,

indicating a potential value in using two media and storage types. This experiment contributed to

a larger culturing effort resulting in a collection of >540 pyrophilous microbes (286 bacteria; 258

fungi) isolated from soil, smoke, and mushrooms isolated from 7 California wildfires (Table S3),

which can be used for future multi-omics, phenotypic, and applied microbiological research.

Indeed, we have already begun investigating the traits and trade-offs of these cultures through

both phenotypic and genomic profiling (52, 53, 77). Sequencing genomes from isolates with

concurrent biophysical assays is a powerful yet under-used tool to gain insight into microbial

traits and connect genomic information to trait expression (78). Obtaining the cultures is the first

step to characterizing their traits via 'omics and phenotypic methods.

Materials & Methods

<u>Site Description:</u> To test the best soil storage method and media to cultivate pyrophilous

microbes, we sampled from The El Dorado wildfire, which burned ~92km^2 Manzanita

(*Arctostaphylos*) dominated chaparral shrublands, in Yucaipa, CA from September through

November of 2020. We examined high-resolution successional data of bacterial and fungal

communities after a similar nearby high-severity Manzanita dominated chaparral wildfire (46),

and compared to other post-fire studies that had successional data (47, 55), to identify 6 months

post-fire as the ideal time to attempt to isolate a wide range of pyrophilous bacteria and fungi.

We specifically identified the timepoint as being a point during microbial succession where

several hypothesized pyrophilous groups (fast-growers, post-fire nutrient acquisitive microbes,

and thermotolerant microbes) (45, 46, 49) were in proportional flux but all present in varying

amounts (46). We thus collected samples from high severity burned soils in June 2021, 6 months

post-fire. Burned sites were characterized as having 50 to 75 percent slopes with dry, sandy soils

in the Springdale family-Lithic Xerorthents association

(https://websoilsurvey.nrcs.usda.gov/app/). At the time of collection soil moisture content was

9.72% (±2.7%) per gram soil as assessed by mass loss after drying.

Soil Collection and Storage: Approximately 250mL of soil from 0-10 cm below the ash

layer of burned soil was collected from 9 plots with ethanol cleaned releasable bulb planters and

placed into whirl-pak bags 6 months post-fire following methods used in a similar ecosystem

(46). After collection, soil was 2mm sieved and either used as fresh inoculum for cultivation or

stored in four different ways: air dried, refrigerated (4ºC), frozen (-80ºC), or suspended 50:50

%w/v soil with glycerol to create a soil slurry and then frozen (-80ºC) (Figure 1).

Culturing Medias Used: Fresh soil samples were cultured immediately onto rich (LB for

bacteria, MYA for fungi), oligotrophic (R2A), or PyOM media. After 3 months of storage, stored

soil samples were cultured onto rich media. This means that the cultures isolated from fresh soils

on rich media served as both the "rich media" condition for the media experiment and the "fresh"

comparison for the soil storage experiment (Figure 1).

We produced LB broth by adding 15g of powdered LB Broth (Fisher Scientific, CAS#

73049-73-7) and MYA broth by adding 5g each of powdered Malt Extract (Research Products

International, Genessee, CAS# 8002-48-0) and Yeast Extract (Sigma-Aldrich, Cat # 70161) per

liter of DI water. For solid agar media, 15g per liter of bacteriological grade agar (Apex

Chemicals, Cat # 20-273) was added to either the LB or MYA broth. R2A agar was produced

using powdered pre-mixed R2A agar (Criterion, Cat # 16721) following manufacturer directions.

We developed an in-house recipe of PyOM media using biochar harvested from the El

Dorado Fire burn sites (Figure S1). Char pieces that were uniformly black in color, brittle, and

appeared fully pyrolyzed were selected from the same plots where soil samples were taken. First,

we ground biochar into a fine powder with a coffee grinder (Chefman, Model: RJ44-OPP-RED)

for 3-5 minutes at maximum speed. Then, we added 50g of ground biochar to a 2L Erlenmeyer

flask, added 1200mL of DI water, covered the mouth of the flask with a watchglass, stirred on a

hotplate for 24 hours at 40°C, then removed from heat to settle for 8 hours. We then siphoned

liquid contents into a clean Erlenmeyer flask, being careful to not disturb the bottom sediment.

We passed the liquid through cheesecloth, and then twice through a coffee filter to remove any

remaining sediment before transferring to clean containers for autoclaving. We added 13g of

bacteriological grade agar (Apex Chemicals, Cat # 20-273) to each liter of PyOM media, since

adding 15g resulted in under-dissolved agar in the final product. This resulted in a media that

should reflect the more complicated carbon sources found in char (PAHs, benzoic compounds,

etc.) and contained almost 10x more nitrate than LB as assessed by GC mass spectroscopy (LB

avg NO_3^- conc. = 0.27mg/L, PyOM avg NO_3^- conc. = 1.2mg/L). We used environmentally

sourced char from the El Dorado burn site, which we believed was the best representative

substrate for pyrophilous microbes, but since there can be fluctuations in chemical content across

char collections, the PYOM media used in this study was derived from a single acquisition of

char. We used DI water for all media types (LB, MYA, R2A, PyOM) and all were autoclaved at

121ºC for 45 minutes.

We added antimicrobials to the liquid agar after autoclaving, once it had cooled to 60ºC.

For media used to isolate fungi and intended to resist bacterial growth, we added filter sterilized

gentamycin and ampicillin (Hardy Diagnostics, Santa Maria, CA) to a final concentration of

10mg/mL each. For media used to isolate bacteria and intended to resist fungal growth, we added

filter sterilized cycloheximide (Research Products International, CAS # 66-81-9) to a final

concentration of 20mg/L.

Experimental Design and Microbial Isolation: We collected 12 soil samples from high-

severity burn locations (3 plots with four 1 m² subplots each). Three grams from each sample

were combined into a single homogenized soil sample, which was then subsampled for culturing

treatments: fresh soil plated on LB, MYA, R2A, or PyOM media, or soil stored for three months

under different conditions (air-dried, refrigerated at 4 °C, frozen at −80 °C, or frozen at −80 °C

with glycerol). For samples stored in glycerol, 1g of fresh soil was mixed with 1mL of glycerol

resulting in an approximately 1:1 w/v soil to glycerol slurry prior to freezing. Stored soil was

only plated onto rich media to not introduce additional confounding variables of media type into

the soil storage experiments. For each treatment, 1 g of soil was suspended in 5 mL sterile DI

water in four replicate Falcon tubes, serially diluted (1:10–1:10,000), and 100 µL of each

dilution was plated onto 10 replicate plates targeting either bacteria or fungi. All fungal plates

contained ampicillin and gentamycin, while bacterial plates contained cycloheximide. In total,

560 plates were prepared (2 kingdoms × 4 dilutions × 10 replicates × 7 treatments plus sterile

controls). Plates were incubated at room temperature (~25°C) for ~1 week, after which colony-

forming units (CFUs) and unique morphologies were counted. Plates containing 30–500 colonies

per plate were considered countable, yielding 6–9 replicate plates per treatment. Unique bacterial

and fungal morphotypes were isolated, DNA extracted, and Sanger sequenced, and isolates

stocked (Figure 1). Diversity was assessed using genus richness (a measurement of α-diversity)

of total or pyrophilous genera captured by each treatment.

Microbes were morphotyped by size, color, and growth rate. For bacteria and yeasts,

individual colonies were picked with sterile pipet tips and streaked onto fresh LB (without

antifungals). For filamentous fungi, a 1 cm² agar square from the colony edge was transferred

onto fresh MYA (without antibacterials). When microbes failed to establish on LB or MYA, they

were maintained on their original media (R2A or PyOM). Isolates were incubated at room

temperature for 1 week and re-isolated until three consecutive clean cultures were obtained for

DNA extraction and stocking.

For long-term storage, bacterial isolates were grown in LB broth (3 mL) at 180 rpm for

24–72 hrs until turbid, then mixed 1:1 with 40% sterile glycerol (final glycerol concentration of

20%) in cryovials and stored at −80 °C. Fungal isolates were grown to within 1 cm of the plate

edge, then 5 mm agar plugs from the colony margin were transferred into screw-cap tubes

containing 1 mL autoclaved tap water (79). Duplicate stock tubes were prepared for each isolate.

DNA Extraction: DNA was extracted from all unique isolates using home-made Extract-

N-Amp Extraction Solution (80) made by combining 10 ml of 1M pH 8.0 Tris-HCl, 1.86g KCl,

0.37g EDTA, and 80 mL DI water, then adding 1M NaOH until pH 9.5 and then diluting to a

total volume of 100mL and filter sterilizing. For bacteria, one well-isolated colony was picked

using a sterile 10μL pipet tip, and for fungi, a 1mm^2 area of the growing edge of the fungus was

scraped with a flame-sterilized probe. The tip or probe was then submerged and swirled in 10μL

of Extraction solution, which was heated at 65ºC for 10 minutes followed by 95ºC for 10 minutes

in a Thermocycler (BioRad). Once finished, 10μL of 3% BSA was added and the samples were

left to chill overnight at 4ºC before being PCR amplified or frozen at -20ºC.

PCR and Sanger Sequencing: The bacterial 16S rRNA region was amplified using a PA

forward and PH reverse primer (Edwards et al., 1989) and the fungal ITS region was amplified

with the ITS1F forward (Gardes & Bruns, 1993) and ITS4 reverse primer (White et al., 1990).

Each 20 μl PCR reaction contained 13.64 μl of molecular grade water, 0.16μL of Taq polymerase

(New England Biolabs, Ipswitch MA) and 2μL of 10x Taq Polymerase Buffer containing $MgCl_2$,

2μL of 10x solution of equally mixed dNTPs, 0.6μL of each primer at 10μM, and 1μL of DNA.

Thermocycler conditions were as follows: 94°C for 2 minutes, then 35 cycles of 57°C (for fungi)

or 62°C (for bacteria) for 1 minute, 72°C for 1 minute, then 94°C for 1 minute, ending with a

final extension of 72°C for 8 minutes (81). The DNA was cleaned using 3μL of diluted Exosap

mastermix (Applied Biosystems, Waltham MA) per 7μL PCR product before diluting and

preparing for Sanger sequencing. Samples were Sanger sequenced in the forward direction at the

UCR Institute for Integrative Genome Biology Genomics Core services.

BLAST Identification: Raw sequences as .ab1 chromatogram files were trimmed in either

SnapGene or Geneious Prime using the same parameters to remove low quality regions (Phred

scores less than 30, preserving the largest continuous region of good quality sequence) then

saved as FASTA files. Taxa were identified using NCBI BLASTn by generating hit tables of the

top 10 matches, with IDs assigned based on query cover, e-value, percent identity, and consensus

among hits of similar quality. Isolates were identified to species if query cover was 100%, e-

value < 0.01, and percent identity >98%; to genus at ~95%, family at 90%, class at 85–90%, and

phylum at <85%. However, given the difficulty of identifying species with only rRNA fragments,

genus level identification is the most reliable and species identifications reflect our best

estimation of taxon identity. All isolate names were preserved from the best-matching BLAST

identification, including any additional modifiers indicating potentially problematic taxonomy.

Statistical Analysis: All statistical analyses and figures were produced in R 4.0.2 (82) and

all scripts are available on GitHub

(https://github.com/DylanEnright93/Culturing_Pyrophilous_Microbes). Normality was tested

with a Shapiro test, then a two-way ANOVA was used to test the effect of soil storage or media

treatment on bacterial and fungal CFU counts, followed by a post-hoc Tukey HSD test.

Heatmap-like visualization of genera and species level diversity were created using the

"geom_tile" function in ggplot2 (83). Diversity among treatments was evaluated using α-

diversity by calculating the richness of genera per treatment and comparing the percent of total

isolates or total pyrophilous isolates (based on literature support) obtained per treatment.

Data Availability: All bacterial 16S and fungal ITS FASTA sequences are accessioned

under accession numbers PV662515 - PV662757 and PV620023 – PV620259, respectively and

will be available at the National Center for Biotechnology at Genbank database at time of

publication. All scripts used to analyze the data are likewise available on GitHub at

https://github.com/DylanEnright93/Culturing_Pyrophilous_Microbes.

Acknowledgements

We thank the San Bernardino National Forest for allowing us to collect the soils needed for this study following the El Dorado wildfire. We thank many individuals who contributed to the Glassman lab pyrophilous microbial culture including lab managers Judy Chung, James Randolph, and Maria Ordonez, UCR PhD students Fabiola Pulido-Chavez, Arik Joukhajian, and Mark Yacoub, and UCR undergraduates Justin Diab, Wine-Jie Lipardo, Audrey Reichard, Priscilla Shultz, Jorge Ponce, and Nathan Heger. This project was funded by the Department of Energy BER Award #DE-SC0023127 and United States Department of Agriculture-NIFA Award #2022-67014-36675 to SIG and the NSF GRFP to DJE. We also thank Dylan Enright's Dissertation Committee members Jason Stajich and Marko Spasojevic and anonymous reviewers for their helpful feedback on the manuscript.

Competing Interests

The authors declare no conflicts of interest or competing financial interests in relation to the work described.

References

1.   Franzosa EA, Hsu T, Sirota-Madi A, Shafquat A, Abu-Ali G, Morgan XC, Huttenhower C.

2015. Sequencing and beyond: integrating molecular "omics" for microbial community

profiling. Nat Rev Microbiol 13:360–372.

2.   Hugenholtz P. 2002. Exploring prokaryotic diversity in the genomic era. Genome Biol

3:reviews0003.1.

3.   Haridas S, Breuill C, Bohlmann J, Hsiang T. 2011. A biologist's guide to *de novo* genome

assembly using next-generation sequence data: A test with fungal genomes. J Microbiol

Methods 86:368–375.

4.   Malik KA, Claus D. 1987. Bacterial Culture Collections: Their Importance to

Biotechnology and Microbiology. Biotechnol Genet Eng Rev 5:137–198.

5.   Kämpfer P. 2014. Chapter 14 - Continuing Importance of the "Phenotype" in the Genomic

Era, p. 307–320. *In* Goodfellow, M, Sutcliffe, I, Chun, J (eds.), Methods in Microbiology.

Academic Press.

6.   Alster CJ, Allison SD, Johnson NG, Glassman SI, Treseder KK. 2021. Phenotypic plasticity

of fungal traits in response to moisture and temperature. ISME Commun 1:43.

7.   Lustenhouwer N, Maynard DS, Bradford MA, Lindner DL, Oberle B, Zanne AE, Crowther

TW. 2020. A trait-based understanding of wood decomposition by fungi. Proc Natl Acad

Sci 117:11551–11558.

8.   Day NJ, Dunfield KE, Johnstone JF, Mack MC, Turetsky MR, Walker XJ, White AL,

Baltzer JL. 2019. Wildfire severity reduces richness and alters composition of soil fungal

576 communities in boreal forests of western Canada. Glob Change Biol

577 https://doi.org/10.1111/gcb.14641.

578 9. Maynard DS, Bradford MA, Covey KR, Lindner D, Glaeser J, Talbert DA, Tinker PJ,

579 Walker DM, Crowther TW. 2019. Consistent trade-offs in fungal trait expression across

580 broad spatial scales. 5. Nat Microbiol 4:846–853.

581 10. Camenzind T, Aguilar-Trigueros CA, Hempel S, Lehmann A, Bielcik M, Andrade-Linares

582 DR, Bergmann J, dela Cruz J, Gawronski J, Golubeva P, Haslwimmer H, Lartey L, Leifheit

583 E, Maaß S, Marhan S, Pinek L, Powell JR, Roy J, Veresoglou SD, Wang D, Wulf A, Zheng

584 W, Rillig MC. 2024. Towards establishing a fungal economics spectrum in soil saprobic

585 fungi. Nat Commun 15:3321.

586 11. Swenson TL, Karaoz U, Swenson JM, Bowen BP, Northen TR. 2018. Linking soil biology

587 and chemistry in biological soil crust using isolate exometabolomics. Nat Commun 9:19.

588 12. Reyes-César A, Absalón ÁE, Fernández FJ, González JM, Cortés-Espinosa DV. 2014.

589 Biodegradation of a mixture of PAHs by non-ligninolytic fungal strains isolated from crude

590 oil-contaminated soil. World J Microbiol Biotechnol 30:999–1009.

591 13. Carini P. 2019. A "Cultural" Renaissance: Genomics Breathes New Life into an Old Craft.

592 mSystems 4.

593 14. Caktu K, Turkoglu E. 2011. Microbial Culture Collections: The Essential Resources for

594 Life. 2. Gazi Univ J Sci 24:175–180.

15. Stackebrandt E, Smith D, Casaregola S, Varese GC, Verkleij G, Lima N, Bridge P. 2014.

Deposit of microbial strains in public service collections as part of the publication process

to underpin good practice in science. SpringerPlus 3:208.

16. Arnold AE, Lutzoni F. 2007. Diversity and Host Range of Foliar Fungal Endophytes: Are

Tropical Leaves Biodiversity Hotspots? Ecology 88:541–549.

17. Davis KER, Joseph SJ, Janssen PH. 2005. Effects of Growth Medium, Inoculum Size, and

Incubation Time on Culturability and Isolation of Soil Bacteria. Appl Environ Microbiol

71:826–834.

18. Stevenson BS, Eichorst SA, Wertz JT, Schmidt TM, Breznak JA. 2004. New Strategies for

Cultivation and Detection of Previously Uncultured Microbes. Appl Environ Microbiol

70:4748–4755.

19. Leadbetter JR. 2003. Cultivation of recalcitrant microbes: cells are alive, well and revealing

their secrets in the 21st century laboratory. Curr Opin Microbiol 6:274–281.

20. Prakash O, Shouche Y, Jangid K, Kostka JE. 2013. Microbial cultivation and the role of

microbial resource centers in the omics era. Appl Microbiol Biotechnol 97:51–62.

21. Martiny AC. 2019. High proportions of bacteria are culturable across major biomes. ISME J

13:2125–2128.

22. Seaver FJ, Clark ED. 1910. Studies in Pyrophilous Fungi: II. Changes Brought about by the

Heating of Soils and Their Relation to the Growth of Pyronema and Other Fungi.

Mycologia 2:109–124.

23.   Pérez-Valera E, Verdú M, Navarro-Cano JA, Goberna M. 2018. Resilience to fire of
phylogenetic diversity across biological domains. Mol Ecol 27:2896–2908.

24.   Pressler Y, Moore JC, Cotrufo MF. 2019. Belowground community responses to fire: meta-
analysis reveals contrasting responses of soil microorganisms and mesofauna. Oikos
128:309–327.

25.   Fox S, Sikes BA, Brown SP, Cripps CL, Glassman SI, Hughes K, Semenova-Nelsen T,
Jumpponen A. 2022. Fire as a driver of fungal diversity — A synthesis of current
knowledge. Mycologia 114:215–241.

26.   Wan S, Hui D, Luo Y. 2001. Fire Effects on Nitrogen Pools and Dynamics in Terrestrial
Ecosystems: A Meta-Analysis. Ecol Appl 11:1349–1365.

27.   Pierson DN, Robichaud PR, Rhoades CC, Brown RE. 2019. Soil carbon and nitrogen
eroded after severe wildfire and erosion mitigation treatments. Int J Wildland Fire 28:814–
821.

28.   Michelotti LA, Miesel JR. 2015. Source Material and Concentration of Wildfire-Produced
Pyrogenic Carbon Influence Post-Fire Soil Nutrient Dynamics. 4. Forests 6:1325–1342.

29.   de Groot WJ, Flannigan MD, Cantin AS. 2013. Climate change impacts on future boreal
fire regimes. For Ecol Manag 294:35–44.

30.   Hantson S, Pueyo S, Chuvieco E. 2015. Global fire size distribution is driven by human
impact and climate. Glob Ecol Biogeogr 24:77–86.

31.   Joint I, Mühling M, Querellou J. 2010. Culturing marine bacteria – an essential prerequisite
for biodiscovery. Microb Biotechnol 3:564–575.

32.   Malosso E, Waite IS, English L, Hopkins DW, O'Donnell AG. 2006. Fungal diversity in
maritime Antarctic soils determined using a combination of culture isolation, molecular
fingerprinting and cloning techniques. Polar Biol 29:552–561.

33.   Černohlávková J, Jarkovský J, Nešporová M, Hofman J. 2009. Variability of soil microbial
properties: Effects of sampling, handling and storage. Ecotoxicol Environ Saf 72:2102–
2108.

34.   Rubin BER, Gibbons SM, Kennedy S, Hampton-Marcell J, Owens S, Gilbert JA. 2013.
Investigating the Impact of Storage Conditions on Microbial Community Composition in
Soil Samples. PLOS ONE 8:e70460.

35.   Caon L, Vallejo VR, Ritsema CJ, Geissen V. 2014. Effects of wildfire on soil nutrients in
Mediterranean ecosystems. Earth-Sci Rev 139:47–58.

36.   Preston CM, Schmidt MWI. 2006. Black (pyrogenic) carbon in boreal forests: a synthesis
of current knowledge and uncertainties. Biogeosciences Discuss 3:211–271.

37.   Woolet J, Whitman T. 2020. Pyrogenic organic matter effects on soil bacterial community
composition. Soil Biol Biochem 141:107678.

38.   Aranda E, Godoy P, Reina R, Badia-Fabregat M, Rosell M, Marco-Urrea E, García-Romera
I. 2017. Isolation of Ascomycota fungi with capability to transform PAHs: Insights into the

biodegradation mechanisms of Penicillium oxalicum. Int Biodeterior Biodegrad 122:141–

150.

39.   Fischer MS, Stark FG, Berry TD, Zeba N, Whitman T, Traxler MF. 2021. Pyrolyzed

Substrates Induce Aromatic Compound Metabolism in the Post-fire Fungus, Pyronema

domesticum. Front Microbiol 12.

40.   Pulido-Barriga MF, Nelson AR, Homyak PM, Wilkins MJ, Glassman SI. 2025. Chaparral

wildfire shifts the functional potential for soil pyrogenic organic matter and nitrogen

cycling. bioRxiv https://doi.org/10.1101/2024.08.18.607989.

41.   Knicker H. 2011. Pyrogenic organic matter in soil: Its origin and occurrence, its chemistry

and survival in soil environments. Quat Int 243:251–263.

42.   González-Pérez JA, González-Vila FJ, Almendros G, Knicker H. 2004. The effect of fire on

soil organic matter—a review. Environ Int 30:855–870.

43.   Lilleskov EA, Kuyper TW, Bidartondo MI, Hobbie EA. 2019. Atmospheric nitrogen

deposition impacts on the structure and function of forest mycorrhizal communities: A

review. Environ Pollut 246:148–162.

44.   Certini G. 2005. Effects of fire on properties of forest soils: a review. Oecologia 143:1–10.

45.   Enright DJ, Frangioso KM, Isobe K, Rizzo DM, Glassman SI. 2022. Mega-fire in redwood

tanoak forest reduces bacterial and fungal richness and selects for pyrophilous taxa that are

phylogenetically conserved. Mol Ecol 31:2475–2493.

46.   Pulido-Chavez MF, Randolph JWJ, Zalman C, Larios L, Homyak PM, Glassman SI. 2023.

673   Rapid bacterial and fungal successional dynamics in first year after chaparral wildfire. Mol

674   Ecol 32:1685–1707.

675   47.   Caiafa MV, Nelson AR, Borch T, Roth HK, Fegel TS, Rhoades CC, Wilkins MJ, Glassman

676   SI. 2023. Distinct fungal and bacterial responses to fire severity and soil depth across a ten-

677   year wildfire chronosequence in beetle-killed lodgepole pine forests. For Ecol Manag

678   544:121160.

679   48.   Petersen PM. 1970. Danish fireplace fungi-an ecological investigation on fungi on burns.

680   Dan Bot Ark 27:1–97.

681   49.   Whitman T, Whitman E, Woolet J, Flannigan MD, Thompson DK, Parisien M-A. 2019. Soil

682   bacterial and fungal response to wildfires in the Canadian boreal forest across a burn

683   severity gradient. Soil Biol Biochem 138:107571.

684   50.   Bárcenas-Moreno G, Bååth E. 2009. Bacterial and fungal growth in soil heated at different

685   temperatures to simulate a range of fire intensities. Soil Biol Biochem 41:2517–2526.

686   51.   Ferrenberg S, O'Neill SP, Knelman JE, Todd B, Duggan S, Bradley D, Robinson T, Schmidt

687   SK, Townsend AR, Williams MW, Cleveland CC, Melbourne BA, Jiang L, Nemergut DR.

688   2013. Changes in assembly processes in soil bacterial communities following a wildfire

689   disturbance. 6. ISME J 7:1102–1111.

690   52.   Sari E, Enright DJ, Ordonez M, Cordova-Ortiz E, Byrd A, Allison SD, Homyak PM,

691   Wilkins MJ, Glassman SI. 2025. Strain-Level Adaptation of Pyrophilous Bacteria through

Gene Fragmentation and Horizontal Gene Transfer. bioRxiv

https://doi.org/10.1101/2025.11.11.687939.

53. Sari E, Enright DJ, Ordonez M, Allison SD, Homyak PM, Wilkins MJ, Glassman SI. 2025.

Gene Duplication, Horizontal Gene Transfer, and Trait Trade-offs Drive Evolution of Post-

Fire Resource Acquisition in Pyrophilous Fungi. bioRxiv

https://doi.org/10.1101/2025.07.17.665399.

54. Kuo H-C, Hui S, Choi J, Asiegbu FO, Valkonen JPT, Lee Y-H. 2014. Secret lifestyles of

Neurospora crassa. Sci Rep 4:5135.

55. Bárcenas-Moreno G, García-Orenes F, Mataix-Solera J, Mataix-Beneyto J, Bååth E. 2011.

Soil microbial recolonisation after a fire in a Mediterranean forest. Biol Fertil Soils 47:261–

272.

56. Brown S, Veach A, Horton J, Ford E, Jumpponen A, Baird R. 2019. Context dependent

fungal and bacterial soil community shifts in response to recent wildfires in the Southern

Appalachian Mountains. For Ecol Manag https://doi.org/10.1016/j.foreco.2019.117520.

57. Whitman T, Woolet J, Sikora M, Johnson DB, Whitman E. 2022. Resilience in soil bacterial

communities of the boreal forest from one to five years after wildfire across a severity

gradient. Soil Biol Biochem 172:108755.

58. Xia Q, Rufty T, Shi W. 2020. Soil microbial diversity and composition: Links to soil texture

and associated properties. Soil Biol Biochem 149:107953.

59.   Pulido Barriga MF, Randolph JWJ, Glassman SI. 2024. Not all is lost: resilience of

712   microbiome samples to freezer failures and long-term storage. mSphere 10:e00603-24.

713   60.   Falconer RE, Otten W, White NA. 2015. Chapter One - Toward Modeling the Resistance

714   and Resilience of "Below-ground" Fungal Communities: A Mechanistic and Trait-Based

715   Approach, p. 1–44. *In* Sariaslani, S, Gadd, GM (eds.), Advances in Applied Microbiology.

716   Academic Press.

717   61.   Castaño C, Parladé J, Pera J, Martínez de Aragón J, Alday JG, Bonet JA. 2016. Soil drying

718   procedure affects the DNA quantification of Lactarius vinosus but does not change the

719   fungal community composition. Mycorrhiza 26:799–808.

720   62.   Cui H, Wang C, Gu Z, Zhu H, Fu S, Yao Q. 2014. Evaluation of soil storage methods for

721   soil microbial community using genetic and metabolic fingerprintings. Eur J Soil Biol

722   63:55–63.

723   63.   Reasoner DJ, Geldreich EE. 1985. A new medium for the enumeration and subculture of

724   bacteria from potable water. Appl Environ Microbiol 49:1–7.

725   64.   Ho A, Di Lonardo DP, Bodelier PLE. 2017. Revisiting life strategy concepts in

726   environmental microbial ecology. FEMS Microbiol Ecol 93:fix006.

727   65.   Hernandez DJ, David AS, Menges ES, Searcy CA, Afkhami ME. 2021. Environmental

728   stress destabilizes microbial networks. ISME J 15:1722–1734.

729   66.   Vieira FCS, Nahas E. 2005. Comparison of microbial numbers in soils by using various

730   culture media and temperatures. Microbiol Res 160:197–202.

67.   Sánchez-Clemente R, Igeño MI, Población AG, Guijo MI, Merchán F, Blasco R. 2018.

Study of pH Changes in Media during Bacterial Growth of Several Environmental Strains.

20. Proceedings 2:1297.

68.   Cray JA, Bell ANW, Bhaganna P, Mswaka AY, Timson DJ, Hallsworth JE. 2013. The

biology of habitat dominance; can microbes behave as weeds? Microb Biotechnol 6:453–

492.

69.   Kobziar LN, Pingree MRA, Larson H, Dreaden TJ, Green S, Smith JA. 2018.

Pyroaerobiology: the aerosolization and transport of viable microbial life by wildland fire.

Ecosphere 9:e02507.

70.   Zeba N, Berry TD, Fischer MS, Traxler MF, Whitman T. 2024. Soil carbon mineralization

and microbial community dynamics in response to pyrogenic organic matter addition. Soil

Biol Biochem 191:109328.

71.   Ben-Jacob E, Cohen I, Gutnick DL. 1998. COOPERATIVE ORGANIZATION OF

BACTERIAL COLONIES: From Genotype to Morphotype. Annu Rev Microbiol 52:779–

806.

72.   Sousa AM, Machado I, Nicolau A, Pereira MO. 2013. Improvements on colony

morphology identification towards bacterial profiling. J Microbiol Methods 95:327–335.

73.   Watrud LS, Martin K, Donegan KK, Stone JK, Coleman CG. 2006. Comparison of

taxonomic, colony morphotype and PCR-RFLP methods to characterize microfungal

diversity. Mycologia 98:384–392.

74.   Collado J, Platas G, Paulus B, Bills GF. 2007. High-throughput culturing of fungi from
752   plant litter by a dilution-to-extinction technique. FEMS Microbiol Ecol 60:521–533.

753   75.   Yang S-J, Kang I, Cho J-C. 2016. Expansion of Cultured Bacterial Diversity by Large-Scale
754   Dilution-to-Extinction Culturing from a Single Seawater Sample. Microb Ecol 71:29–43.

755   76.   Bartelme RP, Custer JM, Dupont CL, Espinoza JL, Torralba M, Khalili B, Carini P. 2020.
756   Influence of Substrate Concentration on the Culturability of Heterotrophic Soil Microbes
757   Isolated by High-Throughput Dilution-to-Extinction Cultivation. mSphere
758   5:10.1128/msphere.00024-20.

759   77.   Enright DJ. 2024. Unraveling Fire Adaptation Traits in Pyrophilous Bacteria and Fungi.
760   Ph.D. University of California, Riverside, United States -- California.

761   78.   Chase AB, Karaoz U, Brodie EL, Gomez-Lunar Z, Martiny AC, Martiny JBH. 2017.
762   Microdiversity of an Abundant Terrestrial Bacterium Encompasses Extensive Variation in
763   Ecologically Relevant Traits. mBio 8:10.1128/mbio.01809-17.

764   79.   McGinnis MR, Padhye AA, Ajello L. 1974. Storage of Stock Cultures of Filamentous
765   Fungi, Yeasts, and Some Aerobic Actinomycetes in Sterile Distilled Water. Appl Microbiol
766   28:218–222.

767   80.   Kreader CA, Corporation S-A, Louis S. The Extract-N-Amp[TM] Blood Kit: Rapid DNA
768   Extraction Coupled with PCR.

769   81.   Bruns TD, Fogel ,Robert, and Taylor JW. 1990. Amplification and Sequencing of Dna from
770   Fungal Herbarium Specimens. Mycologia 82:175–184.

771   82.   R Core Team. 2020. R: A Language and Environment for Statistical Computing. R

772   Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

773   83.   Wickham H, Chang W, Henry L, Pedersen TL, Takahashi K, Wilke C, Woo K, Yutani H,

774   RStudio. 2019. ggplot2: Create Elegant Data Visualisations Using the Grammar of Graphics

775   (3.2.1).

Figures:

Figure 1:

Figure 1: A conceptual diagram of the experimental design used in this experiment. "LB" is
abbreviated for Lysogeny Broth, "MYA" is Malt Yeast Agar, "R2A" is Reasoner's 2 Agar, and
"PyOM" is Pyrogenic Organic Matter. Note that the cultures obtained from fresh soil plated onto
rich media also serve as the "Fresh" storage condition for the soil storage experiment. Green
circles and orange stars symbolize two different morphotypes collected from the same initial
plate.

Figure 2:

Figure 2: A snapshot of the morphological diversity of the cultures obtained from the El Dorado
Fire across all soil storage and media type experiments. Identities were obtained via Sanger
sequencing of the 16S or ITS regions for bacteria and fungi, respectively. Moving left to right
across each row, the isolates shown are A)*Glutamicibacter bergerei, Arthrobacter agilis,*
Streptomyces mauvecolor, Kocuria rosea, B) *Arthrobacter bussei, Holtermaniella festucosa,*
Pedobacter panaciterrae, Curtobacterium oceanosedimentum, C) *Penicillium radiatolobatum,*
Penicillium fagi, Penicillium glabrum, Penicillium thomii, D) *Penicillium chalabudae,*
Penicillium radiatolobatum, Penicillium murcianum, Penicillium adametzii.

Figure 3:

Figure 3: Calculated capturable CFUs per gram soil for A) bacteria and B) fungi among the
tested soil storage methods and C) bacteria and D) fungi across tested media types. Points and
bars are colored by treatment with bars indicating standard error of the mean and black points
indicating the mean CFUs per gram soil for each treatment. Significance was tested using
ANOVA and Tukey Post-Hoc lettering has been applied. LB = Lysogeny Broth; PyOM =
Pyrogenic Organic Matter; R2A = Reasoner's 2 Agar, and G stands for glycerol.

Figure 4:

Figure 4: Summarized counts of A) total bacterial, B) pyrophilous bacterial, C) total fungal, and
D) pyrophilous fungal genera across four soil storage methods compared to fresh soil when
isolated on LB for bacteria or MYA for fungi. Treatments are ordered by highest amount of
genera. G = glycerol.

Figure 5:

Figure 5: Summarized counts of A) total bacterial, B) pyrophilous bacterial, C) total fungal, and
D) pyrophilous fungal genera (green) obtained from fresh soil tested across three media types.
Treatments are ordered by highest number of genera. LB = Lysogeny Broth, MYA = Malt Yeast
Agar, R2A = Reasoner's 2 Agar, PyOM = Pyrogenic Organic Matter.

Figure 6:

Figure 6: Heatmap-style visualization showing the diversity of pyrophilous A) bacterial and B)
fungal genera across the soil storage experiment. A filled blue square indicates the presence of
the genus on the x axis in the soil storage method indicated on the y axis. G = glycerol. Genera
are ordered in alphabetical order, other than [Brevibacterium] which is in brackets due to
controversy over its taxonomy.

Figure 7:

Figure 7: Heatmap-style visualization showing the diversity of pyrophilous A) bacterial and B)
fungal genera across the media types experiment. A filled blue square indicates the presence of
the genus on the x axis in the media indicated on the y axis. LB = Lysogeny Broth, MYA = Malt
Yeast Agar, R2A = Reasoner's 2 Agar, PyOM = Pyrogenic Organic Matter. Genera are ordered in
alphabetical order, other than [Brevibacterium] which is in brackets due to controversy over its
taxonomy.