Evaluating Best Practices for Isolating Pyrophilous Bacteria and Fungi from Burned Soil

Dylan J. Enright[1*], Aishwarya Veerabahu[1], Ryan J. Quaal[1], Marely Vega[1], Anna Nguyen[1], Jenna
Grindeland[1], James W.J. Randolph[1], Maria E. Ordonez[1], and Sydney I. Glassman[1*]

Affiliations:

[1]University of California Riverside, Department of Microbiology and Plant Pathology, 900
University Ave. Riverside CA, 92521, USA

Corresponding authors: sydneyg@ucr.edu and denri009@ucr.edu

Key Words: fire, pyrophilous, culture, bacteria, fungi, soil.

Abstract

15   A live microbial culture is invaluable to assess traits and functions, yet culturing

immediately from fresh soil is logistically challenging and media selection is not trivial. Isolating

pyrophilous bacteria and fungi after wildfires from burned soil, we tested the best: 1) soil storage

method to retain culturable viability; 2) media for highest microbial diversity. We tested four soil

storage methods (dried, stored at 4ºC, stored at -80ºC alone or in glycerol) and compared to fresh

soil obtained 6 months after a severe California shrubland wildfire. We also tested 3 media types:

rich (Lysogeny Broth (LB) for bacteria; Malt Yeast Agar (MYA) for fungi), oligotrophic

(Reasoner's 2 Agar (R2A)) and media made from pyrogenic organic matter (PyOM). For

bacteria, storing soil frozen at -80ºC alone yielded the highest diversity of genera and with

glycerol preserved the most species. For fungi, storing soil at -80ºC preserved the most species

but preserved equivalent diversity of genera as fresh and refrigerated soil. R2A yielded the

highest bacterial and fungal diversity but some species of interest were only captured with

PyOM. Using a combination of methods from 2018-2022, we cultured >500 isolates (286

bacteria; 258 fungi) after 7 California wildfires for testing pyrophilous microbial traits.

Introduction

Although advances in the fields of molecular biology and DNA sequencing have freed

microbiologists from the need to work with culturable species exclusively (Franzosa et al. 2015),

few laboratory resources are as valuable as live cultures. Pure cultures are essential components

for high quality whole-genome assembly (Hugenholtz 2002; Haridas et al. 2011), genetic

engineering experiments (Malik and Claus 1987), and the phenotypic profiling of

microorganisms (Kämpfer 2014; Alster et al. 2021; Lustenhouwer et al. 2020; Day et al. 2019;

Maynard et al. 2019; Camenzind et al. 2024). Additionally, microbial isolates greatly improve

the interpretation of metabolomics data (Swenson et al. 2018), and possession of a diverse

culture collection can unlock a plethora of applied microbiological research directions (Malik

and Claus 1987; Reyes-César et al. 2014; Carini 2019). Once isolated and properly stocked,

culture collections become a permanent resource for future microbiological research and create a

veritable breeding ground for collaborative research (Caktu and Turkoglu 2011; Stackebrandt et

al. 2014; Arnold and Lutzoni 2007). However, there are many hurdles to creating microbial

culture collections and each collection has unique challenges (Caktu and Turkoglu 2011).

The various factors that can be modified when attempting to culture microorganisms are

as varied as there are microbes to isolate. For example, media selection may require grappling

with an endless number of variables including consideration of the proper carbon sources,

nitrogen and phosphorus content, salt concentrations, pH, and a host of other micronutrient

considerations (Carini 2019; Davis, Joseph, and Janssen 2005; Stevenson et al. 2004). One must

also consider incubation temperature and duration, water and oxygen availability, and whether to

alternate light and dark cycles or to make the media nutrient rich or poor (oligotrophic) (Davis,

Joseph, and Janssen 2005; Stevenson et al. 2004; Leadbetter 2003). This complexity contributes

to many microorganisms remaining uncultured (Prakash et al. 2013). Though recent advances in

our understanding of microbiology have begun to make the adage of "99% of microbes are

unculturable" outdated (Carini 2019; Stevenson et al. 2004; Leadbetter 2003; Martiny 2019),

culturing the majority of microorganisms still remains beyond our reach.

One group of understudied microorganisms that is of growing interest are the pyrophilous

(or "fire-loving") microbes. Pyrophilous microbes are the first to increase in abundance

following wildfire, and they can persist in the environment for many years post-fire (Seaver and

Clark 1910; Pérez-Valera et al. 2018; Pressler, Moore, and Cotrufo 2019). While there are many

studies identifying which microbes become more abundant post-fire, there are very few studies

characterizing how pyrophilous microbes interact with their environment (Fox et al. 2022).

Wildfires dramatically change landscapes and create legacy effects that can impact soil

microorganisms and their overall ecosystems for many years following fire (Wan, Hui, and Luo

2001; Pierson et al. 2019; Michelotti and Miesel 2015). With rising global frequency and severity

of wildfires across many ecosystems (de Groot, Flannigan, and Cantin 2013; Hantson, Pueyo,

and Chuvieco 2015), understanding how pyrophilous microbes modify and interact with their

environments is of critical importance. An important step to characterizing the ecology of

pyrophilous microbes is laboratory testing of cultured isolates to provide data on how individual

pyrophilous taxa compete for resources, cycle post-fire nutrients, and interact with the post-fire

environment. However, there are numerous technological and logistical challenges to creating a

culture collection of pyrophilous microbes.

Beyond the well-known hurdles of creating a microbial culture collection, attempting to

culture microbes associated with wildfires presents its own set of challenges. First, there are

logistical challenges associated with working with post-fire environmental samples. Wildfires are

unpredictable natural phenomena. As such, it is often not possible to have the infrastructure for

large-scale culturing efforts (personnel available, media selected and created, broth tubes or petri

dishes ordered, culturing conditions decided and established) in place when a wildfire occurs. As

part of this logistical navigation, post-fire soil samples may have to be stored for varying lengths

of time before they can be used. This is a logistical problem shared among many environmental

microbial culturing efforts ranging from marine systems (Joint, Mühling, and Querellou 2010) to

Antarctic soils (Malosso et al. 2006). While many insights have been gained regarding these

logistical hurdles such as culturing from samples as quickly as possible (Caktu and Turkoglu

2011; Černohlávková et al. 2009), and methods for freezing samples that improve cultural

viability, their efficacy is not always consistent and the viability of culturable microorganisms

may be reduced depending on the soil storage method (Černohlávková et al. 2009; Rubin et al.

2013).

In addition to the logistics, wildfires radically alter environmental nutrient and resource

landscapes (Caon et al. 2014; Pierson et al. 2019). Wildfire pyrolysis turns labile, easily

accessible carbon sources into difficult to degrade pyrogenic organic matter (PyOM) (Michelotti

and Miesel 2015; Preston and Schmidt 2006). This PyOM is often enriched in polycyclic

aromatic hydrocarbons (PAHs) that may require specialized metabolic pathways to process

(Reyes-César et al. 2014; Woolet and Whitman 2020; Aranda et al. 2017; Fischer et al. 2021;

Pulido-Barriga et al. 2025). This can lead to accumulated char persisting in the environment for

up to and exceeding 100 years (Knicker 2011). Additionally, wildfires can deposit an enormous

amount of inorganic nitrogen into the environment when organic nitrogen previously locked in

plant biomass becomes released during burning (Caon et al. 2014; González-Pérez et al. 2004;

Wan, Hui, and Luo 2001). This high-nitrogen environment can radically alter the nutrient

strategies that microorganisms may employ post-fire and limit the microbes capable of surviving.

For example, many species of fungi are known to be nitrophobic (Lilleskov et al. 2019). These

radical nutrient shifts may result in commonly used laboratory media not providing appropriate

resources to cultivate post-fire microbes.

Here, our goal was to create a large culture collection of pyrophilous microbes to use to

test the traits that govern pyrophilous microbial adaptation to fire and the post-fire environment

via biophysical assays and genomics. Keeping in mind the above described logistic and resource

and nutrient-related challenges, we used soil from a local high severity California chaparral

wildfire to test the best 1) methods to store post-fire soil to retain pyrophilous microbial

culturable viability and 2) laboratory media to capture the highest diversity of pyrophilous

microbes. For soil storage, we tested culturing immediately from fresh soils versus soils stored

for 3 months in four ways (air-dried, refrigerated at 4°C, and frozen at -80°C with and without

glycerol). For media type, we tested rich (lysogeny broth (LB) agar for bacteria, malt-yeast agar

(MYA) for fungi), oligotrophic (Reasoner's 2 Agar (R2A)), or a media created in-house from

PyOM. We tested the impact of our treatments on abundance (CFUs: Colony Forming Units) and

microbial richness and composition (assessed via Sanger sequencing of 16S for bacteria or ITS

for fungi; Figure 1). We hypothesized (H1) that air-dried soil would retain the greatest diversity

of pyrophilous bacteria and fungi compared to fresh soil since fires reduce soil moisture, so we

expected air drying soils would be the least dramatic disturbance for soil borne microorganisms.

We further hypothesized (H2) that a combination of the rich and PyOM media would yield the

greatest diversity of pyrophilous microbes as the rich media could grow a wide range of bacteria

and fungi and the PyOM media would capture those pyrophilous species with unique metabolic

requirements specific to post-fire burn scars. Although we specifically sought to create a

pyrophilous culture collection, the insights gleaned from this study not only showcase potential

strategies for culturing fastidious pyrophilous microorganisms but can also aid environmental

microbiologists in decision making about storage and media selection for many types of

environmental culturing. The culmination of this research was the creation of a large and diverse

collection of pyrophilous microorganisms of >540 isolates (286 bacteria; 258 fungi) from burned

soils, mushrooms, and smoke derived from 7 California wildfires.

Methods

Site Description: To test the best soil storage method and media to cultivate pyrophilous

microbes, we sampled from The El Dorado wildfire, which burned ~92km^2 Manzanita

(*Arctostaphylos)* dominated chaparral shrublands, in Yucaipa, CA from September through

November of 2020. We examined high-resolution successional data of bacterial and fungal

communities after a similar nearby high-severity Manzanita dominated chaparral wildfire

(Pulido-Chavez et al. 2023), and compared to other post-fire studies that had successional data

(Bárcenas-Moreno et al. 2011; Caiafa et al. 2023), to identify 6 months post-fire as the ideal time

to attempt to isolate a wide range of pyrophilous bacteria and fungi. We thus collected samples

from high severity burned soils in June 2021, 6 months post-fire. Burned sites were characterized

as having 50 to 75 percent slopes with dry, sandy soils in the Springdale family-Lithic

Xerorthents association (https://websoilsurvey.nrcs.usda.gov/app/).

Soil Collection and Storage: Approximately 250mL of soil from 0-10 cm below the ash

layer of burned soil was collected from 9 plots with ethanol cleaned releasable bulb planters and

placed into whirl-pak bags 6 months post-fire following methods used in a similar ecosystem

(Pulido-Chavez et al. 2023). After collection, soil was 2mm sieved and either used as fresh

inoculum for cultivation or stored in four different ways: air dried, refrigerated (4ºC), frozen (-

80ºC), or suspended 50:50 %w/v soil with glycerol to create a soil slurry and then frozen (-80ºC)

(Figure 1).

Culturing Medias Used: Fresh soil samples were cultured immediately onto rich (LB for

bacteria, MYA for fungi), oligotrophic (R2A), or PyOM media. After 3 months of storage, stored

soil samples were cultured onto rich media. This means that the cultures isolated from fresh soils

on rich media served as both the "rich media" condition for the media experiment and the "fresh"

comparison for the soil storage experiment (Figure 1).

We produced LB broth by adding 15g of powdered LB Broth (Fisher Scientific, CAS#

73049-73-7) and MYA broth by adding 5g each of powdered Malt Extract (Research Products

International, Genessee, CAS# 8002-48-0) and Yeast Extract (Sigma-Aldrich, Cat # 70161) per

liter of DI water. For solid agar media, 15g per liter of bacteriological grade agar (Apex

Chemicals, Cat # 20-273) was added to either the LB or MYA broth. R2A agar was produced

using powdered pre-mixed R2A agar (Criterion, Cat # 16721) following manufacturer directions.

We developed an in-house recipe of PyOM media using biochar harvested from the El

Dorado Fire burn sites (Figure S1). First, we ground biochar into a fine powder with a coffee

grinder (Chefman, Model: RJ44-OPP-RED) for 3-5 minutes at maximum speed. Then, we added

50g of ground biochar to a 2L Erlenmeyer flask, added 1200mL of DI water, covered the mouth

of the flask, stirred on a hotplate for 24 hours at 40°C, then removed from heat to settle for 8

hours. We then siphoned liquid contents into a clean Erlenmeyer flask, being careful to not

disturb the bottom sediment. We passed the liquid through cheesecloth, and then twice through a

coffee filter to remove any remaining sediment before transferring to clean containers for

autoclaving. We added 13g of bacteriological grade agar (Apex Chemicals, Cat # 20-273) to each

liter of PyOM media, since adding 15g resulted in under-dissolved agar in the final product. We

used DI water for all media types (LB, MYA, R2A, PyOM) and all were autoclaved at 121ºC for

45 minutes.

We added antimicrobials to the liquid agar after autoclaving, once it had cooled to 60ºC.

For media used to isolate fungi and intended to resist bacterial growth, we added filter sterilized

gentamycin and ampicillin (Hardy Diagnostics, Santa Maria, CA) to a final concentration of

10mg/mL each. For media used to isolate bacteria and intended to resist fungal growth, we added

filter sterilized cycloheximide (Research Products International, CAS # 66-81-9) to a final

concentration of 20mg/L.

Experimental Design: We selected 12 soil samples collected from high-severity burn

locations (3 plots with four 1m^2 subplots each). We took 3g of soil from each of the 12 samples

and mixed them to create a single homogenous soil sample that was then subsampled for

culturing treatments (fresh soil for LB or MYA culturing, R2A, or PyOM, or stored for 3 months

air-dried, refrigerated, frozen, or frozen in glycerol). For each treatment, we made 4 separate 50

mL falcon tubes by combining 1g soil with 5mL of sterile DI water. We then created dilution

series ranging from 1:10 to 1:10,000 for each treatment and plated 100µL of each dilution onto

10 plates per target organism. We applied ampicillin and gentamycin to all plates targeting fungi

and cycloheximide to all plates targeting bacteria. We had a total of 560 experimental plates (2

Kingdoms (bacteria, fungi) x 4 dilutions (1:10, 1:100, 1:1,000, 1:10,000) x 10 replicates x 7

treatments [4 storage treatments (dried, 4ºC, -80ºC, -80ºC + glycerol) + 3 media types for fresh

soils (Rich, R2A, PyOM)]. After spreading solution onto plates, we wrapped them in parafilm

and kept them in a drawer at room temperature for ~1 week, after which we counted Colony

Forming Units (CFUs) and unique morphologies. Plates containing between 30-500 colonies of

well-defined growths of bacteria or fungi were considered "countable" plates and the initial CFU

concentrations were calculated by multiplying the number of colonies by the dilution factor

resulting in 6-9 countable replicate plates per treatment. We then isolated, extracted DNA,

Sanger sequenced, and stocked each bacterial and fungal unique morphology (Figure 1).

Diversity was assessed based on the percentage (total or pyrophilous) of genera or species

captured by each treatment.

Microbial Isolation Method: Fungi and bacteria were morphotyped using size, color, and

growth rate. For bacteria or yeasts, we used a sterile 10µL pipet tip to pick a well isolated colony

and then used the streak plate method onto either fresh LB plate without antifungals for bacteria

or a fresh MYA plate without antibacterials for fungi. For filamentous fungi, we used a flame-

sterilized scalpel to cut a 1 cm^2 square of the growing edge of a fungus from the agar plate, then

transferred the square growing-side down into the center of a fresh MYA plate without

antibacterials. In cases where microbes would not establish onto LB or MYA, they were

maintained on the media they originated from (either R2A or PyOM). We incubated plates at

room temperature for 1 week before re-isolating following the same procedure. This was

repeated until three clean isolations were obtained prior to DNA extraction and stocking.

Microbial Stocking Method: Each isolate was stocked for future use after a thrice

purified isolate was obtained and DNA was extracted. For bacteria, a colony was picked

aseptically from the last isolation streak plate using a sterile 10µL pipet tip and placed into 3mL

of LB broth in a sterile 15mL Falcon tube, then grown at shaking incubation at 180rpm for 24-

72hrs until broth became turbid. We added 1mL of the turbid culture to a 2mL cryovial and

combined with 1mL of sterile 40% glycerol, resulting in a final 20% glycerol suspension. This

was vortexed briefly and then frozen at -80ºC. For fungi, a 1cm^2 cube of fungus was allowed to

grow until it came within approximately 1cm of the edge of the 100mm Petri dish. Then, using a

sterile transfer tube (Spectrum Laboratories Inc., San Clemente CA), 5mm disks were punched

from the growing edge of the fungus. We transferred 8-10 disks to a 2mL screw-cap tube

containing 1mL of autoclaved tap water (McGinnis, Padhye, and Ajello 1974) then stored at

room temperature. We made duplicate stock tubes for each isolate.

DNA Extraction: DNA was extracted from all unique isolates using home-made Extract-

N-Amp Extraction Solution made by combining 10 ml of 1M pH 8.0 Tris-HCl, 1.86g KCl, 0.37g

EDTA, and 80 mL DI water, then adding 1M NaOH until pH 9.5 and then diluting to a total

volume of 100mL and filter sterilizing. For bacteria, one well-isolated colony was picked using a

sterile 10µL pipet tip, and for fungi, a 1mm^2 area of the growing edge of the fungus was scraped

with a flame-sterilized probe. The tip or probe was then submerged and swirled in 10µL of

Extraction solution, which was heated at 65ºC for 10 minutes followed by 95ºC for 10 minutes in

a Thermocycler (BioRad). Once finished, 10µL of 3% BSA was added and the samples were left

to chill overnight at 4ºC before being PCR amplified or frozen at -20ºC.

PCR and Sanger Sequencing: The bacterial 16S rRNA region was amplified using a PA

forward and PH reverse primer (Edwards et al., 1989) and the fungal ITS region was amplified

with the ITS1F forward (Gardes & Bruns, 1993) and ITS4 reverse primer (White et al., 1990).

Each 20 µl PCR reaction contained 13.64 µl of molecular grade water, 0.16µL of Taq polymerase

(New England Biolabs, Ipswitch MA) and 2µL of 10x Taq Polymerase Buffer containing $MgCl_2$,

2µL of 10x solution of equally mixed dNTPs, 0.6µL of each primer at 10µM, and 1µL of DNA.

Thermocycler conditions were as follows: 94°C for 2 minutes, then 35 cycles of 57°C (for fungi)

or 62°C (for bacteria) for 1 minute, 72°C for 1 minute, then 94°C for 1 minute, ending with a

final extension of 72°C for 8 minutes (Bruns and and Taylor 1990). The DNA was cleaned using

3µL of diluted Exosap mastermix (Applied Biosystems, Waltham MA) per 7µL PCR product

before diluting and preparing for Sanger sequencing. Samples were Sanger sequenced in the

forward direction using either Genewiz (San Diego, CA), Eurofins Genomics (Louisville, KY),

or the UCR Institute for Integrative Genome Biology Genomics Core services.

BLAST Identification: Raw sequences as .ab1 chromatogram files were trimmed in either

SnapGene or Geneious Prime to remove low quality regions then saved as FASTA files. Taxa

were identified using NCBI BLASTn to create hit tables of top 10 hits. IDs were assigned using

a combination of query cover, error value, percent identity, and a consensus of hits with the same

quality scores. In general, an isolate was identified at species level if the query cover was 100%,

max e score was < 0.01, and percent identity was >98%. If percent identity fell to ~95% then

isolate identifications were left to genus, at 90% to family, at 85-90% to class, and at < 85% to

phylum level. All isolate names were preserved from the best-matching BLAST identification,

including any additional modifiers indicating potentially problematic taxonomy.

Statistical Analysis: All statistical analyses and figures were produced in R 4.0.2 (R Core

Team 2020) and all scripts are available on GitHub

(https://github.com/DylanEnright93/Culturing_Pyrophilous_Microbes). Normality was tested

with a Shapiro test, then a two-way ANOVA was used to test the effect of soil storage or media

treatment on bacterial and fungal CFU counts, followed by a post-hoc Tukey HSD test.

Heatmap-like visualization of genera and species level diversity were created using the

"geom_tile" function in ggplot2 (Wickham et al. 2019). Diversity among treatments was

evaluated by calculating the number of species, genera, families, and orders per treatment and

comparing the percent of total or total pyrophilous isolates (based on literature support) obtained

per treatment. All bacterial 16S and fungal ITS FASTA sequences have been deposited to NCBI

GenBank under accession numbers PV662515 - PV662757 and PV620023 – PV620259,

respectively.

Results

Microbes Obtained: Throughout our experiments we isolated a remarkable array of

bacterial and fungal morphotypes (Figure 2). We cultured a total of 170 bacterial isolates

representing 4 phyla, 10 orders, 31 genera and 83 species from burned soils collected 6 months

after the 2020 El Dorado Fire across all our storage and media type experimental treatments. We

also cultured 160 fungal isolates representing 3 phyla, 9 orders, 15 genera and 35 species (Table

S1). Of these isolates, 138/170 bacterial isolates composed of 17 genera and 71 species and

152/160 fungal isolates composed of 8 genera and 30 species had literature support for inclusion

as putative or confirmed pyrophilous taxa (Enright et al. 2022; Pulido-Chavez et al. 2023;

Woolet and Whitman 2020; Aranda et al. 2017; Caiafa et al. 2023; Seaver and Clark 1910;

Petersen 1970; Whitman et al. 2019; Pierson et al. 2019; Bárcenas-Moreno and Bååth 2009;

Ferrenberg et al. 2013; Fox et al. 2022) (Table S2). We used the total (Table S1) and pyrophilous

isolates (Table S2) to test the effect of media type and soil storage method on CFUs and

microbial diversity. In addition to testing storage methods and media selection, here we present

our culture collection of >540 pyrophilous microbes (286 bacteria; 258 fungi) that we isolated

from 2018-2022 from burned soils, mushrooms, and smoke of 7 California wildfires for future

testing of pyrophilous microbial traits (Table S3). For bacteria, we had 286 isolates from 8 phyla,

16 orders, 35 genera, 76 species. For fungi, we had 258 isolates from 3 phyla, 11 orders, 13

genera, and 34 species. The full table includes the fires of origin, collection times, substrate

cultures derived from, isolate names, FASTA sequences, and BLAST summary statistics (Table

S3).

<u>Effect of Storage Method Selection on CFUs</u>: Soil storage method significantly impacted

the number of CFUs obtained per gram soil for both bacteria (ANOVA: $F_{1,4} = 19.97$, $p < 0.001$;

Figure 3A) and fungi (ANOVA: $F_{1,4} = 19.78$, $p < 0.001$; Figure 3B). For both bacteria and fungi,

soil stored air-dried (Bacteria: 2.31×10^5 CFUs/gram, Fungi: 1.33×10^4 CFUs/gram) and stored at -

80ºC with glycerol (Bacteria: 1.31×10^5 CFUs/gram, Fungi: 6.69×10^3 CFUs/gram) retained the

fewest mean CFUs/gram soil. For bacteria, soils stored at 4ºC (6.98×10^5 CFUs/gram) or at -80ºC

without glycerol (6.39×10^5 CFUs/gram) were not significantly different from fresh soils

(5.73×10^5 CFUs). For fungi, fresh soils had the highest mean CFUs/gram soil (3.07×10^4

CFUs/gram), followed by soil frozen at -80ºC without glycerol (2.66×10^4 CFUs/gram), and then

stored at 4ºC (2.16×10^4 CFUs/gram).

<u>Effect of Media Selection on CFUs</u>: Media type significantly impacted the number of

CFUs obtained per gram of fresh soil for both bacteria (ANOVA: $F_{1,2} = 39.43$, $p < 0.001$; Figure

3C) and fungi ($F_{1,2} = 7.77$, $p < 0.01$; Figure 3D). For both bacteria and fungi, PyOM media

yielded the fewest mean CFUs per gram (Bacteria: 1.66×10^5; Fungi: 1.03×10^4), and rich media

(LB for bacteria and MYA for fungi) had the highest average CFUs per gram (Bacteria:

5.73×10^5; Fungi: 3.07×10^4). R2A media had the second highest per gram mean CFU counts for

bacteria (3.20×10^5) and for fungi, CFU counts (2.20×10^4) were statistically equivalent to MYA.

<u>Effect of Storage Method Selection on Bacterial Diversity</u>: Soil storage method strongly

affected diversity of bacterial genera with fresh soil retaining the highest total (Figure 4A) and

frozen soil retaining the highest pyrophilous diversity (Figure 4B). Dried soil retained the lowest

diversity of bacterial genera among both total (Figure 4A) and pyrophilous isolates (Figure 4B).

Figures are ordered by number of genera first, which covers greater phylogenetic diversity, then

by number of species. Across all storage treatments, we cultured a total of 21 bacterial genera

(Figure S2) and 64 species (Figure S3) and we found literature support as pyrophilous for 14

genera (Figure 6A) and 51 species (Figure S4). Considering total bacterial taxa, fresh soil was

the most diverse (62% of genera, 41% of species), followed by frozen soil (62% of genera, 30%

of species), soil frozen with glycerol (57% of genera, 39% of species), refrigerated soil (48% of

genera, 22% of species) with dried soil as least diverse (38% of genera, 23% of species). Only 3

genera (*Arthrobacter, Curtobacterium, Streptomyces*) were isolated across all 5 storage

treatments whereas 8 genera (*Brevibacterium, Fontibacillus, Glutamicibacter, Massilia,*

Ornithinimicrobium, Paeniglutamicibacter, an unidentified Firmicute, and *Pseudoarthobacter*)

were isolated from a single treatment (Figure S2). Finally, 64% of species were isolated one time

with only one species *Curtobacterium oceanosedimentum* observed in all storage treatments

(Figure S3). Among the pyrophilous subset (Figure 4B), frozen soil was the most diverse (85%

of genera, 33% of species), followed by fresh soil (62% of genera, 35% of species), soil frozen

with glycerol (54% of genera, 39% of species), refrigerated soil (54% of genera, 22% of species)

with dried soil as least diverse (38% of genera, 24% of the species).

Effect of Storage Method Selection on Fungal Diversity: Soil storage method strongly

affected fungal diversity with fresh soil retaining the highest total genera (Figure 4C) and fresh,

frozen and refrigerated retaining equal numbers of pyrophilous genera (Figure 4D). Soil frozen

with glycerol retained the lowest diversity of fungal genera among total (Figure 4C) and

pyrophilous isolates (Figure 4D). Across all storage treatments, we cultured a total of 6 fungal

genera (Figure S5) and 23 fungal species (Figure S6), of which we found literature support as

pyrophilous for 3 genera (Figure 6B) and 20 species (Figure S7). For total fungi, fresh soil was

the most diverse, capturing 100% of all 6 genera (*Aspergillus, Holtermaniella,*

Ochrocladiosporium, *Penicillium*, and 2 uncultured Helotiales), followed by frozen and

refrigerated soil capturing 50% of the genera, and dried soil capturing 33% of total fungal genera

(Figure S5). Soil frozen with glycerol performed the worst, capturing only *Penicillium* (17% of

the total; Figure S5). For total fungal species, fresh soil captured 52%, followed by 43% in

frozen, then 30% in refrigerated and dried soil, and finally 26% in frozen with glycerol (Figure

S6). For pyrophilous fungi, fresh, frozen, and refrigerated soil all captured 100% of the 3 genera

(*Aspergillus, Holtermanniella, Penicillium*), whereas dried captured 2 (*Aspergillus, Penicillium*)

and frozen with glycerol only captured 1 (*Pencillium*; Figure 6B). For pyrophilous fungal

species, frozen soil captured 50%, followed closely by 45% in fresh, 35% in refrigerated and

30% in air-dried (Figure S7).

Effect of Media Selection on Bacterial Diversity: Media type strongly affected bacterial

diversity with R2A and LB both capturing the highest numbers of total bacterial genera and

species (Figure 5A), R2A yielding the highest diversity of pyrophilous genera and species

(Figure 5B), and PyOM capturing the fewest total and pyrophilous bacterial genera and species.

Across all media types from fresh soil, we cultured 24 total bacterial genera (Figure S8), 61 total

bacterial species (Figure S9), and 16 pyrophilous bacterial genera (Figure 7A) and 49

pyrophilous bacterial species (Figure S10). Among total bacteria, LB and R2A obtained equally

diverse communities (58% of genera, 44% of species), while PyOM captured the least diverse

(33% of genera, 18% of species). Among pyrophilous bacteria, R2A captured the greatest

diversity (69% of genera, 49% of species), followed by LB (56% of genera, 39% of species) with

PyOM yielding the lowest diversity (44% of genera, 20% of species). Although PyOM media

yielded the lowest diversity, it was the only media to isolate some important pyrophilous

bacterial genera (Whitman et al. 2019; Caiafa et al. 2023; Pulido-Chavez et al. 2023) like

Noviherbaspirillum and *Peribacillus* and interestingly, R2A was the only media to capture

Massilia (Figure 7A).

354   Effect of Media Selection on Fungal Diversity: Media type affected fungal diversity with

R2A capturing the most total (Figure 5C) and pyrophilous (Figure 5D) fungal genera and species

whereas MYA captured the fewest. Across all media types from fresh soil, we cultured 14 total

fungal genera (Figure S11), 29 total fungal species (Figure S12), and 8 pyrophilous fungal

genera (Figure 7B) and 23 pyrophilous fungal species (Figure S13). Among total fungi, R2A

media captured the most diversity (57% of genera, 48% of species) while both MYA and PyOM

were equally diverse (43% of genera, 41% of species). Among pyrophilous fungi, R2A also

captured the most diversity (89% of genera, 57% of species), followed by PyOM (50% of

genera, 43% of species), and MYA was the least diverse (38% of genera, 39% of species). While

PyOM did not capture the most diversity, it was the only media to capture *Neurospora* (Figure

7B), a well-documented post-fire fungus with relatively little knowledge about its ecology (Kuo

et al. 2014).

Discussion

Here, we present results of an experiment to test soil storage and media selection to

obtain the best diversity and abundance of bacterial and fungal isolates, which have informed our

continual culturing efforts resulting in a collection of >540 pyrophilous microbes (286 bacteria;

258 fungi) (Table S3). To our knowledge, this is the first study examining the impact of soil

storage on fungal diversity, and this is also the first large-scale culturing effort of pyrophilous

bacteria, and among the largest soil and mushroom based culturing efforts of pyrophilous fungi

since 1910 (Seaver and Clark 1910). For both bacteria and fungi, freezing soil without glycerol

yielded the highest number of pyrophilous genera and R2A yielded the most genera and species.

Although it did not capture the highest number of CFUs or highest number of taxa, novel solid

agar media we created from field-harvested PyOM was necessary to capture some key

pyrophilous genera of interest like *Neurospora* for fungi and *Noviherbaspirillium* for bacteria.

For bacteria, we conclude that freezing soil at -80°C is the best storage method relative to

fresh soil, yielding the highest number of CFUs and total and pyrophilous bacterial genera.

Interestingly, adding glycerol increased the number of species but reduced the number of total

and pyrophilous genera isolated. While CFU based analyses are useful in giving an

approximation of the total number of culturable cells present in the original sample and a broad

overview of the effectiveness of certain culturing practices at cultivating cells, it does not capture

the important distinction of isolate diversity. However, in this case, freezing soils without

glycerol captured both the highest number of CFUs and yielded the vast majority of the

pyrophilous bacterial genera of interest based on the post-fire bacterial succession literature

(Enright et al. 2022; Pulido-Chavez et al. 2023; Bárcenas-Moreno et al. 2011; Caiafa et al. 2023;

Brown et al. 2019; Whitman et al. 2019). While culturing from fresh soil did capture the greatest

number of total genera and species, freezing soil captured the greatest diversity of the

pyrophilous genera (Figure 4). Interestingly, freezing soil resulted in more total and pyrophilous

bacterial genera being isolated, but adding glycerol increased the number of species and was

necessary to isolate *Massilia*, which is a bacterial genus repeatedly observed post-fire (Pulido-

Chavez et al. 2023; Caiafa et al. 2023; Whitman et al. 2019; 2022), with very little understanding

about its ecology, making it a prime candidate for further study. Therefore, it may be of value for

researchers to freeze some soil with glycerol as well as without, to capture some of these more

potentially vulnerable taxa. Finally, dried soils performed the worst in capturing total and

pyrophilous bacterial diversity, which surprised us given how dry post-fire soils usually are. One

previous study found that refrigeration at 4ºC was better than freezing at -20ºC at retaining

microbial biomass and respiration (Černohlávková et al. 2009). This study examined unburned

grassland and forest soils, which could have different soil chemistry and texture from our burned

chapparal soils. A study examining how storage time (0-14 days) and temperature (room temp,

4°C, and -20°C) affected 16S sequencing results rather than culturing results similarly found that

freezing preserved the most bacterial diversity, though they did not test freezing to -80°C as we

did (Rubin et al. 2013). Both of these studies only froze at -20ºC rather than -80ºC, which may

have resulted in fewer taxa retained (Pulido Barriga, Randolph, and Glassman 2024).

Soil storage method seemed to matter less for pyrophilous fungi than bacterial genera.

While there was one method that stood out as the worst (freezing at -80°C with glycerol), the rest

of the methods captured most of the diversity observed (Figures 4D and 6B). This could be due

to the resilience of fungal spores to a variety of disturbance effects (Falconer, Otten, and White

2015), but it is also very possible that MYA was not the ideal culture medium for pyrophilous

fungi, given the success of R2A and PyOM media at cultivating pyrophilous fungi (Figures 5D

and 7B). Future work repeating the storage experiments with R2A and PyOM might improve

pyrophilous fungal isolate retention. Even with the media selection caveat to our storage

experiments, we were able to draw some very surprising conclusions from our experiments. For

example, our hypothesis that air-drying soil would be the most effective storage method was

fully unsupported by our results. Indeed, air-drying yielded the lowest CFU counts and total and

pyrophilous diversity. This is an important finding since drying soils prior to laboratory use is

not uncommon (Rubin et al. 2013; Castaño et al. 2016; Cui et al. 2014). It is worth noting that

we were unable to find other studies exploring the impact of soil storage on fungal communities

for comparison, indicating a strong need for these methods to be tested across kingdoms.

In assessing the most effective culturing media for capturing pyrophilous bacteria and

fungi, we were surprised at how well the oligotrophic media R2A performed. R2A produced the

second highest number of CFUs after the rich medias for bacteria (LB) and fungi (MYA) (Figure

3) and the most diverse pyrophilous bacterial (Figure 5B) and fungal (Figure 5D) isolates.

Considering that R2A was developed to culture bacteria from potable drinking water (Reasoner

and Geldreich 1985) which is extremely oligotrophic, this would suggest that a large number of

pyrophilous microbes prefer a more oligotrophic environment. While shifting life-strategies of

microbial communities under stress can be extremely varied and nuanced (Ho, Di Lonardo, and

Bodelier 2017), under an increasing stress gradient (such as carbon stress, soil moisture stress, or

heat stress) oligotrophic bacteria and fungi can be favored (Hernandez et al. 2021), which might

be ideal for capturing a wide diversity of microbes thriving in the stressful post-fire environment.

However, this is also a surprising result as post-fire environments tend to be high in nitrogen and

phosphorus, even if the carbon landscape has radically changed (Wan, Hui, and Luo 2001;

Michelotti and Miesel 2015). While previous studies exist examining the role of media

436 conditions on microbial culturability (Vieira and Nahas 2005; Sánchez-Clemente et al. 2018;

437 Malosso et al. 2006; Stevenson et al. 2004), these studies often take either a fine-scale approach

438 to tuning a specific culture medium or focus on very broad-scale patterns of culturability such as

439 relying on CFU-based metrics alone or use only rich media. Continued exploration of media

440 conditions is essential to increasing the number of cultured microbial species, and insights gained

441 from our study may help open new paths in culture media development.

442  Finally, although our home-made PyOM media captured the lowest CFUs of bacteria and

443 fungi, and the lowest bacterial diversity, it captured the second highest fungal diversity and was

444 necessary to isolate key bacterial (i.e. *Noviherbaspirillum soli*) and fungal (i.e. *Neurospora*

445 *crassa*) pyrophilous taxa of interest. In this case, isolating finicky pyrophilous taxa of interest

446 was improved by the low-density growth we observed in PyOM, which likely resulted from the

447 lower nutrient content similar to R2A media. Rich media like LB, MYA, and potato dextrose

448 agar are often plagued by fast growing ruderal species that can overwhelm the growth of taxa

449 that grow at lower density (Cray et al. 2013). These taxa are particularly interesting because

450 *Noviherbaspirillum* has been repeated observed increasing in abundance post-fire (Enright et al.

451 2022; Pulido-Chavez et al. 2023; Caiafa et al. 2023) despite very little understanding of its

452 ecology, and *Neurospora crassa* is famously found post-fire and is widely used as a genetics

453 model organism despite very little understanding about its ecological role in the environment

454 (Kuo et al. 2014). Therefore, we conclude that the best media to use to maximize pyrophilous

455 bacterial and fungal diversity if you can only select one is R2A, but best to also include a PyOM

456 media if specifically interested in isolating the most finicky and perhaps interesting pyrophilous

457 microbes.

While we made significant progress regarding testing effective culturing practices for

environmental microbes and specifically pyrophilous bacteria and fungi, much additional work

remains and caveats exist. Except for the home-made PyOM media, we limited ourselves to

commonly available laboratory culturing media. We acknowledge there is a near endless range of

possible medias to use (Vieira and Nahas 2005) and there may be more effective media available

for the culturing of pyrophilous microorganisms. However, we assert that the broad trends (rich

vs. oligotrophic vs. PyOM media) yield valuable insights into the culturing of these

microorganisms. As for the PyOM media itself, it is a complex undefined media and uses a

water-extraction based method of incorporating nutrients. This method might have resulted in

our media missing some key nutrients required for post-fire metabolism that are less water-

soluble in PyOM than others (Zeba et al. 2024), which could have led to us missing some

pyrophilous microbial taxa. Additionally, the culturing effort in this study all took place

aerobically at room temperature. Further, future work could endeavor to test a wider variety of

culturing and incubation factors including testing a range of growth temperatures (Bárcenas-

Moreno and Bååth 2009; Vieira and Nahas 2005) and media pH (Davis, Joseph, and Janssen

2005; Sánchez-Clemente et al. 2018). Finally, while we acknowledge that separating microbial

cultures based on colony morphology is difficult at best (Ben-Jacob, Cohen, and Gutnick 1998;

Sousa et al. 2013; Watrud et al. 2006), we decided that choosing colonies based on morphotype

and then identifying with Sanger sequencing was an acceptable concession to the technical and

logistic demands of isolating, stocking, and identifying every single colony we grew. This does

mean that we could have missed some species-level diversity since some species can share

extremely similar colony morphologies (Sousa et al. 2013; Watrud et al. 2006) and techniques

like dilution to extinction culturing might capture more isolates identified to species level

(Collado et al. 2007; Yang, Kang, and Cho 2016). Given these caveats, we still obtained a

remarkable diversity of total (Table S1) and pyrophilous (Table S2) bacteria and fungi and found

meaningful differences in media and storage types.

Conclusions

Here, we present the first study to systematically test how soil storage methods impact

fungal cultivability, and the largest scale test of how soil storage and media type affect bacterial

and fungal cultivation. We found that storing soils at -80°C without glycerol and isolating with

oligotrophic R2A media preserved the greatest diversity of culturable pyrophilous bacteria and

fungi compared to fresh soils from burned chaparral. However, certain pyrophilous taxa of

interest were only captured from soil frozen with glycerol, like *Massilia*, and on our novel PyOM

media, like *Neurospora* and *Noviherbaspirillium*, indicating a potential value in using two media

and storage types. This experiment contributed to a larger culturing effort resulting in a

collection of >540 pyrophilous microbes (286 bacteria; 258 fungi) isolated from soil, smoke, and

mushrooms isolated from 7 California wildfires (Table S3), which can be used for future multi-

omics, phenotypic, and applied microbiological research. For example, a culture collection is

necessary to perform biophysical assays and genomics to assess traits that can yield insights into

ecological processes and trade-offs (Maynard et al. 2019). Sequencing genomes from isolates

with concurrent biophysical assays is a powerful yet under-used tool to gain insight into

microbial traits and connect genomic information to trait expression (Chase et al. 2017).

Obtaining the cultures is the first step to characterizing their traits via 'omics and phenotypic

methods.

Acknowledgements

We thank the San Bernardino National Forest for allowing us to collect the soils needed

for this study following the El Dorado wildfire. We thank many individuals who contributed to

the Glassman lab pyrophilous microbial culture including lab managers Judy Chung, James

Randolph, and Maria Ordonez, UCR PhD students Fabiola Pulido-Chavez, Arik Joukhajian, and

Mark Yacoub, and UCR undergraduates Justin Diab, Wine-Jie Lipardo, Audrey Reichard,

Priscilla Shultz, Jorge Ponce, and Nathan Heger. This project was funded by the Department of

Energy BER Award #DE-SC0023127 and United States Department of Agriculture-NIFA Award

#2022-67014-36675 to SIG and the NSF GRFP to DJE. We also thank Dylan Enright's

Dissertation Committee members Jason Stajich and Marko Spasojevic and anonymous reviewers

for their helpful feedback on the manuscript.

Competing Interests

517   The authors declare no conflicts of interest or competing financial interests in relation to

518   the work described.

Data Accessibility and Benefit-Sharing:

521   Sequences are accessioned and will be available at the National Center for Biotechnology

522   at Genbank database at time of publication. All scripts used to analyze the data are likewise

523   available on GitHub at https://github.com/DylanEnright93/Culturing_Pyrophilous_Microbes.

References

Alster, Charlotte J, Steven D Allison, Nels G Johnson, Sydney I Glassman, and Kathleen K
Treseder. 2021. "Phenotypic Plasticity of Fungal Traits in Response to Moisture and
Temperature." *ISME Communications* 1 (1): 43. https://doi.org/10.1038/s43705-021-
00045-9.

Aranda, Elisabet, Patricia Godoy, Rocío Reina, Marina Badia-Fabregat, Mònica Rosell, Ernest
Marco-Urrea, and Inmaculada García-Romera. 2017. "Isolation of Ascomycota Fungi
with Capability to Transform PAHs: Insights into the Biodegradation Mechanisms of
Penicillium Oxalicum." *International Biodeterioration & Biodegradation* 122
(August):141–50. https://doi.org/10.1016/j.ibiod.2017.05.015.

Arnold, A. Elizabeth, and F. Lutzoni. 2007. "Diversity and Host Range of Foliar Fungal
Endophytes: Are Tropical Leaves Biodiversity Hotspots?" *Ecology* 88 (3): 541–49.
https://doi.org/10.1890/05-1459.

Bárcenas-Moreno, Gema, and Erland Bååth. 2009. "Bacterial and Fungal Growth in Soil Heated
at Different Temperatures to Simulate a Range of Fire Intensities." *Soil Biology and
Biochemistry* 41 (12): 2517–26. https://doi.org/10.1016/j.soilbio.2009.09.010.

Bárcenas-Moreno, Gema, Fuensanta García-Orenes, Jorge Mataix-Solera, Jorge Mataix-Beneyto,
and Erland Bååth. 2011. "Soil Microbial Recolonisation after a Fire in a Mediterranean
Forest." *Biology and Fertility of Soils* 47 (3): 261–72. https://doi.org/10.1007/s00374-
010-0532-2.

Ben-Jacob, Eshel, Inon Cohen, and David L. Gutnick. 1998. "COOPERATIVE
ORGANIZATION OF BACTERIAL COLONIES: From Genotype to Morphotype."
Annual Review of Microbiology 52 (Volume 52, 1998): 779–806.
https://doi.org/10.1146/annurev.micro.52.1.779.

Brown, Shawn, Allison Veach, Jonathan Horton, Emerald Ford, Ari Jumpponen, and Richard
Baird. 2019. "Context Dependent Fungal and Bacterial Soil Community Shifts in
Response to Recent Wildfires in the Southern Appalachian Mountains." *Forest Ecology
and Management*, August. https://doi.org/10.1016/j.foreco.2019.117520.

Bruns, Thomas D., Fogel ,Robert, and John W. and Taylor. 1990. "Amplification and Sequencing
of Dna from Fungal Herbarium Specimens." *Mycologia* 82 (2): 175–84.
https://doi.org/10.1080/00275514.1990.12025863.

Caiafa, Marcos V., Amelia R. Nelson, Thomas Borch, Holly K. Roth, Timothy S. Fegel, Charles
C. Rhoades, Michael J. Wilkins, and Sydney I. Glassman. 2023. "Distinct Fungal and
Bacterial Responses to Fire Severity and Soil Depth across a Ten-Year Wildfire
Chronosequence in Beetle-Killed Lodgepole Pine Forests." *Forest Ecology and
Management* 544 (September):121160. https://doi.org/10.1016/j.foreco.2023.121160.

Caktu, Kivilcim, and Emir Turkoglu. 2011. "Microbial Culture Collections: The Essential
Resources for Life." *Gazi University Journal of Science* 24 (2): 175–80.

Camenzind, Tessa, Carlos A. Aguilar-Trigueros, Stefan Hempel, Anika Lehmann, Milos Bielcik,
Diana R. Andrade-Linares, Joana Bergmann, et al. 2024. "Towards Establishing a Fungal
Economics Spectrum in Soil Saprobic Fungi." *Nature Communications* 15 (1): 3321.
https://doi.org/10.1038/s41467-024-47705-7.

Caon, Lucrezia, V. Ramón Vallejo, Coen J. Ritsema, and Violette Geissen. 2014. "Effects of Wildfire on Soil Nutrients in Mediterranean Ecosystems." *Earth-Science Reviews* 139 (December):47–58. https://doi.org/10.1016/j.earscirev.2014.09.001.

Carini, Paul. 2019. "A 'Cultural' Renaissance: Genomics Breathes New Life into an Old Craft." *mSystems* 4 (3). https://doi.org/10.1128/mSystems.00092-19.

Castaño, Carles, Javier Parladé, Joan Pera, Juan Martínez de Aragón, Josu G. Alday, and José Antonio Bonet. 2016. "Soil Drying Procedure Affects the DNA Quantification of Lactarius Vinosus but Does Not Change the Fungal Community Composition." *Mycorrhiza* 26 (8): 799–808. https://doi.org/10.1007/s00572-016-0714-3.

Černohlávková, Jitka, Jiří Jarkovský, Michala Nešporová, and Jakub Hofman. 2009. "Variability of Soil Microbial Properties: Effects of Sampling, Handling and Storage." *Ecotoxicology and Environmental Safety* 72 (8): 2102–8. https://doi.org/10.1016/j.ecoenv.2009.04.023.

Chase, Alexander B., Ulas Karaoz, Eoin L. Brodie, Zulema Gomez-Lunar, Adam C. Martiny, and Jennifer B. H. Martiny. 2017. "Microdiversity of an Abundant Terrestrial Bacterium Encompasses Extensive Variation in Ecologically Relevant Traits." *mBio* 8 (6): 10.1128/mbio.01809-17. https://doi.org/10.1128/mbio.01809-17.

Collado, Javier, Gonzalo Platas, Barbara Paulus, and Gerald F. Bills. 2007. "High-Throughput Culturing of Fungi from Plant Litter by a Dilution-to-Extinction Technique." *FEMS Microbiology Ecology* 60 (3): 521–33. https://doi.org/10.1111/j.1574-6941.2007.00294.x.

Cray, Jonathan A., Andrew N. W. Bell, Prashanth Bhaganna, Allen Y. Mswaka, David J. Timson, and John E. Hallsworth. 2013. "The Biology of Habitat Dominance; Can Microbes Behave as Weeds?" *Microbial Biotechnology* 6 (5): 453–92. https://doi.org/10.1111/1751-7915.12027.

Cui, Hang, Caihuan Wang, Zhenhong Gu, Honghui Zhu, Shenlei Fu, and Qing Yao. 2014. "Evaluation of Soil Storage Methods for Soil Microbial Community Using Genetic and Metabolic Fingerprintings." *European Journal of Soil Biology* 63 (July):55–63. https://doi.org/10.1016/j.ejsobi.2014.05.006.

Davis, Kathryn E. R., Shayne J. Joseph, and Peter H. Janssen. 2005. "Effects of Growth Medium, Inoculum Size, and Incubation Time on Culturability and Isolation of Soil Bacteria." *Applied and Environmental Microbiology* 71 (2): 826–34. https://doi.org/10.1128/AEM.71.2.826-834.2005.

Day, Nicola J., Kari E. Dunfield, Jill F. Johnstone, Michelle C. Mack, Merritt R. Turetsky, Xanthe J. Walker, Alison L. White, and Jennifer L. Baltzer. 2019. "Wildfire Severity Reduces Richness and Alters Composition of Soil Fungal Communities in Boreal Forests of Western Canada." *Global Change Biology*, April. https://doi.org/10.1111/gcb.14641.

Enright, Dylan J., Kerri M. Frangioso, Kazuo Isobe, David M. Rizzo, and Sydney I. Glassman. 2022. "Mega-Fire in Redwood Tanoak Forest Reduces Bacterial and Fungal Richness and Selects for Pyrophilous Taxa That Are Phylogenetically Conserved." *Molecular Ecology* 31 (8): 2475–93. https://doi.org/10.1111/mec.16399.

Falconer, Ruth E., Wilfred Otten, and Nia A. White. 2015. "Chapter One - Toward Modeling the Resistance and Resilience of 'Below-Ground' Fungal Communities: A Mechanistic and Trait-Based Approach." In *Advances in Applied Microbiology*, edited by Sima Sariaslani and Geoffrey Michael Gadd, 93:1–44. Academic Press. https://doi.org/10.1016/bs.aambs.2015.08.001.

Ferrenberg, Scott, Sean P. O'Neill, Joseph E. Knelman, Bryan Todd, Sam Duggan, Daniel Bradley, Taylor Robinson, et al. 2013. "Changes in Assembly Processes in Soil Bacterial

614   Communities Following a Wildfire Disturbance." *The ISME Journal* 7 (6): 1102–11.
615   https://doi.org/10.1038/ismej.2013.11.
616 Fischer, Monika S., Frances Grace Stark, Timothy D. Berry, Nayela Zeba, Thea Whitman, and
617   Matthew F. Traxler. 2021. "Pyrolyzed Substrates Induce Aromatic Compound
618   Metabolism in the Post-Fire Fungus, Pyronema Domesticum." *Frontiers in Microbiology*
619   12. https://www.frontiersin.org/article/10.3389/fmicb.2021.729289.
620 Fox, Sam, Benjamin A. Sikes, Shawn P. Brown, Cathy L. Cripps, Sydney I. Glassman, Karen
621   Hughes, Tatiana Semenova-Nelsen, and Ari Jumpponen. 2022. "Fire as a Driver of
622   Fungal Diversity — A Synthesis of Current Knowledge." *Mycologia* 114 (2): 215–41.
623   https://doi.org/10.1080/00275514.2021.2024422.
624 Franzosa, Eric A., Tiffany Hsu, Alexandra Sirota-Madi, Afrah Shafquat, Galeb Abu-Ali, Xochitl
625   C. Morgan, and Curtis Huttenhower. 2015. "Sequencing and beyond: Integrating
626   Molecular 'omics' for Microbial Community Profiling." *Nature Reviews Microbiology* 13
627   (6): 360–72. https://doi.org/10.1038/nrmicro3451.
628 González-Pérez, José A., Francisco J. González-Vila, Gonzalo Almendros, and Heike Knicker.
629   2004. "The Effect of Fire on Soil Organic Matter—a Review." *Environment International*
630   30 (6): 855–70. https://doi.org/10.1016/j.envint.2004.02.003.
631 Groot, William J. de, Michael D. Flannigan, and Alan S. Cantin. 2013. "Climate Change Impacts
632   on Future Boreal Fire Regimes." *Forest Ecology and Management*, The Mega-fire reality,
633   294 (April):35–44. https://doi.org/10.1016/j.foreco.2012.09.027.
634 Hantson, Stijn, Salvador Pueyo, and Emilio Chuvieco. 2015. "Global Fire Size Distribution Is
635   Driven by Human Impact and Climate." *Global Ecology and Biogeography* 24 (1): 77–
636   86. https://doi.org/10.1111/geb.12246.
637 Haridas, Sajeet, Colette Breuill, Joerg Bohlmann, and Tom Hsiang. 2011. "A Biologist's Guide to
638   *de Novo* Genome Assembly Using next-Generation Sequence Data: A Test with Fungal
639   Genomes." *Journal of Microbiological Methods* 86 (3): 368–75.
640   https://doi.org/10.1016/j.mimet.2011.06.019.
641 Hernandez, Damian J, Aaron S David, Eric S Menges, Christopher A Searcy, and Michelle E
642   Afkhami. 2021. "Environmental Stress Destabilizes Microbial Networks." *The ISME*
643   *Journal* 15 (6): 1722–34. https://doi.org/10.1038/s41396-020-00882-x.
644 Ho, Adrian, D. Paolo Di Lonardo, and Paul L. E. Bodelier. 2017. "Revisiting Life Strategy
645   Concepts in Environmental Microbial Ecology." *FEMS Microbiology Ecology* 93 (3):
646   fix006. https://doi.org/10.1093/femsec/fix006.
647 Hugenholtz, Philip. 2002. "Exploring Prokaryotic Diversity in the Genomic Era." *Genome*
648   *Biology* 3 (2): reviews0003.1. https://doi.org/10.1186/gb-2002-3-2-reviews0003.
649 Joint, Ian, Martin Mühling, and Joël Querellou. 2010. "Culturing Marine Bacteria – an Essential
650   Prerequisite for Biodiscovery." *Microbial Biotechnology* 3 (5): 564–75.
651   https://doi.org/10.1111/j.1751-7915.2010.00188.x.
652 Kämpfer, Peter. 2014. "Chapter 14 - Continuing Importance of the 'Phenotype' in the Genomic
653   Era." In *Methods in Microbiology*, edited by Michael Goodfellow, Iain Sutcliffe, and
654   Jongsik Chun, 41:307–20. New Approaches to Prokaryotic Systematics. Academic Press.
655   https://doi.org/10.1016/bs.mim.2014.07.005.
656 Knicker, Heike. 2011. "Pyrogenic Organic Matter in Soil: Its Origin and Occurrence, Its
657   Chemistry and Survival in Soil Environments." *Quaternary International*, Black Soils
658   and Black Sediments -- Archives of Landscape Evolution, 243 (2): 251–63.
659   https://doi.org/10.1016/j.quaint.2011.02.037.

Kuo, Hsiao-Che, Sun Hui, Jaeyoung Choi, Frederick O. Asiegbu, Jari P. T. Valkonen, and Yong-
Hwan Lee. 2014. "Secret Lifestyles of *Neurospora Crassa*." *Scientific Reports* 4
(May):5135. https://doi.org/10.1038/srep05135.
Leadbetter, Jared R. 2003. "Cultivation of Recalcitrant Microbes: Cells Are Alive, Well and
Revealing Their Secrets in the 21st Century Laboratory." *Current Opinion in*
Microbiology 6 (3): 274–81. https://doi.org/10.1016/S1369-5274(03)00041-9.
Lilleskov, Erik A., Thomas W. Kuyper, Martin I. Bidartondo, and Erik A. Hobbie. 2019.
"Atmospheric Nitrogen Deposition Impacts on the Structure and Function of Forest
Mycorrhizal Communities: A Review." *Environmental Pollution* 246 (March):148–62.
https://doi.org/10.1016/j.envpol.2018.11.074.
Lustenhouwer, Nicky, Daniel S. Maynard, Mark A. Bradford, Daniel L. Lindner, Brad Oberle,
Amy E. Zanne, and Thomas W. Crowther. 2020. "A Trait-Based Understanding of Wood
Decomposition by Fungi." *Proceedings of the National Academy of Sciences* 117 (21):
11551–58. https://doi.org/10.1073/pnas.1909166117.
Malik, Khursheed A., and Dieter Claus. 1987. "Bacterial Culture Collections: Their Importance
to Biotechnology and Microbiology." *Biotechnology and Genetic Engineering Reviews* 5
(1): 137–98. https://doi.org/10.1080/02648725.1987.10647837.
Malosso, Elaine, Ian S. Waite, Lorna English, David W. Hopkins, and Anthony G. O'Donnell.
2006. "Fungal Diversity in Maritime Antarctic Soils Determined Using a Combination of
Culture Isolation, Molecular Fingerprinting and Cloning Techniques." *Polar Biology* 29
(7): 552–61. https://doi.org/10.1007/s00300-005-0088-z.
Martiny, Adam C. 2019. "High Proportions of Bacteria Are Culturable across Major Biomes."
The ISME Journal 13 (8): 2125–28. https://doi.org/10.1038/s41396-019-0410-3.
Maynard, Daniel S., Mark A. Bradford, Kristofer R. Covey, Daniel Lindner, Jessie Glaeser,
Douglas A. Talbert, Paul Joshua Tinker, Donald M. Walker, and Thomas W. Crowther.
2019. "Consistent Trade-Offs in Fungal Trait Expression across Broad Spatial Scales."
Nature Microbiology 4 (5): 846–53. https://doi.org/10.1038/s41564-019-0361-5.
McGinnis, M. R., A. A. Padhye, and L. Ajello. 1974. "Storage of Stock Cultures of Filamentous
Fungi, Yeasts, and Some Aerobic Actinomycetes in Sterile Distilled Water." *Applied*
Microbiology 28 (2): 218–22. https://doi.org/10.1128/am.28.2.218-222.1974.
Michelotti, Lucas A., and Jessica R. Miesel. 2015. "Source Material and Concentration of
Wildfire-Produced Pyrogenic Carbon Influence Post-Fire Soil Nutrient Dynamics."
Forests 6 (4): 1325–42. https://doi.org/10.3390/f6041325.
Pérez-Valera, Eduardo, Miguel Verdú, Jose Antonio Navarro-Cano, and Marta Goberna. 2018.
"Resilience to Fire of Phylogenetic Diversity across Biological Domains." *Molecular*
Ecology 27 (13): 2896–2908. https://doi.org/10.1111/mec.14729.
Petersen, P. M. 1970. "Danish Fireplace Fungi-an Ecological Investigation on Fungi on Burns."
Dansk Botanisk Arkiv 27 (3): 1–97.
Pierson, Derek N., Peter R. Robichaud, Charles C. Rhoades, and Robert E. Brown. 2019. "Soil
Carbon and Nitrogen Eroded after Severe Wildfire and Erosion Mitigation Treatments."
International Journal of Wildland Fire 28 (10): 814–21.
https://doi.org/10.1071/WF18193.
Prakash, Om, Yogesh Shouche, Kamlesh Jangid, and Joel E. Kostka. 2013. "Microbial
Cultivation and the Role of Microbial Resource Centers in the Omics Era." *Applied*
Microbiology and Biotechnology 97 (1): 51–62. https://doi.org/10.1007/s00253-012-
4533-y.

Pressler, Yamina, John C. Moore, and M. Francesca Cotrufo. 2019. "Belowground Community
Responses to Fire: Meta-Analysis Reveals Contrasting Responses of Soil
Microorganisms and Mesofauna." *Oikos* 128 (3): 309–27.
https://doi.org/10.1111/oik.05738.
Preston, C. M., and M. W. I. Schmidt. 2006. "Black (Pyrogenic) Carbon in Boreal Forests: A
Synthesis of Current Knowledge and Uncertainties." *Biogeosciences Discussions* 3 (1):
211–71.
Pulido Barriga, M. Fabiola, James W. J. Randolph, and Sydney I. Glassman. 2024. "Not All Is
Lost: Resilience of Microbiome Samples to Freezer Failures and Long-Term Storage."
mSphere 10 (1): e00603-24. https://doi.org/10.1128/msphere.00603-24.
Pulido-Barriga, M. Fabiola, Amelia R. Nelson, Peter M. Homyak, Michael J. Wilkins, and
Sydney I. Glassman. 2025. "Chaparral Wildfire Shifts the Functional Potential for Soil
Pyrogenic Organic Matter and Nitrogen Cycling." bioRxiv.
https://doi.org/10.1101/2024.08.18.607989.
Pulido-Chavez, M. Fabiola, James W. J. Randolph, Cassandra Zalman, Loralee Larios, Peter M.
Homyak, and Sydney I. Glassman. 2023. "Rapid Bacterial and Fungal Successional
Dynamics in First Year after Chaparral Wildfire." *Molecular Ecology* 32 (7): 1685–1707.
https://doi.org/10.1111/mec.16835.
R Core Team. 2020. *R: A Language and Environment for Statistical Computing*. Vienna, Austria:
R Foundation for Statistical Computing. https://www.R-project.org/.
Reasoner, D J, and E E Geldreich. 1985. "A New Medium for the Enumeration and Subculture of
Bacteria from Potable Water." *Applied and Environmental Microbiology* 49 (1): 1–7.
https://doi.org/10.1128/aem.49.1.1-7.1985.
Reyes-César, Anaisell, Ángel E. Absalón, Francisco J. Fernández, Juan Manuel González, and
Diana V. Cortés-Espinosa. 2014. "Biodegradation of a Mixture of PAHs by Non-
Ligninolytic Fungal Strains Isolated from Crude Oil-Contaminated Soil." *World Journal*
of Microbiology and Biotechnology 30 (3): 999–1009. https://doi.org/10.1007/s11274-
013-1518-7.
Rubin, Benjamin E. R., Sean M. Gibbons, Suzanne Kennedy, Jarrad Hampton-Marcell, Sarah
Owens, and Jack A. Gilbert. 2013. "Investigating the Impact of Storage Conditions on
Microbial Community Composition in Soil Samples." *PLOS ONE* 8 (7): e70460.
https://doi.org/10.1371/journal.pone.0070460.
Sánchez-Clemente, Rubén, María Isabel Igeño, Ana G. Población, María Isabel Guijo, Faustino
Merchán, and Rafael Blasco. 2018. "Study of pH Changes in Media during Bacterial
Growth of Several Environmental Strains." *Proceedings* 2 (20): 1297.
https://doi.org/10.3390/proceedings2201297.
Seaver, Fred J., and Ernest D. Clark. 1910. "Studies in Pyrophilous Fungi: II. Changes Brought
about by the Heating of Soils and Their Relation to the Growth of Pyronema and Other
Fungi." *Mycologia* 2 (3): 109–24. https://doi.org/10.2307/3753620.
Sousa, Ana Margarida, Idalina Machado, Ana Nicolau, and Maria Olívia Pereira. 2013.
"Improvements on Colony Morphology Identification towards Bacterial Profiling."
Journal of Microbiological Methods 95 (3): 327–35.
https://doi.org/10.1016/j.mimet.2013.09.020.
Stackebrandt, Erko, David Smith, Serge Casaregola, Giovanna Cristina Varese, Gerard Verkleij,
Nelson Lima, and Paul Bridge. 2014. "Deposit of Microbial Strains in Public Service

Collections as Part of the Publication Process to Underpin Good Practice in Science."
SpringerPlus 3 (1): 208. https://doi.org/10.1186/2193-1801-3-208.
Stevenson, Bradley S., Stephanie A. Eichorst, John T. Wertz, Thomas M. Schmidt, and John A.
Breznak. 2004. "New Strategies for Cultivation and Detection of Previously Uncultured
Microbes." *Applied and Environmental Microbiology* 70 (8): 4748–55.
https://doi.org/10.1128/AEM.70.8.4748-4755.2004.
Swenson, Tami L., Ulas Karaoz, Joel M. Swenson, Benjamin P. Bowen, and Trent R. Northen.
2018. "Linking Soil Biology and Chemistry in Biological Soil Crust Using Isolate
Exometabolomics." *Nature Communications* 9 (1): 19. https://doi.org/10.1038/s41467-
017-02356-9.
Vieira, F. C. S., and E. Nahas. 2005. "Comparison of Microbial Numbers in Soils by Using
Various Culture Media and Temperatures." *Microbiological Research* 160 (2): 197–202.
https://doi.org/10.1016/j.micres.2005.01.004.
Wan, Shiqiang, Dafeng Hui, and Yiqi Luo. 2001. "Fire Effects on Nitrogen Pools and Dynamics
in Terrestrial Ecosystems: A Meta-Analysis." *Ecological Applications* 11 (5): 1349–65.
https://doi.org/10.1890/1051-0761(2001)011[1349:FEONPA]2.0.CO;2.
Watrud, Lidia S., Kendall Martin, Kelly K. Donegan, Jeffrey K. Stone, and Clarace G. Coleman.
2006. "Comparison of Taxonomic, Colony Morphotype and PCR-RFLP Methods to
Characterize Microfungal Diversity." *Mycologia* 98 (3): 384–92.
https://doi.org/10.1080/15572536.2006.11832673.
Whitman, Thea, Ellen Whitman, Jamie Woolet, Mike D. Flannigan, Dan K. Thompson, and
Marc-André Parisien. 2019. "Soil Bacterial and Fungal Response to Wildfires in the
Canadian Boreal Forest across a Burn Severity Gradient." *Soil Biology and Biochemistry*
138 (November):107571. https://doi.org/10.1016/j.soilbio.2019.107571.
Whitman, Thea, Jamie Woolet, Miranda Sikora, Dana B. Johnson, and Ellen Whitman. 2022.
"Resilience in Soil Bacterial Communities of the Boreal Forest from One to Five Years
after Wildfire across a Severity Gradient." *Soil Biology and Biochemistry* 172
(September):108755. https://doi.org/10.1016/j.soilbio.2022.108755.
Wickham, Hadley, Winston Chang, Lionel Henry, Thomas Lin Pedersen, Kohske Takahashi,
Claus Wilke, Kara Woo, Hiroaki Yutani, and RStudio. 2019. "Ggplot2: Create Elegant
Data Visualisations Using the Grammar of Graphics." https://CRAN.R-
project.org/package=ggplot2.
Woolet, Jamie, and Thea Whitman. 2020. "Pyrogenic Organic Matter Effects on Soil Bacterial
Community Composition." *Soil Biology and Biochemistry* 141 (February):107678.
https://doi.org/10.1016/j.soilbio.2019.107678.
Yang, Seung-Jo, Ilnam Kang, and Jang-Cheon Cho. 2016. "Expansion of Cultured Bacterial
Diversity by Large-Scale Dilution-to-Extinction Culturing from a Single Seawater
Sample." *Microbial Ecology* 71 (1): 29–43. https://doi.org/10.1007/s00248-015-0695-3.
Zeba, Nayela, Timothy D. Berry, Monika S. Fischer, Matthew F. Traxler, and Thea Whitman.
2024. "Soil Carbon Mineralization and Microbial Community Dynamics in Response to
Pyrogenic Organic Matter Addition." *Soil Biology and Biochemistry* 191 (April):109328.
https://doi.org/10.1016/j.soilbio.2024.109328.
793
794
795

Figures:

Figure 1:

Figure 1: A conceptual diagram of the experimental design used in this experiment. "LB" is
abbreviated for Lysogeny Broth, "MYA" is Malt Yeast Agar, "R2A" is Reasoner's 2 Agar, and
"PyOM" is Pyrogenic Organic Matter. Note that the cultures obtained from fresh soil plated onto
rich media also serve as the "Fresh" storage condition for the soil storage experiment. Green
circles and orange stars symbolize two different morphotypes collected from the same initial
plate.

Figure 2:

Figure 2: A snapshot of the morphological diversity of the cultures obtained from the El Dorado
Fire across all soil storage and media type experiments. Identities were obtained via Sanger
sequencing of the 16S or ITS regions for bacteria and fungi, respectively. Moving left to right
across each row, the isolates shown are A)*Glutamicibacter bergerei, Arthrobacter agilis,*
Streptomyces mauvecolor, Kocuria rosea, B) *Arthrobacter bussei, Holtermaniella festucosa,*
Pedobacter panaciterrae, Curtobacterium oceanosedimentum, C) *Penicillium radiatolobatum,*
Penicillium fagi, Penicillium glabrum, Penicillium thomii, D) *Penicillium chalabudae,*
Penicillium radiatolobatum, Penicillium murcianum, Penicillium adametzii.

Figure 3:

Figure 3: Calculated capturable CFUs per gram soil for A) bacteria and B) fungi among the
tested soil storage methods and C) bacteria and D) fungi across tested media types. Points and
bars are colored by treatment with bars indicating standard error of themean and black points
indicating the mean CFUs per gram soil for each treatment. Significance was tested using
ANOVA and Tukey Post-Hoc lettering has been applied. LB = Lysogeny Broth; PyOM =
Pyrogenic Organic Matter; R2A = Reasoner's 2 Agar, and G stands for glycerol.

Figure 4:

Figure 4: Summarized counts of A) total bacterial, B) pyrophilous bacterial, C) total fungal, and
D) pyrophilous fungal genera (green) and species (grey) across four soil storage methods
compared to fresh soil when isolated on LB for bacteria or MYA for fungi. Treatments are
ordered by highest amount of genera first, then species count second. G = glycerol.

Figure 5:

Figure 5: Summarized counts of A) total bacterial, B) pyrophilous bacterial, C) total fungal, and
D) pyrophilous fungal genera (green) and species (grey) obtained from fresh soil tested across
three media types. Treatments are ordered by highest number of genera first, then species count
second. LB = Lysogeny Broth, MYA = Malt Yeast Agar, R2A = Reasoner's 2 Agar, PyOM =
Pyrogenic Organic Matter.

Figure 6:

Figure 6: Heatmap-style visualization showing the diversity of pyrophilous A) bacterial and B)
fungal genera across the soil storage experiment. A filled blue square indicates the presence of
the genus on the x axis in the soil storage method indicated on the y axis. G = glycerol. Genera
are ordered in alphabetical order, other than [Brevibacterium] which is in brackets due to
controversy over its taxonomy.

Figure 7:

Figure 7: Heatmap-style visualization showing the diversity of pyrophilous A) bacterial and B)
fungal genera across the media types experiment. A filled blue square indicates the presence of
the genus on the x axis in the media indicated on the y axis. LB = Lysogeny Broth, MYA = Malt
Yeast Agar, R2A = Reasoner's 2 Agar, PyOM = Pyrogenic Organic Matter. Genera are ordered in
alphabetical order, other than [Brevibacterium] which is in brackets due to controversy over its
taxonomy.