Evaluating Best Practices for Isolating Pyrophilous Bacteria and Fungi from Burned Soil

Dylan J. Enright[1*], Ryan Quaal[1], Aishwarya Verabeehu[1], Anna Nguyen[1], Jenna Maddox[2], and
Sydney I. Glassman[1*]

Affiliations:

[1]University of California Riverside, Department of Microbiology and Plant Pathology, 900
University Ave. Riverside CA, 92521, USA

[2]California State University San Bernardino, Department of Biology, 5500 University Pkwy.,
San Bernardino CA, 92407, USA

Corresponding authors: sydneyg@ucr.edu and denri009@ucr.edu

Abstract

A live microbial culture is invaluable to assess traits and functions via 'omics and
biophysical assays. However, it is not always logistically feasible to culture immediately from
freshly obtained soil, and selecting the proper media for culturing is not trivial. While building a
culture collection of pyrophilous microbes obtained from burnt soil, we tested the best 1) method
of storing soil to retain culturable viability and 2) media to garner the most microbial diversity.
We tested four methods of soil storage (dried, stored at 4ºC, stored at -80ºC alone or in glycerol)
and compared to fresh soil obtained 6 months after a severe California chaparral shrubland
wildfire. For bacteria, soil frozen at -80ºC with glycerol preserved the greatest diversity (25
species, 13 genera) compared to fresh soil (26 species, 13 genera). For fungi, soil stored at -80ºC
alone preserved the greatest diversity (10 species, 3 genera) compared to fresh soil (13 species, 7
genera). We also tested 3 media types: rich media (Lysogeny Broth (LB) for bacteria; Malt Yeast
Agar (MYA) for fungi), oligotrophic media (Reasoner's 2 Agar (R2A) and media made from
pyrogenic organic matter (PyOM). For bacteria, culturing on LB and R2A garnered the greatest
diversity (LB = 26 species, 13 genera, R2A = 27 species, 15 genera). For fungi a combination of
R2A and PyOM captured the greatest diversity (R2A = 15 species, 8 genera, PyOM = 12 species,
6 genera). For both bacteria and fungi, some species of interest were only captured using the
PyOM media. Using a combination of these methods from 2018-2022, we cultured >500 isolates
(286 bacteria; 258 fungi) from burned soils of 7 Southern California wildfires.

Introduction

In microbiological research, few resources are as valuable as live microbial cultures. Though advances in the fields of molecular biology and DNA sequencing have freed microbiologists from the need to work with culturable species exclusively (Franzosa et al. 2015), microbial culture collections remain an invaluable resource. Pure cultures are essential components for high quality whole-genome assembly (Hugenholtz 2002; Haridas et al. 2011), genetic engineering experiments (Malik and Claus 1987), and the phenotypic profiling of microorganisms (Kämpfer 2014; Alster et al. 2021). Additionally, isolated microorganisms greatly improve the interpretation of metabolomics data (Swenson et al. 2018) and possession of a diverse culture collection can unlock a plethora of applied microbiological research directions (Malik and Claus 1987; Reyes-César et al. 2014; Carini 2019). Once isolated and properly stocked, microbial culture collections become a permanent resource for future microbiological research (Caktu and Turkoglu 2011; Stackebrandt et al. 2014; Arnold and Lutzoni 2007). This in turn creates a veritable breeding ground for collaborative research. However, the hurdles of creating microbial culture collections are well documented and cannot be understated (Caktu and Turkoglu 2011).

The various factors that can be modified when attempting to culture microorganisms are as varied as there are microbes to isolate. For example, media selection may require grappling with an endless number of variables including consideration of the proper carbon sources, nitrogen and phosphorus content, salt concentrations, pH, and a host of other micronutrient considerations (Carini 2019; Davis, Joseph, and Janssen 2005; Stevenson et al. 2004). One must also consider whether to make the media nutrient rich or poor (oligotrophic), incubation temperature, water and oxygen availability, whether to alternate light and dark cycles, and the length of time to incubate samples (Davis, Joseph, and Janssen 2005; Stevenson et al. 2004; Leadbetter 2003). All this complexity contributes to so many microorganisms remaining uncultured (Prakash et al. 2013). And though recent advances in our understanding of microbiology have begun to make the adage of "99% of microbes are unculturable" outdated (Carini 2019; Stevenson et al. 2004; Leadbetter 2003; Martiny 2019), the fact remains that many microorganisms currently still lie beyond our reach. Bearing all this in mind, when we set out to create a culture collection of pyrophilous (or "fire-loving") microbes we knew we would face numerous technological and logistical challenges.

In addition to the well-known hurdles of creating a microbial culture collection, attempting to culture microbes associated with wildfires presents its own set of challenges. The first set of challenges are the logistic challenges associated with working with post-fire environmental samples. Wildfires are unpredictable natural phenomena. As such, it may not always be possible to have the infrastructure for large-scale culturing efforts (personnel available, media selected and created, broth tubes or petri dishes ordered, culturing conditions decided and established) in place when a wildfire burn scar is ready to be sampled. As part of this logistical navigation, often times post-fire soil samples may have to be stored for varying lengths of time before they can be used. This is a logistical problem shared among many environmental culturing efforts ranging from culturing of marine microbes (Joint, Mühling, and

Querellou 2010) to culturing from Antarctic soils (Malosso et al. 2006). However, the viability of
culturable microorganisms may be reduced depending on the soil storage method
(Černohlávková et al. 2009; Rubin et al. 2013). In addition to the logistics, wildfires radically
alter environmental nutrient landscapes (Caon et al. 2014; Pierson et al. 2019). Wildfire pyrolysis
turns labile, easily accessible carbon sources into difficult to degrade pyrogenic organic matter
(PyOM) (Michelotti and Miesel 2015; Preston and Schmidt 2006). This PyOM is often enriched
in polycyclic aromatic hydrocarbons (PAHs) that may require specialized metabolic pathways to
process (Reyes-César et al. 2014; Woolet and Whitman 2020; Aranda et al. 2017; Fischer et al.
2021). This can lead to accumulated char persisting in the environment for up to and exceeding
100 years (Knicker 2011). Additionally, wildfires can deposit an enormous amount of inorganic
nitrogen into the environment as organic nitrogen previously locked in plant biomass becomes
released during burning (Caon et al. 2014; Pulido-Chavez et al. 2022; González-Pérez et al.
2004; Wan, Hui, and Luo 2001). This high-nitrogen environment can radically alter the nutrient
strategies that microorganisms may employ post-fire and may limit the microbes capable of
surviving. For example, many species of fungi are known to be nitrophobic (Lilleskov et al.
2019). These radical nutrient shifts may result in commonly used laboratory media not providing
appropriate nutrient conditions to cultivate post-fire microbes.

Here, our goal was to create a large culture collection of pyrophilous microbes to use to
test their traits and determine how they survived or thrived post-fire via biophysical assays and
genomics. Keeping in mind the above described logistic and nutrient-related challenges, we used
soil from a local high severity chaparral wildfire to test Q1) the best methods to store post-fire
soil to retain pyrophilous microbial culturable viability and Q2) which laboratory media best
captured the diversity of pyrophilous microbes. For soil storage we tested soils stored for 3
months in a variety of ways (air-dried, refrigerated at 4°C, and frozen at -80°C with and without
glycerol) versus cultured immediately from the freshly obtained soil samples (Figure 1). For
media type we tested rich media (lysogeny broth (LB) agar for bacteria, malt-yeast agar (MYA)
for fungi), an oligotrophic media (Reasoner's 2 Agar (R2A)), or a media created in-house from
PyOM to test the effect of media selection on the culturability of pyrophilous bacteria and fungi
(Figure 1). We tested the impact of our treatments on abundance (CFUs: Colony Forming Units)
and microbial richness and composition (assessed via Sanger sequencing of 16S for bacteria or
ITS for fungi). We hypothesized (H1) that air-dried soil would retain the greatest diversity of
pyrophilous bacteria and fungi compared to fresh soil since post-fire soils already tend to be
fairly dry in chaparral environments (Pulido-Chavez et al. 2023; Stoof, Wesseling, and Ritsema
2010) thus leaving the soils to air dry would be the least dramatic disturbance for the soil
microorganisms contained within the samples. We further hypothesized (H2) that a combination
of the rich and PyOM media would obtain the greatest diversity of pyrophilous microbes as the
rich media would be able to grow a wide range of bacteria and fungi and the PyOM media would
capture those pyrophilous species with unique metabolic requirements specific to post-fire burn
scars. Although we specifically sought to create a pyrophilous culture collection, the insights
gleaned from this study not only showcase potential strategies for culturing fastidious
pyrophilous microorganisms but can also aid environmental microbiologists in decision making
about storage and media selection for all types of environmental culturing. The culmination of
this research was the creation of a large and diverse collection of pyrophilous microorganisms of

>500 isolates (286 bacteria; 258 fungi) from burned soils, mushrooms, and smoke of 7 Southern
California wildfires.

Methods

Site Description
To test the best soil storage method and culture media to cultivate pyrophilous microbes, we
leveraged a local high-severity wildfire of fire-adapted chaparral shrublands for sampling. The El
Dorado wildfire burned ~92km^2 Manzanita (*Arctostaphylos)* dominated chaparral shrublands in
the area surrounding Yucaipa, CA from September through November of 2020. We examined
high-resolution successional data of bacterial and fungal communities after a similar nearby
high-severity Manzanita dominated chaparral wildfire (Pulido-Chavez et al. 2023), and
compared to other post-fire studies that had successional data (Bárcenas-Moreno et al. 2011;
Caiafa et al. 2023), to identify 6 months post-fire the ideal time to attempt to isolate a wide range
of pyrophilous bacteria and fungi. Therefore, we collected soil samples from high-severity zones
6 months after the El Dorado fire in June 2021. Sites of the fire were characterized as having 50
to 75 percent slopes with soil characteristics described as dry, sandy soils in the Springdale
family-Lithic Xerorthents association.

Soil Collection and Storage

Approximately 250mL of soil from 0-10 cm below the ash layer of burned soil was collected
from 9 plots with ethanol cleaned releasable bulb planters and placed into whirl-pak bags 6
months post-fire following the same methods as used in a similar ecosystem (Pulido-Chavez et
al. 2023). After collection, soil was 2mm sieved and either used as fresh inoculum for cultivation
or stored in four different ways: air dried, refrigerated (4ºC), frozen (-80ºC), or suspended 50:50
soil with glycerol to create a soil slurry and then frozen (-80ºC) (Figure 1).

143
144 Culturing Medias Used
145
Fresh soil samples were cultured immediately onto rich media (lysogeny broth (LB) agar for
bacteria, malt-yeast agar (MYA) for fungi), an oligotrophic media (Reasoner's 2 Agar (R2A)), or
a media created in-house from PyOM to test the effect of media selection on the culturability of
pyrophilous bacteria and fungi. After 3 months of storage, stored soil samples were cultured onto
rich media. This means that the cultures obtained from freshly obtained soil samples on rich
media served as both the "rich media" condition for the media experiment and the "fresh"
comparison for the soil storage experiment (Figure 1).

153
LB agar and broth were produced using powdered LB Broth, Miller (Fisher Scientific, CAS#
73049-73-7). When making solid LB agar, 15g per liter of bacteriological grade agar (Apex
Chemicals, Cat # 20-273) was added to DI water. For each liter of Malt-Yeast Agar (MYA) and
broth 5g of powdered Malt Extract (Research Products International, Genessee, CAS# 8002-48-
0) and 5g of powdered Yeast Extract (Sigma-Aldrich, Cat # 70161) was used. For solid agar

media, 15g per liter of bacteriological grade agar (Apex Chemicals, Cat # 20-273) was added.
R2A agar was produced using powdered R2A agar (Criterion, Cat # 16721) following
manufacturer directions.
162
Pyrogenic Organic Matter (PyOM) Media: We developed an in-house recipe of PyOM agar
media using biochar harvested from the El Dorado Fire burn sites. Biochar was ground into a
fine powder prior to use with a coffee grinder (Chefman, Model: RJ44-OPP-RED). In a 2L
Erlenmeyer flask, 50g of ground biochar was added followed by 1200mL of DI water. The
mouth of the flask was covered and left on a stirring hotplate for 24 hours at 40°C after which it
was removed from heat. The solution was then allowed to settle for 8 hours. Then, the liquid
contents of the flask were carefully siphoned into a clean Erlenmeyer flask, being very careful to
minimally disturb the sediment at the bottom (Figure S1). The liquid was then passed through
cheesecloth, and then twice through a coffee filter to remove any remaining sediment before
being transferred to clean containers for autoclaving. For each liter of PyOM media 13g of
bacteriological grade agar (Apex Chemicals, Cat # 20-273) was added.
174
All media types (LB, MYA, and R2A, after mixing with DI water per manufacturer directions,
and home-made PyOM media) were autoclaved at 121ºC for 45 minutes.

Antibiotics: For media used to isolate fungi and intended to resist bacterial growth, filter
sterilized gentamycin and ampicillin (Hardy Diagnostics, Santa Maria, Ca), were added to the
molten agar after autoclaving, once it had cooled to 60ºC, for a final concentration of 10mg/mL
each. For media used to isolate bacteria and intended to resist fungal growth, cycloheximide
(Research Products International, CAS # 66-81-9) was added to the molten agar, once it had
cooled to 60ºC for a final concentration of 20mg/L.

183
184 Microbial Isolation Method
185
Culturing from Soil: Soil suspension inoculum was prepared by making a slurry of 1g of soil and
5mL of autoclaved deionized water. A dilution series was made from the soil slurry down to
concentrations between 1:10 to 1:1000. This inoculum was spread onto LB, MYA, R2A, or
PyOM media supplemented with cycloheximide if culturing bacteria or ampicillin and
gentamicin if culturing fungi. These plates were then parafilmed and allowed to grow for a week.
Fungi and bacteria were then morphotyped using size, color, and growth rate. Bacteria were
isolated by picking a well isolated colony using a sterile 10µL pipet tip and then streaking for
isolation onto a fresh LB plate without antifungals. Fungi were isolated by using a flame-
sterilized scalpel to cut a 1 cm^2 square of the growing edge of a fungus out of the agar plate. This
square was then transferred to a fresh MYA plate without antibacterials and placed growing-side
down into the center of the plate. In cases where microbes would not establish onto LB or MYA,
they were maintained on the media they originated from (either R2A or PyOM). Plates were
allowed to incubate at room temperature for 1 week before re-isolating following the same
procedure. This was repeated until three clean isolations were obtained prior to DNA extraction
and stocking.
201
202 Microbial Stocking Method
203

Bacteria: Once a thrice purified isolate was obtained and DNA was extracted, each bacterial
isolate was stocked for future use. A colony was picked aseptically from the last isolation streak
plate using a sterile 10µL pipet tip and placed into 3mL of LB broth in a sterile 15mL Falcon
tube. This was then grown under shaking incubation at 180rpm for 24-72hrs until the liquid
culture became turbid. From this liquid culture, 1mL of culture suspension was added to a 2mL
cryovial and combined with 1mL of sterile 40% glycerol, resulting in a final cell suspension that
was 20% glycerol. This was vortexed briefly and then frozen at -80ºC. For each isolate, 2 stock
tubes were made.
212
Fungi: Once a thrice purified isolate was obtained and DNA was extracted, each fungal isolate
was stocked for future use. On a 100mm agar plate, a 1cm^2 cube of fungus was allowed to grow
until it came within approximately 1cm of the edge of the Petri dish. Then, using a sterile
transfer tube, 5mm disks were punched from the growing edge of the fungus. These disks were
then transferred to a 2mL screw-cap tube containing 1mL of autoclaved tap water. For each stock
tube 8-10 disks were obtained, and for each isolate 2 stock tubes were made. These tubes with
then stored at room temperature for future use.
220
Soil Storage Method Experimental Design
For each of 9 soil samples collected, we divided soils from each subsample into either fresh vs
stored for 3 months air-dried, frozen, or frozen in glycerol, then 1g was combined with 5mL of
sterile DI water in 4 separate 50mL falcon tubes to create four initial solutions for each of the
soil samples collected. The initial solutions were then used to create a dilution series ranging
from 1:10 to 1:10,000 for each storage method. 100µL of each solution in the dilution series was
then plated on MYA plates with ampicillin and gentamicin and LB plates with cycloheximide.
The plates were then wrapped in parafilm and placed in a drawer to be kept at room temperature
for approximately one week. Cell counts and unique morphology counts were then done on the
plates. Plates containing between 30-500 colonies were considered "countable" plates and the
initial CFU concentrations were calculated using the number of colonies obtained multiplied by
the dilution factor used. This led to each treatment having between 6 and 9 countable replicate
plates. Morphotypes were determined using size, color, and growth rate. We then isolated,
extracted DNA, and stocked each bacterial and fungal unique morphology (Figure 1).
235
Effect of Media Type Experimental Design
The fresh soil suspension inoculum was prepared by making a slurry of 1 gram of soil and 5mL
of deionized water. The soil suspension was then diluted down to concentrations between 1:10 to
1:10,000. This inoculum was spread onto MYA media with ampicillin and gentamicin, LB media
with cycloheximide, R2A media (with antibacterials or cycloheximide) and PyOM media (with
antibacterials or cycloheximide). The plates were allowed to grow for a week after inoculation
before being morphotyped and CFUs were counted as described in the previous section. Fungi
and bacteria were morphotyped using size, color, and growth rate. Individual morphotypes were
then isolated (Figure 1).
245
246 DNA Extraction & PCR
247
DNA Extraction: DNA was extracted from all unique isolates using 10µL of home-made Extract-
N-Amp Extraction Solution (Tris stock, EDTA, KCl, NaOH, DI H$_2$0; full recipe in supplements).

For bacteria, one well-isolated colony was picked using a sterile 10µL pipet tip, and for fungi, a
1mm^2 area of the growing edge of the fungus was scraped with a flame-sterilized probe. The tip
or probe was then submerged and swirled in the Extraction solution. After the Extraction
Solution was inoculated, the samples were heated at 65ºC for 10 minutes followed by 95ºC for
10 minutes. Once finished, 10µL of a Dilution Solution (3% BSA) was added and the samples
were left to chill overnight at 4ºC. Samples were either frozen at -20ºC or immediately amplified.
256
PCR Amplification of rRNA genes and Sanger sequencing: The bacterial 16S rRNA region was
amplified using a PA forward and PH reverse primer (Edwards et al., 1989) and the fungal ITS
region was amplified with the ITS1F forward (Gardes & Bruns, 1993) and ITS4 reverse primer
(White et al., 1990). Thermocycler conditions were as follows: 94℃ for 2 minutes, then 35
cycles of 57℃ (for fungi) or 62℃ (for bacteria) for 1 minute, 72℃ for 1 minute, then 94℃ for 1
minute, ending with a final extension of 72℃ for 8 minutes. The DNA was cleaned up using 3µL
of diluted Exosap mastermix (Applied Biosystems, Waltham MA) per 7µL PCR product before
submission for diluting and preparing for Sanger sequencing. Samples were Sanger sequenced in
the forward direction using either Genewiz (San Diego, CA), Eurofins Genomics (Louisville,
KY), or the UCR Institute for Integrative Genome Biology Genomics Core services.
267
BLAST Identification: Raw sequences were returned as .ab1 chromatogram files. Sequences were
then trimmed in either SnapGene or Geneious Prime to remove low quality regions and were
saved as FASTA files. Taxa were identified using NCBI BLASTn to create hit tables of top 10
hits. IDs were assigned using a combination of query cover, error value, percent identity, and a
consensus of hits which obtained the same quality scores. In general, an isolate was identified at
species level if the query cover was 100%, max e score was < 0.01, and percent identity was
>98%. If percent identity fell to ~95% then isolates were identified to genus, at 90% isolates
were left at family, at 85-90% isolates were identified to class, and at < 85% isolates were only
identified to phylum level.
277
Data Analysis
All statistical analyses and figures were produced in R 4.0.2 (R Core Team 2020) and all scripts
are available on GitHub. Normality was tested with a Shapiro test, then ANOVA was used to test
the effect of treatment (whether soil storage method or media selection) on bacterial and fungal
CFU counts, followed by a post-hoc Tukey HSD test. Heatmap-like visualization of genera and
species level diversity were created using the "geom_tile" function in ggplot2 (Wickham et al.
2019). All sequences have been deposited to NCBI GenBank.
285
In addition to the test of the best storage methods and media selection for isolating pyrophilous
microbes, here we present our culture collection of pyrophilous microbes. For a table of all
isolates, the fires from which they were sampled, time of collection, location data, and substrate
from which species were isolated please see Supplementary File 2.

Results

Microbes Obtained
From the 2020 El Dorado Fire 6-month timepoint soils we obtained across all experimental
methods a total of 170 bacterial isolates representing 4 phyla, 10 orders, 31 genera and 83

species. We also obtained 160 fungal isolates representing 3 phyla, 9 orders, 15 genera and 35
species. These isolates displayed a wide variety of morphologies (Figure 2) and all isolates are
listed in Table S1. These were the isolates used for further analysis in this study. Of these
isolates, 138/170 bacterial isolates composed of 17 genera and 71 species and 152/160 fungal
isolates composed of 8 genera and 30 species had literature support for inclusion as potential or
confirmed pyrophilous species (Enright et al. 2021; Pulido-Chavez et al. 2023; Woolet and
Whitman 2020; Aranda et al. 2017; Caiafa et al. 2023; Seaver and Clark 1910; Petersen 1970;
Whitman et al. 2019; Brown et al. 2019; Bárcenas-Moreno and Bååth 2009; Ferrenberg et al.
2013). Pyrophilous isolates are listed in Table S2.

<u>Effect of Storage Method Selection on CFUs</u>
Soil storage method significantly impacted the number of CFUs obtained per gram soil for both
bacteria (ANOVA: $F_{1,4} = 19.97$, $p < 0.001$; Figure 3A) and fungi (ANOVA: $F_{1,4} = 19.78$, $p <$
0.001; Figure 3B). For both bacteria and fungi, soil stored air-dried (Bacteria: 2.31×10^5
CFUs/gram, Fungi: 1.33×10^4 CFUs/gram) and stored at -80ºC with glycerol (Bacteria: 1.31×10^5
CFUs/gram, Fungi: 6.69×10^3 CFUs/gram) retained the fewest CFUs/gram soil. For bacteria, soils
stored at 4ºC (6.98×10^5 CFUs/gram) or at -80ºC without glycerol (6.39×10^5 CFUs/gram) were
not significantly different from fresh soils (5.73×10^5 CFUs). For fungi, fresh soils had the highest
average CFUs/gram soil (3.07×10^4 CFUs/gram), followed by soil frozen at -80ºC without
glycerol (2.66×10^4 CFUs/gram), and then stored at 4ºC (2.16×10^4 CFUs/gram).

<u>Effect of Media Selection on CFUs</u>
Choice of growth media significantly impacted the number of CFUs obtained per gram soil for
both bacteria (ANOVA: $F_{1,2} = 39.43$, $p < 0.001$; Figure 3C) and fungi ($F_{1,2} = 7.77$, $p < 0.01$;
Figure 3D). For both bacteria and fungi, PyOM media yielded the fewest CFUs per gram
(Bacteria: 1.66×10^5; Fungi: 1.03×10^4), and rich media (LB for bacteria and MYA for fungi) had
the highest average CFUs per gram (Bacteria: 5.73×10^5; Fungi: 3.07×10^4). Soil slurry plated on
oligotrophic R2A media had the second highest average CFU counts per gram for bacteria
(3.20×10^5). For fungi, CFU counts per gram on R2A (2.20×10^4) were statistically equivalent to
that on MYA.

<u>Effect of Storage Method Selection on Diversity</u>
Bacteria: Method of soil storage strongly affected the diversity of bacteria with fresh soil
retaining the highest diversity and dried soil retaining the lowest diversity (Figure 4A). In
assessing diversity, diversity among genera was given greater consideration than diversity among
species as more diverse genera would cover greater phylogenetic diversity than diversity within
species alone. For this reason, figures are ordered by number of genera first, then number of
species. For bacteria, in the soil storage experiment we obtained a total of 21 bacterial genera
(Figure S2A and S3) and 64 bacterial species (Figure S2A and S4). Fresh soil was the most
diverse, capturing 62% of these genera and 41% of these species. The second most diverse
treatment was frozen soil without glycerol which captured 62% of the genera and 30% of
species. This was followed closely by the frozen with glycerol treatment at 57% of genera and
39% of the species. Next was soil stored at 4ºC with 48% of genera and 22% of the species.
Lastly, the dried soil captured 38% of genera and 23% of the species. Only 3 genera
(*Arthrobacter, Curtobacterium, Streptomyces*) were isolated across all 5 storage treatments
whereas 8 genera (*Brevibacterium, Fontibacillus, Glutamicibacter, Massilia,*

Ornithinimicrobium, Paeniglutamicibacter, Pseudoarthobacter, and an unidentified Firmicute)
were isolated from a single treatment either fresh, refrigerated, frozen, or frozen with glycerol
(Figure S3). It is more difficult to discern additional patterns among the species level diversity
that are not additionally genus level patterns, but this is probably largely due to the fact that most
species (64%) were only observed once and only the species *Curtobacterium oceanosedimentum*
was observed in all treatments (Figure S4).
346
Considering only the isolates that have literature support for being pyrophilous, we isolated from
our soil storage experiment 13 bacterial genera and 51 bacterial species (Figures 4A, 5A; S5). In
this subset, frozen soil with no glycerol actually captured the largest share of the taxa with 85%
of the genera and 33% of the species. Second most diverse was the Fresh soil with 62% of
genera and 35% of species. This was followed closely by soil stored frozen with glycerol (54%
of genera, 39% of species) and soil that was refrigerated (54% of genera, 22% of species). The
least diverse bacterial communities were obtained from the dried soil which accounted for 38%
of genera and 24% of the species.
355
Fungi: For fungi, in total we obtained 6 genera (Figure S2B and S6) and 23 species (Figures S2B
and S7) from the soil storage experiment, of which we found literature support as pyrophilous for
3 genera (Figures 4B and 5B) and 20 species (Figure S8). Fresh soil was the most diverse,
capturing 100% of all 6 genera, whereas frozen and refrigerated soil both captured 50% of the
genera, followed by 33% in dried and soil frozen with glycerol performing the worst, capturing
only *Penicillium* (17% of the total) (Figure S6). For species, fresh soil captured 52%, followed
by 43% in frozen, and 30% in refrigerated and 26% in frozen with glycerol (Figure S7). For the
pyrophilous taxa, fresh, frozen, and refrigerated soil all captured 100% of the 3 fungal genera
whereas dried captured 2 (*Aspergillus, Penicillium*) and frozen with glycerol on captured 1
(*Pencillium*; Figure 5B). For pyrophilous species, frozen soil captured 50%, followed closely by
45% in fresh, 35% in refrigerated and 30% in air-dried (Figure S8).
367
368
369   Overall, for pyrophilous isolates, soil frozen without glycerol contained the most diverse
370   bacterial and fungal populations. The worst storage method for pyrophilous bacterial diversity
371   was air drying soil while freezing with glycerol was the worst storage method for pyrophilous
372   fungal diversity.
373
<u>Effect of Media Selection on Diversity</u>
Choice of growth media strongly affected both bacterial and fungal isolate diversity.
Bacteria: Choice of growth media strongly affected bacterial isolate diversity with R2A and LB
capturing the highest numbers of bacterial genera and species. For bacteria, we obtained a total
of 24 genera and 61 species across all media types (Figures S2C, S9 and S10). Both LB and R2A
obtained equally diverse communities, each capturing 58% of the total genera and 44% of the
total species. Isolates obtained from PyOM were the least diverse, capturing 33% of genera and
18% of species. However, when we subset for isolates with support for being pyrophilous, the
outcome is slightly different. For pyrophilous bacteria we obtained a total of 16 genera and 49
species (Figure 4C, 6A; S11). Among pyrophilous bacteria, R2A captured the greatest diversity
accounting for 69% of total genera and 49% of total species. Second most diverse was LB with
56% of total genera and 39% of total species. Lastly, PyOM obtained the lowest diversity of

pyrophilous bacterial isolates with 44% of total genera and 20% of total species. Notably, while
the PyOM media captured the least diverse bacterial community, the PyOM media was the only
media to capture certain taxa of interest such as *Noviherbaspirillum* (Figure 6A) (Whitman et al.
2019; Caiafa et al. 2023; Pulido-Chavez et al. 2023).
390
Fungi: We obtained a total of 14 genera and 29 species across all media types (Figures S2D,
S12, and S13). R2A media yielded the most diverse fungi, capturing 57% of total genera and
48% of total species. Both MYA and PyOM equally yielded 43% of total genera and 41% of
total species. Considering only pyrophilous fungi, we obtained 8 genera and 23 total species
across all media types (Figure 4D, 6B; S14). The most diverse isolate community was still
isolated from R2A, with R2A capturing 89% of total pyrophilous genera and 57% of the species.
Second most diverse among pyrophilous fungi was PyOM with 50% of total genera and 43% of
total species. Among pyrophilous fungi MYA was the least diverse, capturing 38% of the total
genera and 39% of the total species. Additionally, PyOM was the only media to capture
Neurospora, a well-documented post-fire fungus with relatively little knowledge about its
ecological role (Figure 6B) (Kuo et al. 2014).
402
403 For both bacteria and fungi R2A captured the most diverse isolate population both among total
404 isolates and pyrophilous isolates. Among pyrophilous isolates, PyOM media was the least
405 diverse for bacteria but captured a very interesting and important pyrophilous organism and
406 MYA was the least diverse for fungi.
407
408 **Discussion**
409
Throughout our experiments we managed to obtain a remarkable array of bacterial and
fungal morphotypes (Figure 2). We found that contrary to our hypothesis, freezing soil was the
best method of storing soil to retain the viability of culturable bacteria and fungi (Figures 4 & 5).
Additionally, we found that the best media for culturing pyrophilous microbes was the
oligotrophic R2A media (Figures 4 & 6). We did find that the novel solid agar media we created
from field-harvested PyOM did contribute significantly to the diversity of the fungal isolates and
was the only media to capture certain pyrophilous bacterial and fungal taxa of interest. The
efforts in these experiments has informed our continual culturing efforts and resulted in a
collection of over 500 post-fire microbial isolates (Supplementary File 2).
Here we conclude that from both a CFU and diversity-based perspective, freezing cells at
-80°C is the most effective soil storage method for retaining 85% of the culturable pyrophilous
bacterial genera (Figures 3A and 4A). It is possible that there could have been a slight amount of
heterogeneity in the microbial communities between soil samples (Jung and Kang 2014),
however we believe this to be fairly unlikely since all experiments derived from a single, well-
mixed, sieved soil subsample in order to reduce the effect of heterogeneity on the results. While
CFU based analyses are useful in giving us an approximation of the total number of culturable
cells present in the original sample and a broad overview of the effectiveness of certain culturing
practices at cultivating cells, it does not capture the important distinction of diversity on the
isolates obtained. This method captured the vast majority of the bacterial genera that we were
interested in examining further based on post-fire sequencing studies that reported significant

changes in post-fire bacterial diversity (Enright et al. 2021; Pulido-Chavez et al. 2022; Bárcenas-
Moreno et al. 2011; Caiafa et al. 2023; Brown et al. 2019). We were surprised by this result as
the post-fire soils were very dry to begin with, hence our hypothesis that drying would have the
least impact on microbial viability. One previous study found that freezing at -20ºC was very
detrimental to sandy, dry soils bacterial communities and instead found that refrigeration at 4ºC
was better at retaining microbial biomass and respiration (Černohlávková et al. 2009). While we
did not freeze soils at -20ºC and only froze samples at -80ºC, we did not observe the same
detrimental effect of freezing on our soil bacterial communities, despite the El Dorado Fire soils
being dry and sandy as well. Conversely, Rubin et al found that freezing at -20ºC preserved
bacterial community structure (as assayed using 16S sequencing profiles) the best though they do
note that their study only covered soils of one type from a single climatic region (Rubin et al.
2013). It is possible that the post-fire soils that we collected contain a high concentration of
various solutes released from burning that could have buffered the microbial communities from
the worst effects of freezing. It is also worth noting that the bacterial genus *Massilia* was only
obtained from soil that was preserved by freezing at -80°C with glycerol. This genus has been
repeatedly observed post-fire with very little understanding about its ecology (Pulido-Chavez et
al. 2022; Caiafa et al. 2023; Whitman et al. 2019; 2022), making it a prime candidate for further
study. Therefore, it may be of value for researchers to freeze some soil with glycerol as well as
without, to capture some of these more potentially vulnerable taxa.

Soil storage method seemed to matter less for pyrophilous fungi than bacterial genera.
While there was one method that stood out as the worst (freezing at -80°C with glycerol), the rest
of the methods captured most of the diversity observed (Figures 4B and 5B). This could be due
to the resilience of fungal spores to a variety of disturbance effects (Falconer, Otten, and White
2015), but it is also very possible that MYA was not the ideal culture medium for pyrophilous
fungi, given the success of R2A and PyOM media at cultivating pyrophilous fungi (Figures 4D
and 6B). Future work repeating the storage experiments utilizing R2A and PyOM might improve
pyrophilous fungal isolate retention. We were both interested and surprised to find that our
hypothesis that air-drying soil would be the most effective storage method was fully unsupported
by our results. As previously noted, earlier studies on soil storage with the intent of preserving
microbial communities can be contradictory and thus identifying the best soil storage method can
be both difficult and potentially specific to each soil type (Černohlávková et al. 2009; Rubin et
al. 2013). In fact, both in terms of CFU data and diversity, air-dried soil commonly performed as
one of the worst methods of storing soil for retaining culturable viability of both pyrophilous
fungi and bacteria (Figures 3, 4, and 5). This finding is significant as it is very common for soil
samples to be dried before laboratory use (Rubin et al. 2013; Castaño et al. 2016; Cui et al. 2014)
and if those experiments are to include culture work then researchers could be significantly
reducing their obtainable species by not culturing from fresh soil or by storing soil samples
frozen at -80°C. It is worth noting that we were unable to find studies exploring the impact of
soil storage on fungal communities and as such may indicate a strong need for these methods to
be tested across kingdoms.

In assessing the most effective culturing media for capturing pyrophilous bacteria and
fungi, we were surprised at how well the oligotrophic media R2A performed. While the rich

medias (LB and MYA) produced the most CFUs for both bacteria and fungi, R2A media was the
second-highest producer of CFUs for both groups (Figure 3). Considering that R2A was
developed to culture bacteria from potable drinking water (Reasoner and Geldreich 1985), this
would suggest that a large number of post-fire microbes prefer a more oligotrophic environment.
This was further supported by the diversity of pyrophilous genera obtained by these experiments,
with R2A consistently producing the most diverse pyrophilous bacterial (Figure 4C) and fungal
(Figure 4D) communities. While shifting life-strategies of microbial communities under stress
can be extremely varied and nuanced (Ho, Di Lonardo, and Bodelier 2017), under an increasing
stress gradient (such as carbon stress, soil moisture stress, or heat stress) oligotrophic bacteria
and fungi can be favored (Hernandez et al. 2021). It may be therefore that the low-nutrient
environment of R2A media may actually be ideal for capturing a wide diversity of the heavily-
stressed post-fire environment. This is a surprising result as post-fire environments tend to be
high in nitrogen and phosphorus, even if the carbon landscape has radically changed (Wan, Hui,
and Luo 2001; Michelotti and Miesel 2015). While previous studies exist examining the role of
culture media conditions on microbial culturability (Vieira and Nahas 2005; Sánchez-Clemente
et al. 2018; Malosso et al. 2006; Stevenson et al. 2004), these studies often take either a fine-
scale approach to tuning a specific culture medium or focus on very broad-scale patterns of
culturability such as relying on CFU-based metrics alone or utilizing only rich media. Continued
exploration of media conditions is essential to increasing the number of cultured microbial
species and insights gained from studies like this one will help open new paths in culture media
development. It is worth noting that while our home-made PyOM media only captured 50% of
pyrophilous fungi (Figure 4D) and 44% of bacteria (Figure 4C), it was also the only media to
successfully isolate key pyrophilous taxa of interest both bacteria (i.e. *Noviherbaspirillum soli*)
and fungi (*Neurospora crassa*) (Figure 6). These taxa are particularly interesting because like
Massilia, *Noviherbaspirillum* has been repeated observed increasing in abundance post-fire
(Enright et al. 2021; Pulido-Chavez et al. 2022; Caiafa et al. 2023) despite very little
understanding of its ecology, and *Neurospora crassa* is famously found post-fire and is widely
used as a genetics model organism despite very little understanding about its ecological role in
the environment (Kuo et al. 2014). Therefore, we conclude that the best media to use to
maximize pyrophilous bacterial and fungal diversity if you can only select one is R2A, but best
to also include a PyOM media if specifically interested in isolating the most finicky and perhaps
interesting pyrophilous microbes.

While this study has made significant observations regarding effective culturing practices
for pyrophilous bacteria and fungi, we would be remiss not to acknowledge both some of the
limitations of this study and the additional work still before us. Except for the PyOM media, we
limited ourselves to commonly available laboratory culturing media. We acknowledge there is a
near endless range of possible medias to use (Vieira and Nahas 2005) and there may be more
effective media available for the culturing of pyrophilous microorganisms. However, we assert
that the broad trends (rich media vs. oligotrophic media vs. PyOM) hold valuable insight into the
culturing of these microorganisms. As for the PyOM media itself, it is a complex undefined
media and is uses a water-extraction based method of incorporating nutrients from PyOM.
Recent work has identified that there are some nutrients that are more water-soluble in PyOM
than others (Zeba et al. 2024) so it is possible that our PyOM media does lack some nutrients that

could be playing a role in post-fire metabolic requirements and us such capturing a more limited
sampling of the microbial community. Additionally, the culturing effort in this study all took
place aerobically at room temperature and as such we once again are capturing a narrower
diversity of organisms than exist within our system. Further, future work could endeavor to test a
wider variety of culturing and incubation factors including testing a range of growth
temperatures (Bárcenas-Moreno and Bååth 2009; Vieira and Nahas 2005) and media pH (Davis,
Joseph, and Janssen 2005; Sánchez-Clemente et al. 2018).

Finally, while we acknowledge that separating bacterial and fungal cultures based on
colony morphology is difficult at best (Ben-Jacob, Cohen, and Gutnick 1998; Sousa et al. 2013;
Watrud et al. 2006), we decided that choosing which colonies to isolate based on morphotype
(and when there was any shred of doubt between two cultures, isolating both) and then backing
up identification with Sanger sequencing was an acceptable concession in the face of the
technical and logistic demands of isolating, stocking, and identifying every single colony we
were able to grow. This does mean that we could have missed some species-level diversity since
some species can share extremely similar colony morphologies (Sousa et al. 2013; Watrud et al.
2006), but as we captured 28 bacterial genera and 14 fungal genera across our experiments we
still assert that the isolates cultivated contribute substantially to the diversity of available
culturable species and that insights derived from the diversity of our isolates across treatments
are still meaningful. This is especially poignant as this is the first large-scale culturing effort of
post-fire bacteria we were able to identify, and one of the largest non-mushroom fungal culturing
efforts of pyrophilous fungi since Seaver began classifying pyrophilous species in 1910 (Seaver
and Clark 1910). Future culturing efforts could leverage techniques like dilution to extinction
culturing (Collado et al. 2007; Yang, Kang, and Cho 2016) to capture a broader species-level
number of identified isolates.

In conclusion, our experiments found that from high-severity burn scars in a chaparral
ecosystem that storing soil samples at -80°C without glycerol preserved the greatest diversity of
culturable pyrophilous bacteria and fungi. We found that certain taxa of interest were only
captured from soil frozen with glycerol, indicating a potential value in storing some soil in that
method as well. We found that the best laboratory media for culturing pyrophilous bacteria and
fungi was the oligotrophic media R2A, and the newly created solid agar media created from
field-harvested PyOM also captured a diverse community of isolates including some taxa of
interest that were only captured on PyOM. The methods we employed captured many groups of
interest from previous pyrophilous microbial studies and serve as an additional examination of
prudent laboratory practices when considering future culturing efforts from soil.

Finally, the isolates discussed in this study are one portion of a much larger culturing
effort of pyrophilous bacteria and fungi that leveraged 7 different wildfires, culturing from soil,
smoke, and fungal fruiting bodies, and contains over 500 isolates of bacteria and fungi
(Supplemental File 2). By testing and improving culture methods and undergoing large-scale
culturing efforts, we can significantly increase the diversity of microbial cultures available for
multi-omics, phenotypic, and applied microbiological research. For example, a culture collection
is necessary to perform biophysical assays and genomics to assess traits that can yield insights

into ecological processes and trade-offs (Maynard et al. 2019). Sequencing genomes from
isolates with concurrent biophysical assays is a powerful yet under-used tool to gain insight into
microbial traits and connect genomic information to trait expression (Chase et al. 2017).
Obtaining the cultures is the first step to characterizing their traits via 'omics and phenotypic
methods.

Acknowledgements

We thank the San Bernardino National Forest for allowing us to collect the soils needed
for this study following the El Dorado wildfire. We thank many individuals who contributed to
the Glassman lab pyrophilous microbial culture including lab managers Judy Chung, James
Randolph, and Maria Ordonez, collaborators Tom Bruns and Monika Fischer, UCR PhD students
Fabiola Pulido-Chavez, Arik Joukhajian, and Mark Yacoub, and UCR undergraduates Marely
Vega, Wine-Jie Lipardo, Audrey Reichard, Priscilla Shultz, Jorge Ponce, and Nathan Heger. This
project was funded by the Department of Energy BER Award #DE-SC0023127 and United States
Department of Agriculture-NIFA Award #2022-67014-36675 to SIG and the NSF GRFP to DJE.
We also thank Dylan Enright's Dissertation Committee members Jason Stajich and Marko
Spasojevic and anonymous reviewers for their helpful feedback on the manuscript.

Works Cited

Alster, Charlotte J, Steven D Allison, Nels G Johnson, Sydney I Glassman, and Kathleen K
Treseder. 2021. "Phenotypic Plasticity of Fungal Traits in Response to Moisture and
Temperature." *ISME Communications* 1 (1): 43. https://doi.org/10.1038/s43705-021-
00045-9.
Aranda, Elisabet, Patricia Godoy, Rocío Reina, Marina Badia-Fabregat, Mònica Rosell, Ernest
Marco-Urrea, and Inmaculada García-Romera. 2017. "Isolation of Ascomycota Fungi
with Capability to Transform PAHs: Insights into the Biodegradation Mechanisms of
Penicillium Oxalicum." *International Biodeterioration & Biodegradation* 122
(August):141–50. https://doi.org/10.1016/j.ibiod.2017.05.015.
Arnold, A. Elizabeth, and F. Lutzoni. 2007. "Diversity and Host Range of Foliar Fungal
Endophytes: Are Tropical Leaves Biodiversity Hotspots?" *Ecology* 88 (3): 541–49.
https://doi.org/10.1890/05-1459.
Bárcenas-Moreno, Gema, and Erland Bååth. 2009. "Bacterial and Fungal Growth in Soil Heated
at Different Temperatures to Simulate a Range of Fire Intensities." *Soil Biology and*
Biochemistry 41 (12): 2517–26. https://doi.org/10.1016/j.soilbio.2009.09.010.
Bárcenas-Moreno, Gema, Fuensanta García-Orenes, Jorge Mataix-Solera, Jorge Mataix-Beneyto,
and Erland Bååth. 2011. "Soil Microbial Recolonisation after a Fire in a Mediterranean
Forest." *Biology and Fertility of Soils* 47 (3): 261–72. https://doi.org/10.1007/s00374-
010-0532-2.
Ben-Jacob, Eshel, Inon Cohen, and David L. Gutnick. 1998. "COOPERATIVE
ORGANIZATION OF BACTERIAL COLONIES: From Genotype to Morphotype."

Annual Review of Microbiology 52 (Volume 52, 1998): 779–806.
https://doi.org/10.1146/annurev.micro.52.1.779.

Brown, Shawn, Allison Veach, Jonathan Horton, Emerald Ford, Ari Jumpponen, and Richard
Baird. 2019. "Context Dependent Fungal and Bacterial Soil Community Shifts in
Response to Recent Wildfires in the Southern Appalachian Mountains." *Forest Ecology*
and Management, August. https://doi.org/10.1016/j.foreco.2019.117520.

Caiafa, Marcos V., Amelia R. Nelson, Thomas Borch, Holly K. Roth, Timothy S. Fegel, Charles
C. Rhoades, Michael J. Wilkins, and Sydney I. Glassman. 2023. "Distinct Fungal and
Bacterial Responses to Fire Severity and Soil Depth across a Ten-Year Wildfire
Chronosequence in Beetle-Killed Lodgepole Pine Forests." *Forest Ecology and*
Management 544 (September):121160. https://doi.org/10.1016/j.foreco.2023.121160.

Caktu, Kivilcim, and Emir Turkoglu. 2011. "Microbial Culture Collections: The Essential
Resources for Life." *Gazi University Journal of Science* 24 (2): 175–80.

Caon, Lucrezia, V. Ramón Vallejo, Coen J. Ritsema, and Violette Geissen. 2014. "Effects of
Wildfire on Soil Nutrients in Mediterranean Ecosystems." *Earth-Science Reviews* 139
(December):47–58. https://doi.org/10.1016/j.earscirev.2014.09.001.

Carini, Paul. 2019. "A 'Cultural' Renaissance: Genomics Breathes New Life into an Old Craft."
mSystems 4 (3). https://doi.org/10.1128/mSystems.00092-19.

Castaño, Carles, Javier Parladé, Joan Pera, Juan Martínez de Aragón, Josu G. Alday, and José
Antonio Bonet. 2016. "Soil Drying Procedure Affects the DNA Quantification of
Lactarius Vinosus but Does Not Change the Fungal Community Composition."
Mycorrhiza 26 (8): 799–808. https://doi.org/10.1007/s00572-016-0714-3.

Černohlávková, Jitka, Jiří Jarkovský, Michala Nešporová, and Jakub Hofman. 2009. "Variability
of Soil Microbial Properties: Effects of Sampling, Handling and Storage." *Ecotoxicology*
and Environmental Safety 72 (8): 2102–8. https://doi.org/10.1016/j.ecoenv.2009.04.023.

Chase, Alexander B., Ulas Karaoz, Eoin L. Brodie, Zulema Gomez-Lunar, Adam C. Martiny, and
Jennifer B. H. Martiny. 2017. "Microdiversity of an Abundant Terrestrial Bacterium
Encompasses Extensive Variation in Ecologically Relevant Traits." *mBio* 8 (6):
10.1128/mbio.01809-17. https://doi.org/10.1128/mbio.01809-17.

Collado, Javier, Gonzalo Platas, Barbara Paulus, and Gerald F. Bills. 2007. "High-Throughput
Culturing of Fungi from Plant Litter by a Dilution-to-Extinction Technique." *FEMS*
Microbiology Ecology 60 (3): 521–33. https://doi.org/10.1111/j.1574-6941.2007.00294.x.

Cui, Hang, Caihuan Wang, Zhenhong Gu, Honghui Zhu, Shenlei Fu, and Qing Yao. 2014.
"Evaluation of Soil Storage Methods for Soil Microbial Community Using Genetic and
Metabolic Fingerprintings." *European Journal of Soil Biology* 63 (July):55–63.
https://doi.org/10.1016/j.ejsobi.2014.05.006.

Davis, Kathryn E. R., Shayne J. Joseph, and Peter H. Janssen. 2005. "Effects of Growth
Medium, Inoculum Size, and Incubation Time on Culturability and Isolation of Soil
Bacteria." *Applied and Environmental Microbiology* 71 (2): 826–34.
https://doi.org/10.1128/AEM.71.2.826-834.2005.

Enright, Dylan J., Kerri M. Frangioso, Kazuo Isobe, David M. Rizzo, and Sydney I. Glassman.
2021. "Mega-Fire in Redwood Tanoak Forest Reduces Bacterial and Fungal Richness and
Selects for Pyrophilous Taxa and Traits That Are Phylogenetically Conserved."
https://doi.org/10.1101/2021.06.30.450634.

Falconer, Ruth E., Wilfred Otten, and Nia A. White. 2015. "Chapter One - Toward Modeling the
Resistance and Resilience of 'Below-Ground' Fungal Communities: A Mechanistic and

Trait-Based Approach." In *Advances in Applied Microbiology*, edited by Sima Sariaslani
and Geoffrey Michael Gadd, 93:1–44. Academic Press.
https://doi.org/10.1016/bs.aambs.2015.08.001.
Ferrenberg, Scott, Sean P. O'Neill, Joseph E. Knelman, Bryan Todd, Sam Duggan, Daniel
Bradley, Taylor Robinson, et al. 2013. "Changes in Assembly Processes in Soil Bacterial
Communities Following a Wildfire Disturbance." *The ISME Journal* 7 (6): 1102–11.
https://doi.org/10.1038/ismej.2013.11.
Fischer, Monika S., Frances Grace Stark, Timothy D. Berry, Nayela Zeba, Thea Whitman, and
Matthew F. Traxler. 2021. "Pyrolyzed Substrates Induce Aromatic Compound
Metabolism in the Post-Fire Fungus, Pyronema Domesticum." *Frontiers in Microbiology*
12. https://www.frontiersin.org/article/10.3389/fmicb.2021.729289.
Franzosa, Eric A., Tiffany Hsu, Alexandra Sirota-Madi, Afrah Shafquat, Galeb Abu-Ali, Xochitl
C. Morgan, and Curtis Huttenhower. 2015. "Sequencing and beyond: Integrating
Molecular 'omics' for Microbial Community Profiling." *Nature Reviews Microbiology* 13
(6): 360–72. https://doi.org/10.1038/nrmicro3451.
González-Pérez, José A., Francisco J. González-Vila, Gonzalo Almendros, and Heike Knicker.
2004. "The Effect of Fire on Soil Organic Matter—a Review." *Environment International*
30 (6): 855–70. https://doi.org/10.1016/j.envint.2004.02.003.
Haridas, Sajeet, Colette Breuill, Joerg Bohlmann, and Tom Hsiang. 2011. "A Biologist's Guide to
de Novo Genome Assembly Using next-Generation Sequence Data: A Test with Fungal
Genomes." *Journal of Microbiological Methods* 86 (3): 368–75.
https://doi.org/10.1016/j.mimet.2011.06.019.
Hernandez, Damian J, Aaron S David, Eric S Menges, Christopher A Searcy, and Michelle E
Afkhami. 2021. "Environmental Stress Destabilizes Microbial Networks." *The ISME*
Journal 15 (6): 1722–34. https://doi.org/10.1038/s41396-020-00882-x.
Ho, Adrian, D. Paolo Di Lonardo, and Paul L. E. Bodelier. 2017. "Revisiting Life Strategy
Concepts in Environmental Microbial Ecology." *FEMS Microbiology Ecology* 93 (3):
fix006. https://doi.org/10.1093/femsec/fix006.
Hugenholtz, Philip. 2002. "Exploring Prokaryotic Diversity in the Genomic Era." *Genome*
Biology 3 (2): reviews0003.1. https://doi.org/10.1186/gb-2002-3-2-reviews0003.
Joint, Ian, Martin Mühling, and Joël Querellou. 2010. "Culturing Marine Bacteria – an Essential
Prerequisite for Biodiscovery." *Microbial Biotechnology* 3 (5): 564–75.
https://doi.org/10.1111/j.1751-7915.2010.00188.x.
Jung, Sokhee P., and Hojeong Kang. 2014. "Assessment of Microbial Diversity Bias Associated
with Soil Heterogeneity and Sequencing Resolution in Pyrosequencing Analyses."
Journal of Microbiology 52 (7): 574–80. https://doi.org/10.1007/s12275-014-3636-9.
Kämpfer, Peter. 2014. "Chapter 14 - Continuing Importance of the 'Phenotype' in the Genomic
Era." In *Methods in Microbiology*, edited by Michael Goodfellow, Iain Sutcliffe, and
Jongsik Chun, 41:307–20. New Approaches to Prokaryotic Systematics. Academic Press.
https://doi.org/10.1016/bs.mim.2014.07.005.
Knicker, Heike. 2011. "Pyrogenic Organic Matter in Soil: Its Origin and Occurrence, Its
Chemistry and Survival in Soil Environments." *Quaternary International*, Black Soils
and Black Sediments -- Archives of Landscape Evolution, 243 (2): 251–63.
https://doi.org/10.1016/j.quaint.2011.02.037.

Kuo, Hsiao-Che, Sun Hui, Jaeyoung Choi, Frederick O. Asiegbu, Jari P. T. Valkonen, and Yong-
Hwan Lee. 2014. "Secret Lifestyles of *Neurospora Crassa*." *Scientific Reports* 4
(May):5135. https://doi.org/10.1038/srep05135.
Leadbetter, Jared R. 2003. "Cultivation of Recalcitrant Microbes: Cells Are Alive, Well and
Revealing Their Secrets in the 21st Century Laboratory." *Current Opinion in*
Microbiology 6 (3): 274–81. https://doi.org/10.1016/S1369-5274(03)00041-9.
Lilleskov, Erik A., Thomas W. Kuyper, Martin I. Bidartondo, and Erik A. Hobbie. 2019.
"Atmospheric Nitrogen Deposition Impacts on the Structure and Function of Forest
Mycorrhizal Communities: A Review." *Environmental Pollution* 246 (March):148–62.
https://doi.org/10.1016/j.envpol.2018.11.074.
Malik, Khursheed A., and Dieter Claus. 1987. "Bacterial Culture Collections: Their Importance
to Biotechnology and Microbiology." *Biotechnology and Genetic Engineering Reviews* 5
(1): 137–98. https://doi.org/10.1080/02648725.1987.10647837.
Malosso, Elaine, Ian S. Waite, Lorna English, David W. Hopkins, and Anthony G. O'Donnell.
2006. "Fungal Diversity in Maritime Antarctic Soils Determined Using a Combination of
Culture Isolation, Molecular Fingerprinting and Cloning Techniques." *Polar Biology* 29
(7): 552–61. https://doi.org/10.1007/s00300-005-0088-z.
Martiny, Adam C. 2019. "High Proportions of Bacteria Are Culturable across Major Biomes."
The ISME Journal 13 (8): 2125–28. https://doi.org/10.1038/s41396-019-0410-3.
Maynard, Daniel S., Mark A. Bradford, Kristofer R. Covey, Daniel Lindner, Jessie Glaeser,
Douglas A. Talbert, Paul Joshua Tinker, Donald M. Walker, and Thomas W. Crowther.
2019. "Consistent Trade-Offs in Fungal Trait Expression across Broad Spatial Scales."
Nature Microbiology 4 (5): 846–53. https://doi.org/10.1038/s41564-019-0361-5.
Michelotti, Lucas A., and Jessica R. Miesel. 2015. "Source Material and Concentration of
Wildfire-Produced Pyrogenic Carbon Influence Post-Fire Soil Nutrient Dynamics."
Forests 6 (4): 1325–42. https://doi.org/10.3390/f6041325.
Petersen, P. M. 1970. "Danish Fireplace Fungi-an Ecological Investigation on Fungi on Burns."
Dansk Botanisk Arkiv 27 (3): 1–97.
Pierson, Derek N., Peter R. Robichaud, Charles C. Rhoades, and Robert E. Brown. 2019. "Soil
Carbon and Nitrogen Eroded after Severe Wildfire and Erosion Mitigation Treatments."
International Journal of Wildland Fire 28 (10): 814–21.
https://doi.org/10.1071/WF18193.
Prakash, Om, Yogesh Shouche, Kamlesh Jangid, and Joel E. Kostka. 2013. "Microbial
Cultivation and the Role of Microbial Resource Centers in the Omics Era." *Applied*
Microbiology and Biotechnology 97 (1): 51–62. https://doi.org/10.1007/s00253-012-
4533-y.
Preston, C. M., and M. W. I. Schmidt. 2006. "Black (Pyrogenic) Carbon in Boreal Forests: A
Synthesis of Current Knowledge and Uncertainties." *Biogeosciences Discussions* 3 (1):
211–71.
Pulido-Chavez, M. Fabiola, James W. J. Randolph, Cassandra Zalman, Loralee Larios, Peter M.
Homyak, and Sydney I. Glassman. 2023. "Rapid Bacterial and Fungal Successional
Dynamics in First Year after Chaparral Wildfire." *Molecular Ecology* 32 (7): 1685–1707.
https://doi.org/10.1111/mec.16835.
R Core Team. 2020. *R: A Language and Environment for Statistical Computing*. Vienna, Austria:
R Foundation for Statistical Computing. https://www.R-project.org/.

Reasoner, D J, and E E Geldreich. 1985. "A New Medium for the Enumeration and Subculture of
Bacteria from Potable Water." *Applied and Environmental Microbiology* 49 (1): 1–7.
https://doi.org/10.1128/aem.49.1.1-7.1985.
Reyes-César, Anaisell, Ángel E. Absalón, Francisco J. Fernández, Juan Manuel González, and
Diana V. Cortés-Espinosa. 2014. "Biodegradation of a Mixture of PAHs by Non-
Ligninolytic Fungal Strains Isolated from Crude Oil-Contaminated Soil." *World Journal*
of Microbiology and Biotechnology 30 (3): 999–1009. https://doi.org/10.1007/s11274-
013-1518-7.
Rubin, Benjamin E. R., Sean M. Gibbons, Suzanne Kennedy, Jarrad Hampton-Marcell, Sarah
Owens, and Jack A. Gilbert. 2013. "Investigating the Impact of Storage Conditions on
Microbial Community Composition in Soil Samples." *PLOS ONE* 8 (7): e70460.
https://doi.org/10.1371/journal.pone.0070460.
Sánchez-Clemente, Rubén, María Isabel Igeño, Ana G. Población, María Isabel Guijo, Faustino
Merchán, and Rafael Blasco. 2018. "Study of pH Changes in Media during Bacterial
Growth of Several Environmental Strains." *Proceedings* 2 (20): 1297.
https://doi.org/10.3390/proceedings2201297.
Seaver, Fred J., and Ernest D. Clark. 1910. "Studies in Pyrophilous Fungi: II. Changes Brought
about by the Heating of Soils and Their Relation to the Growth of Pyronema and Other
Fungi." *Mycologia* 2 (3): 109–24. https://doi.org/10.2307/3753620.
Sousa, Ana Margarida, Idalina Machado, Ana Nicolau, and Maria Olívia Pereira. 2013.
"Improvements on Colony Morphology Identification towards Bacterial Profiling."
Journal of Microbiological Methods 95 (3): 327–35.
https://doi.org/10.1016/j.mimet.2013.09.020.
Stackebrandt, Erko, David Smith, Serge Casaregola, Giovanna Cristina Varese, Gerard Verkleij,
Nelson Lima, and Paul Bridge. 2014. "Deposit of Microbial Strains in Public Service
Collections as Part of the Publication Process to Underpin Good Practice in Science."
SpringerPlus 3 (1): 208. https://doi.org/10.1186/2193-1801-3-208.
Stevenson, Bradley S., Stephanie A. Eichorst, John T. Wertz, Thomas M. Schmidt, and John A.
Breznak. 2004. "New Strategies for Cultivation and Detection of Previously Uncultured
Microbes." *Applied and Environmental Microbiology* 70 (8): 4748–55.
https://doi.org/10.1128/AEM.70.8.4748-4755.2004.
Stoof, Cathelijne R., Jan G. Wesseling, and Coen J. Ritsema. 2010. "Effects of Fire and Ash on
Soil Water Retention." *Geoderma* 159 (3): 276–85.
https://doi.org/10.1016/j.geoderma.2010.08.002.
Swenson, Tami L., Ulas Karaoz, Joel M. Swenson, Benjamin P. Bowen, and Trent R. Northen.
2018. "Linking Soil Biology and Chemistry in Biological Soil Crust Using Isolate
Exometabolomics." *Nature Communications* 9 (1): 19. https://doi.org/10.1038/s41467-
017-02356-9.
Vieira, F. C. S., and E. Nahas. 2005. "Comparison of Microbial Numbers in Soils by Using
Various Culture Media and Temperatures." *Microbiological Research* 160 (2): 197–202.
https://doi.org/10.1016/j.micres.2005.01.004.
Wan, Shiqiang, Dafeng Hui, and Yiqi Luo. 2001. "Fire Effects on Nitrogen Pools and Dynamics
in Terrestrial Ecosystems: A Meta-Analysis." *Ecological Applications* 11 (5): 1349–65.
https://doi.org/10.1890/1051-0761(2001)011[1349:FEONPA]2.0.CO;2.
Watrud, Lidia S., Kendall Martin, Kelly K. Donegan, Jeffrey K. Stone, and Clarace G. Coleman.
2006. "Comparison of Taxonomic, Colony Morphotype and PCR-RFLP Methods to

Characterize Microfungal Diversity." *Mycologia* 98 (3): 384–92.
https://doi.org/10.1080/15572536.2006.11832673.
Whitman, Thea, Ellen Whitman, Jamie Woolet, Mike D. Flannigan, Dan K. Thompson, and
Marc-André Parisien. 2019. "Soil Bacterial and Fungal Response to Wildfires in the
Canadian Boreal Forest across a Burn Severity Gradient." *Soil Biology and Biochemistry*
138 (November):107571. https://doi.org/10.1016/j.soilbio.2019.107571.
Whitman, Thea, Jamie Woolet, Miranda Sikora, Dana B. Johnson, and Ellen Whitman. 2022.
"Resilience in Soil Bacterial Communities of the Boreal Forest from One to Five Years
after Wildfire across a Severity Gradient." *Soil Biology and Biochemistry* 172
(September):108755. https://doi.org/10.1016/j.soilbio.2022.108755.
Wickham, Hadley, Winston Chang, Lionel Henry, Thomas Lin Pedersen, Kohske Takahashi,
Claus Wilke, Kara Woo, Hiroaki Yutani, and RStudio. 2019. "Ggplot2: Create Elegant
Data Visualisations Using the Grammar of Graphics." https://CRAN.R-
project.org/package=ggplot2.
Woolet, Jamie, and Thea Whitman. 2020. "Pyrogenic Organic Matter Effects on Soil Bacterial
Community Composition." *Soil Biology and Biochemistry* 141 (February):107678.
https://doi.org/10.1016/j.soilbio.2019.107678.
Yang, Seung-Jo, Ilnam Kang, and Jang-Cheon Cho. 2016. "Expansion of Cultured Bacterial
Diversity by Large-Scale Dilution-to-Extinction Culturing from a Single Seawater
Sample." *Microbial Ecology* 71 (1): 29–43. https://doi.org/10.1007/s00248-015-0695-3.
Zeba, Nayela, Timothy D. Berry, Monika S. Fischer, Matthew F. Traxler, and Thea Whitman.
2024. "Soil Carbon Mineralization and Microbial Community Dynamics in Response to
Pyrogenic Organic Matter Addition." *Soil Biology and Biochemistry* 191 (April):109328.
https://doi.org/10.1016/j.soilbio.2024.109328.

Figures:

Figure 1:

Figure 1: A conceptual diagram of the experimental design used in this experiment. "LB" is
abbreviated for Lysogeny Broth, "MYA" is Malt Yeast Agar, "R2A" is Reasoner's 2 Agar, and
"PyOM" is Pyrogenic Organic Matter. Note that the cultures obtained from fresh soil plated onto
rich media also serve as the "Fresh" storage condition for the soil storage experiment. Green
circles and orange stars symbolize two different morphotypes collected from the same initial
plate.

Figure 2:

Figure 2: A snapshot of the morphological diversity of the cultures obtained from the El Dorado
Fire across all soil storage and media type experiments. Identities were obtained via Sanger
sequencing of the 16S or ITS regions for bacteria and fungi, respectively. Moving left to right
across each row, the isolates shown are A)*Glutamicibacter bergerei, Arthrobacter agilis,*
Streptomyces mauvecolor, Kocuria rosea, B) *Arthrobacter bussei, Holtermaniella festucosa,*
Pedobacter panaciterrae, Curtobacterium oceanosedimentum, C) *Penicillium radiatolobatum,*
Penicillium fagi, Penicillium glabrum, Penicillium thomii, D) *Penicillium chalabudae,*
Penicillium radiatolobatum, Penicillium murcianum, Penicillium adametzii.

Figure 3:

Figure 3: Calculated capturable CFUs per gram soil for bacteria among the soil storage
experiment (A) and the media type experiment (C) and fungal CFUs per gram soil in the soil
storage experiment (B) and the media type experiment (D). Significance was tested using
ANOVA and Tukey Post-Hoc lettering has been applied. Points and bars are colored by
treatment.

Figure 4:

Figure 4: Summarized counts of pyrophilous bacterial and fungal genera (green) and species
(grey) obtained from the soil storage experiment (A, Bacteria & B, Fungi) and from the media
types experiments (C, Bacteria & D, Fungi). Treatments are ordered by highest amount of genera
first, then species count second.

Figure 5:

Figure 5: Heatmap-style visualization showing the diversity of pyrophilous bacterial (A) and
fungal (B) genera across the soil storage experiment. A filled blue square indicates the presence
of the genus on the x axis in the soil storage method indicated on the y axis.

Figure 6:

Figure 6: Heatmap-style visualization showing the diversity of pyrophilous bacterial (A) and
fungal (B) genera across the media types experiment. A filled blue square indicates the presence
of the genus on the x axis in the media indicated on the y axis.