Probing the role of Nogo receptor homolog NgR2 in setting up cochlear connectivity

Nare Karagulyan[1,2,3#], Maja Überegger[4#], Yumeng Qi[5#], Norbert Babai[1,2], Florian Hofer[4], Lejo
Johnson Chacko[6], Fangfang Wang[5], Maria Luque[6], Rudolf Glueckert[6], Anneliese Schrott-
Fischer[6], Yunfeng Hua[5*], Tobias Moser[1,2,3,7*], Christine Bandtlow[4*]

[1]Institute for Auditory Neuroscience, University Medical Center Göttingen, Göttingen, Germany

[2]Auditory Neuroscience and Synaptic Nanophysiology Group, Max Planck Institute for Multidisciplinary
Sciences, Göttingen, Germany

[3]Cluster of Excellence "Multiscale Bioimaging of Excitable Cells", 37075 Göttingen, Germany

[4]Institute of Neurobiochemistry, Biocenter, Medical University of Innsbruck, 6020 Innsbruck, Austria

[5]Shanghai Institute of Precision Medicine, Ninth People's Hospital, Shanghai Jiao Tong University School of
Medicine, Shanghai 200125, China

[6]Department of Otorhinolaryngology, Medical University of Innsbruck, 6020 Innsbruck, Austria

[7]Department for Hearing, Speech and Voice Disorders, Medical University of Innsbruck, 6020 Innsbruck, Austria

Running title: Expression and function of Nogo receptor homolog NgR2 in the cochlea

#equal contribution
*correspondence to
Dr. Christine Bandtlow: christine.bandtlow@i-med.ac.at
Dr. Tobias Moser: tmoser@gwdg.de
Dr. Yunfeng Hua: yunfeng.hua@shsmu.edu.cn

Abstract

Sound encoding depends on the precise and reliable neurotransmission at the afferent
synapses between the sensory inner hair cells (IHCs) and spiral ganglion neurons (SGNs). The
molecular mechanisms contributing to the formation, as well as interplay between the pre-
and postsynaptic components remains largely unclear. Here, we tested the role of the Nogo
receptor homolog NgR2 (RTN4rl2) in the development and function of afferent IHC-SGN
synapses. Upon deletion of NgR2 in mice (NgR2$^{-/-}$), presynaptic IHC active zones showed
enlarged synaptic ribbons and a depolarized shift in the activation of Ca$_V$1.3 Ca^{2+} channels.
The postsynaptic densities (PSDs) of SGNs were smaller and deficient of GluA2/3 despite
maintained Gria2 mRNA expression in SGNs. Next to synaptically engaged PSDs we observed
"orphan" PSDs located away from IHCs. They likely belong to a subset of SGN peripheral
neurites that do not contact the IHCs in NgR2$^{-/-}$ cochleae as found by volume electron

microscopy reconstruction of SGN neurites. Auditory brainstem responses of NgR2$^{-/-}$ mice

showed increased sound thresholds indicating impaired hearing. Together, these findings

suggest that NgR2 contributes to the proper formation and function of auditory afferent

synapses and is critical for normal hearing.

Introduction

Hearing relies upon the correct formation and maturation of cochlear afferent synapses

between postsynaptic type I spiral ganglion neurons (SGNs) and presynaptic inner hair cells

(IHCs; reviewed in Johnson et al., 2019; Bulankina and Moser, 2012; Appler and Goodrich,

2011). In mice, SGN neurites extend and reach IHCs at late embryonic stages (E16.5;

Koundakjian, Appler, and Goodrich, 2007), whereby early synaptic contacts are established

between the two cells at E18 (Michanski et al., 2019). Developmental changes and maturation

of afferent synapses continue until the onset of hearing (p12), with further refinement

occurring till the fourth postnatal week (Wong et al., 2014; Liberman and Liberman, 2016;

Michanski et al., 2019). In mature cochleae, each IHC receives a contact from 5-30 SGNs, while

each SGN contacts only one active zone (AZ) of one IHC in the majority of cases (Meyer and

Moser, 2010; Hua et al., 2021).

To this date, multiple mechanisms have been implicated in SGN neurite guidance and

establishment of IHC innervation by SGNs, including signaling via EphrinA5-EphA4,

Neuropilin2/Semaphorin3F, Semaphorin5B/PlexinA1, Semaphorin3A (Defourny et al., 2013;

Coate et al., 2015; Jung et al., 2019; Cantu-Guerra et al., 2023). Less is known about the

molecules governing the transsynaptic organization at IHC afferent synapses. Similar to the

central conventional synapses, synaptic adhesion proteins such as neurexins and neuroligins

have been suggested to play a role in pre- and postsynaptic assemblies (Ramirez et al., 2022;

Jukic et al., 2024). Furthermore, the Ca$_V$1.3 extracellular auxiliary subunit Ca$_V\alpha_2\delta_2$ was

indicated to be important for proper alignment of presynaptic IHC AZs and postsynaptic

densities (PSD) of SGNs (Fell et al., 2015).

Whether Nogo receptors (NgRs) contribute to setting up afferent connectivity in the

cochlea remained to be investigated. The Nogo receptor family consists of 3 homologous

proteins NgR1-3. NgRs are leucine-rich repeat (LRR) and glycosylphosphatidylinositol (GPI)

anchored cell surface receptors (Figure 1A). Their primary role is thought to limit synaptic

plasticity and axonal outgrowth as well as to restrict axonal regeneration after injury (reviewed in Mironova and Giger, 2013). While NgR1 and NgR3 are involved in axonal guidance (Vaccaro et al., 2022), NgR2 has been proposed to be important for innervation of the epidermis by dorsal root ganglion neurons (Bäumer et al., 2014). Furthermore, NgRs are suggested to control the number and the development of synapses (Wills et al., 2012) and to play a role in transsynaptic signaling (Wang et al., 2021).

Recent single-cell transcriptomic studies of the cochlea detected the expression of NgRs in SGNs and IHCs (Shrestha et al., 2018; Jean et al., 2023). Here, we investigated the role of NgR2 in the cochlea using previously described NgR2 constitutive knock-out mice (NgR2$^{-/-}$, (Wörter et al., 2009). We found NgR2 to be expressed in SGNs and to be required for normal hearing: auditory brainstem responses were impaired upon NgR2 deletion. We discovered both pre- and postsynaptic alterations of IHC-SGN synapses in NgR2$^{-/-}$ mice: presynaptic Ca^{2+} channels of IHCs required stronger depolarization to activate and PSDs seemed deficient of GluA2/3. Additionally, a subset of type I SGN neurites did not contact IHCs but likely still feature "orphan" PSDs.

Results

NgR2 is expressed both in hair cells and spiral ganglion neurons

Recent transcriptomic data had detected NgR2 expression both in hair cells and SGNs (Elkon et al., 2015; Liu et al., 2018; Shrestha et al., 2018; Jean et al., 2023). We verified this by performing RNAscope staining in mid-modiolar sections of the cochlea and detected expression of NgR2 mRNA both in outer and inner hair cells, as well as SGNs (Figure 1B, C). Spiral ganglia of NgR2$^{-/-}$ mice were lacking NgR2-specific mRNA puncta (Figure 1C), demonstrating specific detection of NgR2 expression in hair cells and SGNs. Next, we performed immunostaining and found specific NgR2 immunofluorescence in SGN somata, which was not present in the absence of the primary antibody (Figure 1D, E) or on SGN somata of NgR2$^{-/-}$ mice (data not shown).

Figure 1. NgR2 mRNA and protein expression in IHCs and SGNs of the mouse cochlea. (**A**)
NgR2 is an LRR protein and is anchored to the cell membrane via GPI. In the nervous system
NgR2 has been implicated to interact with MAG, versican, BAI. (**B**) Representative images of
RNAscope ISH for NgR2 mRNA (green dots) in the hair cell region of p1 wild-type animals. Hair
cells are visualized with anti-Myo7a stainings. Scale bar = 10 μm. (**C**) Representative images of
RNAscope ISH for NgR2 mRNA (green dots) combined with immunostaining for neuron-
specific marker βIII-tubulin (grey) in paraffin sections of p40 NgR2$^{+/+}$ and NgR2$^{-/-}$ spiral
ganglion sections. Scale bar = 20 μm. (D) Double immunostaining with anti-NgR2 and βIII-
tubulin on cryosections of p6 spiral ganglia. Immunoreactivity of NgR2 (red) was colocalized
with βIII-tubulin positive neurons (grey). (E) No specific signal was detected by omitting the
primary NgR2 antibody (negative control). Scale bar = 20 μm.

NgR2 is important for the correct development of the auditory afferent synapses

To probe the role of NgR2 in the cochlea, we first studied the numbers of cochlear cells in

NgR2$^{-/-}$ mice. We did not observe any change in SGN density or counts of inner and outer hair

cells at p15, 1-month- and 2-month-old mice (Figure 2-figure supplement 1). Given that NgR2

has been implicated in synapse formation and development (Wills et al., 2012; Borrie et al.,

2014; Wang et al., 2021) we immunolabeled IHC afferent synapses for presynaptic

RIBEYE/Ctbp2 and postsynaptic Homer1 in mice at the age of 3 weeks. While we did not detect

any change in the number of synaptic ribbons in NgR2$^{-/-}$ IHCs, ribbon volumes were bigger

(Figure 2C, D; Figure 2-figure supplement 2). Interestingly, in addition to the Homer1 positive

puncta juxtaposing presynaptic ribbons, we observed additional Homer1 patches, which

appeared to be away from IHCs, potentially marking "orphan" PSDs (Figure 2A, B). Moreover, the Homer1 puncta juxtaposing IHC AZs were significantly smaller in NgR2$^{-/-}$ mice compared to the control IHCs (Figure 2E). While we found the percentage of presynaptic ribbons juxtaposing Homer1 immunofluorescent puncta to be decreased by approximately 7% in NgR2$^{-/-}$ mice (Figure 2F), we cannot exclude that our immunolabeling protocol lacked the sensitivity to detect smaller synaptically engaged PSDs. Out of the four pore-forming AMPA receptor subunits (GluA1-4), mature SGNs express GluA2-4 (Niedzielski and Wenthold, 1995; Matsubara et al., 1996). Our immunolabeling of anti-GluA2/3 revealed a severe reduction of GluA2/3 positive puncta in NgR2$^{-/-}$, indicating possible lack or decrease of GluA2/3 (Figure 2-figure supplement 3A, B). Interestingly, the remaining GluA2/3 patches did not properly juxtapose presynaptic ribbons (Figure 2-figure supplement 3C). Despite this, the expression of Gria2 RNA appeared to be maintained in SGNs, as indicated by RNAscope (Figure 2-figure supplement 3D). To check if the "orphan" PSDs away from IHCs were possibly erroneously engaged with efferent presynaptic terminals we stained the latter for synapsin, which is lacking from IHCs, but did not observe any obvious juxtaposition between the "orphan" Homer1 and synapsin puncta (Figure 2-figure supplement 4).

Figure 2. Pre- and postsynaptic changes at IHCs of NgR2⁻ᐟ⁻ mice. (**A, B**) Maximal intensity projections of representative confocal stacks of IHCs from NgR2$^{+/+}$ (A) and NgR2$^{-/-}$ (B) mice immunolabeled against Vglut3, Homer1, and Ctbp2. Scale bar = 5 μm. Images on the right-hand side are zoomed into the synaptic regions. Scale bar = 2 μm. Some of the putative "orphan" PSDs are marked with the white arrowheads. (**C**) Number of Ctbp2 positive puncta is not changed in IHCs of NgR2$^{-/-}$ mice (NgR2$^{+/+}$: 9.9 ± 0.42, SD = 1.62, n = 15, N = 2 vs NgR2$^{-/-}$: 11.4 ± 0.5, SD = 2.25, n = 20, N = 3; p = 0.06, Mann-Whitney-Wilcoxon test). (**D**) Ribbon volumes are enlarged in NgR2$^{-/-}$ IHCs (NgR2$^{+/+}$: 0.16 ± 0.007 μm³, SD = 0.09 μm³, n = 165, N = 2 vs NgR2$^{-/-}$: 0.21 ± 0.005 μm³, SD = 0.09 μm³, n = 259, N = 3; p < 0.001, Mann-Whitney-Wilcoxon test). (**E**) Homer1 patches which are juxtaposing presynaptic ribbons show decreased volumes in NgR2$^{-/-}$ IHCs (NgR2$^{+/+}$: 0.36 ± 0.01 μm³, SD = 0.16 μm³, n = 160, N = 2 vs NgR2$^{-/-}$: 0.26 ± 0.01 μm³, SD = 0.19 μm³, n = 249, N = 3; p < 0.001, Mann-Whitney-Wilcoxon test). (**F**) Percentage of Ctbp2 puncta juxtaposing Homer1 is slightly decreased in NgR2$^{-/-}$ mice (NgR2$^{+/+}$: 96.7 ± 1.45 %, SD = 5.44 %, n = 14, N = 2 vs NgR2$^{-/-}$: 89.5 ± 2.05 %, SD = 9.18 %, n =

20, N = 3; p = 0.03, Mann-Whitney-Wilcoxon test). Data is presented as mean ± SEM. Box-
whisker plots show the median, 25/75 percentiles (box), and 10/90 percentiles (whiskers).
Individual data points are overlaid.

Additional non-synaptic neurites in the cochlea of NgR2$^{-/-}$ mice

To further examine the afferent cochlear connectivity, we utilized serial block-face scanning

electron microscopy (SBEM) to 3D reconstruct the apical cochlea segments from two NgR2$^{-/-}$

and one littermate NgR2$^{+/+}$ control mice at p36 (Figure 3A). As recently demonstrated (Hua et

al., 2021; Lu et al., 2024), the spatial resolution of SBEM allows reliable ribbon synapse

identification and neurite reconstruction (Figure 3B and C). In these three datasets, we have

traced 133 neural fibers from their postsynaptic ending in the organ Corti towards the

modiolus, of which 115 were classified as peripheral neurites of type I SGN based on their

radial calibers and extensive myelination after entering the habenula perforata (Figure 3D).

We observed a substantial number of neurites that did not engage the IHCs (non-synaptic

neurites) in the NgR2$^{-/-}$ organs of Corti (23 out of 83) in contrast to the wild-type control (3

out of 32, Figure 3E). Although most non-synaptic neurites were found in the inner spiral

bundle, they failed to reach IHC basal lateral poles to form contacts (17 out of 23 in NgR2$^{-/-}$

animals). This result might provide a plausible explanation for the "orphan" Homer1 puncta

shown by immunohistochemistry in NgR2$^{-/-}$ organs of Corti. Note that the presence of non-

synaptic neurites in the NgR2$^{-/-}$ animal was not associated with a profound reduction in ribbon

synapses (Figure 2C, Figure 2-figure supplement 2B, Figure 3-figure supplement 1A),

suggesting a different scenario than deafferentation due to excitotoxic synaptopathy as

recently reported (Moverman et al., 2023). Nevertheless, nearly all traced radial fibers were

found predominantly unbranched in both the NgR2$^{-/-}$ (97%) and the wild-type control mice

(94%; Figure 3F).

Figure 3. Additional non-synaptically engaged SGN neurites in the cochlea of NgR2$^{-/-}$ mice.
(**A**) Workflow of SBEM imaging at the mouse apical cochlear region. (**B**) Example images of
neurites beneath IHCs from the NgR2$^{+/+}$ (left) and NgR2$^{-/-}$ (right) mice. Synaptic ribbons are
indicated with red arrows. The regions of interest were magnified from single sections of
SBEM datasets (insets). Scale bar = 2 μm. (**C**) 3D rendering of afferent fiber reconstruction
with ribbons (red), showing both synaptic (green, type I SGN with ribbon) and non-synaptic
(grey, type I SGN without ribbon) populations in the NgR2$^{-/-}$ mouse. Scale bar = 10 μm. (**D**)
Display of classified radial fibers in the NgR2$^{+/+}$ (left) and NgR2$^{-/-}$ (middle and left) animals. All
fibers were traced from the habenula perforata (cycles) before classification to avoid bias to
terminal types. Scale bar 10 μm. (**E**) Percentage of radial fibers with ribbon per bundle
(NgR2$^{+/+}$: 91.41 ± 8.35 %, n = 3 bundles, N = 1 vs NgR2$^{-/-}$: 74.40 ± 14.58%, n = 6 bundles, N = 2;
p = 0.032, unpaired t-test). (**F**) Percentage of unbranched radial fibers per bundle (NgR2$^{+/+}$:

94.19 ± 5.04 %, n = 3 bundles, N = 1 vs NgR2$^{-/-}$: 96.98 ± 3.49 %, n = 6 bundles, N = 2; p = 0.226,

unpaired t-test).

Deletion of NgR2 results in depolarized shift of the Ca^{2+} channel activation in IHCs

Next, we tested for potential effects of NgR2 deletion on the presynaptic IHC function. We

performed whole-cell patch-clamp recordings from IHCs of p21-29 NgR2$^{-/-}$ and NgR2$^{+/+}$ mice.

First, we recorded voltage-gated Ca^{2+} currents by applying step depolarizations with 5 mV

increment in ruptured patch-clamp configuration (Figure 4A, see Materials and Methods). We

did not observe any change in the maximal Ca^{2+} current amplitude (Figure 4B, C), yet noticed

small (∼ +3 mV) but consistent and statistically significant depolarized shift of the voltage of

half-maximal Ca^{2+} channel activation (V$_{half}$; Figure 4D, E). The voltage sensitivity of the

channels was not changed in NgR2$^{-/-}$ IHCs (Figure 4F). Next, we probed Ca^{2+} influx triggered

exocytosis from IHCs by applying depolarizations of varying durations to voltages saturating

Ca^{2+} influx in IHCs of both genotypes (-17 mV) and recording exocytic membrane capacitance

changes (ΔC$_m$) in the perforated patch-clamp configuration (Figure 4G). Both fast (up to 20 ms

depolarization) and sustained components of exocytosis remained intact in the mutant,

indicating unaffected readily releasable pool and replenishment of the vesicles in NgR2$^{-/-}$ IHCs

(Figure 4H).

Figure 4. Shifted operation range of Ca²⁺ channels but intact exocytosis in IHCs of NgR2⁻ᐟ⁻
mice. (**A**) Representative current traces from IHCs of NgR2⁺ᐟ⁺ (top, black) and NgR2⁺ᐟ⁺ (bottom,
red) evoked by step depolarizations. (**B**) Average Ca²⁺ current-voltage relationships (IV curves)
in NgR2⁺ᐟ⁺ and NgR2⁻ᐟ⁻ IHCs. (**C**) Maximal Ca²⁺ current amplitude is not changed in NgR2⁻ᐟ⁻ IHCs
(NgR2⁺ᐟ⁺: 210 ± 10.9 pA, SD = 40.9 pA, n = 14, N = 5 vs NgR2⁻ᐟ⁻: -200 ± 8.19 pA, SD = 40.9 pA, n
= 25, N = 8; p = 0.47, Student's t-test). (**D**) Fractional activation curves of Ca²⁺ channels
calculated from IVs show depolarized shift in channel activation in NgR2⁻ᐟ⁻ IHCs. (**E**) Voltage of
half maximal activation obtained from Boltzmann fit of the curves from (D) is more positive in

NgR2[-/-] IHCs (NgR2[+/+]: -29.4 ± 0.66 mV, SD = 2.48 mV, n = 14, N = 5 vs NgR2[-/-]: -26.6 ± 0.59 mV, SD = 2.94 mV, n = 25, N = 8; p = 0.004, Student's t-test). (**F**) Voltage sensitivity (k) is not changed in NgR2[-/-] IHCs (NgR2[+/+]: 6.92 ± 0.12 mV, SD = 0.46 mV, n = 14, N = 5 vs NgR2[-/-]: 6.98 ± 0.1 mV, SD = 0.51 mV, n = 25, N = 8; p = 0.93, Mann-Whitney-Wilcoxon test). (**G**) Average current traces evoked by 50 ms depolarization to -17mV (top row) and resulting capacitance response (bottom row) from NgR2[+/+] (left, black) and NgR2[+/+] (right, red) IHCs. Shaded areas represent ± SEM. (**H**) Exocytic capacitance change (ΔC_m, top) and corresponding Ca^{2+} charge (Q_{Ca}^{2+}, bottom) evoked by depolarizations (to -17 mV) of various durations (2, 5, 10, 20, 50, 100 ms). Box-whisker plots show the median, 25/75 percentiles (box) and 10/90 percentiles (whiskers). Individual data points are overlaid.

We then turned to study single AZ function by combining whole cell ruptured patch-clamp combined with spinning disc confocal imaging of presynaptic Ca^{2+} influx, as described previously (Ohn et al., 2016). TAMRA conjugated Ctbp2 binding dimeric peptide and the low affinity Ca^{2+} indicator Fluo4-FF (k_D: 10 µM) were loaded into the cell via the patch pipette and Ca^{2+} influx at single AZs was visualized in the form of hotspots by applying voltage ramp depolarizations to the IHCs and simultaneously imaging Fluo4-FF fluorescence near labeled ribbons (Figure 5A, see Materials and methods). Maximal Ca^{2+} influx at single AZs tended to be higher in IHCs of NgR2[-/-] mice but the difference did not reach statistical significance (Figure 5B, B[i]). The V_{half} of Ca^{2+} channel clusters at single AZs showed depolarized shift ($\sim$ +5 mV), which was statistically significant despite the large variability of V_{half} across different AZs (Figure 5C, C[i]). Furthermore, we did not observe any mismatch in Ctbp2 positive puncta and evoked Ca^{2+} hotspots, indicating the correct localization of Ca^{2+} channels at the AZs. We verified this by performing immunostainings of $Ca_v1.3$, Ctbp2 and Bassoon and did not detect any apparent misalignment of these protein clusters (Figure 5D). In summary, deletion of NgR2 causes a depolarized shift of the activation of presynaptic Ca^{2+} influx in IHCs as demonstrated at the single AZ and whole cell levels. This is expected to elevate the sound pressure levels required to reach a comparable activity of synaptic transmission for sound encoding and predicts elevated auditory thresholds which we tested by recordings of auditory brainstem responses.

Figure 5. Depolarized shift of Ca²⁺ channel activation at single AZs but intact presynaptic organization in NgR2⁻/⁻ IHCs. (**A**) Voltage ramp stimulation protocols (top), evoked whole cell currents (middle) and the presynaptic hotspots of Fluo4-FF fluorescence (bottom) of a representative IHC recording. Black and grey colors represent the two stimulations, one being 5 ms shifted over the other. Images on the right show single image planes of representative NgR2⁺/⁺ (left) and NgR2⁻/⁻ (right) IHCs filled with TAMRA conjugated Ctbp2 binding peptide (Ctbp2 bp) and Fluo4-FF Ca²⁺ dye. Ca²⁺ hotspots are visualized by subtracting the average of baseline planes from the average of 5 planes during stimulation. Scale bar = 2 μm (**B**) Average fluorescence-voltage relationships of Ca²⁺ influx at single AZ from NgR⁺/⁺ and NgR2⁻/⁻ IHCs show no difference in the maximal Ca²⁺ amplitude (**Bⁱ**; NgR2⁺/⁺: 1.6 ± 0.17, SD = 1.2, n = 50 AZs vs NgR2⁻/⁻: 1.7 ± 0.13 pA, SD = 1.09, n = 69 AZs; p = 0.24, Mann-Whitney-Wilcoxon test). Shaded areas represent ± SEM. (**C**) Average fractional activation curves of Ca²⁺ channels at single AZs relationships of single AZ show intact voltage sensitivity (**Cⁱ**; NgR2⁺/⁺: 5.79 ± 0.32 mV, SD = 2.04 mV, n = 42 AZs vs NgR2⁻/⁻: 6.25 ± 0.22 mV, SD = 1.7 mV, n = 59 AZs; p = 0.06, Mann-Whitney-Wilcoxon test) but depolarized shift of V_{half} (**Cⁱⁱ**; NgR2⁺/⁺: -30 ± 1 mV, SD = 6.5 mV, n = 42 AZs vs NgR2⁻/⁻: -25.5 ± 0.98 mV, SD = 7.49 mV, n = 59 AZs; p = 0.002, Student's t-test) in NgR2⁻/⁻ IHCs. Shaded areas represent ± SEM. (**D**) Representative immunolabelings of presynaptic proteins show no apparent mislocalization in NgR2⁻/⁻ IHCs. Scale bar = 5 μm. Box-whisker plots show the median, 25/75 percentiles (box) and 10/90 percentiles (whiskers). Individual data points are overlaid.

NgR2 is essential for normal hearing

In order to evaluate auditory systems function in NgR2$^{-/-}$ mice, we recorded auditory brainstem responses (ABRs) at the age of 2-4 months. ABRs were elicited in response to 4, 8, 16, 32 kHz tone bursts and click stimuli of increasing sound pressure levels. We determined the sound thresholds by identifying the lowest sound pressure level which resulted in detectable ABR waveform. We found ABR thresholds to be significantly increased by around 30-45 dB across all the tested frequencies and in response to the click stimulations in NgR2$^{-/-}$ mice compared to the wild-type controls (Figure 6). NgR2$^{+/-}$ mice showed an intermediate phenotype between the NgR2$^{-/-}$ and NgR2$^{+/+}$ mice with a lower yet significant increase of ABR thresholds of approximately 10-15 dB at 4, 16 and 32 kHz frequencies.

Figure 6. Elevated acoustic thresholds in NgR2$^{-/-}$ mice. ABR thresholds were measured in response to 4, 8, 16, 32 kHz tone bursts and click stimuli. ABR thresholds of individual animals are shown in open circles on top of the mean ± SEM. Statistical significances are reported as *p < 0.05, ***p < 0.001, Kruskal-Wallis followed by Dunn's multiple comparison test.

Discussion

In this study, we investigated the role of the Nogo-66 receptor homolog NgR2 in the cochlea. Consistent with the previous reports, we detected NgR2 expression in IHCs and SGNs. Upon NgR2 deletion, we observed alterations of the IHC-SGN synapses (Figure 7). Presynaptically, Ca^{2+} channel activation was shifted towards depolarized voltages and ribbon size was enlarged. IHC exocytosis was unaltered when probed at saturating depolarizations. At the postsynaptic side, PSDs juxtaposing presynaptic ribbons were smaller compared to the controls and seemed to be deficient of GluA2/3, despite the expression of Gria2 mRNA in

NgR2$^{-/-}$ SGNs. Additionally, we observed PSDs which resided further from the IHC membrane

and did not juxtapose the presynaptic AZs in NgR2$^{-/-}$ mice. These PSDs potentially belong to

the type I SGN neurites that ended in the inner spiral bundle without contacting the IHCs, as

observed in SBEM reconstructions of NgR2$^{-/-}$ SGNs. Finally, ABR thresholds were elevated in

NgR2$^{-/-}$ mice indicating that NgR2 is required for normal hearing.

Figure 7. Schematic illustration of the key structural and functional changes in the auditory
periphery of NgR2$^{-/-}$ mice. NgR2$^{-/-}$ mice display enlarged synaptic ribbons and depolarized
shift in the activation of presynaptic Ca^{2+} channels in IHCs, as well as reduced size of PSDs
juxtaposing presynaptic ribbons. NgR2 deficiency further results in a subset of type I SGN
neurites that reach the inner spiral bundle but do not engage the IHCs.

The changes of presynaptic Ca^{2+} channel properties and ribbons in IHCs could result

from either i) a presynaptic action of NgR2 or ii) from NgR2 acting in the postsynaptic SGNs,

with an impact on transsynaptic signaling to IHCs. Presynaptic changes upon postsynaptic

manipulation have been reported previously. For instance, the deletion of postsynaptically

expressed Pou4f1 transcription factor resulted in hyperpolarized shift of IHC Ca^{2+} channels, as

well as in changes of the presynaptic heterogeneity of IHC AZs (Sherrill et al., 2019). Similarly,

presynaptic ribbon sizes were altered in IHCs of mice, which exhibited conditional deletion of

Runx1 transcription factor from SGNs (Shrestha et al., 2023).

We did not detect NgR2 near the synaptic regions but cannot exclude transient

synaptic localization of NgR2 during development. Indeed, NgR localization has been shown

to be dynamically regulated across different phases of IPSC-derived neuron development

(Wang et al., 2021). We propose that the afferent synaptic changes contribute to the severe

hearing impairment in NgR2$^{-/-}$ mice. The phenotype of increased ABR thresholds could

partially be explained by the depolarized activation of Ca^{2+} channels in IHCs. As it has been

shown, the apparent Ca^{2+} dependence of glutamate release from IHCs on average follows a

near linear relationship (Wong et al., 2014; Özçete and Moser, 2021; Jaime Tobón and Moser,

2023b, 2023a). Therefore, due to the depolarized shift of the Ca^{2+} channel activation, the

glutamate release from IHCs and subsequent spike generation in SGNs would require stronger

stimuli. Likewise, depolarized activation of Ca^{2+} channels has been associated with the higher

firing thresholds and low spontaneous rates in SGNs of Ribeye knock-out mice, albeit less

severe hearing phenotype was observed in those mice (Jean et al., 2018).

In addition or alternatively, the increase in auditory thresholds could be caused by the

deficiency of GluA2/3 AMPA receptors of the SGN postsynapses in NgR2$^{-/-}$ mice. We cannot

exclude that the lack of GluA2/3 subunits can be compensated by the overexpression of GluA4

subunit, which was also recently shown in GluA3 KO mice (Rutherford et al., 2023). Given the

observed synaptic changes, it seems less likely that the hearing impairment would have

resulted from the intrinsic changes of SGN firing. Further experiments, where SGNs would be

stimulated directly via *ex vivo* patch-clamp or optogenetically could help to probe for

functional deficits in SGN of NGR2$^{-/-}$ mice. Another interesting observation was the additional

PSDs not juxtaposing presynaptic ribbons that were detected around IHCs of NgR2$^{-/-}$ mice.

These "orphan" PSDs seem not to be part of the efferent synapses formed between

LOC/MOCs and SGNs as we verified by simultaneous anti-Homer1 and anti-synapsin

immunolabelings. To check the SGN connectivity we performed SBEM in NgR2$^{+/+}$ and NgR2$^{-/-}$

cochlear samples and observed a high percentage of non-synaptic SGN neurites in the

reconstructed neurons of inner spiral bundles in NgR2$^{-/-}$ samples. A sizable subset of the non-

synaptic SGN boutons did not make a contact with IHCs, potentially housing the "orphan"

PSDs observed around the IHCs in our immunolabeled samples. How exactly those non-

synaptic neurites appear remains to be investigated. SGN neurite retraction from IHCs has been demonstrated in noise-exposed animals, followed by the degeneration of the SGN somata and axons (Kujawa and Liberman, 2009, 2015; Moverman et al., 2023). This process however results in up to 50% degeneration of the ribbon synapses. Interestingly, in NgR2[-/-] mice the number of the ribbon synapses as well as the density of the SGN somata was largely intact. This indicates that the non-synaptic neurites do not result from synaptic loss as found in hereditary or acquired synaptopathy (Roux et al., 2006; Ruel et al., 2008; Seal et al., 2008; Kujawa and Liberman, 2009). Although it is plausible that the non-synaptic SGN neurites may fail to reach the IHCs during initial development, it is unlikely that this would selectively impact only a subset of neurons and still allow the Homer1-positive patches to develop in the absence of presynaptic components. Alternatively, it is known that major synaptic pruning takes place in developing SGNs, whereby up to 50% of the synapses are lost between the IHCs and the SGNs (Defourny et al., 2013; Coate et al., 2019), suggesting, that the non-synaptic SGN neurites of NgR2[-/-] mice could be a result of failed pruning of SGNs during the development. Yet how this would then comply with normal synapse and SGN counts remains to be elucidated. Future studies will be required to quantify the type I SGN projections of NgR2[-/-] mice during development and to trace their neurites from the soma to the organ of Corti to probe for potential branching in the spiral lamina.

Future work will be required to identify the NgR2 ligands in the cochlea. To this date, NgR interaction partners are not very well understood. While NgR1 in-trans interactions include NogoA, MAG, OMgp, chondroitin sulfate proteoglycans, the only well established ligand for NgR2 remains MAG (Fournier et al., 2001; Domeniconi et al., 2002; Wang et al., 2002; Venkatesh et al., 2005; Schwab, 2010). Other attractive candidate ligands for NgRs are the brain-specific angiotensin inhibitor (BAI) adhesion GPCRs. BAI-NgR interaction has been suggested to mediate dendritic arborization, axonal elongation, and synapse formation in IPSCs (Wang et al., 2021). Interestingly, a recent study described a hearing impairment in BAI1 deficient mice and suggests that BAI1 is important for correct localization of AMPA receptor subunits in spiral ganglion neurons based on the absence or decrease of different AMPA receptor subunits at the SGN postsynapses in BAI1 deficient mice (Carlton et al., 2024). Similarly, our data from NgR2[-/-] mice show the absense of GluA2/3 positive puncta juxtaposing presynaptic AZs. While BAI1 is suggested to act exclusively in SGNs (Carlton et al., 2024), NgR2 is expressed in both IHCs and SGNs. Along with the evidence for transsynaptic interaction

between NgRs and BAIs (Wang et al., 2021), this raises the possibility that the GluA2/3 AMPA

receptor subunit localization to SGN PSDs requires the interaction of presynaptic NgR2 and

postsynaptic BAI1. Alternatively, NgR2 expression in SGNs might be directly involved in

GluA2/3 tethering.

Finally, NgR2 has been proposed to interact with chondroitin-sulfate proteoglycan

Versican to control the amount of skin innervation by DRG neurites (Bäumer et al., 2014).

While to our knowledge, attempts have yet to be made for detecting Versican signal around

the IHCs, other extracellular matrix proteins such as aggrecan, brevican, Tenascin-R, Tenascin-

C, hyaluron and proteoglycan link proteins 1 and 4 have been shown to form dense, basket-

like structures that surround the base of the IHCs (Kwiatkowska et al., 2016; Sonntag et al.,

2018). Furthermore, deletion of brevican results in alteration of pre- and postsynaptic spatial

coupling at IHC synapses, indicating the possible role of extracellular matrix in transsynaptic

organization. However, while the existing observations suggest that the synaptic changes in

NgR2 deficient animals might reflect derailed interaction of the SGN neurites with the

extracellular matrix, the relevance of such an interaction of NgR2 and Versican in the cochlea

needs to be addressed in the future.

Materials and Methods

Animals

NgR2 null mutant mice on a C57BL/6N background were generated as previously described (Wörter et al., 2009). Knock-out (NgR2$^{-/-}$) and wild-type (NgR2$^{+/+}$) mice were derived from heterozygous matings. Both male and female mice were used in this study. Animals were genotyped as previously described (Wörter et al., 2009). The ages of the mice varied depending on the experiment as noted in the manuscript. The breeding and the experiments were approved by the Institutional Animal Care and Use Committee at the Medical University of Innsbruck, local Animal Welfare Committee of the University Medical Center Göttingen and the Max Planck Institute for Multidisciplinary Sciences, as well as the Animal Welfare Office of the state of Lower Saxony, Germany (LAVES, AZ: 19/3134). The ABR experiments were carried out under the approval of the Austrian Ministry of Education, Science and Research (reference number BMWFW-66.011/0120-WF/V/3b/2016).

Patch-clamp recordings

The apical turn of the organ of Corti was dissected from p21 to p29 animals in HEPES Hanks solution containing (in mM): 5.36 KCl, 141.7 NaCl, 10 HEPES, 0.5 MgSO$_4$, 1 MgCl$_2$, 5.6 D-glucose, and 3.4 L-glutamine (pH 7.2, ~ 300 mOsm/l). IHCs were exposed by gently removing of nearby supporting cells by negative pressure through a glass pipette from either modiolar or pillar side. All experiments were conducted at room temperature (RT, 20–25°C). Patch pipettes were made from GB150-8P or GB150F-8P borosilicate glass capillaries (Science Products, Hofheim, Germany) for perforated and ruptured patch-clamp configurations, respectively. Pipettes were fire polished with a custom-made microforge and coated with Sylgard to minimize the capacitive noise, whenever capacitance recordings were performed. The measurements were performed using EPC-10 amplifier controlled by Patchmaster software (HEKA Elektronik, Germany). The holding potential for IHCs was set to -87 mV across all the experiments.

Ruptured patch-clamp

For ruptured patch-clamp we used extracellular solution containing (in mM): 2.8 KCl, 105 NaCl, 10 HEPES, 1 CsCl, 1 $MgCl_2$, 5 $CaCl_2$, 35 TEA-Cl, and 2 mg/ml D-glucose (pH 7.2, $\sim$ 300 mOsm/l). Intracellular solution contained in mM: 111 Cs-glutamate, 1 $MgCl_2$, 1 $CaCl_2$, 10 EGTA, 13 TEA-Cl, 20 HEPES, 4 Mg-ATP, 0.3 Na-GTP and 1 L-glutathione (pH 7.3, $\sim$ 290 mOsm/l). Additionally, Ca^{2+} indicator Fluo4-FF (0.8 mM, Life Technologies) and TAMRA-conjugated RIBEYE/Ctbp2 binding peptide (10 mM, synthetized by the group of Dr. Olaf Jahn, Göttingen) were added to the intracellular solution for live imaging (Zenisek et al., 2004). Voltage dependency of Ca^{2+} influx was recorded by applying 20 ms long step depoarizations with 5 mV increment. For Ca^{2+} imaging, voltage ramp depolarizations ranging from -87 mV to 63 mV in the course of 150 ms were applied to the cells. Leak correction was performed using P/n protocol and liquid junction potential of 17 mV was corrected offline. Recordings were discarded from the analysis if the series resistance (R_s) exceeded 14 MOhm during the first 3 minutes after rupturing the cell, leak current exceeded -50 pA at the holding potential and Ca^{2+} current rundown was more than 25%.

Recordings were analyzed using Igor Pro 6.3 (Wavemetrics) custom-written programs. Ca^{2+} current-voltage relationships (IV curves) were obtained by averaging approximately 5 ms segments at the maximal activation regions of individual current traces and plotting them against the depolarization voltages. Fractional activation curves of the Ca^{2+} channels were calculated from the IVs and fitted with the Boltzmann function to determine the voltage of half maximal activation (V_{half}) and voltage sensitivity (k).

Perforated patch clamp recordings

To perform simultaneous membrane capacitance (C_m) and Ca^{2+} current measurements we used extracellular solution containing (in mM): 106 NaCl, 35 TEA-Cl, 2.8 KCl, 1 $MgCl_2$, 1 CsCl, 10 HEPES, 3 $CaCl_2$, and 5.6 D-glucose (pH 7.2, $\sim$ 300 mOsm/l). The pipette solution contained (in mM): 137 Cs-gluconate, 15 TEA-Cl, 3 GTP-Na_2, 1 ATP-Mg, 10 HEPES, 1 $MgCl_2$, as well as 300 mg/ml amphotericin B (pH 7.17, $\sim$ 290 mOsm/l). Perforated patch-clamp recordings from IHCs was described previously (Moser and Beutner, 2000). C_m changes were measured using Lindau-Neher technique (Lindau and Neher, 1988). To evoke exocytosis IHCs were stimulated by step depolarizations (to -17 mV) of different durations (2, 5, 10, 20, 50, 100 ms) in randomized manner. Currents were leak-corrected using a p/5 protocol. Recordings were used

only if the leak current was lower than 30 pA and the R_s was lower than 30 MOhm. Recordings

were analyzed using Igor Pro 6.3 (Wavemetrics) custom-written programs. Ca^{2+} charge ($Q_{Ca^{2+}}$)

was calculated by the time integral of the leak-subtracted Ca^{2+} current during the

depolarization step. ΔC_m was calculated as the difference between the average C_m before and

after the depolarization. To measure average C_m we used 400 ms segments and skipped the

initial 100 ms after the depolarization.

Ca^{2+} imaging

Ca^{2+} imaging was performed using spinning disc confocal microscope, as described before

(Ohn et al., 2016). Briefly, the set-up was equipped with a spinning disc confocal unit (CSU22,

Yokogawa) mounted on an upright microscope (Axio Examiner, Zeiss) and scientific CMOS

camera (Andor Neo). Images were acquired using 63x, 1.0 NA objective (W Plan-Apochromat,

Zeiss). The pixel size was measured to be 103 nm.

IHCs were loaded with Fluo4-FF Ca^{2+} dye and TAMRA-conjugated Ctbp2 binding dimeric

peptide via the patch pipette. First, the cells were scanned from bottom to top by imaging

TAMRA fluorescence with 561 nm laser (Jive, Cobolt AB) and exposing each plane for 0.5

seconds. The stack was acquired with 0.5 μm step size using Piezo positioner (Piezosystem).

This allowed us to identify the planes containing synaptic ribbons. Next, we recorded Fluo4-

FF fluorescence increase at individual synapses by imaging ribbon containing planes with 491

nm laser (Calypso, Cobolt AB) at 100 Hz while applying voltage ramp depolarizations to the

cell. 2 voltage ramps were applied at each plane, one being 5 ms shifted relative to the other.

Spinning disk rotation speed was set to 2000 rpm, in order to synchronize with Ca^{2+} imaging.

Images were analyzed using Igor Pro 6.3 (Wavemetrics). Ca^{2+} hotspots were identified by

subtracting the average signal of several baseline frames from the average signal of 5 frames

during stimulation (ΔF image). Since the same Ca^{2+} hotspot appears across multiple planes,

the plane exhibiting the strongest signal was chosen. The intensities of the 3x3 matrix

surrounding the central pixel of the hotspot were averaged across all time points to obtain the

intensity profiles of Ca^{2+} influx over time. Afterwards, the background signal (average of

approximately 60x60 pixel intensities outside the cell) was subtracted from the intensity-time

profiles and $\Delta F/F_0$ traces were calculated. To enhance voltage resolution of Ca^{2+} imaging, two

$\Delta F/F_0$ traces corresponding to two voltage ramp depolarizations (one shifted by 5 ms over the

other) were combined and plotted against the corresponding voltages (FV curves). These curves were subsequently fitted with a modified Boltzmann function. Afterwards, fractional activation curves were calculated by fitting the linear decay of the fluorescence signal from the FV curves with a linear function (G_{max}) and then dividing the FV fit by the G_{max} line. The resulting curves were further fitted with a Boltzmann function. Maximal Ca^{2+} influx ($\Delta F/F_{0\ max}$) was calculated by averaging 5 points during the stimulation.

Immunohistochemistry

Amplified immunohistochemistry for NgR2 was performed on 5 μm dewaxed paraffin sections of NgR2$^{+/+}$ and NgR2$^{-/-}$ inner ears from either sex. All tissue sections were incubated in 10 mM sodium citrate buffer (pH = 6) at 98^{o}C for 10 min, washed in PBS, and endogenous peroxidase quenched using 3% H_2O_2, in 1X PBS for 10 min. Tissue was blocked in 2% normal goat serum, 1% BSA and 0.3% Triton-X100 in PBS for 2 hours at RT, followed by overnight incubation with primary rabbit anti-NgR2/RTN4rl2 (1:400, Invitrogen, PA598577) at 4^{o}C. Biotin-labelled goat anti-rabbit (2 hours at RT; 1:500, Ba-1000, Vector Laboratories) and the ImmPACT DAB Kit (SK-4105, Vector Laboratories) were used for signal detection. After several washes, sections were incubated with TSA plus Cyanine-3 reagent (1:2000, NEL760001KT, Perkin Elmer) for 10 to 15 min at RT in accordance with the manufacturer's protocol. Subsequently, sections were stained overnight with mouse anti-βIII-tubulin (1:1000, 801202, Biolegend) at 4^{o}C, followed by secondary Alexa Fluor 488 conjugated goat anti-mouse (1:1000, Thermo Fisher Scientific), for 2 hours at RT. Finally, sections were cover slipped in Mowiol and stored at 4^{o}C until imaging. For whole mount immunofluorescence, cochleae were fixed in 4% formaldehyde on ice for 45-60 minutes or sometimes overnight. For anti-Ca$_V$1.3 stainings the cochleae were fixed for 10 minutes. After the fixation, the organs of Corti were microdissected in PBS and blocked in GSDB (goat serum dilution buffer; 16% goat serum, 20 mM phosphate buffer (PB), 0.3% Triton X-100, 0.45 M NaCl) for 1 hour at RT. The samples were then incubated in the primary antibody mixture overnight at 4^{o}C. The next day, samples were washed 3 times using wash buffer (20 mM PB, 0.3% Triton X-100, 0.45 M NaCl) followed by incubation in the secondary antibody mixture for 1 hour at RT. Afterwards, the samples were washed 3 times in the washing buffer, one time in PB and mounted using mounting medium (Mowiol).

The following primary antibodies were used: rabbit anti-Homer1 (1:500, 160 002, Synaptic

Systems), mouse anti-Ctbp2 (1:200, 612044, BD Biosciences), rabbit anti-Ca$_V$1.3 (1:100, ACC-

005, Alomone Labs), mouse anti-Bassoon (1:300, ab82958, Abcam), rabbit anti-Myosin7a

(1:800, ab3481, Abcam, or Proteus BioScience), guinea pig anti-RibeyeA (1:500, 192104,

Synaptic Systems), guinea pig anti-Synapsin1/2 (1:500, 106004, Synaptic Systems), guinea pig

anti-Vglut3 (1:500, 135204, Synaptic Systems), chicken anti-parvalbumin (1:200, 195006,

Synaptic Systems), chicken anti-calretinin (1:200, 214 106, Synaptic Systems), guinea pig anti-

VAChT (1:1000, 139105, Synaptic Systems), mouse anti-ATP1A3 (1:300, MA3-915, Thermo

Fisher Scientific).

The following secondary antibodies were used: Alexa Fluor 488 conjugated anti-guinea pig

(1:200, A11073, Thermo Fisher Scientific), Alexa Fluor 488 conjugated anti-rabbit (1:200,

A11008, Thermo Fisher Scientific), Alexa Fluor 568 conjugated anti-chicken (1:200, ab175711,

Abcam), Alexa Fluor 633 conjugated anti-guinea pig (1:200, A21105, Thermo Fisher Scientific),

STAR 580 conjugated anti-mouse (1:200, Abberior, ST635P-1001-500UG), STAR 635

conjugated anti-rabbit (1:200, Abberior, ST635P-1002-500UG).

Confocal stacks were acquired using Leica SP8 confocal microscope or Abberior Instruments

Expert Line STED microscope.

The volumes of the Ctbp2 and Homer1 positive puncta were estimated using the surface

algorithm of Imaris software (version 9.6.0, Bitplane). Following parameters were used to

create the surfaces for both Ctbp2 and Homer1 puncta in all the analyzed stacks: surface detail

0.07 μm, background subtraction 0.562 μm, touching object size 0.4 μm.

The brightness and the contrast of the representative images were adjusted using Fiji (ImageJ)

software for the visualization purposes.

Spiral ganglion neuron densities

NgR2$^{+/+}$ and NgR2$^{-/-}$ cochlea from p40 mice of either sex were processed for SGN counts.

Mouse monoclonal anti-βIII-tubulin (1:1000, 801202, Biolegend) at 4^{o}C, followed by

secondary Alexa Fluor 488 conjugated anti-mouse (1:1000, Thermo Fisher Scientific), was

used to label all SGNs. Nuclei were visualized using a DAPI fluorescent counterstain (1:1000,

Life Technologies). 20X images of mid-modiolar cochlear sections were collected. ImageJ was

used to outline the spiral ganglion and generate the area. Quantitative assessment was

performed on every other mid-modiolar section to reduce chances of double counting.

Labeled cells were counted only if they had a round cell body, presence of nucleus, and

homogenous cytoplasm. Densities were calculated and statistical differences were measure

using a Student's t-test (GraphPad Prism, GraphPad Software Inc.).

Inner and outer Hair cell counts
554

IHC and OHC counts were performed for NgR2$^{+/+}$ and NgR2$^{-/-}$ mice at the ages of p15, 1 month

and 2 months. Images were imported to ImageJ software, where IHC/OHC counts were

performed blinded to genotype using the 'Cell counter' tool. To quantify the cells the nuclear

staining of the DAPI channel was used. The number of IHCs and OHCs was assessed along a

length of 100 μm for each image, and then averaged across all samples to obtain the average

number of IHCs (or OHCs) per 100 μm.

RNAscope
563

RNA *in-situ* hybridization was performed using RNAscope$^{®}$ (Advanced Cell Diagnostics).

Standard mouse probes were used to examine the expression of NgR2/RTN4rl2 (450761, ACD)

and Gria2 (865091, ACD) closely following the manufacturer's instructions. Briefly, 6 mm

paraffin tissue sections underwent deparaffinization with xylene and a series of ethanol

washes. Tissues were heated in kit-provided antigen retrieval buffer and digested by kit-

provided proteinase. Sections were exposed to mFISH target probes and incubated at 40°C in

a hybridization oven for 2 hours. After rinsing, mFISH signal was amplified using company-

provided pre-amplifier and amplifier conjugated to fluorescent dye. Subsequently, sections

were blocked with 1% BSA, 2% normal goat serum in 1xPBS containing 0.3% Triton X-100 for

1 hour at RT. The tissue was incubated in mouse anti-βIII-tubulin antibody (1:500, 801202,

Biolegend) overnight at 4°C. The next day, sections were rinsed with PBS, blocked again before

incubating in secondary Alexa Fluor 488 conjugated anti-mouse antibody (1:1000, Invitrogen)

for 2 hours at RT. Sections were counterstained with DAPI (1:1000, Life Technologies),

mounted, and stored at 4°C until image analysis. mFISH images were captured on a Leica SP8

confocal microscope and processed using ImageJ.

Serial block-face scanning electron microscopy (SBEM)

Sample preparation for SBEM

One NgR2[+/+] and two NgR2[-/-] female mice at p36 were used for the SBEM experiments. Cochleae were processed for SBEM imaging as previously described (Hua et al., 2021). In brief, the animals were decapitated after CO_2 inhalation under anesthesia. The cochleae were dissected from the skulls and immediately fixed by perfusing with ice-cold fixative through the round and oval windows at constant flow speed using an infusion pump (Micro4, WPI). The fixative solution was freshly prepared and made of 4% paraformaldehyde (Sigma-Aldrich), 2.5% glutaraldehyde (Sigma-Aldrich), and 0.08 M cacodylate (pH 7.4, Sigma-Aldrich). After being immersed in the fixative at 4°C for 5 hours, the cochleae were transferred to a decalcifying solution containing the same fixative and 5% ethylenediaminetetraacetic acid (EDTA, Serva) and incubated at 4°C for 5 hours. The samples were then washed twice with 0.15 M cacodylate (pH 7.4) for 30 min each, sequentially immersed in 2% OsO_4 (Sigma-Aldrich), 2.5% ferrocyanide (Sigma-Aldrich), and again 2% OsO_4 at RT for 2, 2, and 1.5 hours. After being washed in 0.15 M cacodylate and distilled water (Sartorius) for 30 min each, the samples were sequentially incubated in filtered 1% thiocarbohydrazide (TCH, Sigma-Aldrich) solution and 2% OsO_4 at RT for 1 and 2 hours, as well as in lead aspartate solution (0.03 M, pH 5.0, adjusted with KOH) at 50 °C for 2 hours with immediate two washing steps with distilled water at RT for 30 min each. For embedding, the samples were dehydrated through graded pre-cooled acetone (Carl Roth) series (50%, 75%, 90%, for 30 min each, all cooled at 4 °C) and then pure acetone at RT (three times for 30 min each). The sample infiltration started with 1:1 and 1:2 mixtures of acetone and Spurr's resin monomer (4.1 g ERL 4221, 0.95 g DER 736, 5.9 g NSA and 1% DMAE; Sigma-Aldrich) at RT for 6 and 12 hours on a rotator. After being impregnated in pure resin for 12 hours, the samples were placed in embedding molds (Polyscience) and hardened in a pre-warmed oven at 70°C for 72 hours.

Sample trimming and SBEM imaging

The sample blocks were mounted upright along the conical center axis on aluminum metal rivets (3VMRS12, Gatan, UK) and trimmed coronally towards the modiolus using a diamond trimmer (TRIM2, Leica, Germany). For each sample, a block face of about ~ 600 x 800 mm²,

centered at the apical segment based on the anatomical landmarks, was created using an
ultramicrotome (UC7, Leica, Germany). The samples were coated with a 30 nm thick gold layer
using a sputter coater (ACE600, Leica, Germany). The serial images were acquired using a field-
emission scanning EM (Gemini300, Carl Zeiss, Germany) equipped with an in-chamber
ultramicrotome (3ViewXP, Gatan, UK) and back-scattered electron detector (Onpoint, Gatan,
UK). Focal charge compensation was set to 100 % with a high vacuum chamber pressure of
2.8×10^3 mbar. The following parameters were set for the SBEM imaging: 12 nm pixel size, 50
nm nominal cutting thickness, 2 keV incident beam energy, and 1.5 ms pixel dwell time.

For the NgR2$^{+/+}$ dataset, 2377 consecutive slices (9000 × 15000 pixels) were collected,
whereas the two NgR2$^{-/-}$ datasets had 3217 slices (16000 × 10000 pixels) and 2425 slices (9000
× 15000 pixels). All datasets were aligned along the z-direction using a custom MATLAB script
based on cross-correlation maximum between consecutive slices (Hua et al., 2022) before
being uploaded to webKnossos (Boergens et al., 2017) for skeleton and volume tracing.

627
628 Identification and quantification of auditory afferent fibers
629
In our SBEM datasets of the cochlea, manual skeleton tracing was carried out on all neurites
that originated from three neighboring habenula perforata (HP) at the center of each dataset.
To search for type I afferent fibers, several morphological features were used, such as
myelination after entering HP, radial and unbranched fiber trajectory, contact with IHCs, as
well as ribbon-associated terminals (Hua et al., 2021). This resulted in 115 putative type I
afferent fibers and further classification was made based on the presence of presynaptic
ribbon, fiber branching, and contact with IHC. For an illustration purpose, a type I afferent
fiber bundle of one NgR2$^{-/-}$ dataset was volume traced and 3D rendered using Amira software
(Thermo Scientific, US).

639
640 Ribbon size measurement and synapse counting
641
642 Ribbon-type synapses were manually annotated in 18 intact IHCs captured by SBEM using
643 webKnossos. The dense core region of individual ribbon synapses was manually contoured,
644 and the associated voxels were counted for ribbon volume measurement. In the case of multi-
645 ribbon synapses, all ribbon bodies at a single active zone were summed up to yield the ribbon
646 volume.

Auditory brainstem responses

Auditory brainstem resposes (ABR) were recorded in 2-4 months old mice of either sex, as described previously (Luque et al., 2021). Briefly, we used a custom-made system in an anechoic chamber in a calibrated open-field configuration. ABRs were recorded via needle electrodes in response to tone bursts of 4, 8, 16, and 32 kHz or a click wide spectrum (2-45,2 KHz, 2 KHz steps) as stimuli. Tone pips of 3 ms duration (1 ms rise and fall time) were presented at a rate of 60/s with alternating phases. Starting with 0 dB, the stimuli were increased in 5 dB steps up to 120 dB. Hearing thresholds were determined as the minimum stimulation level that produced a clearly recognizable potential. Recordings were evaluated by three independent researchers in a blinded manner.

Data analysis and statistics

The data were analyzed using Igor Pro (Wavemetrics), Python and GraphPad Prism (GraphPad Software Inc.) software. For 2 sample comparisons, data were tested for normality and equality of variances using Jarque-Bera and F-test, respectively. Afterwards, two-tailed Student's t-test or Mann-Whitney-Wilcoxon test were performed. The latter was used when normality and/or equality of variances were not met. P-values were corrected for multiple comparisons using Holm-Šídák method. For ABR thresholds Kruskal-Wallis test followed by Dunn's multiple comparison test was used. Data is presented as mean ± standard error of the mean (SEM), unless otherwise stated. Significances are reported as $*p < 0.05$, $**p < 0.01$, $***p < 0.001$. The number of the animals is indicated as N.

Acknowledgements

We thank Sina Langer, Christiane Senger-Freitag, Sandra Gerke for expert technical support. We thank Prof. Dr. Carolin Wichmann, Dr. Susann Michanski, Julius Bahr and Sophia Mutschall for the help with the SBEM sample preparation. We thank Dr. Yi Jiang and Haoyu Wang for the visualization of neurite reconstruction. We would also like to thank Prof. Dr. Olaf Jahn and Lars van Werven for the TAMRA-conjugated Ctbp2 binding peptide synthesis. The study was supported by German Research Foundation through the Cluster of Excellence (EXC2067) Multiscale Bioimaging (TM) and the Leibniz Program (MO896/5 to TM) as well as by the

European Research Council through the Advanced Grant "DynaHear" to TM under the
European Union's Horizon 2020 Research and Innovation program (grant agreement No.
101054467), and by Fondation Pour l'Audition (FPA RD-2020-10), by the SPIN-FWF grant to
CB, and by the National Natural Science Foundation of China (82171133 to YH), Industrial
Support Fund of Huangpu District .in Shanghai (XK2019011 to YH), Innovative Research Team
of High-level Local Universities in Shanghai (SHSMU-ZLCX20211700). NK is a member of the
Hertha Sponer College from the Cluster of Excellence Multiscale Bioimaging (MBExC). TM is a
Max-Planck Fellow at the Max Planck Institute for Multidisciplinary Sciences.
688

689   ## Author contributions
690
691   CB, TM, YH, and NK designed the research. NK, MÜ, YQ, NB, FH, LJC, FW, ML, RG performed
692   experiments and analyzed data. NK and TM wrote the paper with contributions of YH, CB, YQ,
693   NB. CB, TM, YH acquired funding. All authors have reviewed and approved the final version of
694   the manuscript for submission.

References

Appler, J. M., and Goodrich, L. V. (2011). Connecting the ear to the brain: Molecular mechanisms of auditory circuit assembly. *Prog. Neurobiol.* 93, 488–508. doi: 10.1016/j.pneurobio.2011.01.004

Bäumer, B. E., Kurz, A., Borrie, S. C., Sickinger, S., Dours-Zimmermann, M. T., Zimmermann, D. R., et al. (2014). Nogo receptor homolog NgR2 expressed in sensory DRG neurons controls epidermal innervation by interaction with Versican. *J. Neurosci. Off. J. Soc. Neurosci.* 34, 1633–1646. doi: 10.1523/JNEUROSCI.3094-13.2014

Boergens, K. M., Berning, M., Bocklisch, T., Bräunlein, D., Drawitsch, F., Frohnhofen, J., et al. (2017). webKnossos: efficient online 3D data annotation for connectomics. *Nat. Methods* 14, 691–694. doi: 10.1038/nmeth.4331

Borrie, S. C., Sartori, S. B., Lehmann, J., Sah, A., Singewald, N., and Bandtlow, C. E. (2014). Loss of Nogo receptor homolog NgR2 alters spine morphology of CA1 neurons and emotionality in adult mice. *Front. Behav. Neurosci.* 8. Available at: https://www.frontiersin.org/articles/10.3389/fnbeh.2014.00175 (Accessed March 1, 2024).

Bulankina, A. V., and Moser, T. (2012). Neural circuit development in the mammalian cochlea. *Physiology* 27, 100–112. doi: 10.1152/physiol.00036.2011

Cantu-Guerra, H. L., Papazian, M. R., Gorsky, A. L., Alekos, N. S., Caccavano, A., Karagulyan, N., et al. (2023). Cochlear hair cell innervation is dependent on a modulatory function of Semaphorin-3A. *Dev. Dyn.* 252, 124–144. doi: 10.1002/dvdy.548

Carlton, A. J., Jeng, J.-Y., Grandi, F. C., De Faveri, F., Amariutei, A. E., De Tomasi, L., et al. (2024). BAI1 localizes AMPA receptors at the cochlear afferent post-synaptic density and is essential for hearing. *Cell Rep.* 43, 114025. doi: 10.1016/j.celrep.2024.114025

Coate, T. M., Scott, M. K., and Gurjar, M. (2019). Current concepts in cochlear ribbon synapse formation. *Synapse* 73, e22087. doi: 10.1002/syn.22087

Coate, T. M., Spita, N. A., Zhang, K. D., Isgrig, K. T., and Kelley, M. W. (2015). Neuropilin-2/Semaphorin-3F-mediated repulsion promotes inner hair cell innervation by spiral ganglion neurons. *eLife* 4. doi: 10.7554/eLife.07830

Defourny, J., Poirrier, A.-L., Lallemend, F., Mateo Sánchez, S., Neef, J., Vanderhaeghen, P., et al. (2013). Ephrin-A5/EphA4 signalling controls specific afferent targeting to cochlear hair cells. *Nat. Commun.* 4, 1438. doi: 10.1038/ncomms2445

Domeniconi, M., Cao, Z., Spencer, T., Sivasankaran, R., Wang, K. C., Nikulina, E., et al. (2002). Myelin-Associated Glycoprotein Interacts with the Nogo66 Receptor to Inhibit Neurite Outgrowth. *Neuron* 35, 283–290. doi: 10.1016/S0896-6273(02)00770-5

Elkon, R., Milon, B., Morrison, L., Shah, M., Vijayakumar, S., Racherla, M., et al. (2015). RFX
transcription factors are essential for hearing in mice. *Nat. Commun.* 6, 8549. doi:
10.1038/ncomms9549

Fell, B., Eckrich, Stephanie, Hecker, Dietmar, Obermair, Gerald J., Flockerzi, V., Schick, B., et
al. (2015). α2δ2 Controls the Function and Transsynaptic Coupling of Cav1.3 Channels
in Mouse Inner Hair Cells and is Essential for Normal Hearing. *ARO*
Midwintermeeting.

Fournier, A. E., GrandPre, T., and Strittmatter, S. M. (2001). Identification of a receptor
mediating Nogo-66 inhibition of axonal regeneration. *Nature* 409, 341–346. doi:
10.1038/35053072

Hua, Y., Ding, X., Wang, H., Wang, F., Lu, Y., Neef, J., et al. (2021). Electron Microscopic
Reconstruction of Neural Circuitry in the Cochlea. *Cell Rep.* 34, 108551. doi:
10.1016/j.celrep.2020.108551

Hua, Y., Loomba, S., Pawlak, V., Voit, K.-M., Laserstein, P., Boergens, K. M., et al. (2022).
Connectomic analysis of thalamus-driven disinhibition in cortical layer 4. *Cell Rep.* 41,
111476. doi: 10.1016/j.celrep.2022.111476

Jaime Tobón, L. M., and Moser, T. (2023a). Bridging the gap between presynaptic hair cell
function and neural sound encoding. doi: 10.7554/eLife.93749.1

Jaime Tobón, L. M., and Moser, T. (2023b). Ca2+ regulation of glutamate release from inner
hair cells of hearing mice. *Proc. Natl. Acad. Sci.* 120, e2311539120. doi:
10.1073/pnas.2311539120

Jean, P., Morena, D. L. de la, Michanski, S., Tobón, L. M. J., Chakrabarti, R., Picher, M. M., et
al. (2018). The synaptic ribbon is critical for sound encoding at high rates and with
temporal precision. *eLife* 7, e29275. doi: 10.7554/eLife.29275

Jean, P., Wong Jun Tai, F., Singh-Estivalet, A., Lelli, A., Scandola, C., Megharba, S., et al.
(2023). Single-cell transcriptomic profiling of the mouse cochlea: An atlas for
targeted therapies. *Proc. Natl. Acad. Sci.* 120, e2221744120. doi:
10.1073/pnas.2221744120

Johnson, S. L., Safieddine, S., Mustapha, M., and Marcotti, W. (2019). Hair Cell Afferent
Synapses: Function and Dysfunction. *Cold Spring Harb. Perspect. Med.* 9. doi:
10.1101/cshperspect.a033175

Jukic, A., Lei, Z., Cebul, E. R., Pinter, K., Mosqueda, N., David, S., et al. (2024). Presynaptic
Nrxn3 is essential for ribbon-synapse assembly in hair cells. doi:
10.1101/2024.02.14.580267

Jung, J. S., Zhang, K. D., Wang, Z., McMurray, M., Tkaczuk, A., Ogawa, Y., et al. (2019).
Semaphorin-5B Controls Spiral Ganglion Neuron Branch Refinement during
Development. *J. Neurosci.* 39, 6425–6438. doi: 10.1523/JNEUROSCI.0113-19.2019

Koundakjian, E. J., Appler, J. L., and Goodrich, L. V. (2007). Auditory neurons make
stereotyped wiring decisions before maturation of their targets. *J. Neurosci. Off. J.*
Soc. Neurosci. 27, 14078–14088. doi: 10.1523/JNEUROSCI.3765-07.2007

Kujawa, S. G., and Liberman, M. C. (2009). Adding insult to injury: cochlear nerve
degeneration after "temporary" noise-induced hearing loss. *J. Neurosci.* 29, 14077.

Kujawa, S. G., and Liberman, M. C. (2015). Synaptopathy in the noise-exposed and aging
cochlea: Primary neural degeneration in acquired sensorineural hearing loss. *Hear.*
Res. 330, 191–199. doi: 10.1016/j.heares.2015.02.009

Kwiatkowska, M., Reinhard, J., Roll, L., Kraft, N., Dazert, S., Faissner, A., et al. (2016). The
expression pattern and inhibitory influence of Tenascin-C on the growth of spiral
ganglion neurons suggest a regulatory role as boundary formation molecule in the
postnatal mouse inner ear. *Neuroscience* 319, 46–58. doi:
10.1016/j.neuroscience.2016.01.039

Liberman, L. D., and Liberman, M. C. (2016). Postnatal maturation of auditory-nerve
heterogeneity, as seen in spatial gradients of synapse morphology in the inner hair
cell area. *Hear. Res.* 339, 12–22. doi: 10.1016/j.heares.2016.06.002

Lindau, M., and Neher, E. (1988). Patch-clamp techniques for time-resolved capacitance
measurements in single cells. *Pflüg. Arch. Eur. J. Physiol.* 411, 137–146.

Liu, H., Chen, L., Giffen, K. P., Stringham, S. T., Li, Y., Judge, P. D., et al. (2018). Cell-Specific
Transcriptome Analysis Shows That Adult Pillar and Deiters' Cells Express Genes
Encoding Machinery for Specializations of Cochlear Hair Cells. *Front. Mol. Neurosci.*
11. doi: 10.3389/fnmol.2018.00356

Lu, Y., Liu, J., Li, B., Wang, H., Wang, F., Wang, S., et al. (2024). Spatial patterns of noise-
induced inner hair cell ribbon loss in the mouse mid-cochlea. *iScience* 27, 108825.
doi: 10.1016/j.isci.2024.108825

Luque, M., Schrott-Fischer, A., Dudas, J., Pechriggl, E., Brenner, E., Rask-Andersen, H., et al.
(2021). HCN channels in the mammalian cochlea: Expression pattern, subcellular
location, and age-dependent changes. *J. Neurosci. Res.* 99, 699–728. doi:
10.1002/jnr.24754

Matsubara, A., Laake, J. H., Davanger, S., Usami, S., and Ottersen, O. P. (1996). Organization
of AMPA receptor subunits at a glutamate synapse: a quantitative immunogold
analysis of hair cell synapses in the rat organ of Corti. *J. Neurosci.* 16, 4457.

Meyer, A. C., and Moser, T. (2010). Structure and function of cochlear afferent innervation.
Curr. Opin. Otolaryngol. Head Neck Surg. 18, 441–446. doi:
10.1097/MOO.0b013e32833e0586

Michanski, S., Smaluch, K., Steyer, A. M., Chakrabarti, R., Setz, C., Oestreicher, D., et al.
(2019). Mapping developmental maturation of inner hair cell ribbon synapses in the

apical mouse cochlea. *Proc. Natl. Acad. Sci.* 116, 6415–6424. doi:
10.1073/pnas.1812029116

Mironova, Y. A., and Giger, R. J. (2013). Where no synapses go: gatekeepers of circuit
remodeling and synaptic strength. *Trends Neurosci.* 36, 363–373. doi:
10.1016/j.tins.2013.04.003

Moser, T., and Beutner, D. (2000). Kinetics of exocytosis and endocytosis at the cochlear
inner hair cell afferent synapse of the mouse. *Proc. Natl. Acad. Sci.* 97, 883–888. doi:
10.1073/pnas.97.2.883

Moverman, D. J., Liberman, L. D., Kraemer, S., Corfas, G., and Liberman, M. C. (2023).
Ultrastructure of noise-induced cochlear synaptopathy. *Sci. Rep.* 13, 19456. doi:
10.1038/s41598-023-46859-6

Niedzielski, A. S., and Wenthold, R. J. (1995). Expression of AMPA, kainate, and NMDA
receptor subunits in cochlear and vestibular ganglia. *J. Neurosci.* 15, 2338.

Ohn, T.-L., Rutherford, M. A., Jing, Z., Jung, S., Duque-Afonso, C. J., Hoch, G., et al. (2016).
Hair cells use active zones with different voltage dependence of Ca^{2+} influx to
decompose sounds into complementary neural codes. *Proc. Natl. Acad. Sci.* 113,
E4716–E4725. doi: 10.1073/pnas.1605737113

Özçete, Ö. D., and Moser, T. (2021). A sensory cell diversifies its output by varying Ca^{2+} influx-
release coupling among active zones. *EMBO J.* 40, e106010. doi:
10.15252/embj.2020106010

Ramirez, M. A., Ninoyu, Y., Miller, C., Andrade, L. R., Edassery, S., Bomba-Warczak, E., et al.
(2022). Cochlear ribbon synapse maturation requires Nlgn1 and Nlgn3. *iScience* 25,
104803. doi: 10.1016/j.isci.2022.104803

Roux, I., Safieddine, S., Nouvian, R., Grati, M., Simmler, M.-C., Bahloul, A., et al. (2006).
Otoferlin, Defective in a Human Deafness Form, Is Essential for Exocytosis at the
Auditory Ribbon Synapse. *Cell* 127, 277–289. doi: 10.1016/j.cell.2006.08.040

Ruel, J., Emery, S., Nouvian, R., Bersot, T., Amilhon, B., Van Rybroek, J. M., et al. (2008).
Impairment of SLC17A8 Encoding Vesicular Glutamate Transporter-3, VGLUT3,
Underlies Nonsyndromic Deafness DFNA25 and Inner Hair Cell Dysfunction in Null
Mice. *Am. J. Hum. Genet.* 83, 278–292. doi: 10.1016/j.ajhg.2008.07.008

Rutherford, M. A., Bhattacharyya, A., Xiao, M., Cai, H.-M., Pal, I., and Rubio, M. E. (2023).
GluA3 subunits are required for appropriate assembly of AMPAR GluA2 and GluA4
subunits on cochlear afferent synapses and for presynaptic ribbon modiolar-pillar
morphology. *eLife* 12, e80950. doi: 10.7554/eLife.80950

Schwab, M. E. (2010). Functions of Nogo proteins and their receptors in the nervous system.
Nat. Rev. Neurosci. 11, 799–811. doi: 10.1038/nrn2936

Seal, R. P., Akil, O., Yi, E., Weber, C. M., Grant, L., Yoo, J., et al. (2008). Sensorineural Deafness
and Seizures in Mice Lacking Vesicular Glutamate Transporter 3. *Neuron* 57, 263–275.
doi: 10.1016/j.neuron.2007.11.032

Sherrill, H. E., Jean, P., Driver, E. C., Sanders, T. R., Fitzgerald, T. S., Moser, T., et al. (2019).
Pou4f1 Defines a Subgroup of Type I Spiral Ganglion Neurons and Is Necessary for
Normal Inner Hair Cell Presynaptic Ca2+ Signaling. *J. Neurosci.* 39, 5284–5298. doi:
10.1523/JNEUROSCI.2728-18.2019

Shrestha, B. R., Chia, C., Wu, L., Kujawa, S. G., Liberman, M. C., and Goodrich, L. V. (2018).
Sensory Neuron Diversity in the Inner Ear Is Shaped by Activity. *Cell* 174, 1229-
1246.e17. doi: 10.1016/j.cell.2018.07.007

Shrestha, B. R., Wu, L., and Goodrich, L. V. (2023). Runx1 controls auditory sensory neuron
diversity in mice. *Dev. Cell* 58, 306-319.e5. doi: 10.1016/j.devcel.2023.01.008

Sonntag, M., Blosa, M., Schmidt, S., Reimann, K., Blum, K., Eckrich, T., et al. (2018). Synaptic
coupling of inner ear sensory cells is controlled by brevican-based extracellular
matrix baskets resembling perineuronal nets. *BMC Biol.* 16, 99. doi: 10.1186/s12915-
018-0566-8

Vaccaro, G., Dumoulin, A., Zuñiga, N. R., Bandtlow, C. E., and Stoeckli, E. T. (2022). The Nogo-
66 Receptors NgR1 and NgR3 Are Required for Commissural Axon Pathfinding. *J.
Neurosci.* 42, 4087–4100. doi: 10.1523/JNEUROSCI.1390-21.2022

Venkatesh, K., Chivatakarn, O., Lee, H., Joshi, P. S., Kantor, D. B., Newman, B. A., et al. (2005).
The Nogo-66 Receptor Homolog NgR2 Is a Sialic Acid-Dependent Receptor Selective
for Myelin-Associated Glycoprotein. *J. Neurosci.* 25, 808–822. doi:
10.1523/JNEUROSCI.4464-04.2005

Wang, J., Miao, Y., Wicklein, R., Sun, Z., Wang, J., Jude, K. M., et al. (2021). RTN4/NoGo-
receptor binding to BAI adhesion-GPCRs regulates neuronal development. *Cell* 184,
5869-5885.e25. doi: 10.1016/j.cell.2021.10.016

Wang, K. C., Koprivica, V., Kim, J. A., Sivasankaran, R., Guo, Y., Neve, R. L., et al. (2002).
Oligodendrocyte-myelin glycoprotein is a Nogo receptor ligand that inhibits neurite
outgrowth. *Nature* 417, 941–944. doi: 10.1038/nature00867

Wills, Z. P., Mandel-Brehm, C., Mardinly, A. R., McCord, A. E., Giger, R. J., and Greenberg, M.
E. (2012). The Nogo Receptor Family Restricts Synapse Number in the Developing
Hippocampus. *Neuron* 73, 466–481. doi: 10.1016/j.neuron.2011.11.029

Wong, A. B., Rutherford, M. A., Gabrielaitis, M., Pangršič, T., Göttfert, F., Frank, T., et al.
(2014). Developmental refinement of hair cell synapses tightens the coupling of Ca2+
influx to exocytosis. *EMBO J.* 33, 247–264. doi: 10.1002/embj.201387110

Wörter, V., Schweigreiter, R., Kinzel, B., Mueller, M., Barske, C., Böck, G., et al. (2009).
Inhibitory activity of myelin-associated glycoprotein on sensory neurons is largely

independent of NgR1 and NgR2 and resides within Ig-Like domains 4 and 5. *PloS One*
4, e5218. doi: 10.1371/journal.pone.0005218

Zenisek, D., Horst, N. K., Merrifield, C., Sterling, P., and Matthews, G. (2004). Visualizing
synaptic ribbons in the living cell. *J. Neurosci. Off. J. Soc. Neurosci.* 24, 9752–9759.
doi: 10.1523/JNEUROSCI.2886-04.2004

Supplementary Figures

Figure 2-figure supplement 1. Cell densities are not changed in NgR2$^{-/-}$ cochleae. (A)
Mid/Basal-modiolar sections labeled for βIII-tubulin (green, neurons) from p40 NgR2$^{+/+}$ and
NgR2$^{-/-}$ mice Exemplary section for NgR2$^{-/-}$ is the zoomed out image presented in figure 1C.
(**B**) Quantitative analysis shows that the density of SGN cell bodies are similar between NgR2$^{-/-}$
$^{/-}$ and control cochlea. Data is presented as mean ± SD; N = 7 per group. Scale bar = 10 μm. (**C,**
D) IHC (C) and OHC (D) densities are not affected in P15, 1 month and 2 months old NgR2$^{-/-}$
mice.
895

896
897
Figure 2-figure supplement 2. Intact number but enlarged size of the ribbons in NgR2$^{-/-}$ IHCs.
(**A**) Maximal intensity projections of representative IHCs from apical, mid and basal regions of
NgR2$^{+/+}$ (left) and NgR2$^{-/-}$ (right) cochleae of p21-30 mice. Synapses are visualized by staining
against Ctbp2 and Homer1. Scale bar = 5 μm. (**B**) The number of the ribbons is not affected
along the tonotopic axis in NgR2$^{-/-}$ cochleae. (**C**) The size of the ribbons is increased in IHCs of
both apical and middle turns in NgR2$^{-/-}$ cochleae. N = 6 animals/genotype. Box-whisker plots
show the median, 25/75 percentiles (box) and the range (whiskers). Individual data points are
overlaid.
906

907
908
Figure 2-figure supplement 3. Reduced GluA2/3 signal juxtaposing presynaptic ribbons in
IHCs of NgR2$^{-/-}$ mice. (**A**) Maximal intensity projections of representative IHC regions from
NgR2$^{+/+}$ (top) and NgR2$^{-/-}$ (bottom) cochleae immunolabeled against Ctbp2 and GluA2/3. Scale
bar = 5 μm. The zoom-in regions marked with the white rectagles are presented on the right-
hand side. Scale bar = 2 μm. (**B**) The number of the GluA2/3 positive puncta is drastically
reduced in NgR2$^{-/-}$ mice despite the maintained number of presynaptic ribbons. (**C**) Disrupted
colocalization of Ctbp2 and GluA2 positive puncta at IHCs of NgR2$^{-/-}$ mice. N = 6
animals/genotype. (**D**) Representative images of RNAscope ISH show maintained expression
of Gria2 (red dots) in the SGN somata of NgR2$^{-/-}$ mice. Scale bar = 10 μm.
918

Figure 2-figure supplement 4. Efferent innervation pattern in NgR2^{-/-} cochleae. (A) Maximal
intensity projections of confocal stacks of outer and inner hair cell regions at the apical region
of the cochlea from p21-30 NgR2$^{+/+}$ (left) and NgR2$^{-/-}$ (right) mice. SGN fibers/terminals and
efferent terminals are visualized using anti-Na^{2+}/K$^+$ ATPase and anti-vesicular acetylcholine
transporter (VaChT) antibodies, respectively. Scale Bar = 10 μm **(B)** Same as A but zoomed into
the IHC region. Scale Bar = 5 μm **(C)** Maximal intensity projections of confocal stacks of a row
of apical IHC region from p21 NgR2$^{+/+}$ (left) and NgR2$^{-/-}$ (right) mice. Presynaptic efferent
terminals were stained with anti-synapsin1/2 antibody. Scale bar = 5 μm.
929

930
931
Figure 3-figure supplement 1. Quantification of ribbon number and volume in SBEM
reconstructions. (A) Number of ribbons is not changed in NgR2$^{-/-}$ IHCs (NgR2$^{+/+}$: 10.8 ± 0.83,
SD = 2.04, n = 6, N = 1 vs NgR2$^{-/-}$: 10.3 ± 0.61, SD = 2.1, n = 12, N = 2; p = 0.54, Mann-Whitney-
Wilcoxon test). **(B)** Ribbon volume tends to be larger in NgR2$^{-/-}$ IHCs (NgR2$^{+/+}$: 0.021 ± 0.001
μm^3, SD = 0.008 μm^3, n = 65 ribbons in 6 IHCs, N = 1 vs NgR2$^{-/-}$: 0.024 ± 0.001 μm^3, SD = 0.01
μm^3, n = 125 ribbons in 12 IHCs, N = 2; p = 0.06, Mann-Whitney-Wilcoxon test). Box-whisker
plots show the median, 25/75 percentiles (box) and the range (whiskers). Individual data
points are overlaid.

A Apex

SEM

Diamond
knife   e⁻

B NgR2⁺/⁺   NgR2⁻/⁻

C NgR2⁻/⁻

IHC

Ribbon

Type-1 SGN
with ribbon

Type-1 SGN
w/o ribbon

Habenula
perforata

D NgR2⁺/⁺   NgR2⁻/⁻

Type I SGN with ribbon   Type I SGN w/o ribbon

E

% type I SGN with ribbon per bundle

*

NgR2⁺/⁺   NgR2⁻/⁻

F

% unbranched type I SGN per bundle

p=0.226

NgR2⁺/⁺   NgR2⁻/⁻

A

63 mV

-87 mV

50 pA | 100 ms

1 a.u.|

| NgR2$^{+/+}$ | NgR2$^{-/-}$ |

Ctbp2 bp

Fluo4-FF ΔF

B NgR2$^{+/+}$ (50 AZs in 8 IHCs, N=3)

NgR2$^{-/-}$ (69 AZs in 11 IHCs, N=7)

ΔF/F$_0$

Voltage (mV)

B^i

ΔF/F$_{0\,max}$

C NgR2$^{+/+}$ (42 AZs in 8 IHCs, N=3)

NgR2$^{-/-}$ (59 AZs in 11 IHCs, N=7)

Fractional activation

Voltage (mV)

C^i

k (mV)

C^{ii}

**

V$_{half}$ (mV)

D

| NgR2$^{-/-}$ | NgR2$^{-/-}$ |

Ctbp2

Ca$_V$1.3

Bassoon

Merge

NgR2$^{+/+}$

NgR2$^{-/-}$

IHC

SGNs

ribbon size

Ca^{2+} channel

GluA2/3

PSD size

ABR thresholds

depolarized shift