Making yogurt with the ant holobiont uncovers bacteria, acids, and enzymes for food fermentation

Veronica M. Sinotte[1]*, Verónica Ramos-Viana[2], Diego Prado Vásquez[3], Sevgi Mutlu Sirakova[4], Nabila Rodríguez Valerón[3,5], Ana Cuesta-Maté[1], Shannara K. Taylor Parkins[2], Esther Merino Velasco[3], David Zilber[6], Rasmus Munk[3], Sandra B. Andersen[1], Robert R. Dunn[7], Leonie J. Jahn[2]*

*Correspondence**: Veronica M. Sinotte, veronica.sinotte@sund.ku.dk; Leonie J. Jahn, lejj@biosustain.dtu.dk

[1]Centre for Evolutionary Hologenomics, Globe Institute, University of Copenhagen, Copenhagen, Denmark

[2]Novo Nordisk Foundation Centre for Biosustainability, DTU Biosustain, Kgs. Lyngby, Denmark

[3]Alchemist Explore, Research and Development, Alchemist Aps, Copenhagen, Denmark

[4]Rachel Carson Centre for Environment and Society, LMU Munich, Germany

[5]Basque Culinary Centre, Facultad de Ciencias Gastronómicas, Mondragon Unibertsitatea, Donosita - San Sebastián, Spain

[6]Food & Beverages Applied R&D, Novonesis, Hørsholm, Denmark

[7]Department of Applied Ecology, North Carolina State University, Raleigh, North Carolina, USA

Abstract

Milk fermentation has a rich history in which food culture, the environment, and microbes intersect. However, the biocultural origins of fermentation practices and microbes have largely been replaced by industrial processes. Here, we consider a historical fermentation originating from Turkey and Bulgaria – ant yogurt. We revisit the traditional practices and modern gastronomic applications that use red wood ants (*Formica rufa* group) to initiate milk fermentation. Subsequently, we characterize the ants and experimental ant-derived yogurts. We uncover that the ant holobiont, which consists of the ants and their microbes, contributes key acids and enzymes to fermentation. Metabarcoding and culturing revealed that lactic and acetic acid bacteria, including species related to those in conventional yogurt and sourdough, originate from the live ants and proliferate in the milk. The ants and bacteria consequently introduce formic, lactic, and acetic acid, advantageous for yogurt acidification and coagulation. Last, proteases with the potential to act on casein and alter yogurt texture are produced by the ants and bacteria. The ant holobiont thus facilitates fermentation akin to the microbial consortia in other ferments. Our findings highlight the value of integrating traditional, gastronomic, and biological frameworks to uncover the origins and applications of microbes for fermented foods.

Introduction

The fermentation of milk into products such as yogurt, cheese, and kefir originate from ancient practices and has dramatically shaped food cultures. The oldest evidence for dairying dates to 9,000 years ago in Anatolia (modern-day Turkey) (Evershed et al., 2008). Prehistoric dairy fermentation potentially occurred as early as 7,000 years ago, based on fat and protein residues isolated from ceramics resembling cheese strainers (Evans et al., 2023; Salque et al., 2013). In the millennia to follow, diverse dairy practices transformed milk into a preservable, widespread, and nutritious resource (Fisberg & Machado, 2015; Scott et al., 2022; Warinner et al., 2014). Following suit, dairy fermentation became indispensable to regional cuisines and languages (Multu Sirakova, 2023; Öney Tan, 2010). Yogurt, a tangy fermented milk product, was thus a functional cultural adaptation dependent upon interactions between people, dairy animals, the environment, and most importantly, microbes. It is microbes that enter the milk, and through their enzymatic processes, catalyse the fermentation to acidic, viscous yogurt (Nagaoka, 2019). These interspecies relationships are reflected in the Turkish word for a fermentation starter, *maya*, that ultimately "comes from relations within the broader web of life", including microbes, animals, plants, and human culture (Multu Sirakova, 2023).

In the early 1900s, microbiologists characterized the first yogurt culture, laying the foundation for a pivotal shift from the diversity inherent in traditional yogurt to a simplified industrialized yogurt. Stamen Grigorov and Ilya Metchnikoff isolated and popularized a species of bacteria from Bulgarian *maya*, *Lactobacillus delbrueckii* subsp. *bulgaricus* (Grigoroff, 1905; Metchnikoff, 1908). Following Metchnikoff, the industrialization of yogurt focused on a small number of bacterial taxa, predominantly *L. delbrueckii* subsp. *bulgaricus* and *Streptococcus thermophilus* (FAO & WHO, 2018). Both species are lactic acid bacteria, which play an important role in industrial and wild fermentations through food preservation, flavour generation, and potential health benefits (Hu et al., 2022; Wang et al., 2021; Zapasnik et al., 2022). However, the focus on a few bacterial species overlooks the biodiversity embodied in traditional yogurts, which can include multiple species and strains (Demirci et al., 2022; Velikova et al., 2018). Revisiting biocultural origins of yogurt fermentation offers opportunities to explore ecological niches of food microbes for novel starter cultures (Sedó Melina et al., 2022), to understand the interconnectedness of food systems, and to illuminate the history of yogurt and its cultural praxis.

The bacteria in yogurt stem from the multispecies dimensions of *maya*, which we aim to elucidate for a traditional Turkish and Bulgarian yogurt-making practice - ant yogurt. Generally, the bacterial

community that first colonizes the milk and establishes the fermentation ecosystem may stem
from the dairy animal (Wouters et al., 2002), the person making the ferment (Kothe et al., 2024;
Reese et al., 2020), or the environment, such as vegetation (Michaylova et al., 2007). Yogurt
cultures are then propagated by adding a small amount from a primary or old yogurt to new milk,
a process referred to as "backslopping." A historic practice for starting the first or primary yogurt
in Turkey (Öney Tan, 2010) has been documented by the ethnographer Ali Rıza Yalman, "If the
nomads want to make yogurt and cannot find enough starter culture to make yogurt, they crush
the tiny eggs of the ants sheltering under the stones in their palms. When you put this into the
milk […], that milk becomes yogurt." (Yalman, 1977). In neighbouring Bulgaria, a traditional spring
practice involves fermenting yogurt within a red wood ant colony (Multu Sirakova, 2023). Further
oral histories from Sharri mountains of Albania and North Macedonia also recall the use of ants
in traditional yogurt fermentation (F. Demiraj, personal communication, May 2023). These
countries are connected by cultural threads preceding the establishment of national boundaries,
contributing to general continuity between their culinary practices, both today and across
millennia. The ethnographic evidence of ant yogurt across regions suggests ants may play an
overlooked functional role in fermentation.

Here, we test the hypothesis that yogurt fermentation can be initiated by the ant "holobiont", which
includes both the ant and the microbial communities inherent to it (Bordenstein & Theis, 2015;
Margulis & Fester, 1991). The ant holobiont may contribute acids and enzymes key to the
fermentation (Figure 1). First, we explored the potential of red wood ants (*Formica rufa* group)
(Robinson & Stockan, 2020) as starter cultures for yogurt based on ethnographic accounts and
as culinary ingredients for modern gastronomy. Then, we characterized the microbial community
of the ants *F. rufa* and *F. polyctena* and yogurts derived from *F. polyctena*, quantified organic
acids in the yogurts, and assessed the proteases and peptidases originating from ants and
bacteria (Figure 1 - figure supplement 1). The study overall examines an overlooked ecological
niche of bacteria in the context of traditional practices to identify microbes key to past and future
fermented foods.

Results

The ant holobiont serves as a catalysing agent for food applications

We worked towards a holistic understanding of traditional uses of ants as a yogurt starter by
conducting fieldwork in Bulgaria in a community that retained an oral history of the practice (Multu
Sirakova, 2023; Sinotte et al., 2023). This community also is the ancestral home of one of the

authors (S. Mutlu Sirakova). We used an ant colony selected by members of the community,
which was the species *F. rufa*. This is one of four species in the *F. rufa* group that maintain species
ranges within Bulgaria (Antonova & Marinov, 2021). To make the yogurt, four live ants were added
to the jar of warmed raw milk (Figure 2A), a cheese cloth was placed over the top of the glass jar,
and the jar was left to ferment within the colony overnight (Figure 2B). The jar was retrieved after
a day of incubation. The acidity, texture, and flavour were indicative of early stages of yogurt
fermentation. We observed that the milk had acidified to pH 5, coagulated at the bottom of the
container (Figure 2C), and "had a slight tangy taste with mild herbaceousness and pronounced
flavours of grass-fed fat" (D. Zilber).

Culinary applications of the *F. rufa* ants were created by the research and development team at
Restaurant Alchemist, ranked number eight in the world (50 Best, 2024), holding two Michelin
stars (Guide, 2024), and known for science-centred innovation. Co-authors based these
applications on previous experiments where the addition of ants to milk caused coagulation.
Presumably, this coagulation was due to the *F. rufa* ants' formic acid. Discussions on the
traditional uses of ants to coagulate milk (T. Tan, personal communication, April 2022) further
directed the development of culinary applications. Thus, the innovations leverage the potential of
ant-derived formic acid and microbe-derived organic acids to coagulate milk.

Three culinary applications were developed using live ants and ants that were frozen and
subsequently dehydrated. First, the "ant-wich" contained ice cream derived from sheep yogurt
made with live ants as starter culture (Figure 2D). The ants provided a distinct, pungent acidity
that contrasts with the fat of the milk, and a serving temperature of -11ºC balances the desert.
Second, a goat milk "mascarpone" was developed using dehydrated ants to catalyse the milk
coagulation (Figure 2E). The texture was similar to commercial mascarpone, yet the flavour was
pungent and aromatic, like a mature pecorino cheese. Third, a milk wash cocktail was created,
which is a dairy-based cocktail dating to the early 1700s (Wondrich, 2010). Classically, the milk
is curdled with acid from citrus and subsequently filtered to remove the dairy solids, resulting in a
clear beverage with more richness and body. In this case, dehydrated ants induced the curdling
and separation of milk (Figure 2F). The cocktail had fruity notes from the apricot liqueur and the
brandy (raisins, dried figs, and caramelized apples), and a silky texture from the residual milk
whey. Replacing citrus with ants resulted in a milder acidity and a distinct flavour with additional
fruity notes. Despite the potential of these culinary innovations, we note that ants are not included
among the four insects authorized in the European Union according to Regulation 2015/2283 on
Novel Foods.

Given that ants have diverse culinary applications with potential to initiate fermentation, we aimed
to further elucidate the role of the ant holobiont. We hypothesized that the ant holobiont, consisting
of the ant and its microbial partners, contributes to fermentation with bacteria, acids, and
enzymes. To address this hypothesis, we created three elaborations of fermented milk with live,
frozen, or dehydrated ants, per the modern gastronomic applications. The ants were collected in
late spring and early autumn to further determine the impact of the season on the microbiome
and the fermentation.

Live ants provide small, stable, and controlled yogurt microbiomes

We first characterized the bacterial microbiome of red wood ant sister species *F. rufa* and *F.*
polyctena (Borowiec et al., 2021) with 16S *rRNA* metabarcoding. Our aim was to examine how
the ant yogurt practice may be translated across species and geographies. These ants have
ranges spanning Denmark, where experimental yogurts were made, and Bulgaria or Turkey
(Guénard et al., 2017; Janicki et al., 2016), where ethnographic histories originated (Figure 3A).
Both species are found along the edges of pine forests, have thatched pine needle mounds, and
near-identical morphological characteristics (Seifert, 2021). These similarities make them all but
indistinguishable in the field to ant experts (Seifert, 2021), suggesting both may have been used
in traditional and modern culinary applications.

The microbiomes of the two *Formica* species were dominated by lactic acid bacteria
(Lactobacillaceae), acetic acid bacteria (Acetobacteraceae), and obligate intracellular bacteria
(Anaplasmataceae) (Figure 3A). The bacterial families and genera (Figure 3 – figure supplement
1A) align with *Formica*-specific microbiomes associated with species across the Northern
Hemisphere (Jackson et al., 2023; Jackson et al., 2022; Kaczmarczyk-Ziemba et al., 2020; Zheng
et al., 2022). *Formica* ant microbiomes were seasonal. We observed an increase in relative
abundance of lactic and acetic acid bacteria (Figure 3A) and overall microbial biomass (i.e. load)
in the autumn, compared to spring (Figure 3 – figure supplement 1B). Seasonal microbiomes
have not been previously documented in *Formica* ants and are germane to the use of these ants
in fermentation. Overall, the consistent presence of lactic and acetic acid bacteria in the *F. rufa*
group indicates the ant microbiome may be pertinent to food fermentation.

We then tested the hypothesis that lactic and acetic acid bacteria in the ants may transfer to the
yogurt. Based on 16S *rRNA* metabarcoding of experimental yogurts made with *F. polyctena* under
aseptic conditions, we confirmed our hypothesis that bacteria from the ants contribute to the
yogurt microbiome. The preparation of the ants, namely whether they were live, dehydrated, or

frozen, distinctly impacted the bacteria communities in the resulting yogurts (Figure 3B). Like the

ants themselves, the live ant yogurts were consistently dominated by the lactic and acetic acid

bacteria, which displayed seasonal differences in composition. Here, the most abundant bacterial

genus was *Fructilactobacillus* (Figure 3 – figure supplement 1C). We note that these true

abundances may have been higher, considering our metabarcoding positive control indicates we

slight underestimation of the abundance of lactic acid bacteria in samples (Figure 3 – figure

supplement 2). In contrast, dehydrated ant yogurts exhibited variable microbiomes. Frozen ant

yogurts contained two bacteria groups: Bacillaceae, which can proliferate in the yogurt, and

Anaplasmataceae, which obligately lives in ant cells (Buchner, 1965; Jackson et al., 2022) and

thus cannot grow in the yogurt. Here, dehydration and freezing likely reduced bacterial viability

(Santivarangkna et al., 2011) and consequently favoured stochastic community assembly or

freeze-resistant Bacillaceae (Nicholson et al., 2000). The persistence of lactic and acetic acid

bacteria in live ant yogurt suggests it is the best starter for fermentation and provides biological

support for the traditional, live ant-based yogurts.

We determined the consistency of the bacterial microbiome composition across the yogurts.

Alpha and beta diversity metrics were used, where alpha diversity is the number of bacterial

strains per sample (i.e. amplicon sequence variants) and beta diversity is the difference in the

composition of strains from one sample to the next, while weighting relative abundance. The ant

preparations significantly affected the alpha and beta diversity of the yogurts (LM: $F_{2,27}$=5.387,

p=0.0108, based on Shannon index; Adonis: $F_{2,27}$= 8.4402, *p*<0.0001, based on Bray Curtis

distances). The alpha diversity of the live ant yogurts was intermediate (Figure 3C), but the beta

diversity was very low (Figure 3D). Therefore, the live ant yogurts tended to have the same

species and composition from one sample/preparation to the next, as one would hope to see in

predictable and controlled fermentations.

Bacteria from the ants proliferate in the milk

We hypothesized that bacteria from the ant holobiont grew in the milk, facilitating fermentation.

To assess growth, we determined the bacterial load in ants and yogurts with quantitative PCR

(qPCR) and identified viable bacteria with culturomics using 16S *rRNA* Sanger sequencing. Live

ants introduced lactic acid bacteria that proliferated in the yogurt. In the spring, yogurts contained

a higher loads of lactic acid bacteria than the ants introduced as a starter (Figure 4A; *t*(2, 10) = -

2.975, *p*=0.0139). However, in autumn, we observed the opposite pattern (Figure 4A; *t*(2, 10) =

3.269, *p*=0.0084), suggesting that not all lactic acid bacteria from the ants enter the milk, or that

the bacteria associated with the autumn ants are less prolific. While lactic acid bacteria dominated

live ant yogurts, it made up a marginal amount the bacterial load in frozen and dehydrate yogurts

(Figure 4 – figure supplement 1A).

Yogurt made with live ants contained several species of culturable lactic acid bacteria (Figure

4B). Aligned with the dominance of *Fructilactobacillus* in the community analysis, we isolated *F.*

sanfranciscensis, a bacterium not only associated with ants (Zheng et al., 2022) but also

sourdough bread fermentation (Gänzle et al., 2023; Kline & Sugihara, 1971). We further cultured

Lactobacillus delbrueckii subsp*. bulgaricus* and *Lacticaseibacillus casei*, typically found in

fermented dairy products (FAO & WHO, 2018; Zheng et al., 2020). Last, we isolated the acetic

acid bacteria, *Oecophyllibacter saccharovorans* (Figure 4 – figure supplement 2), previously

characterised in association with ant genera closely related to *Formica* (Blaimer et al., 2015;

Brown & Wernegreen, 2016; Chua et al., 2018). The diversity of culturable lactic acid bacteria

thus indicates a niche overlap between ants and fermented foods.

In the yogurts made with dehydrated and frozen ants, the bacterial load disproportionately came

from spore-forming Bacillaceae. Notably, there is little to no Bacillaceae biomass in the ants,

especially when compared to the yogurts (Figure 4C; $t_{frozen}(2, 10)$ = -2.307, *p*=0.0437), and this

Bacillaceae did not proliferate in live ant yogurts (Figure 4 – figure supplement 1B). We cultured

several species of Bacillaceae, stemming largely from dehydrated or frozen ants or yogurts

(Figure 4D). This included the food contaminant, *Bacillus cereus* (Dietrich et al., 2021). Although

we cannot determine if the *Bacillus* load was sufficient to be problematic for consumption, it still

indicates a potential risk. Therefore, we conclude that the microbiome of dehydrated and frozen

ants and their corresponding yogurts are undesirable for food fermentations.

Ants and bacteria contribute to the acidification of yogurt

We hypothesized that the ant holobiont, composed of both the ant and its microbes, may

contribute acids key to fermentation. In yogurt, bacteria metabolize lactose and consequently

produce high amounts of lactic acid and often low amounts of acetic and formic acid (Sieuwerts

et al., 2010). These acids are essential to the tangy flavour, thick texture, and preservation of

yogurt (Fernandez-Garcia & McGregor, 1994). Moreover, *Formica* ants (though, notably, not all

ants) have a venom gland that contains largely formic acid that may be up to 10% its body weight

(O'Rourke, 1950), although the whole ant contains a more complex mélange of chemicals

(Löfqvist, 1976). Thus, the acid from the ant holobiont may have three effects. First, it may

engender yogurt tastes and textures, even without the effects of microbes. Second, it can create

acidic conditions that favour acid-producing microbes. Third, the acid-producing microbes can
themselves alter the acidity and flavour of the ferment. To investigate if the components of the
ant holobiont, body and microbes, contribute organic acids to the yogurt, we quantified formic,
lactic, and acetic acid in yogurts and controls by HPLC and measured pH before and after
fermentation (Figure 5).

The ant holobiont contributed formic acid, which was the most abundant acid found in the yogurts
(Figure 5A). The amount of formic acid was in a similar range to that in the control, in which formic
acid alone was added to pH 4.6 to induce coagulation. The season, but not the type of yogurt
(live, dehydrated, frozen) further dictated the formic acid levels (LM results). The higher levels in
autumn (t-test: t_{live}(2, 10) = -3.500, p= 0.0057; $t_{dehydrated}$(2, 10) =-3.297, p=0.0081) may be due to
the seasonality of formic acid production (O'Rourke 1950). Therefore, the acid from the ants
entered the milk and likely shaped the unique acidic flavour and texture of the culinary
applications, contrasting the rounded flavour of lactic acid typically predominant in fermented
dairy.

Lactic and acetic acid were also detected in the ant yogurts, validating that ant-associated
bacteria likely contributed acids during fermentation (Figure 5A). Acid profiles align with the
microbiome data. Lactic and acetic acid bacteria were the most abundant genera in live ant
yogurts (Figure 3A). Similarly, these acids are highest in live ant yogurts compared to dehydrated
(t_{lactic}(2, 10) =4.270, p=0.0052; t_{acetic}(2, 10) =3.065, p=0.0220) and frozen yogurts (t_{lactic}(2, 10)
=3.842, p=0.0060; t_{acetic}(2, 10) =3.653, p=0.0147). In live yogurts, we observed lactic and acetic
acid were higher in spring (t_{lactic}(2, 10) =2.897, p=0.0231; t_{acetic}(2, 10) =3.325, p=0.012), the season
in which we isolated the greatest diversity of lactic acid bacteria (Figure 4C). This suggests
different strains of bacteria may inherently vary in acid production. Notably, ant yogurts had less
lactic acid, acetic acid (Figure 5A), and bacterial biomass (Figure 5 – figure supplement 1A) than
conventional or homemade yogurts. We did not establish and perform the optimal fermentation
conditions for the ant yogurt, and thus the fermentation was incomplete. However, we can
conclude that the ant microbes contribute organic acids to fermentation.

Ant yogurts contained on average 2.5 g/L total organic acids, while conventional yogurts
contained up to 12 g/L total organic acid. Thus, the ant yogurt remained in a pH range of 5.0-5.9
and did not reduce to a pH of 4.2 of conventional yogurt (Figure 5 – figure supplement 1B), given
the conditions and timing of our experiment. Initially, the pH of the milk dropped upon the addition
of ants, likely due to the formic acid from the ants (Figure 5 – source data 1). The pH continued
to decrease through the fermentation (Figure 5 – figure supplement 1B), likely because of acid-

producing microbes. Typically, in yogurt fermentation, a drop to pH 5.3 initiates coagulation, and

a further acidification to pH 4.6 completes coagulation by causing aggregation of casein micelles

(Robinson et al., 2006) The ant yogurt coagulated yet largely did not reach such a low pH. This

suggests another aspect of the fermentation denatured caseins, namely proteases.

Ants and microbes contribute proteases and peptidases, potentially modifying yogurt texture

Our final hypothesis was that the ant holobiont may contribute enzymes, in particular proteases,

that modify the yogurt texture. Proteases can differentially cleave and degrade casein, the main

protein in milk, and result in a firm or fluid yogurt texture (Amani et al., 2017; Gassem & Frank,

1991). Based on the pH and coagulation results, we predicted the ants themselves may contain

milk-clotting proteases, similar to those previously identified in mealworms (Yang et al., 2022).

Casein proteases and peptidases are also typically produced by the bacteria in yogurt (Amani et

al., 2017). Thus, the ant holobiont, ants and bacteria, may contribute proteases to the yogurt

fermentation. To test this hypothesis, we conducted an untargeted proteomics analysis based on

a self-curated database that contained proteases from *Formica* ants, ant-yogurt bacteria,

conventional yogurt bacteria, and milk proteins.

We confirmed our hypothesis that the ant holobiont contributes proteases and peptidases to the

yogurt fermentation (Figure 5B). Several ant proteases were identified in the yogurt (Figure 5B),

one of which, an ATP-dependent CLP protease, is known to have caseinolytic potential (Maurizi

et al., 1990). Ant-associated bacteria further contributed proteases and peptidases. The lactic

acid bacteria *F. sanfranciscensis* produced two proteases, ATP-dependent CLP protease and

zinc metalloprotease, both known to have caseinolytic homologues (Jia et al., 2015; Maurizi et

al., 1990). Proteases and peptidases typically belonging to bacteria in conventional yogurt, *L.*

*delbruecki*i subsp. *bulgaricus* and *Streptococcus thermophilus*, were found, as well as ant yogurt

bacteria *Paenibacillus* and *Wolbachia pipentis*. Last, we identified milk proteins such as casein,

which made up a high percentage $96.19 \pm 1.07\%$ of the proteome of samples (Figure 5 – source

data 2), potentially because of the low bacterial load compared to conventional yogurt. There

were no significant differences in the relative abundance of all proteins in the proteome between

live and dehydrated ant yogurts or seasons (Figure 5 – figure supplement 2). Thus, proteases in

both live and dehydrated yogurts may, in part, catalyse coagulation in the experimental yogurts

and in the modern culinary applications. Taken together, the ant holobiont, ants and microbes,

contribute proteases potentially relevant to food fermentation.

Discussion

In this study, we explored the traditional practice of using ants as a starter for yogurt fermentation.
We hypothesized that the ant holobiont, both the ant and its microbes, catalysed the fermentation.
First, we uncovered that bacteria from the ants shaped the yogurt microbiome. Using live ants as
a starter, in contrast to frozen or dehydrated ants, resulted in a consistent microbiome containing
lactic and acetic acid bacteria, desirable for fermentation. These ant-bacteria included some
species thought of as canonical fermentation microbes, such as *L. delbrueckii* subsp. *bulgaricus*
and *F. sanfranciscensis*. Second, we discovered that the addition of the ant holobiont to milk
introduced acids and proteases. Formic acid originated from the ant, while lactic and acetic acid
were likely produced by bacteria. The ants and bacteria also contributed proteases with
caseinolytic potential. Our results revealed that the ant holobiont can contribute to fermentation
and thus prompted reconsideration of the microbial consortia in our ferments, their origins, and
applications to fermented food.

We uncovered how ants and their bacteria synergistically shape the fermentation by applying the
holobiont framework. Holobiont theory considers a host and its associated microbes collectively,
where the interplay between animal and microbes shapes the resulting ecology and evolution of
the whole, host plus microbes (Bordenstein & Theis, 2015; Margulis & Fester, 1991; Zilber-
Rosenberg & Rosenberg, 2008). When animals or plants are used in fermentation, their
holobionts may structure that fermentation in ways that reach beyond the effect of the individual
parts. *Formica* ants produce formic acid for defence (O'Rourke, 1950; Tragust et al., 2020) and
feed on the sugar-rich honeydew (Ivens & Kronauer, 2022) and protein-rich insects (Robinson &
Stockan, 2020). In response, the lactic and acetic acid bacteria are hypothesized to metabolize
the honeydew (Zheng et al., 2022), are highly acid tolerant (Tragust et al., 2020) and are potential
contributors to proteolysis. Once ants are introduced into milk, the properties of the ant holobiont
extend to the yogurt ecosystem. Formic acid initially acidifies the milk, ant-bacteria break down
milk sugars producing additional acid, and proteases liberate peptides and amino acids for
bacterial growth. By looking through the lens of the holobiont, we could clarify why *Formica* ants
and bacteria are pre-adapted for the fermentation of yogurt and potentially other foods.

The effect of the ant holobiont on yogurt fermentation parallels the role of microbes in other
fermented foods. In most ferments, multiple species of microbes offer complementary
contributions to the fermentation process (Canon et al., 2020; Sieuwerts et al., 2008). In
conventional yogurt, the bacteria *S. thermophilus* produces formic acid that is then metabolized
by *L. delbrueckii* subsp. *bulgaricus*, and in return, *L. delbrueckii* subsp. *bulgaricus* produces
proteases*.* Collectively, these bacteria then metabolize lactose and produce lactic acid to

transform milk into tangy yogurt (Sieuwerts, 2016; Sieuwerts et al., 2010). In sourdough, yeast
and lactic acid bacteria can secrete factors that promote one another's growth (Sieuwerts et al.,
2018), and overall have complementary roles in the fermentation of sugars and protein derivatives
(Gänzle, 2014; Gänzle et al., 2023). Together, sourdough microbes create leavened bread with
acidic, kokumi, and umami flavours (Gänzle et al., 2023). In the ant yogurt, our results suggest
the separate elements of the ant holobiont, the ant body and microbes, act synergistically.
Consequently, the ant holobiont introduces acids, proteases, and potentially other compounds
that contribute to the organoleptic characteristics such as distinct acidity, coagulation, and flavour.

Intriguingly, we isolated two species of lactic acid bacteria from the ant yogurt that are commonly
found in fermented foods, *L. delbrueckii* subsp. *bulgaricus* and *F. sanfranciscensis*. *L. delbrueckii*
subsp. *bulgaricus* was first scientifically isolated from Bulgarian yogurt (Grigoroff, 1905;
Metchnikoff, 1908) and is presently used in conventional yogurt around the world (FAO & WHO,
2018). The phylogeny indicates that related species are associated with bees and vertebrates
such as pigs and hamsters (Duar et al., 2017). It has also been identified on the leaves of the
snowdrop flower, *Galanthus nivalis*, endemic to Bulgaria (Michaylova et al., 2007). We now add
the ant holobiont to the list of potential natural habitats of these bacteria. The second bacterial
species, *F. sanfranciscensis,* is one of the most prevalent bacteria in sourdough bread globally
(Gänzle & Zheng, 2019; Landis et al., 2021). The *Fructilactobacillus* clade appears to have
diversified with ants (Jackson et al., 2023) and related social insects (Duar et al., 2017). We
suspect that *F. sanfranciscensis* has evolved in ants over millions of years and was only
introduced into fermented foods with the advent of bread making in the last several thousand
years (Arranz-Otaegui et al., 2018; Carretero et al., 2017). These results lay the groundwork for
unveiling the diversity of fermentative bacteria residing in ants and for elucidating the bacterial
evolution from ant holobionts to fermented foods.

Our results bring forth new questions we believe are central to understanding the ecological and
ancient origins of fermentative microbes. The ecological origins and assembly of ferment
microbiomes, which colonize fermentations in lieu of backslopping, are only being to be
characterized (Louw et al., 2023). Recently, it has been convincingly argued that the wine yeasts
can originate from social wasps, where the symbiosis between wasps and yeasts is essential for
the microbial transfer to wine ecosystems (Di Paola et al., 2023; Madden et al., 2018; Stefanini et
al., 2012). Similarly, our results are reconcilable with the idea that bacteria for yogurt fermentation
may originate from ants. This poses the question whether the ant holobiont was influential to the
historic origins of yogurt or bread fermentation. Beyond traditional practices using ants in yogurt

(Multu Sirakova, 2023; Öney Tan, 2010; Yalman, 1977), ant colony materials may be integrated
in sourdough fermentation (Mutlu Sirakova 2023), providing further avenues for microbial transfer.
Conceivably, the ant bacteria then could be backslopped in traditional fermentations, successively
propagating and evolving in yogurts. Although it is not clear when these traditions arose, it is
noteworthy that the earliest evidence of yogurt and architectural depictions of ants and bees were
found in prehistoric remains from our study region (Hendy et al., 2018; Walter, 2021). While these
fragmentary anecdotes do not reconcile or resolve the origins of fermentative microbes, they
highlight cultural threads that may connect ants and ferments.

As a group of co-authors, including anthropologists, culinary innovators, and food scientists, we
differentially perceive the implications of the ant holobiont on yogurt-making. From an
anthropological viewpoint, it challenges the ethos of conventional yogurt fermentation. Unlike
traditional yogurts, large-scale industrial yogurt production relies on a limited number of bacterial
strains. Conversely, the lactic acid bacteria found in ant yogurt are maintained in the ants, yogurt,
and environment, and are cultured through traditional dairy practices passed down over
generations (Multu Sirakova, 2023). To preserve this microbial diversity, we must uphold its
cultural heritage, along with the conservation of red wood ants (Balzani et al., 2022). In relation
to modern gastronomy, elaborations on the use of ants in milk, along with their microbes and
enzymes, may hold wider purchase for fermented foods. It opens new culinary possibilities for
other dairy products and fermented beverages that are sour or effervescent. Beyond the
possibility of imagining new foods, exposing the public to familiar foods made with microbes and
insects could conceivably help change consumer perceptions of entomophagy. From the food
science perspective, microbes or enzymes from ant yogurts can be integrated into the food
system to explore novel applications and flavours. For example, these ant microbes hold potential
promise for new plant-based foods, such as dairy-free yogurts. While we differ in what the future
of ant holobiont-facilitated fermentation may be, we agree it provides a powerful example that can
broaden our thinking, as a society, about what is possible.

Materials and Methods

Traditional and modern culinary applications of ants

We travelled to Nova Mahala, Bulgaria to recreate ant yogurt with communities that retained
memories and oral histories of the ant yogurt tradition. We based the yogurt fermentation on
freshly acquired raw cow milk and *F. rufa* ants, from a colony found just outside of the village. The
milk was warmed until nearly scalding, such that it could "bite your pinky finger" (Multu Sirakova,

2023). Four live *F. rufa* ants were added to 400 mL of milk. A cheesecloth was secured over the

milk, and fabric was wrapped around the glass container for insulation. The milk jar was then

buried inside the ant colony, covering it completely with the mound material. The milk was left

within the colony and retrieved the next day, 26 hours later. The yogurt was then tasted, checked

with pH strips, and stirred to observe coagulation.

Three culinary applications of the *Formica rufa* ants were developed by the research and

development team at Restaurant Alchemist.

Ant yogurt ice cream sandwich: The "ant-wich" consisted of ant yogurt ice cream, ant gel filling,

and ant tuile. To make the ant yogurt ice cream, organic 7% fat sheep's milk (Hårbølle Mejeri,

Møn, Denmark) was fermented with 1.6% live ants. Live ants were crushed and mixed in an

aliquot of milk, which was passed through a strainer with pore size 100 µmM (Sigma Aldrich,

Denmark) to remove ant body parts and potential parasites that may occur in low prevalence

(Jensen et al., 2017). The strained aliquot was mixed with the rest of the milk, followed by an

incubation period of 8 h at 40°C and subsequent storage at 4°C overnight. To sweeten and thicken

the yogurt into ice cream, 300 g of sheep's cream (Hårbølle Mejeri Møn, Denmark), and 22 g of

sugar (Nordic Sugar, Denmark) was added to 380 g of ant yogurt, followed by 3 g of gelatine

sheets (Condi, bloom 220, Denmark) and 5 g of ice cream stabilizer. The ant gel filling, based on

an ant infusion, was made for the ice cream sandwich. For the infusion, dehydrated ants (*Formica*

rufa) and water (filtered, pH 7) were mixed at the ratio of 10/100 (v/w), vacuum infused at 85°C

for 8 h, strained through a fine mesh, and kept at 4°C. Subsequently, the ant gel was made with

100 g of ant infusion, 7.5 g of powdered glucose 33DE (Sosa, Spain), 70 g of inverted sugar

(Trimoline, Cristalco, France), and 5 g of NH pectin (Sosa, Denmark). These were mixed, cooked

for 5 min, and set to rest at 4°C for 24 h. Finally, an ant tuile was made to sandwich the ant ice

cream and filling. The tuile was made by mixing 100 g of sweetened condensed milk (Tørsleffs,

Denmark), 100 g of 00 wheat flour (Condi, Denmark), 100 g of egg whites, 3 g of baking powder

(Condi, Denmark), 2 g of salt (Maldon), and 8 g of charcoal-black colorant (Sosa, Spain). The

mixture was blended for 5 min, spread to measure 2 mm in height, and baked in the oven at 155

°C for 15 min. The final "ant-wich" was assembled to take the shape of an ant using a laser-cut

stencil and served at -11 °C.

Ant mascarpone-like cheese: A mascarpone-like cheese was developed using ants as the

coagulant. In common practice, lemon juice and milk or cream is added to make mascarpone,

where the citric acid coagulates the dairy. Here, the acid from the ants likely contributed to the

coagulation. Pasteurized sheep's milk (7% fat) and cream (64% fat) (Hårbølle Mejeri, Møn,

Denmark) were mixed with the proportions being 30% and 70%, respectively. The dairy mixture

was homogenized and heated to 95°C. Then, 10 mL ant water infusion (described above) was

added for each kilogram of this dairy mixture, stirring until complete coagulation (Munieweg et al.,

2021). The curd was cooled to 4°C for 6 h and strained through a cheesecloth. The mascarpone-

like cheese was stored at 4°C.

Cocktail clarified with ant milk-wash: Milk-wash is a technique commonly used in cocktails to

clarify a liquid. Milk Punch, the common drink prepared with this technique, is a dairy-based drink

dating to the late 1600s and early 1700s. The earliest written recipe of clarified milk punch comes

from a 1711 cookbook by Mary Rockett (Wondrich, 2010). To do so, the milk is curdled by lowering

the pH, which makes it possible to strain out the dairy solids. The result is a clear beverage with

more richness and body. It is also a popular modern technique at cocktail bars, normally produced

in large amounts (Wondrich, 2010). Using this technique, a cocktail was developed, but by

inducing the milk coagulation with dehydrated ants. The formic acid replaced other products

commonly used to lower the pH, such as lemon, lime, or any other acids like citric or ascorbic

acid. For the cocktail, an alcoholic base was made using 181 g of brandy (Ximenez Spinola

Battonage Brandy, Spain), 101 g of genepi liqueur (Genepi Grand Tetras, France), and 118 g of

apricot liqueur (Bitter Truth Apricot Liqueur, Germany) mixed in a glass jar. 10 g of dehydrated

ants (*Formica rufa*) were added and infused for 2 h at RT. The alcoholic mix plus dehydrated ants

were poured into a glass jar with 200 g of milk (5% fat), stirred, and kept at 2 °C for 6 h to curdle

the milk. Subsequently, the mixture was filtered through a coffee filter and kept at 2 °C until

serving. The cocktail was garnished with 4 frozen ants. The cocktail is in the "aperitif" style of

cocktails.

Ant collection and processing for experimental yogurt

Experimental yogurts were made in spring, according to the tradition (Multu Sirakova, 2023), and

in autumn, to test the seasonality of the fermentation. We used the red wood ant *Formica*

polyctena for microbiome characterization and experimental yogurt fermentation, and *F. rufa* for

microbiome characterization. *F. polyctena* is the sister species of *F. rufa*, where the later was

used in the ethnographic account and at the Alchemist R&D. *F. polyctena* worker ants were

collected at the end of May and the start of October 2022*,* from a colony in Rude Skov, Denmark

(55.8334720, 12.474938410). Similarly, worker ants from a colony of *F. rufa*, found in

Bøllemosen, Denmark (55.827676, 12.534667), were collected at the end of May and late

September 2023. Ants were taken without disturbing the mound structure, considering several

species within the group have experienced species declines within Europe (Balzani et al., 2022).

Ants were gathered with sterilized equipment and kept on ice while transported to the lab.

F. polyctena worker ants were then processed to make the 3 elaborations of ant yogurt, live,

frozen, and dehydrated. Live ants were provided sterile 15% sucrose as food and used to make

ant yogurt within the week of collection. Prior to freezing, ants were aseptically removed from the

nest material, then snap-frozen and stored at -20°C. Dehydrated ants were prepared by placing

frozen ants in a vented Petri dish with sterile filter paper and incubated at 40°C for 16 h and were

used within a week. Live and dehydrated ants were frozen at -20°C for later microbiome

characterization.

The ant species were identified by barcoding of the *cytochrome c oxidase I gene (COXI)*. DNA

was extracted with Chelex (Sigma Aldrich, Denmark), amplified, cleaned, and sequenced (Sinotte

et al., 2021). PCRs were performed with primers LCO1490 5´-

GGTCAACAAATCATAAAGATATTGG-3´and HCO2198 5´-

TAAACTTCAGGRTGACCAAAAAATCA-3´ (Zuccon et al., 2012). PCR products were cleaned

with PureIT ExoZap (Amplicon, Denmark) and sequenced at Macrogen (the Netherlands).

Sequences were trimmed and aligned using default settings in Geneious (Biomatters Ltd., New

Zealand), and the consensus sequence was compared to NCBI's nucleotide database with

BLASTn.

Live, dehydrated, and frozen elaborations of experimental ant-yogurt

Ant yogurts were made under sterile conditions within a laminar flow hood, to assure that the ants

were the sole source of microbes. The yogurt was made with 10% w/v reconstituted milk powder

in distilled water (Sigma Aldrich, Denmark) and sterilized at 121 °C for 5 min to avoid

caramelization. Yogurts were made in Falcon tubes with 30 mL of sterile milk and 0.5 g live ants,

crushed with a sterile pestle, 0.5 g frozen ants, or 0.4 g dehydrated ants. Yogurts and negative

controls were incubated at 42°C for 8 h and then stored at 4°C overnight, according to standard

yogurt-making practices (Nagaoka, 2019; Robinson et al., 2006). The following day yogurts were

subsampled for microbial culturomics, acid quantification, DNA extraction, and proteomics.

Subsamples were stored at -20 °C.

In the spring, six samples of each of the live, frozen, and dehydrated ant-yogurts were made as

well as 2 negative controls of milk alone, one incubated and one not incubated. In the autumn,

the same design was used, but frozen ant-yogurt was not made as it was found to contain
pathogens.

DNA extraction of yogurt and ants

DNA was extracted from the yogurts and ants to assess the microbial community composition. All
extractions were performed under sterile conditions in a laminar air flow. The yogurt samples were
first homogenized by vortexing, then passed through a Corning cell strainer with pore size 100
µm (Sigma Aldrich, Denmark) to remove ant body parts. The yogurt underwent three washes prior
to lysis to remove excess fats that may act as inhibitors and to pellet microbial cells. First, 3 g of
yogurt and 10 mL of sterile 2% sodium citrate solution pH 7.4 were incubated at room temperature
for 10 min, then centrifuged at 5,000 g for 10 min at 4°C. Second, the pellet was retained and
washed in 15 mL TES buffer (50 mM Tris, 1 mM EDTA, 20% sucrose) and centrifuged at 5,000 g
for 10 min at 4°C. A final wash was done in 1 mL TES, which was pelleted, and supernatant
removed. For the ants, two ants were used per DNA extraction. Prior to extraction, 200 µL
molecular water was added to the ants and they were crushed with a sterile pestle.

DNA extraction was performed with the DNeasy Blood and Tissue Kit (Qiagen, Germany)
according to manufacturer's instructions for gram positive bacteria with the following
modifications. Physical lysis of ants and yogurts was performed via bead-beating with 0.1 mm G2
DNA/RNA Enhancer beads (VWR, Denmark) in a TissueLyser (Qiagen, Denmark) at 30 Hz for 3
min. Samples then underwent chemical lysis, including 30 min lysis in lysozyme with an addition
of 2 µL of mutanolysin (25 U/µL) followed by 1 h proteinase K lysis. The final elution of the DNA
extract was 200 µL in AE buffer, passed twice through the column to maximize yield. Due to
exceptionally low DNA concentrations in some experimental yogurts, three extractions were
performed per sample, and thus 9 g of yogurt in total was washed, extracted, and pooled per
sample. Similarly, two ant extractions were pooled for a final sample representing four ants. The
elutes were pooled and concentrated with AMP Pure XP Reagent (Beckman Coulter, Denmark)
according to manufacturer's instructions, then underwent a clean-up for inhibitors with the DNeasy
PowerClean Pro kit (Qiagen, Denmark) according to manufacturer's instructions.

The final samples for sequencing included yogurts, ants, and several controls. Yogurt samples
thus included 20 samples from spring and 13 from autumn, as previously described. From the
ants, there were six final samples for live and for dehydrated *F. polyctena* each from spring and
autumn, and six samples of live *F. rufa* in spring and autumn. In addition to the experimental

yogurts and ants, a mock community of standard bacteria (Zymobiomics, Nordic Biosite,

Denmark) was included to determine any community bias generated by extraction and later

metabarcoding. Further, three conventional plain yogurts were included as positive controls. Last,

five blank DNA extractions were included as negative controls for DNA extraction.

Quantification of bacterial load

The bacterial microbial loads of the yogurt, ant, and control samples were assessed with real-

time PCR (qPCR). To quantify the bacteria in samples, the Femto Bacterial DNA Quantification

Kit (Zymobiomics, Denmark) was used according to the manufacturer's instructions. The Femto

Bacterial Kit targets the 16S *rRNA* gene of bacteria and estimates the amount of bacterial DNA

(ng/μL) in a sample based on a series of standards and non-template control. qPCR of samples

was run in duplicate on Mx3005p (Agilent, Germany). The technical replicates demonstrated

minimal variation and were thus averaged. Subsequently, the amount of DNA/mL of yogurt, in the

total yogurt, or ant starter, was back calculated based on the standards and experimental

parameters.

Bacterial community metabarcoding and analysis

The bacterial community of the ants, yogurts, and controls was characterized with bacterial

metabarcoding. Library preparation and targeted sequencing of the 16S *rRNA* V3-V4 region was

performed with the primers 341F (5' -CCTAYGGGRBGCASCAG- 3') and 806R (5' -

GGACTACNNGGGTATCTAAT- 3') at Novogene Europe (Cambridge, United Kingdom). In brief,

amplicons for the targeted region underwent size selection, then equivalent amounts of PCR

product were end-repaired, A-tailed, and ligated with Illumina adapters. Library concentrations

were checked with Qubit and qPCR, and size distribution was checked with BioAnalyzer. Libraries

were pooled and sequenced on Illumina for pair-ended 250 bp at a depth of 30k per sample. After

sequencing, primers and adaptors were removed by Novogene, and raw reads were provided for

analysis.

The analysis of sequences was carried out in R (v. 4.3.1) (Team, 2023), which included

determination of amplicon sequence variants (ASVs), taxonomic assignment, visualization, and

statistical analyses. The dada2 pipeline (v. 1.30.0) (Callahan et al., 2016) was used for ASV and

taxonomic assignment. Reads were not trimmed due to the overall high quality of sequences, but

were filtered with the default parameters, with *maxEE* adjusted to (2,4). Reads were then paired

with *mergePairs*, *minOverlap* set to 16, and chimeras were removed. Taxonomy was assigned to

the species level where possible with the SILVA database (v. 138.1) (Glöckner et al., 2017).
Potential contaminants were then removed using SCRuB (v.0.0.1) (Austin et al., 2023) based on
the blank negative controls that were successfully sequenced. Mitochondria, chloroplasts, and
Eukarya were removed, resulting in 63.87% of ASVs with genus-level classification and 82.23%
with family-level. The mock community was then assessed to determine any potential bias created
by DNA extraction and sequencing. All eight bacterial genera present in the mock community
were validated.

The metabarcoding data was then analysed and visualized using the packages phyloseq
(v.1.46.0) (McMurdie & Holmes, 2013), microviz (v.0.12.0) (Barnett et al., 2021), and ggplot2
(v.3.4.4) (Wickham, 2011). The composition of the microbiome of ants and yogurts was visualized
with microviz. The beta diversity of the yogurts was visualized with phyloseq through a NMDS
ordination of their Bray-Curtis distances, resulting in a plot ordinated by yogurt treatments. The
alpha diversity of the yogurts was further determined with phyloseq. Last, the microbial loads of
Lactobacillaceae and Bacillaceae in the ant starters and yogurts were estimated by multiplying
the relative abundance of each genus, based on the metabarcoding data, by the quantified
bacterial load, based on 16S qPCR. This was visualized using ggplot2.

We assessed the effect of treatment (live, dehydrated, or frozen) on the beta diversity of the
yogurts using Bray-Curtis distances and an adonis PERMANOVA using the package vegan
(v.2.6.4) (Oksanen et al., 2013) and function *adonis2*. Subsequently, we performed a pairwise-
comparison between the three treatments based on the PERMANOVA using the package
pairwiseAdonis (v.0.4.1) (Martinez, 2017), which performs *p*-value adjustment for multiple
comparisons. The alpha diversity of the three yogurts was based on the Shannon index. We
performed a general linear model to determine the effect of treatment on the alpha diversity,
confirming the parametric assumptions of the model. Pairwise comparisons were then performed
with a Tukey HSD. Pairwise comparisons of the Lactobacillaceae and Bacillaceae load in the ant
starters and yogurts was performed with two sample t-tests for equal or unequal variance
depending upon the data.

Bacterial culturomics of milk ferments and ants

Bacteria from the ant yogurts and ants were cultured and identified with Sanger sequencing. 100
µL of yogurt sample was inoculated and plated in De Man, Rogosa and Sharpe (MRS) (VWR
Chemicals, US), Gifu Anaerobic Medium, modified (GAM) (HyServe GmbH & Co. KG, Germany),

yeast peptone dextrose (YPD), and Luria-Bertani (LB) media. YPD contained 10 g/L yeast extract,
20 g/L peptone, and 20 g/L D-glucose. LB contained 10 g/L peptone, 5 g/L yeast extract and 0.5
g/L sodium chloride. A total of 20 g/L agar was added for solid cultivations. For ants, individual
ants were inoculated into liquid media. Tubes and plates were incubated at 30°Cover 2 days, or
until growth was visible. All four media were cultivated in aerobic conditions and MRS and GAM
were also incubated anaerobically. Subcultures were plated and incubated in the same
conditions. Distinct bacterial colonies from plates were identified with Sanger sequencing of the
conserved 16S *rRNA* gene, which included colony PCR (cPCR) with primers 5´-
AGAGTTTGATCCTGGCTCAG-3´ and 5´-CCGTCAATTCCTTTRAGTTT-3´ (Cebeci & Gürakan,
2008). All cPCR reactions consisted of 12.5 µL of Red-Taq Mastermix (VWR, Denmark), 0.5 µL
of forward 10 µM primer, 0.5 µL of reverse 10 µM primer, 11.5 µL of water and a bacterial colony.
The reactions followed the same thermocycler program that consisted of an initial 4 min at 94 °C,
followed by 35 cycles of 94 °C for 30 s, 51 °C for 30 s, 72 °C for 1 min with a final 72 °C for 10
min. PCR products were Sanger sequenced (Eurofins Scientific SE). Results from sequencing
were taxonomically classified using BLAST and presented in a Sankey diagram
(app.rawgraphs.io).

Quantifying pH, lactic, acetic, and formic acid in yogurt

The pH and coagulation of all yogurts were assessed before and after fermentation. After
fermentation, the pH was measured from one sample of each of the groups, and the coagulation
of the treatment group was assessed. Given the abundance of formic acid found in *Formica* ants
(O'Rourke, 1950), we also examined the effect of formic acid on coagulation. Formic acid (Sigma
Aldrich, Denmark) was added to the sterile milk until pH 4.6, 5.15, 5.9, and 6.6, which represented
the range of pHs observed in conventional yogurts, ant yogurts, and sterile milk. The pH of the
negative control was also measured. Additionally, pH of all the samples was measured prior to
DNA extraction and enzyme analysis, after they had undergone a cycle of freezing and thawing.
This was performed on a separate pH reader than that which provided the initial measurement
before and after fermentation.

The amount of formic, lactic, and acetic acid in the samples was quantified with high-performance
liquid chromatography (HPLC). Yogurt samples were centrifuged at 5000 rpm for 5 min at 4°C,
and their supernatant was passed through a 0.2 µm syringe filter (Sigma Aldrich, Denmark). The
filtrate was analysed by a Dionex Ultimate 3000 HPLC system (Thermo Fisher Scientific, USA)
equipped with a refractive index detector and a Bio-Rad Aminex HPx87 column (Bio-Rad, USA).

The column oven temperature was set at 30°C. The flow rate was set at 0.6 mL/min with 5 mM
H_2SO_4 (Sigma Aldrich, Denmark) as an eluent. The injection volume was 10 μL. Peaks were
identified by comparison to the prepared standards, and integration of the peak areas was used
to quantify acids from obtained standard curves using the software Chromeleon7 (Thermo Fisher
Scientific, USA). Plots were visualised in GraphPad Prism. T-tests were used to compare organic
acid levels between samples, with Benjamini-Hochberg correction for multiple testing.

Proteomics of ant yogurts

Sample preparation for proteomics analysis was performed directly from the supernatant of the
yogurt samples after centrifugation for 10 min at 14,000 g and further processed as described
earlier (Kozaeva et al., 2021). After centrifugation, the supernatants were subjected to the
bicinchoninic acid (BCA) assay to estimate protein concentrations. Trypsin and LysC digestion
mix (Promega) was added to 20 μg protein of each sample and incubated for 8 h. Trifluoroacetic
acid was added to halt digestion and the samples were desalted using C18 resin (Empore, 3M)
before HPLC-MS analysis.

HPLC-MS analysis of the samples was performed on an Orbitrap Exploris 480 instrument
(Thermo Fisher Scientific) preceded by an EASY-nLC 1200 HPLC system (Thermo Fisher
Scientific). For each sample, 1 μg of peptides was captured on a 2 cm C18 trap column (Thermo
Fisher 164946). Subsequently separation was executed using a 70 min gradient from 8% (v/v) to
48% (v/v) of acetonitrile in 0.1% (v/v) formic acid on a 15-cm C18 reverse-phase analytical column
(Thermo EasySpray ES904) at a flow rate of 250 nL/min. For data-independent acquisition, the
mass spectrometer was run with the HRMS1 method (Xuan et al., 2020) preceded by the FAIMS
Pro Interface (Thermo Fisher Scientific) with a compensation voltage (CV) of -45 V. Full MS1
spectra were collected at a resolution of 120,000 and scan range of 400-1,000 m/z, with the
maximum injection time set to auto. MS2 spectra were obtained at a resolution of 60,000, with
the maximum injection time set to auto and the collision energy set to 32. Each cycle consisted
of three DIA experiments each covering a range of 200 m/z with a window size of 6 m/z and a 1
m/z overlap, while a full MS scan was obtained in between experiments.

To assign the proteases to their potential organisms of origin, we created a database consisting
of the proteases, peptidases and proteins based on UniProt reference proteomes, the gold
standard in proteomics. The proteomes consisted of the ant host, bacteria from the ant yogurt,
and two conventional yogurt bacteria. For the ant, we used the *F. execta* proteome, since the *F.*

polyctena proteome was not publicly available, and *F. execta* is a closely related species. From the *F. execta* proteome, we selected the proteases specifically. For all the bacteria included, we refined our databased to only include proteases and peptidases in the proteome. We selected bacteria most prevalent among the bacteria isolated from the live ant yogurt bacterial metabarcoding and culturomics: *Fructilactobacillus sanfranciscensis*, *Oecophylibacter saccharovorans*, and *Paenibacillus sp.* We also included *Wolbachia pipentis*, which is closely related to the obligate intracellular bacteria found in *Formica* ants (Jackson et al., 2022) and is represented in our analysis. We included the conventional yogurt bacteria *Streptococcus thermophilus* and *Lactobacillus delbrueckii* subsp. *bulgaricus.* *Streptococcus* was only present in low abundance in the bacteria metabarcoding data, and *L. delbrueckii* subsp. *bulgaricus* was isolated in one instance. We included them to determine if the ant yogurt contained proteases with the same or similar functionality. Last, we included common milk proteins, such as caseins, found in the *Bos taurus* proteome. For DIA data analysis, Spectronaut v18 (Biognosys, Switzerland) was used for protein identification and relative quantification of peptides. The default settings for "directDIA" were applied with an FDR cut-off of 1%, except for MS1 quantification for the peptides. Protein abundances were inferred from the peptide abundances using the MaxLFQ algorithm available within Spectronaut.

In the analysis of the results, first we determined if proteases, peptides, and milk proteins, changed in relative abundance between spring or autumn and live or dehydrated samples. We were unable to determine the abundance of a subset of proteases identified, likely due to their low abundance, and they were therefore not included in the analyses. For each protein, the average relative abundance was computed across all autumn samples and all spring samples. Similarly, the average relative abundance was computed across all live samples and all dehydrated samples.

The log2 fold change (log2FC) was calculated using the formula:

log2FC=$\log_2$(average abundance in condition 1/ average abundance in condition 2)

P-values were obtained by comparing the relative abundances of each protein across all samples and their replicates in the autumn versus spring conditions. Similarly, *p*-values were calculated by comparing the relative abundances of each protein across all samples and their replicates in the alive versus dehydrated conditions. The Benjamini-Hochberg procedure was applied to adjust the *p*-values for multiple comparisons, controlling the false discovery rate (FDR). The adjusted *p*-values were then transformed to their -log10 values for further analysis and visualization.

Second, we visualized the composition of the proteases and peptidases in GraphPad Prism. To

do so, we removed the milk proteins originating from *Bos taurus*, given they made up a high

proportion of the proteome. The aim was thus to clarify ant or bacterial contributions to the

enzymatic profile. The relative abundance of each protease or peptidase visualized was thus

based on the total abundance of proteases and peptidases in each sample, and not the proteome.

Data availability

Amplicon sequences from metabarcoding and isolated bacterial strains will be made available on

the SRA archive in GenBank upon publication. Proteome data will be made available on the RCSB

Protein Database Archive upon publication. Additional data frames will be made available on

FigShare upon publication.

Code availability

The code for analyses had been deposited in Zenodo upon publication.

Author contributions

Conceptualization: V.M.S., L.J.J., R.R.D., V.R-V., D.P.V., S.M.S; Methodology: V.M.S., V.R.-V.,

D.P.V, N.R.V, S.K.T.P, E.M.V, S.B.A; Formal analysis: V.M.S., V.R.-V., S.K.T.P; Investigation:

V.M.S., V.R.-V. D.P.V., S.M.S., N.R.V., A.C-M., S.K.T.P., E.M.V., D.Z.; Resources: L.J.J., S.B.A,

R.M., S.M.S.; Data curation: V.M.S.; Writing – original draft preparation: V.M.S., V.R-V., L.J.J.,

R.R.D; Writing – review & editing: all authors; Visualisation: V.M.S., V.R.-V.; Supervision: L.J.J.,

R.D.D., S.B.A., R.M.; Project administration: V.M.S., L.J.J.; Funding acquisition: S.B.A., L.J.J.,

R.R.D., R.M.

Acknowledgments

We thank Rasmus Stenbak Larsen for the assistance in collecting the ant colonies. We

acknowledge myrmecologist Dr. Vera Anatova from the Bulgaria Academy of Sciences, who

provided knowledge and assisted us with the permit НС ЗП-122/18.04.2023 from the Ministry of

Environment and Water to collect wood ants in Bulgaria. We thank villagers from Nova Mahala

for sharing their knowledge, especially Musserem, Mehmet, and Fatma. We acknowledge

gastronomers Fejsal Demiraj and Tangör Tan for their personal communications around the

traditional use of ants in dairy fermentations. Last, we thank the laboratory staff at the Globe

Institute for their expert support and guidance, especially Lasse Vinner, Tina Blumensaadt Brand,

and Pernille Selmer Olsen. V.M.S., A.C.M. and S.B.A. were supported by the Danish National

Research Foundation Centre for Evolutionary Hologenomics DNRF 143. V.R.V, and L.J.J. were

supported by the Novo Nordisk Foundation, NNF Grant number: NNF20CC0035580.

References

50 Best. (2024). *Alchemist*. William Reed Ltd. Retrieved 3 September 2024 from

706   https://www.theworlds50best.com/discovery/Establishments/Denmark/Copenhagen/Alch

707   emist.html

708   Amani, E., Eskandari, M. H., & Shekarforoush, S. (2017). The effect of proteolytic activity of starter

709   cultures on technologically important properties of yogurt. *Food Science & Nutrition*, *5*(3),

710   525-537. https://doi.org/10.1002/fsn3.427

711   Antonova, V., & Marinov, M. P. (2021). Red wood ants in Bulgaria: distribution and density related

712   to habitat characteristics. *Journal of Hymenoptera Research*, *85*, 135-159.

713   https://doi.org/10.3897/jhr.85.61431

714   Arranz-Otaegui, A., Carretero, L. G., Ramsey, M. N., Fuller, D. Q., & Richter, T. (2018).

715   Archaeobotanical evidence reveals the origins of bread 14,400 years ago in northeastern

716   Jordan. *Proc Natl Acad Sci U S A*, *115*(31), 7925-7930.

717   https://doi.org/10.1073/pnas.1801071115

718   Austin, G. I., Park, H., Meydan, Y., Seeram, D., Sezin, T., Lou, Y. C., Firek, B. A., Morowitz, M.

719   J., Banfield, J. F., Christiano, A. M., Pe'er, I., Uhlemann, A. C., Shenhav, L., & Korem, T.

720   (2023). Contamination source modeling with SCRuB improves cancer phenotype

721   prediction from microbiome data. *Nature Biotechnology*, *41*(12), 1820-+.

722   https://doi.org/10.1038/s41587-023-01696-w

723   Balzani, P., Dekoninck, W., Feldhaar, H., Freitag, A., Frizzi, F., Frouz, J., Masoni, A., Robinson,

724   E., Sorvari, J., & Santini, G. (2022). Challenges and a call to action for protecting European

725   red wood ants. *Conservation Biology*, *36*(6). https://doi.org/10.1111/cobi.13959

726   Barnett, D. J., Arts, I. C., & Penders, J. (2021). microViz: an R package for microbiome data

727   visualization and statistics. *Journal of Open Source Software*, *6*(63), 3201.

728   Blaimer, B. B., Brady, S. G., Schultz, T. R., Lloyd, M. W., Fisher, B. L., & Ward, P. S. (2015).

729   Phylogenomic methods outperform traditional multi-locus approaches in resolving deep

730   evolutionary history: a case study of formicine ants. *Bmc Evolutionary Biology*, *15*.

731   https://doi.org/10.1186/s12862-015-0552-5

732   Bordenstein, S. R., & Theis, K. R. (2015). Host Biology in Light of the Microbiome: Ten Principles

733   of Holobionts and Hologenomes. *PLoS Biol*, *13*(8).

734   https://doi.org/10.1371/journal.pbio.1002226

Borowiec, M. L., Cover, S. P., & Rabeling, C. (2021). The evolution of social parasitism in *Formica*
ants revealed by a global phylogeny. *Proc Natl Acad Sci U S A*, *118*(38).
https://doi.org/10.1073/pnas.2026029118

Brown, B. P., & Wernegreen, J. J. (2016). Deep divergence and rapid evolutionary rates in gut-
associated Acetobacteraceae of ants. *BMC Microbiol*, *16*(1), 140.
https://doi.org/10.1186/s12866-016-0721-8

Buchner, P. (1965). *Endosymbiosis of animals with plant microorgansims: revised English*
version. Interscience Publishers.

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P.
(2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature*
Methods, *13*(7), 581-+. https://doi.org/10.1038/Nmeth.3869

Canon, F., Nidelet, T., Guédon, E., Thierry, A., & Gagnaire, V. (2020). Understanding the
Mechanisms of Positive Microbial Interactions That Benefit Lactic Acid Bacteria Co-
cultures. *Front Microbiol*, *11*. https://doi.org/10.3389/fmicb.2020.02088

Carretero, L. G., Wollstonecroft, M., & Fuller, D. Q. (2017). A methodological approach to the
study of archaeological cereal meals: a case study at Catalhoyuk East (Turkey).
Vegetation History and Archaeobotany, *26*(4), 415-432. https://doi.org/10.1007/s00334-
017-0602-6

Cebeci, A., & Gürakan, G. C. (2008). Molecular methods for identification of *Lactobacillus*
delbrueckii subsp *bulgaricus* and *Streptococcus thermophilus* using methionine
biosynthesis and 16S rRNA genes. *Journal of Dairy Research*, *75*(4), 392-398.
https://doi.org/10.1017/S0022029908003543

Chua, K. O., Song, S. L., Yong, H. S., See-Too, W. S., Yin, W. F., & Chan, K. G. (2018). Microbial
Community Composition Reveals Spatial Variation and Distinctive Core Microbiome of the
Weaver Ant *Oecophylla smaragdina* in Malaysia. *Sci Rep*, *8*.
https://doi.org/10.1038/s41598-018-29159-2

Demirci, T., Akin, N., Öztürk, H. I., & Ogul, A. (2022). A metagenomic approach to homemade
back-slopped yogurts produced in mountainous villages of Turkey with the potential next-
generation probiotics. *Lwt-Food Science and Technology*, *154*.
https://doi.org/10.1016/j.lwt.2021.112860

Di Paola, M., Gori, A., Stefanini, I., Meriggi, N., Renzi, S., Nenciarini, S., Cerasuolo, B., Moriondo,
M., Romoli, R., Pieraccini, G., Baracchi, D., Turillazzi, F., Turillazzi, S., & Cavalieri, D.
(2023). Using wasps as a tool to restore a functioning vine grape mycobiota and preserve
the mycobial "terroir". *Sci Rep*, *13*(1). https://doi.org/10.1038/s41598-023-43541-9

Dietrich, R., Jessberger, N., Ehling-Schulz, M., Märtlbauer, E., & Granum, P. E. (2021). The Food
Poisoning Toxins of *Bacillus cereus*. *Toxins*, *13*(2).
https://doi.org/10.3390/toxins13020098

Duar, R. M., Lin, X. X. B., Zheng, J. S., Martino, M. E., Grenier, T., Pérez-Muñoz, M. E., Leulier,
F., Gänzle, M., & Walter, J. (2017). Lifestyles in transition: evolution and natural history of
the genus *Lactobacillus*. *FEMS Microbiol Rev*, *41*, S27-S48.
https://doi.org/10.1093/femsre/fux030

Evans, M., Lundy, J., Lucquin, A., Hagan, R., Kowalski, L., Wilczynki, J., Bickle, P., Adamczak,
K., Craig, O. E., Robson, H. K., & Hendy, J. (2023). Detection of dairy products from
multiple taxa in Late Neolithic pottery from Poland: an integrated biomolecular approach.
R Soc Open Sci, *10*(3). https://doi.org/10.1098/rsos.230124

Evershed, R. P., Payne, S., Sherratt, A. G., Copley, M. S., Coolidge, J., Urem-Kotsu, D., Kotsakis,
K., Ozdogan, M., Ozdogan, A. E., Nieuwenhuyse, O., Akkermans, P. M., Bailey, D.,
Andeescu, R. R., Campbell, S., Farid, S., Hodder, I., Yalman, N., Ozbasaran, M., Bicakci,
E., . . . Burton, M. M. (2008). Earliest date for milk use in the Near East and southeastern
Europe linked to cattle herding. *Nature*, *455*(7212), 528-531.
https://doi.org/10.1038/nature07180

FAO, & WHO. (2018). *Standard for Fermented Milks*.

Fernandez-Garcia, E., & McGregor, J. U. (1994). Determination of Organic-Acids during the
Fermentation and Cold-Storage of Yogurt. *Journal of Dairy Science*, *77*(10), 2934-2939.
https://doi.org/10.3168/jds.S0022-0302(94)77234-9

Fisberg, M., & Machado, R. (2015). History of yogurt and current patterns of consumption.
Nutrition Reviews, *73*, 4-7. https://doi.org/10.1093/nutrit/nuv020

Gänzle, M. G. (2014). Enzymatic and bacterial conversions during sourdough fermentation. *Food
Microbiology*, *37*, 2-10. https://doi.org/10.1016/j.fm.2013.04.007

Gänzle, M. G., Qiao, N. Z., & Bechtner, J. (2023). The quest for the perfect loaf of sourdough
bread continues: Novel developments for selection of sourdough starter cultures.
International Journal of Food Microbiology, *407*.
https://doi.org/10.1016/j.ijfoodmicro.2023.110421

Gänzle, M. G., & Zheng, J. S. (2019). Lifestyles of sourdough lactobacilli - Do they matter for
microbial ecology and bread quality? *International Journal of Food Microbiology*, *302*, 15-
23. https://doi.org/10.1016/j.ijfoodmicro.2018.08.019

Gassem, M. A., & Frank, J. F. (1991). Physical-Properties of Yogurt Made from Milk Treated with
Proteolytic-Enzymes. *Journal of Dairy Science*, *74*(5), 1503-1511.
https://doi.org/10.3168/jds.S0022-0302(91)78310-0

Glöckner, F. O., Yilmaz, P., Quast, C., Gerken, J., Beccati, A., Ciuprina, A., Bruns, G., Yarza, P.,
Peplies, J., Westram, R., & Ludwig, W. (2017). 25 years of serving the community with
ribosomal RNA gene reference databases and tools. *Journal of Biotechnology*, *261*, 169-
176. https://doi.org/10.1016/j.jbiotec.2017.06.1198

Grigoroff, S. (1905). Etude sur le lait fermenté comestible : le "Kissélo-mléko" de Bulgarie. *Revue*
médicale de la Suisse romande : organe officiel de la Société médicale de la Suisse
romande.

Guénard, B., Weiser, M. D., Gómez, K., Narula, N., & Economo, E. P. (2017). The Global Ant
Biodiversity Informatics (GABI) database: synthesizing data on the geographic distribution
of ant species (Hymenoptera: Formicidae). *Myrmecological News*, *24*, 83-89. <Go to
ISI>://WOS:000395966600005

Guide, M. (2024). *Full List of Micehlin Star Restaurants in the Nordic Countries 2024*. Michelin
Guide. Retrieved 3 September 2024 from
https://guide.michelin.com/dk/en/article/michelin-guide-ceremony/full-list-of-michelin-
star-restaurants-in-the-nordic-countries

Hendy, J., Colonese, A. C., Franz, I., Fernandes, R., Fischer, R., Orton, D., Lucquin, A., Spindler,
L., Anvari, J., Stroud, E., Biehl, P. F., Speller, C., Boivin, N., Mackie, M., Jersie-
Christensen, R. R., Olsen, J. V., Collins, M. J., Craig, O. E., & Rosenstock, E. (2018).
Ancient proteins from ceramic vessels at Catalhoyuk West reveal the hidden cuisine of
early farmers. *Nat Commun*, *9*. https://doi.org/10.1038/s41467-018-06335-6

Hu, Y. Y., Zhang, L., Wen, R. X., Chen, Q., & Kong, B. H. (2022). Role of lactic acid bacteria in
flavor development in traditional Chinese fermented foods: A review. *Critical Reviews in*
Food Science and Nutrition, *62*(10), 2741-2755.
https://doi.org/10.1080/10408398.2020.1858269

Ivens, A. B. F., & Kronauer, D. J. C. (2022). Aphid-farming ants. *Current biology*, *32*(15), R813-
R817. https://doi.org/10.1016/j.cub.2022.06.072

Jackson, R., Patapiou, P. A., Golding, G., Helanterä, H., Economou, C. K., Chapuisat, M., &
Henry, L. M. (2023). Evidence of phylosymbiosis in *Formica* ants. *Front Microbiol*, *14*.
https://doi.org/10.3389/fmicb.2023.1044286

Jackson, R. A. H. E. L., Monnin, D., Patapiou, P. A., Golding, G. M., Helanterä, H., Oettler, J.,
Heinze, J., Wurm, Y. N., Economou, C. K., Chapuisat, M., & Henry, L. M. (2022).

Convergent evolution of a labile nutritional symbiosis in ants. *Isme Journal*, *16*(9), 2114-
2122. https://doi.org/10.1038/s41396-022-01256-1

Janicki, J., Narula, N., Ziegler, M., Guénard, B., & Economo, E. P. (2016). Visualizing and
interacting with large-volume biodiversity data using client-server web-mapping
applications: The design and implementation of antmaps.org. *Ecological Informatics*, *32*,
185-193. https://doi.org/10.1016/j.ecoinf.2016.02.006

Jensen, A. B., Malagocka, J., Eilenberg, J., & Fredensborg, B. L. (2017). Viability of *Dicrocoelium*
dendriticum metacercariae in *Formica polyctena* ants after exposure to different
treatments. *Journal of Insects as Food and Feed*, *3*(1), 15-20.
https://doi.org/10.3920/Jiff2016.0042

Jia, B. L., Li, Z. Q., Liu, J. L., Sun, Y., Jia, X. M., Xuan, Y. H., Zhang, J. Y., & Jeon, C. O. (2015).
A Zinc-Dependent Protease AMZ-tk from a Thermophilic Archaeon is a New Member of
the Archaemetzincin Protein Family. *Front Microbiol*, *6*.
https://doi.org/10.3389/fmicb.2015.01380

Kaczmarczyk-Ziemba, A., Zagaja, M., Wagner, G. K., Pietrykowska-Tudruj, E., & Staniec, B.
(2020). First Insight into Microbiome Profiles of Myrmecophilous Beetles and Their Host,
Red Wood Ant *Formica polyctena* (Hymenoptera: Formicidae)-A Case Study. *Insects*,
11(2). https://doi.org/10.3390/insects11020134

Kline, L., & Sugihara, T. F. (1971). Microorganisms of the San Francisco Sour Dough Bread
Process. *Applied Microbiology*, *21*(3), 459-465. https://doi.org/doi:10.1128/am.21.3.459-
465.1971

Kothe, C. I., Rasmussen, J. A., Mak, S. S. T., Gilbert, M. T. P., & Evans, J. (2024). Exploring the
microbial diversity of novel misos with metagenomics. *Food Microbiology*, *117*.
https://doi.org/10.1016/j.fm.2023.104372

Kozaeva, E., Volkova, S., Matos, M. R. A., Mezzina, M. P., Wulff, T., Volke, D. C., Nielsen, L. K.,
& Nikel, P. I. (2021). Model-guided dynamic control of essential metabolic nodes boosts
acetyl-coenzyme A-dependent bioproduction in rewired. *Metabolic Engineering*, *67*, 373-
386. https://doi.org/10.1016/j.ymben.2021.07.014

Landis, E. A., Oliverio, A. M., McKenney, E. A., Nichols, L. M., Kfoury, N., Biango-Daniels, M.,
Shell, L. K., Madden, A. A., Shapiro, L., Sakunala, S., Drake, K., Robbat, A., Booker, M.,
Dunn, R. R., Fierer, N., & Wolfe, B. E. (2021). The diversity and function of sourdough
starter microbiomes. *Elife*, *10*. https://doi.org/10.7554/eLife.61644

Löfqvist, J. (1976). Formic acid and saturated hydrocarbons as alarm pheromones for the ant
Formica rufa. *Journal of Insect Physiology*, *22*(10), 1331-1346.

Louw, N. L., Lele, K., Ye, R., Edwards, C. B., & Wolfe, B. E. (2023). Microbiome Assembly in
Fermented Foods. *Annual Review of Microbiology*, *77*, 381-402.
https://doi.org/10.1146/annurev-micro-032521-041956

Madden, A. A., Epps, M. J., Fukami, T., Irwin, R. E., Sheppard, J., Sorger, D. M., & Dunn, R. R.
(2018). The ecology of insect-yeast relationships and its relevance to human industry.
Proceedings of the Royal Society B-Biological Sciences, *285*(1875).
https://doi.org/10.1098/rspb.2017.2733

Margulis, L., & Fester, R. (1991). *Symbiosis as a source of evolutionary innovation: speciation*
and morphogenesis. MIT press.

Martinez, A. P. (2017). Pairwiseadonis: Pairwise multilevel comparison using Adonis. R package
version 1.

Maurizi, M. R., Clark, W. P., Katayama, Y., Rudikoff, S., Pumphrey, J., Bowers, B., & Gottesman,
S. (1990). Sequence and Structure of Clp-P, the Proteolytic Component of the Atp-
Dependent Clp Protease of *Escherichia coli*. *Journal of Biological Chemistry*, *265*(21),
12536-12545. <Go to ISI>://WOS:A1990DP45800065

McMurdie, P. J., & Holmes, S. (2013). phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. *PLoS One*, *8*(4).
https://doi.org/10.1371/journal.pone.0061217

Metchnikoff, I., I. . (1908). *The Prolongation of Life*. Putnman.

Michaylova, M., Minkova, S., Kimura, K., Sasaki, T., & Isawa, K. (2007). Isolation and
characterization of *Lactobacillus delrueckii* ssp *bulgaricus* and *Streptococcus*
thermophilus from plants in Bulgaria. *FEMS microbiology letters*, *269*(1), 160-169.
https://doi.org/10.1111/j.1574-6968.2007.00631.x

Multu Sirakova, S. (2023). Forgotten Stories of Yogurt: Cultivating Multispecies Wisdom. *Journal*
of Ethnobiology, *43*(3), 250-261. https://doi.org/10.1177/02780771231194779

Munieweg, F. R., Gaviao, E. R., Czarnobay, M., Dilda, A., Stefani, L. D. M., & Nespolo, C. R.
(2021). Mascarpone cheese from sheep's milk - a new option for the consumer. *Food*
Science and Technology, *41*, 568-575. https://doi.org/10.1590/fst.32420

Nagaoka, S. (2019). Yogurt Production. In M. Kanauchi (Ed.), *Lactic Acid Bacteria: Methods and*
Protocols (pp. 45-54). Springer New York. https://doi.org/10.1007/978-1-4939-8907-2_5

Nicholson, W. L., Munakata, N., Horneck, G., Melosh, H. J., & Setlow, P. (2000). Resistance of
Bacillus endospores to extreme terrestrial and extraterrestrial environments. *Microbiology*
and Molecular Biology Reviews, *64*(3), 548-+. https://doi.org/10.1128/Mmbr.64.3.548-
572.2000

O'Rourke, F. J. (1950). Formic acid production among the Formicidae. *Annals of the*
Entomological Society of America, *43*(3), 437-443.

Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'hara, R., Simpson, G. L.,
Solymos, P., Stevens, M. H. H., & Wagner, H. (2013). Package 'vegan'. *Community*
ecology package, version, *2*(9), 1-295.

Öney Tan, A. (2010). Yoghurt in the Turkish Kitchen. In H. Saberi (Ed.), *Cured, Smoked, and*
Fermented: Proceedings of the Oxford Symposium on Food and Cooking (pp. 322-333).
Prospect Books.

Reese, A. T., Madden, A. A., Joossens, M., Lacaze, G., & Dunn, R. R. (2020). Influences of
Ingredients and Bakers on the Bacteria and Fungi in Sourdough Starters and Bread.
Msphere, *5*(1). https://doi.org/10.1128/mSphere.00950-19

Robinson, E. J., & Stockan, J. A. (2020). Wood Ants (*Formica rufa* Species Group). In C. Starr
(Ed.), *Encyclopedia of Social Insects*. Switzerland: Springer Nature.

Robinson, R. K., Lucey, J. A., & Tamime, A. Y. (2006). Manufacture of Yogurt. In A. Tamime (Ed.),
Fermented Milks (pp. 53-75). Blackwell Science.

Salque, M., Bogucki, P. I., Pyzel, J., Sobkowiak-Tabaka, I., Grygiel, R., Szmyt, M., & Evershed,
R. P. (2013). Earliest evidence for cheese making in the sixth millennium BC in northern
Europe. *Nature*, *493*(7433), 522-525. https://doi.org/10.1038/nature11698

Santivarangkna, C., Kulozik, U., & Foerst, P. (2011). Storing Lactic Acid Bacteria: Current
Methodologies and Physiological Implications. *Stress Responses of Lactic Acid Bacteria*,
479-504. https://doi.org/10.1007/978-0-387-92771-8_20

Scott, A., Reinhold, S., Hermes, T., Kalmykov, A. A., Belinskiy, A., Buzhilova, A., Berezina, N.,
Kantorovich, A. R., Maslov, V. E., Guliyev, F., Lyonnet, B., Gasimov, P., Jalilov, B., Eminli,
J., Iskandarov, E., Hammer, E., Nugent, S. E., Hagan, R., Majander, K., . . . Warinner, C.
(2022). Emergence and intensification of dairying in the Caucasus and Eurasian steppes.
Nat Ecol Evol, *6*(6), 813-+. https://doi.org/10.1038/s41559-022-01701-6

Sedó Melina, G. E., Shetty, R., Xiao, H., Watjen, A. P., Tovar, M., & Bang-Berthelsen, C. H.
(2022). Development of a novel lactic acid bacteria starter culture approach: From insect
microbiome to plant-based fermentations. *Lwt-Food Science and Technology*, *167*.
https://doi.org/10.1016/j.lwt.2022.113797

Seifert, B. (2021). A taxonomic revision of the Palaearctic members of the *Formica rufa* group
(Hymenoptera: Formicidae) - the famous mound-building red wood ants. *Myrmecological*
News, *31*, 133-179. https://doi.org/10.25849/myrmecol.news_031:133

Sieuwerts, S. (2016). Microbial interactions in the yoghurt consortium: current status and product
implications. *SOJ Microbiology & Infectious Diseases*, *4*(2), 01-05.

Sieuwerts, S., Bron, P. A., & Smid, E. J. (2018). Mutually stimulating interactions between lactic
acid bacteria and *Saccharomyces cerevisiae* in sourdough fermentation. *Lwt-Food*
Science and Technology, *90*, 201-206. https://doi.org/10.1016/j.lwt.2017.12.022

Sieuwerts, S., de Bok, F. A. M., Hugenholtz, J., & Vlieg, J. E. T. V. (2008). Unraveling microbial
interactions in food fermentations: from classical to genomics approaches. *Appl Environ*
Microbiol, *74*(16), 4997-5007. https://doi.org/10.1128/Aem.00113-08

Sieuwerts, S., Molenaar, D., van Hijum, S. A. F. T., Beerthuyzen, M., Stevens, M. J. A., Janssen,
P. W. M., Ingham, C. J., de Bok, F. A. M., de Vos, W. M., & Vlieg, J. E. T. V. (2010). Mixed-
Culture Transcriptome Analysis Reveals the Molecular Basis of Mixed-Culture Growth in
Streptococcus thermophilus and *Lactobacillus bulgaricus*. *Appl Environ Microbiol*, *76*(23),
7775-7784. https://doi.org/10.1128/Aem.01122-10

Sinotte, V. M., Conlon, B. H., Seibel, E., Schwitalla, J. W., de Beer, Z. W., Poulsen, M., & Bos, N.
(2021). Female-biased sex allocation and lack of inbreeding avoidance in termites.
Ecology and Evolution, *11*(10), 5598-5605. https://doi.org/10.1002/ece3.7462

Sinotte, V. M., Multu Sirakova, S., & Zilber, D. (2023). Preserving the Past in the Present:
Forgotten Flavors of Yogurt. *Serviette*, (03).

Stefanini, I., Dapporto, L., Legras, J. L., Calabretta, A., Di Paola, M., De Filippo, C., Viola, R.,
Capretti, P., Polsinelli, M., Turillazzi, S., & Cavalieri, D. (2012). Role of social wasps in
Saccharomyces cerevisiae ecology and evolution. *Proc Natl Acad Sci U S A*, *109*(33),
13398-13403. https://doi.org/10.1073/pnas.1208362109

Team, R. C. (2023). *R: A Language and Environment for Statistical Computing*. In R Foundation
for Statistical Computing. https://www.R-project.org

Tragust, S., Herrmann, C., Hafner, J., Braasch, R., Tilgen, C., Hoock, M., Milidakis, M. A., Gross,
R., & Feldhaar, H. (2020). Formicine ants swallow their highly acidic poison for gut
microbial selection and control. *Elife*, *9*. https://doi.org/10.7554/eLife.60287

Velikova, P., Petrov, K., Lozanov, V., Tsvetanova, F., Stoyanov, A., Wu, Z. J., Liu, Z. M., &
Petrova, P. (2018). Microbial diversity and health-promoting properties of the traditional
Bulgarian yogurt. *Biotechnology & Biotechnological Equipment*, *32*(5), 1205-1217.
https://doi.org/10.1080/13102818.2018.1475255

Walter, S. B., Norbert. (2021). Emerging Bees: Identification and Possible Meanings of Insect
Figures at Göbekli Tepe. In J. H. Daujat, Angelos; Berthon, Rémi; Chahoud, Jwana;

Kassianidou, Vasiliki; Vigne, Jean-Denis (Ed.), *Archaeozoology of Southwest Asia and*
Adjacent Areas XIII (pp. 233-244). Lockwood Press.

Wang, Y. Q., Wu, J. T., Lv, M. X., Shao, Z., Hungwe, M., Wang, J. J., Bai, X. J., Xie, J. L., Wang,
Y. P., & Geng, W. T. (2021). Metabolism Characteristics of Lactic Acid Bacteria and the
Expanding Applications in Food Industry. *Frontiers in Bioengineering and Biotechnology*,
9. https://doi.org/10.3389/fbioe.2021.612285

Warinner, C., Hendy, J., Speller, C., Cappellini, E., Fischer, R., Trachsel, C., Arneborg, J.,
Lynnerup, N., Craig, O. E., Swallow, D. M., Fotakis, A., Christensen, R. J., Olsen, J. V.,
Liebert, A., Montalva, N., Fiddyment, S., Charlton, S., Mackie, M., Canci, A., . . . Collins,
M. J. (2014). Direct evidence of milk consumption from ancient human dental calculus. *Sci*
Rep, *4*. https://doi.org/10.1038/srep07104

Wickham, H. (2011). ggplot2. *Wiley interdisciplinary reviews: computational statistics*, *3*(2), 180-
185.

Wondrich, D. (2010). *Punch: The Delights (and Dangers) of the Flowing Bowl*. Penguin.

Wouters, J. T. M., Ayad, E. H. E., Hugenholtz, J., & Smit, G. (2002). Microbes from raw milk for
fermented　dairy　products.　*International　Dairy　Journal*,　*12*(2-3),　91-109.
https://doi.org/10.1016/S0958-6946(01)00151-0

Xuan, Y., Bateman, N. W., Gallien, S., Goetze, S., Zhou, Y., Navarro, P., Hu, M., Parikh, N., Hood,
B. L., Conrads, K. A., Loosse, C., Kitata, R. B., Piersma, S. R., Chiasserini, D., Zhu, H.
W., Hou, G. X., Tahir, M., Macklin, A., Khoo, A., . . . Conrads, T. P. (2020). Standardization
and harmonization of distributed multi-center proteotype analysis supporting precision
medicine studies. *Nat Commun*, *11*(1). https://doi.org/10.1038/s41467-020-18904-9

Yalman, A. R. (1977). *Cenupta Türkmen oymakları* (Vol. II). Külür bakanlığı yayınları.

Yang, X., Zhang, Z. M., Zhang, W. B., Qiao, H. J., Wen, P. C., & Zhang, Y. (2022). Proteomic
analysis, purification and characterization of a new milk-clotting protease from *Tenebrio*
molitor larvae. *Journal of Functional Foods*, *89*. https://doi.org/10.1016/j.jff.2022.104944

Zapasnik, A., Sokolowska, B., & Bryla, M. (2022). Role of Lactic Acid Bacteria in Food
Preservation and Safety. *Foods*, *11*(9). https://doi.org/10.3390/foods11091283

Zheng, J. S., Wittouck, S., Salvetti, E., Franz, C. M. A. P., Harris, H. M. B., Mattarelli, P., O'Toole,
P. W., Pot, B., Vandamme, P., Walter, J., Watanabe, K., Wuyts, S., Felis, G. E., Gänzle,
M. G., & Lebeer, S. (2020). A taxonomic note on the genus *Lactobacillus*: Description of
23 novel genera, emended description of the genus *Lactobacillus* Beijerinck 1901, and
union of Lactobacillaceae and Leuconostocaeae. *International Journal of Systematic and*
Evolutionary Microbiology, *70*(4), 2782-2858. https://doi.org/10.1099/ijsem.0.004107

Zheng, Z., Zhao, M. Q., Zhang, Z. J., Hu, X., Xu, Y., Wei, C., & He, H. (2022). Lactic Acid Bacteria
Are Prevalent in the Infrabuccal Pockets and Crops of Ants That Prefer Aphid Honeydew.
Front Microbiol, *12*. https://doi.org/10.3389/fmicb.2021.785016

Zilber-Rosenberg, I., & Rosenberg, E. (2008). Role of microorganisms in the evolution of animals
and plants: the hologenome theory of evolution. *FEMS Microbiol Rev*, *32*(5), 723-735.
https://doi.org/10.1111/j.1574-6976.2008.00123.x

Zuccon, D., Brisset, J., Corbari, L., Puillandre, N., Utge, J., & Samadi, S. (2012). An optimised
protocol for barcoding museum collections of decapod crustaceans: a case-study for a 10-
40-years-old collection. *Invertebrate Systematics*, *26*(5-6), 592-600.
https://doi.org/10.1071/Is12027

1013

1014

Figure 1. Hypothesized contributions of the ant holobiont to yogurt fermentation. *Formica* red
wood ants create characteristic thatched mounds (left), which may have been used in traditional
yogurt fermentation practices (Multu Sirakova 2023). The traits of the ant holobiont, the composite
of both ant and the microbes found within it, are hypothesized to act as the starter of the yogurt
fermentation (centre). These include the microbes (lactic acid and acetic acid bacteria) that are
found in and on the ants, acids that are produced by the ant and the ant-associated microbes,
and enzymes that have ant or microbial origins.
Figure supplement 1. Experimental workflow to characterize microbes, acids, and proteases in
ant yogurt.

1024

Figure 2. Ants as culinary catalysts of milk fermentation and coagulation. (**A-C**) Photographs of
traditional yogurt fermentation initiated by *Formica* ants taken during field work in Bulgaria. (**A**)
Ants were added to warmed milk that was then (**B**) buried within the ant colony and left to ferment
overnight. (**C**) The resulting milk had started to coagulate and acidify, indicative of early stages of
yogurt fermentation. (**D-F**) Culinary applications created by restaurant Alchemist's research and
development team using *F. rufa* ants: (**D**) ant yogurt ice cream sandwich, (**E**) ant "mascarpone-
like" cheese, and (**F**) milk-wash cocktail (Photos a-c by David Zilber, and d-f by Søren
Gammelmark and Kåre Knudsen of Restaurant Alchemist).

Figure 3. *Formica* ants host a stable, seasonal microbiome, and preparation of the ants affects the corresponding yogurt microbiome. (**A**) European species ranges of *F. polyctena* and *F. rufa* (top), both within the *F. rufa* group of red wood ants (maps modified from antmaps.org). The microbiome composition of ant species across seasons (bottom). Bar represents four pooled ants from a colony of each respective species, collected in Denmark. (**B**) The bacterial microbiome of the three ant preparations and corresponding ant yogurts made in spring and autumn. Bars represent replicates. Live ant microbiomes are also represented in panel (**A**), and live and frozen ants are identical. They are shown for direct comparison to the corresponding yogurts. (**C**) The alpha diversity of ant yogurts. Letters indicate pairwise statistical differences (Tukey HSD: *p*<0.05), and boxes illustrate standard deviations. (**D**) The beta diversity of ant yogurts based on

1044   Bray-Curtis distances and Non-Metric Multidimensional Scaling (NMDS) analysis. The grouping

1045   is supported by a PERMANOVA.

1046   **Figure supplement 1.** Genus-level microbiomes and seasonal bacterial loads of ants.

1047   **Figure supplement 2.** Detection bias of bacterial abundances based on mock community control.

1048

1050

Figure 4. Bacterial biomass and culturable diversity of lactic acid and spore-forming bacteria vary based on the preparation of ant yogurts. (**A**) The estimated amount of lactic acid bacteria (Lactobacillaceae) DNA, representative of lactic acid bacterial load, in the live ants and resulting yogurts across seasons. (**B**) The diversity of culturable species of lactic acid bacteria from the ants and yogurts, where the majority of isolates stem from live ants and live ant yogurts. The thickness of each line approximates abundance of culturable bacteria by representing the number of microbial media plates where the bacteria grew compared to other samples. (**C**) The amount of DNA from spore-forming Bacillaceae, indicative of Bacillaceae bacterial load, is exceptionally low in dehydrated and frozen ants and increases in the respective yogurt fermentations. (**D**) Culturable species of Bacillaceae isolated from ants and yogurts. Frozen ant yogurts were not made in autumn. Line thickness represents an approximation of culturable bacteria as above. (* = $p \leq .05$; ** = $p \leq .01$; *** = $p \leq .001$; n.s. = p>0.05; y-axis present on a logarithmic scale).

Figure supplement 1. Bacterial load and relative abundance of Lactobacillaceae and Bacillaceae.

Figure supplement 2. All bacteria cultured from ants and ant yogurts in spring and autumn.

1066

Figure 5. Organic acids, proteases, and peptidases in ant yogurts. (**A**) Concentration of metabolites formic acid, lactic acid, and acetic acid in the yogurts from spring (S) and autumn (A). The samples include yogurts made with live, dehydrated, and frozen ants, formic acid, and three conventional yogurts as controls (C1-3). Data points represent biological replicates and error bars represent the standard deviation. Statistical analysis was performed using Student's t-test (two-sample unequal variance; * = P ≤ .05; ** = P ≤ .01; *** = P ≤ .001). (**B**) Proteases and peptidases detected in live and dehydrated ant yogurt samples. Proteases and peptides originate from the ant holobiont, including the ant and its bacteria. It also includes bacteria found in conventional yogurt, that were potentially present in low abundances. The rows represent biological replicates. Milk proteins were removed to improve the visualization of the proteases and peptidases. Milk proteins made up 96.19 ± 1.07% of the samples. The absolute abundance of all proteins in the proteomes can be found Figure 5 – source data 1.

Figure supplement 1. Bacterial load and pH of ant yogurts, conventional yogurts, and controls.

1080   **Figure supplement 2.** Differential abundance of ant yogurt proteases across seasons and

1081   elaborations.

1082   **Source data 1.** pH of yogurts and formic acid controls before and after incubation.

1083   **Source data 2.** Absolute abundance of milk proteins, proteases, and peptidases in proteomes.

Ant Holobiont

Ant
Microbes

Acids

Enzymes

Formica ant colony

Yogurt

A

Lactobacillaceae ng DNA

Spring | Autumn

*

**

ants / yogurt

B

Spring

Frozen yogurt
Dehydrated yogurt
Crushed dehydrated ants

Live yogurt

Live ants

Fructilactobacillus sanfranciscensis

Lactobacillus delbrueckii subsp. bulgaricus

Lactiplantibacillus pentosus

Lactobacillus salivarus

Lactobacillus plantarum

Isolation source | Cultured **Lactobacillaceae**

Autumn

Live Yogurt

Live ants
Dehydrated ants
Crushed dehydrated ants

Fructilactobacillus sanfranciscensis

Lacticaseibacillus casei

Isolation source | Cultured **Lactobacillaceae**

C

Bacillaceae ng DNA

Dehydrated spring | Dehydrated autumn | Frozen spring

n.s. | n.s. | *

ants / yogurt

D

Spring

Dehydrated ants
Dehydrated yogurt
Live yogurt

Live ants

Frozen yogurt

Crushed dehydrated ants
Frozen ants

Margalitia shackletonii
Niallia sp.
Bacillus weihenstephanensis
Bacillus mycoides
Bacillus paramycoides
Lysinbacillus fusiformis
Viridibacillus sp.
Bacillus sp.
Bacillus subtilis
Bacillus cereus
Bacillus thuringiensis

Isolation source | Cultured **Bacillaceae**

Autumn

Dehydrated yogurt

Crushed dehydrated ants

Dehydrated ants

Live Ants

Bacillus tequilensis
Bacillus subtilis
Bacillus velezensis
Bacillus thuringiensis
Bacillus sp.
Bacillus cereus
Bacillus paramycoides

Isolation source | Cultured **Bacillaceae**

A

Formic acid

Lactic acid

Acetic acid

Concentration (g/L)

Seasonal yogurts and controls

Yogurt type

- Live ants
- Dehydrated ants
- Frozen ants
- Formic acid control
- Conventional yogurt

B

Origin	Protease or peptidase
Streptococcus thermophilus	Foldase protein PrsA
Lactobacillus delbrueckii subsp. *bulgaricus*	Alpha beta-hydrolase
	Beta-hexosaminidase
Paenibacillus sp.	LD-carboxypeptidase
Fructolactobacillus sanfranciensis	ATP-dependent Clp-protease ∗
	Zinc metalloprotease ∗
Wolbachia pipientis	Family phage prohead protease
	Family peptidase
Formica ants	ATP-dependent Clp-protease ∗
	Other proteases or peptidases

∗ putatively caseinolytic

Relative abundance of proteases and peptidases