1   The consequences of host heterogeneity for parasite transmission depend on how host

2   infectiousness is determined.

Jacob A. Cohen[1,2] (jacob.cohen@biology.ox.ac.uk), Andrew D. Dean[1] (andrew.dean@liverpool.ac.uk),

Mark Viney[1] (mark.viney@liverpool.ac.uk), Andy Fenton[1] (a.fenton@liverpool.ac.uk)

1. Department of Biology, University of Oxford, 11a Mansfield Road, Oxford OX1 3SZ, UK.

2. Department of Evolution, Ecology and Behaviour, Institute of Infection, Veterinary and Ecological

Sciences, University of Liverpool, Liverpool L69 7ZB, UK.

Correspondence to: Jacob Cohen. University of Oxford, Peter Medawar Building for Pathogen

Research, 3 South Parks Road, Oxford OX1 3SY. jacob.cohen@biology.ox.ac.uk.

Acknowledgements: JAC was funded by a PhD studentship from the 'Adapting to the Challenges of a

Changing Environment (ACCE)' Doctoral Training Partnership funded by the Natural Environment

Research Council (NERC) (Grant NE/S00713X/1). ADD was funded by a NERC Pushing the Frontiers

grant (Grant NE/X01424X/1).

Abstract

Host heterogeneities in susceptibility and infectiousness can affect parasite transmission, but standard

analyses typically consider a limited range of epidemiological scenarios. We propose two contrasting

phenomenological scenarios that capture a wide range of processes by which host infectiousness is

determined, and explore how these scenarios affect the impact of host heterogeneities on parasite

transmission. Specifically, we contrast 'Recipient dependence' (RD), where a host's infectiousness is a

fixed characteristic of the recipient host being infected, with 'Donor dependence' (DD), where a host's

infectiousness is determined by the donor host that infected them. Contrasting Susceptible-Infected

models of these two phenomenological scenarios show that under RD, R_0 is driven by population-level

covariance between susceptibility and infectiousness, but for DD scenarios it is driven by the maximum

infectiousness present in the population. Our results show that these different scenarios, which

capture different ways by which host infectiousness is determined, should be considered explicitly

when exploring the consequences of host heterogeneity on transmission.

Introduction

Individuals can vary substantially in their propensities to be infected by and transmit parasites (VanderWaal *et al.* 2016). These individual-level host heterogeneities can have significant effects on the transmission of parasites and so affect patterns of infection in host populations. Superspreaders – hosts that are responsible for a disproportionate number of transmission events – are a well-known, if extreme, example of such host heterogeneity (Woolhouse *et al.* 1997; Lloyd-Smith *et al.* 2005; Lemieux *et al.* 2021). Despite the commonly accepted importance of host heterogeneity, there has been somewhat limited investigation of this, particularly in understanding the epidemiological consequences of how a host's infectiousness is determined.

Parasite transmission is a function of three factors: (i) host susceptibility, which is the host's propensity to become infected following parasite exposure, (ii) host infectiousness, the capacity of an infected host to transmit parasites, and (iii) host contact rate, the rate of transmission-relevant contacts, dependent on the transmission mode of the parasite in question (McCallum *et al.* 2017). Variation among individual hosts in any of these traits can alter parasite transmission dynamics in a host population (Dwyer *et al.* 1997; Coutinho *et al.* 1999; Barlow 2000; Matthews *et al.* 2006; Streicker *et al.* 2013; Kong *et al.* 2016; Stephenson *et al.* 2017; White *et al.* 2018). Importantly, heterogeneities in these three traits can exist simultaneously (so-called 'coupled heterogeneities'), and potentially interact with each other, altering transmission dynamics (Yates *et al.* 2006; Miller 2007; Hickson *et al.* 2014; Vazquez-Prokopec *et al.* 2016). For instance, it has been shown that covariation between host heterogeneities can increase or decrease the parasite's basic reproduction number, R_0, depending on the traits considered and whether the covariation is positive or negative (Vazquez-Prokopec *et al.* 2016; Lloyd *et al.* 2020).

To date, most studies investigating the consequences of host heterogeneities on parasite transmission dynamics implicitly assume that a host's infectiousness is a fixed characteristic of the 'recipient' individual being infected (Coutinho *et al.* (1999); Yates *et al.* (2006); Miller (2007); Lloyd *et al.* (2020)).

We call this 'Recipient dependence' (RD) because the infectiousness of the newly infected host (the

recipient) is solely determined by characteristics of that individual. There are a range of processes

which may lead to RD determination of host infectiousness. For example, hosts might differ in terms

of genotype (Anacleto *et al.* 2019), nutritional status (Cornet *et al.* 2014), behavioural phenotype

(Araujo *et al.* 2016), or social connectedness (Mantzaris *et al.* 2013; Adelman *et al.* 2015), all of which

result in levels of infectiousness linked to those characteristics of the recipient host. Generally, most

studies of host heterogeneities implicitly assume RD-based scenarios (*e.g.* Coutinho *et al.* (1999); Yates

et al. (2006); Miller (2007); Lloyd *et al.* (2020)). In such systems, heterogeneity can either increase or

decrease R_0, depending on whether the covariance between host susceptibility and infectiousness is

positive or negative (Lloyd *et al.* 2020).

We contrast the RD scenario with scenarios where the infectiousness of a newly infected host depends

on characteristics of the 'donor' host that infected it. We call this 'Donor dependence' (DD) because

the infectiousness of the newly infected host is determined by the infecting host (the donor). DD may

occur, for example, when the parasite load received by a newly infected host determines how

infectious that host becomes (Strong *et al.* 2015), such as when highly infectious hosts generate other

highly infectious hosts (Beldomenico 2020; Wanelik *et al.* 2023). It can also capture situations where

co-circulating parasite strains differ in their infectivity (Brault *et al.* 2004; Dai *et al.* 2013; Medina *et al.*

2018), so that hosts infected with highly infective parasite strains then become highly infectious hosts

themselves. Critically, and in contrast to the RD scenario, in the DD scenario a host's infectiousness is

determined solely by the identity of the donor host that infected them. We propose that RD and DD

are two epidemiologically distinct phenomenological ways to conceptualise various processes by

which a host's infectiousness is determined, distinguished by the identity of the host (the recipient or

the donor) that determines the infectiousness of a newly infected host.

We suggest that the differentiation of the RD and DD scenarios will lead to different patterns of host

infectiousness in a population and, crucially, alter the impact of host heterogeneities on parasite

transmission. Previous work (e.g., Vazquez-Prokopec *et al.* (2016); Lloyd *et al.*

(2023)) has implicitly used one or the other of these scenarios but has not systematically explored

how they differ in affecting the impact of coupled heterogeneities in susceptibility and infectiousness

on parasite transmission. To investigate this, we develop Susceptible-Infected (SI) compartmental

models that incorporate heterogeneity in susceptibility and infectiousness under both RD and DD

scenarios. We then analyse these models to determine how RD and DD affect the relationships

between those host heterogeneities, parasite transmission and host population dynamics. We find

that how infectiousness is determined – RD *vs*. DD – can have substantial effects on parasite

transmission, particularly affecting the primary drivers of the basic reproduction number (R_0).

Material and methods

Model Framework

The standard density-dependent SI model in a homogeneous population of size *N* divided into

susceptible (S) and infected (I) sub-populations (Anderson *et al.* 1991) is given by

$$\frac{dS}{dt} = bN - \beta SI - dS, \tag{1}$$

$$\frac{dI}{dt} = \beta SI - (d + \alpha)I, \tag{2}$$

95 where $N = S + I$, b is the birth rate, d the baseline mortality rate, and α the parasite-induced

96 mortality rate. The transmission coefficient β, while often written as a simple constant, actually

97 incorporates both contact rate (κ) and infection probability given a contact (ν) (Begon *et al.* 2002),

98 yielding

$$\beta = \kappa\nu. \tag{3}$$

100 The infection probability ν can be further partitioned into the product of recipient susceptibility (σ)

101 and donor infectiousness (ι),

$$\nu = \sigma\iota, \tag{4}$$

where σ and ι take values in [0,1], with higher values representing greater susceptibility or greater

infectiousness, respectively. Thus, host heterogeneity in susceptibility and infectiousness can be

incorporated by dividing the population into sub-populations comprising individuals that share the

same values of susceptibility and infectiousness. Precisely how this is done depends on whether

infectiousness is determined by a RD or DD scenario.

Recipient dependence (RD)

Under this scenario recipient hosts are assigned to an infected sub-population based on a fixed, pre-

determined trait inherent to that individual. Supposing that there are n unique pairs of trait values

(σ_j, ι_j), $1 \leq j \leq n$, we divide the population into n susceptible sub-populations S_j, and n infected sub-

populations I_j, where the jth sub-populations share the jth trait pair. Thus, a RD heterogeneous

analogue of Equations (1)-(2) is

$$\frac{dS_j}{dt} = \frac{bN}{n} - \kappa \sigma_j S_j \sum_{m=1}^{n} \iota_m I_m - dS_j, \tag{5}$$

$$\frac{dI_j}{dt} = \kappa \sigma_j S_j \sum_{m=1}^{n} \iota_m I_m - (d + \alpha)I_j, \tag{6}$$

where we have additionally assumed that birth rate b, baseline mortality rate d, and parasite-induced

mortality rate α, are equal across sub-populations. For the sake of simplicity, births are evenly

distributed across susceptible sub-populations, effectively assuming non-inherited host

heterogeneity, a phenomenon that has been previously observed, for example in *Daphnia magna*

(Ben-Ami *et al.* 2008). We emphasise that in the RD scenario, the fixed traits of individuals determine

the susceptible and infected sub-population to which they belong, so that all individuals in a

susceptible sub-population move to the same infected sub-population upon becoming infected.

Donor dependence (DD)

In this scenario, recipient hosts acquire their infectiousness trait when they become infected. Like in

the RD scenario, we divide the susceptible sub-population into n_S sub-populations S_j, each with

susceptibility σ_j, $1 \leq j \leq n_S$. Under the DD scenario, however, individuals are assigned the

infectiousness of the donor host that infected them. We therefore define $I_{j,k}$ to be those individuals

with susceptibility σ_j that were infected by an individual with infectiousness ι_k, $0 \leq k \leq n_I$, and so

now share that same infectiousness value. Thus, a DD heterogeneous analogue of Equations (1)-(2) is

$$\frac{dS_j}{dt} = \frac{bN}{n_S} - \kappa\sigma_j S_j \sum_{m=1}^{n_I} \sum_{l=1}^{n_S} \iota_m I_{l,m} - dS_j, \tag{7}$$

$$\frac{dI_{j,k}}{dt} = \kappa\sigma_j \iota_k S_j \sum_{l=1}^{n_S} I_{l,k} - (d + \alpha)I_{j,k}. \tag{8}$$

b, d and α are again assumed to be equal for all infected sub-populations. Unlike the RD model, the

infected sub-population that the recipient host will join is not pre-determined before infection, so

individuals in the same susceptible sub-population do not always move to the same infected sub-

population upon infection. As such, the DD model has n_s susceptible sub-populations and $n_S n_I$

infectious sub-populations, whereas the RD model has equal numbers of both susceptible and infected

sub-populations.

Equations (7)-(8), which will be useful when quantifying population-level heterogeneity, can be

simplified by defining $I_k = \sum_{j=1}^{n_S} I_{j,k}$, *i.e.*, the sum of all individuals with the same infectiousness value,

regardless of their initial susceptibility value. Summing Equation (8) over $0 \leq j \leq n_S$ yields

$$\frac{dS_j}{dt} = \frac{bN}{n_S} - \kappa\sigma_j S_j \sum_{m=1}^{n_I} \iota_m I_m - dS_j, \tag{9}$$

$$\frac{dI_k}{dt} = \kappa\iota_k I_k \sum_{m=1}^{n_S} \sigma_m S_m - (d + \alpha)I_k, \tag{10}$$

describing the dynamics of all infected individuals with infectiousness ι_k. This simplified form of the

system is useful for calculating R_0. Note that we have used the same notation in the RD and the DD

models to define similar but not precisely equivalent variables, and therefore we rely on context to

provide clarity about which is being referred to throughout the remainder of this work.

Quantifying population-level heterogeneity

We quantified population-level heterogeneity (hereafter referred to simply as 'heterogeneity') after (Laliberté *et al.* 2010), which is applicable to a wide range of systems and can deal with multiple traits and missing values (Olusoji *et al.* 2023). We undertook heterogeneity calculations in the context of two-dimensional σ, ι trait space, in which susceptibility (σ) and infectiousness (ι) form the two axes.

We first calculated the abundance-weighted centroid (c, hereafter referred to as the 'centroid') of the population, given simply as the population mean of each trait ($\bar{\sigma}$ and $\bar{\iota}$, Figure 1A & C). We then calculated the heterogeneity score, h, by finding the mean abundance-weighted, Euclidean distance to the centroid (e.g. z_1 in Figure 1B, z_{11} in Figure 1D) of all sub-populations. We calculated initial heterogeneity using the initial abundances of the sub-populations and final (equilibrium) heterogeneity using equilibrium sub-population abundances.

The calculation of heterogeneity differs between the RD and DD scenarios in how sub-populations were grouped, and abundances calculated. Specifically, for RD, all individuals in the same susceptible sub-population move to the same infected sub-population, and so we summed the abundances of the corresponding susceptible and infected sub-populations. These grouped sub-populations were then used to calculate h. For DD, each infected sub-population's Euclidean distance to the centroid was calculated using both its σ and ι values (e.g. I_{11} in Figure 1D) while, because susceptible sub-populations only had a σ value, their distance to the centroid was calculated in a single dimension (e.g. S_1 in Figure 1D), and so there were no grouped sub-populations used in calculating h. The formulae for heterogeneity calculations for both the RD and DD scenarios can be found in the Supplementary Information (SI).

Figure 1. Schematic representation for calculating heterogeneity for the RD (A, B) and DD (C, D) scenarios. For both the RD and DD scenarios the population mean susceptibility ($\bar{\sigma}$) and population mean infectiousness ($\bar{\iota}$) are calculated in a single dimension (A and C). The sizes of the black dots indicate the relative abundances of the relevant sub-populations. (B) Heterogeneity for the RD scenario is the mean Euclidean distance (z_j), weighted by the abundance of each sub-population, to the centroid (c). (D) Heterogeneity for the DD scenario is the mean of the Euclidean distances (z_{jk}) of the infected sub-populations to the centroid, and the single dimension distance (σ) of the susceptible sub-populations, weighted by the abundance of each sub-population; here the values for S_1 and S_2 form lines rather than points in σ, ι trait space because they have no infectiousness values. In all panels the diameters of the black circles represent the abundance of the sub-population.

Model Analyses

We analysed both the RD and DD scenarios under three heterogeneity contexts:

1) *Bipartite heterogeneity*

Where the population is divided into sub-populations corresponding to two distinct pairs of

susceptibility and infectiousness values.

2) *Tripartite isometric heterogeneity*

Where the population is divided into sub-populations corresponding to three distinct pairs of

susceptibility and infectiousness values, equidistant from each other in σ, ι trait space.

3) *Tripartite non-isometric heterogeneity*

Where the population is divided into sub-populations corresponding to three distinct pairs of

susceptibility and infectiousness values, but which are not necessarily equidistant in σ, ι trait

space.

Here we present analyses relating to context 1; contexts 2 and 3 are presented in the SI.

For the bipartite heterogeneity context, we set the initial number of susceptibles per sub-population

to 49 and the number of infecteds per sub-population to 1 (RD, 2 sub-populations) or 0.5 (DD, 4 sub-

populations). Initial mean population susceptibility and infectiousness trait values were 0.5; thus $\sigma_2 =$

$1 - \sigma_1$ and $\iota_2 = 1 - \iota_1$. All other parameter values were the same for all analyses (Table 1). By varying

σ_1 and ι_1, we varied the initial population heterogeneity while maintaining the same initial mean

population trait values. Hence, the effects of changing heterogeneity were decoupled from the effects

of changing initial mean population trait values.

We generated 2,601 unique combinations of σ and ι trait values, and used these to analyse both the

RD and DD models to understand how heterogeneity in susceptibility and infectiousness affected

three key descriptors of epidemiological dynamics: (i) the basic reproduction number (R_0), a measure

of epidemic potential (Anderson *et al.* 1991); (ii) the equilibrium host population abundance, which

quantifies the impact of the parasite on the host population; and (iii) the change in heterogeneity

between the initial and final state of the system, indicating how heterogeneity changes with epidemic

progression (presented in the SI).

Table 1. Parameter definitions and values for all analyses of the RD and DD models.

Model parameter	Definition	Value
κ	Contact rate per individual per time	0.5
b	Birth rate per time	1.5
d	Mortality rate per time	1
α	Parasite-induced mortality rate per time	0.7

Calculating R_0

R_0 predicts the risk of an epidemic occurring, as well as the size and severity of that epidemic

(Anderson *et al.* 1991; Heffernan *et al.* 2005), and also the effort needed to control and eliminate a

parasite from a population (Roberts 2007). Thus, understanding the effect of heterogeneity on R_0

provides considerable insight into how host heterogeneity in susceptibility and infectiousness affects

parasite transmission through a population.

Using next generation matrices (Diekmann *et al.* 2010) (shown in full in the SI), we calculated R_0 for

the each scenario to be

RD:

$$R_0 = \frac{\kappa}{d + \alpha} \sum_{j=1}^{n} \sigma_j \iota_j S_j , \tag{11}$$

225 **DD:**

$$R_0 = \frac{\kappa \iota_{\max}}{d + \alpha} \sum_{j=1}^{n_S} \sigma_j S_j, \tag{12}$$

where $\iota_{\max}$ is the maximum infectiousness value across all sub-populations.

Equilibrium Analyses

When possible we calculated the equilibrium solutions of Equations (5)-(8) analytically. For parameter

values for which this was not possible, we solved the system numerically over 50,000 time steps, which

was sufficiently long to reach equilibrium. Any sub-population with susceptibility $\sigma = 0$ (*i.e.*,

completely resistant to infection) experiences unbounded growth, and so such cases were omitted

from equilibrium analyses.

We also used these equilibrium solutions to calculate equilibrium population-level heterogeneity

(following the method described above) to test whether the population-level heterogeneity changed

with epidemiological progress.

Software packages

Plots were generated in R (R Core Team 2022) using packages ggplot2 (Wickham 2016), ggforce

(Pedersen 2022a), scales (Wickham *et al.* 2022), showtext (Qiu 2022), pBrackets (Schulz 2021),

patchwork (Pedersen 2022b) and latex2exp (Meschiari 2022). Equilibrium analyses were conducted

using Mathematica (Wolfram Research Inc. 2022). Heterogeneity and R_0 values were calculated in R

(R Core Team 2022).

Results

Here we present analysis of the bipartite heterogeneity context, where there are two sub-populations. The results for the tripartite heterogeneities are broadly consistent with the bipartite analyses, highlighting the generality of the findings; we describe those results in the SI.

(i) How heterogeneity affects R_0 under RD v DD scenarios

Recipient dependence

In this scenario as heterogeneity increases R_0 can increase (Figure 2A, red points), decrease (Figure 2A, blue points) or remain the same as the homogenous case (Figure 2A, yellow points). Whether it increases, decreases or remains the same depends on the population-level covariance between susceptibility and infectiousness. Mapping R_0 values onto σ, ι trait space (Figure 2B) (where the sub-populations are mirrored across the centre point ($\sigma_2 = 1 - \sigma_1$ and $\iota_2 = 1 - \iota_1$)), shows that if there is heterogeneity in one trait only, R_0 is the same as in the homogeneous population ($h = 0$) (Figure 2B yellow shading). This remains true even at high levels of heterogeneity. However, with positive covariance between susceptibility and infectiousness, R_0 increases compared with the homogeneous population (Figure 2B, Supplementary Equation (S13)). R_0 is at its maximum when one sub-population has maximal infectiousness and susceptibility values of 1, while the other sub-population has values of 0 for both; *i.e.* when the sub-populations lie at extremes of the positive diagonal in σ, ι trait space (Figure 2B). Conversely, with a negative covariance between susceptibility and infectiousness, R_0 is lower than in the homogenous case (Figure 2B). In the extreme case, when one sub-population consists of only completely resistant hosts (susceptibility = 0, infectiousness = 1), and the other sub-population of completely dead-end hosts (susceptibility = 1, infectiousness = 0), such that they lie at extremes of the negative diagonal in σ, ι trait space, no individuals can both be infected and transmit onwards, resulting in an R_0 of 0 (Figure 2B).

When there is a high degree of heterogeneity, there is a bifurcating distribution of points (Figure 2A)

due to the boundaries of σ, ι trait space that restrict the number of viable trait combinations (*i.e.*,

where both susceptibility and infectiousness are between 0 and 1 for all sub-populations). For

example, with maximum heterogeneity there are only two possible configurations of the sub-

populations such that they lie at the opposite extremes of the diagonals in σ, ι trait space, resulting in

just two points (Figure 2A).

In summary, increasing heterogeneity in the RD scenario can lead to increasingly divergent R_0,

compared to the homogeneous population. But, whether R_0 is greater or less than the homogeneous

population is determined by whether the population-level covariance between susceptibility and

infectiousness is positive or negative, respectively. When the covariance is zero (heterogeneity in

either susceptibility or infectiousness, but not both), R_0 is the same as the homogeneous population,

regardless of the degree of heterogeneity.

Donor dependence

In the DD scenario, R_0 is affected in a qualitatively different fashion than in the RD scenario.

Specifically, increasing heterogeneity does not lower R_0 compared to the homogenous population

(Figure 2C). Heterogeneity generally has little effect on R_0, and changes in susceptibility alone do not

affect R_0, but changes in infectiousness do (Figure 2D).

In the DD scenario R_0 is driven by the maximum infectiousness in the population (Figure 2C,

Supplementary Equation (S13)). R_0 is independent of susceptibility because, for equal sized susceptible

sub-populations, there is a fixed mean value across the population (details in SI). Hence, R_0 changes

only along the infectiousness axis (Figure 2D), not along the susceptibility axis. For the same overall

degree of heterogeneity (*i.e.*, vertical slice in Figure 2C), increasing heterogeneity in infectiousness

(and a necessary decrease in heterogeneity in susceptibility) increases R_0. Susceptibility can only affect

R_0 when the initial susceptible sub-population abundances are not equal (see SI).

Summary of effects of heterogeneity on R_0

In both the RD and DD scenarios increasing heterogeneity can lead to increasingly divergent R_0 values

compared to the homogeneous case. But, what drives this effect on R_0 differs between the two

scenarios. In the RD scenario it is the population-level covariance between susceptibility and infectiousness; in the DD scenario it is the maximum infectiousness in the population.

Figure 2. The effect of initial heterogeneity on R_0 for the RD (A, B) and DD (C, D) scenarios. The dotted line in (A, C) is where $R_0 = 1$. For RD (A) R_0 changes as initial heterogeneity increases, and with the covariance between susceptibility (σ) and infectiousness (ι), as indicated by the colour bar. (B) shows R_0 (the colour scale) plotted in σ, ι trait space with the centroid (c) at $\sigma = \iota = 0.5$, and concentric dashed-line circles showing increasing heterogeneity; the positions of the sub-populations are mirrored across the centroid ($\sigma_2 = 1 - \sigma_1$ and $\iota_2 = 1 - \iota_1$). For DD (C) R_0 does not change as initial heterogeneity increases, but scales with maximum infectiousness (ι_{max}), as the green colour bar. (D)

is the DD version of panel (B), but note that the R_0 and heterogeneity scales differ between (B) and

(D).

(ii) How heterogeneity affects host abundance under RD v DD scenarios

Equilibrium total host abundance generally increases with increasing initial heterogeneity, as does the

variability in equilibrium abundance, for both the RD and DD scenarios (Figure 3). However, specific

aspects of these relationships differ between the two scenarios.

Recipient dependence

In the RD scenario there is a complex relationship between initial heterogeneity and host equilibrium

abundance (Figure 3A & C, and see Figure S2 for an enlarged version at the lower host abundance

range). Equilibrium abundances are grouped into parabolic 'clusters' where each cluster has

equilibrium abundances that both increase and decrease from an initial baseline abundance as

heterogeneity increases. Each cluster is determined by the minimum susceptibility in the population;

all points in a cluster have the same population-level minimum susceptibility value (and so also the

same maximum susceptibility value) (Figure 3C). Clusters are ordered by these minimum susceptibility

values; those with the lowest minimum susceptibility values (σ_{min}, Figure 3C) have the highest

abundances. This is because in populations with low minimum susceptibility fewer individuals become

infected, so that fewer hosts are exposed to parasite-induced mortality (α), which thereby increases

overall host abundance.

The within-cluster divergence in host abundance that occurs with increasing heterogeneity arises due

to increasingly divergent infectiousness values in the population. Since all points within a cluster have

the same pair of σ values for the component sub-populations, an increase in heterogeneity is achieved

by divergence in the two infectiousness values. This divergence, and so increase in heterogeneity,

leads to changes in R_0 within a cluster that changes equilibrium host abundance. High R_0 values lead

to lower abundances (the lower red tail of a cluster in Figure 3A), while low R_0 values lead to higher

abundances (the upper blue tail of a cluster in Figure 3A).

Donor dependence

In the DD scenario there are also clusters in equilibrium host abundance determined by the minimum

susceptibility value (σ_{min}) within the population (Figure 3D), but the points within each cluster are

arranged in near-vertical lines, suggesting that heterogeneity has little effect on equilibrium host

abundance within a cluster. As with the RD scenario, equilibrium host abundance is maximised when

R_0 is minimised (Figure 3B) which, for the DD scenario occurs when the maximum infectiousness in

the population (ι_{max}) is at its lowest (Figure 2C). Hence within each cluster, increasing maximum

infectiousness increases R_0, reducing equilibrium host abundance. The DD scenario generally shows

lower equilibrium host abundances than the RD scenario because the minimum R_0 values for the RD

scenario are lower than the DD scenario.

Figure 3. The effect of initial heterogeneity on equilibrium total host abundance for RD (A, C) and DD

(B, D) scenarios. (A) and (C) are the same RD plots, but differently coloured, (A) for R_0 and (C) for

minimum population-level susceptibility (σ_{min}). (B) and (D) are the same DD plots, but differently

coloured, (B) for R_0 and (D) for σ_{min}. Note that the R_0 scales differ between (A) and (B). An enlarged

version of (A) and (C) can be found in the SI (Figure S2).

Discussion

While existing theory shows that host heterogeneity in susceptibility and infectiousness can affect population-level parasite transmission (*e.g.* Lloyd *et al.* (2020)), our findings demonstrate that these effects differ considerably depending how host infectiousness is determined. We show that the contrasting RD and DD scenarios result in different relationships between host heterogeneity in susceptibility and infectiousness and epidemiological outcomes, specifically R_0 and host abundance.

While we find that R_0 changes with increasing heterogeneity in both scenarios, there is a notable contrast between the two scenarios in what drives these effects, and the direction of the effects. The RD scenario shows divergent R_0 values as heterogeneity increases, which is determined by the covariance between susceptibility and infectiousness. This finding is consistent with previous modelling: three models of vector-borne infections, where infectiousness was an inherent host trait (*i.e.* implicitly assuming RD), incorporated host heterogeneity in susceptibility and infectiousness and found that positive covariance between heterogeneities led to an increase in R_0 relative to the homogeneous case, while negative covariance led to a decrease (Dietz 1980; Koella 1991; Vazquez-Prokopec *et al.* 2016). Thus, these models showed that in a RD scenario, population-level covariance between susceptibility and infectiousness determines R_0, consistent with our findings.

In contrast, we find that in the DD scenario R_0 is determined by the maximum infectiousness in the population, such that R_0 increases as heterogeneity in infectiousness increases. Heterogeneity in susceptibility is largely irrelevant for the DD scenario, only becoming so if abundances are very different between sub-populations. There are fewer other studies that consider DD-like scenarios, but one example is the hypothesis that patterns of SARS-CoV-2 transmission may be due to superspreaders tending to generate new superspreaders, for example through a dose-dependent effect (Beldomenico 2020). A model exploring how R_0 responded to this scenario (compared to the null model where superspreaders appeared randomly) showed that R_0 increases with an increase in the probability that a superspreader generates additional superspreaders (Wanelik *et al.* 2023). Thus,

moving from the null model to a DD-like scenario increased R_0, suggesting that the DD scenario tends

to increase R_0, consistent with our findings.

The RD and DD scenarios also differ in how host heterogeneity affects equilibrium total host

abundance. While host abundance tends to increase with increasing heterogeneity in both scenarios,

the effect is more pronounced in the RD scenario than in the DD scenario. Further, for a given

susceptibility value, RD and DD differ by generating divergent (RD) or monotonic (DD) changes in

equilibrium host abundance, a pattern that becomes more pronounced with increasing heterogeneity.

These RD and DD scenarios are phenomenological characterisations that each encompass a variety of

real-world processes. Empirical examples matching assumptions of the RD scenario include the finding

that canaries' nutritional status can affect their infectiousness with avian malaria (Cornet *et al.* 2014);

rabbit myxoma virus infection status determines its infectiousness for co-infecting nematodes

(Cattadori *et al.* 2007); several examples of different host strains exhibiting varying levels of

infectiousness when infected with the same parasite isolates (Bolas-Fernandez *et al.* 1989; Jørgensen

et al. 1998; Dorfman *et al.* 2024); a genetic basis for observed variation in host infectiousness in

Scophthalmus maximus (Turbot) infected with a ciliate parasite (Anacleto *et al.* 2019). In addition,

critical to understanding parasite transmission in a RD scenario, is covariance between susceptibility

and infectiousness, and there are empirical examples of this. For example, it has been shown that

juvenile *Cygnus columbianus bewickii* (Bewick's swan) are more susceptible to avian influenza virus

than adults and also have higher shedding rates than adults, demonstrating positive covariance

between host susceptibility and infectiousness (Hoye *et al.* 2012). Alternatively, negative covariance

between host susceptibility and infectiousness has been proposed as a mechanism for social insects

avoiding parasitism by investing more in anti-parasitic behaviours to reduce their infectiousness in a

colony with high susceptibility (Cremer *et al.* 2007), such that a highly susceptible colony would be

less infectious than a less susceptible colony. At the individual level, negative covariance might occur

when highly susceptible individuals acquire high levels of disease, thus increasing sickness behaviours

that reduce host infectiousness (Araujo *et al.* 2016).

Empirical examples matching assumptions of the DD scenario are calves infected with three different

doses of bovine viral diarrhoea virus (BVDV), where the most infectious individuals were those given

the highest viral dose, due to a longer infectious period (Strong *et al.* 2015). Similar patterns have been

found with a number of other host-parasite systems (Gaskell *et al.* 1979; Mumford *et al.* 1990; Zarkov

2012).

In reality, many host-parasite systems are unlikely to fit solely under either RD or DD scenarios, instead

falling somewhere between the two. For example, in the BVDV calf infection study, while calves

infected with the highest BVDV doses were the most infectious, there was still within-dose group

heterogeneity in infectiousness (Strong *et al.* 2015), which may have been caused by traits inherent

to individual calves. Thus, while the overall system may be best described by the DD scenario, there

are still aspects of the RD scenario at play. The reverse can also be true. For example, although

myxoma-infected rabbits may be more nematode infectious (Cattadori *et al.* 2007), aligning with the

RD scenario, there may still be some DD involved. Specifically, the nematode transmits between hosts

via a faecal-oral pathway (Cattadori *et al.* 2007). Therefore, a rabbit might become more infectious

when infected by a host with a high parasite burden, because it will acquire more parasites in turn.

These examples demonstrate that R_0 will often be affected by both the covariance between

susceptibility and infectiousness (RD) as well as the maximum infectiousness in the population (DD),

though which of these two measures is more influential will depend on where on the spectrum of RD

to DD that specific host-parasite system exists.

Previous work has typically treated how a host's infectiousness is determined as a black box, generally

assuming it is a fixed property of the recipient host. This overlooks the importance of how a host's

infectiousness is determined in influencing the consequences of host heterogeneity for parasite

transmission. Yet this process can have real-world consequences. For instance, there is interest in

breeding parasite resistant livestock to reduce the substantial economic and climatic costs caused by

parasites in livestock systems (Knap *et al.* 2020). However, to determine the livestock traits that are

most appropriate to select, it will be important to consider where the host-parasite system falls on

the RD-DD spectrum. For example, in a RD scenario, selecting for reduced parasite susceptibility (*i.e.*,

for resistance), versus in a DD scenario selecting for reduced parasite shedding. The work we present

here shows the importance of explicitly considering how infectiousness is determined, and that

ignoring it can lead to an incomplete understanding of the effects of host heterogeneities on parasite

transmission. A failure to do so could have consequences for both future theoretical and empirical

work.

Acknowledgements

439   JAC was funded by a PhD studentship from the 'Adapting to the Challenges of a Changing Environment

440   (ACCE)' Doctoral Training Partnership funded by the Natural Environment Research Council (NERC)

441   (Grant NE/S00713X/1). ADD was funded by a NERC Pushing the Frontiers grant (Grant NE/X01424X/1).

References

Adelman, J.S., Moyers, S.C., Farine, D.R. & Hawley, D.M. (2015) Feeder use predicts both acquisition and transmission of a contagious pathogen in a North American songbird. *Proceedings of the Royal Society B: Biological Sciences,* **282,** 20151429.

Anacleto, O., Cabaleiro, S., Villanueva, B., Saura, M., Houston, R.D., Woolliams, J.A. & Doeschl-Wilson, A.B. (2019) Genetic differences in host infectivity affect disease spread and survival in epidemics. *Scientific Reports,* **9,** 4924.

Anderson, R.M. & May, R.M. (1991) *Infectious Diseases of Humans*. Oxford University Press.

Araujo, A., Kirschman, L. & Warne, R.W. (2016) Behavioural phenotypes predict disease susceptibility and infectiousness. *Biology Letters,* **12,** 20160480.

Barlow, N.D. (2000) Non-linear transmission and simple models for bovine tuberculosis. *journal of animal ecology,* **69,** 703-713.

Begon, M., Bennett, M., Bowers, R.G., French, N.P., Hazel, S. & Turner, J. (2002) A clarification of transmission terms in host-microparasite models: numbers, densities and areas. *Epidemiology & Infection,* **129,** 147-153.

Beldomenico, P.M. (2020) Do superspreaders generate new superspreaders? A hypothesis to explain the propagation pattern of COVID-19. *International Journal of Infectious Diseases,* **96,** 461-463.

Ben-Ami, F., Regoes, R.R. & Ebert, D. (2008) A quantitative test of the relationship between parasite dose and infection probability across different host–parasite combinations. *Proceedings of the Royal Society B: Biological Sciences,* **275,** 853-859.

Bolas-Fernandez, F. & Wakelin, D. (1989) Infectivity of *Trichinella* isolates in mice is determined by host immune responsiveness. *Parasitology,* **99,** 83-88.

Brault, A.C., Langevin, S.A., Bowen, R.A., Panella, N.A., Biggerstaff, B.J., Miller, B.R. & Komar, N. (2004) Differential virulence of West Nile strains for American crows. *Emerging infectious diseases,* **10,** 2161.

Cattadori, I.M., Albert, R. & Boag, B. (2007) Variation in host susceptibility and infectiousness generated by co-infection: the myxoma–*Trichostrongylus retortaeformis* case in wild rabbits. *Journal of The Royal Society Interface,* **4,** 831-840.

Cornet, S., Bichet, C., Larcombe, S., Faivre, B. & Sorci, G. (2014) Impact of host nutritional status on infection dynamics and parasite virulence in a bird-malaria system. *journal of animal ecology,* **83,** 256-265.

Coutinho, F.A.B., Massad, E., Lopez, L.F., Burattini, M.N., Struchiner, C.J. & Azevedo-Neto, R.S.d. (1999) Modelling heterogeneities in individual frailties in epidemic models. *Mathematical and computer modelling,* **30,** 97-115.

Cremer, S., Armitage, S.A. & Schmid-Hempel, P. (2007) Social immunity. *Current biology,* **17,** R693-R702.

Dai, Y., Liu, M., Cheng, X., Shen, X., Wei, Y., Zhou, S., Yu, S. & Ding, C. (2013) Infectivity and pathogenicity of Newcastle disease virus strains of different avian origin and different virulence for mallard ducklings. *Avian Diseases,* **57,** 8-14.

Diekmann, O., Heesterbeek, J. & Roberts, M.G. (2010) The construction of next-generation matrices for compartmental epidemic models. *Journal of The Royal Society Interface,* **7,** 873-885.

Dietz, K. (1980) Models for vector-borne parasitic diseases. *Vito Volterra Symposium on Mathematical Models in Biology*, pp. 264-277. Springer.

Dorfman, B., Marcos-Hadad, E., Tadmor-Levi, R. & David, L. (2024) Disease resistance and infectivity of virus susceptible and resistant common carp strains. *Scientific Reports,* **14,** 4677.

Dwyer, G., Elkinton, J.S. & Buonaccorsi, J.P. (1997) Host heterogeneity in susceptibility and disease dynamics: tests of a mathematical model. *The American Naturalist,* **150,** 685-707.

Gaskell, R. & Povey, R. (1979) The dose response of cats to experimental infection with feline viral
rhinotracheitis virus. *Journal of Comparative Pathology, 89,* 179-191.
Heffernan, J.M., Smith, R.J. & Wahl, L.M. (2005) Perspectives on the basic reproductive ratio. *Journal*
of The Royal Society Interface, 2, 281-293.
Hickson, R. & Roberts, M. (2014) How population heterogeneity in susceptibility and infectivity
influences epidemic dynamics. *Journal of Theoretical Biology, 350,* 70-80.
Hoye, B.J., Fouchier, R.A. & Klaassen, M. (2012) Host behaviour and physiology underpin individual
variation in avian influenza virus infection in migratory Bewick's swans. *Proceedings of the*
Royal Society B: Biological Sciences, 279, 529-534.
Jørgensen, L., Leathwick, D., Charleston, W., Godfrey, P., Vlassoff, A. & Sutherland, I. (1998) Variation
between hosts in the developmental success of the free-living stages of trichostrongyle
infections of sheep. *International Journal for Parasitology, 28,* 1347-1352.
Knap, P.W. & Doeschl-Wilson, A. (2020) Why breed disease-resilient livestock, and how? *Genetics*
Selection Evolution, 52, 1-18.
Koella, J.C. (1991) On the use of mathematical models of malaria transmission. *Acta tropica, 49,* 1-
25.
Kong, L., Wang, J., Han, W. & Cao, Z. (2016) Modeling heterogeneity in direct infectious disease
transmission in a compartmental model. *International journal of environmental research and*
public health, 13, 253.
Laliberté, E. & Legendre, P. (2010) A distance-based framework for measuring functional diversity
from multiple traits. *Ecology, 91,* 299-305.
Lemieux, J.E., Siddle, K.J., Shaw, B.M., Loreth, C., Schaffner, S.F., Gladden-Young, A., Adams, G., Fink,
T., Tomkins-Tinch, C.H. & Krasilnikova, L.A. (2021) Phylogenetic analysis of SARS-CoV-2 in
Boston highlights the impact of superspreading events. *Science, 371,* eabe3261.
Lloyd, A.L., Kitron, U., Perkins, T.A., Vazquez-Prokopec, G.M. & Waller, L.A. (2020) The basic
reproductive number for disease systems with multiple coupled heterogeneities.
Mathematical biosciences, 321, 108294.
Lloyd-Smith, J.O., Schreiber, S.J., Kopp, P.E. & Getz, W.M. (2005) Superspreading and the effect of
individual variation on disease emergence. *Nature, 438,* 355-359.
Mantzaris, A. & Higham, D. (2013) Dynamic communicatbility predicts infectiousness. *Temporal*
Networks (eds P. Holme & J. Saramäki). Springer, Berlin, Heidelberg.
Matthews, L., McKendrick, I.J., Ternent, H., Gunn, G., Synge, B. & Woolhouse, M. (2006) Super-
shedding cattle and the transmission dynamics of *Escherichia coli* O157. *Epidemiology &*
Infection, 134, 131-142.
McCallum, H., Fenton, A., Hudson, P.J., Lee, B., Levick, B., Norman, R., Perkins, S.E., Viney, M.,
Wilson, A.J. & Lello, J. (2017) Breaking beta: deconstructing the parasite transmission
function. *Philosophical Transactions of the Royal Society B: Biological Sciences, 372,*
20160084.
Medina, L., Castillo, C., Liempi, A., Herbach, M., Cabrera, G., Valenzuela, L., Galanti, N., de los
Angeles Curto, M., Schijman, A.G. & Kemmerling, U. (2018) Differential infectivity of two
Trypanosoma cruzi strains in placental cells and tissue. *Acta tropica, 186,* 35-40.
Meschiari, S. (2022) latex2exp: Use LaTex Expression in Plots.
Miller, J.C. (2007) Epidemic size and probability in populations with heterogeneous infectivity and
susceptibility. *Physical Review E, 76,* 010101.
Mumford, J.A., Hannant, D. & Jessett, D. (1990) Experimental infection of ponies with equine
influenza (H3N8) viruses by intranasal inoculation or exposure to aerosols. *Equine Veterinary*
Journal, 22, 93-98.
Olusoji, O.D., Barabás, G., Spaak, J.W., Fontana, S., Neyens, T., De Laender, F. & Aerts, M. (2023)
Measuring individual-level trait diversity: a critical assessment of methods. *Oikos, 2023,*
e09178.
Pedersen, T.L. (2022a) ggforce: Accelerating 'ggplot2'.

Pedersen, T.L. (2022b) patchwork: The Composer of Plots.

Qiu, Y. (2022) showtext: Using Fonts More Easily in R Graphs.

R Core Team (2022) R: A language and environment for statistical computing. R Foundation for
544   Statistical Computing, Vienna, Austria.

545   Roberts, M. (2007) The pluses and minuses of R0. *Journal of The Royal Society Interface,* **4,** 949-961.

546   Schulz, A. (2021) pBrackets: Plot Brackets.

547   Stephenson, J.F., Young, K.A., Fox, J., Jokela, J., Cable, J. & Perkins, S.E. (2017) Host heterogeneity
548   affects both parasite transmission to and fitness on subsequent hosts. *Philosophical*
549   *Transactions of the Royal Society B: Biological Sciences,* **372,** 20160093.

550   Streicker, D.G., Fenton, A. & Pedersen, A.B. (2013) Differential sources of host species heterogeneity
551   influence the transmission and control of multihost parasites. *Ecology letters,* **16,** 975-984.

552   Strong, R., La Rocca, S.A., Paton, D., Bensaude, E., Sandvik, T., Davis, L., Turner, J., Drew, T., Raue, R.
553   & Vangeel, I. (2015) Viral dose and immunosuppression modulate the progression of acute
554   BVDV-1 infection in calves: evidence of long term persistence after intra-nasal infection.
555   *PLoS One,* **10,** e0124689.

556   VanderWaal, K.L. & Ezenwa, V.O. (2016) Heterogeneity in pathogen transmission: mechanisms and
557   methodology. *Functional Ecology,* **30,** 1606-1622.

558   Vazquez-Prokopec, G.M., Perkins, T.A., Waller, L.A., Lloyd, A.L., Reiner Jr, R.C., Scott, T.W. & Kitron,
559   U. (2016) Coupled heterogeneities and their impact on parasite transmission and control.
560   *Trends in parasitology,* **32,** 356-367.

561   Wanelik, K.M., Begon, M., Fenton, A., Norman, R.A. & Beldomenico, P.M. (2023) Positive feedback
562   loops exacerbate the influence of superspreaders in disease transmission. *iScience,* **26**.

563   White, L.A., Forester, J.D. & Craft, M.E. (2018) Covariation between the physiological and behavioral
564   components of pathogen transmission: host heterogeneity determines epidemic outcomes.
565   *Oikos,* **127,** 538-552.

566   Wickham, H. (2016) ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

567   Wickham, H. & Seidel, D. (2022) scales: Scale Functions for Visualization.

568   Wolfram Research Inc. (2022) Mathematica. Wolfram Research Inc., Champaign, Illinois.

569   Woolhouse, M.E., Dye, C., Etard, J.-F., Smith, T., Charlwood, J., Garnett, G., Hagan, P., Hii, J., Ndhlovu,
570   P. & Quinnell, R. (1997) Heterogeneities in the transmission of infectious agents:
571   implications for the design of control programs. *Proceedings of the National Academy of*
572   *Sciences,* **94,** 338-342.

573   Yates, A., Antia, R. & Regoes, R.R. (2006) How do pathogen evolution and host heterogeneity
574   interact in disease emergence? *Proceedings of the Royal Society B: Biological Sciences,* **273,**
575   3075-3083.

576   Zarkov, I. (2012) The importance of inoculation dose of avian H6N2 influenza a virus on virus
577   shedding of *Anas plathyrynchos* ducks after induced generalised infection. *Trakia Journal of*
578   *Sciences,* **10,** 58-61.