Peripheral opioid receptor antagonism alleviates fentanyl-induced
cardiorespiratory depression and is devoid of aversive effects

[1,2,3]Brian C. Ruyle, [1,2,3]Sarah Masud, [1,2,3]Rohith Kesaraju, [1,2,3]Mubariz Tahirkheli, [1,2,3]Juhi Modh,
[1,2,3]Sofia Angulo-Lopera, [1,2,3]Tania Lintz, [1,2,3]Jessica A. Higginbotham, [1,2,3]Nicolas Massaly,
[1,2,3,4,5]Jose A Moron*

Affiliations:
[1]Department of Anesthesiology, Washington University in St Louis; St. Louis, MO, USA

[2]Pain Center, Washington University in St Louis; St. Louis, MO, USA

[3]School of Medicine, Washington University in St. Louis; St. Louis, MO, USA

[4]Department of Neuroscience, Washington University in St. Louis; St. Louis, MO, USA

[5]Department of Psychiatry, Washington University in St. Louis; St. Louis, MO, USA

*corresponding author Email: jmoron-concepcion@wustl.edu

ABSTRACT

Millions of Americans suffering from Opioid Use Disorders (OUD) face a high risk of fatal
overdose due to opioid-induced respiratory depression (OIRD). Fentanyl, a powerful synthetic
opioid, is a major contributor to the rising rates of overdose deaths. Reversing fentanyl-induced
respiratory depression has proved to be challenging due to both its high potency and
lipophilicity. We assessed the contributions of central and peripheral mu opioid receptors
(MORs) in mediating fentanyl-induced physiological responses. The peripherally-restricted MOR
antagonist naloxone methiodide (NLXM) both prevented and reversed OIRD to a comparable
extent as naloxone (NLX), indicating substantial involvement of peripheral MORs to OIRD.
Interestingly, NLXM-mediated OIRD reversal did not produce aversive behaviors observed after
NLX. We show that neurons in the nucleus of the solitary tract (nTS), the first central synapse of
peripheral afferents, exhibit biphasic patterns of activity following fentanyl exposure. NLXM
pretreatment attenuates this activity, suggesting that these responses are mediated by
peripheral MORs. Together, these findings establish a critical role for peripheral MOR, including
ascending inputs to the nTS, as sites of dysfunction during OIRD. Furthermore, selective
peripheral MOR antagonism could be a promising therapeutic strategy for managing OIRD by
sparing CNS-driven acute opioid-associated withdrawal and aversion observed after NLX.

Keywords
Opioids, brainstem, respiratory depression, opioid receptors, fentanyl

Significance Statement:

In this study, we compare the central versus peripheral components underlying fentanyl-induced cardiorespiratory depression to prevent overdose deaths. Our data indicates that these effects are, at least partially, due to the activation of mu opioid receptors present in peripheral sites. These findings provide insight into peripheral contributions to fentanyl-induced overdoses and could potentially lead to the development of treatments selectively targeting the peripheral system, sparing individuals from the CNS-driven acute opioid withdrawal generally observed with the use of naloxone.

INTRODUCTION

Millions of Americans live with Opioid Use Disorders (OUD) and face a high risk of opioid-induced respiratory depression (OIRD), the leading cause of opioid related deaths (*1*). The number of fatalities attributed to OIRD has been exacerbated by the rise of distribution and use of synthetic opioids, such as fentanyl (*2–5*). Fentanyl is a highly lipophilic opioid that readily crosses the blood-brain barrier and binds tightly to mu opioid receptors (MORs), which are abundant in the respiratory centers of the medulla and pons (*1, 6, 7*). Compared to other opioids such as heroin and morphine, fentanyl exhibits both a faster onset of OIRD and higher potency for MOR binding (*8, 9*), minimizing successful prevention of lethal outcomes using naloxone (NLX), a competitive and preferential MOR antagonist (*10, 11*). Despite its high efficacy in reversing OIRD, NLX also precipitates unpleasant withdrawal symptoms and aversion, as reported in patients and preclinical models (*11–13*), making its implementation for managing OUD challenging. Therefore, a better understanding of the pharmacology, specific brain regions and pathways, and physiological responses induced by high doses of fentanyl are necessary to develop effective prevention and intervention strategies, ultimately saving lives of patients suffering from OUD.

Several regions have been implicated as key sites of triggering OIRD, including the Pre-Botzinger Complex in the brainstem and the parabrachial nucleus and Kolliker-Fuse nucleus in the pons (*1, 14–19*). Opioids act directly within these networks to alter inspiratory and expiratory phase duration, leading to overall reductions in respiratory rate (*17, 20*). In addition, the nucleus of the solitary tract (nTS), located in the dorsal brainstem, is the first central site that receives sensory afferent information via cranial nerves related to cardiorespiratory, gustatory and gastrointestinal function (*21*). Acute hypoxia robustly activates nTS neurons (*22, 23*) to initiate

appropriate autonomic and cardiorespiratory responses that facilitate a return to homeostatic

physiological states. MOR is highly expressed in the nTS, primarily on vagal afferent fibers that

terminate within the nTS(24). Interestingly, opioids induce systemic and brain hypoxia (25, 26)

and produce an increase in Fos-immunoreactivity in the nTS (27, 28). However, engaging MOR

signaling in the nTS impairs hypoxic ventilatory responses (7), suggesting that OIRD may result

from nTS dysfunction failing to engage appropriate cardiorespiratory responses, leading to

prolonged OIRD. However, the specific contributions of nTS MOR, including ascending MOR-

expressing inputs to this region, in driving aberrant cardiorespiratory depression are still not fully

understood.

In addition to their high expression throughout brainstem respiratory nuclei, MORs are also

expressed in the periphery, including sensory ganglia, lung afferents, cardiac tissue and cranial

nerves that terminate within the nTS (29, 30). Naloxone Methiodide (NLXM) is a quaternary

derivative of NLX that does not cross the blood brain barrier at low doses, making it a useful tool

to assess the peripheral vs central contribution of MORs in opioid-induced physiology and

behavior (10, 31). Despite its lower binding affinity for MOR, NLXM has been shown to reverse

opioid-induced hypoventilation and brain hypoxia in rodent models (12, 31, 32), suggesting that

peripheral opioid receptors may play a greater role in OIRD than previously thought. However,

no studies to date have provided a comprehensive assessment of opioid-induced

cardiorespiratory depression and systemic hypoxia combined with a real-time assessment of

MOR-mediated activity of nTS neurons.

In the present study, we examined the relative contributions of central and peripheral MORs

mediating fentanyl-induced depression of cardiorespiratory parameters and investigated

potential mechanisms of opioid-induced dysfunction within the nTS. We report that fentanyl-

induced cardiorespiratory depression and prolonged systemic hypoxia can be prevented or

reversed by NLXM to a degree similar to NLX. As compared to NLX, NLXM-mediated reversal

of OIRD does not produce aversive-like behaviors. Fentanyl induced robust Fos-IR expression

in the nTS in a dose-dependent manner, and this activation can be attributed to a combination

of direct effects of opioids acting at MOR located within the nTS and the subsequent hypoxia

that develops after fentanyl exposure. Given that the majority of nTS MOR is located on vagal

afferent fibers, we utilized various techniques to evaluate opioid-induced activity within the nTS

and manipulate nTS MOR signaling during OIRD. Using wireless *in vivo* fiber photometry, we

show that fentanyl induces a biphasic pattern of activity in nTS, characterized by a transient

increase followed by a prolonged decrease in neuronal excitability. NLXM pretreatment strongly

attenuated this biphasic response, indicating that fentanyl-induced nTS activity is influenced by

MOR-expressing peripheral afferents. Together, these findings provide novel insights into
peripheral mechanisms mediating OIRD and support peripheral MOR antagonism as a
promising therapeutic strategy for managing OIRD for patients suffering from OUD while sparing
the CNS-driven acute aversive behaviors that are observed with the use of NLX.
103

RESULTS

Intravenous fentanyl induces respiratory depression and activates nTS neurons in a dose-dependent manner

We first examined the effect of intravenous fentanyl to induce cardiorespiratory depression and
activate neurons in the nucleus tractus solitarius (nTS). Catheterized rats received intravenous
saline (1 ml/kg) or fentanyl at a range of doses (2, 20, or 50 µg/kg; **Figure 1A**), and
cardiorespiratory parameters were measured up to 60 minutes. The first 20 minutes of these
responses are shown in **Figure 1B-D**. Neither saline nor 2 µg/kg fentanyl evoked a significant
change in cardiorespiratory parameters. In comparison, both 20 and 50 µg/kg fentanyl produced
rapid decreases in oxygen saturation, heart rate, and respiratory rate, with the largest decrease
occurring after 50 µg/kg fentanyl. The nadir (defined as the lowest point measured after
fentanyl) and recovery (calculated as 90% of baseline values prior to fentanyl) for each
physiological parameter were assessed in all groups (**Figure 1E-J**). There was no significant
difference in the nadir or recovery values between saline- and 2 µg/kg fentanyl-treated rats. In
comparison, rats given 50 µg/kg fentanyl exhibited a significantly lower nadir in oxygen
saturation (**Figure 1B,E**) and respiratory rate compared to 20 µg/kg (**Figure 1D,I**). In addition,
the recovery time was significantly longer in rats that received 50 µg/kg fentanyl compared to 20
µg/kg fentanyl (**Figure 1B,E, D,I**). The nadir of bradycardia was similar between 20- and 50
µg/kg rats (**Figure 1G**), although the duration of the response was significantly longer in rats
that received the highest dose of fentanyl (**Figure 1H**). We next examined nTS activation (Fos-
IR) following intravenous fentanyl (**Figure 1K)**. A significant increase in Fos-IR was observed
after 2 µg/kg fentanyl, (**Figure 1L**), a dose that did not induce OIRD. Further dose-dependent
increases in Fos-IR were observed after 20 and 50 µg/kg fentanyl. We evaluated a
subpopulation of catecholaminergic nTS neurons, which are critical for full expression of
hypoxia-evoked cardiorespiratory responses (*22, 33*) for activation after intravenous fentanyl
(**Figure 1M**). These cells were similarly activated at 2 and 20 µg/kg fentanyl, with a further
increase observed at the highest dose (50 µg/kg).
131

Peripheral opioid receptor antagonism prevents fentanyl-induced respiratory depression

The above data provide a comprehensive assessment of the dose-dependent changes in cardiorespiratory parameters and nTS neuronal activation evoked by intravenous fentanyl. To gain insight into the extent to which central and peripheral opioid receptors contribute to these effects, a separate cohort of rats received intravenous administration of the non-selective opioid receptor antagonist naloxone (NLX; 1 mg/kg) or the peripherally-restricted opioid receptor antagonist naloxone methiodide (NLXM, 1 and 5 mg/kg) prior to intravenous fentanyl administration at 20 µg/kg (**Figure 2A**) or 50 µg/kg (**Figure S2A**). It has been previously shown that NLXM does not cross the blood brain barrier at low doses (*31*, *34*). Our data are in agreement with these reports, as we show that neither dose of NLXM reached a significant level within the central nervous system at either time point evaluated (**Figure 6**). Thus, we utilized NLXM to further investigate the roles of peripheral and central MOR in OIRD.

As shown in **Figure 2B-D**, saline pretreatment followed by 20 µg/kg fentanyl produced an onset and duration of cardiorespiratory depression similar to **Figure 1**. Similar responses were observed in rats that received saline pretreatment followed by 50 µg/kg fentanyl (**Figure S1B-D**). As expected, NLX pretreatment completely blocked the decrease in cardiorespiratory parameters at both 20 µg/kg and 50 µg/kg doses of fentanyl. In rats that received 20 µg/kg fentanyl, 1 mg/kg NLXM pretreatment prevented the bradycardic and hypoventilatory responses to fentanyl (**Figure 2I-D**). For all parameters, saline-pretreated rats exhibited lower nadir values than NLX- and NLXM-pretreated rats, and there was no significant difference in the nadir or duration between NLX (1mg/kg) and NLXM (1mg/kg) pretreated rats that received 20 µg/kg fentanyl (**Figure 2E-J**). There was a slight decrease in the nadir of oxygen saturation in NLXM-pretreated rats vs NLX-pretreated rats, although this was not significant (**Figure 2E**, p=0.0876). In rats that received 50 µg/kg fentanyl, 1 mg/kg NLXM pretreatment was unable to prevent the initial decrease in all cardiorespiratory parameters (**Fig S1E,G,I**). However, the duration of these responses was significantly shorter compared to saline-treated rats (**Figure S1F,H,J**). A higher dose of NLXM (5 mg/kg), which does not cross the blood brain barrier (**Figure 5D**), was then used in a separate group of animals that received 50 µg/kg fentanyl. Pretreatment with 5 mg/kg NLXM fully prevented fentanyl-induced bradycardia and hypoventilation but did not completely prevent a decrease in oxygen saturation, although the recovery time was similar to the NLX-pretreated group (p = 0.7151) (**Figure S1B-J**). Together, these data demonstrate that peripheral MOR antagonism is sufficient to prevent 20 µg/kg fentanyl-induced cardiorespiratory depression and strongly attenuates OIRD evoked by 50 µg/kg fentanyl.

Neuronal activation, as measured by Fos-IR, was examined in the nTS of rats pretreated

with saline or opioid receptor antagonists followed by 20 µg/kg (**Figure 2K**) or 50 µg/kg fentanyl

(**Figure S1K**). For fentanyl doses, saline pretreatment evoked the highest degree of Fos-IR in

the nTS, including in a subpopulation of TH-IR neurons. Blocking peripheral MOR signaling with

NLXM did not decrease fentanyl-induced nTS neuronal activation, while NLX, which blocks both

central and peripheral MORs resulted in a significant reduction in nTS activation (**Figure 2L,M**).

In rats that received the highest dose of fentanyl, the number of Fos-IR was similar between rats

pretreated with saline and NLXM at both doses, with the only significant reduction occurring in

NLX-pretreated animals (**Figure S1L**). This pattern of activation was also observed in

catecholaminergic cells (**Figure S1M**), although around 30% of TH-IR neurons also displayed

Fos-IR in NLX-pretreated rats.

Peripheral opioid receptor antagonism reverses fentanyl-induced respiratory depression
The above data demonstrate that peripheral MOR antagonism largely prevents fentanyl-induced

respiratory depression but only minimally reduces nTS neuronal activation. Given that reversal

of OIRD has more relevance as a therapeutic intervention, we next evaluated the extent to

which NLXM reverses cardiorespiratory depression after it has been induced by 20 µg/kg

(**Figure 3A**) and 50 µg/kg fentanyl (**Figure S2A**). Both 20 and 50 µg/kg fentanyl induced rapid

decreases in oxygen saturation, heart rate and respiratory rate, similar to what was observed in

Figure 1 (**Figure 3B-D; Figure S2B-D**). The fentanyl-induced nadir in all physiological

parameters was similar prior to intravenous administration of saline, NLX or NLXM. In rats that

received 20 µg/kg fentanyl, both intravenous NLX (1 mg/kg) and NLXM (1 mg/kg) quickly

restored cardiorespiratory values to baseline significantly faster than saline-pretreated animals

(**Figure 3F,H,J**). This reversal of cardiorespiratory parameters was similar between NLX- and

NLXM-treated animals. In rats that received 50 µg/kg fentanyl, NLX quickly restored all

physiological parameters to their baseline values (**Figure S2F,H,J**). While 1 mg/kg NLXM did

not sufficiently reverse the OIRD induced by 50 µg/kg fentanyl (**Figure S2F**)**,** a higher dose of 5

mg/kg NLXM fully reversed the fentanyl-induced bradycardia and hypoventilation, similar to

NLX. Importantly, this dose of NLXM is not detectable in the central nervous system (**Figure**

5D). Thus, blocking peripheral MORs is sufficient to fully reverse OIRD.

Because acute hypoxia activates nTS neurons (*23*), we investigated the impact of

peripheral and central MOR antagonism on fentanyl-induced nTS neuronal activation (**Figure**

3K). At both doses of fentanyl administered, a similar degree of nTS neuronal activation was

observed in all groups regardless of MOR antagonist administered after fentanyl (**Figure 3L,M;**

Figure S2L,M). Together, these data demonstrate that selective antagonism of peripheral

opioid receptors is sufficient to reverse cardiorespiratory depression induced by both moderate

and high doses of fentanyl. Moreover, these data suggest that even a brief period of fentanyl-

induced cardiorespiratory depression is sufficient to induce robust activation of nTS neurons,

and this subsequent MOR antagonism has no additional effect on nTS activation.

Reversal of Fentanyl-Induced Respiratory Depression with NLXM is not aversive

NLX rapidly reverses OIRD, but produces unwanted side effects including the immediate

onset of severe withdrawal (*12*, *13*). The intense symptoms induced during NLX-precipitated

withdrawal present additional challenges in managing OUD in patients. Given that peripheral

MOR antagonism sufficiently reverses fentanyl-induced cardiorespiratory depression to a

degree that is similar to NLX, we then used a conditioned place aversion (CPA) approach to

assess whether selective peripheral antagonism of MOR signaling also mediates aversive

behaviors. Rats underwent a 3-session CPA challenge that consisted of a Pre-Conditioning test,

Conditioning test, and Post-Conditioning test (**Figure 4A**). Horizontal locomotion in both

compartments of the CPA boxes was assessed during both pre- and post-conditioning tests

(**Figure 4B**). During the pre-conditioning test, rats exhibited no natural preference or aversion

for either compartment. On the Conditioning day (Day 2), a divider was placed in the middle of

the CPA box. Rats received intravenous fentanyl injection (50 µg/kg) in their home cage. Four

minutes later, rats were placed into the paired compartment of the CPA box and immediately

given intravenous saline (1 m1/kg), NLX (1 mg/kg) or NLXM (1 or 5 mg/kg). During the Post-

conditioning test, the divider was removed and rats were allowed to explore both compartments

of the CPA box. **Figure 4C** shows the time spent on the conditioned side of the CPA box for all

groups. During the Post-Conditioning test, NLX-treated animals spent significantly less time in

the NLX-paired compartment, highlighting the aversive responses that are evoked by NLX

infused after fentanyl treatment. In contrast, NLXM did not produce aversive behaviors to the

NLXM-paired compartment, but rather induced a preference at the higher dose (**Figure 4D**), an

effect that may be mediated by fentanyl acting at central MORs. Importantly, a single

conditioning session with fentanyl administered in the home cage followed by saline in the

compartment instead of a MOR antagonist did not produce a preference or aversion to the

saline-paired compartment. As shown in **Figure S3**, neither NLX nor NLXM significantly alters

physiological parameters from baseline values. To rule out any potential effects of endogenous

opioids influencing the behavioral responses to opioid receptor antagonism, we performed an

additional test in which rats were given saline in their home cage, followed by NLX or NLXM in

one of the CPA compartments (**Figure 4E,F**). Neither antagonist produced a preference nor

aversion to the MOR antagonist-paired compartment (**Figure 4G**). There was no difference in

CPA scores between these groups (**Figure 4H**).

Fentanyl induces a biphasic pattern of activity in nTS neurons

As shown in Figure 1, fentanyl induced robust Fos-IR expression in the nTS. While Fos-IR

quantification provides insight into the anatomical activation of brainstem subregions in

response to fentanyl administration and OIRD reversal, this approach does not provide any

temporal resolution of the activity of these cells. Therefore, we used wireless *in vivo* fiber

photometry to assess dynamic changes in nTS calcium transient activity after intravenous

fentanyl administration in anesthetized rats (**Figure 5A**). Catheterized rats underwent bilateral

injections of an AAV expressing a calcium sensor (pGP-AAV-syn-jGCaMP8m-WPRE) into the

nTS. On the day of the experiment, an optic fiber was lowered on the surface of the brainstem

above the nTS and calcium transient activity was recorded (**Figure 5B**). GCaMP viral

expression was observed throughout the caudal-rostral extent of the nTS in all rats (**Figure 5C**).

Fentanyl (20 µg/kg) infusion induced a biphasic pattern of activity in the nTS, characterized by

an initial robust increase in activity (Early Response) followed by a decrease below baseline

(Late Response, **Figure 5D**). Pretreatment with NLXM (5 mg/kg) attenuated both phases of the

fentanyl-induced changes in neuronal activity. As shown in **Figure 5E**, the area under the cure

(AUC) was significantly elevated during the Early and Late Responses after fentanyl. In

contrast, no significant fentanyl-induced change in AUC was observed in NLXM-pretreated rats

at either time point, indicating that changes in fentanyl-induced nTS neuronal activity are

strongly mediated by peripheral MORs. We next examined the extent to which MOR antagonists

restore the fentanyl-induced reductions in activity. After MOR antagonist administration,

fentanyl-induced suppression of nTS activity returned to baseline during the early recovery in

rats that received NLX, but not NLXM. However, both NLX and NLXM enhanced nTS activity

above baseline during the late recovery period (**Figure 5F,G).** During the Late Recovery period,

nTS activity was significantly enhanced above baseline in both NLX and NLXM-treated animals.

Together, these data demonstrate that peripheral MOR antagonism is sufficient to reverse the

fentanyl-induced decrease in nTS neuronal activity.

DISCUSSION

OIRD is a complex problem that continues to be at the forefront of the opioid epidemic

(*1*, *35*). The increased potency and faster onset of synthetic opioids like fentanyl has been

attributed to the rising wave of opioid-related deaths in the US (*2*, *11*). Numerous studies have

evaluated opioid-induced dysfunction within key regulatory sites in the central nervous system
that contribute to the severity of OIRD (*1*, *14*, *15*, *17*, *27*). Peripheral MORs have also been
examined for their roles in impacting OIRD (*12*, *32*). However, the specific contributions of
central and peripheral MORs to the onset and duration of fentanyl-induced cardiorespiratory
depression, along with the role of the nTS in these effects, are not completely understood. In
this study, we show that blocking peripheral MORs sufficiently prevents and reverses fentanyl-
induced cardiorespiratory depression and systemic hypoxia. Furthermore, we demonstrate that
peripheral opioid receptors mediate fentanyl-induced activity of nTS neurons. These data
suggest that the nTS may be an area impacted by fentanyl via peripheral MOR activation.
Lastly, we demonstrate that in addition to reversing fentanyl-induced cardiorespiratory
depression, NLXM does not induce aversive behaviors, as compared to NLX. Together, these
findings provide evidence that peripheral MOR antagonism may be a potential novel target to
reverse OIRD without inducing withdrawal, anxiety, and aversion, all of which can contribute to
further relapse in drug-seeking behaviors (*2*, *13*).

While numerous studies have focused on evaluating the effects of opioids acting at
MORs present within brainstem and pontine networks, less attention has been given to
identifying peripheral MOR mechanisms contributing to OIRD. NLX is the most common
reversal agent used to treat OIRD (*13*, *35*). Despite its efficacy in treating OIRD, NLX induces
withdrawal symptoms and aversion in both patients and preclinical models (*12*, *13*, *36*). In this
study, we demonstrate that NLX-induced reversal of fentanyl-induced respiratory depression, as
shown by a significant decrease in time spent in the NLX-associated compartment. This is likely
the result of the induction of withdrawal symptoms that occur following antagonism of MOR in
central reward pathways (*12*, *13*, *37*). In contrast, this aversive response was not observed in
rats that received NLXM; NLXM-treated rats exhibited a preference for the associated
compartment. These data suggest that alleviating fentanyl-induced respiratory distress without
antagonizing central MOR may mitigate the aversive state produced by NLX. While previous
reports have shown that NLX can be detected centrally after a subcutaneous injection of NLXM
(*31*), we report here that no detectable amount of NLX or NLXM was found in brain tissue at the
doses used in these studies (1 and 5 mg/kg, intravenous). Similar to NLX, peripheral MOR
antagonism with NLXM partially reverses opioid-induced analgesia (*12*, *38*) which may pose
limitations for pain management. Future investigations are needed to evaluate mechanisms
underlying the therapeutic potential of NLXM as an intervention for managing OIRD in patients
with OUDs.

The nTS is a region with a high degree of MOR expression, and appears to be primarily

located on afferent fibers arising from the vagus nerve (*7*, *24*, *41*). While MOR has been shown

to be present in nTS somas and dendrites (*24*, *27*), these MOR-expressing cells do not express

Fos following exposure to opioids or hypoxia (*27*), which suggests a mechanism of presynaptic

action of MOR signaling in the nTS. Likewise, the nodose ganglion (NG), which contains the cell

bodies of nTS-projecting vagal sensory afferents, has high expression of MOR (*30*), and intra-

nodose injection of fentanyl induces hypoventilation, although to a much lesser degree

compared to systemic fentanyl (*42*).

In the present study, we used fiber photometry to characterize nTS neuronal responses

to show that fentanyl induces an initial excitatory response in nTS cells. However, this response

was short-lasting compared to the subsequent decrease in activity below baseline, suggesting

that the primary effect of MOR signaling produce generalized inhibition of nTS neurons. This

prolonged suppression of nTS neuronal activity may contribute to the overall duration of OIRD.

Furthermore, our photometry data demonstrate that while the NLXM-mediated reversal of

fentanyl-induced inhibition of nTS neurons was slower compared to NLX, activity was eventually

restored. A similar finding was observed in OIRD experiments, as NLXM induced a rate of

recovery in all parameters that was comparable to NLX. Taken together, this suggests that

peripheral MOR antagonism may sufficiently override simultaneous central effects of fentanyl

acting at MOR present in cardiorespiratory nuclei, including the nTS.

It is important to note that while all our experiments assessing fentanyl-induced

cardiorespiratory depression were conducted in conscious rats breathing room air, photometry

experiments were performed under isoflurane anesthesia. Due to the location of the nTS,

chronic fiber implantation and subsequent in vivo photometry recordings in conscious,

unrestrained animals is not possible. In the nTS, Isoflurane has been reported to suppress

visceral afferent glutamatergic transmission and enhance GABAergic transmission via

postsynaptic mechanisms (*39*), as well as induce cFos expression in the nTS (*40*). Precautions

were taken to use the lowest dose of isoflurane needed to maintain sufficient anesthesia.

However, we cannot rule out the possible influence of isoflurane anesthesia to influence the

observed biphasic responses evoked by fentanyl. Nevertheless, our data support a mechanism

by which fentanyl induces OIRD via disruption of visceral afferent signaling to the nTS, thereby

leading to prolonged OIRD due to suppression of physiological reflex responses that would be

normally engaged in response to systemic hypoxia.

The nTS plays an essential role in the regulation of basal-and reflex-evoked

cardiorespiratory function (*21*). Activation of MOR in the nTS has been shown to blunt hypoxic

ventilatory responses following MOR agonism in the nTS (*7*). Our data are consistent with

previous work demonstrating opioid-induced activation of nTS neurons (*27*). The degree of

fentanyl induced Fos-IR in the nTS resembles what is observed after acute exposure to hypoxic

air (*20, 46*). Interestingly, we also observed increased Fos in the nTS at a dose of fentanyl that

was below the threshold to induce respiratory depression. It is unclear whether the Fos-IR

observed after low or high doses of fentanyl is, in part, the result of increased glutamatergic

transmission stemming from enhanced chemoreceptor afferent signaling to the nTS or via MOR

signaling within the nTS itself.

Glutamate is the primary neurotransmitter released from sensory afferent fibers into the

nTS (*21, 43*). While intra-nTS application of opioid agonists have been shown to reduce

glutamate release and calcium channels via presynaptic mechanisms (*44, 45*), opioids also

have been shown to induce a similar biphasic action of Ca2+-dependent spikes in the nodose

ganglion (*46*) that resembles what we observed in GCaMP-expressing nTS cells after

intravenous fentanyl. NLXM strongly attenuated the fentanyl-induced excitatory wave observed

in GCaMP-expressing nTS neurons, suggesting that MOR-expressing peripheral afferents

mediate this transient activation of nTS neurons, even if the primary effect of fentanyl is to inhibit

the nTS and suppress cardiorespiratory reflex function.

One possibility is that the increase in nTS activity may be mediated by opioid-induced

disinhibition within the nTS. The nTS contains a large population of second order GABAergic

interneurons that receive direct inputs from visceral afferents (*47*), and hypoxia activates these

cells(*22*). nTS MOR agonists suppress the activity of these cells, primarily via presynaptic

mechanisms (*48*), although postsynaptic mechanisms have been reported (*49*). Similarly,

visceral afferents also activate catecholaminergic nTS neurons (*50*), which are critical for the

generation of hypoxia-evoked cardiorespiratory responses(*33, 51*). Opioids inhibit visceral

afferent transmission to this cell group, primarily via a presynaptic mechanism (*44*). In the

present study, we observed robust Fos expression in these cells at all doses of fentanyl

examined. These findings raise the possibility that nTS dysfunction, via MOR-expressing

peripheral inputs that terminate within this region, represents the first central site of fentanyl-

induced dysfunction, ultimately leading to downstream effects in other brain regions that

contribute to diminished autonomic and cardiorespiratory responses that contribute to the

duration of OIRD. Future studies are needed to fully evaluate the exact opioid-MOR effects in

the nTS, including specific MOR afferent inputs and the nTS phenotypes, to determine their

relative contributions to OIRD.

In summary, these studies highlight a significant involvement of peripheral MORs

contribute to the rapid cardiorespiratory depression induced by intravenous fentanyl. We have

provided extensive characterization into how fentanyl elicits cardiorespiratory depression and

examined the relationship between peripheral MORs and nTS activity in contributing to the

duration of OIRD. Moreover, our findings suggest that selectively blocking peripheral ORs could

be a promising therapeutic strategy for managing OIRD. Notably, peripheral opioid receptor

antagonists are already prescribed for patients undergoing chemotherapy to alleviate side

effects such as opioid-induced constipation (*52, 53*). Importantly, our data demonstrate that

peripheral MOR antagonism prevents and reverses OIRD without triggering unwanted effects of

withdrawal that occur following NLX administration, suggesting that peripheral MOR antagonists

like NLXM could potentially be utilized in treatment strategies for managing OUD.

Materials and Methods:

All surgical and experimental procedures were approved by Washington University Committee

in accordance with the National Institutes of Health Guidelines for the Care and Use of

Laboratory Animals and Animal Research: Reporting In Vivo Experiments (ARRIVE) guidelines.

Rats were initially group-housed with two to three animals per cage on a 12/12-h dark/light cycle

(lights on at 7:00) and acclimated to the animal facility holding rooms for at least 7 days before

any manipulation. Following catheterization surgery, rats were single-housed and remained on

an identical 12/12-h dark/light cycle. The temperature for the holding rooms of all animals

ranged from 21 to 24 °C while the humidity was between 30 and 70%. Rats received food and

water ad libitum for the duration of all experiments. All experiments were performed during the

light cycle. For all experiments, male and female Sprague Dawley rats (250-350g) were used. In

all data sets, male rats are represented by filled symbols and female rats are represented by

open symbols.

IV catheterization procedure:

Animals were deeply anesthetized using Isoflurane (3% induction, 2% maintenance). A small

incision was made in the dorsal surface of the neck. Another small incision was made on the

ventral surface of the neck, and underlying tissue was carefully dissected to expose the jugular

vein. An indwelling catheter was inserted into the jugular vein and sutured in place using non-

absorbable silk suture. The catheter was then tunneled subcutaneously and exited the body

through the first small hole in neck on the dorsal side. The ventral incision was closed using

non-absorbable silk suture. The exposed catheter was connected to a backpack device

(Instech) containing a port for drug administration. Animals were given Carprofen (2 mg/kg sc),

Baytril (8 mg/kg sc) and bupivacaine (5 mg/kg, site of incision). In addition, rats were given

Carprofen tablets (Bio Serv, MD150-2) for 2 days after surgery to assist in wound healing and

analgesia. Rats were singly housed following surgery, and were allowed to recover for one

week prior to the commencement of experiments. Catheter patency was maintained with daily

flushing of 0.3 mL sterile saline containing gentamicin (1.33 mg/mL, i.v.)

Respiratory depression experiments:

Following recovery from IV catheterization surgery, conscious, freely-moving rats in their home

cages were fitted with a non-invasive pulse oximeter collar (Starr LifeSciences).

Cardiorespiratory parameters (oxygen saturation, heart rate and respiratory rate) were collected

(MouseOx v2.0, Starr LifeSciences) at baseline conditions in animals breathing room air, and

after intravenous administration of fentanyl citrate (2, 20 or 50 µg/kg). A subset of animals

received weight-adjusted intravenous administration of the nonselective opioid receptor

antagonist naloxone hydrochloride (NLX; 1 mg/kg; Sigma 51481-60-8) or the peripherally-

restricted opioid receptor antagonist naloxone methiodide (NLXM; 1 or 5 mg/kg; Sigma N129)

before or after fentanyl administration. Cardiorespiratory parameters were then measured for up

to 60 minutes. For all OIRD experiments, the time course graphs consist of 30 seconds of data

averaged into a single data point. The nadir was defined as the lowest point measured after

fentanyl administration (for all graphs, red arrow indicates time of fentanyl administration,

time=0 minutes). In addition, the recovery rate was defined as the time to reach 90% baseline of

pre-fentanyl values for each cardiorespiratory parameter. Two hours after fentanyl

administration, rats were transcardially perfused with 0.01M PBS followed by paraformaldehyde

(4% in 0.01M PBS). Brains were removed and stored overnight at 4 °C in 4% paraformaldehyde

for post-fixation, followed by at least 72 h incubation in sucrose (30% in 0.01M PBS) solution.

Isopentane was used to flash-freeze brains. Brainstem sections containing the nTS were sliced

in the coronal plane (40 µm) using a cryostat (Leica CM 1950).

Biodistribution

A subset of previously catheterized drug-naïve rats was deeply anesthetized and then received

intravenous administration of NLX (1 mg/kg) or NLXM (1 or 5 mg/kg). While still under

anesthesia, rats were quickly decapitated two or ten minutes after intravenous administration of

MOR antagonists. MOR antagonists were dosed in all rats at an experimentally relevant

concentration with triplicate blood and brain samples collected at two time points. The early time

point at 2 minutes evaluated conditions when the plasma concentration of the OR antagonist is

high and reflects the identical time point of intravenous administration for OIRD pretreatment

experiments. The second collection time point (10 minutes) was intended to evaluate steady-

state conditions and represent physiological experiments when reversal of OIRD was achieved

by both antagonists. Brains were quickly removed and flash frozen in liquid nitrogen. Trunk

blood was collected into capillary collection tubes containing lithium heparin (Greiner Bio-One;

Fisher Scientific). The blood was spun on a centrifuge for 10 min at 4 °C, and plasma was

carefully extracted from the supernatant. Brains and plasma samples underwent PK analyses

for detection of NLX or NLXM at both time points. Standard rat PK studies dosed the candidate

MOR antagonists and sample blood over multiple time points. A general description of tissue

distribution can be obtained during PK modeling by calculating the volume of distribution.

Samples underwent a direct measurement of peripheral restriction by measuring compound

concentration in plasma and brain.

Conditioned Place Aversion

The aversive effects following reversal of OIRD were measured using Conditioned Place

Aversion (CPA) (*54*). During a pre-conditioning session (Day 1), rats were allowed to explore

both sides of a two compartment CPA box for 15 minutes. On the Conditioning Day (Day 2), a

divider was placed in the middle of the box to restrict movement to one side on the box. Rats

were then randomly assigned to one side of the CPA box, and this side was paired with either

saline (1 ml/kg), NLX (1 mg/kg) or NLXM (1 or 5 mg/kg). Rats were given intravenous fentanyl

(50 μg/kg) or saline in their home cages. Four minutes later, they were placed in the paired side

of the box and immediately given saline or a MOR antagonist. Rats were confined to this

compartment for 15 minutes. For all CPA experiments, measures were taken to balance the

side of CPA box by drug and by sex. On the post-conditioning test (Day 3), the divider was

removed. Rats were then placed in the CPA box and allowed to explore both sides of the CPA

box for 15 minutes. The total time spent in the paired chamber and CPA scores were calculated

for all groups.

In vivo Fiber Photometry

To selectively measure nTS neuronal calcium transient activity, we injected an AAV containing a

calcium sensor expressed under the synapsin promoter (pGP-AAV-syn-jGCaMP8m-WPRE) into

the nTS (500 nl per side; 0.3-0.5 mm anterior, ± 0.4 mm lateral and 0.4 mm ventral to brain

surface, relative to calamus scriptorius). Rats were allowed at least 3 weeks to recover from

nanoinjection surgery to allow for robust viral expression in the nTS before IV catheters were

implanted as described above. On the day of the experiment, rats were deeply anesthetized

with isoflurane (3-4% induction, 2% maintenance), placed in the stereotaxic apparatus, and the

nTS was exposed as described above. To overcome limitations associated with chronic fiber

implants in the nTS, a wireless photometry headstage (TeleFipho, Amuza Inc.) was secured to

a 5 mm flat-tipped silica optic fiber cannula (400 μm c.d./470 μm o.d., NA 0.37, 2.5 mm

receptacle; Doric) and mounted the stereotaxic arm with an apparatus 3D printed in house to

permit simultaneous nTS exposure and photometry recordings. The optic fiber was lowered to

the site of the dorsal brainstem and firmly pressed on the surface above the nTS until a signal

was detected (~0.3-0.5 mm ventral from the surface). An LED generated blue light was

bandpass-filtered (445-490 nm) to excite jGCaMP8m and emission fluorescence was detected

by an internal photodiode detector for green light (bandpass filtered at 500-550 nm). An internal

DC amplifier transmitted the data (16-bit arbitrary unit; AU) to a wireless receiver (TeleFipho,

Amuza Inc.) and the data was extracted in real-time using TeleFipho software (Amuza Inc.) at a

sampling rate of 100 Hz. The offset was set to 90° and the LED power was adjusted to achieve

an optimal signal output (30-40k a.u.) not to exceed 36 mW power, and remained constant

throughout the recording. Baseline activity of GCaMP fluorescence nTS-infected cells was

measured after at least 5 min to mitigate effects of baseline drift in signal due to slow

photobleaching artifacts. Then, baseline activity was measured for at least 30 s prior to

administration of either fentanyl (20 µg/kg, i.v.) or NLXM (5 mg/kg, i.v.). After response activity

was recorded, rats received a second intravenous injection of either an opioid receptor

antagonist (NLX,1 mg/kg or NLXM, 5 mg/kg) or fentanyl (20 µg/kg), respectively. Each

intravenous administration was manually time locked to calcium transient recordings by a

second experimenter and response activity following each injection was recorded for 60 s

before administration of the second injection or conclusion of the experiment. Rats were

transcardially perfused with PFA at the end of the experiment as described above and viral

expression was validated for all animals.

For analysis of calcium transient activity, the raw data was fit to a double-exponential curve and

subtracted from the raw data to account for any residual effects of baseline drift. Matlab Signal

Analyzer was used to extract regions of interest (ROIs) in the signal in 90-s traces (30 s pre-

treatment, 60 s post-treatment) and traces were aligned to the time of injection. ROIs were

standardized as z-scores using the sample standard deviation, S, where $S = \sqrt{\frac{\sum_{i=1}^{n}(x_i - \bar{X})^2}{n-1}}$ such

that signals were centered to have a mean of 0 and scaled to have a standard deviation of 1.

Based on this, the area under the curve (AUC) of ROI z-scores were calculated in 30-s bins

using the trapezoidal rule (*trapz* function in Matlab) and one-sample t-tests were used to

determine differences relative to baseline (theoretical mean = 0). For presentation, transients

were down sampled by a factor of 10 with a phase offset of 2 and the resultant 10 Hz traces

were smoothed using a moving mean duration of 5% of total ROI.

Immunohistochemistry (IHC):

Immunohistochemical procedures were performed as previously described (*55*). Brainstem

sections containing the nTS were sliced using a cryostat (Leica) at a thickness of 40 um. Free-

floating sections were rinsed in 0.01M PBS, blocked in 10% Normal Donkey Serum (Sigma) in

0.01M PBS-triton, and then incubated with the following antibodies: cFos (Rabbit anti-cFos;

ab190289, Abcam; AB_2737414); Tyrosine Hydroxylase (TH, Mouse anti-TH; MAB318,

Millipore; AB_2201528). The following day, sections were rinsed with 0.01M PBS, incubated

with Donkey anti-Rabbit Cy3 and Donkey anti-Mouse Alexa fluor 488 (1:200; in 3% NDS and

0.3% Triton in 0.01M PBS) and used to visualize targets of interest (Jackson ImmunoResearch).

Image acquisition:

512 Immunoreactivity (IR) and viral expression were examined with a Leica DMR microscope at 20X

513 magnification. 20 μm thick z-stacks of the nTS (2 μm per plane) were collected. ImageJ (v. 1.48,

514 NIH) software was used for post-processing and analysis. In all groups, three sections of the

515 nTS (corresponding to -360, 0, +360 μm relative to calamus scriptorius) were collected.

Brainstem immunohistochemical analyses

517 Using ImageJ software, the regions containing the nTS were outlined, and unilateral counts of

518 Fos-IR, TH-IR and co-labeled Fos- and TH-IR neurons were performed using a custom-made

519 plugin (Cell Counter) for each of the 3 nTS sections per animal. The total number of each

520 phenotype was summed from these 3 sections.

Statistics

522 For all experiments, the normality of sample data was determined using D'Agostino and

523 Pearson tests and Shapiro–Wilk tests. Statistical significance was taken as * $p<0.05$, ** $p<0.01$,

524 *** $p<0.001$ and ****$p<0.0001$, as determined by two-way repeated measures analysis of

525 variance (ANOVA) followed by Sidak's post hoc test; one-way ANOVA with Tukey post-hoc test,

526 two-tailed unpaired t-test, two-tailed paired t-test, one sample t-test.

References

References

1.   J.-M. Ramirez, N. J. Burgraff, A. D. Wei, N. A. Baertsch, A. G. Varga, H. A. Baghdoyan, R.
Lydic, K. F. Morris, D. C. Bolser, E. S. Levitt, Neuronal mechanisms underlying opioid-
induced respiratory depression: our current understanding. *Journal of Neurophysiology*
125, 1899–1919 (2021).

2.   S. D. Comer, C. M. Cahill, Fentanyl: Receptor pharmacology, abuse potential, and
implications for treatment. *Neuroscience & Biobehavioral Reviews* **106**, 49–57 (2019).

3.   C. L. Mattson, Trends and Geographic Patterns in Drug and Synthetic Opioid Overdose
Deaths — United States, 2013–2019. *MMWR Morb Mortal Wkly Rep* **70** (2021).

4.   S. Parida, J. P. De Aquino, M. Sofuoglu, High potency synthetic opioids: Curbing the third
wave of the opioid crisis. *Neuroscience & Biobehavioral Reviews* **106**, 9–10 (2019).

5.   N. Wilson, M. Kariisa, P. Seth, H. Smith, N. L. Davis, Drug and Opioid-Involved Overdose
Deaths — United States, 2017–2018. *MMWR Morb Mortal Wkly Rep* **69**, 290–297 (2020).

6.   A. Mansour, C. A. Fox, S. Burke, F. Meng, R. C. Thompson, H. Akil, S. J. Watson, Mu,
delta, and kappa opioid receptor mRNA expression in the rat CNS: An in situ hybridization
study. *Journal of Comparative Neurology* **350**, 412–438 (1994).

7.   J. Zhuang, X. Gao, F. Gao, F. Xu, Mu-opioid receptors in the caudomedial NTS are critical
for respiratory responses to stimulation of bronchopulmonary C-fibers and carotid body in
conscious rats. *Respir Physiol Neurobiol* **235**, 71–78 (2017).

8.   R. Hill, R. Santhakumar, W. Dewey, E. Kelly, G. Henderson, Fentanyl depression of
respiration: Comparison with heroin and morphine. *Br J Pharmacol* **177**, 254–265 (2020).

9.   R. C. N. Marchette, E. R. Carlson, E. V. Frye, L. E. Hastings, J. C. M. Vendruscolo, G.
Mejias-Torres, S. J. Lewis, A. Hampson, N. D. Volkow, L. F. Vendruscolo, G. F. Koob,
Heroin- and Fentanyl-Induced Respiratory Depression in a Rat Plethysmography Model:
Potency, Tolerance, and Sex Differences. *J Pharmacol Exp Ther* **385**, 117–134 (2023).

10. T. Lewanowitsch, R. J. Irvine, Naloxone and its quaternary derivative, naloxone
methiodide, have differing affinities for mu, delta, and kappa opioid receptors in mouse
brain homogenates. *Brain Res* **964**, 302–305 (2003).

11. N. Fairbairn, P. O. Coffin, A. Y. Walley, Naloxone for heroin, prescription opioid, and illicitly
made fentanyl overdoses: Challenges and innovations responding to a dynamic epidemic.
International Journal of Drug Policy **46**, 172–179 (2017).

12. T. Lewanowitsch, R. J. Irvine, Naloxone methiodide reverses opioid-induced respiratory
depression and analgesia without withdrawal. *European Journal of Pharmacology* **445**, 61–
67 (2002).

13. J. T. Lai, C. E. Goldfine, B. P. Chapman, M. M. Taylor, R. K. Rosen, S. P. Carreiro, K. M.
Babu, Nobody Wants to Be Narcan'd: A Pilot Qualitative Analysis of Drug Users'
Perspectives on Naloxone. *West J Emerg Med* **22**, 339–345 (2021).

14.   I. Bachmutsky, X. P. Wei, E. Kish, K. Yackle, Opioids depress breathing through two small
brainstem sites. *eLife* **9**, e52694 (2020).

15.   J. T. Bateman, E. S. Levitt, Opioid suppression of an excitatory pontomedullary respiratory
circuit by convergent mechanisms. *eLife* **12**, e81119 (2023).

16.   A. Haji, H. Yamazaki, Y. Ohi, R. Takeda, Distribution of μ receptors in the ventral
respiratory group neurons; immunohistochemical and pharmacological studies in
decerebrate cats. *Neuroscience Letters* **351**, 37–40 (2003).

17.   E. S. Levitt, A. P. Abdala, J. F. R. Paton, J. M. Bissonnette, J. T. Williams, μ opioid
receptor activation hyperpolarizes respiratory-controlling Kölliker–Fuse neurons and
suppresses post-inspiratory drive. *The Journal of Physiology* **593**, 4453–4469 (2015).

18.   B. Palkovic, V. Marchenko, E. J. Zuperku, E. A. E. Stuth, A. G. Stucke, Multi-Level
Regulation of Opioid-Induced Respiratory Depression. *Physiology (Bethesda)* **35**, 391–404
(2020).

19.   G. Montandon, W. Qin, H. Liu, J. Ren, J. J. Greer, R. L. Horner, PreBötzinger Complex
Neurokinin-1 Receptor-Expressing Neurons Mediate Opioid-Induced Respiratory
Depression. *J. Neurosci.* **31**, 1292–1301 (2011).

20.   K. T. S. Pattinson, Opioids and the control of respiration. *British Journal of Anaesthesia*
100, 747–758 (2008).

21.   M. C. Andresen, D. L. Kunze, Nucleus tractus solitarius--gateway to neural circulatory
control. *Annu Rev Physiol* **56**, 93–116 (1994).

22.   T. L. King, C. M. Heesch, C. G. Clark, D. D. Kline, E. M. Hasser, Hypoxia activates nucleus
tractus solitarii neurons projecting to the paraventricular nucleus of the hypothalamus.
American Journal of Physiology-Regulatory, Integrative and Comparative Physiology **302**,
R1219–R1232 (2012).

23.   L. J. Teppema, J. G. Veening, A. Kranenburg, A. Dahan, A. Berkenbosch, C. Olievier,
Expression of c-fos in the rat brainstem after exposure to hypoxia and to normoxic and
hyperoxic hypercapnia. *Journal of Comparative Neurology* **388**, 169–190 (1997).

24.   S. A. Aicher, A. Goldberg, S. Sharma, V. M. Pickel, μ-opioid receptors are present in vagal
afferents and their dendritic targets in the medial nucleus tractus solitarius. *Journal of
Comparative Neurology* **422**, 181–190 (2000).

25.   A. Disney, K. M. Olson, A. M. Shafer, S. C. Moore, J. P. Anand, J. R. Traynor, S. M.
Husbands, Opioid Antagonists from the Orvinol Series as Potential Reversal Agents for
Opioid Overdose. *ACS Chem Neurosci* **13**, 3108–3117 (2022).

26.   E. Solis, K. T. Cameron-Burr, Y. Shaham, E. A. Kiyatkin, Fentanyl-Induced Brain Hypoxia
Triggers Brain Hyperglycemia and Biphasic Changes in Brain Temperature.
Neuropsychopharmacol. **43**, 810–819 (2018).

27. S. N. Maletz, B. T. Reid, A. G. Varga, E. S. Levitt, Nucleus Tractus Solitarius Neurons
Activated by Hypercapnia and Hypoxia Lack Mu Opioid Receptor Expression. *Front Mol*
Neurosci **15**, 932189 (2022).

28. E. Salas, E. Alonso, M. J. Polanco, M. V. Cano, M. Ruiz-Gayo, L. F. Alguacil, Differential
regulation of CDK5 and c-Fos expression by morphine in the brain of Lewis and Fischer
344 rat strains. *Neuroscience* **230**, 151–156 (2013).

29. P. J. Cabot, P. R. Dodd, T. Cramond, M. T. Smith, Characterization of non-conventional
opioid binding sites in rat and human lung. *Eur J Pharmacol* **268**, 247–255 (1994).

30. J.-L. Li, T. Kaneko, N. Mizuno, Effects of peripheral nerve ligation on expression of μ-
opioid receptor in sensory ganglion neurons: an immunohistochemical study in dorsal root
and nodose ganglion neurons of the rat. *Neuroscience Letters* **214**, 91–94 (1996).

31. D. Perekopskiy, A. Afzal, S. N. Jackson, L. Muller, A. S. Woods, E. A. Kiyatkin, The Role of
Peripheral Opioid Receptors in Triggering Heroin-induced Brain Hypoxia. *Sci Rep* **10**, 833
(2020).

32. F. Henderson Jr, W. J. May, R. B. Gruber, A. P. Young, L. A. Palmer, B. Gaston, S. J.
Lewis, Low-dose morphine elicits ventilatory excitant and depressant responses in
conscious rats: Role of peripheral μ-opioid receptors. *Open Journal of Molecular and*
Integrative Physiology **3**, 111–124 (2014).

33. T. L. King, B. C. Ruyle, D. D. Kline, C. M. Heesch, E. M. Hasser, Catecholaminergic
neurons projecting to the paraventricular nucleus of the hypothalamus are essential for
cardiorespiratory adjustments to hypoxia. *Am J Physiol Regul Integr Comp Physiol* **309**,
R721-731 (2015).

34. S. McGaraughty, P. Honore, C. T. Wismer, J. Mikusa, C. Z. Zhu, H. A. McDonald, B.
Bianchi, C. R. Faltynek, M. F. Jarvis, Endogenous opioid mechanisms partially mediate
P2X3/P2X2/3-related antinociception in rat models of inflammatory and chemogenic pain
but not neuropathic pain. *Br J Pharmacol* **146**, 180–188 (2005).

35. J. T. Bateman, S. E. Saunders, E. S. Levitt, Understanding and countering opioid-induced
respiratory depression. *British Journal of Pharmacology* **n/a** (2021).

36. J. Neale, J. Strang, Naloxone—does over-antagonism matter? Evidence of iatrogenic harm
after emergency treatment of heroin/opioid overdose. *Addiction* **110**, 1644–1652 (2015).

37. M. J. Katovich, J. W. Simpkins, J. O'Meara, Effects of opiod antagonists and their
quaternary analogs on temperature changes in morphine-dependent rats. *Life Sciences*
39, 1845–1854 (1986).

38. V. Spahn, G. Del Vecchio, D. Labuz, A. Rodriguez-Gaztelumendi, N. Massaly, J. Temp, V.
Durmaz, P. Sabri, M. Reidelbach, H. Machelska, M. Weber, C. Stein, A nontoxic pain killer
designed by modeling of pathological receptor conformations. *Science* **355**, 966–969
(2017).

39. J. H. Peters, S. J. McDougall, D. Mendelowitz, D. R. Koop, M. C. Andresen, Isoflurane
Differentially Modulates Inhibitory and Excitatory Synaptic Transmission to the Solitary
Tract Nucleus. *Anesthesiology* **108**, 675–683 (2008).

40. Y. Hu, J. T. Qi, Z. Zhang, M. Q. Luo, Y. W. Wang, Comparative Brain-Wide Mapping of
Isoflurane and Ketamine-Activated Nuclei and Functional Networks. *eLife* **12** (2023).

41. S. Nomura, Y.-Q. Ding, T. Kaneko, J.-L. Li, N. Mizuno, Localization of µ-opioid receptor-
like immunoreactivity in the central components of the vagus nerve: A light and electron
microscope study in the rat. *Neuroscience* **73**, 277–286 (1996).

42. Z. Zhang, C. Zhang, M. Zhou, F. Xu, Activation of Opioid µ-Receptors, but not δ- or κ-
Receptors, Switches Pulmonary C-Fiber-Mediated Rapid Shallow Breathing into An Apnea
in Anesthetized Rats. *Respir Physiol Neurobiol* **183**, 211–217 (2012).

43. W. T. Talman, M. H. Perrone, D. J. Reis, Evidence for L-glutamate as the neurotransmitter
of baroreceptor afferent nerve fibers. *Science* **209**, 813–815 (1980).

44. R. J. Cui, B. L. Roberts, H. Zhao, M. C. Andresen, S. M. Appleyard, Opioids inhibit visceral
afferent activation of catecholamine neurons in the solitary tract nucleus. *Neuroscience*
222, 181–190 (2012).

45. H. Rhim, R. J. Miller, Opioid receptors modulate diverse types of calcium channels in the
nucleus tractus solitarius of the rat. *J. Neurosci.* **14**, 7608–7615 (1994).

46. H. Higashi, P. Shinnick-Gallagher, J. P. Gallagher, Morphine enhances and depresses
Ca2+-dependent responses in visceral primary afferent neurons. *Brain Research* **251**,
186–191 (1982).

47. T. W. Bailey, S. M. Appleyard, Y.-H. Jin, M. C. Andresen, Organization and properties of
GABAergic neurons in solitary tract nucleus (NTS). *J Neurophysiol* **99**, 1712–1722 (2008).

48. A. J. Boxwell, Y. Yanagawa, S. P. Travers, J. B. Travers, The µ-opioid receptor agonist
DAMGO presynaptically suppresses solitary tract-evoked input to neurons in the rostral
solitary nucleus. *J Neurophysiol* **109**, 2815–2826 (2013).

49. N. R. Glatzer, B. N. Smith, Modulation of synaptic transmission in the rat nucleus of the
solitary tract by endomorphin-1. *J Neurophysiol* **93**, 2530–2540 (2005).

50. S. M. Appleyard, D. Marks, K. Kobayashi, H. Okano, M. J. Low, M. C. Andresen, Visceral
afferents directly activate catecholamine neurons in the solitary tract nucleus. *J Neurosci*
27, 13292–13302 (2007).

51. C. S. Bathina, A. Rajulapati, M. Franzke, K. Yamamoto, J. T. Cunningham, S. Mifflin,
Knockdown of tyrosine hydroxylase in the nucleus of the solitary tract reduces elevated
blood pressure during chronic intermittent hypoxia. *Am J Physiol Regul Integr Comp*
Physiol **305**, R1031–R1039 (2013).

52. E. Floettmann, K. Bui, M. Sostek, K. Payza, M. Eldon, Pharmacologic Profile of Naloxegol,
a Peripherally Acting µ-Opioid Receptor Antagonist, for the Treatment of Opioid-Induced
Constipation. *J Pharmacol Exp Ther* **361**, 280–291 (2017).

53.   J. V. Pergolizzi, P. J. Christo, J. A. LeQuang, P. Magnusson, The Use of Peripheral μ-
683   Opioid Receptor Antagonists (PAMORA) in the Management of Opioid-Induced
684   Constipation: An Update on Their Efficacy and Safety. *Drug Des Devel Ther* **14**, 1009–
685   1025 (2020).

686   54.   T. M. Tzschentke, Measuring reward with the conditioned place preference (CPP)
687   paradigm: update of the last decade. *Addict Biol* **12**, 227–462 (2007).

688   55.   B. C. Ruyle, D. Martinez, C. M. Heesch, D. D. Kline, E. M. Hasser, The PVN enhances
689   cardiorespiratory responses to acute hypoxia via input to the nTS. *Am J Physiol Regul
690   Integr Comp Physiol* **317**, R818–R833 (2019).

Acknowledgements

693   We wish to thank Justin Meyer for breeding the rat colonies that were used in these

694   experiments and for general assistance throughout the experiments. We also thank Mike

695   Cameron (Scripps Institute) for mass spectrometry experiments used for all biodistribution

696   experiments. We also wish to thank Kristine Yoon and Thomas Schodl for their technical

697   assistance.

Author Contributions

Conceptualization: BCR NM and JAM

Methodology: BCR, NM and JAM

Formal analysis of all data: BCR, SM, RK, MT and JAH

Photometry acquisition and extraction: BCR and JAH

Respiratory depression experiments: BCR, SM, RK, MT, JM, SAL

Conditioned Place Aversion: BCR, SM, MT, JM

Biodistribution experiments: BCR and TL

Immunohistochemistry, microscopy and cell counting: BCR, SM, RK, MT, JM, SAL

Catheterization surgeries: BCR, SM, RK, MT, JM, SAL

Writing of original draft: BCR

Writing (review and editing): BCR, RK, JM, SAL, TL, JAH, NM, JAM

Funding Acquisition, BR, JM

Resources: JM

Supervision: BR and JM

Data Availability Statement

The authors confirm that all relevant data are included in the paper and its Supplementary Information files. All data generated in this study are provided in the Supplementary Information/Source Data file. Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Jose A. Morón (jmoron-concepcion @wustl.edu).

Competing Interests

We have no competing interests to declare.

Funding:

National Institute of Health grant R01DA042499 (JMC)

National Institute of Health grant R01DA045463 (JMC)

National Institute of Health grant R01DA054900(JMC)

National Institute of Health grant R56DA059067 (JMC)

Washington University McDonnell Small Grants Program GF0012452 (BCR)

National Institute of Health grant T32DA007261 (BCR)

National Institute of Health grant S10OD030332-01(Michael Cameron, Scripps Research)

FIGURES

Figure 1: Fentanyl Induces Cardiorespiratory Depression and nTS neuronal activation in a dose-dependent manner

A. Schematic showing the timeline for OIRD collar experiments. Male and Female SD rats received jugular catheterization surgery. The following week, conscious rats in their home cage received intravenous administration of either saline (n=7, triangles) or fentanyl (2 µg/kg: (n=10, squares); 20 µg/kg, (n=13, circles); 50 µg/kg, (n=7, diamonds). Cardiorespiratory parameters were measured for 3 minutes prior to IV administration and up to 60 minutes after saline or fentanyl. At the conclusion of the experiment, brains were collected from a subset of animals in each group and processed for Fos- and TH- immunoreactivity. Time course data showing oxygen saturation (**B**), heart rate (**C**) and respiratory rate (**D**) data measurements collected in all groups before and after saline or fentanyl. Changes in oxygen saturation were assessed as nadir (lowest value; **E,G,I**) and time to recover to 90% of baseline (pre-saline/fentanyl; **F,H,J**) administration. There was a dose-dependent decrease in the nadir for oxygen saturation (One-way ANOVA, $F_{(3,34)}$ = 157.4, $p < 0.0001$; Tukey's post-hoc ****$p < 0.0001$), heart rate (One-way ANOVA F$_{(3, 34)}$ = 64.96, $p < 0.0001$; Tukey's post-hoc ****$p < 0.0001$) and respiratory rate (One-way ANOVA, F$_{(3, 34)}$ = 43.94, $p < 0.0001$; Tukey's post-hoc ****$p < 0.0001$; *$p < 0.05$). Time to recover to 90% baseline Oxygen Saturation (One-way ANOVA, F$_{(3, 34)}$ = 25.16, $p < 0.0001$; Tukey's post-hoc ****$p < 0.0001$, **$p < 0.01$, *$p < 0.05$) heart rate (One-way ANOVA, $F_{(3, 34)}$ = 23.75, $p < 0.0001$; Tukey's post-hoc ****$p < 0.0001$, ***$p < 0.001$ **$p < 0.01$) and respiratory rate (One-way ANOVA, $F_{(3, 34)}$ = 12.61, $p < 0.0001$; Tukey's post-hoc ****$p < 0.0001$, *$p < 0.05$). **K.** Merged photomicrographs of a coronal brainstem section showing representative Fos- and TH-immunoreactivity in the nTS of rats that received intravenous saline or fentanyl. Scale bar = 200 µm. **L.** Mean data show that 20 and 50 µg/kg fentanyl induced a significantly greater number of Fos-IR cells in the nTS compared to saline and 2 µg/kg fentanyl. (One-way ANOVA, $F_{(3,25)}$ = 37.87, $p < 0.0001$; Tukey's post hoc ****$p < 0.0001$, **$p < 0.001$, *$p < 0.05$) **M.** Mean data showing the percentage of TH-IR nTS neurons that displayed Fos-IR. Rats that were given 2 µg/kg fentanyl displayed a significantly higher increase in activated TH-IR neurons compared to saline-treated rats. 50 µg/kg fentanyl induced a significantly higher percentage of activation in TH cells compared to all other groups. (One-way ANOVA, $F_{(3,25)}$ = 30.99, $p < 0.0001$; Tukey's post-hoc ****$p < 0.0001$, ***$p < 0.001$; **$p < 0.01$)

Figure 2. NLXM Attenuates Respiratory Depression and Limits nTS Neuronal Activation Induced by 20 µg/kg Fentanyl: A. Schematic showing the timeline for OIRD collar experiments. Male and Female SD rats received jugular catheterization surgery. The following week, conscious rats in their home cage received intravenous administration (pretreatment) of either saline (1 ml/kg; n=6), NLX (1 mg/kg; n=7) or NLXM (1 mg/kg; n=13). Two minutes later, all rats received intravenous fentanyl (20 µg/kg). Cardiorespiratory parameters were measured up to 60 minutes fentanyl. At the conclusion of the experiment, brains were collected from a subset of animals in each group and processed for Fos- and TH- immunoreactivity. Time course data showing oxygen saturation (**B**), heart rate (**C**) and respiratory rate (**D**) data measurements collected in all groups before and after saline/OR-X pretreatment and after fentanyl. Changes in oxygen saturation were assessed as nadir (lowest value; **E,G,I**) and time to recover to 90% of baseline (pre-saline/OR-X; **F,H,J**) administration. Mean nadir values were assessed in saline- NLX- and NLXM-pretreated rats (Oxygen Saturation: One-way ANOVA, $F_{(2,23)}$ = 171.7, p <0.0001;Tukey's post-hoc test, **** p<0.0001; Heart Rate: One-way ANOVA, $F_{(2,23)}$ = 37.43, p <0.0001;Tukey's post-hoc test, **** p<0.0001; Respiratory Rate: One-way ANOVA, $F_{(2,23)}$ = 15.93, p <0.0001;Tukey's post-hoc test, ****p<0.0001, ***p<0.05). Time to reach 90% baseline in saline-, NLX- and NLXM-pretreated rats. Mean time to reach 90% baseline was assessed for all parameters. Oxygen Saturation: (One-way ANOVA, $F_{(2,23)}$ = 104.6, p <0.0001; Tukey's post-hoc test ****p<0.0001; Heart rate: (One-way ANOVA, $F_{(2,23)}$ = 32.47, p <0.0001; Tukey's post-hoc test ****p<0.0001); Respiratory rate: (One-way ANOVA, $F_{(2,23)}$ = 36.43, p <0.0001; Tukey's post-hoc test). **K.** Merged photomicrographs of a coronal brainstem section showing representative Fos- and TH- immunoreactivity in the nTS of rats that received fentanyl with or without saline or NLX/NLXM pretreatment. Scale bar = 200 µm. Fos-IR and the percentage of TH-IR neurons expressing Fos-IR was evaluated in 3 nTS sections per rat. **L.** Mean data show no significant difference in the number of Fos-IR cells between saline- and NLXM-pretreated rats. NLX-pretreated rats displayed the lowest degree of Fos-IR in the nTS. Number Fos-IR cells: One-way ANOVA, $F_{(2,18)}$ = 30.52, p <0.0001; Tukey's post-hoc p<0.0001. **M.** Mean data of the percentage of TH-IR cells expressing Fos-IR. The percentage of Fos-IR TH-IR cells was similar between saline- and NLXM-pretreated rats. NLX-pretreated rats displayed the lowest percentage of Fos-IR in TH-IR cells. Fos+TH/TH: (One-way ANOVA, $F_{(2,18)}$ = 21.33, p <0.0001; Tukey's post-hoc **** p<0.0001; *** p<0.05).

A Jugular Vein Catheterization — Fentanyl (20 µg/kg; iv) — **OIRD Reversal** Saline or OR-X — **2 Hours Later** Brains collected for Immunohistochemistry — Assessment of Neuronal Activation

Bregma -14.52 mm
Bregma -14.40 mm
Bregma -14.28 mm

B Oxygen Saturation
C Heart Rate
D Respiratory Rate

Fentanyl + Saline
Fentanyl + NLXM (1 mg/kg)
Fentanyl + NLX (1 mg/kg)

E Nadir (% O2 Sat)
F Min to reach 90%BL
G Nadir (beats/min)
H Min to reach 90% BL
I Nadir (breaths/min)
J Min to reach 90% BL

K Fentanyl + Saline (1ml/kg) — Fentanyl + NLXM (1 mg/kg) — Fentanyl + NLX (1mg/kg)

L #Fos-IR nTS neurons
M %TH-IRnTS cells that are Fos-IR

Figure 3 NLXM Reversal of 20 μg/kg Fentanyl-Induced Respiratory Depression is Comparable to NLX. A. Schematic showing the timeline for OIRD reversal experiments. Male and Female SD rats received jugular catheterization surgery. The following week, conscious rats in their home cage received intravenous fentanyl (20 μg/kg). Four minutes later, rats received intravenous administration (reversal) of either saline (1 ml/kg; n=6), NLX (1 mg/kg; n=7) or NLXM (1 mg/kg; n=9). Cardiorespiratory parameters were measured up to 60 minutes after fentanyl. At the conclusion of the experiment, brains were collected from a subset of animals in each group and processed for Fos- and TH- immunoreactivity. Time course data showing oxygen saturation (**B**), heart rate (**C**) and respiratory rate (**D**) data measurements collected in all groups before and after fentanyl and saline/OR-X administration. Changes in oxygen saturation were assessed as nadir (lowest value; **E,G,I**) and time to recover to 90% of baseline (pre-saline/OR-X; **F,H,J**) administration. Mean nadir values were assessed in saline- NLX- and NLXM-pretreated rats (Oxygen Saturation: One-way ANOVA, $F_{(2,19)}$ = 0.2854, p=0.7549; Heart Rate: One-way ANOVA, $F_{(2,19)}$ = 1.076, p=0.3609; Respiratory Rate: One-way ANOVA, $F_{(2,19)}$ = 5.823, p=0.0107. Time to reach 90% baseline in saline-, NLX- and NLXM-pretreated rats. Mean time to reach 90% baseline was assessed for all parameters. Oxygen Saturation: One-way ANOVA, $F_{(2,19)}$ = 34.38, p <0.0001;Tukey's post-hoc test, **** p<0.0001; Heart Rate: One-way ANOVA, $F_{(2,19)}$ = 24.50, p <0.0001;Tukey's post-hoc test, **** p<0.0001; Respiratory Rate: One-way ANOVA, $F_{(2,19)}$ = 14.68, p=0.0001;Tukey's post-hoc test, ***p<0.05). **K.** Merged photomicrographs of a coronal brainstem section showing representative Fos- and TH- immunoreactivity in the nTS of rats that received fentanyl followed by saline or NLX/NLXM. Scale bar = 200 μm. **L.** Mean data of the number of Fos-IR cells evaluated in 3 nTS sections per rat Number Fos-IR cells: One-way ANOVA, $F_{(2,15)}$ = 0.07269, p=0.9302. **M.** Mean data of the percentage of TH-IR cells that expressing Fos-IR: One-way ANOVA, $F_{(2,16)}$ = 3.507, p=0.0563. There were no significant differences in the number of Fos-IR cells or the percentage of TH-IR cells expressing Fos-IR between groups.

Figure 4: NLXM reversal of OIRD is not aversive. A. Schematic of the Conditioned Place

Aversion (CPA) experimental design for rats given fentanyl followed by saline or OR-X.

Catheterized rats underwent a pretest (Day 1). On the test day (Day 2), rats received fentanyl

(50 µg/kg) in their home cages. Four minutes later, rats received intravenous saline, NLX or

NLXM in the paired side of the box. During the post-test (Day 3), rats were placed in the box

and allowed to explore both compartments. **B.** Track plot data showing movement in both sides

of the CPA box during the Pre-test (day 1) and Post-test (Day 3). For each condition, the Pre-

and Post-test data is from the same animal. In these representative examples, the saline or OR-

X paired side is shown on the right side of the box. However, the paired side was balanced

between left and right sides of the box for all groups. **C.** Mean data with individual plots showing

changes in the time spent on the saline- and OR-X paired side. **D.** The CPA score (calculated

as time spent in the saline or OR-X paired compartment during posttest minus the time spent in

the same compartment during the pretest. For CPA scores, one-sample t-test from hypothetical

value (no aversion). There was a significant decrease in the CPA score for NLX treated rats

(t=3.706 (df=4, p<0.05). 1 mg NLXM: t=2.340, df=6 p=0.578; 5 mg/kg NLXM: t=2.225, df=7,

p=0.0614. Saline, t=0.6113 (df=5, p=0.5677). **E**. Schematic of the Conditioned Place Aversion

(CPA) experimental design for saline-pretreated animals in combination with saline or OR-X. **F.**

Track plot data showing movement in both sides of the CPA box during the Pre-test (day 1) and

Post-test (Day 3). For each condition, the Pre- and Post-test data is from the same animal. In

these representative examples, the saline or OR-X paired side is shown on the right side of the

box. However, the paired side was balanced between left and right sides of the box for all

groups. **G.** Mean data with individual plots showing changes in the time spent on the saline- and

OR-X paired side. There was no significant difference between the time spent in NLX or NLXM

paired side in these animals **H.** The CPA score (calculated as time spent in the saline or OR-X

paired compartment during posttest minus the time spent in the same compartment during the

pretest. For CPA scores, one-sample t-test from hypothetical value (no aversion). There was no

significant difference between NLX or NLXM treated animals. NLX: t=0.2812, df=7, p=0.7867.

NLXM, t=0.005032, d=6=. P=0.9961.

FIGURE 5: Fentanyl-induced nTS activity is mediated by peripheral MORs.

A. Experimental timeline. Male and female rats underwent jugular catheterization surgery,

followed by bilateral nanoinjection of jGCaMP8m into the nTS and then allowed 3-5 weeks for

viral expression. On the test day, rats were anesthetized and placed in a stereotax. The nTS

was exposed and a fiber was lowered into the nTS. **B.** Representative example of eYFP

expression and fiber placement in the nTS. Scale bar = 200 μm. **C.** Spread of overlay of

individual animal viral expression across the caudal-rostral extent of the nTS. **D.** nTS cell

response (mean z-score) aligned to 20 μg/kg fentanyl administration (red arrow) at Baseline (-

30-0 seconds), early response (0-30 seconds) and late response (30-60 seconds) in rats that

received 20 μg/kg fentanyl only (red trace) or 5 mg/kg NLXM followed by 20 μg/kg fentanyl

(green trace). **E.** Area under the curve (AUC) was calculated during baseline (-30 to 0 seconds),

Early Response (0 to 30 seconds) or Late Response (30 to 60 seconds). For AUC graphs, one-

sample t-test from hypothetical zero value. In rats that received fentanyl only, there was a

significant increase in the AUC during the Early Phase (t=6.747, df=4 ** p=0.0025) followed by a

significant decrease in the Late Phase (t=14.35, df=4 *** p=0.0001). In rats that received 5

mg/kg NLXM prior to 20 μg/kg fentanyl, no differences in AUC were observed at any time point

(Baseline; t=0.2916, df=6 p=0.7804; Early Phase; t=1.760, df=6 p=0.1288; 30-60 seconds;

t=1.002, df=6, p=0.3548). **F.** nTS cell response (mean z-score) aligned to MOR antagonists

(black arrow). Rats received iv 20 μg/kg fentanyl (Post-Fentanyl phase, -30 to 0 seconds) prior

to MOR antagonist infusion of NLX (1 mg/kg, blue trace) or NLXM (5 mg/kg, green trace). Data

during Early Recovery (0 to 30 seconds) and Late Recovery (30 to 60 seconds) are shown. **G.**

Mean data showing AUC analyses for both groups. For all graphs, one-sample t-test from

hypothetical zero value. NLXM-treated rats had a significant decrease in activity from Post-

Fentanyl baseline prior to NLXM (t=4.799, df=2 * p=0.0408) and during the Early Recovery

phase (t=6.683, df=2 p=0.0217), and exhibited enhanced activity above baseline during the Late

Recovery phase (t=6.716, df=2, * p=0.0215). NLX-treated rats displayed a similar decrease

below baseline prior to NLX. No difference from baseline was observed during the Early

Recovery phase (t=0.01238, df=4 p=0.9907), and a significant increase above baseline was

observed during the Late Recovery phase (t=3.318, df=4, * p=0.0215).

Figure 6: NLXM does not cross the blood brain barrier. **A**. Timeline of Biodistribution

experiments. Male and female SD rats received intravenous jugular catheterization surgery. The

following week, rats were anesthetized and received intravenous NLX (1 mg/kg) or NLXM (1 or

5 mg/kg). Rats were sacrificed 2- and 10- minutes after IV admin. Plasma was extracted from

trunk blood and intact brains were removed and rapidly frozen with liquid nitrogen. **B**. Amount of

NLX detected in plasma and brain samples at 2 and 10 minutes after intravenous NLX (1

mg/kg). One-way ANOVA, $F_{(3,21)}$ = 66.05, p <0.0001; Tukey's post-hoc test, **** p<0.0001; **

p<0.01. **C**. Amount of NLXM detected in plasma and brain samples at 2 and 10 minutes after

intravenous NLXM (1 mg/kg). One-way ANOVA, $F_{(3,21)}$ = 125.6, p <0.0001; Tukey's post-hoc

test, **** p<0.0001; * p<0.05. **D**. Amount of NLXM detected in plasma and brain samples at 2

and 10 minutes after intravenous NLXM (5 mg/kg). One-way ANOVA, $F_{(3,18)}$ = 144.0, p <0.0001;

Tukey's post-hoc test, **** p<0.0001; *** p<0.001.